Nighttime sleep promotes mechanosensory habituation in *Drosophila*

Simon A. Lowe[1*], Abigail D. Wilson[1], Ko-Fan Chen[2], and James E.C. Jepson[1*]

[1]Department of Epilepsy, UCL Queen Square Institute of Neurology, London, UK

[2]Division of Genetics and Genome Biology, University of Leicester, UK

*To whom correspondence should be addressed: s.lowe@ucl.ac.uk, j.jepson@ucl.ac.uk

ABSTRACT

Sleep is a highly conserved behaviour that facilitates episodic, procedural, and/or

associative modes of memory across species [1-3], a process hypothesised to involve

structural downscaling of synapses potentiated during waking episodes [4-6]. Habituation, an

ancient form of non-associative memory in which repetition of a stimulus leads to diminishing

behavioural responses, is considered a prerequisite for such complex forms of learning [7,8].

Yet whether sleep also influences the capacity of organisms to habituate is poorly

understood. Here we examine this question using the fruit fly, *Drosophila*. We describe an

automated system that yields long-term analogue measurements of habituation to

mechanosensory stimuli in adult flies. Using this platform, we find that *Drosophila* lacking

neuronal calcium sensor Neurocalcin – which exhibit reduced sleep specifically during the

night [9] – display impaired mechanosensory habituation during the day. Mimicking a method

used to treat insomnia [10], we show that compressing nighttime duration both restores

consolidated night sleep and renormalises habituation in *Neurocalcin* mutants. Conversely,

in wild type flies without sleep deficits, decreasing night length reduces total night sleep and

disrupts daytime habituation. In wild type flies, exposure to longer nights – which increases

overall night sleep – enhances synaptic downscaling in sensory circuits. Strikingly, this effect

is abolished in *Neurocalcin* mutants, which do not exhibit increased sleep under these

conditions. Collectively, our findings reveal a role for nighttime sleep in facilitating habituation

and suggest that this may occur by promoting synaptic downscaling.

RESULTS AND DISCUSSION

Induction of mechanosensory habituation in adult *Drosophila*

Numerous assays have been developed to induce habituation of adult *Drosophila* to visual, olfactory, and gustatory, stimuli [11-14]. However, while recently examined in *Drosophila* larvae [15], habituation to mechanical stimuli has been comparatively understudied in adult fruit flies. More broadly, whether sleep quality modulates the ability of adult flies (or other organisms) to habituate to environmental stimuli remains an open question.

We therefore sought to investigate potential links between sleep and habituation to mechanosensory stimuli in adult *Drosophila*. The synaptic homeostasis hypothesis stipulates that sleep facilitates downscaling of synapses potentiated during waking experience, thus enabling memory by preventing neuronal hyper-metabolism [16] and enhancing signal-to-noise ratios across circuits [5]. Ongoing locomotor activity occurring during wakefulness could induce an opposite change, potentiating synaptic connections in circuits mediating movement [17,18] and mechanosensory feedback [19-21]. We therefore hypothesised that enhancing this potentiation by limiting sleep (thus prolonging locomotion) might reduce sleep-induced synaptic downscaling and impair the ability of habituation-promoting inhibitory mechanisms [22,23] to suppress motor circuit activity in response to repeated mechanical stimuli.

To test this hypothesis, we developed a simple, high-throughput habituation assay using the DART (*Drosophila* ARousal Tracking) system (Fig. 1A) [24], which combines continuous video-tracking of individual flies with mechanical stimuli delivered via platform-associated motors. DART can be programmed to deliver consistent mechanosensory stimuli that elicit robust, transient, locomotor responses in control flies (Figure S1A), the magnitude and dynamics of which can be quantified. We repurposed the DART system to test for mechanosensory habituation in adult flies by delivering multiple identical stimuli at intervals of 10 min (Fig. 1B). Flies responded to repeated mechanical stimuli with locomotor responses that gradually attenuated in magnitude (Fig. 1C), quantified as the mean peak speed in the 1 min following each stimulus (Fig. 1D).

If this decrement represents true habituation rather than generalised sensory

adaptation, fatigue, or structural damage, it should exhibit key properties that define

habituation: spontaneous recovery, potentiation of habituation over repeated bouts, and

stimulus-specificity [7,8]. We sought to validate the assay by testing for these characteristics.

After attenuation of locomotion following 20 mechanical stimuli, locomotor speeds

exhibited spontaneous recovery over time in response to single mechanical stimuli, with ~

80% recovery occurring following 24 h (Figure S1B-F). We found that the rate of

mechanosensory habituation could be significantly enhanced by applying a habituation-

inducing paradigm of 20 identical mechanical stimuli on three consecutive days (Fig. 1E, F).

Finally, after 20 consecutive mechanical stimuli, we delivered a distinct 'lights-off' visual

stimulus expected to similarly induce a startle response (Fig. 1G). Despite the robust

decrease in locomotor response to repeated mechanical stimuli (Fig. 1G, H), locomotor

responses to the visual stimulus did not differ in magnitude between naïve and mechanically

habituated flies (Fig. 1G, I). Hence, progressive reductions in locomotor responses in our

paradigm are consistent with well-defined characteristics of habituation [7,8].

A number of platforms analogous to DART can track locomotor activity of populations

of flies using video- or infrared beam-based analysis, and can deliver mechanosensory

stimuli or be modified to do so. These include the *Drosophila* Activity Monitor [25], Ethoscope,

[26] and Zantiks [27] systems. We designed a similar habituation protocol using the Zantiks

automated-tracking system (Figure S1F; see Methods). The Zantiks system could detect and

quantify robust locomotor responses of adult flies to single motor-driven stimuli (Figure

S1G), as well as a progressive decline in locomotor response to repeated, identical

mechanosensory stimuli that was qualitatively similar to that observed using the DART

system (Figure S1H, I and Fig. 1D). These data provide cross-platform validation of our

findings using the DART.

Neurocalcin promotes mechanosensory habituation in *Drosophila*

Similarly to humans, deep sleep in *Drosophila* primarily occurs during the night [28]. We

therefore sought to test whether night sleep influences mechanosensory habituation induced

during the day. Mechanically disrupting sleep provides a potential means to do so. However,

homeostatic rebound sleep induced by sleep deprivation [29] could potentially cause

confounding changes in stimulus-induced movement during the day. Hence, we utilised a

genetic manipulation that selectively disrupts night sleep whilst leaving daytime sleep intact:

loss of the highly conserved neuronal calcium sensor Neurocalcin [9].

In contrast to mutations such as *shaker*, *sleepless*, *fumin*, and *insomniac* – which

disrupt both day and night sleep in *Drosophila* [30-34] – loss of Neurocalcin specifically impairs

sleep during the night in oscillating light: dark conditions [9], providing a means to examine the

relationship between night sleep and daytime habituation. Indeed, using our paradigm to

measure the rate and amplitude of mechanosensory habituation in control flies and flies

homozygous for a *Neurocalcin* null allele (*Nca*[KO1]) [9], we found that loss of Neurocalcin

significantly reduced the degree of mechanosensory habituation, an effect that was

particularly prominent after ~ 10 mechanical stimuli (Fig. 2A, B). We observed similarly

reduced mechanosensory habituation in flies homozygous for an independent *Neurocalcin*

null allele (*Nca*[KO2]) (Figure S2A-B), confirming the above genotype-phenotype correlation.

Neurocalcin could potentially modulate mechanosensory habituation by enhancing

sensitivity to mechanical stimuli. We therefore tested locomotor responses in control and

Neurocalcin knockout flies to mechanical stimulus strengths ranging from 1% to 100% of the

strength used in the habituation protocol. *Neurocalcin* knockout flies showed statistically

similar amplitudes of locomotor responses to isogenic controls at 6 out of 7 stimulus

strengths tested, and a slightly enhanced response to stimuli 10% of the strength used in the

habituation protocol (Figure S2C). These data therefore did not support a general increase in

mechanosensory sensitivity in *Neurocalcin* knockout flies, and confirmed that the habituation

protocol utilised a stimulus strength that was beyond the plateau at which there was no

difference between control and *Neurocalcin* knockout flies.

Since our habituation protocol yielded long-term quantitative recordings of individual

flies, we were able to examine whether the reduction in habituation in *Neurocalcin* knockout

flies was due to a change in number of flies responding to mechanical stimuli, or a decrease

in the magnitude of the individual flies' responses. To separate response to a stimulus from

background noise, we defined a threshold for a "response" to each stimulus as $\geq$ 10% of the

response to the mean naïve response. We found that for both control and *Neurocalcin*

knockout flies, the attenuation in response came entirely from the proportion of flies

responding (Fig. 2C). When only "responding" flies were included in analysis, there was no

decrease in the amplitude of locomotor responses across repetitive stimuli, nor any

difference between genotype (p = 0.46, main effect of Genotype & of Stimulus #, 2-way

ANOVA) (Fig. 2D). The all-or-nothing nature of locomotor responses in our paradigm

supports the premise that the decrease in mean amplitude across repeated stimuli

represents a learning event i.e an altered "decision" of whether to respond or not to a given

stimulus, as opposed to a motor defect affecting the ability to sustain robust reactions over

multiple stimuli.

Short nights restore consolidated sleep and mechanosensory habituation in

***Neurocalcin* knockout flies**

Hippocalcin, the mammalian homologue of Neurocalcin, promotes synaptic plasticity and

memory in mice [35-39], and cognitive development in humans [40,41]. Thus, we sought to test

whether Neurocalcin facilitates habituation through its effect on night sleep, or via sleep-

independent processes potentially linked to memory.

We previously documented that night sleep loss in *Neurocalcin* mutant flies was

particularly prominent under long night, winter-like, conditions [9], suggesting that night length

can modulate sleep loss in *Neurocalcin* mutant flies. Interestingly, increasing sleep drive by

restricting the period of consolidated night sleep is utilised in cognitive behavioural therapy

for insomnia, and has similarly been shown to rescue sleep loss in a variety of *Drosophila*

sleep mutants [10]. We therefore examined sleep architecture in control and *Neurocalcin*

knockout flies under short and long night conditions using a multibeam monitoring system

that facilitates high-resolution quantification of sleep in *Drosophila* [42] (Figure S3A).

Control and *Neurocalcin* knockout flies were raised under standard 12 h light: 12 h

dark conditions then, after 3-5 days of adulthood, were shifted to three days of long (8 h

light: 16 h dark) or short (16 h light: 8 h dark) night conditions, with sleep quantified on the

third day of each light: dark cycle. Consistent with prior results [9], we confirmed reduced

nighttime but not daytime sleep in *Neurocalcin* knockout flies under long night conditions

(Fig. 3A, C and Figure S3B-C), and found that this decrease was associated with

fragmentated sleep, characterised by a significant increase in the number of sleep bouts and

a concomitant reduction in sleep bout duration (Fig. 3D, E). In contrast, under short night

conditions, overall night sleep (Fig. 3B, C) and the degree of sleep consolidation (Fig. 3D, E)

were not significantly different between *Neurocalcin* knockout flies and controls. Day sleep

was also slightly increased in *Neurocalcin* knockout flies under short night conditions, likely

due to reduced locomotor activity towards lights-off (Fig. 3B, Figure S3B-C). Thus, changing

night length provides a means to tune night sleep quality on-demand in *Neurocalcin*

knockout flies, allowing us to disentangle sleep-dependent versus -independent functions of

Neurocalcin.

Taking advantage of this finding, we quantified mechanosensory habituation in

control and *Neurocalcin* knockout flies entrained to short or long night conditions (Fig. 3F).

Strikingly, while mechanosensory habituation was reduced in *Neurocalcin* knockout flies

compared to controls under long night conditions (Fig. 3F), this difference was abolished

under short night conditions (Fig. 3G). These data strongly suggest that Neurocalcin

regulates habituation directly through its effect on night sleep.

Loss of Neurocalcin decorrelates night length and synaptic protein expression

Intriguingly, within-genotype comparisons revealed that control flies exposed to short nights

exhibited a reduced rate of mechanosensory habituation compared to control flies exposed

to long nights (Figure S4A), indicating that limiting night sleep (Fig. 3C) disrupts

mechanosensory habituation during the following day. Conversely, *Neurocalcin* knockout

flies, which exhibit improved night sleep in short versus long nights (Fig. 3C), exhibited the

opposite correlation: an enhanced rate of habituation in short versus long night conditions

(Figure S4B).

How might night duration modulate mechanosensory habituation occurring after

nighttime sleep? Prior studies have shown that sleep deprivation (SD) in adult flies impairs

synaptic downscaling, resulting in increased levels of the presynaptic active zone protein

Bruchpilot [43] that can be visualised through immuno-fluorescent microscopy. Prolonged (16

h) SD was initially shown to increase presynaptic Bruchpilot expression by ~ 75% across

Drosophila brain neuropil regions [44]. More recently, 12 h night-specific SD was shown to

modulate Bruchpilot to a more limited degree, enhancing Bruchpilot expression in cholinergic

neurons by ~ 25%, and in GABAergic neurons by ~ 10% [45]. Loss of the *Drosophila*

Insomniac protein, which reduces overall day and night sleep by ~ 65% in male flies [32], also

leads to a 40-50% increase in presynaptic Bruchpilot expression [46].

Extending our analysis in the context of the synaptic homeostasis hypothesis [5] and

the above studies, we investigated whether altering night length was sufficient to modify

presynaptic Bruchpilot expression in control flies, and if so, whether loss of Neurocalcin

affected this relationship. We immuno-labelled synaptic Bruchpilot in isolated brains of

control and *Neurocalcin* knockout flies following three days of entrainment to short or long

night conditions, then quantified fluorescent intensity of synaptic Bruchpilot through confocal

microscopy, normalising fluorescence to a control DNA marker (DAPI, see Methods).

Consistent with the relative subtlety of this manipulation compared to mechanical

sleep deprivation [44,45] or loss of Insomniac [46], we observed a trend towards a small increase

(~ 16%) in Bruchpilot throughout the brain neuropil region of control flies subjected to short

versus long nights, which was non-significant (Figure S4C). However, in the neuropil-rich

antennal lobe regions, which process olfactory information in insects, we observed a

significant increase (~ 40%) in synaptic Bruchpilot levels in short versus long night

conditions (Fig. 4A, B).. Notably, we did not observe a similar increase in antennal lobe

Bruchpilot expression in *Neurocalcin* mutants exposed to short versus long nights (Fig. 4C,

D), nor in whole-brain Bruchpilot immuno-fluorescence (Figure S4D), suggesting that the

impaired sleep quality of *Neurocalcin* exposed to long nights (Fig. 3A, C) blocks enhanced

synaptic downscaling that occurs in wild types flies under this condition.

Collectively, the above data support a model in which reduced night sleep in control

flies exposed to short nights perturbs synaptic downscaling in specific neuropil domains

compared to flies experiencing long nights. We hypothesise that such domains may include

circuits relevant to mechano-sensation and movement, where impaired downscaling could

limit the ability of habituation-promoting mechanisms (such as enhanced synaptic

transmission from local inhibitory neurons [23]) to dampen functional connectivity between

mechanosensory neurons and downstream motoneuron outputs. The improvement of sleep

quality in *Neurocalcin* mutants exposed to short nights (Fig. 3A, C-E) breaks the normal

correlation between night length and synaptic Bruchpilot expression (Fig. 4), leading to

improved rather than impaired mechanosensory habituation in short compared to long night

conditions (Figure S4).

Further studies will be required to test this model – in particular, by identifying

mechanosensory and/or pre-motor circuits in which synaptic output is modulated by

habituation-promoting mechanisms. Intriguingly, recent work has shown that continuous or

repetitive mechanosensory stimulation, akin to the gentle rocking of a human baby,

increases daytime sleep in *Drosophila* through a habituation-like mechanism [47]. In concert

with our findings, these data suggest an intertwined and potentially bi-directional relationship

between mechano-sensation, sleep, and habituation, across both day-night and seasonal

cycles. Finally, we note that sleep and habituation are dysregulated in a variety of human

neurodevelopmental disorders [48-52] and corresponding animal models [49,53-57]. Since

habituation promotes attention to environmental novelties that form the substrates of

memory-related coincidence detection mechanisms [58], our work has implications for

understanding the common co-morbidity of sleep, habituation, and memory defects in these

clinically important conditions.

ACKNOWLEDGEMENTS

We thank Dr. Gabriel Aughey for comments on the manuscript, members of BKFLab for

technical assistance with the DART system, and members of the Zantiks team for technical

assistance with the Zantiks MWP system. This study was funded by a MRC Senior Non-

Clinical Fellowship (MR/V03118X/1) and a BBSRC Research Grant (BB/R02281X/1) to

J.E.C.J; a BBSRC Research Grant (BB/X00094X/1) to S.L and J.E.C.J; a Wellcome Early

Career Fellowship to A.D.W (308298/Z/23/Z); and a BBSRC Research Grant

(BB/W014939/1), and a Startup and CRE fund from the Division of Genetics and Genome

Biology of University of Leicester (RM33077K), to K-F.C.

AUTHOR CONTRIBUTIONS

S.A.L – Conceptualization, Methodology, Investigation, Writing - Original Draft, Writing –

Review & Editing, Visualization, Funding acquisition; A.D.W – Methodology, Investigation,

Writing - Original Draft, Writing – Review & Editing, Visualization. K-F.C – Investigation, Writing

– Review & Editing. J.E.C.J – Conceptualization, Investigation, Visualization, Writing - Original

Draft, Writing – Review & Editing, Supervision, Project administration, Funding acquisition.

DECLARATION OF INTERESTS

The authors declare no competing interests.

MATERIALS AND METHODS

Experimental model and subject details

Experimental models were fruit flies of the species *Drosophila melanogaster*. Genotypes

used were iso31 (an isogenized strain of w^{1118} flies and a kind gift from Prof. Kyunghee Koh,

Thomas Jefferson University), w^{1118};+; Nca^{KO1}/TM2, and w^{1118};+; Nca^{KO2}/TM2 [9] To control for

genetic background, these lines were outcrossed into the iso31 background for 5

generations. Flies were raised and maintained on standard fly food under 12 h: 12 h light-

dark cycles (12L: 12D) at a constant 25°C. Experiments took place at 25°C, and unless

otherwise indicated, in 12L: 12D conditions. Experiments were conducted on male flies.

Habituation protocol

Drosophila ARousal Tracking (DART) analysis was performed as described previously [1].

Briefly, five- to seven-day old male flies were individually loaded into glass behavioural tubes

(Trikinetics inc., MA, USA) containing 4% sucrose and 2% agar, and left for 24 h to

acclimatise in the indicated L:D conditions. The mechanical stimulus consisted of a

mechanical vibration in the form of a train of five 200 ms pulses separated by 800 ms

intervals, set at the DART system's maximum intensity, which elicited a robust movement

response. The habituation protocol consisted of 20 mechanical stimuli delivered at ten-

minute intervals, beginning at Zeitgeber time 3 (ZT3). Videos were taken using a USB

webcam (Logitech). Speed (mm / s) was analysed by the DART system from absolute

position, and binned in one min intervals. Flies were excluded from analysis if they exceeded

an average of 1 mm / s speed in the five one-minute bins preceding the first stimulus.

Response to the stimulus was defined as the mean speed (mm / s) in the one-minute bin

following the stimulus. Speed was normalised within genotype to the mean of the response

to the first stimulus. To derive the % of flies responding to stimuli, a "response" was

distinguished from background movement by the (arbitrary) threshold of a change in mean

speed between the one-minute bins preceding and following a stimulus of $\geq$10% of the mean

response to the first stimulus.

A distinct automated *Drosophila* tracking system, the Zantiks MWP unit (Zantiks Ltd.,

Cambridge, UK) was used to generate an equivalent habituation protocol. 5-7 day old male

iso31 control flies were loaded into individual wells filled with 4% sucrose and 2% agar in a

48-well plate, and left to acclimatise for 24 h in a 12 h light: 12 h dark cycle at 25°C.

Drosophila were loaded into the Zantiks unit at ZT0, and the habituation protocol

commenced at ZT3. The habituation protocol again consisted of 20 mechanical stimuli

delivered at 10 min intervals, each stimulus consisting of a single 1 s pulse set at medium

strength, generated by the Zantiks mechanical stimulation stand. Distance travelled is

automatically tracked by the Zantiks system, and total distance travelled was binned in 1 min

intervals. Scripts used for the habituation protocol on the Zantiks MWP are available at

github.com/abigaildwilson/zanscripts.

Sleep analysis

Sleep analysis was performed using the *Drosophila* Activity Monitor 5 system (DAM5H,

Trikinetics inc., MA, USA), which improves on the fidelity of the earlier DAM2 system by

using 15 infrared beams per tube, rather than a single one, to track movement [2]. As above,

three- to five-day old male flies were loaded in behavioural tubes, and were left for two full

days to acclimatise at the relevant light cycle. On the third day the number of beam breaks

was measured using the *Drosophila* Activity Monitor. Sleep was defined using the standard

definition of > 5 mins of inactivity, which has been shown to correlate with postural changes,

increased arousal threshold and altered brain activity patterns [2]. Sleep bout number and

duration was analysed using the open-source software suite phaseR

(https://github.com/abhilashlakshman/phaseR/tree/main) [3].

Immuno-histochemistry and confocal microscopy

Brains were dissected in phosphate-buffered saline (PBS) (Sigma Aldrich) between ZT1-2,

fixed by 20 min room temperature incubation in 4% paraformaldehyde (MP biomedicals)

diluted in PBS + 0.3% Triton-X (Sigma-Aldrich) (PBT), washed three times in PBT, and

blocked for 1 h in 5% normal goat serum in PBT. Brains were incubated for 48 hours at 4°C

in primary antibody (mouse anti-BRP [nc82], Developmental Studies Hybridoma Bank, at a

concentration of 1:200); then washed three times in PBT and incubated overnight at 4°C in

secondary antibody (goat anti-mouse AlexaFluor 647, ThermoFisher Scientific, at a

concentration of 1:500) and counterstain (DAPI, ThermoFisher Scientific, at a concentration

of 1:500). Brains were washed three final times before mounting in SlowFade Gold anti-fade

mountant (ThermoFisher). Images were taken with a Zeiss LSM 980 confocal microscope

with an EC 'Plan-Neofluar' 20x/0.50 M27 air objective, taking z-stacks through the entire

brain with step sizes of 1 µm. Images were visualised and analysed using ImageJ; maximum

intensity z-stack projections were generated, and following background subtraction, regions

of interest (ROIs) were manually drawn and mean pixel intensity calculated. BRP intensity

readings from both the whole central brain and limited to the antennal lobes were normalised

to DAPI readings from the whole brain. Data is pooled from three separate experiments, with

brains within experiments dissected, immuno-stained and imaged in parallel. Values are

normalised to the mean values of the long night (L8: D16) condition within an experiment.

 FIGURE LEGENDS

Fig. 1. Mechanosensory habituation in adult *Drosophila*. (**A**) Schematic illustrating the

Drosophila ARousal Tracking (DART) system. (**B**) Protocol used to induce mechanosensory

habituation. (**C**) Mean speed over time in one-minute bins, and (**D**) mean speed in the bin

following each stimulus, of N = 24 control (iso31) male flies, exposed to 20 successive

mechanical stimuli. Data are normalised to the mean in the bin following the first stimulus,

i.e. the naïve response. (**E-F**) Mean speed of locomotion over time (E) and mean response

in the one-minute bin following each stimulus (F) in N = 25 control adult male flies, in

response to an identical habituation protocol applied on three consecutive days. Data from

Day 1 and Day 3 are shown. Data are normalised to the response to the first stimulus within

each day. Interaction between stimulus number and day: ***p<0.0005, two-way ANOVA. (**G**)

Mean speed of locomotion over time as a "lights-off" stimulus is delivered to naïve flies

(black, N = 24) and immediately following a train of 20 mechanosensory stimuli (blue, N =

17). (**H**) Normalised locomotor responses of control adult male flies to the 1st and 20th

mechanical stimulus shows a clear decline in response to repeated stimulus: N = 17.

***p<0.0005, Mann-Whitney U-test. (**I**) Response to lights-off does not differ between naïve

flies (N = 24) and those habituated to the distinct mechanosensory stimuli (N = 17). p =

0.7187, Mann-Whitney U-test. Error bars: standard error of the mean (SEM).

Fig. 2. *Neurocalcin* knockout flies exhibit impaired mechanosensory habituation. (**A**) Mean

speed of locomotion over time, and (**B**) mean speed in one-minute bin following each

stimulus of control (N = 87) and *Nca* knockout (*Nca*^{KO1}) (N = 58) adult male flies exposed to

20 consecutive mechanosensory stimuli. Data are normalised within each genotype to the

naïve response. *Nca* knockout flies habituate more slowly than controls (interaction between

stimulus number and genotype: p<0.0001, two-way ANOVA). Error bars: SEM. (**C**)

Proportion of flies "responding" decreased over multiple stimuli, with the rate of decrease

slower for *Nca*^{KO1} flies. (**B**) Mean speed of flies "responding" did not decrease in response to

repeated stimuli, nor did it differ between genotype (main effect of stimulus number:

p=0.2407, and of genotype: p=0.3761; interaction between stimulus and genotype:

p=0.4639, two-way ANOVA). Error bars: SEM.

Fig. 3. *Neurocalcin* modulates mechanosensory habituation through a sleep-dependent

mechanism. (**A-B**) Mean sleep levels of adult male flies over 24 h under (A) short day, long

night (L8: D16) conditions (control: N = 32; Nca^{KO1}: N = 32), and (B) long day, short night

(L16: D8) conditions (control: N = 31; Nca^{KO1}: N = 20). (**C-E**) Total minutes of sleep during

subjective night (C), number of individual sleep bouts occurring during the night normalised

to hours of night (D), and mean length of sleep bouts occurring during the night (E). Kruskal

Wallis test with Dunn's multiple comparison test. (**F**) Protocol for exposure to short or long

night conditions prior to habituation assay. (**G-H**) Nca^{KO1} flies habituate more slowly than

controls in (G) long night conditions: N = 60 (control), 66 (Nca^{KO1}), interaction between

stimulus and genotype: p = 0.0472. Performance does not differ between the genotypes in

(H) short night conditions: N = 61 (control), 61 (Nca^{KO1}), interaction between stimulus and

genotype: p = 0.2395. Two-way ANOVA with Dunn's multiple comparison tests. Error bars:

SEM.

Fig. 4. Night sleep duration modulates Bruchpilot immunofluorescence in sensory circuits.

(**A, C**) Representative images of (A) iso31 and (C) Nca^{KO1} whole brains immuno-stained for

anti-BRP (top), with a DAPI counterstain (bottom), in long night (L8: D16) vs short night (L16:

D8) conditions. Scale bars: 100µM. (**B, D**). Mean BRP immunofluorescence within antennal

lobe neuropil regions (two data points per brain), normalised to whole-brain DAPI

immunofluorescence, is increased in short- versus long-night conditions in controls (p =

0.0018, unpaired t-test with Welch's correction), but unchanged in Nca^{KO1} brains (p = 0.81,

Mann-Whitney U-test). Values are normalised to the mean value in long night conditions.

Error bars: SEM.

 SUPPLEMENTARY FIGURE LEGENDS

Supp. Fig. 1. (**A**) Quantification of mean locomotor speed over time in response to a single

mechanical stimulus. Mean speed of N = 11 control (iso31) male flies are shown. (**B-D**)

Mean speed over time as a further single mechanosensory stimulus is delivered following an

interval of (B) 1 h (N = 87), (C) 4 h (N = 36), or (D) 24 h (N = 27). (**E**) Mean speed in the 1

min bin following a single stimulus at the indicated interval after the initial habituation

protocol. Data are normalised to the naïve response to the first stimulus in the 20-stimulus

train (N = 87). Responses progressively return towards the baseline level. (**F**) Schematic of

the Zantiks system. Individual flies are loaded into a 48-well plate containing a solidified agar

and sucrose food surface. Mechanical stimuli are delivered via a motor beneath the plate,

with locomotor responses recorded via videography. (**G**) Mean locomotor responses over

time of N = 24 flies to a single mechanical stimulus (solid line), quantified as distance

travelled in a 1 min interval. (**H-I**) Mean distance travelled per 1 min (**H**; line represents mean

value over time), and mean distance travelled in the 1 min following each mechanical

stimulus (**I**), in N = 48 control (iso31) male flies exposed to 20 successive mechanical stimuli

in the Zantiks system. Error bars in E, I: SEM

Supp. Fig. 2. (**A**) Mean speed of locomotion over time, and (**B**) mean speed in one-minute

bin following each stimulus of control (iso31) (N = 41) and a second independently

generated Nca knockout line (Nca^{KO2}) (N = 30) adult male flies exposed to 20 consecutive

mechanosensory stimuli. Data are normalised within each genotype to the naïve response.

Nca^{KO2} flies habituate more slowly than controls (interaction between stimulus number and

genotype: $p < 0.0001$, two-way ANOVA). Error bars: SEM. (**C**) Mean speed of control and

Nca^{KO1} adult male flies in the one-minute bin following a single mechanosensory stimulus of

varying strength (values are normalised to the strength used in the habituation assay).

Control: N = 22-52; Nca^{KO1}: N = 15-41. *$p < 0.05$, ns: $p > 0.05$, Mann-Whitney U-test.

Supp. Fig. 3. (**A**) Schematic illustrating components of the *Drosophila* Activity Monitor 5

(DAM5) system. Flies are individually housed within glass behavioural tubes. Locomotion is

tracked via breaking of 15 infrared beams across the tubes. Sleep is derived from periods

without locomotion (see Methods). (**B-D**) Total minutes of sleep during subjective day (B),

number of individual sleep bouts during subjective day normalised to hours of day (C), and

mean length of sleep bouts occurring during the day (D), for populations presented in Fig.

3A-B. Kruskal Wallis test with Dunn's multiple comparison test. $*p < 0.05$, $**p < 0.01$, $***p <$

0.001. Error bars: SEM.

Supp. Fig. 4. (**A-B**) Within-genotype comparisons of the effect of light cycle on habituation.

(A) Iso31 flies habituate more quickly in long-night conditions (interaction between genotype

and light cycle: $p = 0.0027$). In contrast, Nca^{KO1} flies (B) habituate more quickly in short-night

conditions (interaction between genotype and light cycle: $p = 0.0494$). (**C-D**) Mean Bruchpilot

(BRP) immunofluorescence within the whole central brain, normalised to DAPI, is not

significantly changed between short- versus long-night conditions in control (C: $p = 0.1648$),

or Nca^{KO1} brains (D: $p = 0.5461$). Unpaired t-test with Welch's correction (C) or Mann

Whitney U-test (D). Values are normalised to the mean value in long night conditions. Error

bars: SEM.

REFERENCES

1. Kanaya, H.J., Park, S., Kim, J.H., Kusumi, J., Krenenou, S., Sawatari, E., Sato, A., Lee, J., Bang, H., Kobayakawa, Y., et al. (2020). A sleep-like state in Hydra unravels conserved sleep mechanisms during the evolutionary development of the central nervous system. Sci Adv *6*. 10.1126/sciadv.abb9415.

2. Lakhiani, R., Shanavas, S., and Melnattur, K. (2023). Comparative biology of sleep in diverse animals. J Exp Biol *226*. 10.1242/jeb.245677.

3. Jaggard, J.B., Wang, G.X., and Mourrain, P. (2021). Non-REM and REM/paradoxical sleep dynamics across phylogeny. Curr Opin Neurobiol *71*, 44–51. 10.1016/j.conb.2021.08.004.

4. Miyamoto, D., Marshall, W., Tononi, G., and Cirelli, C. (2021). Net decrease in spine-surface GluA1-containing AMPA receptors after post-learning sleep in the adult mouse cortex. Nat Commun *12*, 2881. 10.1038/s41467-021-23156-2.

5. Tononi, G., and Cirelli, C. (2014). Sleep and the price of plasticity: from synaptic and cellular homeostasis to memory consolidation and integration. Neuron *81*, 12–34. 10.1016/j.neuron.2013.12.025.

6. de Vivo, L., Bellesi, M., Marshall, W., Bushong, E.A., Ellisman, M.H., Tononi, G., and Cirelli, C. (2017). Ultrastructural evidence for synaptic scaling across the wake/sleep cycle. Science *355*, 507–510. 10.1126/science.aah5982.

7. Thompson, R.F., and Spencer, W.A. (1966). Habituation: a model phenomenon for the study of neuronal substrates of behavior. Psychol Rev *73*, 16–43. 10.1037/h0022681.

8. Rankin, C.H., Abrams, T., Barry, R.J., Bhatnagar, S., Clayton, D.F., Colombo, J., Coppola, G., Geyer, M.A., Glanzman, D.L., Marsland, S., et al. (2009). Habituation revisited: an updated and revised description of the behavioral characteristics of habituation. Neurobiol Learn Mem *92*, 135–138. 10.1016/j.nlm.2008.09.012.

9. Chen, K.F., Lowe, S., Lamaze, A., Kratschmer, P., and Jepson, J. (2019). Neurocalcin regulates nighttime sleep and arousal in Drosophila. Elife *8*. 10.7554/eLife.38114.

10. Belfer, S.J., Bashaw, A.G., Perlis, M.L., and Kayser, M.S. (2021). A Drosophila model of sleep restriction therapy for insomnia. Mol Psychiatry *26*, 492–507. 10.1038/s41380-019-0376-6.

11. Engel, J.E., and Wu, C.F. (2009). Neurogenetic approaches to habituation and dishabituation in Drosophila. Neurobiol Learn Mem *92*, 166–175. 10.1016/j.nlm.2008.08.003.

12. Engel, J.E., and Wu, C.F. (1996). Altered habituation of an identified escape circuit in Drosophila memory mutants. J Neurosci *16*, 3486–3499. 10.1523/JNEUROSCI.16-10-03486.1996.

13. Asztalos, Z., Arora, N., and Tully, T. (2007). Olfactory jump reflex habituation in Drosophila and effects of classical conditioning mutations. J Neurogenet *21*, 1–18. 10.1080/01677060701247508.

14. Duerr, J.S., and Quinn, W.G. (1982). Three Drosophila mutations that block associative learning also affect habituation and sensitization. Proc Natl Acad Sci U S A *79*, 3646–3650. 10.1073/pnas.79.11.3646.

15. Berne, A., Zhang, T., Shomar, J., Ferrer, A.J., Valdes, A., Ohyama, T., and
Klein, M. (2023). Mechanical vibration patterns elicit behavioral transitions and
habituation in crawling Drosophila larvae. Elife *12*. 10.7554/eLife.69205.
16. Feeney, S.P., McCarthy, J.M., Petruconis, C.R., and Tudor, J.C. (2025). Sleep
loss is a metabolic disorder. Sci Signal *18*, eadp9358.
10.1126/scisignal.adp9358.
17. Brezovec, B.E., Berger, A.B., Hao, Y.A., Chen, F., Druckmann, S., and
Clandinin, T.R. (2024). Mapping the neural dynamics of locomotion across the
Drosophila brain. Curr Biol *34*, 710–726 e714. 10.1016/j.cub.2023.12.063.
18. Aimon, S., Cheng, K.Y., Gjorgjieva, J., and Grunwald Kadow, I.C. (2023).
Global change in brain state during spontaneous and forced walk in
Drosophila is composed of combined activity patterns of different neuron
classes. Elife *12*. 10.7554/eLife.85202.
19. Mamiya, A., Gurung, P., and Tuthill, J.C. (2018). Neural Coding of Leg
Proprioception in Drosophila. Neuron *100*, 636–650 e636.
10.1016/j.neuron.2018.09.009.
20. Pratt, B.G., Dallmann, C.J., Chou, G.M., Siwanowicz, I., Walling-Bell, S.,
Cook, A., Sustar, A., Azevedo, A., and Tuthill, J.C. (2026). Proprioceptive limit
detectors contribute to sensorimotor control of the Drosophila leg. Nat
Commun. 10.1038/s41467-026-69333-z.
21. Tuthill, J.C., and Wilson, R.I. (2016). Parallel Transformation of Tactile Signals
in Central Circuits of Drosophila. Cell *164*, 1046–1059.
10.1016/j.cell.2016.01.014.
22. Larkin, A., Karak, S., Priya, R., Das, A., Ayyub, C., Ito, K., Rodrigues, V., and
Ramaswami, M. (2010). Central synaptic mechanisms underlie short-term
olfactory habituation in Drosophila larvae. Learn Mem *17*, 645–653.
10.1101/lm.1839010.
23. Das, S., Sadanandappa, M.K., Dervan, A., Larkin, A., Lee, J.A., Sudhakaran,
I.P., Priya, R., Heidari, R., Holohan, E.E., Pimentel, A., et al. (2011). Plasticity
of local GABAergic interneurons drives olfactory habituation. Proc Natl Acad
Sci U S A *108*, E646–654. 10.1073/pnas.1106411108.
24. Faville, R., Kottler, B., Goodhill, G.J., Shaw, P.J., and van Swinderen, B.
(2015). How deeply does your mutant sleep? Probing arousal to better
understand sleep defects in Drosophila. Sci Rep *5*, 8454. 10.1038/srep08454.
25. Pfeiffenberger, C., Lear, B.C., Keegan, K.P., and Allada, R. (2010). Locomotor
activity level monitoring using the Drosophila Activity Monitoring (DAM)
System. Cold Spring Harb Protoc *2010*, pdb prot5518. 10.1101/pdb.prot5518.
26. Geissmann, Q., Garcia Rodriguez, L., Beckwith, E.J., French, A.S., Jamasb,
A.R., and Gilestro, G.F. (2017). Ethoscopes: An open platform for high-
throughput ethomics. PLoS Biol *15*, e2003026. 10.1371/journal.pbio.2003026.
27. Allen, T., and Budenberg, W.J. (2021). Replicating Light-Off Startle
Responses in Drosophila melanogaster. bioRxiv. 10.1101/2021.02.24.432669.
28. van Alphen, B., Yap, M.H., Kirszenblat, L., Kottler, B., and van Swinderen, B.
(2013). A dynamic deep sleep stage in Drosophila. J Neurosci *33*, 6917–6927.
10.1523/JNEUROSCI.0061-13.2013.
29. Huber, R., Hill, S.L., Holladay, C., Biesiadecki, M., Tononi, G., and Cirelli, C.
(2004). Sleep homeostasis in Drosophila melanogaster. Sleep *27*, 628–639.
10.1093/sleep/27.4.628.

30.　Kume, K., Kume, S., Park, S.K., Hirsh, J., and Jackson, F.R. (2005).
Dopamine is a regulator of arousal in the fruit fly. J Neurosci *25*, 7377–7384.
10.1523/JNEUROSCI.2048-05.2005.

31.　Cirelli, C., Bushey, D., Hill, S., Huber, R., Kreber, R., Ganetzky, B., and
Tononi, G. (2005). Reduced sleep in Drosophila Shaker mutants. Nature *434*,
1087–1092. 10.1038/nature03486.

32.　Stavropoulos, N., and Young, M.W. (2011). insomniac and Cullin-3 regulate
sleep and wakefulness in Drosophila. Neuron *72*, 964–976.
10.1016/j.neuron.2011.12.003.

33.　Koh, K., Joiner, W.J., Wu, M.N., Yue, Z., Smith, C.J., and Sehgal, A. (2008).
Identification of SLEEPLESS, a sleep-promoting factor. Science *321*, 372–
376. 10.1126/science.1155942.

34.　Afonso, D.J., Liu, D., Machado, D.R., Pan, H., Jepson, J.E., Rogulja, D., and
Koh, K. (2015). TARANIS Functions with Cyclin A and Cdk1 in a Novel
Arousal Center to Control Sleep in Drosophila. Curr Biol *25*, 1717–1726.
10.1016/j.cub.2015.05.037.

35.　Saitoh, S., Takamatsu, K., Kobayashi, M., and Noguchi, T. (1994). Expression
of hippocalcin in the developing rat brain. Brain Res Dev Brain Res *80*, 199–
208. 10.1016/0165-3806(94)90105-8.

36.　Jo, J., Son, G.H., Winters, B.L., Kim, M.J., Whitcomb, D.J., Dickinson, B.A.,
Lee, Y.B., Futai, K., Amici, M., Sheng, M., et al. (2010). Muscarinic receptors
induce LTD of NMDAR EPSCs via a mechanism involving hippocalcin, AP2
and PSD-95. Nat Neurosci *13*, 1216–1224. 10.1038/nn.2636.

37.　Tzingounis, A.V., Kobayashi, M., Takamatsu, K., and Nicoll, R.A. (2007).
Hippocalcin gates the calcium activation of the slow afterhyperpolarization in
hippocampal pyramidal cells. Neuron *53*, 487–493.
10.1016/j.neuron.2007.01.011.

38.　Palmer, C.L., Lim, W., Hastie, P.G., Toward, M., Korolchuk, V.I., Burbidge,
S.A., Banting, G., Collingridge, G.L., Isaac, J.T., and Henley, J.M. (2005).
Hippocalcin functions as a calcium sensor in hippocampal LTD. Neuron *47*,
487–494. 10.1016/j.neuron.2005.06.014.

39.　Kobayashi, M., Masaki, T., Hori, K., Masuo, Y., Miyamoto, M., Tsubokawa, H.,
Noguchi, H., Nomura, M., and Takamatsu, K. (2005). Hippocalcin-deficient
mice display a defect in cAMP response element-binding protein activation
associated with impaired spatial and associative memory. Neuroscience *133*,
471–484. 10.1016/j.neuroscience.2005.02.034.

40.　Atasu, B., Hanagasi, H., Bilgic, B., Pak, M., Erginel-Unaltuna, N., Hauser,
A.K., Guven, G., Simon-Sanchez, J., Heutink, P., Gasser, T., and Lohmann, E.
(2018). HPCA confirmed as a genetic cause of DYT2-like dystonia phenotype.
Mov Disord *33*, 1354–1358. 10.1002/mds.27442.

41.　Magrinelli, F., Bhatia, K.P., Beiraghi Toosi, M., Arab, F., Karimiani, E.G.,
Sedighzadeh, S., Ansari, B., Neshatdoust, M., Rocca, C., Houlden, H., and
Maroofian, R. (2023). Childhood-Onset Choreo-Dystonia Due to a Recurrent
Novel Homozygous Nonsense HPCA Variant: Case Series and Literature
Review. Mov Disord Clin Pract *10*, 101–108. 10.1002/mdc3.13529.

42.　Sitaraman, D., Vecsey, C.G., and Koochagian, C. (2024). Activity Monitoring
for Analysis of Sleep in Drosophila melanogaster. Cold Spring Harb Protoc
2024, pdb top108095. 10.1101/pdb.top108095.

43.　Kittel, R.J., Wichmann, C., Rasse, T.M., Fouquet, W., Schmidt, M., Schmid, A.,
Wagh, D.A., Pawlu, C., Kellner, R.R., Willig, K.I., et al. (2006). Bruchpilot

promotes active zone assembly, Ca2+ channel clustering, and vesicle
release. Science *312*, 1051–1054. 10.1126/science.1126308.

44. Gilestro, G.F., Tononi, G., and Cirelli, C. (2009). Widespread changes in
synaptic markers as a function of sleep and wakefulness in Drosophila.
Science *324*, 109–112. 10.1126/science.1166673.

45. Weiss, J.T., Blundell, M.Z., Singh, P., and Donlea, J.M. (2024). Sleep
deprivation drives brain-wide changes in cholinergic presynapse abundance
in Drosophila melanogaster. Proc Natl Acad Sci U S A *121*, e2312664121.
10.1073/pnas.2312664121.

46. Huang, S., Piao, C., Beuschel, C.B., Gotz, T., and Sigrist, S.J. (2020).
Presynaptic Active Zone Plasticity Encodes Sleep Need in Drosophila. Curr
Biol *30*, 1077–1091 e1075. 10.1016/j.cub.2020.01.019.

47. Ozturk-Colak, A., Inami, S., Buchler, J.R., McClanahan, P.D., Cruz, A., Fang-
Yen, C., and Koh, K. (2020). Sleep Induction by Mechanosensory Stimulation
in Drosophila. Cell Rep *33*, 108462. 10.1016/j.celrep.2020.108462.

48. Pellicano, E., Rhodes, G., and Calder, A.J. (2013). Reduced gaze aftereffects
are related to difficulties categorising gaze direction in children with autism.
Neuropsychologia *51*, 1504–1509. 10.1016/j.neuropsychologia.2013.03.021.

49. Coll-Tane, M., Eidhof, I., Han, J., Raun, N., van Renssen, L.V., Fisher, S.E.,
Kayser, M.S., Kleefstra, T., Pillen, S., Hudac, C.M., et al. (2026). Conserved
sleep disturbances in FOXP1 syndrome originate from developmental
dysregulation of peptidergic signaling. J Clin Invest *136*. 10.1172/JCI193475.

50. Bruni, O., Breda, M., Mammarella, V., Mogavero, M.P., and Ferri, R. (2025).
Sleep and circadian disturbances in children with neurodevelopmental
disorders. Nat Rev Neurol *21*, 103–120. 10.1038/s41582-024-01052-9.

51. Ethridge, L.E., White, S.P., Mosconi, M.W., Wang, J., Byerly, M.J., and
Sweeney, J.A. (2016). Reduced habituation of auditory evoked potentials
indicate cortical hyper-excitability in Fragile X Syndrome. Transl Psychiatry *6*,
e787. 10.1038/tp.2016.48.

52. Ewbank, M.P., Rhodes, G., von dem Hagen, E.A., Powell, T.E., Bright, N.,
Stoyanova, R.S., Baron-Cohen, S., and Calder, A.J. (2015). Repetition
Suppression in Ventral Visual Cortex Is Diminished as a Function of
Increasing Autistic Traits. Cereb Cortex *25*, 3381–3393.
10.1093/cercor/bhu149.

53. Fenckova, M., Blok, L.E.R., Asztalos, L., Goodman, D.P., Cizek, P., Singgih,
E.L., Glennon, J.C., IntHout, J., Zweier, C., Eichler, E.E., et al. (2019).
Habituation Learning Is a Widely Affected Mechanism in Drosophila Models of
Intellectual Disability and Autism Spectrum Disorders. Biol Psychiatry *86*,
294–305. 10.1016/j.biopsych.2019.04.029.

54. Restivo, L., Ferrari, F., Passino, E., Sgobio, C., Bock, J., Oostra, B.A., Bagni,
C., and Ammassari-Teule, M. (2005). Enriched environment promotes
behavioral and morphological recovery in a mouse model for the fragile X
syndrome. Proc Natl Acad Sci U S A *102*, 11557–11562.
10.1073/pnas.0504984102.

55. Lovelace, J.W., Wen, T.H., Reinhard, S., Hsu, M.S., Sidhu, H., Ethell, I.M.,
Binder, D.K., and Razak, K.A. (2016). Matrix metalloproteinase-9 deletion
rescues auditory evoked potential habituation deficit in a mouse model of
Fragile X Syndrome. Neurobiol Dis *89*, 126–135. 10.1016/j.nbd.2016.02.002.

56. Coll-Tane, M., Gong, N.N., Belfer, S.J., van Renssen, L.V., Kurtz-Nelson, E.C.,
Szuperak, M., Eidhof, I., van Reijmersdal, B., Terwindt, I., Durkin, J., et al.

(2021). The CHD8/CHD7/Kismet family links blood-brain barrier glia and
serotonin to ASD-associated sleep defects. Sci Adv 7.
10.1126/sciadv.abe2626.

57. Shelton, A.R., and Malow, B. (2021). Neurodevelopmental Disorders
Commonly Presenting with Sleep Disturbances. Neurotherapeutics *18*, 156–
169. 10.1007/s13311-020-00982-8.

58. Tsien, J.Z. (2000). Linking Hebb's coincidence-detection to memory formation.
Curr Opin Neurobiol *10*, 266–273. 10.1016/s0959-4388(00)00070-2.

Figure 1

Figure 2

Figure 3

Figure 4