Engineering directional phosphoryl flow enables programmable signaling dynamics in bacteria

Short title: Reprogramming cellular decisions by engineering directional information flow

Authors: Marcos Nieves[1,†], Juan Manuel Valle[1,2,†], Joaquín Dalla Rizza[1], Federico Carrión[3], Pablo Naranjo-Meneses[4,5], Nicole Larrieux[1], Ilka B. Bischofs[4,5], Alejandro Buschiazzo[1,6], Felipe Trajtenberg[1,2]*

Affiliations:

[1]Laboratory of Molecular and Structural Microbiology, Institut Pasteur de Montevideo, Montevideo, Uruguay

[2]Locbio, Montevideo, Uruguay

[3]Laboratory of Immunovirology, Institut Pasteur de Montevideo, Montevideo, Uruguay,

[4]BioQuant Center, Heidelberg University, Heidelberg, Germany

[5]Max Planck Institute for Terrestrial Microbiology, Marburg, Germany

[6]Département de Microbiologie, Institut Pasteur, Paris., France

*Corresponding author. Email: felipet@pasteur.edu.uy

Abstract:

Cells must make critical decisions by integrating information from a constantly changing environment to ensure their survival. They rely on intricate signaling networks to detect external and internal cues to trigger specific responses, yet these systems are generally viewed as components wired with simpler linear connections. Bacterial phosphorelay systems offer a versatile framework for studying more complex connections and engineering new biological circuits. Here, using the well-studied *Bacillus subtilis* sporulation phosphorelay, we demonstrate that the directionality of information flow can be reprogrammed to generate different dynamic responses. We show that phosphoryl-transfer reversibility is an evolvable trait encoded in conserved, surface-exposed motifs of two-component system proteins. Unidirectional phosphoryl-transfer generates a short-term information storage mechanism, enabling signal integration over time and allowing phosphatases, acting at different levels, to produce different outcomes. In contrast, a bidirectional system enhanced the action of phosphatase activity early in the pathway. The ability to control phosphoryl-transfer equilibria opens exciting avenues for designing sophisticated synthetic signaling systems with enhanced decision-making capabilities.

INTRODUCTION

Cells are constantly evaluating their environment and collecting information about their internal status, performing complex biological computations to make decisions. The adaptive response triggered by sensory systems allows for survival under highly fluctuating environments, including

cell communication, nutrient limitation, temperature shifts or desiccation. TCSs are widely spread
signaling pathways in bacteria and archaea, but also present in fungi and plants [1]. Information
travels from a sensory histidine kinase (HK) to an effector response regulator (RR) by means of
phosphoryl-transfer reactions between specific histidine and aspartate residues [2].

An interesting feature of TCS chemistry, is that phosphoryl-transfer equilibria can be displaced
towards the histidine or the aspartate residues (S1 Fig). Based purely on energetic grounds, the
phosphoryl group *en transfer* tends to go to the histidine's imidazole, and away from the
carboxylate side chain of aspartate [3]. However, different mechanisms can overcome this
energetic uphill [4, 5], eventually favoring aspartate phosphorylation. Among such mechanisms,
the donor:acceptor interatomic distance within the reaction center [4, 5], HK allosteric
conformational rearrangements, as well as RR dimerization [4], have all been shown to contribute
to determining the overall directionality of the reaction. In the simplest TCS configuration, with a
HK and a RR, a strongly unidirectional system is needed to promote turnover and efficient
signaling [4].

Phosphorelays are a complex type of TCS, comprising several intermediate proteins between the
sensory HK and the effector RR. Longer phosphoryl-transfer cascades as this enables a rich
integrative signaling processes that regulate important biological phenomena [6, 7]. The
phosphoryl group must travel through multiple reaction centers with disparate directions: His to
Asp, as well as Asp to His. This raises several fundamental questions: Are phosphoryl-transfer
reactions fully reversible in these pathways? Can the equilibrium of these reactions be engineered
to tune their directionality? What properties can emerge in these networks?

To answer these questions, we turned our attention to the sporulation regulatory pathway from *B.*
subtilis [6]. Sporulation is an important developmental process that generates a highly specialized
cell, the dormant endospore, which sustains a stagnated metabolism that renders them extremely
resistant to environmental stress [8]. Several HKs (KinA, KinB, KinC, KinD and KinE) [9]
monitor internal and external signals in the *B. subtilis* cells (S2 Fig). This sensory information is
channeled through intermediate components, Spo0F and Spo0B, that transfer the phosphoryl
moiety to the downstream master regulator Spo0A [6]. Spo0F and Spo0A are RRs, while Spo0B
is a phosphotransferase. The phosphorelay is also affected by dedicated phosphatases, such as
receptor aspartyl phosphatases (Raps) and Spo0E-like proteins, that drain the phosphoryl group
from Spo0F and Spo0A, respectively [10, 11]. Spo0A controls different outcomes, including
sporulation [6], biofilm formation [12], cannibalism [13] and cellular competence [14]. Once a
certain level of phosphorylated Spo0A is reached [15], a complex phenotypic reprogramming is
triggered, ultimately ending with the formation of the endospore [15]. In addition to other
downstream target genes, Spo0A is also involved in regulating the expression of components of
the phosphorelay pathway itself, thus shaping complex behaviours under different conditions [16].
The *B. subtilis* sporulation phosphorelay is very well studied both experimentally and
computationally [14, 17, 18], an ideal model to address our questions.

Here we show that the sporulation phosphorelay from *B. subtilis* can be reprogrammed by specific
mutations at key protein:protein interfaces, reshaping the output dynamics of the signaling system.
We demonstrate that natural phosphoryl-transfer equilibria are displaced towards downstream
components in each reaction pair along the sporulation path. A defined His-Asp distance is
determined by conserved motifs on the two RRs that interact with Spo0B. These interface contacts
position each RR in a different pose, suggesting that directionality determinants are encoded within
the RR structure and they are evolutionarily selected. Through experimental approaches and

mathematical modelling, we show that the *B. subtilis* sporulation phosphorelay can transiently
store information in the form of phosphorylated Spo0A, enabling a rheostat-like behavior of the
system depending on the kinase/phosphatase activity ratio. In contrast, imposing a reversible
phosphoryl-transfer equilibrium enhances the action of Rap phosphatases. These findings provide
a new understanding of how protein-level mechanisms can shape network-wide signaling
dynamics and emergent properties.

RESULTS

Unidirectional phosphoryl-transfer towards downstream phosphorelay components

Considering that 3D structures of the Spo0F:Spo0B complex from *B. subtilis* have been known for
decades, to better understand the phosphorelay's signaling mechanism we determined the crystal
structure of the Spo0B:Spo0A binary complex. Its structure was solved in two different crystal
forms (S1 Table) at 2.1 and 3.5 Å resolution. The refined atomic model shows the expected 2:2
phosphotransferase:RR stoichiometry (Fig 1A), similarly to Spo0F:Spo0B. The transferase
interacts with Spo0A mainly through helices α1 and α2 on the DHp (***D***imerization and ***H***istidine
phosphotransfer) domain with the α1α5 face of Spo0A, analogous to many other HPt:RR (HPt
stands for ***H***istidine ***P***hospho*t*ransferase) and HK:RR complexes [5, 19-23]. In each of the
independently refined Spo0B:Spo0A complexes (7 in total), one of the two Spo0A molecules
displays better defined electron density, resulting in lower refined B factors. Interestingly, these
can be explained by additional contacts established between Spo0A and the pseudo-ATP-binding
domain (pseudoABD) of Spo0B, breaking the symmetric structure of Spo0B.

Despite overall similar organizations of Spo0A and Spo0F with respect to Spo0B, the
superposition of each complex using Spo0B's α1/α2 helices as reference, reveals a significant shift
in the position of Spo0A, involving translation and rotation relative to Spo0B (Fig 1B). As a result,
the phosphorylatable Asp of Spo0A is placed significantly farther away from Spo0B's His (6.6 Å)
compared to the equivalent Asp in Spo0F (5.00 Å), as shown in Fig. 1C. Interestingly, the
Spo0F:Spo0B complex exhibits a larger REC:pseudoABD interface, which buries a surface of
~300 Å^2 and includes hydrogen-bonds and van der Waals contacts. In contrast, the equivalent
Spo0B:Spo0A interface totals ~90 Å^2 with only a few van der Waals interactions (Fig 1D). From
a structural point of view, the receiver domain of Spo0A and Spo0F are very similar (Fig 1E).
However, both complexes show specific sets of contacts with Spo0B that are differential and
engage evolutionarily conserved residues [24]. Notably, key differences lie on the DHp:REC
interface. The highly conserved motif REXX (including Arg18 and Glu19 at the beginning of helix
α1) in Spo0A, is substituted to XGGR in Spo0F (Fig 1E and S3). This effectively moves the
position of the Arg by one helix turn (Arg16$_{Spo0F}$ → Arg19$_{Spo0A}$), while introducing a bulkier
adjacent residue (Gly14$_{Spo0F}$ → Glu18$_{Spo0A}$). In addition, the conserved Phe92 from Spo0A is
packed against Spo0B residues Met37 from α1, and Val63, Ala65 and Lys67 from α2. On the other
hand, Spo0F includes a Tyr in the same position, which is solvent exposed. Taking all together,
the data imply an evolutionary pressure to correctly place each RR protein with respect to the
Spo0B partner, placing the phosphorylatable His and Asp residues at specific and distinct
distances. A longer interatomic distance between phosphoryl-donor and -acceptor are known to
correlate with a stronger directionality from P~His to Asp [4]. This suggests that the Spo0F:Spo0B
and Spo0B:Spo0A phosphotransfer equilibriums could be differentially displaced towards either
the His or the Asp, respectively.

To confirm our prediction, we studied the reaction of the phosphorylated phosphotransferase
Spo0B (P~Spo0B) with the two RR *in vitro* using radiolabeled ATP[γ-^{33}P]. A fixed concentration
of P~Spo0B [15 μM] was exposed to varying concentrations of Spo0A (or Spo0F) from 0 to 50
μM. Equilibrium was reached after 1 min. The distribution of phosphate among the two proteins
was analyzed by autoradiography and showing an accumulation of phosphorylated RR [RR~P]
with a concomitant decrease of Spo0B~P as the concentration of RR increases (S4 Fig). The
fraction of phosphate on the RR over all phosphorylated proteins (ξ_p, defined as
[RR~P]/([RR~P]+[Spo0B~P])) for both Spo0A (red symbols) and Spo0F (blue symbols) was well
described by a Hill function (Fig 1F). The Spo0B-Spo0F phosphoryl-transfer showed low levels
of cooperativity (Hill coefficient n = 1.2 for Spo0F) while the reaction between Spo0B and SpoA
showed strong cooperativity (n = 2.2). Moreover, when compared to the equilibrium distribution
of an ideal fully reversible phosphoryl-transfer reaction (black line), it is clear that phosphoryl-
transfer between Spo0B and Spo0F was shifted towards the phosphotransferase (Asp to His
direction), while the phosphoryl-transfer between Spo0B and Spo0A is shifted towards the RR
(His to Asp direction). Together these findings indicate that the *B. subtilis* sporulation
phosphorelay is configured to facilitate a directional phosphoryl-flux from upstream to
downstream signaling components.

Reconfiguring reversibility in TCS phosphorelays

Next, we tested whether we could modulate the phosphoryl-transfer directionality by mutating
residues in the binding interface of Spo0B with Spo0F and Spo0A. Informed by the structure of
the complexes, we chose eleven Spo0F (F1-11) and three Spo0A (A1-3) variants by grafting five
different interface motifs (S5 Fig). For example, Spo0F variants F1 and F2 involve grafting of one
or more residues from motif 1 of Spo0A to Spo0F, F3 and F4 are modifications at motif 3, and F5
to F7 are combinations of modifications in motifs 1 and 3 (see full details in S2 Table and S5 Fig).
For Spo0A, we included the double mutant Y108A and H122A that interferes with dimerization
(A1), a point mutant E14G (A2), which is part of the conserved motif interacting with Spo0B and
a truncated variant on Spo0A's, including only the receiver domain (A3).

For all proteins variants we measured the distribution of phosphoryl-groups after incubation with
P~Spo0B as before (Fig. 2A). For Spo0F (blue bars), substitutions at motif 3 (present in F3, F4,
F6, F7 and F8) resulted in a significant shift of the phosphoryl-equilibria towards the RR compared
to wild-type Spo0F, losing directionality. The strongest effects were seen for F3 and F4, both
containing a K56I substitution that is located two residues after the phosphorylatable Asp. In
contrast, mutant F9, which includes substitutions at motifs 1, 2 and 4, shifted the reaction towards
phosphorylation of the phosphotransferase Spo0B. This effect was lost by including K56I and
E86Q substitutions on motif 3 (F10 and F11), suggesting a compensatory effect between motifs.
Notably, all three Spo0A mutants shifted the phosphoryl-transfer equilibrium towards Spo0B (red
bars).

We next performed titration assays with two Spo0F variants F3 (green line) and F6 (cyan line) and
compared it to wild type Spo0F (F0, blue line) as shown in Fig 2B. At equimolar concentrations
of Spo0B and Spo0F, the phosphoryl-group were almost evenly split between both proteins
(dashed lines), indicative of a fully reversible phosphoryl-transfer equilibrium. Interestingly, a
single substitution K56I on Spo0F (F3) was enough to make the phosphoryl-transfer fully
reversible

Taken together these data show that directionality of both phosphotransferase reactions can be
shifted from the directional phosphoryl-transfer, as observed in wild type, to a fully reversible
equilibrium.

Phosphoryl flow directionality is essential for phosphorylation-dependent short-term memory

Collectively, our results suggest that phosphoryl-transfer directionality in the sporulation
phosphorelay is a trait that can be accessed by evolution. We thus wondered whether the
directionality of phosphoryl-transfer reactions could confer specific properties to a signaling
system. The overall organization of the sporulation regulation pathway (S2 Fig) led us to speculate
that phosphoryl-transfer reversibility could affect the action of phosphatases at the level of Spo0F.
To investigate this possibility, we used a deterministic mass-action kinetic model of the signaling
system (Fig 3A) based on reported data [18, 25] . The kinetic model included KinA, Spo0F, Spo0B,
Spo0A and RapA. Feed-forward loops (in KinA, Spo0F, Spo0A and RapA) were modelled as Hill
equations, dependent on the levels of P~Spo0A (full details in Material and Methods). Under a
regime of unidirectional phosphoryl-transfer reactions, as observed in our biochemical
characterizations (Fig 1F), kinetic modelling indicated that even at very high concentration of
phosphatase activity, information accumulates over time (Fig 3B, left panel), in the form of total
amount of P~Spo0A.

To test the predictions derived from the kinetic model, we constructed *B. subtilis* strains that
included a super folder green fluorescent protein (GFP) reporter gene under the *spoIIE* high-
threshold promoter. This was generated with a *spo0F*, *spo0E* and *rapA* triple knockout strain.
RapA was inserted ectopically under an IPTG-inducible *Phyperspank* promoter, to enable the
control of phosphatase expression in the absence of the PhrA inhibitor. On this GFP reporter strain,
we then reverted the *spo0F* knockout by homologous recombination at the endogenous locus with
wild type *spo0F* (strain BSS117). Exponentially growing *B. subtilis* cells were switched from a
rich to a poor medium to induce sporulation at 37 ºC, and fluorescence was measured over time.
Different IPTG concentrations were used to control RapA expression at different steady state
levels. The back-complemented strain showed an increase in fluorescence over time, even at the
highest IPTG concentrations (Fig 3B, right panel). Although higher RapA levels slowed down the
activation process, eventually cells reached the maximum signal (Fig 3B, right panel).

On the other hand, simulations of a phosphorelay with a fully reversible Spo0F:Spo0B phosphoryl-
transfer step, as measured for the Spo0F-K56I mutant (F3, Fig 2B), showed that a high enough
RapA concentration prevents accumulation of P~Spo0A and, therefore, precludes the activation
of high-threshold promoters (Fig 3C, left panel). Note that, by including the Spo0F_K56I variant,
only the Spo0F:Spo0B step is being modified, such that the pathway retains some directionality
towards Spo0A phosphorylation, albeit to a lesser extent. As predicted by the model, the back-
complemented strain with Spo0F-K56I variant was unable to activate the Spo0A-controlled
pathway at the highest RapA concentrations (Fig 3C, left panel). These results indicate that
phosphoryl-transfer reversibility is relevant, and it is configured in such a way that the sporulation
regulation system can slowly accumulate information over time, in the form of P~Spo0A, even
under high RapA phosphatase activity.

In a different experimental setup, where RapA activity is induced after the system was activated,
the signaling pathway with wild type Spo0F stored information in the form of P~Spo0A (Fig 3D).
In contrast, the *spo0F-K56I* strain showed a complete reset of the system at high RapA
concentrations (Fig 3E), indicating that pathway activation can be reversed, consistent with our

simulations. Our kinetic model further predicts that, unlike the observed effect of RapA
phosphatase in the *spo0F* wild-type background (Fig. 3B), phosphatase activity acting at the
Spo0A level should reset the system (Fig. S6A). This prediction was confirmed experimentally by
monitoring *spoIIE* promoter activity in a *spo0E rapA* double knockout strain, in which Spo0E was
expressed under IPTG control (S6B Fig). Altogether, these results indicate that the directionality
of phosphoryl flow in the sporulation phosphorelay is organized to transiently record and store
information, effectively functioning as a short-term memory system that can be reset by
downstream phosphatase activity.

DISCUSSION

This study reveals that the directionality of phosphoryl-transfer reactions plays a pivotal role in
information storage, signal integration and shaping of output responses. Challenging a common
view of signaling networks as simple and linear pathways, phosphorelays offers the opportunity
to create more complex network connections. We now show that by modifying conserved, surface-
exposed motifs, we can engineer networks with distinct phosphoryl-transfer reversibility,
highlighting how the inherent biochemistry of protein-protein interactions can be manipulated to
generate sophisticated, non-intuitive network behaviors.

The phosphotransfer reaction between Spo0F and Spo0B is shifted towards histidine
phosphorylation, as expected on energetic grounds. However, the opposite is observed in the
Spo0B:Spo0A phosphotransfer equilibrium, indicating that each RR protein encodes the
directionality determinants in its 3D structure. The crystal structure of the Spo0B:Spo0A complex
revealed a longer His-Asp distance compared to that of Spo0F:Spo0B. The distance between the
phosphoryl donor and acceptor residues has been proposed to be a key directionality determinant
[4, 5]. Spo0A dimerization is also a directionality determinant, a feature shared with other RRs
[4]. Thus, RR dimerization not only triggers an effector response, but in addition plays a role in
minimizing back-transfer of the phosphoryl group to the HK, and thereby, maximizes a
unidirectional, downstream phosphoryl flux.

The structural comparison between Spo0B:Spo0A and Spo0F:Spo0B uncovered specific
interfaces that seem key for the proper positioning of the RR within the complex. We had
previously shown that HK:RR interactions are "slippery" [5] and that such plasticity allows for the
relocation of the RR within the complex guided by interactions with additional interfaces, such as
those between the ABD and REC domains [4]. In the case of the TCS DesK:DesR from *B. subtilis*,
the ABD:REC contacts shift the position of the RR, bringing the phosphorylated Asp residue closer
to the conserved Gln [4]. Analogously, we now reveal that the interactions between the
pseudoABD domain of Spo0B and Spo0F are tighter than with Spo0A. In addition, Spo0F and
Spo0A show conserved differences in positions that are directly involved in the DHp:REC
interaction. These two traits, end up placing Spo0F in a shifted position with respect to Spo0A (Fig
1) in complex with Spo0B, resulting in a shorter His-Asp distance, and suggesting that
phosphotransfer directionality is encoded in the protein:protein interaction surfaces. Sequence
conservation indicates that directionality might also be an evolvable property, a matter for future
studies. The fact that directionality determinants between Spo0F and Spo0A can be grafted and
exchanged, indeed suggests that directionality is transferrable, and thus evolvable under selective
pressure.

Notably, the identified *spo0F_K56I* mutant, which shifts the phosphoryl-transfer equilibrium,
allowed us to assess the relevance of phosphorelays with different directional transmission of

information. As we now show, unidirectional phosphoryl-transfer in the sporulation pathway
allows a mechanism of short-term memory, whereby P~Spo0A can be accumulated over time.
This stored information is reset by the action of Spo0A-specific phosphatases like Spo0E (S6 Fig).
All things considered, a conceptual model implies a system that can switch between recording,
storing and resetting phases (Fig 4). Moreover, the system can work as a rheostat, offering
a gradual and controlled response to changes in kinase/phosphatase ratios (S7 Fig). This rheostat-
like behavior is enabled by the phosphoryl-transfer unidirectionality and means that rather than
simply turning "on" or "off," the pathway can fine-tune its activation dynamics across a continuous
spectrum. This allows it to effectively integrate the diverse signals from various sensor HKs and
phosphatases.

The sporulation pathway exhibits distinct behaviors under varying growth conditions. In some
cases, it follows a pulsatile pattern [18, 26], while in others, cells gradually accumulate P~Spo0A
until reaching a specific threshold that triggers pathway activation [15, 27]. The promoter
controlling *rapA* can also show pulsatile dynamics [28]. Such different dynamics could be
generated due to different kinase/phosphatase ratios, as well as by triggering alternative dedicated
phosphatase activities, like RapA or Spo0E. However, further studies are needed in less-
engineered strains and in different culture scenarios to fully understand how *B. subtilis* makes the
decision to sporulate.

A striking aspect of the sporulation pathway is the ability to generate different cellular outcomes
while integrating a myriad of signals [29]. This has been further supported by recent observations
showing that different HKs can trigger alternate cell fates [30]. The presence of different Spo0A
promoters with distinct thresholds [16] could imply, not only the ability to control sequential steps
along the sporulation program, but also the capacity to trigger alternate genetic programs due to
different activation kinetics. The latter, as we see in our study, are coupled to time-sensitive
activation or repression of phosphatases in the sporulation phosphorelay. The modularity and
signal-integrating features of this pathway represent a blueprint for building complex, multi-input
genetic circuits with tailored behaviors.

The significance of phosphoryl-transfer reversibility extends beyond the *B. subtilis* sporulation
pathway. The chemical versatility of His to Asp phosphotransfer allows generating reactions with
varying degree of reversibility (S1 Fig). Phosphotransfer directionality has been proposed to
warrant efficient signal transmission [4], and also to allow for "phosphate sinks" in several
phosphorelay systems [31-33]. In summary, we have shown that phosphotransfer directionality is
a key determinant to shape how information is integrated and processed. We also demonstrated
that directionality can be reprogrammed and can thus be subject to evolutionary pressure. This
adds a completely new layer of complexity in rewiring signaling pathways, involving not only the
inter-protein specificity determinants [34], but also the directionality of the phosphoryl-transfer
reactions. Engineering signaling pathways could benefit from the ability to modulate the
reversibility of such equilibria in order to generate more efficient and tunable information-
processing systems for Synthetic Biology.

MATERIALS AND METHODS

Cloning and protein purification

Cloning and protein purification were performed as described previously [35]. The
plasmid for recombinant production of KinA (pETM_KinA) was constructed by cloning into a
pETM-14 plasmid (kindly provided by Frank Lehman) using primers KinA_F and KinA_R by

restriction-free cloning [36]. Similar procedure was followed to generate plasmids pQE80_Spo0A
and pQE80_Spo0A_REC, using primers Spo0A_F, Spo0A_R and Spo0A_REC_R. The primers
spo0E_up_F, spo0E_up_R and Spo0E_down_F, spo0E_down_R were used to amplify 500-bp
fragments upstream and downstream of spo0E, respectively. The DNA fragments were fused by
PCR and cloned into the pMAD vector by RELC using the enzymes SalI and BglII.
Plasmids pQE80_Spo0F, pQE80_Spo0B, pET_Spo0B_strep, and all the Spo0F and Spo0A
mutants tested, were obtained by synthesis and subcloning services (Genscript). Tables of strains,
plasmids and oligonucleotide primers are included as supplementary data (S3-S5 Tables).

Recombinant proteins were expressed as N-terminally His6-tagged fusions in *E. coli* strain BL21
(DE3) pLysS as previously performed for other TCS proteins [37]. Transformed bacteria were
grown in TB medium, and recombinant protein expression was induced during exponential growth
by adding 1 mM IPTG for 18 hours at 20 ºC. Cells were harvested by centrifugation and
resuspended in lysis buffer (50 mM Tris-HCl pH 8.0, 0.5 M NaCl, 1 mg/mL lysozyme, 0.01%
Triton X-100, 20 mM imidazole). Cell lysis was performed using either sonication or an
Emulsiflex homogenizer. The lysates were centrifuged, and the soluble fraction was loaded onto a
5 mL HisTrap HP column (Cytiva) equilibrated with buffer A (50 mM Tris-HCl, pH 8.0, 500 mM
NaCl, 20 mM imidazole) for IMAC purification. After this initial step, the His6-tag was removed
by proteolysis with TEVdt protease [38]. Imidazole from the IMAC step was removed by dialysis
against 50 mM Tris-HCl pH 8.0, 0.5 M NaCl. Undigested protein and protease were removed by
reverse IMAC. The final purification step was size exclusion chromatography using a HiLoad
16/60 Superdex 75 prep grade column (Cytiva), equilibrated with 20 mM Tris-HCl pH 8.0, 0.1 M
NaCl (SEC buffer). All proteins were concentrated to ~10 mg/mL and stored at -80°C.

Protein crystallization, data collection and model building/refinement

The crystals of Spo0B:Spo0A complex were obtained by sitting-drop vapor-diffusion at 293 K in
96 well plates (Greiner) assisted by a Honeybee 963 robotic station (Digilab). The protein complex
was prepared by mixing purified and concentrated Spo0B (57 mg/mL) and Spo0A-REC (50
mg/mL) at 1:1.1 equivalents. The complex was prepared in the presence of 10 mM MgCl$_2$ and
diluted in SEC buffer to a final concentration of 25 mg/mL. Drops were set up with 300 nL of
protein solution and 300 nL of reservoir, equilibrated against 100 µL of reservoir solution (0.2 M
NaF, 20% (w/v) polyethylene glycol 3350). Crystals were cryoprotected by quick-soaking in a
solution containing 0.155 M NaF, 22% (w/v) PEG 3350, 25% (v/v) glycerol, and flash frozen in
liquid nitrogen.

Single crystal X-ray diffraction data collection was performed at 100 K using either a copper
rotating anode home source (Protein Crystallography Facility, Institut Pasteur de Montevideo), or
synchrotron radiation at Beamline I-03 (Diamond Light Source, UK). Data processing was carried
out with Xia2 [39] or autoPROC [40]. The Spo0B:Spo0A complex was solved by molecular
replacement [41] using the Spo0B dimer (PDB ID 1IXM) and the Spo0A receiver domain (PDB
ID 1QMP) as search probes. Model rebuilding, refinement and validation were carried out using
Coot [42], Buster [43], Phenix [44] and MolProbity tools [45]. Protein visualization, structural
analyses, and figure rendering were performed with Pymol (Schrodinger, LLC, 2015). The
software used for data processing, structure determination, and analysis was provided by the
SBGrid Consortium [46].

Phosphotransferase assays

Phosphotransfer reactions were performed with P~Spo0B radiolabeled with ^{33}P and different Spo0F and Spo0A variants. P~Spo0B was prepared by incubating a His-tagged Spo0B with KinA and Spo0F in the presence of 150 µCi [γ-^{33}P]-ATP (10 mCi/mL, American Radiolabelled Chemicals, Inc.), unlabeled ATP (400 µM final concentration) and $MgCl_2$ for 30 min at room temperature in reaction buffer (100 mM NaCl, 20 mM Tris-HCl pH 8 and 5 mM $MgCl_2$). P~Spo0B was then purified using a Histrap column (Cytiva), concentrated and buffer-exchanged to reaction buffer using ultrafiltration. Proteins were analyzed by SDS-PAGE, quantified, aliquoted and stored at -20 ˚C. Evaluation of the phosphoryl-transfer equilibria was done by incubating 15 µM P~Spo0B with 20 µM of each of the RRs (Spo0F and Spo0A variants), for 10 min at 25 °C. Reactions were quenched with SDS-PAGE sample buffer and subsequently ran in polyacrylamide gels (Cytiva). Gels were exposed to phosphor screen (for 1 h) and imaged in a Typhoon FLA 7000 imaging system (GE Healthcare). Band intensities were analyzed by densitometry with ImageJ [47] and plotted in Matlab R2020a (MathWorks Inc.). Titration assays were performed as above but with varying concentrations of RRs (from 0 to 50 µM). Each reaction was performed in triplicate.

Phosphotransfer assays between P~Spo0F and Spo0A, with limiting amounts of Spo0B, were performed following similar procedures as previously [4]. Briefly, Spo0F was phosphorylated with KinA in the presence of ATP and $MgCl_2$, and then purified and buffer-exchanged in a Superdex75 10/300 column (Cytiva) equilibrated with reaction buffer. Equimolar amounts of P~Spo0F and Spo0A (15 µM) were incubated in the presence of 0.3 µM Spo0B. At different time points, the reactions were stopped by adding SDS-PAGE sample buffer and loaded in a Phostag SDS-PAGE. Coomassie blue-stained gels were scanned, and quantification was done by densitometry using ImageJ.

B. subtilis strain construction

All strains are derivatives of *B. subtilis* 168 1A700 and derived from strain BIB415 (*ΔrapA ΔphrA Δspo0F* [48] by a series of genetic modifications. First, *spo0E* was deleted with the help of a pMAD_spo0E plasmid constructed with primers ST94-99 following established protocols [48]. Next, *rapA* was placed under the control of the P*hyperspank* promoter by transformation with pDR111_rapA [49]. Subsequently, the P*spoIIE*_GFP and Pspo0F_mCherry constructs were inserted into the *yhdG_yhdH* and *bglS* loci, respectively. Finally, the *spo0F* knockout was reverted to either the wild-type allele or the *spo0F-K56I* mutant using plasmids pMAD_Spo0F [48] and pMAD_Spo0F-K56I, obtaining strains BSS117 *ΔrapAΔphrA Δspo0E amyE::[specR Phyperspank_rapA] yhdG_yhdH::[phleoR Psdp_mCherry] bglS::[kanR PspoIIE_gfp]* and BSS118 *ΔrapAΔphrA Δspo0E amyE::[specR Phyperspank_rapA] yhdG_yhdH::[phleoR Psdp_mCherry] bglS::[kanR PspoIIE_gfp] spo0F-K56I*). Similarly, starting with the *spo0E* knockout of strain BIB415, *spo0E* was reinserted in the genome under the control of the P*hyperspank* promoter by transformation with pDR111_spo0E. Subsequently, reporter constructs insertion and *spo0F* knockout restoration were performed identically as for the previous two strains. Thus, obtaining strain BSS119 *ΔrapAΔphrA Δspo0E amyE::[specR Phyperspank_spo0E] yhdG_yhdH::[phleoR Psdp_mCherry] bglS::[kanR PspoIIE_gfp]*. All strains were confirmed by PCR analysis using primers flanking the integration sites, followed by sequencing to verify accurate integration. The resulting strains were stored at -80°C in 15% glycerol until further use.

Sporulation assay

For sporulation assays under constant RapA and Spo0E levels, strains were grown in LB medium with kanamycin (10 µg/mL) and IPTG (1 mM) overnight at 30ºC with shaking at 200 rpm. Fresh cultures were started at an OD_{600} of 0.015 by dilution in LB medium with kanamycin (10 µg/mL) and different concentrations of IPTG (from 0 to 500 µM). Cultures were grown for 3 h at 37ºC and 220 rpm, or until they reached an OD_{600} of 0.3. Then, cultures were washed twice in buffer Tris-HCl (10 mM pH 7.4) and resuspended in sporulation medium (SM) with corresponding IPTG concentrations and at a final OD_{600} of 0.1. For each strain and IPTG concentration, 200 µL of each condition was loaded in a 96-well plate with black walls and a clear bottom. Cells were incubated and monitored in a Varioskan (Thermo) with a regime of 30 s of shaking every 4.5 min at 480 rpm, 37ºC. Fluorescence (λ_{ex} = 481 nm, λ_{em} = 507 nm) and OD (450 nm) were continuously monitored. Assays were performed in triplicate.

For assays with either RapA or Spo0E induction after sporulation pathway activation, a similar assay was conducted in the absence of IPTG. For this, 180 µL of cultures at an OD of 0.1 in SM medium were inoculated into the 96-well plate, then incubated and monitored using the Varioskan until GFP signal activation was detected and incubated for 30 min. Then, phosphatase expression was induced by adding 20 µL of SM medium with different IPTG concentrations (10-fold relative to the desired final concentration) and cells were monitored under the same conditions.

Kinetic model and model fitting

Kinetic models were constructed as a set of differential equations based on mass action kinetics using custom scripts in Matlab R2020a (MathWorks Inc.). The model equations were solved numerically using the ode15s solver. Parameters were globally optimized by minimizing the sum of squared residuals (SSRs) of the dataset with the simplex search method as implemented in fminsearch with boundaries [50].

The kinetic model of the sporulation pathway of figure 3 assumes KinA as constitutively active HK. Additional modulators, like Sda, or yet other dedicated phosphatases besides RapA or Spo0E, were not included. We considered cellular quantification and dynamics measured by Eswaramoorthy et al. to set the starting levels and the ratio between synthesis and degradation for each phosphorelay component [25]. Kinetic constants describing phosphoryl-transfer reactions and binding between proteins were taken from Narula et al. [18] and fitting kinetic models to phosphoryl-transfer titration experimental data (S8 Fig). All parameters are publicly available in a Zenodo repository under the following DOI: https://doi.org/10.5281/zenodo.15869743. Reporter expression in Fig. 3 and 4 was modelled using an activation Hill function dependent on P~Spo0A concentration.

References

1.   Wuichet K, Cantwell BJ, Zhulin IB. Evolution and phyletic distribution of two-component signal transduction systems. Curr Opin Microbiol. 2010;13(2):219-25. Epub 20100203. doi: 10.1016/j.mib.2009.12.011. PubMed PMID: 20133179; PubMed Central PMCID: PMCPMC3391504.

2.   Buschiazzo A, Trajtenberg F. Two-Component Sensing and Regulation: How Do Histidine Kinases Talk with Response Regulators at the Molecular Level? Annu Rev Microbiol. 2019;73:22.1-. doi: 10.1146/annurev-micro-091018- 054627.

3.   Stock JB, Stock AM, Mottonen JM. Signal transduction in bacteria. Nature.
1990;344(6265):395-400. Epub 1990/03/29. doi: 10.1038/344395a0. PubMed PMID: 2157156.
4.   Lima S, Blanco J, Olivieri F, Imelio JA, Nieves M, Carrion F, et al. An allosteric switch
ensures efficient unidirectional information transmission by the histidine kinase DesK from
Bacillus subtilis. Sci Signal. 2023;16(769):eabo7588. Epub 20230124. doi:
10.1126/scisignal.abo7588. PubMed PMID: 36693130.
5.   Trajtenberg F, Imelio JA, Machado MR, Larrieux N, Marti MA, Obal G, et al. Regulation
of signaling directionality revealed by 3D snapshots of a kinase:regulator complex in action. eLife.
2016;5. doi: 10.7554/eLife.21422. PubMed PMID: 27938660; PubMed Central PMCID:
PMC5231405.
6.   Burbulys D, Trach KA, Hoch JA. Initiation of sporulation in B. subtilis is controlled by a
multicomponent phosphorelay. Cell. 1991;64(3):545-52. PubMed PMID: 1846779.
7.   Kieber JJ, Schaller GE. Cytokinin signaling in plant development. Development.
2018;145(4). Epub 20180227. doi: 10.1242/dev.149344. PubMed PMID: 29487105.
8.   Setlow P. Germination of spores of Bacillus species: what we know and do not know.
Journal of bacteriology. 2014;196(7):1297-305. Epub 20140131. doi: 10.1128/JB.01455-13.
PubMed PMID: 24488313; PubMed Central PMCID: PMCPMC3993344.
9.   Jiang M, Shao W, Perego M, Hoch JA. Multiple histidine kinases regulate entry into
stationary phase and sporulation in Bacillus subtilis. Mol Microbiol. 2000;38(3):535-42. doi:
10.1046/j.1365-2958.2000.02148.x. PubMed PMID: 11069677.
10.   Perego M. Kinase-phosphatase competition regulates Bacillus subtilis development.
Trends Microbiol. 1998;6(9):366-70. doi: 10.1016/s0966-842x(98)01350-x. PubMed PMID:
9778730.
11.   Stephenson SJ, Perego M. Interaction surface of the Spo0A response regulator with the
Spo0E phosphatase. Mol Microbiol. 2002;44(6):1455-67. doi: 10.1046/j.1365-
2958.2002.02974.x. PubMed PMID: 12067336.
12.   Hamon MA, Lazazzera BA. The sporulation transcription factor Spo0A is required for
biofilm development in Bacillus subtilis. Mol Microbiol. 2001;42(5):1199-209. doi:
10.1046/j.1365-2958.2001.02709.x. PubMed PMID: 11886552.
13.   Gonzalez-Pastor JE, Hobbs EC, Losick R. Cannibalism by sporulating bacteria. Science.
2003;301(5632):510-3. Epub 20030619. doi: 10.1126/science.1086462. PubMed PMID:
12817086.
14.   Schultz D, Wolynes PG, Ben Jacob E, Onuchic JN. Deciding fate in adverse times:
sporulation and competence in Bacillus subtilis. Proceedings of the National Academy of Sciences
of the United States of America. 2009;106(50):21027-34. Epub 20091207. doi:
10.1073/pnas.0912185106. PubMed PMID: 19995980; PubMed Central PMCID:
PMCPMC2795487.
15.   Fujita M, Losick R. Evidence that entry into sporulation in Bacillus subtilis is governed by
a gradual increase in the level and activity of the master regulator Spo0A. Genes Dev.
2005;19(18):2236-44. doi: 10.1101/gad.1335705. PubMed PMID: 16166384; PubMed Central
PMCID: PMCPMC1221893.
16.   Fujita M, Gonzalez-Pastor JE, Losick R. High- and low-threshold genes in the Spo0A
regulon of Bacillus subtilis. Journal of bacteriology. 2005;187(4):1357-68. doi:
10.1128/JB.187.4.1357-1368.2005. PubMed PMID: 15687200; PubMed Central PMCID:
PMCPMC545642.

17.   Chastanet A, Vitkup D, Yuan GC, Norman TM, Liu JS, Losick RM. Broadly heterogeneous activation of the master regulator for sporulation in Bacillus subtilis. Proceedings of the National Academy of Sciences of the United States of America. 2010;107(18):8486-91. Epub 20100419. doi: 10.1073/pnas.1002499107. PubMed PMID: 20404177; PubMed Central PMCID: PMCPMC2889527.

18.   Narula J, Kuchina A, Lee DD, Fujita M, Suel GM, Igoshin OA. Chromosomal Arrangement of Phosphorelay Genes Couples Sporulation and DNA Replication. Cell. 2015;162(2):328-37. Epub 20150709. doi: 10.1016/j.cell.2015.06.012. PubMed PMID: 26165942; PubMed Central PMCID: PMCPMC4506695.

19.   Willett JW, Herrou J, Briegel A, Rotskoff G, Crosson S. Structural asymmetry in a conserved signaling system that regulates division, replication, and virulence of an intracellular pathogen. Proceedings of the National Academy of Sciences of the United States of America. 2015;112(28):E3709-18. doi: 10.1073/pnas.1503118112. PubMed PMID: 26124143; PubMed Central PMCID: PMC4507200.

20.   Casino P, Rubio V, Marina A. Structural insight into partner specificity and phosphoryl transfer in two-component signal transduction. Cell. 2009;139(2):325-36. Epub 20091001. doi: 10.1016/j.cell.2009.08.032. PubMed PMID: 19800110.

21.   Zapf J, Sen U, Madhusudan, Hoch JA, Varughese KI. A transient interaction between two phosphorelay proteins trapped in a crystal lattice reveals the mechanism of molecular recognition and phosphotransfer in signal transduction. Structure. 2000;8(8):851-62. PubMed PMID: 10997904.

22.   Xu Q, Porter SW, West AH. The Yeast YPD1/SLN1 Complex. Structure. 2003;11(12):1569-81. doi: 10.1016/j.str.2003.10.016.

23.   Yamada S, Sugimoto H, Kobayashi M, Ohno A, Nakamura H, Shiro Y. Structure of PAS-linked histidine kinase and the response regulator complex. Structure. 2009;17(10):1333-44. doi: 10.1016/j.str.2009.07.016. PubMed PMID: 19836334.

24.   Davidson P, Eutsey R, Redler B, Hiller NL, Laub MT, Durand D. Flexibility and constraint: Evolutionary remodeling of the sporulation initiation pathway in Firmicutes. PLoS Genet. 2018;14(9):e1007470. Epub 20180913. doi: 10.1371/journal.pgen.1007470. PubMed PMID: 30212463; PubMed Central PMCID: PMCPMC6136694.

25.   Eswaramoorthy P, Dinh J, Duan D, Igoshin OA, Fujita M. Single-cell measurement of the levels and distributions of the phosphorelay components in a population of sporulating Bacillus subtilis cells. Microbiology (Reading). 2010;156(Pt 8):2294-304. Epub 20100422. doi: 10.1099/mic.0.038497-0. PubMed PMID: 20413551.

26.   Levine JH, Fontes ME, Dworkin J, Elowitz MB. Pulsed feedback defers cellular differentiation. PLoS biology. 2012;10(1):e1001252. Epub 20120131. doi: 10.1371/journal.pbio.1001252. PubMed PMID: 22303282; PubMed Central PMCID: PMCPMC3269414.

27.   Russell JR, Cabeen MT, Wiggins PA, Paulsson J, Losick R. Noise in a phosphorelay drives stochastic entry into sporulation in Bacillus subtilis. The EMBO journal. 2017;36(19):2856-69.

28.   Bischofs IB, Hug JA, Liu AW, Wolf DM, Arkin AP. Complexity in bacterial cell-cell communication: quorum signal integration and subpopulation signaling in the Bacillus subtilis phosphorelay. Proc Natl Acad Sci U S A. 2009;106(16):6459-64. Epub 2009/04/22. doi: 10.1073/pnas.0810878106. PubMed PMID: 19380751; PubMed Central PMCID: PMCPMC2672556.

29.   Lopez D, Kolter R. Extracellular signals that define distinct and coexisting cell fates in Bacillus subtilis. FEMS Microbiol Rev. 2010;34(2):134-49. Epub 20091123. doi: 10.1111/j.1574-6976.2009.00199.x. PubMed PMID: 20030732.

30.   Chen Z, Srivastava P, Zarazua-Osorio B, Marathe A, Fujita M, Igoshin OA. Bacillus subtilis Histidine Kinase KinC Activates Biofilm Formation by Controlling Heterogeneity of Single-Cell Responses. mBio. 2022;13(1):e0169421. Epub 20220111. doi: 10.1128/mbio.01694-21. PubMed PMID: 35012345; PubMed Central PMCID: PMCPMC8749435.

31.   Kennedy EN, Barr SA, Liu X, Vass LR, Liu Y, Xie Z, et al. Azorhizobium caulinodans chemotaxis is controlled by an unusual phosphorelay network. Journal of bacteriology. 2021:JB0052721. Epub 20211129. doi: 10.1128/JB.00527-21. PubMed PMID: 34843377.

32.   Sourjik V, Schmitt R. Phosphotransfer between CheA, CheY1, and CheY2 in the chemotaxis signal transduction chain of Rhizobium meliloti. Biochemistry. 1998;37(8):2327-35. doi: 10.1021/bi972330a. PubMed PMID: 9485379.

33.   Amin M, Kothamachu VB, Feliu E, Scharf BE, Porter SL, Soyer OS. Phosphate sink containing two-component signaling systems as tunable threshold devices. PLoS Comput Biol. 2014;10(10):e1003890. Epub 20141030. doi: 10.1371/journal.pcbi.1003890. PubMed PMID: 25357192; PubMed Central PMCID: PMCPMC4214558.

34.   Skerker JM, Perchuk BS, Siryaporn A, Lubin EA, Ashenberg O, Goulian M, et al. Rewiring the specificity of two-component signal transduction systems. Cell. 2008;133(6):1043-54. doi: 10.1016/j.cell.2008.04.040. PubMed PMID: 18555780; PubMed Central PMCID: PMC2453690.

35.   Lima S, Blanco J, Olivieri F, Imelio JA, Carrión F, Alvarez B, et al. Molecular basis of unidirectional information transmission in two-component systems: lessons from the DesK-DesR thermosensor. bioRxiv. 2021:2021.11.12.468217. doi: 10.1101/2021.11.12.468217.

36.   Unger T, Jacobovitch Y, Dantes A, Bernheim R, Peleg Y. Applications of the Restriction Free (RF) cloning procedure for molecular manipulations and protein expression. Journal of structural biology. 2010;172(1):34-44. doi: 10.1016/j.jsb.2010.06.016. PubMed PMID: 20600952.

37.   Trajtenberg F, Grana M, Ruetalo N, Botti H, Buschiazzo A. Structural and enzymatic insights into the ATP binding and autophosphorylation mechanism of a sensor histidine kinase. J Biol Chem. 2010;285(32):24892-903. doi: 10.1074/jbc.M110.147843. PubMed PMID: 20507988; PubMed Central PMCID: PMC2915725.

38.   Rizza JD, Ortega C, Carrion F, Flo M, Correa A. Production, purification and characterization of a double-tagged TEV protease. Protein Expr Purif. 2022;191:106021. Epub 20211116. doi: 10.1016/j.pep.2021.106021. PubMed PMID: 34798273.

39.   Winter G. xia2: an expert system for macromolecular crystallography data reduction. Journal of Applied Crystallography. 2010;43(1):186-90. doi: doi:10.1107/S0021889809045701.

40.   Vonrhein C, Flensburg C, Keller P, Sharff A, Smart O, Paciorek W, et al. Data processing and analysis with the autoPROC toolbox. Acta Crystallogr D Biol Crystallogr. 2011;67(Pt 4):293-302. doi: 10.1107/S0907444911007773. PubMed PMID: 21460447; PubMed Central PMCID: PMCPMC3069744.

41.   McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. Phaser crystallographic software. J Appl Crystallogr. 2007;40(Pt 4):658-74. Epub 2007/08/01. doi: 10.1107/S0021889807021206. PubMed PMID: 19461840; PubMed Central PMCID: PMC2483472.

42.   Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. Acta Crystallogr D Biol Crystallogr. 2010;66(Pt 4):486-501. Epub 2010/04/13. doi:

10.1107/S0907444910007493. PubMed PMID: 20383002; PubMed Central PMCID: PMC2852313.

43.   Bricogne G, Blanc E, Brandl M, Flensburg C, Keller P, Paciorek W, et al. BUSTER. 2.8.0. ed. Cambridge: Global Phasing Ltd. United Kingdom; 2009.

44.   Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, et al. Towards automated crystallographic structure refinement with phenix.refine. Acta Crystallogr D Biol Crystallogr. 2012;68(Pt 4):352-67. Epub 20120316. doi: 10.1107/S0907444912001308. PubMed PMID: 22505256; PubMed Central PMCID: PMCPMC3322595.

45.   Chen VB, Arendall WB, 3rd, Headd JJ, Keedy DA, Immormino RM, Kapral GJ, et al. MolProbity: all-atom structure validation for macromolecular crystallography. Acta Crystallogr D Biol Crystallogr. 2010;66(Pt 1):12-21. doi: 10.1107/S0907444909042073. PubMed PMID: 20057044; PubMed Central PMCID: PMCPMC2803126.

46.   Morin A, Eisenbraun B, Key J, Sanschagrin PC, Timony MA, Ottaviano M, et al. Collaboration gets the most out of software. Elife. 2013;2:e01456. Epub 2013/09/17. doi: 10.7554/eLife.01456. PubMed PMID: 24040512; PubMed Central PMCID: PMCPMC3771563.

47.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nature methods. 2012;9(7):671-5. PubMed PMID: 22930834.

48.   Babel H, Naranjo-Meneses P, Trauth S, Schulmeister S, Malengo G, Sourjik V, et al. Ratiometric population sensing by a pump-probe signaling system in Bacillus subtilis. Nat Commun. 2020;11(1):1176. Epub 2020/03/07. doi: 10.1038/s41467-020-14840-w. PubMed PMID: 32132526; PubMed Central PMCID: PMCPMC7055314.

49.   Mutlu A, Trauth S, Ziesack M, Nagler K, Bergeest JP, Rohr K, et al. Phenotypic memory in Bacillus subtilis links dormancy entry and exit by a spore quantity-quality tradeoff. Nat Commun. 2018;9(1):69. Epub 2018/01/06. doi: 10.1038/s41467-017-02477-1. PubMed PMID: 29302032; PubMed Central PMCID: PMCPMC5754360.

50.   D'Errico J. fminsearchbnd, fminsearchcon. MATLAB Central File Exchange, (2021). 2021;https://mathworks.com/matlabcentral/fileexchange/8277-fminsearchbnd-fminsearchcon.

51.   Helliwell JR. Error estimates in atom coordinates and B factors in macromolecular crystallography. Curr Res Struct Biol. 2023;6:100111. Epub 20231115. doi: 10.1016/j.crstbi.2023.100111. PubMed PMID: 38058355; PubMed Central PMCID: PMCPMC10695842.

Acknowledgments:

The authors thank Dayana Benchoam and Victor de Lorenzo for helpful discussions and comments. We thank Stephanie Trauth for cloning the pMAD_Spo0E plasmid as part of project ERC-StG 260860. We acknowledge computational and storage services (Maestro cluster) provided by the Institut Pasteur (Paris). AB and FT are associated researchers from PEDECIBA (Programa de Desarrollo de las Ciencias Básicas). IBB is a member of the DFG priority program SPP2389 (BI1213/6-1) and the Marburg cluster of excellence M4C.

Funding:

Agencia Nacional de Investigación e Innovación grants FCE_1_2017_1_13629 and

FCE_1_2021_1_166888

DAAD fellowship

Max Planck Society.

Competing interests: FT is the founder and holds equity in Locbio. Locbio is developing
technologies related to generating biosensors and live biotherapeutics. The other authors declare
no competing interests.

Data and materials availability: Code and models used to generate Figure 3 are publicly
available in the Zenodo repository under the following
DOI: https://doi.org/10.5281/zenodo.15869743. All data are available in the main text or the
supplementary materials.
619

Fig. 1. Crystal structure of the Spo0B:Spo0A complex. A) Ribbon representation of the complex. Two protomers of Spo0B (depicted in green and yellow) interact with two receiver domains of Spo0A (pink and magenta). B) Structural alignment of Spo0B:Spo0A and Spo0F:Spo0B complexes, using the RR-binding site of Spo0B as reference (Spo0B residues 30 to 67). Spo0F is shown as white ribbon. Red arrows indicate the composite movement that relates the position of Spo0F and Spo0A in the complexes. C) Close-up of the active sites showing the distance between His and Asp within the complexes. The distance between the reactive His and Asp in the Spo0B:Spo0A complex is 6.6 Å ± 0.2 (standard deviation estimated using per atom-corrected DPI [51]), whereas in the Spo0F complex the distance is 5.0 ± 0.5 Å. D) Protein:protein interacting surfaces between Spo0B and Spo0F (white) or Spo0A (magenta) is shown. Selected set of residue pair interactions are depicted as sticks. E) Sequence conservation logo of the β1 α1 loop and helix α1 region of Spo0F and Spo0A. F) The degree of phosphorylation (ξ_p) was calculated as the relative amount of RR~P over all phosphorylated proteins (RR~p+Spo0B~P) as obtained by densitometric quantification of phosphotransfer titration assays of Spo0F (blue) and Spo0A (red), respectively. Circles show mean and standard deviation from three independent experiments. Lines represent fits of the data to a Hill function. Green dashed lines indicate the reactions in equimolar conditions. The thin black line represents the expected behavior in a fully reversible equilibrium.

Fig. 2. Mutant grafting and phosphotransfer directionality analyses. A) Level of phosphorylation of RR
variants (20 μM) after reaching equilibrium with P~Spo0B (15 μM), normalized against wild type Spo0F
(F0) or Spo0A (A0). Reactions were analyzed by autoradiography. Spo0F wild type (F0) and its variants
are shown in blue, whereas Spo0A wild type (A0) and its variants are depicted in red. B) Phosphotransfer
equilibrium titration experiments for wild type Spo0F (F0) and its variants F3 and F6. P~Spo0B (15μM)
was incubated with varying concentrations of the RR (from 0 to 50 μM) and analyzed after reaching
equilibrium. Phosphorylation levels were determined (circles) and Hill functions were fitted to the data
obtained (solid lines).

Fig. 3: Directionality in phosphoryl-transfer affects signal integration and information storage. A) Schematic representation of the ordinary differential equations included in the kinetic model. Red arrows indicate phosphoryl-transfer reactions. Dashed arrows depict transcriptional activation and are modelled as an activating Hill function dependent on P~Spo0A concentration. B) Left panel shows the simulations of the sporulation system under activating conditions with wild type spo0F and under different steady state RapA concentrations. The reporter concentration is plotted as a function of time. The parameters that describe the phosphotransfer reactions were approximated to describe the observed displacements of the equilibria according to the titration experiments. Right panel are the time courses of sporulation assays in

engineered B. subtilis strains expressing wild type Spo0F (strain BSS117). Sporulation was induced by
incubating cells in poor media under different IPTG concentrations (setting distinct RapA intracellular
levels). Thick lines represent the mean GFP/DO ratio, and the shaded region indicates the one standard
deviation. C) Similar to B but using Spo0F_K56I variant (left simulations and right the sporulation assay
with strain BSS118). Note that reporter levels in strain carrying wild type Spo0F increase even at the highest
IPTG concentrations, while in these conditions, the pathway remains turned off in the strain with
spo0F_K56I mutant. D) Similar to B, but RapA was induced (indicated with red arrows) after the system
showed signs of activation.　In the right panel IPTG was added at different concentrations after
approximately 5 hours of incubation. E) Similar to D but using the BSS118 strain with the spo0F_K56I
mutant.
667

Figure 4. Information recording, storage and reset in the B. subtilis sporulation regulation pathway.
P~Spo0A accumulation time courses obtained by kinetic simulations in systems with fully reversible
equilibria (red line) or with unidirectional phosphoryl-transfer transmission (blue line). During the first 60
min, the reactions proceed in the absence of phosphatases, with KinA activity recording information as
P~Spo0A. When a pulse of RapA phosphatase (green bar) was included, the information was stored only
in the unidirectional system, establishing a short-term memory. A pulse of Spo0E phosphatase (magenta
bar), after a second recording phase, resulted in a reset of the information.
676

Fig. S1. TCS phosphoryl-transfer directionality

Fig. S2: Sporulation phosphorelay pathway. Different signals are integrated through the action of
multiple HKs (KinA to KinE) that funnel the information to the RR Spo0F. Information (in the form of a
phosphoryl group) is then transmitted to the effector RR Spo0A through the phosphotransferase Spo0B.
Dedicated Spo0F- and Spo0A-specific phosphatases are shown (Raps and Spo0E-like phosphatases,
respectively). Red arrows indicate phosphoryl-transfer reactions. Dashed arrows represent the effect of
transcriptional activation exerted by phosphorylated Spo0A on the other components of the pathway.

Fig. S3: High structural similarity between Spo0F (from PDB Id 2FTK) and Spo0A.

Fig. S4; Phosphoryl-transfer equilibria in the B. subtilis sporulation phosphorelay pathway. A and B) Representative gel of a phosphotransfer titration assay between phosphorylated phosphotransferase Spo0B (P~Spo0B) and the RRs Spo0F (A) or Spo0A (B) at equilibrium. Reactions were carried out with varying concentrations of RR from 0 to 50 µM at fixed level of Spo0B (15 µM) at pH 8 and 25 °C and analysed by SDS-PAGE and autoradiography.

		1	2	3	4	5
1.	Spo0F G14E					
2.	Spo0F G14E+Y13R+R16V+A34Y					
3.	Spo0F			+K56I		
4.	Spo0F			+K56I+E86Q		
5.	Spo0F G14E+Y13R+R16V+A34Y			+E86Q		
6.	Spo0F G14E+Y13R+R16V+A34Y			+E86H		
7.	Spo0F G14E+Y13R+R16V+A34Y		+K56I+E86Q			
8.	Spo0F G14E+Y13R+R16V+A34Y		+K56I+E86Q			+L37Q+Q38E+D41S+G59H
9.	Spo0F G14E+Y13R+R16V+A34Y+Y84F+L87E			+V22Y+I108M+D109E+R112V+D113G		
10.	Spo0F G14E+Y13R+R16V+A34Y+Y84F+L87E+K56I+E86Q+V22Y+I108M+D109E+R112V+D113G+L37Q+Q38E+D41S+G59H					
11.	Spo0F G14E+Y13R+R16V+A34Y+Y84F+L87E+K56I+E86Q+V22Y+I108M+D109E+R112V+D113G+L37Q+Q38E+D41S+G59Y					

Fig. S5: Grafting of motifs from Spo0A to Spo0F. A) View of the binding interface of Spo0F or Spo0A, to Spo0B. Spo0F and Spo0A are depicted in solvent-accessible surface representation. The surface is coloured according to the mapping of electrostatic potential (red=negative, blue=positive). Green patches indicate the selected sectors to graft motifs among different mutants. Table including all the substitutions included in Spo0F variants.

Fig. S6: Resetting of the sporulation phosphorelay by phosphatase activity acting at the level of
Spo0A. (A) Simulations of the sporulation system under activating conditions with wild type Spo0F, and
under different steady state Spo0E concentrations, are shown. The reporter concentration is plotted as a
function of time. (B) Time courses of sporulation in a *B. subtilis* strain expressing wild type Spo0F.
Sporulation was induced by incubating cells in nutrient-deprived media under different IPTG
concentrations (determining different Spo0E intracellular concentrations). Thick lines represent mean
GFP/DO ratios, and the shaded region indicates one standard deviation
713

Fig. S7. A Rheostat-like mechanism regulates Spo0A phosphorylation, leading to more graded responses. (A) Schematic representation of a rheostat-like regulatory mechanism for a signaling pathway. Simulations predicting the rate of Spo0A phosphorylation as a function of KinA and RapA concentrations, under unidirectional (B) or reversible (C) regimes. To visualize a clearer picture, these simulations were based on simplified models, where KinA was assumed to be constitutively active, and feedback loops regulating different sporulation components were not considered. The simulations were performed for 30 min. The colour ramp, from blue to red, indicates the concentration of phosphorylated Spo0A at the end of the simulation.

Parameter	Fitted value	Lower bound	Upper bound	Parameter	Fitted value	Lower bound	Upper bound	Parameter	Fitted value	Lower bound	Upper bound	Parameter	Fitted value	Lower bound	Upper bound
KeqOF	2,945	1,775	4,115	KeqOF	1,782	1,373	2,191	KeqOF	4,010	3,350	4,707	KeqOF	3,591	3,243	3,939
KeqOA	0,353	0,213	0,492	KeqOA	0,666	0,345	0,988	KeqOA	0,915	0,748	1,084	KeqOA	1,340	1,317	1,364
[OBp]i	15,000	10,632	19,368	[OBp]i	14,851	14,850	14,852	[OBp]i	14,910	14,900	14,912	[OBp]i	7,500	7,416	7,584
KdOBA	100,000	-553,286	753,286	Kddim	0,510	-0,206	1,226	Kddim	0,100	-0,063	0,263	Kddim	0,145	0,074	0,216
SSE	0,801			KdOBA	98,125	-176,845	373,095	KdOBA	2,214	-0,456	4,881	KdOBA	0,379	0,367	0,392
				SSE	0,613			SSE	0,271			SSE	0,169		

Fig. S8: Fitting of kinetic models to phosphoryl-transfer titration experimental data. Each model was constructed as a custom script in MATLAB R2020a (MathWorks Inc.), consisting of a set of differential equations that describe each reaction. Continuous traces (red, magenta and blue lines) in (A) to (D) depict the prediction of the best-fitted model. The black continuous line shows the expected behaviour under a fully reversible equilibrium regime. Green lines indicate the reactions in equimolar concentrations.

731   **Table S1: X ray diffraction data collection and refinement statistics.**

Data collection	Spo0B:Spo0A$_{REC}$	Spo0B:Spo0A$_{REC}$
Space group	P2$_1$2$_1$2$_1$	C2$_1$
Cell dimensions		
a, b, c (Å)	76.361 125.504 149.472	221.240 103.145 66.461
α, β, γ (°)	-	90.0 96.91 90.0
Wavelength (Å)	0.97622	1.54179
Resolution (Å)	76.36 - 2.05 (2.08 - 2.05)	28.72 - 3.50 (3.58- 3.50)
Unique reflections	90767 (4368)	18675 (859)
R$_{meas}$	0.071 (1.506)	0.143 (1.121)
R$_{pim}$	0.028 (0.667)	0.073 (0.580)
CC$_{½}$	0.999 (0.466)	0.998 (0.626)
I/σI	11.15 (1.2)	8.9 (1.5)
Completeness	100 (97.7)	91.3 (69.0)
Redundancy	6.2 (5.0)	3.8 (3.5)
Refinement		
Resolution (Å)	74.74 – 2.05 (2,06 – 2.05)	24.02 - 3.50 (3.53 – 3.50)
Number of refls used (N in the free set)	4472 (85)	915 (21)
R$_{work}$ / R$_{free}$	0.2235 / 0.2461	0.2426/0.292
Number of atoms	10209	6389
Protein	9749	6389
Ligands + ions	66 (GOL)	-
Water	394	-
B factors (Å^2)		
Wilson plot	49.05	100.7
Mean (overall)	70.83	125.9
R.m.s. deviations		
Bond lengths (Å)	0.008	0.010
Bond angles (°)	0.84	1.378
Number of residues in Ramachandran plot[§] (favored/allowed/outliers)	1188/19/1	749/27/0
PDB ID	pdb 00009PI0	pdb 00009PHZ

732
733
734

735 **Table S2: Spo0F and Spo0A variant tested**

F0	Spo0F wild type
F1	Spo0F G14E
F2	Spo0F G14E+Y13R+R16V+A34Y
F3	Spo0F+K56I
F4	Spo0F+K56I+E86Q
F5	Spo0F G14E+Y13R+R16V+A34Y+E86Q
F6	Spo0F G14E+Y13R+R16V+A34Y+E86H
F7	Spo0F G14E+Y13R+R16V+A34Y+K56I+E86H
F8	Spo0F G14E+Y13R+R16V+A34Y+K56I+E86Q+L37Q+Q38E+D41S+G59H
F9	Spo0F G14E+Y13R+R16V+A34Y+Y84F+L87E+V22Y+I108M+D109E+R112V+D113G
F10	Spo0F G14E+Y13R+R16V+A34Y+Y84F+L87E+K56I+E86Q+V22Y+I108M+D109E+R112V+D113G+L37Q+Q38E+D41S+G59H
F11	Spo0F G14E+Y13R+R16V+A34Y+Y84F+L87E+K56I+E86Q+V22Y+I108M+D109E+R112V+D113G+L37Q+Q38E+D41S+G59Y
A0	Spo0A wild type
A1	Spo0A+Y108A+H122A
A2	Spo0A+E14G
A3	Receiver domain of Spo0A

737 **Table S3. Plasmids used in this work**

Name	Source
pMAD_spo0E (EIB214)	This work
pMAD_spo0F (EIB221)	Babel et al.
pMAD_Spo0F-K56I	This work
pDR111_rapA (EIB297)	Mutlu et al.
pPspoiiE_GFP_BglS	This work
pRB117_Pspo0F_mCherry	This work
pET-28b_Spo0F_F0	This work
pET-28b_Spo0F_F1	This work
pET-28b_Spo0F_F2	This work
pET-28b_Spo0F_F3	This work
pET-28b_Spo0F_F4	This work
pET-28b_Spo0F_F5	This work
pET-28b_Spo0F_F6	This work
pET-28b_Spo0F_F8	This work
pET-28b_Spo0F_F9	This work
pET-28b_Spo0F_F10	This work
pET-28b_Spo0F_F11	This work
pQE80_Spo0A_A0	This work
pQE80_Spo0A_A1	This work
pQE80_Spo0A_A2	This work
pQE80_Spo0A_A3	This work
pQE80_Spo0A	This work
pETM_KinA	This work
pQE80_Spo0B	This work
pET_Spo0B_strep	This work

739 **Table S4. Primers used in this work**

Name	Sequence (5' -> 3')
KinA_F	CTGGAAGTTCTGTTCCAGGGGCCCACAAAAACCGAT ATTCATGCAGTC
KinA_R	GGCTTTGTTAGCAGCCGGATCTCACTTTTTTGGAAAT GAAATTTTAAACGCTGTG
Spo0A_F	CCTGTATTTTCAGGGATCCGGTTCAATGGAGAAAATT AAAGTTTGTGTTGC
Spo0A_R	GCTGCAGGTCGACGCCAAGATCCTTTTTTAAGAAGCC TTATGCTCTAACC
Spo0A_REC_R	GCTGCAGGTCGACGCCAAGATCCTTTTTTAGCTGGCA TTTCCGCTGACCTG
spo0E_up_F	CCAGTCGACGAGCATCACCATACAGACGATGAGG
spo0E_up_R	GCATCTGATAGGCCGCTATTTGCCCATTCACTTCACCC CTCG
Spo0E_down_F	AAATAGCGGCCTATCAGATGC
spo0E_down_R	CATAGATCTACTGACGATCTCCCTAGGAGC
spo0E_up_F (ST94)	CCAGTCGACGAGCATCACCATACAGACGATGAGG
spo0E_up_R (ST95)	GCATCTGATAGGCCGCTATTTGCCCATTCACTTCACCCCTCG
spo0E_down_F (ST96)	AAATAGCGGCCTATCAGATGC
spo0E_down_R (ST97)	CATAGATCTACTGACGATCTCCCTAGGAGC
spo0E_F (ST98)	ATGGGCGGTTCTTCTGAACAAG
spo0E_R (ST99)	CTATTTATTTGCATCATATGCTGG

741 **Table S5. Strains used in this work**

Name		Source
BIB415	*ΔrapA ΔphrA Δspo0F*	Parental strain from Babel et al.
BSS093	*ΔrapA::phrA Δspo0F Δspo0E*	This work
BSS094	*ΔrapA::phrA Δspo0F Δspo0E amyE::[specR Phyperspank_rapA]*	This work
BSS101	*ΔrapA::phrA Δspo0F Δspo0E amyE::[specR Phyperspank_rapA] yhdG_yhdH::[phleoR Psdp_mCherry]*	This work
BSS105	*ΔrapA::phrA Δspo0F Δspo0E amyE::[specR Phyperspank_rapA] yhdG_yhdH::[phleoR Psdp_mCherry] bglS::[kanR PspoIIE_gfp]*	This work
BSS117	*ΔrapA::phrA Δspo0E amyE::[specR Phyperspank_rapA] yhdG_yhdH::[phleoR Psdp_mCherry] bglS::[kanR PspoIIE_gfp]*	This work
BSS118	*ΔrapA::phrA Δspo0E amyE::[specR Phyperspank_rapA] yhdG_yhdH::[phleoR Psdp_mCherry] bglS::[kanR PspoIIE_gfp] spo0F-K56I*	This work
BSS119	*ΔrapA::phrA Δspo0E amyE::[specR Phyperspank_spo0E] yhdG_yhdH::[phleoR Psdp_mCherry] bglS::[kanR PspoIIE_gfp]*	This work