Dual origins for neural cells during development of the *Clytia* planula larva

Antonella Ruggiero*[1], Anna Ferraioli[2]*, Sandra Chevalier*, Pascal Lapébie , Romain Girard[3], Tsuyoshi Momose, Carine Barreau[#] and Evelyn Houliston[#]

 Sorbonne Université, CNRS, Laboratoire de Biologie du Développement de Villefranche-sur-mer (LBDV), 06230 Villefranche-sur-mer, France.

* Equal contributions

[1] Current address: European Molecular Biology Laboratory (EMBL), Meyerhofstraße 1 , 69117, Heidelberg, Germany.

[2] Current address: Michael Sars Centre, University of Bergen, 5008 Bergen, Norway.

[3] Current address: Collège de France - CIRB, UMR 7241 - INSERM U1050, 11 place Marcelin Berthelot, 75231 Paris, France.

[#] Corresponding authors**:** Evelyn Houliston (+33) 493763983, houliston@imev-mer.fr; Carine Barreau: carine.barreau@imev-mer.fr

Abstract

Adult hydrozoan cnidarians undergo extensive tissue turnover, generating neural cell types including nematocytes (stinging cells) from interstitial stem cells (i-cells) expressing stemness proteins such as Piwi and Nanos. The contribution of i-cells during embryogenesis, however, has been unclear. Here we address the origin of neural cells during development of the *Clytia hemisphaerica* 'planula' larva. Marker gene in situ hybridisation revealed that Piwi/Nanos1-expressing cells within the presumptive endoderm of the early gastrula generate a substantial pool of nematoblasts, some of which migrate and differentiate into the planula ectoderm. Neurogenic transcription factor expression showed a markedly distinct pattern, predominantly in a basal layer of the aboral/lateral ectoderm during gastrula stages. Embryo bisection and lineage tracing experiments confirmed that neurosensory and secretory cell types derive from gastrula ectoderm, while nematocytes and at least some ganglionic cells derive from i-cells. Knockdown and inhibitor treatments revealed steps of both neurogenesis and nematogenesis regulated by Wnt-β-catenin. We conclude that two distinct neurogenesis pathways operate in parallel during *Clytia* embryogenesis, one involving aboral ectoderm delamination, and one generating mainly nematocytes from i-cell-like precursors.

Introduction

Neural cells are one of the key animal innovations, and their developmental and evolutionary origins are thus much debated (Arendt et al., 2015; Burkhardt and Jékely, 2021). Cnidarian species, including jellyfish and sea anemones, have a range of neural cell types and nervous systems, which have evolved in parallel to those of their sister clade Bilateria (Bosch et al., 2017; Kelava et al., 2015; Rentzsch et al., 2017). These animals thus provide a valuable perspective on the underlying mechanisms of neurogenesis through comparisons with bilaterian model species. Current knowledge of neurogenesis in cnidarians is founded on extensive studies using adult *Hydra* polyps and during development of the sea anemone *Nematostella vectensis,* members respectively of the two main cnidarian clades Hydrozoa and Anthozoa. Hydra polyps propagate mainly by budding and undergo continuous cell turnover. The epithelia of their two body layers, ectoderm and gastroderm, renew themselves independently, while neural cells, defined broadly to include ganglion and sensory type cnidarian neurons (Martin, 2000) and stinging cells (nematocytes), along with secretory gland cells and germ line stem cells, derive from a distinct population of multipotent stem cells called interstitial cells (i-cells) (Bosch et al., 2010; David, 2012; Frank et al., 2009; Siebert et al., 2019). i-cells are a specific feature of Hydrozoa. They typically have a rounded morphology and relatively large nucleus, and occupy spaces between the basolateral surfaces of the epitheliomuscular cells of the two adult body layers. They express orthologues of many genes associated with totipotency and germ cells, such as Nanos, Piwi, PL10 and Vasa (Genikhovich et al., 2006; Leclère et al., 2012; Rebscher et al., 2008). In the hydra polyp, neuron and nematocyte precursors derived from i-cells migrate into the ectoderm and gastroderm layers to take up their positions along the body axis and complete differentiation. In another well-studied hydrozoan model, *Hydractinia*, single i-cells transplanted between polyps similarly give rise to neurons, nematocytes, gland cells and germ line, but unlike *Hydra* can also generate epithelial cells types (Varley et al., 2023). The i-cell to neuron, to nematocyte and to gland cell developmental trajectories in *Hydra* and *Hydractinia* polyps have been deduced from scRNAseq data (Cazet et al., 2023; Salamanca-Díaz et al., 2025; Siebert et al., 2019; Song et al., 2025) and tracked via expression of neurogenic transcription factors such as SoxBs (Chrysostomou et al., 2022). scRNAseq data have demonstrated equivalent trajectories in *Clytia* adult jellyfish (Chari et al., 2021).

While the i-cell derived pathways for neurogenesis in these adult hydrozoans are well documented, the extent to which they also operate during initial embryonic development remains relatively poorly explored. Histological studies have described i-cells in the larval stage, the planula, largely confined to the endodermal region (Bodo and Bouillon, 1968; Frank et al., 2009; Leclère et al., 2012), although whether they arise first in the endoderm or ectoderm may differ between species (see discussion in (Genikhovich et al., 2006)). Ganglionic-type and sensory-type neurons have been described in planulae, their neurites extending respectively from cell bodies close to the mesoglea or from the basal domain of the cell, to form a network between the two tissue layers of the larva (Attenborough et al., 2019; Martin, 2000; Piraino et al., 2011; Ramon-Mateu et al., 2025). Martin and Thomas provided experimental evidence from studies in hydrozoan planulae from various species that ganglionic cells derive from i-cells in the endoderm (Martin, 1988; Martin, 1991; Martin and Thomas, 1980; Martin and Thomas, 1981; Thomas et al., 1987). These experimental studies also indicated, however, that at least some planula neural types derive directly from the ectodermal layer. Notably, neurosensory cells were identified by electron microscopy in planulae derived from aboral halves of *Clytia* (*Phialidium) gregarium* gastrula-stage embryos, which lack i-cells as well as presumptive gastroderm (Thomas et al., 1987). In contrast, nematocytes and ganglionic neurons were found only in larvae derived from oral halves. Similarly, when i-cells were depleted from developing larvae from the hydrozoan polyp species *Pennaria* by colchicine treatment, planulae were preferentially depleted in nematocytes and ganglionic cells (Martin and Thomas, 1981).

 Unlike hydrozoans, anthozoan cnidarians do not have a distinct i-cell population, although recent studies in *Nematostella* have identified cells with similar characteristics in the gastrodermal mesenteries of the adult (Denner et al., 2024; Miramón-Puértolas et al., 2024), and shown that neural and gland cell types have common progenitors (Steger et al., 2022). During embryo development, neural precursors arise in a salt and pepper fashion within the two epithelial body layers. A first wave of neurogenesis during gastrulation generates sensory cells and ganglion cells in the ectoderm, and in a second wave, during the planula stage, precursors arise similarly in the gastroderm (Kelava et al., 2015; Lemaître et al., 2023; Nakanishi et al., 2012; Rentzsch et al., 2017; Richards and Rentzsch, 2014; Richards and Rentzsch, 2015).

We have addressed the origin of neural cell types during larval development in *Clytia hemisphaerica.* In *Clytia* species, gastrulation proceeds by ingression of presumptive endoderm cells from the oral pole (Byrum, 2001) carrying with them the i-cells (Leclère et al., 2012). We first mapped nematogenesis and neurogenesis by in situ hybridisation using molecular markers. We then tested the contribution of the ectoderm to neurogenesis by gastrula bisection and lineage tracing experiments. Finally we assessed the influence of Wnt-β-catenin signalling on neural development using antisense Morpholino oligonucleotides targeting Wnt3 (Wnt3-MO; Momose et al., 2008) and timed treatments with the pharmacological Wnt-β-catenin pathway inhibitor PKF118-310 (Lepourcelet et al., 2004). Our findings demonstrate that the neural cell types of the *Clytia* planula larva develop during gastrulation by two parallel pathways, one generating sensory neurons along with secretory cells types from a basal layer of aboral/lateral ectoderm, and the other operating within the gastrodermal region to mainly produce nematocytes from i-cell like precursors.

Materials and Methods

Manipulation of Eggs and Embryos

Eggs and embryos were obtained from laboratory-raised Z strain medusae (Leclère et al., 2019) and cultured in 0.2 μm Millipore filtered ASW (Red Sea Salt brand; (Lechable et al., 2020). Medusae were kept on daily dark-light cycles with gametes collected two hours after light. Blastula bisection and blastomere separation were performed using fine Tungsten wire loops in agar-lined petri dishes as described previously (Leclère et al., 2012). PKF118-310, which interferes with the interaction between β-catenin and the conserved transcription factor Tcf/LEF required to regulate gene expression via this pathway (Lepourcelet et al., 2004), was used at 0.8 μM in FASW, diluted from a 15 mM stock immediately before use.

Dendra2 lineage tracing

The coding sequence for Dendra2 (Chudakov et al., 2007) was adapted to the codon usage of *Clytia* (Weissbourd, 2021). A T3 promoter-Dendra2 ORF DNA template was obtained by PCR and mRNA was synthesised using mMessage mMachine™ T3 Transcription Kit (Ambion), followed by poly-A addition using the Poly(A) Tailing Kit (InVitroGen). 0.4 - 1.0μg/μl mRNA was injected into freshly spawned eggs prior to fertilisation using an Eppendorf nanoject injection system as described in detail previously (Uveira et al., 2025), aiming at 3-5% egg volume. Embryos were allowed to develop at 18°C until the 4 or 8 cell stage then cultured 16.5°C overnight (approx 18 hours total) until the early gastrula stage. They were then temporarily mounted between slides and coverslips separated by plasticine spacers compressed to immobilise nearly completely the larvae during photoconversion, performed on a Leica Stellaris confocal microscope using the LasX software FRAP mode with a 10X objective. Each defined region, either oral or aboral, was exposed to 30 seconds scanning of the 405 nm Laser beam, at 100% power. Embryos were then cultured at 17°C in MFSW until planula stage before fixation for immunofluorescence staining.

Fluorescence microscopy

For immunofluorescence, samples were fixed for 2 hours at room temperature in IF fix [0.1 M Hepes (pH 6.9), 50 mM EGTA (pH 7.2), 10 mM MgSO4, 80 mM maltose, 0.2% Triton X-100, 4% paraformaldehyde] before proceeding to three washes in PBS-0.01% Triton X-100 for 10 min or longer, extraction in PBS-0.2% Triton for 10 minutes, two washes in PBS-0.01% Triton X-100, and one rinse in PBS. Primary antibodies were an affinity purified rabbit

polyclonal antibody raised against the peptide GRFamide (gift of Gáspár Jékely) and an anti-LWamide mouse monoclonal antibody (gift of O. Koizumi (Koizumi et al., 2015)). Samples were incubated overnight at 4°C with primary antibodies and/or Alexa488- or Rhodamine-Phalloidin (1:100 from stocks in methanol; Fisher Scientific) to visualise the actin-rich cell cortices, followed by washes in PBS-0.01% Triton X-100 (three times for 10 min or longer) and then incubation in rhodamine, Alexa555 or Alexa647 coupled anti-rabbit or anti-mouse IgGs (Jackson ImmunoResearch, dilution 1:100) for 3 hours at room temperature, with Hoechst dye 33342 (Bisbenzimide from Sigma; 0.3 μg/ml final) included to stain DNA. After washes in 0.01% PBS-Triton X-100 (three times each for 10 min or longer) and a PBS rinse, specimens were mounted in Citifluor mounting medium (Citifluor Ltd) and imaged using Leica SP5, SP8 or Stellaris confocal microscopes.

Gene cloning

In situ hybridisation probes were synthesised from cDNA clones retrieved from our EST collection when available (Chevalier et al., 2006). For the remaining sequences, cDNAs were cloned by PCR using the pGEM-T Easy system (Promega). Accession numbers for the newly identified sequences, and XLOC identifiers (Chari et al., 2021; Leclère et al., 2019) for all genes analysed, are provided in Supplementary File S1.

In situ hybridisation

DIG-labeled and fluorescein-labeled antisense RNA probes for in situ hybridisation were synthesised using Promega T3/T7/Sp6 RNA polymerases, purified using ProbeQuant G-50 MicroColumns (GE Healthcare) and stored at -20°C in 50% formamide. Planulae were fixed for in situ hybridisation in a cold solution of 3.7% formaldehyde and 0.2% glutaraldehyde in PBS on ice for 40 minutes to 2 hours, fixation time depending on the developmental stage. Specimens were then washed with PBST (1X PBS + 0.1% Tween 20), dehydrated stepwise in methanol on ice, and stored in 100% methanol at -20°C. Samples were fixed after culture for 24, 48 or 72 hours post fertilisation (hpf) at 17–18°C.

In situ hybridisation was performed either using the InsituPro robot (Intavis®) as described previously (Lapébie et al., 2014, modified as in Chari et al., 2021) or by hand as described in Sinigaglia et al. (2020).

Results

Nematoblast production from i-cells during gastrulation

We tracked the formation of neural cell types, including nematocytes, during *Clytia* embryogenesis using in situ hybridisation (Figure 1). We targeted genes identified previously as expressed in i-cells and/or during neurogenesis/ nematogenesis by trajectory inference analysis of scRNAseq data from *Clytia* adult medusa (Chari et al., 2021). We included *Piwi* and *Nanos1* probes as markers of i-cells (Leclère et al., 2012). Note that the operational identification of 'i-cells' used in this study includes stem cells along with cells that have already embarked on nematogenesis/neural programmes, as shown by the onset of lineage-specific gene expression (Chari et al., 2021; Denker et al., 2008; Mochizuki et al., 2000; Siebert et al., 2008; Varley et al., 2023). As found previously (Leclère et al., 2012), cells expressing high levels of Nanos1, Piwi and Vasa were detected in the oral pole region at the early gastrula stage, then concentrated within the oral part of the inner gastrodermal region at the parenchymula (P1) stage, and finally spread within this region along the oral-aboral axis of fully formed planuae (P2 and P3 stages; Fig. 1A). Planulae are competent to metamorphose by the third day following fertilisation.

The formation of stinging cells, or nematogenesis, can be separated into two distinct phases as characterised from *Clytia* medusa scRNAseq data (Chari et al., 2021). A 'nematoblast' phase involving construction of the nematocyst capsule is followed by a 'maturation' phase during which a functional nematocil is assembled. These phases have almost non-overlapping transcriptome signatures, so can be distinguished by in situ hybridisation using specific markers (Fig. 1B). Transcripts for the early-expressed nematoblast genes *Znf845* and *Mos3* were strongly detected at the oral pole of the early gastrula embryo (Fig. 1B), consistent with their original identification as orally-expressed early zygotic genes (Lapébie et al., 2014). As they develop, nematoblasts move into the blastocoel along with the mass of mesenchymal presumptive gastrodermal cells. The expression patterns of *Znf845* and *Mos3* are very similar to those of *Piwi* and *Nanos1*, consistent with their coexpression in the 'i-cell' transcriptome-based clusters extracted from *Clytia* medusa and planula scRNAseq datasets (Chari et al., 2021; Ferraioli, 2022). Together these observations suggest that many of the *Piwi*/*Nanos1* expressing cells detected during planula development are early stage nematoblasts rather than stem cells.

Figure 1. Expression of nematogenesis/neurogenesis genes during planula development
In situ hybridisation images of embryos fixed at early gastrula (EG), parenchymula (P1; corresponding to the end of gastrulation; see Kraus et al., 2020), 2-day (P2) and 3-day (P3) settlement-competent planula larvae, detecting the following mRNAs: A) Piwi and Nanos1 as markers of "i-cells" (which under this operational definition includes also early stages of derivative cell types); B) Znf845, Mos3 and Minicollagen 3/4a expressed during the first phase 'nematoblast formation' as indicated by the top arrow, and M14 peptidase and Nematocilin during the second phase (nematocyte maturation; bottom arrow); C) Transcription factors Neurogenin and Hlh6, and neural markers ELAV, PP5 (RFamide precursor) and PP2 (GLWamide precursor). Arrows indicate the extent of ingression of the presumptive endoderm in EG images. Oral pole is at the top in all panels. Scale bars in A, B and C: 100μm.

Expression of *Minicollagen 3/4a* mRNA, which codes for a structural component of the nematocyst capsule (Denker et al., 2008) expressed during nematoblast formation but strongly downregulated during the subsequent phase of nematocyte maturation and migration (Chari et al., 2021; Condamine et al., 2019), was detected from the parenchymula (P1) stage. In the planula, nematoblasts are positioned towards the basal side of the developing gastroderm epithelium, occupying a central "trunk" position along the oral-aboral axis (Fig. 1B; Bodo and Bouillon, 1968).

Despite the presence of abundant nematoblasts in P2 and P3 planula larvae, we detected relatively few cells undergoing nematocyte maturation. mRNA for the M14 peptidase gene, whose expression marks the transition from nematoblast to terminal differentiation, could be detected in some central cells within the gastroderm at P2. We interpret these as a sub-population of the nematoblasts starting to differentiate at this stage. Fewer such cells were observed at P3 suggesting that nematocyte differentiation occurs as a wave. Reciprocally, mRNA for Nematocilin, a component of the nematocil trigger structure and a marker of mature nematocytes, became clearly detectable in by P3 in the ectodermal layer, scattered in the along the trunk, and clustered at the oral pole (Fig. 1B). These expression patterns confirm that nematocyte differentiation accompanies migration into the ectoderm, whilst indicating also that only a small subpopulation of nematoblasts continue development to form mature nematocytes prior to larval metamorphosis.

Overall these profiles of nematogenesis gene expression indicate i) that nematoblasts start to develop at the onset of gastrulation as an abundant Piwi/Nanos positive cell population in the presumptive gastroderm, and ii) that development of most of this nematocyte pool becomes arrested at the end of the nematoblast formation phase.

Mixed expression profiles for neurogenic and neural markers
We similarly tracked by in situ hybridisation the formation of non-nematocyte neural cells (Fig. 1C). We included probes for two transcription factors expressed along the i-cell to neuron trajectory in the scRNAseq analyses from *Clytia* medusae (Chari et al., 2021): the atonal family Hlh6, orthologue to *Nematostella* Ath-like (Richards and Rentzsch, 2014), and Neurogenin. These two transcription factors showed very distinct expression profiles (Fig. 1C). Neurogenin mRNA could be detected in a few scattered cells within the endoderm from 24h, consistent with the development of neural precursors from i-cells. In contrast, Hlh6

showed strong expression much earlier, at the gastrula stage, in cells lining the blastocoel, covering lateral and aboral regions. The expression profile of Hlh6 is reminiscent of that of the SoxB2 family transcription factor Sox10 (Kraus et al., 2020), suggestive of neurogenic identity. Supporting this hypothesis, strong expression of an ELAV family RNA binding protein, commonly used across species as markers of neural identity, was detected in cells with a largely similar distribution. In the *Clytia* planula most neural cells express this ELAV gene, (identifier XLOC_030971; Ramon-Mateu et al, 2025). At the beginning of gastrulation, ELAV positive cells were detected scattered across the inner surface of the blastocoel, stronger expression observed closest to the oral pole first. As gastrulation progressed, ELAV expressing cells accumulated in a corresponding layer between the ingressing endoderm and the ectoderm, as well as within both tissue layers. As reported previously for Sox10 (Kraus et al., 2020), expression of Hlh6 fell sharply after gastrulation.

These expression patterns suggest that neurogenesis occurs by stratification of the ectoderm during late blastula and gastrula stages, involving sequential expression of Sox10, hlh6 and then ELAV in the basal layer. Stratification (or pseudo-stratification) was previously suggested by our study of cell morphogenesis during *Clytia* gastrulation based on confocal and electron microscopy (Kraus et al., 2020). Nuclei were found to appear progressively on the basal side of the ectoderm to line the blastocoel during gastrulation, in parallel with the EMT (Epithelio Mesenchymal Transition)-based ingression of presumptive gastroderm cells from the oral pole. We re-examined confocal z-stacks of gastrula stage embryos stained to highlight cell boundaries and nuclei (Figure 2), constructing maximum-projection images to visualise the emerging layer of cells (coloured orange in Fig. 2D and 2H). This layer mirrors the distribution of Hlh6/ELAV expressing cells at these stages (Fig. 1C), supporting the idea that it is neurogenic. We cannot tell from this analysis whether the stratification involves complete or incomplete cell delamination, as illustrated in Fig. 2A and 2E).

Figure 2. Formation of an inner ectodermal layer during gastrulation
Early (A-D) and mid (E-H) gastrula stage embryos, illustrating stratification of the presumptive ectoderm. Presumptive gastroderm cells are ingressing from the oral pole (top right in both embryos). A and E: Cartoons of the two embryos indicating the planes of confocal sections shown in B,C,F and G. Undulations of the blastocoel roof are observed after fixation and mounting. The position of the putative neurogenic layer is shown in pink, and of the apical nuclei of the ectodermal cells in blue. The insets illustrate stratified organisation in the boxed zones; the pink basal cells may retain thin apical projections reaching the embryo outer surface ('pseudo-stratification'). B,C,F and G: confocal images each from z stacks. Actin-rich cell cortices were stained with rhodamine-phalloidin (magenta) and DNA with Hoechst 33342 (cyan). Distance between planes B and C is 14.8μm, between F and G is 32.2 μm. Arrowheads indicate some examples of the nuclei of putative delaminating cells. bc= blastocoel. Ingressing presumptive endoderm cells are outlined in pink. Apical nuclei of ectodermal folds are outlined in green. D and H: maximum projection images of the same z stacks (5 planes over a total thickness of 9.6μm for D; 5 planes over a total thickness of 32.2μm for H). The layer of basal ectodermal cells is overlaid in orange. Scale bars 25μm.

To detect differentiated neural cells at planula stages we monitored expression of the RFamide neuropeptide precursor PP5 and the GLWamide precursor PP2. RFamides are widely expressed in cnidarian neurons of various types including ganglionic type cells and in putative planula sensory cells in some species (Attenborough et al., 2019; Hayakawa et al., 2007; Piraino et al., 2011; Weissbourd, 2021). GLWamides, involved in mediating the larval settlement response in many species of cnidarian (Takahashi and Hatta, 2011), have been detected in sensory cells distributed at or around the aboral pole in *Hydractinia* and *Clytia*, and in a more widely distributed ganglionic cell population in *Clava* (Gajewski et al., 1998; Piraino et al., 2011; Ramon-Mateu et al., 2025). We found that during development *Clytia* PP2 mRNA is first detected in a few cells at the parenchymula stage, and then more strongly

in P2 and P3 planulae, exclusively in sensory-type neurons in the aboral-pole ectoderm (Fig. 1C). Cells expressing PP5 mRNA were detected in P2 and P3 planulae in ganglionic-type cell bodies in the trunk region of the ectoderm close to the mesoglea (Fig. 1C). These two neuropeptide precursors in *Clytia* thus allow subpopulations of neurons positioned differentially along the body axis with aboral secretory versus ganglionic morphologies to be distinguished.

Ectodermal neurogenesis demonstrated by gastrula bisection experiments
To test whether some neural cell types derive from the gastrula ectoderm during *Clytia* gastrulation we used two approaches. Firstly, we bisected early-mid gastrula stage embryos and fixed them at the planula stages to assess neural cell development by immunofluorescence or in situ hybridisation (Figure 3).

Phalloidin staining confirmed that aboral-derived planulae lacked gastroderm, but also showed that the ectoderm contained differentiated mucous cells and aboral secretory cells, easily distinguishable by their cytoplasm-poor morphology and nuclei positioned respectively at basal or medial positions within the ectoderm (Fig. 3B). Immunofluorescence detected both RFamide and LWamide expressing cells in planulae derived from oral halves, but only LWamide expressing cells in planulae derived from aboral halves (Fig. 3C). This result was confirmed by in situ hybridisation with probes detecting the precursor mRNAs for PP5 (Fig. 3D) and PP2 (Fig. 3E) respectively.

These embryo bisection experiments thus demonstrate clearly that mucous cells, secretory cells and GLWamide sensory neurons, can form from early gastrula ectoderm. They also are consistent with the conclusion that RFamide neurons originate from the oral pole of the gastrula, and specifically from i-cells, although the absence of an oral Wnt signalling centre in aboral-derived embryos could account, at least in part, for the lack of detectable PP5/RFamide expression, an idea developed further below.

Figure 3. Gastrula bisection demonstrates ectodermal origin for some neural cells

A) Cartoon of experimental design. Unmanipulated early-mid gastrulae were allowed to develop for 48-60 hpf. Other embryos were bisected into oral and aboral parts and then allowed to develop in parallel. Orange dots indicate i-cells positioned in the ingressing presumptive endoderm region (grey) at the oral pole. B) Overlaid confocal images of Hoechst staining of nuclei (blue) and Alexa488-Phalloidin staining (yellow). Endogenous GFP in oral and lateral ectoderm is also visible in this channel (Fourrage et al., 2014). Uncut, Oral and Aboral derived planulae from left to right, all fixed on the second day after fertilisation. White arrows show examples of mucous cells whereas white asterisks point out the aboral secretory cells area. C) Confocal images of the same planulae as in B stained with anti-RFa (green) and anti-LWa (magenta). Examples of neurites stained with each antibody are shown with green arrows and magenta arrowheads respectively. D) In situ hybridisation to detect PP5 and PP2 expression in planula fixed at 24 hpf. Arrows show examples of stained cells. Uncut, Oral and Aboral derived planulae from left to right. Oral is top left in all panels. Scale bars 50μm.

Dendra2 tracing supports dual origins for neural cells

To confirm the ectodermal origin of GLWamide neurons and test the origin of RFamide neurons, we performed lineage tracing using the photoconvertible fluorescent protein Dendra2 (Chudakov et al., 2007). We injected mRNA coding for Dendra2 into eggs prior to fertilisation and photoconverted the protein from green to red in patches of cells at either the aboral or oral poles of the embryo at early-mid gastrula stages using the 405nm laser on a confocal microscope. The embryos were then cultured to the planula stage (P2) and the identity of Dendra2 red positive cells analysed. As illustrated in Fig. 4A, we reason that in cases of aboral photoconversion the target zone in the early gastrula does not include presumptive gastroderm or i-cells, implying that any Dendra2 red cells detected at planula stages must derive from the aboral ectoderm. Conversely, oral Dendra2 photoconversion zones at early gastrula stages are predicted to include part of the presumptive oral ectoderm and gastroderm, and some of the i-cells. During gastrulation these will ingress aborally with the developing gastroderm to occupy the central region of the developing planula, before some of their neural and nematocyte descendants migrate out into the ectoderm. We thus anticipate finding Dendra2-red positive nematocytes and neural cells in the ectoderm at planula stages, both within the oral Dendra2-red positive patch, but also individually in more aboral positions separated spatially from this patch.

As expected, the progeny of aboral photoconverted Dendra2 cells formed coherent aboral patches of aboral ectoderm by the planula stage, while the progeny of oral photoconverted cells included oral ectoderm and gastroderm, the exact time and zone of photoconversion in each embryo determining the precise pattern observed (Fig. 4B-J). Confirming the results of the gastrula bisection experiments, the patches of photoconverted Dendra2 progeny in planulae derived from aboral (ie ectoderm-restricted) irradiation included mucous and secretory cells, characterised by their typical morphologies and nuclear positioned respectively in basal or medial positions (Fig. 4B-C). In addition, some Dendra2-red cells in the aboral ectoderm were stained by the anti-LWamide antibody (Fig. 4D-E).

In planulae derived from oral photoconversion, Dendra2-red nematocytes were clearly detected in the ectoderm far from the oral patch of photoconverted epidermal cells, validating this technique as an indicator of i-cell origins (Fig. 4F-H). Furthermore we could detect RFamide in Dendra2 red cells at positions detached from the oral ectoderm patch, but not scattered as widely as the nematocytes (Fig. 4I-J).

Figure 4. Dendra2 photoconversion traces neural fates from the gastrula stage

A) Experimental design; Regions of the aboral or oral poles of early gastrula stage embryos expressing Dendra2 from mRNA injected into the egg were photoconverted to red using the 405nm laser of a Leica Stellaris confocal microscope. B-J) Confocal microscope images of planulae derived from either aboral (B-E) or oral (F-J) conversions. B, C, D, E, F, G, H, I and J show the distribution of cells containing Dendra2-red. Other panels show the same planes with the Dendra2-red image in magenta overlaid with Phalloidin-Alexa488/endogenous GFP (green) and Hoechst staining of nuclei (cyan) (B', C', F' and G'; with Hoechst (cyan) only in H'; with anti-LWamide staining (yellow) and Hoechst (cyan) in D', E', E''; with anti-RFamide staining (yellow) and Hoechst (cyan) in I', J', J''. (B-C): examples of aborally derived mucous cells (arrowheads) showing characteristic cytoplasm-free cell bodies and basal nuclei. Nuclei in the mucous cells (m), aboral secretory cells (s) and epithelial cells

(e) are all positive for Dendra2-red. (D-E): examples of LWamide positive cells in the aboral ectoderm (arrows and arrowheads) positive for Dendra2-red. (F-H): examples of nematocytes (arrows and arrowheads) in ectodermal regions far from the oral-irradiated epithelial clone, likely derived from gastrodermal i-cells. (I-J): examples of RFamide positive cells in the trunk ectoderm (arrows and arrowheads) positive for Dendra2-red. Oral pole is at the top in all panels. Each image is through a central plane of the planula except F/ F' which is closer to the surface, within the ectoderm layer. Scale bars all 50µm.

We conclude from these two approaches that during *Clytia* planula development from the gastrula stage, neural cell types have two origins. I-cells in the gastrodermal region provide nematocytes and at least some of the RFamide ganglionic cells, whereas GLWamide sensory cells, along with mucous and aboral secretory cells, derive mainly from gastrula ectoderm.

Wnt-β-catenin signalling promotes nematogenesis and neurogenesis

Wnt-β-catenin signalling has widespread roles in regulating neurogenesis across animals as well as body axis specification. In *Clyta* it determines the oral-aboral axis, with a key role for the ligand Wnt3, expressed orally (Momose et al., 2008; Uveira et al., 2025), while in *Hydractinia* it also regulates i-cell differentiation into neurons (Duffy et al., 2010; Duffy et al., 2012). Conversely, pharmacological stimulation of the Wnt-β-catenin pathway in *Hydractinia* and *Hydra* polyps, causes excess formation of nematocytes and of some nerve cell types (Hensel et al., 2014; Khalturin et al., 2007; Teo et al., 2006). In *Hydra* it has been proposed that Wnt activity promotes nematocyte differentiation for the mouth and tentacle structures at the oral tip of the polyp where this pathway is active (Bosch and David, 1991; Hobmayer et al., 2000; Khalturin et al., 2007; Lengfeld et al., 2009).

To test the role of Wnt-β-catenin signalling in *Clytia* nematogenesis/neurogenesis, we used two approaches: Wnt3-MO injection, prior to fertilisation, which prevents oral territory specification and delays gastrulation (Lapébie et al., 2014; Momose et al., 2008), or timed incubations in PKF118-310, which interferes with TCF/β-catenin interactions (Chudekov et al., 2007; Figure 5). Early PKF118-310 treatments (from cleavage stage to 24 hpf) have been shown to mimic the morphological aboralising effects of Wnt-β-catenin signalling inhibition on axis establishment (Sinigaglia et al., 2020), which is completed by the parenchymula stage; Late treatments (24 to 48 hpf) specifically targeted the period of neural cell differentiation. These late treatments did not shift the established expression territories for Wnt3, WegA1 or FoxQ2a, but did abolish Bra1 expression at the oral tip of the planula (Supplementary File S2), indicating that once established axial identities along the oral-aboral axis can be

maintained independent of Wnt-β-catenin signalling. We monitored neural development in these experiments using the panel of in situ hybridisation probes presented in Figure 1. The staining was assessed for each group by counting the embryos/larvae showing similar numbers of stained cells as untreated embryos (scored as "positive"), either indistinct staining or of just a few cells (scored as "trace"), or no cellular staining (scored as "none") (Supplementary File S3).

Following Wnt3-MO injection (Fig. 5A) or early treatments with PKF118-310 (Fig. 5B), the first phase of nematogenesis was disrupted, such that most Wnt3-MO embryos showed only a few nematoblasts clearly positive for Minicollagen3/4a. Mos3 expression was barely detectable and Znf845 cells reduced to trace levels or absent. This effect on nematoblast formation was reflected by reductions in the cell populations expressing Nanos1 and Piwi (Fig. 5; Supplementary File S3); by 48 hpf the Nanos1-expressing population was reduced to trace levels in many Wnt3-MO embryos, as expected given the continued expression of these mRNAs during early nematogenesis. In contrast we observed no obvious reductions in cell populations expressing Nanos1 (Fig. 5A,B) in Wnt3-MO or PKF118-310 treated embryos at 24 hpf, suggesting that initial i-cell formation and maintenance are not directly affected by Wnt signalling.

The early steps of neural cell development by both i-cell and ectodermal neurogenesis pathways were found to be sensitive to Wnt-β-catenin inhibition. Both Neurogenin and Hlh6 expression were strongly reduced in PKF118-310 treated embryos at 24 hpf, and ELAV expression was markedly reduced by both early and late PKF118-310 treatments. A further requirement for Wnt signalling in neural differentiation steps, including the nematocyte maturation phase, was shown by late treatments with PKF118-310 (Fig. 5C). This had no obvious effect on existing Znf845 or Minicollagen expressing cell populations, but blocked their differentiation into nematocytes: Nematocilin expression was detected only weakly, and no nematocytes were found in the ectoderm. Furthermore, after these late PKF118-310 treatments, the number of cells expressing ELAV was greatly reduced, indicating that neural differentiation is not maintained in the absence of Wnt signalling. Consistently, both PP5/RFamide and PP2/GLWamide populations were reduced to trace levels in most embryos following PKF118-310 treatment from 24 hpf. In contrast, in Wnt3-MO planulae at 48h, we detected no RFamide cells but did detect PP2/GLWamide cells

despite the strong aboralised phenotype (Fig 5A). One possible explanation for this apparent discrepancy is that the planula aboral territory requires low levels of Wnt-β-catenin signalling for its development, as shown in *Nematostella* (Leclère et al., 2016). Given that several Wnt ligands are expressed in the *Clytia* planula in overlapping domains (Momose et al., 2008), we suggest that low level Wnt-β-catenin signalling may operate in Wnt3-MO treated planulae. These levels would mimic normal aboral conditions and promote GLWamide neuron development, whereas PKF118-310 would completely abolish both.

Taken together these results indicate programmes of neurogenesis, including nematogenesis, both ectodermal and i-cell derived, are sensitive to Wnt signalling. Wnt signalling promotes both the formation of nematoblasts and their differentiation into nematocytes, and likewise early neurogenic processes as well as terminal differentiation, particularly in oral and trunk regions of the developing planula.

Discussion

Clytia hemisphaerica offers an excellent opportunity to address the embryological origin of neural cells during hydrozoan larval development. By combining gene expression profiling, bisection experiments and lineage tracing, we have uncovered two parallel pathways, schematized in Figure 6. At the onset of gastrulation, Piwi/Nanos1 positive cells arising in the oral pole presumptive endoderm domain of the early gastrula rapidly embark on a programme of nematogenesis. This pathway, like all the neurogenesis steps we examined, is strongly affected by disruption of Wnt-β-catenin signalling, explaining the previously observed strong downregulation of Znf845 and Mos3, which we now know are early nematogenesis markers (Chari et al., 2021), in Wnt3-MO embryos (Lapébie et al., 2014). During normal planula development a large pool of Minicollagen3/4a-expressing nematoblasts accumulates in the gastrodermal region as early as the parenchymula stage. Only a small subpopulation of these subsequently migrate into the ectoderm and differentiate into mature nematocytes. We speculate that most of the 'stalled' nematoblasts will undergo rapid differentiation during metamorphosis to provide nematocytes for the primary polyp, enabling it to catch its first prey, as has been proposed for other hydrozoan planulae (Burmistrova et al., 2018; Chrysostomou et al., 2022). I-cells also generate some other neural cells including RFamide positive ganglionic cells.

Figure 6. Dual neurogenesis pathways during *Clytia* larval development
Cartoon illustrating successive developmental stages from egg to planula. At the early gastrula, parenchymula and planula stages, cell types specific to the ectodermal pathway and i-cell pathways are depicted on the left and right sides respectively. See text for detailed discussion.

In parallel to this i-cell pathway, a distinct neurogenesis process occurs in a basal layer of the aboral/lateral gastrula ectoderm generating GLWamide and possibly other neurosensory cells. Mucous cells and aboral secretory cells ("cellules speumeuses" and "cellules spheruleuses" respectively; Bodo and Bouillon, 1968) also derive from this layer. These two secretory cell types share molecular characteristics with neurons present in the planula and medusa (Ramon-Mateu 2025; Ferraioli et al, in preparation), and can be considered as part of a wider group of neural-secretory cell types, possibly with a common origin (Arendt et al., 2015; Babonis, 2025).

I-cell formation and fate during Clytia embryogenesis

The work reported here allows us to reassess the fate during *Clytia* larval development of i-cells, defined here as cells expressing genes typical of germ cell/stem cell pluripotency programmes (Juliano et al., 2010) such as Piwi, Vasa and Nanos1. Previously we showed that maternal "germ plasm" like granules including mRNAs of several of these genes is positioned around the female pronucleus at the egg animal pole (Leclère et al., 2012) as illustrated in orange in Figure 6, but found that Piwi/Nanos1 expressing i-cells could develop within the gastroderm from embryo fragments irrespective of whether or not they contained this germ plasm. We now know that most of the "i-cells" of the developing planula form de novo at the oral pole from the early gastrula stage, and that from this early stage they have already embarked upon nematogenesis as indicated by expression of Znf845, Mos3 and later Minicollagen (this study; (Chari et al., 2021). This validates previous histological descriptions noting the appearance of distinguishable nematoblasts during gastrulation, well ahead of the appearance of any cells with morphologies typical of totipotent i-cells (Bodo and Bouillon, 1968). The previous report that Piwi/Nanos1 positive cells can develop in larvae derived from vegetal as well as animal fragments of early embryos (Leclère et al., 2012) can, in this new light, be accounted for by the onset of nematogenesis in these embryos, and does not necessarily represent de novo formation of totipotent i-cells. Some i-cells also generate RFamide neurons and possibly other neural cell types. Expression of various potential molecular regulators of neural development have been reported in cells both scattered within the endodermal region and in the developing aboral ectoderm (this study, (Jager et al., 2011; Kraus et al., 2020; Lapébie et al., 2014)). Consistently, shRNA-mediated downregulation of a SoxB gene, expressed in an i-cell type pattern in the gastrodermal region during *Hydractinia* planula development, prevented formation of RFamide expressing cells (Chrysostomou et al., 2022; Flici et al., 2017).

Whether pluripotent i-cells, not yet engaged in neurogenesis or nematogenesis, and/or cells committed to a germline fate, are present in *Clytia* embryos and larvae remains an important open question. To resolve this issue will require the identification of specific markers for totipotent and germ line cells in *Clytia*, something that has so far remained elusive.

Alternative neurogenesis pathways in hydrozoans

Our bisection and Dendra2 tracing experiments in *Clytia* showed that several neural cell types of the planula larva, including GLWamide cells, mucous cells and secretory cells, derive from the ectoderm. This was also the conclusion from previous studies based on EM of hydrozoan planulae manipulated in various ways to eliminate i-cells (Martin and Thomas, 1981; Thomas et al., 1987). This ectodermal neurogenesis pathway is characterised by expression in a basal, neurogenic, layer of the blastula and gastrula ectoderm of the transcription factor genes Sox10 (Kraus et al., 2020) and then Hlh6 expression (this study). Also expressed in this layer are a number of genes from families implicated in modulating neurogenesis in other species such as bZip as well as Wnt and Notch pathway regulators, identified by their upregulation in Wnt-3 MO early gastrula-stage embryos (Lapébie et al., 2014). Strong expression of the neural gene ELAV in this layer, even before gastrulation is complete, emphasises the important contribution of ectodermal neurogenesis during planula formation. This picture of embryo neurogenesis differs from that characterised in the adult stages, i.e. polyps, of *Hydractinia* (Song et al., 2025) and *Hydra,* where i-cells give rise to all neurons including GLWamide positive sensory cells (Mitgutsch, 1999; Siebert et al., 2008). scRNAseq data suggest that i-cells generate a similar range of neural cell types in *Clytia* medusa stages (Chari et al., 2021).

Ectodermal neurogenesis is typical of bilaterian embryogenesis and has also been described in anthozoan cnidarians such as *Nematostella,* where both sensory and ganglionic cells arise in the ectodermal epithelium (Lemaître et al., 2023; Nakanishi et al., 2012; Rentzsch et al., 2017), and in the scyphozoan *Aurelia* (Nakanishi et al., 2008). In *Nematostella,* neurogenesis in the gastroderm follows that in the ectoderm at the planula stage (Nakanishi et al., 2012). Furthermore, juvenile and adult gastrodermal tissues harbour Vasa2 /Piwi1 expressing cells with similarities to hydrozoan i-cells, generating at least some of the neural precursors in addition to germ cells (Miramón-Puértolas et al., 2024). In hydrozoans, endodermal positioning of i-cells appears to be a particularity of the planula stage (Bodo and Bouillon, 1968; Leclère et al., 2012; Martin, 1991; Rebscher et al., 2008), perhaps relating to a requirement to stockpile nematoblasts for rapid deployment in the primary polyp following

metamorphosis (see above). At other life cycle stages, i-cells are mainly found between cells at the base of the ectoderm, with the notable exception at the medusa stage of the gonad (Bosch et al., 2010; Hou et al., 2021; Jessus et al., 2020).

More generally, our study reinforces the hypothesis that neurogenesis from the ectodermal epithelium evolved in a common ancestor of Cnidaria and Bilateria (Nakanishi et al., 2012). Ectodermal neurogenesis would have persisted during evolution of Hydrozoa, at least during development of the planula larva. Under this scenario, i-cells may have emerged in Hydrozoa as a complementary mechanism for neurogenesis/nematogenesis. In the *Clytia* larva, i-cells give rise mainly to nematocytes but also at least some of the RFamide cells, while in the adult medusa both nematocytes and most other neural cell types derive from i-cells (Chari et al., 2021; Denker et al., 2008). We can thus speculate the marked stem-like capacity of hydrozoan i-cells to proliferate locally may have facilitated continuous rapid production of nematocytes at particular sites in adult hydrozoans.

Acknowledgements :

We thank D. Houston (Univ. Iowa) for his participation in pilot embryo cutting experiments, T. Lamonerie (Univ. Côte d'Azur) for participation in the Dendra2 experiments, our research group colleagues and R. Copley for helpful discussions, G. Jékely for donating the anti-RFamide antibody and O. Koizumi for the anti-LWamide antibody.

Microscopy facilities were provided by the Imaging Platform and *Clytia* by the Service Aquariologie of Centre de Ressources Biologiques Marines (IMEV – FR 3761) of Institut de la Mer de Villefranche, both supported by EMBRC-France (ANR-10-INBS-02).

The work was funded by the H2020/Marie Skłodowska-Curie ITN "EvoCell" (grant agreement no. 766053) to E.H and CNRS core funding to LBDV.

SUPPLEMENTARY FILES

S1) Gene names and accession numbers
Details of the genes followed in this study.

S2) Polarity maintenance in PKF treated larvae

In situ hybridisation images of planula larvae (48 hpf) with probes to detect the genes indicated to the left of each panel, as markers for oral (Wnt3,Bra1) and Aboral (WgA1, FoxQ2a) polarity. Left column: Untreated planulae (cont.). Right column: incubated from 24-48 hpf in PKF118-310 (PKF). Oral pole is at the top in all panels. All panels are the same magnification, scale bars 100μm.

S3) Regulation of nematogenesis and neurogenesis by Wnt-β-catenin signalling
Numbers of each type of in situ hybridisation patterns obtained with each probe. Yellow/ orange cells indicate the main phenotype for each group, as illustrated in Figure 5. Counts are from visual scoring of each stained planula in the corresponding group from one or two (*) experiment(s): 'positive' = clearly stained cells detected; 'trace' = staining observed but indistinct and/or very few cells. ND = not done.

References

Arendt, D., Benito-Gutierrez, E., Brunet, T. and Marlow, H. (2015). Gastric pouches and the mucociliary sole: setting the stage for nervous system evolution. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 20150286–20150286.

Attenborough, R. M. F., Hayward, D. C., Wiedemann, U., Forêt, S., Miller, D. J. and Ball, E. E. (2019). Expression of the neuropeptides RFamide and LWamide during development of the coral Acropora millepora in relation to settlement and metamorphosis. *Dev. Biol.* **446**, 56–67.

Babonis, L. S. (2025). That Stinging Sensation: Modularity and the Origin of the Stinging Cell. *Integr. Comp. Biol.* **65**, 661–675.

Bodo, F. and Bouillon, J. (1968). Étude histologique du développement embryonnaire de quelques hydroméduses de Roscoff: Phialidium hemisphaericum (L.), Obelia sp. Péron et Lesueur, Sarsia eximia (Allman), Podocoryne carnea (Sars), Gonionemus vertens Agassiz. *Cah. Biol. Mar.* **9**, 69-104.

Bosch, T. C. and David, C. N. (1991). Decision Making in Interstitial Stem Cells of Hydra. *Vivo Athens Greece* **5**, 515–520.

Bosch, T. C. G., Anton-Erxleben, F., Hemmrich, G. and Khalturin, K. (2010). The Hydra polyp: nothing but an active stem cell community. *Dev. Growth Differ.* **52**, 15–25.

Bosch, T. C. G., Klimovich, A., Domazet-Lošo, T., Gründer, S., Holstein, T. W., Jékely, G., Miller, D. J., Murillo-Rincon, A. P., Rentzsch, F., Richards, G. S., et al. (2017). Back to the Basics: Cnidarians Start to Fire. *Trends Neurosci.* **40**, 92–105.

Burkhardt, P. and Jékely, G. (2021). Evolution of synapses and neurotransmitter systems: The divide-and-conquer model for early neural cell-type evolution. *Curr. Opin. Neurobiol.* **71**, 127–138.

Burmistrova, Y. A., Osadchenko, B. V., Bolshakov, F. V., Kraus, Y. A. and Kosevich, I. A. (2018). Embryonic development of thecate hydrozoan Gonothyraea loveni (Allman, 1859). *Dev. Growth Differ.* **60**, 483–501.

Byrum, C. A. (2001). An Analysis of Hydrozoan Gastrulation by Unipolar Ingression. *Dev. Biol.* **240**, 627–640.

Cazet, J. F., Siebert, S., Little, H. M., Bertemes, P., Primack, A. S., Ladurner, P., Achrainer, M., Fredriksen, M. T., Moreland, R. T., Singh, S., et al. (2023). A chromosome-scale epigenetic map of the Hydra genome reveals conserved regulators of cell state. *Genome Res.* **33**, 283–298.

Chari, T., Weissbourd, B., Gehring, J., Ferraioli, A., Leclère, L., Herl, M., Gao, F., Chevalier, S., Copley, R. R., Houliston, E., et al. (2021). Whole-animal multiplexed single-cell RNA-seq reveals transcriptional shifts across Clytia medusa cell types. *Sci. Adv.* **7**, eabh1683.

Chrysostomou, E., Flici, H., Gornik, S. G., Salinas-Saavedra, M., Gahan, J. M.,

McMahon, E. T., Thompson, K., Hanley, S., Kilcoyne, M., Schnitzler, C. E., et al. (2022). A cellular and molecular analysis of SoxB-driven neurogenesis in a cnidarian. *eLife* **11**, e78793.

Chudakov, D. M., Lukyanov, S. and Lukyanov, K. A. (2007). Using Photoactivatable Fluorescent Protein Dendra2 to Track Protein Movement. *BioTechniques* **42**, 553–563.

Condamine, T., Jager, M., Leclère, L., Blugeon, C., Lemoine, S., Copley, R. R. and Manuel, M. (2019). Molecular characterisation of a cellular conveyor belt in Clytia medusae. *Dev. Biol.*

David, C. N. (2012). Interstitial stem cells in Hydra: multipotency and decision-making. *Int. J. Dev. Biol.* **56**, 489–497.

Denker, E., Manuel, M., Leclère, L., Le Guyader, H. and Rabet, N. (2008). Ordered progression of nematogenesis from stem cells through differentiation stages in the tentacle bulb of Clytia hemisphaerica (Hydrozoa, Cnidaria). *Dev. Biol.* **315**, 99–113.

Denner, A., Steger, J., Ries, A., Morozova-Link, E., Ritter, J., Haas, F., Cole, A. G. and Technau, U. (2024). Nanos2 marks precursors of somatic lineages and is required for germline formation in the sea anemone Nematostella vectensis. *Sci. Adv.* **10**, eado0424.

Duffy, D. J., Plickert, G., Kuenzel, T., Tilmann, W. and Frank, U. (2010). Wnt signaling promotes oral but suppresses aboral structures in Hydractinia metamorphosis and regeneration. *Dev. Camb. Engl.* **137**, 3057–3066.

Duffy, D. J., Millane, R. C. and Frank, U. (2012). A heat shock protein and Wnt signaling crosstalk during axial patterning and stem cell proliferation. *Dev. Biol.* **362**

Ferraioli, A. (2022). Comparison of cell types across life cycle stages of the hydrozoan Clytia hemisphaerica. PhD Thesis, Sorbonne Université.

Flici, H., Schnitzler, C. E., Millane, R. C., Govinden, G., Houlihan, A., Boomkamp, S. D., Shen, S., Baxevanis, A. D. and Frank, U. (2017). An Evolutionarily Conserved SoxB-Hdac2 Crosstalk Regulates Neurogenesis in a Cnidarian. *Cell Reports* **18**, 1395–1409.

Fourrage, C., Swann, K. and Open, J. G. G. (2014). An endogenous green fluorescent protein–photoprotein pair in Clytia hemisphaerica eggs shows co-targeting to mitochondria and efficient bioluminescence energy *Open Biol.* **9**;4(4):130206.

Frank, U., Plickert, G. and Müller, W. A. (2009). Cnidarian Interstitial Cells: The Dawn of Stem Cell Research.pp. 33–59. Dordrecht: Springer Netherlands.

Gajewski, M., Schmutzler, C. and Plickert, G. (1998). Structure of neuropeptide precursors in cnidaria. *Ann. N. Y. Acad. Sci.* **839**, 311–315.

Genikhovich, G., Kürn, U., Hemmrich, G. and Bosch, T. C. G. (2006). Discovery of genes expressed in Hydra embryogenesis. *Dev. Biol.* **289**, 466–481.

Hayakawa, E., Takahashi, T., Nishimiya-Fujisawa, C. and Fujisawa, T. (2007). A novel

neuropeptide (FRamide) family identified by a peptidomic approach in Hydra magnipapillata. *FEBS J.* **274**, 5438–5448.

Hensel, K., Lotan, T., Sanders, S. M., Cartwright, P. and Frank, U. (2014). Lineage-specific evolution of cnidarian Wnt ligands. *Evol. Dev.* **16**, 259–269.

Hobmayer, B., Rentzsch, F., Kuhn, K., Happel, C. M., von Laue, C. C., Snyder, P., Rothbächer, U. and Holstein, T. W. (2000). WNT signalling molecules act in axis formation in the diploblastic metazoan Hydra. *Nature* **407**, 186–189.

Hou, S., Zhu, J., Shibata, S., Nakamoto, A. and Kumano, G. (2021). Repetitive accumulation of interstitial cells generates the branched structure of Cladonema medusa tentacles. *Development* **148**, dev199544.

Jager, M., Quéinnec, E., Le Guyader, H. and Manuel, M. (2011). Multiple Sox genes are expressed in stem cells or in differentiating neuro-sensory cells in the hydrozoan Clytia hemisphaerica. *EvoDevo* **2**, 12.

Jessus, C., Munro, C. and Houliston, E. (2020). Managing the Oocyte Meiotic Arrest—Lessons from Frogs and Jellyfish. *Cells* **9**, 1150.

Juliano, C. E., Swartz, S. Z. and Wessel, G. M. (2010). A conserved germline multipotency program. *Dev. Camb. Engl.* **137**, 4113–4126.

Kelava, I., Rentzsch, F. and Technau, U. (2015). Evolution of eumetazoan nervous systems: insights from cnidarians. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 20150065.

Khalturin, K., Anton-Erxleben, F., Milde, S., Plötz, C., Wittlieb, J., Hemmrich, G. and Bosch, T. C. G. (2007). Transgenic stem cells in Hydra reveal an early evolutionary origin for key elements controlling self-renewal and differentiation. *Dev. Biol.* **309**, 32–44.

Koizumi, O., Hamada, S., Minobe, S., Hamaguchi-Hamada, K., Kurumata-Shigeto, M., Nakamura, M. and Namikawa, H. (2015). The nerve ring in cnidarians: its presence and structure in hydrozoan medusae. *Zoology* **118**, 79–88.

Kraus, Y., Chevalier, S. and Houliston, E. (2020). Cell shape changes during larval body plan development in *Clytia hemisphaerica*. *Dev. Biol.* **468**, 59–79.

Lapébie, P., Ruggiero, A., Barreau, C., Chevalier, S., Chang, P., Dru, P., Houliston, E. and Momose, T. (2014). Differential responses to Wnt and PCP disruption predict expression and developmental function of conserved and novel genes in a cnidarian. *PLoS Genet.* **10**, e1004590.

Lechable, M., Jan, A., Duchene, A., Uveira, J., Weissbourd, B., Gissat, L., Collet, S., Gilletta, L., Chevalier, S., Leclère, L., et al. (2020). An improved whole life cycle culture protocol for the hydrozoan genetic model Clytia hemisphaerica. *Biol. Open* **9**, bio051268.

Leclère, L., Jager, M., Barreau, C., Chang, P., Le Guyader, H., Manuel, M. and Houliston, E. (2012). Maternally localized germ plasm mRNAs and germ cell/stem cell formation in the cnidarian Clytia. *Dev. Biol.* **364**, 236–248.

Leclère, L., Horin, C., Chevalier, S., Lapébie, P., Dru, P., Peron, S., Jager, M., Condamine, T., Pottin, K., Romano, S., et al. (2019). The genome of the jellyfish Clytia hemisphaerica and the evolution of the cnidarian life-cycle. *Nat. Ecol. Evol.* **3**, 801–810.

Lemaître, Q. I. B., Bartsch, N., Kouzel, I. U., Busengdal, H., Richards, G. S., Steinmetz, P. R. H. and Rentzsch, F. (2023). NvPrdm14d-expressing neural progenitor cells contribute to non-ectodermal neurogenesis in Nematostella vectensis. *Nat. Commun.* **14**, 4854.

Lengfeld, T., Watanabe, H., Simakov, O., Lindgens, D., Gee, L., Law, L., Schmidt, H. A., Ozbek, S., Bode, H. and Holstein, T. W. (2009). Multiple Wnts are involved in Hydra organizer formation and regeneration. *Dev. Biol.* **330**, 186–199.

Lepourcelet, M., Chen, Y.-N. P., France, D. S., Wang, H., Crews, P., Petersen, F., Bruseo, C., Wood, A. W. and Shivdasani, R. A. (2004). Small-molecule antagonists of the oncogenic Tcf/beta-catenin protein complex. *Cancer Cell* **5**, 91–102.

Martin, V. J. (1988). Development of nerve cells in hydrozoan planulae: I. Differentiation of ganglionic cells. *Biol. Bull.* **174**, 319–329.

Martin, V. J. (1991). Differentiation of the interstitial cell line in hydrozoan planulae. *Hydrobiologia* **216**, 75–82

Martin, V. J. (2000). Reorganization of the nervous system during metamorphosis of a hydrozoan planula. *Invertebr. Biol.* **119**, 243–253.

Martin, V. J. and Thomas, M. B. (1980). Nerve elements in the planula of the hydrozoanPennaria tiarella. *J. Morphol.* **166**, 27–36.

Martin, V. J. and Thomas, M. B. (1981). The origin of the nervous system in Pennaria tiarella, as revealed by treatment with colchicine. *Biological Bulletin* **160**,303-310

Miramón-Puértolas, P., Pascual-Carreras, E. and Steinmetz, P. R. H. (2024). A population of Vasa2 and Piwi1 expressing cells generates germ cells and neurons in a sea anemone. *Nat. Commun.* **15**, 8765.

Mitgutsch, C. et al (1999). Expression and Developmental Regulation of the Hydra-RFamide and Hydra-LWamide Preprohormone Genes in Hydra: Evidence for Transient Phases of Head Formation. *Dev. Biol.* **207**, 189-203.

Mochizuki, K., Sano, H., Kobayashi, S., Nishimiya-Fujisawa, C. and Fujisawa, T. (2000). Expression and evolutionary conservation of nanos-related genes in Hydra. *Dev. Genes Evol.* **210**, 591–602.

Momose, T., Derelle, R. and Houliston, E. (2008). A maternally localised Wnt ligand required for axial patterning in the cnidarian Clytia hemisphaerica. *Dev. Camb. Engl.* **135**, 2105–2113.

Nakanishi, N., Yuan, D., Jacobs, D. K. and Hartenstein, V. (2008). Early development, pattern, and reorganization of the planula nervous system in Aurelia (Cnidaria, Scyphozoa). *Dev. Genes Evol.* **218**, 511–524.

Nakanishi, N., Renfer, E., Technau, U. and Rentzsch, F. (2012). Nervous systems of the sea anemone Nematostella vectensis are generated by ectoderm and endoderm and shaped by distinct mechanisms. *Development* **139**, 347–357.

Piraino, S., Zega, G., Di Benedetto, C., Leone, A., Dell'Anna, A., Pennati, R., Carnevali, D. C., Schmid, V. and Reichert, H. (2011). Complex neural architecture in the diploblastic larva of Clava multicornis (Hydrozoa, Cnidaria). *J. Comp. Neurol.* **519**, 1931–1951.

Ramon-Mateu, J., Ferraioli, A., Teixidó, N., Domart-Coulon, I., Houliston, E. and Copley, R. R. (2025). Aboral cell types of Clytia and coral larvae have shared features and link taurine to the regulation of settlement. *Sci. Adv.* **11**, eadv1159.

Rebscher, N., Volk, C., Teo, R. and Plickert, G. (2008). The germ plasm component vasa allows tracing of the interstitial stem cells in the cnidarian Hydractinia echinata. *Dev. Dyn.* **237**, 1736–1745.

Rentzsch, F., Layden, M. and Manuel, M. (2017). The cellular and molecular basis of cnidarian neurogenesis. *Wiley Interdiscip. Rev. Dev. Biol.* **6**, e257.

Richards, G. S. and Rentzsch, F. (2014). Transgenic analysis of a SoxB gene reveals neural progenitor cells in the cnidarian Nematostella vectensis. *Dev. Camb. Engl.* **141**, 4681–4689.

Richards, G. S. and Rentzsch, F. (2015). Regulation of Nematostella neural progenitors by SoxB, Notch and bHLH genes. *Dev. Camb. Engl.* **142**, 3332–3342.

Salamanca-Díaz, D. A., Horkan, H. R., García-Castro, H., Emili, E., Salinas-Saavedra, M., Pérez-Posada, A., Rossi, M. E., Álvarez-Presas, M., Mac Gabhann, R., Hillenbrand, P., et al. (2025). The Hydractinia cell atlas reveals cellular and molecular principles of cnidarian coloniality. *Nat. Commun.* **16**, 2121.

Siebert, S., Anton-Erxleben, F. and Bosch, T. C. G. (2008). Cell type complexity in the basal metazoan Hydra is maintained by both stem cell based mechanisms and transdifferentiation. *Dev. Biol.* **313**, 13–24.

Siebert, S., Farrell, J. A., Cazet, J. F., Abeykoon, Y., Primack, A. S., Schnitzler, C. E. and Juliano, C. E. (2019). Stem cell differentiation trajectories in Hydra resolved at single-cell resolution. *Science* **365**, eaav9314.

Sinigaglia, C., Peron, S., Eichelbrenner, J., Chevalier, S., Steger, J., Barreau, C., Houliston, E. and Leclère, L. (2020). Pattern regulation in a regenerating jellyfish. *eLife* **9**, E3365.

Song, J., de Jong, D., Waletich, J., Baxevanis, A. D. and Schnitzler, C. E. (2025). An updated and spatially validated somatic single-cell atlas of Hydractinia symbiolongicarpus. *BioRxiv.* 2025.06.03.657738.

Steger, J., Cole, A. G., Denner, A., Lebedeva, T., Genikhovich, G., Ries, A., Reischl, R., Taudes, E., Lassnig, M. and Technau, U. (2022). Single-cell transcriptomics identifies conserved regulators of neuroglandular lineages. *Cell Rep.* **40**, 111370.

Takahashi, T. and Hatta, M. (2011). The Importance of GLWamide Neuropeptides in Cnidarian Development and Physiology. *J. Amino Acids* **2011**, 424501.

Teo, R., Möhrlen, F., Plickert, G., Müller, W. A. and Frank, U. (2006). An evolutionary conserved role of Wnt signaling in stem cell fate decision. *Dev. Biol.* **289**, 91–99.

Thomas, M. B., Freeman, G. and Martin, V. J. (1987). The Embryonic Origin of Neurosensory Cells and the Role of Nerve Cells in Metamorphosis in Phialidium gregarium(Cnidaria, Hydrozoa). *Int. J. Invertebr. Reprod. Dev.* **11**, 265–285.

Uveira, J., Donati, A., Léria, M., Lechable, M., Lahaye, F., Vesque, C., Houliston, E. and Momose, T. (2025). Planar cell polarity coordination in a cnidarian embryo provides clues to animal body axis evolution. *eLife* **14**,.

Varley, Á., Horkan, H. R., McMahon, E. T., Krasovec, G. and Frank, U. (2023). Pluripotent, germ cell competent adult stem cells underlie cnidarian regenerative ability and clonal growth. *Curr. Biol.* **33**, 1883-1892.e3.

Weissbourd, B., Momose, T., Nair, A., Kennedy, A., Hunt, B. and Anderson, D.J. (2021) A genetically tractable jellyfish model for systems and evolutionary neuroscience. *Cell* **184**, 5854-5868