Parallel and non-parallel features of adaptive radiation in Yucatán pupfishes

Matthew C. Kustra[1,2,3,*], David Tian[1,2], M. Fernanda Palominos[1,2], Feifei Guo[1,2], Dylan Chau[1,2], Oskar Golwala[1,2], HoWan Chan[4], Andrés Alvarez Zapata[5], Reyna Guadalupe Cetz Paredes[5], Frida Ximena Cortés Sánchez[6], Sonia Gabriela Hernández[6], Adan Fernando Mar-Silva[6], Fernando Mex[7], Charles Tralka[1,2], Maribel Badillo-Alemán[7], Juan J. Schmitter-Soto[8], Carlos A. Gracida-Juárez[9], Christopher M. Martinez[10], Jairo Arroyave[6], and Christopher H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley, CA, United States

[2]Museum of Vertebrate Zoology, University of California, Berkeley, CA, United States

[3]Miller Institute for Basic Research in Science, University of California, Berkeley, CA, United States

[4]Department of BioSciences, Rice University, Houston, TX, United States

[5]Instituto Tecnologico Superior de Conkal, Yucatan, Mexico

[6]Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, Mexico

[7]Unidad Multidisciplinaria de Docencia e Investigación, Unidad Sisal, Universidad Nacional Autónoma de México, Sisal, Yucatán, Mexico

[8] Departamento de Sistemática y Ecología Acuática, El Colegio de la Frontera Sur, Chetumal, Quintana Roo, Mexico

[9]Subdirección de Posgrado e Investigación, Tecnológico Nacional de México, Campus Felipe Carrillo Puerto.Carretera a Vigía Chico km 1.5, 77200 Felipe Carrillo Puerto, Quintana Roo.

[10]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA, United States

*Corresponding author: matthewckustra@berkeley.edu

Abstract

Understanding the extent of parallelism across adaptive radiations remains a central

problem in evolutionary biology. We used whole-genome resequencing of 123 individuals to

study an adaptive radiation of *Cyprinodon* pupfishes in Laguna Chichancanab, Mexico,

compared to previous research on an independent radiation of San Salvador Island (SSI)

pupfishes in the Bahamas, to assess the repeatability of their genetic architecture, sources of

adaptive variation, and stages of selection. Despite rapid craniofacial divergence of trophic

specialists within 8-15 kya, only two candidate genes (3.6%) were shared between Caribbean

radiations. Although adaptive introgression played a major role in SSI, we found minimal

evidence of adaptive introgression in Chichancanab, likely due to the higher geographical

isolation of this inland lake. Instead, de novo mutations provided a substantial source of adaptive

variation (30.6%) for the endemic zooplanktivore, nearly 10 times higher than the scale-eater in

SSI. We found strong evidence that selection occurred in stages, first on regulatory and standing

genetic variation, then on de novo and nonsynonymous mutations, parallel to SSI. Unique to

Chichancanab, adaptive variants near opsin and spermatogenesis genes are consistent with our

findings of greater visual acuity and differences in sperm morphology, respectively, within the

zoooplanktivore. Our study suggests that de novo mutations may contribute more to rapid

adaptive radiations than previously appreciated, particularly in isolated environments.

Introduction

Much of biodiversity results from adaptive radiation—when a clade diversifies rapidly into three or more new species (1–5). Young adaptive radiations provide excellent systems for investigating speciation and adaptation (3, 6–9). However, it remains unclear to what degree radiations occur in parallel given similar starting positions of genetic diversity, gene flow, and environments (10, 11).

Beneficial de novo mutations may provide novel variation to reach new fitness peaks on the adaptive landscape during adaptive radiations (12, 13); however, the time needed for de novo beneficial mutations to arise contrasts with the rapid divergence of adaptive radiations (11, 14, 15). Alternatively, hybridization could introduce large amounts of genetic variation and novel phenotypes (transgressive segregation) through hybrid swarms, enabling populations to reach previously inaccessible fitness peaks (16). Support for the role of hybridization in adaptive radiations has been demonstrated both theoretically (17, 18) and empirically in many systems (3, 19–27). However, we still do not know the importance of different sources of genetic variation across time and space.

Caribbean pupfishes are a remarkable system for assessing the repeatability of adaptive radiations due to two independent, recent adaptive radiations that contain a generalist and multiple trophic specialists within isolated lake environments that have historically lacked predators. The San Salvador Island (SSI), Bahamas radiation proceeded predominantly via the reassembly of pre-existing standing genetic variation found throughout the Caribbean, with additional contributions of adaptive introgression in both molluscivore (durophage) and scale-eating (lepidophage) specialists, and an additional 2% of de novo mutations within the scale-eater (26, 28–32). However, it is unclear how a second radiation in Laguna Chichancanab,

Mexico, drawing from the same Caribbean pool of standing genetic variation over the same

timeframe unfolded, allowing us to assess the degree to which radiations are parallel.

Here, we use whole-genome resequencing of 123 individuals to provide insight into the

genomic architecture and genetic origins of the Laguna Chichancanab *Cyprinodon* pupfish

radiation (33). This recent adaptive radiation (~8000 years old (34)) consisted of at least five

species historically (Strecker, 2005; Humphries and Miller 1981), including a detritivore (*C.*

beltrani), piscivore (*C. maya*), rocky substrate specialist (*C. verecundus*), bivalve specialist (*C.*

labiosus), and zooplanktivore (*C. simus*). However, due to recent colonization by native fishes

not historically present in the lake and invasive colonization by African tilapia (*Oreochromis* sp.)

(36, 37), all trophic specialist species now appear to be extinct since our collections in 2022, with

only the generalist, *C. beltrani*, remaining in 2024 (MK pers. obs.) and 2025 (CG, JSS pers.

obs.). Despite exhibiting one of the fastest rates of trait diversification of any vertebrate in tooth

length (130 times faster than background rates across *Cyprinodon* (29)) and rapid divergence in

other craniofacial morphology (Figure 1A), previous genetic studies of microsatellite data and

mtDNA suggest that this radiation has low genetic differentiation (38, 39), potentially enabling

the identification of highly differentiated loci among species that may be involved in trophic

adaptation.

Results

Demographic history

To understand the demographic history and population structure of the Yucatán

Cyprinodon pupfishes, we collected specimens from the wild in 2022 (Figure 1B) and conducted

whole genome resequencing of the species flock and the coastal sister species (*n* = 15 *C.*

artifrons, *n* = 35 *C. beltrani*, *n* = 35 *C. labiosus*, and *n* = 17 *C. simus*). We aligned these samples,

as well as eleven other previously sequenced Caribbean pupfish species, to the *Cyprindon*

nevadensis mionectes reference genome (GCA_030533445.1). After calling and filtering

variants, our dataset included 23,960,536 SNPs (12.4x mean depth).

103   Using MSMC2 on Chichancanab species on a linkage disequilibrium pruned set of SNPs

(*n* =5,476,855 SNPs) and a mutation rate of 1.56×10^{-8} substitutions per base pair (estimated

from a Caribbean *Cyprinodon* species (26)), we found that the three Chichancanab species (*C.*

beltrani, *C. labiosus*, and *C. simus*) diverged from the coastal sister species *C. artifrons* at the

end of the Pleistocene, 10-15 kya, consistent with receding sea levels resulting in the formation

of the Chichancanab lake (34). Since divergence from *C. artifrons*, the Chichancanab species

flock shows a similar demographic history, characterized by a sharp decline in effective

population sizes, consistent with endemic sympatric species within an isolated lake (Figure 1C).

Genetic differentiation of the species flock

113   Using a linkage-pruned set of SNPs and the subset of data restricted to Yucatán pupfish,

we used principal component analysis and ADMIXTURE to visualize population structure. We

found that all species formed distinct, non-overlapping genetic clusters, with *C. artifrons* being

the most genetically divergent from and sister to the Chichancanab species (Figure 1D). Within

the Chichancanab species flock, *C. simus* was more genetically divergent from *C. labiosus* and

C. beltrani (Figure 1D). These results were also supported by genome-wide F_{ST} estimates and a

phylogenetic tree constructed with *ADMIXTOOLS2* (Table S1; Figure S1). Four clusters

corresponded to the four different species, as determined by ADMIXTURE (Figure 1E). All *C.*

artifrons individuals showed minimal evidence of admixture with Chichancanab species (Figure

1E).

Figure 1. Yucatán *Cyprinodon* pupfishes show strong genetic differentiation. (A) µCT scans
of Yucatán pupfish. (B) Google Earth image of the Yucatán Peninsula showing the sampling site
for *C. artifrons*, with an inset map of Laguna Chichancanab. The pie chart represents the relative
frequency of the different Chichancanab species in our dataset. (C) Estimated demographic
history with *MSMC2* using linkage disequilibrium pruned SNPs. Each colored line represents an
individual, with colors corresponding to different species. (D) Principal component analysis
using a set of linkage disequilibrium pruned SNPs. (E) Estimated ancestry proportions for each
individual inferred using *ADMIXTURE* with K = 4 (see Figure S2 for K=2-3) from the linkage-
pruned dataset.

Adaptive loci within trophic specialists

Although genome-wide differentiation among Chichancanab species was moderately low

(F_{ST} ranging from 0.032 to 0.09; Figure 2A; Table S1), we were able to identify highly

differentiated regions of the genome within the trophic specialist zooplanktivore *C. simus*. We

identified 19 fixed and 1,127 nearly-fixed ($F_{ST} >0.95$) SNPs in *C. simus* relative to other

Chichancanab pupfishes (Table S2, 3). In contrast, we found zero fixed or nearly-fixed SNPs

within the generalist *C. beltrani* or bivalve-specialist C. *labiosus*. However, we were able to

identify some moderately differentiated SNPs within *C. labiosus* ($n = 97$) and *C. beltrani* ($n =$

13) relative to the other Chichancanab species at an F_{ST} threshold of 0.8 (Table S2).

To search for signals of selection, we calculated the normalized four-taxon population

branch statistic (PBS_{nj}; Schmidt et al., 2019) across 10-kb non-overlapping windows because

linkage disequilibrium decayed at this distance (Figure S3). Simulation studies demonstrate that

normalized *PBS* is effective in identifying both hard and soft selective sweeps (40). Out of the

1,146 SNPs fixed or nearly fixed within *C. simus*, 661 were within a strongly supported selective

sweep window (top 1% PBS_{nj} outlier (40)), including 425 SNPs within 20 kb of the first or last

exon of an annotated gene. These 425 candidate adaptive SNPs ("adaptive variants") were

located in the proximity of or within 55 genes (Figure 2A), distributed across 19 out of 24

chromosomes (Figure 2A). Adaptive variants associated with the same gene were often

distributed across multiple introns or exons and spaced apart by 1kb or more (Figure S4, Table

S4).

Source and function of adaptive genetic variation in the zooplanktivore

We next characterized the location of the 425 candidate adaptive variants near or within a

gene in the zooplanktivore *C. simus*. More than 95% were in intronic or flanking regions (20-kb

upstream or downstream of a gene; Figure 2B). Within coding regions, 13 out of 20 adaptive

variants were nonsynonymous mutations (Figure 2B). The number of adaptive variants per gene

followed an exponential distribution, with a handful of genes containing many variants and most

genes with only a single adaptive variant (Figure 2B). We functionally characterized these genes

by manually checking online databases and doing a literature search (Table S3; Extended Table

S1). 32 out of the 55 candidate genes were associated with craniofacial development (n =11;

20%) or sensory/neural function (n = 21; 38.2%). We manually checked if any of these genes

overlapped with the 176 unique candidate genes identified in the SSI radiation (26), and we

found only two overlapping candidate genes (*DNM1* and *SPTLC3*), both of which are associated

with motor neuron development (41, 42) and are scale-eater-specific candidate genes.

Next, we characterized the source of the variants and found that 63% of the adaptive

SNPs in *C. simus* also existed as standing genetic variation in the coastal Yucatan population of

C. artifrons ("standing variation"), 31% were *de novo* variants found only in Chichancanab, and

6% were detected as standing genetic variation in other pupfish species but not *C. artifrons*

(Figure 2C).

Figure 2. Architecture and source of adaptive genetic variation in the zooplanktivore. (A,
top row) F_{st} between *C. simus* and *C. beltrani* + *C. labiosus* calculated in non-overlapping 10-kb
windows. F_{st} values for candidate SNPs (red circles) were calculated separately per locus to
show the location of fixed variants. (A, lower row) Population branch statistics (PBS_{nj}) with *C.*
simus as the target population versus *C. beltrani*, *C. labiosus*, and *C. artifrons* in 10-kb non-
overlapping windows. Candidate genes within 20 kb of fixed or nearly-fixed (Fst > 0.95) SNPs
and within PBS_{nj} 1% outlier windows are highlighted in red and labeled. Alternating colors
represent different chromosomes. (B) Candidate adaptive variants near or within a gene were
defined as SNPs that were fixed ($n = 17$) or nearly fixed ($n = 1,129$, $F_{st} > 0.95$) and occurred
within a 1% population branch statistic (PBS_{nj}) outlier window for *C. simus* (Venn diagram). Pie
chart and bar graphs indicate the proportion of these 425 adaptive variants that were found
within the flanking, intronic, or coding regions. (C) Pie chart and bar graphs indicate the
proportion of these 425 variants that were detected only within Chichancanab (de novo), as
standing genetic variation in coastal sister species *C. artifrons*, or in other Caribbean species, but
not *C. artifrons*. * Indicates genes with a craniofacial annotation; ‡ indicates genes with neural or
sensory annotation.

Limited adaptive introgression within Chichancanab pupfishes

To test for introgression into the Chichancanab radiation, we used *f*-statistics calculated in

Dsuite (43) to examine gene flow from 11 additional *Cyprinodon* species from across the clade

and the sister species to the genus *Megupsilon aporus*. Within Yucatán pupfishes, we found

evidence of introgression between *C. artifrons* and both *C. beltrani* and *C. labiosus*, with the

strongest signal in *C. beltrani* (Figure 3A, B). We also found evidence for introgression between

C. beltrani and most outgroup species examined, including the sister species to *Cyprinodon*,

Megupsilon aporus. Relative to *C. beltrani*, *C. labiosus* and *C. simus* also showed evidence of

introgression with pupfishes endemic to the Chihuahua desert (*C. eximius* and *C. fontinalis*;

Figure 3A, B). Furthermore, we also found evidence of introgression between the two sympatric

radiations (*C. variegatus* from SSI) and *C. simus* (Figure 3A, B). However, additional *f*-branch

statistics indicate that all patterns of excess allele sharing between these outgroup species besides

C. artifrons and the Chichancanab species flock likely result from ancient introgression events

arising before the colonization of Laguna Chichancanab (Figure 3C; S5).

To test for adaptive introgression into *C. simus*, we next examined whether the set of

adaptive variants (Figure 2) occurred within outlier introgression windows using sliding window

scans of distance fraction (d_f; Pfeifer & Kapan, 2019) for species that had significant genome-

wide introgression into *C. simus* (Figure 3A , B). We found that none of the outlier introgression

windows overlapped with *C. simus* adaptive variants (Figure 3D).

Next, we performed *df* scans for introgression between *C. artifrons* and Chichancanab

species. Although there were outlier introgression regions between *C. simus* and *C. artifrons*

(positive values d_f), these regions did not overlap with *C. simus* adaptive variants (Figure 3D).

However, the position of several of these *C. simus* adaptive variants was found in outlier

introgression windows in *C. labiosus* and *C. beltrani* (Figure 3D). The number of outlier

introgression windows was much lower in *C. simus* relative to *C. labiosus* and *C. beltrani*

(Figure 3E, F). Furthermore, both *C. labiosus* and *C. beltrani* genomes contained several

introgression tracts that were much longer than *C. simus* tract lengths, indicative of much more

recent introgression events in these species (Figure 3E, F).

Figure 3. Limited adaptive introgression into *C. simus*. Heatmaps of (A) Patterson's *D*-statistics and (B) F_4 ratios for P1-P2 species combinations within the Chichancanab radiation along the x-axis and P3 focal species along the y-axis. All four-taxon trees used *Cualac tessellatus* as the outgroup (P4). Only significant (*Z*-score > 3) results are colored; combinations not relevant to the Chichancanab radiation are not shown. (C) *f*-branch (f_b) statistics indicate introgression occurring before the Chichancanab radiation. Red boxes indicate significant introgression between P3 species (top row) and the focal species or internal branch (dashed lines) on the left. Non-significant f_b statistics are indicated by white boxes. Grey boxes represent comparisons that cannot be made since f_b cannot be calculated for introgression between sister taxa. See Figure S5 for full *f*-branch graphs. (D) Genome-wide plot of the distance fraction (d_f, an introgression test statistic*;* Pfeifer & Kapan, 2019) calculated in non-overlapping windows of 91 SNPs (approximately 10 kb) for species trios that showed evidence of introgression with *C. simus*. d_f outlier windows that contained a candidate gene under selection in *C. simus* are colored red and labeled with the gene name. The y-axis label gives the introgression topology: (P1, P2), P3. Positive values indicate regions of introgression between *C. simus* and P3 (from top to bottom: *C. eximius*, *C. fontinalis*, *C. variegatus* (SSI), and *C. artifrons* in the bottom two subplots); negative values indicate regions of introgression between P1 (*C. beltrani* or *C. labiosus*) and P3. Red dashed lines indicate outlier cutoffs, which were determined empirically by the percentage of genome-wide introgression estimated by F_4 ratios in panel B. (E, F) Distribution of the size of d_f outlier introgression tract lengths (in 10 kb windows) between *C. artifrons* and Chichancanab species. The introgression tracts containing candidate adaptive genes in *C. simus* are labeled with gene names. SSI: San Salvador Island, Bahamas; NC: North Carolina, USA.

The stages of adaptive divergence of the zooplanktivore

We used *starTRMCA* (45) to estimate the timing of selective sweeps for the set of

candidate adaptive variants near genes within the zooplanktivore *C. simus* (Fig. 4). We estimated

that most selective sweeps occurred in a staggered sequence and not simultaneously after

colonization of Laguna Chichancanab (Figure 5A)(34). There were also a few older sweeps that

we estimate may predate the age of Laguna Chichancanab and divergence from the coastal sister

species *C. artifrons* (Figure 1B). Overall, we observed no clear temporal stages of adaptive

divergence related to craniofacial genes (asterisks: Figure 5) or neural/sensory genes (‡: Figure

5); instead, we found that selection on these two broad functional categories occurred throughout

the period of divergence of *C. simus* following the colonization of Laguna Chichancanab.

Next, we used generalized linear regressions to test if sweep age was correlated with the

proportion of nonsynonymous adaptive variants, de novo adaptive variants, or the overall

number of adaptive variants per gene region. We found that both nonsynonymous adaptive

variants ($\chi^2 = 5.852$, $p = 0.016$) and de novo mutations ($\chi^2 = 6.221$, $p = 0.013$) were significantly

more likely to occur among the more recent selective sweeps during the final stage of adaptive

divergence in *C. simus* relative to other members of the species flock (Figure 5B, C). However,

nonsynonymous mutations were not significantly more likely to be de novo mutations (All

adaptive variants: $\chi^2 = 0.102$, $P = 0.749$; singleton adaptive variants: $\chi^2 = 2.010$, $p = 0.156$), and

there was no correlation between estimated sweep age and the number of candidate adaptive

variants per gene region ($\chi^2 = 0.111$, $p = 0.740$).

Figure 4. Stages of adaptive divergence in the zooplanktivore. (A) Points indicate median
sweep age with 95% credible intervals. The dashed line indicates the approximate age of Laguna
Chichancanab (34). For each sweep, the first column of pie charts shows the proportions of
flanking/intronic/synonymous/nonsynonymous, second column of pie charts shows the
proportions of de novo/standing variation, and third column of squares shows the number of
candidate adaptive variants per gene (from Fig. 2). For genes with multiple adaptive variants, we
used the variant with the median position as the focal position. * Indicates genes with a
craniofacial annotation; ‡ indicates genes with neural or sensory annotation. (B) The proportion
of nonsynonymous adaptive variants is significantly larger in more recent selective sweeps,
indicated by the best-fit quasibinomial regression line; each pie chart represents a single
candidate gene. (C) The proportion of de novo variants is significantly larger in more recent
selective sweeps; each pie chart represents a single candidate gene.

Increased visual acuity in the zooplanktivore is consistent with vision-related candidate adaptive

genes

We identified many *C. simus* adaptive variants associated with eye development (*ASIC3*,

CRIM1, *MAB21L2*) and opsin genes (*OPN1LW*, *SWS2*). Because the trophic specialization of *C.*

simus requires visually locating zooplankton in the open water before targeted suction-feeding

strikes, we hypothesized that this species would have higher visual acuity than the detritivore, *C.*

beltrani. We conducted an optomotor response experiment on lab-reared individuals from both

species (F1 *C. beltrani* and long-term laboratory colony of *C. simus* (F10+)) and found that *C.*

simus has a much stronger optomotor response than *C. beltrani* in response to black-and-white

vertical bars spinning at a constant rate of ~89.4 rotations per minute (Figure 5A, B; Wilcoxon

signed-rank test: $W = 1$, $p = 0.009$). There was no significant difference in rotations per minute

between the species for the same individuals during control observation periods of no spinning

(Figure S6A; $W = 14$, $p = 0.3105$) or during a control treatment with an all-black background

spinning at the same frequency (Figure S6B; $W = 9$, $p = 0.147$).

Figure 5. Zooplanktivore *C. simus* has significantly greater visual acuity and divergent sperm morphology relative to the generalist *C. beltrani*. (A) Diagram of the optomotor response experimental apparatus (adapted from Neuhauss et al., 1999). (B) Boxplot and jittered points of the number of complete rotations fish swam per minute, indicative of their optomotor response to black-and-white vertical bars spinning at a constant rate. (C) Representative labeled image of a single *C. beltrani* sperm at 100x stained with rose bengal. (D) Boxplots with jittered points of sperm head diameter for *C. artifrons, beltrani,* and *simus*. Significant Tukey's post hoc differences are shown with an asterisk.

Divergent sperm morphology of the zooplanktivore is consistent with spermatogenesis-related

candidate adaptive genes

Four candidate genes (*C4orf22*, *STK35*, *ERC2*, and *TBC1D25*) showed annotations relating to spermatogenesis/fertility. We therefore examined whether three Yucatan species, raised in a common garden laboratory environment (F1 *C. artifrons,* F1 *beltrani,* and a long-term laboratory colony of C. *simus* F10+), differed in sperm morphology. We found that species significantly differed in sperm head length (Figure 5D; $F = 5.724$, $p = 0.025$), but not sperm midpiece length (Figure S7A; $F = 2.403$, $p = 0.147$) or flagellum length (Figure S7B; $F = 0.414$, $p = 0.673$). Tukey's post hoc pairwise comparisons indicated that *C. simus* had significantly larger sperm heads than *C. beltrani* ($q = 3.367$; $p = 0.0207$; Fig. 5D).

Discussion

Here, we investigated the genetic architecture, source of adaptive variation, and temporal dynamics of selection underlying speciation and trophic specialization in a young radiation of *Cyprinodon* pupfishes endemic to Laguna Chichancanab, Mexico. By comparing these patterns to our previous work on a recent radiation of *Cyprinodon* pupfishes in SSI, Bahamas (26, 28–32), we can examine the parallel and non-parallel features of these independent radiations in similar environments. Despite replicate radiations of Caribbean pupfishes in predator-free, large, isolated saline lakes with few competitors, we found little parallelism between the radiations. Although craniofacial morphology evolved rapidly within trophic specialists in both radiations, we found almost zero overlap in the adaptive genes underlying trophic specialization, potentially reflecting the divergent specialists within each radiation. There was no behavior-first stage of adaptive divergence in Chichancanab species and gene categories potentially relevant to speciation were not shared, such as pigmentation genes in SSI versus opsin and spermatogenesis

genes in Chichancanab. Unlike the SSI radiation and most other well-studied adaptive radiations

(e.g., Malawi cichlids (20), Galápagos finches (47)), adaptive introgression likely played a

minimal role in the Chichancanab radiation. Instead, over 30% of the candidate adaptive variants

in the zooplanktivore, *C. simus*, likely arose from de novo mutations compared to 2% in the SSI

scale-eater specialist. The one parallel genetic feature was a clear final stage of refinement in the

adaptive divergence of the zooplanktivore, marked by increased rates of de novo and

nonsynonymous adaptive mutations occurring later in the radiation.

Adaptive introgression did not contribute to adaptive radiation of Chichancanab pupfishes

Genomic analyses have shown that hybridization and introgression played a major role in

many adaptive radiations (3, 3, 16, 19–27, 48, 49), including the *Cyprinodon* SSI radiation (26,

31). However, in the Chichancanab radiation, we find little evidence for introgression playing a

major role, and did not detect any evidence of adaptive introgression following the colonization

of Laguna Chichancanab. Although our analyses indicate extensive introgression among

Cyprinodontidae lineages, we did not find any introgression events into the root of the radiation,

nor has there been gene flow from species other than the sister-species to the radiation, *C.*

artifrons. This secondary gene flow with *C. artifrons* is mostly between *C. beltrani* and, to a

lesser extent, *C. labiosus.* Gene flow may be higher with these species because *C. artifrons,* a

benthic feeding detritivore, is less ecologically divergent from both *C. beltrani,* a benthic feeding

detritivore, and *C. labiosus*, a benthic feeding bivalve specialist. This secondary gene flow

between *C. artifrons* and *C. beltrani* and *C. labiosus* overlaps with many genes under selection in

C. simus, suggesting that secondary gene flow is potentially helping maintain divergence

between *C. simus* and the benthic Chichancanab species not by bringing in new adaptive

variation for trophic specialization, but by maintaining the generalist/omnivore species.

These results are remarkably different from the *Cyprinodon* adaptive radiation of SSI,

where there is strong evidence of introgression at the root of the radiation, evidence of ongoing

introgression, and the presence of many introgressed adaptive variants in both trophic specialists

(26, 31). Laguna Chichancanab is a landlocked lake with no above-ground river connections and

is geographically more isolated compared to SSI, a small island in the approximate center of the

Caribbean. Furthermore, the sister species, *C. artifrons*, is abundant along the entire coast of the

Yucatán, meaning that any gene flow into Chichancanab would likely first have to pass through

C. artifrons. An additional, non-mutually exclusive explanation could be that the water chemistry

of Laguna Chichancanab, with near-saturation of calcite and gypsum, is a greater barrier to non-

adapted pupfish species than the hypersaline lakes of SSI. Thus, geography and ecology may

play an important role in determining the relative role that introgression may play in radiations.

De novo mutations play the dominant role in driving adaptive divergence of the zooplanktivore

The importance of de novo mutations in adaptive radiations remains contentious, in part

due to the time needed for de novo beneficial mutations to arise (11, 15). Interestingly, we find

that over 30% of the zooplanktivore adaptive variants were detected only within Chichancanab

species, suggesting their de novo origins within the basin, approximately 10 times greater than

the role of de novo mutations in the adaptive divergence of the scale-eater in the SSI radiation. A

primary factor could be that the estimated effective population size for the Chichancanab

radiation is approximately twice that of the SSI radiation (26), meaning that selection on de novo

mutations would be more effective, with a higher probability of fixation, all else being equal

(14). There may also be a lack of existing standing genetic variation relevant to the trophic

specialization of zooplanktivory. Although radiations should proceed along the genetic lines of

least resistance (50), the morphological axis along which the zooplanktivore has diverged is

orthogonal to almost all morphological variation that exists in Caribbean pupfish species (29).

Evolving along the line of greatest resistance instead of the line of least resistance may require de

novo mutations. While the SSI scale-eater is also highly divergent, it is closer to the existing

morphological space of Caribbean pupfish species and in the direction of greatest variance, at

least within the two phenotypic dimensions of greatest variance (29). Finally, an alternative

possibility could be that many of these de novo mutations do exist as standing variation and were

missed in our dataset, but this is unlikely to explain all de novo mutations, given our extensive

sampling of *Cyprinodon* species across the clade and spanning the Caribbean. Thus, our results

highlight that de novo mutations may play a more important role in driving young adaptive

radiations than currently appreciated (12, 13), especially when ecotypes are highly divergent

from other species within the radiation.

Extremely limited overlap in the genes underlying adaptive radiations of trophic specialists

418   We found only two candidate genes (*DNM1* and *SPTLC3*) among the 55 zooplanktivore

candidate genes in the Chichancanab radiation that overlapped with the hundreds of candidate

genes identified so far in the SSI radiation (26, 28, 30, 32, 51–55). Consistent with both trophic

specializations requiring the pursuit of mobile prey, both genes are involved in motor neuron

development. *DNM1* is a dynamin with nervous tissue-specific expression that is crucial for the

proper formation of motor neuron axons (41). *SPTLC3* is involved in the biosynthesis of

sphingolipids (56), and suppression of *SPTLC3* in zebrafish results in motor neuron axon defects

(42).

The limited number of shared adaptive genes is somewhat surprising given the high

conservation of gene function in both morphological development (57, 58) and behavior (59),

and the frequent reuse of the same candidate genes among vertebrate adaptive radiations,

particularly for craniofacial genes. For example, *BMP4* is involved in diversification in both

Galápagos finches (60) and African cichlids (61, 62). Further, within the SSI radiation, many of

the same genes are involved in craniofacial divergence between divergent trophic specialists (26,

53). Selection may still be acting on the same genetic pathways and genetic regulatory networks

missed by our direct candidate comparisons.

Another potential reason for this non-parallelism could be that trophic specialists in each

radiation have diverged in different sets of traits (29). Specifically, the Chichancanab radiation

has diversified in tooth length over 130 times faster than background rates, driven by the

extremely short teeth of the zooplanktivore (29). Consistent with divergence in tooth

morphology, one of the candidate craniofacial genes with many de novo mutations in flanking

and intronic regions was *WNT10A*. Mutations in *WNT10A* are associated with tooth

developmental disorders in humans (63). Furthermore, knockout and overexpression experiments

in sticklebacks and zebrafish have demonstrated *WNT10A*'s role in teeth development and

regeneration (64, 65).

The parallel role of vision but the non-parallelism of adaptive candidate genes

One of the more striking features of both *Cyprinodon* radiations is the evolution of novel

trophic specialists from a generalist ancestor: the scale-eater (*C. desquamator*) in the SSI

radiation and the zooplanktivore (*C. simus*) in the Chichancanab radiation. Although these are

distinct dietary specializations, they share a similarity in that they require tracking free-

swimming prey. Indeed, four of the top 15 enriched GO terms of candidate genes in the SSI

scale-eater were related to eye development (26). Similarly, several of the candidate genes we

identified in the zooplanktivore were related to eye development (*ASIC3*, *MAB21L2*, and

CRIM1). In zebrafish, a two-basepair deletion of *CRIM1* results in misshapen lenses as well as a

large eye-to-head ratio (66). Interestingly, one of the defining morphological features of *C. simus*

is its large eye-to-head ratio (33) (Figure 1A). *CRIM1* is also a promising candidate because it is

one of the few candidate genes with multiple nonsynonymous mutations (Figure 2B).

Consistent with candidate genes related to eye development and trophic specialization,

C. simus does indeed exhibit a stronger optomotor response compared to *C. beltrani*, indicating

that this specialist has greatly improved visual acuity to detect rapidly rotating black-and-white

bars.

Although both radiations contained many vision-related candidate genes, there was zero

overlap in these genes. Most notably, we found several adaptive variants associated with opsins

(*SWS2*, blue-sensitive, and *OPN1LW*, red-sensitive) in the zooplanktivore. However, in the SSI

radiation, opsins do not appear to be under selection in any species (26). This result is somewhat

surprising given that opsins appear to be evolutionarily labile and rapidly evolve in other fish

lineages (e.g., sticklebacks (67), cichlids (68, 69), wrasses (70), reviewed in (71)). One plausible

explanation could be that SSI lake environments are relatively shallow, so differences in depth—

a common selective pressure on opsin evolution (71)—are likely not a strong selective force on

SSI. Alternatively, the main selective force acting on opsins may be prey capture efficiency. In

many fish species, zooplanktivory is associated with higher expression of shorter wavelength

opsins (e.g., *SWS1* (71)), whereas piscivory is not often associated with changes in opsin

expression (68, 69). Interestingly, both opsins contain many candidate de novo mutations, and

SWS2 is the youngest estimated sweep in the zooplanktivore. One explanation could be that most

Caribbean *Cyprinodon* species are benthic and live in similar habitats, resulting in limited

standing genetic variation in opsins.

Non-parallelism of sexual selection: the relative role of pre- and post-mating sexual selection

Although the role that premating sexual selection plays in speciation has historically been

the main focus (72, 73), there is both theoretical and empirical support for the role of postmating

sexual selection in maintaining reproductive isolation via postmating prezygotic reproductive

barriers (74–79). In the SSI radiation, species differ in pigmentation, and a handful of the

candidate adaptive variants fall in regulatory regions of genes associated with pigmentation (26).

Furthermore, there is evidence for premating isolation by female preference in the scale-eater

(80). However, in Chichancanab, *C. labiosus* does not visually distinguish between different

species in the radiation (81). Unlike the SSI radiation, we did not find any candidate adaptive

variants associated with pigmentation; instead, we identified adaptive variants associated with

spermatogenesis (*ERC2*, *TBC1D25*, *C4orf22*, *STK35*). Consistent with previous work

demonstrating the importance of these genes in spermatogenesis (82–85), we find that *C. simus*

does indeed have larger sperm heads than *C. beltrani*, suggesting a potential role for post-mating

prezygotic reproductive barriers.

Parallelism in the final stages of adaptation: refinement with de novo and nonsynonymous

mutations

Adaptation may occur in stages, with less pleiotropic cis-regulatory changes occurring

before more pleiotropic coding changes (26, 86, 87). Selection is also predicted to act on

standing variation first due to the waiting time for de novo mutations to arise within a population

(14, 88). In both radiations (Richards et al., 2021), we observe strong evidence for these stages of

adaptation.

The simplest explanation for why we see parallelism in de novo mutations sweeping later

could be due to the waiting time required for a beneficial mutation to (1) arise and (2) reach a

substantial frequency within the population (11, 14). Although de novo and nonsynonymous

mutations were not statistically associated across all candidate variants, many of the late

sweeping adaptive variants are both de novo and nonsynonymous. These variants may also

sweep late because nonsynonymous mutations are often pleiotropic and deleterious in an

ancestral genetic background (14, 89). Thus, they may not be advantageous until later in the

adaptive walk, where compensatory epistasis may mediate negative pleiotropic effects (90–94).

Our results highlight that this final refinement stage is likely a general pattern to be expected

across adaptive radiations.

Methods

Sampling

We collected Laguna Chichancanab specimens in 2022 over two days using a 5 x 1.3 m

seine net with 1.6 mm mesh size. We sampled two sites, an entrance road by La Presumida, and a

bridge crossing the narrow section of the lake on the road to San Diego. *C. labiosus* and *C.*

beltrani were sampled from both sites for this study, whereas *C. simus* was only collected at La

Presumida. *C. artifrons* were collected by cast net from the coastal estuary at Sisal. Additional

specimens were collected from the Bahamas (Lake Cunningham, New Providence Island and

Crescent Pond, San Salvador Island), the Dominican Republic (Laguna Bavaro), or Fort Fisher,

North Carolina as described previously (Martin 2016) or sourced from the American Killifish

Association, the London Zoological Society, the Dallas Children's Aquarium, and Frans

Vermuelen. All specimens were euthanized in an overdose of buffered MS-222 (Fiquel, Inc.)

following approved animal care and use protocols from the University of California, Berkeley

and the University of California, Davis.

Sequencing, Variant calling and filtering

Individual DNA samples were resequenced using Illumina Hiseq 4000 and Novaseq. Raw

reads were trimmed with *fastp* (v0.23.4) (95). Reads were mapped to the UCB_CyNevMio_1.0

Cyprindon nevadensis mionectes reference genome (GCA_030533445.1) with *bwa mem*

(v0.7.17) (96). Duplicate reads in the bam files were marked with *Picard MarkDuplicates*

(GATK v4.5.0.0) (118). Coverage and mapping quality were assessed with *qualimap* (v2.2.2)

(97). We followed the *GATK* (v.4.5.0.0) genotyping pipeline to call variants (119). We used

HaplotypeCaller (-ERC GVCF) v4.5.0.0 to call variants for each individual and stored variants

in the GenomicsDB datastore format prior to using *GenotypeGVCFs* to perform joint

genotyping. We restricted our analyses to biallelic SNPs and applied the recommended *GATK*

hard filters (QD < 2, QUAL < 30, SOR > 3, FS > 60, MQ < 40, MQRankSum < -12.5,

ReadPosRankSum < -8) to filter SNPs (118, 120). Missing genotypes were reset to ./. from 0/0

due to how *GATK* represents missing genotypes in version 4.5.0.0 (101).

After hard-filtering and variant calling, we filtered the data set further by removing SNPs

with minor allele frequency < 0.05, more than 10% missing, depth less than 10x, and GQ less

than 20 using *BCFtools* (v1.16)(102). This resulted in a final data set that included 23,960,536

SNPs.

Population structure of Yucatán Cyprinodon pupfishes

To assess population structure, we first pruned SNPs in linkage disequilibrium using

PLINK (v. 1.9) with the following parameters: "-indep-pairwise 50 5 0.2" (103). This filtered our

data set from 23,960,536 SNPs to 5,476,855 SNPs. With this linkage disequilibrium pruned data

set, we first filtered the full VCF file to only include relevant Yucatán *Cyprinodon* species (i.e.,

C. artifrons, C. beltrani, C. labiosus, and *C. simus*), then conducted a PCA using *PLINK*. Next,

we used *ADMIXTURE* (v1.3) to determine the optimal number of population clusters that best fit

the data and to assign individuals to their corresponding population clusters (104). We performed

this analysis and calculated cross-validation error for the number of clusters K = 1-6. Models in

ADMIXTURE with K=2-4 were equally supported (Figure S2A). However, given morphological

distinctness and the results of the PCA, we present the results for when K = 4 (Figure 1D).

Demographic history of Yucatán Cyprinodon pupfish

We estimated demographic history using *MSMC2* (105). For each Yucatán pupfish

species, we selected three individuals with the highest mean depth and generated single-sample

VCF files, individual mask files, and "mapability" mask files. We then ran *MSMC2* with default

parameters except we set phasing to "unphased" and "P_PAR=8*1+25*1+1*2+1*3." For

plotting, we used a generation time of 1 year and a mutation rate of 1.56×10^{-8} substitutions per

base pair, estimated from Caribbean *Cyprinodon* species (26).

Identifying candidate genes

We calculated F_{st} genome-wide, per site, and in non-overlapping 10-kb windows using

the "--weir-fist-pop" function in *VCFtools*(v0.1.16) for all pairwise combinations of Yucatán

Cyprinodon species (106). We chose 10-kb windows because linkage disequilibrium was

substantially decayed at this distance (Figure S3), and these windows allowed us to quantify

finer-scale genomic variation. With the 10-kb windowed F_{st} values, we calculated a modified

population branch statistic (PBS) for four taxa, PBS_{nj} (107). This statistic is useful because it

makes no assumptions about species topology or polarization (107). We then normalized PBS_{nj}

by the total tree length, as normalized *PBS* is most effective and specific at identifying selective

sweeps (40). We classified SNPs as candidates if they were fixed or nearly fixed (per-site F_{st}

>0.95) compared to other species within the radiation and were among the top 1% outliers out of

all PBS_{nj} windows, following the threshold used in (40).

Introgression

Using the full VCF file, which contained a wide range of species in the family

Cyprinodontidae, we tested for evidence of introgression in Yucatán *Cyprinodon* species using

Dsuite (43). We first excluded any samples/species that had an average depth < 4. We then

calculated all possible trios with a species tree based on the most recent phylogeny (108) with

Cualac tessellatus as the outgroup. We calculated Z-scores to assess the significance of *D*

statistics using 1,000 jackknife blocks and considered a Z-score greater than three as evidence

for significant introgression (109). We only focused our primary analysis on species trios where

P1-P2 included species within the Laguna Chichancanab radiation. We used the other trio

calculations to calculate f-branch (f_b) statistics to account for correlated f_4-ratio scores and to

inform the timing of introgression events (43).

For species trios that showed evidence of introgression where *C. simus* was either P1 or

P2, we then calculated the distance fraction (d_f), a robust metric of introgression for windowed

analyses (44), in non-overlapping windows of 91 informative SNPs. We chose this number of

SNPs because it resulted in windows of approximately 10 kb. We considered windows to be

introgressed if they were in the top X% of absolute d_f value, where X% was the percentage of

estimated genome wide introgression (e.g, f_4-ratio) (110, 111). To estimate introgression block

sizes, we merged neighboring windows that were both introgression outliers and took the total

size of the blocks.

Characterizing candidate genes

After identifying candidate SNPs, we then used *Bedtools* (v2.31) to see if SNPs occurred

near (20-kb) or within a gene (112) with a GFF annotation file for the *C. nev. mionectes*

reference genome (Tian et al. in prep). We classified SNPs within 20 kb of a gene (upstream or

downstream) as flanking. For SNPs within a gene, we further classified them into intronic,

synonymous, or nonsynonymous using *SnpEff* (113).

To characterize the origin of candidate SNPs, we extracted the alleles from every species

for each candidate SNP. We classified variants that were found in any other species as "standing

variation". We further subdivided this group into variants that were found in other species but not

in *C. artifrons*. If "standing variation" variants fell within an introgression window, we classified

them as "introgressed." Finally, if the alternative variant was only found in Chichancanab

species, we classified it as de novo.

Timing of selective sweeps

To gain insight into the timing of selective sweeps in *C. simus*, we used *starTRMCA* (45).

Because many genes contained multiple candidate SNPs, for this analysis, we selected the

variant that was in the median position of each gene. We then extracted a 1-Mb region

surrounding that variant and removed all sites that contained missing data. For simplification, we

only used the generalist/detritivore, *C. beltrani*, as the "reference" population and the

zooplanktivore, *C. simus*, as the "selected" population. We used a fixed recombination rate of

$2\text{x}10^{-8}$ (swordtail fishes; (114)) and a mutation rate of $1.56\text{x}10^{-8}$ substitutions per base pair

(Caribbean *Cyprinodon* species; 26). We then ran five separate Markov chains for 30,000

iterations with a proposal standard deviation of 150 (preliminary analyses changing this

parameter had minimal influence on results). We then discarded the first 10,000 iterations of each

chain as burn-in.

To test for stages of adaptation, we conducted separate generalized linear regressions to

test if the sweep age was correlated with the proportion of nonsynonymous SNPs, proportion of

de novo SNPs, and the number of SNPs. For the generalized linear regressions of proportion of

nonsynonymous SNPs and proportion of de novo SNPs, we used a quasi-binomial family to

account for overdispersion with a logit link function. For the number of SNPs, we used a quasi-

Poisson family with a log link function. To test for the significance of sweep age, we used a

likelihood ratio test comparing a model with the estimated sweep age as an effect compared to a

null model.

Optomotor response

We tested for differences in visual acuity between the generalist/detritivore, *C. beltrani*,

and the zooplanktivore, *C. simus*, by conducting an optomotor response behavioral assay. We

tested lab-reared *C. beltrani* (F1, $n = 7$) and lab-reared *C. simus* (long-term laboratory colony

exceeding ten generations in the lab, $n = 5$). Each fish was tested consecutively with each control

or treatment for 1-minute observation periods, with the control observation occurring first,

followed by presentation of all-black and black-and-white spinning bars. The order of

presentation for the latter to treatments was alternated in each trial. Fish were placed in a

suspended cylindrical plastic container 0.3 m in diameter that remained stationary during the

course of the trials. Spinning black-and-white or all-black paper was rotated outside the clear

plastic container at an RPM of ~90rpm during each observation period.

Due to the data being nonparametric (count, zero-inflated, unequal variance between

groups), we conducted a Wilcoxon rank sum test for the on-banded (all black) and on-banded

portions separately. Because we had an a priori hypothesis that *C. simus* would display a stronger

optomotor response (greater visual acuity), we calculated *p*-values with a one-tailed test.

Sperm morphology

To test for differences in sperm morphology, we collected sperm samples from five

individuals per species in the lab (*C. simus, C. artifrons, C. beltrani*). We first anesthetized the

fish with a solution of MS-222. After the fish was anesthetized, we collected 1μL of milt (fish

semen) with a 1μL glass microcapillary tube and fixed the sample overnight in a 4% PFA

solution stained with Rose Bengal. We then plated sperm on a positively charged slide with a

sealed coverslip. We imaged sperm with oil immersion with a 63x objective and measured 10-30

sperm per individual in ImageJ (115). Due to poor sperm samples from some individuals (little

sperm), we ended up with the following sample sizes: three *simus*, four *artifrons*, and five

beltrani.

To retain information about variation within individuals, we fit separate linear mixed-

effects models for each trait of interest (sperm head, midpiece, and flagellum lengths) with

population as a fixed effect and a random intercept for individual ID, using *LME4* (116). To

assess the significance of the model, we used a Type II ANOVA with a Wald F test using

Kenward-Roger degrees of freedom. If there was a significant effect of population, we conducted

post hoc pairwise tests using the *Emmeans* package (117) and corrected for multiple comparisons

using the Tukey method. We did not correct for phylogeny due to how closely related the species

are, the limited sample size, and the fact that the only significant differences were between sister

species.

Acknowledgments

MCK was supported by a Miller Postdoctoral Research Fellowship from the Miller

Institute for Basic Research in Science, University of California Berkeley. This research was

funded by NSF CAREER 1749764, NIH 5R01DE027052-02, Daphne and Ted Pengelley Award

from the Center for Population Biology, and ARCS Foundation awards to CHM. We thank the

Gerace Research Center and Troy Day for logistical support in the Bahamas, and the governments

of Mexico and the Bahamas BEST commission for permission to collect and export samples

(Permit No. PPF/DGOPA-001/20). All research procedures and animal care protocols (AUP-2021-

02-14062-1 and AUP-2021-07-14515) were approved by the University of California, Berkeley

Animal Care and Use committee and the UC Davis Institutional Animal Care and Use committee.

References

1. R. G. Gillespie, *et al.*, Comparing Adaptive Radiations Across Space, Time, and Taxa. *J*
Hered. **111**, 1–20 (2020).

2. C. H. Martin, E. J. Richards, The Paradox Behind the Pattern of Rapid Adaptive
Radiation: How Can the Speciation Process Sustain Itself Through an Early Burst? *Annu.*
Rev. Ecol. Evol. Syst. **50**, 569–593 (2019).

3. Y. Naciri, H. P. Linder, The genetics of evolutionary radiations. *Biol. Rev.* **95**, 1055–1072
(2020).

4. D. Schluter, *The Ecology of Adaptive Radiation* (OUP Oxford, 2000).

5. J. T. Stroud, J. B. Losos, Ecological Opportunity and Adaptive Radiation. *Annu. Rev. Ecol.*
Evol. Syst. **47**, 507–532 (2016).

6. R. Butlin, *et al.*, What do we need to know about speciation? *Trends Ecol. Evol.* **27**, 27–39
(2012).

7. J. A. Coyne, H. A. Orr, *Speciation* (Oxford University Press, 2004).

8. D. R. Matute, B. S. Cooper, Comparative studies on speciation: 30 years since Coyne and
Orr. *Evolution* **75**, 764–778 (2021).

9. D. Schluter, L. H. Rieseberg, Three problems in the genetics of speciation by selection.
Proc. Natl. Acad. Sci. U.S.A. **119**, e2122153119 (2022).

10. D. Berner, W. Salzburger, The genomics of organismal diversification illuminated by
adaptive radiations. *Trends Genet.* **31**, 491–499 (2015).

11. D. A. Marques, J. I. Meier, O. Seehausen, A Combinatorial View on Speciation and
Adaptive Radiation. *Trends Ecol. Evol.* **34**, 531–544 (2019).

12. J. B. Pease, D. C. Haak, M. W. Hahn, L. C. Moyle, Phylogenomics Reveals Three Sources
of Adaptive Variation during a Rapid Radiation. *PLOS Biol.* **14**, e1002379 (2016).

13. B. W. Stone, C. A. Wessinger, Ecological Diversification in an Adaptive Radiation of
Plants: The Role of De Novo Mutation and Introgression. *Mol. Biol. Evol.* **41**, msae007
(2024).

14. R. D. H. Barrett, D. Schluter, Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38–44 (2008).

15. M. Rousselle, *et al.*, Is adaptation limited by mutation? A timescale-dependent effect of genetic diversity on the adaptive substitution rate in animals. *PLOS Genet.* **16**, e1008668 (2020).

16. O. Seehausen, Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19**, 198–207 (2004).

17. K. Kagawa, G. Takimoto, O. Seehausen, Transgressive segregation in mating traits drives hybrid speciation. *Evolution* **77**, 1622–1633 (2023).

18. K. Kagawa, G. Takimoto, Hybridization can promote adaptive radiation by means of transgressive segregation. *Ecol. Lett.* **21**, 264–274 (2018).

19. R. De-Kayne, *et al.*, Genomic architecture of adaptive radiation and hybridization in Alpine whitefish. *Nat. Commun.* **13**, 4479 (2022).

20. D. A. Joyce, *et al.*, Repeated colonization and hybridization in Lake Malawi cichlids. *Curr. Biol.* **21**, R108–R109 (2011).

21. S. Lamichhaney, *et al.*, Evolution of Darwin's finches and their beaks revealed by genome sequencing. *Nature* **518**, 371–375 (2015).

22. C. H. Martin, *et al.*, Complex histories of repeated gene flow in Cameroon crater lake cichlids cast doubt on one of the clearest examples of sympatric speciation. *Evolution* **69**, 1406–1422 (2015).

23. J. I. Meier, *et al.*, Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat. Commun.* **8**, 14363 (2017).

24. C. Pouchon, *et al.*, Phylogenomic Analysis of the Explosive Adaptive Radiation of the Espeletia Complex (Asteraceae) in the Tropical Andes. *Syst. Biol.* **67**, 1041–1060 (2018).

25. E. J. Richards, J. W. Poelstra, C. H. Martin, Don't throw out the sympatric speciation with the crater lake water: fine-scale investigation of introgression provides equivocal support for causal role of secondary gene flow in one of the clearest examples of sympatric speciation. *Evol. Lett.* **2**, 524–540 (2018).

26. E. J. Richards, *et al.*, A vertebrate adaptive radiation is assembled from an ancient and disjunct spatiotemporal landscape. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2011811118 (2021).

27. N. Rosser, *et al.*, Hybrid speciation driven by multilocus introgression of ecological traits. *Nature* **628**, 811–817 (2024).

28. M. E. S. John, J. C. Dunker, E. J. Richards, S. Romero, C. H. Martin, Parallel evolution of integrated craniofacial traits in trophic specialist pupfishes. *Ecol. Evol.* **14**, e11640 (2024).

738   29.   C. H. Martin, P. C. Wainwright, Trophic novelty is linked to exceptional rates of
739   morphological diversification in two adaptive radiations of cyprinodon pupfish. *Evolution*
740   **65**, 2197–2212 (2011).

741   30.   A. H. Patton, E. J. Richards, K. J. Gould, L. K. Buie, C. H. Martin, Hybridization alters
742   the shape of the genotypic fitness landscape, increasing access to novel fitness peaks
743   during adaptive radiation. *eLife* **11**, e72905 (2022).

744   31.   E. J. Richards, C. H. Martin, Adaptive introgression from distant Caribbean islands
745   contributed to the diversification of a microendemic adaptive radiation of trophic
746   specialist pupfishes. *PLOS Genet.* **13**, e1006919 (2017).

747   32.   E. J. Richards, C. H. Martin, We get by with a little help from our friends: shared adaptive
748   variation provides a bridge to novel ecological specialists during adaptive radiation. *Proc.
749   R. Soc. B.* **289**, 20220613 (2022).

750   33.   J. M. Humphries, R. R. Miller, A Remarkable Species Flock of Pupfishes, Genus
751   Cyprinodon, from Yucatán, México. *Copeia* **1981**, 52–64 (1981).

752   34.   D. A. Hodell, J. H. Curtis, M. Brenner, Possible role of climate in the collapse of Classic
753   Maya civilization. *Nature* **375**, 391–394 (1995).

754   35.   U. Strecker, Description of a new species from Laguna Chichancanab, Yucatan, Mexico:
755   Cyprinodon suavium (Pisces: Cyprinodontidae). *Hydrobiologia* **541**, 107–115 (2005).

756   36.   C. A. Gracida-Juárez, J. J. Schmitter-Soto, M. J. Genner, Community structure of
757   indigenous fishes relative to habitat variation and invasive tilapia in lakes of Quintana
758   Roo, Mexico. *Environ. Biol. Fishes* **107**, 401–414 (2024).

759   37.   U. Strecker, The impact of invasive fish on an endemic Cyprinodon species flock
760   (Teleostei) from Laguna Chichancanab, Yucatan, Mexico. *Ecol. Freshw. Fish* **15**, 408–418
761   (2006).

762   38.   U. Strecker, Genetic differentiation and reproductive isolation in a *Cyprinodon* fish species
763   flock from Laguna Chichancanab, Mexico. *Mol. Phylogenetics Evol.* **39**, 865–872 (2006).

764   39.   U. Strecker, C. G. Meyer, C. Sturmbauer, H. Wilkens, Genetic Divergence and Speciation
765   in an Extremely Young Species Flock in Mexico Formed by the
766   Genus*Cyprinodon*(Cyprinodontidae, Teleostei). *Mol. Phylogenetics Evol.* **6**, 143–149
767   (1996).

768   40.   M. Shpak, K. N. Lawrence, J. E. Pool, The Precision and Power of Population Branch
769   Statistics in Identifying the Genomic Signatures of Local Adaptation. *Genome Biol. Evol.*
770   **17**, evaf080 (2025).

771   41.   C. Bragato, *et al.*, Zebrafish dnm1a gene plays a role in the formation of axons and
772   synapses in the nervous tissue. *J. Neurosci. Res.* **101**, 1345–1359 (2023).

42.   C. Gonzaga-Jauregui, *et al.*, Exome Sequence Analysis Suggests that Genetic Burden
Contributes to Phenotypic Variability and Complex Neuropathy. *Cell Reports* **12**, 1169–
1183 (2015).

43.   M. Malinsky, M. Matschiner, H. Svardal, Dsuite - Fast D-statistics and related admixture
evidence from VCF files. *Mol. Ecol. Res.* **21**, 584–595 (2021).

44.   B. Pfeifer, D. D. Kapan, Estimates of introgression as a function of pairwise distances.
BMC Bioinform. **20**, 207 (2019).

45.   J. Smith, G. Coop, M. Stephens, J. Novembre, Estimating Time to the Common Ancestor
for a Beneficial Allele. *Mol. Biol. Evol.* **35**, 1003–1017 (2018).

46.   S. C. F. Neuhauss, *et al.*, Genetic Disorders of Vision Revealed by a Behavioral Screen of
400 Essential Loci in Zebrafish. *J. Neurosci.* **19**, 8603–8615 (1999).

47.   E. D. Enbody, *et al.*, Community-wide genome sequencing reveals 30 years of Darwin's
finch evolution. *Science* **381**, eadf6218 (2023).

48.   J. V. Peñalba, *et al.*, The Role of Hybridization in Species Formation and Persistence. *Cold*
Spring Harb. Perspect. Biol. **16**, a041445 (2024).

49.   O. Seehausen, Conditions when hybridization might predispose populations for adaptive
radiation. *J. Evol. Biol.* **26**, 279–281 (2013).

50.   D. Schluter, Adaptive radiation along genetic lines of least resistance. *Evolution* **50**, 1766–
1774 (1996).

51.   J. A. McGirr, C. H. Martin, Few Fixed Variants between Trophic Specialist Pupfish
Species Reveal Candidate Cis-Regulatory Alleles Underlying Rapid Craniofacial
Divergence. *Mol. Biol. Evol.* **38**, 405–423 (2021).

52.   J. A. McGirr, C. H. Martin, Novel Candidate Genes Underlying Extreme Trophic
Specialization in Caribbean Pupfishes. *Mol. Biol. Evol.* **34**, 873 (2017).

53.   J. A. McGirr, C. H. Martin, Parallel evolution of gene expression between trophic
specialists despite divergent genotypes and morphologies. *Evol. Lett.* **2**, 62–75 (2018).

54.   J. A. McGirr, C. H. Martin, Ecological divergence in sympatry causes gene misexpression
in hybrids. *Mol. Ecol.* **29**, 2707–2721 (2020).

55.   M. F. Palominos, V. Muhl, C. H. Martin, Craniofacial-specific transcriptomics uncovers
novel genes underlying jaw divergence in dietary specialist pupfishes. *Genetics* iyaf207
(2025).

56.   T. Hornemann, *et al.*, The SPTLC3 Subunit of Serine Palmitoyltransferase Generates
Short Chain Sphingoid Bases. *J. of Biol. Chem.* **284**, 26322–26330 (2009).

57. M. E. Bronner, N. M. LeDouarin, Development and evolution of the neural crest: An
overview. *Dev. Biol.* **366**, 2–9 (2012).

58. F. Hirth, H. Reichert, Conserved genetic programs in insect and mammalian brain
development. *BioEssays* **21**, 677–684 (1999).

59. C. J. Reaume, M. B. Sokolowski, Conservation of gene function in behaviour. *Phil. Tran.*
R. Soc. B. **366**, 2100–2110 (2011).

60. A. Abzhanov, M. Protas, B. R. Grant, P. R. Grant, C. J. Tabin, Bmp4 and Morphological
Variation of Beaks in Darwin's Finches. *Science* **305**, 1462–1465 (2004).

61. R. C. Albertson, J. T. Streelman, T. D. Kocher, P. C. Yelick, Integration and evolution of
the cichlid mandible: The molecular basis of alternate feeding strategies. *Proc. Nat. Acad.*
Sci. U.S.A. **102**, 16287–16292 (2005).

62. Y. Terai, N. Morikawa, N. Okada, The Evolution of the Pro-Domain of Bone
Morphogenetic Protein 4 (Bmp4) in an Explosively Speciated Lineage of East African
Cichlid Fishes. *Mol. Biol. Evol.* **19**, 1628–1632 (2002).

63. S. Nawaz, *et al.*, WNT10A missense mutation associated with a complete Odonto-
Onycho-Dermal Dysplasia syndrome. *Eur. J. Hum. Genet.* **17**, 1600–1605 (2009).

64. E. L. Benard, *et al.*, wnt10a is required for zebrafish median fin fold maintenance and
adult unpaired fin metamorphosis. *Dev. Dyn.* **253**, 566–592 (2024).

65. T. A. Square, *et al.*, Modulation of tooth regeneration through opposing responses to Wnt
and BMP signals in teleosts. *Development* **150**, dev202168 (2023).

66. T. Le, *et al.*, A zebrafish model of crim1 loss of function has small and misshapen lenses
with dysregulated clic4 and fgf1b expression. *Front. Cell Dev. Biol.* **13** (2025).

67. D. A. Marques, *et al.*, Convergent evolution of SWS2 opsin facilitates adaptive radiation
of threespine stickleback into different light environments. *PLOS Biol.* **15**, e2001627
(2017).

68. C. M. Hofmann, *et al.*, The Eyes Have It: Regulatory and Structural Changes Both
Underlie Cichlid Visual Pigment Diversity. *PLOS Biol.* **7**, e1000266 (2009).

69. K. E. O'Quin, C. M. Hofmann, H. A. Hofmann, K. L. Carleton, Parallel Evolution of
Opsin Gene Expression in African Cichlid Fishes. *Mol. Biol. Evol.* **27**, 2839–2854 (2010).

70. G. A. C. Phillips, K. L. Carleton, N. J. Marshall, Multiple Genetic Mechanisms Contribute
to Visual Sensitivity Variation in the Labridae. *Mol. Biol. Evol.* **33**, 201–215 (2016).

71. K. L. Carleton, D. Escobar-Camacho, S. M. Stieb, F. Cortesi, N. J. Marshall, Seeing the
rainbow: mechanisms underlying spectral sensitivity in teleost fishes. *J. Exp. Biol.* **223**,
jeb193334 (2020).

72.   M. G. Ritchie, Sexual Selection and Speciation. *Annu. Rev. Ecol. Evol. Syst.* **38**, 79–102
(2007).

73.   M. R. Servedio, J. W. Boughman, The Role of Sexual Selection in Local Adaptation and
Speciation. *Annu. Rev. Ecol. Evol. Syst.* **48**, 85–109 (2017).

74.   A. Devigili, *et al.*, Possible glimpses into early speciation: the effect of ovarian fluid on
sperm velocity accords with post-copulatory isolation between two guppy populations. *J.*
Evol. Biol. **31**, 66–74 (2018).

75.   M. D. Garlovsky, *et al.*, Synthesis and Scope of the Role of Postmating Prezygotic
Isolation in Speciation. *Cold Spring Harb. Perspect. Biol.* a041429 (2023).

76.   M. C. Kustra, M. R. Servedio, S. H. Alonzo, Cryptic female choice can maintain
reproductive isolation. *Evolution* **79**, 2259–2273 (2025).

77.   M. C. Kustra, S. H. Alonzo, The coevolutionary dynamics of cryptic female choice. *Evol.*
Lett. **7**, 191–202 (2023).

78.   P. D. Lorch, M. R. Servedio, The evolution of conspecific gamete precedence and its effect
on reinforcement. *J. Evol. Biol.* **20**, 937–949 (2007).

79.   J. L. Marshall, M. L. Arnold, D. J. Howard, Reinforcement: the road not taken. *Trends*
Ecol. Evol. **17**, 558–563 (2002).

80.   A. Kodric-Brown, R. J. D. West, Asymmetries in premating isolating mechanisms in a
sympatric species flock of pupfish (Cyprinodon). *Behav. Ecol.* **25**, 69–75 (2014).

81.   U. Strecker, A. Kodric-Brown, Mate recognition systems in a species flock of Mexican
pupfish. *J. Evol. Biol.* **12**, 927–935 (1999).

82.   M. H. K. Chau, *et al.*, Investigation of the genetic etiology in male infertility with
apparently balanced chromosomal structural rearrangements by genome sequencing. *Asian*
J. Androl. **24** (2022).

83.   H. Li, Y. Dai, Z. Luo, D. Nie, Cloning of a new testis-enriched gene C4orf22 and its role
in cell cycle and apoptosis in mouse spermatogenic cells. *Mol. Biol. Rep.* **46**, 2029–2038
(2019).

84.   Y. Miyamoto, *et al.*, The STK35 locus contributes to normal gametogenesis and encodes a
lncRNA responsive to oxidative stress. *Biol. Open* **7**, bio032631 (2018).

85.   S. Nawaz, S. Hussain, S. Basit, W. Ahmad, First evidence of involvement of *TBC1D25* in
causing human male infertility. *Eur. J. Med. Genet.* **64**, 104142 (2021).

86.   L. Campagna, *et al.*, Repeated divergent selection on pigmentation genes in a rapid finch
radiation. *Sci. Adv.* **3**, e1602404 (2017).

87.   F. C. Jones, *et al.*, The genomic basis of adaptive evolution in threespine sticklebacks.
874   *Nature* **484**, 55–61 (2012).

875   88.   Z. D. Blount, R. E. Lenski, J. B. Losos, Contingency and determinism in evolution:
876   Replaying life's tape. *Science* **362**, eaam5979 (2018).

877   89.   P. W. Hedrick, Adaptive introgression in animals: examples and comparison to new
878   mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* **22**, 4606–
879   4618 (2013).

880   90.   S. G. Das, M. Mungan, J. Krug, Epistasis-mediated compensatory evolution in a fitness
881   landscape with adaptational tradeoffs. *Proc. Natl. Acad. Sci. U.S.A.* **122**, e2422520122
882   (2025).

883   91.   M. J. Harms, J. W. Thornton, Historical contingency and its biophysical basis in
884   glucocorticoid receptor evolution. *Nature* **512**, 203–207 (2014).

885   92.   M. Karageorgi, *et al.*, Genome editing retraces the evolution of toxin resistance in the
886   monarch butterfly. *Nature* **574**, 409–412 (2019).

887   93.   R. D. Tarvin, *et al.*, Interacting amino acid replacements allow poison frogs to evolve
888   epibatidine resistance. *Science* **357**, 1261–1266 (2017).

889   94.   D. M. Weinreich, N. F. Delaney, M. A. DePristo, D. L. Hartl, Darwinian Evolution Can
890   Follow Only Very Few Mutational Paths to Fitter Proteins. *Science* **312**, 111–114 (2006).

891   95.   S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
892   *Bioinformatics* **34**, i884–i890 (2018).

893   96.   H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
894   [Preprint] (2013). Available at: http://arxiv.org/abs/1303.3997.

895   97.   K. Okonechnikov, A. Conesa, F. García-Alcalde, Qualimap 2: advanced multi-sample
896   quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292–294 (2016).

897   98.   A. McKenna, *et al.*, The Genome Analysis Toolkit: A MapReduce framework for
898   analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

899   99.   M. A. DePristo, *et al.*, A framework for variation discovery and genotyping using next-
900   generation DNA sequencing data. *Nat. Genet.* **43**, 491–498 (2011).

901   100.   G. A. Van der Auwera, *et al.*, From FastQ Data to High-Confidence Variant Calls: The
902   Genome Analysis Toolkit Best Practices Pipeline. *Curr. Prot. Bioinf.* **43**, 11.10.1-11.10.33
903   (2013).

904   101.   D. Caetano-Anolles, GenotypeGVCFs and the death of the dot (obsolete as of GATK
905   4.6.0.0). *GATK* (2024).

102. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008
(2021).

103. S. Purcell, *et al.*, PLINK: A Tool Set for Whole-Genome Association and Population-
Based Linkage Analyses. *Amer. J. Hum. Genet.* **81**, 559–575 (2007).

104. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in
unrelated individuals. *Genome. Res.* **19**, 1655–1664 (2009).

105. S. Schiffels, K. Wang, "MSMC and MSMC2: The Multiple Sequentially Markovian
Coalescent" in *Statistical Population Genomics*, J. Y. Dutheil, Ed. (Springer US, 2020),
pp. 147–166.

106. P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

107. J. M. Schmidt, M. de Manuel, T. Marques-Bonet, S. Castellano, A. M. Andrés, The impact
of genetic adaptation on chimpanzee subspecies differentiation. *PLOS Genet.* **15**,
e1008485 (2019).

108. S. G. Hernández-Ávila, C. W. Hoagstrom, W. A. Matamoros, Historical biogeography of
North American killifishes (Cyprinodontiformes) recapitulates geographical history in the
Gulf of México watershed. *Zool. J. Linn. Soc.* **202**, zlae105 (2024).

109. E. Y. Durand, N. Patterson, D. Reich, M. Slatkin, Testing for Ancient Admixture between
Closely Related Populations. *Mol. Biol. Evol.* **28**, 2239–2252 (2011).

110. C. Feng, J. Wang, A. Liston, M. Kang, Recombination Variation Shapes Phylogeny and
Introgression in Wild Diploid Strawberries. *Mol. Biol. Evol.* **40**, msad049 (2023).

111. A. Morales-Cruz, *et al.*, Introgression among North American wild grapes (Vitis) fuels
biotic and abiotic adaptation. *Genome Biol.* **22**, 254 (2021).

112. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).

113. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster
strain w1118; iso-2; iso-3. *Fly* **6**, 80–92 (2012).

114. M. Schumer, *et al.*, Natural selection interacts with recombination to shape the evolution
of hybrid genomes. *Science* **360**, 656–660 (2018).

115. C. T. Rueden, *et al.*, ImageJ2: ImageJ for the next generation of scientific image data.
BMC Bioinform. **18**, 529 (2017).

116. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using
lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

117. R. V. Lenth, *emmeans: Estimated Marginal Means, aka Least-Squares Means* (2021).

118. G. A. Van der Auwere & B. D. O' Connor. *Genomics in the cloud: using docker, GATK, and*
WDL in Terra. (Sebastopol, CA O'Reilly Media, 2020).

119. A. McKenna *et al.,* The genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res.* **20**, 1297-1303 (2010).

120. M. A. DePristo *et al.,* A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat. Genet.* **43**, 491-498 (2011).

967   **Supplementary materials**

969   **Table S1.** Mean genome-wide pairwise F_{ST} statistics within the Chichancanab radiation and with
970   sister outgroup (*artifrons*).

	labiosus	*beltrani*	*artifrons*
simus	0.090	0.068	0.182
labiosus		0.032	0.184
beltrani			0.168

972   **Table S2.** F_{ST} statistics within the Chichancanab radiation.

Comparison	$F_{ST} = 1$ (fixed)	$1 > F_{ST} \geq 0.95$ (nearly fixed)	$0.95 > F_{ST} \geq 0.9$	$0.9 > F_{ST} \geq 0.8$	Other	Mean F_{ST}
simus vs *beltrani* + *labiosus*	19	1127	3304	20173	10359605	0.07
labiosus vs *beltrani* + *simus*	0	0	3	94	10384131	0.04
beltrani vs *simus* + *labiosus*	0	0	0	13	10384215	0.02

986 **Table S3.** Fixed SNPs between *C. simus* and *C. beltrani* + *C. labiosus*. Bold rows indicate SNPs within top 1% of PBS windows.

987 Bolded genes only are genes that have regions that fall within the top 1% PBS percentile.

Position	Gene	Distance to gene	PBS percentile	Annotation
Chr 1: 5290600	**STK35**	12594	1.85	
Chr 1:5290602	**STK35**	12592	1.85	Fertility, eye development and oxidative stress (Miyamoto et al 2018).
Chr 1:5290604	**STK35**	12590	1.85	
Chr 2: 17498913	SLC17A8	Intron	1.24	Glutamate neurotransmitter, inner hair cell dysfunction
Chr 3: **11687002**	**AGO3**	**Intron**	**0.34**	**RNA regulator**
Chr 3: 32286614	RBMS3	14363	22.93	
Chr 3: 32286627	RBMS3	14350	22.93	Involved in chondrogenic craniofacial development of zebrafish
Chr 3: 32286638	RBMS3	14339	22.93	(Jayasena and Bronner 2012).
Chr 3: 32286614	RBMS3	14363	22.93	
Chr 11: 27324192	GRIA3	Intron	2.85	
Chr 11: 27336469	GRIA3	2772	3.18	Glutamate receptor strongly expressed in zebrafish retina
Chr 11: 27336472	GRIA3	2775	3.18	development.
Chr 11: 27336476	GRIA3	2779	3.18	
Chr 11: **28255536**	**NA**	**3026**	**0.98**	**NA**
Chr 11: **28255616**	**NA**	**3106**	**0.98**	**NA**
Chr 11: **28256343**	**NA**	**3833**	**0.98**	**NA**
Chr 11: **28257367**	**NA**	**4857**	**0.98**	**NA**
Chr 11: 28260047	NA	7537	2.87	NA
Chr 12: 13854139	**C4orf22**	11903	1.30	Spermatogenesis, cilia, flagellum
Chr 24: **12693158**	**KDM6A**	**Intron**	**0.05**	**Kabuki syndrome (craniofacial development)**

989　**Table S4.** Summary statistics of the distance between closest adaptive candidate SNPs within the same

990　gene

Gene	Median Distance	Mean Distance	SD	
TM6SF2	7430	7430	NA	2
C19orf10	4789	4789	NA	2
KDM6A	2892.5	2892.5	2410.52702	3
KCNB2	1803	1738.75	1614.92299	5
UBE2T	1742	1742	NA	2
TMEM230	1734	3039	3599.52427	4
ABCA1	1037.5	1097.5	959.375665	5
HAPLN4	974	1608.21739	9927.05273	24
GRIN2A	904	1270.6667	4273.38008	4
DPH1	608	1224.5	1999.25934	7
WNT10A	402	1442.17857	3986.39626	29
OPN1LW	386.5	359.5	225.229513	5
NIPSNAP3A	253	1451.71429	2819.19065	8
GALNT15	246.5	675.166667	3156.86746	13
SWS2	246	1381.25	2423.03892	5
STK35	217	793.366667	1992.8543	31
CRIM1	201	47.829268	2176.05381	42
PDYN	187	326.371429	355.925561	36
ADCYAP1R1	186.5	833	1304.35501	11
DNM1	186.5	459.708333	677.261781	25
DNM1	186.5	459.708333	677.261781	25
KCNH8	131.5	584.366667	1030.65532	31
TOP1	124	332.375	6687.63393	17
AGO3	115	357.625	552.097284	9
GMEB2	112	319.490196	2485.05809	52
GPNMB	57	57	77.781746	3
IGFBP4	26	2254	3880.67997	4
FAM83D	23	2523.66667	4347.74658	4
EXOSC3	17	17	NA	2
C4orf22	15	15	9.899495	3
TMEM106B	7	11.4	10.358571	6
THOC3	6	5.833333	3.600926	7
MID1	1	1	NA	2

Figure S1. Best fitting admixture graph assuming zero admixture events. Admixture graph

was found using the *findGraphs* function in *ADMIXTOOLS2* (Maier et al., 2023). Number on

edges represent drift weights.

Figure S2. ADMIXTURE results support genetic differentiation of Yucatán Cyprinodon

pupfish species. (A) Cross-validation error for different values of K clusters, equally support 2,

3, and 4 clusters. (B) ADMIXTURE plots showing the clusters at different values of K.

1000

1001

1002

1003

1016

1017   **Figure S3. Linkage decay occurs rapidly in *Cyprinodon simus*.** Linkage decay was calculated

1018   in windows of 1000KB using PLINK (v1.9). The results graphed are a random sample of all

1019   calculated r^2 values taken for plotting purposes ($n = 9{,}645$ r^2 values).

1033

Figure S4. Adaptative SNPs associated with the same gene are generally spaced apart.

Distribution of the distance to nearest adaptive SNP within a gene. Note the log scale of the x-axis.

1037

1038

1039

1040

1041

1042

1043

Figure S5. *f*-branch (*f*_b) show that introgression with desert pupfish occurred before the

Chichancanab radiation. Boxes with red color represent significant introgression between P3

species (top) and the species/internal branch on the left. For visualization, non-significant f_b were

changed to zero and are completely white boxes. Grey boxes represent comparisons that cannot

be made, as f_b cannot be calculated for introgression between sister taxon. Dashed lines on the

left are internal branches. SSI: San Salvador Island, Bahamas; NC: North Carolina, USA.

1050

Figure S6. **Both species did not have a strong response to either control scenario in the optomotor response experiment. (A)** Number of complete rotations fish swam when the machine was off and **(B)** when the machine was on but the apparatus was all black. Boxplots and points of the number of complete rotations individuals performed during the "off-banded" portion of the experiment. Note that some boxplots are compressed to a single line because most individuals had no response.

1068

1069 **Figure S7**. **Species do not differ in sperm midpiece size or sperm flagellum length.** Boxplots

1070 and points of the mean sperm morphological estimate for an individual.

1071