Parallel and non-parallel features of adaptive radiation in Yucatán pupfishes

Matthew C. Kustra[1,2,3,*], David Tian[1,2], M. Fernanda Palominos[1,2], Feifei Guo[1,2], Dylan Chau[1,2], Oskar Golwala[1,2], HoWan Chan[4], Andrés Alvarez Zapata[5], Reyna Guadalupe Cetz Paredes[5], Frida Ximena Cortés Sánchez[6], Sonia Gabriela Hernández[6], Adan Fernando Mar-Silva[6], Fernando Mex[7], Charles Tralka[1,2], Maribel Badillo-Alemán[7], Juan J. Schmitter-Soto[8], Carlos A. Gracida-Juárez[9], Christopher M. Martinez[10], Jairo Arroyave[6], and Christopher H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley, CA, United States

[2]Museum of Vertebrate Zoology, University of California, Berkeley, CA, United States

[3]Miller Institute for Basic Research in Science, University of California, Berkeley, CA, United States

[4]Department of BioSciences, Rice University, Houston, TX, United States

[5]Instituto Tecnologico Superior de Conkal, Yucatan, Mexico

[6]Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, Mexico

[7]Unidad Multidisciplinaria de Docencia e Investigación, Unidad Sisal, Universidad Nacional Autónoma de México, Sisal, Yucatán, Mexico

[8] Departamento de Sistemática y Ecología Acuática, El Colegio de la Frontera Sur, Chetumal, Quintana Roo, Mexico

[9]Subdirección de Posgrado e Investigación, Tecnológico Nacional de México, Campus Felipe Carrillo Puerto.Carretera a Vigía Chico km 1.5, 77200 Felipe Carrillo Puerto, Quintana Roo.

[10]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA, United States

*Corresponding author: matthewckustra@berkeley.edu

Abstract

Understanding the extent of parallelism across adaptive radiations remains a central

problem in evolutionary biology. We used whole-genome resequencing of 123 individuals to

compare an adaptive radiation of *Cyprinodon* pupfishes in Laguna Chichancanab, Mexico, to an

independent radiation of San Salvador Island (SSI) pupfishes in the Bahamas, and assess the

repeatability of adaptive genetic architecture, sources of adaptive variation, and stages of

selection. Despite rapid craniofacial divergence of trophic specialists within 8-15 kya, only two

candidate genes (3.6%) were shared between Caribbean radiations. Although adaptive

introgression played a major role on SSI, we found minimal evidence of adaptive introgression

in Chichancanab, likely due to geographic isolation of this inland lake. Instead, de novo

mutations provided a substantial source of adaptive variation (30.6%) for the endemic

zooplanktivore, 15.3 times higher than the endemic scale-eater on SSI. We found strong evidence

that selection occurred in stages, first on regulatory and standing genetic variation, then on de

novo and nonsynonymous mutations, parallel to SSI. We then verified that adaptive variants near

vision-development and spermatogenesis genes, uniquely within Chichancanab, are consistent

with greater visual acuity and divergent sperm morphology in the zoooplanktivore. Consistent

with extensive adaptive de novo mutations in *WNT10A* and rapid diversification of tooth size in

the zooplanktivore, we found that experimental inhibition of this pathway in generalists results in

narrower teeth. We conclude that de novo mutations, not introgression, can drive rapid adaptive

radiations in isolated environments.

Introduction

Much of biodiversity results from adaptive radiation—when a clade diversifies rapidly into three or more new species (1–5). Young adaptive radiations provide excellent systems for investigating speciation and adaptation (3, 6–9). However, it remains unclear to what degree radiations occur in parallel given similar starting positions on the adaptive landscape with similar levels of genetic diversity, gene flow, and ecological opportunity (10, 11).

Beneficial de novo mutations can provide novel variation to reach new fitness peaks on the adaptive landscape during adaptive radiation (12, 13); however, the longer times needed for de novo beneficial mutations to arise contradicts the short timescales of rapid divergence (11, 14, 15). Hybridization can introduce large amounts of genetic variation and novel phenotypes (transgressive segregation) through adaptive introgression and hybrid swarms, enabling populations to reach previously inaccessible fitness peaks (16). Support for an important role of hybridization in adaptive radiations has been demonstrated both theoretically (17, 18) and empirically in many systems (3, 19–27). However, we still do not know the relative importance of different sources of genetic variation for triggering and sustaining adaptive radiations across time and space.

Caribbean pupfishes are a remarkable system for assessing the repeatability of adaptive radiations due to two independent, recent adaptive radiations each containing a generalist and multiple trophic specialists within isolated lake environments that have historically lacked predatory fishes and most competitors except *Gambusia*. The San Salvador Island (SSI), Bahamas radiation proceeded predominantly via the reassembly of pre-existing standing genetic variation found throughout the Caribbean, with substantial contributions of adaptive introgression to both the molluscivore (durophage) and scale-eating (lepidophage) specialists.

Only 0% and 2% of adaptive mutations in the molluscivore and scale-eater, respectively, were de

novo mutations unique to SSI (26, 28–32). However, it previously was unknown how a second

radiation in Laguna Chichancanab, Mexico, drawing from the same Caribbean pool of standing

genetic variation unfolded over the same timeframe.

Here, we use whole-genome resequencing of 123 individuals to provide insight into the

genomic architecture and genetic origins of the Laguna Chichancanab *Cyprinodon* pupfish

radiation (33). This recent adaptive radiation (~8000 years old (34)) consisted of at least five

species historically(33, 35, 36), including a detritivore (*C. beltrani*), piscivore (*C. maya*), rocky

substrate specialist (*C. verecundus*), bivalve specialist (*C. labiosus*), and zooplanktivore (*C.*

simus). However, due to recent colonization by native fishes not historically present in the lake

and invasive colonization by African tilapia (*Oreochromis niloticus x mossambicus*) (37, 38), all

trophic specialist species now appear to be extinct since our collections in 2022, with only the

generalist, *C. beltrani*, remaining in 2024 (MK, CM pers. obs.) and 2025 (CG, JSS pers. obs.).

Despite exhibiting one of the fastest rates of trait diversification of any vertebrate in tooth size

(130 times faster than background rates across *Cyprinodon* (29)) and rapid divergence in other

craniofacial morphology (Figure 1A; S1), previous genetic studies of microsatellite data and

mtDNA suggest that this radiation has low genetic differentiation (39, 40), enabling the

identification of highly differentiated loci among these species.

Results

Genetic differentiation of the species flock

To measure population structure of the Yucatán *Cyprinodon* pupfishes, we collected

specimens from the wild in 2022 (Figure 1B) and conducted whole genome resequencing of the

species flock and the coastal sister species ($n = 15$ C. artifrons, $n = 35$ C. beltrani, $n = 35$ C.

labiosus, and $n = 17$ C. simus). We aligned these samples, as well as eleven other previously

sequenced Caribbean pupfish species, to the high-quality *Cyprindon nevadensis mionectes*

reference genome (GCA_030533445.1). After calling and filtering variants, our dataset included

23,960,536 SNPs (12.4x mean depth). Using a linkage disequilibrium pruned set of SNPs ($n =$

5,476,855 SNPs), we used principal component analysis and ADMIXTURE to visualize

population structure within Yucatán pupfishes. We found that all species formed distinct, non-

overlapping genetic clusters, with *C. artifrons* being the most genetically divergent from and

sister to the Chichancanab species (Figure 1C). Within the Chichancanab species flock, *C. simus*

was more genetically divergent from *C. labiosus* and *C. beltrani* (Figure 1C). These results were

also supported by genome-wide F_{ST} estimates and a phylogenetic tree constructed with

ADMIXTOOLS2 (Table S1; Figure S2). Four clusters corresponded to the four different species,

as determined by ADMIXTURE (Figure 1D). All *C. artifrons* individuals showed minimal

evidence of admixture with Chichancanab species (Figure 1D).

Demographic history

Using MSMC2 for Chichancanab species with a mutation rate of 1.56×10^{-8} substitutions

per base pair (estimated from a Caribbean *Cyprinodon* species (26)), we found that the three

Chichancanab species (*C. beltrani*, *C. labiosus*, and *C. simus*) diverged from the coastal sister

species *C. artifrons* at the end of the Pleistocene 10-15 kya, consistent with receding sea levels

resulting in the formation of the Chichancanab lake (34). After divergence from *C. artifrons*, the

Chichancanab species flock shows a similar demographic history, characterized by a sharp

decline in effective population sizes, consistent with endemic sympatric species within an

isolated lake (Figure 1E).

127
Figure 1. Yucatán *Cyprinodon* pupfishes show strong genetic differentiation. (A) μCT scans
of Yucatán pupfish with maxilla colored in blue, premaxilla in pink, dentary in green, and
articular in orange (see Figure S1 for different views). (B) Map of the Caribbean showing the
sampling sites for all species in our dataset with an inset map of Lake Chichancanab and an
aerial view image from one of the sampling sites. (C) Principal component analysis using a set of
linkage disequilibrium pruned SNPs. (D) Estimated ancestry proportions for each individual
inferred using *ADMIXTURE* with K = 4 (see Figure S3 for K=2-3) from the linkage-pruned
dataset. (E) Estimated demographic history with *MSMC2* on a log scale. Each colored line
represents an individual, with colors corresponding to different species.

Adaptive loci within trophic specialists

Although genome-wide differentiation among Chichancanab species was moderately low

(F_{ST} ranging from 0.032 to 0.09; Figure 2A; Table S1), we were able to identify highly

differentiated regions of the genome within the trophic specialist zooplanktivore *C. simus*. We

identified 19 fixed and 1,127 nearly-fixed ($F_{ST} > 0.95$) SNPs in *C. simus* relative to other

Chichancanab pupfishes (Tables S2-S3). In contrast, we found zero fixed or nearly-fixed SNPs

within the generalist *C. beltrani* or bivalve-specialist C. *labiosus*. However, we were able to

identify some moderately differentiated SNPs within *C. labiosus* ($n = 97$) and *C. beltrani* ($n =$

13) relative to the other Chichancanab species at an F_{ST} threshold of 0.8 (Table S2).

To search for signals of selection, we calculated the normalized four-taxon population

branch statistic (PBS_{nj}; Schmidt et al., 2019) across 10-kb non-overlapping windows because

linkage disequilibrium decayed at this distance (Figure S4). Simulation studies demonstrate that

normalized *PBS* is effective for identifying both hard and soft selective sweeps (41). Out of the

1,146 SNPs fixed or nearly fixed within *C. simus*, 661 were within a strongly supported selective

sweep window (top 1% PBS_{nj} outlier (41)), including 425 SNPs within 20 kb of the first or last

exon of an annotated gene. These 425 candidate adaptive SNPs ("adaptive variants") were

located in the proximity of or within 55 genes (Figure 2A), distributed across 19 out of 24

chromosomes (Figure 2A). Adaptive variants associated with the same gene were often

distributed across multiple introns or exons and spaced apart by 1kb or more (Figure S5, Table

S4).

Source and function of adaptive genetic variation in the zooplanktivore

We next characterized the location of the 425 candidate adaptive variants near or within a

gene in the zooplanktivore *C. simus*. More than 95% were in intronic or flanking regions (20-kb

upstream or downstream of a gene; Figure 2B). Within coding regions, 13 out of 20 adaptive

variants were nonsynonymous mutations (Figure 2B). The number of adaptive variants per gene

followed an exponential distribution, with a handful of genes containing many variants and most

genes with only a single adaptive variant (Figure 2B). We functionally characterized these genes

by manually checking online databases and doing a literature search (Table S3; Extended Table

S1). 32 out of the 55 candidate genes were associated with craniofacial development ($n =11$;

20%) or sensory/neural function ($n = 21$; 38.2%). We manually checked if any of these genes

overlapped with the 426 unique candidate genes identified in the SSI radiation (26, 28, 30, 32,

42–46), and we found only two overlapping candidate genes (*DNM1* and *SPTLC3*), both of

which are associated with motor neuron development (47, 48) and are scale-eater-specific

candidate genes.

Next, we characterized the source of the variants and found that 63% of the adaptive

SNPs in *C. simus* also existed as standing genetic variation in the coastal Yucatan population of

C. artifrons ("standing variation"), 31% were *de novo* variants found only in Chichancanab, and

6% were detected as standing genetic variation in other pupfish species but not *C. artifrons*

(Figure 2C).

Figure 2. Architecture and source of adaptive genetic variation in the zooplanktivore. (A, top row) F_{st} between *C. simus* and *C. beltrani* + *C. labiosus* calculated in non-overlapping 10-kb windows. F_{st} values for candidate SNPs (red circles) were calculated separately per locus to show the location of fixed variants. (A, lower row) Population branch statistics (PBS_{nj}) with *C. simus* as the target population versus *C. beltrani*, *C. labiosus*, and *C. artifrons* in 10-kb non-overlapping windows. Candidate genes within 20 kb of fixed or nearly-fixed (Fst > 0.95) SNPs and within PBS_{nj} 1% outlier windows are highlighted in red and labeled. Alternating colors represent different chromosomes. (B) Candidate adaptive variants near or within a gene were defined as SNPs that were fixed ($n = 17$) or nearly fixed ($n = 1,129$, $F_{st} > 0.95$) and occurred within a 1% population branch statistic (PBS_{nj}) outlier window for *C. simus* (Venn diagram). Pie chart and bar graphs indicate the proportion of these 425 adaptive variants that were found within the flanking, intronic, or coding regions. (C) Pie chart and bar graphs indicate the proportion of these 425 variants that were detected only within Chichancanab (de novo: purple), as standing genetic variation in coastal sister species *C. artifrons* (pink), or in other Caribbean species, but not *C. artifrons*. * Indicates genes with a craniofacial annotation; ‡ indicates genes with neural or sensory annotation.

Limited adaptive introgression within Chichancanab pupfishes

To test for introgression into the Chichancanab radiation, we used *f*-statistics calculated in

Dsuite (49) to examine gene flow from 11 additional *Cyprinodon* species from across the clade

and the sister species to the genus *Megupsilon aporus*. Within Yucatán pupfishes, we found

evidence of introgression between *C. artifrons* and both *C. beltrani* and *C. labiosus*, with the

strongest signal in *C. beltrani* (Figure 3A, B). We also found evidence for introgression between

C. beltrani and most outgroup species examined, including the sister species to *Cyprinodon*,

Megupsilon aporus. Relative to *C. beltrani*, *C. labiosus* and *C. simus* also showed evidence of

introgression with pupfishes endemic to the Chihuahua desert (*C. eximius* and *C. fontinalis*;

Figure 3A, B). Furthermore, we also found evidence of introgression between the two sympatric

radiations (*C. variegatus* from SSI) and *C. simus* (Figure 3A, B). However, additional *f*-branch

statistics indicate that all patterns of excess allele sharing between these outgroup species besides

C. artifrons and the Chichancanab species flock likely resulted from ancient introgression events

before the colonization of Laguna Chichancanab (Figure 3C; S6).

To test for adaptive introgression into *C. simus*, we next examined whether the set of

adaptive variants (Figure 2) occurred within outlier introgression windows using sliding window

scans of distance fraction (d_f; Pfeifer & Kapan, 2019) for species that had significant genome-

wide introgression into *C. simus* (Figure 3A , B). We found that none of the outlier introgression

windows overlapped with *C. simus* adaptive variants (Figure 3D).

Next, we performed *df* scans for introgression between *C. artifrons* and Chichancanab

species. Although there were outlier introgression regions between *C. simus* and *C. artifrons*

(positive values d_f), these regions did not overlap with *C. simus* adaptive variants (Figure 3D).

However, the position of several of these *C. simus* adaptive variants overlapped with outlier

introgression windows for both *C. labiosus* and *C. beltrani* but not *C. simus* (Figure 3D). The

number of outlier introgression windows was much lower in *C. simus* relative to *C. labiosus* and

C. beltrani (Figure 3E, F). Furthermore, both *C. labiosus* and *C. beltrani* populations contained

several introgression tracts that were much longer than *C. simus* tract lengths, indicating much

more recent introgression events in these species (Figure 3E, F).

Figure 3. Minimal evidence of adaptive introgression into Chichancanab pupfishes.
Heatmaps of (A) Patterson's D-statistics and (B) F_4 ratios for P1-P2 species combinations within
the Chichancanab radiation along the x-axis and P3 focal species along the y-axis. All four-taxon
trees used *Cualac tessellatus* as the outgroup (P4). Only significant (Z-score > 3) results are
colored; combinations not relevant to the Chichancanab radiation are not shown. (C) *f*-branch (f_b)
statistics indicate introgression occurring before the Chichancanab radiation. Red boxes indicate
significant introgression between P3 species (top row) and the focal species or internal branch
(dashed lines) on the left. Non-significant f_b statistics are indicated by white boxes. Grey boxes
represent comparisons that cannot be made since f_b cannot be calculated for introgression
between sister taxa. See Figure S5 for full *f*-branch graphs. (D) Genome-wide plot of the distance
fraction (d_f, an introgression test statistic*;* Pfeifer & Kapan, 2019) calculated in non-overlapping
windows of 91 SNPs (approximately 10 kb) for species trios that showed evidence of
introgression with *C. simus*. d_f outlier windows that contained a candidate gene under selection in
C. simus are colored red and labeled with the gene name. The y-axis label gives the introgression
topology: (P1, P2), P3. Positive values indicate regions of introgression between *C. simus* and P3
(from top to bottom: *C. eximius*, *C. fontinalis*, *C. variegatus* (SSI), and *C. artifrons* in the bottom
two subplots); negative values indicate regions of introgression between P1 (*C. beltrani* or *C.
labiosus*) and P3. Red dashed lines indicate outlier cutoffs, which were determined empirically
by the percentage of genome-wide introgression estimated by F_4 ratios in panel B. (E, F)
Distribution of the size of d_f outlier introgression tract lengths (in 10 kb windows) between *C.
artifrons* and Chichancanab species. The introgression tracts containing candidate adaptive genes
in *C. simus* are labeled with gene names. SSI: San Salvador Island, Bahamas; NC: North
Carolina, USA.

The stages of adaptive divergence of the zooplanktivore

We used *starTRMCA* (51) to estimate the timing of selective sweeps for the set of

candidate adaptive variants near genes within the zooplanktivore *C. simus* (Fig. 4). We estimated

that most selective sweeps occurred in a staggered sequence and not simultaneously after

colonization of Laguna Chichancanab (Figure 5A)(34). There were also a few older sweeps that

we estimate may predate the age of Laguna Chichancanab and divergence from the coastal sister

species *C. artifrons* (Figure 1B). Overall, we observed no clear temporal stages of adaptive

phenotypic divergence related to craniofacial genes (asterisks: Figure 5) or neural/sensory genes

(‡: Figure 5). Instead, we found that selection on these two broad functional categories occurred

throughout the period of divergence of *C. simus* following the colonization of Laguna

Chichancanab, contrary to expectations of functional stages of adaptation (52) found in the SSI

radiation (26).

Next, we used generalized linear regressions to test if sweep age was correlated with the

proportion of nonsynonymous adaptive variants, de novo adaptive variants, or the overall

number of adaptive variants per gene region. We found that both nonsynonymous adaptive

variants ($\chi^2 = 5.852$, $p = 0.016$) and de novo mutations ($\chi^2 = 6.221$, $p = 0.013$) were significantly

more likely to occur in more recent selective sweeps during the final stage of adaptive

divergence in *C. simus* (Figure 5B, C). However, nonsynonymous mutations were not

significantly more likely to be de novo mutations found only in Chichancanab species (All

adaptive variants: $\chi^2 = 0.102$, $P = 0.749$; singleton adaptive variants: $\chi^2 = 2.010$, $p = 0.156$), and

there was no correlation between estimated sweep age and the number of candidate adaptive

variants per gene region ($\chi^2 = 0.111$, $p = 0.740$).

Figure 4. Stages of adaptive divergence in the zooplanktivore. (A) Points indicate median
sweep age with 95% credible intervals. The dashed line indicates the approximate age of Laguna
Chichancanab (34). For each sweep, the first column of pie charts shows the proportions of
flanking/intronic/synonymous/nonsynonymous, second column of pie charts shows the
proportions of de novo/standing variation, and third column of squares shows the number of
candidate adaptive variants per gene (from Fig. 2). For genes with multiple adaptive variants, we
used the variant with the median position as the focal position. * Indicates genes with a
craniofacial annotation; ‡ indicates genes with neural or sensory annotation. (B) The proportion
of nonsynonymous adaptive variants is significantly larger in more recent selective sweeps,
indicated by the best-fit quasibinomial regression line; each pie chart represents a single
candidate gene. (C) The proportion of de novo variants is significantly larger in more recent
selective sweeps; each pie chart represents a single candidate gene.

311   *Increased visual acuity in the zooplanktivore is consistent with vision-related candidate adaptive*

312   *genes*

313   We identified several *C. simus* adaptive variants associated with eye development

(*ASIC3*(53), *CRIM1*(54), *MAB21L2*(55)) and opsin (*OPN1LW*, *SWS2*) genes (Table S1). Because

the trophic specialization of *C. simus* requires visually locating zooplankton in the open water

before targeted suction-feeding strikes, we hypothesized that this species would have higher

visual acuity than the detritivore, *C. beltrani*. We conducted optomotor response trials on lab-

reared individuals from both species (F1 *C. beltrani* and long-term laboratory colony of *C. simus*

(F10+)) and found that *C. simus* has a much stronger optomotor response than *C. beltrani* in

response to black-and-white vertical bars spinning at a constant rate of approximately 90

rotations per minute (Figure 5A, B; Wilcoxon signed-rank test: $W = 1$, $P = 0.009$). There was no

significant difference in rotations per minute between the species for the same individuals during

control observation periods with no spinning (Figure S7A; $W = 14$, $P = 0.3105$) or during positive

control observation periods with an all-black background spinning at the same frequency (Figure

S7B; $W = 9$, $P = 0.147$).

Figure 5. Divergence in visual acuity, sperm morphology, and tooth development reflect
candidate adaptive genes in the zooplanktivore. (A) Diagram of the optomotor response
experimental apparatus (adapted from Neuhauss et al., 1999). (B) Boxplot and jittered points of
the number of complete rotations fish swam per minute, indicative of their optomotor response to
black-and-white vertical bars spinning at a constant rate. (C) Representative labeled image of a
single *C. beltrani* sperm at 100x stained with rose bengal. (D) Boxplots with jittered points of
sperm head diameter for *C. artifrons, beltrani,* and *simus.* Significant Tukey's post hoc
differences are shown with an asterisk. (E) Manhattan plot showing per-site F_{st} of SNPs within
50 kb of WNT10A. Black rectangles indicate exons of WNT10A, and dashed lines indicate
introns. Red points represent nearly fixed SNPs ($F_{st} > 0.95$), and black points represent all other
SNPs. The black line is the smoothed average F_{st} in this region, with grey shading being the 95%
confidence intervals. (F) Representative image of *C. beltrani* larvae that were treated with
DMSO (control; top) or with iCRT14 Wnt inhibitor + DMSO (bottom) and stained with alizarin
red (bone) and alcian blue (cartilage). (G) Boxplot and jittered points of average tooth base width
of the left and right teeth closest to the mandibular symphysis for *C. beltrani* larvae treated with
either a control or Wnt inhibitor (iCRT14). Shapes represent different split-brood replicates.

Divergent sperm morphology of the zooplanktivore is consistent with spermatogenesis-related

candidate adaptive genes

Four candidate genes (*C4orf22*, *STK35*, *ERC2*, and *TBC1D25*) are known to affect

spermatogenesis/fertility (57–60). We therefore examined whether three Yucatan species, raised

in a common garden laboratory environment (F1 *C. artifrons,* F1 *beltrani,* and a long-term

laboratory colony of C. *simus* F10+), differed in sperm morphology using differential

interference contrast microscopy. We found that species significantly differed in sperm head

length (Figure 5D; $F = 5.724$, $p = 0.025$), but not sperm midpiece length (Figure S7A; $F = 2.403$,

$p = 0.147$) or flagellum length (Figure S8B; $F = 0.414$, $p = 0.673$). Tukey's post hoc pairwise

comparisons indicated that *C. simus* had significantly larger sperm heads than *C. beltrani* ($q =$

3.367; $p = 0.0207$; Fig. 5D).

Inhibition of WNT10A alters tooth morphology in Chichancanab pupfish

We identified a large number of highly differentiated SNPs surrounding the well-characterized *WNT10A*, which controls tooth, hair, and skin development within the Wnt signaling pathway (Figure 2; Figure 5E (61–63)). This is consistent with the most exceptional morphological difference between *C. simus* and *C. beltrani,* extreme divergence in tooth size (100 times faster than background rates (29)).

To test for the importance of *WNT10A* in tooth development within pupfishes, we used iCRT14 to inhibit the Wnt / β catenin pathway downstream of WNT10A during metamorphosis in *C. beltrani* lab-reared fry (64, 65). Chemical inhibition from 8 dpf to 23 dpf resulted in narrower teeth than control fry treated with DMSO ($\chi^2 = 4.840$, $P = 0.0278$; Figure 5F, G). There were no significant differences in mean tooth length ($\chi^2 = 0.455$, $p = 0.500$; Figure S9) or number of ossified teeth when stained with alizarin red ($\chi^2 = 0.114$, $p = 0.7357$; Figure S10).

Discussion

Here, we investigated the genetic architecture, sources of adaptive variation, and temporal dynamics of selection underlying speciation and trophic specialization in a young adaptive radiation of *Cyprinodon* pupfishes endemic to Laguna Chichancanab, Mexico. By comparing these patterns to our previous work on a recent radiation of *Cyprinodon* pupfishes in SSI, Bahamas (26, 28–32), we discovered parallel and non-parallel features of these independent radiations in similar environments. Despite replicate radiations of Caribbean pupfishes in large, isolated, predator-free, saline lakes with few competitors, we found little parallelism between the radiations. Although craniofacial morphology diverged rapidly within trophic specialists in both radiations, we found almost zero overlap in the adaptive genes underlying trophic specialization,

potentially reflecting the divergent specialists within each radiation. In contrast to SSI, there was

no behavior-first stage of adaptive divergence in Chichancanab specialists and, with the

exception of craniofacial development, candidate adaptive gene ontologies potentially relevant to

speciation were not shared, such as pigmentation genes in SSI versus opsin and spermatogenesis

genes in Chichancanab. Unlike the SSI radiation and most other well-studied adaptive radiations

(e.g., Malawi cichlids (20), Galápagos finches (66)), adaptive introgression likely played a

minimal role in the Chichancanab radiation. We found no overlap between adaptive variants in

the specialists and tracts of introgression. Instead, over 30% of the candidate adaptive variants in

the zooplanktivore, *C. simus*, likely arose from de novo mutations compared to only 2% in the

SSI scale-eater specialist and 0% in the molluscivore specialist. One outstanding parallel feature

was a clear stage of refinement in the adaptive divergence of the zooplanktivore, marked by

increased rates of de novo and nonsynonymous adaptive mutations occurring among the most

recent selective sweeps in this species, similar to the scale-eating specialist on SSI.

Adaptive introgression did not contribute to speciation of Chichancanab pupfishes

Genomic analyses have revealed that hybridization and adaptive introgression played a

major role in many adaptive radiations (3, 3, 16, 19–27, 67, 68), including the *Cyprinodon* SSI

radiation (26, 31). However, in Chichancanab, we found minimal evidence of introgression

overall and did not detect any signatures of adaptive introgression following the colonization of

Laguna Chichancanab. Although our analyses indicate extensive introgression among

Cyprinodontidae lineages, we did not find any introgression events into the root of the radiation,

nor has there been secondary gene flow from outgroup species other than the sister species to the

radiation, *C. artifrons*. Secondary gene flow from *C. artifrons* was almost entirely detected with

C. beltrani and, to a lesser extent, *C. labiosus*. Gene flow may be higher with these species

because *C. artifrons,* a coastal benthic feeding detritivore, is less ecologically divergent from

both *C. beltrani,* a benthic feeding detritivore, and *C. labiosus*, a benthic feeding bivalve

specialist. Secondary gene flow between *C. artifrons* and *C. beltrani* and *C. labiosus* overlapped

with many genes under selection in *C. simus*, suggesting that secondary gene flow is potentially

helping to increase similarities between the benthic detritivores within the lake and the coast

rather than by bringing in new adaptive variation for trophic specialization in *C. simus*.

These results are remarkably different from the *Cyprinodon* adaptive radiation of SSI,

where there is strong evidence of introgression into the root of the radiation and substantial

evidence of secondary adaptive introgression into both trophic specialists from at least three

different biogeographic regions of the Caribbean and Atlantic coast (26, 31). Laguna

Chichancanab is an endorheic basin with no above-ground hydrological connections to the ocean

and is substantially more geographically isolated than SSI, a small island in the approximate

center of the Caribbean. Furthermore, the sister species to the Chichancanab species flock, *C.*

artifrons, is abundant along the entire coast of the Yucatán peninsula, suggesting that any gene

flow into Chichancanab would likely first have to pass through *C. artifrons* populations. An

additional factor is the water chemistry of Laguna Chichancanab which is near-saturation with

calcite and gypsum and may be a greater physiological barrier to marine pupfish colonists than

the slightly hypersaline lakes of SSI. Thus, geography and ecology play an important role in

determining the relative importance of adaptive introgression in radiations.

De novo mutations primarily drive adaptive divergence of the zooplanktivore

The importance of de novo mutations in adaptive radiations remains contentious, in part

due to the time needed for de novo beneficial mutations to arise (11, 15). Interestingly, we find

that 30.6% of the zooplanktivore adaptive variants were detected only within Chichancanab

species, suggesting their de novo origins within the basin. This proportion is 15.3 times greater

than the contribution of de novo mutations to adaptive divergence of the scale-eater in the SSI

radiation. A primary factor could be that the estimated effective population size for the

Chichancanab radiation is approximately twice that of the SSI radiation (26), indicating that

selection on de novo mutations is more effective, with a higher probability of fixation, all else

being equal (14). There may also be a lack of existing standing genetic variation within

Caribbean pupfish populations that is adaptive for zooplanktivory. In support of this hypothesis,

although radiations should proceed along genetic lines of least resistance (69), the primary axis

of craniofacial morphological variation along which the zooplanktivore has diverged is

orthogonal to the major axes of variation that within Caribbean pupfish species (29). Evolving

along the line of greatest resistance instead of least resistance may require de novo mutations.

While the SSI scale-eater is also highly divergent, it is closer to the existing morphological space

of Caribbean pupfish species and in the direction of greatest variance along the first two

principal component axes (29). Similarly, mutational target size for the craniofacial features of a

zooplanktivore may be much smaller than the features of a scale-eater, resulting in less standing

genetic variation available for adaptation to this trophic niche. Alternatively, we may not have

detected these de novo mutations in other populations, but this is unlikely to explain all de novo

mutations given our extensive sampling of *Cyprinodon* species across the clade and spanning the

Caribbean (Fig. 1). Thus, our results highlight that de novo mutations may play a more important

role in driving young adaptive radiations than currently appreciated (12, 13), especially when

ecotypes are highly divergent from other species within the radiation.

Minimal parallelism in the genes underlying adaptive divergence of trophic specialists

We found only two candidate genes (*DNM1* and *SPTLC3*) among the 55 zooplanktivore

candidate genes in the Chichancanab radiation that overlapped with the hundreds of candidate

genes identified so far in the SSI radiation (26, 28, 30, 32, 42–46). Consistent with trophic

specialization requiring the pursuit of mobile prey, both genes are involved in motor neuron

development. *DNM1* is a dynamin with nervous tissue-specific expression that is crucial for the

proper formation of motor neuron axons (47). *SPTLC3* is involved in the biosynthesis of

sphingolipids (70), and suppression of *SPTLC3* in zebrafish results in motor neuron axon defects

(48).

The extremely limited number of shared adaptive genes is surprising given the high

conservation of gene function in both morphological development (71, 72) and behavior (73),

and the frequent parallelism of candidate genes across vertebrate adaptive radiations, particularly

for craniofacial genes. For example, *BMP4* is involved in diversification in both Galápagos

finches (74) and African cichlids (75, 76). Furthermore, within the SSI radiation, many of the

same genes show parallel patterns of differential expression even within divergent trophic

specialists (26, 43). Nonetheless, parallelism of shared regulatory networks, rather than specific

genes, is still a possibility.

Rapid divergence in tooth morphology of the zooplanktivore through WNT10a

Another reason for this non-parallelism is that trophic specialists in each radiation have

diverged in different sets of traits during adaptation to different ecological niches (29).

Specifically, the Chichancanab radiation has diversified in tooth size over 130 times faster than

background rates, driven by the extremely short and narrow teeth of the zooplanktivore (Figure

S1; (29). We identified a well-characterized tooth development gene in the Wnt pathway,

WNT10A, with many de novo mutations in flanking (n = 2) and intronic (n = 13) regions.

Mutations in *WNT10A* are associated with tooth developmental disorders in humans (62) and

both knockout and overexpression experiments in sticklebacks and zebrafish have demonstrated

WNT10A's role in teeth development and regeneration (61, 63). Consistent with *WNT10A*

playing a role in rapid divergence of tooth morphology, we found that inhibiting the Wnt

pathway in the *C. beltrani* generalist during pre-metamorphosis development resulted in

narrower teeth, similar to the reduced tooth sizes in the zooplanktivore *C. simus* (Fig. 4c). This

indicates that the substantial number of de novo mutations in WNT10a may be suppressing the

Wnt pathway, resulting in a large phenotypic effect on tooth size reduction in the zooplanktivore.

Increased visual acuity in the zooplantkivore is consistent with unique selection on vision-related

genes

One of the more striking features of both *Cyprinodon* radiations is the evolution of novel

trophic specialists from a generalist ancestor: the scale-eater (*C. desquamator*) in the SSI

radiation and the zooplanktivore (*C. simus*) in the Chichancanab radiation. Although these are

distinct dietary specializations, they share a similarity in that they require tracking free-

swimming prey. Indeed, four of the top 15 enriched GO terms of candidate genes in the SSI

scale-eater were related to eye development (26). Similarly, several of the candidate genes we

identified in the zooplanktivore were related to eye development, including *ASIC3*, *MAB21L2*,

and *CRIM1*. *CRIM1* is the most promising candidate because it is one of the few candidate genes

with multiple nonsynonymous mutations (Figure 2B). In zebrafish, a two-base pair deletion in

CRIM1 resulted in abnormal lenses as well as a large eye-to-head ratio (54). Interestingly, one of

the defining morphological features of *C. simus* is its large eye-to-head ratio (33) (Figure 1A).

Consistent with candidate genes related to eye development, we confirmed that *C. simus* exhibits

a much stronger optomotor response than *C. beltrani*, indicating that this specialist has greatly

improved visual acuity, even when reared in a common laboratory environment.

Although both radiations contained many vision-related candidate genes, selection on

opsin genes was unique to Chichancanab. Most notably, we found several adaptive variants

associated with *SWS2* (blue-sensitive), and *OPN1LW* (red-sensitive) in the zooplanktivore.

Interestingly, both opsins contain many candidate de novo mutations, and *SWS2* is the youngest

selective sweep in the zooplanktivore. One explanation could be that most Caribbean

Cyprinodon species are benthic and live in similar habitats, resulting in limited standing genetic

variation in opsins. Additionally, most environments (including SSI and Chichancanab) are

relatively shallow, so differences in depth—a common selective pressure on opsin evolution

(77)—are likely not a strong selective force on SSI. Thus, the main selective force acting on

vision in Chichancanab may be prey capture efficiency. In many fish species, zooplanktivory is

associated with higher expression of shorter wavelength opsins (e.g., *SWS1* (77)), whereas

piscivory is not often associated with changes in opsin expression (78, 79).

Divergent sperm morphology in Chichancanab species consistent with selection on

spermatogenesis genesis

Although the role that premating sexual selection plays in speciation has historically been

the main focus (80, 81), there is both theoretical and empirical support for the role of postmating

sexual selection in maintaining reproductive isolation via postmating prezygotic reproductive

barriers (82–87). In the SSI radiation, species differ in pigmentation, and a handful of the

candidate adaptive variants fall in regulatory regions of genes associated with pigmentation (26),

but not spermatogenesis. Furthermore, there is evidence for premating isolation by female

preference in the scale-eater (88). However, in Chichancanab, the two species tested in our data

set (*C. labiosus* and *C. beltrani*) were not found to visually distinguish between different species

in the radiation (89).

Unlike the SSI radiation, we did not find any candidate adaptive variants associated with

pigmentation in Chichanancab; instead, we identified adaptive variants associated with four

spermatogenesis genes, *ERC2*, *TBC1D25*, *C4orf22*, and *STK35*. Consistent with previous work

demonstrating the importance of these genes in spermatogenesis (57–60), we found that *C. simus*

does indeed have larger sperm heads than *C. beltrani*, suggesting a unique role for post-mating

prezygotic reproductive barriers in this radiation (Fig. 4b).

Parallelism in the stages of adaptation: refinement with de novo and nonsynonymous mutations

Adaptation may occur in stages, with less pleiotropic cis-regulatory changes occurring

before potentially more pleiotropic amino acid changes (26, 90, 91). Selection is also predicted

to act on standing variation first (14, 92). In both radiations, we observed strong evidence for a

refinement stage of coding substitutions occurred last. In Chichancanab, these late-sweeping

coding substitutions were also de novo mutations (Fig. 3). The simplest explanation for

parallelism in more recent ages of adaptive de novo mutations is the waiting time required for

beneficial mutations to (1) arise and (2) reach a substantial frequency within the population (14,

92). These variants may only be selected in the later stages of an adaptive walk because

nonsynonymous mutations are often pleiotropic and likely deleterious in the ancestral genetic

background (14, 93). Thus, they may not be advantageous until later in the adaptive walk, where

sign epistasis may reverse negative pleiotropic effects (94–98). Our results highlight that this

final refinement stage is likely a general pattern during adaptive divergence.

Methods

Sampling

We collected Laguna Chichancanab specimens in 2022 over two days using a 5 x 1.3 m

seine net with 1.6 mm mesh size. We sampled two sites, an entrance road by La Presumida, and a

bridge crossing the narrow section of the lake on the road to San Diego. *C. labiosus* and *C.*

beltrani were sampled from both sites for this study, whereas *C. simus* was only collected at La

Presumida. *C. artifrons* were collected by cast net from the coastal estuary at Sisal. Additional

specimens were collected from the Bahamas (Lake Cunningham, New Providence Island and

Crescent Pond, San Salvador Island), the Dominican Republic (Laguna Bavaro), or Fort Fisher,

North Carolina as described previously (Martin 2016) or sourced from the American Killifish

Association, the London Zoological Society, the Dallas Children's Aquarium, and Frans

Vermuelen. All specimens were euthanized in an overdose of buffered MS-222 (Fiquel, Inc.)

following approved animal care and use protocols from the University of California, Berkeley

and the University of California, Davis.

Sequencing, Variant calling and filtering

Individual DNA samples were resequenced using Illumina Hiseq 4000 and Novaseq. Raw

reads were trimmed with *fastp* (v0.23.4) (99). Reads were mapped to the UCB_CyNevMio_1.0

Cyprindon nevadensis mionectes reference genome (GCA_030533445.1) with *bwa mem*

(v0.7.17) (100). Duplicate reads in the bam files were marked with *Picard MarkDuplicates*

(GATK v4.5.0.0) (118). Coverage and mapping quality were assessed with *qualimap* (v2.2.2)

(101). We followed the *GATK* (v.4.5.0.0) genotyping pipeline to call variants (119). We used

HaplotypeCaller (-ERC GVCF) v4.5.0.0 to call variants for each individual and stored variants

in the GenomicsDB datastore format prior to using *GenotypeGVCFs* to perform joint

genotyping. We restricted our analyses to biallelic SNPs and applied the recommended *GATK*

hard filters (QD < 2, QUAL < 30, SOR > 3, FS > 60, MQ < 40, MQRankSum < -12.5,

ReadPosRankSum < -8) to filter SNPs (118, 120). Missing genotypes were reset to ./. from 0/0

due to how *GATK* represents missing genotypes in version 4.5.0.0 (105).

After hard-filtering and variant calling, we filtered the data set further by removing SNPs

with minor allele frequency < 0.05, more than 10% missing, depth less than 10x, and GQ less

than 20 using *BCFtools* (v1.16)(106). This resulted in a final data set that included 23,960,536

SNPs.

Population structure of Yucatán Cyprinodon pupfishes

To assess population structure, we first pruned SNPs in linkage disequilibrium using

PLINK (v. 1.9) with the following parameters: "-indep-pairwise 50 5 0.2" (107). This filtered our

data set from 23,960,536 SNPs to 5,476,855 SNPs. With this linkage disequilibrium pruned data

set, we first filtered the full VCF file to only include relevant Yucatán *Cyprinodon* species (i.e.,

C. artifrons, C. beltrani, C. labiosus, and *C. simus*), then conducted a PCA using *PLINK*. Next,

we used *ADMIXTURE* (v1.3) to determine the optimal number of population clusters that best fit

the data and to assign individuals to their corresponding population clusters (108). We performed

this analysis and calculated cross-validation error for the number of clusters K = 1-6. Models in

ADMIXTURE with K=2-4 were equally supported (Figure S2A). However, given morphological

distinctness and the results of the PCA, we present the results for when K = 4 (Figure 1D).

Demographic history of Yucatán Cyprinodon pupfish

We estimated demographic history using *MSMC2* (109). For each Yucatán pupfish

species, we selected three individuals with the highest mean depth and generated single-sample

VCF files, individual mask files, and "mapability" mask files. We then ran *MSMC2* with default

parameters except we set phasing to "unphased" and "P_PAR=8*1+25*1+1*2+1*3." For

plotting, we used a generation time of 1 year and a mutation rate of 1.56 x 10^{-8} substitutions per

base pair, estimated from Caribbean *Cyprinodon* species (26).

Identifying candidate genes

We calculated F_{st} genome-wide, per site, and in non-overlapping 10-kb windows using

the "--weir-fist-pop" function in *VCFtools*(v0.1.16) for all pairwise combinations of Yucatán

Cyprinodon species (110). We chose 10-kb windows because linkage disequilibrium was

substantially decayed at this distance (Figure S3), and these windows allowed us to quantify

finer-scale genomic variation. With the 10-kb windowed F_{st} values, we calculated a modified

population branch statistic (PBS) for four taxa, PBS_{nj} (111). This statistic is useful because it

makes no assumptions about species topology or polarization (111). We then normalized PBS_{nj}

by the total tree length, as normalized *PBS* is most effective and specific at identifying selective

sweeps (41). We classified SNPs as candidates if they were fixed or nearly fixed (per-site F_{st}

>0.95) compared to other species within the radiation and were among the top 1% outliers out of

all PBS_{nj} windows, following the threshold used in (41).

Introgression

Using the full VCF file, which contained a wide range of species in the family

Cyprinodontidae, we tested for evidence of introgression in Yucatán *Cyprinodon* species using

Dsuite (49). We first excluded any samples/species that had an average depth < 4. We then

calculated all possible trios with a species tree based on the most recent phylogeny (112) with

Cualac tessellatus as the outgroup. We calculated Z-scores to assess the significance of *D*

statistics using 1,000 jackknife blocks and considered a Z-score greater than three as evidence

for significant introgression (113). We only focused our primary analysis on species trios where

P1-P2 included species within the Laguna Chichancanab radiation. We used the other trio

calculations to calculate *f*-branch (f_b) statistics to account for correlated f_4-ratio scores and to

inform the timing of introgression events (49).

For species trios that showed evidence of introgression where *C. simus* was either P1 or

P2, we then calculated the distance fraction (d_f), a robust metric of introgression for windowed

analyses (50), in non-overlapping windows of 91 informative SNPs. We chose this number of

SNPs because it resulted in windows of approximately 10 kb. We considered windows to be

introgressed if they were in the top X% of absolute d_f value, where X% was the percentage of

estimated genome wide introgression (e.g, f_4-ratio) (114, 115). To estimate introgression block

sizes, we merged neighboring windows that were both introgression outliers and took the total

size of the blocks.

Characterizing candidate genes

After identifying candidate SNPs, we then used *Bedtools* (v2.31) to see if SNPs occurred

near (20-kb) or within a gene (116) with a GFF annotation file for the *C. nev. mionectes*

reference genome (Tian et al. in prep). We classified SNPs within 20 kb of a gene (upstream or

downstream) as flanking. For SNPs within a gene, we further classified them into intronic,

synonymous, or nonsynonymous using *SnpEff* (117).

To characterize the origin of candidate SNPs, we extracted the alleles from every species

for each candidate SNP. We classified variants that were found in any other species as "standing

variation". We further subdivided this group into variants that were found in other species but not

in *C. artifrons*. If "standing variation" variants fell within an introgression window, we classified

them as "introgressed." Finally, if the alternative variant was only found in Chichancanab

species, we classified it as de novo.

Timing of selective sweeps

To gain insight into the timing of selective sweeps in *C. simus*, we used *starTRMCA* (51).

Because many genes contained multiple candidate SNPs, for this analysis, we selected the

variant that was in the median position of each gene. We then extracted a 1-Mb region

surrounding that variant and removed all sites that contained missing data. For simplification, we

only used the generalist/detritivore, *C. beltrani*, as the "reference" population and the

zooplanktivore, *C. simus*, as the "selected" population. We used a fixed recombination rate of

2×10^{-8} (swordtail fishes; (118)) and a mutation rate of 1.56×10^{-8} substitutions per base pair

(Caribbean *Cyprinodon* species; 26). We then ran five separate Markov chains for 30,000

iterations with a proposal standard deviation of 150 (preliminary analyses changing this

parameter had minimal influence on results). We then discarded the first 10,000 iterations of each

chain as burn-in.

To test for stages of adaptation, we conducted separate generalized linear regressions to

test if the sweep age was correlated with the proportion of nonsynonymous SNPs, proportion of

de novo SNPs, and the number of SNPs. For the generalized linear regressions of proportion of

nonsynonymous SNPs and proportion of de novo SNPs, we used a quasi-binomial family to

account for overdispersion with a logit link function. For the number of SNPs, we used a quasi-

Poisson family with a log link function. To test for the significance of sweep age, we used a

likelihood ratio test comparing a model with the estimated sweep age as an effect compared to a

null model.

Optomotor response

We tested for differences in visual acuity between the generalist/detritivore, *C. beltrani*,

and the zooplanktivore, *C. simus*, by conducting an optomotor response behavioral assay. We

tested lab-reared *C. beltrani* (F1, $n = 7$) and lab-reared *C. simus* (long-term laboratory colony

exceeding ten generations in the lab, $n = 5$). Each fish was tested consecutively with each control

or treatment for 1-minute observation periods, with the control observation occurring first,

followed by presentation of all-black and black-and-white spinning bars. The order of

presentation for the latter to treatments was alternated in each trial. Fish were placed in a

suspended cylindrical plastic container 0.3 m in diameter that remained stationary during the

course of the trials. Spinning black-and-white or all-black paper was rotated outside the clear

plastic container at an RPM of ~90rpm during each observation period.

Due to the data being nonparametric (count, zero-inflated, unequal variance between

groups), we conducted a Wilcoxon rank sum test for the on-banded (all black) and on-banded

portions separately. Because we had an a priori hypothesis that *C. simus* would display a stronger

optomotor response (greater visual acuity), we calculated *p*-values with a one-tailed test.

Sperm morphology

To test for differences in sperm morphology, we collected sperm samples from five

individuals per species in the lab (*C. simus, C. artifrons, C. beltrani*). We first anesthetized the

fish with a solution of MS-222. After the fish was anesthetized, we collected 1μL of milt (fish

semen) with a 1μL glass microcapillary tube and fixed the sample overnight in a 4% PFA

solution stained with Rose Bengal. We then plated sperm on a positively charged slide with a

sealed coverslip. We imaged sperm with oil immersion with a 63x objective and measured 10-30

sperm per individual in ImageJ (119). Due to poor sperm samples from some individuals (little

sperm), we ended up with the following sample sizes: three *simus*, four *artifrons*, and five

beltrani.

To retain information about variation within individuals, we fit separate linear mixed-

effects models for each trait of interest (sperm head, midpiece, and flagellum lengths) with

population as a fixed effect and a random intercept for individual ID, using *LME4* (120). To

assess the significance of the model, we used a Type II ANOVA with a Wald *F* test using

Kenward-Roger degrees of freedom. If there was a significant effect of population, we conducted

post hoc pairwise tests using the *Emmeans* package (121) and corrected for multiple comparisons

using the Tukey method. We did not correct for phylogeny due to how closely related the species

are, the limited sample size, and the fact that the only significant differences were between sister

species.

Wnt inhibition experiment

703   To confirm the role of the Wnt pathway in proper jaw and teeth development, we

conducted an experiment chemically inhibiting the Wnt pathway using iCRT14 during

metamorphosis in *C. beltrani*. To control the exact time of development, we first performed in

vitro fertilizations and incubated developing embryos at 24 °C until hatching (~8dpf). On the day

of hatching, we haphazardly split broods into control and experimental groups to control for

batch and family effects. We raised experimental groups in tank water with a 100 nanomolar

concentration of iCRT14 and control groups in a 100 nanomolar concentration of DMSO

(solution used to dilute stock iCRT14). We ran the experiment for two weeks until individuals

were (22dpf), changing the solution every 3-4 days. We performed this experiment on four

separate broods (~12 individuals/treatment). One brood was excluded as there was improper

development (no ossification of the spine) in the controls.

714   After the experiment, we euthanized fish with MS-222 and stained fish with alizarin red

(bone) and alcian blue (cartilage) following a modified published protocol (122). After staining,

we prepared whole mounts and imaged fish with fluorescent microscopy (Zeiss LSM880 FCS)

to quantify ossification of the jaw and teeth (123). We then dissected out the lower jaw and

prepared flat mounts to measure teeth morphology. Specifically, we measured the tooth length

and base width for the left and right teeth closest to the mandibular symphysis. For statistical

analyses, we averaged these values.

We fit separate linear mixed-effects models for tooth length and base width with

experimental treatment as a fixed effect and a random intercept for brood/replicate, using *LME4*

(120). For the number of ossified teeth, we used a generalized linear mixed-effect model with a

Poisson family and log-link function. To assess the significance of these models, we used a

likelihood ratio test comparing it to a null model without treatment as a fixed effect.

Acknowledgments

MCK was supported by a Miller Postdoctoral Research Fellowship from the Miller

Institute for Basic Research in Science, University of California Berkeley. This research was

funded by NSF CAREER 1749764, NIH 5R01DE027052-02, Daphne and Ted Pengelley Award

from the Center for Population Biology, and ARCS Foundation awards to CHM. We thank the

Gerace Research Center and Troy Day for logistical support in the Bahamas, and the governments

of Mexico and the Bahamas BEST commission for permission to collect and export samples

(Permit No. PPF/DGOPA-001/20). All research procedures and animal care protocols (AUP-2021-

02-14062-1 and AUP-2021-07-14515) were approved by the University of California, Berkeley

Animal Care and Use committee and the UC Davis Institutional Animal Care and Use committee.

References

1.   R. G. Gillespie, *et al.*, Comparing Adaptive Radiations Across Space, Time, and Taxa. *J*
Hered **111**, 1–20 (2020).

2.   C. H. Martin, E. J. Richards, The Paradox Behind the Pattern of Rapid Adaptive
Radiation: How Can the Speciation Process Sustain Itself Through an Early Burst? *Annual*
Review of Ecology, Evolution, and Systematics **50**, 569–593 (2019).

3. Y. Naciri, H. P. Linder, The genetics of evolutionary radiations. *Biological Reviews* **95**,
1055–1072 (2020).

4. D. Schluter, *The Ecology of Adaptive Radiation* (OUP Oxford, 2000).

5. J. T. Stroud, J. B. Losos, Ecological Opportunity and Adaptive Radiation. *Annual Review*
of Ecology, Evolution, and Systematics **47**, 507–532 (2016).

6. R. Butlin, *et al.*, What do we need to know about speciation? *Trends in Ecology and*
Evolution **27**, 27–39 (2012).

7. J. A. Coyne, H. A. Orr, *Speciation* (Oxford University Press, 2004).

8. D. R. Matute, B. S. Cooper, Comparative studies on speciation: 30 years since Coyne and
Orr. *Evolution* **75**, 764–778 (2021).

9. D. Schluter, L. H. Rieseberg, Three problems in the genetics of speciation by selection.
Proceedings of the National Academy of Sciences **119**, e2122153119 (2022).

10. D. Berner, W. Salzburger, The genomics of organismal diversification illuminated by
adaptive radiations. *Trends in Genetics* **31**, 491–499 (2015).

11. D. A. Marques, J. I. Meier, O. Seehausen, A Combinatorial View on Speciation and
Adaptive Radiation. *Trends in Ecology & Evolution* **34**, 531–544 (2019).

12. J. B. Pease, D. C. Haak, M. W. Hahn, L. C. Moyle, Phylogenomics Reveals Three Sources
of Adaptive Variation during a Rapid Radiation. *PLOS Biology* **14**, e1002379 (2016).

13. B. W. Stone, C. A. Wessinger, Ecological Diversification in an Adaptive Radiation of
Plants: The Role of De Novo Mutation and Introgression. *Mol Biol Evol* **41**, msae007
(2024).

14. R. D. H. Barrett, D. Schluter, Adaptation from standing genetic variation. *Trends in*
Ecology & Evolution **23**, 38–44 (2008).

15. M. Rousselle, *et al.*, Is adaptation limited by mutation? A timescale-dependent effect of
genetic diversity on the adaptive substitution rate in animals. *PLOS Genetics* **16**, e1008668
(2020).

16. O. Seehausen, Hybridization and adaptive radiation. *Trends in Ecology & Evolution* **19**,
198–207 (2004).

17. K. Kagawa, G. Takimoto, O. Seehausen, Transgressive segregation in mating traits drives
hybrid speciation. *Evolution* **77**, 1622–1633 (2023).

18. K. Kagawa, G. Takimoto, Hybridization can promote adaptive radiation by means of
transgressive segregation. *Ecology Letters* **21**, 264–274 (2018).

19.   R. De-Kayne, *et al.*, Genomic architecture of adaptive radiation and hybridization in
778   Alpine whitefish. *Nat Commun* **13**, 4479 (2022).

779   20.   D. A. Joyce, *et al.*, Repeated colonization and hybridization in Lake Malawi cichlids.
780   *Current Biology* **21**, R108–R109 (2011).

781   21.   S. Lamichhaney, *et al.*, Evolution of Darwin's finches and their beaks revealed by genome
782   sequencing. *Nature* **518**, 371–375 (2015).

783   22.   C. H. Martin, *et al.*, Complex histories of repeated gene flow in Cameroon crater lake
784   cichlids cast doubt on one of the clearest examples of sympatric speciation. *Evol* **69**,
785   1406–1422 (2015).

786   23.   J. I. Meier, *et al.*, Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat*
787   *Commun* **8**, 14363 (2017).

788   24.   C. Pouchon, *et al.*, Phylogenomic Analysis of the Explosive Adaptive Radiation of the
789   Espeletia Complex (Asteraceae) in the Tropical Andes. *Syst Biol* **67**, 1041–1060 (2018).

790   25.   E. J. Richards, J. W. Poelstra, C. H. Martin, Don't throw out the sympatric speciation with
791   the crater lake water: fine-scale investigation of introgression provides equivocal support
792   for causal role of secondary gene flow in one of the clearest examples of sympatric
793   speciation. *Evolution Letters* **2**, 524–540 (2018).

794   26.   E. J. Richards, *et al.*, A vertebrate adaptive radiation is assembled from an ancient and
795   disjunct spatiotemporal landscape. *Proceedings of the National Academy of Sciences* **118**,
796   e2011811118 (2021).

797   27.   N. Rosser, *et al.*, Hybrid speciation driven by multilocus introgression of ecological traits.
798   *Nature* **628**, 811–817 (2024).

799   28.   M. E. S. John, J. C. Dunker, E. J. Richards, S. Romero, C. H. Martin, Parallel genetic
800   changes underlie integrated craniofacial traits in an adaptive radiation of trophic specialist
801   pupfishes. [Preprint] (2021). Available at:
802   https://www.biorxiv.org/content/10.1101/2021.07.01.450661v2 [Accessed 21 November
803   2023].

804   29.   C. H. Martin, P. C. Wainwright, Trophic novelty is linked to exceptional rates of
805   morphological diversification in two adaptive radiations of cyprinodon pupfish. *Evolution*
806   **65**, 2197–2212 (2011).

807   30.   A. H. Patton, E. J. Richards, K. J. Gould, L. K. Buie, C. H. Martin, Hybridization alters
808   the shape of the genotypic fitness landscape, increasing access to novel fitness peaks
809   during adaptive radiation. *eLife* **11**, e72905 (2022).

810   31.   E. J. Richards, C. H. Martin, Adaptive introgression from distant Caribbean islands
811   contributed to the diversification of a microendemic adaptive radiation of trophic
812   specialist pupfishes. *PLOS Genetics* **13**, e1006919 (2017).

32.   E. J. Richards, C. H. Martin, We get by with a little help from our friends: shared adaptive
variation provides a bridge to novel ecological specialists during adaptive radiation. *Proc*
Biol Sci **289**, 20220613 (2022).

33.   J. M. Humphries, R. R. Miller, A Remarkable Species Flock of Pupfishes, Genus
Cyprinodon, from Yucatán, México. *Copeia* **1981**, 52–64 (1981).

34.   D. A. Hodell, J. H. Curtis, M. Brenner, Possible role of climate in the collapse of Classic
Maya civilization. *Nature* **375**, 391–394 (1995).

35.   U. Strecker, Description of a new species from Laguna Chichancanab, Yucatan, Mexico:
Cyprinodon suavium (Pisces: Cyprinodontidae). *Hydrobiologia* **541**, 107–115 (2005).

36.   J. M. Humphries, Cyprinodon verecundus, n. sp., a Fifth Species of Pupfish from Laguna
Chichancanab. *Copeia* **1984**, 58–68 (1984).

37.   C. A. Gracida-Juárez, J. J. Schmitter-Soto, M. J. Genner, Community structure of
indigenous fishes relative to habitat variation and invasive tilapia in lakes of Quintana
Roo, Mexico. *Environmental Biology of Fishes* **107**, 401–414 (2024).

38.   U. Strecker, The impact of invasive fish on an endemic Cyprinodon species flock
(Teleostei) from Laguna Chichancanab, Yucatan, Mexico. *Ecology of Freshwater Fish* **15**,
408–418 (2006).

39.   U. Strecker, Genetic differentiation and reproductive isolation in a *Cyprinodon* fish species
flock from Laguna Chichancanab, Mexico. *Molecular Phylogenetics and Evolution* **39**,
865–872 (2006).

40.   U. Strecker, C. G. Meyer, C. Sturmbauer, H. Wilkens, Genetic Divergence and Speciation
in an Extremely Young Species Flock in Mexico Formed by the
Genus*Cyprinodon*(Cyprinodontidae, Teleostei). *Molecular Phylogenetics and Evolution* **6**,
143–149 (1996).

41.   M. Shpak, K. N. Lawrence, J. E. Pool, The Precision and Power of Population Branch
Statistics in Identifying the Genomic Signatures of Local Adaptation. *Genome Biol Evol*
17, evaf080 (2025).

42.   J. A. McGirr, C. H. Martin, Novel Candidate Genes Underlying Extreme Trophic
Specialization in Caribbean Pupfishes. *Molecular Biology and Evolution* **34**, 873 (2017).

43.   J. A. McGirr, C. H. Martin, Parallel evolution of gene expression between trophic
specialists despite divergent genotypes and morphologies. *Evol Lett* **2**, 62–75 (2018).

44.   J. A. McGirr, C. H. Martin, Ecological divergence in sympatry causes gene misexpression
in hybrids. *Molecular Ecology* **29**, 2707–2721 (2020).

45. J. A. McGirr, C. H. Martin, Few Fixed Variants between Trophic Specialist Pupfish
Species Reveal Candidate Cis-Regulatory Alleles Underlying Rapid Craniofacial
Divergence. *Mol Biol Evol* **38**, 405–423 (2021).

46. M. F. Palominos, V. Muhl, C. H. Martin, Craniofacial-specific transcriptomics uncovers
novel genes underlying jaw divergence in dietary specialist pupfishes. *Genetics* iyaf207
(2025). https://doi.org/10.1093/genetics/iyaf207.

47. C. Bragato, *et al.*, Zebrafish dnm1a gene plays a role in the formation of axons and
synapses in the nervous tissue. *Journal of Neuroscience Research* **101**, 1345–1359 (2023).

48. C. Gonzaga-Jauregui, *et al.*, Exome Sequence Analysis Suggests that Genetic Burden
Contributes to Phenotypic Variability and Complex Neuropathy. *Cell Reports* **12**, 1169–
1183 (2015).

49. M. Malinsky, M. Matschiner, H. Svardal, Dsuite - Fast D-statistics and related admixture
evidence from VCF files. *Molecular Ecology Resources* **21**, 584–595 (2021).

50. B. Pfeifer, D. D. Kapan, Estimates of introgression as a function of pairwise distances.
BMC Bioinformatics **20**, 207 (2019).

51. J. Smith, G. Coop, M. Stephens, J. Novembre, Estimating Time to the Common Ancestor
for a Beneficial Allele. *Mol Biol Evol* **35**, 1003–1017 (2018).

52. J. T. Streelman, P. D. Danley, The stages of vertebrate evolutionary radiation. *Trends in
Ecology & Evolution* **18**, 126–131 (2003).

53. W.-G. Li, T.-L. Xu, ASIC3 Channels in Multimodal Sensory Perception. *ACS Chem.
Neurosci.* **2**, 26–37 (2011).

54. T. Le, *et al.*, A zebrafish model of crim1 loss of function has small and misshapen lenses
with dysregulated clic4 and fgf1b expression. *Front. Cell Dev. Biol.* **13** (2025).

55. B. N. Kennedy, *et al.*, Zebrafish *rx3* and *mab21l2* are required during eye morphogenesis.
Developmental Biology **270**, 336–349 (2004).

56. S. C. F. Neuhauss, *et al.*, Genetic Disorders of Vision Revealed by a Behavioral Screen of
400 Essential Loci in Zebrafish. *J. Neurosci.* **19**, 8603–8615 (1999).

57. M. H. K. Chau, *et al.*, Investigation of the genetic etiology in male infertility with
apparently balanced chromosomal structural rearrangements by genome sequencing. *Asian
Journal of Andrology* **24** (2022).

58. H. Li, Y. Dai, Z. Luo, D. Nie, Cloning of a new testis-enriched gene C4orf22 and its role
in cell cycle and apoptosis in mouse spermatogenic cells. *Mol Biol Rep* **46**, 2029–2038
(2019).

59. Y. Miyamoto, *et al.*, The STK35 locus contributes to normal gametogenesis and encodes a
lncRNA responsive to oxidative stress. *Biol Open* **7**, bio032631 (2018).

60. S. Nawaz, S. Hussain, S. Basit, W. Ahmad, First evidence of involvement of *TBC1D25* in
causing human male infertility. *European Journal of Medical Genetics* **64**, 104142 (2021).

61. T. A. Square, *et al.*, Modulation of tooth regeneration through opposing responses to Wnt
and BMP signals in teleosts. *Development* **150**, dev202168 (2023).

62. S. Nawaz, *et al.*, WNT10A missense mutation associated with a complete Odonto-
Onycho-Dermal Dysplasia syndrome. *Eur J Hum Genet* **17**, 1600–1605 (2009).

63. E. L. Benard, *et al.*, wnt10a is required for zebrafish median fin fold maintenance and
adult unpaired fin metamorphosis. *Developmental Dynamics* **253**, 566–592 (2024).

64. J. L. Catrow, Y. Zhang, M. Zhang, H. Ji, Discovery of Selective Small-Molecule Inhibitors
for the β-Catenin/T-Cell Factor Protein–Protein Interaction through the Optimization of
the Acyl Hydrazone Moiety. *J. Med. Chem.* **58**, 4678–4692 (2015).

65. F. C. Gonsalves, *et al.*, An RNAi-based chemical genetic screen identifies three small-
molecule inhibitors of the Wnt/wingless signaling pathway. *Proceedings of the National
Academy of Sciences* **108**, 5954–5963 (2011).

66. E. D. Enbody, *et al.*, Community-wide genome sequencing reveals 30 years of Darwin's
finch evolution. *Science* **381**, eadf6218 (2023).

67. J. V. Peñalba, *et al.*, The Role of Hybridization in Species Formation and Persistence. *Cold
Spring Harb Perspect Biol* **16**, a041445 (2024).

68. O. Seehausen, Conditions when hybridization might predispose populations for adaptive
radiation. *Journal of Evolutionary Biology* **26**, 279–281 (2013).

69. D. Schluter, ADAPTIVE RADIATION ALONG GENETIC LINES OF LEAST
RESISTANCE. *Evol* **50**, 1766–1774 (1996).

70. T. Hornemann, *et al.*, The SPTLC3 Subunit of Serine Palmitoyltransferase Generates
Short Chain Sphingoid Bases *. *Journal of Biological Chemistry* **284**, 26322–26330
(2009).

71. M. E. Bronner, N. M. LeDouarin, Development and evolution of the neural crest: An
overview. *Developmental Biology* **366**, 2–9 (2012).

72. F. Hirth, H. Reichert, Conserved genetic programs in insect and mammalian brain
development. *BioEssays* **21**, 677–684 (1999).

73. C. J. Reaume, M. B. Sokolowski, Conservation of gene function in behaviour.
Philosophical Transactions of the Royal Society B: Biological Sciences **366**, 2100–2110
(2011).

74.   A. Abzhanov, M. Protas, B. R. Grant, P. R. Grant, C. J. Tabin, Bmp4 and Morphological
914   Variation of Beaks in Darwin's Finches. *Science* **305**, 1462–1465 (2004).

915   75.   R. C. Albertson, J. T. Streelman, T. D. Kocher, P. C. Yelick, Integration and evolution of
916   the cichlid mandible: The molecular basis of alternate feeding strategies. *Proceedings of*
917   *the National Academy of Sciences* **102**, 16287–16292 (2005).

918   76.   Y. Terai, N. Morikawa, N. Okada, The Evolution of the Pro-Domain of Bone
919   Morphogenetic Protein 4 (Bmp4) in an Explosively Speciated Lineage of East African
920   Cichlid Fishes. *Mol Biol Evol* **19**, 1628–1632 (2002).

921   77.   K. L. Carleton, D. Escobar-Camacho, S. M. Stieb, F. Cortesi, N. J. Marshall, Seeing the
922   rainbow: mechanisms underlying spectral sensitivity in teleost fishes. *J Exp Biol* **223**,
923   jeb193334 (2020).

924   78.   C. M. Hofmann, *et al.*, The Eyes Have It: Regulatory and Structural Changes Both
925   Underlie Cichlid Visual Pigment Diversity. *PLOS Biology* **7**, e1000266 (2009).

926   79.   K. E. O'Quin, C. M. Hofmann, H. A. Hofmann, K. L. Carleton, Parallel Evolution of
927   Opsin Gene Expression in African Cichlid Fishes. *Mol Biol Evol* **27**, 2839–2854 (2010).

928   80.   M. G. Ritchie, Sexual Selection and Speciation. *Annual Review of Ecology, Evolution, and*
929   *Systematics* **38**, 79–102 (2007).

930   81.   M. R. Servedio, J. W. Boughman, The Role of Sexual Selection in Local Adaptation and
931   Speciation. *Annual Review of Ecology, Evolution, and Systematics* **48**, 85–109 (2017).

932   82.   A. Devigili, *et al.*, Possible glimpses into early speciation: the effect of ovarian fluid on
933   sperm velocity accords with post-copulatory isolation between two guppy populations.
934   *Journal of Evolutionary Biology* **31**, 66–74 (2018).

935   83.   M. D. Garlovsky, *et al.*, Synthesis and Scope of the Role of Postmating Prezygotic
936   Isolation in Speciation. *Cold Spring Harb Perspect Biol* a041429 (2023).
937   https://doi.org/10.1101/cshperspect.a041429.

938   84.   M. C. Kustra, M. R. Servedio, S. H. Alonzo, Cryptic female choice can maintain
939   reproductive isolation. *Evol* **79**, 2259–2273 (2025).

940   85.   M. C. Kustra, S. H. Alonzo, The coevolutionary dynamics of cryptic female choice.
941   *Evolution Letters* **7**, 191–202 (2023).

942   86.   P. D. Lorch, M. R. Servedio, The evolution of conspecific gamete precedence and its effect
943   on reinforcement. *Journal of Evolutionary Biology* **20**, 937–949 (2007).

944   87.   J. L. Marshall, M. L. Arnold, D. J. Howard, Reinforcement: the road not taken. *Trends in*
945   *Ecology & Evolution* **17**, 558–563 (2002).

88. A. Kodric-Brown, R. J. D. West, Asymmetries in premating isolating mechanisms in a
sympatric species flock of pupfish (Cyprinodon). *Behavioral Ecology* **25**, 69–75 (2014).

89. U. Strecker, A. Kodric-Brown, Mate recognition systems in a species flock of Mexican
pupfish. *Journal of Evolutionary Biology* **12**, 927–935 (1999).

90. L. Campagna, *et al.*, Repeated divergent selection on pigmentation genes in a rapid finch
radiation. *Science Advances* **3**, e1602404 (2017).

91. F. C. Jones, *et al.*, The genomic basis of adaptive evolution in threespine sticklebacks.
Nature **484**, 55–61 (2012).

92. Z. D. Blount, R. E. Lenski, J. B. Losos, Contingency and determinism in evolution:
Replaying life's tape. *Science* **362**, eaam5979 (2018).

93. P. W. Hedrick, Adaptive introgression in animals: examples and comparison to new
mutation and standing variation as sources of adaptive variation. *Molecular Ecology* **22**,
4606–4618 (2013).

94. S. G. Das, M. Mungan, J. Krug, Epistasis-mediated compensatory evolution in a fitness
landscape with adaptational tradeoffs. *Proceedings of the National Academy of Sciences*
122, e2422520122 (2025).

95. M. J. Harms, J. W. Thornton, Historical contingency and its biophysical basis in
glucocorticoid receptor evolution. *Nature* **512**, 203–207 (2014).

96. M. Karageorgi, *et al.*, Genome editing retraces the evolution of toxin resistance in the
monarch butterfly. *Nature* **574**, 409–412 (2019).

97. R. D. Tarvin, *et al.*, Interacting amino acid replacements allow poison frogs to evolve
epibatidine resistance. *Science* **357**, 1261–1266 (2017).

98. D. M. Weinreich, N. F. Delaney, M. A. DePristo, D. L. Hartl, Darwinian Evolution Can
Follow Only Very Few Mutational Paths to Fitter Proteins. *Science* **312**, 111–114 (2006).

99. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

100. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
[Preprint] (2013). Available at: http://arxiv.org/abs/1303.3997 [Accessed 22 September
2025].

101. K. Okonechnikov, A. Conesa, F. García-Alcalde, Qualimap 2: advanced multi-sample
quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292–294 (2016).

102. A. McKenna, *et al.*, The Genome Analysis Toolkit: A MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

103. M. A. DePristo, *et al.*, A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat Genet* **43**, 491–498 (2011).

104. G. A. Van der Auwera, *et al.*, From FastQ Data to High-Confidence Variant Calls: The
Genome Analysis Toolkit Best Practices Pipeline. *Current Protocols in Bioinformatics* **43**,
11.10.1-11.10.33 (2013).

105. D. Caetano-Anolles, GenotypeGVCFs and the death of the dot (obsolete as of GATK
4.6.0.0). *GATK* (2024). Available at: https://gatk.broadinstitute.org/hc/en-
us/articles/6012243429531-GenotypeGVCFs-and-the-death-of-the-dot-obsolete-as-of-
GATK-4-6-0-0 [Accessed 27 July 2025].

106. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008
(2021).

107. S. Purcell, *et al.*, PLINK: A Tool Set for Whole-Genome Association and Population-
Based Linkage Analyses. *The American Journal of Human Genetics* **81**, 559–575 (2007).

108. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

109. S. Schiffels, K. Wang, "MSMC and MSMC2: The Multiple Sequentially Markovian
Coalescent" in *Statistical Population Genomics*, J. Y. Dutheil, Ed. (Springer US, 2020),
pp. 147–166.

110. P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

111. J. M. Schmidt, M. de Manuel, T. Marques-Bonet, S. Castellano, A. M. Andrés, The impact
of genetic adaptation on chimpanzee subspecies differentiation. *PLOS Genetics* **15**,
e1008485 (2019).

112. S. G. Hernández-Ávila, C. W. Hoagstrom, W. A. Matamoros, Historical biogeography of
North American killifishes (Cyprinodontiformes) recapitulates geographical history in the
Gulf of México watershed. *Zool J Linn Soc* **202**, zlae105 (2024).

113. E. Y. Durand, N. Patterson, D. Reich, M. Slatkin, Testing for Ancient Admixture between
Closely Related Populations. *Mol Biol Evol* **28**, 2239–2252 (2011).

114. C. Feng, J. Wang, A. Liston, M. Kang, Recombination Variation Shapes Phylogeny and
Introgression in Wild Diploid Strawberries. *Mol Biol Evol* **40**, msad049 (2023).

115. A. Morales-Cruz, *et al.*, Introgression among North American wild grapes (Vitis) fuels
biotic and abiotic adaptation. *Genome Biology* **22**, 254 (2021).

116. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).

117.   P. Cingolani, *et al.*, A program for annotating and predicting the effects of single
1014   nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster
1015   strain w1118; iso-2; iso-3. *Fly* **6**, 80–92 (2012).

1016   118.   M. Schumer, *et al.*, Natural selection interacts with recombination to shape the evolution
1017   of hybrid genomes. *Science* **360**, 656–660 (2018).

1018   119.   C. T. Rueden, *et al.*, ImageJ2: ImageJ for the next generation of scientific image data.
1019   *BMC Bioinformatics* **18**, 529 (2017).

1020   120.   D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using
1021   lme4. *Journal of Statistical Software* **67**, 1–48 (2015).

1022   121.   R. V. Lenth, *emmeans: Estimated Marginal Means, aka Least-Squares Means* (2021).

1023   122.   M. Walker, C. Kimmel, A two-color acid-free cartilage and bone stain for zebrafish larvae.
1024   *Biotechnic & Histochemistry* **82**, 23–28 (2007).

1025   123.   R. Koita, S. Oikawa, T. Tani, M. Matsuda, A. Kawamura, Live Visualization of Calcified
1026   Bones in Zebrafish and Medaka Larvae and Juveniles Using Calcein and Alizarin Red S.
1027   *Bio-protocol* **14** (2024).

1028   124.   R. Maier, *et al.*, On the limits of fitting complex models of population history to f-
1029   statistics. *eLife* **12**, e85492 (2023).

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043 **Supplementary materials**

1044

1045 **Table S1.** Mean genome-wide pairwise F_{ST} statistics within the Chichancanab radiation and with
1046 sister outgroup (*artifrons*).

	labiosus	*beltrani*	*artifrons*
simus	0.090	0.068	0.182
labiosus		0.032	0.184
beltrani			0.168

1047

1048   **Table S2.** F_{ST} statistics within the Chichancanab radiation.

Comparison	$F_{ST} = 1$ (fixed)	$1 > F_{ST} \geq 0.95$ (nearly fixed)	$0.95 > F_{ST} \geq 0.9$	$0.9 > F_{ST} \geq 0.8$	Other	Mean F_{ST}
simus vs *beltrani* + *labiosus*	19	1127	3304	20173	10359605	0.07
labiosus vs *beltrani* + *simus*	0	0	3	94	10384131	0.04
beltrani vs *simus* + *labiosus*	0	0	0	13	10384215	0.02

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062 **Table S3.** Fixed SNPs between *C. simus* and *C. beltrani* + *C. labiosus*. Bold rows indicate SNPs within top 1% of PBS windows.

1063 Bolded genes only are genes that have regions that fall within the top 1% PBS percentile.

Position	Gene	Distance to gene	PBS percentile	Annotation
Chr 1: 5290600	**STK35**	12594	1.85	
Chr 1:5290602	**STK35**	12592	1.85	Fertility, eye development and oxidative stress (Miyamoto et al 2018).
Chr 1:5290604	**STK35**	12590	1.85	
Chr 2: 17498913	SLC17A8	Intron	1.24	Glutamate neurotransmitter, inner hair cell dysfunction
Chr 3: **11687002**	**AGO3**	**Intron**	**0.34**	**RNA regulator**
Chr 3: 32286614	RBMS3	14363	22.93	
Chr 3: 32286627	RBMS3	14350	22.93	Involved in chondrogenic craniofacial development of zebrafish
Chr 3: 32286638	RBMS3	14339	22.93	(Jayasena and Bronner 2012).
Chr 3: 32286614	RBMS3	14363	22.93	
Chr 11: 27324192	GRIA3	Intron	2.85	
Chr 11: 27336469	GRIA3	2772	3.18	Glutamate receptor strongly expressed in zebrafish retina
Chr 11: 27336472	GRIA3	2775	3.18	development.
Chr 11: 27336476	GRIA3	2779	3.18	
Chr 11: **28255536**	**NA**	**3026**	**0.98**	**NA**
Chr 11: **28255616**	**NA**	**3106**	**0.98**	**NA**
Chr 11: **28256343**	**NA**	**3833**	**0.98**	**NA**
Chr 11: **28257367**	**NA**	**4857**	**0.98**	**NA**
Chr 11: 28260047	NA	7537	2.87	NA
Chr 12: 13854139	**C4orf22**	11903	1.30	Spermatogenesis, cilia, flagellum
Chr 24: **12693158**	**KDM6A**	**Intron**	**0.05**	**Kabuki syndrome (craniofacial development)**

1064

1065 **Table S4.** Summary statistics of the distance between closest adaptive candidate SNPs within the same

1066 gene

Gene	Median Distance	Mean Distance	SD	
TM6SF2	7430	7430	NA	2
C19orf10	4789	4789	NA	2
KDM6A	2892.5	2892.5	2410.52702	3
KCNB2	1803	1738.75	1614.92299	5
UBE2T	1742	1742	NA	2
TMEM230	1734	3039	3599.52427	4
ABCA1	1037.5	1097.5	959.375665	5
HAPLN4	974	1608.21739	9927.05273	24
GRIN2A	904	1270.6667	4273.38008	4
DPH1	608	1224.5	1999.25934	7
WNT10A	402	1442.17857	3986.39626	29
OPN1LW	386.5	359.5	225.229513	5
NIPSNAP3A	253	1451.71429	2819.19065	8
GALNT15	246.5	675.166667	3156.86746	13
SWS2	246	1381.25	2423.03892	5
STK35	217	793.366667	1992.8543	31
CRIM1	201	47.829268	2176.05381	42
PDYN	187	326.371429	355.925561	36
ADCYAP1R1	186.5	833	1304.35501	11
DNM1	186.5	459.708333	677.261781	25
DNM1	186.5	459.708333	677.261781	25
KCNH8	131.5	584.366667	1030.65532	31
TOP1	124	332.375	6687.63393	17
AGO3	115	357.625	552.097284	9
GMEB2	112	319.490196	2485.05809	52
GPNMB	57	57	77.781746	3
IGFBP4	26	2254	3880.67997	4
FAM83D	23	2523.66667	4347.74658	4
EXOSC3	17	17	NA	2
C4orf22	15	15	9.899495	3
TMEM106B	7	11.4	10.358571	6
THOC3	6	5.833333	3.600926	7
MID1	1	1	NA	2

1067

1068

Figure S1. μCT scans of Yucatán pupfish with maxilla colored in blue, premaxilla in pink,

1070   dentary in green, and articular in orange.

1071

Figure S2. Best fitting admixture graph assuming zero admixture events. Admixture graph

was found using the *findGraphs* function in *ADMIXTOOLS2* (124). Number on edges represent

drift weights.

1072

1073

1074

1075

1076

1077

1078   **Figure S3. ADMIXTURE results support genetic differentiation of Yucatán *Cyprinodon***

1079   **pupfish species.** (A) Cross-validation error for different values of K clusters, equally support 2,

1080   3, and 4 clusters. (B) ADMIXTURE plots showing the clusters at different values of K.

1081

1082

1095

Figure S4. Linkage decay occurs rapidly in *Cyprinodon simus*. Linkage decay was calculated

in windows of 1000KB using PLINK (v1.9). The results graphed are a random sample of all

calculated r^2 values taken for plotting purposes ($n = 9{,}645$ r^2 values).

1112

Figure S5. Adaptative SNPs associated with the same gene are generally spaced apart.

Distribution of the distance to nearest adaptive SNP within a gene. Note the log scale of the x-axis.

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

Figure S6. *f*-branch (*f_b*) **show that introgression with desert pupfish occurred before the**

Chichancanab radiation. Boxes with red color represent significant introgression between P3

species (top) and the species/internal branch on the left. For visualization, non-significant *f_b* were

changed to zero and are completely white boxes. Grey boxes represent comparisons that cannot

be made, as *f_b* cannot be calculated for introgression between sister taxon. Dashed lines on the

left are internal branches. SSI: San Salvador Island, Bahamas; NC: North Carolina, USA.

1129

Figure S7. **Both species did not have a strong response to either control scenario in the**

optomotor response experiment. (A) Number of complete rotations fish swam when the

machine was off and **(B)** when the machine was on but the apparatus was all black. Boxplots and

points of the number of complete rotations individuals performed during the "off-banded"

portion of the experiment. Note that some boxplots are compressed to a single line because most

individuals had no response.

1147

1148   **Figure S8**. **Species do not differ in sperm midpiece size or sperm flagellum length.** Boxplots

1149   and points of the mean sperm morphological estimate for an individual.

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170   **Figure S9. Tooth length was not affected by Wnt inhibition.** Boxplot and jittered points of
1171   average tooth length of the left and right teeth closest to the mandibular symphysis for *C.*
1172   *beltrani* larvae treated with either a control or Wnt inhibitor (iCRT14). Shapes represent different
1173   split-brood replicates.

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185 **Figure S10. Number of ossified teeth was not affected by Wnt inhibition.** Boxplot and jittered
1186 points of number of ossified teeth in *C. beltrani* larvae treated with either a control or Wnt
1187 inhibitor (iCRT14). Shapes represent different split-brood replicates.

1188