Parallel and non-parallel features of adaptive radiation in Yucatán pupfishes

Matthew C. Kustra[1,2,3,*], David Tian[1,2], M. Fernanda Palominos[1,2], Feifei Guo[1,2], Dylan Chau[1,2], Oskar Golwala[1,2], HoWan Chan[4], Andrés Alvarez Zapata[5], Reyna Guadalupe Cetz Paredes[5], Frida Ximena Cortés Sánchez[6], Sonia Gabriela Hernández[6], Adan Fernando Mar-Silva[6], Fernando Mex[7], Charles Tralka[1,2], Maribel Badillo-Alemán[7], Juan J. Schmitter-Soto[8], Carlos A. Gracida-Juárez[9], Christopher M. Martinez[10], Jairo Arroyave[6], and Christopher H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley, CA, United States
[2]Museum of Vertebrate Zoology, University of California, Berkeley, CA, United States
[3]Miller Institute for Basic Research in Science, University of California, Berkeley, CA, United States
[4]Department of BioSciences, Rice University, Houston, TX, United States
[5]Instituto Tecnologico Superior de Conkal, Yucatan, Mexico
[6]Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, Mexico
[7]Unidad Multidisciplinaria de Docencia e Investigación, Unidad Sisal, Universidad Nacional Autónoma de México, Sisal, Yucatán, Mexico
[8]Departamento de Sistemática y Ecología Acuática, El Colegio de la Frontera Sur, Chetumal, Quintana Roo, Mexico
[9]Subdirección de Posgrado e Investigación, Tecnológico Nacional de México, Campus Felipe Carrillo Puerto, Felipe Carrillo Puerto, Quintana Roo, Mexico.
[10]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA, United States

*Corresponding author: Matthew Kustra

Email: matthewckustra@berkeley.edu

Keywords: Speciation, De novo mutations, Introgression, Genomics, Adaptation

This PDF file includes:

      Main Text
      Figures 1 to 5
      Supporting Information (Supplemental Methods; Tables S1-S4; Figures S1-S10)

Abstract

Understanding the extent of parallelism across adaptive radiations remains a central

problem in evolutionary biology. We used whole-genome resequencing of 123 individuals to

compare the adaptive radiation of *Cyprinodon* pupfishes in Lake Chichancanab, Mexico, to an

independent radiation of San Salvador Island (SSI) pupfishes in the Bahamas, and assess the

repeatability of adaptive genetic architecture, sources of adaptive variation, and stages of

selection. Despite rapid craniofacial divergence of trophic specialists within 8-15 kya, only two

candidate genes (0.5%; 2/426) were shared between Caribbean radiations. Although adaptive

introgression played a major role in SSI, we found minimal evidence of adaptive introgression in

Chichancanab, likely due to the geographic isolation of this inland lake. Instead, de novo

mutations provided a substantial source of adaptive variation (30.6%) for the endemic

zooplanktivore, 15 times higher than the endemic scale-eater on SSI. However, in parallel with

SSI, we found strong evidence that adaptive divergence occurred in stages, first on regulatory

and standing genetic variation, then on de novo and nonsynonymous mutations. Consistent with

adaptive variants near opsin and spermatogenesis genes, functional categories unique to

Chichancanab, we found greater visual acuity and divergent sperm morphology in lab-reared

zooplanktivores relative to generalists using laboratory assays. Consistent with extensive

adaptive de novo mutations in *WNT10A* and rapid diversification of tooth size in the

zooplanktivore, we found that experimental inhibition of the Wnt pathway in generalists resulted

in narrower oral teeth. We conclude that de novo mutations, not introgression, can drive rapid

adaptive radiations in isolated environments.

Introduction

Much of biodiversity results from adaptive radiation—when a clade diversifies rapidly into three or more new species (1–5). Young adaptive radiations provide excellent systems for investigating speciation and adaptation (3, 6–9). However, it remains unclear to what degree radiations occur in parallel given similar starting positions on the adaptive landscape with similar levels of genetic diversity, gene flow, and ecological opportunity (10, 11).

Beneficial de novo mutations can provide novel variation to reach new fitness peaks on the adaptive landscape during adaptive radiation (12, 13). However, the longer time needed for de novo beneficial mutations to arise contradicts the short timescales of rapid divergence (11, 14, 15). Hybridization can introduce large amounts of genetic variation and novel phenotypes (transgressive segregation) through adaptive introgression and hybrid swarms, enabling populations to reach previously inaccessible fitness peaks (16). Support for an important role of hybridization in adaptive radiations has been demonstrated both theoretically (17, 18) and empirically in many systems (3, 19–27). However, we still do not know the relative importance of different sources of genetic variation for triggering and sustaining adaptive radiations across time and space.

Caribbean pupfishes are a remarkable system for assessing the repeatability of adaptive radiations due to two independent, recent adaptive radiations, each containing a generalist and multiple trophic specialists within isolated lake environments that have historically lacked predatory fishes and most competitors except *Gambusia*. The San Salvador Island (SSI), Bahamas radiation proceeded predominantly via the reassembly of pre-existing standing genetic variation found throughout the Caribbean, with substantial contributions of adaptive introgression to both the molluscivore (durophage) and scale-eating (lepidophage) specialists.

Only 0% and 2% of adaptive mutations in the molluscivore and scale-eater, respectively, were de

novo mutations unique to SSI (26, 28–32). However, it was previously unknown how a second

radiation in Lake Chichancanab, Mexico, drawing from the same Caribbean pool of standing

genetic variation, unfolded over the same timeframe.

Here, we use whole-genome resequencing of 123 individuals to provide insight into the

genomic architecture and genetic origins of the Lake Chichancanab *Cyprinodon* pupfish

radiation (33). This recent adaptive radiation (~8000 years old (34)) consisted of at least five

species (or up to seven (35)) historically (33, 35, 36), including a detritivore (*C. beltrani*),

piscivore (*C. maya*), rocky substrate specialist (*C. verecundus*), bivalve specialist (*C. labiosus*),

and zooplanktivore (*C. simus*). At the time of description (33), the relative abundances of these

endemic species were 68-85% *C. beltrani*, 6-18% *C. maya*, 2-13% *C. labiosus*, and <1% *C.*

simus and *C. verecundus*. However, due to recent colonization by invasive African tilapia

(*Oreochromis niloticus x mossambicus*) (37, 38) and by several species native to surrounding

basins but not Chichancanab (*Astyanax angustifrons, Mayaheros urophthalmus, Poecilia*

mexicana; pers. obs.), all trophic specialist species now appear to be extinct since our collections

in 2022, with only the generalist, *C. beltrani*, remaining in 2024. Despite exhibiting one of the

fastest rates of trait diversification of any vertebrate in tooth size (130 times faster than

background rates across *Cyprinodon* (29)) and rapid divergence in other craniofacial morphology

(Figure 1A; S1), previous genetic studies of microsatellite data and mtDNA suggest that this

radiation has low genetic differentiation (39, 40), enabling the identification of highly

differentiated loci among these species.

Figure 1. Yucatán *Cyprinodon* pupfishes show strong genetic differentiation. (A) μCT scans
of Yucatán pupfish with maxilla colored in blue, premaxilla in pink, dentary in green, and
articular in orange (see Figure S1 for different views). (B) Map of the Caribbean showing the
sampling sites for all species in our dataset with an inset map of Lake Chichancanab and an
aerial view image from one of the sampling sites. (C) Principal component analysis using a set of
linkage disequilibrium pruned SNPs. (D) Estimated ancestry proportions for each individual
inferred using *ADMIXTURE* with K = 4 (see Figure S3 for K=2-3) from the linkage-pruned
dataset. (E) Estimated demographic history with *MSMC2* on a log scale. Each colored line
represents an individual, with colors corresponding to different species.

Results

Genetic differentiation of the species flock

To measure population structure of the Yucatán *Cyprinodon* pupfishes, we conducted

whole genome resequencing of the species flock and the coastal sister species ($n = 15$ *C.*

artifrons, $n = 35$ *C. beltrani*, $n = 35$ *C. labiosus*, and $n = 17$ *C. simus*) using specimens collected

from the wild in 2022 (Figure 1B). We aligned these samples, as well as eleven other previously

sequenced Caribbean pupfish species, to the high-quality *Cyprindon nevadensis mionectes*

reference genome (GCA_030533455.1). After calling and filtering variants, our dataset included

23,960,536 SNPs (12.4x mean depth). Using a linkage disequilibrium pruned set of SNPs ($n =$

5,476,855 SNPs), we used principal component analysis and *ADMIXTURE* to visualize

population structure within Yucatán pupfishes. We found that all species formed distinct, non-

overlapping genetic clusters, with *C. artifrons* being the most genetically divergent from and

sister to the Chichancanab species (Figure 1C). Within the Chichancanab species flock, *C. simus*

was more genetically divergent from *C. labiosus* and *C. beltrani* (Figure 1C). These results were

also supported by genome-wide F_{ST} estimates and a phylogenetic tree constructed with

ADMIXTOOLS2 (Table S1; Figure S2). Four clusters corresponded to the four different species,

as determined by *ADMIXTURE* (Figure 1D). All *C. artifrons* individuals showed minimal

evidence of admixture with Chichancanab species (Figure 1D; Figure S3).

Demographic history

Using *MSMC2* for Chichancanab species with a mutation rate of 1.56×10^{-8} substitutions

per base pair (estimated from a Caribbean *Cyprinodon* species (26)), we found that the three

Chichancanab species (*C. beltrani*, *C. labiosus*, and *C. simus*) diverged from the coastal sister

species *C. artifrons* at the end of the Pleistocene 10-15 kya, consistent with receding sea levels

resulting in the formation of Lake Chichancanab (34). After divergence from *C. artifrons*, the

Chichancanab species flock showed a similar demographic history, characterized by a sharp

decline in effective population sizes, consistent with endemic sympatric species within an

isolated lake (Figure 1E).

Adaptive loci within trophic specialists

Although genome-wide differentiation among Chichancanab species was moderately low

(F_{ST} ranging from 0.032 to 0.09; Figure 2A; Table S1), we were able to identify highly

differentiated regions of the genome within the trophic specialist zooplanktivore *C. simus*. We

identified 19 fixed and 1,127 nearly-fixed ($F_{ST} > 0.95$) SNPs in *C. simus* relative to other

Chichancanab pupfishes (Tables S2-S3). In contrast, we found zero fixed or nearly-fixed SNPs

within the generalist *C. beltrani* or bivalve-specialist C. *labiosus*. However, we were able to

identify some moderately differentiated SNPs within *C. labiosus* ($n = 97$) and *C. beltrani* ($n =$

13) relative to the other Chichancanab species at an F_{ST} threshold of 0.8 (Table S2).

To search for signals of selection, we calculated the normalized four-taxon population

branch statistic (PBS_{nj}) across 10-kb non-overlapping windows because linkage disequilibrium

decayed at this distance (Figure S4). Simulation studies demonstrate that normalized *PBS* is

effective for identifying both hard and soft selective sweeps (41). Out of the 1,146 SNPs fixed or

nearly fixed within *C. simus*, 661 were within a strongly supported selective sweep window (top

1% PBS_{nj} outlier (41)), including 425 SNPs within 20 kb of the first or last exon of an annotated

gene. These 425 candidate adaptive SNPs ("adaptive variants") were located in the proximity of,

or within, 55 genes (Figure 2A), distributed across 19 out of 24 chromosomes (Figure 2A).

Adaptive variants associated with the same gene were often distributed across multiple introns or

exons and spaced apart by 1kb or more (Figure S5, Table S4).

Source and function of adaptive genetic variation in the zooplanktivore

172   We next characterized the location of the 425 candidate adaptive variants near or within a

gene in the zooplanktivore *C. simus*. More than 95% were in intronic or flanking regions (20-kb

upstream or downstream of a gene; Figure 2B). Within coding regions, 13 out of 20 adaptive

variants were nonsynonymous mutations (Figure 2B). The number of adaptive variants per gene

followed an exponential distribution, with a handful of genes containing many variants and most

genes with only a single adaptive variant (Figure 2B). We functionally characterized these genes

by manually checking online databases and doing a literature search (Table S3; Figure 2B). 32

out of the 55 candidate genes were associated with craniofacial development ($n =11$; 20%) or

sensory/neural function ($n = 21$; 38.2%). We manually checked if any of these genes overlapped

with the 426 unique candidate genes identified in the SSI radiation (26, 28, 30, 32, 42–46), and

we found only two overlapping candidate genes (*DNM1* and *SPTLC3*), both of which are

associated with motor neuron development (47, 48) and are scale-eater-specific candidate genes.

184   Next, we characterized the source of the variants and found that 63% of the adaptive

185   SNPs in *C. simus* also existed as standing genetic variation in the coastal Yucatan population of

186   *C. artifrons* ("standing variation"), 31% were *de novo* variants found only in Chichancanab, and

187   6% were detected as standing genetic variation in other pupfish species but not *C. artifrons*

188   (Figure 2C).

Figure 2. Architecture and source of adaptive genetic variation in the zooplanktivore. (A, top row) F_{st} between *C. simus* and *C. beltrani* + *C. labiosus* calculated in non-overlapping 10 kb windows. F_{st} values for candidate SNPs (red circles) were calculated separately per locus to show the location of fixed variants. (A, lower row) Population branch statistics (PBS_{nj}) with *C. simus* as the target population versus *C. beltrani*, *C. labiosus*, and *C. artifrons* in 10-kb non-overlapping windows. Candidate genes within 20 kb of fixed or nearly-fixed (Fst > 0.95) SNPs and within PBS_{nj} 1% outlier windows are highlighted in red and labeled. Alternating colors represent different chromosomes. (B) Candidate adaptive variants near or within a gene were defined as SNPs that were fixed ($n = 17$) or nearly fixed ($n = 1,129$; $F_{st} > 0.95$) and occurred within a 1% population branch statistic (PBS_{nj}) outlier window for *C. simus* (Venn diagram). Pie chart and bar graphs indicate the proportion of these 425 adaptive variants that were found within the flanking, intronic, or coding regions. (C) Pie chart and bar graphs indicate the proportion of these 425 variants that were detected only within Chichancanab (de novo: purple), as standing genetic variation in coastal sister species *C. artifrons* (pink), or in other Caribbean species, but not *C. artifrons*. * Indicates genes with a craniofacial annotation; ‡ indicates genes with neural or sensory annotation.

Limited adaptive introgression within Chichancanab pupfishes

To test for introgression into the Chichancanab radiation, we used f-statistics calculated in

Dsuite (49) to examine gene flow from 11 additional *Cyprinodon* species from across the clade

and the sister species to the genus *Megupsilon aporus*. Within Yucatán pupfishes, we found

evidence of introgression between *C. artifrons* and both *C. beltrani* and *C. labiosus*, with the

strongest signal in *C. beltrani* (Figure 3A, B). We also found evidence for introgression between

C. beltrani and most outgroup species examined, including the sister species to *Cyprinodon*,

Megupsilon aporus. Relative to *C. beltrani*, *C. labiosus* and *C. simus* also showed evidence of

introgression with pupfishes endemic to the Chihuahua desert (*C. eximius* and *C. fontinalis*;

Figure 3A, B). Furthermore, we also found evidence of introgression between the two sympatric

radiations (*C. variegatus* from SSI) and *C. simus* (Figure 3A, B). However, additional *f*-branch

statistics indicate that all patterns of excess allele sharing between these outgroup species besides

C. artifrons and the Chichancanab species flock likely resulted from ancient introgression events

before the colonization of Lake Chichancanab (Figure 3C; S6).

To test for adaptive introgression into *C. simus*, we next examined whether the set of

adaptive variants (Figure 2) occurred within outlier introgression windows using sliding window

scans of distance fraction (d_f(50)) for species that had significant genome-wide introgression

into *C. simus* (Figure 3A , B). We found that none of the outlier introgression windows

overlapped with *C. simus* adaptive variants (Figure 3D).

Next, we performed d_f scans for introgression between *C. artifrons* and Chichancanab

species. Although there were outlier introgression regions between *C. simus* and *C. artifrons*

(positive values d_f), these regions did not overlap with *C. simus* adaptive variants (Figure 3D).

However, the position of several of these *C. simus* adaptive variants overlapped with outlier

introgression windows for both *C. labiosus* and *C. beltrani* but not *C. simus* (Figure 3D). The

number of outlier introgression windows was much lower in *C. simus* relative to *C. labiosus* and

C. beltrani (Figure 3E, F). Furthermore, both *C. labiosus* and *C. beltrani* populations contained

several introgression tracts that were much longer than *C. simus* tract lengths, indicating much

more recent introgression events in these species (Figure 3E, F).

Figure 3. Minimal evidence of adaptive introgression into Chichancanab pupfishes.
Heatmaps of (A) Patterson's D-statistics and (B) F_4 ratios for P1-P2 species combinations within
the Chichancanab radiation along the x-axis and P3 focal species along the y-axis. All four-taxon
trees used *Cualac tessellatus* as the outgroup (P4). Only significant (Z-score > 3) results are
colored; combinations not relevant to the Chichancanab radiation are not shown. (C) f-branch (f_b)
statistics indicate introgression occurring before the Chichancanab radiation. Red boxes indicate
significant introgression between P3 species (top row) and the focal species or internal branch
(dashed lines) on the left. Non-significant f_b statistics are indicated by white boxes. Grey boxes
represent comparisons that cannot be made since f_b cannot be calculated for introgression
between sister taxa. See Figure S6 for full f-branch graphs. (D) Genome-wide plot of the distance
fraction (d_f, an introgression test statistic (50)) calculated in non-overlapping windows of 91
SNPs (approximately 10 kb) for species trios that showed evidence of introgression with *C.*
simus. d_f outlier windows that contained a candidate gene under selection in *C. simus* are colored
red and labeled with the gene name. The y-axis label gives the introgression topology: (P1, P2),
P3. Positive values indicate regions of introgression between *C. simus* and P3 (from top to
bottom: *C. eximius*, *C. fontinalis*, *C. variegatus* (SSI), and *C. artifrons* in the bottom two
subplots); negative values indicate regions of introgression between P1 (*C. beltrani* or *C.*
labiosus) and P3. Red dashed lines indicate outlier cutoffs, which were determined empirically
by the overall percentage of genome-wide introgression estimated from the F_4 ratios in panel B.
(E, F) Distribution of the size of d_f outlier introgression tract lengths (in 10 kb windows)
between *C. artifrons* and Chichancanab species. The introgression tracts containing candidate
adaptive genes in *C. simus* are labeled with gene names. SSI: San Salvador Island, Bahamas;
NC: North Carolina, USA.

The stages of adaptive divergence of the zooplanktivore

We used *starTMRCA* (51) to estimate the timing of selective sweeps for the set of

candidate adaptive variants near genes within the zooplanktivore *C. simus*. We estimated that

most selective sweeps occurred in a staggered sequence and not simultaneously after

colonization of Lake Chichancanab (Figure 4A)(34). There were also a few older sweeps that we

estimate may predate the age of Lake Chichancanab and divergence from the coastal sister

species *C. artifrons* (Figure 1B). Overall, we observed no clear temporal stages of adaptive

phenotypic divergence related to craniofacial genes (asterisks) or neural/sensory genes (‡; Figure

4A). Instead, we found that selection on these two broad functional categories occurred

throughout the period of divergence of *C. simus* following the colonization of Lake

Chichancanab, contrary to behavior-first, then craniofacial stages of adaptation (52) found in the

SSI radiation (26).

Next, we used generalized linear regressions to test if sweep age was correlated with the

proportion of nonsynonymous adaptive variants, de novo adaptive variants, or the overall

number of adaptive variants per gene region. We found that both nonsynonymous adaptive

variants ($\chi^2 = 5.852$, $p = 0.016$) and de novo mutations ($\chi^2 = 6.221$, $p = 0.013$) were significantly

more likely to occur in more recent selective sweeps during the final stage of adaptive

divergence in *C. simus* (Figure 4B, C). However, nonsynonymous mutations were not

significantly more likely to be de novo mutations found only in Chichancanab species (All

adaptive variants: $\chi^2 = 0.102$, $p = 0.749$; singleton adaptive variants: $\chi^2 = 2.010$, $p = 0.156$), and

there was no correlation between estimated sweep age and the number of candidate adaptive

variants per gene region ($\chi^2 = 0.111$, $p = 0.740$).

Figure 4. Stages of adaptive divergence in the zooplanktivore. (A) Points indicate median sweep age with 95% credible intervals. The dashed line indicates the approximate age of Lake Chichancanab (34). For each sweep, the first column of pie charts shows the proportions of flanking/intronic/synonymous/nonsynonymous, second column of pie charts shows the proportions of de novo/standing variation, and third column of squares shows the number of candidate adaptive variants per gene (from Figure 2). For genes with multiple adaptive variants, we used the variant with the median position as the focal position. * Indicates genes with a craniofacial annotation; ‡ indicates genes with neural or sensory annotation. (B) The proportion of nonsynonymous adaptive variants is significantly larger in more recent selective sweeps, indicated by the best-fit quasibinomial regression line; each pie chart represents a single candidate gene. (C) The proportion of de novo variants is significantly larger in more recent selective sweeps; each pie chart represents a single candidate gene.

Increased visual acuity in the zooplanktivore is consistent with vision-related candidate adaptive

genes

We identified several *C. simus* adaptive variants associated with eye development

(*ASIC3*(53), *CRIM1*(54), *MAB21L2*(55)) and opsin (*OPN1LW*, *SWS2*) genes. Because the trophic

specialization of *C. simus* requires visually locating zooplankton in the water column before

precise suction-feeding strikes, we hypothesized that this species would have higher visual acuity

than the detritivore, *C. beltrani*. We conducted optomotor response trials on lab-reared

individuals from both species (F1 *C. beltrani* and long-term laboratory colony of F10+ *C. simus*)

and found that *C. simus* has a much stronger optomotor response than *C. beltrani* in response to

black-and-white vertical bars spinning at a constant rate of approximately 90 rotations per

minute (Figure 5A, B; Wilcoxon signed-rank test: $W = 1$, $p = 0.009$). There was no significant

difference in rotations per minute between the species for the same individuals during control

observation periods with no spinning (Figure S7A; $W = 14$, $p = 0.3105$) or during positive control

observation periods with an all-black background spinning at the same frequency (Figure S7B;

$W = 9$, $p = 0.147$).

Figure 5. Divergence in visual acuity, sperm morphology, and tooth development reflects
candidate adaptive genes in the zooplanktivore. (A) Diagram of the optomotor response
experimental apparatus (adapted from (56)). (B) Boxplot and jittered points of the number of
complete rotations fish swam per minute, indicative of their optomotor response to black-and-
white vertical bars spinning at a constant rate. (C) Representative labeled image of a single *C.*
beltrani sperm at 100x stained with rose bengal. (D) Boxplots with jittered points of sperm head
diameter for *C. artifrons, beltrani,* and *simus.* Significant Tukey's post hoc differences are shown
with an asterisk. (E) Manhattan plot showing per-site F_{st} of SNPs within 50 kb of *WNT10A*.
Black rectangles indicate exons of *WNT10A*, and dashed lines indicate introns. Red points
represent nearly fixed SNPs ($F_{st} > 0.95$), and black points represent all other SNPs. The black
line is the smoothed average F_{st} in this region, with grey shading being the 95% confidence
intervals. (F) Representative images of *C. beltrani* larvae that were treated with DMSO (control;
top) or with iCRT14 Wnt inhibitor + DMSO (bottom) and stained with alizarin red (bone) and
alcian blue (cartilage). (G) Boxplot and jittered points of average tooth base width of the left and
right teeth closest to the mandibular symphysis for *C. beltrani* larvae treated with a DMSO
control or Wnt inhibitor (iCRT14) + DMSO. Shapes represent three independent split-brood
replicates.

Divergent sperm morphology of the zooplanktivore is consistent with spermatogenesis-related

candidate adaptive genes

Four candidate genes (*C4orf22*, *STK35*, *ERC2*, and *TBC1D25*) are known to affect

spermatogenesis/fertility (57–60). We therefore examined whether three Yucatan species, raised

in a common garden laboratory environment (F1 *C. artifrons,* F1 *beltrani,* and a long-term

laboratory colony of F10+ C. *simus*), differed in sperm morphology using differential

interference contrast microscopy. We found that species significantly differed in sperm head

length (Figure 5D; $F = 5.724$, $p = 0.025$), but not sperm midpiece length (Figure S8A; $F = 2.403$,

$p = 0.147$) or flagellum length (Figure S8B; $F = 0.414$, $p = 0.673$). *C. simus* had significantly

larger sperm heads than *C. beltrani* using Tukey's post hoc pairwise comparisons ($q = 3.367$; $p = 0.0207$; Fig. 5D).

Inhibition of WNT10A alters tooth morphology in Chichancanab pupfish

We identified a large number of highly differentiated SNPs surrounding the well-

characterized *WNT10A*, which controls tooth, hair, and skin development within the Wnt

signaling pathway (Figure 2; Figure 5E (61–63)). This is consistent with the most exceptional

morphological difference between *C. simus* and *C. beltrani,* extreme divergence in tooth size

(130 times faster than background rates (29)).

To test for the importance of *WNT10A* in tooth development within pupfishes, we used

iCRT14 to inhibit the Wnt / β catenin pathway downstream of WNT10A during metamorphosis

in *C. beltrani* lab-reared fry (64, 65). Chemical inhibition from 8 dpf to 22 dpf resulted in

narrower teeth than control fry treated with DMSO ($\chi^2 = 4.840$, $p = 0.0278$; Figure 5F, G). There

were no significant differences in mean tooth length ($\chi^2 = 0.455$, $p = 0.500$; Figure S9) or

number of ossified teeth when stained with alizarin red ($\chi^2 = 0.114$, $p = 0.7357$; Figure S10).

Discussion

Here, we investigated the genetic architecture, sources of adaptive variation, and

temporal dynamics of selection underlying speciation and trophic specialization in a young

adaptive radiation of *Cyprinodon* pupfishes endemic to Lake Chichancanab, Mexico. By

comparing these patterns to our previous work on a recent radiation of *Cyprinodon* pupfishes in

SSI, Bahamas (26, 28–32), we discovered parallel and non-parallel features of these independent

radiations in similar environments. Despite replicate radiations of Caribbean pupfishes in large,

isolated, predator-free, saline lakes with few competitors, we found little parallelism between the

radiations. Although craniofacial morphology diverged rapidly within trophic specialists in both

radiations, we found almost zero overlap in the adaptive genes underlying trophic specialization

(0.5%), potentially reflecting the divergent specialists within each radiation. In contrast to SSI,

there was no behavior-first stage of adaptive divergence in Chichancanab specialists and, with

the exception of craniofacial development, candidate adaptive gene ontologies potentially

relevant to speciation were not shared between radiations, such as pigmentation genes in SSI

versus opsin and spermatogenesis genes in Chichancanab. Unlike the SSI radiation and most

other well-studied adaptive radiations (e.g., Malawi cichlids (20), Galápagos finches (66)),

adaptive introgression likely played a minimal role in the Chichancanab radiation. We found no

overlap between adaptive variants in the specialists and tracts of introgression. Instead, over 30%

of the candidate adaptive variants in the zooplanktivore, *C. simus*, likely arose from de novo

mutations compared to only 2% in the SSI scale-eater specialist and none in the molluscivore

specialist. One outstanding parallel feature was a clear stage of refinement in the adaptive

divergence of the zooplanktivore, marked by increased rates of de novo and nonsynonymous

adaptive mutations occurring among the most recent selective sweeps in this species, similar to

the scale-eating specialist on SSI. In contrast to *C. simus*, we identified no sufficiently

differentiated SNPs within *C. labiosus*, the bivalve specialist, to identify candidate genes.

Nonetheless, minimal divergence of this specialist in Chichancanab parallels the minimal

divergence of the molluscivore on SSI, each in contrast to a more divergent trophic specialist.

Adaptive introgression did not contribute to speciation of Chichancanab pupfishes

Genomic analyses have revealed that hybridization and adaptive introgression played a

major role in many adaptive radiations (3, 11, 16, 19–27, 67, 68), including the *Cyprinodon* SSI

radiation (26, 31). However, in Chichancanab, we found minimal evidence of introgression

overall and did not detect any signatures of adaptive introgression following the colonization of

Lake Chichancanab. Although our analyses indicate extensive introgression among *Cyprinodon*

and *Megupsilon* lineages, we did not find any introgression events into the root of the radiation,

nor has there been secondary gene flow from outgroup species other than the sister species to the

radiation, *C. artifrons*. Secondary gene flow from *C. artifrons* was almost entirely detected with

C. beltrani and, to a lesser extent, *C. labiosus.* Gene flow may be higher with these species

because *C. artifrons,* a coastal benthic feeding detritivore, is less ecologically divergent from

both *C. beltrani,* a benthic feeding detritivore, and *C. labiosus*, a benthic feeding bivalve

specialist. Secondary gene flow between *C. artifrons* and both *C. beltrani* and *C. labiosus*

overlapped with many genes under selection in *C. simus*, suggesting that secondary gene flow is

helping increase similarities between the benthic detritivores within the lake and the coastal *C.*

artifrons population rather than by bringing in adaptive variation for trophic specialization in *C.*

simus.

These results are remarkably different from the *Cyprinodon* adaptive radiation of SSI,

where there is strong evidence of introgression into the root of the radiation and substantial

evidence of secondary adaptive introgression into both trophic specialists from at least three

different biogeographic regions of the Caribbean and Atlantic coast (26, 31). Lake Chichancanab

is an endorheic basin with no above-ground hydrological connections to the ocean and is

substantially more geographically isolated than SSI, a small island in the approximate center of

the Caribbean. Furthermore, the sister species to the Chichancanab species flock, *C. artifrons*, is

abundant along the entire coast of the Yucatán peninsula, suggesting that any gene flow into

Chichancanab would likely first have to pass through *C. artifrons* populations. An additional

factor is the water chemistry of Lake Chichancanab which is near-saturation with calcite and

gypsum and may be a greater physiological barrier to marine pupfish colonists than the slightly

hypersaline lakes of SSI (33-36). Thus, geography and ecology play an important role in

determining the relative importance of adaptive introgression in radiations.

De novo mutations primarily drive adaptive divergence of the zooplanktivore

The importance of de novo mutations in adaptive radiations remains contentious, in part

due to the time needed for de novo beneficial mutations to arise (11, 15). Interestingly, we found

that 30.6% of the zooplanktivore adaptive variants were detected only within Chichancanab

species, suggesting their de novo origins within the basin. This proportion is 15.3 times greater

than the contribution of de novo mutations to adaptive divergence of the scale-eater in the SSI

radiation. A primary factor could be that the estimated effective population size for the

Chichancanab radiation is approximately twice that of the SSI radiation (26), indicating that

selection on de novo mutations is more effective, with a higher probability of fixation, all else

being equal (14). There may also be a lack of existing standing genetic variation within

Caribbean pupfish populations that is adaptive for zooplanktivory. In support of this hypothesis,

although radiations should proceed along genetic lines of least resistance (69), the primary axis

of craniofacial morphological variation along which the zooplanktivore diverged is far removed

from the first two major axes of craniofacial variation within Caribbean pupfish species (29).

Evolving along the line of greatest resistance instead of least resistance may require de novo

mutations. While the SSI scale-eater is also highly divergent, it is closer to existing craniofacial

variation within Caribbean pupfish species and in the direction of greatest variance along the first

two principal component axes (29). Similarly, mutational target size for the craniofacial features

of a zooplanktivore may be much smaller than the features of a scale-eater, resulting in less

standing genetic variation available for adaptation to this trophic niche. Alternatively, we may

not have detected these de novo mutations in other populations, but this is unlikely to explain all

de novo mutations given our extensive sampling of *Cyprinodon* species across the clade and

spanning the Caribbean (Figure 1B). Thus, our results highlight that de novo mutations may play

a more important role in driving young adaptive radiations than currently appreciated (12, 13),

especially when ecotypes are highly divergent from other species within the radiation.

Minimal parallelism in the genes underlying adaptive divergence of trophic specialists

We found only two candidate genes (*DNM1* and *SPTLC3*) among the 55 zooplanktivore

candidate genes in the Chichancanab radiation that overlapped with the hundreds of candidate

genes identified so far in the SSI radiation (26, 28, 30, 32, 42–46). Consistent with trophic

specialization requiring the pursuit of mobile prey, both genes are involved in motor neuron

development. *DNM1* is a dynamin with nervous tissue-specific expression that is crucial for the

proper formation of motor neuron axons (47). *SPTLC3* is involved in the biosynthesis of

sphingolipids (70), and suppression of *SPTLC3* in zebrafish results in motor neuron axon defects

(48).

The extremely limited number of shared adaptive genes is surprising given the high

conservation of gene function in both morphological development (71, 72) and behavior (73),

and the frequent parallelism of candidate genes across vertebrate adaptive radiations, particularly

for craniofacial genes. For example, *BMP4* is involved in diversification in both Galápagos

finches (74) and African cichlids (75, 76). Furthermore, within the SSI radiation, many of the

same genes show parallel patterns of differential expression even within divergent trophic

specialists (26, 43). Nonetheless, parallelism of shared regulatory networks, rather than specific

genes, is still a possibility.

Rapid divergence in tooth morphology of the zooplanktivore through WNT10A

Another reason for non-parallelism is that trophic specialists in each radiation have

diverged in different sets of traits during adaptation to different ecological niches (29).

Specifically, the Chichancanab radiation has diversified in tooth size over 130 times faster than

background rates, driven by the extremely short and narrow teeth of the zooplanktivore (Figure

S1; (29). We identified a well-characterized tooth development gene in the Wnt pathway,

WNT10A, with many de novo mutations in flanking ($n = 2$) and intronic ($n = 13$) regions.

Mutations in *WNT10A* are associated with tooth developmental disorders in humans (62) and

both knockout and overexpression experiments in sticklebacks and zebrafish have demonstrated

WNT10A's role in teeth development and regeneration (61, 63). Consistent with *WNT10A*

playing a role in rapid divergence of tooth morphology, we found that inhibiting the Wnt

pathway in the *C. beltrani* generalist during pre-metamorphosis development resulted in

narrower teeth, similar to the reduced tooth sizes in the zooplanktivore *C. simus* (Figure 5C).

This indicates that the substantial number of de novo mutations in *WNT10A* may be suppressing

the Wnt pathway, resulting in a substantial reduction in tooth size in the zooplanktivore.

Increased visual acuity in the zooplanktivore is consistent with unique selection on vision-related

genes

One of the more striking features of both *Cyprinodon* radiations is the evolution of novel

trophic specialists from a generalist ancestor: the scale-eater (*C. desquamator*) in the SSI

radiation and the zooplanktivore (*C. simus*) in the Chichancanab radiation. Although these are

distinct dietary specializations, they share a similarity in that they require tracking free-

swimming prey. Indeed, four of the top 15 enriched GO terms of candidate genes in the SSI

scale-eater were related to eye development (26). Similarly, several of the candidate genes we

identified in the zooplanktivore were related to eye development, including *ASIC3*, *MAB21L2*,

and *CRIM1*. *CRIM1* is the most promising candidate because it is one of three candidate genes

with multiple nonsynonymous mutations (Figure 2). In zebrafish, a two-base pair deletion in

CRIM1 resulted in abnormal lenses as well as a large eye-to-head ratio (54). Interestingly, one of

the defining morphological features of *C. simus* is its large eye-to-head ratio (33) (Figure 1A).

Consistent with candidate genes related to eye development, we confirmed that *C. simus* exhibits

a much stronger optomotor response than *C. beltrani* (Figure 5B), indicating that this specialist

has greatly improved visual acuity, even when reared in a common laboratory environment.

Although both radiations contained many vision-related candidate genes, selection on

opsin genes was unique to Chichancanab. Most notably, we found several adaptive variants

associated with *SWS2* (blue-sensitive), and *OPN1LW* (red-sensitive) in the zooplanktivore.

Interestingly, both opsins contain many candidate de novo mutations, and *SWS2* is the youngest

selective sweep in the zooplanktivore. One explanation could be that most Caribbean

Cyprinodon species are benthic and live in similar habitats, resulting in limited standing genetic

variation in opsins. Additionally, most environments (including SSI and Chichancanab) are

relatively shallow, so differences in depth—a common selective pressure on opsin evolution

(77)—are likely not a strong selective force on SSI. Thus, the main selective force acting on

vision in Chichancanab may be prey capture efficiency. In many fish species, zooplanktivory is

associated with higher expression of shorter wavelength opsins (e.g., *SWS1* (77)), whereas

piscivory is not often associated with changes in opsin expression (78, 79).

Divergent sperm morphology in Chichancanab species is consistent with selection on

spermatogenesis genesis

Although the role that premating sexual selection plays in speciation has historically been

the main focus (80, 81), there is both theoretical and empirical support for the role of postmating

sexual selection in maintaining reproductive isolation via postmating prezygotic reproductive

barriers (82–87). In the SSI radiation, species differ in pigmentation, and a handful of the

candidate adaptive variants fall in regulatory regions of genes associated with pigmentation (26),

but not spermatogenesis. Furthermore, there is evidence for premating isolation by female

preference in the scale-eater (88). However, in Chichancanab, the two species tested in our data

set (*C. labiosus* and *C. beltrani*) were not found to visually distinguish between different species

in the radiation in laboratory mate choice trials (89).

Unlike the SSI radiation, we did not find any candidate adaptive variants associated with

pigmentation in Chichancanab; instead, we identified adaptive variants associated with four

spermatogenesis genes, *ERC2*, *TBC1D25*, *C4orf22*, and *STK35*. Consistent with previous work

demonstrating the importance of these genes in spermatogenesis (57–60), we found that *C. simus*

does indeed have larger sperm heads than *C. beltrani*, suggesting a unique role for post-mating

prezygotic reproductive barriers in this radiation (Figure 5D).

Parallelism in the stages of adaptation: refinement with de novo and nonsynonymous mutations

Adaptation may occur in stages, with less pleiotropic cis-regulatory changes occurring

before potentially more pleiotropic amino acid changes (26, 90, 91). Selection is also predicted

to act on standing variation first (14, 92). In both radiations, we observed strong evidence for a

refinement stage in which coding substitutions were selected on last. In Chichancanab, these

late-sweeping coding substitutions were also de novo mutations (Figure 4). The simplest

explanation for parallelism in more recent ages of adaptive de novo mutations is the waiting time

required for beneficial mutations to (1) arise and (2) reach a substantial frequency within the

population (14, 92). These variants may only be selected in the later stages of an adaptive walk

because nonsynonymous mutations are often pleiotropic and likely deleterious in the ancestral

genetic background (14, 93). Thus, they may not be advantageous until later in the adaptive

walk, when sign epistasis may reverse negative pleiotropic effects (94–98). Our results highlight

that this final refinement stage is likely a general pattern during adaptive divergence.

Materials and Methods

Sampling and genotyping

In 2022, we collected Lake Chichancanab specimens (*C. labiosus*, *C. beltrani*, and *C. simus*) and *C. artifrons* (Figure 1C). Additional species were collected from the wild or sourced from captive breeding populations in previous years. All specimens were euthanized in an overdose of buffered MS-222 (Fiquel, Inc.) following approved animal care and use protocols from the University of California, Berkeley and the University of California, Davis. All specimens used in this study are catalogued in the Museum of Vertebrate Zoology Fishes collection (MVZ:Fish:1410-1499).

Individual DNA samples were sequenced using Illumina HiSeq 4000 and NovaSeq. We mapped reads to the UCB_CyNevMio_1.0 *Cyprindon nevadensis mionectes* reference genome (GCA_030533455.1). Variants were called using *GATK* (v.4.5.0.0) and filtered according to best practices (99), resulting in 23,960,536 SNPs. See SI for more details.

Population structure and demographic history of Yucatán Cyprinodon *pupfishes*

To assess population structure, we first pruned SNPs in linkage disequilibrium using *PLINK* (v. 1.9), resulting in 5,476,855 SNPs. With this linkage disequilibrium pruned data set, we visualized population structure with a PCA (*PLINK*) and *ADMIXTURE* (v1.3) (100).

We estimated demographic history using *MSMC2* (101) using three individuals with the highest mean depth. See SI for more details.

Identifying and characterizing candidate genes

We calculated F_{st} genome-wide, per site, and in non-overlapping 10-kb windows in

VCFtools (v0.1.16) for all pairwise combinations of Yucatán *Cyprinodon* species (102). With the

10-kb windowed F_{st} values, we calculated a modified, normalized population branch statistic

(PBS), PBS_{nj} (103). We classified SNPs as candidate adaptive variants if they were fixed or

nearly fixed (per-site $F_{st} >0.95$) compared to other species within the radiation and were among

the top 1% outliers out of all PBS_{nj} windows, following the threshold used in (41).

We classified variants within 20 kb of a gene (upstream or downstream) as flanking. For

variants within a gene, we further classified them into intronic, synonymous, or nonsynonymous

using *SnpEff* (104). To characterize the origin of candidate variants, we extracted the alleles from

every species for each candidate variant. We classified variants that were found in any other

populations outside of Lake Chichancanab as standing genetic variation. We further subdivided

this group into variants that were found in other species but not in *C. artifrons*. Finally, if the

alternative variant was only detected within Chichancanab species, we classified it as de novo.

See SI for more details.

Introgression

We tested for evidence of introgression in Yucatán *Cyprinodon* species using *Dsuite* with

Cualac tessellatus as the outgroup, calculating f_4-ratios, D statistics, and f-branch statistics (49).

For species trios that showed evidence of introgression where *C. simus* was either P1or P2, we

then looked for introgression regions across the genome by calculating the distance fraction (d_f

(50)), in non-overlapping windows of 91 informative SNPs. To estimate introgression block

sizes, we merged neighboring windows that were introgression outliers and summed the total

size of the adjacent block. See SI for more details.

Timing of selective sweeps

For each candidate gene, we used *starTMRCA* (51) to estimate the age of selective

sweeps, extracting a 1-Mb region surrounding the candidate variant (51). We used a fixed

recombination rate of $2\text{x}10^{-8}$ (swordtail fishes; (105)) and a mutation rate of $1.56 \text{ x } 10^{-8}$

substitutions per base pair (Caribbean *Cyprinodon* species; 26). We then conducted separate

generalized linear models to test if the sweep age was correlated with the proportion of

nonsynonymous SNPs, proportion of de novo SNPs, and the number of SNPs. See SI for more

details.

Optomotor response, sperm morphology, and Wnt inhibition experiment

We tested for differences in visual acuity between the generalist/detritivore, *C. beltrani*,

and the zooplanktivore, *C. simus*, by conducting an optomotor response behavioral assay (56) on

lab-reared *C. beltrani* (F1; $n = 7$) and lab-reared *C. simus* (F10+; $n = 5$). Because the data were

nonparametric, we used a Wilcoxon rank-sum test to assess differences in optomotor response.

To test for differences in sperm morphology, we collected sperm samples from five

individuals per species in the lab (F10+*C. simus,* F1 *C. artifrons,* F1 *C. beltrani*). After fixing the

sperm sample in a 4% PFA solution stained with Rose Bengal, we plated and imaged sperm

under oil immersion with a 63x objective. We measured 10-30 sperm per individual in ImageJ

(106). We fit separate linear mixed-effects models for each trait of interest (sperm head,

midpiece, and flagellum lengths) with species as a fixed effect and a random intercept for

individual ID.

To confirm the role of the Wnt pathway in proper jaw and teeth development, we

conducted an experiment chemically inhibiting the Wnt pathway using iCRT14 during

metamorphosis (8dpf to 22 dpf) in *C. beltrani*. To control the exact time of development, we first

performed in vitro fertilizations and incubated developing embryos at ~26 °C until hatching

(~8dpf). On the day of hatching, we haphazardly split broods into control (DMSO) and

experimental groups (100 nanomolar iCRT14) to control for batch and family effects. After

euthanizing and staining fish with alizarin red (bone) and alcian blue (cartilage; (107)), we

quantified the number of ossified teeth and measured the tooth length and base width for the left

and right teeth closest to the mandibular symphysis using ImageJ (106). We fit separate linear

mixed-effects models for average tooth length and base width with experimental treatment as a

fixed effect and a random intercept for brood/replicate. For the number of ossified teeth, we used

a generalized linear mixed-effect model with a Poisson family and log-link function. See SI for

more details.

Acknowledgments

MCK was supported by a Miller Postdoctoral Research Fellowship from the Miller

Institute for Basic Research in Science, University of California Berkeley. This research was

funded by NSF CAREER 1749764 and NIH 5R01DE027052-02 awards to CHM. We thank the

Gerace Research Center and Troy Day for logistical support in the Bahamas, and the governments

of Mexico and the Bahamas BEST commission for permission to collect and export samples

(Permit No. PPF/DGOPA-001/20). All research procedures and animal care protocols (AUP-2021-

02-14062-1 and AUP-2021-07-14515) were approved by the University of California, Berkeley

Animal Care and Use committee and the UC Davis Institutional Animal Care and Use committee.

Specimens are deposited in the Museum of Vertebrate Zoology (MVZ:Fish:1410-1499).

References

1.   R. G. Gillespie, *et al.*, Comparing adaptive radiations across space, time, and taxa. *J.*
Hered. **111**, 1–20 (2020).

2.   C. H. Martin, E. J. Richards, The paradox behind the pattern of rapid adaptive radiation:
how can the speciation process sustain itself through an early burst? *Annu. Rev. Ecol. Evol.*
Syst. **50**, 569–593 (2019).

3.   Y. Naciri, H. P. Linder, The genetics of evolutionary radiations. *Biol. Rev.* **95**, 1055–1072
(2020).

4.   D. Schluter, *The ecology of adaptive radiation* (OUP Oxford, 2000).

5.   J. T. Stroud, J. B. Losos, Ecological opportunity and adaptive radiation. *Annu. Rev. of*
Ecol. Evol. Syst. **47**, 507–532 (2016).

6.   R. Butlin, *et al.*, What do we need to know about speciation? *Trends Ecol. Evol.* **27**, 27–39
(2012).

7.   J. A. Coyne, H. A. Orr, *Speciation* (Oxford University Press, 2004).

8.   D. R. Matute, B. S. Cooper, Comparative studies on speciation: 30 years since Coyne and
Orr. *Evolution* **75**, 764–778 (2021).

9.   D. Schluter, L. H. Rieseberg, Three problems in the genetics of speciation by selection.
Proc. Natl. Acad. Sci. U.S.A. **119**, e2122153119 (2022).

10. D. Berner, W. Salzburger, The genomics of organismal diversification illuminated by
adaptive radiations. *Trends Genet.* **31**, 491–499 (2015).

11. D. A. Marques, J. I. Meier, O. Seehausen, A combinatorial view on speciation and adaptive
radiation. *Trends Ecol. Evol.* **34**, 531–544 (2019).

12. J. B. Pease, D. C. Haak, M. W. Hahn, L. C. Moyle, Phylogenomics reveals three sources
of adaptive variation during a rapid radiation. *PLOS Biol.* **14**, e1002379 (2016).

13. B. W. Stone, C. A. Wessinger, Ecological diversification in an adaptive radiation of plants:
The role of de novo mutation and introgression. *Mol. Biol. Evol.* **41**, msae007 (2024).

14. R. D. H. Barrett, D. Schluter, Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38–44 (2008).

15. M. Rousselle, *et al.*, Is adaptation limited by mutation? A timescale-dependent effect of genetic diversity on the adaptive substitution rate in animals. *PLOS Genet.* **16**, e1008668 (2020).

16. O. Seehausen, Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19**, 198–207 (2004).

17. K. Kagawa, G. Takimoto, O. Seehausen, Transgressive segregation in mating traits drives hybrid speciation. *Evolution* **77**, 1622–1633 (2023).

18. K. Kagawa, G. Takimoto, Hybridization can promote adaptive radiation by means of transgressive segregation. *Ecol. Lett.* **21**, 264–274 (2018).

19. R. De-Kayne, *et al.*, Genomic architecture of adaptive radiation and hybridization in Alpine whitefish. *Nat. Commun.* **13**, 4479 (2022).

20. D. A. Joyce, *et al.*, Repeated colonization and hybridization in Lake Malawi cichlids. *Curr. Biol.* **21**, R108–R109 (2011).

21. S. Lamichhaney, *et al.*, Evolution of Darwin's finches and their beaks revealed by genome sequencing. *Nature* **518**, 371–375 (2015).

22. C. H. Martin, *et al.*, Complex histories of repeated gene flow in Cameroon crater lake cichlids cast doubt on one of the clearest examples of sympatric speciation. *Evolution* **69**, 1406–1422 (2015).

23. J. I. Meier, *et al.*, Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat. Commun.* **8**, 14363 (2017).

24. C. Pouchon, *et al.*, Phylogenomic analysis of the explosive adaptive radiation of the *Espeletia* complex (Asteraceae) in the tropical Andes. *Syst. Biol.* **67**, 1041–1060 (2018).

25. E. J. Richards, J. W. Poelstra, C. H. Martin, Don't throw out the sympatric speciation with the crater lake water: fine-scale investigation of introgression provides equivocal support for causal role of secondary gene flow in one of the clearest examples of sympatric speciation. *Evol. Lett.* **2**, 524–540 (2018).

26. E. J. Richards, *et al.*, A vertebrate adaptive radiation is assembled from an ancient and disjunct spatiotemporal landscape. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2011811118 (2021).

27. N. Rosser, *et al.*, Hybrid speciation driven by multilocus introgression of ecological traits. *Nature* **628**, 811–817 (2024).

28. M. E. S. John, J. C. Dunker, E. J. Richards, S. Romero, C. H. Martin, Parallel evolution of integrated craniofacial traits in trophic specialist pupfishes. *Ecol. Evol.* **14**, e11640 (2024).

728   29.   C. H. Martin, P. C. Wainwright, Trophic novelty is linked to exceptional rates of
729   morphological diversification in two adaptive radiations of *Cyprinodon* pupfish. *Evolution*
730   **65**, 2197–2212 (2011).

731   30.   A. H. Patton, E. J. Richards, K. J. Gould, L. K. Buie, C. H. Martin, Hybridization alters
732   the shape of the genotypic fitness landscape, increasing access to novel fitness peaks
733   during adaptive radiation. *eLife* **11**, e72905 (2022).

734   31.   E. J. Richards, C. H. Martin, Adaptive introgression from distant Caribbean islands
735   contributed to the diversification of a microendemic adaptive radiation of trophic
736   specialist pupfishes. *PLOS Genet.* **13**, e1006919 (2017).

737   32.   E. J. Richards, C. H. Martin, We get by with a little help from our friends: shared adaptive
738   variation provides a bridge to novel ecological specialists during adaptive radiation. *Proc.*
739   *Biol. Sci.* **289**, 20220613 (2022).

740   33.   J. M. Humphries, R. R. Miller, A Remarkable Species Flock of Pupfishes, Genus
741   Cyprinodon, from Yucatán, México. *Copeia* **1981**, 52–64 (1981).

742   34.   D. A. Hodell, J. H. Curtis, M. Brenner, Possible role of climate in the collapse of Classic
743   Maya civilization. *Nature* **375**, 391–394 (1995).

744   35.   U. Strecker, Description of a new species from Laguna Chichancanab, Yucatan, Mexico:
745   *Cyprinodon suavium* (Pisces: Cyprinodontidae). *Hydrobiologia* **541**, 107–115 (2005).

746   36.   J. M. Humphries, *Cyprinodon verecundus,* n. sp., a fifth species of pupfish from Laguna
747   Chichancanab. *Copeia* **1984**, 58–68 (1984).

748   37.   C. A. Gracida-Juárez, J. J. Schmitter-Soto, M. J. Genner, Community structure of
749   indigenous fishes relative to habitat variation and invasive tilapia in lakes of Quintana
750   Roo, Mexico. *Environ. Biol. Fish.* **107**, 401–414 (2024).

751   38.   U. Strecker, The impact of invasive fish on an endemic *Cyprinodon* species flock
752   (Teleostei) from Laguna Chichancanab, Yucatan, Mexico. *Ecol. Freshw. Fish* **15**, 408–418
753   (2006).

754   39.   U. Strecker, Genetic differentiation and reproductive isolation in a *Cyprinodon* fish species
755   flock from Laguna Chichancanab, Mexico. *Mol. Phyl. and Evol.* **39**, 865–872 (2006).

756   40.   U. Strecker, C. G. Meyer, C. Sturmbauer, H. Wilkens, Genetic divergence and speciation
757   in an extremely young species flock in Mexico formed by the genus *Cyprinodon*
758   (Cyprinodontidae, Teleostei). *Mol. Phyl. Evol.* **6**, 143–149 (1996).

759   41.   M. Shpak, K. N. Lawrence, J. E. Pool, The precision and power of population branch
760   statistics in identifying the genomic signatures of local adaptation. *Genome Biol. Evol.* **17**,
761   evaf080 (2025).

42. J. A. McGirr, C. H. Martin, Novel candidate genes underlying extreme trophic
specialization in Caribbean pupfishes. *Mol. Biol. Evol.* **34**, 873 (2017).

43. J. A. McGirr, C. H. Martin, Parallel evolution of gene expression between trophic
specialists despite divergent genotypes and morphologies. *Evol. Lett.* **2**, 62–75 (2018).

44. J. A. McGirr, C. H. Martin, Ecological divergence in sympatry causes gene misexpression
in hybrids. *Mol. Ecol.* **29**, 2707–2721 (2020).

45. J. A. McGirr, C. H. Martin, Few fixed variants between trophic specialist pupfish species
reveal candidate cis-regulatory alleles underlying rapid craniofacial divergence. *Mol. Biol.*
Evol. **38**, 405–423 (2021).

46. M. F. Palominos, V. Muhl, C. H. Martin, Craniofacial-specific transcriptomics uncovers
novel genes underlying jaw divergence in dietary specialist pupfishes. *Genetics* **231**,
iyaf207 (2025).

47. C. Bragato, *et al.*, Zebrafish dnm1a gene plays a role in the formation of axons and
synapses in the nervous tissue. *J. Neurosci. Res.* **101**, 1345–1359 (2023).

48. C. Gonzaga-Jauregui, *et al.*, Exome sequence analysis suggests that genetic burden
contributes to phenotypic variability and complex neuropathy. *Cell Rep.* **12**, 1169–1183
(2015).

49. M. Malinsky, M. Matschiner, H. Svardal, Dsuite - Fast D-statistics and related admixture
evidence from VCF files. *Mol. Ecol. Res.* **21**, 584–595 (2021).

50. B. Pfeifer, D. D. Kapan, Estimates of introgression as a function of pairwise distances.
BMC Bioinform. **20**, 207 (2019).

51. J. Smith, G. Coop, M. Stephens, J. Novembre, Estimating time to the common ancestor for
a beneficial allele. *Mol. Biol. Evol.* **35**, 1003–1017 (2018).

52. J. T. Streelman, P. D. Danley, The stages of vertebrate evolutionary radiation. *Trends Ecol.*
Evol. **18**, 126–131 (2003).

53. W.-G. Li, T.-L. Xu, ASIC3 Channels in multimodal sensory perception. *ACS Chem.*
Neurosci. **2**, 26–37 (2011).

54. T. Le, *et al.*, A zebrafish model of crim1 loss of function has small and misshapen lenses
with dysregulated clic4 and fgf1b expression. *Front. Cell Dev. Biol.* **13** (2025).

55. B. N. Kennedy, *et al.*, Zebrafish *rx3* and *mab21l2* are required during eye morphogenesis.
Dev. Biol. **270**, 336–349 (2004).

56. S. C. F. Neuhauss, *et al.*, Genetic disorders of vision revealed by a behavioral screen of
400 essential loci in zebrafish. *J. Neurosci.* **19**, 8603–8615 (1999).

57.   M. H. K. Chau, *et al.*, Investigation of the genetic etiology in male infertility with
796   apparently balanced chromosomal structural rearrangements by genome sequencing. *Asian*
797   *J. of Androl.* **24** (2022).

798   58.   H. Li, Y. Dai, Z. Luo, D. Nie, Cloning of a new testis-enriched gene C4orf22 and its role
799   in cell cycle and apoptosis in mouse spermatogenic cells. *Mol. Biol. Rep.* **46**, 2029–2038
800   (2019).

801   59.   Y. Miyamoto, *et al.*, The STK35 locus contributes to normal gametogenesis and encodes a
802   lncRNA responsive to oxidative stress. *Biol. Open* **7**, bio032631 (2018).

803   60.   S. Nawaz, S. Hussain, S. Basit, W. Ahmad, First evidence of involvement of *TBC1D25* in
804   causing human male infertility. *Eur. J. Med. Genet.* **64**, 104142 (2021).

805   61.   T. A. Square, *et al.*, Modulation of tooth regeneration through opposing responses to Wnt
806   and BMP signals in teleosts. *Development* **150**, dev202168 (2023).

807   62.   S. Nawaz, *et al.*, WNT10A missense mutation associated with a complete odonto-onycho-
808   dermal dysplasia syndrome. *Eur. J. Hum. Genet.* **17**, 1600–1605 (2009).

809   63.   E. L. Benard, *et al.*, WNT10A is required for zebrafish median fin fold maintenance and
810   adult unpaired fin metamorphosis. *Dev. Dyn.* **253**, 566–592 (2024).

811   64.   J. L. Catrow, Y. Zhang, M. Zhang, H. Ji, Discovery of selective small-molecule inhibitors
812   for the β-catenin/T-cell factor protein–protein interaction through the optimization of the
813   acyl hydrazone moiety. *J. Med. Chem.* **58**, 4678–4692 (2015).

814   65.   F. C. Gonsalves, *et al.*, An RNAi-based chemical genetic screen identifies three small-
815   molecule inhibitors of the Wnt/wingless signaling pathway. *Proc. Natl. Acad. Sci. U.S.A.*
816   **108**, 5954–5963 (2011).

817   66.   E. D. Enbody, *et al.*, Community-wide genome sequencing reveals 30 years of Darwin's
818   finch evolution. *Science* **381**, eadf6218 (2023).

819   67.   J. V. Peñalba, *et al.*, The role of hybridization in species formation and persistence. *Cold*
820   *Spring Harb. Perspect. Biol.* **16**, a041445 (2024).

821   68.   O. Seehausen, Conditions when hybridization might predispose populations for adaptive
822   radiation. *J. Evol. Biol.* **26**, 279–281 (2013).

823   69.   D. Schluter, Adaptive radiation along genetic lines of least resistance. *Evolution* **50**, 1766–
824   1774 (1996).

825   70.   T. Hornemann, *et al.*, The SPTLC3 subunit of serine palmitoyltransferase generates short
826   chain sphingoid bases. *J. of Biol. Chem.* **284**, 26322–26330 (2009).

827   71.   M. E. Bronner, N. M. LeDouarin, Development and evolution of the neural crest: An
828   overview. *Dev. Biol.* **366**, 2–9 (2012).

72.   F. Hirth, H. Reichert, Conserved genetic programs in insect and mammalian brain
development. *BioEssays* **21**, 677–684 (1999).

73.   C. J. Reaume, M. B. Sokolowski, Conservation of gene function in behaviour. *Philos.*
Trans. R. Soc. B. **366**, 2100–2110 (2011).

74.   A. Abzhanov, M. Protas, B. R. Grant, P. R. Grant, C. J. Tabin, Bmp4 and morphological
variation of beaks in Darwin's finches. *Science* **305**, 1462–1465 (2004).

75.   R. C. Albertson, J. T. Streelman, T. D. Kocher, P. C. Yelick, Integration and evolution of
the cichlid mandible: The molecular basis of alternate feeding strategies. *Proc. Natl. Acad.*
Sci. U.S.A. **102**, 16287–16292 (2005).

76.   Y. Terai, N. Morikawa, N. Okada, The evolution of the pro-domain of bone morphogenetic
protein 4 (Bmp4) in an explosively speciated lineage of East African cichlid fishes. *Mol.*
Biol. Evol. **19**, 1628–1632 (2002).

77.   K. L. Carleton, D. Escobar-Camacho, S. M. Stieb, F. Cortesi, N. J. Marshall, Seeing the
rainbow: mechanisms underlying spectral sensitivity in teleost fishes. *J. Exp. Biol.* **223**,
jeb193334 (2020).

78.   C. M. Hofmann, *et al.*, The eyes have it: regulatory and structural changes both underlie
cichlid visual pigment diversity. *PLOS Biol.* **7**, e1000266 (2009).

79.   K. E. O'Quin, C. M. Hofmann, H. A. Hofmann, K. L. Carleton, Parallel evolution of opsin
gene expression in African cichlid fishes. *Mol. Biol. Evol.* **27**, 2839–2854 (2010).

80.   M. G. Ritchie, Sexual selection and speciation. *Annu. Rev. Ecol. Evol. Syst.* **38**, 79–102
(2007).

81.   M. R. Servedio, J. W. Boughman, The role of sexual selection in local adaptation and
speciation. *Annu. Rev. Ecol. Evol. Syst.* **48**, 85–109 (2017).

82.   A. Devigili, *et al.*, Possible glimpses into early speciation: the effect of ovarian fluid on
sperm velocity accords with post-copulatory isolation between two guppy populations. *J.*
Evol. Biol. **31**, 66–74 (2018).

83.   M. D. Garlovsky, *et al.*, Synthesis and scope of the role of postmating prezygotic isolation
in speciation. *Cold Spring Harb. Perspect. Biol.* **16**, a041429 (2023).

84.   M. C. Kustra, M. R. Servedio, S. H. Alonzo, Cryptic female choice can maintain
reproductive isolation. *Evolution* **79**, 2259–2273 (2025).

85.   M. C. Kustra, S. H. Alonzo, The coevolutionary dynamics of cryptic female choice. *Evol.*
Lett. **7**, 191–202 (2023).

86.   P. D. Lorch, M. R. Servedio, The evolution of conspecific gamete precedence and its effect
on reinforcement. *J. Evol. Biol.* **20**, 937–949 (2007).

87.   J. L. Marshall, M. L. Arnold, D. J. Howard, Reinforcement: the road not taken. *Trends Ecol. Evol.* **17**, 558–563 (2002).

88.   A. Kodric-Brown, R. J. D. West, Asymmetries in premating isolating mechanisms in a sympatric species flock of pupfish (*Cyprinodon*). *Behav. Ecol.* **25**, 69–75 (2014).

89.   U. Strecker, A. Kodric-Brown, Mate recognition systems in a species flock of Mexican pupfish. *Journal of Evolutionary Biology* **12**, 927–935 (1999).

90.   L. Campagna, *et al.*, Repeated divergent selection on pigmentation genes in a rapid finch radiation. *Science Advances* **3**, e1602404 (2017).

91.   F. C. Jones, *et al.*, The genomic basis of adaptive evolution in threespine sticklebacks. *Nature* **484**, 55–61 (2012).

92.   Z. D. Blount, R. E. Lenski, J. B. Losos, Contingency and determinism in evolution: Replaying life's tape. *Science* **362**, eaam5979 (2018).

93.   P. W. Hedrick, Adaptive introgression in animals: examples and comparison to new mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* **22**, 4606–4618 (2013).

94.   S. G. Das, M. Mungan, J. Krug, Epistasis-mediated compensatory evolution in a fitness landscape with adaptational tradeoffs. *Proc. Natl. Acad. Sci. U.S.A.* **122**, e2422520122 (2025).

95.   M. J. Harms, J. W. Thornton, Historical contingency and its biophysical basis in glucocorticoid receptor evolution. *Nature* **512**, 203–207 (2014).

96.   M. Karageorgi, *et al.*, Genome editing retraces the evolution of toxin resistance in the monarch butterfly. *Nature* **574**, 409–412 (2019).

97.   R. D. Tarvin, *et al.*, Interacting amino acid replacements allow poison frogs to evolve epibatidine resistance. *Science* **357**, 1261–1266 (2017).

98.   D. M. Weinreich, N. F. Delaney, M. A. DePristo, D. L. Hartl, Darwinian evolution can follow only very few mutational paths to fitter proteins. *Science* **312**, 111–114 (2006).

99.   A. McKenna, *et al.*, The genome analysis toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

100. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

101. S. Schiffels, K. Wang, "MSMC and MSMC2: The multiple sequentially markovian coalescent" in *Statistical Population Genomics*, J. Y. Dutheil, Ed. (Springer US, 2020), pp. 147–166.

102. P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

103. J. M. Schmidt, M. de Manuel, T. Marques-Bonet, S. Castellano, A. M. Andrés, The impact
of genetic adaptation on chimpanzee subspecies differentiation. *PLOS Genet.* **15**,
e1008485 (2019).

104. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster*
strain w1118; iso-2; iso-3. *Fly* **6**, 80–92 (2012).

105. M. Schumer, *et al.*, Natural selection interacts with recombination to shape the evolution
of hybrid genomes. *Science* **360**, 656–660 (2018).

106. C. T. Rueden, *et al.*, ImageJ2: ImageJ for the next generation of scientific image data.
BMC Bioinform. **18**, 529 (2017).

107. M. Walker, C. Kimmel, A two-color acid-free cartilage and bone stain for zebrafish larvae.
Biotech. Histochem. **82**, 23–28 (2007).

Supporting Information for

Parallel and non-parallel features of adaptive radiation in Yucatán pupfishes

Matthew C. Kustra [*], David Tian, M. Fernanda Palominos, Feifei Guo, Dylan Chau, Oskar Golwala, HoWan Chan, Andrés Alvarez Zapata, Reyna Guadalupe Cetz Paredes, Frida Ximena Cortés Sánchez, Sonia Gabriela Hernández, Adan Fernando Mar-Silva, Fernando Mex, Charles Tralka, Maribel Badillo-Alemán, Juan J. Schmitter-Soto, Carlos A. Gracida-Juárez, Christopher M. Martinez, Jairo Arroyave, and Christopher H. Martin

[*]**Corresponding author:** Matthew Kustra

Email: matthewckustra@berkeley.edu

The supporting information includes:

Supporting text (SI methods)

Tables S1 to S4

Figures S1 to S10

SI References

SI Methods

Sampling

We collected Lake Chichancanab specimens in 2022 over two days using a 5 x 1.3 m seine net with 1.6 mm mesh size. We sampled two sites, an entrance road by La Presumida, and a bridge crossing the narrow section of the lake on the road to San Diego (Fig. 1C). *C. labiosus* and *C. beltrani* were sampled from both sites for this study, whereas *C. simus* was only found at La Presumida. *C. artifrons* were collected by cast net from a coastal estuary at Sisal. Additional specimens were collected from the Bahamas (Lake Cunningham, New Providence Island and Crescent Pond, San Salvador Island), the Dominican Republic (Laguna Bavaro), or Fort Fisher, North Carolina as described previously (Martin 2016) or sourced from the American Killifish Association, the London Zoological Society, the Dallas Children's Aquarium (*Megupsilon aporus*), and Frans Vermuelen (*Cyprinodon dearborni*). All specimens were euthanized in an overdose of buffered MS-222 (Fiquel, Inc.) following approved animal care and use protocols from the University of California, Berkeley and the University of California, Davis. All specimens used in this study are catalogued in the Museum of Vertebrate Zoology Fishes collection (MVZ:Fish:1410-1499).

The London Zoological Society provided our live colony of *C. simus* in 2015, originating from a much earlier field collection, which we subsequently raised for ten generations. We collected live colonies of *C. beltrani* and *C. artifrons* in 2024 from La Presumida, Lake Chichancanab and Sisal, respectively, over four days using a 5 x 1.3 m seine net with 1.6 mm mesh size.

Sequencing, Variant calling and filtering

Individual DNA samples were resequenced using Illumina Hiseq 4000 and Novaseq. Raw

reads were trimmed with *fastp* (v0.23.4) (1). Reads were mapped to the UCB_CyNevMio_1.0

Cyprindon nevadensis mionectes reference genome (GCA_030533455.1) with *bwa mem*

(v0.7.17) (2). We followed the *GATK* (v.4.5.0.0) genotyping pipeline to call variants (119).

Duplicate reads in the bam files were marked with *Picard MarkDuplicates* (GATK v4.5.0.0)

(118). Coverage and mapping quality were assessed with *qualimap* (v2.2.2) (4). We used

HaplotypeCaller (-ERC GVCF) v4.5.0.0 to call variants for each individual and stored variants

in the GenomicsDB datastore format prior to using *GenotypeGVCFs* to perform joint

genotyping. We restricted our analyses to biallelic SNPs and applied the recommended *GATK*

hard filters (QD < 2, QUAL < 30, SOR > 3, FS > 60, MQ < 40, MQRankSum < -12.5,

ReadPosRankSum < -8) to filter SNPs (118, 120). Missing genotypes were reset to ./. from 0/0

due to how *GATK* represents missing genotypes in version 4.5.0.0 (7).

After hard-filtering and variant calling, we filtered the data set further by removing SNPs

with minor allele frequency < 0.05, more than 10% missing, depth less than 10x, and GQ less

than 20 using *BCFtools* (v1.16)(8). This resulted in a final data set that included 23,960,536

SNPs.

Population structure of Yucatán Cyprinodon *pupfishes*

To assess population structure, we first pruned SNPs in linkage disequilibrium using

PLINK (v. 1.9) with the following parameters: "-indep-pairwise 50 5 0.2" (9). This filtered our

data set from 23,960,536 SNPs to 5,476,855 SNPs. With this linkage disequilibrium pruned data

set, we first filtered the full VCF file to only include relevant Yucatán *Cyprinodon* species (i.e.,

C. artifrons, C. beltrani, C. labiosus, and *C. simus*), then conducted a PCA using *PLINK*. Next,

we used *ADMIXTURE* (v1.3) to determine the optimal number of population clusters that best fit

the data and to assign individuals to their corresponding population clusters (10). We performed

this analysis and calculated cross-validation error for the number of clusters K = 1-6. Models in

ADMIXTURE with K=2-4 were equally supported (Figure S2A). However, given morphological

distinctness and the results of the PCA, we present the results for when K = 4 (Figure 1D).

1004

Demographic history of Yucatán Cyprinodon *pupfish*

We estimated demographic history using *MSMC2* (11). For each Yucatán pupfish species,

we selected three individuals with the highest mean depth and generated single-sample VCF

files, individual mask files, and "mapability" mask files. We then ran *MSMC2* with default

parameters except we set phasing to "unphased" and "P_PAR=8*1+25*1+1*2+1*3." For

plotting, we used a generation time of 1 year and a mutation rate of 1.56×10^{-8} substitutions per

base pair, estimated from lab-reared pedigrees of Caribbean *Cyprinodon* species (12).

1012

Identifying candidate genes

We calculated F_{st} genome-wide, per site, and in non-overlapping 10-kb windows using

the "--weir-fist-pop" function in *VCFtools*(v0.1.16) for all pairwise combinations of Yucatán

Cyprinodon species (13). We chose 10-kb windows because linkage disequilibrium substantially

decayed at this distance (Figure S3), and these windows allowed us to quantify fine-scale

genomic variation. With the 10-kb windowed F_{st} values, we calculated a modified population

branch statistic (PBS) for four taxa, PBS_{nj} (14). This statistic is useful because it makes no

assumptions about species topology or polarization (14). We then normalized PBS_{nj} by the total

tree length, as normalized *PBS* is most effective and specific at identifying selective sweeps (15).

We classified SNPs as candidate adaptive variants if they were fixed or nearly fixed (per-site F_{st}

>0.95) compared to other species within the radiation and were among the top 1% outliers out of

all PBS_{nj} windows, following the threshold used in (15).

1025

Introgression

Using the full VCF file, which contained a wide range of species in the family

Cyprinodontidae, we tested for evidence of introgression in Yucatán *Cyprinodon* species using

Dsuite (16). We first excluded any samples/species that had an average depth < 4. We then

calculated all possible trios with a species tree based on the most recent phylogeny (17) with

Cualac tessellatus as the outgroup. We calculated Z-scores to assess the significance of D

statistics using 1,000 jackknife blocks and considered a Z-score greater than three as evidence

for significant introgression (18). We only focused our primary analysis on species trios where

P1-P2 included species within the Lake Chichancanab radiation. We used the other trio

calculations to calculate *f*-branch (f_b) statistics to account for correlated f_4-ratio scores and to

inform the timing of introgression events (16).

For species trios that showed evidence of introgression where *C. simus* was either P1or

P2, we then calculated the distance fraction (d_f), a robust metric of introgression for windowed

analyses (19), in non-overlapping windows of 91 informative SNPs. We chose this number of

SNPs because it resulted in windows of approximately 10 kb. We considered windows to be

introgressed if they were in the top X% of absolute d_f value, where X% was the percentage of

estimated genome wide introgression (e.g, f_4-ratio) for that focal group, following (20, 21). To

estimate introgression block sizes, we merged neighboring windows that were introgression

outliers and summed the total size of the adjacent blocks.

1045

1046 *Characterizing candidate genes*

1047   After identifying candidate variants, we then used *Bedtools* (v2.31) to see if variants

1048 occurred near a gene (within 20 kb) or within a gene (22) with a GFF annotation file for the *C.*

1049 *nev. mionectes* reference genome (Tian et al. in prep). We classified variants within 20 kb of a

1050 gene (upstream or downstream) as flanking. For SNPs within a gene, we further classified them

1051 into intronic, synonymous, or nonsynonymous using *SnpEff* (23).

1052   To characterize the origin of candidate variants, we extracted the alleles from every

1053 species for each candidate variant. We classified variants that were found in any other

1054 populations outside of Lake Chichancanab as standing genetic variation. We further subdivided

1055 this group into variants that were found in other species but not in *C. artifrons*. If variants fell

1056 within an introgression window, we classified them as "introgressed." Finally, if the alternative

1057 variant was only detected within Chichancanab species, we classified it as de novo.

1058

1059 *Timing of selective sweeps*

1060   To gain insight into the timing of selective sweeps in *C. simus*, we used *starTMRCA* (24).

Because many genes contained multiple candidate variants, for this analysis, we selected the

variant that was in the median position of each gene. We then extracted a 1-Mb region

surrounding that variant and removed all sites that contained missing data. For simplification, we

only used the generalist/detritivore, *C. beltrani*, as the "reference" population and the

zooplanktivore, *C. simus*, as the "selected" population. We used a fixed recombination rate of

$2\text{x}10^{-8}$ (swordtail fishes; (25)) and a mutation rate of $1.56 \text{ x } 10^{-8}$ substitutions per base pair

(Caribbean *Cyprinodon* species; 26). We then ran five separate Markov chains for 30,000

iterations with a proposal standard deviation of 150 (preliminary analyses changing this

parameter had minimal influence on results). We then discarded the first 10,000 iterations of each

chain as burn-in.

To test for stages of adaptation, we conducted separate generalized linear models to test if

the sweep age was correlated with the proportion of nonsynonymous SNPs, proportion of de

novo SNPs, and the number of SNPs. For the generalized linear regressions of proportion of

nonsynonymous SNPs and proportion of de novo SNPs, we used a quasi-binomial family to

account for overdispersion with a logit link function. For the number of SNPs, we used a quasi-

Poisson family with a log link function. To test for the significance of sweep age, we used a

likelihood ratio test comparing a model with the estimated sweep age as an effect compared to a

null model.

1079

Optomotor response

We tested for differences in visual acuity between the generalist/detritivore, *C. beltrani*,

and the zooplanktivore, *C. simus*, by conducting an optomotor response behavioral assay (26).

We tested lab-reared *C. beltrani* (F1, $n = 7$) and lab-reared *C. simus* (long-term laboratory colony

exceeding ten generations in the lab, $n = 5$). Each fish was tested consecutively with each control

or treatment for 1-minute observation periods, with the control observation occurring first,

followed by presentation of all-black (positive control) and black-and-white spinning bars

(treatment: acuity trial). The order of presentation for positive control and treatment was

alternated in each trial. Fish were placed in a suspended cylindrical plastic container 0.3 m in

diameter that remained stationary during the course of the trials. Spinning black-and-white or all-

black paper was rotated outside the clear plastic container at ~90 RPM during each observation

period.

Due to the data being nonparametric (count, zero-inflated, unequal variance between

groups), we conducted a Wilcoxon rank sum test for the on-banded (all black) and on-banded

portions separately. Because we had an a priori hypothesis that *C. simus* would display a stronger

optomotor response (greater visual acuity), we calculated *p*-values with a one-tailed test.

1096

Sperm morphology

To test for differences in sperm morphology, we collected sperm samples from five

individuals per species in the lab (*C. simus, C. artifrons, C. beltrani*). We first anesthetized the

fish with a solution of MS-222. After the fish was anesthetized, we collected 1μL of milt (fish

semen) with a 1μL glass microcapillary tube and fixed the sample overnight in a 4% PFA

solution stained with Rose Bengal. We then plated sperm on a positively charged slide with a

sealed coverslip. We imaged sperm with oil immersion with a 63x objective and measured 10-30

sperm per individual in ImageJ (27). Due to poor sperm samples from some individuals (not

sufficient quantities), we ended up with the following sample sizes: three *simus*, four *artifrons*,

and five *beltrani*.

To retain information about variation within individuals, we fit separate linear mixed-

effects models for each trait of interest (sperm head, midpiece, and flagellum lengths) with

population as a fixed effect and a random intercept for individual ID, using *LME4* (28). To assess

the significance of the model, we used a Type II ANOVA with a Wald *F* test using Kenward-

Roger degrees of freedom. If there was a significant effect of population, we conducted post hoc

pairwise tests using the *Emmeans* package (29) and corrected for multiple comparisons using the

Tukey method. We did not correct for phylogeny due to how closely related the species are, the

limited sample size, and the fact that the only significant differences were between sister species.

1115

WNT experiment

To confirm the role of the WNT pathway in proper jaw and teeth development, we

conducted an experiment chemically inhibiting the WNT pathway using iCRT14 during

metamorphosis in *C. beltrani*. To control the exact time of development, we first performed in

vitro fertilizations and incubated developing embryos at ~26 until hatching (~8dpf). On the day

of hatching, we haphazardly split broods into control and experimental groups to control for

batch and family effects. We raised experimental groups in tank water with a 100 nanomolar

concentration of iCRT14 and control groups in a 100 nanomolar concentration of DMSO

(solution used to dilute stock iCRT14). We ran the experiment for two weeks until individuals

were 22dpf, changing the solution every 3-4 days. We performed this experiment on four

separate broods (~12 individuals/treatment). One brood was excluded due to improper

development (no ossification of the spine) in the controls.

After the experiment, we euthanized fish with MS-222 and stained fish with alizarin red

(bone) and alcian blue (cartilage) following a modified published protocol (30). After staining,

we prepared whole mounts and imaged fish with fluorescent microscopy (Zeiss LSM880 FCS)

to quantify ossification of the jaw and teeth (31). We then dissected out the lower jaw and

prepared flat mounts to measure teeth morphology. Specifically, we measured the tooth length

and base width for the left and right teeth closest to the mandibular symphysis using ImageJ (27).

For statistical analyses, we took the mean of the left and right tooth length and base width.

We fit separate linear mixed-effects models for tooth length and base width with experimental

treatment as a fixed effect and a random intercept for brood/replicate, using *LME4* (28). For the

number of ossified teeth, we used a generalized linear mixed-effect model with a Poisson family

and log-link function. To assess the significance of these models, we used a likelihood ratio test

comparing it to a null model without treatment as a fixed effect.

1140

1141 **SI Tables**

1142

1143 **Table S1.** Mean genome-wide pairwise F_{ST} statistics within the Chichancanab radiation and with
1144 sister outgroup (*artifrons*).

	labiosus	*beltrani*	*artifrons*
simus	0.090	0.068	0.182
labiosus		0.032	0.184
beltrani			0.168

1145

1146   **Table S2.** F_{ST} statistics within the Chichancanab radiation.

Comparison	$F_{ST} = 1$ (fixed)	$1 > F_{ST} \geq 0.95$ (nearly fixed)	$0.95 > F_{ST} \geq 0.9$	$0.9 > F_{ST} \geq 0.8$	Other	Mean F_{ST}
simus vs *beltrani* + *labiosus*	19	1127	3304	20173	10359605	0.07
labiosus vs *beltrani* + *simus*	0	0	3	94	10384131	0.04
beltrani vs *simus* + *labiosus*	0	0	0	13	10384215	0.02

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160   **Table S3.** Fixed SNPs between *C. simus* and *C. beltrani* + *C. labiosus*. Bold rows indicate SNPs within top 1% of PBS windows.

1161   Bolded genes only are genes that have regions that fall within the top 1% PBS percentile.

Position	Gene	Distance to gene	PBS percentile	Annotation
Chr 1: 5290600	**STK35**	12594	1.85	
Chr 1:5290602	**STK35**	12592	1.85	Fertility, eye development and oxidative stress (32).
Chr 1:5290604	**STK35**	12590	1.85	
Chr 2: 17498913	SLC17A8	Intron	1.24	Glutamate neurotransmitter, inner hair cell dysfunction
Chr 3: **11687002**	**AGO3**	**Intron**	**0.34**	**RNA regulator**
Chr 3: 32286614	RBMS3	14363	22.93	
Chr 3: 32286627	RBMS3	14350	22.93	Involved in chondrogenic craniofacial development of zebrafish (33).
Chr 3: 32286638	RBMS3	14339	22.93	
Chr 3: 32286614	RBMS3	14363	22.93	
Chr 11: 27324192	GRIA3	Intron	2.85	
Chr 11: 27336469	GRIA3	2772	3.18	
Chr 11: 27336472	GRIA3	2775	3.18	Glutamate receptor
Chr 11: 27336476	GRIA3	2779	3.18	
Chr 11: **28255536**	**NA**	**3026**	**0.98**	**NA**
Chr 11: **28255616**	**NA**	**3106**	**0.98**	**NA**
Chr 11: **28256343**	**NA**	**3833**	**0.98**	**NA**
Chr 11: **28257367**	**NA**	**4857**	**0.98**	**NA**
Chr 11: 28260047	NA	7537	2.87	NA
Chr 12: 13854139	**C4orf22**	11903	1.30	Spermatogenesis, cilia, flagellum (34)
Chr 24: **12693158**	**KDM6A**	**Intron**	**0.05**	**Kabuki syndrome (craniofacial development; 35)**

1162

Table S4. Summary statistics of the distance between closest adaptive candidate SNPs within the same

gene

Gene	Median Distance	Mean Distance	SD	
TM6SF2	7430	7430	NA	2
C19orf10	4789	4789	NA	2
KDM6A	2892.5	2892.5	2410.52702	3
KCNB2	1803	1738.75	1614.92299	5
UBE2T	1742	1742	NA	2
TMEM230	1734	3039	3599.52427	4
ABCA1	1037.5	1097.5	959.375665	5
HAPLN4	974	1608.21739	9927.05273	24
GRIN2A	904	1270.6667	4273.38008	4
DPH1	608	1224.5	1999.25934	7
WNT10A	402	1442.17857	3986.39626	29
OPN1LW	386.5	359.5	225.229513	5
NIPSNAP3A	253	1451.71429	2819.19065	8
GALNT15	246.5	675.166667	3156.86746	13
SWS2	246	1381.25	2423.03892	5
STK35	217	793.366667	1992.8543	31
CRIM1	201	47.829268	2176.05381	42
PDYN	187	326.371429	355.925561	36
ADCYAP1R1	186.5	833	1304.35501	11
DNM1	186.5	459.708333	677.261781	25
DNM1	186.5	459.708333	677.261781	25
KCNH8	131.5	584.366667	1030.65532	31
TOP1	124	332.375	6687.63393	17
AGO3	115	357.625	552.097284	9
GMEB2	112	319.490196	2485.05809	52
GPNMB	57	57	77.781746	3
IGFBP4	26	2254	3880.67997	4
FAM83D	23	2523.66667	4347.74658	4
EXOSC3	17	17	NA	2
C4orf22	15	15	9.899495	3
TMEM106B	7	11.4	10.358571	6
THOC3	6	5.833333	3.600926	7
MID1	1	1	NA	2

1166   **SI Figures**

1167

1168   **Figure S1.** μCT scans of Yucatán pupfish with maxilla colored in blue, premaxilla in pink,

1169   dentary in green, and articular in orange.

1170

1171 **Figure S2. Best fitting admixture graph assuming zero admixture events.** Admixture graph

1172 was found using the *findGraphs* function in *ADMIXTOOLS2* (36). Number on edges represent

1173 drift weights.

1174

1175

1176

Figure S3. ADMIXTURE results support genetic differentiation of Yucatán Cyprinodon

pupfish species. (A) Cross-validation error for different values of K clusters, equally support 2,

3, and 4 clusters. (B) *ADMIXTURE* plots showing the clusters at different values of K.

1177

1178

1179

1180

1181

1194

Figure S4. Linkage decay occurs rapidly in *Cyprinodon simus*. Linkage decay was calculated

in windows of 1000 kb using *PLINK* (v1.9). The results graphed are a random sample of all

calculated r^2 values taken for plotting purposes (n = 9,645 r^2 values).

1195

1196

1197

1211

Figure S5. Adaptative SNPs associated with the same gene are generally spaced apart.

Distribution of the distance to nearest adaptive SNP within a gene. Note the log scale of the x-axis.

1215

1216

1217

1218

1219

1220

1221

Figure S6. *f*-branch (*f*$_b$) show that introgression with desert pupfish occurred before the

Chichancanab radiation. Boxes with red color represent significant introgression between P3

species (top) and the species/internal branch on the left. For visualization, non-significant *f*$_b$ were

changed to zero and are completely white boxes. Grey boxes represent comparisons that cannot

be made, as *f*$_b$ cannot be calculated for introgression between sister taxon. Dashed lines on the

left are internal branches. SSI: San Salvador Island, Bahamas; NC: North Carolina, USA.

1228

Figure S7. **Both species did not have a strong response to either control scenario in the**

optomotor response experiment. (A) Number of complete rotations fish swam when the

machine was off and **(B)** when the machine was on but the apparatus was all black**.** Boxplots and

points of the number of complete rotations individuals performed during the "off-banded"

portion of the experiment. Note that some boxplots are compressed to a single line because most

individuals had no response.

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247 **Figure S8**. **Species do not differ in sperm midpiece size or sperm flagellum length.** Boxplots

1248 and points of the mean sperm morphological estimate for an individual.

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1268

1269 **Figure S9. Tooth length was not affected by Wnt inhibition.** Boxplot and jittered points of
1270 average tooth length of the left and right teeth closest to the mandibular symphysis for *C.*
1271 *beltrani* larvae treated with either a control or Wnt inhibitor (iCRT14). Shapes represent different
1272 split-brood replicates.

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284 **Figure S10. Number of ossified teeth was not affected by Wnt inhibition.** Boxplot and jittered
1285 points of number of ossified teeth in *C. beltrani* larvae treated with either a control or Wnt
1286 inhibitor (iCRT14). Shapes represent different split-brood replicates.

SI References:

1. S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

2. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. [Preprint] (2013). Available at: http://arxiv.org/abs/1303.3997.

3. A. McKenna, *et al.*, The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

4. K. Okonechnikov, A. Conesa, F. García-Alcalde, Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292–294 (2016).

5. M. A. DePristo, *et al.*, A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* **43**, 491–498 (2011).

6. G. A. Van der Auwera, *et al.*, From FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit Best Practices Pipeline. *Curr. Protoc. Bioinform.* **43**, 11.10.1-11.10.33 (2013).

7. D. Caetano-Anolles, GenotypeGVCFs and the death of the dot (obsolete as of GATK 4.6.0.0). *GATK* (2024). Available at: https://gatk.broadinstitute.org/hc/en-us/articles/6012243429531-GenotypeGVCFs-and-the-death-of-the-dot-obsolete-as-of-GATK-4-6-0-0.

8. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008 (2021).

9. S. Purcell, *et al.*, PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Amer. J. Hum. Genet.* **81**, 559–575 (2007).

10. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

11. S. Schiffels, K. Wang, "MSMC and MSMC2: The Multiple Sequentially Markovian Coalescent" in *Statistical Population Genomics*, J. Y. Dutheil, Ed. (Springer US, 2020), pp. 147–166.

12. E. J. Richards, *et al.*, A vertebrate adaptive radiation is assembled from an ancient and disjunct spatiotemporal landscape. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2011811118 (2021).

13. P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

1329   14.   J. M. Schmidt, M. de Manuel, T. Marques-Bonet, S. Castellano, A. M. Andrés, The impact
1330   of genetic adaptation on chimpanzee subspecies differentiation. *PLOS Genet.* **15**,
1331   e1008485 (2019).

1332   15.   M. Shpak, K. N. Lawrence, J. E. Pool, The precision and power of population branch
1333   statistics in identifying the genomic signatures of local adaptation. *Genome Biol. Evol.* **17**,
1334   evaf080 (2025).

1335   16.   M. Malinsky, M. Matschiner, H. Svardal, Dsuite - Fast D-statistics and related admixture
1336   evidence from VCF files. *Mol. Ecol. Res.* **21**, 584–595 (2021).

1337   17.   S. G. Hernández-Ávila, C. W. Hoagstrom, W. A. Matamoros, Historical biogeography of
1338   North American killifishes (Cyprinodontiformes) recapitulates geographical history in the
1339   Gulf of México watershed. *Zool J. Linn. Soc.* **202**, zlae105 (2024).

1340   18.   E. Y. Durand, N. Patterson, D. Reich, M. Slatkin, Testing for ancient admixture between
1341   closely related populations. *Mol. Biol. Evol.* **28**, 2239–2252 (2011).

1342   19.   B. Pfeifer, D. D. Kapan, Estimates of introgression as a function of pairwise distances.
1343   *BMC Bioinform.* **20**, 207 (2019).

1344   20.   C. Feng, J. Wang, A. Liston, M. Kang, Recombination variation shapes phylogeny and
1345   introgression in wild diploid strawberries. *Mol. Biol. Evol.* **40**, msad049 (2023).

1346   21.   A. Morales-Cruz, *et al.*, Introgression among North American wild grapes (*Vitis*) fuels
1347   biotic and abiotic adaptation. *Genome Biol.* **22**, 254 (2021).

1348   22.   A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
1349   features. *Bioinformatics* **26**, 841–842 (2010).

1350   23.   P. Cingolani, *et al.*, A program for annotating and predicting the effects of single
1351   nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster*
1352   strain w1118; iso-2; iso-3. *Fly* **6**, 80–92 (2012).

1353   24.   J. Smith, G. Coop, M. Stephens, J. Novembre, Estimating time to the common ancestor for
1354   a beneficial allele. *Mol. Biol. Evol.* **35**, 1003–1017 (2018).

1355   25.   M. Schumer, *et al.*, Natural selection interacts with recombination to shape the evolution
1356   of hybrid genomes. *Science* **360**, 656–660 (2018).

1357   26.   S. C. F. Neuhauss, *et al.*, Genetic disorders of vision revealed by a behavioral screen of
1358   400 essential loci in zebrafish. *J. Neurosci.* **19**, 8603–8615 (1999).

1359   27.   C. T. Rueden, *et al.*, ImageJ2: ImageJ for the next generation of scientific image data.
1360   *BMC Bioinform.* **18**, 529 (2017).

1361   28.   D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models using
1362   lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

29.   R. V. Lenth, *emmeans: Estimated Marginal Means, aka Least-Squares Means* (2021).

30.   M. Walker, C. Kimmel, A two-color acid-free cartilage and bone stain for zebrafish larvae.
Biotech. Histochem. **82**, 23–28 (2007).

31.   R. Koita, S. Oikawa, T. Tani, M. Matsuda, A. Kawamura, Live visualization of calcified
bones in zebrafish and medaka larvae and juveniles using calcein and alizarin red S. *Bio-*
protocol **14** (2024).

32.   Y. Miyamoto, *et al.*, The STK35 locus contributes to normal gametogenesis and encodes a
lncRNA responsive to oxidative stress. *Biol. Open* **7**, bio032631 (2018).

33.   C.S. Jayasena, M.E. Bronner, Rbms3 functions in craniofacial development by
posttranscriptionally modulating TGF-β signaling. *J. Cell. Biol.* **199**(2012)

34.   H. Li, Y. Dai, Z. Luo, D. Nie, Cloning of a new testis-enriched gene C4orf22 and its role
in cell cycle and apoptosis in mouse spermatogenic cells. *Mol. Biol. Rep.* **46**, 2029–2038
(2019).

35.   P.M. Van Laarhoven, Kabuki syndrome genes KMT2D and KDM6A: functional analyses
demonstrate critical roles in craniofacial, heart and brain development. *Hum. Mol.*
Genet. **24**, (2015).

36.   R. Maier, *et al.*, On the limits of fitting complex models of population history to f-
statistics. *eLife* **12**, e85492 (2023).

1381

1382