Binary vector origin predictably determines *Agrobacterium*-mediated transformation outcome across eukaryotic kingdoms

Authors: Matthew J. Szarzanowicz[1,2,3], Michael Busche[4], Ziyu Dai[1,5], Nathan Lapp,[1,2] Gina M. Geiselman[6], Jiayuan Jia[1,5], Lucas M. Waldburger[1,2,7,8,9], Yan Liang[1,9], Guoliang Yuan[5], Rylan Duong[5], Kyle R. Pomraning[5], Nathan J. Hillson[1,9], Jacob O. Brunkard[4], Blake A. Simmons[1,9], John M. Gladden[1,6], Joonhoon Kim[1,5], Mitchell G. Thompson[1,2,*], Patrick M. Shih[1,2,3,*]

Affiliations:

[1]Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.

[2]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA

[4]Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI, USA

[5]Energy and Environment Directorate, Pacific Northwest National Laboratory, 902 Battelle Boulevard, Richland, WA 99354, USA

[6]Sandia National Laboratories, Livermore, CA, USA.

[7]Department of Electrical Engineering and Computer Science, University of California, Berkeley, CA 94720, USA

[8]Department of Bioengineering, University of California, Berkeley, California, USA

[9]Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

*Correspondence should be addressed to either Mitchell G. Thompson (mthompson@lbl.gov) or Patrick M. Shih (pmshih@berkeley.edu)

Abstract

Agrobacterium-mediated transformation (AMT) is the primary means of genetic engineering in plants and many fungi, but the factors that control transformation outcomes—efficiency, transgene insertion number, and transgene integrity—remain poorly characterized. Although transformation outcomes dictate an event's potential utility in both industrial and academic contexts, AMT remains largely unoptimized for these metrics. Here, we systematically analyze the impact of the transgene-harboring binary vector on transformation outcomes across plant and fungal species. Through a comparison of different plasmid origin of replication (ORI) families and engineered copy number variants, our results reveal that the ORI family—not plasmid copy number—dictates T-DNA insertion number, backbone inclusion, and transformation efficiency, while plasmid copy number tuning alters efficiency without changing ORI family-specific

signatures. Independent of plasmid copy number across kingdoms, the most widely used pVS1
ORI-based vectors (e.g. pCambia) result in significantly more insertions per transformant and high
levels of transgene silencing compared to the less-utilized pSa ORI family, which enriches for
more uniform single insertion events. Furthermore, we demonstrate that ORI-dependent
transformation outcomes in yeast predictably reflect those in *Arabidopsis*. Together, these results
lay the foundation for future binary vector design aimed at achieving more predictable,
controllable, and optimized transformation outcomes across diverse eukaryotic hosts.

Introduction
Agrobacterium-mediated transformation (AMT) is the predominant method for introducing
transgenes into plants and fungi across academia and industry. Despite its utility, AMT remains
an imprecise tool, producing transformants with unpredictable insertion number,
concatemerization, and vector backbone inclusion—collectively termed event outcome. These
outcomes directly determine a transformant's suitability for specific engineering goals, and due to
the wide range of systems that utilize AMT, ideal event outcomes can vary substantially. For
example, single-copy, backbone-free insertions are preferred in conventional crop engineering
for regulatory compliance and transgene stability, whereas multi-copy or concatemerized
insertions are desirable in metabolic or gene engineering applications to increase enzyme
abundance or maximize delivery of homology-directed repair templates. Such diversity of target
outcomes makes controllable transformation highly coveted yet largely unachievable with current
legacy tools. Thus, unlocking the full potential of AMT will require developing and characterizing
components that enable predictable, tunable event outcomes across diverse plant and fungal
hosts.

The primary chassis of AMT is derived from the plant pathogen *Agrobacterium tumefaciens* which
has a natural capacity to horizontally transfer DNA to plants. A variety of strains and transgene-
harboring binary vectors were developed in the 1980s and 1990s, and they continue to form the
core toolsets used for AMT today, often fully unchanged from their original design[1]. While AMT
generally produces cleaner, lower-copy insertions than sister techniques—e.g., biolistics—
reported rates of transgene insertion and integrity vary widely within the literature, making it
difficult to determine how each legacy component affects transformation outcome. The sources
of variance contributing to these disparate results are numerous and could include host genetic
background, host tissue source, *Agrobacterium* strain, transgene architecture, transformation
protocol, and the binary vector itself[2–5].

As the carrier of the transgene, the binary vector plays a major role in influencing transformation
outcome. Being a fully synthetic component, each binary vector must contain a user-selected ORI
to enable proper replication within *Agrobacterium*. Different ORIs have variable copy number
maintenance, stability and partitioning characteristics, potentially changing how the transgene is
maintained and processed within the bacterium. Although multiple studies have reported that the
binary vector ORI influences transformation outcomes, their conclusions vary—particularly
regarding the role of high-copy origins[5–7]. These disparities can likely be attributed to some

combination of differences in the plant genetic background, bacterial strain, use of standard vs.
super binary vectors, transformation protocol and T-DNA payload. As a result, direct comparisons
from the literature are often convoluted for plant transformation due to variations in experimental
design.

To systematically assess the impact of binary vector copy number on AMT, we previously
constructed a library of 71 copy number variants across four ORIs, each differing by single SNPs
in the *repA* gene[8]. These variants were built into otherwise identical binary vectors and evaluated
for their influence on transient and stable transformation, demonstrating that both could be
significantly improved with single SNPs. Our findings revealed that copy number alone could not
explain the observed transformation changes—highlighting that each ORI likely has intrinsic,
origin-specific properties that influence transformation. However, a key question remains
regarding how plasmid copy number, transformation efficiency, and event outcome are
interrelated—factors that collectively define the functional utility of any engineered variant.

To dissect this relationship in a controlled manner across eukaryotic kingdoms, we pursued three
primary objectives. First, we characterize the relationship between ORI family, variant-specific
copy number, transgene integration number, transgene integrity and backbone inclusion rates
using a controlled experimental design in both the model plant *Arabidopsis thaliana* and
oleaginous yeast *Rhodosporidium toruloides*, along with validation in the filamentous fungus
Aspergillus niger. This establishes a foundation for understanding how ORI identity and copy
number jointly influence transformation efficiency and outcome. Second, we assess whether GFP
expression can serve as a reliable proxy for transgene insertion number in stable transformants,
providing a scalable alternative to labor-intensive methods for quantifying insertions. Finally, we
test whether transformation outcomes in *R. toruloides* can predict corresponding results in *A.*
thaliana, thereby evaluating the potential of using a high-throughput fungal system to predict
transformation outcomes in a plant. Our results reveal a key means to improve AMT efficiency
while controlling for transformation outcomes across eukaryotic kingdoms. Additionally, we
demonstrate a method for high-throughput predictive evaluation of binary vector performance,
laying the groundwork for the systematic exploration and engineering of ORI sequence space.

Results

ORI family and copy number influence transformation efficiency and GFP expression
107
As the transgene-harboring plasmid, the copy number of the binary vector within *A. tumefaciens*
and the ORI used to control plasmid replication have both been shown to alter stable and transient
expression efficiencies[8]. To determine how both the ORI family and plasmid copy number
influence transformation outcomes, we selected two ORI families for analysis: pVS1 and pSa.
The pVS1 ORI was selected due to its widespread use in the plant community through vectors
such as the pCambia series along with its generally high transformation efficiency while pSa was
chosen for its low efficiency WT form that had the largest improvement from copy tuning in our
previous study[8]. For these families, three variants spanning a low, medium, and high copy number
were generated using pVS1 (2, 10, and 29 copies) and pSa (5, 18, and 31 copies) backbones,
varying within a family by a single SNP. These vectors were otherwise identical in architecture,

with each containing a selectable marker (kanamycin for *A. thaliana*, nourseothricin for *R.*
toruloides) near the left border (LB), and a constitutively expressed GFP cassette near the right
border (RB). To control for sequence context effects near the borders, randomized 100 bp
buffering sequences were added between functional elements and border sequences and
standardized across all constructs. These vectors were then introduced into standard *A.*
tumefaciens strains for their respective host (GV3101::pMP90 for *A. thaliana* and EHA105 for *R.*
toruloides) and used to obtain ~96 independent transformants per ORI variant, yielding 559
transformants for *A. thaliana* and 576 for *R. toruloides*. These transformants were then used to
analyze the relationship between ORI family, vector copy number, population GFP expression,
and transgene insertion number following the schematic outlined in **Fig. 1**.

128
Fig. 1: Pipeline analyzing ORI impact on AMT outcomes. **(a)** Six binary vectors were built per organism,
consisting of a selectable marker flanking the LB and a constitutive GFP cassette flanking the RB. These binary vectors
are maintained in *A. tumefaciens* by one of two ORIs—pVS1 and pSa—which vary in copy number due to single SNPs
in the *repA* ORF. **(b)** Strains of *A. tumefaciens* harboring these vectors were used to transform *R. toruloides* and *A.*
thaliana, with ~96 transformants collected per ORI variant. **(c)** These transformants were then analyzed for GFP
expression, transgene copy number, and vector backbone inclusion. **(d)** Relationships between GFP expression,
transgene copy number, backbone inclusion, and ORI variant were then analyzed, **(e)** and these were used to compare
outcome between organisms to evaluate potential predictive capacity between a high-throughput yeast system and a
low-throughput plant.

138
As all transformants for a given species were derived from identical T-DNA sequences harboring
a constitutive GFP cassette, variability in GFP expression is likely the product of genomic insertion
location, transgene insertion number, transgene silencing, and potentially T-DNA degradation or
damage. As AMT-mediated insertions have been well-characterized to occur randomly
throughout the genome[9,10], GFP expression levels across a population would be expected to not
differ significantly between binary vector ORI variants if this factor does not influence
transformation outcome. Similarly, if differences are observed between ORI variants, it is likely
that such variance is due to the impact of the ORI on either transgene insertion number or T-DNA
degradation as insertion location and silencing are host-mediated responses. To evaluate
potential differences in gene expression, GFP output was measured in both systems using a plate
reader assay. For *A. thaliana*, four leaves per transformant were selected, and a single leaf disk
per leaf was hole-punched and placed into a 96-well plate for quantification. The disk values were
then averaged to produce a final expression value to better account for leaf to leaf variability in
expression (**Fig. S1a**). For *R. toruloides*, liquid cultures were grown in a 96-well block overnight

to an OD_{600} between 0.1 and 1.0, and an aliquot of this culture was transferred to a 96-well plate
for GFP quantification normalized to OD.
155
The two selected ORI families vary in their WT copy numbers, with pSa maintaining the binary
vector at approximately five copies per cell and pVS1 at ten copies[8]. Through the use of single
repA SNPs, this number can be manipulated (**Fig. 2a**). Even when comparing variants between
ORI families with relatively similar copy numbers, transformation efficiencies tended to vary
significantly. This is particularly true for *R. toruloides* (**Fig. 2b**) which typically has more consistent
transformation efficiencies between replicates than the highly variable floral dipping of *A. thaliana*
(**Fig. 2c**). Transformation efficiencies from *A. thaliana* floral dipping have been well documented
to vary by an order of magnitude or more between batches, complicating direct comparisons
across variables[11]. For this reason, we used a large sample size of seeds (~38,000-80,000 seeds
per variant) derived from 27-39 independent plants. Due to this high degree of variability across
floral dips, a generalized linear model (GLM) was used to estimate transformation efficiencies,
accounting for both the seed number and transformation efficiency (**Fig. 2c**). Results from both
R. toruloides and *A. thaliana* show that the ORI family and sub-variants influence transformation
efficiency, corroborating a trend previously seen in[8]. ORI variants with similar estimated copy
numbers from different ORI families—e.g. pSa V152F (31 copies) and pVS1 R106H (29 copies)—
tended to produce significantly different transformation efficiencies from one another, particularly
in *R. toruloides* where this pairing differed by an average efficiency of 16.5-fold across replicates.
This provides further evidence that intrinsic ORI family properties are a major source of variance
in AMT outcomes, extending beyond copy number alone.

Fig. 2. Transformation efficiency and GFP expression vary by ORI family. (a) The six ORI variants varied in copy number using *repA* SNPs as characterized in [8]. Both ORI families and some variants had significant differences in transformation efficiency as measured by total colony number for *R. toruloides* **(b)** and percent transformed seed for *A. thaliana* **(c)**. Significant differences were determined by a two-tailed ANOVA followed by a Tukey's post hoc test for *R. toruloides* (p < 0.05 for significance letters) and a binomial GLM with estimated marginal means and Wald 95% CIs for *A. thaliana*. Transformants from the same transformation varied dramatically in GFP expression **(d)**, and this variance was found to be primarily ORI family—not vector copy number—dependent for both *R. toruloides* **(e)** and *A. thaliana* **(f)**. Population level expression differences were determined by a two-tailed ANOVA followed by a Tukey's HSD test (p < 0.05 for significance letters). When rank-ordering the samples by GFP expression, it was observed that the pVS1 ORI family had significantly more transformants above the pSa-population maximum (19.4%, p = 7.16 * 10^{-16}) for *R. toruloides* **(g)** but significantly fewer than the lower boundary (2nd percentile) of the continuous pSa family range for *A. thaliana* **(h)** (20.7%, p = 4.03 * 10^{-13}) as measured by a hypergeometric test.

When analyzing population-level GFP outputs, ORI-dependent differences became even more
pronounced. Across both systems, GFP expression distributions were highly consistent within
each ORI family, despite differences observed on the copy number and transformation efficiency
levels. Within the *R. toruloides* distributions, transformants derived from any pSa variant exhibited
lower average and less variable GFP expression than pVS1-derived transformants. Contrastingly,
pVS1-derived samples displayed a broader range of expression, with a long upper tail extending
toward highly and visually-distinct GFP expressing colonies (**Fig. 2e and 2d**). The results in *A.*
thaliana were intriguingly the opposite: pVS1-derived lines were found to have a long lower tail
comprising transformants with little to effectively no GFP expression (**Fig. 2f**). When rank-ordering
all samples by GFP output, 19.4% (56/288) of *R. toruloides* pVS1-derived transformants
exceeded the maximum pSa expression value across all pSa variants (n = 288, **Fig. 2g**).
Conversely, 20.7% (56/271) of pVS1-derived lines in *A. thaliana* were found to fall below the
general pSa population's minimum, with only 1.9% (5/264) of pSa outliers doing the same (**Fig.**
2h). In both species, around 20% of the pVS1-derived samples notably deviated beyond the
extremes of the pSa population. This species-dependent expression variability may reflect gene
silencing in *A. thaliana* that is triggered upon hyper-expression of a transgene[12]—a process that
may be absent or attenuated in some fungi compared to plant systems.
205
To further explore this observation, a filamentous fungal system, *Aspergillus niger*, was
transformed with the same six ORI variants, and 96 transformants were evaluated per vector.
These parallel experiments reproduced the family-level patterns with pVS1 variants yielding
markedly higher transformation efficiencies and a broader, right-shifted GFP distribution
compared to pSa (**Fig. S2, b–c**). Thus, across all systems, the ORI family was observed to exhibit
significant impacts on both transformation efficiency and transgene expression, and notably,
these expression trends remained even when controlling for plasmid copy number across ORI
families.
214
215   **Transgene insertion number is ORI family-dependent and influences GFP expression**
216
The differences observed in GFP expression between ORI families are likely due to different
transgene insertion numbers across the population. To determine if the ORI family and copy-
tuned variants altered transformation outcome, gDNA was extracted from all samples and
prepared for transgene insertion quantification via dPCR. For each transformant, two probes
targeting the GFP cassette and a genomic target were used within the same reaction, and the
transgene insertion number was estimated from the ratio of GFP to genomic copies, adjusting for
ploidy. dPCR enables absolute quantification of starting template copies, offering greater
accuracy and precision than qPCR and higher-throughput multiplexing compared to Southern
blotting, making it well-suited for large-scale transgene studies[13].
226
Similar to the GFP data, notable ORI-dependent differences in insertion number distributions were
observed for both organisms (**Fig. 3a and 3b**). All pSa ORI variants produced significantly lower
average insertion numbers and higher rates of single insertions than pVS1 binary vectors across
systems (63% single insertions for combined pSa samples in *R. toruloides* vs 40% for pVS1, and
36% vs 13% in *A. thaliana*). A notable difference in the total number of insertions was observed

between organisms, with 95% of transformants having fewer than six insertions in *R. toruloides*
compared to 35 copies in *A. thaliana* (**Fig. 3c** and **Fig. 3d**). Of particular note, all pVS1 vectors in
A. thaliana produced high-insertion transformants that reached an estimated 140 insertions,
coupled with low rates of single-insertion events. Such high insertion numbers are likely
attributable to concatemerization of T-DNAs rather than exclusive insertion into independent loci.
Transformation of germline tissues, such as floral dipping, is known to produce higher rates of T-
DNA concatemerization compared to somatic transformation, and large concatemerized T-DNA
insertions have been previously reported in floral dipped *A. thaliana*[14–16].
240

241
242
243

Fig. 3: Transgene insertion number and GFP expression vary by ORI family and organism.
Transgene insertion numbers were estimated with dPCR and grouped by 0, 1, 2-4, and 5+ insertions for *R. toruloides* **(a)** and *A. thaliana* **(b)**. Population percentage averages for each species were calculated from the estimated marginal means of a binomial GLM along with Wald 95% CIs, depicted as error bars. The ORI family significantly altered the rate of transgene insertions, with pSa producing both lower average insertion numbers and higher rates of single insertions than pVS1 for *R. toruloides* **(c)** ($p < 3.75 * 10^{-14}$) and *A. thaliana* **(d)** ($p < 2.2 * 10^{-16}$) as computed by a Wilcoxon test. GFP followed a linear gene dosage regression in *R. toruloides* **(e)** ($R^2 = 0.82$, $p < 2.2 * 10^{-16}$ from the F statistic) that fit both ORI families. **(f)** In *A. thaliana*, the pVS1-derived population had a substantial proportion of low-expressing high-insertion transformants, reducing the variance explained by insertion number to effectively zero (adjusted $R^2 = 0.03$, $p = 2.04 * 10^{-3}$) while the pSa-derived population retained some of the linear relationship ($R^2 = 0.42$, $p < 2.2*10^{-16}$). When looking only at transformants with fewer than 8 insertions in *A. thaliana*, the pSa **(g)** and pVS1 **(h)** populations differed significantly in the linear relationship between GFP and insertion number despite being transformed with the same T-DNA sequence (pSa: adjusted $R^2 = 0.67$, $p < 2.2 * 10^{-16}$ and pVS1: adjusted $R^2 = 0.21$, $p = 3.61*10^{-8}$).

Such high insertion numbers were not observed in *R. toruloides* for either ORI, nor for the pSa variants in *A. thaliana*, making this an exclusive feature of pVS1 vectors in *A. thaliana*, regardless of plasmid copy number. Moreover, as the pVS1 ORI family is broadly used in plant transformation within vector series such as pCambia and pPZP, this finding is notable and cautionary for plant transformation. As further validation of higher insertion rates for pVS1, we quantified GFP transgene insertion number in *A. niger* transformants derived from WT pSa (n = 93) and pVS1 vectors (n = 92). Consistent with the other systems, pVS1-derived transformants had significantly higher average insertion numbers than pSa (**Fig. S2 d–e**, $p < 0.001$), further corroborating this ORI-family level trend.

Within *A. thaliana* and *R. toruloides*, all pVS1 copy number variants produced single insertions at non-significantly different rates (**Fig. S3**, $p > 0.05$). A similar lack of significant intra-family differences was observed among pSa variants in both systems, although in *A. thaliana* the high-copy pSa V152F variant trended toward a reduction in single-insertion rate when compared to pSa WT (**Fig. S3**, $p = 0.069$). Taken together, these results demonstrate that plasmid copy number, transformation efficiency, and single-insertion generation rates are not inextricably linked, suggesting possible future engineering to improve efficiency without compromising single-insertion event generation.

To evaluate the relationship between gene dosage and expression, estimated GFP insertion numbers were plotted against GFP expression for each transformant. For *R. toruloides*, a strong linear relationship was observed between insertion number and expression, indicating a predictable gene dosage effect. A Cook's D-adjusted linear regression yielded an R² of 0.82, indicating that insertion number accounts for the majority of expression variance, with minimal influence from positional effects or silencing (**Fig. 3e, Fig. S4).** Both pSa- and pVS1-derived transformants aligned along this trendline, suggesting minimal influence of ORI family on transgene expression outcomes within *R. toruloides*.

In contrast, *A. thaliana* exhibited a far more complex pattern with pronounced ORI-dependent effects on transgene expression. Transformants with insertion numbers exceeding 16 almost exclusively exhibited low but detectable GFP expression, likely indicating transgene silencing which is a well documented phenomenon in plants. Across all insertion number ranges, there was

at least a small proportion of individuals with similarly low levels of expression caused either by
silencing, transgene degradation, or positional effects within the genome (e.g. heterochromatic
insertion). Despite the >200-fold variance in expression levels, a general relationship between
gene dosage and expression could be defined at lower insertion numbers, declining after seven
insertions as silencing became more prevalent. Among transformants with seven or fewer
insertions, a weaker gene dosage relationship was observed (R^2 = 0.43) relative to *R. toruloides*,
demonstrating greater variability attributable to positional effects or silencing in the plant system.
Intriguingly, splitting the data by ORI family increased the predictive capacity dramatically for pSa
(R^2 = 0.67, n = 243) but decreased it for pVS1 (R^2 = 0.21, n = 132) (**Fig. 3g and 3h**). This
difference can largely be attributed to a greater number of low expressing—likely silenced—
transformants derived from pVS1 binary vectors. As both the pSa and pVS1 vectors carried
identical T-DNA sequences across all variants, these differences are ORI-dependent,
demonstrating that the binary vector ORI can profoundly influence not only transgene insertion
rate but also downstream expression levels within *A. thaliana*. Why identical T-DNA sequences
launched from different binary vector backbones yield variable expression outcomes for a given
insertion number remains unknown; however, one could speculate that pVS1-launched T-DNA
undergoes greater rates of concatemerization or degradation compared to pSa, leading to
changes in host silencing or transgene integrity.

Taken as a whole, these results show that the binary vector ORI family—not the vector copy
number—profoundly impacts transgene insertion rates across systems. Within *A. thaliana*, the
ORI family exhibits an additional effect of influencing transgene expression for a given insertion
number, making the selection of the binary vector ORI important not just for controlling insertion
rates but also transgene expression quality.

ORI family influences backbone insertion but not T-DNA integrity in *A. thaliana*

The impact of the ORI family on transgene expression suggests this feature of the binary vector
may influence transgene mobilization or integrity. To further investigate how the ORI family
changes transformation outcome, we evaluated T-DNA integrity and backbone insertion
frequencies within the *A. thaliana* transformant population (**Fig. 4a**). T-DNA degradation has been
widely reported within the literature and results in truncated insertions that do not span the entire
RB-LB sequence[17,18]. As mobilized T-DNAs are covalently bound to VirD2 at their 5' RB end,
there is a general assumption within the community that LB degradation is more common due to
susceptibility to exonuclease activity[3]. As such, it is common practice to place the selectable
marker near the LB to help purge LB-degraded transformants during selection.

We investigated LB and RB border degradation by quantifying insertion numbers of the
kanamycin resistance (Kan) and GFP cassettes, positioned near the LB and RB, respectively.
Using the same DNA templates as the GFP analysis, we performed additional dPCR assays to
determine Kan insertion number, with probes targeting Kan and a genomic reference. Each probe
provides an absolute template count with an associated 95% confidence interval based on
Poisson partitioning. Thus, for each sample we obtained four independent error estimates (GFP,
Kan, and two genomic controls). To reduce false positives, we only considered transformants
where both targets had 95% CIs confined within a single integer insertion number. After filtering,

215 transformants were evaluable for pSa and 134 for pVS1, with no event exceeding 12
insertions of either cassette due to the higher quantification error associated with elevated
insertion number.

When comparing the insertion number of Kan to GFP within this high confidence population, we
found significantly higher rates of Kan:GFP insertion number discordance within pVS1-derived
transformants than pSa-derived transformants (odds ratio 2.52x, p = 0.00045), suggesting higher
rates of T-DNA degradation within the pVS1 population (**Fig. 4b**). When considering only events
with single insertions of Kan, discordance was similar between the two ORI families (n.s., p >
0.05), implicating the higher average insertion rate of all pVS1-variants to the higher discordance
on the population level. As this analysis only measures the ratio of the two genes, the rates of
detected RB or LB degradation are likely underestimated due to masking in multi-insertion
transformants with equal numbers of LB and RB degraded T-DNAs.

For each transformant with an unequal Kan:GFP insertion ratio, an estimate of LB or RB
degradation can be made based on the surplus of Kan or GFP, indicating RB and LB degradations
or truncations, respectively. To test the LB-degradation hypothesis from exonuclease activity, we
analyzed the rate of LB degradation using a generalized linear model across ORI variants. This
analysis found no enrichment of LB degradation for any ORI variant (p > 0.05), suggesting
incomplete T-DNA insertions occur at similar rates from both the RB and LB borders and that ORI
families and sub-variants do not significantly influence this property (**Fig. 4c**). These unanticipated
results challenge the notion of greater LB degradation and suggest the common design practice
of localizing the selectable marker to the LB deserves reconsideration.

In addition to T-DNA degradation, numerous reports have characterized the inclusion of the binary
vector backbone (BB) in transformants. This phenomenon is most commonly observed as a LB
readthrough, in which termination at the LB sequence does not properly occur[19]. Within the
literature, backbone inclusion rates vary widely, typically ranging between 20-80% of
transformants[7,20–24]. The wide variability in this reported range likely reflects variation in
transformed species, *Agrobacterium* strain, binary vector architecture, transformation protocol
and sampling size, making it difficult to identify significant sources of variance related to backbone
mobilization.

To analyze the impact of the binary vector ORI on backbone inclusion, we quantified inclusion
rates for all *A. thaliana* transformants using qPCR for a backbone target ~300bp from the LB
along with a gDNA internal control. Transformants cleanly separated into backbone-positive and
negative groups when comparing the Cq values for the backbone and gDNA targets (**Fig. S5a**).
When comparing backbone inclusion and GFP insertion number, both ORI families showed
significant enrichment of transgene insertions in backbone positive lines, suggesting the
probability of backbone inclusion increases in multi-insertion lines (**Fig. 4d**, p < 0.001 for both pSa
and pVS1). Despite the differences in average insertion numbers between ORI families, however,
backbone inclusion rates for all ORI variants did not significantly differ and ranged from 58-79%
(**Fig. S5b,** p > 0.05 with Holm-corrected GLM). As a transformant needs only one backbone
insertion to be classified as backbone-positive, it is reasonable to hypothesize that transformants
with higher total insertion numbers (due to more independent T-DNA insertions or

concatemerization) also have a higher chance of having at least one included backbone. To
investigate this, we implemented a generalized linear model incorporating backbone inclusion,
ORI family, and measured GFP insertions to analyze the probability of backbone inclusion as a
function of GFP insertion number. This analysis found that the pSa ORI family had a significantly
higher chance of backbone inclusion per copy of GFP than pVS1 (**Fig. 4e**, odds ratio: 6.13x, p <
0.001). As pSa variants tend to produce transformants with notably lower insertion numbers than
the pVS1 family, this difference is masked on the population level as abundant high-insertion
pVS1-derived transformants tend to have backbone.

When comparing backbone inclusion rate probabilities for each ORI variant at different GFP
insertion numbers, variants from each ORI family clustered tightly and were not significantly
different (**Fig. 4f**). This suggests that the differences observed in backbone inclusion are an ORI
family-dependent property and are not significantly impacted by manipulations in plasmid copy
number that affect other properties such as transformation efficiency. Why certain ORI families
influence the rate of backbone inclusion on otherwise identical binary vectors is unknown. There
is literature precedence of certain ORIs producing lower rates of backbone inclusion such as the
pRi ORI compared to RK2[7], but to date, there is no validated mechanistic explanation to describe
this phenomenon. This study provides strong evidence that such differences are derived from
intrinsic properties of the ORI itself in a manner that is independent of binary vector copy number.

Fig. 4: Evaluation of T-DNA integrity and backbone inclusion in *A. thaliana*. a) Schematic of the T-DNA integrity and backbone inclusion analysis. T-DNA integrity was assessed by dPCR of the LB-proximal *Kan* and RB-proximal *GFP* cassettes, each normalized to a gDNA control. Backbone inclusion was quantified by qPCR of a locus ~300 bp from the LB and normalized to the same gDNA control. For integrity analyses, only samples with 95% CI ranges from the dPCR Poisson quantification confined to a single integer copy for both *Kan* and *GFP* were retained. **b)** Incidence of discordant *Kan:GFP* ratios was modeled with a binomial GLM. The predicted probabilities of discordance from the estimated marginal mean of the GLM + 95% Wald CI are shown. Across the full population, pVS1 transformants (n = 134) showed higher discordance than pSa (n = 215; odds ratio = 2.52, *p* = 0.00045). In single-insertion lines, rates did not differ (pSa n = 101, pVS1 n = 30; *p* > 0.05). **c)** Border truncations were inferred from *Kan:GFP* imbalance and analyzed by ORI variant using a binomial GLM. Predicted probabilities from emmeans were compared against the null expectation of equal LB and RB degradation (50%) via Wald tests. No variant (pSa n = 70–73, pVS1 n = 34–52) deviated significantly (*p* > 0.05). **d)** Backbone inclusion vs. log2 GFP insertion number compared by a Wilcoxon rank sum test for pSa (n = 243, p = 2.35 * 10^{-12}) and pVS1 (n = 267, p = 1.37 * 10^{-23}). **e)** A GLM including ORI family, GFP insertion number, and backbone inclusion was used to test family-level effects. Wald tests of emmeans contrasts revealed that pSa transformants (n = 243) had a significantly greater odds of backbone inclusion per GFP insertion than pVS1 (n = 267; odds ratio = 6.13, *p* = 2.15 * 10^{-12}). **f)** Predicted probabilities of backbone inclusion at defined GFP insertion numbers were estimated for each ORI variant using emmeans from the GLM; error bars represent 95% Wald CIs.

GFP expression is a functional predictor of transgene insertion number

All core analyses in this study relied upon large-population transformation, gDNA extraction, and dPCR-based quantification. This pipeline is highly labor intensive, costly, and not scalable to screening thousands of ORI variant candidates, even within most industrial settings. To address these limitations in throughput, we investigated whether GFP expression could serve as a reliable proxy for predicting the number of inserted transgenes. Given the broad and irregular distribution of expression levels observed in the *A. thaliana* dataset, we implemented an ensemble of machine learning models to better capture nonlinear trends and improve performance across a variably expressing population. Models were trained using either GFP expression alone or GFP combined with ORI identity to assess the added predictive value of the ORI family.

To enhance resolution, transformants were grouped into three biologically relevant categories based on transgene insertion number: single insertions, 2–4 insertions, and five or more insertions. A similar strategy was applied to *R. toruloides*, with the addition of a zero-copy group to account for transformants with no detectable GFP insertions despite viable growth on nourseothricin.

For both species, we initially evaluated a diverse set of individual classification models (**Fig. S6**). From these, the top three performers—each utilizing distinct classification strategies—were selected to form a majority-vote ensemble in which predictions for unknown samples are determined by consensus. In *A. thaliana*, the selected models were Support Vector Classification (SVC), Naive Bayes (NB), and Quadratic Discriminant Analysis. For *R. toruloides*, the ensemble consisted of SVC, NB, and a multinomial logistic regression model.

Each model was evaluated under both GFP-only and GFP+ORI conditions using a stratified 70:30 training-to-testing split, repeated over 100 bootstrap iterations to estimate average model performance. Classification outcomes are summarized in the confusion matrices shown in **Fig. 5b-c and S7**. These matrices provide the basis for calculating precision and recall metrics—

enabling quantification of false positive and negative rates—which were combined to yield the F1
scores shown in **Fig. 5d and S7.**

The results of these ensembles show that *R. toruloides* insertion number can be predicted reliably
from GFP alone (Mean F1 value of 0.82), with single insertions in particular achieving high
predictability (F1 = 0.89). Importantly, including the ORI family did not substantially improve model
performance, supporting the notion that the insertion number-GFP trend is ORI-independent in
this species, enabling prediction without *a priori* knowledge about an ORI family.

Fig. 5: Ensemble prediction of GFP insertion number in *R. toruloides*. **(a)** An ensemble of
predictive models was trained to predict the GFP insertion number of a transformant based on either GFP output alone
or GFP + ORI identity as predictive factors. These data were used to conduct 100 bootstrapped 70:30 training-test
splits across three independent models, which were then combined in a majority voting ensemble. Confusion matrices
were generated from the bootstrapped runs to analyze the ensemble's performance and rates of false positive and
negative results. *R. toruloides* showed high predictive capacity of insertion number from GFP alone **(b)** that was not
improved by the addition of ORI identity **(c)**. *R. toruloides* ensemble F1 statistics are reported in **(d)**, showing predictive
capacity for all groups from GFP alone (F1: 0 insertions, 0.78; 1 insertion, 0.89; 2-4 insertions, 0.82; 5+ insertions,
0.78). Error bars represent the SEM from the bootstrapped runs.

Prediction in *A. thaliana* was more challenging due to greater complexity between GFP
expression, insertion number, and silencing. Within this system, GFP expression alone enabled
modest classification of single-insertion and five+ insertion transformants (F1 scores = 0.61 and
0.49, respectively) but showed no ability to identify the intermediate 2-4 insertion group (F = 0.01).
Including the ORI family identity in the training model substantially improved predictive
performance, increasing F1 scores to 0.75, 0.62, and 0.75 for the 3 groups (**Fig. S7**). This further
supports the previous observation of ORI-dependent expression outcomes within *A. thaliana*.

In summary, GFP expression can serve as an excellent proxy for transgene insertion number in
R. toruloides, providing a promising avenue for high-throughput insertion number estimation using
simple fluorescence measurements. In *A. thaliana*, GFP alone provides some, but more limited,
information about insertion number extremes (single vs five+ insertions), and inclusion of ORI
identity is needed to achieve finer resolution. As it is possible that ORIs beyond the two used in
this study will perform differently cased on the variance observed, *a priori* information about the
ORI will be needed to train ORI-specific models in *A. thaliana* to maximize performance in this
species.

Intra- and interspecies modeling using GFP for copy number prediction

Having shown that GFP is a viable proxy for insertion number in *R. toruloides*, we next evaluated
the model's external validity and generalization. To do this, we took the same six ORI variants
and conducted another transformation, selecting 96 transformants per variant for evaluation. In
the same manner as before, these transformants were grown overnight and measured for GFP
fluorescence, which was then used for ensemble prediction of the insertion number group. These
predicted population-level insertional distributions were compared to their measured reference
populations to determine if there was significant similarity with previous results. For all ORI
variants across all insertion groups, predicted class proportions fell within the 95% posterior-
predicted confidence intervals of the measured insertion number classes (Dirichlet–multinomial),
and there were no significant differences between the measured and predicted population
outcomes (**Fig. 6a,** Fisher's Exact Test with Monte-Carlo, FDR-adjusted; all q > 0.05). This
demonstrates that GFP-based modeling recapitulates *R. toruloides* population compositions in
new transformations without gDNA/dPCR, validating fluorescent measurements and ensemble
prediction as an accurate alternative to manual quantification.

Fig. 6. Intra- and inter-organism modeling of transformation outcomes. **(a)** *R. toruloides* model validation by ORI variant. For each variant, bars show the measured population proportion for each GFP insertion-number class with 95% simultaneous multinomial intervals (Sison–Glaz). Markers denote the measured (o) or new population predicted (x) proportion for each group. Each group was evaluated with a Fisher's Exact Test followed by Monte-Carlo resampling (FDR adjusted; all q > 0.05). The inset reports the total-variation distance (TV) between measured and predicted outcomes. **(b)** Multidimensional scaling (MDS) of ORI-variant copy number distributions using 1D embeddings from Kolmogorov–Smirnov (KS) and Earth Mover's Distance (EMD) dissimilarities. Points show measured and predicted populations for pSa and pVS1 variants; RK2 is shown for predicted only. Ellipses indicate significant ORI-family clusters ($R^2 = 0.89$, $p = 4.2*10^{-6}$). **c)** Linear regression of *A. thaliana* versus *R. toruloides* single GFP insertion frequencies across the six shared ORI variants; the shaded band is the 95% CI of the fit, derived from the standard errors of the slope and intercept. (Pearson $r^2 = 0.85$, $p = 0.0087$; Spearman $\rho = 0.89$, $p = 0.0048$). RK2 predicted points are shown for reference but excluded from the regression. **(d)** Procrustes analysis comparing the pairwise KS-distances of ORI variants between organisms (measured data only). Lines connect matched ORI variants between organisms and ellipses denote ORI families. A Procrustes Randomization Test (PROTEST) indicates significant congruence of the two configurations (p = 0.026). **(e)** ORI–ORI pairwise KS distances within versus between ORI families in each organism. Boxplots summarize the distribution of within-family and between-family distances; overlaid p-values are from ANOVA contrasts. The between-within ORI variant trends are concordant across organisms and do not differ significantly (see **Fig. S8**).

Having validated the reproducibility of model results on ORI variants that were used to train the ensemble, we next evaluated ensemble performance on a new unseen ORI family, RK2. Using the WT form (~1 copy per cell) and two *repA* copy number variants—S20F (~5 copies) and R11G (~15 copies)–-we transformed *R. toruloides* with binary vectors containing identical T-DNA cargo as previous experiments. Measuring the GFP expression from 96 transformants per ORI variant, we estimated the population GFP insertion distribution using the ensemble. We then embedded

all populations—the original pSa and pVS1 measured transformants, their prediction-only
replicates, and novel RK2 data—via multi-dimensional scaling on pairwise 1D KS and EMD
distances (**Fig. 6b**). Measured and prediction-only populations co-localized, and all three ORI
families separated significantly (Permutational ANOVA, $p = 4.2 * 10^{-6}$), confirming that
fluorescence-based predictions recapitulate ORI-family level clustering. All RK2 variants
clustered more closely to the pSa family than pVS1 variants, strongly suggesting that RK2 binary
vectors produce transformation outcomes that are more similar to pSa than the pVS1 family, with
greater numbers of single insertions and lower average insertion number. Together, these results
provide a proof-of-concept pipeline in which new ORIs can be rapidly screened from fluorescence
in ~1 week in *R. toruloides* without the need for costly and laborious gDNA extraction and
transgene quantification for exploratory analyses.

With the *R. toruloides* modeling validated, we next asked whether the population-level signals in
R. toruloides were reflected in *A. thaliana*. To explore this, we analyzed the measured single-
insertion event generation frequency for all ORI variants in both systems. Single insertion events
are a valued transformation outcome for most plant engineering, making this a metric of particular
importance. Plotting the population outcomes against one another found a strong and significant
correlation between the two systems (**Fig. 6d,** Pearson $r^2 = 0.85$, $p = 0.0087$; Spearman $\rho = 0.89$,
$p = 0.0048$).

To expand the analysis beyond single-insertion transformants, we summarized whole-population
differences by MDS of pairwise KS distances among ORIs for each organism (**Fig. S8**). These
species-level MDS configurations were then compared using a Procrustes analysis which found
significant alignment between ORI variants across species (**Fig. 6e**, PROTEST $p = 0.026$). This
analysis implies that inter-ORI outcomes are conserved across organisms, despite possible
differences in overall effect size. To further explore how population variance between all pairwise
ORI-ORI groupings correlated between organisms, pairwise KS distances were calculated and
plotted against one another, resulting in 15 total comparisons per species. This comparison
showed significant concordance of ORI-ORI variant differences across organisms, particularly for
differentiating ORI-family level differences (**Fig. S9**, $p < 1.0 * 10^{-6}$). We further divided these
pairwise comparisons into within-ORI-family and between-family groups and found consistent
differences between organisms with between-family comparisons having greater KS distances
than within-family pairings (**Fig. 6f**, $p < 0.001$). On the species level, *A. thaliana* was found to have
a generally higher insertion rate and greater variability for all ORI variants than transformants in
R. toruloides. This is reflected in both the insertion spread and larger KS distance magnitudes,
but comparing within- and between-family distances showed no significant differences between
organisms, further suggesting a general cross-species trend (**Fig. S9**).

Taken as a whole, this modeling suggests two core results: 1) GFP can accurately predict
transformation outcomes in *R. toruloides*, and 2) population data in *R. toruloides* has a significant
predictive capacity on transformation outcomes in *A. thaliana* on both the single-insertion level
and ORI-ORI variance level more broadly. This pipeline of modeling GFP fluorescence to estimate
population insertional distributions in *R. toruloides* followed by transformation outcome prediction
in *A. thaliana* opens the door to streamlined screening of ORI variants, bypassing bottlenecks that
typically limit plant-based systems.

Discussion

In this study, we systematically evaluated how the binary vector ORI and plasmid copy number influence AMT outcomes across eukaryotic kingdoms. Our results demonstrate that ORI family—not copy number—is the primary determinant of transformation efficiency, transgene insertion number, transgene expression, and vector backbone inclusion frequency. ORI-dependent differences persisted even when plasmid copy number was substantially altered, revealing intrinsic, ORI-specific properties that strongly shape transformation outcomes.

While binary vector copy number was observed to influence transformation efficiency, its impact on transgene insertion frequency was dramatically lower compared to the influence of the underlying ORI family. These findings are particularly important given legitimate concerns that engineering ORIs to improve transformation efficiency could compromise quality by increasing transgene insertion number and backbone inclusion rates. In this study, we did not observe consistent or substantial tradeoffs between plasmid copy number and transformation quality across the variants and organisms tested. While modest effects were detectable—for example the pSa V152F variant in *Arabidopsis*—such impacts were not a general feature of the dataset. As such, there appears to be an engineerable space where single-insertion rates or transformation efficiency can be improved without substantially compromising other transformation variables, although broader testing across additional ORI families and transformation contexts will be necessary to fully validate this approach.

Since similar plasmid copy number variants across different ORI families produced highly variable results, intrinsic ORI characteristics appear to be major determinants of AMT outcomes. Our findings suggest that these ORI-specific features—such as potential differences in T-DNA supercoiling, plasmid localization within *Agrobacterium*, or interaction with host factors—retain their influence even when plasmid copy number is extensively modified. Notably, disparities in transgene expression in *A. thaliana* among transformants with identical insertion numbers underscore an underappreciated role for the binary vector ORI's influence on transformant quality. These results carry significant implications for users of pCambia vectors, which are widely used in academia for plant transformation but rely on the pVS1 ORI. In *A. thaliana*, pVS1-based vectors consistently produced high multi-insertion events—reaching up to 140 copies per transformant—and were associated with elevated rates of transgene silencing. These outcomes raise concerns about the reliability of pCambia-derived transformants in *Arabidopsis* research. Given these findings, we recommend reevaluating the use of pCambia vectors in *A. thaliana* and conducting a more systematic assessment of alternative ORIs for this system. For any engineering effort that would benefit from higher proportions of single insertions and functional transformation efficiency, utilizing the pSa K155N variant in *A. thaliana* and pSa K155N or V152F for *R. toruloides* are recommended from the variants tested in this study. For high insertion rates across systems, the pVS1 R106H vector can be deployed to combine high efficiency transformation with high rates of insertion.

Given the notable effects observed at the ORI family level, expanding bioprospecting efforts to explore the diverse natural repertoire of ORIs is a critical next step for improving AMT. To date,

conducting such a screen and evaluating transformation outcomes directly in plants—even
efficient models like *Arabidopsis*—remains prohibitively low-throughput. To address this
bottleneck, we developed a high-throughput fungal proxy system using *R. toruloides* coupled with
GFP fluorescence as a predictive indicator of transformation outcome. Comparison of results
between this yeast system and *A. thaliana* revealed significant mirrored trends across organisms
for both rates of single insertions and general ORI-ORI variant differences. This proxy system
represents a promising advance to enable scalable, predictive evaluation of binary vector
performance without the throughput limitations imposed by direct plant assays. Future research
should prioritize validating the generalizability of this system across a wider range of ORIs, host
organisms, and vector architectures. By implementing a high-throughput proxy system, this
approach has the potential to accelerate the exploration of ORI sequence space and advance
AMT toward becoming a more predictive and controllable platform for transgene insertion.

Methods

Media, chemicals and culture conditions
Bacterial cultures were grown in LB Miller medium (BD Biosciences) shaking at 200 rpm at 37 °C
for *E. coli* and 30 °C for *A. tumefaciens* unless otherwise noted. Cultures were supplemented with
kanamycin (50 mg L^{-1}; Sigma-Aldrich), gentamicin (30 mg L^{-1}; Thermo Fisher Scientific), or
rifampicin (100 mg L^{-1}; Teknova) as indicated. All compounds unless otherwise noted were
purchased from Sigma-Aldrich.
611
612

Strains and plasmids
All bacterial strains and plasmids used in this study are listed in Supplementary Table 1 and are
viewable through the public instance of the Joint BioEnergy Institute (JBEI) registry (https://public-
registry.jbei.org/folders/931). All strains and plasmids created in this work can be requested from
the strain archivist at JBEI with a signed material transfer agreement. All plasmids were Gibson
assembled using standard protocols[25] or Golden Gate assembled using standard protocols[26].
Plasmids were routinely isolated using the QIAprep spin miniprep kit (Qiagen), and all primers
were purchased from Integrated DNA Technologies (IDT). Plasmid sequences were verified using
whole-plasmid sequencing (Plasmidsaurus). *Agrobacterium* was routinely transformed by
electroporation as described previously using a 1-mm cuvette and a 2.4-kV, 25-µF, 200-Ω pulse[27].
623

Arabidopsis thaliana **transformation**
The Columbia-0 (Col-0) ecotype of *Arabidopsis thaliana* was grown in Sunshine no. 4 medium
supplemented with Osmocote at 22C under long-day conditions (16 h light, 8 h darkness;
150 µmol m^{-2} s^{-1} PAR) for 2 weeks after sowing numerous seeds onto the surface of a pot.
Seedlings were then transplanted to 4-inch pots, with 3 seedlings per pot, and grown under the
same conditions until flowering. The primary floral inflorescence was removed after reaching ~2
cm to encourage axillary shooting. When the inflorescences had grown and set their first few
siliques, the pots of plants were transformed via floral dip as previously described[28].
632
Briefly, the binary vectors used in this study were transformed into *A. tumefaciens* GV3101

(pMP90), which was grown to saturation in LB medium + kanamycin, gentamicin and rifampicin,
shaking at 200 rpm at 30 °C overnight. 100 mL of culture was centrifuged at 3,200g for 15 minutes
and resuspended in 200 mL of floral dip medium, consisting of 5% sucrose and 0.02% Silwet. All
formed siliques were removed with scissors, and then all inflorescences per pot were submerged
into this solution for 30 seconds. For buds that could not reach the solution, a pipette was used
to eject 50 µL of bacterial solution onto the bud surface. Up to 4 pots that had been dipped into
the same solution were then placed into a tray sideways and covered with a plastic dome that
had been misted with water. The following day, the pots were uncovered, returned to an upright
position, and placed into a growth chamber with identical growth conditions as above. One week
later, this process was repeated without removing developing siliques. Following the second dip,
the plants were allowed to continue growing with standard watering for 2 weeks followed by
cessation of watering to allow for full desiccation.
646
***A. thaliana* selection and evaluation**
Fully desiccated plants were harvested by sifting seeds through progressively finer sieves to
remove chaff. As seed weight was used to estimate total seed number, care was taken to remove
even fine chaff through >20 progressive sifts. ~150 mg of seed—corresponding to roughly 7500
seeds[29]—were placed into 1.5 mL Eppendorf tubes and sterilized. First, 1 mL of 70% ethanol was
placed into each tube for 1 minute, inverting. The ethanol was removed, and 1 mL of 50%
commercial bleach (12.5% hypochlorite) + 1 drop of Triton-20 surfactant was added, and the tube
was inverted for 7.5 minutes. This sterilization solution was then removed, and the seeds were
rinsed 3 times with sterile water. Seeds were then evenly spread onto the surface of 6.5-inch
plates containing Murashige and Skoog salts + $1\,g\,L^{-1}$ MES brought to a pH of 5.7 along with
$50\,mg\,L^{-1}$ kanamycin and $300\,mg\,L^{-1}$ timentin to kill residual *A. tumefaciens* and $8\,g\,L^{-1}$ agar for
solidification. These plates were placed into growth chambers with identical conditions used for
A. thaliana and were allowed to grow for 3 weeks.
660
Following germination and kanamycin selection, 96 transgenic *A. thaliana* were transplanted per
construct into flats—1 plant per cell, 24 cells per tray, 4 trays per construct. Plants were allowed
to grow under identical long-day conditions for 3 weeks. Following this in a systematic plant-by-
plant manner, 4 leaves were removed per plant, and a single 6 mm hole punch was taken per
leaf. These leaf discs were placed abaxial side up in a 96-well culture plate containing 330 µL
water per well. The remaining tissue was then placed into a separate 96-well block for storage
and later genomic DNA extraction. Each 96-well plate containing the leaf discs was then
measured for GFP fluorescence using a BioTek Synergy H1 microplate reader set for conditions
of 483-nm excitation and 512-nm emission. Raw disc-level *A. thaliana* meta data can be found in
Supplementary Table 2. Genomic DNA from each sample was extracted using a 96-sample
DNeasy kit (Qiagen, Product 69181).
672
***Rhodosporidium toruloides* transformation and evaluation**
The binary vectors used in this study were transformed into *A. tumefaciens* EHA105 (pTiBo542
ΔT-DNA). Cultures of *A. tumefaciens* were grown overnight in LB + kanamycin and rifampicin at
30 °C shaking at 200 rpm. Cultures were suspended at an OD_{600} of 1.0 in an AB salts induction

medium for 24 hours [30]. On the same day, *R. toruloides* that had been grown to saturation in YPD
medium (BD Biosciences) at 30 °C shaking at 200 rpm was diluted into YPD, such that the culture
would reach an OD_{600} between 0.5-1.0 the following morning.
680
After the induction period, 1 mL of OD_{600} = 1.0 *A. tumefaciens* and 1 mL of OD_{600} = 1.0 *R.*
toruloides were mixed and centrifuged at 4000g for 5 minutes. The supernatant was removed,
and the pellet resuspended in 1 mL of induction medium. This resuspension was then vacuumed
using a central laboratory vacuum line through a Millipore 0.45-micron membrane, which was
then placed onto induction medium that had been solidified with 2% agar. The cells were allowed
to co-cultivate for 4 days in the dark at 26 °C. Following this period, the pellet was resuspended
by vortexing the filter in 5 mL of YPD, and cells were plated on YPD + 2% agar with 100 mg L^{-1}
nourseothricin and 300 mg L^{-1} timentin. Selected colonies emerged after 2 days of incubation at
30 °C, and these were quantified using an AnalytikJena UVP GelSolo for imaging and Fiji for
counting (https://imagej.net/software/fiji/downloads).
691
For each construct, 96 colonies were then randomly selected and used to inoculate a 96-well
block (500 μL YPD + 20 mg L^{-1} nourseothricin + 300 mg L^{-1} timentin per well) for overnight growth
at 30 °C shaking at 200 rpm. The following day, 150 μL of culture was added to a 96-well plate
along with 150 μL of YPD, and each plate was measured for both OD_{600} and GFP fluorescence
using a BioTek Synergy H1 microplate reader set for conditions of 483-nm excitation and 512-nm
emission. Final GFP fluorescence was normalized to culture concentration by dividing measured
fluorescence by culture OD_{600}. Data for this experiment can be found in Supplementary Table 3.
Genomic DNA for all cultures was extracted from the pellet of 1 mL culture grown to an OD_{600} of
~1.0 using a Quick-DNA Fungal/Bacterial 96-sample kit (Zymo Research, Product D6006).
701
***Aspergillus niger* transformation and evaluation**
The six binary vectors used in this study were transformed into the EHA105 strain of *A.*
tumefaciens by electroporation. Fresh spores of the *albAD* mutant of *Aspergillus niger* derived
from the ATCC 11414 stock were produced on *Aspergillus* Complete Medium (CM) and diluted in
0.4% Tween-80 to a concentration of 10^5-10^6 spores per 125 mL. This mutant was used to enable
better GFP quantification by removing melanin biosynthesis. The *Agrobacterium* cultures were
grown in induction medium [31] to an OD_{600} of 0.6-1.0 and were then mixed in a 1:1 ratio with the
spore solution. This mixture was spread onto Amersham TM hybond-N nylon membranes and
placed on induction medium plates, followed by incubation for 2 days at 20 °C. The membranes
were then transferred to minimal medium agar plates containing 150 μg/mL hygromycin B and
250 μg/mL cefotaxime and were incubated at 30 °C for 3-4 days.
713
Transformed colonies were randomly picked and transferred to 13x100 mm sterile glass tubes
containing a 1.5 mL CM agar slant with the proper antibiotics. After 3-4 days of growth at 30 °C,
spores were released via vortexing and pipetting with 1 mL sterile 0.4% Tween-80 and passed
through one-layer Miracloth (EMD Millipore Corp, Burlington, MA, USA) into 1.5 mL
microcentrifuge tubes. Spore concentrations were determined by ORFLO Moxi V cell analyzer
(ORFLO Technologies, El Cajon, CA, USA), and the GFP fluorescence of each spore culture was
measured with a SpectraMax M5 (Molecular Devices, San Jose, CA, USA) using 395 nm

excitation and 509 nm emission spectra. GFP fluorescence was normalized to 10^6 spores for each
strain. Data can be found in Supplementary Table 3.
723
For genomic DNA extraction, spores of each transformant were inoculated into 1.5 mL CM liquid
medium with the proper antibiotics in 13 x 100 mm culture tubes and incubated without shaking
at 30 °C for 30-40 hours. Biomass was then removed and transferred into 1.5 mL tubes and
lyophilized followed by homogenization with glass beads using a mini-beadbeater-16 (BioSpec
Products, Inc, Bartlesville, OK, USA). DNA was purified using the Maxwell® RSC plant DNA kit
by the Maxwell® RSC instrument (Promega, Madison, WI, USA).
730
Transgene insertion number quantification
Transgene insertion number was quantified for all systems using nanoplate digital PCR with the
QIAcuity One system (Qiagen, Product 911001). Genomic DNA concentrations were quantified
using a Qubit fluorometric system (ThermoFisher Scientific, Product Q33238) and were diluted to
a concentration of 10,000-100,000 genomic copies per µL, followed by digestion with organism-
specific restriction enzymes that are specified below. All primary quantifications targeted the GFP
cassette within the T-DNA and a genomic control target using the primer-probe pairings specified
below:
A. thaliana **(Digested with NciI):**
· GFP (F: TTGTACTCCAGCTTGTGC, R: GACGGCAACTACAAGACC, FAM probe:
TTCAGCTCGATGCGGTTCACCAGG)
· gDNA (F: CCATCTACGGCATCAATATCC, R: AAACATTCCCAAGCAGTCC, HEX
probe: ACCACTTCCAAGAACATCTGCCACA)
· Kanamycin cassette (additional *A. thaliana* target) (F: TACCGTAAAGCACGAGGA, R:
GGACCGCTATCAGGACATA, FAM probe: TCGCCGCCAAGCTCTTCAGCAATA).
R. toruloides **(Digested with SacI)**
· GFP (F: GGTGTTCTGCTGGTAGT, R: CAGAAGAACGGCATCAAG, FAM Probe:
CCTCGATGTTGTGGCGGATCTTGA)
· gDNA (F: CCAGTCACCAACGGGAACG, R: CGACTCCGCTCACGAAGC, HEX probe:
AGCGCTATGTCGACGAGCGCTTCCCGTA)
A. niger **(Digested with AluI)**
· GFP (F: CAGGAACGCACGATCTTC, R: CGATACGGTTAACCAGTGTATC, FAM
probe: ACTATAAGACGCGGGCGGAGGTGA)
· gDNA (F: GTACACCCGCAACTTTACC, R: GCAGAGTGCCATTGTATGT, HEX probe:
TGGAGGCATTTGACAGCCATGCAGT)
756
All samples were run in 96-well 8,500-partition QIAcuity nanoplates (Product ID: 250021) using
the QIAcuity Probe PCR Mastermix (Product ID: 250102) with 40 2-step cycles of 15 seconds
denaturing at 95 °C and 30 seconds annealing and extension at 60 °C. Each reaction contained
primers and probes for both the T-DNA target (FAM probe) and gDNA target (HEX probe),
enabling absolute quantification of template numbers for both targets in each sample using
Poisson statistical methods via the QIAcuity Software Suite. The transgene insertion number was
calculated as the ratio of T-DNA:gDNA templates, adjusting for ploidy (*A. thaliana* diploid, fungal
systems haploid). Data can be found in Supplementary Table 3.

Backbone analysis

The inclusion of the binary vector backbone within *A. thaliana* transformants was analyzed using
qPCR on a CFX96 (Bio-Rad, discontinued) using SsoAdvanced™ Universal SYBR® Green
Supermix (Bio-Rad, Product ID: 1725271). A sequence 40 bp from the left border was targeted
with the following primers: F: AAATCACCACTCGATACAGG, R: ACAAGACGAACTCCAATTCA,
and the same primer set as the dPCR gDNA target was used for the gDNA control. Each target
was run in a separate reaction using 40 2-part cycles consisting of 15 seconds of 95 °C for
denaturing and 30 seconds of 60 °C for annealing and extension. Backbone inclusion was defined
as any sample with a Backbone Cq – Control Cq that was ≥2, which is indicative of samples with
notably lower backbone templates compared to gDNA control target templates. Backbone positive
and negative samples separated into distinct groups using this threshold as shown in Fig. S5a.
Backbone signal found in all samples was attributed to lysed residual *Agrobacterium* harboring
binary vectors that contained the backbone template sequence.
779

Insertion number modeling

All modeling analyses were conducted in R using the packages caret, MASS, e1071,
randomForest, class, nnet, and glmnet. Data from the combined transformation dataset were
filtered by organism, rounded to the nearest integer GFP insertion number, and grouped into three
or four categorical classes. For *R. toruloides*, insertions were classified into "Zero," "One," "Two–
Four," and "Five+" insertion groups, while *A. thaliana* was divided into "One," "Two–Four," and
"Five+," as zero-insertion events were virtually absent in this system. Two predictor sets were
evaluated for each organism: one using GFP fluorescence alone, and another combining
fluorescence with binary vector ORI family identity.

Eight supervised learning algorithms were benchmarked: quadratic and linear discriminant
analysis, k-nearest neighbors (k = 4), random forest, multinomial logistic regression, linear
support vector machine (SVM), naïve Bayes, and penalized multinomial regression. Models were
trained and tested over 100 iterations of stratified Monte Carlo cross-validation, in which 70% of
data were randomly assigned to training and 30% to testing using caret::createDataPartition. For
each iteration, model predictions on held-out data were compared to true class labels using
caret::confusionMatrix, from which overall and per-class balanced accuracies were extracted.
Numeric predictors were scaled where appropriate, and factor levels were standardized across
partitions. Mean per-class and overall accuracies were computed across iterations, with the
standard error of the mean reported for overall accuracy.

From these benchmark results, a three-model ensemble was constructed to improve prediction
robustness. The ensemble combined a linear SVM, naïve Bayes, and multinomial logistic
regression classifier for *R. toruloides*, and a linear SVM, naïve Bayes, and quadratic discriminant
analysis model for *A. thaliana*. Each model was trained independently using GFP fluorescence (±
ORI family identity) as input features, and predictions were aggregated through majority voting,
where each model cast one vote for a predicted insertion class. When all three models agreed,

the classification was unanimous; when two agreed, the majority class was selected. In the rare
case of a three-way tie (each model predicting a different class), the sample was assigned to the
"Zero" category to ensure deterministic resolution. This ensemble framework leveraged the
complementary strengths of discriminative and probabilistic learners, producing more stable and
accurate predictions of GFP insertion number from fluorescence data.

Statistical Analyses

All statistical analysis was conducted in R using the following packages: core parametric and
nonparametric tests—including t, Wilcoxon, Kolmogorov–Smirnov, Shapiro–Wilk, and correlation
tests—were conducted with the stats package. The emmeans package provided estimated
marginal means and pairwise contrasts for generalized linear models. Distributional differences
were quantified using Earth Mover's Distance from emdist, and rstatix was used for pairwise
Wilcoxon tests with Benjamini–Hochberg correction. Linear and logistic model summaries were
tidied using broom, and multcompView generated compact letter displays summarizing multiple
comparisons. For machine learning–based classification, the caret framework was used to
benchmark eight algorithms spanning MASS (LDA/QDA), e1071 (Naïve Bayes, SVM),
randomForest, class (kNN), and glmnet (penalized regression), with MLmetrics providing F1
score evaluation.

Acknowledgements:

We would like to thank Stanton Gelvin and N. Louise Glass for productive discussions related to
this manuscript. Figure cartoon representations sourced from www.biorender.com. M.B. and
J.O.B. were supported by NIH R01-GM145814. L.M.W. was funded through the National Science
Foundation Graduate Research Fellowship Program. *Aspergillus* work was performed by the
Pacific Northwest National Laboratory, operated for the U. S. Department of Energy (DOE) by
Battelle under Contract DE-AC06-76RL01830 and supported by the U.S. DOE's Bioenergy
Technologies Office under agreement 2.5.3.106. Other work was part of the JBEI
(https://www.jbei.org) supported by the US Department of Energy (DOE), Office of Science, Office
of Biological and Environmental Research, through contract DE-AC02-05CH11231 between
Lawrence Berkeley National Laboratory and the DOE. The funders had no role in manuscript
preparation or the decision to publish. The views and opinions of the authors expressed herein
do not necessarily state or reflect those of the United States government or any agency thereof.
Neither the United States government nor any agency thereof, nor any of their employees, makes
any warranty, expressed or implied, assumes any legal liability or responsibility for the accuracy,
completeness or usefulness of any information, apparatus, product or process disclosed or
represents that its use would not infringe privately owned rights. The United States government
retains and the publisher, by accepting the article for publication, acknowledges that the United
States government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or
reproduce the published form of this manuscript or allow others to do so for United States
government purposes. The DOE provides public access to these results of federally sponsored
research in accordance with the DOE public access plan (http://energy.gov/downloads/doe-
public-access-plan).

Conflicts of Interest: M.J.S., M.G.T., and P.M.S. have financial interest in BasidioBio. P.M.S. has financial interest in Totality Biosciences. N.J.H. declares financial interests in TeselaGen Biotechnologies and Ansa Biotechnologies. B.A.S. has a financial interest in Illium Technologies, Caribou Biofuels, and Erg Bio.

Contributions: Conceptualization, M.J.S., M.G.T., P.M.S.; methodology, M.J.S., M.B., Z.D., N.L., G.M.G., J.J., M.G.T; investigation, M.J.S., M.B., Z.D., J.J., L.M.W., Y.L., M.G.T., P.M.S.; writing—original draft, M.J.S.; writing—review and editing, all authors; resources and supervision, N.J.H., J.O.B., B.S., J.M.G., J.K., M.G.T., P.M.S.

Bibliography

1. Hellens, R., Mullineaux, P. & Klee, H. Technical Focus:a guide to Agrobacterium binary Ti vectors. *Trends Plant Sci.* 5, 446–451 (2000).

2. Chetty, V. J. *et al.* Evaluation of four Agrobacterium tumefaciens strains for the genetic transformation of tomato (Solanum lycopersicum L.) cultivar Micro-Tom. *Plant Cell Rep.* 32, 239–247 (2013).

3. Gelvin, S. B. Integration of Agrobacterium T-DNA into the Plant Genome. *Annu. Rev. Genet.* 51, 195–217 (2017).

4. Grevelding, C., Fantes, V., Kemper, E., Schell, J. & Masterson, R. Single-copy T-DNA insertions in Arabidopsis are the predominant form of integration in root-derived transgenics, whereas multiple insertions are found in leaf discs. *Plant Mol. Biol.* 23, 847–860 (1993).

5. Zhi, L. *et al.* Effect of Agrobacterium strain and plasmid copy number on transformation frequency, event quality and usable event quality in an elite maize cultivar. *Plant Cell Rep.* 34, 745–754 (2015).

6. Oltmanns, H. *et al.* Generation of backbone-free, low transgene copy plants by launching T-DNA from the Agrobacterium chromosome. *Plant Physiol.* 152, 1158–1166 (2010).

7. Ye, X. *et al.* Enhanced production of single copy backbone-free transgenic plants in multiple crop species using binary vectors with a pRi replication origin in Agrobacterium

tumefaciens. *Transgenic Res.* 20, 773–786 (2011).

8. Szarzanowicz, M. J. *et al.* Binary vector copy number engineering improves Agrobacterium-

mediated transformation. *Nat. Biotechnol.* 43, 1708–1716 (2025).

9. Francis, K. E. & Spiker, S. Identification of Arabidopsis thaliana transformants without

selection reveals a high occurrence of silenced T-DNA integrations. *Plant J.* 41, 464–477

(2005).

10. Kim, S.-I., Veena & Gelvin, S. B. Genome-wide analysis of Agrobacterium T-DNA

integration sites in the Arabidopsis genome generated under non-selective conditions. *Plant*

J. 51, 779–791 (2007).

11. Ghedira, R., De Buck, S., Nolf, J. & Depicker, A. The efficiency of Arabidopsis thaliana

floral dip transformation is determined not only by the Agrobacterium strain used but also

by the physiology and the ecotype of the dipped plant. *Mol. Plant Microbe Interact.* 26, 823–

832 (2013).

12. Napoli, C., Lemieux, C. & Jorgensen, R. Introduction of a chimeric chalcone synthase gene

into Petunia results in reversible co-suppression of homologous genes *in trans*. *Plant Cell*

2, 279–289 (1990).

13. Głowacka, K. *et al.* An evaluation of new and established methods to determine T-DNA

copy number and homozygosity in transgenic plants. *Plant Cell Environ.* 39, 908–917

(2016).

14. De Buck, S., Podevin, N., Nolf, J., Jacobs, A. & Depicker, A. The T-DNA integration pattern

in Arabidopsis transformants is highly determined by the transformed target cell. *Plant J.*

60, 134–145 (2009).

15. Jupe, F. *et al.* The complex architecture and epigenomic impact of plant T-DNA insertions.

PLoS Genet. 15, e1007819 (2019).

16. Pucker, B., Kleinbölting, N. & Weisshaar, B. Large scale genomic rearrangements in

selected Arabidopsis thaliana T-DNA lines are caused by T-DNA insertion mutagenesis.

BMC Genomics 22, 599 (2021).

17. Tinland, B. The integration of T-DNA into plant genomes. *Trends Plant Sci.* 1, 178–184

(1996).

18. Kleinboelting, N. *et al.* The Structural Features of Thousands of T-DNA Insertion Sites Are

Consistent with a Double-Strand Break Repair-Based Insertion Mechanism. *Mol. Plant* 8,

1651–1664 (2015).

19. Thomson, G., Dickinson, L. & Jacob, Y. Genomic consequences associated with

Agrobacterium-mediated transformation of plants. *Plant J.* 117, 342–363 (2024).

20. Nicolia, A., Ferradini, N., Veronesi, F. & Rosellini, D. An Insight into T-DNA Integration

Events in Medicago sativa. *Int. J. Mol. Sci.* 18, (2017).

21. Olhoft, P. M., Flagel, L. E. & Somers, D. A. T-DNA locus structure in a large population of

soybean plants transformed using the Agrobacterium-mediated cotyledonary-node method.

Plant Biotechnol. J. 2, 289–300 (2004).

22. De Buck, S. T-DNA vector backbone sequences are frequently integrated into the genome

of transgenic plants obtained by Agrobacterium-mediated transformation. *Mol. Breed.* 6,

459–468 (2000).

23. Afolabi, A. S., Worland, B., Snape, J. W. & Vain, P. A large-scale study of rice plants

transformed with different T-DNAs provides new insights into locus composition and T-DNA

linkage configurations. *Theor. Appl. Genet.* 109, 815–826 (2004).

24. Kuraya, Y. *et al.* Suppression of transfer of non-T-DNA vector backbone sequences by

multiple left border repeats in vectors for transformation of higher plants mediated by

Agrobacterium tumefaciens. *Mol. Breeding* 14, 309–320 (2004).

25. Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred

kilobases. *Nat. Methods* 6, 343–345 (2009).

26. Engler, C., Kandzia, R. & Marillonnet, S. A one pot, one step, precision cloning method with

high throughput capability. *PLoS ONE* 3, e3647 (2008).

27. Kámán-Tóth, E., Pogány, M., Dankó, T., Szatmári, Á. & Bozsó, Z. A simplified and efficient

Agrobacterium tumefaciens electroporation method. *3 Biotech* 8, 148 (2018).

28. Clough, S. J. & Bent, A. F. Floral dip: a simplified method for *Agrobacterium*-mediated

transformation of *Arabidopsis thaliana*. *Plant J.* 16, 735–743 (1998).

29. Rivero, L. *et al.* Handling *Arabidopsis* plants: growth, preservation of seeds, transformation,

and genetic crosses. *Methods Mol. Biol.* 1062, 3–25 (2014).

30. Zhang, S. *et al.* Engineering *Rhodosporidium toruloides* for increased lipid production.

Biotechnol. Bioeng. 113, 1056–1066 (2016).

31. de Groot, M. J., Bundock, P., Hooykaas, P. J. & Beijersbergen, A. G. *Agrobacterium*

tumefaciens-mediated transformation of filamentous fungi. *Nat. Biotechnol.* 16, 839–842

(1998).