Human airway epithelia clear rhinovirus-infected cells using two waves of cell extrusion

Faith Fore[1,2,5], Dustin Bagley[1,2,5], Rocio Teresa Martinez-Nunez[3,5], Julia Aniscenko[4,5], Sebastian Johnston[4,5] and Jody Rosenblatt[1,2,5,6]

1. The Randall Centre for Cell & Molecular Biophysics, School of Basic & Medical Biosciences, King's College London, London.
2. The Francis Crick Institute, London, UK.
3. King's College London, Department of Infectious Diseases, School of Immunology & Microbial Sciences, Faculty of Life Sciences & Medicine, King's College London, Guy's Hospital, London, UK
4. National Heart and Lung Institute, Imperial College London, United Kingdom
5. Asthma UK Centre in Allergic Mechanisms of Asthma, London, United Kingdom
6. School of Cancer and Pharmaceutical Sciences, King's College London, London SE1 1UL, UK.

Abstract

Rhinoviruses (RV) cause most common colds and asthma attacks. While RV typically infects airway epithelia, most studies infect cell lines that cannot form proper epithelial barriers. Cell extrusion is a highly conserved process that drives most epithelial cell death by seamlessly squeezing out excess live cells to maintain constant cell densities and a tight barrier. We discovered that healthy bronchial epithelia clear RV within 24 hours by selectively extruding infected cells, a process called virus-induced cell extrusion (VICE). In contrast, barrier-defective cells fail to clear the virus and accumulate infection. VICE occurs in two phases: an early wave triggered by mechanical signals, and a later wave dependent on viral replication. Although extrusion removes infection, virus-carrying expelled cells can still spread RV. These findings highlight extrusion's dual role in both clearing and transmitting rhinovirus, independent of typical inflammation.

Main

Rhinoviruses (RV), the leading cause of the common cold, typically cause mild illness but can lead to severe symptoms in young children and adults with underlying conditions such as asthma, chronic obstructive pulmonary disease (COPD), or cystic fibrosis. RVs are non-enveloped positive-sense single-stranded RNA (ssRNA) viruses [1–3] of which there are three classes: RV-A, RV-B, and RV-C. The RV-A and RV-B classes subdivide into major and minor groups, where the major group binds the intercellular adhesion molecule 1 (ICAM-1) receptor and the minor group binds the low-density lipoprotein receptor (LDLR). RV-C instead uses the cadherin-related family member 3 (CDHR3) receptor [2,4].

Human airway epithelia (HAE) provide a physical barrier to the outside world that filters out toxins and pathogens, acting as a fundamental, primary step in innate immune responses [2]. Despite epithelia being the primary sites of viral infection, most virology studies infect cultured single cells that do not form a tight epithelial barrier that viruses would normally encounter, thereby overlooking this critical first step. Because HAEs are constantly exposed to pathogens, pollutants, and irritants, they are adept at repairing and renewing damaged cells. Moreover, as epithelial cells turn over, they must match the number of cells that divide with those that die; otherwise tumours or barrier defects and inflammation could rapidly occur [5]. We previously found that mechanics regulate cell number homeostasis: stretching from too few cells activates division, and crowding from too many triggers cells to die through an ancient mechanism, conserved from sea sponges to humans, termed cell extrusion [6]. To extrude, crowding activates Piezo1 in live cells to form and contract a basolateral ring of actin and myosin that seamlessly squeezes them apically out of the layer [7–11].

Additionally, apoptotic and damaged cells also extrude through a conserved mechanism that does not require Piezo1 to ensure defective cells are seamlessly removed without disrupting the barrier[9]. Given the fundamental role of epithelia in protecting the body from pathogens, we investigated how epithelial sheets cope with RV infections.

Results

Human rhinovirus infection leads to extrusion of infected epithelial cells

To investigate how intact epithelial monolayers respond to RV infection, we infected human bronchial epithelial cells (16HBE14o⁻) with RVA01B or RVA02 (minor group), RVB14 or RVA16 (major group) at a multiplicity of infection (MOI) of 3 for 24 hours. RV-infected cells were identified by immunostaining for the capsid protein VP3 (major group) or double-stranded RNA (dsRNA) (minor group), actin with phalloidin, and nuclei with DAPI. Confocal imaging revealed that both major and minor RV strains triggered extrusion of infected cells, characterised by a constricted actin-myosin ring beneath VP3⁺ or dsRNA⁺ cells (**Fig. 1a**). Because all RV strains tested induced extrusion, we focused on RVA16, a major group, which could be tracked from the earliest stages of infection using VP3 immunostaining.

Remarkably, extrusion of RVA16-infected cells was detectable at the earliest time point of 2 hours post-infection (hpi) (**Fig. 1b**), when the VP3 immunostaining was restricted to the cell periphery, suggesting that viral attachment or early post-attachment events may be sufficient to trigger extrusion. In contrast, by 24 hpi VP3 localised throughout the cell during extrusion. While some non-infected cells extrude, as expected during homeostatic cell turnover, RVA16-infected VP3⁺ cells were preferentially extruded, since cell extrusion rates were significantly higher in infected monolayers compared to mock-infected controls (**Fig. 1c**). We termed this epithelial response to RV Virus-Induced Cell Extrusion (VICE).

By 24 hpi, few VP3⁺ cells remained, steadily declining from ~25% at 2 hpi to < 5% at 24 hpi (**Fig. 1d**). Additionally, quantification of RV RNA by RT-qPCR in both extruded and monolayer-associated cells (**Fig. 1e**) showed very few RV copies remaining in the infected monolayer by 24 hpi. Notably, both extruded and monolayer cells had equal RV RNA levels at 8hpi, indicating that roughly half of the virus had been cleared by VICE by this time, consistent with immunostaining data in **Fig. 1b** and **1d.** While RT-qPCR measures total viral RNA and immunofluorescence quantifies infected cells, both methods showed the same trend in viral reduction following RVA16 infection. Importantly, the percentage of infected cells progressively declined as the rate of extrusion increased (**Fig. 1f**), supporting a direct role of VICE in RV clearance. Thus, 16HBE14o⁻ cells eliminate RV infection through VICE in the absence of other innate immune cells.

To confirm if VICE occurs in real epithelial tissues, we infected *ex vivo* mouse lung slices with RVA01B, a minor RV strain which can infect mice. Infection of mouse lung slices with RVA01B induced high rates of extrusion by 24 hpi (**Fig. 1g**). Because we lacked a structural RV antibody to RVA01B, we immunostained infected slices with dsRNA antibody at 24 hpi, at which point the extrusions were extensive enough to cause sheet detachment (**Fig. 1g(ii)**), as seen in bronchoconstricted airways[12]. The high extrusion rates may reflect a higher effective MOI, which is more difficult to calculate in tissue samples.

An intact human airway epithelium clears rhinovirus-infected cells by VICE

As we found that extrusion alone could eliminate RV, we investigated why this may not have been noted in previous studies. To do so, we infected BEAS-2B cells, a human epithelial cell line established in 1988 from normal human bronchial epithelial cells[13] that is widely used in RV studies. Unlike 16HBE14o⁻ monolayers (**Fig. 1f**), BEAS-2Bs could not extrude and failed to eliminate RV, resulting in RV accumulation over time (**Fig. 2a**). We reasoned that the inability of BEAS-2Bs to clear RV (**Fig. 2c**) might stem from their inability to form a tight epithelium with proper tight zonula adherens(ZO-1) and adherens junctions (E-cadherin) as seen in 16HBE14o⁻ monolayers (**Fig. 2b**). Thus, although BEAS-2Bs derive from bronchial epithelia, they behave more like mesenchymal cells [14]. To test if BEAS-2Bs inability to eliminate RV infection by extrusion was due to their lack of proper cell-cell junctions, we disrupted 16HBE14o⁻ junctions by calcium removal or chelation with EGTA (**Extended Data Fig.1**). Both ways of disrupting junctions prevented VICE-induced RV clearance, resulting in RV accumulation at rates seen in BEAS-2Bs at 24 hpi (**Fig. 2c&d**). These findings

demonstrate that VICE-mediated clearance of RV requires an intact, extrusion-competent epithelial monolayer, revealing why studies using non-extruding cell lines may overlook this critical early antiviral defence **(Fig. 2e)**.

Fig. 1 : Human rhinovirus infection leads to extrusion of infected epithelial cells. a, Confocal projections of 16HBE14o- monolayers infected for 24 hours with major-group (RVA16, RVB14) or minor-group (RVA01, RVA02) rhinoviruses. Infected cells were stained for VP3 (major-group viruses) or dsRNA (minor-group viruses). F-actin was labelled with phalloidin; nuclei with DAPI. **b**, Time course of RVA16 infection (MOI = 3) at 2, 6 and 24 h post-infection (hpi). **c**, Quantification of extrusions per 1000 cells in RVA16-infected and mock monolayers. **d**, Percentage of VP3+ cells in 16HBE14o- monolayers over time after RVA16 infection. **e**, RV RNA levels in extruded and monolayer-associated cells at 8 and 24 hpi, normalised to GAPDH and SDHA. **f**, Rate of extrusion and the proportion of VP3+ cells in RVA16-infected monolayers. **g**, Confocal projections of mouse precision-cut lung slices (PCLS) infected with RVA01 for 24 h and mock controls. Infected cells stained with dsRNA; F-actin with phalloidin; nuclei with DAPI . Data are from ≥ 3 biological replicates. Transparency distinguishes independent biological experiments (d and e); bars represent the mean with s.e.m. (c–f). Statistical significance was determined by Welch's ANOVA with Games–Howell post hoc test (d) and Mann–Whitney test (e). **** p < 0.0001, *** p < 0.001, ** p < 0.01, * p < 0.05. White arrowheads indicate extruding cells. Data are with DAPI.

Fig. 2: An intact human airway epithelium clears rhinovirus-infected cells by VICE. All infections were performed with RVA16 at an MOI of 3 unless otherwise indicated. **a**, Confocal images of Beas-2B cells infected with RVA16 for 2, 6 and 24 hours. **b**, Confocal projections showing F-actin, E-cadherin and ZO-1 staining in 16HBE14o⁻ and Beas-2B monolayers. **c**, Percentage of infected Beas-2B cells quantified from VP3⁺ cells at different time points after RVA16 infection. **d**, Percentage of VP3⁺ cells remaining at 24 hours in 16HBE14o⁻ monolayers with intact or disrupted junctions. **e**, Schematic illustrating clearance of infected cells in extrusion-competent (16HBE14o⁻) versus non-extruding (Beas-2B) epithelia. **f**, Kinetics of cell extrusion in RVA16-infected 16HBE14o⁻ monolayers compared with mock infected or JTE-013–treated cells. Extrusions were quantified every 2 hours for 12 hours by live-cell imaging (Extended Data SMovie 1–3). **g**, Percentage of VP3⁺ cells remaining at 24 hours in RVA16-infected 16HBE14o⁻ monolayers with or without JTE-013 treatment. Data are from ≥ 3 biological replicates. Transparency distinguishes independent biological experiments (c, d and g); bars represent the mean with s.e.m. (**c, d** and **g**); error bars show s.e.m. Statistical significance was determined by one-way ANOVA with Tukey post hoc test (**c**), Kruskal–Wallis test (**d** and **g**), and **** p < 0.0001.

To confirm whether cell extrusion causes viral clearance, we blocked canonical extrusion signalling during infection. To extrude, a cell emits the lipid sphingosine-1 phosphate (S1P), which binds the G-protein-coupled receptor S1P receptor 2 (S1P$_2$) to activate Rho-mediated contraction of an actomyosin ring basally and circumferentially that squeezes the cell out apically[9,11,15–17]. Because nearly all extrusions resulting from RV infection were due to VICE from our fixed, immunostained samples (**Fig. 1c**), we reasoned that we could score VICE rates over time using time-lapse phase imaging ± the S1P$_2$ antagonist JTE-013[9]. JTE-

013 reduced the VICE rate to low extrusion levels seen in mock-infected cells (**Fig. 2f & Extended Data SMovie.1-3**), confirming that VICE requires the canonical extrusion signalling pathway. JTE-013 also blocked RV clearance from the monolayer at 24 hpi (**Fig. 2g**), confirming that RV elimination requires VICE. Together, these findings indicate that VICE eliminates RV-infected epithelial cells using the canonical extrusion pathway.

Extended Data Fig. 1: Disruption of tight junctions in 16HBE14o⁻ monolayers. Representative confocal images showing disruption of tight junctions in 16HBE14o⁻ monolayers. Junctions were disrupted by incubation in calcium-free medium or by chelating extracellular calcium with EGTA. Tight junctions were visualised using ZO-1 (zonula occludens-1) staining; nuclei were counterstained with DAPI. Scale bar, 25 µm.

RV induces two waves of cell extrusion, a live and an apoptotic one

While all apical extrusion pathways require the canonical signalling tested above, different stimuli can trigger this signalling via different upstream pathways. To maintain homeostatic cell densities, most cells extrude while alive in response to crowding via Piezo1[10], however, apoptotic stimuli can also induce extrusion to ensure that dying cells do not compromise the integrity of the monolayer [18]. To investigate whether RV induces apoptotic or live-cell extrusion, we infected 16HBE14o⁻ monolayers with RVA16 and immunostained for the viral structural protein VP3, active caspase-3 (a marker of apoptosis), and phalloidin (actin) to visualise extrusion rings. We found that VICE was predominantly non-apoptotic during the first 8 hpi, shifting to mainly apoptotic extrusions for the next 12 hours (**Fig. 3a&b**). These two separate waves of live and apoptotic extrusion likely account for the two peaks of total VICE over 24 hours, seen in **Fig. 1c.**

Given that apoptotic extrusion requires caspase activity,[18] we tested whether the second wave of VICE requires caspases by treating infected 16HBE14o⁻ cells with z-VAD-FMK, an irreversible pan-caspase inhibitor[19,20]. Time-lapse imaging revealed that extrusion rates were significantly reduced from 8 hours onward in z-VAD-FMK treated cells compared to untreated controls, without largely impacting the first wave of extrusion (**Fig. 3c**). Since the apoptotic extrusion wave initiates at 8 hpi, the canonical time of RV replication[2], we tested whether the second apoptotic wave of VICE requires RV replication using remdesivir, a cell-permeable monophosphate that blocks viral RNA replication in RV, Ebola virus, Middle East Respiratory Syndrome Coronavirus (MERS-CoV) 2, severe acute respiratory syndrome (SARS)-CoV and SARS-CoV 2 [21,22,23]. We confirmed that remdesivir blocks dsRNA synthesis following RV infection in HeLa-H1 cells using immunostaining ± remdesivir (**Extended Data. Fig. 2**). Notably, remdesivir treatment greatly reduced RV clearance through VICE at 24 hpi (**Fig. 3d**). Live imaging showed that VICE rates in RVA16-infected 16HBE14o⁻ cells treated with remdesivir were similar to untreated cells during the first 8 hours but declined significantly thereafter (**Fig. 3e**). Additionally, remdesivir reduced apoptosis in RV-infected 16HBE14o⁻ monolayers, compared with untreated infected cells (**Fig. 3f**), leading to fewer apoptotic extrusions (**Fig. 3g**). Thus, viral replication triggers the second apoptotic wave of VICE.

Fig. 3:   RV induces two waves of cell extrusion, a live and an apoptotic one. All infections were performed in 16HBE14o⁻ monolayers with RVA16 at an MOI of 3 unless otherwise indicated. **a**, Confocal images of apoptotic (caspase-3–positive) and non-apoptotic (caspase-3–negative) extrusions at 6 h post infection. Actin (red), nuclei (blue), VP3 (green) and active caspase-3 (magenta). White arrowheads indicate extruding cells. **b**, Quantification of apoptotic and non-apoptotic extrusions over time. **c**, Extrusion kinetics compared with mock- or z-VAD-FMK–treated cells, measured every 2 h for 12 h by live-cell imaging. **d**, Percentage of VP3⁺ cells at 24 h with or without remdesivir treatment. **e**, Extrusion kinetics ± remdesivir, quantified every 2 h for 12 h by live-cell imaging. **f**, Apoptosis with or without remdesivir. **g,** Apoptotic extrusions with or without remdesivir. Data are from ≥ 3 biological replicates. Transparency distinguishes independent biological experiments (**c and e**); bars represent the mean with s.e.m. (**b, d, f and g**); error bars show s.e.m. Statistical significance was determined by two-tailed t-test (**b**), Kruskal–Wallis test (**d**), Welch's t-test (**f**) and Mann–Whitney test (**g**); **** p < 0.0001, *** p < 0.001, ** p < 0.01.

Extended Data Fig. 2: Remdesivir inhibits replication of RVA16 in HeLa-H1 cells. Representative confocal images showing inhibition of RVA16 replication in HeLa-H1 cells treated with remdesivir. Cells were infected with RVA16 for 24 hours ± of remdesivir. Viral replication was visualised by dsRNA staining (red); nuclei were counterstained with DAPI (blue). Scale bar, 25 μm

The first VICE wave is mechanical and independent of dynamin-dependent RV endocytosis

Since the first VICE wave is predominantly live cell extrusion, we tested if stretch-activated channels (SACs), which typically control live cell extrusion during homeostasis upstream of S1P2 signalling, control this first wave. To do so, we infected 16HBE14o⁻ in the presence or absence of gadolinium (Gd⁺³), a generic SAC blocker or GsMTx4, a tarantula venom peptide toxin that inhibits the transient receptor potential channels TRPC1, TRPC6 [24,25],[26] and Piezo1 [24,26]. Both Gd⁺³ and GsMTx4 significantly reduce RV clearance by 24 hpi **(Fig. 4a)**, albeit to a lesser extent than blocking S1P₂ with JTE-013. Live imaging demonstrated that both Gd⁺³ and GsMtx4 reduced VICE during the first 8 hours of infection **(Fig. 4b),** indicating that SACs control the first wave of VICE, accounting for reduced RV clearance.

Fig.4: The first extrusion wave is mechanical and independent of dynamin-dependent RV endocytosis or cell death. All infections were performed in 16HBE14o⁻ monolayers with RVA16 at an MOI of 3 unless otherwise indicated. **a**, Percentage of VP3⁺ cells remaining at 24 hpi with or without Gd³⁺ or GsMtx4. **b**, Extrusion kinetics compared with mock- or Gd³⁺/GsMtx4-treated monolayers, measured every 2 hours for 12 hours by live-cell imaging. **c**, Scanning confocal projections at 1 hpi with or without Dynole 34-2 (scale bar, 25 μm). **d**, Extrusion kinetics compared with mock- or Dynole-treated monolayers, measured every 2 hours for 12 hours by live-cell imaging. **e**, Percentage of VP3⁺ cells remaining at 24 h with or without Dynole 34-2. **f,** Total extrusion events at 0 hpi in mock- and Dynole-treated monolayers. **g**, Schematic model of the VICE (virus-induced cell extrusion) pathway summarising the proposed mechanism of RV-driven extrusion and the pharmacological targets tested in this study, including Gd³⁺ and GsMtx4 (stretch-activated/TRP channels), z-VAD-FMK (apoptosis), remdesivir (viral replication) and JTE-013 (S1P₂ receptor). Data are from ≥ 3 biological replicates. Transparency distinguishes independent biological experiments (**a, e and f**); bars represent the mean with s.e.m. (**a, e and f**); error bars show s.e.m. Statistical significance was determined by Kruskal–Wallis test, ns p > 0.05; **** p < 0.0001; ** p < 0.01.

In both cases, GsMTx4 had a greater impact on VICE and virus clearance (**Fig. 4a&b**), confirming the role of VICE in RV clearance. These results confirm that VICE occurs in at least two waves, with the first live cell extrusion wave being controlled by SACs and the second apoptotic cell extrusion wave dependent on viral replication.

Extended Data Fig. 3: The early VICE wave is initiated during rhinovirus adsorption. a, Schematic of the infection model illustrating the adsorption and infection initiation steps (0 hpi). **b**, Representative confocal images showing inhibition of transferrin uptake by Dynole 34-2. Early endosomes were identified by EEA1 (early endosome antigen 1) staining. **c**, Scanning confocal z-stack of a VICE cell at 0 hpi following RVA16 adsorption at room temperature. Images are shown from the basal to apical plane, numbered 1–16. **d**, Representative confocal images of RVA16-infected cells ± Dynole 34-2. Scale bar, 25 μm

Given that RV localised to the cell borders during VICE in the earliest extrusions measured (2 hpi, **Fig. 1b**), we investigated if VICE might be activated before/during virus internalisation during the virus adsorption incubation

step **(Extended Data Fig 3a)**. Using high-resolution confocal imaging, following a one-hour adsorption step at room temperature (see **Extended Data Fig 3a**), we found all RV remained colocalised with cortical actin at the plasma membrane during VICE (Fig. 1c, n=8). To test if VICE requires RV uptake by receptor-mediated endocytosis[27], we infected 16HBE14o- monolayers with RVA16 ± dynole 34-2, a cell-permeable allosteric inhibitor of dynamin-mediated endocytosis[28,29]. A transferrin uptake assay confirmed that dynole 34-2 inhibits dynamin-dependent endocytosis in our system **(Extended Data Fig 3b)**. Indeed, we found the same peripheral VP3 staining in dynole-treated cells as seen in early VICE (**Fig. 4c**). The VICE scored during this early time frame with RV restricted to the cell periphery could result from homeostatic background extrusions that had inadvertently been coated with RV. However, we found that the number of extrusions during RV absorption ± dynole was markedly higher than not only background mock controls but also than occurs during later VICE waves (**Fig. 4d**, ~40 per 1000 cells versus ~30 per 1000 cells at 12 hpi typically). Thus, the early RV binding robustly promotes VICE. Additionally, inhibiting endocytosis with dynole 34-2 did not alter VICE rates over time (**Fig. 4e**) or significantly reduce RV clearance at 24 hpi, compared with control infections (**Fig. 4f**). While RV localises predominantly to plasma membrane during VICE between 2-8 hpi, the few extruding infected cells observed at 24 hpi had internalised virus (by VP3 staining) **(Extended Data Fig. 4d)**, potentially by macropinocytosis[30]. Thus, epithelia use two strategies to ensure extrusion can eliminate RV infections: by activating live cell extrusion before the virus enters the cell, which is potentially mechanically activated, and by apoptotic cell extrusion in cells where RV has replicated (**Fig. 4g**).

Extruded virus-infected cells can infect other cells

While epithelia eliminate RV by VICE, the cells that extrude are laden with virus, especially at later timepoints where virus has replicated, which could be highly infectious. To test if the cells extruded during VICE could act as 'viral bombs' that could infect other cells, we collected extruded cells at 16 hpi, washed them with medium, and added them to fresh BEAS-2Bs, HeLa-H1s, and 16HBE14o- monolayers for 24 hours (**Fig 5a**). We found that extruded VICE cells could infect fresh monolayers by 6 hours post-VICE cell addition, and that this infection increased only in BEAS-2Bs and HeLa-H1 cells, so that they were nearly all infected by 24 hours (**Fig. 5b&c**). By contrast, 16HBE14o- eliminate RV by VICE by 24 hours, as expected (**Fig. 5b&c**). Notably, viral bombs/infected extruded cells harbour significantly more viral units than cell-free virus: as ~100% of BEAS-2Bs were infected with VICE cells (**Fig. 5d**), compared to only ~20% when infected with purified RV by 24 hours (**Fig. 2h**). While the viral bombs likely contain more RV than the purified virus, making it hard to compare actual MOI, our results reveal that viral bombs produced during VICE are competent to infect other cells and provide a more physiological route of virus spread than purified virus.

Discussion

Here, we found that epithelia rapidly clear rhinovirus by inducing virus-induced cell extrusion (VICE). VICE limits RV spread, providing a rapid protective response for airway epithelia without the need for other innate immune cells. We believe VICE represents an ancient, primordial, innate immune mechanism originating from many marine organisms, such as sea sponges, jellyfish, and corals, that are comprised of epithelia alone that have robust immune systems without the benefit of a blood system. We have identified two separate extrusion mechanisms that work in tandem to ensure the elimination of rhinovirus: an early endocytosis-independent wave of live extrusions, followed by a viral replication wave of apoptotic cells. Although highly efficient, this system represents a double-edged sword, eliminating infected cells but facilitating rapid dissemination of infectious virions. Additionally, VICE in excess, as may be the case in asthmatics, could damage the epithelium, leading to an inflammatory response.

Fig. 5: Extruded rhinovirus-infected cells can infect new monolayers. All infections were performed in 16HBE14o⁻ monolayers with RVA16 at an MOI of 3 unless otherwise indicated. **a**, Schematic showing the experimental design: extruded infected cells were collected from RVA16-infected monolayers and applied to naïve 16HBE14o⁻, Beas-2B, or HeLa-H1 monolayers. **b**, Confocal projections of 16HBE14o⁻, Beas-2B, and HeLa-H1 monolayers at 6 h and 24 h after exposure to harvested extruded cells. Infected cells were identified by VP3 staining (red); nuclei were counterstained with DAPI (blue). **c**, Quantification of VP3⁺ cells in 16HBE14o⁻, Beas-2B, and HeLa-H1 monolayers 24 h after exposure to harvested extruded cells. **d**, Model illustrating that RVA16-infected monolayers clear infection via extrusion, and that extruded cells can initiate infection in fresh naïve monolayers.Data are from ≥ 3 biological replicates. Transparency distinguishes independent biological experiments (**c**); bars represent the mean with s.e.m. (**c**); error bars show s.e.m. Statistical significance was determined by Kruskal–Wallis test; *** p < 0.001, **** p < 0.0001.

VICE-induced viral clearance is likely not limited to RV and likely represents the most common first step in innate immunity. Our findings support early observations by Boyde and Reed (1972), who used scanning electron

microscopy to show that bovine tracheal epithelia infected with rhinovirus sloughed off, whereas those infected with parainfluenza did not [31]. The increased severity of parainfluenza symptoms, compared to RV, may suggest that the inability to eliminate virus infections by VICE would lead to more systemic infections, needing other innate and adaptive immune responses. Conversely, the excess extrusion we noted in response to RV infections of mouse lung slices could trigger asthma attacks, as rhinovirus is the most common trigger of asthma exacerbations, whereas the lack of VICE response from parainfluenza may account for its link to milder asthma exacerbations [32,33]. Similar extrusion or 'sloughing' has been noted in response to infection with RSV [32], enteroviruses [34], and *S. Tm* bacterial infections [35,36]. Interestingly, RSV infection in infants, while common, can occasionally lead to excess sloughing of infected cells into airway lumens, causing obstruction and inflammation[32,37], and likely airway remodelling that can lead to asthma development [12]. Thus, excessive or insufficient VICE can damage the airway barrier, leading to inflammation and other immune responses to clear the virus, both of which could promote asthma, COPD, and cystic fibrosis down the line [38–40]. Thus, VICE must be tightly regulated to strike a balance between viral clearance and tissue integrity.

Our findings on VICE have added to a growing body of research revealing different pathways that activate extrusion. Here, the first rapid pathway requires mechanosensitive channels, also implicated in enterovirus-induced extrusion [34]. While we refer to this as a mechano-sensitive response, it may represent a non-specific irritant response akin to that from pollution, irritants, or allergens [41,42] that could perturb the membrane during attachment or initial cell membrane fusion, than a response to crowding, as originally identified during homeostatic live cell extrusion[10]. Both GsMTX-4 or Gd³⁺ which block Piezo-1-dependent crowding-induced extrusion, can block transient receptor channels that trigger calcium influx in response to pollutants [42] and could also act to trigger extrusion. Viruses acting in a non-specific mechanical mode may explain why VICE occurs independent of host-viral receptor type (**Fig. 1**) and virus internalisation (**Fig. 4**). This mechanism would also allow a generic response to new viruses or variants arising in the community. Because VICE clears RV before productive uncoating [43,44], it would also contain viral spread, making it a critical early antiviral defence.

Further, our findings highlight the importance of viral replication in the second, apoptotic wave of extrusion. Our experiments suggest that any cells that fail to trigger the first rapid response will endocytose the virus that can replicate, triggering the second apoptotic VICE wave. Viral replication driving virus clearance through VICE could explain the paradox of why remdesivir was not as effective as expected during the SARS CoV2 pandemic, if it also prevented elimination through VICE [23,45,46]. Additionally, because virus replication induces type I & III interferons (IFNs) that are known to trigger apoptosis, we might expect the IFN activation to control this second step [47]. Importantly, as asthma and COPD have deficient IFNs responses [48–50] a nd apoptosis [48], VICE could be instead be defective in asthma, leading to longer infections and more epithelial damage.

Finally, our finding that the extruded cells from VICE can efficiently infect other cells suggests two important points. One, extruded virus-laden cells can infect naïve monolayers efficiently, suggesting a physiologically relevant transmission route whereby extrusion acts both as a host-protective mechanism and a vehicle for viral dissemination. Presumably, extruded VICE cells can leave one epithelia region where infection starts to infect another area or individual. Two, we were struck by the fact that while these viral bombs could infect new cells, the infected 16HBE14o⁻ monolayers do not appear to become reinfected from the viral bombs floating in their medium. This suggests a form of learned epithelia-specific innate immune response, which will need further investigation. Practically, this aligns with what is known about human infections, which generally migrate from our throats to our nasal passages. Importantly, the same epithelia do not repetitively experience reinfection or they would rapidly disintegrate into a massive wound with every infection. Identifying such a mechanism could lead to new therapeutic avenues aimed at preventing viral infections altogether, rather than merely managing their symptoms.

Clearly, understanding this ancient, yet emerging form of innate immunity will be of great importance to our management of virus spread and symptoms. We believe that using cell lines that do not model tissues viruses normally encounter has led to oversight of this fundamental, primary step in innate immunity. Without understanding more about the mechanisms and signalling driving VICE, we will be stuck managing only later, downstream symptoms from viral infections, instead of preventing them.

ACKNOWLEDGEMENTS

We thank Michael J. Redd (University College London) for assistance with scanning confocal imaging, Pippa Hawes (The Francis Crick Institute) and Rui Galao Pedro for reviewing the manuscript, and members of the Rosenblatt laboratory for help with experimental design and figure layouts feedback.

FUNDING

The Wellcome Investigator Award 221908/Z/20/Z and an Academy of Medical Sciences Professorship (APR2\1007) to J.R, a Darwin Trust Studentship to F.F, and a National Institute of Health Research (NIHR) Biomedical Research Centre funding scheme, Medical Research Council, Asthma UK Centre grant G1000758, and Asthma UK grant CH11SJ to SLJ.

AUTHOR CONTRIBUTIONS

J.R., F.F. and D.C.B. conceived the study and designed the experiments. F.F. performed all experiments and data analyses. F.F. and J.R. interpreted the data and wrote the manuscript. All authors edited and approved the final manuscript.

CONFLICT OF INTERESTS

All authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY

All data are available in the main text or the extended data. The source material link will be provided upon request.

Extended Data Figure Legends

Extended Data SMovie1: Time-lapse imaging of mock-infected 16HBE14o⁻ monolayers. Representative time-lapse sequence of mock-infected 16HBE14o⁻ monolayers imaged for 12 hours showing few extrusion events. Scale bar, 25 μm.

Extended Data SMovie2: Time-lapse imaging of RVA16-infected 16HBE14o⁻ monolayers. Representative time-lapse sequence of RVA16-infected 16HBE14o⁻ monolayers imaged for 12 hours, showing frequent extrusion events. Scale bar, 25 μm.

Extended Data SMovie3: Time-lapse imaging of RVA16-infected 16HBE14o⁻ monolayers treated with JTE-013. Representative time-lapse sequence of RVA16-infected 16HBE14o⁻ monolayers treated with the $S1P_2$ receptor antagonist JTE-013, imaged for 12 hours showing extrusion events relative to untreated infected monolayers. Scale bar, 25 μm.

Methods

Table 1: List of reagents and chemicals

Cell culture growth medium			
Cell/Tissue	**Growth medium**	**Catalogue Number**	**Supplier**
HeLa-H1	Dulbecco's modified Eagle Medium (DMEM)	D6429	Sigma
Beas-2B	Dulbecco's modified Eagle Medium DMEM	31966-024	GIBCO
16HBE14o⁻	Minimum Essential Medium (MEM)	31095-029	GIBCO

List of medium/buffer recipes		
Buffer/Medium	**Recipe**	**Cell line**
HBE growth medium	MEM supplemented with 10% FBS, 1% P/S	16HBE14o⁻
Beas-2B growth medium	DMEM-gibco supplemented with 10% FBS, 1% P/S).	Hela-H1
HeLa growth medium	DMEM-sigma supplemented with 10% FBS, 2% 1M HEPES, 1% Glutamax and 1% $NaHCO_3$.	Beas-2B
infection medium	Growth medium supplemented with 2% FBS, 2% 1M HEPES, 1% Glutamax and 1% $NaHCO_3$	Various
IF blocking buffer	(2% bovine serum albumin (BSA), 0.1% Triton X-100, 0.1% sodium azide	Various
Freezing medium	10% DMSO and 90% FBS	Various

List of antibodies					
Protein/Epitope	**Antibody Type**	**Host**	**Catalog Number**	**Supplier**	**Dilution**
VP3	Primary	Mouse	MA5-18250	Invitrogen	1:100
dsRNA	Primary	Mouse	MABE1134	Sigma	1:100
MDA5	Primary	Rabbit	NBP1-76760	Novus	1:100
Active-caspase 3	Primary	Rabbit	9669S	Cell Signalling	1:100
Piezo1	Primary	Rabbit	NBP1-78537	Novus	1:200
E-cadherin	Primary	Rabbit	3195S	Cell Signalling	1:200
ZO-1	Primary	Rabbit	617300	Invitrogen	1:200
Phalloidin 488	Secondary	N/A	A12379	Invitrogen	1:200
Phalloidin 568	Secondary	N/A	A12380	Invitrogen	1:200
Phalloidin 647	Secondary	N/A	A22287	Invitrogen	1:200
488 anti-mouse	Secondary	Goat	A32723	Invitrogen	1:200
568 anti-rabbit	Secondary	Goat	A11011	Invitrogen	1:200
568 anti-mouse	Secondary	Goat	A11004	Invitrogen	1:200
647 anti-rabbit	Secondary	Goat	A32733	Invitrogen	1:200
647 anti-rabbit	Secondary	Goat	A21235	Invitrogen	1:200
DAPI-4′,6-diamidino-2-phenylindole	N/A	N/A	D1306	Invitrogen	1:1000

List of reagents		
Reagent	**Catalogue number**	**Supplier**
MEM	31095-029	GIBCO
DMEM-sigma	D6429	Sigma
DMEM-gibco	31966-024	GIBCO
Foetal Bovine Serum (FBS)	10270106	GIBCO
Penicillin-streptomycin (P/S)	11580626	GIBCO
Dulbecco's phosphate-buffered saline (DPBS)	14190250	GIBCO
Trypsin	25300-096	GIBCO
Hanks' Balanced Salt Solution (HBSS)	14025	GIBCO
PFA	047317.9M	Thermo Fisher
Egtazic acid (EGTA)	324626	MERCK
Triton X-100	X100RS	Sigma
BSA	A9418	Sigma
Sodium Azide	S8032	Sigma
Pro Long Gold	P36930	Thermo Fisher
dNTP	DNTP100-1KT	Sigma
Random Hexamer	SO142	Thermo Fisher
Ribolock RNAse Inhibitor	EO0382	Thermo Fisher
Revertaid H Minus RT	EP0452	Thermo Fisher
TriZol	15596026	Thermo Fisher
Glycoblue	AM9516	Thermo Fisher
Luna Buffer	M3004E	New England Labs
Chloroform	C/920/15	Fisher Scientific
Dynole	AB120463	Abcam
Dimethyl sulfoxide	276855	MERCK
Low melt agarose	Bp1360-100	Sigma
Rat collagen	C3867	Sigma
Trypan Blue	T8154-100ML	MERCK
Nail varnish	N/A	Boots
Cell counting kit-8 (CCK-8)	96992	MERCK
MTT assay growth kit	CT02	MERCK

List of drugs or inhibitors used and in RVA16 infection				
Drug/Inhibitor	Concentration	Target	Catalogue Number	Supplier
JTE-013	10 μM	S1P2 receptor		
GsMtx4	50 μM	Piezo1, TRPC1, and TRPC6	J4080	Sigma
z-VAD-FMK	50 μM	Pan-caspase inhibitor (apoptosis)	V116-2MG	MERCK
Remdesivir	25 μM	Viral replication	AB120463-1001	Stratech
GdCl$_3$	10 μM	Stretch-activated channels	G7532	Sigma
Dynole	10 μM	Dynamin-dependent endocytosis	AB120463-1001	ABCAM

Reaction components used for cDNA synthesis.	
Reagent	Volume/reaction
10 mM Random hexamer primers	0.5 μL
5X Buffer RT	2 μL
dNTP mix (10mM each dNTP)	1 μL
40U/μL Ribolose RNAse inhibitor	0.25 μL
200U/μL Revertaid H minus reverse transcriptase	0.5 μL
300ng of RNA Template	varies
Total reaction volume	10 μL

Primers and probes used for quantitative PCR	
Gene	Primer/Probe/set
Rhinovirus forward (900 nM)	GTGAAGAGCCSCRTGTGCT
Rhinovirus reverse (900 nM)	GCTSCAGGGTTAAGGTTAGCC
Rhinovirus probe	TGAGTCCTCCGGCCCCTGAATG
GAPDH	Hs02786624_g1
SDHA	HK-DD-hu-900

Cell culture

16HBE14o (Merck, SCC150), HeLa-H1 (ATCC, CRL-1958) and BEAS-2Bs were grown in respective growth medium (**Table 1**) supplemented with 10% foetal bovine serum (FBS; 10270106) and 1% penicillin/ streptomycin (P/S; 15070063) in a cell culture incubator at 37°C with 5% CO_2. For all experiments, cells were seeded in growth medium and cultured until confluent. At confluency, the growth medium was removed, cells were washed twice with infection medium (**Table 1**) and incubated overnight in the same medium. All cell lines tested negative for mycoplasma contamination in periodic tests (Sartorius, 20-700-20).

Rhinovirus stock production and titration

Viral stocks of RVA01B, RVA02, RVB14, and RVA16 (all from ATCC) were generated in HeLa-H1 cells and their titres were determined by measuring 50% tissue culture infectious dose ($TCID_{50}$), as described previously [51]. Briefly, cells at ~90% confluency were washed twice with DMEM infection medium and inoculated with the virus diluted in infection medium. After 1 hour of gentle agitation at room temperature, cultures were incubated at 37 °C, 5% CO_2 until ~90% cytopathic effect (CPE) was observed. Cells were then dislodged, subjected to two freeze–thaw cycles at –80 °C, and clarified by centrifugation (4000 rpm, 15 min, 4 °C). The supernatant was filtered (0.2 μm; catalogue no. S2GPU01RE) and stored in aliquots at -80 °C.

For $TCID_{50}$ determination, HeLa-H1 cells were seeded at 0.5×10^5 cells/mL (150 μL/well, 96-well plate). Undiluted virus or 10-fold serial dilutions were added, and plates were incubated for 4-5 days at 37 °C, 5% CO_2. CPE was scored microscopically, and $TCID_{50}$/mL values were calculated using the Spearman–Kärber method [52].

Rhinovirus infection of cell cultures

Cells were seeded at equal densities into 4- or 8-well IBIDI chambers or 24-well plates with size-1.5 coverslips and cultured at 37 °C with 5% CO_2 until ~95% confluency, followed by serum-starvation overnight in medium containing 2% FBS.

Drug cytotoxicity was evaluated using MTT (Merck, CT02) or CCK-8 (Merck, 96992) assays, according to the manufacturer's instructions. For experiments with Remdesivir or Dynole 34-2, cells were pre-incubated with the drug for 2 h at 37 °C with 5% CO_2 before infection. Viral stocks were diluted in infection medium to achieve a multiplicity of infection (MOI) of 3, with inoculum volumes adjusted for each format (150 μL for 24-well plates, 150 μL for 8-well chambers, 500 μL for 6-well plates, 200 μL for 4-well chambers). For drug treatments, the inoculum contained the drug at the corresponding concentration. Cells were inoculated for 1 h at room temperature with gentle agitation, washed three times with growth medium, and maintained at 37 °C, 5% CO_2 in fresh medium with or without drug.

Cultures were harvested at 0–96 hpi for downstream applications, including ELISA from supernatants, immunofluorescence (IF) from coverslips, or protein/RNA extraction. IBIDI chambers or glass-bottom 24-well plates were used for time-lapse imaging.

Disruption of cell junctions
Calcium free medium

Cells were seeded at equal densities and cultured in HBE growth medium without calcium (MEM, no calcium; Thermo Fisher Scientific, 11380037) supplemented with 2% Glutamax, 10% FBS, and 1% P/S for 36 h. The medium was then replaced with calcium-free HBE infection medium and incubated overnight. Infection was performed at MOI 3 in calcium-free medium for 1 h at room temperature with gentle agitation. Cells were washed three times and cultured in calcium-free growth medium at 37 °C, 5% CO_2 until harvest.

Egtazic acid (EGTA) treatment

Cells were grown to ~90% confluency and serum-starved overnight in infection medium. The following day, cells were pre-incubated in infection medium containing 1 mM EGTA. Infection was performed at MOI 3 in EGTA-supplemented infection medium for 1 hour at room temperature with gentle agitation. After inoculum removal, cells were washed three times and maintained in growth medium containing 1 mM EGTA at 37 °C with 5% CO_2 until harvest.

Cells were grown to ~90% confluency and serum-starved overnight in infection medium. The next day, cells were pre-incubated in infection medium containing 1 mM EGTA, then inoculated at an MOI of 3 in EGTA-supplemented medium for 1 h at room temperature with gentle agitation. Following inoculum removal, cells were washed three times and cultured in growth medium containing 1 mM EGTA at 37 °C, 5% CO_2 until harvest.

Immunofluorescence staining
Cells grown on coverslips were fixed with 4% paraformaldehyde (PFA) at designated time points and rinsed three times with DPBS. After blocking for 1 h at room temperature in IF blocking buffer (**Table**), cells were incubated overnight at 4 °C with primary antibodies (**Table 1**). Following three rinses and three 10-min washes in DPBS, cells were incubated with secondary antibodies (Table) for 2 h at room temperature. Nuclei were counterstained with DAPI (1:1000, v/v) for 10 min, washed, mounted in ProLong Gold (**Table 1**), and cured for 24 h before imaging.

For PCLSs, the procedure was identical, except secondary antibodies were incubated overnight at 4 °C, and washes were extended to three 30-min steps.

Harvesting extruded cells
16HBE14o cells infected with RVA16 were processed for recovery of extruded cells. Culture medium was collected into a 15 mL Falcon tube, and the remaining monolayer was treated with 500 μL trypsin at room temperature for ~2 min, with detachment monitored by light microscopy. Trypsinisation was quenched with 2 mL HBE infection medium, and detached cells were pooled with the collected medium. Suspensions were centrifuged at 1200 rpm for 5 min at 4 °C to pellet extruded cells, which were washed three times and resuspended in infection medium.

Microscopy
Live-cell phase contrast and immunofluorescence imaging were performed on a Nikon Eclipse Ti2 microscope. For time-lapse phase imaging, a Plan Fluor 20× Ph1 DLL NA 0.50 objective, a Photometrics Iris 15 16-bit camera, and a CoolLED pE-4000 light source were used, controlled by NIS-Elements software (Nikon, v5.30.02). For immunofluorescence imaging, Plan Fluor 20×, 40× water-immersion, or 60× NA 1.40 oil-immersion objectives were used in combination with an iXon 888 Andor 16-bit camera, a Yokogawa CSU-W1 spinning disk confocal unit, and Toptica Photonics laser excitation, controlled by NIS-Elements (Nikon, v5.21.03).

RNA extraction and cDNA synthesis

RNA from extruded or monolayer cells was isolated using TRIzol Reagent (Thermo Fisher Scientific) with minor modifications. Cells were washed with DPBS and lysed in 250 μL TRIzol per 1×10^5-10^7 cells. Chloroform (200 μL per 1 mL TRIzol) was added, and phase separation was performed by centrifugation (10,000 rpm, 30 min, 4 °C). The aqueous phase was transferred to tubes containing 1 μL glycogen blue (30 μg/mL), and RNA was precipitated with 0.5 vol isopropanol at -80 °C overnight. Pellets were recovered by centrifugation (10,000 rpm, 60 min, 4 °C), washed twice with 75% ethanol, air-dried, and resuspended in 20-50 μL nuclease-free water. RNA concentration and purity were assessed by NanoDrop spectrophotometry, and samples were stored at -80 °C.

cDNA was synthesised from total RNA using the reagents and volumes listed in **Table 1**. Reaction mixtures were incubated at 25 °C for 10 min, 42 °C for 60 min, and 70 °C for 10 min in a Bio-Rad T100 thermal cycler. cDNA was diluted with RNase-free water and stored at -20 °C.

Quantitative PCR (qPCR)

qPCR was performed in 10 μL reactions containing 1 μL cDNA, using the ViiA7 RUO Thermal Cycler (Thermo Fisher Scientific) with specific probes and primers (**Table 1**). Cycling conditions were initial denaturation at 95 °C for 1 min, followed by 40 cycles of 95 °C for 15 s and 60 °C for 30 s. Samples were run in duplicate, and amplification plots were used to confirm quantification within the exponential phase. Cycle threshold (Ct) values were averaged, with thresholds set within the linear lower third of each amplification curve. Target gene expression was normalised to the housekeeping genes GAPDH and SDHA, and presented as normalised RV copies.

RVA01B infection of Precision Cut Lung Slices (PCLSs)

PCLSs were infected with RVA01B using a high virus titre of 5.5×10^7 PFU/mL, corresponding to 1.65×10^7 infectious units in the 300 μL inoculum applied per slice.

Data quantification

Cellular events, including extrusions and total cell numbers, were quantified using NIS-Elements software (Nikon, v5.21.03). For time-lapse imaging, frames corresponding to 2 h intervals were analysed. Extruding cells were distinguished from dividing cells by the absence of daughter cell formation and by their floating morphology tethered to the monolayer. Cell counts were obtained using GA Analysis in NIS-Elements, with thresholds for brightness and size applied automatically and verified visually.

For fixed immunostained samples, quantification was performed by Z-stack inspection in NIS-Elements. Extruded cells were identified by actin rings at the basal plane and nuclei positioned above the monolayer. Nuclei were segmented automatically using GA Analysis, with segmentation accuracy confirmed visually. Images were further processed using Fiji (ImageJ).

Image and statistical analysis

Image presentation and statistical analyses were performed in GraphPad Prism (v10.3.1). Data represent the mean ± SEM from at least three independent biological replicates (n = 3). Statistical differences were evaluated using independent t-tests or one-way ANOVA with Dunnett's or Tukey's post-hoc tests, as specified in figure legends.

References

1. Jacobs, S. E., Lamson, D. M., St. George, K. & Walsh, T. J. Human Rhinoviruses. *Clin Microbiol Rev* **26**, 135–162 (2013).

2. Ganjian, H., Rajput, C., Elzoheiry, M. & Sajjan, U. Rhinovirus and Innate Immune Function of Airway Epithelium. *Front. Cell. Infect. Microbiol.* **10**, 277 (2020).

3. Esneau, C., Bartlett, N. & Bochkov, Y. A. Rhinovirus structure, replication, and classification. in *Rhinovirus*

Infections 1–23 (Elsevier, 2019). doi:10.1016/B978-0-12-816417-4.00001-9.

4. Blaas, D. & Fuchs, R. Mechanism of human rhinovirus infections. *Mol Cell Pediatr* **3**, 21 (2016).

5. Eisenhoffer, G. T. & Rosenblatt, J. Bringing balance by force: live cell extrusion controls epithelial cell numbers. *Trends in Cell Biology* **23**, 185–192 (2013).

6. Ohsawa, S., Vaughen, J. & Igaki, T. Cell Extrusion: A Stress-Responsive Force for Good or Evil in Epithelial Homeostasis. *Developmental Cell* **44**, 284–296 (2018).

7. Rosenblatt, J., Raff, M. C. & Cramer, L. P. An epithelial cell destined for apoptosis signals its neighbors to extrude it by an actin- and myosin-dependent mechanism. *Current Biology* **11**, 1847–1857 (2001).

8. Slattum, G., Gu, Y., Sabbadini, R. & Rosenblatt, J. Autophagy in Oncogenic K-Ras Promotes Basal Extrusion of Epithelial Cells by Degrading S1P. *Current Biology* **24**, 19–28 (2014).

9. Gu, Y., Forostyan, T., Sabbadini, R. & Rosenblatt, J. Epithelial cell extrusion requires the sphingosine-1-phosphate receptor 2 pathway. *Journal of Cell Biology* **193**, 667–676 (2011).

10. Eisenhoffer, G. T. *et al.* Crowding induces live cell extrusion to maintain homeostatic cell numbers in epithelia. *Nature* **484**, 546–549 (2012).

11. Slattum, G., McGee, K. M. & Rosenblatt, J. P115 RhoGEF and microtubules decide the direction apoptotic cells extrude from an epithelium. *Journal of Cell Biology* **186**, 693–702 (2009).

12. Bagley, D. C. *et al.* Bronchoconstriction damages airway epithelia by crowding-induced excess cell extrusion. *Science* **384**, 66–73 (2024).

13. Reddel, R. R. *et al.* Transformation of Human Bronchial Epithelial Cells by Infection with SV40 or Adenovirus-12 SV40 Hybrid Virus, or Transfection via Strontium Phosphate Coprecipitation with a Plasmid Containing SV40 Early Region Genes. *Cancer Research* **48**, 1904–1909 (1988).

14. Han, X., Na, T., Wu, T. & Yuan, B.-Z. Human lung epithelial BEAS-2B cells exhibit characteristics of mesenchymal stem cells. *PLoS ONE* **15**, e0227174 (2020).

15. Gudipaty, S. A. & Rosenblatt, J. Epithelial cell extrusion: Pathways and pathologies. *Seminars in Cell & Developmental Biology* **67**, 132–140 (2017).

16. Fadul, J. & Rosenblatt, J. The forces and fates of extruding cells. *Current Opinion in Cell Biology* **54**, 66–71 (2018).

17. Gu, Y. *et al.* Defective apical extrusion signaling contributes to aggressive tumor hallmarks. *eLife* **4**, e04069 (2015).

18. Andrade, D. & Rosenblatt, J. Apoptotic regulation of epithelial cellular extrusion. *Apoptosis* **16**, 491–501 (2011).

19. Deszcz, L., Gaudernak, E., Kuechler, E. & Seipelt, J. Apoptotic events induced by human rhinovirus infection. *Journal of General Virology* **86**, 1379–1389 (2005).

20. Slee, E. A. *et al.* Benzyloxycarbonyl-Val-Ala-Asp (OMe) fluoromethylketone (Z-VAD.FMK) inhibits apoptosis by blocking the processing of CPP32. *Biochemical Journal* **315**, 21–24 (1996).

21. Malin, J. J., Suárez, I., Priesner, V., Fätkenheuer, G. & Rybniker, J. Remdesivir against COVID-19 and Other

Viral Diseases. *Clin Microbiol Rev* **34**, e00162-20 (2020).

22. Simonis, A., Theobald, S. J., Fätkenheuer, G., Rybniker, J. & Malin, J. J. A comparative analysis of remdesivir and other repurposed antivirals against SARS-CoV2. *EMBO Mol Med* **13**, e13105 (2021).

23. Mozaffari, E. *et al.* Remdesivir is Associated with Reduced Mortality in Patients Hospitalized for COVID-19 Not Requiring Supplemental Oxygen. *Open Forum Infectious Diseases* **11**, ofae202 (2024).

24. Bae, C., Sachs, F. & Gottlieb, P. A. The Mechanosensitive Ion Channel Piezo1 Is Inhibited by the Peptide GsMTx4. *Biochemistry* **50**, 6295–6300 (2011).

25. Suchyna, T. M. *et al.* Identification of a Peptide Toxin from *Grammostola spatulata* Spider Venom That Blocks Cation-Selective Stretch-Activated Channels. *The Journal of General Physiology* **115**, 583–598 (2000).

26. Tadala, L. *et al.* Infection-induced membrane ruffling initiates danger and immune signaling via the mechanosensor PIEZO1. *Cell Reports* **40**, 111173 (2022).

27. Fuchs, R. & Blaas, D. Productive Entry Pathways of Human Rhinoviruses. *Advances in Virology* **2012**, 1–13 (2012).

28. ABCAM. *Dynole® 34-2, Dynamin I and Dynamin II Inhibitor*. https://www.abcam.com/en-us/products/biochemicals/dynole-34-2-dynamin-i-and-dynamin-ii-inhibitor-ab120463?srsltid=AfmBOopqI8onEM49HUZI2DXq4W3oDzw3qa9ZFiqaHoaBRnrZ9P7aT_IN#.

29. Hill, T. A. *et al.* Inhibition of Dynamin Mediated Endocytosis by the *Dynoles* —Synthesis and Functional Activity of a Family of Indoles. *J. Med. Chem.* **52**, 3762–3773 (2009).

30. Gundu, C. *et al.* Dynamin-Independent Mechanisms of Endocytosis and Receptor Trafficking. *Cells* **11**, 2557 (2022).

31. Reed, S. E. & Boyde, A. Organ cultures of respiratory epithelium infected with rhinovirus or parainfluenza virus studied in a scanning electron microscope. *Infection and Immunity* **6**, 68–76 (1972).

32. Liesman, R. M. *et al.* RSV-encoded NS2 promotes epithelial cell shedding and distal airway obstruction. *J. Clin. Invest.* **124**, 2219–2233 (2014).

33. Lee, W. S. *et al.* The Association Between Respiratory Viruses and Asthma Exacerbation in Children Visiting Pediatric Emergency Department: A Retrospective Cohort Study. *JCM* **14**, 1311 (2025).

34. Moshiri, J., Craven, A. R., Mixon, S. B., Amieva, M. R. & Kirkegaard, K. Mechanosensitive extrusion of Enterovirus A71-infected cells from colonic organoids. *Nat Microbiol* **8**, 629–639 (2023).

35. Knodler, L. A. *et al.* Dissemination of invasive *Salmonella* via bacterial-induced extrusion of mucosal epithelia. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 17733–17738 (2010).

36. Bastounis, E. E. *et al.* Mechanical competition triggered by innate immune signaling drives the collective extrusion of bacterially infected epithelial cells. *Developmental Cell* **56**, 443-460.e11 (2021).

37. Gudipaty, S. A. & Rosenblatt, J. Epithelial cell extrusion: Pathways and pathologies. *Seminars in Cell and Developmental Biology* **67**, 132–140 (2017).

38. Gern, J. E. Rhinoviruses and the onset of asthma. in *Rhinovirus Infections* 121–136 (Elsevier, 2019). doi:10.1016/B978-0-12-816417-4.00005-6.

39. Kelly, J. T. & Busse, W. W. Host immune responses to rhinovirus: Mechanisms in asthma. *Journal of Allergy and Clinical Immunology* **122**, 671–682 (2008).

40. Hershenson, M. B. Rhinovirus-Induced Exacerbations of Asthma and COPD. *Scientifica* **2013**, 1–12 (2013).

41. Grace, M. S., Baxter, M., Dubuis, E., Birrell, M. A. & Belvisi, M. G. Transient receptor potential (TRP) channels in the airway: role in airway disease. *British Journal of Pharmacology* **171**, 2593–2607 (2014).

42. Deering-Rice, C. E. *et al.* Characterization of Transient Receptor Potential Vanilloid-1 (TRPV1) Variant Activation by Coal Fly Ash Particles and Associations with Altered Transient Receptor Potential Ankyrin-1 (TRPA1) Expression and Asthma. *Journal of Biological Chemistry* **291**, 24866–24879 (2016).

43. Conzemius, R., Ganjian, H., Blaas, D. & Fuchs, R. ICAM-1 Binding Rhinoviruses A89 and B14 Uncoat in Different Endosomal Compartments. *J Virol* **90**, 7934–7942 (2016).

44. Ganjian, H., Zietz, C., Mechtcheriakova, D., Blaas, D. & Fuchs, R. ICAM-1 Binding Rhinoviruses Enter HeLa Cells via Multiple Pathways and Travel to Distinct Intracellular Compartments for Uncoating. *Viruses* **9**, 68 (2017).

45. Grundeis, F. *et al.* Remdesivir for the treatment of COVID-19. *Cochrane Database of Systematic Reviews* **2023**, (2023).

46. Veronese, N. *et al.* Real life experience on the use of Remdesivir in patients admitted to COVID-19 in two referral Italian hospital: a propensity score matched analysis. *Sci Rep* **14**, 9303 (2024).

47. Balachandran, S. *et al.* Alpha/Beta Interferons Potentiate Virus-Induced Apoptosis through Activation of the FADD/Caspase-8 Death Signaling Pathway. *J Virol* **74**, 1513–1523 (2000).

48. Wark, P. A. B. *et al.* Asthmatic bronchial epithelial cells have a deficient innate immune response to infection with rhinovirus. *The Journal of Experimental Medicine* **201**, 937–947 (2005).

49. Mallia, P. *et al.* Experimental Rhinovirus Infection as a Human Model of Chronic Obstructive Pulmonary Disease Exacerbation. *Am J Respir Crit Care Med* **183**, 734–742 (2011).

50. Contoli, M. *et al.* Role of deficient type III interferon-λ production in asthma exacerbations. *Nat Med* **12**, 1023–1026 (2006).

51. Papi, A. & Johnston, S. L. Rhinovirus Infection Induces Expression of Its Own Receptor Intercellular Adhesion Molecule 1 (ICAM-1) via Increased NF-κB-mediated Transcription. *Journal of Biological Chemistry* **274**, 9707–9720 (1999).

52. Lei, C., Yang, J., Hu, J. & Sun, X. On the Calculation of TCID50 for Quantitation of Virus Infectivity. *Virologica Sinica* **36**, 141–144 (2021).