Soil trace gas oxidizers divergently respond to short- and long-term warming

Thanh Nguyen-Dinh[1,#], Vojtěch Tláskal[2,#] Pok Man Leung[1,3], Magdalena Wutkowska[2],

Justus Amuche Nweze[2,4,5], Katri Ylä-Soininmäki[6], Johanna Kerttula[6], Mathilde Borg

Dahl[7], Andrea Söllinger[8], Tilman Schmider[8], Lukas Kohl[6], Christina Biasi[6,9], Andreas

Richter[10], Tim Urich[7], Alexander T. Tveit[8], Chris Greening[1,3], Bjarni D. Sigurdsson[11],

Anne Daebeler[2]*

1 - Department of Microbiology, Biomedicine Discovery Institute, Monash University,

Clayton, VIC 3800, Australia.

2 - Institute of Soil Biology and Biogeochemistry, Biology Centre CAS, České

Budějovice, Czech Republic.

3 - Securing Antarctica's Environmental Future, Monash University, Clayton, VIC 3800,

Australia.

4 - Department of Ecosystem Biology, Faculty of Science, University of South Bohemia,

České Budějovice, Czech Republic.

5 - Department of Science Laboratory Technology, Faculty of Physical Sciences,

University of Nigeria, Nsukka, Nigeria.

6 - Department of Environmental and Biological Sciences, University of Eastern Finland,

Kuopio, Finland.

7 - Institute of Microbiology, University of Greifswald, Greifswald, Germany.

8 - Department of Arctic and Marine Biology, UiT The Arctic University of Norway,

Tromsø, Norway.

9 - Institute of Ecology, University of Innsbruck, Innsbruck, Austria

10 - Centre for Microbiology and Environmental Systems Science, University of Vienna,

Vienna, Austria.

11 - Faculty of Environmental and Agricultural Sciences, Agricultural University of

Iceland, Reykjavík, Iceland

# These authors contributed equally to the study

*Correspondence: Anne Daebeler, anne.daebeler@bc.cas.cz

Keywords: climate change, global warming, greenhouse gases, grasslands

Abstract

The upland soil microbiome is dominated by aerobic bacteria that oxidize atmospheric

trace gases, including CO, H_2, and CH_4. As a result, soils are the largest biological sink

for these climate-active gases. Whether global warming will enhance or suppress these

processes remains unclear. Here, we studied the warming responses of soil trace gas

oxidizers by profiling natural geothermal gradients in a subarctic grassland with over 60

years of field warming at +6°C. We integrate field flux measurements, *ex situ*

biogeochemical assays, metagenomics, and metatranscriptomics to determine

ecosystem and cellular-level responses. Our results show that the oxidation of

atmospheric CO and H_2, but not CH_4, increased with long-term warming due to higher

cell numbers. However, at the cellular level, trace gas oxidizers, especially

methanotrophs, tended to reduce gas consumption and transcription of gas-

metabolizing enzymes in response to long-term warming. Our findings suggest that soils

may remain a robust sink for trace gases despite lower per-cell activity. This work

establishes a framework for interpreting the relationships between temperature and

microbial trace gas oxidation on timescales relevant to Earth's climate system.

Introduction

Carbon monoxide (CO), hydrogen (H_2), and methane (CH_4) are the most abundant

climate-forcing reduced gases in Earth's atmosphere. CO and H_2 are indirect

greenhouse gases owing to their reactivity with atmospheric hydroxyl radicals; they

have a 100-year global warming potential equivalent to 2 and 12 times that of carbon

dioxide (CO_2), respectively (1, 2). CH_4 is the second most abundant greenhouse gas

and has a 30-fold stronger warming potential than CO2 on a 100-year timescale (3).

Anthropogenic activities have increased emissions of all three gases since

industrialization, accelerating global warming (4–6). Soil microorganisms play a key role

in regulating the atmospheric budgets of these gases, acting as both sources and sinks.

They constitute the single most important biological sink through microbial oxidation,

contributing with 70–80%, 10–20%, and 3–5% of the annual atmospheric removal of H_2

(7–9), CO (10, 11), and CH_4 (12–14), respectively. While this microbial sink has helped

to keep CO and H_2 levels in the atmosphere relatively constant [mixing ratios of 0.09

ppmv for CO (11) and 0.53 ppmv for H_2 (9)], the relatively weak sink for CH_4 has not

effectively counterbalanced emissions and atmospheric levels have increased almost

threefold (to 1.91 ppmv in 2021) compared to preindustrial concentrations (5). The

microbial trace gas sink, therefore, provides a significant, albeit not complete, mitigation

of the climate impact of these gases.

Trace gas oxidizers are taxonomically diverse and abundant inhabitants of different

soils across the globe, with an average of 56, 39, and 1% of the total soil microbial

community predicted to use atmospheric CO, H_2, and CH_4 as energy sources,

respectively (15, 16). They utilize specialized enzymes, specifically form I [MoCu]-CO

dehydrogenases (17), high-affinity [NiFe]-hydrogenases (18), and particulate CH_4

monooxygenases (19), to extract energy from the ubiquitous yet diluted atmospheric

substrates. Atmospheric CO and H_2 primarily provide microbes an important continuous

energy supply to persist under energy limitation (20–22); in-depth studies have revealed

that microbes upregulate expression of high-affinity hydrogenases and CO

dehydrogenases in direct response to organic carbon deprivation, and that these

enzymes greatly enhance survival during starvation (23–26). Most trace gas oxidizers

use trace gases at atmospheric concentrations to support mixotrophic growth with

organic or alternative energy sources (22, 27, 28). Remarkably, some oligotrophic

bacteria are even able to grow on atmospheric energy sources alone, as recently

demonstrated for several atmospheric methanotrophs (27, 29). Together, their activities

are responsible for a net annual uptake of 250 Tg of CO, 60 Tg of H_2, and 31 Tg of CH_4

from the atmosphere [(30) and references therein; (14) for CH_4].

Despite the abundant and active nature of soil microbes in regulating atmospheric trace

gas levels, the impact of climate change on this important soil ecosystem service is

unknown. Global warming is one of the largest threats to crucial soil functions mediated

by microorganisms, including organic carbon storage (31, 32). Previous studies have

shown significant and persistent soil organic carbon (SOC) depletion after warming of

≤6°C (33–36). This SOC loss is driven by warming-induced acceleration of microbial

activity, but is stabilised by a decline in microbial biomass in the long term as the

microbial biomass equilibrates with the depleted SOC stores (33, 35, 37). The new long-

term steady state of topsoil in a warmer world is therefore predicted to be characterized

by reduced microbial biomass, species richness, and SOC stocks (35). The shifts in

microbial community structure of SOC decomposers and in resource availability may

also influence the abundance and activity of trace gas oxidizers, yet this link remains

poorly understood. Very little research is available to predict if the soil sink for climate-

active trace gases will be maintained, decline, or even increase in a future climate (38,

39). Any change in this regard is likely to have far-reaching climatic, ecological, and

biogeochemical ramifications, as outlined above. This is especially crucial as emission

of climate-related gases are projected to increase, for example due to the transition to a

hydrogen economy (fugitive H_2 emissions) (40), intensifying wildfires (CO release from

incomplete combustion) (41), positive feedback dynamics from wetlands (42), and

growing agricultural demand (CH_4 emissions from ruminants and rice paddies) (43).

Moreover, a depletion of the chemical sink for trace gases (hydroxyl radicals) has been

inferred to cause increased levels of atmospheric CH_4 (44).

Here, we leveraged an *in situ*, natural warming study site (> 60 years) (45) to determine

the response of soil CO, H_2, and CH_4 oxidizers to long-term warming. The study site

harnesses the natural geothermal heat from the Mid-Atlantic Ridge, which is transmitted

to the bedrock beneath a subarctic grassland, creating temperature gradients ranging

from unwarmed to +6°C and above. This represents an unparalleled research site to

assess long-term microbial responses of distinct functional groups and of the entire

ecosystem in a region that holds substantial carbon stocks and is vulnerable to rapid

temperature change (46). In this study, we systematically assessed the response of

trace CO, H_2, and CH_4 oxidizers to both short-term (one month) and long-term

(decades) soil warming. We quantified the fluxes of climate-active atmospheric gases in

the field and used metagenomics and metatranscriptomics to characterize the present

and active trace gas oxidizers in unwarmed and long-term warmed soil. We then used

the same soil for a laboratory experiment to quantify the potential soil uptake rates of

trace levels of CO, H_2, and CH_4 in unwarmed, short-term warmed (with only days of

warming in the lab), and long-term warmed soil (with decades of warming *in situ*). We

hypothesized that i) trace gas oxidizers will show increased oxidation rates in both the

short- and the long-term warming condition due to accelerated activity and as a

response to decreased SOC availability, respectively; ii) trace gas oxidizers will

increase in relative abundance in the long-term warmed soil, as they are able to sustain

themselves better under SOC depletion caused by warming than microbes not able to

utilize trace gases; and iii) at the ecosystem level, warmed soil will show an increased

sink capacity for CO, H_2, and CH4 in both the short- and the long-term.

Results

Long-term warming enhances the CO and H_2 but not the CH_4 sink

Results

Long-term warming enhances the CO and H_2 but not the CH_4 sink

We profiled four replicated temperature transects, comparing unwarmed and long-term

warmed soil (6°C above ambient, Fig. S1) for differences in key microbial (cell count,

microbial carbon, microbial nitrogen) and physico-chemical (pH, moisture, dissolved

organic carbon, total dissolved nitrogen) parameters (Fig. S2, Table S1). The number of

bacteria and archaea, as determined by 16S rRNA qPCR quantification integrated with

metagenome data, was statistically similar between the unwarmed and warmed soils;

however, three out of the four investigated warming transects showed an increase (Fig.

S2). Additionally, the warmed soil exhibited a higher respiration activity, with *in situ* soil

CO_2 emission (751.7±123.2 mg CO_2 m^{-2} h^{-1}) at a significant 1.6-fold faster rate than

from unwarmed soil (460.7±9.8 mg CO_2 m^{-2} h^{-1}) (Fig. 1A, Table S2). However, soil CO_2

concentrations along the soil depth profile were comparable in the long-term warmed

and unwarmed soil with significantly higher concentrations in the deepest analysed

layer of 20 cm (Fig. 1C). The corresponding δ¹³C-CO_2 values became progressively

lighter with increasing depth with the exception of two warmed plots at which we

observed a slight increase in δ¹³C-CO_2 values with depth suggesting a slight inflow of

geogenic CO_2 (Fig. 1E, Table S3).

In contrast to the communities mediating organic carbon respiration, soil

methanotrophic activity was not stimulated by long-term warming. There was an overall

22% decrease in *in situ* atmospheric net CH_4 uptake in warmed (0.032±

0.007 mg CH_4 m^{-2} h^{-1}) compared to unwarmed plots (0.041±0.012 mg CH_4 m^{-2} h^{-1}),

though rates varied considerably between plots (Fig. 1B, Table S2). CH_4 concentrations

decreased similarly with depth in the soil pore space of both unwarmed and warmed

soil, and δ¹³C-CH_4 signatures became increasingly heavier with soil depth (Fig. 1D, F,

Table S3). A model estimate of δ^{13}C-CH$_4$ values along the soil profile based on

methanotrophic δ^{13}C-CH$_4$ fractionation agreed well with the obtained data, indicating

that CH$_4$ oxidation can be assumed as the reason for the observed decrease in CH$_4$ in

deeper soil layers (Fig. 1F). These patterns suggest no biotic or geogenic CH$_4$ sources

were present below the study plots. Therefore, the measured net CH$_4$ uptake likely

corresponds to the gross CH$_4$ oxidation activity of atmospheric CH$_4$ oxidizers in the soil.

This is further corroborated by the negligible abundance and transcription of key

methanogenesis genes in the obtained metagenomes and metatranscriptomes (Fig. S3,

Table S4–S7). Logistical challenges prevented an equivalent quantification of soil CO

and H$_2$ fluxes in the field; however, we were able to detect *in situ* uptake of both gases

by unwarmed and warmed soil (Table S8). As a support to this unreplicated

observation, a diffusion model for the studied trace gases was built, showing the

possible uptake of CO and H$_2$ by the upper soil layer and likely depletion of these gases

within the top 10 cm, while CH$_4$ was predicted to diffuse deeper and only be depleted at

a depth of >20 cm (Table S9). This aligns well with our measured CH$_4$, H$_2$, and CO

uptake data (Tables S2, S8) and provides confidence in the estimated gas flux across

the soil profile.

Soil microcosms were prepared to measure the *ex situ* oxidation of trace-level CO, H$_2$,

and CH$_4$, with soil from unwarmed and long-term warmed plots incubated at their

respective average summer temperature of 13°C and 19°C, respectively. A short-term

warming treatment (incubation at 19°C for 40 days) with soil from unwarmed plots was

also included to evaluate the effect of a warming pulse. Both atmospheric H$_2$ and CO

oxidation rates were significantly faster in long-term warmed soil (0.72 and 0.076 nmol h^{-1}

^{1} gDW^{-1}, respectively) than in unwarmed soil (0.25 and 0.034 nmol h^{-1} gDW^{-1},

respectively) (Fig. 2A–B, Table S10–S12). In contrast, CH$_4$ oxidation rates in the

microcosms remained similar (Fig. 2C), in line with the unchanged *in situ* soil CH_4

uptake of unwarmed and warmed soil (Fig. 1A). These data imply increased soil sink

capacities for CO and H_2, but not for CH_4 uptake in the long term. Short-term warming of

soil, on the other hand, did not lead to changes in the consumption rates of any of the

three gases (Fig. 2A–C, Table S11–S12), suggesting that the size of the soil trace gas

sink is resilient to short-term warming.

Warming impacts the relative abundance and community composition of trace

gas oxidizers

To investigate how long-term warming modulates the populations of trace gas oxidizers,

we obtained metagenomes from samples of both unwarmed and long-term warmed

plots. Additionally, we included data from previously obtained metagenomes of the

same plots into the analyses. A gene-centric approach was applied to profile the

community abundance of 51 metabolic marker genes, specifically form I CO

dehydrogenases, hydrogenases, and CH_4 monooxygenases, which catalyze the

oxidation of CO, H_2, and CH_4, respectively (Fig. 3, Fig. S3, Table S4–S5). Consistent

with our prior survey on temperate soils (*15*), hydrogenases (primarily group 1h [NiFe]-

hydrogenases) and [MoCu]-CO dehydrogenases (*coxL*) that mediate atmospheric

uptake during long-term survival were encoded by most microbial community members,

whereas CH_4 monooxygenases (*pmoA*, *mmoX*) were relatively rare, under both

temperature conditions (Fig. S3). Although not significant, warmed soil harboured a

higher abundance of [NiFe]-hydrogenases (warmed: 0.69, unwarmed: 0.56 copies/cell,

based on the abundance of hydrogenase genes related to the summed count of 14

single-copy marker genes), whereas CH_4 monooxygenases decreased by an average of

32% (warmed: 0.012, unwarmed: 0.017 copies/cell) (**F**ig. 3A). No notable relative

abundance shifts in the CO dehydrogenases (**F**ig. 3A) or any of the other profiled

metabolic marker genes were observed (Fig. S3, Table S5).

We additionally analysed shifts in the relative abundance of major soil prokaryotic taxa

mediating trace gas oxidation by profiling the taxonomic affiliation of the trace gas

oxidation marker genes from the metagenomic assembly. Based on this analysis, the

richness of CO (180±27 unique best hit genomes), H_2 (160±36), and CH_4 (4.0±0.8)

oxidizer communities were not affected by warming. [NiFe]-hydrogenases and CO

dehydrogenases were widespread, encoded by 22 and 13 phyla, respectively, and

predominantly associated with the bacterial classes *Actinomycetia*, *Thermoleophilia*,

and *Alphaproteobacteria* (Fig. 3B, Table S13). Actinomycetal CO and H_2 oxidizers

showed a trend of increased relative abundance with long-term soil warming, while

Thermoleophilia groups overall decreased. CH_4 monooxygenases were mainly assigned

to alphaproteobacterial (primarily *Methylocella* spp.; *Methylocapsa* spp. in the recent

GTDB v226) and to a lesser extent to gammaproteobacterial taxa, and both groups

showed a trend of decreased relative abundance with warming (Fig. 3B). This

observation is consistent with metagenomic community profiling based on 16S rRNA

genes by phyloFlash (Table S14).

To gain deeper insights into the mediators of trace gas oxidation, we performed

metagenomic assembly and binning, yielding 250 high- and medium-quality

metagenome-assembled genomes (MAGs; dereplicated at 95% ANI) (Table S15). Of

these, 110 bacterial MAGs spanning 12 phyla held the genomic potential for trace gas

oxidation, namely 88, 45, and 1 MAGs encoding the enzymes for atmospheric CO, H_2,

and CH_4 oxidation, respectively (Fig. 4, Table S16). To determine whether long-term

warming affected the abundance of distinct trace gas oxidizers, we mapped the MAGs

to the metagenomic short reads from the unwarmed and warmed soils (Table S17). In

line with the results of the short-read and assembled-read analyses outlined above,

more than half of the CO- and H_2-oxidizing MAGs were relatively enriched in the

warmed soil (50 and 27 MAGs, respectively), while the sole CH_4-oxidizing MAG

(*Methylocella* spp.) decreased by 4.3-fold (Fig. 4). However, few of these differences

were statistically significant (after correcting for multiple comparisons) given the

considerable variability in community composition between the plots. Finally, there was

an increase of summed relative abundance of H_2-oxidising MAGs ($1.3\pm0.2\%$

unwarmed, $2.0\pm0.1\%$ warmed) in warmed soil, whereas the summed relative

abundance of all MAGs was unchanged ($8.7\pm2.0\%$ unwarmed, $11.3\pm0.6\%$ warmed,

Fig. S4). Altogether, these analyses suggest diverse CO and H_2 oxidizers increased in

relative abundance with warming, whereas methanotrophs decreased, which aligns well

with the *in situ* and *ex situ* activity data.

Cell-specific gene transcription and activity of trace gas oxidizers decrease with

warming

We obtained the metatranscriptomes of the same soil samples to further determine the

relative transcriptional activities of soil trace gas oxidizers (Fig. 5). As done for the

metagenome analyses, we also included previously obtained metatranscriptomes from

the same plots. For H_2 oxidation, [NiFe]-hydrogenases were abundantly transcribed at

both temperature conditions (Fig. S3, Table S18), but transcription ratios (average

RNA:DNA ratios) were significantly lower in warmed soil, as compared to unwarmed soil

(0.22 and 0.39, respectively) (Fig. 5B, Table S19–S20). For CO and CH_4 oxidation,

transcripts and transcription ratios for CO dehydrogenases and CH_4 monooxygenases

likewise decreased in warmed soil, albeit not significantly (Fig. 5, Table S7, S19). The

metatranscriptomic analyses further revealed a trend of warming-induced decreases in

transcripts for NuoF, responsible for NADH oxidation, and a drop in transcription of

genes related to aerobic respiration and oxidative phosphorylation (Fig. S3, Table S7).

To evaluate if the decrease of *in situ* transcription of trace gas oxidation genes can be

related to the measured *ex situ* oxidation rates, we converted the latter rates per cell by

normalizing the bulk soil consumption rates for each gas by the cell count (as based on

qPCR data integrated with metagenome data) and the proportion of respective trace

gas oxidizers estimated by metagenomic analysis (Fig. 2D–F, Table S10). The *ex situ*

cellular rates closely matched the *in situ* gene transcription, with a general 1.83-, 1.73-,

and 3.26-fold decrease in cellular gas uptake for CO, H_2, and CH_4, respectively, in long-

term warmed soil. Together, our data suggest long-term soil warming decreased trace

gas oxidation on a per-cell basis.

Shifts in relative abundance of trace gas oxidizers are consistent in long-term

warmed tundra and temperate forest soils

To determine if our findings are generalisable to other soils experiencing warming, we

retrieved and reanalysed metagenomes from two other controlled warming studies,

namely tundra soil from the Carbon in Permafrost Experimental Heating Research

(CiPEHR) site (warming at 2–5°C above ambient for 1.5 years; AK, USA) (47) and

temperate forest soil at the Harvard Forest Long Term Ecological Research (LTER) site

(warming at 5°C above ambient for 5, 8, and 20 years; MA, USA) (48) (Table S21). In

the tundra soil metagenomes, we detected a 23.1% and 12.6% higher relative

abundance of CO and H_2 oxidation genes per cell, respectively (Fig. 6A, B).

Additionally, we observed a 2.1-fold decrease in CH_4 monooxygenase genes under

warming (Fig. 6C), which replicates our previous findings. Similar trends were also

observed for the LTER temperate forest soil warmed for 5 and 8 years; however, in this

soil, methanotroph abundance was 2-fold higher after 20 years of warming (Fig. 6).

Together, these results suggest that long-term warming promotes soil H_2 and CO

oxidizers, while it does not seem to consistently promote methanotrophs.

Discussion

Soil warming has been shown to increase microbial respiration and growth (35), and

nitrogen mineralization (49), eventually leading to cumulative carbon and nitrogen

losses in topsoils (34, 47, 48, 50). In light of the ability of heterotrophic trace gas

oxidizers to supplement energy demand from atmospheric substrates, we initially

hypothesized that trace gas oxidizers would increase their gas consumption under

carbon- and nitrogen-depleted conditions in long-term warmed soil. Contrary to this

hypothesis, our study did not show an increase *in situ* sink capacity for CO, H_2, and CH_4

of warmed soils. Moreover, *ex situ* cell-specific oxidation rates of CO, H_2, and CH_4 did

not increase with long-term (> 60 years) warming of 6°C (Fig. 2D–F). Furthermore, *in*

situ community-level transcription of marker genes encoding enzymes for trace gas

oxidation showed decreasing trends (Fig. 5). This may be due to the observed non-

decreased levels of dissolved organic carbon and nitrogen (Fig. S2), which likely

continued to sufficiently support heterotrophy in organisms capable of atmospheric CO

and H_2 oxidation. Although decreased dissolved organic carbon and nitrogen levels

have been reported at the same site previously (51, 52), our results indicate that such

depletion is not constant, and yearly or possibly seasonal fluctuations occur. We can

only speculate that this is connected to seasonal variations in plant activity. The non-

decreased levels of dissolved organic carbon and nitrogen are consistent with our *in*

situ gas measurements, which show significantly higher respiration rates in warmed soil

compared to those in unwarmed soil.

Despite the declining cellular-level responses, the overall *ex situ* sink capacity of

warmed soil increased for CO and H_2. We demonstrated that this effect was possibly

driven by the increase of total microbial cell number and biomass in three out of the four

studied warming transects, thereby compensating for reduced cellular gas uptake

activity under long-term warming (Fig. S2). It should be noted that previous work on the

same study site has, in contrast, observed a decrease in microbial biomass and

abundance (51), while yet another study shows no change in microbial biomass carbon

with warming (52). Thus, the dominant soil CO and H_2 oxidizing populations appear to

retain their function with long-term warming, but multi-season studies across a larger

geographical range are needed to definitively confirm this observation. Moreover, the

fact that their relative abundance, especially those affiliated with *Actinomycetia*,

increased across the three investigated long-term warmed soil ecosystems (Fig. 6)

suggests that trace gases likely remain a highly valuable energy source for soil

microbes to persist under warmer conditions (23, 53). Given the high capacity for CO

and H_2 uptake across different soil types and the substantial community size of trace

gas oxidizers (15), our results suggest that soils can serve as a resilient and reliable

sink for these trace gases in a warmer future.

In contrast to CO and H_2 uptake, the soil sink capacity for the potent greenhouse gas

CH4 remained stable, with some indications of weakening due to prolonged warming.

This is suggested by multiple lines of evidence, which, although individually not

statistically significant, all showed the same pattern of decrease. Firstly, while the

aerated soil maintained a net consumption of atmospheric CH_4, in two out of the four

studied warming transects, both *in situ* and *ex situ* rates of CH_4 uptake declined with

long-term warming (Fig. 1B, 2C). Likewise, relative abundance, gene transcription, and

DNA:RNA ratios of methanotrophs in two to three out of the four warming transects

declined with warming. Finally, in conjunction with the observation of decreased

methanotroph abundance in warmed Alaskan tundra soil, all data suggest that the soil

CH_4 sink in upland soils of polar regions is likely to weaken under climate warming.

Even medium-term warming of subarctic grassland soil showed a unimodal,

temperature-dependent decrease in soil CH_4 uptake with warming of >4°C (39). Yet, we

note that some previous studies on soil warming in Arctic and in alpine grassland soils

have reported increased soil CH_4 uptake (38, 54). In line with these studies, permafrost

communities were recently shown to contain relatively more methanotrophs capable of

atmospheric CH_4 oxidation with warming (55). Importantly, previous pure-culture work

has demonstrated that responses of methanotrophic bacteria to warming depend on the

available methane concentration (56), suggesting that temperature and substrate

availability may interact to determine uptake rates. Collectively, this highlights that

methanotrophic populations and the soil CH_4 sink are sensitive and respond variably to

warming. Future studies should therefore address potential interactive effects of global

warming and atmospheric methane concentrations. Our work highlights the need to

profile the crucial CH_4 sink function of soils in relation to a warmer future climate.

Materials and Methods

Experimental sites and soil sampling

The experimental site is a grassland in Grændalur (the "green valley") near Hveragerði,

Iceland (64.0272328°N, 21.1964731°W), which has been naturally warmed for a

minimum of 60 years (45) or possibly even before 1708 (57). At this site, geothermal

heat within the site's bedrock has created a natural radiative warming gradient through

the soil, which is exploited in permanent study plots along replicated 50 m long

transects ranging from unwarmed soil temperatures to +20°C warming at a 10 cm soil

depth. In this study, we only focus on the soil temperature range from unwarmed to

+6°C. Temperature logger measurements taken in 2013 reveal a constant warming

trend following the seasonal temperature cycle at an elevated level (45), as illustrated in

Fig. S1 (58). The ecosystem is a grassland dominated by *Agrostis capillaris* on a slightly

acidic Silandic Andosol. Soil samples were collected in July 2022 from a depth of 0–

10 cm from four unwarmed plots (16°C at time of sampling) and four warmed plots

(approximately 22°C at time of sampling), *i.e.*, eight soil samples in total. All samples

were homogenized by mixing in sterile sampling bags on site. In parallel, samples for

metatranscriptomics were taken (0–10 cm layer) (59), immediately frozen and kept on

dry ice until arrival in the lab, where they were stored at -70°C until RNA extraction.

Samples for metagenomics, qPCR, and microcosm incubations were cooled until they

reached the laboratory, where the former two were frozen at -20°C and the latter kept at

4°C until incubation within one week.

Microbial biomass, dissolved organic carbon and nitrogen, and pH

Soil pH was measured in a 1:5 (w/v) soil to 10 mM $CaCl_2$ suspension using a pH meter.

Dissolved organic carbon (DOC) and total dissolved nitrogen (TDN) concentrations

were quantified in 1 M KCl extracts (1:7.5 w/v) with a TOC/TN analyzer (TOC-L

CPH/CPN, Shimadzu). Microbial biomass C (Cmic) and N (Nmic) were determined

using the chloroform fumigation-extraction method (60): Briefly, soils were fumigated

with chloroform for 48 h in the dark at room temperature, then extracted with 1 M KCl

(1:7.5 w/v) and analyzed as above. Cmic and Nmic were calculated as the difference in

extractable C and N between fumigated and unfumigated samples and corrected for

incomplete extraction by dividing Cmic by 0.45 and Nmic by 0.54.

Surface gas fluxes

In situ gas fluxes (CO_2 and CH_4) at the eight study plots (*i.e.*, four unwarmed

temperature and four +6°C warming plots, see above) were measured using a LI-COR

gas analyzer with cavity ringdown technique (LI-7810, LI-COR Biotechnology, USA). A

PVC chamber (11 cm in diameter × 10 cm in height; 0.95 L total volume) was attached

to a collar fixed in the soil to a depth of 2 cm for at least 1 hour in advance, with the

remaining aboveground plant parts removed from the plots before the collar placement.

The chamber was connected to the gas analyzer. After chamber closure and initial

stabilization, the linear concentration increase of CO_2 and CH_4 was recorded for 15

minutes at each plot on two separate days to account for day-to-day variation (2022-07-

24 and 2022-07-26). Linearity of concentration increase/decrease was monitored in the

field. Fluxes of CO_2 and CH_4 were calculated from the slope of the linear concentration

change in the chamber (R2 > 0.95) as a function of time.

Due to logistical constraints, samples for quantifying *in situ* fluxes of CO and H_2 were

collected from only one of the four warming transects in October 2025 and analyzed in

the laboratory using a trace gas analyzer (see Supplementary Materials for a detailed

description).

Soil gas profiles and δ¹³C measurements

Gas samples of 25 mL were taken from the eight study plots with a stainless-steel

sampling probe (ø = 3 mm, l = 40 cm) at three soil depths (5, 10, and 20 cm) and

injected into pre-evacuated 12 mL vials (Labco Ltd., UK). Two replicates were taken in

each depth of every plot, one for analysis of CO_2 and CH_4 concentrations and one for

analysis of $\delta^{13}C$ values in CO_2 and CH_4. Additionally, an atmospheric gas sample was

taken by drawing air above each plot, resulting in a total of 56 samples. Soil

temperatures at 5, 10, 15, and 20 cm were measured using a metal probe and a manual

thermometer (TM-80N with a K-type thermocouple probe, Tenmars Electronics, Taiwan)

during sampling at each plot. Soil moisture was measured in triplicate at all eight plots

at a distance of 10 cm using a moisture meter ML-3 Thetaprobe connected to a HH2

moisture sensor (Eijkelkamp, The Netherlands). Gas samples for quantification and

$\delta^{13}C$ measurement of CO_2 and CH_4 were treated as follows. Samples were stored at

room temperature before analysis of gas concentration by gas chromatography with

thermal conductivity (CO_2) and flame ionization detection (Agilent 7890B) and cavity

ring-down spectroscopy (CRDS, Picarro G2201-I with SAM autosampler, Picarro Inc.,

USA). For CRDS, samples were diluted with synthetic air to <2,000 ppmv CO_2. The

measured isotope ratios were corrected for drift and concentration effects based on the

repeated analysis of an in-house standard gas at different dilutions. To correct for

concentration effects, we fitted a linear regression between the measured isotope ratio

and the inverse concentration, with individual analyses weighted inversely to the

standard deviation during measurement to account for the lower precision of

measurements from more dilute solutions (61). $\delta^{13}C$ values of CH_4 samples with a

concentration < 1.2 ppmv after dilution were excluded as reliable measurements are not

possible below this concentration. CO, H_2, and CH_4 soil concentration values were fitted

into a diffusion model to estimate gas concentration across the soil profile

(Supplementary Materials). The Rayleigh distillation function, with a fractionation factor

(α) of 1.020, was used to predict $\delta^{13}C–CH_4$ values.

Microcosm incubations

Soil microcosms were set up to determine the ability of soil microbial communities to

oxidize trace levels of CO, H_2, and CH_4, under control, short-term warming, and long-

term warming scenarios. Soil samples from four unwarmed plots and four warmed

(+6°C) plots were used (sampling and storage described above). To establish the

control and short-term warming treatments, approximately 2.5 g of soil from unwarmed

plot samples was placed in 160 mL serum vials and incubated at 13°C for the control

soil treatment (n=4) and at 19°C for the short-term warmed treatment (n=4). Likewise,

soil from the warmed plot samples was incubated at 19°C for the long-term warmed

treatment (n=4). Both 13°C and 19°C fall within the range of the respective average

summer temperatures in unwarmed plots and warmed plots over the past nine years

(Fig. S1). Each serum vial was amended with CO, H_2, and CH_4 to initial mixing ratios of

≈10 ppmv for each gas. At the start of the experiment, the serum vials were also

adjusted to 15% overpressure with ambient laboratory air to compensate for the gas

sampled during the experiment. 2 mL of headspace gas were sampled using a gas-tight

syringe at different time intervals (eight times in the first three days and eight times in

the remaining incubation time) and up to 40 days. Gas samples were measured using a

VICI gas chromatograph with a pulsed discharge helium ionization detector calibrated

against standard ultra-pure CO, H_2, and CH_4 gas mixtures as previously described (22).

Pooled autoclaved soil (60 min at 120°C) and empty serum vials were prepared as

negative controls.

Kinetic analysis

For kinetic analysis, measurement time points with individual gas concentrations

exceeding 0.4 ppmv were used. The gas consumption patterns were fitted with both

linear and exponential models. In all cases, the exponential model provided a better fit

to the data based on Akaike information criterion scores; thus, the first-order reaction

rate constants were calculated and used for kinetic modeling. The bulk atmospheric gas

oxidation rates of each sample were calculated with respect to the mean atmospheric

mixing ratio of the corresponding gas (CO: 0.09 ppmv; H_2: 0.53 ppmv; CH_4: 1.9 ppmv)

(9, 11, 13). Cell-specific gas oxidation rates were determined under the assumption that

all cells are viable and active. The soil cell abundance was determined using 16S rRNA

gene copy numbers from quantitative PCR (qPCR, see details below) corrected to the

average 16S rRNA gene copy numbers derived from the metagenomic short reads (see

below). The cell-specific gas oxidation rates were then inferred by dividing the bulk

oxidation rate by the product of the estimated relative abundance of trace gas oxidizers

derived from the metagenomic short reads (the average gene copy numbers, assuming

one copy per cell; see details below) and the soil cell abundance.

DNA and RNA extraction, qPCR, and sequencing

Community DNA was extracted from ≈0.25 g of soil using the DNeasy PowerSoil Pro kit

following the manufacturer's instructions. The number of bacterial and archaeal cells in

each sample was estimated using quantitative PCR (qPCR) of the 16S rRNA gene with

the universal primer pair F515 and R806, as described previously (62). Further details

of processing are described in the Supplementary Materials.

The RNA extraction and metatranscriptomic sequencing were described in (52).

Detailed steps are described in the Supplementary Materials.

Functional annotation of unassembled metagenome and metatranscriptome reads

High-quality DNA short reads and mRNA reads underwent metabolic annotation using DIAMOND v.2.0.15 (63). Reads were aligned against a custom database of 51 metabolic marker proteins (64), covering the major pathways for aerobic and anaerobic respiration, energy conservation from organic and inorganic compounds, carbon fixation, nitrogen fixation, and phototrophy (65). The alignments were filtered to retain sequences of at least 40 amino acids in length or with at least 80% query or subject coverage. They were further filtered based on an identity threshold of 80% (PsaA), 75% (HbsT), 70% (PsbA, IsoA, AtpA, YgfK, and ARO), 60% (CoxL, MmoX, AmoA, NxrA, RbcL, NuoF, [FeFe]-hydrogenases and [NiFe]-hydrogenases group 4), or 50% (all other genes). For metagenomic data, the gene abundance was converted to reads per kilobase million (RPKM) as follows: RPKM = X / total sample reads $\times$ 10^6, where X = reads aligned to a gene / gene length (kbp) (66). This value was divided by the mean abundance of 14 universal single-copy ribosomal marker genes (also in RPKM) provided by SingleM v0.13.2 (67) to obtain the 'average gene copies per cell'. This value corresponds to the proportion of community members that encode the gene. For metatranscriptomic data, the gene transcription was expressed in RPKM as described above. Heatmaps showing the metagenomic data and RPKM for metatranscriptomes were generated in R version 4.3.1 (68) using the ggplot2 package (69).

Metagenome assembly and taxonomic analysis

Metagenome reads were processed using Trimmomatic v0.39 (70) to remove adapter

contamination, trim low-quality ends of reads, and remove sequences shorter than 36

bp. The quality of the filtered reads was rechecked using Fastp v0.23.4 (71). Filtered

reads were assembled individually using MEGAHIT v1.2.9 (72). Gene calling was done

using Prodigal v2.6.3 (73) with the mode "-meta". The same custom database of 51

metabolic marker proteins as mentioned above was used for a DIAMOND blastp search

v2.1.9 (*63*) with metagenome gene calls as queries; best hits with percent identity >75

and query coverage >70 to the CoxL, MmoX, PmoA, [NiFe]-hydrogenases, [FeFe]-

hydrogenases, [Fe]-hydrogenases were retrieved. These hits were further compared

using DIAMOND blastp v2.0.14 against the bacterial RefSeq database to assign

taxonomy to the target gene calls. The taxonomy search utilized filtering for only the

best hits, with the following parameters in descending priority: max bitscore > min e-

value > max percent identity. The RefSeq r220 database was accessed on 2023-11-20

(*74*) using the NCBI Datasets v15.29.0 tool (http://www.ncbi.nlm.nih.gov/datasets) to

retrieve bacterial genomes and MAGs, which were classified by the GTDB taxonomy

(bac120_taxonomy_r214). Some of the metagenome gene calls had hits to RefSeq

genomes (or MAGs) not included in the GTDB taxonomy; therefore, GTDB-Tk v2.4 (75)

was used to assign taxonomy to these genomes. In this way, it was possible to link

trace gas oxidation genes assembled in metagenomes with their closest relatives of

known taxonomic identity. The relative abundance of target trace gas oxidation genes

was expressed as the number of target genes per million total assembled genes in each

metagenome. Richness of the closest genomes with respective trace gas oxidation

genes was calculated for each sample. Data were processed in R 4.3.1 (68) using the

packages tidyverse 2.0.0 (76), tidylog 1.0.2 (*77*), and vegan 2.6-4 (78).

Metagenomic binning, functional annotation, and phylogenetic analysis

A total of eight metagenomes (4× from unwarmed plots and 4× from warmed plots),
which were quality controlled using BBTools (79) as described above and co-assembled
using MEGAHIT v1.2.9 (72) with the minimum contig length of 500 bp. Coverage profiles
for each contig were generated by mapping the short reads to the assemblies using
BBMAP v38.81 (79) and SAMtools v1.19.3 (80). Genome binning was performed with
MetaBat2 v2.15.5 (81), MaxBin2 v2.2.7 (82), CONCOCT v1.1.0 (83), and SemiBin2
v2.1.0 (84). The four bin sets were consolidated using the bin refinement module of
MetaWrap v1.3.2 (85) and dereplicated using dRep v3.5.0 (86) with a minimum
completeness of 50% and contamination of 10%. The completeness, contamination,
and heterogeneity of the bins were assessed using CheckM2 v1.0.1 (87), with the
quality thresholds selected as previously described (88). Only medium (completeness >
50%, contamination < 10%) and high (completeness > 90%, contamination < 5%)
quality bins were retained for further processing and were termed metagenome-
assembled genomes (MAGs). Taxonomy was assigned to each MAG using GTDB-Tk
v2.4.0 (75) with GTDB release 220 (89). Relative abundance of each MAG in each
sample was calculated using the 'genome' function (--min-covered-fraction 0, --min-
read-aligned-percent 0.75, and –min-read-percent-identity 0.95) of CoverM v0.7.0 (90).
Recovered MAGs were then metabolically annotated against the custom database
using Prodigal v2.6.3 (73) with the mode "-single" to predict open reading frames
(ORFs) and DIAMOND v2.0.15 (63) to identify metabolic marker genes as previously
described. A maximum likelihood tree was constructed for MAGs containing trace gas
oxidizing genes using GTDB-tk v2.4.0 with the commands '*identify*' and '*align*' (75). The
tree was constructed using IQ-TREE v2.3.4 (91) with 1,000 ultrafast bootstrap using the
LG+F+I+G4 model. The tree was visualized and mid-rooted using the Interactive Tree of

Life (iTOL) (92).

Analysis of publicly available metagenomes from long-term soil warming studies

Given the limited number of samples, we included analyses of public metagenomic

(BioProject ID PRJNA746424) and metatranscriptomic (BioProject ID PRJNA663238)

data obtained from the same studied site (49). To compare the effects of warming on

trace gas oxidation capacity of different types of soils, publicly available metagenomes

from temperate forest soil at the Harvard Forest Long Term Ecological Research

(LTER) site (MA, USA) (48) and from tundra soil from the Carbon in Permafrost

Experimental Heating Research (CiPEHR) site (Alaska, USA) (47) were retrieved. The

chosen LTER soil was sampled from 0–20 cm depths, and the warmed soil had been

heated consistently 5°C above the temperatures observed in the nearby control soil for

5, 8, and 20 years (48). The soil from the CiPEHR sites was collected from depths of

15–25 cm, with the warmed soil experimentally heated between 2–5°C above the

control soil temperature for 1.5 years (47). All datasets were quality-assessed and

annotated against 51 metabolic marker proteins, as described above.

Statistical analysis

All analyses were performed using the statistical software R version 4.3.1 (68) and

GraphPad Prism version 10.4.1. Linear mixed models generated with the function lme

of the package nlme v.3.1-164 (93) were used to evaluate the influence of short and

long-term warming on *in situ* gas fluxes and *ex situ* CO, H_2, and CH_4 oxidation rates.

The models included the field transect as a random effect to account for the paired

sampling design and were checked for meeting the model assumptions by the function

check_model of the package performance (94). Oxidation rate values were log-

transformed to normalize residuals where necessary. Likewise, linear mixed models

were used to test for significant differences in the relative abundance of transcripts and

transcription ratios of marker genes for trace gas oxidation, as well as in the relative

abundance of MAGs mapped against our metagenomes. Statistical significance was set

at ≤ 0.05, and a trend at ≤ 0.1. Detailed statistical evaluation and outputs are provided in

Table S11–S12. For public databases, differences in marker gene abundances were

determined using the Mann-Whitney test in GraphPad Prism.

Acknowledgements

We thank Páll Sigurðsson for help during field sampling and Christoph Gall for help with

the analysis of soil parameters.

Funding

VT, JAN, MW, and AD were supported by the JuniorStar project 21-17322M from the

Czech Science Foundation to AD.

MBD was funded by the German Science Foundation (DFG, BO 5559/1-1 nr.

433256088).

ATT and AS acknowledge funding from the Research Council of Norway (FRIPRO

projects LoAir, 315129, and SHRINK, 344999).

PML acknowledges an Australian Research Council DECRA Fellowship

(DE250101210) for salary support.

CG acknowledges an Australian Research Council Future Fellowship (FT240100502)

for salary support.

LK acknowledges a Research Council of Finland Fellowship (354501) for salary

support.

AR acknowledges funding from the Austrian Science Fund (FWF), grant

https://doi.org/10.55776/COE7.

Author contributions

Conceptualization: AD, CG, VT, DN, PML, MW

Methodology: VT, DN, PML, MW, JAN, KYS, JK, MBD, AS, TS, LK, CB, AR, TU, ATT,

CG, BDS, AD

Investigation: DN, VT, MW, JAN, MBD, AS, TS, LK, CG, AD

Visualization: VT, DN

Supervision: AT, CG, BDS, AD

Writing—original draft: DN, PML, VT, CG, AD

Writing—review & editing: VT, DN, PML, CB, AR, TU, ATT, CG, BDS, AD, MW, AS

Competing interests

The authors declare that they have no competing interests.

Data availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or

the Supplementary Materials. The raw sequencing data, together with sample

metadata, have been deposited at the ENA under the accession numbers PRJEB98590

(metagenome data) and PRJNA1130687 (metatranscriptome data, Dahl et al., 2025),

run accessions SRR29850129, SRR29850130, SRR29850145, SRR29850146,

SRR29850149, SRR29850150, SRR29850151, SRR29850152. Processing scripts are

deposited   at   a   public   repository

https://github.com/TlaskalV/trace_gas_oxidation_and_warming/.

References

1. P. Forster, V. Ramaswamy, P. Artaxo, T. Berntsen, R. Betts, D. W. Fahey, J. Haywood, J. Lean, D. C. Lowe, G. Myhre, J. Nganga, R. Prinn, G. Raga, M. Schulz, R. Van Dorland, *Changes in Atmospheric Constituents and in Radiative Forcing Chapter 2* (Cambridge University Press, United Kingdom, 2007; http://inis.iaea.org/search/search.aspx?orig_q=RN:39002468).

2. M. Sand, R. B. Skeie, M. Sandstad, S. Krishnan, G. Myhre, H. Bryant, R. Derwent, D. Hauglustaine, F. Paulot, M. Prather, D. Stevenson, A multi-model assessment of the Global Warming Potential of hydrogen. *Commun. Earth Environ.* 4, 203 (2023).

3. IPCC, *Global Warming of 1.5°C: IPCC Special Report on Impacts of Global Warming of 1.5°C above Pre-Industrial Levels in Context of Strengthening Response to Climate Change, Sustainable Development, and Efforts to Eradicate Poverty* (Cambridge University Press, ed. 1, 2022; https://www.cambridge.org/core/product/identifier/9781009157940/type/book).

4. J. D. Patterson, M. Aydin, A. M. Crotwell, G. Pétron, J. P. Severinghaus, P. B. Krummel, R. L. Langenfelds, V. V. Petrenko, E. S. Saltzman, Reconstructing atmospheric H_2 over the past century from bi-polar firn air records. *Clim. Past* 19, 2535–2550 (2023).

5. R. B. Skeie, Ø. Hodnebrog, G. Myhre, Trends in atmospheric methane concentrations since 1990 were driven and modified by anthropogenic emissions. *Commun. Earth Environ.* 4, 317 (2023).

6.   X. Faïn, S. Szopa, V. Naïk, P. Martinerie, D. M. Etheridge, R. H. Rhodes,

C. M. Trudinger, V. V. Petrenko, K. Fourteau, P. Place, Preindustrial to present-day

changes in atmospheric carbon monoxide: agreements and gaps between ice

archives and global model reconstructions. Gases/Atmospheric Modelling and Data

Analysis/Troposphere/Chemistry (chemical composition and reactions) [Preprint]

(2024). https://doi.org/10.5194/egusphere-2024-653.

7.   R. Conrad, W. Seiler, Decomposition of atmospheric hydrogen by soil

microorganisms and soil enzymes. *Soil Biol. Biochem.* 13, 43–49 (1981).

8.   T. S. Rhee, C. A. M. Brenninkmeijer, T. Röckmann, The overwhelming

role of soils in the global atmospheric hydrogen cycle. *Atmospheric Chem. Phys.* 6,

1611–1625 (2006).

9.   D. H. Ehhalt, F. Rohrer, The tropospheric cycle of H_2: a critical review.

Tellus B Chem. Phys. Meteorol. 61, 500 (2009).

10.   G. W. Bartholomew, M. Alexander, Soil as a sink for atmospheric carbon

monoxide. *Science* 212, 1389–1391 (1981).

11.   M. A. K. Khalil, R. A. Rasmussen, Atmospheric carbon monoxide:

Latitudinal distribution of sources. *Geophys. Res. Lett.* 17, 1913–1916 (1990).

12.   M. Bender, R. Conrad, Kinetics of CH_4 oxidation in oxic soils exposed to

ambient air or high CH_4 mixing ratios. *FEMS Microbiol. Lett.* 101, 261–270 (1992).

13. S. Kirschke, P. Bousquet, P. Ciais, M. Saunois, J. G. Canadell, E. J.

Dlugokencky, P. Bergamaschi, D. Bergmann, D. R. Blake, L. Bruhwiler, P.

Cameron-Smith, S. Castaldi, F. Chevallier, L. Feng, A. Fraser, M. Heimann, E. L.

Hodson, S. Houweling, B. Josse, P. J. Fraser, P. B. Krummel, J.-F. Lamarque, R.

L. Langenfelds, C. Le Quéré, V. Naik, S. O'Doherty, P. I. Palmer, I. Pison, D.

Plummer, B. Poulter, R. G. Prinn, M. Rigby, B. Ringeval, M. Santini, M. Schmidt, D.

T. Shindell, I. J. Simpson, R. Spahni, L. P. Steele, S. A. Strode, K. Sudo, S. Szopa,

G. R. Van Der Werf, A. Voulgarakis, M. Van Weele, R. F. Weiss, J. E. Williams, G.

Zeng, Three decades of global methane sources and sinks. *Nat. Geosci.* 6, 813–

823 (2013).

14. M. Saunois, A. Martinez, B. Poulter, Z. Zhang, P. A. Raymond, P. Regnier,

J. G. Canadell, R. B. Jackson, P. K. Patra, P. Bousquet, P. Ciais, E. J.

Dlugokencky, X. Lan, G. H. Allen, D. Bastviken, D. J. Beerling, D. A. Belikov, D. R.

Blake, S. Castaldi, M. Crippa, B. R. Deemer, F. Dennison, G. Etiope, N. Gedney, L.

Höglund-Isaksson, M. A. Holgerson, P. O. Hopcroft, G. Hugelius, A. Ito, A. K. Jain,

R. Janardanan, M. S. Johnson, T. Kleinen, P. B. Krummel, R. Lauerwald, T. Li, X.

Liu, K. C. McDonald, J. R. Melton, J. Mühle, J. Müller, F. Murguia-Flores, Y. Niwa,

S. Noce, S. Pan, R. J. Parker, C. Peng, M. Ramonet, W. J. Riley, G. Rocher-Ros,

J. A. Rosentreter, M. Sasakawa, A. Segers, S. J. Smith, E. H. Stanley, J.

Thanwerdas, H. Tian, A. Tsuruta, F. N. Tubiello, T. S. Weber, G. R. Van Der Werf,

D. E. J. Worthy, Y. Xi, Y. Yoshida, W. Zhang, B. Zheng, Q. Zhu, Q. Zhu, Q.

Zhuang, Global methane budget 2000–2020. *Earth Syst. Sci. Data* 17, 1873–1958

(2025).

15.   S. K. Bay, X. Dong, J. A. Bradley, P. M. Leung, R. Grinter, T.

Jirapanjawat, S. K. Arndt, P. L. M. Cook, D. E. Larowe, P. A. Nauer, E. Chiri, C.

Greening, Trace gas oxidizers are widespread and active members of soil

microbial communities. *Nat. Microbiol.* 6, 246–256 (2021).

16.   Y. Xu, Y. Teng, S. Dai, J. Liao, X. Wang, W. Hu, Z. Guo, X. Pan, X. Dong,

Y. Luo, Atmospheric trace gas oxidizers contribute to soil carbon fixation driven by

key soil conditions in terrestrial ecosystems. *Environ. Sci. Technol.* 58, 21617–

21628 (2024).

17.   A. Kropp, D. L. Gillett, H. Venugopal, M. A. Gonzálvez, J. P. Lingford, S.

Jain, C. K. Barlow, J. Zhang, C. Greening, R. Grinter, Quinone extraction drives

atmospheric carbon monoxide oxidation in bacteria. *Nat. Chem. Biol.*, doi: 10.1038/

s41589-025-01836-0 (2025).

18.   R. Grinter, A. Kropp, H. Venugopal, M. Senger, J. Badley, P. R. Cabotaje,

R. Jia, Z. Duan, P. Huang, S. T. Stripp, C. K. Barlow, M. Belousoff, H. S. Shafaat,

G. M. Cook, R. B. Schittenhelm, K. A. Vincent, S. Khalid, G. Berggren, C.

Greening, Structural basis for bacterial energy extraction from atmospheric

hydrogen. *Nature*, doi: 10.1038/s41586-023-05781-7 (2023).

19.   A. T. Tveit, A. G. Hestnes, S. L. Robinson, A. Schintlmeister, S. N.

Dedysh, N. Jehmlich, M. Von Bergen, C. Herbold, M. Wagner, A. Richter, M. M.

Svenning, Widespread soil bacterium that oxidizes atmospheric methane. *Proc.*

Natl. Acad. Sci. 116, 8515–8524 (2019).

20.   P. Constant, S. P. Chowdhury, J. Pratscher, R. Conrad, Streptomycetes

contributing to atmospheric molecular hydrogen soil uptake are widespread and

encode a putative high☐affinity [NiFe]☐hydrogenase. *Environ. Microbiol.* 12, 821–

829 (2010).

21.   C. Greening, M. Berney, K. Hards, G. M. Cook, R. Conrad, A soil

actinobacterium scavenges atmospheric H_2 using two membrane-associated,

oxygen-dependent [NiFe] hydrogenases. *Proc. Natl. Acad. Sci.* 111, 4257–4261

(2014).

22.   Z. F. Islam, P. R. F. Cordero, J. Feng, Y. C. Sean, K. B. Thanavit, R. M.

Gleadow, C. R. Carere, M. B. Stott, E. Chiri, C. Greening, Two Chloroflexi classes

independently evolved the ability to persist on atmospheric hydrogen and carbon

monoxide. *ISME J.* 13, 1801–1813 (2019).

23.   M. Berney, G. M. Cook, Unique flexibility in energy metabolism allows

mycobacteria to combat starvation and hypoxia. *PLoS ONE* 5, e8614 (2010).

24.   M. Berney, C. Greening, K. Hards, D. Collins, G. M. Cook, Three different

[NiFe] hydrogenases confer metabolic flexibility in the obligate aerobe

Mycobacterium smegmatis. *Environ. Microbiol.* 16, 318–330 (2014).

25.   Q. Liot, P. Constant, Breathing air to save energy – new insights into the

ecophysiological role of high☐affinity [NiFe]☐hydrogenase in *Streptomyces*

avermitilis. *MicrobiologyOpen* 5, 47–59 (2016).

26.   P. R. F. Cordero, K. Bayly, P. Man Leung, C. Huang, Z. F. Islam, R. B.

Schittenhelm, G. M. King, C. Greening, Atmospheric carbon monoxide oxidation is

a widespread mechanism supporting microbial survival. *ISME J.* 13, 2868–2881

(2019).

27.   A. T. Tveit, T. Schmider, A. G. Hestnes, M. Lindgren, A. Didriksen, M. M.

Svenning, Simultaneous oxidation of atmospheric methane, carbon monoxide and

hydrogen for bacterial growth. *Microorganisms* 9, 153 (2021).

28.   P. M. Leung, A. Daebeler, E. Chiri, I. Hanchapola, D. L. Gillet, R. B.

Schittenhelm, H. Daims, C. Greening, A nitrite-oxidising bacterium constitutively

consumes atmospheric hydrogen. *ISME J.*, doi: 10.1038/s41396-022-01265-0

(2022).

29.   T. Schmider, A. G. Hestnes, J. Brzykcy, H. Schmidt, A. Schintlmeister, B.

R. K. Roller, E. J. Teran, A. Söllinger, O. Schmidt, M. F. Polz, A. Richter, M. M.

Svenning, A. T. Tveit, Physiological basis for atmospheric methane oxidation and

methanotrophic growth on air. *Nat. Commun.* 15, 4151 (2024).

30.   C. Greening, R. Grinter, Microbial oxidation of atmospheric trace gases.

Nat. Rev. Microbiol. 20, 513–528 (2022).

31.   T. W. Crowther, J. van den Hoogen, J. Wan, M. A. Mayes, A. D. Keiser, L.

Mo, C. Averill, D. S. Maynard, The global soil community and its influence on

biogeochemistry. *Science* 365 (2019).

32.   P. García-Palacios, T. W. Crowther, M. Dacal, I. P. Hartley, S. Reinsch, R.

Rinnan, J. Rousk, J. Van Den Hoogen, J.-S. Ye, M. A. Bradford, Evidence for large

microbial-mediated losses of soil carbon under anthropogenic warming. *Nat. Rev.*

Earth Environ. 2, 507–517 (2021).

33.   J. M. Melillo, S. D. Frey, K. M. DeAngelis, W. J. Werner, M. J. Bernard, F.

P. Bowles, G. Pold, M. A. Knorr, A. S. Grandy, Long-term pattern and magnitude of

soil carbon feedback to the climate system in a warming world. *Science* 358, 101–

105 (2017).

34.   S. Marañón-Jiménez, J. Peñuelas, A. Richter, B. D. Sigurdsson, L.

Fuchslueger, N. I. W. Leblans, I. A. Janssens, Coupled carbon and nitrogen losses

in response to seven years of chronic warming in subarctic soils. *Soil Biol.*

Biochem. 134, 152–161 (2019).

35.   T. W. N. Walker, I. A. Janssens, J. T. Weedon, B. D. Sigurdsson, A.

Richter, J. Peñuelas, N. I. W. Leblans, M. Bahn, M. Bartrons, C. De Jonge, L.

Fuchslueger, A. Gargallo-Garriga, G. E. Gunnarsdóttir, S. Marañón-Jiménez, E. S.

Oddsdóttir, I. Ostonen, C. Poeplau, J. Prommer, D. Radujković, J. Sardans, P.

Sigurðsson, J. L. Soong, S. Vicca, H. Wallander, K. Ilieva-Makulec, E. Verbruggen,

A systemic overreaction to years versus decades of warming in a subarctic

grassland ecosystem. *Nat. Ecol. Evol.* 4, 101–108 (2020).

36.   N. Verbrigghe, N. I. W. Leblans, B. D. Sigurdsson, S. Vicca, C. Fang, L.

Fuchslueger, J. L. Soong, J. T. Weedon, C. Poeplau, C. Ariza-Carricondo, M.

Bahn, B. Guenet, P. Gundersen, G. E. Gunnarsdóttir, T. Kätterer, Z. Liu, M.

Maljanen, S. Marañón-Jiménez, K. Meeran, E. S. Oddsdóttir, I. Ostonen, J.

Peñuelas, A. Richter, J. Sardans, P. Sigurðsson, M. S. Torn, P. M. Van Bodegom,

E. Verbruggen, T. W. N. Walker, H. Wallander, I. A. Janssens, Soil carbon loss in

warmed subarctic grasslands is rapid and restricted to topsoil. *Biogeosciences* 19,

3381–3393 (2022).

37.   B. Bond-Lamberty, V. L. Bailey, M. Chen, C. M. Gough, R. Vargas,

Globally rising soil heterotrophic respiration over recent decades. *Nature* 560, 80–

83 (2018).

38.   Y. Oh, Q. Zhuang, L. Liu, L. R. Welp, M. C. Y. Lau, T. C. Onstott, D.

Medvigy, L. Bruhwiler, E. J. Dlugokencky, G. Hugelius, L. D'Imperio, B. Elberling,

Reduced net methane emissions due to microbial methane oxidation in a warmer

Arctic. *Nat. Clim. Change* 10, 317–321 (2020).

39.   A. S. Nielsen, L. M. Avila, B. D. Sigurdsson, K. S. Larsen, J. R.

Christiansen, Taking the heat: Soil warming optimum of CH_4 uptake in subarctic

mineral soils. *Biogeochemistry* (in press).

40.   M. B. Bertagni, S. W. Pacala, F. Paulot, A. Porporato, Risk of the

hydrogen economy for atmospheric methane. *Nat. Commun.* 13, 7706 (2022).

41.   M. M. Bela, N. Kille, S. A. McKeen, J. Romero☐Alvarez, R. Ahmadov, E.

James, G. Pereira, C. Schmidt, R. B. Pierce, S. M. O'Neill, X. Zhang, S.

Kondragunta, C. Wiedinmyer, R. Volkamer, Quantifying carbon monoxide

emissions on the scale of large wildfires. *Geophys. Res. Lett.* 49, e2021GL095831

(2022).

42.   S. Peng, X. Lin, R. L. Thompson, Y. Xi, G. Liu, D. Hauglustaine, X. Lan, B.

Poulter, M. Ramonet, M. Saunois, Y. Yin, Z. Zhang, B. Zheng, P. Ciais, Wetland

emission and atmospheric sink changes explain methane growth in 2020. *Nature*

612, 477–482 (2022).

43.   C. Arndt, A. N. Hristov, W. J. Price, S. C. McClelland, A. M. Pelaez, S. F.

Cueva, J. Oh, J. Dijkstra, A. Bannink, A. R. Bayat, L. A. Crompton, M. A. Eugène,

D. Enahoro, E. Kebreab, M. Kreuzer, M. McGee, C. Martin, C. J. Newbold, C. K.

Reynolds, A. Schwarm, K. J. Shingfield, J. B. Veneman, D. R. Yáñez-Ruiz, Z. Yu,

Full adoption of the most effective strategies to mitigate methane emissions by

ruminants can help meet the 1.5 °C target by 2030 but not 2050. *Proc. Natl. Acad.*

Sci. 119, e2111294119 (2022).

44.   M. Rigby, S. A. Montzka, R. G. Prinn, J. W. C. White, D. Young, S.

O'Doherty, M. F. Lunt, A. L. Ganesan, A. J. Manning, P. G. Simmonds, P. K.

Salameh, C. M. Harth, J. Mühle, R. F. Weiss, P. J. Fraser, L. P. Steele, P. B.

Krummel, A. McCulloch, S. Park, Role of atmospheric oxidation in recent methane

growth. *Proc. Natl. Acad. Sci.* 114, 5373–5377 (2017).

45.   B. D. Sigurdsson, N. I. W. Leblans, S. Dauwe, E. Guðmundsdóttir, P.

Gundersen, G. E. Gunnarsdóttir, M. Holmstrup, K. Ilieva-Makulec, T. Kätterer, B.

Marteinsdóttir, M. Maljanen, E. S. Oddsdóttir, I. Ostonen, J. Peñuelas, C. Poeplau,

A. Richter, P. Sigurðsson, P. Van Bodegom, H. Wallander, J. Weedon, I. Janssens,

Geothermal ecosystems as natural climate change experiments: The ForHot

research site in Iceland as a case study. *Icel. Agric. Sci.* 29, 53–71 (2016).

46.   IPCC, *Climate Change 2022 – Impacts, Adaptation and Vulnerability:*

Working Group II Contribution to the Sixth Assessment Report of the

Intergovernmental Panel on Climate Change (Cambridge University Press, ed. 1,

2023; https://www.cambridge.org/core/product/identifier/9781009325844/type/

book).

47.   K. Xue, M. M. Yuan, Z. J. Shi, Y. Qin, Y. Deng, L. Cheng, L. Wu, Z. He, J.

D. Van Nostrand, R. Bracho, S. Natali, Edward. A. G. Schuur, C. Luo, K. T.

Konstantinidis, Q. Wang, J. R. Cole, J. M. Tiedje, Y. Luo, J. Zhou, Tundra soil

carbon is vulnerable to rapid microbial decomposition under climate warming. *Nat.*

Clim. Change 6, 595–600 (2016).

48.   G. Pold, A. F. Billings, J. L. Blanchard, D. B. Burkhardt, S. D. Frey, J. M.

Melillo, J. Schnabel, L. T. A. Van Diepen, K. M. DeAngelis, Long-term warming

alters carbohydrate degradation potential in temperate forest soils. *Appl. Environ.*

Microbiol. 82, 6518–6530 (2016).

835 49. J. Séneca, A. Söllinger, C. W. Herbold, P. Pjevac, J. Prommer, E.

836 Verbruggen, B. D. Sigurdsson, J. Peñuelas, I. A. Janssens, T. Urich, A. T. Tveit, A.

837 Richter, Increased microbial expression of organic nitrogen cycling genes in long-

838 term warmed grassland soils. *ISME Commun.* 1 (2021).

839 50. F. Protti☐Sánchez, L. M. Avila, K. Meeran, J. Ingrisch, P. Sigurðsson, K.

840 S. Larsen, B. D. Sigurdsson, I. Janssens, M. Bahn, Soil respiration responses to

841 prolonged warming vary seasonally in a subarctic grassland. *Glob. Change Biol.*

842 31, e70388 (2025).

843 51. T. W. N. Walker, C. Kaiser, F. Strasser, C. W. Herbold, N. I. W. Leblans,

844 D. Woebken, I. A. Janssens, B. D. Sigurdsson, A. Richter, Microbial temperature

845 sensitivity and biomass change explain soil carbon loss with warming. *Nat. Clim.*

846 *Change* 8, 885–889 (2018).

847 52. D. Metze, J. Schnecker, C. L. N. De Carlan, B. Bhattarai, E. Verbruggen, I.

848 Ostonen, I. A. Janssens, B. D. Sigurdsson, B. Hausmann, C. Kaiser, A. Richter,

849 Soil warming increases the number of growing bacterial taxa but not their growth

850 rates. *Sci. Adv.* 10, eadk6295 (2024).

851 53. C. Greening, S. G. Villas-Bôas, J. R. Robson, M. Berney, G. M. Cook, The

852 growth and survival of *Mycobacterium smegmatis* is enhanced by co-metabolism of

853 atmospheric H_2. *PLoS ONE* 9, e103034 (2014).

54.   Q. Qi, J. Zhao, R. Tian, Y. Zeng, C. Xie, Q. Gao, T. Dai, H. Wang, J. He,

K. T. Konstantinidis, Y. Yang, J. Zhou, X. Guo, Microbially enhanced methane

uptake under warming enlarges ecosystem carbon sink in a Tibetan alpine

grassland. *Glob. Change Biol.* 28, 6906–6920 (2022).

55.   H. Wang, E. Lindemann, P. Liebmann, M. Varsadiya, M. M. Svenning, M.

Waqas, S. Petters, A. Richter, G. Guggenberger, J. Barta, T. Urich, Methane-

cycling microbiomes in soils of the pan-Arctic and their response to permafrost

degradation. *Commun. Earth Environ.* 6, 748 (2025).

56.   A. T. Tveit, A. Söllinger, E. M. Rainer, A. Didriksen, A. G. Hestnes, L.

Motleleng, H.-J. Hellinger, T. Rattei, M. M. Svenning, Thermal acclimation of

methanotrophs from the genus *Methylobacter*. *ISME J.* 17 (2023).

57.   Á. Magnússon, P. Vídalín, *Jarðabók. Annað Bindi [The Farm Register of*

Iceland. 2nd Volume] (Copenhagen, 1918).

58.   A. Söllinger, L. S. Ahlers, M. B. Dahl, P. Sigurðsson, C. Le Noir De Carlan,

B. Bhattarai, C. Gall, V. S. Martin, C. Rottensteiner, L. L. Motleleng, E. M. Breines,

E. Verbruggen, I. Ostonen, B. D. Sigurdsson, A. Richter, A. T. Tveit,

Microorganisms in subarctic soils are depleted of ribosomes under short-,

medium-, and long-term warming. *ISME J.* 18, wrae081 (2024).

59. M. B. Dahl, S. Brachmann, A. Söllinger, M. Schnell, L. Ahlers, M.

Wutkowska, K. J. Hoff, N. Nath, V. Groß, H. Wang, M. Weil, M. Piecha, M.

Schaffer, C. Jensen, A. W. Kuss, C. Gall, E. Wimmer, T. Pribasnig, A. T. Tveit, B.

D. Sigurdsson, C. Schleper, A. Richter, T. Urich, Quantifying soil microbiome

abundance by metatranscriptomics and complementary molecular techniques—

Cross☐validation and perspectives. *Mol. Ecol. Resour.*, e14130 (2025).

60. P. C. Brookes, A. Landman, G. Pruden, D. S. Jenkinson, Chloroform

fumigation and the release of soil nitrogen: A rapid direct extraction method to

measure microbial biomass nitrogen in soil. *Soil Biol. Biochem.* 17, 837–842

(1985).

61. C. Voigt, A.-M. Virkkala, G. Hould Gosselin, K. A. Bennett, T. A. Black, M.

Detto, C. Chevrier-Dion, G. Guggenberger, W. Hashmi, L. Kohl, D. Kou, C.

Marquis, P. Marsh, M. E. Marushchak, Z. Nesic, H. Nykänen, T. Saarela, L.

Sauheitl, B. Walker, N. Weiss, E. J. Wilcox, O. Sonnentag, Arctic soil methane sink

increases with drier conditions and higher ecosystem respiration. *Nat. Clim.*

Change 13, 1095–1104 (2023).

62. Y.-J. Chen, P. M. Leung, P. L. M. Cook, W. W. Wong, T. Hutchinson, V.

Eate, A. J. Kessler, C. Greening, Hydrodynamic disturbance controls microbial

community assembly and biogeochemical processes in coastal sediments. *ISME J.*

16, 750–763 (2022).

63. B. Buchfink, K. Reuter, H.-G. Drost, Sensitive protein alignments at tree-

of-life scale using DIAMOND. *Nat. Methods* 18, 366–368 (2021).

64.   P. M. Leung, C. Greening, Greening lab metabolic marker gene

databases, (2020); https://doi.org/10.26180/14431208.v1.

65.   R. Lappan, G. Shelley, Z. F. Islam, P. M. Leung, S. Lockwood, P. A.

Nauer, T. Jirapanjawat, G. Ni, Y.-J. Chen, A. J. Kessler, T. J. Williams, R.

Cavicchioli, F. Baltar, P. L. M. Cook, S. E. Morales, C. Greening, Molecular

hydrogen in seawater supports growth of diverse marine bacteria. *Nat. Microbiol.* 8,

581–595 (2023).

66.   C. Martínez-Pérez, C. Greening, S. K. Bay, R. J. Lappan, Z. Zhao, D. De

Corte, C. Hulbe, C. Ohneiser, C. Stevens, B. Thomson, R. Stepanauskas, J. M.

González, R. Logares, G. J. Herndl, S. E. Morales, F. Baltar, Phylogenetically and

functionally diverse microorganisms reside under the Ross Ice Shelf. *Nat.*

Commun. 13, 117 (2022).

67.   B. J. Woodcroft, S. T. N. Aroney, R. Zhao, M. Cunningham, J. A. M.

Mitchell, R. Nurdiansyah, L. Blackall, G. W. Tyson, Comprehensive taxonomic

identification of microbial species in metagenomic data using SingleM and

Sandpiper. *Nat. Biotechnol.*, doi: 10.1038/s41587-025-02738-1 (2025).

68.   R Core Team, R: A language and environment for statistical computing,

Foundation for Statistical Computing (2025); http://www.r-project.org/.

69.   H. Wickham, *Ggplot2: Elegant Graphics for Data Analysis* (Springer-

Verlag New York, 2016; https://ggplot2.tidyverse.org).

70.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: A flexible trimmer for

Illumina sequence data. *Bioinformatics* 30, 2114–2120 (2014).

71.   S. Chen, Ultrafast one‑pass FASTQ data preprocessing, quality control,

and deduplication using fastp. *iMeta* 2, e107 (2023).

72.   D. Li, C. M. Liu, R. Luo, K. Sadakane, T. W. Lam, MEGAHIT: An ultra-fast

single-node solution for large and complex metagenomics assembly via succinct

de Bruijn graph. *Bioinformatics* 31, 1674–1676 (2015).

73.   D. Hyatt, G. L. Chen, P. F. LoCascio, M. L. Land, F. W. Larimer, L. J.

Hauser, Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* 11, 119 (2010).

74.   D. H. Haft, A. Badretdin, G. Coulouris, M. DiCuccio, A. S. Durkin, E.

Jovenitti, W. Li, M. Mersha, K. R. O'Neill, J. Virothaisakun, F. Thibaud-Nissen,

RefSeq and the prokaryotic genome annotation pipeline in the age of

metagenomes. *Nucleic Acids Res.*, gkad988 (2023).

75.   P.-A. Chaumeil, A. J. Mussig, P. Hugenholtz, D. H. Parks, GTDB-Tk v2:

memory friendly classification with the genome taxonomy database. *Bioinformatics*

38, 5315–5316 (2022).

76.   H. Wickham, M. Averick, J. Bryan, W. Chang, L. D. A. Mcgowan, R.

François, G. Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. Lin, E. Miller,

S. M. Bache, K. Müller, J. Ooms, D. Robinson, D. P. Seidel, V. Spinu, K.

Takahashi, D. Vaughan, C. Wilke, K. Woo, Welcome to the Tidyverse. *J. Open*

Source Softw. 4, 1686 (2019).

77.   B. Elbers, "tidylog: Logging for 'dplyr' and 'tidyr' Functions" (manual,

2020); https://CRAN.R-project.org/package=tidylog.

78.   J. Oksanen, G. L. Simpson, F. G. Blanchet, R. Kindt, P. Legendre, P. R.

Minchin, R. B. O'Hara, P. Solymos, M. H. H. Stevens, E. Szoecs, H. Wagner, M.

Barbour, M. Bedward, B. Bolker, D. Borcard, G. Carvalho, M. Chirico, M. De

Caceres, S. Durand, H. B. A. Evangelista, R. FitzJohn, M. Friendly, B. Furneaux,

G. Hannigan, M. O. Hill, L. Lahti, D. McGlinn, M.-H. Ouellette, E. Ribeiro Cunha, T.

Smith, A. Stier, C. J. F. Ter Braak, J. Weedon, "vegan: Community ecology

package" (manual, 2022); https://CRAN.R-project.org/package=vegan.

79.   B. Bushnell, BBMap: A fast, accurate, splice-aware aligner, (2014); https://

bbmap.org.

80.   P. Danecek, J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard,

A. Whitwham, T. Keane, S. A. McCarthy, R. M. Davies, H. Li, Twelve years of

SAMtools and BCFtools. *GigaScience* 10, giab008 (2021).

81.   D. D. Kang, F. Li, E. Kirton, A. Thomas, R. Egan, H. An, Z. Wang,

MetaBAT 2: an adaptive binning algorithm for robust and efficient genome

reconstruction from metagenome assemblies. *PeerJ* 7, e7359 (2019).

82.   Y.-W. Wu, B. A. Simmons, S. W. Singer, MaxBin 2.0: an automated

binning algorithm to recover genomes from multiple metagenomic datasets.

Bioinformatics 32, 605–607 (2016).

83.   J. Alneberg, B. S. Bjarnason, I. de Bruijn, M. Schirmer, J. Quick, U. Z. Ijaz,

L. Lahti, N. J. Loman, A. F. Andersson, C. Quince, Binning metagenomic contigs

by coverage and composition. *Nat. Methods* 11, 1144–1146 (2014).

84.   S. Pan, X.-M. Zhao, L. P. Coelho, SemiBin2: self-supervised contrastive

learning leads to better MAGs for short- and long-read sequencing. *Bioinformatics*

39, i21–i29 (2023).

85.   G. V. Uritskiy, J. DiRuggiero, J. Taylor, MetaWRAP—a flexible pipeline for

genome-resolved metagenomic data analysis. *Microbiome* 6, 158 (2018).

86.   M. R. Olm, C. T. Brown, B. Brooks, J. F. Banfield, dRep: a tool for fast and

accurate genomic comparisons that enables improved genome recovery from

metagenomes through de-replication. *ISME J.* 11, 2864–2868 (2017).

87.   A. Chklovski, D. H. Parks, B. J. Woodcroft, G. W. Tyson, CheckM2: a

rapid, scalable and accurate tool for assessing microbial genome quality using

machine learning. *Nat. Methods* 20, 1203–1212 (2023).

88.   R. M. Bowers, N. C. Kyrpides, R. Stepanauskas, M. Harmon-Smith, D.

Doud, T. B. K. Reddy, F. Schulz, J. Jarett, A. R. Rivers, E. A. Eloe-Fadrosh, S. G.

Tringe, N. N. Ivanova, A. Copeland, A. Clum, E. D. Becraft, R. R. Malmstrom, B.

Birren, M. Podar, P. Bork, G. M. Weinstock, G. M. Garrity, J. A. Dodsworth, S.

Yooseph, G. Sutton, F. O. Glöckner, J. A. Gilbert, W. C. Nelson, S. J. Hallam, S. P.

Jungbluth, T. J. G. Ettema, S. Tighe, K. T. Konstantinidis, W.-T. Liu, B. J. Baker, T.

Rattei, J. A. Eisen, B. Hedlund, K. D. McMahon, N. Fierer, R. Knight, R. Finn, G.

Cochrane, I. Karsch-Mizrachi, G. W. Tyson, C. Rinke, L. Schriml, P. Hugenholtz, P.

Yilmaz, F. Meyer, A. Lapidus, D. H. Parks, A. M. Eren, J. F. Banfield, T. Woyke,

Minimum information about a single amplified genome (MISAG) and a

metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat.*

Biotechnol. 35, 725–731 (2017).

89.   D. H. Parks, M. Chuvochina, P.-A. Chaumeil, C. Rinke, A. J. Mussig, P.

Hugenholtz, A complete domain-to-species taxonomy for Bacteria and Archaea.

Nat. Biotechnol. 38, 1079–1086 (2020).

90.   S. T. N. Aroney, R. J. P. Newell, J. N. Nissen, A. P. Camargo, G. W.

Tyson, B. J. Woodcroft, CoverM: read alignment statistics for metagenomics.

Bioinformatics 41, btaf147 (2025).

91.   B. Q. Minh, H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams,

A. Von Haeseler, R. Lanfear, IQ-TREE 2: New models and efficient methods for

phylogenetic inference in the genomic era. *Mol. Biol. Evol.* 37, 1530–1534 (2020).

92.   I. Letunic, P. Bork, Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. *Nucleic Acids Res.* 49, W293–W296

(2021).

93.   J. Pinheiro, D. Bates, R Core Team, "nlme: Linear and nonlinear mixed

effects models" (manual, 2023); https://CRAN.R-project.org/package=nlme.

94.   D. Lüdecke, M. S. Ben-Shachar, I. Patil, P. Waggoner, D. Makowski,

performance: An R package for assessment, comparison and testing of statistical

models. *J. Open Source Softw.* 6, 3139 (2021).

95.   E. Kopylova, L. Noé, H. Touzet, SortMeRNA: fast and accurate filtering of

ribosomal RNAs in metatranscriptomic data. *Bioinformatics* 28, 3211–3217 (2012).

1001

Figures

Figure 1. Soil-atmosphere gas fluxes, soil gas concentrations, and C isotope

signatures. *In situ* flux of CO_2 (**A**) and CH4 (**B**) from unwarmed and long-term warmed

soil. Individual data points from two consecutive days are shown, with lines connecting

pairs within a single warming transect used for statistical testing. Concentrations of CO_2

(**C**) and CH_4 (**D**) in unwarmed and long-term warmed soil at different depths (5, 10, and

20 cm, mean±SE are depicted by black lines). Vertical dotted lines denote the average

atmospheric CO2 and CH4 concentration measured on site. Relationship between CO_2

(**E**) and CH_4 (**F**) concentrations and C isotope values in unwarmed and long-term

warmed soil at different depths. The dot-dashed line in (**F**) represents a model following

a Rayleigh distillation function with a fractionation factor (alpha) of 1.020. The vertical

dashed line in (**F**) denotes the detection limit for δ^{13}C-CH_4 in samples with <1.2 ppmv

CH_4. Significant differences and trends are depicted by square brackets and an asterisk

or dot, respectively.

1017

Figure 2. *Ex situ* trace gas oxidation rates. Bulk soil oxidation rates (**A, B, C**) and cell-specific oxidation rates (**D, E, F**) of CO, H_2, and CH_4 obtained from unwarmed, short-term warmed, and long-term warmed incubation treatments. Individual data points are shown together with lines connecting samples from the same warming transect. DW stands for dry weight; asterisks denote significant differences between treatments.

1023

Figure 3. Relative abundance shifts of trace gas oxidizing gene composition

caused by warming. (**A**) Relative abundance of marker genes, calculated as gene

copies per cell. (**B**) Relative abundance values of marker genes in relative proportion to

total assembled gene counts for the most abundant taxa are displayed. Low-abundant

taxa are grouped as "Other" for *coxL* (overall abundance average <3.2% per taxon) and

[NiFe]-hydrogenases (<4.3%). Data shown are from this study (circles; n=4) and from

public metagenomic short reads (BioProject ID PRJNA746424, triangles; n=1). Lines

indicate pairs within a single warming transect that were used for statistical testing.

Significant trends are depicted by square brackets and a dot.

1033

Figure 4. Phylogenomics and differential abundance of non-redundant trace gas oxidizer genomes. (A) The phylogenomic tree of 120 medium- and high-quality metagenome-assembled genomes (MAGs) was constructed using approximately

maximum likelihood estimation with the LG+F+I+G4 model. The scale bar represents

10% sequence divergence, and grey circles at nodes indicate bootstrap support. High-

quality MAGs (completeness > 90%, contamination < 5%) are labeled with stars. All

other MAGs have medium quality (completeness > 50%, contamination < 10%). **(B)**

Completeness and contamination values of MAGs. **(C)** Average relative abundance (%)

of trace gas oxidizing MAGs in unwarmed and warmed soils. **(D)** Log_2 fold change in

relative abundance between warmed and unwarmed soil (positive values = enrichment

in warmed soil). **(E)** Presence/absence of genes involved in trace gas oxidation in

MAGs. Type I CO dehydrogenase (CoxL) for CO oxidation (yellow), high-affinity [NiFe]-

hydrogenases (group 1f, 1l, 1h and 2a) (red), and CH_4 monooxygenases

(PmoA+MmoX) for CH_4 oxidation (blue).

1048

Figure 5. **Transcription of trace gas oxidizing genes encoded by the soil**

community. Transcript abundance of marker genes, calculated as RPKM (**A**), from this

study (circles; n=4) and from public metatranscriptomic short reads (BioProject ID

PRJNA663238, triangles; n=4). Black lines show the mean±SE (n=8). Transcription

ratio (RNA:DNA) of marker genes based on metagenomic and metatranscriptomic short

reads from this study (**B**). Lines indicate pairs within a single warming transect that were

used for statistical testing.

1056

Figure 6. Relative abundance of trace gas oxidizers in global, long-term experimentally warmed soils. Abundances of genes encoding CO dehydrogenases (*coxL*) (**A**),high-affinity [NiFe]-hydrogenases (group 1f, 1h, 1l and 2a)(**B**)*,* and CH_4 monooxygenases (*pmoA + mmoX*) (**C**) in tundra and forest soils under different long-term warming scenarios. The abundance of metabolic marker genes is shown based on metagenomic short reads from two forest soils (48) warmed by +5°C for 8 and 20 years, and a tundra soil (47) warmed by 2–5°C for 1.5 years. The asterisk denotes a significant difference between warmed and control soil (Mann-Whitney test , $p < 0.05$; n=4). Black

1065　lines show the mean±SE.

 Supplementary Materials

 Supplementary Methods

 Gas diffusion model

1069 We fitted a diffusion and model assuming steady state conditions (Eq. 1) where dC_i/dt is

1070 the concentration change of gas i over time in a soil volume [mol m^{-3} sec^{-1}], Di_i is the

1071 concentration change due to diffusion [mol m^{-3} sec^{-1}] and R_i is the concentration change

1072 due to microbial production or consumption [mol m^{-3} sec^{-1}].

1073
$$(1)\ \frac{dC_i}{dt} = Di_i - R_i = 0$$

Di_i was calculated following Fick's second law of diffusion (Eq. 2) adjusted for air-filled

porosity (ε_a; dimensionless) and soil tortuosity (D_S/D_0, i.e., diffusivity in soil relative to air;

dimensionless), where C_i stands for the concentration of gas i [mol m^{-3}], z for depth [m]

and D$_i$ for the diffusion coefficient of gas i in air [m^2 sec^{-1}]. In this initial simulation, we

assume that soil tortuosity and diffusivity are constant throughout the simulated soil

profile.

1080
$$(2)\ Di_i = \frac{d^2 C_i}{dz^2} \cdot D_i \cdot \frac{D_S}{D_0} \cdot \frac{1}{\varepsilon_a}$$

1081

R_i was calculated as a pseudo first-order reaction with a known reaction rate at

atmospheric concentration (eq 3), where R_{atm} is the reaction rate at atmospheric

concentration [mol kg^{-1} d.w.], $C_{i,atm}$ is the concentration of gas i in the atmosphere [mol

m^{-3}], and BD is the bulk density of the soil [kg m^{-3}]. This means that we assume that the

potential soil oxidation rate remains constant with depth, but that the *realized* oxidation

rate with changes with depth in response to changing in gas concentration.

$$(3) \ R_i = R_{atm} \cdot \frac{C_i}{C_{i,atm}} \cdot BD \cdot \frac{1}{\varepsilon_a}$$

1089

1090 Substituting Eq. 2–3 into Eq. 1 gives

1091 $$(4) \ \frac{d^2 C_i}{dz^2} \cdot D_i \cdot \frac{D_S}{D_0} \cdot \frac{1}{\varepsilon_a} - R_{atm} \cdot \frac{C_i}{C_{i,atm}} \cdot BD \cdot \frac{1}{\varepsilon_a} = 0$$

1092 which can be simplified to

1093 $$(5) \ \frac{d^2 C_i}{dz^2} \cdot \frac{1}{C_i} = \frac{R_{atm} \cdot BD}{C_{i,atm} \cdot D_i \cdot \frac{D_S}{D_0}}$$

1094

1095 Substituting the right-hand side of Eq. 5 ($\frac{R_{atm} \cdot BD}{C_{i,atm} \cdot D_i \cdot \frac{D_S}{D_0}}$) for the constant k gives a second

1096 order differential equation

1097 $$(6) \ \frac{d^2 C_i}{dz^2} \cdot \frac{1}{C_i} = k$$

1098 which resolves to

1099 $$(7) \ C_i(z) = c1 \cdot e^{\sqrt{k} \cdot z}$$

1100

1101 We apply the following constraints, (i) $C_i(z=0) = C_{i,atm}$, i.e., the concentration at the soil

1102 surface is the known atmospheric concentration and (ii) $C_i'(z) < 0$, the concentrations

1103 decrease with depth. These conditions are only met when $c1=0$ and $c2=C_{i,atm}$. The depth

1104 profile of a trace gas was therefore described as

1105
$$(8)\ C_i(z) = C_{i,atm} \cdot e^{-\sqrt{k} \cdot z} = C_{i,atm} \cdot e^{-\sqrt{\frac{\frac{R_{atm} \cdot BD}{C_{i,atm} \cdot D_i \cdot D_S}}{D_0}} \cdot z}$$

1106

1107 The flux at depth z can be calculated as:

1108
$$(9)\ F = \frac{dC_i}{dz} \cdot D_i \cdot \frac{D_S}{D_0} = -C_{i,atm} \cdot \sqrt{k} \cdot e^{-\sqrt{k} \cdot z} \cdot D_i \cdot \frac{D_S}{D_0}$$

1109

1110 The surface flux (z = 0) is therefore

1111
$$(8)\ F = -C_{i,atm} \cdot \sqrt{\frac{\frac{R_{atm} \cdot BD}{C_{i,atm} \cdot D_i \cdot D_S}}{D_0}} \cdot D_i \cdot \frac{D_S}{D_0}$$

1112

1113 Parameters applied:

Parameter [unit]		CH$_4$	H$_2$	CO	source / comment
D_i [m^2 sec^{-1}]		$21.0 \cdot 10^{-6}$	$75.6 \cdot 10^{-6}$	$20.8 \cdot 10^{-6}$	
BD [g m^{-3}]		$0.6 \cdot 10^6$ (control) / $0.7 \cdot 10^6$ (warmed)			(reference 1)
$Ds/D0$		0.15 [0.1, 0.2]			fitted to data
ε_a		0.76 (control) / 0.72 (warmed)			calculated from BD based on particle density $2.5 \cdot 10^6$ g m^{-3}
c_{air} [ppmv]		2.1	0.53	0.08	measured for CH$_4$, assumed for H$_2$ and CO
R [mol s^{-1} g^{-1} d.w.]	control, warmed	2.48 ± 0.72 ·10^{-14}, 2.05 ± 0.83 ·10^{-14}	0.94 ± 0.22 ·10^{-14}, 2.11 ± 0.52 ·10^{-14}	7.22 ± 2.18 ·10^{-14}, 20.09 ± 7.74 ·10^{-14}	This study

1114

1115

In situ CH$_4$, H$_2$ and CO uptake

For the _in situ_ H$_2$ and CO uptake measurements, a metal frame was fixed in the soil of a

study plot and the plant material within the frame was removed. Then, an acrylic glass

chamber (edge length 20 cm × 20 cm, total volume of 3,600 mL) was placed on the

metal frame. Gas samples from the chamber were taken after 0, 2, 4, 8, 16, and 32 min

via an embedded butyl rubber stopper using a gas tight syringe (25MDR-LL-GT, SGE

International Pty Ltd, Australia). The gas samples were injected into 12 mL vials (738W,

Labco, UK) prefilled with ultrapure water, allowing the gas to displace the water. 2 mL of

each the gas sample were analyzed using a gas chromatograph (ThermoScientific

Trace 1300, Thermo Fisher Scientific) equipped with a sample loop, a Hayesep column

(SU12875, RESTEK, Bellefonte, Pennsylvania, USA), a Molsieve 5 A column

(PKC17080, RESTEK), a pulsed discharge detector, and a flame ionization detector. A

high-quality gas containing 2.5 ppmv CH$_4$, 2.5 ppmv H$_2$, and 2.5 ppmv CO in N$_2$ served

as standard (HiQ, AGA, Sweden) (29).

1130

DNA qPCR, and sequencing

The qPCR assays were performed in triplicate using a 96-well plate in a QuantStudio 7

Flex Real-Time PCR System (Applied Biosystems) and 16S rRNA gene copy numbers

were quantified against a serially diluted pMA plasmid containing the Escherichia coli

16S rRNA gene. Metagenomic shotgun libraries, including an extraction blank control,

were prepared with the Illumina DNA Prep Tagmentation kit and sequenced on an

Illumina NovaSeq 6000 SP at the Ramaciotti Centre for Genomics, University of New

South Wales. Two samples (Unwarmed 1 and Warmed 1) were deeply sequenced with

an average of 131,415,352 read pairs whereas an average of 35,238,799 read pairs

was generated from the other samples. The negative control returned only 139 read

pairs, suggesting minimal contaminants. Raw metagenomic data were quality assessed

using FastQC v0.11.7 and MultiQC v1.12. The BBDuk function of the BBTools suite

v38.96 (79) was used to remove the 151st base, trim adapters, filter PhiX reads, trim the

3' end at quality threshold of 15 and discard reads below 50 bp in length.

1145

RNA extraction and sequencing

Soil samples (≈40g) were homogenized (CryoMill, Retsch GmbH, Haan, Germany)

under frozen conditions (via liquid nitrogen) and RNA was extracted from each sample

(≈0.5g) using the RNeasy PowerSoil Total RNA Kit (Qiagen, Venlo, Netherlands).

Extractions were done following the manufacturer's protocol, but with the following

modifications: each soil sample was subjected to two cycles of bead-beating (for lysing

and homogenization) with a FastPrep-24 5G (MP Biomedicals, Irvine, CA, USA) to

maximize the recovery of RNA per sample. Extracts from both bead-beating steps were

combined directly after phase separation in a joint capture tube. Extracts were cleaned

with the MEGAclear Transcription Clean-Up Kit (Thermo Fisher Scientific, Waltham,

MA, USA). The RNA concentration was determined with the Qubit RNA BR Assay Kit

(Thermo Fisher Scientific, Waltham, MA, USA) and the RNA quality was analysed with

a Bioanalyzer using the Agilent RNA 6000 Nano Kit (Agilent Technologies, Santa Clara,

CA, USA). Sequence libraries were prepared using the NEBNext Ultra II RNA Library

Prep Kit for Illumina (New England BioLabs, Ipswich, MA, USA) following the

manufacturer's protocol which included reverse transcription. cDNA input was 100–120

ng. cDNA was fragmented and size selection for 250bp was performed with HighPrep

PCR beads (MagBio Genomics Inc., Gaithersburg, MD, USA). The samples were PE

sequenced using a NextSeq 550 System using the NextSeq 500/550 High Output Kit

v2.5 (300 Cycles; Illumina, San Diego, CA, USA) at the sequence facility at Greifswald

University. Metatranscriptome data were filtered for putative mRNA transcripts using

SortMeRNA v.2.1 (95).

1168

1169 **Figure S1. Soil temperatures over the past nine years before sampling.** Average

1170 annual soil temperature at 10 cm depth for unwarmed (green) and long-term warmed

1171 (orange) soil. Lines represent mean temperatures (n=4), and shaded areas indicate

1172 standard deviation across years 2013–2022. Hourly measurements (24.96±0.03 per

1173 day, n=23,837) were first averaged to daily values before calculating annual means.

1174

Figure S2. Edaphic characteristics of unwarmed and long-term warmed soil. DOC

– dissolved organic carbon (**A**), TDN – total dissolved nitrogen (**B**), C_{mic} – microbial

carbon (**C**), N_{mic} – microbial nitrogen (**D**), pH of soil (**E**), field soil moisture content (**F**),

cell number per gram of dry soil estimated based on qPCR integrated with metagenome

data (**G**). Lines connect pairs within a single warming transect that were used for

statistical testing, transect ID is indicated by numbers.

1181

1182

Figure S3. Abundance and transcription of metabolic marker genes encoded by soil community. The abundance of metabolic marker genes is shown based on the metagenomic short reads (A) and metatranscriptomic short reads (B) across unwarmed (n=4) and warmed plots (n=4). Publicly available metagenomic data (BioProject PRJNA746424; unwarmed (n=1) and warmed (n=1)) and metatranscriptomic data generated in 2016 [BioProject PRJNA663238; unwarmed (n=4) and warmed (n=4)] were included.

Figure S4. Summed relative abundances of recovered MAGs. Relative abundance

of MAGs encoding CoxL (A), high affinity [NiFe] hydrogenases (including groups 1h, 1f,

1l and 2a) (B), and PmoA/MmoX (C) were considered as CO oxidizers, H_2 oxidizers,

and CH_4 oxidizers, respectively. The rest of MAGs did not encode any of these marker

enzymes (D), all MAGs included in the dataset encompass both oxidizers and

non-oxidizers (E). Significant differences and trends are depicted by square brackets

and asterisk or dot, respectively. Lines connect pairs within a single warming transect

that were used for statistical testing.