**DNMT1 Coordinates PV Interneuron-Glia Coupling to Maintain Cortical Network Stability and Regulate Behavior**

Jenice Linde[1,2§], Can Bora Yildiz[1,2§], Katharina Vöhringer[1,2], Gerion Nabbefeld[2,3], Severin Graff[2,4], Daniel Pensold[1], Julia Reichard[1,2], Marie Hermanns[1], Christoph Weber-Hamacher[5], Marc Spehr[2,5], Björn Kampa[2,3], Anja Urbach[6], Simon Musall[2,4‡], Geraldine Zimmer-Bensch[1,2 ‡*]

[1] Department for Neuroepigenetics, Institute for Biology II, RWTH Aachen University

[2] Research Training Group RTG2416 "MultiSenses – MultiScales", RWTH Aachen University

[3] Department for Molecular and Systemic Neurophysiology, Institute for Biology II, RWTH Aachen University

[4] Institute of Biological Information Processing (IBI-3) Bioelectronics, Forschungszentrum Jülich

[5] Department of Chemosensation, Institute for Biology II, RWTH Aachen University; 52074 Aachen, Germany

[6] Department of Neurology, Jena University Hospital

[§, ‡] Equal contribution

[*] Corresponding author

# Abstract

Parvalbumin (PV) interneurons are central to cortical network stability and psychiatric vulnerability. Here, we identify DNA methyltransferase 1 (DNMT1) as a key epigenetic regulator linking PV interneuron function to glial and extracellular matrix remodeling. Conditional PV-specific *Dnmt1* deletion combined with single-cell RNA-seq, in vivo electrophysiology, histology, and behavioral analyses revealed that loss of DNMT1 increases PV spiking activity but reduces inhibitory efficacy, leading to network desynchronization and depression- and anxiety-like behavior in mice. These physiological alterations were accompanied by broad, non-cell-autonomous transcriptional changes in astrocytes and oligodendroglial populations, prominently affecting pathways involved in perineuronal-net (PNN) organization and neuron-glia communication. Cell-cell interaction analyses revealed disrupted NRXN-NLGN, TNR-integrin, and semaphoring signaling, consistent with weakened perisomatic adhesion and PNN integrity. Together, our findings demonstrate that DNMT1 maintains inhibitory circuit stability through cell-autonomous regulation of PV interneuron function, which secondarily shapes glial transcriptional states and extracellular scaffolds to preserve cortical network synchronization and emotional behavior.

## Introduction

GABAergic interneurons are critical regulators of cortical circuit dynamics, ensuring precise control over pyramidal neuron activity and maintaining the excitation/inhibition (E/I) balance essential for healthy brain function. Disruption of inhibitory signaling contribute to maladaptive network states and have been implicated in a range of neuropsychiatric disorders, including schizophrenia and major depressive disorder (MDD)[1–3]. However, the etiology of such disorders is inherently multifactorial, due to interactions between genetic predispositions, environmental stressors, and epigenetic mechanisms that converge to shape disease vulnerability[3]. The exact nature of GABAergic dysfunction in MDD therefore remains debated, with contradictory findings further complicating the establishment of a unified mechanistic framework[4,5].

Epigenetic modifications, such as DNA methylation mediated by DNA methyltransferases (DNMTs), form a crucial interface between external stimuli and neuronal function. Aberrant DNA methylation has been documented across multiple neuronal subtypes in psychiatric and neurological conditions, including schizophrenia, epilepsy, and MDD[6,7]. Among DNMTs, DNMT1 has emerged as a key regulator of inhibitory circuit function: altered DNMT1 expression in cortical interneurons has been linked to schizophrenia and disrupted GABAergic signaling[3].

Within the diverse population of inhibitory neurons, parvalbumin-positive (PV) interneurons are especially important to regulate network activity. As fast-spiking interneurons, they synchronize network oscillations and gate excitatory inputs, thereby maintaining cortical E/I balance[8]. Alterations in PV interneuron number or function have been causally linked to neuropsychiatric diseases[9,10], and their vulnerability to chronic stress makes them prime candidates for involvement in MDD pathology[11]. Our previous work also demonstrated that DNMT1 exerts transcriptional control in postmitotic cortical interneurons[12] and regulates synaptic transmission in PV interneurons by modulating endocytosis-dependent GABA reuptake at presynaptic terminals, thereby fine-tuning inhibitory output[13].

Beyond their intrinsic properties, PV interneurons are profoundly shaped by interactions with surrounding glial cells and the extracellular matrix. These include astrocytes and oligodendrocytes, as well as extracellular matrix components, such as perineuronal nets (PNNs). PNNs tightly ensheathe PV interneurons, stabilize synaptic contacts, and limit plasticity, thereby influencing both inhibitory efficacy and network stability[14,15]. Disruption of glial support or PNN integrity has therefore been increasingly recognized as an important contributor to PV interneuron dysfunction[16] and psychiatric disease vulnerability[17].

In this study, we investigated the cell-autonomous and non-cell-autonomous consequences of DNMT1 function in PV interneurons to reveal how epigenetic dysregulation contributes to cortical circuit instability and disease vulnerability. To this end, we used conditional PV-specific *Dnmt1* knockout mice and combined single-cell transcriptomics, electrophysiology, histological analyses, and behavioral

assays. *Dnmt1* deletion reduced the inhibitory impact of PV interneurons on network
activity in vivo and altered oligodendroglial transcriptional programs converging on
PNN regulation and neuron-glia communication. This dual effect identifies DNMT1
as a central epigenetic coordinator of inhibitory circuit function, linking intrinsic PV
interneuron activity with extrinsic glial and extracellular matrix support. Together, our
findings reveal how epigenetic perturbations in inhibitory neurons propagate across
cellular domains, thereby destabilizing cortical networks and promoting behavioral
phenotypes reminiscent of MDD.

# Results

### *Dnmt1* deletion in PV interneurons alters cortical network activity across
### different cortical areas

90
91

Building on a PV-specific conditional *Dnmt1* knockout mouse line[13,18], we already
demonstrated that DNMT1 regulates GABAergic transmission in PV interneurons
through regulation of endocytosis-associated gene expression[13]. Given that cortical
interneuron dysfunction and dysregulated *Dnmt1* expression have both been
implicated in neuropsychiatric diseases[19], we next examined the functional
consequences of *Dnmt1* deletion in PV interneurons at the network level. To this
end, we performed in vivo electrophysiological recordings in the primary visual cortex
(V1) of awake, behaving *Pvalb-Cre/tdTomato/Dnmt1 loxP$^2$* knockout (KO) and *Pvalb-
Cre/tdTomato* control mice using Neuropixels probes (Fig. 1a).

To selectively modulate PV interneuron activity, we used viral injections to induce
Cre-dependent expression of the blue light-sensitive opsin Channelrhodopsin-2
(ChR2) in V1 PV interneurons. This allowed us to isolate PV spiking activity within
the recorded neural population based on their short-latency responses to 10-ms-long
blue light pulses (Fig. 1b). Spontaneous firing rates of PV interneurons in *Pvalb-
Cre/Dnmt1*-KO mice were elevated relative to controls, suggesting that DNMT1 is
involved in restraining PV interneuron excitability and confirming our previous in vitro
results[13].

When considering all recorded cortical neurons, including excitatory cells,
spontaneous firing rates were also increased in KO mice (Fig. 1c). Moreover,
optogenetic activation of PV interneurons strongly inhibited cortical activity in
controls but was markedly less effective in KO mice (Fig. 1d). This indicates that the
inhibitory impact of PV interneurons on cortical networks was reduced despite their
hyperactivity, potentially due to homeostatic network adaptation.

Visual stimulation further revealed altered cortical responses in KO mice. Compared
to controls, stimulus-evoked activity in V1 was weaker, temporally delayed, and less
precise, with reduced stimulus specificity across cortical layers (Fig. 1a, e–h,
Extended Data Fig. 1a–c). In line with earlier findings[12,20], visual responses in control
mice showed a well-defined laminar organization, with the earliest and strongest
responses in layer 4 (0.3 – 0.5 mm) and subsequent activation of the superficial and

infragranular layers, as evident in the local field potentials (LFPs; Fig 1g, upper panels) and current source densities (CSD; Fig. 1g, lower panels). In contrast, excitatory responses across all cortical layers were reduced in KO mice and the inhibitory rebound component typically associated with feedback inhibition was absent (Fig. 1h). These alterations underscore an impaired contribution of PV interneurons to cortical information processing in KO mice. Consistent with the weakened sensory responses in KO mice (Fig. 1e), we also found fewer stimulus-responsive neurons with a lower response specificity (Extended Data Fig. 1c).

To assess whether these impairments were specific to the visual cortex or generalize to other sensory modalities, we also recorded cortical activity in the primary somatosensory cortex (S1) during whisker stimulation. In controls, tactile input evoked robust and temporally precise responses, peaking in layer 5, followed by a characteristic inhibitory rebound reflecting feedback inhibition (Extended Data Fig. 1d, e). In contrast, *Dnmt1*-KO mice showed markedly weaker and less temporally precise responses that were prolonged and lacked the inhibitory rebound that is typically mediated by PV interneurons (Extended Data Fig. 1d, e). Optogenetic activation of PV interneurons during tactile stimulation further confirmed their reduced efficacy: while PV activation strongly suppressed peak responses in controls, it had little effect in KO mice (Extended Data Fig. 1f). These findings therefore match our observations in V1 and indicate that DNMT1 loss in PV interneurons impairs their inhibitory contribution across primary cortical areas.

Next, we extended our recordings to examine higher-order cortical dynamics, focusing on the anterolateral motor cortex (ALM), a prefrontal region engaged in decision-making and evidence accumulation. Because ALM is strongly recruited during goal-directed behaviors, we sought to determine whether DNMT1 loss in PV interneurons also affects higher-order cortical dynamics during task engagement. We therefore trained mice on a visual evidence accumulation task, in which they reported the side with the higher number of cues by licking a corresponding waterspout (Extended Data Fig. 2e). To exclude general learning deficits, we first examined spatial learning and memory by conducting a Morris water maze test. KO and control mice located the hidden platform with similar accuracy and efficiency, showing no genotype-dependent behavioral impairments (Extended Data Fig. 2a–d). Similarly, KO mice acquired the visual evidence accumulation task as efficiently as controls, with no differences in learning curves or performance (Extended Data Fig. 2f, g).

Electrophysiological recordings nevertheless revealed clear alterations in network activity. Similar to V1, ALM neurons in KO mice showed attenuated neural responses to visual stimulation (Extended Data Fig. 2h) and optogenetic activation of PV interneurons again confirmed their reduced inhibitory efficacy (Extended Data Fig. 2i). Importantly, optogenetic activation of PV interneurons impaired decision-making in both genotypes, demonstrating that ALM function remained essential for task performance (Extended Data Fig. 2j).

Together, these findings demonstrate that DNMT1 loss in PV interneurons reduces
their inhibitory efficacy across cortical regions, spanning from primary sensory areas
to prefrontal circuits critical for decision-making. However, despite these robust
physiological alterations, basic perceptual, learning, and memory capacities were
preserved, suggesting that compensatory network mechanisms can buffer the
functional consequences of impaired PV interneuron control.

**DNMT1 deficiency in PV interneurons impairs gamma synchrony and induces**
**depression-like behavior**

PV interneurons are critically involved in the generation of cortical gamma
oscillations[21,22]. Strikingly, alongside the altered firing patterns in V1 (Fig. 1g, h)
*Dnmt1*-KO mice also displayed severely reduced visually evoked gamma oscillations
(Fig. 1i, j). Because gamma synchrony depends on the precise timing of PV
interneuron output, our findings suggest that DNMT1 is also required to maintain the
rhythmic inhibitory drive underlying fast cortical coordination. Importantly, impaired
gamma oscillations are a well-documented hallmark of neuropsychiatric disorders,
including schizophrenia, bipolar disorder, and major depressive disorder[23]. In
patients with depression, reduced gamma power has been associated with cognitive
inflexibility, impaired attention, and diminished affective regulation, while animal
models consistently link PV dysfunction and gamma desynchronization to
depression-like behaviors[22,24,25].

Notably, bulk RNA sequencing also revealed transcriptional changes in
*Pvalb☐Cre/Dnmt1*-KO mice resembling those in models of depression and anxiety
(Extended Data Tab. 1, 2). This raised the question whether DNMT1 loss in PV
interneurons might trigger mood disorder-related behaviors. Spontaneous home-
cage activity provides a sensitive measure of innate motivation and has been widely
used to detect depression- and apathy-like phenotypes in rodents[26]. Monitoring
home-cage behavior revealed consistently reduced movements in KO mice (Fig.
2a, b), a change not explained by visual deficits (Extended Data Fig. 2) or motor
deficits. KO mice performed normally in rotarod (Fig. 2c) and ladder rung tests[18], but
showed a shorter latency to fall in the wire hang test (Fig. 2d), which coincided with
significantly increased body weight (Fig. 2e). Increased body weight is often
comorbid with depressive symptoms[27], and the weight gain of KO mice could
therefore reflect reduced physical activity and energy expenditure. In support of this,
food intake was unchanged (Extended Data Fig. 3a–c), and appetite-related
hormone levels, including ghrelin, insulin, and PYY, were unaffected (Extended Data
Fig. 3d). We also detected elevated leptin levels likely reflecting increased adipose
tissue rather than altered endocrine regulation[28,29]. Moreover, immunostaining of the
arcuate nucleus confirmed that PV-positive cells are largely distinct from the major
neuronal populations regulating feeding, showing no overlap with POMC neurons
and only partial overlap with AgRP neurons (Extended Data Fig. 4, 5)[30]. This makes
it unlikely that *Dnmt1* deletion in PV cells directly interferes with hypothalamic
feeding circuits as a driver of weight gain. Together, the obesity and diminished

motor activity in KO mice suggest lower energy expenditure and diminished innate
motivation consistent with an apathy-like state[31].

Additional behavioral assays further supported an apathetic and anhedonic
phenotype of *Dnmt1* KO mice. Nest-building, an innate behavior essential for heat
conservation and shelter, is a sensitive readout of apathetic states in rodents[32].
Using a 1–5 ordinal scale, we found significantly lower nest scores in *Dnmt1* KO
mice (Fig. 2f), typically shredding but failing to assemble the nesting material within
16 hours. Assessment of coat state, a marker of self-care and motivation[33,34], also
revealed systematic neglect in *Dnmt1*-KO mice, with significantly higher cumulative
scores compared to controls indicating a deterioration of their overall grooming and
coat care (Fig. 2g).

Consistent with lower self-motivated activity, voluntary wheel running was strongly
reduced (Fig. 3a), particularly during the dark phase, suggesting diminished
motivation for rewarding behaviors[35,36]. KO males also displayed reduced sexual
motivation, mounting females less frequently (Fig. 3b), and showing no attraction to
female urine (Fig. 3c), although general social interaction remained intact (Fig. 3b).
Taken together, *Dnmt1*-deficient mice show a cluster of behavioral traits, including
apathy, anhedonia, and reduced spontaneous activity, that collectively reflect a
dysphoric state resembling depressive symptoms.

**Dnmt1 deletion in PV interneurons leads to increased anxiety in mice**

To assess whether *Dnmt1* KO mice also display altered anxiety levels, we performed
the novelty-suppressed feeding (NSF) test, which measures the conflict between
food-seeking motivation and anxiety in a novel environment (Fig. 3d). After 24 hours
of food deprivation, both genotypes showed similar weight loss (Extended Data Fig.
6a), implying a similarly strong motivation to obtain food.

Nevertheless, *Dnmt1*-KO mice took significantly longer to approach and consume a
food pellet centrally placed in an open arena (Fig. 3e) and foraged closer to the walls
of the arena, reflected by an increased mean distance to the pellet (Fig. 3d). To
confirm that this effect reflected anxiety rather than altered motivation, mice were
transferred to their home-cages immediately after they started biting the pellet. With
the food deprivation still serving as an active motivator in this familiar environment,
KO mice approached the pellet faster than controls (Fig. 3f) and consumed equal
amounts within 5 minutes (Extended Data Fig. 6b). These results demonstrate that
DNMT1 loss in PV interneurons increases novelty-induced anxiety, further
complementing the depression-like phenotype of KO mice.

**Single-nucleus transcriptomics reveals non–cell-autonomous glial remodeling**

The combined electrophysiological and behavioral results show that DNMT1 loss in
PV interneurons disrupts inhibitory control, weakens gamma synchrony, and gives
rise to a spectrum of depression- and anxiety-related behaviors. Notably, DNMT1
loss not only increased firing of PV interneurons but also elevated spontaneous

activity across the broader neuronal population, consistent with an adaptive network
response to altered inhibition. These findings highlight the cell-autonomous role of
DNMT1 in PV interneurons but also the need to understand how such local
dysfunctions reverberate through cortical circuits.

Given that interactions between inhibitory and excitatory neurons are also tightly
regulated by glia cells, we next turned to single-nucleus RNA sequencing to assess
whether *Dnmt1* deletion in PV interneurons exerts non-cell-autonomous effects on all
surrounding cell types. This approach enabled us to capture transcriptional
adaptations across neurons and glial populations, providing a systems-level view of
how DNMT1-dependent dysfunction in PV interneurons destabilizes cortical
networks.

Unsupervised clustering of single nuclei isolated from the cortical tissue of adult
*Dnmt1*-KO and control mice revealed the expected diversity of neuronal and glial cell
types, including excitatory neurons, PV and non-PV interneurons, oligodendrocytes,
astrocytes, and microglia (Fig. 4a, b). Interestingly, although *Dnmt1* deletion was
restricted to PV interneurons, relative proportions of excitatory subclasses were also
altered. Population analysis revealed increased proportions of layer 2/3
intratelencephalic (IT) and layer 4 sensory neurons, combined with a marked
reduction in layer 5/6 IT neurons (Extended Data Fig. 7). This underrepresentation of
layer 5 neurons was also observed in earlier studies and likely reflects a combination
of factors, including slower fixation and RNA degradation in deeper cortical layers,
mechanical fragility of large nuclei with low nucleus-to-cytoplasm ratios, and
inefficient recovery of long, chromatin-associated transcripts characteristic of layer 5
pyramidal cells[37–40]. Consequently, apparent shifts in excitatory subclass proportions
in split-pool snRNA-seq datasets should be interpreted with caution, as they may
partly arise from these technical biases rather than genuine biological differences[37].

We further detected broad and robust transcriptional alterations across multiple cell
types between samples from control and KO mice, with prominent changes in
excitatory neurons of layers 4–6 (Fig. 4c). Notably, the differentially expressed genes
of excitatory neurons in layer 4, the cortex's principal thalamorecipient layer which is
also particularly populated by PV interneurons[41], were enriched for gene ontology
(GO) terms related to synaptic transmission, structure and organization. These
results are consistent with altered input-layer gating due to disrupted inhibitory
control after *Dnmt1* deletion (Fig. 4d). Since DNMT1 loss was selective to PV
interneurons, these alterations likely reflect secondary, non-cell-autonomous effects.

A plausible mechanism is that the increased GABAergic transmission we observed
electrophysiologically (Fig. 1b) directly affects the surrounding network through
altered synaptic signalling. To systematically explore such effects, we next applied
cell–cell communication analysis (CellChat) to identify changes in signaling
pathways linking PV interneurons to surrounding neurons and glial populations.
Differential interaction analyses showed redistributed communication strength and
altered signaling pathways between PV interneurons and excitatory subclasses as
well as glial populations in the *Dnmt1*-KO mice (Fig. 4e, f).

Communication between excitatory neurons and PV interneurons showed a
layer specific redistribution. Outputs from PV cells were prominently increased
toward layer 4 sensory neurons, aligning the strongest inhibitory reweighting with the
primary sensory input gateway. Modest increases occurred toward layer 5/6 IT and
layer 6 corticothalamic neurons, whereas projections to layer 2/3 IT and deep-layer
extra-telencephalic neurons were slightly reduced (Fig. 4e, f). This laminar
imbalance suggests a shift toward input layer-centric inhibition that may favor rapid
sensory gating but weaken distributed recurrent stability. Conversely, inputs from
multiple excitatory neurons to PV interneurons declined, potentially explaining the
reduced inhibitory efficacy despite PV hyperactivity (Fig. 4e, f).

The decline in inputs to PV interneurons suggests a potential compromise in the
integrity of perineuronal nets (PNNs), a specialized extracellular matrix structure that
is critical for stabilizing synaptic contacts onto PV cells and supports their high-
frequency firing[14,15]. As PNNs require continuous support from the oligodendrocyte
lineage, we next looked at their interactions with PV interneurons. Although overall
oligodendrocyte lineage proportions were unchanged (Fig. 5a), we detected a
decoupling of the PV–oligodendrocyte precursor cells (OPC) axis with decreased
signaling in both directions as well as a selective reduction in newly mature
*Pdgfc* positive perineuronal oligodendrocytes[42,43] that synthesize, organize, and
anchor extracellular matrix (ECM) constituents (Fig. 4d, Fig. 5b). Concordantly,
perineuronal oligodendrocytes in *Dnmt1*-KO mice downregulated genes implicated in
stress resilience, ECM remodeling, and structural programs (*Sgk1*, *Fkbp5*, *Bcl2l1*,
*Htra1*, *Mobp*, *Zbtb16*), indicating a reduced capacity to sustain high demand
perisomatic coverage and ECM organization (Extended Data Fig. 8). Notably, PV
interneuron–oligodendrocyte and reciprocal oligodendrocyte–PV connections were
significantly reweighted in *Dnmt1*-KO mice, indicating a disruption in PV–
oligodendrocyte coupling (Fig. 5b–d).

Therefore, we next examined which specific ligand–receptor pairs contributed most
strongly to these altered communication probabilities. Outgoing signaling from PV
interneurons to oligodendroglial cells and incoming signaling from oligodendroglial
populations to PV interneurons are shown in Figure 5d and 5e, respectively. Overall,
more extensive alterations were detected in the incoming direction, suggesting that
PV interneurons become particularly sensitive to glia-derived cues when DNMT1 is
lost. In contrast, PV-to-glia signaling showed fewer but distinct changes, mainly
involving PV-OPC interactions. Specifically, NRXN2–NLGN1 signaling was
increased in *Dnmt1*-KO samples, whereas CNTN1–NRCAM communication was
reduced, implying weakened contact-mediated coupling[44].

Among glia-to-PV pathways, several axes were markedly affected across OPCs and
mature oligodendrocyte subtypes. NRG1–ERBB4 signaling from OPCs to PV
interneurons was significantly diminished, indicating a loss of trophic support
normally promoting PV excitability and metabolic stability[45,46]. In contrast, SEMA–
PLXN signaling was enhanced, particularly from differentiated oligodendrocyte
populations, potentially reflecting a structural remodeling response. Notably,

semaphorins have been reported as constituents of PNNs[47]. PTPRM–PTPRM
signaling from perineuronal oligodendrocytes to PV cells, as well as NRXN–NGLN
signaling from OPCs, was reduced (Fig. 5e). PTPRM can mediate homophilic
interactions, known to be involved in cell-cell adhesion and synapse formation[48]
while NRXN–NGLN signaling operates as an axo-glial communication pathway
between axonal neurexin and oligodendrocytic neuroligin. NGLN3 is expressed in
oligodendrocyte precursor and mature cells, where it interacts with axonal NRXN to
promote myelin formation, maintenance, and oligodendrocyte differentiation[49].
Together with the reduced CNTN1–NRCAM interactions (Fig 5d) and attenuated
TNR–integrin inputs onto PV somata (Fig. 5e), these changes indicate deficiencies in
perisomatic ECM anchoring and PNN cohesion[50–52]. Collectively, these alterations
indicate a reorganization of neuron-glia communication that compromise PNN
stabilization and PV synaptic integration, leading to a reduction in inhibitory efficacy
of PV interneurons.

**Histological validation confirms perineuronal net deficits**

To validate the results of the cell-cell interaction analyses, we next aimed to visualize
potential changes in the PNN formation. Indeed, consistent with the predicted
disruptions in neuron–glia signaling, histological analysis revealed a reduction of
Wisteria floribunda agglutinin (WFA)-labeled PNNs in the *Dnmt1*-KO mice (Fig. 6a–
d, Supplementary Video 1, 2). This points to a disruption of the structural support
mechanisms that normally stabilize PV interneuron function. While the fraction of PV
cells enwrapped by PNNs remained unchanged, WFA labeling intensity was
significantly decreased (Fig. 6e, f). Given that PNNs stabilize PV firing properties and
gamma synchrony[15,53], these findings implicate glial and extracellular matrix
abnormalities as amplifiers of PV interneuron deficits, contributing to both the
physiological (reduced PV efficacy, weakened gamma) and behavioral
(apathy/anhedonia/anxiety) phenotypes in *Dnmt1*-KO mice.

Together, these findings provide a mechanistic bridge from DNMT1 loss in PV
interneurons to disrupted inhibitory signaling, defective PV-glia communication and
impaired PNN integrity, ultimately destabilizing the cellular ecosystem required for
cortical network balance and gamma synchrony.

# Discussion

Here we show that deleting *Dnmt1* specifically in cortical PV interneurons reshapes
inhibition across multiple scales, including cellular, network, and behavior,
culminating in depression- and anxiety-like phenotypes. In vivo Neuropixels
recordings revealed increased spontaneous firing of PV cells and elevated baseline
activity in surrounding cortical populations, accompanied by a reduced inhibitory
impact of PV activation on local circuits. The dominant consequence of *Dnmt1*
deletion is therefore not reduced PV interneuron activity but diminished synaptic
efficacy and a corresponding loss of inhibitory control. This is consistent with our

previous finding that *Dnmt1* loss enhances GABAergic vesicle replenishment and
release at PV terminals[13]. We interpret the global rise in excitatory neuron firing and
the blunted response to PV photostimulation as a homeostatic network adaptation to
counteract persistently elevated GABAergic drive by down-weighting inhibitory
influence, a phenomenon observed in vitro and in vivo when networks strive to
stabilize E/I balance[54]. Moreover, because PV interneurons form GABAergic
autapses that regulate their fast-spiking dynamics and contribute to gamma
synchrony[55], the observed increase in PV firing may also represent an adaptive
response to altered self-inhibition. These network adjustments likely contribute to the
smaller, delayed, and less precise visually evoked responses and the loss of a
feedback-inhibitory rebound across layers.

Despite these physiological changes, basic perceptual and learning capacities
remained intact, suggesting a circuit-level compensation potentially via subcortical
pathways that sustain task performance even when cortical precision is reduced.
Nevertheless, DNMT1 loss degraded PV-dependent network dynamics across
sensory (V1, S1) and prefrontal (ALM) cortical areas, PV-mediated inhibition was
weakened, and visually evoked gamma oscillations were markedly reduced, in line
with the central role of PV cells in generating gamma rhythms[21,22].

*Dnmt1*-KO mice exhibited an apathy- and anhedonia-like phenotype, including
reduced home-cage activity, poor nest construction, deteriorated coat condition,
diminished wheel running and sexual motivation, and prolonged latency in novelty-
suppressed feeding. Food intake and circulating ghrelin, insulin, and PYY levels
remained unaltered (Extended Data Fig. 4a–d), whereas body weight and leptin
were elevated (Fig. 2e, Extended Data Fig. 4d), indicating reduced energy
expenditure rather than primary hypothalamic dysregulation. Correspondingly, PV
cells in the arcuate nucleus showed no overlap with POMC and only partial overlap
with AgRP populations. These findings point to cortical, rather than hypothalamic,
mechanisms underlying the dysphoric phenotype, in line with evidence linking PV
interneuron dysfunction and impaired gamma synchrony to mood disorders in both
humans and rodent models[11,22,27,56,57].

A central point of this study is that a PV-restricted, cell-autonomous manipulation of
*Dnmt1* expression propagates across the network, leading to widespread, non-cell-
autonomous alterations. Single-nucleus RNA-seq uncovered a remodeling of
excitatory neuron composition as well as their transcriptional rewiring of genes
involved in synaptic structure and transmission. Consistent with these molecular
changes, circuit-level analyses demonstrated a laminar redistribution of inhibitory
signaling. Communication between excitatory neurons and PV interneurons showed
a marked layer-specific reorganization, with PV outputs prominently reweighted
toward layer 4 while connections to layer 2/3 and deep-layer extratelencephalic
neurons were modestly reduced. Such preferential strengthening of feedforward
inhibition within the sensory input layer resembles canonical thalamocortical PV
architectures, where fast-spiking PV cells mediate rapid inhibitory gating of sensory
signals[58,59]. Modest increases in PV targeting of layer 5/6 subtypes may reinforce

corticothalamic feedback loops, whereas diminished supragranular inhibition could
weaken recurrent excitatory–inhibitory stability across cortical columns[60,61].
Interestingly, excitatory drive onto PV cells was reduced, in line with the notion that
PV interneurons can become hyperactive yet functionally decoupled when excitatory
afferents weaken, resulting in ineffective inhibition despite elevated firing[56,62].
Collectively, these laminar rebalancing effects suggest an input layer-centric
inhibitory bias that overemphasizes sensory gating at the cost of distributed network
stability.

Similarly, communication between PV interneurons and oligodendroglial cells was
particularly reweighted in the cortex of KO mice, accompanied by a selective
depletion of *Pdgfc*-expressing perineuronal oligodendrocytes and reduced
expression of genes maintaining ECM integrity. Oligodendrocyte lineage cells are
crucial signaling partners for interneurons, responding dynamically to inhibitory
signals via GABA receptors[63]. The coordinated loss of key ligand–receptor
interactions between PV cells and oligodendroglial cells, such as TNR–integrin, and
NRXN–NLGN, points to a breakdown of perisomatic adhesion and PNN cohesion,
leading to impaired network connectivity and plasticity[14].

Genetic deletion of TNR disrupts PNN morphology, causing irregular net structuring
around cortical interneurons and altered conduction velocity, highlighting its
importance for cohesive perisomatic ECM[64]. Integrin receptors, in turn, contribute to
synaptic stabilization by binding ECM components such as TNR, reinforcing PNN
architecture and modulating excitatory–inhibitory balance[17]. On the synaptic
adhesion side, neurexins (NRXNs) and neuroligins (NLGNs) form transsynaptic
complexes that coordinate synapse formation and maintain structural adhesion and
communication between neurons[65,66]. Perturbations in NRXN–NLGN signaling have
been shown to destabilize synaptic alignment and weaken the structural cohesion of
perisomatic synapses[67,68]. Moreover NRXN–NGLN signaling mediates axo-glial
communication by coupling presynaptic neurexins on axons to neuroligin-3 (NGLN3)
on oligodendrocyte lineage cells. NGLN3 is expressed in both precursor and mature
oligodendrocytes, where its interaction with axonal NRXNs facilitates differentiation,
myelin formation and stability[49].

As a consequence of these disrupted neuron-glia interactions, elevated PV activity
and altered spiking patterns may drive activity-dependent glial plasticity that
reshapes PNN and neuro-glial positioning. Histological validation confirmed this
systems-level implication, showing diminished net intensity of WFA-labeled PNNs in
KO mice. This is consistent with the established role of PNNs in maintaining PV
interneuron excitability, synaptic integration, and gamma synchrony[14]. Indeed, PNN
degradation has been shown to weaken gamma oscillations and cognitive
precision[14,69], providing a mechanistic link between synaptic inhibition and network
disruptions.

Because PNNs stabilize PV excitability and support oscillatory coherence, their loss
provides a functional bridge between reduced PV efficacy, weakened gamma power,
and mood-related behavior. We propose a feed-forward model in which DNMT1 loss

increases presynaptic GABA release, triggering homeostatic desensitization of
inhibitory signaling and glial remodeling that culminates in ECM degradation and
network desynchronization. The resulting disruption of PNNs may further
destabilizes PV function, amplifying network desynchronization and dysphoric
behavior.

This cascade reconciles diverse GABAergic findings in mood disorders: alterations in
PV efficacy, gamma coherence, and PNN integrity can occur without cell loss,
reflecting broader neuron-glia-ECM dysregulation. By positioning DNMT1 as an
upstream regulator of this axis, our results highlight an epigenetic mechanism by
which inhibitory microcircuits coordinate with myelination and extracellular scaffolds
to sustain cortical timing.

## Methods

### Animals

Transgenic *Pvalb-Cre/tdTomato/Dnmt1 loxP²*-mice were used as knockout animals
and *Pvalb-Cre/tdTomato*-mice as controls. Unless stated differently, all experiments
were conducted using male mice aged 3–6 months. Both mouse lines were
established on a C57BL/6-background by crossing in the *Pvalb-Cre* line (obtained
from Christian Huebner, University Hospital Jena, Germany and described in
Hippenmeyer et al.[70]) and the *tdTomato* transgenic reporter mouse line (obtained
from Christian Hübner, University Hospital Jena, Germany and described in Madisen
et al.[71]). To obtain *Dnmt1*-knockout mice, *Dnmt1* floxP² mice (B6;129Sv-
Dnmt1[tm4Jae]/J, Jaenisch laboratory, Whitehead Institute; USA; Jackson-Grusby et
al.[72]) were crossed in, in which LoxP-sites are flanking exons 4 and 5 of the *Dnmt1*
gene. This results in a null *Dnmt1* allele upon Cre-mediated excision[72]. Correct
genotypes were validated with the following primers: *Dnmt1* forward 5′-
GGGCCAGTTGTGTGACTTGG and reverse 3′-CCTGGGCCTGGATCTTGGGGA
(results in a 334 bp wildtype and 368 bp mutant band); *tdTomato* wildtype forward 5′-
AAGGGAGCTGCAGTGGAGTA, wildtype reverse 3′-
CCGAAAATCTGTGGGAAGTC, mutant forward 5′-CTGTTCCTGTACGGCATGG
and mutant reverse 3′-CTGTTCCTGTACGGCATGG (results in a 297 bp wildtype
and 196 bp mutant band); *Pvalb-Cre* forward 5′-
AAACGTTGATGCCGGTGAACGTGC and reverse 3′-
TAACATTCTCCCACCGTCAGTACG (resulting in a 214 bp fragment). All animal
procedures were performed in strict compliance with the EU directives 86/609/EWG
and 2007/526/EG guidelines for animal experiments and were approved by the local
governments (Thueringer Landesamt, Bad Langensalza, Germany and Landesamt
für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen, Recklinghausen,
Germany). Unless indicated otherwise, animals were housed under standard
housing conditions: mice were kept in compatible groups, under 12 h light/dark
conditions, with ad libitum access to food and water, and nestlets and tunnels as

environmental enrichments. The weight of individual animals was determined prior to
behavioral experiments or prior to sacrificing and brain preparation.

**Electrophysiological recordings**

**Surgeries**

For all surgeries isoflurane in oxygen was used as anesthesia (3% isoflurane for
induction, 1–2% for maintenance), and subcutaneous injections of carprofen (4
mg/kg, Rimadyl, Zoetis GmbH) and buprenorphine (0.1 mg/kg, Buprenovet sine,
Bayer Vital GMBH) for analgesia. Eye ointment (Bepanthen, Bayer Vital GmbH) was
applied to their eyes to prevent damage from drying out. After medially incising the
dorsal skin of the head, it was pushed outward and fixed in place utilizing tissue
adhesive (Vetbond, 3M). Dental cement (C&B Metabond, Parkell; Ortho-Jet, Lang
Dental) was applied on top of the tissue adhesive to create a support structure for a
custom-built circular headbar.

Viral injections (AAV1.shortCAG.dlox.hChR2(H134R).WPRE.hGHp, Zurich Vector
Core, viral titer = ~8x1012 vg/ml) were carried out according to the experimental
scheme. In naive mice (Fig. 1a–I, 2a–c), S1 and V1 were targeted unilaterally,
whereas in trained animals (Fig. 2g–i, Extended Data Fig. 2b, c) bilateral injections in
ALM were conducted. The positions of all injections were chosen by stereotactic
coordinates (ALM: AP 2.5/ ML -1.5; V1: AP -4/ ML -2; S1: AP -2/ ML -3.5). Injections
were conducted using a thin glass pipette with a micropump (Nanoject3, Drummond
Scientific) after thinning out the skull with a dental drill. At each site, two quantities of
200 nl were injected at different cortical depths (300/600 μm), adding up to a total
volume of 400 nl virus per target region. The 200 nl were injected in 20 steps using
10 nl portions at 14 s intervals. For electrophysiological recordings, circular
craniotomies were performed using a biopsy puncher and then covered with a round
coverslip of similar size using light-curable dental cement (DE Flowable composite,
Nordenta) to fix it in place. The coverslips contained a small opening to later allow
access for the Neuropixels probe. Between recordings, this opening was covered
with silicon. A thin layer of cyanoacrylate (Zap-A-Gap CA+, Pacer technology) was
used to cover the skull. After the surgery, buprenorphine (0.1 mg/kg body weight)
was again injected subcutaneously. The animal's home cage was kept on a heating
pad following the surgery until the mouse recovered from anesthesia. To prevent
inflammation of the surgical wounds, the animals received a subcutaneous injection
of carprofen on post‑op days 1 and 2 while the animal's drinking water was
supplemented with antibiotics (enrofloxacin 0.025 mg/ml) and analgesics
(buprenorphine 0.0125 mg/ml) for at least three days. Progression of wound healing
and the animal's general health were scored daily following the surgery.

To record from naive animals, two mice per genotype were utilized; all of them were
approx. 6 months of age during surgery. For behavioral analysis and recordings in
trained mice, 7 animals (3 control/4 knockout) were used; they were approx. 8 weeks
old during surgeries.

## Neuropixels recordings

Electrophysiological recordings were carried out in head-fixed mice using Neuropixels 1.0 probes. Recordings were carried out on four consecutive days. Probes were painted with DiD cell labeling solution (Invitrogen V22887) before each recording to later identify recording sites in fixed brain slices.

The stimulation protocol was started 5–10 min after insertion and we recorded high pass-filtered data above 300 Hz at 30 kHz and low pass filtered signals between 0.5–100 Hz at 2.5 kHz from the bottom 384 channels of the Neuropixels probe (~3.8 mm active recording area). Signals were acquired with an external Neuropixels PXIe card (IMEC, Belgium) used in a PXIe-Chassis (PXIe-1071, National Instruments, USA). Triggers and control signals for different stimuli were separately recorded as analogue and digital signals using the SpikeGLX software (Janelia Farm Research Campus, USA; Bill Karsh, https://github.com/billkarsh/SpikeGLX).

The visual stimulus to test sensory responses in V1 and ALM consisted of a 0.1-s long full-field low-pass filtered Gaussian noise patterns with a cutoff at 0.12 cycles per degree and a temporal frequency of 1 Hz[73]. In each trial, the stimulus was presented twice, with an inter-stimulus interval of 0.5 s. Visual stimulation to induce gamma oscillations consisted of 5-s-long full field drifting square wave gratings at horizontal orientation and a spatial frequency of 0.04 cycles per degree. All visual stimuli were presented at 17-cm distance from the right eye on a gamma-corrected LED backlit LCD monitor (Viewsonic VX3276-2K-MHD-2, 32", 60Hz refresh rate). The overall stimulation protocol consisted of 50 trials of each type, presented in randomized order with inter-trial intervals between 3–5 seconds.

For optogenetic stimulation, we used a fiber-coupled laser system (ReadyBeam Bio1, FISBA AG), connected to a 105 μm glass fiber (0.22 NA) that was positioned over the cortical recording site. Optogenetic stimulation consisted of a 3-s long pulse at 488 nm with a peak power of 10 mW from the fiber tip. To avoid transient changes in PV activity, the pulse intensity was first increasing for 1 second, then remained at a peak intensity for 1 second, and lastly decreased back to zero for 1 second. The sequence of visual, optogenetic, and combined trials was fully randomized. For optotagging of PV interneurons, we presented 20 optogenetic stimulus sequences at 5 Hz for 1 second, with an individual pulse length of 10 ms and a peak power of 2 mW from the fiber tip. PV interneurons were then identified by short-latency responses (<2 ms) to individual laser pulses and an increase in firing rate during the 3-s-long optogenetic stimulation period.

To extract spiking activity, channels that were broken or outside of the brain were detected and removed from the recording using the Spike Interface analysis toolbox[74]. The remaining channels were then time-shifted and median-subtracted across channels and time. Corrected channels were then spike-sorted with the Kilosort 2.5 software and manually curated using phy (https://github.com/cortex-lab/phy). All sorted spiking and local field potential data were then analyzed using custom Matlab code (2020b, MathWorks).

Peri-event time histograms (PETHs) for each cluster were computed with a bin size
of 10 ms and baseline-subtracted by the mean activity within 1 second before the
laser onset. Trial-averaged local field potentials (LFPs) in response to visual
stimulation were similarly baseline-corrected. To compute current source densities
(CSDs) from LFP signals, we used the inverse CSD method by Pettersen et al.[75].
We applied the spline iCSD method, assuming a smoothly varying CSD between
electrode contacts, based on polynomial interpolation. We assumed a
homogeneous, isotropic conductivity of $\sigma = 0.3$ S/m within and directly above
cortex[75]. To reduce spatial noise, the estimated CSD was subsequently convolved
with a Gaussian spatial filter with a standard deviation of 0.1 mm. For LFP and CSD
analyses, the signals from neighboring contacts at the same depth were averaged
together to improve signal quality.

**Behavioral Assays**

**Evidence accumulation task**

After full recovery from surgeries, animals were water-deprived for one day before
introducing them to the behavioral task. To obtain rewards, mice needed to
accumulate visual evidence in form of visual grating-like cues and indicate the side
with the higher number of cues. Experiments were conducted in a custom-built setup
with stimulus presentation, lick-detection and water reward controlled by a
microcontroller (Teensy 3.2, PJRC) and Python code (Python 3.7). In general, the
task consisted of three periods: 3 s sensory stimulation with randomized sequences
of up to 6 cues on one or both sides, 0.5 s delay and a 2-s response window. In
between trials was a 3.5-s inter trial interval, which was prolonged after missed trials
or incorrect responses. Mice first learned to reliably detect visual stimuli, by
indicating the side were all 6 cues (spatial frequency of 0.018 cpd, temporal
frequency of 2 Hz) were shown. After mice performed more than 75% of trials
correctly in 3 out of the last 4 sessions, distractor stimuli were added on the non-
rewarded side. The number of cues on both the target and distractor side varied,
resulting in different levels of difficulty depending on the absolute difference in cues
shown on either side. Training was conducted using 3 control and 4 knockout
animals.

**Morris water maze (MWM)**

Visual navigation and learning capacities were investigated using an MWM. The
setup consisted of a cylindric water-filled pool (Ø 1 m) surrounded by an opaque
curtain, four distinct landmarks on the pool wall, and a transparent platform (Ø 10
cm). The setup was divided into four virtual quadrants with the platform being
consistently in the same quadrant. The start position of the mouse was pseudo-
randomly alternated between trials. On the first day of testing, mice started once per
quadrant for habituation. The platform slightly protruded from water level to facilitate

quicker locating. Upon finding the platform, the animal was required to remain on top
of it for 5 s for the trial to be counted successful. If an animal did not find the platform
or left it after less than 5 s and did not return onto it within the 90-s period of the trial,
the mouse was manually guided to the platform before taking it back to their cages.
Between trials, mice were kept under red light for at least 10 min. For the actual
learning task, the platform was submerged (1–2 cm) and the maximum duration per
trial was set to 60 s. Each mouse started three times per quadrant in a pseudo-
random order, equaling 12 trials per day. This training scheme was conducted on
five consecutive days. Mice were recorded from above and behavioral data were
processed using ANY-maze (Stoelting Co., Wood Dale, USA). We tested 8 control
and 8 knockout mice.

**Home cage activity**

To assess the animals' activity in their home cages, cameras were set up in front of
cages of group-housed animals. Each cage was filmed on several separate days
with a framerate of 25 frames per second. From each of these recordings, eight
uninterrupted 30 min periods were used for further analysis. For each time period, a
pixel-wise comparison with ImageJ was utilized to calculate the difference between
every fifth frame of a video. For this, we used the ImageJ Image Calculator and
adjusted the threshold to exclude background noise. Subsequently, we measured
the area of differing pixels. We filmed 3 cages per genotype housing 2 mice each (6
control/6 knockout).

**Rotarod**

A rotarod was used to assess balance and motor performance of mice. Initially, the
mice (13 control/11 knockout) were individually habituated for 4 min at a constant
speed of 4 rpm. The subsequent test phase comprised three consecutive trials, each
lasting a maximum of 5 min, with continuous acceleration from 4 to 40 rpm.

**Wire hang test**

To record neuromuscular strength, mice were placed on a metal wire cage lid, which
was then slowly inverted and positioned over a type IV Makrolon cage (Fig. 3d).
Each mouse was given three trials of max. 5 min, with inter trial intervals of approx.
3–4 min. In total, 24 mice (13 control/11 knockout) conducted this experiment.

**Food intake**

Food intake of group-housed mice was quantified by assessing the change in
cumulative pellet weight after either 24 h or 72 h. Animals remained in their familiar
cage settings and pellet weight measurements were calculated relative to the
number of animals per cage as the consumed amount per day. Tests were
conducted on 4 control and 4 knockout cages (each cage containing 2 mice: 8

control/8 knockout). For each cage two 24-h periods and eight 72-h periods were
recorded.

To measure food and water intake in single-housed mice, mice were kept individually
for at least two weeks, before measuring pellet and water bottle weight of each cage
on five consecutive days. For this, we used 4 control and 6 knockout mice.

**Nest-building test**

Nest-building tests were executed as described by Deacon[78] and as visualized in
Fig. 3f. In brief, mice were weighed and transferred to individual testing cages (IVC
cages, Tecniplast) with ad libitum access to food and water, but without any
additional environmental enrichment. One nestlet of known weight was placed in
each cage. The test was initiated at 5 pm and nest-building skills were scored the
next morning at 9 am. Nests were scored according to the following criteria: Score 1
= no noticeable interaction with the nestlet (more than 90% of the nestlet remained
intact), Score 2 = nestlet was partially torn (50–90% of the nestlet remained intact),
Score 3 = nestlet was largely torn, but no identifiable nest site was build (10–50% of
the nestlet remained intact, but less than 90% of shredded material is gathered
within a quarter of the cage area), Score 4 = nestlet was used to build a flat nest
(less than 10% of the nestlet remained intact, less than 50% of the nest
circumference is build higher than the resting mouse), Score 5 = nestlet was used to
build a (near) perfect nest (less than 10% of the nestlet remained intact, more than
50% of the nest circumference is build higher than the resting mouse). Each mouse
was tested twice with tests being one week apart from each other. In the time
between tests, mice remained single-housed without access to further enrichments
beside the handed nestlet. Overall, 49 mice (24 control/25 knockout) at ages
between 3–6 months were tested.

**Coat state**

The animals' coat state was assessed in a blinded manner. The scoring system was
adapted from Nollet et al.[34] as visualized in Fig. 3g. Seven different regions of the
animal's body were scored individually with one of three distinct ranks. In case of a
good coat state (smooth and shiny fur, no tousled or spiky patches) rank 0 was
assigned. Strongly matted, oily or even stained fur was assigned with a rank 1. Slight
deviations from the optimum coat state were ranked with a 0.5. Among the
investigated areas were both dorsal (head, neck, back, tail) and ventral areas (fore
paws, abdomen, hind paws). Overall, 12 control and 12 knockout mice (2–7 months)
were investigated.

**Voluntary wheel running**

Single-housed mice were kept in cages with access to running wheels (TSE
Systems). Animals were housed for up to three days in these cages. While food and
water were provided ad libitum, no further enrichments despite the running wheels

were provided. Each animals' wheel usage was continuously recorded and
subsequently evaluated using a custom-written Python script. In total, 11 control and
9 knockout mice were utilized.

**Interaction test**

To test for social interest in female conspecifics, virgin C57BL6J-females were
transferred to cages of virgin transgenic, single-housed males. These encounters
were filmed for 10 min to evaluate the behavior of the transgenic males. Behaviors
such as sniffing, grooming or mounting of the female intruder were evaluated
second-wise. The same female encountered two males per day, one per genotype.
Between these encounters were at least 6 hours to ensure that the females did not
transfer any noticeable odors from the first encounter during the second session of
the day. This way we were able to compare the reaction of males of different
genotypes to the same female during the same time of her estrus cycle. A total of 16
animals (8 controls/8 knockouts) were utilized.

**Female urine sniffing test**

The reaction of transgenic males towards female olfactory cues was tested using
female urine. For this, urine from 5–10 C57BL6J-females from different cages was
collected and mixed. To circumvent reactions of the tested males towards the
presentation medium, pieces of parafilm were kept in the animals' cages for one
hour prior to testing them. Then, 40 µl tap water were applied to the parafilm as
neutral cue, and the time spent sniffing at the application site was assessed for 3
min. After a 45 min break, the same test was conducted using urine instead of water.
A total of 16 animals (8 controls/ 8 knockouts) were utilized.

**Novelty-suppressed feeding test**

Experiments were conducted according to Samuels & Hen[79] and as visualized in
Fig. 5a. In brief, mice were weighed and transferred to individual cages (IVC cages,
Tecniplast) with environmental enrichments and *ad libitum* access to water. After 24
h of food deprivation, the novelty suppressed feeding test was conducted. For this,
mice were individually placed in an arena (i.e., a mouse shipping container with
bedding on the ground) with a food pellet attached to a platform in the center. Mice
were filmed from above to later analyze their movement patterns using
DeepLabCut[80]. Once a mouse started consuming the presented pellet, the animal
was transferred to its home cage, and a food pellet of known weight was placed in
the food tray. After the test, all animals were weighed again. In total, 20 mice (10
control/ 10 knockout) aged 3–10 months were tested.

**Luminex blood analysis**

To test for hormonal blood concentrations, blood samples were collected *post-*
*mortem*. For each mouse, two samples of 50 µl blood serum were processed.

Samples were purified and analyzed using Luminex System (Millipore). Using
antibody-coated beads, a multiplex assay was performed to assess changes in
hormone levels being quantified using the MAGPIX detection system. The
MILLIPLEX Analyst 5.1 software then calculated hormone levels of ghrelin, GLP-1,
glucagon, insulin, leptin, PYY, and TNFα in pg/ml. A total of 6 control and 6 knockout
mice, all aged 6 months, were used.

**Brain tissue preparation**

Mice were anesthetized using 5% (v/v) isoflurane before decapitation. For
immunohistochemistry, mice were transcardially perfused first with phosphate-
buffered saline (PBS; pH 7.4), then with 4% paraformaldehyde (PFA) in PBS (pH
7.4). Subsequently, brains were prepared and kept in 4% PFA for 24 h at 4 °C for
post-fixation. Cryoprotection was achieved by incubating the samples in 10% (v/v)
and then in 30% (v/v) sucrose in PBS for 24 h each. Brains were subsequently
frozen on dry ice and stored at -80 °C.

**Single-nucleus RNA sequencing**

Neocortical tissue of *Pvalb-Cre/tdTomato* and *Pvalb-Cre/tdTomato/Dnmt1 loxP²* mice
was dissected from 1-mm-thick coronal slices corresponding to the region between
Bregma +1 and –2. For each genotype, tissue from the three mice was pooled prior
to nuclei extraction to obtain sufficient material and minimize inter-individual
variability. The samples were snap-frozen on dry ice and stored at –80 °C until nuclei
extraction to preserve RNA integrity. The nuclei were extracted, fixed and stored until
library preparation as previously described[12]. The snRNA-seq libraries were
generated using the Evercode™ WT kit (Parse Biosciences, USA). The quality of the
cDNA and the final libraries was verified using the High Sensitivity D5000 and D1000
ScreenTape kits (Agilent Technologies, USA). Libraries were equimolarly pooled and
sequenced together with 5% PhiX spike-in library on an Illumina NovaSeq machine
running in paired-end mode with the S2 Reagent Kit v1.5 (200 cycles; Illumina,
USA).

The sequencing dataset was processed and filtered using the default parameters in
Trailmaker™(https://app.trailmaker.parsebiosciences.com/; Parse Biosciences,
2025). The generated Leiden-clustered dataset was downloaded as a Seurat object
and analyzed further using the Seurat[81] package (5.3.0) in R (4.4.3). For
visualization, a uniform manifold approximation and projection (UMAP) embedding
was computed to represent cell clusters in a low-dimensional space. Cell clusters
containing fewer than 0.3% of the total nuclei were removed to minimize potential
artifacts arising from low-quality clusters. Cluster-specific marker genes were
identified using the presto package implementation of the Wilcoxon rank-sum test,
comparing each cluster against all other cells. The top 50 upregulated and top 50
downregulated marker genes per cluster were cross-referenced with established cell

type-specific markers from published literature and publicly available reference
databases.

The annotated clusters were subsequently used to infer cell–cell communication
probabilities using the CellChat[82] database and the corresponding R package
(CellChat version 1.6.1).

**Immunohistochemistry**

For immunohistochemistry of free-floating, 40-µm-thick, adult brain sections, slices
were first washed three times in PBS/0.05% Triton X-100. GABA-staining required
incubation in 1 N HCl for 10 min at 4 °C, then in 2 N HCl for 10 min at room
temperature and lastly in 2 N HCl for 20 min at 37 °C. Afterwards, all sections were
placed in citrate buffer (0.05% Tween20, pH 6) for 15 min at 90–100 °C. After three
washing steps with PBS/0.05% Triton X-100, sections were blocked for 2 h using
blocking solution (4% BSA and 10% NGS in PBS/0.05% Triton X-100). Incubation
with primary antibodies was conducted for 60 h at 4 °C (rabbit anti-GABA (A2052,
1:750), rabbit anti-β-Endorphin (Ab5028, 1:1000), mouse anti-RFP (MA5-15257,
1:500)). Secondary antibodies were applied for 3 h at room temperature (Cy5-
donkey anti-rabbit (A10523, 1:1000), Cy3-goat anti-mouse (115035062, 1:1000)).
Sections were then stained with DAPI (100 ng/mL in PBS; Sigma-Aldrich) for 5 min.
Afterwards, they were mounted onto object slides in pre-warmed 0.5% gelatine
supplemented with 0.05% $KCrS_2O_8$ x $H_2O$ before being embedded in Mowiol (Carl
Roth).

For the staining of perineuronal nets, 30-µm-thick adult brain cryosections were
collected on glass slides. Sections were washed three times in PBS/0.2% Triton X-
100. After blocking for 2 h using blocking solution (4% BSA and 10% NGS in
PBS/0.2% Triton X-100), the slices were incubated with Wisteria floribunda lectin
(WFA; L32481, Thermo Fisher Scientific, 1:200) at 4 °C overnight. The slices were
then washed three times with PBS/0.2% Triton X-100 before they were stained with
DAPI (100 ng/mL in PBS, Carl Roth) for 15 min. After two washing steps with PBS,
the slices were embedded in Mowiol (Sigma-Aldrich).

**Microscopy and image analysis**

The brain slices were imaged using a Leica DMi8 fluorescent microscope in
combination with a THUNDER® imager unit (Leica, Germany) and the
corresponding software LASX (Leica, Germany). Post-processing was performed in
the LASX software using the "Mosaic Merge" and "Thunder Lightning" tools. A
maximum intensity projection of the merged tile scans was processed using Fiji[83]
(ImageJ, NIH, USA). Additionally, PV interneurons and surrounding PNNs were
captured using a Leica SP8 Tau-STED3X confocal microscope (Leica, Germany),
and the software LASX (Leica, Germany). The resulting z-stacks were processed
with the Fiji software (ImageJ, NIH, USA), applying a 50-pixel sliding paraboloid
background subtraction.

Fluorescence intensity measurements were performed using Fiji (ImageJ, NIH, USA). Regions of Interest (ROI) were manually defined, and the mean corrected density of each ROI was calculated by normalizing the corrected integrated density to the ROI area. For the analysis of PV interneurons enwrapped in PNNs, QuPath[84] version 0.6.0 was used. Based on a groovy script for QuPath developed for this study, PV interneurons in the primary somatosensory cortex (S1) were identified, and the surrounding region was analyzed for the presence of WFA signals.

**Statistics**

Statistical tests were carried out using Matlab (2022b), Python (2.7), and R (4.4.3). Normally distributed data was tested using two-tailed Student's *t*-test, one-way ANOVA or two-way ANOVA. Non-normally distributed or ordinal data was tested using a Wilcoxon *rank-sum* test. Tukey's test was conducted as a post-hoc test. The conducted tests for individual experiments are indicated in the text.

# Acknowledgments

We thank Nilufar Nojavan Lahiji and Hendro Langecker for helping with the Morris water maze experiment, Annalena Dobbert for support with the histological analysis, Mira Jakovcevski for methodical assistance in the coat state analysis, Sandra Brill for assistance in the coat state analysis, support in animal breeding/caretaking as well as fruitful scientific discussions.

# Author contributions

J.L.: conceptual design, performed experiments, data analysis, figure illustration, wrote the manuscript; C.B.Y.: conceptual design, performed experiments, data analysis, figure illustration, wrote the manuscript; K.V.: performed experiments, data analysis, figure illustration, manuscript editing; G.N.: method development, experimental supervision, data analysis, figure illustration, manuscript editing; S.G.: performed experiments, data analysis; D.P.: experimental supervision, data discussion, manuscript editing; J.R.: performed experiments, manuscript editing; M.H.: performed experiments, manuscript editing; B.K.: data discussion, manuscript editing; A.U.: performed experiments, data analysis, data discussion, manuscript editing; S.M.: performed experiments, data analysis, data discussion, manuscript editing; G.Z.B.: conceptual design, data discussion, wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

# Funding

This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—368482240/GRK2416, ZI-1224/13-1, ZI-1224/19-1, ZI-1224/20-1, the Excellence Strategy of the Federal Government and the Länder

848 (OPSF812, SFASIA002); the IZKF Jena, the Helmholtz association (VH-NG-1611),
849 and the state of North Rhine-Westphalia through the iBehave initiative.
850

## Competing interests

The authors declare no competing interests.

## References

1. Fatemi, S. H., Folsom, T. D. & Thuras, P. D. Deficits in GABAB receptor system in schizophrenia and mood disorders: A postmortem study. *Schizophrenia Research* **128**, 37–43 (2011).

2. Jahangir, M., Zhou, J.-S., Lang, B. & Wang, X.-P. GABAergic System Dysfunction and Challenges in Schizophrenia Research. *Front. Cell Dev. Biol.* **9**, (2021).

3. Linde, J. & Zimmer-Bensch, G. DNA Methylation-Dependent Dysregulation of GABAergic Interneuron Functionality in Neuropsychiatric Diseases. *Front. Neurosci.* **14**, 586133 (2020).

4. Sanacora, G. Cortical Inhibition, Gamma-Aminobutyric Acid, and Major Depression: There Is Plenty of Smoke but Is There Fire? *Biological Psychiatry* **67**, 397–398 (2010).

5. Della Vecchia, A., Arone, A., Piccinni, A., Mucci, F. & Marazziti, D. GABA System in Depression: Impact on Pathophysiology and Psychopharmacology. *CMC* **29**, 5710–5730 (2022).

6. Feng, J. & Fan, G. The role of DNA methylation in the central nervous system and neuropsychiatric disorders. *Int Rev Neurobiol* **89**, 67–84 (2009).

7. Gräff, J., Kim, D., Dobbin, M. M. & Tsai, L.-H. Epigenetic regulation of gene expression in physiological and pathological brain processes. *Physiol Rev* **91**, 603–649 (2011).

8. Moore, A. K., Weible, A. P., Balmer, T. S., Trussell, L. O. & Wehr, M. Rapid Rebalancing of Excitation and Inhibition by Cortical Circuitry. *Neuron* **97**, 1341-1355.e6 (2018).

9. Ji, M.-H. *et al.* Overinhibition mediated by parvalbumin interneurons might contribute to depression-like behavior and working memory impairment induced by lipopolysaccharide challenge. *Behav Brain Res* **383**, 112509 (2020).

10. Perlman, G., Tanti, A. & Mechawar, N. Parvalbumin interneuron alterations in stress-related mood disorders: A systematic review. *Neurobiol Stress* **15**, 100380 (2021).

11. Ghosal, S., Hare, B. & Duman, R. S. Prefrontal Cortex GABAergic Deficits and
Circuit Dysfunction in the Pathophysiology and Treatment of Chronic Stress and
Depression. *Curr Opin Behav Sci* **14**, 1–8 (2017).

12. Reichard, J. *et al.* DNMT1-mediated regulation of somatostatin-positive
interneuron migration impacts cortical architecture and function. *Nat Commun* **16**,
6834 (2025).

13. Pensold, D. *et al.* DNA methylation-mediated modulation of endocytosis as
potential mechanism for synaptic function regulation in murine inhibitory cortical
interneurons. *Cerebral Cortex* **30**, 3921–3937 (2020).

14. Wingert, J. C. & Sorg, B. A. Impact of Perineuronal Nets on Electrophysiology of
Parvalbumin Interneurons, Principal Neurons, and Brain Oscillations: A Review.
*Front Synaptic Neurosci* **13**, 673210 (2021).

15. Sorg, B. A. *et al.* Casting a Wide Net: Role of Perineuronal Nets in Neural
Plasticity. *J. Neurosci.* **36**, 11459–11468 (2016).

16. Wen, T. H., Binder, D. K., Ethell, I. M. & Razak, K. A. The Perineuronal 'Safety'
Net? Perineuronal Net Abnormalities in Neurological Disorders. *Front Mol
Neurosci* **11**, 270 (2018).

17. Auer, S. *et al.* The Role of Perineuronal Nets in Physiology and Disease: Insights
from Recent Studies. *Cells* **14**, 321 (2025).

18. Hahn, A. *et al.* DNA methyltransferase 1 (DNMT1) function is implicated in the
age-related loss of cortical interneurons. *Frontiers in cell and developmental
biology* **8**, 639 (2020).

19. Linde, J. & Zimmer-Bensch, G. DNA methylation-dependent dysregulation of
GABAergic interneuron functionality in neuropsychiatric diseases. *Frontiers in
Neuroscience* **14**, 982 (2020).

20. Senzai, Y., Fernandez-Ruiz, A. & Buzsáki, G. Layer-Specific Physiological
Features and Interlaminar Interactions in the Primary Visual Cortex of the Mouse.
*Neuron* **101**, 500-513.e5 (2019).

21. Kriener, B., Hu, H. & Vervaeke, K. Parvalbumin interneuron dendrites enhance
gamma oscillations. *Cell Rep* **39**, 110948 (2022).

22. Sauer, J.-F., Strüber, M. & Bartos, M. Impaired fast-spiking interneuron function
in a genetic mouse model of depression. *Elife* **4**, e04979 (2015).

23. Herrmann, C. S. & Demiralp, T. Human EEG gamma oscillations in
neuropsychiatric disorders. *Clin Neurophysiol* **116**, 2719–2733 (2005).

24. Zhao, Y. *et al.* Parietal gamma oscillations decreased in cognitive biotype of
depression. *Sci Rep* **15**, 37100 (2025).

25. Liu, Q., Shi, C., Wang, F. & Tong, J. Neuroinflammation-induced parvalbumin
interneuron and oscillation deficits might contribute to neurobehavioral
abnormities in a two-hit model of depression. *Heliyon* **9**, e18468 (2023).

26. Fureix, C. *et al.* Do greater levels of in-cage waking inactivity in laboratory mice
reflect a spontaneous depression-like symptom? A pharmacological
investigation. *Pharmacol Biochem Behav* **212**, 173311 (2022).

27. Mueller, J. *et al.* The relationship of within-individual and between-individual
variation in mental health with bodyweight: An exploratory longitudinal study.
*PLoS One* **19**, e0295117 (2024).

28. Ostlund, R. E., Yang, J. W., Klein, S. & Gingerich, R. Relation between plasma
leptin concentration and body fat, gender, diet, age, and metabolic covariates. *J*
*Clin Endocrinol Metab* **81**, 3909–3913 (1996).

29. Halaas, J. L. *et al.* Physiological response to long-term peripheral and central
leptin infusion in lean and obese mice. *Proc Natl Acad Sci U S A* **94**, 8878–8883
(1997).

30. Timper, K. & Brüning, J. C. Hypothalamic circuits regulating appetite and energy
homeostasis: pathways to obesity. *Dis Model Mech* **10**, 679–689 (2017).

31. Valles, K., Ayers, E., Verghese, J. & Ceïde, M. Behavioral Correlates of Apathy in
Community Dwelling Older Adults. *The American Journal of Geriatric Psychiatry*
**31**, S42–S43 (2023).

32. Baumann, A. *et al.* Preliminary Evidence of Apathetic-Like Behavior in Aged
Vesicular Monoamine Transporter 2 Deficient Mice. *Front Hum Neurosci* **10**, 587
(2016).

33. Mineur, Y. S., Prasol, D. J., Belzung, C. & Crusio, W. E. Agonistic behavior and
unpredictable chronic mild stress in mice. *Behav Genet* **33**, 513–519 (2003).

34. Nollet, M., Le Guisquet, A.-M. & Belzung, C. Models of depression: unpredictable
chronic mild stress in mice. *Curr Protoc Pharmacol* **Chapter 5**, Unit 5.65 (2013).

35. Belke, T. W. Running and responding reinforced by the opportunity to run: effect
of reinforcer duration. *J Exp Anal Behav* **67**, 337–351 (1997).

36. Pizzagalli, D. A. Depression, Stress, and Anhedonia: Toward a Synthesis and
Integrated Model. *Annual Review of Clinical Psychology* **10**, 393–423 (2014).

37. Bakken, T. E. *et al.* Single-nucleus and single-cell transcriptomes compared in
matched cortical cell types. *PLOS ONE* **13**, e0209648 (2018).

38. Lake, B. B. *et al.* Integrative single-cell analysis of transcriptional and epigenetic
states in the human adult brain. *Nat Biotechnol* **36**, 70–80 (2018).

39. Yao, Z. *et al.* A taxonomy of transcriptomic cell types across the isocortex and
hippocampal formation. *Cell* **184**, 3222-3241.e26 (2021).

40. Luo, C. *et al.* Single-cell methylomes identify neuronal subtypes and regulatory
elements in mammalian cortex. *Science* **357**, 600–604 (2017).

41. Kanigowski, D. & Urban-Ciecko, J. Conditioning and pseudoconditioning
differently change intrinsic excitability of inhibitory interneurons in the neocortex.
*Cereb Cortex* **34**, bhae109 (2024).

42. Szuchet, S. *et al.* The genetic signature of perineuronal oligodendrocytes reveals
their unique phenotype. *Eur J Neurosci* **34**, 1906–1922 (2011).

43. Tian, Y., Zhan, Y., Jiang, Q., Lu, W. & Li, X. Expression and function of PDGF-C
in development and stem cells. *Open Biology* **11**, 210268 (2021).

44. Falk, J., Bonnon, C., Girault, J.-A. & Faivre-Sarrailh, C. F3/contactin, a neuronal
cell adhesion molecule implicated in axogenesis and myelination. *Biol Cell* **94**,
327–334 (2002).

45. Borges, B. C., Meng, X., Long, P., Kanold, P. O. & Corfas, G. Loss of
oligodendrocyte ErbB receptor signaling leads to hypomyelination, reduced
density of parvalbumin‑expressing interneurons, and inhibitory function in the
auditory cortex. *Glia* **71**, 187–204 (2023).

46. Li, K.-X. *et al.* Neuregulin 1 regulates excitability of fast-spiking neurons through
Kv1.1 and acts in epilepsy. *Nat Neurosci* **15**, 267–273 (2011).

47. Vo, T. *et al.* The chemorepulsive axon guidance protein semaphorin3A is a
constituent of perineuronal nets in the adult rodent brain. *Molecular and Cellular*
*Neuroscience* **56**, 186–200 (2013).

48. Mo, X. *et al.* PTPRM Is Critical for Synapse Formation Regulated by Zinc Ion.
*Front. Mol. Neurosci.* **15**, (2022).

49. Proctor, D. T. *et al.* Axo-glial communication through neurexin-neuroligin
signaling regulates myelination and oligodendrocyte differentiation. *Glia* **63**,
2023–2039 (2015).

50. Sinha, A., Nickerson, G., Bouyain, S. & Matthews, R. T. Contactin-1 is a critical
neuronal cell surface receptor for perineuronal net structure. *bioRxiv*
2024.11.05.622114 (2024) doi:10.1101/2024.11.05.622114.

51. Kurolap, A. *et al.* Bi-allelic variants in neuronal cell adhesion molecule cause a
neurodevelopmental disorder characterized by developmental delay, hypotonia,
neuropathy/spasticity. *Am J Hum Genet* **109**, 518–532 (2022).

52. Mueller-Buehl, C. *et al.* Brevican, Neurocan, Tenascin-C, and Tenascin-R Act as
Important Regulators of the Interplay Between Perineuronal Nets, Synaptic
Integrity, Inhibitory Interneurons, and Otx2. *Front Cell Dev Biol* **10**, 886527
(2022).

53. Carulli, D. *et al.* Cerebellar plasticity and associative memories are controlled by
perineuronal nets. *Proc Natl Acad Sci U S A* **117**, 6855–6865 (2020).

54. Turrigiano, G. G. Homeostatic plasticity in neuronal networks: the more things
change, the more they stay the same. *Trends Neurosci* **22**, 221–227 (1999).

55. Deleuze, C. *et al.* Strong preference for autaptic self-connectivity of neocortical
PV interneurons facilitates their tuning to γ-oscillations. *PLoS Biol* **17**, e3000419
(2019).

56. Ferguson, B. R. & Gao, W.-J. PV Interneurons: Critical Regulators of E/I Balance
for Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. *Front*
*Neural Circuits* **12**, 37 (2018).

57. Rajkowska, G., O'Dwyer, G., Teleki, Z., Stockmeier, C. A. & Miguel-Hidalgo, J. J.
GABAergic neurons immunoreactive for calcium binding proteins are reduced in
the prefrontal cortex in major depression. *Neuropsychopharmacology* **32**, 471–
482 (2007).

58. Espinoza, C., Guzman, S. J., Zhang, X. & Jonas, P. Parvalbumin+ interneurons
obey unique connectivity rules and establish a powerful lateral-inhibition
microcircuit in dentate gyrus. *Nat Commun* **9**, 4605 (2018).

59. Rupert, D. D. & Shea, S. D. Parvalbumin-Positive Interneurons Regulate Cortical
Sensory Plasticity in Adulthood and Development Through Shared Mechanisms.
*Front. Neural Circuits* **16**, (2022).

60. Hadler, M. D., Alle, H. & Geiger, J. R. P. Parvalbumin interneuron cell-to-network
plasticity: mechanisms and therapeutic avenues. *Trends Pharmacol Sci* **45**, 586–
601 (2024).

61. D'Souza, R. D. & Burkhalter, A. A Laminar Organization for Selective Cortico-
Cortical Communication. *Front Neuroanat* **11**, 71 (2017).

62. Perova, Z., Delevich, K. & Li, B. Depression of Excitatory Synapses onto
Parvalbumin Interneurons in the Medial Prefrontal Cortex in Susceptibility to
Stress. *J Neurosci* **35**, 3201–3206 (2015).

63. Bai, X., Kirchhoff, F. & Scheller, A. Oligodendroglial GABAergic Signaling: More
Than Inhibition! *Neurosci Bull* **37**, 1039–1050 (2021).

64. Weber, P. *et al.* Mice Deficient for Tenascin-R Display Alterations of the
Extracellular Matrix and Decreased Axonal Conduction Velocities in the CNS. *J*
*Neurosci* **19**, 4245–4262 (1999).

65. Arias-Aragón, F., Robles-Lanuza, E., Sánchez-Gómez, Á., Martinez-Mir, A. &
Scholl, F. G. Analysis of neurexin-neuroligin complexes supports an isoform-
specific role for beta-neurexin-1 dysfunction in a mouse model of autism. *Mol*
*Brain* **18**, 20 (2025).

66. Dean, C. & Dresbach, T. Neuroligins and neurexins: linking cell adhesion,
synapse formation and cognitive function. *Trends in Neurosciences* **29**, 21–29
(2006).

67. Connor, S. A. *et al.* Loss of Synapse Repressor MDGA1 Enhances Perisomatic
Inhibition, Confers Resistance to Network Excitation, and Impairs Cognitive
Function. *Cell Reports* **21**, 3637–3645 (2017).

68. Luo, F., Sclip, A., Merrill, S. & Südhof, T. C. Neurexins regulate presynaptic
GABAB-receptors at central synapses. *Nat Commun* **12**, 2380 (2021).

69. Hadler, M. D., Tzilivaki, A., Schmitz, D., Alle, H. & Geiger, J. R. P. Gamma
oscillation plasticity is mediated via parvalbumin interneurons. *Science Advances*
**10**, eadj7427 (2024).

70. Hippenmeyer, S. *et al.* A developmental switch in the response of DRG neurons
to ETS transcription factor signaling. *PLoS Biol* **3**, e159 (2005).

71. Madisen, L. *et al.* A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. *Nat Neurosci* **13**, 133–140
(2010).

72. Jackson-Grusby, L. *et al.* Loss of genomic methylation causes p53-dependent
apoptosis and epigenetic deregulation. *Nat Genet* **27**, 31–39 (2001).

73. Niell, C. M. & Stryker, M. P. Highly selective receptive fields in mouse visual
cortex. *J Neurosci* **28**, 7520–7536 (2008).

74. Buccino, A. P. *et al.* SpikeInterface, a unified framework for spike sorting. *Elife* **9**,
e61834 (2020).

75. Pettersen, K. H., Devor, A., Ulbert, I., Dale, A. M. & Einevoll, G. T. Current-
source density estimation based on inversion of electrostatic forward solution:
effects of finite extent of neuronal activity and conductivity discontinuities. *J*
*Neurosci Methods* **154**, 116–133 (2006).

76. Prusky, G. T., Alam, N. M., Beekman, S. & Douglas, R. M. Rapid quantification of
adult and developing mouse spatial vision using a virtual optomotor system.
*Invest Ophthalmol Vis Sci* **45**, 4611–4616 (2004).

77. Wiesbrock, C., Musall, S. & Kampa, B. M. A flexible Python-based touchscreen
chamber for operant conditioning reveals improved visual perception of cardinal
orientations in mice. *Front. Cell. Neurosci.* **16**, (2022).

78. Deacon, R. M. J. Assessing nest building in mice. *Nat Protoc* **1**, 1117–1119
(2006).

79. Samuels, B. A. & Hen, R. Novelty-Suppressed Feeding in the Mouse. in *Mood*
*and Anxiety Related Phenotypes in Mice: Characterization Using Behavioral*
*Tests, Volume II* (ed. Gould, T. D.) 107–121 (Humana Press, Totowa, NJ, 2011).
doi:10.1007/978-1-61779-313-4_7.

80. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body
parts with deep learning. *Nat Neurosci* **21**, 1281–1289 (2018).

81. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-
cell analysis. *Nat Biotechnol* **42**, 293–304 (2024).

82. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat.
*Nat Commun* **12**, 1088 (2021).

83. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis.
*Nat Methods* **9**, 676–682 (2012).

84. Bankhead, P. *et al.* QuPath: Open source software for digital pathology image
analysis. *Sci Rep* **7**, 16878 (2017).

1078

1079

**Figure 1: *Dnmt1*-knockout in PV interneurons alters spontaneous and induced neuronal activity.**

We performed electrophysiological recordings in the primary visual cortex (V1) of awake *Pvalb-Cre/tdTomato* control (Ctrl) and *Pvalb-Cre/tdTomato/Dnmt1 loxP²* (KO) mice using Neuropixels probes ($n = 8$ recordings from 2 mice per genotype). **(a)** Schematic illustration of the experimental setup and recording sites. **(b)** Fast-spiking PV interneurons were identified via optotagging (see example spike waveforms, left). Their spontaneous firing rates were significantly increased in KO mice ($n_{Ctrl} = 54$ and $n_{KO} = 37$ neurons; Wilcoxon *rank-sum* test: $p = 0.0297$). **(c)** Spontaneous firing rates of V1 neurons were significantly higher in KO mice compared to control mice ($n_{Ctrl} = 838$ and $n_{KO} = 448$ neurons; Wilcoxon *rank-sum* test: $p = 5.72 \times 10^{-19}$). **(d)** Relative change in the firing rate of V1 neurons inhibited by optogenetic activation of PV interneurons. PV-induced inhibition was significantly weaker in KO mice ($n_{Ctrl}=332$ and $n_{KO}=238$ neurons; Wilcoxon *rank-sum* test: $p = 3 \times 10^{-6}$). **(e)** Average firing rates of V1 neurons in response to visual stimulation in control and knockout mice. Visual stimuli were 0.1-s-long, full-field presentations of low-frequency noise patterns, presented at time point 0 and after 0.5 seconds. Shading shows the standard error of means (SEM) across neurons. **(f)** Peak response firing rate of V1 neurons after visual stimulation. KO mice had significantly weaker sensory responses compared to control mice ($n_{Ctrl} = 293$ and $n_{KO} = 74$ neurons; Wilcoxon *rank-sum* test: $p = 1.1 \times 10^{-15}$). **(g)** Local field potentials (LFP, upper panels) and current source densities (CSD, lower panels) of control and *Dnmt1*-KO mice across the whole depth of the cortex. White lines indicate visual stimulus onset. **(h)** LFPs in V1 in response to visual stimulation at time point 0, averaged across cortical layers. Shading shows SEM across recordings. **(i–j)** KO mice displayed strongly diminished gamma oscillations (30–90 Hz) in comparison to control mice. **(i)** Relative change in spectral power compared to baseline activity. Visually evoked oscillations in control mice matched the gamma band (30–90 Hz), while in KO mice oscillations of much lower frequencies (< 20 Hz) occurred. **(j)** Change in spectral power within the cortex displayed as mean ± SEM. The direct comparison between control and *Dnmt1*-KO

1111    mice revealed the shift towards lower-frequency oscillations in KO mice. Panels b–f

1112    display individual data points with median (centre line) and interquartile range (upper

1113    and lower lines). Ctrl = control, CSD = current source density, KO = knockout, LFP =

1114    local field potential, PV = parvalbumin, V1 = primary visual cortex; * $p < 0.5$,

1115    ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

1116

**Figure 2: *Pvalb-Cre/tdTomato/Dnmt1 loxP²* (KO) mice display apathetic behavior.**

**(a–b)** Spontaneous activity within the home-cage was consistently diminished in KO mice compared to *Pvalb-Cre/tdTomato* (control/Ctrl) mice ($n = 6$ mice per genotype). **(a)** Mice were monitored in their home-cages to evaluate their spontaneous, undisturbed behavior. By comparing every fifth frame from the resulting videos (i, ii), the change in pixel values between frames (iii) was binarized (iv) to calculate the change in position of the contained animals. **(b)** Comparison of multiple 30-min time intervals revealed consistently lower activity levels of KO mice (Two-way ANOVA: $p_{(Genotype)} = 1.9x10^{-5}$; $p_{(Time\ bin)} = 0.98$, $p_{(Genotype\ x\ Time\ bin)} = 0.91$). **(c)** Average latency to fall from a rotating rod ('Rotarod') with increasing velocity showed no apparent motor deficits in knockout mice ($n_{(Control)} = 13$, $n_{(Knockout)} = 11$; ANOVA: $p = 0.31$). (**d**) A wire hang test showed a significantly reduced latency to fall in KO mice, which can indicate a difference in motor strength but is highly dependent on the animals' body weight ($n_{(Control)} = 13$, $n_{(Knockout)} = 11$; ANOVA: $p = 3x10^{-6}$). **(e)** Body weights were elevated for KO mice compared to controls ($n_{(Control)} = 43$, $n_{(Knockout)} = 33$; ANOVA: $p = 0.0068$). **(f)** Nest-building capacities were significantly decreased in KO ($n = 25$) versus control mice ($n = 24$; Generalized estimating equation model: $p_{(Genotype)} = 0.006$). Nest quality was assessed using an ordinal scale from 1–5 as visualized by the schematic in the middle. As the representative images (right side) show, knockout mice were shredding the provided material but failed to assemble visible nest sites. **(g)** Assessment of the animals' coat state as indicated in the illustration on the bottom right showed significantly higher rankings of the knockout mice, indicating worsened fur care ($n = 12$ mice per genotype; Mann-Whitney U test: $p = 0.026$). Panels b–g display individual data points with median (centre line) and

1143  interquartile range (upper and lower lines). Ctrl = control, KO = knockout; * $p < 0.5$, **
1144  $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

1145

1146
**Figure 3:** ***Pvalb-Cre/tdTomato/Dnmt1 loxP[2]* (KO) mice display traits of an**
**anhedonic, anxious phenotype.**

**(a)** Voluntary running on a wheel provided within the home-cage was strongly
reduced in KO (*n* = 9) versus *Pvalb-Cre/tdTomato* (control/Ctrl) mice (*n* = 11).
Running wheels were provided in the home cages of single-housed mice for up to
three nights. Almost all mice displayed fluctuating utilization of the running wheel
(upper heat maps) based on the daytime (middle graph), but in general, KO mice
were running less frequently than controls. The overall reduced activity of knockout

mice is discernible in the mean running time averaged per genotype (lower graph).
**(b)** Evaluating the interaction of male KO and control mice ($n$ = 8 mice per genotype)
with female C57BL/6J-mice in a 10-min time frame revealed that KO males spent
significantly less time trying to mount the females (left graph, Student's $t$-test: $p$ =
0.019), while they spent similar times sniffing their conspecifics (right graph,
Student's $t$-test: $p$ = 0.44). **(c)** Presentation of pooled urine from female C57B/6J
mice significantly increased time spent sniffing in control but not in KO mice ($n$ = 8
mice per genotype; Linear mixed effects model: $p_{(Genotype\ x\ stimulant\ type)}$ = 0.048, Tukey
post-hoc test: $p_{(Ctrl:\ water\ vs\ urine)}$ = 0.0017, $p_{(KO:\ water\ vs.\ urine)}$ = 0.25, $p_{(Water:\ Ctrl\ vs\ KO)}$ = 1.0,
$p_{(Urine:\ Ctrl\ vs\ KO)}$ = 0.0413). **(d)** Conditional *Dnmt1* deletion in PV cells induces anxiety
in a novelty-suppressed feeding (NSF) test ($n$ = 10 per genotype). Upper row:
Scheme and schedule of the NSF test. Mice were food deprived for 24 h before
being transferred into an arena with new bedding and a centrally attached food pellet
of known weight. Once the animals started chewing the presented food, their latency
was recorded, and they were immediately removed to their home-cage.
Subsequently, the pellet from the arena was presented in the home-cage and the
animals' latency to approach and consume it was recorded. Videos taken during the
NSF test were labeled using DeepLabCut, to evaluate the animals' behavior until
they approached the food pellet (lower graphs). **(e)** KO mice displayed a significantly
increased latency to approach the presented food pellet in the arena compared to
controls, suggesting increased levels of anxiety. **(f)** In their home cages, KO mice
approached the same food pellet significantly faster than control mice. Panels b, c, e,
and f display individual data points with median (centre line) and interquartile range
(upper and lower lines). Ctrl = control, KO = knockout; * $p$ < 0.5, ** $p$ < 0.01, *** $p$ <
0.001, **** $p$ < 0.0001.

**Figure 4: Single-nucleus RNA sequencing reveals transcriptomic shifts in excitatory neurons as well as a redistribution of interaction strength between PV interneurons and excitatory subtypes in *Pvalb⊠Cre/tdTomato/Dnmt1 loxP²* (KO) mice.**

**(a)** Uniform manifold approximation and projection (UMAP) embedding reveals the expected cellular composition of the neocortex. **(b)** Dot blot of marker genes across the identified clusters. **(c)** Volcano plots illustrate the extent of upregulated and downregulated genes in excitatory neurons from layers 4–6 of KO mice (log₂FC ≥

0.26, adjusted *p* value < 0.05). **(d)** Gene ontology analysis of differentially expressed
genes (DEGs) in the layer 4 sensory neurons reveal an enrichment for biological
processes related to synaptic transmission, organization, and activity. **(e)** CellChat-
based mapping demonstrates altered interaction networks among neuronal and glial
cell clusters in the KO cortex. The senders, i.e., the clusters the signaling event is
initiated from, are listed in the y-axis while the clusters receiving the signal are found
in the x-axis. **(f)** The circular network diagram depicts altered connectivity patterns
between excitatory neurons and PV interneurons in KO mice. Increased and
decreased communication probabilities in KO mice are depicted in red and blue,
respectively.

1198

**Figure 5: DNMT1 deficiency in PV interneurons reshapes PV-oligodendroglial communication and alters ligand–receptor interactions crucial for perineuronal nets (PNNs) in *Pvalb⊡Cre/tdTomato/Dnmt1 loxP²* (KO) mice.**

**(a)** The *Pdgfc*-expressing population within newly mature oligodendrocytes, representing perineuronal oligodendrocytes, is significantly reduced in the KO cortex (two-sample proportion test, $p = 0.005$). **(b)** The circular network diagram illustrates differential interactions between oligodendroglial cells and PV interneurons in KO mice. Increased and decreased communication probabilities in KO mice are depicted in red and blue, respectively. **(c)** Chord diagrams depict the altered interactions grouped by ligand/receptor family, separated based on directionality. Upper diagram:

1209   PV interneurons → oligodendroglial cells; lower diagram: Oligodendroglial
1210   cells → PV interneurons. Ligand–receptor pairs with known roles in stabilizing PNNs,
1211   the perisomatic niche and cell–cell interactions have altered communication
1212   probability in **(d)** outgoing signaling from PV interneurons to oligodendroglial cells as
1213   well as **(e)** incoming signaling from oligodendroglial cells to PV interneurons.

**Figure 6: Loss of DNMT1 in PV interneurons disrupts perineuronal nets (PNNs) in the neocortex of *Pvalb☐Cre/tdTomato/Dnmt1 loxP²* (KO) mice.**

**(a–b)** Representative microphotographs of the primary somatosensory cortex (S1) in coronal brain sections of *Pvalb-Cre/tdTomato* (Ctrl; **a**) and *Pvalb☐Cre/tdTomato/Dnmt1 loxP²* (KO, **b**) mice stained for PNNs using Wisteria floribunda lectin (WFA). Scale bars: 200 μm. **(c–d)** Orthogonal XZ und YZ projections show the three-dimensional localization of a PV interneuron soma and surrounding PNNs stained with WFA. Scale bars: 10 μm. **(e)** Mean corrected WFA signal density is reduced in the neocortex of KO mice (box plots: median, interquartile range, min-to-max; paired Student's *t*-test, $p = 0.046$; $N = 3$ biological replicates with $n = 3$ slices each per group). **(f)** Proportion of PV interneurons enwrapped by PNNs in S1 of KO mice is unchanged (box plots: median, interquartile range, min-to-max; paired Student's *t*-test, $p = 0.277$; $N = 3$ biological replicates with n = 8 slices per genotype). * $p < 0.05$.