A three amino acid sequence gates protein stability control of bHLH104 and iron

uptake in *Arabidopsis thaliana*

Dibin Baby[1,2], Sarah Weldi[1], Marie Knopf[1], Vinitha Venkadasamy[1], Jessica Lee

Erickson[3], Petra Bauer[1,2*]

[1]Institute of Botany, Heinrich Heine University, Düsseldorf 40225, Germany

[2]Cluster of Excellence on Plant Science (CEPLAS), Heinrich Heine University,

Düsseldorf 40225, Germany

[3] Zentrum für Molekularbiologie der Pflanzen (ZMBP), Allgemeine Genetik, University

of Tübingen, 72076 Tübingen, Germany.

*Corresponding author: Petra Bauer, petra.bauer@hhu.de

Dibin Baby- https://orcid.org/0000-0003-0592-6067

Petra Bauer- https://orcid.org/0000-0002-0404-4532

Abstract

Iron (Fe) homeostasis is precisely regulated to prevent iron imbalance. The basic

helix-loop-helix (bHLH) transcription factor bHLH104, a member of bHLH subgroup

IVc, activates Fe uptake. bHLH104 is negatively controlled by the Fe-binding E3 ligase

BRUTUS (BTS) that interacts with the C-terminus of bHLH104, according to

predictions with involvement of the last three amino acids proline-alanine-alanine

(PAA). Yet, experimental evidence for the PAA interaction site and importance for post-

translational regulation are lacking. Here, we demonstrate the role of PAA for plant

growth and bHLH104 regulation. Transgenic plants expressing a bHLH104 variant

lacking the terminal PAA (*b104^dPAA*) have a constitutive Fe deficiency response.

Strong accumulation of Fe causes *b104^dPAA* plants to have abnormal shoot and root

development and delayed flowering, accompanied by mis-regulated Fe response and

upregulated Fe acquisition genes. The severity of Fe accumulation phenotypes

coincides with bHLH104 protein amounts. In plants, bHLH104 levels are controlled by

proteasomal degradation in a PAA-dependent manner. Cell assays show that the PAA

sequence is required for bHLH104 interaction with BTS and subsequent ubiquitination.

Taken together, bHLH104 requires the terminal PAA to be post-translationally

controlled by BTS. Manipulation of bHLHIVc protein PAA sequence represents a biofortification strategy for crop plants.

60-word Plain language summary

Plants need iron but excess is harmful. Iron uptake transcription factor bHLH104 is negatively controlled by Fe-binding E3 ligase BTS. We show that three–amino acid terminal sequence of bHLH104 controls its breakdown by the BTS pathway. Removing PAA keeps bHLH104 active, driving constant iron uptake and iron accumulation phenotypes. Targeting this motif can be a strategy for crop biofortification.

Highlights of this study

- A conserved C-terminal PAA sequence in bHLH104 is required for its post-translational control in *Arabidopsis thaliana.*
- Expression of bHLH104 lacking the PAA sequence (*b104^dPAA*) perturbs Fe regulation, resulting in constitutive iron accumulation in plants depending on *b104^dPAA* protein levels.
- The PAA sequence mediates interaction with E3 ligases BTS and is essential for bHLH104 ubiquitination.

Keywords: *Arabidopsis thaliana*, iron, bHLHIVc, transcription factor, ubiquitination, BTS,

Introduction

Iron (Fe) is an essential micronutrient required for plant growth and development. It plays critical roles in photosynthesis, respiration, chlorophyll biosynthesis, and redox reactions (Briat et al., 2007; Jeong and Guerinot, 2009). Although Fe is abundant in the Earth's crust, it is mostly present in the ferric (Fe^{3+}) form, which is poorly soluble and difficult for plant roots to absorb—especially in alkaline or calcareous soils (Marschner, 2012). At the same time, excess Fe can be toxic due to its ability to generate reactive oxygen species (ROS) (Panda and Yamamoto, 2019). Therefore,

plants must tightly regulate Fe uptake, distribution, and storage to maintain internal balance.

In *Arabidopsis thaliana*, a group of basic helix–loop–helix (bHLH) IVc transcription factors—including bHLH34, bHLH104, ILR3 (bHLH105), and bHLH115—acts as an early regulatory module in the Fe deficiency (-Fe) response. These proteins activate the expression of internal Fe allocation and Fe deficiency response regulatory genes in roots and shoots (Schwarz and Bauer, 2020). Among their targets are also genes coding for bHLH Ib transcription factors that, together with the bHLH transcription factor FIT (FER-LIKE IRON DEFICIENCY-INDUCED TRANSCRIPTION FACTOR), induce root Fe uptake genes such as *IRT1* and *FRO2* (Long et al., 2010; Selote et al., 2015; Zhang et al., 2015; Li et al., 2016; Liang et al., 2017).

The regulation of bHLH IVc transcription factors is not fully understood. Notably, transcript levels of *bHLH IVc* genes do not change substantially under - Fe conditions (Zhang et al., 2015; Li et al., 2016; Liang et al., 2017), indicating that they are activated during the - Fe response primarily at protein level. One proposed mechanism involves post-translational regulation via the 26S proteasome mediated by the E3 ubiquitin ligase proteins, BRUTUS (BTS)/BTS-LIKE (BTSL) proteins. It is proposed that these E3 ligases target bHLHIVc proteins such as bHLH105/ILR3 and bHLH115 for degradation (Selote et al., 2015; Hindt et al., 2017). This hypothesis is particularly intriguing since stability of the BTS/BTSL E3 ligases is controlled by their N-terminus which contains hemerythrin domains for Fe and oxygen sensing (Selote et al., 2015; Pullin et al., 2025). It is possible, and supported by genetic studies that, BTS become destabilized under - Fe, resulting in their inactivation and allowing downstream targets such as bHLH IVc transcription factors to activate Fe uptake. However, molecular mechanisms are not yet fully resolved.

One bHLH IVc group protein is bHLH104 (Zhang et al., 2015; Li et al., 2016). The protein-level regulatory mechanisms controlling its function remain incompletely characterized. The C-terminus of bHLH104 can interact with BTS/L proteins (Lichtblau et al., 2022). It possesses a conserved C-terminal sequence. In our previous study, we showed that deleting this 25-amino-acid-long stretch abolished bHLH104 interaction with BTS (Lichtblau et al., 2022). T-DNA loss-of-function lines have an Fe deficiency-sensitive phenotype (Li et al., 2016). Previous studies have shown that

mutations at the C-terminal region of bHLH IVc proteins, including large deletions (60-
64 amino acids) or point mutations at the terminal alanine residue, cause Fe
accumulation phenotypes or disrupt interaction with proteosome components
(Rampey et al., 2006; Zhao et al., 2016; Sharma and Yeh, 2020; Li et al., 2021).
Structural predictions further revealed that a conserved three–amino acid short
sequence—proline–alanine–alanine (PAA)—is located near the BTS-binding interface
(Lichtblau et al., 2022). Hereafter, we refer to this as the PAA short sequence.

So far, it has not yet been demonstrated whether the PAA short sequence is important
for bHLH104 regulation. The key open questions of this study were whether this minimal
motif is necessary and sufficient for BTS engagement and ubiquitination, whether it
governs bHLH104 protein abundance, and how its loss impacts Fe homeostasis and plant
development. We show that the PAA short sequence plays a critical role for all these
steps and in regulating Fe homeostasis properly in *planta*. Our work provides
mechanistic insight into how bHLH104 is regulated and highlights the broad impact of
post-translational control in nutrient signaling networks.

Results

Plants with bHLH104 lacking PAA (*b104^dPAA*) have distinct growth phenotypes

At first, we tested the hypothesis that the PAA short sequence plays a role in the
functionality of bHLH104 in plants (Figure 1). We studied the growth of transgenic
Arabidopsis thaliana lines in the wild-type Col-0 background expressing either regular
full-length bHLH104 (*pb104::b104*) or a mutant deletion variant lacking the conserved
PAA short sequence (*pb104::b104^dPAA*), both driven by the native *BHLH104*
promoter (Figure 2A). Interestingly, obvious phenotypic variation was already visible
in the T1 population of the *pb104::b104^dPAA* transgenics: While some plants
displayed severely reduced growth with brown necrotic spots on leaves, others
appeared indistinguishable from wild type (WT) (Supplementary Figure S1). To
capture these contrasting phenotypes, we selected and propagated four
pb104::b104^dPAA lines, two with severe phenotypes (#1, #2) and two with no obvious
macroscopic phenotype (#3, #4), in addition to *pb104::b104 #1* and *#2* controls.

To quantify phenotypic variation across the life cycle, detailed morphological and physiological assessments were made with T3 plants grown on agar plates or soil. In 12-day-old seedlings, *pb104::b104^dPAA* lines *#1* and *#2* showed significantly shorter primary roots than #3, #4 or *pb104::b104* control lines (Figure 2B, 2C). Root Fe(III) reductase activity is necessary for Fe uptake in roots (Yi and Guerinot, 1996). Reductase activity measurements revealed that *pb104::b104^dPAA* lines *#1* and *#2* have high Fe reductase activity compared to #3, #4 and the control plants (Figure 2D). The machine-aided Phenospex PlantEye phenotyping platform can reliably depict minor variation in leaf color and rosette size in *A. thaliana* (Knopf and Bauer, 2025). By four weeks, *pb104::b104^dPAA* lines *#1* and *#2* displayed low rosette area (Figures 2E, 2F) and low normalized difference vegetation index (NDVI) values (Figure 2G), which indicate low chlorophyll content, photosynthetic efficiency and growth performance compared with other lines. Flowering was delayed compared to other lines (Figure 2H) and seed yield was significantly lower for *pb104::b104^dPAA* lines *#1* and *#2* compared to other lines (Figure 2I).

Having established that pb104::b104^dPAA lines exhibited distinct growth phenotypes (Figure 2), we selected among them representative lines for detailed molecular and physiological characterization. We focused subsequent analyses on pb104::b104^dPAA line #2, which showed strong phenotypes, and line #3, which showed mild phenotypes, to comprehensively analyze Fe content, gene expression, and the molecular basis of these phenotypes.

Reduced root and shoot growth can be a consequence of Fe toxicity in plants (Naranjo-Arcos et al., 2017). Thus, we tested whether the growth defects observed in *pb104::b104^dPAA* plants were alleviated in conditions of Fe deficiency. We grew the representative lines on alkaline calcareous soil (ACS; pH 8.1), which provides Fe-limiting conditions (Supplementary Figure S2; Knopf and Bauer, 2025). Notably, *pb104::b104^dPAA line #2*, which exhibited pronounced growth defects in control soil (pH 5.9), showed increased 3D leaf area in ACS, indicating partial recovery under Fe-limited ACS conditions (Supplementary Figure S2C). NDVI values, which were reduced when grown in control soil, were restored to near-WT levels in ACS, suggesting improved leaf health (Supplementary Figure S2D). While flowering remained delayed relative to WT, it occurred earlier in ACS than in control soil (Supplementary Figure S2E). These findings show that the growth defects observed

in control soil are largely alleviated when Fe availability is limited, linking the
b104^dPAA-associated phenotypes to Fe status.

Together, these results show that removal of the PAA short sequence in bHLH104 can
lead to drastic dominant growth phenotypes across the life cycle. To explore whether
these phenotypes reflect altered Fe homeostasis, we next examined tissue Fe content
and expression of Fe-responsive genes in both seedlings and plants at the
reproductive stage.

b104^dPAA can Cause Constitutive Iron Uptake and Transcriptional Misregulation of Fe Deficiency Response genes in Seedlings

After few days of germination, *A. thaliana* seedlings rely on external Fe supply. To
assess whether b104^dPAA affects Fe homeostasis during early development, we
measured total Fe content via ICP-MS and analyzed gene expression profiles in 10-
day-old seedlings of WT, control and contrasting *pb104::b104^dPAA* lines #2 and #3
in both +Fe and -Fe conditions (Figure 3A, B). As an additional control, *pUBQ10::b104*
lines express full-length bHLH104 constitutively from the strong UBIQUITIN10
promoter, allowing us to assess whether high transcript levels alone affect Fe
homeostasis. Phenotypically, *pUBQ10::b104* lines #1 and #2 are indistinguishable
from WT (Supplementary Figure S3).

Elemental profiling revealed that *pb104::b104^dPAA* line #2 accumulated substantially
more Fe than all other genotypes under both +Fe and –Fe conditions (Figure 3C). This
increase was observed in both roots and shoots, with roots showing higher Fe levels
than shoots. In contrast, *pb104::b104^dPAA* line #3 and the *pb104::bHLH104* and
pUBQ10::b104 lines had Fe contents similar to WT. All lines exhibited reduced Fe
contents under –Fe, confirming the effectiveness of the treatment (Figure 3C).

Differences were also detected for manganese (Mn), zinc (Zn), and copper (Cu) levels
(Supplementary Figure S4). Uptake of these metal ions is closely connected with Fe
homeostasis (Connolly et al., 2002). In *pb104::b104^dPAA* line #2, Zn and Cu were
elevated in roots under –Fe, and Cu was also increased in shoots. Notably, Zn levels
were elevated even under +Fe in *pb104::b104^dPAA* roots compared to WT. Roots of
pUBQ10::b104 lines also showed increased Mn under –Fe compared to WT. Metal
concentrations in the other lines were similar to WT. Hence, these findings suggest
that *pb104::b104^dPAA* line #2 has increased Fe accumulation and affects the uptake

of other micronutrients like Zn and Cu. This highlights the importance of proper
regulation of bHLH104 function in maintaining metal homeostasis under varying Fe
conditions.

To elucidate the transcriptional response, we analyzed gene expression patterns via
RNA-seq following -Fe treatment at the seedling stage in both root and shoot tissues.
The highest number of differentially expressed genes (DEGs) was observed in shoot
tissue of *pb104:b104^dPAA #2* line, with 4,645 DEGs under +Fe conditions and 2,857
DEGs under -Fe conditions after 10 days of treatment, compared to WT. There was
also substantial differential expression in roots, with 1,210 DEGs under +Fe conditions
and 111 DEGs under -Fe conditions (Supplementary Figure S5A).

Gene Ontology (GO) analysis revealed that the upregulated genes in shoot tissue of
the *pb104:b104^dPAA* #2 plants were enriched in terms related to hypoxia and
oxidative stress, while genes associated with photosynthesis were downregulated. In
pb104:b104^dPAA root tissue, upregulated gene categories were additionally
enriched for terms related to metal ion transport, while downregulated categories
showed enrichment for cell wall modification-related GO terms (Supplementary Figure
S5B). Hence, the *pb104:b104^dPAA #2* plants indeed show symptoms of Fe overload.

We examined the expression patterns of selected genes involved in Fe uptake and
homeostasis regulation to characterize their specific regulatory expression profiles.
BHLHIVc genes, *BHLH104, ILR3*, and *BHLH115*, were upregulated in the roots of the
pb104::b104^dPAA line #2 under +Fe compared to other lines (Figure 3D). Their target
genes, such as *BHLH39, BHLH100, IMA1*, *PYE, BTS, BTSL1* and BTSL2, were also
upregulated, indicating activation of the Fe deficiency response signaling. Likewise,
the further downstream targets in roots, like Fe acquisition genes *IRT1, FRO2, FIT,*
were up-regulated (Figure 3D), consistent with the elevated Fe uptake and Fe content.
Interestingly, expression of negative regulators of the Fe deficiency response and of
Fe sufficiency markers, such as *BHLH11* and FER1 were also increased, suggesting
a proper response to sufficient Fe, but a failure in feedback repression mechanisms
(Figure 3D). In the shoot tissue, the Fe storage related gene *FER1* and transcription
factor bHLH121 were strongly upregulated in *pb104::b104^dPAA line #2* compared to
controls, regardless of Fe availability, in agreement with the elevated Fe content in this
line (Figure 3D). Targeted expression profiling using reverse transcription quantitative

PCR (RT-qPCR) in seedling-stage samples confirmed expression patterns to those observed in the transcriptome analysis (Supplementary Figure S6).

To summarize, the misregulation of Fe response genes indicates that bHLH104 with deleted PAA disrupts normal Fe homeostasis by constitutively activating Fe acquisition while preventing proper feedback inhibition during seedling development.

b104^dPAA Causes Constitutive Fe Uptake and Transcriptional Misregulation at a Reproductive Growth Stage

Plants redistribute nutrients during reproduction to support flowering and seed development (Waters and Grusak, 2008; Pottier et al., 2018). To assess whether b104^dPAA affects Fe homeostasis during later developmental stages, we analyzed 30-day-old Arabidopsis plants grown hydroponically and subjected to a 3-day ±Fe treatment (Figure 4A). This time, we focused on *pb104::b104^dPAA #3*. Due to its severe developmental delay and low biomass regarding very severe growth defects, *pb104::b104^dPAA #2* had to be excluded from this analysis and was examined separately, as described later (Figure 4B). Old leaves, young leaves, and roots were collected for gene expression analysis, as described in (Ngigi et al., 2025) (Figure 4C-D).

Elemental profiling showed that although there were no differences from WT at the seedling stage, *p104::b104^dPAA line #3* accumulated more Fe than control lines at the late developmental stage (Figure 4E). Under +Fe, roots of *pb104::b104^dPAA #3* had increased Mn and Zn levels compared to WT (Supplementary Figure S7). Cu levels were similar as WT.

In transcriptome analysis, *pb104:b104^dPAA #3* plants had the highest DEG counts in young leaves (3,877 DEGs) and old leaves (1,895 DEGs) under +Fe conditions compared with controls, while root tissue exhibited minimal differential expression differences (71 DEGs) (Supplementary Figure S8A). GO term enrichment analysis revealed that genes associated with immune response were significantly upregulated in both old and young leaves of *pb104:b104^dPAA #3* plants under +Fe conditions. In contrast, 3-day -Fe treatment resulted in the upregulation of genes related to Fe homeostasis in young leaves and roots (Supplementary Figure S8B). This indicates a tissue-specific transcriptional response to iron availability at the reproductive stage.

Specifically, expression analysis showed that in roots of *pb104::b104^dPAA line #3* genes were upregulated under +Fe, similar as we reported above for the #2 line at the seedling stage, namely upregulation of *BHLHIVc* genes, Fe deficiency response, Fe sufficiency marker and negative regulator genes (Figure 4F). Interestingly, Fe deficiency response genes were not upregulated under –Fe in *pb104::b104^dPAA line #3*, suggesting that the –Fe response may have been suppressed due to negative feedback signaling caused by elevated internal Fe levels in this line, in contrast to controls where these genes were upregulated (Figure 4F). In line with this notion, the Fe storage gene *FER1* was upregulated in the leaves of *pb104::b104^dPAA* line #2 compared to controls. Targeted expression profiling by RT-qPCR confirmed again the patterns (Supplementary Figure S9).

Finally, as *pb104::b104^dPAA* line #2 grew poorly in +Fe hydroponic conditions, we modified the hydroponic growth system and changed to a Fe resupply assay. Seedlings were transferred to -Fe medium for 27 days, followed by a 3-day +/- Fe resupply. Here, we found interesting phenotypes. While controls remained chlorotic and accumulated Fe mostly in the roots following the Fe resupply, *pb104::b104^dPAA* line #2 showed substantial Fe accumulation in root and leaves despite persistent leaf chlorosis, suggesting very high root-to-shoot Fe translocation. Interestingly, *pb104::b104^dPAA* line #2 accumulated significantly less Mn, Zn, Cu compared to control plants (Supplementary Figure S10). This shows that at the reproductive stage, there is a particularly elevated Fe uptake and translocation towards shoots in *pb104::b104^dPAA* #2 compared to the control.

Since *pb104::b104^dPAA* lines accumulated not only Fe but also Mn, Zn, and Cu, we examined the possibility that this might have been caused by altered expression of metal transporter genes. We found elevated *NRAMP1*, *COPT2*, *MTP8*, and *ZIP* family members in *pb104::b104^dPAA* roots under +Fe at the seedling stage (Supplementary Figure S11). Interestingly, under –Fe, transporter gene expression levels were lower, yet Mn, Zn and Cu accumulation were higher, suggesting that in the absence of Fe, other metals are taken up more efficiently (Supplementary Figures S7 and S11). At the reproductive stage, *pb104::b104^dPAA* roots maintained high expression of metal transporters including *NRAMP1, ZIP3, COPT2, MTP8,* and *ZIP9*, yet only Fe accumulated, but not other elements, indicating Fe is the primary target despite activation of multiple metal transporters (Supplementary Figure S11). Interestingly, Fe

acquisition and mobilization genes were constitutively upregulated in *pb104::b104^dPAA* lines regardless of Fe availability at the seedling stage (Supplementary Figure S12).

To summarize, Fe accumulation is elevated in *pb104::b104^dPAA* lines during the reproductive stage, consistent with patterns observed at the seedling stage and with iron and metal transporter gene regulation patterns. Overall, these findings show that metal accumulation and associated phenotypes due to *b104^dPAA* expression remain consistent from seedling to reproductive plant developmental stages.

The PAA short sequence is required for 26S Proteasome-mediated degradation of bHLH104

The severity of phenotypes in *pb104::b104^dPAA* lines was not explained by differences in transgene transcript levels (Supplementary Figures S6, S9). Additionally, *pUBQ10::b104* lines showed no significant phenotype compared to WT despite elevated *bHLH104* transgene expression (Supplementary Figure S3, Supplementary Figure S6). Therefore, we predicted that variation in bHLH104 protein accumulation underlies the observed phenotypic differences.

We thus examined *in vivo* accumulation of mTurquoise2-tagged bHLH104 and b104^dPAA proteins, expressed under either the UBQ10 or bHLH104 promoter (Figure 5A, B). Immunoblotting of root tissue proteins revealed high accumulation of b104^dPAA protein in *pb104::b104^dPAA* lines #1 and #2, correlated with their strong phenotypes (Figure 5B). In contrast, full-length bHLH104 protein was barely detectable in both *pb104::bHLH104* and *pUBQ10::b104* lines. *pb104::b104^dPAA* lines #3 and #4 showed lower protein abundance, corresponding to weaker phenotypes.

To check whether the low abundance of bHLH104 is due to proteasomal degradation, we treated samples with proteosome inhibitor MG132. MG132 treatment stabilized full-length bHLH104 in *pUBQ10::b104* lines by more than two-fold. However, MG132 treatment did not affect the abundance of bHLH104 in any native promoter lines (Figure 5B-C). To confirm that bHLH104 is subject to proteasomal degradation, a cell-free degradation assay was performed. When bHLH104 protein was incubated with whole protein extracts from WT seedlings, it was rapidly degraded. This degradation

was inhibited by the proteasome inhibitor MG132, supporting the involvement of the 26S proteasome (Figure 5D).

Confocal imaging of root meristems further confirmed the increased abundance of b104^dPAA, as indicated by strong fluorescence signals in *pb104::b104^dPAA* lines, whereas fluorescence was undetectable in lines expressing full-length bHLH104 (Figure 5E).

Altogether, these findings show that the PAA short sequence is most required for degradation of bHLH104 by the 26S proteasome. Removing this short sequence prevents normal turnover of the protein and is correlated with increased Fe uptake and associated phenotypes.

The C-terminal PAA sequence of bHLH104 mediates interaction with BTS E3 ligases and enables ubiquitination

To understand how the PAA sequence contributes to bHLH104 degradation, we examined its role in mediating interaction with BTS.

AlphaFold2 multimer modeling (Evans et al., 2021; Jumper et al., 2021) and electrostatic surface analysis (Pettersen et al., 2004) revealed a charge-based interface with high pLDDT scores, indicating strong confidence in the predicted structure (Figure 6A-B). These models suggest that the PAA sequence forms a stable and specific docking surface for BTS, likely critical for E3-substrate recognition.

To further study protein interaction in the presence and absence of PAA experimentally, we made use of established protein expression systems. Due to the difficulties of detecting bHLH104 and BTS in plant cell assays, we opted for the yeast two hybrid (Y2H) assay (Lichtblau et al., 2022). BTS interacts with the C-terminal part of bHLH104 (bHLH104-C) in yeast, and bHLH104-C does not auto-activate in the yeast-two-hybrid (Y2H) system (Figure 6C) (Lichtblau et al., 2022). We now tested whether interaction is different with bHLH104-C^dPAA. Indeed, deletion of the PAA sequence abolished the interaction with BTS (Figure 6D). bHLH104 is known to homodimerize (Zhang et al., 2015; Li et al., 2016; Lichtblau et al., 2022). bHLH104-C^dPAA still dimerized with bHLH104-C, confirming that bHLH104-C^dPAA is present and properly folded in yeast cells. A similar loss of interaction was observed between bHLH104-C-dPAA and and

BTSL1 and BTSL2 (Supplementary Figure S13), indicating that the PAA sequence is broadly required for recognition by BTS/ BTSL E3 ligases.

To test whether bHLH104 is ubiquitinated by BTS with requirement of the short sequence PAA, we established the bacterial UbiGate system (Kowarschik et al., 2018; Ortmann et al., 2023) to reconstitute a BTS-dependent ubiquitin-conjugation cascade for bHLH104 (Figure 6E). Previous reports on the ubiquitination activity of BTS have been based on *in vitro* ubiquitination assays (Selote et al., 2015; Rodríguez-Celma et al., 2019). However, *in vitro* studies are often ambiguous as purified target proteins are contained in a reaction mix. In this study, we employed the UbiGate system, in which all components of the ubiquitination cascade—including ubiquitin, E1, E2, E3 ligases, and the substrate protein—are co-expressed in bacterial cells, a far more complex environment than an *in vitro* reaction. The resulting ubiquitination patterns are analyzed directly from bacterial cell lysates. Each component is fused to a distinct epitope tag, allowing individual tracking and detection of the proteins involved. Since full-length BTS is unstable in *E. coli*, we used a truncated version (BTS-HC) lacking the first two hemerythrin domains. In the presence of E1, E2 (UBC8), and BTS-HC, bHLH104-C was polyubiquitinated, while, in contrast, this was not the case for b104-C^dPAA (Figure 6F). BTS ubiquitination served as a control. It has been reported that BTS undergoes autoubiquitination (Selote et al., 2015), which we also observed in our assay. bHLH115 has been previously proposed as a target of BTS-dependent degradation (Selote et al., 2015; Xing et al., 2021). As a validation of the UbiGate assay, we confirmed that BTS directly ubiquitinated bHLH115 (Supplementary Figure S14).

These results provide direct biochemical evidence that bHLH104 is a substrate of BTS inside living cells and that the PAA short sequence is required for both physical interaction and ubiquitination. Deletion of PAA sequence disrupts recognition and ubiquitination by BTS.

Discussion

Our findings in this study demonstrate that the terminal PAA short sequence of bHLH104 is most relevant for bHLH104 control via proteasome-mediated turnover. Lack of PAA leads to higher abundance and a stabilization of bHLH104, constitutive

Fe uptake, and transcriptional misregulation across developmental plant growth stages. The PAA sequence is directly involved in protein interaction and ubiquitination by BTS, highlighting its importance as an E3 ligase–interaction interface in maintaining Fe homeostasis through post-translational regulation of transcription factors and transcriptional output.

bHLH104 abundance is controlled via the proteasome

The regulation of bHLH104 activity at protein level had been largely unknown, but regarded relevant (Zhang et al., 2015; Li et al., 2016; Liang et al., 2017). Our results show that wild-type bHLH104 protein is susceptible to degradation in a plant cell environment, and inhibition of the proteasome pathway increases bHLH104 abundance. Interestingly, despite having high *BHLH104* transcript levels, pUBQ10::b104 lines showed no phenotype, implying that protein stability, rather than transcript level, determines bHLH104 activity.

The direct interaction of bHLH104 with the BTS E3 ligase has been reported previously (Long et al., 2010; Selote et al., 2015; Lichtblau et al., 2022), and genetic evidence supported a regulatory relationship between BTS and bHLH104 (Zhang et al., 2015). The *bts-2* mutant exhibits increased Fe uptake similar to bHLH104 overexpression, whereas *bts-2 bhlh104-2* double mutants lose this phenotype, confirming that BTS negatively regulates bHLH104 function (Zhang et al., 2015). Recent studies have identified that BTS interacts with the C-terminal region of bHLHIVc proteins (Li et al., 2021; Lichtblau et al., 2022). Interestingly, C-terminal mutations in bHLH IVc proteins in other plant species also result in Fe accumulation (Rampey et al., 2006; Zhao et al., 2016; Sharma and Yeh, 2020), suggesting the functional importance of this conserved region. We tested this hypothesis and identified that PAA is needed for BTS interaction. We propose a model (Figure 7) in which bHLH104 protein is maintained at low abundance through proteasome-mediated degradation. Lack of the PAA sequence increases bHLH104 protein abundance, which results in increased Fe uptake in plants. A limitation of this study is the difficulty in visualizing mTurquoise2-tagged bHLH104 in planta under standard conditions, and the relatively low abundance of b104^dPAA protein suggests that additional regulatory mechanisms beyond PAA-mediated degradation may exist. Future studies should investigate other post-translational modifications and identify conditions that promote bHLH104 stabilization.

b104^dPAA perturbs Fe regulation and metal homeostasis

bHLHIVc transcription factors are functionally redundant (Zhang et al., 2015; Li et al., 2016; Liang et al., 2017). Our study showed that gain-of-function of bHLH104 alone caused a physiological effect in *pb104::b104^dPAA* lines where bHLH104 protein abundance was increased. Notably, *pb104::b104^dPAA* lines accumulated not only Fe but also Mn, Zn, and Cu, which coincided with elevated expression of *NRAMP1*, *COPT2*, *MTP8*, and *ZIP* family members in *pb104::b104^dPAA* roots under +Fe. This indicated that indeed excessive metal acquisition was a consequence of elevated gene expression and resulting transporter activity in these cases. The lower metal transporter gene expression at -Fe when elevated Mn, Zn and Cu accumulation was noted, was indeed surprising, suggesting that in the absence of Fe, other metals are taken up more efficiently, perhaps through the regular -Fe-induced Fe acquisition routes involving IRT1 (Vert et al., 2002). Indeed, the multi-metal accumulation can be explained by persistent activation of broad-specificity transporters IRT1 and NRAMP1, which transport Fe, Mn, Zn, and Cu (Connolly et al., 2002; Vert et al., 2002; Cailliatte et al., 2010). This pattern is also consistent with reports that heavy metals like Zn can induce Fe deficiency responses at the level of Fe sensing rather than by directly inhibiting Fe transport (Lešková et al., 2017), suggesting that accumulated metals may reinforce the Fe deficiency signal in *pb104::b104^dPAA* plants. Similar phenotypes where Fe accumulation disrupts plant growth have been reported in *BTS* loss-of-function mutants and in overexpression lines of *BHLH39* or *BHLH105* (Zhang et al., 2015; Li et al., 2016; Hindt et al., 2017; Naranjo-Arcos et al., 2017). A still unresolved question is why the known negative post-translational regulation mechanisms, e.g. IRT1 ubiquitination and degradation under low Fe and excessive Zn and Mn, are not in place here (Dubeaux et al., 2018; Abuzeineh et al., 2022). Perhaps this type of regulation requires that Zn and Mn are supplied in the rhizosphere rather than inside the root. In our experimental settings, excessive Zn and Man were not supplied and Zn and Mn accumulated due to plant internal misbalance.

We observed that b104^dPAA protein abundance in seedling roots correlated with phenotype severity among *pb104::b104^dPAA* lines. A plausible explanation is that this variation in protein abundance across lines results from differences in transgene

copy number arising from the floral dip transformation method (Gelvin, 2003). Notably, *pb104::b104^dPAA line #3*, which showed negligible b104^dPAA protein in seedling roots, grew normally during early development but accumulated Fe in leaves at the reproductive stage. This suggests that minimal b104^dPAA activity can increase Fe content without causing severe growth defects, highlighting potential for Fe biofortification strategies. Additional investigations are required to determine the precise threshold at which bHLH104 protein levels affect plant function without compromising growth. In sum, our discovery that a three–residue PAA sequence gates BTS-mediated turnover of bHLH104 delivers a precise target for tuning Fe acquisition in plants. The findings integrate with emerging work on BTS/BTSL hemerythrin-based Fe/oxygen sensing (Selote et al., 2015; Pullin et al., 2025) and clarify how post-translational control translates Fe status into transcriptional outputs.

Pleiotropic and tissue-specific effects of b104^dPAA

The phenotypes observed in *pb104::b104^dPAA* plants extend beyond simple Fe accumulation, suggesting that constitutive Fe deficiency signaling has broader physiological consequences. Several phenotypes observed in these lines may reflect activation of multiple stress regulatory pathways. First, the delayed flowering phenotype may result from -Fe response signaling interfering with flowering time control through epigenetic, circadian, and hormone pathways (Cheng et al., 2004; Salomé et al., 2012; Chen et al., 2013; Hong et al., 2013; Wild et al., 2016; Dutta et al., 2017; Maruoka et al., 2022). Second, gene ontology analysis of *pb104::b104^dPAA* leaves showed enrichment of immune signaling and salicylic acid signaling terms. Fe deficiency treatment is known to elevate salicylic acid levels and induce salicylic acid-responsive genes (Shen et al., 2016), and bHLH105 can integrate Fe homeostasis with plant immune responses against pathogens (Rampey et al., 2006; Aparicio and Pallás, 2017; Samira et al., 2018). This suggests that the Fe deficiency response is interconnected with immune signaling pathways.

Expression of b104^dPAA in *pb104::b104^dPAA* lines results in distinct tissue-specific consequences. The absent upregulation of internal Fe allocation genes such as *BTS, PYE, IMA1*, *BHLH100, BHLH39* under -Fe condition in the leaves of mature *pb104::b104^dPAA* lines suggests that elevated local Fe levels in the leaves must normally suppress systemic Fe deficiency signaling upstream of bHLH104, consistent with recent evidence that leaf Fe responses are determined mainly by local Fe status

rather than root-derived systemic signals (Nguyen et al., 2022; Ngigi et al., 2025). In contrast, in roots of *pb104::b104^dPAA* lines, -Fe acquisition genes such as *FIT*, *IRT1* and *FRO2* were upregulated. This indicates that, rather than systemic signaling, an active and constitutive Fe acquisition response is maintained in the roots. In wild-type plants, bHLH104 is likely stabilized only under Fe deficiency, enabling transient activation of uptake genes. Lack of PAA prevents this regulation, resulting in constitutive bHLH104 activity and continuous Fe uptake regardless of Fe status. Interestingly, we observed enhanced Fe mobilization to young leaves in *pb104::b104^dPAA line #2* after 3 days of Fe supply following growth in –Fe. In contrast, wild-type plants retained most Fe in roots. Increased *IRT1* expression when b104^dPAA is expressed aligns with recent findings that IRT1 protein, independent of its Fe(II) transport activity, is necessary for effective root-to-shoot Fe partitioning (Quintana et al., 2022). This shows that bHLH104 protein stability serves as a critical control point linking Fe sensing to transcriptional responses in roots, whereas leaves respond primarily to local Fe availability through independent regulatory mechanisms. Grafting experiments would help determine whether the effects of stabilized bHLH104 are root-autonomous or involve systemic signaling to shoots.

The short sequence PAA is critical for protein interaction

The present study found that the terminal PAA sequence of bHLH104 mediates critical interactions. Additionally, small proteins IMA1 and IMA3, which interact with BTS and BTSL1, also contain a conserved C-terminal PAA sequence, indicating that this sequence is broadly conserved for E3 ligase recognition (Li et al., 2021; Lichtblau et al., 2022). Other members of the bHLH IVc family have instead a PVA terminal sequence. It has not yet been examined which role PVA plays and whether it is similar to PAA. One possible explanation is that the middle alanine in the PAA sequence of bHLH104 may not be essential. Proline is structurally distinctive because its cyclic side chain imparts rigidity and promotes tight turns in protein structure, suggesting that the proline residue plays a critical structural role (Melnikov et al., 2016). Previous studies have shown the significance of terminal alanine residue in bHLH IVc protein regulation. The terminal alanine in bHLH115 and bHLH105 is essential for BTS interaction (Li et al., 2021), while overexpression of bHLH34 with its terminal alanine substituted to valine resulted in a metal accumulation phenotype similar to our observations (Sharma and Yeh, 2020). Taken together, these observations support the conclusion that the

PAA sequence is required for the recognition of bHLH104 by the proteasomal degradation system. The conservation of this sequence in bHLH104 orthologues, and its presence with minor variations in other bHLH IVc members, suggests the existence of a conserved BTS(L) recognition interface within the family. In future studies, the specificity and position effects of each amino acid in the PAA and PVA short sequences can be further examined. It will also be interesting to test whether artificial addition of PAA to transcription factors can target them for degradation by BTS/L proteins under Fe deficiency. If so, this can be exploited in the future in synthetic and biotechnological approaches.

Concluding remarks and perspectives

This study provides evidence that a short three–amino acid PAA sequence in bHLH104 is required for post-translational regulation of protein abundance and activity.

The conserved PAA sequence provides opportunities for biotechnological applications. As bHLH IVc proteins are conserved across plant species (Jiang et al., 2024), genome editing of this sequence in crop orthologues may enhance Fe accumulation for biofortification. In contrast to whole-gene overexpression, precise PAA editing offers potential to increase Fe content while minimizing pleiotropic effects. However, because the PAA sequence is also present in IMA proteins, which positively regulate Fe uptake, editing approaches must specifically target bHLH IVc genes without affecting *IMA* genes. Testing whether PAA addition to other transcription factors targets them for BTS/BTSL-mediated degradation could enable Fe-responsive protein degradation systems for synthetic biology applications.

Overall, our findings provide comprehensive evidence which shows that post-translational regulation of bHLH104 is necessary to maintain a balance between Fe acquisition and normal growth and development, and that disrupting this control can lead to wide-ranging physiological effects. This research has direct impact for improving crop productivity on alkaline/calcareous soils and under climate stress. Identifying sequence motifs that control master regulators of Fe uptake such as bHLH104 provides paths for crop improvement and biofortification. Our discovery that a three–residue PAA motif gates BTS-mediated turnover of bHLH104 delivers a precise target for tuning Fe acquisition in plants. Given current efforts to enhance

micronutrient use efficiency and nutritional quality, these insights are both mechanistically fundamental and directly translatable.

Materials and methods

Gene Accession numbers

The following *Arabidopsis thaliana* genes were the focus of this study: bHLH104 (AT4G14410), BTS/BRUTUS (AT3G18290), BTSL1 (AT1G74770), and BTSL2 (AT1G18910).

Plant Material and Growth Conditions

Arabidopsis thaliana Columbia-0 (Col-0) was used as the wild type in this study. The transgenic lines pbHLH104::mTurquoise2:bHLH104, pUBQ10::mTurquoise2:bHLH104, and pbHLH104::mTurquoise2:b104^dPAA were used for phenotypic and molecular analyses.

Seeds were surface-sterilized in 6% (v/v) sodium hypochlorite solution supplemented with 0.1% (v/v) Triton X-100 for 8 min, followed by five washes with sterile distilled water. Sterilized seeds were stratified in the dark at 4°C for 48 h before sowing.

For plate-based experiments, seeds were germinated and grown vertically on modified Hoagland agar medium containing 1.5 mM $Ca(NO_3)_2 \cdot 4H_2O$, 0.5 mM KH_2PO_4, 1.25 mM KNO_3, 0.75 mM $MgSO_4 \cdot 7H_2O$, 50 µM H_3BO_3, 25 µM KCl, 10 µM $MnSO_4 \cdot H_2O$, 2 µM $ZnSO_4 \cdot 7H_2O$, 1.5 µM $CuSO_4 \cdot 5H_2O$, and 0.075 µM $(NH_4)_6Mo_7O_{24}$, supplemented with 1% (w/v) sucrose and 1.4% (w/v) plant agar, pH adjusted to 5.8. Iron was supplied as 50 µM Fe(III)-EDTA for Fe-sufficient conditions (+Fe) or omitted for Fe-deficient (-Fe) conditions. Seedlings were grown under long-day photoperiods (16 h light/8 h dark) at 21°C, under white light at 120 µmol m^{-2} s^{-1} (CLF Plant Climatics). For Fe deficiency experiments, seedlings were grown for 10 days before tissue collection. For root length measurements, an independent set of seedlings was grown on Hoagland plates for 12 days prior to measurement.

For hydroponic experiments, seedlings were initially grown on Hoagland agar plates for 12 days before being transferred to a hydroponic setup based on a quarter-strength

Hoagland solution (Nguyen et al., 2016) containing 0.75 mM $Ca(NO_3)_2$, 0.625 mM

KNO_3, 0.25 mM KH_2PO_4, 0.375 mM $MgSO_4$, 25 µM H_3BO_3, 25 µM KCl, 5 µM $MnSO_4$,

1 µM $ZnSO_4$, 0.75 µM $CuSO_4$, and 0.0375 µM $(NH_4)_6Mo_7O_{24}$. Fe was supplied as 25

µM FeNaEDTA (+Fe). The nutrient solution was replaced every three days and

continuously aerated using air pumps. Plants were grown under the same

photoperiod, light intensity, and temperature conditions as plate-grown seedlings.

Hydroponic growth continued until 27 days after germination, at which point plants

were divided into +Fe and −Fe conditions by either maintaining or omitting FeNaEDTA.

After three days of treatment (day 30), tissues were harvested, immediately frozen in

liquid nitrogen, and stored at −80°C.

For soil-based phenotyping experiments, seedlings were transferred to soil on the

eighth day after germination. Plants were grown in growth cabinets (Polyklima, True

Daylight + LED) under long-day conditions (16 h light/8 h dark) with a light intensity of

98–112 $µmol·m^{-2}·s^{-1}$. The temperature was maintained at 21°C during the light period

and 19°C during the dark period. Humidity was not controlled. Soil was prepared using

a peat-based substrate, and the pH was determined as described below. Eight plants

per genotype were grown, with eight pots fitted per tray. Phenotypic measurements

were taken after three weeks of soil growth.

Generation of Transgenic Lines

Constructs for generating transgenic lines were assembled using the GreenGate

cloning   system   (Lampropoulos   et   al.,   2013).   For   the

pbHLH104::mTurquoise2:bHLH104   and   pbHLH104::mTurquoise2:bHLH104^dPAA

constructs, a 1636-bp promoter region upstream of the bHLH104 start codon was

amplified and inserted into the pGGA000 entry vector. The coding sequences of

bHLH104 (852 bp) and bHLH104^dPAA (843 bp) were separately cloned into the

pGGC000 entry vector. The bHLH104 terminator sequence (975 bp) was cloned into

the pGGE000 entry vector. For both constructs, an N-terminal mTurquoise2

fluorescent tag from pGGB009 and a dummy linker sequence from pGGD002 were

included. The selection cassette (pUBQ10::HygromycinR::tOCS) was provided by

pGGF005. All entry modules were combined into the pGGZ003 destination vector

through a single GreenGate assembly reaction. Primers used are listed in

Supplementary Table 1.

For the pUBQ10::mTurquoise2:bHLH104 construct, the UBQ10 promoter was cloned from the pGGA006 entry vector to drive expression of the mTurquoise2-tagged bHLH104 coding sequence. The remaining modules, including the coding sequence, dummy linker, terminator, and selection cassette, were assembled as described above. Primers used in this study is listed in Supplementary table S1.

All final plasmids were verified by sequencing and introduced into Rhizobium radiobacter strain GV3103 (pSOUP). Arabidopsis thaliana Col-0 plants were transformed using the floral dip method (Clough and Bent, 1998). Transgenic seedlings were selected on hygromycin-containing medium. Independent T1 lines were established and propagated and used for phenotypic and molecular analyses.

Phenotyping and Image Acquisition

Automated phenotyping was performed using the MicroScan system equipped with a PlantEye F600 sensor (Phenospex, Heerlen, The Netherlands) and controlled by the implemented software HortControl (Phena version 2.0, HortControl version 3.85). To minimize interference from non-plant background, image analysis was optimized by removing color information corresponding to Hue values between 200° and 360° (blue to purple) as previously described (Arend et al., 2016; Hüther et al., 2020; Knopf and Bauer, 2025).

Plants were individually measured by placing the pots into a blue holder, which fixed the plant height and ensured positioning within the scanning unit. For technical replication, plant pots were rotated by 90° between successive measurements.

The PlantEye system recorded three-dimensional and spectral parameters. The three-dimensional surface area of the plants (3D leaf area, in mm²) was determined. Spectral data included, the normalized difference vegetation index (NDVI), calculated as (near-infrared reflectance [NIR] − red reflectance [RED])/(NIR + RED). For each parameter, both the average value per plant and the percentage of voxels falling within predefined ranges were calculated automatically using the preset analysis settings implemented in the HortControl software.

Root length was manually measured from plate-grown seedlings using Fiji (ImageJ).

Flowering time was determined by measuring the height of the primary inflorescence shoot. A plant was considered to have reached flowering if the inflorescence shoot

exceeded 1.5 cm. If a shoot was greater than 2 cm on the previous day, or greater than 3 cm two days earlier, the respective earlier day was recorded as the flowering time.

Seed number was estimated based on seed weight. For each biological replicate, 100 seeds were manually counted and weighed to determine the mean weight per 100 seeds. The total seed weight per sample was then measured, and the total seed number was calculated as (total seed weight/weight of 100 seeds)×100. Three biological replicates were analyzed for each genotype.

Fe reductase activity assay

Fe reductase activity was measured using a modified protocol based (Gratz et al., 2019). Ten-day-old Arabidopsis thaliana seedlings grown under Fe-sufficient or Fe-deficient conditions were used. For each replicate, roots from five plants were rinsed with 100 mM $Ca(NO_3)_2$ and transferred into 1.5 ml of assay buffer containing 100 μM FeNaEDTA and 300 μM ferrozine. The samples were incubated for 3 h in the dark. Formation of the Fe(II)–ferrozine complex was quantified by measuring absorbance at 562 nm using an Infinite 200 PRO plate reader (Tecan). Enzyme activity was calculated using the ferrozine extinction coefficient (ε = 28.6 mM^{-1} cm^{-1}) and normalized to total root length. Each assay included at least three biological replicates.

Plant Protein Extraction and Immunoblotting

Total proteins were extracted from roots of Arabidopsis thaliana seedlings. Root tissues were dried, weighed, frozen in liquid nitrogen, and ground using a Precellys homogenizer (three cycles at 6500 rpm). Proteins were solubilized in 2× SDG buffer (125 mM Tris-HCl, pH 6.8, 5% SDS, 4% DTT, 20% glycerol, 0.02% bromophenol blue) at a ratio of 10 μl buffer per 5 mg tissue. After rotation for 10 minutes and centrifugation (13,000 rpm, 10 minutes, 4°C), supernatants were collected and boiled at 60°C for 10 minutes.

Proteins were separated by SDS-PAGE using 4–20% Mini-PROTEAN TGX Stain-Free gels (Bio-Rad) and transferred to 0.45 μm nitrocellulose membranes (Bio-Rad) with the Trans-Blot Turbo Transfer System (Bio-Rad). Gels were imaged before transfer using the ChemiDoc MP Imaging System (Bio-Rad).

Membranes were blocked for 5 minutes in EveryBlot Blocking Buffer (Bio-Rad) and

incubated for 1 hour at room temperature with primary antibodies against mTurquoise2

(Biozol, NBS-ASJ-L9MMEL-60; 1:1000 dilution) or Actin (Agrisera, A0480; 1:5000

dilution) in 2.5% TBST with blocking buffer. After three washes in 1× TBST,

membranes were incubated with HRP-conjugated secondary antibodies (mouse anti-

goat for mTurquoise2 [Invitrogen, G-21040; 1:1000 dilution] and anti-mouse for Actin

[Jackson ImmunoResearch; 1:10,000 dilution in TBST with 5% milk]) for 1 hour.

Chemiluminescent signals were developed with ECL substrate (Amersham ECL

Select or SuperSignal West Femto) and imaged using the ChemiDoc MP Imaging

System.

To assess the effect of proteasome inhibition on bHLH104 protein stability, 10-day-old

Arabidopsis seedlings were transferred to liquid Hoagland medium and incubated for

18 hours in a plant growth chamber under the same growth conditions described

above. Seedlings were treated with either 50 µM MG132 or an equal volume of DMSO

(control). After treatment, total protein from roots was extracted and analyzed by

immunoblotting. Band intensities were quantified using Biorad-Image Lab™ Software,

and normalized to total protein staining. For each genotype, three independent

biological replicates were used. Relative protein abundance was calculated as the

ratio of normalized band intensity in MG132-treated versus DMSO-treated samples

(MG132/DMSO), representing the fold increase in bHLH104 accumulation upon

proteasome inhibition.

Cell-Free Protein Degradation Assay

The coding sequence of bHLH104 was cloned into an expression vector containing an

N-terminal His tag and transformed into Escherichia coli Rosetta (DE3) cells. Single

colonies were cultured overnight in LB medium with ampicillin, and secondary cultures

were inoculated at 1% volume, grown at 37°C to OD_{600} = 0.6–0.8, and induced with 1

mM IPTG at 28°C for 10–12 hours. Cells were harvested by centrifugation and stored

at –70°C.

For purification, frozen cell pellets were thawed on ice, resuspended in lysis buffer (30

mM Tris-HCl, pH 8.0, 150 mM NaCl) supplemented with protease inhibitors (Roche),

and sonicated. Lysates were clarified by centrifugation (14,000 rpm, 30 minutes, 4°C),

and His-tagged bHLH104 protein was purified using a Ni-NTA column (Cytiva) on a

Bio-Rad NGC chromatography system. Bound proteins were batch-eluted with 250
mM imidazole in lysis buffer, and fractions containing the target protein were pooled,
concentrated using Amicon centrifugal filters, and quantified using absorbance at 280
nm with a NanoQuant plate (Tecan). Protein concentrations were calculated using the
molar extinction coefficient predicted by ExPASy ProtParam, and proteins were
aliquoted and stored at –70°C.

We followed protocol from Wang et al., 2009, for cell-free degradation assay. Col-0
seedlings were grown for 10 days on Fe-deficient MS agar plates. Approximately 500
mg of root tissue was flash-frozen, ground using a Precellys homogenizer (6500 rpm,
three cycles of 20 seconds), and extracted in 500 µl degradation buffer (25 mM Tris-
HCl pH 7.5, 10 mM NaCl, 10 mM MgCl$_2$, 4 mM PMSF, 5 mM DTT, 10 mM ATP).
Extracts were clarified by centrifugation (15,000 rpm, 15 minutes, 4°C).

Degradation reactions were initiated by incubating 50 µl of plant extract with purified
His-bHLH104 protein at 22°C. Protein extracts were treated with either 160 µM MG132
or an equal volume of DMSO (control). Reactions were stopped at indicated time
points by adding SDS loading buffer and boiling at 95°C for 10 minutes. Proteins were
separated by SDS-PAGE, transferred to nitrocellulose membranes, and detected by
immunoblotting using an HRP-conjugated anti-His antibody (Thermo Fisher Scientific,
MA1-21315-HRP).

Confocal Microscopy

Confocal imaging was performed using a Zeiss LSM 880 laser-scanning confocal
microscope equipped with a 40× C-Apochromat water immersion objective. Two-day-
old Arabidopsis thaliana seedlings grown under Fe-sufficient (+Fe) conditions were
used for imaging. Excitation of mTurquoise2 fluorescence was carried out using a 458
nm laser line, and emitted fluorescence was collected between 493 nm and 613 nm.
Images were processed and analyzed using the manufacturer's software ZEN lite
(Zeiss).

ICP-MS Analysis

723 Tissue collection for inductively coupled plasma mass spectrometry (ICP-MS) was
724 performed with separate ceramic tools for Fe-sufficient and Fe-deficient samples to
725 prevent metal contamination. Roots, young leaves, and old leaves from hydroponically

grown plants, and shoots and roots from plate-grown seedlings, were harvested as pooled biological replicates (three replicates per line and condition). Tissues were washed sequentially in Tris-EDTA buffer and deionized water, dried on tissue paper, transferred into pre-weighed 5-ml Falcon tubes, and oven-dried at 65°C for up to two weeks.

Dry weights were recorded, and tissue digestion was performed with 67% nitric acid (ICP-MS grade) at volumes adjusted to sample mass (1 ml for 20–35 mg, 500 µl for 10–20 mg, 350 µl for <10 mg). Samples were incubated at room temperature overnight and then heated at 95°C until clear. After centrifugation (4000 rpm, 30 minutes, 4°C), supernatants were diluted with deionized water to maintain a final acid concentration below 5% (v/v). Dilution factors were calculated based on sample weight.

ICP-MS measurements were corrected for background signal and normalized by sample dilution factors. Statistical analysis was performed in R using two-way ANOVA followed by Tukey's post hoc test ($p < 0.05$), and data were visualized using ggPlantmap scripts as described in (Jo and Kajala, 2024).

RNA Isolation and RT-qPCR Analysis

Total RNA was extracted from roots and shoots of 10-day-old seedlings and from roots, young leaves, and old leaves of 30-day-old hydroponically grown plants using the RNeasy Plant Mini Kit (Qiagen) with on-column DNase digestion. Tissues were homogenized under liquid nitrogen with a Heidolph RZR 2020 and Precellys 24 homogenizer. RNA quality and concentration were assessed by absorbance (260/280 nm) using a TECAN Infinite M200 PRO plate reader; samples with ratios between 2.1 and 2.25 and concentrations >90 ng/µl were used.

First-strand cDNA synthesis was performed with the RevertAid Reverse Transcriptase kit (Thermo Fisher Scientific) using 300–500 ng RNA per reaction. RNA was treated with DNase I, followed by heat inactivation with EDTA and oligo(dT)$_{18}$ primer annealing. Reverse transcription was carried out at 42°C for 1 hour. cDNA was diluted 1:10 and 1:100 for downstream applications and stored at –20°C.

Quantitative PCR (qPCR) was performed using SYBR Green chemistry (Bio-Rad) in a CFX Opus 384 Real-Time PCR System (Bio-Rad) and according to protocol in (Ngigi and Bauer, 2023). Each sample was measured in technical duplicates alongside a six-

point standard curve (10^7–10^2 copies) for each gene. The PCR program included an initial denaturation (95°C, 3 min) followed by 40 cycles of 95°C denaturation (10 sec), annealing at gene-specific temperature (15 sec), and extension (5 sec). Melt curve analysis was conducted from 65°C to 95°C.

Starting quantities (SQ) were calculated from the standard curves, and data were normalized in a two-step procedure: first by correcting gene of interest SQ values using EFg normalization (to correct residual genomic DNA), and second by adjusting across samples using an EFc coefficient to ensure cross-tissue comparability. Negative normalized values were adjusted to zero. Normalized expression values were visualized as z-score heatmaps using R.

RNA Sequencing and analysis

RNA sequencing (RNA-seq) was performed by BMKGENE (Biomarker Technologies). Library preparation and sequencing were conducted according to the service provider's standard protocols. Briefly, mRNA was enriched using poly-A selection, fragmented, and reverse transcribed to generate cDNA libraries. Libraries were sequenced on the Illumina NovaSeq platform, generating 150 bp paired-end reads. Data was processed following protocol in (Mai, 2024).

RNA-seq matrices were analyzed in R (v4.3+) with limma (v3.58+) using the limma-trend workflow appropriate for TPM/CPM-like data. Expression values were log2-transformed as $\log_2(x + 0.5)$. For each comparison (line vs WT), linear models were fit with empirical Bayes moderation, and contrasts were estimated to obtain log2 fold-changes (log2FC) and BH-adjusted P values. Genes were called differentially expressed if FDR ≤ 0.05 and $|\log_2 FC| \geq 0.58$ (~1.5-fold). Direction (up/down) was assigned by the sign of log2FC; the resulting DEG_significant tables were used for enrichment and count summaries.

RNA-seq data will be made available in a public database upon acceptance.

Gene Ontology enrichment

GO Biological Process enrichment was performed with clusterProfiler (v4.10+) against the appropriate org.*.eg.db annotation. Up- and down-regulated DEG sets were tested separately. The gene universe for each test comprised all genes assessed in the corresponding comparison. Over-representation p values were computed by the

788 hypergeometric test and adjusted by Benjamini–Hochberg; terms with FDR ≤ 0.05
789 were considered significant. When indicated, redundant terms were condensed with
790 simplify() (Wang similarity; cutoff = 0.7), retaining the most significant representative

Structural Predictions and Visualization

Protein structures for bHLH104, BTS and its interaction interfaces were predicted
using AlphaFold Multimer (Evans et al., 2021; Jumper et al., 2021) to model the full-
length and interacting protein complexes. The protein sequences of bHLH104 and its
interacting partner, BTS were input into AlphaFold Multimer to generate accurate
structural models of their interactions. The models were assessed for confidence using
the predicted Local Distance Difference Test (pLDDT) and analyzed for structural
integrity.

The resulting structures were visualized and analyzed using UCSF ChimeraX
(Pettersen et al., 2004). ChimeraX was used for detailed structural analysis, including
the electrostatic surface modeling to visualize potential protein-protein interaction
interfaces. Electrostatic potentials were calculated based on the predicted protein
structures, and interaction hotspots were identified by analyzing the electrostatic
properties of the interface regions.

Yeast Two-Hybrid Assay

Yeast two-hybrid assays were performed as described by (Gietz and Schiestl, 2007)
with modifications. C-terminal fragments of bHLH104 and bHLH104^dPAA were
cloned into the Gateway-compatible yeast vectors pACT2-GW (activation domain
fusion, AD) and pGBKT7-GW (DNA-binding domain fusion, BD). Primers used are
listed in Supplementary Table 1. Full-length BTS, BTSL1, and BTSL2 constructs were
obtained from (Lichtblau et al., 2022). All protein pairs were assayed in reciprocal
combinations (AD–X with BD–Y and BD–X with AD–Y), and empty vector controls
were included.

Yeast strain AH109 was co-transformed with plasmid pairs using the lithium acetate
(LiAc) method and selected on synthetic dropout (SD) medium lacking leucine and
tryptophan (SD–Leu–Trp). Colonies were cultured overnight in liquid SD–Leu–Trp
medium, adjusted to an OD_{600} of 1.0, and spotted (10 µl) in tenfold dilution series onto
SD medium lacking leucine, tryptophan, and histidine (SD–Leu–Trp–His). To suppress

autoactivation, SD–Leu–Trp–His plates supplemented with 15 mM 3-amino-1,2,4-
triazole (3-AT) were used specifically for assays involving BD–BTSL2. Plates were
incubated at 30°C, and growth was monitored for up to 14 days.

UbiGate Assay

Ubiquitination assays were performed using the UbiGate system as described
previously (Kowarschik et al., 2018; Ortmann et al., 2023) with adaptations for
bHLH104 substrates. The coding sequences of codon optimized bHLH104 and
bHLH104^dPAA were cloned into the pGG-TP1 vector (6×HA–TwinStrepII backbone)
by GoldenGate assembly. Complete ubiquitination modules were assembled into the
pGG-pET28 destination vector and included 6×His-tagged ubiquitin (pGG-TP2), E1
enzyme (UBA1, pGG-P3), E2 conjugating enzyme (GST–UBC8, pUG-TP4), and a
truncated BTS E3 ligase (BTS-HC, pGG-TP5), using the linker plasmid pICH41800.
Primers used are listed in Supplementary Table 1.

For each E3-substrate combination, full ubiquitination constructs and substrate-minus,
E2-minus, and E3-minus control constructs were generated. Constructs were
transformed into Escherichia coli Rosetta (DE3) and expressed by induction with 1
mM IPTG and 3% ethanol at 28°C for 3 hours. Cells were harvested by centrifugation
and stored at –80°C.

For protein extraction, bacterial pellets were resuspended in phosphate-buffered
saline (PBS) to $OD_{600} = 13$, treated with lysozyme (10 mg/mL), subjected to four
freeze-thaw cycles, and digested with RNase A and DNase I (1 mg/mL each). Lysates
were mixed with 2× Laemmli sample buffer and boiled at 95°C for 10 minutes.

Proteins were separated by SDS-PAGE using TGX precast gels (Bio-Rad) and
transferred to 0.2 μm nitrocellulose membranes with the Trans-Blot Turbo system (Bio-
Rad). Membranes were blocked with 5% (w/v) non-fat dry milk in TBST and probed
with the following primary antibodies: HRP-conjugated anti-Myc (Thermo Fisher
Scientific, R951-25-HRP), anti-GST (clone 8-326, Thermo Fisher Scientific) followed
by HRP-conjugated secondary antibody (Jackson ImmunoResearch), HRP-
conjugated Strep-Tactin (Fisher Scientific), and HRP-conjugated anti-6×His (Thermo
Fisher Scientific, MA1-21315-HRP). Signals were detected using the Amersham ECL
Prime kit (Cytiva) and visualized with the ChemiDoc MP Imaging System (Bio-Rad).

Statistical Analysis and Data Visualization

All statistical analyses and data visualization were performed using RStudio (version 2025.04.1). Phenotyping data were visualized using ggplot2, where violin plots with overlaid box plots were generated to compare data distributions across genotypes and experimental conditions. Gene expression data were Z-score normalized using the scale() function in R, and gene expression heatmaps were created using the pheatmap package to visualize normalized expression levels across tissues and experimental conditions. For ICP-MS data visualization, the ggPlantMap package (Jo and Kajala, 2024) was used.

Statistical comparisons were performed using two-way ANOVA followed by Tukey's Honest Significant Difference (HSD) test for multiple comparisons. A p-value threshold of 0.05 was considered significant.

Acknowledgements

We thank Elke Wieneke for excellent technical assistance. The authors thank the Biocenter-MS Platform, Sabine Ambrosius and Sabine Metzger, University of Cologne, for conducting and providing support in the ICP-MS measurements. The authors thank Center for Advanced Imaging (CAi) at Heinrich Heine University for providing advanced imaging facility. DB and VV were members of the international graduate school iGRAD-Plant, Düsseldorf (the DFG International Research Training group 2466). Chat GPT 4.0 has been used to edit and shorten text, and correct and improve language.

Author contributions

DB and PB planned and designed the research. DB, SW, MK, and VV performed the experiments and analyzed the data. DB wrote the manuscript. JLE generated and provided materials. DB and PB supervised the research. PB acquired funding and agreed to serve as the author responsible for contact and ensures communication. All authors read, commented and approved the submitted version.

Funding

This project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy – EXC-2048/1 – project ID 390686111. This project received funding through the DFG International Research

Training group 2466 (GRK F020512056 (NextPlant)). Funded by the Deutsche Forschungsgemeinschaft (DFG) – TRR 341/1 – 456082119.

References

Abuzeineh A, Vert G, Zelazny E (2022) Birth, life and death of the Arabidopsis IRT1 iron transporter: the role of close friends and foes. Planta **256**: 112

Aparicio F, Pallás V (2017) The coat protein of *Alfalfa mosaic virus* interacts and interferes with the transcriptional activity of the bHLH transcription factor ILR3 promoting salicylic acid-dependent defence signalling response. Mol Plant Pathol **18**: 173–186

Arend D, Lange M, Pape J-M, Weigelt-Fischer K, Arana-Ceballos F, Mücke I, Klukas C, Altmann T, Scholz U, Junker A (2016) Quantitative monitoring of Arabidopsis thaliana growth and development using high-throughput plant phenotyping. Sci Data **3**: 160055

Briat J-F, Curie C, Gaymard F (2007) Iron utilization and metabolism in plants. Curr Opin Plant Biol **10**: 276–282

Cailliatte R, Schikora A, Briat J-F, Mari S, Curie C (2010) High-Affinity Manganese Uptake by the Metal Transporter NRAMP1 Is Essential for *Arabidopsis* Growth in Low Manganese Conditions. Plant Cell **22**: 904–917

Chen Y-Y, Wang Y, Shin L-J, Wu J-F, Shanmugam V, Tsednee M, Lo J-C, Chen C-C, Wu S-H, Yeh K-C (2013) Iron Is Involved in the Maintenance of Circadian Period Length in Arabidopsis. Plant Physiol **161**: 1409–1420

Cheng H, Qin L, Lee S, Fu X, Richards DE, Cao D, Luo D, Harberd NP, Peng J (2004) Gibberellin regulates *Arabidopsis* floral development via suppression of DELLA protein function. Development **131**: 1055–1064

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J **16**: 735–743

Connolly EL, Fett JP, Guerinot ML (2002) Expression of the IRT1 Metal Transporter Is Controlled by Metals at the Levels of Transcript and Protein Accumulation. Plant Cell **14**: 1347–1357

Dubeaux G, Neveu J, Zelazny E, Vert G (2018) Metal Sensing by the IRT1 Transporter-Receptor Orchestrates Its Own Degradation and Plant Metal Nutrition. Mol Cell **69**: 953-964.e5

Dutta A, Choudhary P, Caruana J, Raina R (2017) JMJ 27, an Arabidopsis H3K9 histone demethylase,
modulates defense against *Pseudomonas syringae* and flowering time. Plant J **91**: 1015–1028

Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J
(2021) Protein complex prediction with AlphaFold-Multimer. biorxiv 2021.10. 04.463034

Gelvin SB (2003) *Agrobacterium* -Mediated Plant Transformation: the Biology behind the "Gene-
Jockeying" Tool. Microbiol Mol Biol Rev **67**: 16–37

Gietz RD, Schiestl RH (2007) Large-scale high-efficiency yeast transformation using the LiAc/SS carrier
DNA/PEG method. Nat Protoc **2**: 38–41

Gratz R, Manishankar P, Ivanov R, Köster P, Mohr I, Trofimov K, Steinhorst L, Meiser J, Meiser J, Mai
H-J, et al (2019) CIPK11-Dependent Phosphorylation Modulates FIT Activity to Promote
Arabidopsis Iron Acquisition in Response to Calcium Signaling. Dev Cell. doi:
10.1016/j.devcel.2019.01.006

Hindt MN, Akmakjian GZ, Pivarski KL, Punshon T, Baxter I, Salt DE, Guerinot ML (2017) BRUTUS and
its paralogs, BTS LIKE1 and BTS LIKE2, encode important negative regulators of the iron
deficiency response in Arabidopsis thaliana. Metallomics. doi: 10.1039/c7mt00152e

Hong S-Y, Seo PJ, Ryu JY, Cho S-H, Woo J-C, Park C-M (2013) A Competitive Peptide Inhibitor KIDARI
Negatively Regulates HFR1 by Forming Nonfunctional Heterodimers in Arabidopsis
Photomorphogenesis. Mol Cells **35**: 25–31

Hüther P, Schandry N, Jandrasits K, Bezrukov I, Becker C (2020) ARADEEPOPSIS, an Automated
Workflow for Top-View Plant Phenomics using Semantic Segmentation of Leaf States. Plant
Cell **32**: 3674–3688

Jeong J, Guerinot ML (2009) Homing in on iron homeostasis in plants. Trends Plant Sci **14**: 280–285

Jiang M, Niu Y, Wen G, Zhao C, Gao Q, Li G (2024) Evolutionary studies of the basic helix–loop–helix (
BHLH) IVC gene family in plants and the role of *ATILR3* in *Arabidopsis* response to ABA
stress. Physiol Plant **176**: e14128

Jo L, Kajala K (2024) ggPlantmap: an open-source R package for the creation of informative and
quantitative ggplot maps derived from plant images. J Exp Bot **75**: 5366–5376

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,
Žídek A, Potapenko A (2021) Highly accurate protein structure prediction with AlphaFold.
nature **596**: 583–589

Knopf MC, Bauer P (2025) Affordable Non-Invasive Machine-Aided Phenotyping Identifies
Phenotypic Variation to Soil Stress Across the *ARABIDOPSIS THALIANA* Life Cycle. Physiol Plant
177: e70427

Kowarschik K, Hoehenwarter W, Marillonnet S, Trujillo M (2018) UbiGate: a synthetic biology
toolbox to analyse ubiquitination. New Phytol **217**: 1749–1763

Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU, Forner J (2013) GreenGate-a novel,
versatile, and efficient cloning system for plant transgenesis. PloS One **8**: e83043

Lešková A, Giehl RFH, Hartmann A, Fargašová A, Von Wirén N (2017) Heavy Metals Induce Iron
Deficiency Responses at Different Hierarchic and Regulatory Levels. Plant Physiol **174**: 1648–
1668

Li X, Zhang HM, Ai Q, Liang G, Yu D (2016) Two bHLH Transcription Factors, bHLH34 and bHLH104,
Regulate Iron Homeostasis in Arabidopsis thaliana. Plant Physiol. doi: 10.1104/pp.15.01827

Li Y, Li Y, Li Y, Li Y, Lu C, Lu C, Li C, Lei RH, Pu MN, Zhao JH, et al (2021) IRON MAN interacts with
BRUTUS to maintain iron homeostasis in Arabidopsis. Proc Natl Acad Sci U S A. doi:
10.1073/pnas.2109063118

Liang G, Zhang H, Li X, Ai Q, Yu D (2017) bHLH transcription factor bHLH115 regulates iron
homeostasis in Arabidopsis thaliana. J Exp Bot **68**: 1743–1755

Lichtblau DM, Schwarz B, Baby D, Endres C, Sieberg C, Bauer P (2022) The Iron Deficiency-Regulated
Small Protein Effector FEP3/IRON MAN1 Modulates Interaction of BRUTUS-LIKE1 With bHLH
Subgroup IVc and POPEYE Transcription Factors. Front Plant Sci **13**: 930049

Long TA, Tsukagoshi H, Busch W, Lahner B, Salt DE, Benfey PN (2010) The bHLH Transcription Factor
POPEYE Regulates Response to Iron Deficiency in Arabidopsis Roots. Plant Cell. doi:
10.1105/tpc.110.074096

Mai H-J (2024) PlugNSeq: An Easy, Rapid, and Streamlined mRNA-Seq Data Analysis Pipeline
Empowering Insightful Exploration with Well-Annotated Organisms, Requiring Minimal
Bioinformatic Expertise v1. doi: 10.17504/protocols.io.4r3l2qjo3l1y/v1

Marschner H (2012) Marschner's mineral nutrition of higher plants. Academic press

Maruoka T, Gan E-S, Otsuka N, Shirakawa M, Ito T (2022) Histone Demethylases JMJ30 and JMJ32
Modulate the Speed of Vernalization Through the Activation of FLOWERING LOCUS C in
Arabidopsis thaliana. Front Plant Sci **13**: 837831

Melnikov S, Mailliot J, Rigger L, Neuner S, Shin B, Yusupova G, Dever TE, Micura R, Yusupov M
(2016) Molecular insights into protein synthesis with proline residues. EMBO Rep **17**: 1776–
1784

Naranjo-Arcos MA, Maurer F, Meiser J, Meiser J, Pateyron S, Fink-Straube C, Fink-Straube C, Bauer
P (2017) Dissection of iron signaling and iron accumulation by overexpression of subgroup Ib
bHLH039 protein. Sci Rep. doi: 10.1038/s41598-017-11171-7

Ngigi M, Khan M, Remus R, Gupta SK, Bauer P (2025) Age-dependent differential iron deficiency
responses of rosette leaves during reproductive stages in *Arabidopsis thaliana*. J Exp Bot
eraf207

Ngigi MN, Bauer P (2023) High-Throughput Plant Gene Expression Analysis by 384-Format Reverse
Transcription-Quantitative PCR for Investigating Plant Iron Homeostasis. *In* J Jeong, ed, Plant
Iron Homeost. Methods Protoc. Springer US, New York, NY, pp 1–22

Nguyen NT, Khan MA, Castro–Guerrero NA, Chia J-C, Vatamaniuk OK, Mari S, Jurisson SS, Mendoza-
Cozatl DG (2022) Iron Availability within the Leaf Vasculature Determines the Magnitude of
Iron Deficiency Responses in Source and Sink Tissues in Arabidopsis. Plant Cell Physiol **63**:
829–841

Nguyen NT, McInturf SA, Mendoza-Cózatl DG (2016) Hydroponics: A Versatile System to Study
Nutrient Allocation and Plant Responses to Nutrient Availability and Exposure to Toxic
Elements. J Vis Exp 54317

Ortmann S, Marx J, Lampe C, Handrick V, Ehnert T-M, Zinecker S, Reimers M, Bonas U, Erickson JL
(2023) A conserved microtubule-binding region in Xanthomonas XopL is indispensable for
induced plant cell death reactions. PLoS Pathog **19**: e1011263

Panda SK, Yamamoto YY (2019) Redox homeostasis in plants: from signalling to stress tolerance.
Springer

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE (2004) UCSF

Chimera—a visualization system for exploratory research and analysis. J Comput Chem **25**:

1605–1612

Pottier M, Dumont J, Masclaux-Daubresse C, Thomine S (2018) Autophagy is essential for optimal

translocation of iron to seeds in Arabidopsis. J Exp Bot. doi: 10.1093/jxb/ery388

Pullin J, Rodríguez-Celma J, Franceschetti M, Mundy JEA, Svistunenko DA, Bradley JM, Le Brun NE,

Balk J (2025) Iron-sensing and redox properties of the hemerythrin-like domains of

Arabidopsis BRUTUS and BRUTUS-LIKE2 proteins. Nat Commun **16**: 3865

Quintana, J., Bernal, M., Scholle, M., Holländer-Czytko, H., Nguyen, N. T., Piotrowski, M., Mendoza-

Cózatl, D. G., Haydon, M. J. & Krämer, U. 2022. Root-to-shoot iron partitioning in Arabidopsis

requires IRON-REGULATED TRANSPORTER1 (IRT1) protein but not its iron(II) transport

function. *Plant J,* **109,** 992-1013.

Rampey RA, Woodward AW, Hobbs BN, Tierney MP, Lahner B, Salt DE, Bartel B (2006) An

Arabidopsis Basic Helix-Loop-Helix Leucine Zipper Protein Modulates Metal Homeostasis and

Auxin Conjugate Responsiveness. Genetics **174**: 1841–1857

Rodríguez-Celma J, Connorton JM, Kruse I, Green RT, Franceschetti M, Chen Y-T, Cui Y, Ling H-Q, Yeh

K-C, Balk J (2019) Arabidopsis BRUTUS-LIKE E3 ligases negatively regulate iron uptake by

targeting transcription factor FIT for recycling. Proc Natl Acad Sci U S A. doi:

10.1073/pnas.1907971116

Salomé PA, Oliva M, Weigel D, Krämer U (2012) Circadian clock adjustment to plant iron status

depends on chloroplast and phytochrome function. EMBO J **32**: 511–523

Samira R, Li B, Kliebenstein D, Li C, Davis E, Gillikin JW, Long TA (2018) The bHLH transcription factor

ILR3 modulates multiple stress responses in Arabidopsis. Plant Mol Biol **97**: 297–309

Schwarz B, Bauer P (2020) FIT, a regulatory hub for iron deficiency and stress signaling in roots, and

FIT-dependent and -independent gene signatures. J Exp Bot **71**: 1694–1705

Selote D, Selote D, Samira R, Matthiadis A, Gillikin JW, Long TA (2015) Iron-Binding E3 Ligase

Mediates Iron Response in Plants by Targeting Basic Helix-Loop-Helix Transcription Factors.

Plant Physiol. doi: 10.1104/pp.114.250837

Sharma R, Yeh K (2020) The dual benefit of a dominant mutation in Arabidopsis *IRON DEFICIENCY*
TOLERANT1 for iron biofortification and heavy metal phytoremediation. Plant Biotechnol J
18: 1200–1210

Shen C, Yang Y, Liu K, Zhang L, Guo H, Sun T, Wang H (2016) Involvement of endogenous salicylic
acid in iron-deficiency responses in Arabidopsis. J Exp Bot **67**: 4179–4193

Vert G, Grotz N, Dédaldéchamp F, Gaymard F, Guerinot ML, Briat J-F, Curie C (2002) IRT1, an
Arabidopsis Transporter Essential for Iron Uptake from the Soil and for Plant Growth. Plant
Cell **14**: 1223–1233

Wang F, Zhu D, Huang X, Li S, Gong Y, Yao Q, Fu X, Fan L-M, Deng XW (2009) Biochemical Insights on
Degradation of *Arabidopsis* DELLA Proteins Gained From a Cell-Free Assay System. Plant Cell
21: 2378–2390

Waters BM, Grusak MA (2008) Whole-plant mineral partitioning throughout the life cycle in
Arabidopsis thaliana ecotypes Columbia, Landsberg *erecta* , Cape Verde Islands, and the
mutant line *ysl1ysl3*. New Phytol **177**: 389–405

Wild M, Davière J-M, Regnault T, Sakvarelidze-Achard L, Carrera E, Lopez Diaz I, Cayrel A, Dubeaux
G, Vert G, Achard P (2016) Tissue-Specific Regulation of Gibberellin Signaling Fine-Tunes
Arabidopsis Iron-Deficiency Responses. Dev Cell **37**: 190–200

Xing Y, Xu N, Bhandari DD, Lapin D, Sun X, Luo X, Wang Y, Cao J, Wang H, Coaker G, et al (2021)
Bacterial effector targeting of a plant iron sensor facilitates iron acquisition and pathogen
colonization. Plant Cell **33**: 2015–2031

Yi Y, Guerinot ML (1996) Genetic evidence that induction of root Fe(III) chelate reductase activity is
necessary for iron uptake under iron deficiency[†]. Plant J **10**: 835–844

Zhang J, Zhang J, Liu B, Li M, Feng D, Jin H, Wang P, Liu J, Liu J, Liu J, et al (2015) The bHLH
Transcription Factor bHLH104 Interacts with IAA-LEUCINE RESISTANT3 and Modulates Iron
Homeostasis in Arabidopsis. Plant Cell. doi: 10.1105/tpc.114.132704

Zhao Q, Ren Y-R, Wang Q-J, Wang X-F, You C-X, Hao Y-J (2016) Ubiquitination-Related MdBT Scaffold
Proteins Target a bHLH Transcription Factor for Iron Homeostasis. Plant Physiol **172**: 1973–
1988

1049

Figure 1

PAA sequence in bHLH104 enables E3 ligase interaction and degradation

Figure 1. Conceptual model of bHLH104 regulation by its conserved PAA sequence.
In the full-length bHLH104 protein, the PAA sequence forms the interaction interface with an E3 ligase, likely enabling ubiquitination and proteasomal degradation of bHLH104.

Figure 2

Figure 2. Transgenic expression of bHLH104 lacking the PAA sequence leads to growth alterations and physiological changes in Arabidopsis.

((A) Schematic representation of constructs used to generate transgenic lines expressing either full-length bHLH104 (pb104::b104) or a C-terminal deletion variant lacking the conserved three–amino acid PAA sequence (pb104::b104^dPAA). Both constructs are driven by the native bHLH104 promoter (2,069 bp) and fused at the N-terminus to an mTurquoise2 fluorescent tag. The deleted PAA sequence in b104^dPAA is indicated by a red cross at the C-terminus. Scale bar = 20 bp.

(B) Representative images of 10-day-old seedlings grown on +Fe medium (scale bar = 2 cm).

(C) Primary root length of 12-day-old seedlings grown vertically on Hoagland medium supplemented with Fe.

(D) Fe reductase activity in roots of 10-day-old seedlings grown under +Fe. Activity was quantified spectrophotometrically as Fe(II)–ferrozine complex formation and normalized to total root length.

(E) Representative images of four-week-old soil-grown plants (scale bar = 2 cm). Shown are Col-0 (WT), two independent pb104::b104 lines (#1, #2), and four independent pb104::b104^dPAA lines (#1–4).

(F–G) Quantitative analysis of aboveground phenotypes using the Phenospex PlantEye system: (F) 3D leaf area (mm²); (G) NDVI (Normalized Difference Vegetation Index)

(H) Flowering time, recorded as days to bolting.

(I) Seed number per plant, estimated from total seed weight based on mean weight per 100 seeds.

Data represent means ± SD (n = 7–12 biological replicates per line). Statistical analysis was performed using two-way ANOVA followed by Tukey's post hoc test. Different letters indicate statistically significant differences at $p < 0.05$.

Figure 3

A

shoot

root

Harvest

0 → 10 DAS

Grown in **+/- Fe** media

B

L1 - WT

L2 - *pb104::b104 #1*

L3 - *pb104::b104 #2*

L4 - *pb104::b104^dPAA#2*

L5 - *pb104::b104^dPAA#3*

L6 - *pUBQ10::b104#1*

L7 - *pUBQ10::b104#2*

C

+ Fe

- Fe

L1   L2   L3   L4   L5   L6   L7

Fe content
(µg/g D. W.)

6000

D

Figure 3. Transgenic lines expressing bHLH104^dPAA exhibit increased iron accumulation and altered iron homeostasis gene expression in seedling stage.

(A) Schematic representation of the experimental design. *Arabidopsis* seedlings were grown for 10 days after sowing (DAS) under either iron-sufficient (+Fe) or iron-deficient (–Fe) conditions. Shoot and root tissues were collected separately for ICP-MS and gene expression analyses. (B) Genotypes used in this study. (C) Iron content in shoot and root tissues of seedlings grown under +Fe and –Fe conditions, measured by ICP-MS. Data represent the mean of three biological replicates (n = 3). Heat map intensity reflects iron concentration (μg/g dry weight). Color shading was rendered using the R package ggplantmap, with darker blue indicating higher Fe accumulation. Notably*, pb104::b104^dPAA* lines show increased Fe accumulation in roots and shoots under +Fe conditions.(D) Clustered heatmaps showing the expression profiles (RNA sequencing) of selected Fe homeostasis genes. Heatmaps were generated in RStudio version 4.4.1 using calculated Z-scores of normalized gene expression values across tissues for both +Fe and –Fe conditions. Purple indicates low expression and yellow indicates high expression. Asterisks indicate genotypes that differ significantly from WT within the same tissue and treatment condition ($p < 0.05$), as determined by two-way ANOVA and Tukey's post hoc test (n = 3 for each group).

Figure 4

A

L1 - WT
L2 - *pb104::b104 #1*
L3 - *pb104::b104 #2*
L4 - *pb104::b104^dPAA#2*
L5 - *pb104::b104^dPAA#3*
L6 - *pUBQ10::b104#1*
L7 - *pUBQ10::b104#2*

B

L1 L2 L3 L4 L5 L6 L7

C

3-day +/- Fe treatment

Harvest

0   30 DAS

Grown in **+ Fe** media

D

Young leaves

old leaves

root

E

+ Fe

- Fe

L1   L2   L3   L5   L6   L7

Fe content (µg/g D. W.)

4000

F

young leaves		old leaves		root	
+Fe	-Fe	+Fe	-Fe	+Fe	-Fe

BHLH34
BHLH39
PYE
BHLH100
IMA1
BTS
FER1
bHLH104
bHLH115
bHLH11
BTSL2
FIT
URI
ILR3
BTSL1
FRO2
IRT1

L1 L2 L5   L1 L2 L5   L1 L2 L5   L1 L2 L5   L1 L2 L5   L1 L2 L5

Z-score

3
2
1
0
-1

Figure 4. Elevated shoot iron accumulation and altered Fe deficiency responses in reproductive-stage bHLH104^dPAA transgenic lines.

(A) Plant lines used In this experiment

(B) Representative images of 30-days-old Arabidopsis plants grown in hydroponic system (scale bar = 5 cm). Among grown plant lines, *pb104::b104^dPAA #2* (L4) grew poorly and was not used in ICPMS and gene expression analysis.

(C) Experimental design for the dataset shown. Arabidopsis plants were grown hydroponically under iron-sufficient (+Fe) conditions for 27 days, followed by a 3-day treatment with either +Fe or –Fe media.

(D) Schematic representation of tissue separation used for both ICP-MS and RT-qPCR analysis. Young and old leaves were separated from the roots to examine tissue-specific iron accumulation and gene expression. The same tissue layout is used to represent heatmaps in the ICP-MS dataset.

(E) Iron content in young leaves, old leaves, and roots under +Fe and –Fe conditions, measured by ICP-MS. Data represent mean values from three biological replicates (n = 3). Heat map intensity reflects iron concentration (μg/g dry weight). Color shading was generated using the R package ggplantmap, with deeper blue indicating higher Fe content. Asterisks indicate genotypes that differ significantly from WT within the same tissue and treatment condition ($p < 0.05$), as determined by two-way ANOVA and Tukey's post hoc test (n = 3 for each group).

(F) Clustered heatmaps showing the expression profiles (RNA sequencing) of selected Fe homeostasis genes in young leaves, old leaves, and roots under both +Fe and –Fe conditions. Data represent the mean of three biological replicates (n = 3). Z-scores were calculated from normalized RNA expression values and visualized in RStudio version 4.4.1. Yellow indicates high expression; purple indicates low expression.

Figure 5

A

L1 - WT
L2 - *pb104::b104 #1*
L3 - *pb104::b104 #2*
L4 - *pb104::b104^dPAA#1*
L5 - *pb104::b104^dPAA#2*
L6 - *pb104::b104^dPAA#3*
L7 - *pb104::b104^dPAA#4*
L8 - *pUBQ10::b104#1*
L9 - *pUBQ10::b104#2*

B

C

D

E

Figure 5. The PAA short sequence mediates bHLH104 protein stability and affects its abundance in Arabidopsis.

(A) Genotypes and transgenic lines used in this experiment.

(B) Immunoblot analysis of mTurquoise2-tagged bHLH104 and b104^dPAA in different transgenic lines treated with either DMSO (control) or 50 µM MG132. Total protein and anti-mTurquoise2 blots are shown. Samples are labeled from #L1 to #L9. The transgenic lines exhibiting increased protein abundance correspond to those that displayed altered growth phenotypes in Figure 1, suggesting a link between protein stability and the observed developmental responses.

(C) Relative protein levels of mTurquoise-tagged bHLH104 were determined by Western blot. Band intensities were normalized to total protein, and fold-change was calculated as the ratio of MG132-treated to DMSO-treated samples. Bars represent mean ± SEM; individual data points are shown.

(D) Cell-free degradation assay showing in vitro degradation of His-tagged bHLH104 protein when incubated with protein extracts from 10-day-old Col-0 (WT) *Arabidopsis thaliana* seedlings. His-bHLH104 was expressed in *E. coli* and remained stable on its own, but degraded upon incubation with the plant extracts, indicating proteasome-mediated degradation. Total protein from whole seedlings was extracted using standard protein extraction procedures. Samples were collected at the indicated time points and analyzed by immunoblotting with anti-His and anti-Actin antibodies. MG132 treatment stabilizes bHLH104, indicating proteasome-dependent degradation.

(E) Confocal microscopy of mTurquoise2 fluorescence in roots of transgenic Arabidopsis seedlings expressing full-length bHLH104 (left) or b104^dPAA (right). Images show two representative seedlings (S1 and S2) per genotype. Arabidopsis thaliana seedlings were at the germination stage, 5 days old, corresponding to 2 days after stratification, and grown on +Fe medium. TM, transmitted light; Overlay, merged mTurq2 and TM channels. Scale bar = 25 µm; excitation at 458 nm; emission collected from 493–613 nm.

Figure 6

Figure 6. The PAA short sequence is required for bHLH104 interaction with BTS and its ubiquitination.

(A) Structural prediction of the bHLH104–BTS complex using AlphaFold2 Multimer (Evans et al., 2021; Jumper et al., 2021). Left: Chain-colored model showing BTS (rose color) and bHLH104 (bronze color). Right: Per-residue confidence model with pLDDT scores color-coded as very high (pLDDT > 90), confident (90 > pLDDT > 70), low (70 > pLDDT > 50), and very low (pLDDT < 50).

(B) Electrostatic surface potential modeling (Pettersen et al., 2004) reveals a negatively charged region near the proline–alanine–alanine (PAA) sequence in bHLH104 that complements a positively charged surface in the leucine–cysteine–arginine (LCR) domain of BTS. The electrostatic scale ranges from –10 (purple) to +10 (cyan) kT/e.

(C) Schematic representation of full-length bHLH104, bHLH104-C and bHLH104-C-dPAA used for Y2H and UbiGate assays. C-terminal parts lack the N-terminus and the DNA-binding basic Helix Loop Helix (bHLH) domain. Numbers indicate amino acid positions.

(D)Yeast two-hybrid assays showing protein–protein interactions between bHLH104-C or bHLH104-C^dPAA and BTS. bHLH104 homodimerization serves as a control. Full length bHLH104 shows autoactivation in Y2H. Hence bHLH104-C and bHLH104-C-dPAA were used.

(E) Schematic overview of the ubiquitin/ 26s proteosome pathway. Each component is expressed in *E.coli* for UbiGate assay. Recombinant bHLH104 or bHLH104^dPAA proteins (Strep-tagged) were used as substrates. BTS-HC served as the E3 ligase (Myc-tagged), in combination with UBC8 (E2) (GST-tagged) and an E1 activating enzyme.

(F) Ubiquitination assay showing that bHLH104 undergoes polyubiquitination in the presence of E1, E2, E3, and ubiquitin (Ub), whereas bHLH104^dPAA fails to be ubiquitinated. Immunoblots with anti-GST (E2), anti-Myc (E3), anti-Strep (substrate), and anti-UBQ (ubiquitin) confirm protein expression and substrate modification

Figure 7

Figure 7. Working model of bHLH104 regulation through the C-terminal PAA sequence and its impact on Fe homeostasis.
The C-terminal PAA sequence in bHLH104 mediates interaction with BTS-type E3 ligases, targeting the protein for proteasomal degradation. Deletion of the PAA motif (b104^dPAA) disrupts this interaction, leading to protein stabilization. The increased abundance of stabilized b104^dPAA enhances Fe acquisition responses in *Arabidopsis*.

Supplementary Figure S1

WT   *pb104::b104*   *pb104::b104^dPAA*   *pUBQ10::b104*

Supplementary Figure S1. Preliminary phenotypic screen of independent transgenic *Arabidopsis* lines expressing bHLH104 or bHLH104^dPAA. 26 days old plants of Col-0 wild type (WT), *pb104::b104*, *pb104::b104^dPAA*, and *pUBQ10::b104* lines were grown under same conditions. Lines selected for further analysis in the main study are indicated: *pb104::b104 lines* #1 and #2, and *pb104::b104^dPAA lines* #1–#4 and *pUBQ10::b104* lines #1 and #2, scale bar = 2 cm.

Supplementary Figure S2

Supplementary Figure S2. Response of bHLH104^dPAA transgenic lines to iron-limiting conditions in alkaline calcareous soil (ACS).
Arabidopsis thaliana transgenic lines were grown in either control soil (pH 5.9) or alkaline calcareous soil (ACS, pH 8.1).(A) Photos of four-week-old plants grown in control and ACS soil conditions.(B) List of plant lines used in this experiment, numbered from L1 to L7. scale bar = 2 cm (C) 3D leaf area measurements: Plants were evaluated for leaf expansion and overall growth in both conditions.(D) NDVI (Normalized Difference Vegetation Index): NDVI values were measured to assess photosynthetic activity in response to ACS conditions.(E) Flowering time: Flowering time was compared between control and ACS conditions, with letters indicating statistical differences. Statistical significance was assessed by two-way ANOVA followed by Tukey's HSD test ($p < 0.05$). N = 8 plants per line.

Supplementary Figure S3

A   *pUBQ10::b104*

B

WT   *pUBQ10::b104*
       #1   #2

C   Root length (in cm)

D

WT   *pUBQ10::b104*
       #1   #2

E   3D Leaf Area (in mm^2)

F   NDVI

WT   *pUBQ10::b104*
       #1   #2

G   Flowering time (in days)

WT   *pUBQ10::b104*
       #1   #2

Supplmentary Figure S3. Constitutive expression of bHLH104 did not alter growth and physiological traits in Arabidopsis.

 (A) Schematic representation of the construct used to generate transgenic lines expressing full-length bHLH104 (*pUBQ10::bHLH104*). The construct is driven by the ubiquitin 10 (UBQ10) promoter and is fused at the N-terminus to the mTurquoise2 fluorescent tag. Scale bar = 20 bp.

 (B-C) Photos and root length quantification data of 12-day-old seedlings grown vertically on Hoagland medium supplemented with iron (+Fe). Scale bar =2 cm

 (D) Representative images of 4-week-old Arabidopsis plants grown in soil. Shown are Col-0 (WT) and two independent *pUBQ10::b104* lines (#1, #2). Scale bar = 2 cm

 (E–F) Quantitative analysis of aboveground phenotypes acquired using the Phenospex PlantEye system: (E) 3D leaf area (mm²); (F) NDVI (Normalized Difference Vegetation Index).

 (G) Flowering time, measured manually as the number of days to bolting.

 Data represent means ± SD (n = 7–12 biological replicates per line). Statistical analysis was performed using two-way ANOVA followed by Tukey's post hoc test. Different letters indicate statistically significant differences at $p < 0.05$.

Supplementary Figure S4. Elemental content analysis of Mn, Zn, and Cu in 10-day old Arabidopsis seedlings.
Heatmap showing the distribution of Mn, Zn, and Cu content (in μg/g dry weight) across different *Arabidopsis* seedlings, including wild-type (WT), pb104::b104 (lines #1, #2), *pb104::b104^dPAA* (lines #2, #3), *pUBQ10::b104* (lines #1, #2). The color scale indicates the range of elemental concentrations from low (light blue) to high (dark blue). Seedlings were grown in Hoagland media with or without 50 mM Fe supplementation. Data were collected from 10-day-old seedlings, with elemental content measurements obtained via ICP-MS analysis. Three biological replicates per line were used. The plot was generated using the ggPlantMap package in RStudio. Asterisks indicate genotypes that differ significantly from WT within the same tissue and treatment condition (p < 0.05), as determined by two-way ANOVA and Tukey's post hoc test (n = 3 for each group).

Supplementary Figure S5

A

B

Supplementary Figure S5: Differential expression and GO enrichment across seedling tissues.

(A) Counts of differentially expressed genes (DEGs) for each line vs WT in root and shoot under Fe(+) and Fe(−). Orange = up-regulated; blue = down-regulated; numbers above/below bars indicate counts. DEGs were defined at FDR ≤ 0.05 and |log2FC| ≥ 0.58. X-axis labels show the abbreviated line IDs.

(B) GO Biological Process enrichment for the up-regulated (left panel) and down-regulated (right panel) DEG sets. Columns correspond to tissue/Fe combinations (Root Fe(+), Root Fe(−), Shoot Fe(+), Shoot Fe(−)). Each dot represents a significant term (BH FDR ≤ 0.05); dot size = number of genes in the term, fill = −log10(FDR), and ring color = GeneRatio. Terms plotted are the top-ranked per panel; the gene universe was all genes tested in the corresponding comparison.

Supplementary Figure S6

A

B

L1 - WT
L2 - *pb104::b104 #1*
L3 - *pb104::b104 #2*
L4 - *pb104::b104^dPAA#2*
L5 - *pb104::b104^dPAA#3*
L6 - *pUBQ10::b104#1*
L7 - *pUBQ10::b104#2*

C

Supplementary Figure S6. Targeted gene expression profiling in seedlings.
(A) Schematic representation of the experimental design. Arabidopsis seedlings were grown for 10 days after sowing (DAS) under either iron-sufficient (+Fe) or iron-deficient (–Fe) conditions. Shoot and root tissues were collected separately for RT-qPCR analyses. (B) Genotypes used in this study. (C) Clustered heatmaps showing the expression profiles of selected Fe homeostasis genes. Heatmaps were generated in RStudio version 4.4.1 using calculated Z-scores of normalized gene expression values across tissues for both +Fe and –Fe conditions. Purple indicates low expression and yellow indicates high expression. Asterisks indicate genotypes that differ significantly from WT within the same tissue and treatment condition ($p < 0.05$), as determined by two-way ANOVA and Tukey's post hoc test (n = 3 for each group).

Supplementary Figure S7

Mn content
(µg/g D. W.)

0   1250   2500

Zn content
(µg/g D. W.)

0   300   600

Cu content
(µg/g D. W.)

0   600   1200

WT

pb104::b104
#1

pb104::b104
#2

pb104::b104^dPAA
#3

pUBQ10::b104
#1

pUBQ10::b104
#2

+Fe   -Fe   +Fe   -Fe   +Fe   -Fe   Treatment

Supplementary Figure S7. Elemental content analysis of Mn, Zn, and Cu in mature *Arabidopsis thaliana*.
Heatmap showing the distribution of iron (Fe) content (in µg/g dry weight) in young leaves, old leaves, and roots of *Arabidopsis* transgenic lines, including wild-type (WT), *pb104::b104* (lines #1, #2), *pb104::b104^dPAA* (line #3), *pUBQ10::b104* (lines #1, #2). Plants were grown hydroponically under iron-sufficient (+Fe) conditions for 27 days, followed by a 3-day treatment with either +Fe or –Fe media. The color scale indicates the iron content, with darker blue shades representing higher iron concentrations. Data represent the mean values from three biological replicates (n = 3). Heatmap intensity reflects iron concentration (µg/g dry weight) measured by ICP-MS. The plot was generated using the ggPlantMap package in RStudio. Asterisks indicate genotypes that differ significantly from WT within the same tissue and treatment condition (p < 0.05), as determined by two-way ANOVA and Tukey's post hoc test (n = 3 for each group).

Supplementary Figure S8

A

B

Supplementary Figure S8: Differential expression and GO enrichment across reproductive stage tissues

(A) Numbers of differentially expressed genes (DEGs) for each line vs WT in root (Root), young leaves (YL), and old leaves (OL) under Fe(+) and Fe(−). Orange bars show up-regulated DEGs; blue bars show down-regulated DEGs; values above/below bars are counts. DEGs were defined at FDR ≤ 0.05 and |log2FC| ≥ 0.58.

(B) GO Biological Process enrichment for the up-regulated (left panel) and down-regulated (right panel) DEG sets. Columns correspond to tissue/Fe combinations (Root, YL, OL × Fe(+), Fe(−)). Each dot represents a significant term (BH FDR ≤ 0.05); dot size = number of genes in the term, fill = −log10(FDR), and ring color = GeneRatio. Terms shown are the top-ranked per panel; the gene universe comprised all genes tested in the corresponding comparison. Abbreviations: YL, young leaves; OL, old leaves.

A

L1 - WT
L2 - *pb104::b104 #1*   L5 - *pb104::b104^dPAA#3*
L3 - *pb104::b104 #2*   L6 - *pUBQ10::b104#1*
L4 - *pb104::b104^dPAA#2*   L7 - *pUBQ10::b104#2*

B

3-day +/- Fe
treatment

Harvest

0   30 DAS

Grown in **+ Fe** media

C

Young
leaves

old leaves

root

D

Supplementary Figure S9. Targeted gene expression profiling in reproductive-stage plants
(A) Plant lines used In this experiment
(B) Experimental design for the dataset shown. *Arabidopsis* plants were grown hydroponically under iron-sufficient (+Fe) conditions for 27 days, followed by a 3-day treatment with either +Fe or –Fe media.
(C) Schematic representation of tissue separation used for RT-qPCR analysis. Young and old leaves were separated from the roots to examine tissue-specific iron accumulation and gene expression.
(D) Clustered heatmaps showing the expression profiles of selected Fe homeostasis genes in young leaves, old leaves, and roots under both +Fe and –Fe conditions. Data represent the mean of three biological replicates (n = 3). Z-scores were calculated from normalized RT-qPCR expression values and visualized in RStudio version 4.4.1. Yellow indicates high expression; purple indicates low expression.

Supplementary Figure S10

Supplementary figure S10. Tissue-specific metal accumulation in dPAA line #2 grown under iron-deficient conditions.

(A) Experimental setup: Col-0 and *pb104::b104^dPAA line #2* seedlings were grown hydroponically in Fe-deficient (–Fe) media for 27 days, followed by a 3-day treatment with either +Fe or –Fe media. Tissue collection occurred at 30 days after sowing (DAS).(B) Tissue layout: Young leaves, old leaves, and roots were separated to examine tissue-specific metal accumulation (Fe, Mn, Zn, Cu).(C) Heatmap of metal content: The distribution of Fe, Mn, Zn, and Cu content (in μg/g dry weight) in dPAA line #2 and Col-0. The color scale reflects metal concentrations, with darker blue shades indicating higher content. *pb104::b104^dPAA line #2* exhibited significant Fe accumulation in young leaves after Fe resupply, in contrast to Col-0, where most Fe remained in the roots. Data represent the mean values from three biological replicates (n = 3). Elemental content was measured by ICP-MS. The plot was generated using the ggPlantMap package in RStudio. Asterisks indicate genotypes that differ significantly from WT within the same tissue and treatment condition (p < 0.05), as determined by two-way ANOVA and Tukey's post hoc test (n = 3 for each group).

Supplementary Figure S11

A

L1 - WT
L2 - *pb104::b104 #1*
L3 - *pb104::b104 #2*
L4 - *pb104::b104^dPAA#2*
L5 - *pb104::b104^dPAA#3*

B

C

Supplementary Figure S11. Expression profiles of genes involved in Mn, Zn, and Cu homeostasis.
Clustered heatmap showing expression profiles (RNA sequencing) of genes involved in manganese (Mn), zinc (Zn), and copper (Cu) homeostasis across tissues in pb104::b104^dPAA lines under +Fe and –Fe conditions. Genes include metal transporters and the Cu homeostasis regulator SPL7 (Guerinot, 2000; Sancenón et al., 2004; Socha and Guerinot, 2014). Heatmap was generated in RStudio version 4.4.1 using calculated Z-scores of normalized gene expression values. Purple indicates low expression and yellow indicates high expression. Asterisks indicate genotypes that differ significantly from WT within the same tissue and treatment condition ($p < 0.05$), as determined by two-way ANOVA and Tukey's post-hoc test (n = 3 for each group).

Supplementary Figure S12

A

L1 - WT
L2 - *pb104::b104 #1*
L3 - *pb104::b104 #2*
L4 - *pb104::b104^dPAA#2*
L5 - *pb104::b104^dPAA#3*

B

C

Supplementary Figure S12. Expression patterns of 23 selected Fe homeostasis genes involved in Fe acquisition and mobilization.
Clustered heatmaps showing expression profiles of 23 Fe homeostasis genes (gene selection based on Ngigi et al., 2025; Schwarz et al., 2020). Heatmaps were generated in RStudio version 4.4.1 using Z-scores of normalized gene expression values across tissues for +Fe and –Fe conditions. Purple indicates low expression; yellow indicates high expression. Asterisks indicate significant differences from WT within the same tissue and treatment (p < 0.05, two-way ANOVA with Tukey's post-hoc test, n = 3).

Supplementary Figure S13

AD	BD	AD	BD				
bHLH104-C	BTSL1	BTSL1	bHLH104-C				
bHLH104-C	BTSL2	BTSL2	bHLH104-C				15
bHLH104-C-dPAA	BTSL1	BTSL1	bHLH104-C-dPAA				
bHLH104-C-dPAA	BTSL2	BTSL2	bHLH104-C-dPAA				15
Empty	BTSL1	BTSL1	Empty				
Empty	BTSL2	BTSL2	Empty				15
bHLH104-C	Empty	Empty	bHLH104-C				
bHLH104-C-dPAA	Empty	Empty	bHLH104-C-dPAA				

10^{-5} 10^{-2}

SD - LW   SD - LWH   3-AT [mM]

Supplementary Figure S13: Yeast two-hybrid assay results showing protein–protein interactions between bHLH104 BTSL1 and BTSL2.

bHLH104 interacts with BTSL1 and BTSL2, but not bHLH104dPAA. bHLH104^ΔPAA retains the ability to heterodimerize with full-length bHLH104, serving as a control for the presence and expression of the PAA-deleted variant. Full length bHLH104 shows autoactivation in Y2H. Hence bHLH104-C and bHLH104-C-dPAA were used.

Yeast co-transformed with the AD and BD combinations were spotted in 10-fold dilution series (A600 = 10–1–10–4) on SD-LW (transformation control) and SD-LWH plates supplemented with different 3AT concentrations (conc.) as indicated (selection for protein interaction). Negative controls: empty vectors. Arrows indicate interaction.

Supplementary Figure S14

α-GST-E2-ligase   α-Myc-E3 ligase   α-Strep-substrate   α-UBQ

Supplementary Figure S14: Ubiquitination assay showing that bHLH115 undergoes ubiquitination

Ubigate assay showing ubiquitination of bHLH105 in the presence of E1, E2, E3, and ubiquitin (Ub). Immunoblots with anti-GST (E2), anti-Myc (E3), anti-Strep (substrate), and anti-His-UBQ (ubiquitin) confirm protein expression and substrate modification

Supplementary Table 1. Primers used in this study.

fw = forward, rev = reverse, d = delta a.k.a. deletion, -C = C-term, (att)B1/B2 = Gateway B attachment sites for BP reaction.

Primer name	→ Sequence (5' to 3')	Purpose	Origin
Amplification of full-length CDS or deletion constructs for Y2H			
bHLH104-C_B1 fw	GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGGCTCT	cloning of bHLH104-C (attB1)	lichtblau 2023
bHLH104-C-d_B2 rev	GGGGACCACTTTGTACAAGAAAGCTGGGTCTTAAGGCCTGAGTTCTTGAT	cloning of bHLH104-C-d (attB2)	This study
bHLH104_B2 rev	GGGGACCACTTTGTACAAGAAAGCTGGGTCTTAAGCAGCAGGAGGCCTG	cloning of bHLH104-C (attB2)	lichtblau 2022
Cloning bHLH104 promoter and CDS for transgenic lines			
bHLH104 fw	aacaggtctcaggctggatgtatccttctctcgacg	Cloning of bHLH104/bHLH104-d CDS	This study
bHLH104 rev	aacaggtctcactgaAGCAGCAGGAGGCCTGA	Cloning of bHLH104 CDS	This study
bHLH104-d rev	aacaggtctcactgaAGGCCTGAGTTCTTGAT	Cloning of bHLH104-d CDS	This study
pbHLH104 fw	AACAGGTCTCAACCTCTAGCTCTGAGAAGCGGCATC	Cloning of bHLH104 promoter	
pbHLH104 rev	AACAGGTCTCATGTTTTGAGAGTCTCCAAACAAATT	Cloning of bHLH104 promoter	This study
Amplification of full-length CDS or deletion constructs for Ubigate			
BTS-HC fw	ttgaagacaaaATGGCAGAACG	Cloning BTS HC fragment	This study
BTS-HC_rev	ttgaagacaaaagcTTAGCTGCTGGTG	Cloning BTS HC fragment	This study
optbhlh104-C fw	GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGACCGATAAACCGG	Cloning bHLH104 in pUGTP1 (attB1)	This study
optbhlh104 rev	GGGGACCACTTTGTACAAGAAAGCTGGGTCTTATGCTGCAGGCGGA	Cloning bHLH104 in pUGTP1 (attB2)	This study
optbhlh104d rev	GGGGACCACTTTGTACAAGAAAGCTGGGTCTTACGGACGCAGTTCT	Cloning bHLH104 in pUGTP1 (attB2)	This study