PLANT MICROTECHNIQUE WITH RESIN - TOWARDS PLANT HISTOLOMICS

Ivan T. Cerritos-Castro, Araceli Patrón-Soberano, Ana P. Barba de la Rosa[*].

IPICYT, Instituto Potosino de Investigación Científica y Tecnológica A.C., San Luis Potosí,

México.

*For correspondence: apbarba@ipicyt.edu.mx

Tel: 52 (444) 8342000 Ext. 2082

ABSTRACT

Plant microtechnique; a laborious discipline that provides little data. The difficulty

in handling paraffin sections and the lack of a strategy for analyzing micrographs

discourage its use. Nevertheless, it provides the fundamental cellular context for

understanding the function of biomolecules. In this work, we standardized a resin

embedding method, a chemical treatment for slides, a trichrome staining, and an

immunostaining. We also developed a strategy for micrograph segmentation,

morphometric and compositional analysis, and proposed the basis for a histolomic field.

The resin-based microtechnique was easier than the paraffin-based ones, revealed a

wealth of morphological and compositional information, and was reproducible in a variety

of tissues of plant species. Histolomics allowed numerical comparisons of primary

morphometric and compositional features such as size or protein content, as well as

composite features such as the C_4 kranz anatomy. This method takes the cellular context

to the field of omics.

INTRODUCTION

Tissues are the result of the interaction of millions of ordered molecules. Our understanding of their nature and function is largely based on their molecular disassembling for *in vitro* studies. This approach has been strengthened by omics, which correlate *in vitro* molecular abundances with the phenotype of tissues, organs and even whole organisms[1]. However, the abundance of a molecule is just as important as its location, as it is related to its function and defines the composition and morphology of tissues[2,3].

Plant histology describes the morphology and composition of tissues and cells[4]. Along with immunostaining, it localizes the abundance of molecules in tissues[3]. Despite its importance, plant histology is underutilized in plant research. The main reasons may be the dexterity required by their methods (microtechnique), and the lack of a harmonized strategy for analyzing their results. Compared to *in vitro* methods, plant microtechnique is more artisanal[5]. Classic microtechnique consist of sample fixation, dehydration, paraffin embedding, sectioning, affixing (attachment to glass slides), staining, immunostaining, dehydration, mounting (resin and coverslip placement), and microscopy[6]. Each step is susceptible to failure, but especially from affixing to mounting, which we call the dexterity-dependent steps. Paraffin sections are fragile; they may be wrinkled or broken during affixing and may become detached or scratched during subsequent steps. In addition, paraffin is removed from the section after the affixing step, making the tissue even more susceptible to damage.

If we replace the paraffin with a permanent and more stable support, the dexterity-dependent steps will be easier. LR White resin, an acrylic resin, is an embedding medium used primarily in Transmission Electron Microscopy (TEM), but is compatible with light microscopy[5]. Unlike paraffin, its hydrophilic nature allows analysis on sections without removing the resin. That provides mechanical support to the section during dexterity-dependent steps, reducing section breakage and scratches. On the other hand, effective glass slide cleaning can prevent section wrinkles, and effective "gluing" can

prevent section detachment. Various acidic and oxidizing solutions have been formulated

to clean and, at the same time, "activate" the slides[7–9]. Activation creates hydroxyl groups

that increase the wettability of the glass surface. The hydroxyl groups can then react with

(3-aminopropyl)triethoxysilane (AES) in a "gluing" process to increase the adhesiveness of

the slide[10].

Once the section is affixed, staining and immunostaining are the basic methods of

plant microtechnique to study the morphology, general composition, and specific location

of proteins in a tissue.[6] Johansen's staining has been the classic method for plant paraffin

sections since 1940 and has been modified several times[6,11–14]. It is based on safranin,

which stains mainly cell wall lignin and suberin in magenta, and fast green, which stains

mainly cytoplasmic proteins in turquoise[15–17]. Johansen's staining gives a picture of the

general composition of the whole tissue, while immunostaining gives a picture of the

abundance of specific proteins at specific locations in the tissue[3,6]. In contrast, resin

sections are usually stained only with toluidine blue for light microscopy, which stains the

entire tissue blue[18].

After a successful microtechnique, scientists often don't know what to do with the

micrographs and are limited to pointing out tissue abnormalities or the location of

immunolabel. If we define and systematize a histological analysis, scientists will be able to

exploit the information from a micrograph. The primary features of a tissue are described

in terms of morphology, morphometry and composition. Morphology qualitatively

describes the overall shape of a plant, organ, or tissue; morphometry quantitatively

describes the number, size and shape of an element; and composition describes the

overall molecular abundance in cell structures[4,19,20]. Tissue staining allows for boundary-

based morphological and morphometric analysis, as well as color-based compositional

analysis, which is typically performed manually and with the naked eye. However, color

measurement and image segmentation are digital tools that can provide comprehensive

numerical descriptions of a tissue that go beyond the naked eye[21]. On the whole, an easier

and standardized microtechnique along with a harmonized and comprehensive digital

analysis of micrographs would take us from histology to the "histolomics" research.

Here, we present a resin-based plant microtechnique to facilitate the dexterity-dependent steps, and propose a strategy for analyzing micrographs: the histolomics. We standardised a method for resin embedding, tested several solutions for cleaning, activating, and gluing of glass slides, developed a trichrome staining for resin sections, and combined it with immunostaining. We then applied on amaranth and several C_3 and C_4 plant leaves commonly used in plant research. C_4 plants have an evolved leaf morphology called "Kranz anatomy" that enhances carbon fixation that C_3 plants lack, providing a framework to study[22]. Finally, we defined and measured morphometric and compositional features to quantitatively characterize and compare the tissues.

RESULTS

Glass slides activation and gluing

We tested several solutions for cleaning and activating glass slides. Sulphocromic and Piranha solutions gave the best results: the highest wettability and correspondingly the lowest contact angle. We then "glued" the activated slides with AES. All slides showed a decrease in wettability and an increase in contact angle (Fig 1a, c). Finally, a resin section was affixed in each slide and subjected to strong washes. Again, sulphocromic and piranha activated slides gave the best results, retaining the entire section, while others detached fragments or the entire section (Fig. 1 b).

Staining of resin sections

Classic Johansen's staining did not work on resin sections; we reinvented it and made it trichrome. We tested toluidine blue, Johansen's, and Coomassie's staining methods and measured the color acquired by the cell walls (CW), protoplasm (PR) and resin background in the HSB color space (Hue, Saturation and Brightness). For more details on tissue and cell morphology, see Supplementary Fig. 1 and 2. Hue values depended on the dye, and saturation values on the staining performance. In contrast, brightness values depended mainly on the microscope and camera exposure settings and

were very similar for all micrographs. Toluidine blue tended to be monochromatic (similar

hue values), and Johansen's staining polychromatic (different hue values) but uneven and

poorly saturated (Fig. 2). Achieving different hue values between cell structures is

important for their identification and micrograph segmentation. The same is true for the

saturation between all the tissue and the unstained resin background. By testing

combinations, we found that Coomassie's solvent allows staining of resin sections with

Coomassie's dye itself and Johansen's dyes (Safranin and Fast Green), producing

polychromatic, uniform, and well-saturated colors (Fig 2). We therefore reformulated the

Johnanse's staining and added a third dye: Lugol's iodine, creating a trichrome staining

(Fig. 3).

Trichrome staining revealed the morphology and general composition of amaranth

leaf (Supplementary Fig. 3). Individually, fast green stained mainly the protoplasm

turquoise, safranin stained most of the tissue magenta, and iodine stained only the starch

granules mauve. None stained vacuoles and resin background, and fast green and safranin

did not stain starch granules. The protoplasm (PR) encompasses the cytosol and all

organelles, including the vacuole. For an accurate description, we coined the term

"proteoplasm (PT)" to refer only to the turquoise (proteinaceous) structures of the

protoplasm: cytosol and all organelles except the vacuole and starch granules. A

differentiating and synergistic effect was observed in the trichrome combination: safranin

now selectively stained cell walls, and hue and saturation values increased. The hue

change (veering) in the proteoplasm shifted the color from turquoise to blue hues.

Differently, the veering in the cell wall divided the color into magenta and reddish hues

(Supplementary Fig. 3). We measured the color of cell walls and proteoplasm separately

in the epidermis, mesophyll, and bundle sheath. Proteoplasm had similar hues in all

tissues. However, the bundle sheath cell wall had magenta, reddish, and even orange

hues, while the epidermis and mesophyll cell walls had only magenta hues. In addition,

the bundle sheath showed stained and unstained starch granules. This pattern was

confirmed by TEM microscopy, which showed electrodense and electrolucent starch

granules (Fig. 4).

The order and timing of each dye in trichrome staining were important for the
color selectivity. A section stained with fast green could be correctly counterstained with
safranin for 5 to 10 s, after which safranin replaced fast green (Supplementary Fig. 4). A
section stained with safranin could not be correctly counterstained with fast green
(Supplementary Fig. 5). Finally, iodine staining (alone or in a counterstain) produced
random intensities independent of time: null, intermediate or intense, even between
different sections of the same sample on the same slide; we could not identify the reason.
However, along the same section, the random intensity obtained was always uniform
(Supplementary Fig. 6).

For a successful staining and micrograph interpretation, there were some technical
issues that need to be taken into account. The orange hue of the bundle sheath cell wall
was best observed with the 40X objective and a diaphragm aperture equivalent to the 10X
objective. Mounting with entellan improves the sharpness from the start and as it dries.
On the other hand, the iodine staining of the starch granules may fade over time, from
days to months. For that reason, we recommend that micrographs be taken one to three
days after mounting. Finally, it is important to be aware of common artifacts in resin
sections for proper histological interpretation (Supplementary Fig. 7).

Resin-based microtechnique resolution

The resin-based microtechnique allowed light microscopy to resolve small
organelles and even ultrastructures that would normally require TEM microscopy to
observe. Trichrome staining identified the cell wall in magenta to orange, the proteoplasm
in turquoise, and the starch granules in mauve. In fact, the proteoplasm showed different
saturation values, allowing the identification of the cytosol, chloroplasts, mitochondria,
nucleus, and nucleolus, if present (Fig. 5 and Supplementary Fig. 8 and 9). Furthermore, an
intensely iodine counterstained paradermal section showed two layers in the epidermal
cell wall: an internal orange and an external magenta. This section also showed
ultrastructural details of what appeared to be: membranes within vacuoles, chloroplasts

with protein granules, unknown rod-shaped organelles, and vesicles transporting cell wall

material, among others (Supplementary Fig. 10).

Histolomic analysis

The resin-based microtechnique required a strategy for the analysis of results and

towards histolomics research we proposed a pipeline of four steps: visual analysis, color

analysis, micrograph segmentation, and data analysis. First, visual analysis consisted of

classical qualitative identification of tissues and overall morphology (Supplementary Fig. 1

and 2). We also added the identification of organelles and other relevant features visible

thanks to the high resolution of resin-based microtechnique. Cross sections of a young and

a senescent amaranth leaf were compared. When looking at the entire micrograph,

general differences were observed in the senescent leaf such as decreased proteoplasm,

increased starch granules, and rounding of the palisade mesophyll. When looking at the

enlarged micrograph, small size differences, such as reduction of cell wall thickness,

appearance of starch granules in the palisade mesophyll, disappearance of cytosol, and

others, were observed in the senescent leaf (Supplementary Fig. 11).

Second, the color analysis consisted of measuring the color of the cell structures

and assigning them a hue range for color segmentation (Fig. 4). Third, segmentation of the

micrographs consisted of digitally isolating cell structures from the background into layers

for further data analysis (Supplementary Video 1). The color contrast provided by the

trichrome staining allowed to digitally segmenting tissues based on color (composition)

and cell boundaries. We semi-automatically segmented cell walls, proteoplasm, and

unstained elements (vacuoles, starch granules, and background) based on their hue and

saturation differences. The proteoplasm could be further segmented into low-saturated

elements (nucleus and cytosol) and high-saturated elements (the rest of the organelles).

The layers show tissue composition and localization more clearly than the original

micrograph. However, the thin cell walls of the mesophyll lacked continuity, making

boundaries-based morphometric analysis impossible (Supplementary Fig. 12). Therefore,

we manually segmented the micrograph cell by cell. The previous cross-sections of a

young and senescent amaranth leaf were segmented for data analysis (Supplementary Fig.
13 and 14).

Fourth, data analysis consisted of defining and measuring primary and derived
morphometric and compositional features on the layers and comparing them.
Morphometric features involved the quantitative measurement of physical traits such as:
number, size, perimeter, area, thickness, and circularity. Compositional features involved
the semi-quantitative measurement of stained molecules by densitometry (as an SDS-
PAGE gel): proteins, lignin and suberin, starch granules, and immunolabels. Smaller and
more rounded palisade mesophyll cells, lower cell wall thickness, lower proteoplasm
content and significantly higher starch granule content were found in the senescent leaf.
On the other hand, the protein content showed a decrease and the lignin content showed
no differences (Supplementary Fig. 15).

After all, the more colors, the more information to analyze. Intense iodine
counterstaining allowed furthering semi-automatically segment the starch granules, as
well as an inner and outer wall, based on their color veering (Supplementary Fig. 16).

Performance of resin-based microtechnique in different plant species and tissues

The resin-based microtechnique produced comparable results between different
plant species and tissues. To test the versatility of the resin-based microtechnique and the
potential of histolomic analysis, we applied them to three C_3 plant leaves: *Arabidopsis*
thaliana (arabidopsis), *Triticum aestivum* (wheat), *Glycine Max* (soybean), and two C_4
plants leaves: *Zea mays* (maize) and *Amaranthus cruentus* (amaranth). Resin-based
microtechnique worked on all tissues, dye selectivity was the same as previously observed
for amaranth leaves, and randomness of iodine counterstaining was also observed.
Histolomic analysis revealed a number of similarities and differences between the leaf
sections: the same fundamental tissues but different morphology, morphometry and
composition (Supplementary Fig. 17 to 23). In order to deal with data and to obtain a
physiologically and evolutionarily meaningful value, we proposed a composite feature: the
"C_4 kranz level". This feature condensed some primary and derived morphometric and

compositional features, related to the C$_4$ Kranz anatomy[23] (Supplementary Fig. 24). *A.*

thaliana had the least developed C$_4$ Kranz anatomy and *A. cruentus* the most developed.

We also tested the resin-based microtechnique to follow histological changes during

germination and cotyledon maturation of amaranth seeds. The selectivity of the dyes was

the same as for leaves and revealed the development of organelles and cell wall

thickening (Supplementary Fig. 25 and 26). Finally, trichrome staining also worked on

paraffin sections, even without deparaffinization (Supplementary Fig. 27).

Trichrome staining and immunostaining of resin sections

Trichrome staining can be used along with immunostaining on resin sections for

precise protein localization. We performed an immunostaining in an amaranth leaf section

to detect the RuBisCO large subunit (RbcL) responsible for carbon fixation. We used the

BCIP/NBT chromogen for signal development and then stained with trichrome staining.

BCIP/NBT gave an intense blue signal and did not interfere with the performance and hues

of the trichrome staining, resulting in a tetrachrome stain. Micrograph was segmented

based on the hue ranges of each dye, and protein and RbcL concentrations were

measured. RbcL was detected in almost all tissues, but mainly in the bundle sheath and

mesophyll. The bundle sheath contained about two times more RbcL than the mesophyll

(Fig. 6).

DISCUSSION

The resin-based microtechnique facilitated the dexterity-dependent steps and

offered advantages over the classic paraffin-based microtechnique. First, resin provided

mechanical support for the tissue in all steps, preventing breakage and scratching, and

preserving tissue structure better than paraffin (Supplementary video 1, and

Supplementary Fig. 11 and 27). Also, multiple thin and ultrathin sections can be obtained

simultaneously for light and electron microscopy analysis of the same cells, or for testing

multiple antibodies. Second, glass slide activation and gluing perfected the section

adhesion. Pirahna or sulphocromic activated and AES-glued slides eliminated section

detachment in all the slides that we used in this work (Fig. 1). Sections could only be detached by deliberate scraping. This allows multiple analyses to be performed without the risk of losing sections (Fig. 6). Due to the high toxicity of Cr^{6+}, we recommend using only the piranha solution for slides activation. On the other hand, it is worth to note that classic microtechnique do not take into account the activation of slides and recommend a short gluing time with AES[24], which in our experience did not work. Finally, the activation also deeply cleans the slides, reducing wrinkles and artifacts caused by dirt (Supplementary Fig. 7); however, there are also some drawbacks. Resin-based microtechnique requires an ultramicrotome, which may not be available in some laboratories. Despite thorough slide cleaning, resin sections naturally tend to wrinkle during affixing as water evaporates, so the best affixed sections has to be selected.

Our trichrome staining method (Figure 3) significantly reduced steps, time, reagents, toxicity, and variability compared to previous methods[6,11–14]. Classic staining methods such as Johnanse's rely on multiple steps, repeated use of several reagents on staining jars, and exposure to toxic or even explosive reagents such as picric acid[6]. Our trichrome method reduced the 18 steps of the Johansen's method to 8 steps, the 2-48 hours to 5-10 min, and 6 reagents instead of 9. In addition, our trichrome method uses small volumes of less toxic and more accessible reagents, making them easier to dispose of, reducing variability due to repeated use (Supplementary Video 1).

Trichrome staining revealed the general molecular composition of the cell structures. According to Johansen, fast green stains cytoplasm and cellulosic walls, whereas safranin stains lignified and/or suberized walls, as well as nucleus and chromosomes[6]. Precisely, fast green stains proteins and is even used to quantitatively stain SDS-PAGE gels and nitrocellulose membranes[15,16]. Safranin stains lignin and suberin, but its effectiveness in staining chromatin is not well-documented[17,25]. Iodine reacts with various polysaccharides, showing the highest affinity for amylose[26,27]. So, trichrome staining was consistent with the reported molecular specificity of the dyes, except for the nucleus and chromatin. Overall, the nucleus, nucleolus, and heterochromatin stained turquoise (Fig. 5). *T. aestivum* and *Z. mays* nuclei showed heterochromatin, but only those

of *T. aestivum* stained magenta (Supplementary Fig. 19 and 20). Safranin can interact with

DNA[28]. However, we found that safranin alone stained almost all tissue structures; its

specificity was defined by counterstaining with fast green (Supplementary Fig. 3 and 4).

Therefore, there should be specific molecules responsible for the safranin staining in *T.*

aestivum heterochromatin. Similarly, trichrome staining revealed stainable and non-

stainable starch granules. The amylose content could explain the differences. However,

the unstained granules were whiter and more electrolucent than the background in light

and electron microscopy respectively. On the other hand, the evidence is suggestive but

not conclusive if non-stained granules are the same as electrolucent granules. We can only

conclude that the amaranth leaf has two types of starch granules with different

compositions and interactions with LR white resin. Strikingly, both types can be formed in

the same chloroplast (Fig. 4).

Iodine counterstaining also differentiated the molecular composition of the cell

wall. It caused safranin-stained cell walls veering from magenta to reddish and orange

hues (Fig. 4). Our evidence suggests that this occurs gradually and selectively beginning in

the bundle sheath wall (Fig. 4) and continues until most of the wall of all tissues veers

(Supplementary Fig. 10). However, the epidermal wall showed an internal orange wall and

an external thicker magenta wall. Moreover, color segmentation showed that all cell walls

retained a small magenta portion (Supplementary Fig. 16). The cell wall consists of several

layers from the outside to the inside: middle lamella, primary wall, secondary wall, and

plasma membrane. The primary wall is composed of cellulose, hemicellulose, proteins,

and pectin; and the secondary wall is composed of cellulose, hemicellulose, and lignin[29].

The epidermal wall has additional outer layers of cutin and wax[30]. The primary wall is

about 0.1 μm thick, while the lignified secondary wall is thicker, about 2 to 10 μm[31]. Our

evidence suggests that safranin stains lignified and non-lignified walls, and that the orange

walls correspond to secondary walls (thicker) and the magenta walls to primary walls

(thinner). In addition, safranin stains lignin and suberin. Lignin can form complexes with

iodine, changing its UV-VIS absorption spectrum, while suberin interaction with iodine is

poorly understood[32–34]. The lignin-iodine reaction could cause the lignified secondary

walls veering while the non-lignified primary walls and the cutinized epidermal walls

remain magenta. Nevertheless, the ring structures of the tracheids are also lignified and

did not veer to orange hues[4] (Supplementary Fig. 10). Iodine counterstaining left several

questions about its reactions with starch granules and cell walls, as well as the

composition of these structures themselves. More research is needed to understand them

and also to reduce the randomness of iodine staining.

Our resin-based microtechnique resolved tissues, cells, organelles, and even

ultrastructural details, enabling large-scale histolomic analysis. The smaller resolved

ultrastructures were membranes within the bundle sheath vacuoles approximately ≈ 336-

560 nm thick (Supplementary Fig. 10). These membranes corresponded to cytosol

invaginations (310-900 nm thick) as proved by TEM microscopy (Fig. 5). The resolution of

widefield microscopy is limited to ≈ 200 nm in the X-Y axis (resolution), and 500–700 in the

z axis (depth of field)[35]. Resin-based microtechnique yielded ultrastructural detail close to

the resolution limit of widefield microscopy and close to the ultrastructural detail

provided by TEM microscopy at low magnifications (Fig. 5 and Supplementary Fig. 8-10). It

is in this "bridge" where the histolomic potential of the resin-based microtechnique lies:

to know a tissue in detail with TEM microscopy and then to analyze large areas at the

tissular, cellular and ultrastructural levels with light microscopy. It is conceptually similar

to correlative microscopy, but technically simpler and histologically more comprehensive.

On the other hand, we used resin sections about ≈500 nm thick to be within the depth of

field of widefield microscopy and to avoid interference from out-of-focus planes; the

essence of confocal microscopy. Thus, resin-based microtechnique complement the range

of high-resolution techniques that rely on fluorescent labeling, which tend to show only

what is immunolabeled and lose information from the rest of the tissue.

Histolomics requires a standardised strategy for micrograph analysis and

reporting. The nature of micrographs raises the need for an online platform with

interactive annotations for convenient sharing and tracking of plant tissue research

(Supplementary Fig. 11). On the other hand, colors in plant research are usually described

only by their "intuitive name". Accurate color description is important for micrograph

analysis. The RGB color space is widely used, even for lignin quantification[36]. However, the

HSB color space is the most intuitive and effectively separated the influence of dye (hue),

staining performance (saturation), and microscope setup (brightness) (Fig. 2). Hue

differences allowed segmentation of up to four cell structures, leaving some space in the

color wheel for more dyes. However, as with fluorescence microscopy, there is a risk of

color overlap[36], as was observed with cell walls and starch granules (Fig. 6).

Digital micrographs are data and should be analyzed beyond the naked eye. The

morphometric features were useful in quantifying the morphological differences between

C_3 and C_4 leaves, except for circularity, which did not show significant differences despite

being evident in the micrographs (Supplementary Fig. 22). Compositional features

revealed important physiological trends, such as the same protein concentration in young

and senescent amaranth leaves despite a reduction in proteoplasm content

(Supplementary Fig. 15). The primary and derived morphometric and compositional

features provide a basic description of the tissue, but can be selected, weighted, and

combined to investigate a particular phenomenon. At the "C_4 kranz level", we selected

primary and derived features related to the morphological traits of the C_4 Kranz

anatomy[23], weighted them according to their importance, and summed them to obtain a

score (Supplementary Fig. 24). This score condensed a complex micrograph, allowed

quantitative comparisons between plants, and could also be correlated with other

physiological or evolutionary issues such as carbon fixation efficiency or C_3-C_4 evolution.

Different combinations of primary and derived features can be used to quantitatively

study the effect of other phenomena such as biotic and abiotic stresses, genetic

mutations, nutrition, etc. on the phenotype of plant tissues.

In this work, we presented some primary, derived and composite features, but

more should be developed. Reliable and different cell shape formulas are needed, as

circularity failed. Positional features of cells and organelles will also be useful: e.g. the

position of organelles in bundle sheath cells to weigh their influence at the "C_4 kranz

level"; the position of organelles during cell division, among others. In addition, the tissue

area and the number of individual cells or organelles that can be measured are limited by

the need for time-consuming manual segmentation. Deep learning strategies should be

developed to automate micrograph segmentation and measurements of more individual

elements and larger areas to improve statistical power and leverage the resolution of

resin-based microtechnique.

Histolomics provided large-scale histologic descriptions, but took into account the

phenotypic individuality of each cell. In *A. thaliana* was observed a unique proteinaceous

cell vascular bundles (VBPC) with a smooth morphology and 26% of the protein content of

vascular bundles (Supplementary Fig. 17). This suggests a highly specialized function, and

although *A. thaliana* is the most studied plant, we could not find any information about

this cell. From ultrastructure to whole section, and from germination to senescence,

histolomics revealed histological similarities that species and tissues share, as well as

peculiarities that define them.

Histolomics challenged the C_4 category of the amaranth plant. RuBisCO fixes

carbon dioxide to form carbohydrates including starch. However, RuBisCO can also fix

oxygen, reducing the efficiency of carbon fixation. C_4 plants accumulate RuBisCO mainly in

bundle sheath cells, where carbon dioxide is enriched and thick lignified walls prevent

escape[23]. In amaranth leaves, we found increasing amounts of RbcL, starch granules, cell

wall thickness, and lignin concentration in palisade mesophyll, spongy mesophyll, and

bundle sheath. The amaranth leaf has an almost histolomically "perfect" Kranz anatomy,

but its RuBisCO have a distribution close to C_2 plants[23].

Omics have developed thanks to technological advances such as next-generation

sequencing and mass spectrometry[37], whereas plant histology still uses techniques that

are almost a century old[6]. In this work, plant microtechnique was reinvented to simplify its

methods and generate reproducible micrographs that allowed histolomic analysis. We

define histolomics as the large-scale, systematic, high-resolution microscopic study of

tissue composition, morphology, and morphometry from the ultrastructural to the whole

tissue level. Here we presented the first database of histolomes of plants commonly used

in research, obtaining data and images that make us rethink the perspective from which

we study life and its beauty.

ONLINE METHODS

Glass slides preparation

The glass slides treatments (2947-25X75, Corning) were performed in three steps: washing, chemical activation, and silanization. Washing: glass slides were soaked in 2% soap (Hyclin Plus Neutral, Hycel) for 30 min, and then both sides of each slide were scrubbed with 15% soap with streak-free fiber (3068257, Scotch-Brite). They were rinsed with tap water and then with Milli-Q water. Chemical activation: Six different solutions were tested: MeOH:HCl 1:1 (MeOH-HCl); suflochromic mixture, $K_2Cr_2O_7 \approx 4\%$ in concentrated H_2SO_4 (H_2CrO_4); piranha solution, 30 ml concentrated H_2O_2 in 70 ml concentrated H_2SO_4 (Piranha); glacial acetic acid (CH_3COOH); concentrated sulfuric acid (H_2SO_4) or 1 M sodium hydroxide (NaOH). The sulfochromic mixture was prepared by adding a little water to $K_2Cr_2O_7$ until a paste was made, and then the acid was added little by little. Warning! All solutions are dangerous; particularly the sulfochromic mixture and piranha solution are strong oxidizers and can react violently with organic material. In addition, the sulfochromic mixture is highly toxic, so it must be handled with care and disposed of according to your country's regulations. One group of slides was immersed in staining jars for each of the solutions for 30 min at room temperature, and another was kept without chemical activation (None). Slides were rinsed with Milli-Q water 3 times, excess water was wrung out on a paper towel, and dried in an oven at 60 °C for 30 min. Silanization: Dried slides were immersed in 4% (3-aminopropyl)triethoxysilane (AES, 440140, Sigma-Aldrich) solution in absolute ethanol for 1 h at room temperature, drained and allowed to stand in 1 mM acetic acid in absolute ethanol for 10 min. They were rinsed with Milli-Q water and dried as above, increasing the oven time at 60 °C to 1 h. It is important that after the last rinse, the slides are kept in dust-free containers to avoid artifacts in microscopy.

To evaluate the effectiveness of the activation, the contact angle of the slides with water was measured. A slide was placed on a flat, level surface behind of a camera (T5i with EF 100mm f/2.8L Macro IS USM lens, Canon) and in front of a backlit flash (newer S-

400N). 1 µL of Milli-Q water was added to the slide, allowed to stand for 30 s, and a

photograph was taken. The contact angle was measured on the photographs using the

ruler tool in Adobe Photoshop CS6.

To evaluate the effectiveness of silanization of the slides, their ability to retain a

resin section under strong washes was tested. Thin (500 nm) cross sections of *Amaranthus*

cruentus midrib were attached to each group of slides at 70°C for two minutes and

subjected to three aggressive washing procedures. First, they were digested with

proteinase K (10 µg/ml in PBS), denatured at 95°C for 2 min, and rinsed with a wash bottle

for 10 s. Second, they were heated in water at 95°C for 15 min, followed by rinsing with a

water jet for 10 s. Finally, they were subjected to heating in MeOH at 95°C for 3 min,

followed by rinsing with a water jet for 10 s. The sections were stained with safranin (0.5%

in water) and micrographs were taken. To prevent drying, the slides were covered with

the lid of a microtube during the incubation steps (See supplementary Video 1).

Plant material

Seeds of *Amaranthus cruentus* L cv. Amaranteca (Amaranth), *Arabidopsis thaliana*

(Arabidopsis), *Triticum aestivum* (Wheat), *Glycine max* (Soy bean), and *Zea mays* (Maize)

were germinated in 500 g of BM2 HP germination substrate (Berger, Saint-Modeste,

Quebec, CA). After 4 weeks, one leaf per plant was harvested. For *A. cruentus*, one leaf

was harvested at 4 weeks (young) and one at 8 weeks (senescent). In addition, during

germination of *A. cruentus*, seedlings were harvested at different stages: from seed to

seedling with mature cotyledons.

General sample processing and microscopy

Approximately 3 x 4 mm samples were taken from the midrib, a primary vein and

the leaf blade. In addition, one seed, one germinating seed, and complete cotyledons at

various stages of development were sampled from *A. cruentus*. Samples were fixed in 3%

glutaraldehyde, 2% formaldehyde, 0.5% Tween 20 in PBS (5 to 10 times the sample

volume) under vacuum for 20 min, and then shaken for 1 h, all on ice. Fixation was

continued overnight at 4°C and then rinsed three times with PBS while shaking for 1 hour

each time at room temperature. Samples were dehydrated in 30%, 50%, 70%, 95% and

absolute ethanol with shaking for 1 h at each step. Pre-inclusion in LR White

resin:absolute ethanol 1:1 with shaking was performed overnight, followed by inclusion in

pure resin (#14381, Electron Miscorscopy Sciences, PA, USA) another night. Pre-

embedding can be done in less time (4 h) for light dense samples such as leaves and

overnight for dense samples such as seeds. Usually, all samples were left overnight to

ensure complete and homogeneous resin penetration. Finally, samples were transferred

to gelatin capsules #2 (70103, Electron Miscorscopy Sciences, PA, USA) with pure resin

and polymerized at 55 °C for 48 h. For light and electron microscopy, thin and ultrathin

sections, respectively, were obtained (PowerTome PC RMC, Boeckeler Instruments). 3 x 4

mm samples can be difficult to handle; alternatively, slightly larger samples (≈ 5 x 6 mm)

can be obtained and the excess cut off with a scalpel or microscissors prior to

encapsulation. It is important that the sample does not exceed these dimensions so that it

can be positioned inside the capsule. Further details on embedding, sectioning and

staining are shown in Supplementary Video 1.

Thin sections were transferred to slides using a perfect loop, affixed in hotplate at

70 °C for 2 min and subjected to different procedures (described below). Finally,

micrographs were obtained with an Axio Imager M2 microscope (Zeiss) and a Canon T5i

camera. Ultrathin sections were placed on Formvar Carbon Support film, 150 Cupper

mesh (FCF-150 Cu) and counterstained with 2% uranyl acetate for 10 min, followed by 2%

lead citrate for 4 min. Samples were examined on a JEM-200 CX transmission electron

microscope (JEOL, Ltd.) at 100 kV equipped with a digital camera (SIA, Duluth).

For paraffin embedding, an *A. cruentus* leaf sample was fixed and dehydrated in

ethanol as described previously, but with two absolute ethanol steps. Sample was then

infiltrated in absolute alcohol:xylene 1:1 for 1 h, xylene for 1 h, xylene:paraffin (Tissue-Tek

VIP 4005 Embedding Medium, Sakura) 1:1 for 1 h at 60 °C, paraffin 1 for 1 h at 60 °C and

finally paraffin 2 for 1 h at 60°C. 10 µm sections (CUT 6062, SLEE, Nieder-Olm, Germany)

were obtained and placed in a water bath at 32 °C, collected on slides and affixed on

hotplate at 60 °C for 2 min. The sections were then stained and micrographs were taken.

Staining

Preliminary assays: Paradermal sections were stained with different methods.

Toluidine blue O (70103, Ted Pella, CA, SUA) for 5 min and rinse with water. Johansen's

method (Fig. 3a)[6]. Coomassie blue (Coomassie brilliant blue 0.025% in MeOH 40% Acetic

Acid 10%) for 5 min and rinse with water. Safranin (safranin O 1% in MeOH 40% and acetic

acid 10%) for 10 s, rinse with water, Coomassie blue for 10 s and rinse with water. Fast

green (fast green fcf 1% in 40% MeOH and 10% acetic acid) for 5 min, rinse with water,

Coomassie blue for 2'30'' and rinse with water. Finally, fast green for 5 min, rinse with

water, safranin for 5s and rinse with water. After the last water rinse, the slides were

dried on hotplate at 70 °C for 2 min and mounted with acrylic resin (Entellan, Merck) and a

coverslip.

Trichrome staining. The sections in Supplementary Figs. 4 to 6 were stained as

indicated in Fig. 3b, making modifications to the staining steps, order, and times as

indicated in each Figure. The sections in the remaining Figures (except Fig. 1 and 2) were

stained as indicated in Fig. 3b without modification.

Immunohistochemical staining

Resin sections were subjected to antigen retrieval with 100 µl of proteinase k at 50

µg/ml in PBS at 37°C for 30 min. The sections were then rinsed with TBST (Tris 20 mM,

NaCl 150 mM, Tween 20 1% pH 7.6). Blocking was performed with 200 µl of 1% BSA in

TBST (blocking solution) at 37 °C for 30 min. The blocking solution was removed and 200

µl of rabbit anti-RbcL (RuBisCO large subunit) antibody (Rabit anti-RbcL, Agrisera) diluted

1:500 in blocking solution was added. Sections were then incubated at 37 °C for 1 h, and

rinsed twice with TBST for 5 min each time. Alkaline phosphatase-conjugated goat anti-

rabbit antibody (200 µl) (Merk-Millipore) diluted 1:500 in blocking solution was added and

incubated at 37 °C for 30 min. Sections were rinsed twice with TBST and color developed

with fresh BCIP/NBT solution at 37 °C for 2 h. BCIP/NBT solution: 0.02% BCIP, 0.03% NBT,

in alkaline phosphatase buffer (100 mM Tris, 100 mM NaCl, 5 mM $MgCl_2$, 0.05% Tween 20

pH 9.5). Finally, the sections were rinsed twice with TBST, twice with Milli-Q water, and

dried on hotplate at 70 °C for 2 min. A negative control was performed without the anti-

RbcL antibody, and a positive sample was stained with trichrome staining. Incubation

steps were performed by covering the section with a microtube lid, as shown in

Supplementary Video 1.

529   Color measurement

530   Different areas (n=30) of the micrographs were measured with the Adobe

531   Photoshop CS6 "eyedropper" tool with a sample size of 3 x 3 pixels, and statistical analysis

532   was performed with GraphPad Prism 9.

534   Semi-automatic image segmentation with Matlab R2024a

535   The micrographs were segmented using the "Color Tresholder" tool in the "HSV"

536   color space. The segmented images were exported in black and white. To make accurate

537   adjustments, the HSV values were monitored in the segmentation function, and the

538   sliders were adjusted until the desired values were obtained. Images were exported at a

539   resolution close to the original micrograph (See supplementary video 1).

541   Manual image segmentation and morphometric and compositional measurement with

542   Adobe Photoshop CS6

Morphometric analysis. A micrograph layer with a layer mask was generated for

each element to be measured. The density of the mask was adjusted to 50% and each

element was manually contoured with the brush tool. The workspace was calibrated using

the "set scale to size" tool and the micrograph scale bar. The selections of each element to

be analyzed were retrieved through the layer masks and measured with the "record

measurements" function. The data were exported and analyzed with GraphPad Prism 9.

549 Compositional analysis by densitometric analysis of lignin, proteins and

550 immunolabel. The micrograph was converted to grayscale/negative, and a baseline

551 adjustment was performed. The black and white segmented image was resized to the size

552 of the micrograph and used as a mask. Finally, image analysis was performed on the

553 masked grayscale/negative micrograph and the data was saved for statistical analysis in

554 GraphPad Prism 9.

555 A detailed tutorial on image segmentation and analysis using Matlab and Adobe

556 Photoshop CS6 is shown in the second part of Supplementary Video 1.

Supplementary information

559 Supplementary video 1: https://www.youtube.com/watch?v=di2SDqe49rY

Acknowledgments

562 Ivan Takeshi Cerritos Castro thanks to CONACYT fellowship 719217, and the grant

563 CNCA.FONCA.04S.04.ACT.PR.ACT.178.18. from the National Fund for Culture and Arts

564 (FONCA), Mexico. The authors thank to Laboratory for Nanoscience and Nanotechnology

565 Research, LINAN, IPICYT.

Author contributions

568 **Ivan Takeshi Cerritos-Castro:** Conceptualization, investigation, fund acquisition,

569 methodology, and writing the original draft. **Olga Araceli Patrón-Soberano:** Investigation,

570 methodology, review and editing. **Ana Paulina Barba de la Rosa:** Supervision, funds,

571 resources, review, and editing.

Competing financial interest

574 The authors declare no competing financial interests.

FIGURE CAPTIONS

Figure 1. Activation and gluing treatments increase the wettability and adhesiveness of

glass slides.

a Contact angle photographs after chemical activation and gluing (silanization). b Resin

sections in the different slides after strong washing. Arrows indicate detachments. Scale

bar = 500 μm. c Contact angle mean (n=3) ± SD. One-way ANOVA and Holm-Šídák's post-

tests were performed, letters indicate significant differences at p<0.01. Activation

treatment is indicated at the top. Activation creates hydroxyl groups that make the glass

hydrophilic, causing water droplets to spread widely (wettability) and reducing the contact

angle. The hydroxyl groups then react with the AES during the gluing process, improving

the adhesiveness of the glass.

Figure 2. Classic Johansen's staining does not work on resin sections.

a Light micrographs of *A. cruentus* leaf paradermal sections stained with different

methods. Scale bar = 50 μm. b Violin plots (n=30) with median, interquartile range and

hue ranges (filled, only for saturation ≤ 15%) observed in cell wall (CW), protoplasm (PR)

and background (BK). Kruskal-Wallis ANOVA and Dunn's post-test were performed;

differences are shown at p<0.01. HSB graphs correspond to the top micrograph.

Hue (H) represents the visible light spectrum on the color wheel, with values ranging from

0 to 360°. Saturation indicates the amount of color and brightness the amount of light,

and their values range from 0 to 100%. Low saturation tends to gray and high saturation

tends to vivid colors. Low brightness tends to black and high brightness tends to white.

The hue depended on the dye, the saturation on the staining performance, and the

brightness on the illumination settings of the microscope and camera (standardized).

Figure 3. Reinvented Johansen's staining for resin sections and made it trichrome.

a Johansen's 1940 staining method with safranin and fast green for paraffin sections. **b**

Trichrome staining method with fast green, safranin, and Lugol's iodine for both paraffin

and acrylic resin sections. The Johansen method has been a reference in plant histology

since 1940. It requires many steps, time, and toxic, explosive, and expensive reagents.

Johansen's method did not work on resin sections, so we completely reformulated its dyes

and steps, and added Lugol's iodine. The trichrome method greatly reduced the steps,

time, toxicity, added a new dye, and worked on both resin and paraffin sections.

Figure 4. Trichrome staining differentiates cellular composition beyond Johansen's dyes.

a Light micrograph of a cross-section of *A. cruentus* leaf stained with trichrome staining.

Arrows indicate the hue range of some structures. **b** Violin plots (n=30) with median,

interquartile range, data distribution (circles), and hue range observed in the epidermal

wall (E-CW), the mesophyll wall (M-CW), bundle sheath wall (B-CW), bundle sheath

proteoplasm (B-PT), mesophyll proteoplasm (M-PT), mesophyll starch granules (M-ST),

and bundle sheath starch granules (B-ST). **c** Enlargement of the white box in a. Mauve

arrows point to stained and white arrows to unstained granules. d TEM of an *A. cruentus*

leaf cross section. Mauve arrows point to electrodense granules and white arrows to

electrolucent granules. ST, starch; CL, chloroplast; CW, cell wall; MI, mitochondrion; NU,

nucleus; and NL, nucleolus. Addition of iodine counterstaining to a tissue stained with

Johansen's dyes caused hue changes (veering). Veering occurred mainly in the CW in a

tissue-dependent manner. M-CW hardly veered, E-CW veered to reddish, and B-CW

veered to reddish and even orange hues, suggesting compositional differences in their cell

walls. The proteoplasm, on the other hand, showed no significant veering between

tissues. Finally, some ST stained mauve, while others remained white, even whiter than

the background itself. This pattern was repeated when the section was examined by TEM,

suggesting compositional differences in the starch granules.

Figure 5. Resin-based microtechnique allows light microscopy to resolve structures that

would normally require TEM.

Digitally isolated bundle sheath cells from light and electron micrographs (not the same

cell), from a young (4 weeks), and a senescent (8 weeks) *A. cruentus* leaf. Scale bar= 5 μm.

High-resolution light micrographs revealed several organelles and ultrastructures that

were identified by TEM. The characteristic cell wall thinning of PITS was observed in the

bundle sheath-mesophyll cell junctions. Vacuoles were easily identified by their lack of

color and texture; however, some cytoplasmic invaginations were detected. All

proteoplasm stained turquoise, but saturation differences revealed several organelles and

cytosol. Although these structures were not sharply resolved by light microscopy, it was

sufficient to identify them and follow their behavior during the senescence process. The

senescent cell has lost its cytoplasm, its chloroplasts have lost their centripetal position,

contain more starch and are undergoing degradation, but the cell still retains the nucleus

and nucleolus. This behavior can be distinguished by both optical and TEM microscopy.

Figure 6. Trichrome staining and immunostaining revealed the precise localization of

RbcL in cells and organelles.

Light micrographs of *A. cruentus* leaf sections immunostained for RbcL. **a** No signal was

observed in the negative control. **b** RbcL and background (BK) signal was observed in the

anti-RbcL sample. **c** Trichrome staining stained the starch but did not cause significant

veering in the cell wall. **d** The combination of immunostaining and trichrome staining

summed the colors of both techniques without affecting each other. AE, adaxial

epidermis; PM, palisade mesophyll; VB, vascular bundles; BS, bundle sheath; SM, spongy

mesophyll; BE, abaxial epidermis. To differentiate the immunostaining signal: a weak

signal outside the proteoplasm was considered as background and a strong signal inside

the proteoplasm as RbcL. Tetrachrome staining (**d**) facilitated this differentiation by

showing the immunostaining signal and the stained tissue, in contrast to (**b**) where the

tissue lacks contrast. **e-h** Color segmentation of (**d**) showing separately the cellular

elements and the RbcL signal. **i** Violin plots (n=30) with median, interquartile range, and

observed color range. Kruskal-Wallis ANOVA and Dunn's post-test were performed;

differences are shown with $p<0.01$. Although there were significant differences, the hue

ranges overlapped, so in practice they were adjusted (in Matlab) to intermediate values

for color segmentation (**j**). Densitometric analysis showed the highest RbcL content in the

BS, but similar concentrations in all tissues (**l**).

REFERENCES

1.   Zivy, M. *et al.* The quest for tolerant varieties: the importance of integrating

"omics" techniques to phenotyping. *Front. Plant Sci.* **6**, (2015).

2.   Xu, H. & Bassel, G. W. Linking Genes to Shape in Plants Using Morphometrics. *Annu.*

Rev. Genet. **54**, 417–437 (2020).

3.   Millar, A. H., Carrie, C., Pogson, B. & Whelan, J. Exploring the Function-Location

Nexus: Using Multiple Lines of Evidence in Defining the Subcellular Location of Plant

Proteins. *Plant Cell* **21**, 1625–1631 (2009).

4.   Taylor, T. N., Taylor, E. L. & Krings, M. 7 - Introduction to Vascular Plant Morphology

and Anatomy. in (eds. Taylor, T. N., Taylor, E. L. & Krings, M. B. T.-P. (Second E.)

201–222 (Academic Press, 2009). doi:https://doi.org/10.1016/B978-0-12-373972-

8.00007-3.

5.   McCann, M. T., Ozolek, J. A., Castro, C. A., Parvin, B. & Kovacevic, J. Automated

Histology Analysis: Opportunities for signal processing. *IEEE Signal Process. Mag.*

32, 78–87 (2015).

6.   Johansen, D. A. *Plant microtechnique*. (McGraw-Hill Book company, Inc., 1940).

7.   Labit, H. *et al.* A simple and optimized method of producing silanized surfaces for

FISH and replication mapping on combed DNA fibers. *Biotechniques* **45**, 649–658

(2008).

8.   Han, Y., Mayer, D., Offenhäusser, A. & Ingebrandt, S. Surface activation of thin

silicon oxides by wet cleaning and silanization. *Thin Solid Films* **510**, 175–180 (2006).

9.   Jakša, G., Štefane, B. & Kovač, J. Influence of different solvents on the morphology

of APTMS-modified silicon surfaces. *Appl. Surf. Sci.* **315**, 516–522 (2014).

10.   Agnihotri, S., Mukherji, S. & Mukherji, S. Immobilized silver nanoparticles enhance

contact killing and show highest efficacy: elucidation of the mechanism of

bactericidal action of silver. *Nanoscale* **5**, 7328 (2013).

11.   O'Brien, T. P. & McCully, M. E. *The Study of plant structure : principles and selected*
695   *methods*. (Termarcarphi Pty. Ltd, 1981).

696   12.   Ma, Y., Sawhney, V. K. & Steeves, T. A. Staining of paraffin-embedded plant material
697   in safranin and fast green without prior removal of the paraffin. *Can. J. Bot.* **71**,
698   996–999 (1993).

699   13.   Fowke, L. C. & Rennie, P. J. Botanical Microtechnique for Plant Cultures. in *Plant*
700   *Cell, Tissue and Organ Culture* 217–228 (Springer Berlin Heidelberg, 1995).
701   doi:10.1007/978-3-642-79048-5_17.

702   14.   Zenchyzen, B. *et al.* Comparative Nectary Morphology across Cleomaceae
703   (Brassicales). *Plants* **12**, 1263 (2023).

704   15.   Kurien, B. T. & Scofield, R. H. Ultrarapid Electrophoretic Transfer of High and Low
705   Molecular Weight Proteins Using Heat. in 181–190 (2009). doi:10.1007/978-1-
706   59745-542-8_20.

707   16.   Bertolini, M. J., Tankersley, D. L. & Schroeder, D. D. Staining and destaining
708   polyacrylamide gels: A comparison of Coomassie blue and fast green protein dyes.
709   *Anal. Biochem.* **71**, 6–13 (1976).

710   17.   Kitin, P., Nakaba, S., Hunt, C. G., Lim, S. & Funada, R. Direct fluorescence imaging of
711   lignocellulosic and suberized cell walls in roots and stems. *AoB Plants* **12**, plaa032
712   (2020).

713   18.   Koga, D., Kusumi, S., Shodo, R., Dan, Y. & Ushiki, T. High-resolution imaging by
714   scanning electron microscopy of semithin sections in correlation with light
715   microscopy. *Microscopy* **64**, 387–394 (2015).

716   19.   Kaplan, D. R. The science of plant morphology: definition, history, and role in
717   modern biology. *Am. J. Bot.* **88**, 1711–1741 (2001).

718   20.   Ashburner, J. & Ridgway, G. R. Tensor-Based Morphometry. in *Brain Mapping* 383–
719   394 (Elsevier, 2015). doi:10.1016/B978-0-12-397025-1.00309-2.

720   21.   Vu, Q. D. *et al.* Methods for Segmentation and Classification of Digital Microscopy
721   Tissue Images. *Front. Bioeng. Biotechnol.* **7**, (2019).

22.   Khoshravesh, R., Stata, M., Adachi, S., Sage, T. L. & Sage, R. F. Evolutionary

Convergence of C4 Photosynthesis: A Case Study in the Nyctaginaceae. *Front. Plant*

Sci. **11**, (2020).

23.   Sage, R. F., Khoshravesh, R. & Sage, T. L. From proto-Kranz to C4 Kranz: building the

bridge to C4 photosynthesis. *J. Exp. Bot.* **65**, 3341–3356 (2014).

24.   Bagasra, O. Protocols for the in situ PCR-amplification and detection of mRNA and

DNA sequences. *Nat. Protoc.* **2**, 2782–95 (2007).

25.   Smith, K., Steinberger, A. & Perloff, W. SAFRANIN AS A STAIN FOR CHROMOSOMES.

Lancet **281**, 1379–1380 (1963).

26.   Jideani, I. A., Takeda, Y. & Hizukuri, S. Structures and physicochemical properties of

starches from acha (Digitaria exilis), iburu (D. iburua), and tamba (Eleusine

coracana). *Cereal Chem.* **73**, 677–685 (1996).

27.   Hubbe, M. A., Chandra, R. P., Dogu, D. & van Velzen, S. T. J. Analytical staining of

cellulosic materials: A Review. *BioResources* **14**, 7387–7464 (2019).

28.   Long, J., Wang, X., Xu, D. & Ding, L. Spectroscopic studies on the interaction

mechanisms of safranin T with herring sperm DNA using acridine orange as a

fluorescence probe. *J. Mol. Recognit.* **27**, 131–137 (2014).

29.   Loix, C. *et al.* Reciprocal Interactions between Cadmium-Induced Cell Wall

Responses and Oxidative Stress in Plants. *Front. Plant Sci.* **8**, (2017).

30.   Chen, H. & Wang, L. Pretreatment Strategies for Biochemical Conversion of

Biomass. in *Technologies for Biochemical Conversion of Biomass* 21–64 (Elsevier,

2017). doi:10.1016/B978-0-12-802417-1.00003-X.

31.   Megías, P. M. Atlas of plant and animal histology. *Universidad De Vigo*

https://mmegias.webs.uvigo.es/02-english/5-celulas/ampliaciones/2-pared-

celular.php (2023).

32.   Deng, Y. *et al.* Investigation of aggregation and assembly of alkali lignin using iodine

as a probe. *Biomacromolecules* **12**, 1116–25 (2011).

33.   Kosma, D. K. *et al.* At <scp>MYB</scp> 41 activates ectopic suberin synthesis and

assembly in multiple plant species and cell types. *Plant J.* **80**, 216–229 (2014).

34.   Scott, F. M. Internal Suberization of Tissues. *Bot. Gaz.* **111**, 378–394 (1950).

35.   Sims, J., Schlögelhofer, P. & Kurzbauer, M.-T. From Microscopy to Nanoscopy:

753   Defining an Arabidopsis thaliana Meiotic Atlas at the Nanometer Scale. *Front. Plant*

754   *Sci.* **12**, (2021).

755   36.   Liao, K. *et al.* A novel in-situ quantitative profiling approach for visualizing changes

756   in lignin and cellulose by stained micrographs. *Carbohydr. Polym.* **297**, 119997

757   (2022).

758   37.   Dai, X. & Shen, L. Advances and Trends in Omics Technology Development. *Front.*

759   *Med.* **9**, (2022).

PLANT MICROTECHNIQUE WITH RESIN - TOWARDS PLANT HISTOLOMICS

Ivan T. Cerritos-Castro[1], Araceli Patrón-Soberano[1], Ana P. Barba de la Rosa[1*].

[1]Molecular Biology Division, Instituto Potosino de Investigación Científica y Tecnológica A.C., IPICYT, San Luis Potosí, México.

FIGURES

The images are high resolution, please zoom in to see the details.

GRAPHICAL ABSTRACT

FIGURE 1 | **Activation and gluing treatments increase the wettability and adhesiveness of glass slides**

a Contact angle photographs after chemical activation and gluing (silanization). **b** Resin sections in the different slides after strong washing. Arrows indicate detachments. Scale bar = 500 μm. **c** Contact angle mean (n=3) ± SD. One-way ANOVA and Holm-Šídák's post-tests were performed, letters indicate significant differences at p<0.01. Activation treatment is indicated at the top. Activation creates hydroxyl groups that make the glass hydrophilic, causing water droplets to spread widely (wettability) and reducing the contact angle. The hydroxyl groups then react with the AES during the gluing process, improving the adhesiveness of the glass.

a Light micrographs of *A. cruentus* leaf paradermal sections stained with different methods. Scale bar = 50 µm. **b** Violin plots (n=30) with median, interquartile range and hue ranges (filled, only for saturation ≤ 15%) observed in cell wall (CW), protoplasm (PR) and background (BK). Kruskal-Wallis ANOVA and Dunn's post-test were performed, differences are shown at p<0.01. HSB graphs correspond to the top micrograph. Hue (H) represents the visible light spectrum on the color wheel, with values ranging from 0 to 360°. Saturation indicates the amount of color and brightness the amount of light, and their values range from 0 to 100%. Low saturation tends to gray and high saturation tends to vivid colors. Low brightness tends to black and high brightness tends to white. The hue depended on the dye, the saturation on the staining performance, and the brightness on the illumination settings of the microscope and camera (standardized).

a JOHANSEN'S METHOD

PARAFIN SECTION (Dewaxing process) → Xylene 1 15 s → Xylene 2 15 s → Absolute EtOH 1 10 s → Absolute EtOH 2 10 s → EtOH 96% 10 s → EtOH 70% 10 s → (Staining process) → Safranin 2 - 48 h → Water 5 s → Picro-alcohol 10 s → Ammoniacal alcohol 10 s → Absolute EtOH 10 s → Fast green 15 s → Used clearing solution 5 s → Fresh clearing solution 5 s → Xylene:Absolute EtOH 99:1 5 s → Xylene 1 5 s → Xylene 2 5 s → Mount in balsam

Safranin: 0.98% Safranin O, 0.98% sodium acetate, 49% methoxyethanol and 23% EtOH.
Pirco-alcohol: 0.5% picric acid (explosive) in 95% ethanol.
Ammoniacal alcohol: 4 to 5 drops of ammonia in 95% EtOH.
Fast green: Solution 1: Saturated solution of fast green fcf in methoxyethanol:absolute EtOH 1:1.
Solution 2: 25% absolute EtOH ans 75% clove oil.
Working solution: Add solution 1 to solution 2 until the "desired coloration" is obtained.
Clearing soluton: 50% clove oil, 25% absolute EtOH and 25% xylene.
Used clearing solution: Clearing solution used until saturation of fast green, diluted with absolute EtOH:Xilene 1:1.

Time required 2 - 48 h

b OUR TRICHROME METHOD

RESIN SECTION → Fast green 2 - 5 min → Water 5 s → Safranin 10 s → Water 5 s → Lugol's iodine 30 S → Water 5 s → Dry at 70°C 1 - 2 min → Mount in acrylate medium

Safranin: 1% Safranin O, 40% MeOH and 10% acetic acid in water.
Fast green: 1% fast green fcf, 40% MeOH and 10% acetic acid in water.
Lugol's iodine: 1% KI, and 0.5% I$_2$ in water.

Time required: 5 - 10 min

FIGURE 3   **We reinvented Johansen's staining for resin sections and made it trichrome**

a Johansen's 1940 staining method with safranin and fast green for paraffin sections. **b** Trichrome staining method with fast green, safranin, and Lugol's iodine for both paraffin and acrylic resin sections. The Johansen method has been a reference in plant histology since 1940. It requires many steps, time, and toxic, explosive, and expensive reagents. Johansen's method did not work on resin sections, so we completely reformulated its dyes and steps, and added Lugol's iodine. The trichrome method greatly reduced the steps, time, toxicity, added a new dye, and worked on both resin and paraffin sections.

a Light micrograph of a cross-section of *A. cruentus* leaf stained with trichrome staining. Arrows indicate the hue range of some structures. **b** Violin plots (n=30) with median, interquartile range, data distribution (circles), and hue range observed in the epidermal wall (E-CW), the mesophyll wall (M-CW), bundle sheath wall (B-CW), bundle sheath proteoplasm (B-PT), mesophyll proteoplasm (M-PT), mesophyll starch granules (M-ST), and bundle sheath starch granules (B-ST). **c** Enlargement of the white box in a. Mauve arrows point to stained and white arrows to unstained granules. **d** TEM of an *A. cruentus* leaf cross section. Mauve arrows point to electrodense granules and white arrows to electrolucent granules. ST, starch; CL, chloroplast; CW, cell wall; MI, mitochondrion; NU, nucleus; and NL, nucleolus. Addition of iodine counterstaining to a tissue stained with Johansen's dyes caused hue changes (veering). Veering occurred mainly in the CW in a tissue-dependent manner. M-CW hardly veered, E-CW veered to reddish, and B-CW veered to reddish and even orange hues, suggesting compositional differences in their cell walls. The proteoplasm, on the other hand, showed no significant veering between tissues. Finally, some ST stained mauve, while others remained white, even whiter than the background itself. This pattern was repeated when the section was examined by TEM, suggesting compositional differences in the starch granules.

LIGHT MICROSCOPY
(TRICHROME STAINING)

ELECTRON MICROSCOPY
(FALSE COLOR)

YOUNG LEAF

MESOPHYLL

PITS

Cell wall (CW)

Cytosol and membrane invaginations

Vacuole (VC)

VC

Nucleolus(NL)

Nucleus (NU)

Chloroplas (CL)

Starch granules (ST)

Cytosol (CY)

Mitochondria (MI)

VASCULAR BUNDLES

SENESCENT LEAF

MESOPHYLL

CW

NU

VC

NL

VC

CL

ST

MI

VASCULAR BUNDLES

FIGURE 5 | Resin-based microtechnique allows light microscopy to resolve structures that would normally require TEM

Digitally isolated bundle sheath cells from light and electron micrographs (not the same cell), from a young (4 weeks), and a senescent (8 weeks) *A. cruentus* leaf. Scale bar= 5 μm. High resolution light micrographs revealed several organelles and ultrastructures that were identified by TEM. The characteristic cell wall thinning of PITS was observed in the bundle sheath-mesophyll cell junctions. Vacuoles were easily identified by their lack of color and texture; however, some cytoplasmic invaginations were detected. All proteoplasm stained turquoise, but saturation differences revealed several organelles and cytosol. Although these structures were not sharply resolved by light microscopy, it was sufficient to identify them and follow their behavior during the senescence process. The senescent cell has lost its cytoplasm, its chloroplasts have lost their centripetal position, contain more starch and are undergoing degradation, but the cell still retains the nucleus and nucleolus. This behavior can be distinguished by both optical and TEM microscopy.

FIGURE 6 — **Trichrome staining and immunostaining revealed the precise localization of RbcL in cells and organelles**

Light micrographs of *A. cruentus* leaf sections immunostained for RbcL. **a** No signal was observed in the negative control. **b** RbcL and background (BK) signal was observed in the anti-RbcL sample. **c** Trichrome staining stained the starch but did not cause significant veering in the cell wall. **d** The combination of immunostaining and trichrome staining summed the colors of both techniques without affecting each other. AE, adaxial epidermis; PM, palisade mesophyll; VB, vascular bundles; BS, bundle sheath; SM, spongy mesophyll; BE, abaxial epidermis. To differentiate the immunostaining signal: a weak signal outside the proteoplasm was considered as background and a strong signal inside the proteoplasm as RbcL. Tetrachrome staining (d) facilitated this differentiation by showing the immunostaining signal and the stained tissue, in contrast to (b) where the tissue lacks contrast. **e-h** Color segmentation of (d) showing separately the cellular elements and the RbcL signal. **i** Violin plots (n=30) with median, interquartile range, and observed color range. Kruskal-Wallis ANOVA and Dunn's post-test were performed, differences are shown with p<0.01. Although there were significant differences, the hue ranges overlapped, so in practice they were adjusted (in Matlab) to intermediate values for color segmentation (**j**). Densitometric analysis showed the highest RbcL content in the BS (**k**), but similar concentrations in all tissues (**l**).

a SECTIONING PLANES

Longitudinal section
Paradermal section
Tertiary veins (not visible)
Secondary vein
Primary vein
Cross section
Leaf blade
Midrib

b PANORAMA OF A CROSS SECTION

Adaxial side (Upper)
LEAF BLADE
TERTIARY VEIN
PRIMARY VEIN
SECONDARY VEIN
100 μm
Abaxial side (Lower)

c LEAF BLADE MORPHOLOGY

Adaxial epidermis
Palisade mesophyll
Bundle sheath
Tertiary vein
Air spaces (Background)
Spongy mesophyll
Abaxial epidermis
50 μm

d CELL MORPHOLOGY

CELL WALL
PROTOPLASM
Nucleus
Vacuole
Chloroplasts
Starch granules
Mitochondrian, peroxisome, cytosol, etc.
10 μm

e COLOR CODE (FALSE COLOR)

Epidermis
Palisade mesophyll cells
Bundle sheath cells
Vascular bundles
Spongy mesophyll cells
Parenchyma cells
Crystal sand idioblast
Druse idioblast

SUPPLEMENTARY FIGURE 1 | Sectional planes of a leaf and panorama of a cross section

a Anatomy of a leaf and sectioning planes. **b** Light micrograph of a cross-section of a primary vein of an *A. cruentus* leaf. The micrograph was colorized and the background digitally removed. **c** Enlargement of the leaf blade. **d** Enlargement of palisade mesophyll and bundle sheath cells. **e** Color coding (false color) to indicate the different tissues. Plant leaves consist of a blade that is supplied by veins that decrease in diameter as they move away from the midrib. If performed through the midrib, a primary or secondary vein, the cross section can show all leaf tissues as it extends from the adaxial to the abaxial epidermis. Tertiary veins are embedded in the center of the leaf and lack parenchyma associated with vascular bundles, responsible for vein protuberance. Tertiary veins are tightly surrounded by a layer of large bundle sheath cells, and this in turn by a single layer of mesophyll cells, forming a wreath shape called "Kranz anatomy". Mesophyll is divided into palisade (receive direct sunlight) and spongy. The plant cell has a cell wall, and its protoplasm contains a large vacuole and other organelles. Kranz anatomy and differentiated mesophyll tissue (palisade and spongy) are not present in all plant leaves. Kranz anatomy enhances carbon fixation in plants known as C_4, and plants lacking it are known as C_3.

a PANORAMA OF A PARADERMAL SECTION

LEAF BLADE

TERTIARY VEIN

PRIMARY VEIN
(Outside the cutting plane)

50 μm

b LEAF BLADE

Spongy mesophyll

Air spaces
Background

Vacuoles
"Empty" spaces within cells

Chloroplasts

Starch grains
Inside chloroplasts

50 μm

c COLOR CODE (FALSE COLOR)

- Epidermis
- Bundle sheath cells
- Vascular bundles
- Spongy mesophyll cells
- Druse idioblast

SUPPLEMENTARY FIGURE 2 | Panorama of a paradermal section

a Light micrograph of a paradermal section of an *A. cruentus* leaf blade. The micrograph was colorized and the background digitally removed. **b** Magnification of a. **c** Color coding (false color) to indicate different tissues. The paradermal section is useful to study the distribution of tertiary veins, a greater number of minority cells such as druse idioblasts, and a greater extent of vascular bundles. Cross section can show all tissues of the leaf, but its extent is limited by the leaf thickness. Paradermal section can show a greater extent of leaf tissues, but the variety is limited. Because the leaf is not flat, the paradermal section usually leaves gaps where the major veins (midrib and primary and secondary veins) used to be. The mid-thickness is the best zone to study because it contains the greatest cellular diversity. A superficial section will show only epidermal cells, a ¾ thickness section will show only mesophyll cells.

a FAST GREEN STAINING FOR 5 MIN + SAFRANIN COUNTERSTAINING FOR

5 s   10 s   20 s   30 s   60 s

b COLOR CHANGES

CELL WALL (CW)   PROTEOPLASM (PT)   STARCH GRAINS (ST)

SUPPLEMENTARY FIGURE 4 | **Fast green-stained sections can be correctly counterstained with safranin for 5 to 10 s**

a Light micrograph of *A. cruentus* leaf sections stained with fast green for 5 min and counterstained with safranin at different times. Scale bar = 50 µm. **b** Violin plots (n=30) with median, interquartile range and color range observed in cell wall (CW), proteoplasm (PT) and starch granules (ST) at different counterstaining times. Safranin has an affinity for cell wall lignin and suberin, whereas fast green has an affinity for proteoplasm proteins. Safranin rapidly and selectively displaced fast green from the cell wall; its area of affinity (counterstaining). However, after 10 s it began to displace fast green from its affinity area: the proteoplasm (replacement).

a Light micrograph of *A. cruentus* leaf sections stained with safranin for 5 min and counterstained with fast green at different times. Scale bar = 50 µm. **b** Violin plots (n=30) with median, interquartile range, and range of color observed in cell wall (CW), proteoplasm (PT), and starch granules (ST) at different counterstaining times. Safranin has an affinity for cell wall lignin and suberin, whereas fast green has an affinity for proteoplasm proteins. Fast green slowly displaced safranin from the proteoplasm, its affinity area, but in an unsaturated and uneven manner. After 20 s, it also began to displace safranin from the cell wall, producing a poorly saturated result. This order of staining is the original order of Johansen's method. However, only the reverse order gave us good results in resin sections.

SUPPLEMENTARY FIGURE 6 | **Iodine counterstaining produces hue veering in stained sections**

a Light micrograph of *A. cruentus* leaf sections stained with fast green for 5 min, counterstained with safranin for 10 s, and with iodine at different times. Scale bar = 50 μm. **b** Violin plots (n=30) with median, interquartile range, and range of color observed in cell wall (CW), proteoplasm (PT), and starch granules (ST) at different counterstaining times. Iodine has an affinity for starch granules and produces color veering mainly in safranin-stained cell walls. Iodine staining, alone or as a counterstain, produced random intensities of starch granule staining: null, intermediate, or intense. That was independent of time and even between sections of the same sample on the same slide. However, when it occurred, the staining intensity was always homogeneous along the same section. So far, we could not identify the reason for this, and we had to perform many replicates to obtain the desired result. In these sections, the starch granules were not stained. However, a positive time-dependent cell wall veining was observed, as well as a small and random veering in the proteoplasm.

a Light micrograph of a paradermal section of an *A. cruentus* leaf. **b-i** Magnification of the artifacts listed in a. **b** Wrinkles are the most common and difficult artifact to avoid; they can cause staining problems and excessive background in immunostaining (**1-3**). **c** Bubbles are small wrinkles that can be recognized by their birefringence (**4,5**). However, they are not always birefringent and can be confused with cellular elements. (**6**). **d** Dirt causes spots, wrinkles and bubbles in the section (**7-9**). Cleanliness is of paramount importance when handling microscopy specimens. **e** Dyes may leave residues on the specimen if not filtered out, or may form precipitates when interacting with other dyes (**10-12**). **f** Inhomogeneous staining is often caused by wrinkles, bubbles, or dirt. However, as with iodine, sometimes there are no clear reasons for its occurrence. In areas **13-15**, there are no wrinkles, bubbles, or dirt, and the section is properly adhered to the glass. However, from there (white line) and upwards, the staining was not homogeneous. **g** Even an area detached from the glass (**16**) was stained homogeneously. **h** Resin sections are prone to tearing, deforming the original histology (**17**). **i** Some artifacts and even the specimen itself may show different colors depending on the focal point (**18**), due to birefringence or chromatic aberration, respectively. Therefore, it is important to define a focusing criterion for all sections.

Palisade and spongy mesophyll cells digitally isolated from light and electron micrographs (not the same cell) of a young (4 weeks old) and a senescent (8 weeks old) *A. cruentus* leaf. Color in light micrograph is from trichrome staining, and in TEM is false color mimicking trichrome staining colors. Scale bar = 5 *µ*m. High-resolution light micrographs revealed several organelles and ultrastructures identified by TEM. The vacuole, identified by its lack of color and texture, covers large areas in the cells. The proteoplasm stained turquoise, but differences in saturation revealed the presence of various organelles identified by TEM. Larger and/or more numerous organelles are usually always present in the micrographs because they are more likely to be in the plane of the section. The opposite is true for small and/or few organelles, such as the nucleus, which are not always observed in the micrographs. The young leaf has a defined cytosol, while the senescent leaf has lost it and its organelles are reduced in size. On the other hand, different amounts of starch granules were observed inside the mesophyll chloroplast.

SUPPLEMENTARY FIGURE 9 | **Resin-based microtechnique resolution: epidermis and vascular bundle cells**

Epidermis and vascular bundle cells digitally isolated from light and electron micrographs (not the same cell) of a young (4 weeks old) A. cruentus leaf. Color in light micrograph is from trichrome staining, and in TEM is false color mimicking trichrome staining colors. Scale bar = 5 µm. Vacuole (VC), chloroplast (CL), amyloplasts (AL), starch granule (ST), nucleus (NU), nucleolus (NL), cytosol (CY) and mitochondrion (MI). High-resolution light micrographs revealed several organelles and ultrastructures identified by TEM. The epidermis and vascular bundles are formed by different cells. The epidermis consists of epidermal cells, stomata and trichomes. The vascular bundles are formed by the conducting cells xylem and phloem and a variety of accessory cells. The epidermal cells are almost completely devoid of proteoplasm; they mainly retain some traces of amyloplasts with starch granules. Stomata, on the other hand, contain organelle-nourished proteoplasm like a mesophyll cell, as well as significant amounts of starch granules. Xylem and phloem cells are usually observed as hollow and their accessory cells with proteoplasm nourished by organelles, cytosol and starch. Vacuoles and chloroplasts or alternatively amyloplasts are observed in almost all leaf cells.

a PARADERMAL SECTION

b DIFFERENTIAL CELL WALL STAINING

HSB 330-355
HSB 355-20
≈ 0.3 - 2 μm
≈ 1 - 4 μm
HSB 1-26
Stomata
Epidermis
CL
ST
NU
NL
NU

c DIFFERENTIAL STAINING OF TRACHEIDS

HSB 300-335
Vascular bundles
ST CL NL
NU
HSB 5-25
① ②
CL ST NU
NL
Bundle sheath

d PROTEIN BODIES WITHIN THE TRACHEIDS

Vascular bundles

e CELL WALL METABOLISM

④ ④ ④
Vascular bundles
④
③
③
Bundle sheath
Spongy mesophyll

100 μm

SUPPLEMENTARY FIGURE 10 | Intense iodine counterstaining reveals cell wall composition and ultrastructures

a Paradermal section of an *A. cruentus* leaf stained with trichrome stain. **b - e** Magnifications of areas marked in a. Scale bars = 10 μm. Vacuole (VC), chloroplast (CL), starch granule (ST), nucleus (NU), nucleolus (NL), and cytosol (CY). a Iodine counterstaining was intense; it stained starch granules, almost all cell walls veered, and ultrastructural details were further contrasted. **b** Epidermal cell walls showed two layers: an internal orange wall and an external magenta wall. Also, the external magenta wall showed a saturation gradient, from inner saturated to outer colorless, ending in a saturated boundary. **c** Almost all cell walls veered to orange, but xylem tracheids remained magenta as the outer layer of the epidermal wall, and some protein bodies were observed inside them (**d**). Various ultrastructural details were observed such as: chloroplasts with granular texture (**1**), membranes inside vacuoles (**2**), rod-shaped organelles (**3**), and vesicular trafficking of cell wall material (**e, 4**).

Polynucleolarity

Unknown

Atypical wall thickenings

PITS and plasmodesmata

Decreased protein concentration

Decreased wall thickness

Starch granule appearance in PM

Increased starch granules in BS

PM became rounded

Increased cell layers in PM

Disappearance of the cytosol

BS CL lost centripetal position

≈200 μm

SUPPLEMENTARY FIGURE 11 | Histolomic analysis – Visual analysis

Light micrographs of cross sections of young (4 weeks) and senescent (8 weeks) leaves of *A. cruentus* stained with trichrome staining. Scale bar = 50 μm. Adaxial epidermis (AE), palisade mesophyll (PM), bundle sheath (BS), vascular bundles (VB), spongy mesophyll (SM) and, abaxial epidermis (BE). Cell wall (CW), vacuole (VC), chloroplast (CL), starch granule (ST), nucleus (NU), nucleolus (NL), cytosol (CY) and mitochondrion (MI). Visual analysis is the first step in histolomic analysis, the main aim of which is to identify tissues, general morphology, organelles and some ultrastructural details. It helps to define the boundaries for micrograph segmentation, relevant features for data analysis, as well as unique traits to highlight. The *A. cruentus* leaf has an average thickness of about 200 μm and 8 to 10 cells (not including VB), a clear kranz anatomy and a clear differentiation between PM and SM. BS, VB, PM and SM are well nourished with organelles, mainly chloroplasts, while the epidermis barely has proteoplasm. Several ultrastructural details were observed: Some nuclei with apparent polynucleolarity (**1**), unknown proteinaceous structures (perhaps mitochondrion) with atypical size and/or location (**2**), atypical wall thickenings in SM (**3**) and PITS-plasmodesmata between PM and BS (**4**). Also, compared to the young leaf, the senescent leaf had reduced protein concentration (**5**), reduced cell wall thickness (**6**), increased ST in PM (**7**) and BS (**8**), PM became rounded (**9**) and had more cell layers (**10**), the cytosol diapered (**11**) and the CL of BS lost its centripetal position (**12**).

SUPPLEMENTARY FIGURE 12 | Histolomic analysis — Semi-automated segmentation

Light micrograph of a trichrome-stained cross section of a young amaranth leaf and the layers obtained after color-based segmentation of its tissue components. Scale bar = 50 μm. The micrograph was digitally segmented in MatLab using the "color threshold" app and the "segmentation code" settings to isolate the stained and unstained tissue components. Thin and lightly stained mesophyll cell walls lacked continuity (**1**), as did the outer epidermal walls (**2**). Differences in saturation of the proteoplasm allowed visual identification of organelles and cytosol, as well as secondary segmentation into low-saturated proteoplasm (**3**), such as cytosol and nucleus, and high-saturated proteoplasm, such as chloroplasts and mitochondria (**4**). Iodine counterstaining was null in the starch granules of this section (**5**), so they were segmented along with other unstained components such as vacuoles (**6**), intercellular air spaces (**7**), and resin background (**8**). Semi-automatic segmentation is convenient, but fails in thin and poorly stained cell walls, and also requires further manual segmentation to sub-segment the layers per tissue.

CELL WALLS	PROTEOPLASM	STARCH GRAINS	VACUOLES

YOUNG LEAF

SENESCENT LEAF

SUPPLEMENTARY FIGURE 13 | **Histolomic analysis - Manual segmentation of tissue components of an entire section**

Manually segmented layers form light micrographs of a trichrome-stained cross section of a young and senescent amaranth leaf. Scale bar = 50 μm. Manual segmentation is time-consuming, but necessary to delineate thin and/or poorly stained cell walls that are lost in semi-automated segmentation. Each layer is more than just an attractive image; it contains size, perimeter, area and shape information that can be analyzed for morphometric measurements, as well as color intensity by area (densitometry) information that can be analyzed for compositional measurements. Clear and continuous cell boundaries are important for further morphometric measurements. In addition, manual segmentation is necessary to separate layers per tissue for independent study.

SUPPLEMENTARY FIGURE 14 Histolomic analysis - Manual segmentation of tissue components per tissue

Manually segmented layers form light micrographs of a trichrome-stained cross section of a young and senescent amaranth leaf. Scale bar = 50 μm. Histolomics aims at the individualized phenotypic study of the tissues of an organ. We visualize cell-by-cell segmentation and analysis as the future of histolomics, which is currently very laborious to perform manually.

MORPHOMETRY

a RELATIVE AREA

$$\%Area_{tissue} = \frac{Area_{tissue}}{Area_{total}} * 100\%$$

b AREA PER SINGLE CELL

c # OF CELLS

d CIRCULARITY

$$Circularity = 4\pi \frac{Area_{cell}}{(Perimeter_{cell})^2}$$

A value of 1.0 indicates a perfect circle. A value of 0 indicates an elongated polygon.

e CELL WALL THICKNESS

f PROTEOPLASM CONTENT

$$Proteoplasm\ content = \frac{Area_{proteoplasm}}{Area_{tissue}} * 100\%$$

g CELL WALL CONTENT

$$Cell\ wall\ content = \frac{Area_{cell\ wall}}{Area_{tissue}} * 100\%$$

h STARCH GRANULES CONTENT

$$Starch\ granules\ content = \frac{Area_{starch\ granules}}{Area_{tissue}} * 100\%$$

i VACUOLE CONTENT

$$Vacuole\ content = \frac{Area_{vacuoles}}{Area_{tissue}} * 100\%$$

COMPOSITION

j PROTEIN CONTENT

k LIGNIN CONTENT

l PROTEIN CONCENTRATION

$$Protein\ concentration = \frac{Integrated\ gray\ value_{protoplasm}}{Area_{protoplasm}}$$

m LIGNIN CONCENTRATION

$$Lignin\ concentration = \frac{Integrated\ gray\ value_{cell\ wall}}{Area_{cell\ wall}}$$

SUPPLEMENTARY FIGURE 15 | **Histolomic analysis – Data analysis**

Morphometric and compositional analysis of a trichrome-stained cross section of a young and senescent amaranth leaf. Adaxial epidermis (AE), abaxial epidermis (BE), adaxial inner epidermis (IAE), adaxial outer epidermis (EAE), abaxial inner epidermis (IBE), abaxial outer epidermis (EBE), palisade mesophyll (PM), spongy mesophyll (SM), bundle sheath (BS), vascular bundles (VB), air spaces (AR), young leaf (Y), and senescent leaf (S). For measurements with replicates, a t-test was performed comparing young vs. senescent leaf, differences are shown with p<0.05. Histolomics needs its own features and formulas for in-depth study of morphometric and compositional traits beyond the naked eye. We propose three types of histolomic features: primary, derived, and composite. Primary features are individual measurements on the layers, such as cell count, thickness, areas, perimeters and total protein/lignin content. Derived features are ratios or combinations of two or more primary features, such as protein/lignin concentrations and organelles content. Composite features are combinations of different derived and primary features, can be weighted and are useful for quantitative comparison of complex tissue behavior (Supplementary Fig. 24). In this figure, we present primary and derived features and their formula (when apply). Their values gave a quantitative view of the differences observed in the visual analysis of the senescent leaf with respect to the young leaf: decreased protein concentration (**l**), decreased cell wall thickness (**e**), more starch granules in the PM (**h**), the PM became rounded (**d**), and increased layers of the PM (**c**). Other important features related to kranz anatomy were observed between BS, PM and SM. BS was by far the largest tissue area (**a**), had larger cells (**b**), had less vacuole content (**i**), and had the largest content of proteoplasm (**f**), protein (**j**), and starch granules (**h**). The bundle sheath also had slightly higher lignin concentration (m) and had thicker cell walls (**g, e**). Protein and lignin measurements are semi-quantitative as they are not related to a calibration curve (**j-m**). However, in comparative studies, the differences between quantities are usually more relevant than the absolute value itself.

SEMI-AUTOMATICALLY SEGMENTED LAYERS OF AN INTENSELY IODINE COUNTERSTAINED TISSUE

TRICHROME STAINING — **INTERNAL WALL** — **EXTERNAL WALL**

STARCH — **PROTEOPLASM** — **SEGMENTATION CODE**

HUE (°)

EXTERNAL WALL
290-350

INTERNAL WALL
351-60

STARCH
252-289

COLOR WHEEL
Segmentation code
Saturation: 10 - 100%
Brightness: 0 - 100%

PROTEOPLASM
147-251

SUPPLEMENTARY FIGURE 16 | Intense iodine counterstaining allows further color (composition) segmentation

Semi-automatic color segmentation of a paradermal section of an *A. cruentus* leaf stained with trichrome staining. Scale bar = 100 *µ*m. The color wheel conveniently shows the hue range assigned to each tissue component, as well as the saturation and brightness settings for segmentation. Intense iodine counterstaining veered almost all cell walls to orange, except for epidermal walls, which had an internal orange wall and an external magenta wall. The inner wall showed a wide range of reddish, orange, and yellow hues, while the outer wall showed magenta to reddish hues. Color segmentation showed that although the majority of the walls veered toward orange, they also retained some magenta hues. This suggests the presence of two layers of different composition on the cell wall. On the other hand, intense iodine counterstaining allowed segmentation of starch granules and did not produce significant veering in the hue range of proteoplasm as occurred with cell walls. However, the color range of starch granules slightly overlapped with the color range of the outer wall, making the starch granules and outer wall layers "impure". Nevertheless, intense iodine counterstaining differentiated the cell wall color and improved the staining of the thin mesophyll cell walls, thus extending the molecular composition information and improving boundary detection.

SUPPLEMENTARY FIGURE 17 | Histolome of *Arabidopsis thaliana* leaf

a Cross section of a trichrome-stained *Arabidopsis thaliana* leaf. **b** Magnification of the squared area in a. **c** Derived (morphometry and composition) and composite (C_4 Kranz level) features of the tissue (same scales for all plant species). **d** Derived features of the vascular bundle proteinaceous cell (VBPC). Adaxial epidermis (AE), abaxial epidermis (BE), palisade mesophyll (PM), spongy mesophyll (SM), bundle sheath (BS), vascular bundles (VB), air spaces (AR). Cell wall (CW), vacuole (VC), chloroplast (CL), starch granule (ST), nucleus (NU), nucleolus (NL), cytosol (CY) and mitochondrion (MI). Although not all plants have a developed BS or differentiated PM and SM, we have highlighted these tissues for comparison purposes. We have increased the brightness of the BS and decreased the brightness of the SM to distinguish them from the other tissues. The leaf of *A. thaliana* had a small BS tightly surrounded by two layers of PM and three layers of SM: not wreath-shaped. Also, all their cells had low proteoplasmic content (c), except for a unique VBPC in the VB with a smooth and well-nourished proteoplasm, covering almost the entire cell cytoplasm (**1**). One VBPC was detected in the VB of the primary vein and two in the secondary vein and midrib, but with less proteoplasm and protein concentration than in the primary vein (a). VBPC had up to 26% of the protein content of the VB in the primary vein (d). On the other hand, some unknown ultrastructural details were observed: organelles with purple dots (**2**), extracellular proteoplasm (**3**), dark purple and rod-shaped organelles in VB (**4**), intermittent coloring in VB cell walls (**5**) and apparently broken cells (**6**).

a

Midrib

Parenchyma

100 μm

b LEAF MORPHOLOGY

⑤

①

⑤

Secondary vein

Section damage

Primary vein

100 μm

c LEAF BLADE MORPHOLOGY

AE

PM

③ ②

VB

BS

① IM ③ ③

③ ④

⑤ AR

NU?

SM

CL Starch CY

CW MI? VC

BE

≈160 μm

50 μm

136 μm

d C₃

C₄ KRANZ LEVEL

MORPHOMETRY

RELATIVE AREA

PROTEOPLASM   CELL WALL

COMPOSITION

[PROTEIN]   [LIGNIN]

SY

SUPPLEMENTARY FIGURE 18 | Histolome of *Glycine max* leaf

Cross section of a trichrome-stained *Glycine max* leaf midrib (**a**) and leaf blade (**b**). **c** Magnification of the squared area in b. **d** Derived (morphometry and composition) and composite (C₄ Kranz level) features of the tissue (same scales for all plant species). The leaf of *G. max* had a small BS loosely surrounded by a layer of PM and two layers of SM: not wreath-shaped. We also observed a rod-like mesophyll similar to PM but perpendicular to it, which we called intermediate mesophyll (IM, **1**). Intermediate mesophyll was observed along the blade as well as other rounded PM cells (**2**). Intermediate mesophyll was also compositionally different, as it did not accumulate starch granules like BS and (**2**), and unlike PM and SM (**3**). In addition, despite being a C₃ plant, BS had centrifugally oriented CL; a C₄ characteristic (**4**). Finally, the most prominent morphological feature was the large number of intercellular air spaces (**5**).

SUPPLEMENTARY FIGURE 19 | Histolome of *Triticum aestivum* leaf

Cross section of a trichrome-stained *T. aestivum* leaf midrib (**a**) and leaf blade (**b**). **c** Magnification of the squared area in b. **d** Derived (morphometry and composition) and composite (C_4 Kranz level) features of the tissue (same scales for all plant species). The leaf of *T. aestivum* had a large BS tightly surrounded by a layer of undifferentiated PM and two layers of SM: wreath-shaped. Other unique features were observed as: magenta-stained granules in turquoise-stained nuclei (**1**), intermittent magenta-purple staining in VB walls (**2**), VB barely had proteoplasm (**3**), and the protein content of some organelles was polarized (**4**).

a

100 µm   Midrib   Parenchyma

b   LEAF MORPHOLOGY

100 µm   Secondary vein   Primary vein

d

C_4

MORPHOMETRY

RELATIVE AREA

PROTEOPLASM   CELL WALL

COMPOSITION

[PROTEIN]   [LIGNIN]

C_4 KRANZ LEVEL

MZ

c   LEAF BLADE MORPHOLOGY

AE   PM   Dust   NU   CL No starch   CW   VC   MI?   BS   VB   SM   BE   AR

≈110 µm

50 µm

164 µm

SUPPLEMENTARY FIGURE 20 | Histolome of _Zea mays_ leaf

Cross section of a trichrome-stained _Z. mays_ leaf midrib (**a**) and leaf blade (**b**). **c** Magnification of the squared area in b. **d** Derived (morphometry and composition) and composite (C_4 Kranz level) features of the tissue (same scales for all plant species). The leaf of _Z. mays_ had a very large BS tightly surrounded by a layer of undifferentiated PM and a layer of SM: wreath-shaped. SM between VB had an elongated shape with sharp edges (**1**), and BS chloroplasts had a centripetal position (**2**). On the other hand, nuclei had granules (**3**), VB had little proteoplasm (**4**), and there were very few intercellular air spaces (**5**).

a Primary vein — Parenchyma — 100 μm

b LEAF MORPHOLOGY — 100 μm — Tertiary vein

c LEAF BLADE MORPHOLOGY

d C₄ — MORPHOMETRY — RELATIVE AREA — PROTEOPLASM — CELL WALL — COMPOSITION — [PROTEIN] — [LIGNIN]

C₄ KRANZ LEVEL

AE — PM — BS — VB — SM — AR — BE — ≈200 μm — 50 μm — 92 μm

SUPPLEMENTARY FIGURE 21 | Histolome of *Amaranthus cruentus* leaf

Cross section of a trichrome-stained *A. cruentus* leaf midrib (**a**) and leaf blade (**b**). **c** Magnification of the squared area in b. **d** Derived (morphometry and composition) and composite (C₄ Kranz level) features of the tissue (same scales for all plant species). The leaf of *A. cruentus* had a large BS tightly surrounded by a layer of differentiated PM and two layers off SM: wreath-shaped. Additional details are reviewed in Supplementary Figure 11.

HISTOLOMIC DATA ANALYSIS - LEAF SPECIES COMPOSITION

PROTEIN CONTENT

a PALISADE MESOPHYLL **b** BUNDLE SHEATH **c** SPONGY MESOPHYLL

PROTEIN CONCENTRATION

d **e** **f**

LIGNIN CONTENT

g **h** **i**

LIGNIN CONCENTRATION

j **k** **l**

SUPPLEMENTARY FIGURE 23 | Histolomic compositional analysis of different plant leaves.

Morphometric analysis of *A. thaliana* (AT), *G. max* (SY), *T. aestivum* (WH), *Z. mays* (MZ), and *A. cruentus* (AM) leaves. Results are presented in ascending order of C_4 Kranz level (AT, SY, WH, MZ, and AM), with the left being "most C_3" and the right being "most C_4". We put special emphasis on the mainly photosynthetic tissues: PM, SM and BS, which define the wreath shape and the global C_4 kranz anatomy. Some trends related to the increasing C_4 anatomy were observed: decreased protein content in PM and SM (**a, c**) and increased in BS (**b**), while protein concentration showed no trend (**d-f**). In contrast, lignin content and concentration showed no trend (**g - l**). However, AM tissues had the highest lignin concentration.

a

#	Primary/ Derived Feature	Feature Score	Primary Features	AT Measurements	AT Absolute value	AT Weighted score	SY Measurements	SY Absolute value	SY Weighted score	WH Measurements	WH Absolute value	WH Weighted score	MZ Measurements	MZ Absolute value	MZ Weighted score	AM Measurements	AM Absolute value	AM Weighted score	Maximum absolute value
1	BS Size ratio	20	BS area (μm^2) / M area (μm^2)	8026 / 59276	0.12	5.4	1266 / 7395	0.15	6.6	4521 / 12201	0.27	12.2	3120 / 3921	0.44	20.0	4477 / 6336	0.41	18.7	0.44
2	BS organelle size ratio	20	BS PT area (μm^2) / M PT area (μm^2)	246 / 2843	0.08	2.9	277 / 2689	0.09	3.5	400 / 1993	0.17	6.2	798 / 675	0.54	20.0	1664 / 1587	0.51	18.9	0.54
3	Position of BS mitochondria	15	YES (1)/NO (0)	NO	0	0	NO	0	0	INTERMEDIATE	0.5	7.5	NO	0	0.0	YES	1	15	1
4	Position of BS chloroplast	15	YES (1)/NO (0)	NO	0	0	YES	1	15	NO	0	0	YES	1	15	YES	1	15	1
5	Wreath shape	15	YES (1)/NO (0)	NO	0	0	NO	0	0	YES	1	15	YES	1	15	YES	1	15	1
6	BS CW size ratio	5	BS CW area (μm^2) / M CW area (μm^2)	527 / 2289	0.19	2.4	125 / 731	0.15	1.9	222 / 616	0.26	3.4	206 / 430	0.32	4.1	453 / 709	0.39	5	0.39
7	BS CW lignin ratio	5	BS Lignin (IGV/area) / M Lignin (IGV/area)	5185 / 2981	1.74	3.7	6577 / 2772	2.37	5.0	4082 / 2648	1.54	3.2	4980 / 2504	1.99	4.2	8018 / 7885	1.02	2.1	2.37
8	BS air insulation	5	BS-M Contact perimeter (μm) / BS-AR Contact perimeter (μm)	493 / 77	0.86	5.0	62 / 140	0.31	1.8	153 / 246	0.38	2.2	175 / 114	0.61	3.5	228 / 61	0.79	4.6	0.86
	Total	100	C$_4$ Kranz level		3.0	19.4		4.1	33.7		4.1	49.7		5.9	81.8		6.1	94.3	

b

$$BS\ size\ ratio = \frac{Area_{BS}}{Area_{BS} + Area_M}$$

$$BS\ organelle\ size\ ratio = \frac{Area_{BS\ PT}}{Area_{BS\ PT} + Area_{M\ PT}}$$

$$BS\ CW\ size\ ratio = \frac{Area_{BS\ cw}}{Area_{BS\ cw} + Area_{M\ cw}}$$

$$BS\ CW\ lignin\ ratio = \frac{Lignin\ concentration_{BS}}{Lignin\ concentratio_M}$$

$$BS\ air\ insulation = \frac{Contact\ P_{BS-M}}{Contact\ P_{BS-M} + Contact\ P_{BS-AR}}$$

$$Weighted\ score = \frac{Absolute\ value}{Maximun\ absolute\ value} * Trait\ score$$

SUPPLEMENTARY FIGURE 24 | Calculation of the C$_4$ kranz anatomy level

Calculation of the C$_4$ kranz level for *A. thaliana* (AT), *G. max* (SY), *T. aestivum* (WH), *Z. mays* (MZ), and *A. cruentus* (AM) leaves (**a**) and the formulas used (**b**). Bundle sheath (BS), Mesophyll (M), Cell Wall (CW), Proteoplasm (PT), Air Spaces (AR), Mitochondrion (MI), Chloroplast (CL) and Integrated Gray Value (IGV). Because individual comparisons of primary and derived features did not yield clear conclusions, they must be combined into a single value that allows for simple and quantitative comparisons. The primary features describe the individual characteristics of each tissue, such as: position of BS/mitochondria/chloroplast, which describes whether mitochondria/chloroplast are located centripetally (MI and CL) or centrifugally (CL) in BS, and wreath shape, which describes whether BS are large and tightly surrounded by a single layer of PM and SM (sometimes two PM layers). Derived features compare the primary features of two tissues, such as BS size ratio, which describes the size of BS tissue compared to M tissue. The same logic is used to compare organelle size, CW, lignin concentration, and air insulation. This set of morphometric and compositional features constitutes the C$_4$ kranz anatomy[1]. However, from our perspective, each of these features has a different degree of importance, so we have assigned them a score to calculate a weighted sum that reflects the overall "C$_4$ kranz level", a composite feature. It should be noted that the "C$_4$ Kranz level" is a histolomically based measurement and is used as an example of a histolomic application. It does not pretend to be an exhaustive description of C$_4$ photosynthesis, as it also implies the specific accumulation of some key enzymes such as Glycine decarboxylase complex in the mitochondrion; PEP carboxylase in the M cytosol, and RuBisCO and NADP-malic enzyme in the BS chloroplasts, and their contribution is not weighted here[1].

[1]Sage, R. F., Khoshravesh, R. & Sage, T. L. From proto-Kranz to C4 Kranz: building the bridge to C$_4$ photosynthesis. J. Exp. Bot. 65, 3341–3356 (2014)

SUPPLEMENTARY FIGURE 25 | Histolomic visual analysis of *A. cruentus* seed germination

a Polarization micrograph of amaranth seed at different germination stages. Days after sowing are indicated. **b** Light micrograph of a whole seed section (0 days), and in germination (1.7 days). **c** Magnification of the white boxed area in b, and a cross section of seedling cotyledon. The sections were stained with trichrome staining. Adaxial epidermis (AE), abaxial epidermis (BE), palisade mesophyll (PM), spongy mesophyll (SM), bundle sheath (BS), vascular bundles (VB), air spaces (AR). Cell wall (CW), vacuole (VC), chloroplast (CL), starch granule (ST), nucleus (NU), nucleolus (NL), cytosol (CY) and mitochondrion (MI). The resin-based microtechnique worked on amaranth seeds and seedlings. The embryo (cotyledons, hypocotyl and radicle) was stained turquoise, the testa magenta and the perisperm was not stained. The embryo was composed of fusiform cells whose proteoplasm contains mainly protein bodies, nucleus and cytosol (**1**). Their CW stained weakly, making them almost indistinguishable (**2**). After germination, the fusiform cells began to degrade their protein bodies (**3**), thickened their cell wall (**4**) and differentiated into conducting tissues such as tracheids (**5**). After three days, the cotyledon showed all the basic tissues of a leaf: AE, PM, BS, VB, SM, BE and some AR. Also, the protein bodies disappeared, but smaller and denser ones appeared (**6**), starch granules accumulated in almost all tissues (**7**), the epidermis formed vacuoles (**8**), and the BE formed a thicker outer wall than the AE outer wall.

SUPPLEMENTARY FIGURE 26 | Histolomic visual analysis of *A. cruentus* cotyledon development

a Polarization micrograph of amaranth cotyledons at different developmental stages. Days after sowing are indicated. **b** Light micrograph of a cross sections of the cotyledon in a. The sections were stained with trichrome staining. Adaxial epidermis (AE), abaxial epidermis (BE), palisade mesophyll (PM), spongy mesophyll (SM), bundle sheath (BS), vascular bundles (VB), air spaces (AR). Cell wall (CW), vacuole (VC), chloroplast (CL), starch granule (ST), nucleus (NU), nucleolus (NL), cytosol (CY) and mitochondrion (MI). After 3.5 days, all cells showed a clear development of organelles. Epidermal cells were the first to produce VC (**1**), followed by SM and PM (**2**), and finally BS (**3**). All cells except VB formed starch granules (**4**), the BS CL began to position themselves centripetally (**5**), and some PM began to acquire their characteristic rod-like shape (**6**). After 4.9 days, the cotyledon had a mature leaf-like morphology, all its ST disappeared (**7**) and the epidermal cells lost almost all their proteoplasm (**8**). Finally, after 7.9 days, the true leaf began to form, the cotyledon entered senescence and its chloroplasts took on a slender, curved shape (**9**).

50 μm

SUPPLEMENTARY FIGURE 27 | **Trichrome staining also works on paraffin-embedded sections**

Light micrograph of a paraffin-embedded *A. cruentus* leaf section stained with trichrome staining. The section was not deparaffinized. Trichrome staining also worked on paraffin-embedded tissue. This section is 10 um thick, 20 times thicker than the resin sections. Therefore, the staining is more intense and the color of the starch granules masks the green of the chloroplasts. In addition, the cells of PM and SM contain important extensions of the cell wall, which is why they are mostly observed in a reddish color.