De novo assembly of complete *Plasmodium falciparum* isolate genomes using PacBio HiFi sequencing technology

Prince B. Nyarko[1]*, Mathieu Quenu[1]*, Marc-Antoine Guery[1], Camille Cohen[1], William L. Hamilton[2,3,4], Sophia T. Girgis[4], Alex Makunin[5], Shane A. McCarthy[5], Mara K. N. Lawniczak[5], Antoine Claessens[1]

1 LPHI, CNRS, INSERM, University of Montpellier, Montpellier, France

2. University of Cambridge, Department of Medicine, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0QQ, United Kingdom

3. Cambridge University Hospitals NHS Foundation Trust, Cambridge Biomedical Campus, Hills Road, Cambridge CB2 0QQ, United Kingdom

4. Wellcome Sanger Institute, Wellcome Trust Genome Campus, Hinxton, CB10 1RQ, United Kingdom

5. Tree of Life, Wellcome Sanger Institute, Hinxton CB10 1SA, UK.

* Equal contributors

Correspondence antoine.claessens@umontpellier.fr

Abstract

Plasmodium falciparum possesses a highly structured genome with extensive sequence diversity concentrated in Variant Surface Antigen (VSA) families. These genes—*var*, *rif*, and *stevor*—play key roles in immune evasion and pathogenesis and are difficult to assemble using short-read sequencing technologies. Here, we applied PacBio HiFi long-read sequencing to generate high-quality *de novo* genome assemblies from 43 *P. falciparum* parasite cultures originating from community cases in The Gambia. Parasites were culture-adapted, cloned by limiting dilution where possible, and sequenced using high molecular weight DNA extracts. Assemblies from single-genotype lineages were constructed using hifiasm, producing complete chromosomal-length scaffolds with high base accuracy without requiring short-read polishing. We recovered full repertoires of *var*, *rif*, and *stevor* genes and classified them into known subgroups. Together, our results demonstrate that PacBio HiFi sequencing enables accurate assembly of complex *P. falciparum* genomes from natural infections. This work provides a valuable genomic resource for future studies of parasite evolution, transmission dynamics, and antigenic diversity, and suggests that VSA repertoires can serve as reliable proxies of genetic relatedness across infections.

Introduction

Malaria remains a global leading cause of morbidity and mortality, affecting millions and causing hundreds of thousands of deaths annually (*World Malaria Report 2024*, n.d.) The disease is caused by unicellular apicomplexan parasites of the genus *Plasmodium*, with *P. falciparum* responsible for most of the severe cases and fatalities. The first *P. falciparum* genome sequenced was the laboratory strain 3D7 in 2002 (Gardner et al., 2002). Since then, numerous studies have increased the number of *P. falciparum* genome sequences.

Most genome sequencing efforts have relied on Illumina short-read technologies (MalariaGen., 2021), but long-read platforms, such as Oxford Nanopore Technology or PacBio's single-molecule real-time (SMRT) sequencing have been increasingly applied to *Plasmodium* genomes (Delandre et al., 2024; Holzschuh et al., 2024; Vembar et al., 2016). Historically, the lower base-calling accuracy of long-read platforms necessitated complementary short-read sequencing to achieve assemblies that were both accurate and complete (Higgins et al., 2024). However, advances in PacBio HiFi (High Fidelity) sequencing using circular consensus sequencing (CCS) now promise long and highly accurate reads (Hon et al., 2020; Wenger et al., 2019). While this technology has been used to produce high-quality assemblies for some *Plasmodium* species (*P. yoelii* - Godin et al., 2023), it has not yet been widely applied to *P. falciparum* field isolates.

The genome of *P. falciparum* consists of a conserved 'core genome' and a hypervariable non-core genome, the latter harboring large multigene families encoding Variant Surface Antigens (VSAs) (Otto et al., 2018). The three VSA families - *var*, *rif* and *stevor* - are important for pathogenesis and immune evasion and are located in subtelomeric and internal gene clusters (Harrison et al., 2020; Jespersen et al., 2016; Rowe et al., 2009; Sakoguchi et al., 2025). Their sequences are highly polymorphic with each parasite isolate typically having a unique set of VSAs whose genetic diversity can be generated by non-homologous recombination events (Claessens et al., 2014). Their classification into biologically relevant subgroups (e.g., *var* A, B, and C; *rif* A and B) provides insights into parasite biology and virulence. For example, group A *var* genes are implicated in cerebral malaria (Claessens et al., 2012). Analyses of VSA repertoires in field isolates also bear epidemiological relevance, enabling lineage tracking across host infections (Day et al., 2025; Tonkin-Hill et al., 2021). However, assembling these hypervariable regions with short-read data has proven difficult due to repetitive sequences and limited homology between isolates, often resulting in incomplete sets of VSA sequences.

In this study, we apply PacBio HiFi sequencing to generate *de novo* assemblies of *P. falciparum* parasites from The Gambia. Using *in vitro* culture to obtain high-yield high molecular weight (HMW) DNA, we sequenced both clonal lineages (obtained via limiting dilution) and bulk field-adapted populations, resulting in 43 assemblies. We then used bioinformatics tools to assemble and annotate genomes, extract, classify and analyze VSA sequences.

Methods

Study design and sample collection

All *P. falciparum* field isolates reported here have been described previously (Collins et al., 2022; Fogang et al., 2024; Guery et al., 2025). Briefly, 5mL venous blood samples were collected from *P. falciparum* positive participants in 2016 and 2017 in The Gambia. Plasma, PBMCs and red blood cells were immediately separated by centrifugation. Red blood cells

were cryopreserved with glycerolyte in liquid nitrogen. The study protocol was reviewed and
approved by the Gambia Government/MRC Joint Ethics Committee (SCC 1476, SCC 1318,
L2015.50) and by the London School of Hygiene & Tropical Medicine Ethics Committee (Ref.
10982). The field studies were also approved by local administrative representatives, the
village chiefs. Written informed consent was obtained from participants over 18 years old and
from parents/guardians for participants under 18 years. Written assent was obtained from all
individuals aged 12–17 years. Approval under the Nagoya Protocol was granted by the
National Focal Point of The Gambia.

Parasite culturing

Thawed parasites were cultured in RPMI-1640 (sigma) supplemented with 25mM HEPES,
2mM L-glutamine, 0.5% Albumax II (sigma), 50µg/L gentamicin (sigma) and 5% AB$^+$ normal
human serum (NHS) in static conditions. Parasites were cultured in 10mL volumes at 2%
haematocrit in a blood gas environment of 90% N_2, 5% CO_2 and 5% oxygen (O_2).

Cloning by limiting dilution

Once field isolates were culture-adapted, typically after 2 to 3 weeks, clones were isolated by
limiting dilution in 96-well plates at a theoretical density of 1 parasite per well (ppw) – a 3ppw
was included as positive control – and maintained at 1.25% haematocrit. Plates were kept in a
modular chamber with blood gas and incubated at 37°C. On day 7, exhausted media was
removed and replaced with fresh media containing uninfected RBCs (uRBCs) at 1.25%
haematocrit. Culture media was changed every two days from this point onwards. From day
14, 3ppw wells were screened by microscopy for viable parasites. If parasites were absent,
plates were screened every two days until day 21, at which point the cloning assay was
considered failed and discarded. If viable parasites were present, the entire plate was screened
for parasites in the 1ppw wells. Positive wells were initially transferred into 6-well plates with
100µL uRBCs and 3mL media, and grown until parasitaemia was visible by microscopy
(usually 2 to 4 replication cycles). Parasites were then genotyped by MSP1 and MSP2 (see
details in supplementary methods, Supplementary Figure 1) to assess the number of parasite
genotypes present in cultures. Wells with confirmed unique single clones were selected for
expansion in 25mL cultures at 2% haematocrit, while all other wells were terminated. Parasites
were then grown to higher parasitaemia (~10%) and harvested at the schizont stage for high
molecular weight DNA extraction.

High molecular weight DNA extraction and PacBio sequencing
High molecular weight (HMW) DNA was extracted from both cloned parasites and 'bulk'
populations (parasite cultures directly adapted from field isolates) using the Monarch® HMW
DNA Extraction Kit for Tissue (New England Biolabs). The tissue kit was chosen over the
blood kit due to the protein separation step being able to remove residual cell debris and
haemozoin contaminants. Several modifications were made to the extraction procedure.
Briefly, RBCs were initially lysed with RBC Lysis Buffer from the Monarch HMW DNA
Extraction Kit for Cells & Blood. Free parasite pellets were then washed with 1X PBS until
the supernatant was clear. All subsequent steps followed the tissue kit's protocol. The agitation
speed for lysis was reduced to 1500 rpm and in special cases where parasite material was
limited, precipitation enhancer was added prior to binding DNA to the glass beads. DNA was
solubilised in 200µL of elution buffer, incubated at 37°C for 1 hour to enhance dissolution,
stored at 4°C overnight and then frozen at -20°C until sequencing. Prior to sequencing, DNA
was cleaned using SPRI (Solid Phase Reversible Immobilization) beads, sheared with g-TUBE

(Covaris) aiming for a size of 10kb and assessed for integrity using FEMTO Pulse (Agilent).
Samples were sequenced using the Pacific Biosciences (PacBio) high fidelity (HiFi) circular
consensus sequencing technology (Wenger *et al.*, 2019) on the Sequel IIe plaform, with up to
10 samples per SMRT cell.

Genome assemblies and annotation

We assembled genomes from PacBio HiFi reads datasets using Hifiasm (H. Cheng et al., 2021),
using the -l0 option to assemble haploid genome. Contigs were split, oriented and arranged to
the 3D7 chromosome configuration using RagTag, with default parameters (Alonge et al.,
2022). Assemblies coming from polyclonal isolates were left in the post-Hifiasm assembly
state, with no re-arrangements to the 3D7 chromosome configuration. Companion (Steinbiss
et al., 2016) was used to annotate clone and bulk assemblies with the following options:
pseudogene detection activated, AUGUSTUS (Stanke et al., 2004) for structural annotation
and RATT (Otto et al., 2011) for transfer of highly conserved genes from 3D7.

VSAs extraction

Var genes were retrieved from all assemblies by screening for genes annotated as potentially
encoding PfEMP1 by the Companion pipeline and filtering for a nucleotide length higher than
2500bp. Each *var* gene had its different domains annotated by submitting its amino acid
sequences to the varDOM server (Rask et al., 2010). Copies of *var2csa* were extracted from
the list of *var* genes by screening for genes with DBLpam domains. Upstream group affiliation
of the remaining *var* genes was performed as in the method described in Pangilinan et al., 2025.
Briefly, a phylogenetic tree was constructed using the 500bp upstream nucleotide sequences of
newly assembled *var* genes plus reference upstream sequences previously annotated as group
A, B or C (Lavstsen et al., 2003; Rask et al., 2010). Sequences of all *var* genes were aligned
using MAFFT v7.520 and a phylogenetic tree was inferred from the alignment using FastTree
v2.1.11. Upstream sequences were annotated based on the clade of the reference upstream
group. The HMM-based software STRIDE (Zhou et al., 2022) was used to retrieve *rif* and
stevor from the list of Companion proteins, and further annotate the *rif* into *rif* groups A or *rif*
group B.

Estimation of complexity of infection across bulk parasite populations

To infer whether bulk parasite populations were single- or poly-genotype we looked at average
heterozygosity levels over a panel of 72,020 core-genome variable loci. This set of variable
loci was generated by mapping PacBio HiFi reads to the reference 3D7 genome using
minimap2 v2.28 (Li, 2018). SNP calling and generation of VCF files were performed with
GATK v4.1.0.5.1 (Van der Auwera et al., 2020), using the HaplotypeCaller pipeline. Variants
were filtered based on their quality using bcftools v1.9 (Danecek et al., 2021) and imported in
an R environment with the R package vcfR (Knaus & Grünwald, 2017). Genome assemblies
were considered to be originating from a single-genotype population if their average
heterozygosity level was lower than 0.02 over the entire set of 72,020 variants, which
corresponded to the highest heterozygosity level found in clone genome assemblies.

Estimation of base-calling error rates in hifiasm assemblies

To estimate base-calling error rates level in newly-generated assemblies, we used two *ex vivo*
isolates (i.e. without any culturing), DC03m1 and DC10m3, sequenced with Illumina in a
previous study (Guery et al., 2025). Illumina reads were mapped to their respective long read
assembly using bwa MEM v0.7.17 (Li & Durbin, 2009). The output sam files were then
converted and sorted using samtools v1.10 (Li et al., 2009). GATK v4.1.0.5.1 (Van der Auwera
et al., 2020) was used again for SNPs calling and generation of VCF files, using the
HaplotypeCaller pipeline. Additionally, nucleotide level accuracies in assemblies were also

estimated by comparing the k-mer decomposition of PacBio HiFi assemblies to illumina reads
datasets via the software merqury (Rhie et al., 2020). Here we used the quality value (qv) score
to assess base-calling accuracy, a metric that is generated by looking at k-mers present in high-
quality reads dataset (Illumina) but not in the PacBio assemblies.

Estimation of pairwise Identity By Descent (IBD) and VSAs sharing amongst samples
186

To estimate core genome relatedness between parasites genomes we calculated the pairwise
IBD values between all genomes. Chromosomal segments are in IBD state if they descend from
a recent common ancestor, and a pairwise IBD value indicate the average proportion of two
genomes that are in IBD state. In practise, PacBio HiFi reads were mapped to the reference
3D7 genome using minimap2 v2.28 (Li, 2018). SNP calling was performed as described above.
From this, VCF files were parsed with the R package vcfR to generate a tab-delimited table
containing SNPs position and status for each genome. We used hmmIBD v2.0.0 (Schaffner et
al., 2018) to calculate the fraction of the genome in an IBD state, from here referred as pairwise
IBD values, between each pair of genomes.

In parallel, we assessed the similarity of VSAs repertoires between each single-genotype
genome pair by clustering *var*, *rif*, and *stevor* sequences using CD-HIT (Fu et al., 2012).
Nucleotide sequences in different isolate genomes were considered identical if their sequence
similarity was higher than 99%. After counting the number of shared VSAs between two
genomes we calculated the three normalised pairwise VSA sharing indexes *var* sharing, *rif*
sharing and *stevor* sharing using the following formula:

$VSA sharing index$
$$= \frac{2 * Number of shared VSA genes}{Number of VSA genes \in genome1 + Number of VSA genes \in genome2}$$

To account for the non-independence of pairwise estimates, overall correlations between core
genome IBD values and VSA sharing indexes were then evaluated using Mantel tests, as
implemented in the R package vegan (v2.6-10). Distance matrices of pairwise IBD values and
normalized *var, rif* and *stevor* sharing were compared via the mante() function, with 9999
permutations and p-values computed using the Spearman rank correlation method.
211
212

Results

Blood isolates were collected in four neighboring villages from The Gambia, following the
sample design presented in Collins et al., 2022 and Fogang et al., 2024. Briefly, samples were
collected during the wet season (October 2016 – November 2016) from both asymptomatic and
symptomatic cases, and from asymptomatic carriers at monthly timepoints during the following
dry season (December 2016 to May 2017). From this sampling design, isolates were separated
into three infection groups: the Wet season Symptomatics (WS), corresponding to parasites
collected from symptomatic cases in the wet season; the Wet season Asymptomatics (WA),
corresponding to parasites collected from asymptomatic cases in the wet season and the Dry
season asymptomatic Cohort samples (DC), corresponding to parasites collected at monthly
timepoints from asymptomatic cases during the dry season. Sample identifiers incorporate
group, timepoint, and sequencing type (e.g., bulk 'b' vs clone 'c'; e.g., WA01b, DC01m1c).
Metadata and group assignments are detailed in Table 1.

Culture adaptation, cloning, and high molecular weight DNA extraction

Parasites were culture-adapted to generate the high DNA yields necessary for long-read sequencing. Cloning by limiting dilution allowed the isolation of single-genotype populations (Supplementary Figure 1). Two extraction kits (Qiagen Genomic-tip 100/G and Monarch HMW DNA extraction kit) were tested for high molecular weight (HMW) DNA recovery, with the Monarch kit performing better (Supplementary Figure 2). Overall, a total set of 43 DNA extractions derived from 30 isolates were all successfully sequenced with PacBio HiFi (Table 1). Twenty three (23) genomes were derived from cloning dilutions and were strictly single-genotype parasite populations. The other 20 DNA samples were bulk populations (not cloned) with an unknown number of parasite genotypes. Based on their level of heterozygosity over a set of 72,020 informative loci, we inferred 8 bulk genomes to be single-genotype and 12 to consist of multiple parasite genotypes (Table 1). Across infection groups, we sequenced 17 genomes from asymptomatic infections in the dry season (DC group), 15 genomes from symptomatic infections in the wet season (WS group) and 11 genomes from asymptomatic infections in the wet season (WA group).

Pacbio HiFi enables chromosome-scale genome assemblies

Raw assemblies from the 43 Pacbio HiFi reads datasets contained between 17 and 185 contigs (Table 2), with a median number of 54. The N50 statistics in all genomes (range 1,198,427 to 1,693,536 bp, median = 1,571,854 bp) were indicative of the presence of large chromosome-size contigs, which was further supported by syntenic analyse comparing newly assembled genomes to the 3D7 reference genome (Figure 2). The number of complete telomere-to-telomere contigs was however systematically lower than the 14 expected *P. falciparum* chromosomes (range 2 to 9, median = 4), demonstrating difficulties in assembling hypervariable sub-telomeric regions and complete telomeric contigs.

Across the 31 single-genotype genomes, some of the Hifiasm assemblies contained contigs with regions syntenic to more than one chromosome. Mapping PacBio HiFi reads onto their assemblies revealed that the apparent chromosome breaks coincided with regions of low coverage or low sequence complexity, with no evidence of breaks supported by high read depth. Accordingly, raw contigs were split and reorganized into the 3D7 chromosome structure using ragtag (Alonge et al., 2022). After the re-contiguation step, all single-genotype assemblies had 14 large contigs corresponding to the 14 *P. falciparum* chromosomes, alongside 2 contigs for the mitochondrial and apicoplast genomes and a variable number of small extra contigs (median = 32, range 4 to 171; Figure 2, Table 2).

Pacbio Hifi P. falciparum genome assemblies showed high base-calling accuracy

To assess base-calling accuracy of genome assemblies, we aligned Illumina reads from two ex vivo isolates, DC03m1 and DC10m3, (Guery et al., 2025) to their respective Pacbio HiFi assemblies from clonal genomes. We found only 55 SNPs over the entire DC03m1 genome (total length 23.6 Mbp) and 40 SNPs over the entire DC10m1 genome (total length 23.5 Mbp). We also computed k-mer-based quality scores using Merqury (Rhie et al., 2020), which yielded quality value (qv) scores of 64.5 for DC03m1 and 65.5 for DC10m3, indicating error rates ranging between 1 in 1,000,000 bp and 1 in 10,000,000 bp for both genomes. Those results confirmed the high accuracy and quality of genome assemblies generated by hifiasm for Pacbio HiFi reads, and their use in generating reliable *P. falciparum* genome assemblies.

IBD analysis reveals genotypic relatedness

To investigate genetic relatedness, we calculated identity-by-descent (IBD) fractions using core-genome SNPs mapped to the 3D7 reference genome (Supplementary Figure 3). In some cases, the same parasite genotypes were observed in different hosts (e.g. DC03m1c and WS11b), highlighting the presence and persistence of certain parasite genotypes across infection groups and time. In addition to shared parasite genotypes, we also observed groups of parasites with intermediate pairwise IBD values ranging from 0.1 to 0.9, highlighting groups of inter-related parasites that share a common recent ancestry (*e.g.* DC07m1c and DC10m3c).

Full VSAs repertoires retrieved from PacBio HiFi assemblies

We extracted 3,288 *var* genes from all genomes (1,504 unique sequences) using Companion annotations. Clonal assemblies contained 56–84 *var* genes per genome (median = 69), consistent with expectations for single-genotype isolates. Bulk assemblies had 65–130 *var* genes, suggesting up to two co-infecting strains in some samples. The number of extracted *var* genes strongly correlated with the number of protein-coding DNA sequences (CDS) containing the LARSFADIG PfEMP1 motif ($p < 2e-16$), consistent with earlier findings (Otto et al., 2019).

Using the information on shared genotypes across samples deduced from pairwise IBD values (Supplementary Figure 3), we defined an unbiased dataset of unique single-genotype isolates (N = 17). Within this dataset, the numbers and proportions of each *var* upstream groups were relatively stable (Figure 3 and Table 2). Group A *var* genes ranged from 8 to 16 genes per genome with a median number of 11 per genome, group B *var* genes ranged from 29 to 46 per genome with a median number of 37 genes per genome and group C *var* genes ranged from 8 to 19 per genome with a median number of 12 per genome. In genomes sequenced from poly-genotype isolates the proportions of groups A, B and C were similar to the ones observed in clones, with a larger overall number of *var* genes (Table 2). Some *var* genes could not be reliably classified into any upstream group. This included genes with very divergent upstream sequences and *var* genes sequences located in small contigs where their 500bp upstream sequence could not be isolated.

As previously reported (Benavente et al., 2018; Sander et al., 2009), multiple copies of the conserved *var* gene *var2csa* were observed in multiple genomes (Table 2). Across the 17 unique single-genotype, 9 assemblies had a single *var2csa* copy and 8 assemblies had two or more *var2csa* copies (median = 2, range 1 to 3). In single-genotype genomes where copies of *var2csa* could be localised there was always a copy on chromosome 12, as in 3D7 and other reference genomes. Extra copies of the gene were found on sub-telomeric regions of chromosomes 1, 3, 8, 12, 13 and 14.

Similar to the pattern observed with *var* genes, the numbers and proportions of *rif* and *stevor* stayed relatively constant across different isolates (Figure 2). In the dataset of unique single-genotype genomes (N = 17) the total number of *rif* varied from 117 to 196 per genome with a median number of 168 genes. From those, *rif* group A ranged from 85 to 139 per genome with a median number of 116 genes and *rif* group B ranged from 32 to 65 with a median number of 48 genes per genome. The *stevor* were less numerous, ranging from 26 to 51 with a median number of 39 genes per genome.

326 There was no correlation between the number of *var* genes and the number of *rif* (p = 0.61), or
327 the number of *stevor* (p = 0.83), and the numbers of *rif* and *stevor* were mildly correlated (p =
328 0.03; see Supplementary Figure 4).
329
330 ***Pairwise core genome IBD and VSAs sharing indexes are highly correlated***
331
We quantified pairwise sharing indexes for *var*, *rif*, and *stevor* gene repertoires in the single-
genotype dataset (N = 31, excluding technical replicates). The pairwise sharing indexes reflects
a measure of genetic similarity for non-core *P. falciparum* genomes. Overall, sharing patterns
closely paralleled core-genome IBD values—genome pairs with high IBD exhibited similar
VSA repertoires (Figure 4A). Matrices permutation tests revealed significant correlations
between core-genome IBD and VSA sharing for all three families (Figure 4B) (p < 0.01). Of
note, we also observed most pairs of genomes seemed to share some *stevor* sequences, with an
average *stevor* sharing value of 0.03 in pairs of samples with IBD < 0.25, against a lower
average sharing value of 0.008 for *rif* and 0.003 for *var*. This is in line with previous
observations that reported *stevor* as the least hypervariable of the three VSAs families (Q.
Cheng et al., 1998; Lawton et al., 2025).
343
Discussion
345
In total, we successfully culture-adapted and long-read sequenced parasite genomes from 43
parasite cultures, originating from 30 distinct field isolates. To the best of our knowledge this
is the largest dataset of *P. falciparum* long read assemblies generated from natural infections
to date. With PacBio HiFi reads and using the hifiasm assembler, we demonstrated long and
accurate assembly contigs could be generated without the need for complementary illumina
libraries, as is necessary with other long reads sequencing technologies (Delandre et al., 2024;
Higgins et al., 2024). Particularly, our method proved useful in assembling *P. falciparum* sub-
telomeric regions and complete sets of VSA families which are highly polymorphic and hard
to assemble using short-read datasets (Andradi-Brown et al., 2024).
355
356 Using cloning by limiting dilutions, parasite lineages could be delimited to ensure sequences
357 were retrieved from single-genotype populations. Bulk populations (without the cloning step)
358 also proved useful in retrieving complete VSA sequences, regardless of the number of parasite
359 genotypes present.
360
The risk of culture-adaptation–induced mutations, although well established for parasites
subjected to extended *in vitro* passages (Claessens et al., 2017, 2023; Kemp et al., 1992), is
minimal here, as all genomes were derived from parasites maintained in culture only for the
few weeks required for DNA extraction. This makes the generated assemblies a reliable
genomic resource for analysis of circulating West African *P. falciparum*—across both
symptomatic and asymptomatic infections.
367
The analysis of VSA variation among genome assemblies confirmed patterns found in previous
studies, with ratios of each *var* and *rif* groups stable across isolates. The presence of multiple
copies of *var2csa* was observed in approximately half of the genomes generated here, which
seems to be common in field isolates (Sander et al., 2009). There however seem to be an
evolutionary constraint on maintaining at least one copy of the gene at the 5' telomeric end of
chromosome 12, as this copy was found in all genomes.
374

Contrary to a previous study which sampled *P. falciparum* reference genomes from widespread worldwide locations (Otto et al., 2018) we have generated a set of genomes originating from infections that were both close in collection dates and geographical locations. With this sampling design, we collected parasites with a wide range of genetic relatedness, from completely unrelated parasites to identical parasite genotypes sampled across different hosts or timepoints of a chronic infection. The relationship between core genome and non-core genome estimates of relatedness have seldom been studied (Neafsey et al., 2021), and the generation of genetic diversity in these two genomic regions stems from different biological processes. In the core genome, homologous recombination events occurring during meiosis lead to an uneven repartitioning of genetic variants and SNPs in offspring, with the proportion of the genome regions containing similar genetic variants tightly linked to the kinship value between parasites (Guo et al., 2025; Schaffner et al., 2018). In the non-core genome, variability is additionally generated through non-homologous recombination events, leading to sequence hypervariability and rapid generation of new VSAs (Claessens et al., 2014). Despite these differences, we observed a remarkably close relationship between core genome pairwise IBD values and the three sharing indexes for *var*, *rif*, *stevor*. These results indicate that in epidemiological settings similar to the one observed in The Gambia, measures of VSA sharing could effectively be used as a proxy for core genome relatedness, and vice-versa. These results support the validity of the growing number of studies relying on mass amplification of conserved *var* domains (e.g. DBLα) for population genetic analyses, and highlight VSA sharing as valid measure of genetic relatedness (Day et al., 2025).

Overall, the genome and VSA sequence datasets provided in this study presents a great resource for future research. Increasing the number of long read sequences generated from field isolated parasites will enable analyses of large structural variants occurring in *P. falciparum* genomes at a large scale, including regions with high repeats or high AT content. Furthermore, the complete and high-quality *var* gene sequences extracted here will prove a useful resource and training dataset for algorithm linking partial *var* sequences to full *var* domain composition (Mackenzie et al., 2022), or upstream group (Pangilinan et al., 2025). Finally, obtaining full-length VSA sequences will enable robust analyses of *var* gene expression in samples coming from The Gambia, and will enable differentiating *var* expression between asymptomatic and symptomatic infections.

Data availability statement

All PacBio HiFi raw reads datasets are publicly available through the NCBI accession number indicated in Table 1. Assembled genomes are available on NCBI, BioProject PRJNA1368450. Annotation, VSA sequences and classification are available at the public repository https://zenodo.org/records/17551160. A series of codes and scripts used for this paper can be found on the github repository https://github.com/Mquenu/Falcigenomes-toolkit, https://zenodo.org/records/17724288.

Acknowledgements

We thank Craig Corton for helpful advice on sample processing. Funding: ANR 18-CE15-0009-01. Fondation pour la Recherche Médicale FRM EQU202303016290. Service Programme d'Excellence I-Site, MUSE Pre-ERC, MOTIVATOR.

References

Alonge, M., Lebeigle, L., Kirsche, M., Jenike, K., Ou, S., Aganezov, S., Wang, X., Lippman, Z. B., Schatz, M. C., & Soyk, S. (2022). Automated assembly scaffolding using RagTag elevates a new tomato system for high-throughput genome editing. *Genome Biology*, *23*(1), 1–19. https://doi.org/10.1186/S13059-022-02823-7/FIGURES/2

Andradi-Brown, C., Wichers-Misterek, J. S., von Thien, H., Höppner, Y. D., Scholz, J. A. M., Hansson, H., Hocke, E. F., Gilberger, T. W., Duffy, M. F., Lavstsen, T., Baum, J., Otto, T. D., Cunnington, A. J., & Bachmann, A. (2024). A novel computational pipeline for var gene expression augments the discovery of changes in the Plasmodium falciparum transcriptome during transition from in vivo to short-term in vitro culture. *ELife*, *13*. https://doi.org/10.7554/ELIFE.87726

Benavente, E. D., Oresegun, D. R., de Sessions, P. F., Walker, E. M., Roper, C., Dombrowski, J. G., de Souza, R. M., Marinho, C. R. F., Sutherland, C. J., Hibberd, M. L., Mohareb, F., Baker, D. A., Clark, T. G., & Campino, S. (2018). Global genetic diversity of var2csa in Plasmodium falciparum with implications for malaria in pregnancy and vaccine development. *Scientific Reports 2018 8:1*, *8*(1), 1–8. https://doi.org/10.1038/s41598-018-33767-3

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nature Methods 2021 18:2*, *18*(2), 170–175. https://doi.org/10.1038/s41592-020-01056-5

Cheng, Q., Cloonan, N., Fischer, K., Thompson, J., Waine, G., Lanzer, M., & Saul, A. (1998). stevor and rif are Plasmodium falciparum multicopy gene families which potentially encode variant antigens. *Molecular and Biochemical Parasitology*, *97*(1–2), 161–176. https://doi.org/10.1016/S0166-6851(98)00144-3

Claessens, A., Adams, Y., Ghumra, A., Lindergard, G., Buchan, C. C., Andisi, C., Bull, P. C., Mok, S., Gupta, A. P., Wang, C. W., Turner, L., Arman, M., Raza, A., Bozdech, Z., & Rowe, J. A. (2012). A subset of group A-like var genes encodes the malaria parasite ligands for binding to human brain endothelial cells. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(26). https://doi.org/10.1073/PNAS.1120461109

Claessens, A., Affara, M., Assefa, S. A., Kwiatkowski, D. P., & Conway, D. J. (2017). Culture adaptation of malaria parasites selects for convergent loss-of-function mutants. *Scientific Reports 2017 7:1*, *7*(1), 1–8. https://doi.org/10.1038/srep41303

Claessens, A., Hamilton, W. L., Kekre, M., Otto, T. D., Faizullabhoy, A., Rayner, J. C., & Kwiatkowski, D. (2014). Generation of Antigenic Diversity in Plasmodium falciparum by Structured Rearrangement of Var Genes During Mitosis. *PLOS Genetics*, *10*(12), e1004812. https://doi.org/10.1371/JOURNAL.PGEN.1004812

Claessens, A., Stewart, L. B., Drury, E., Ahouidi, A. D., Amambua-Ngwa, A., Diakite, M., Kwiatkowski, D. P., Awandare, G. A., & Conway, D. J. (2023). Genomic variation during culture adaptation of genetically complex Plasmodium falciparum clinical isolates. *Microbial Genomics*, *9*(5), 001009. https://doi.org/10.1099/MGEN.0.001009/CITE/REFWORKS

Collins, K. A., Ceesay, S., Drammeh, S., Jaiteh, F. K., Guery, M. A., Lanke, K., Grignard, L., Stone, W., Conway, D. J., D'Alessandro, U., Bousema, T., & Claessens, A. (2022). A Cohort Study on the Duration of Plasmodium falciparum Infections During the Dry Season in The Gambia. *The Journal of Infectious Diseases*, *226*(1), 128–137. https://doi.org/10.1093/INFDIS/JIAC116

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., & Davies, R. M. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2). https://doi.org/10.1093/GIGASCIENCE/GIAB008

Day, K. P., Tan, M. H., He, Q., Ruybal-Pesántez, S., Zhan, Q., Tiedje, K. E., & Pascual, M. (2025). Var genes, strain hyperdiversity, and malaria transmission dynamics. *Trends in Parasitology*, *41*(6), 471–485. https://doi.org/10.1016/J.PT.2025.04.010

Delandre, O., Lamer, O., Loreau, J. M., Papa Mze, N., Fonta, I., Mosnier, J., Gomez, N., Javelle, E., & Pradines, B. (2024). Long-Read Sequencing and De Novo Genome Assembly Pipeline of Two Plasmodium falciparum Clones (Pf3D7, PfW2) Using Only the PromethION Sequencer from Oxford Nanopore Technologies without Whole-Genome Amplification. *Biology*, *13*(2), 89. https://doi.org/10.3390/BIOLOGY13020089/S1

Fogang, B., Lellouche, L., Ceesay, S., Drammeh, S., Jaiteh, F. K., Guery, M. A., Landier, J., Haanappel, C. P., Froberg, J., Conway, D., D'Alessandro, U., Bousema, T., & Claessens, A. (2024). Asymptomatic Plasmodium falciparum carriage at the end of the dry season is associated with subsequent infection and clinical malaria in Eastern Gambia. *Malaria Journal*, *23*(1). https://doi.org/10.1186/S12936-024-04836-Y

Fu, L., Niu, B., Zhu, Z., Wu, S., & Li, W. (2012). CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics*, *28*(23), 3150. https://doi.org/10.1093/BIOINFORMATICS/BTS565

Gardner, M. J., Hall, N., Fung, E., White, O., Berriman, M., Hyman, R. W., Carlton, J. M., Pain, A., Nelson, K. E., Bowman, S., Paulsen, I. T., James, K., Eisen, J. A., Rutherford, K., Salzberg, S. L., Craig, A., Kyes, S., Chan, M. S., Nene, V., … Barrell, B. (2002). Genome sequence of the human malaria parasite Plasmodium falciparum. *Nature*, *419*(6906), 498. https://doi.org/10.1038/nature01097

Gen, M., Ahouidi, A., Ali, M., Almagro-Garcia, J., Amambua-Ngwa, A., Amaratunga, C., Amato, R., Amenga-Etego, L., Andagalu, B., Anderson, T. J. C., Andrianaranjaka, V., Apinjoh, T., Ariani, C., Ashley, E. A., Auburn, S., Awandare, G. A., Ba, H., Baraka, V., Barry, A. E., … Ye, H. (2021). An open dataset of Plasmodium falciparum genome variation in 7,000 worldwide samples. *Wellcome Open Research*, *6*, 42. https://doi.org/10.12688/WELLCOMEOPENRES.16168.2

Godin, M. J., Sebastian, A., Albert, I., & Lindner, S. E. (2023). Long-read genome assembly and gene model annotations for the rodent malaria parasite Plasmodium yoelii 17XNL. *Journal of Biological Chemistry*, *299*(7). https://doi.org/10.1016/J.JBC.2023.104871/ATTACHMENT/BE8F4C96-D7A6-465F-A437-A435642F188E/MMC9.PDF

Guery, M.-A., Ceesay, S., Drammeh, S., Jaiteh, F. K., d'Alessandro, U., Bousema, T., Conway, D. J., & Claessens, A. (2025). Household clustering and seasonal genetic variation of Plasmodium falciparum at the community-level in The Gambia. *ELife*, *13*. https://doi.org/10.7554/ELIFE.103047.2

Guo, B., Rowley, E., O'Connor, T. D., & Takala-Harrison, S. (2025). Potential and pitfalls of using identity-by-descent for malaria genomic surveillance. *Trends in Parasitology*, *41*(5), 387–400. https://doi.org/10.1016/J.PT.2025.03.012/ATTACHMENT/04E8641E-AA9E-41CF-959F-32D78B225F93/MMC1.DOCX

Harrison, T. E., Mørch, A. M., Felce, J. H., Sakoguchi, A., Reid, A. J., Arase, H., Dustin, M. L., & Higgins, M. K. (2020). Structural basis for RIFIN-mediated activation of LILRB1 in malaria. *Nature*, *587*(7833), 309–312. https://doi.org/10.1038/S41586-020-2530-3

Higgins, M., Manko, E., Ward, D., Phelan, J. E., Nolder, D., Sutherland, C. J., Clark, T. G., & Campino, S. (2024). New reference genomes to distinguish the sympatric malaria parasites, Plasmodium ovale curtisi and Plasmodium ovale wallikeri. *Scientific Reports 2024 14:1*, *14*(1), 1–10. https://doi.org/10.1038/s41598-024-54382-5

Holzschuh, A., Lerch, A., Fakih, B. S., Aliy, S. M., Ali, M. H., Ali, M. A., Bruzzese, D. J., Yukich, J., Hetzel, M. W., & Koepfli, C. (2024). Using a mobile nanopore sequencing lab for end-to-end genomic surveillance of Plasmodium falciparum: A feasibility study. *PLOS*

Global Public Health, *4*(2 February), e0002743.
https://doi.org/10.1371/JOURNAL.PGPH.0002743/OG_IMAGE.JPG

Hon, T., Mars, K., Young, G., Tsai, Y. C., Karalius, J. W., Landolin, J. M., Maurer, N., Kudrna, D., Hardigan, M. A., Steiner, C. C., Knapp, S. J., Ware, D., Shapiro, B., Peluso, P., & Rank, D. R. (2020). Highly accurate long-read HiFi sequencing data for five complex genomes. *Scientific Data 2020 7:1*, *7*(1), 1–11. https://doi.org/10.1038/s41597-020-00743-4

Jespersen, J. S., Wang, C. W., Mkumbaye, S. I., Minja, D. T., Petersen, B., Turner, L., Petersen, J. E., Lusingu, J. P., Theander, T. G., & Lavstsen, T. (2016). Plasmodium falciparum var genes expressed in children with severe malaria encode CIDRα1 domains. *EMBO Molecular Medicine*, *8*(8), 839–850. https://doi.org/10.15252/EMMM.201606188

Kemp, D. J., Thompson, J., Barnes, D. A., Triglia, T., Karamalis, F., Petersen, C., Brown, G. V., & Day, K. P. (1992). A chromosome 9 deletion in Plasmodium falciparum results in loss of cytoadherence. *Memorias Do Instituto Oswaldo Cruz*, *87 Suppl 3*, 85–89. https://doi.org/10.1590/S0074-02761992000700011

Knaus, B. J., & Grünwald, N. J. (2017). vcfr: a package to manipulate and visualize variant call format data in R. *Molecular Ecology Resources*, *17*(1), 44–53. https://doi.org/10.1111/1755-0998.12549

Lavstsen, T., Salanti, A., Jensen, A. T., Arnot, D. E., & Theander, T. G. (2003). Sub-grouping of Plasmodium falciparum 3D7 var genes based on sequence analysis of coding and non-coding regions. *Malaria Journal*, *2*(1), 27. https://doi.org/10.1186/1475-2875-2-27

Lawton, J. G., Zhou, A. E., Stucke, E. M., Takala-Harrison, S., Silva, J. C., & Travassos, M. A. (2025). Diamonds in the rif: Alignment-free comparative genomics analysis reveals strain-transcendent Plasmodium falciparum antigens amidst extensive genetic diversity. *Infection, Genetics and Evolution*, *129*, 105725. https://doi.org/10.1016/J.MEEGID.2025.105725

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*, *34*(18), 3094–3100. https://doi.org/10.1093/BIOINFORMATICS/BTY191

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*, *25*(14), 1754. https://doi.org/10.1093/BIOINFORMATICS/BTP324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079. https://doi.org/10.1093/BIOINFORMATICS/BTP352

Mackenzie, G., Jensen, R. W., Lavstsen, T., & Otto, T. D. (2022). Varia: a tool for prediction, analysis and visualisation of variable genes. *BMC Bioinformatics*, *23*(1). https://doi.org/10.1186/S12859-022-04573-6

Neafsey, D. E., Taylor, A. R., & MacInnis, B. L. (2021). Advances and opportunities in malaria population genomics. *Nature Reviews. Genetics*, *22*(8), 502. https://doi.org/10.1038/S41576-021-00349-5

Otto, T. D., Assefa, S. A., Böhme, U., Sanders, M. J., Kwiatkowski, D. P., Berriman, M., & Newbold, C. (2019). Evolutionary analysis of the most polymorphic gene family in Falciparum malaria. *Wellcome Open Research*, *4*. https://doi.org/10.12688/WELLCOMEOPENRES.15590.1/DOI

Otto, T. D., Böhme, U., Sanders, M., Reid, A., Bruske, E. I., Duffy, C. W., Bull, P. C., Pearson, R. D., Abdi, A., Dimonte, S., Stewart, L. B., Campino, S., Kekre, M., Hamilton, W. L., Claessens, A., Volkman, S. K., Ndiaye, D., Amambua-Ngwa, A., Diakite, M., … Berriman, M. (2018). Long read assemblies of geographically dispersed Plasmodium falciparum isolates reveal highly structured subtelomeres. *Wellcome Open Research*, *3*, 52. https://doi.org/10.12688/WELLCOMEOPENRES.14571.1

Otto, T. D., Dillon, G. P., Degrave, W. S., & Berriman, M. (2011). RATT: Rapid Annotation
Transfer Tool. *Nucleic Acids Research*, *39*(9), e57–e57.
https://doi.org/10.1093/NAR/GKQ1268

Pangilinan, E. A., Quenu, M., Claessens, A., & Otto, T. D. (2025). upsAI: A high-accuracy
machine learning classifier for predicting Plasmodium falciparum var gene upstream groups.
BioRxiv, 2025.05.19.654848. https://doi.org/10.1101/2025.05.19.654848

Rask, T. S., Hansen, D. A., Theander, T. G., Pedersen, A. G., & Lavstsen, T. (2010). Plasmodium
falciparum erythrocyte membrane protein 1 diversity in seven genomes--divide and conquer.
PLoS Computational Biology, *6*(9). https://doi.org/10.1371/JOURNAL.PCBI.1000933

Rhie, A., Walenz, B. P., Koren, S., & Phillippy, A. M. (2020). Merqury: Reference-free quality,
completeness, and phasing assessment for genome assemblies. *Genome Biology*, *21*(1), 1–27.
https://doi.org/10.1186/S13059-020-02134-9/FIGURES/6

Rowe, J. A., Claessens, A., Corrigan, R. A., & Arman, M. (2009). Adhesion of Plasmodium
falciparum-infected erythrocytes to human cells: Molecular mechanisms and therapeutic
implications. *Expert Reviews in Molecular Medicine*, *11*.
https://doi.org/10.1017/S1462399409001082

Sakoguchi, A., Chamberlain, S. G., Mørch, A. M., Widdess, M., Harrison, T. E., Dustin, M. L.,
Arase, H., Higgins, M. K., & Iwanaga, S. (2025). RIFINs displayed on malaria-infected
erythrocytes bind KIR2DL1 and KIR2DS1. *Nature 2025*, 1–9.
https://doi.org/10.1038/s41586-025-09091-y

Sander, A. F., Salanti, A., Lavstsen, T., Nielsen, M. A., Magistrado, P., Lusingu, J., Ndam, N. T.,
& Arnot, D. E. (2009). Multiple var2csa-Type PfEMP1 Genes Located at Different
Chromosomal Loci Occur in Many Plasmodium falciparum Isolates. *PLOS ONE*, *4*(8),
e6667. https://doi.org/10.1371/JOURNAL.PONE.0006667

Schaffner, S. F., Taylor, A. R., Wong, W., Wirth, D. F., & Neafsey, D. E. (2018). HmmIBD:
Software to infer pairwise identity by descent between haploid genotypes. *Malaria Journal*,
17(1), 1–4. https://doi.org/10.1186/S12936-018-2349-7/FIGURES/1

Stanke, M., Steinkamp, R., Waack, S., & Morgenstern, B. (2004). AUGUSTUS: a web server for
gene finding in eukaryotes. *Nucleic Acids Research*, *32*(Web Server issue), W309.
https://doi.org/10.1093/NAR/GKH379

Tonkin-Hill, G., Ruybal-Pesántez, S., Tiedje, K. E., Rougeron, V., Duffy, M. F., Zakeri, S.,
Pumpaibool, T., Harnyuttanakorn, P., Branch, O. L. H., Ruiz-Mesía, L., Rask, T. S.,
Prugnolle, F., Papenfuss, A. T., Chan, Y. B., & Day, K. P. (2021). Evolutionary analyses of
the major variant surface antigen-encoding genes reveal population structure of Plasmodium
falciparum within and between continents. *PLOS Genetics*, *17*(2), e1009269.
https://doi.org/10.1371/JOURNAL.PGEN.1009269

Van der Auwera, G., O'Connor, B., & Safari. (2020). Genomics in the Cloud: Using Docker,
GATK, and WDL in Terra. *O'Reilly Media*, 300.
https://www.oreilly.com/library/view/genomics-in-the/9781491975183/

Vembar, S. S., Seetin, M., Lambert, C., Nattestad, M., Schatz, M. C., Baybayan, P., Scherf, A., &
Smith, M. L. (2016). Complete telomere-to-telomere de novo assembly of the Plasmodium
falciparum genome through long-read (>11 kb), single molecule, real-time sequencing. *DNA
Research*, *23*(4), 339–351. https://doi.org/10.1093/DNARES/DSW022

Wenger, A. M., Peluso, P., Rowell, W. J., Chang, P. C., Hall, R. J., Concepcion, G. T., Ebler, J.,
Fungtammasan, A., Kolesnikov, A., Olson, N. D., Töpfer, A., Alonge, M., Mahmoud, M.,
Qian, Y., Chin, C. S., Phillippy, A. M., Schatz, M. C., Myers, G., DePristo, M. A., …
Hunkapiller, M. W. (2019). Accurate circular consensus long-read sequencing improves
variant detection and assembly of a human genome. *Nature Biotechnology 2019 37:10*,
37(10), 1155–1162. https://doi.org/10.1038/s41587-019-0217-9

World malaria report 2024. (n.d.). Retrieved July 25, 2025, from
https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2024
Zhou, A. E., Shah, Z. V., Bradwell, K. R., Munro, J. B., Berry, A. A., Serre, D., Takala-Harrison,
S., O'Connor, T. D., Silva, J. C., & Travassos, M. A. (2022). STRIDE: a command-line
HMM-based identifier and sub-classifier of Plasmodium falciparum RIFIN and STEVOR
variant surface antigen families. *BMC Bioinformatics*, *23*(1), 1–9.
https://doi.org/10.1186/S12859-021-04515-8/TABLES/2

627
628

Figures

630
631
Figure 1: Schematic overview of the genome assembly method used to generate high quality genomes and extract Variant Surface Antigens (VSAs) from parasite cultures.
634

635
636
637
Figure 2: PacBio HiFi sequencing and hifiasm assembly allows chromosome-scale genome assemblies. Each circos plot represents chromosomes of the reference 3D7 genome

(top) and the DC01m1 contigs (bottom) as a representative genome generated from single-
genotype parasite lineage. Grey lines between top and bottom sequence assemblies indicate
syntenic regions of high sequence similarity. The outer ring indicates position of genes
annotated by the companion pipeline, genes in the forward strand are indicated in blue and
genes in the reverse strand indicated in red. Gaps in synteny can be seen over internal *var*
clusters on chromosomes 4, 7, 8 and 12.
646
647

648
649
650
651
652 **Figure 3: Numbers and proportions of VSA groups are stable across unique single-**
653 **genotype assemblies (N = 17).** The upper panel indicates the number of *var* gene per genome
654 assembly, with colours indicating the different *var* groups. The lower panel indicates the
655 number of *rif* and *stevor* per genome assembly, with colours indicating the different groups.
656
657

Figure 4: Groups of related parasites share common *var*, *rif* and *stevor* genes.
(A) Pairwise matrices of IBD values and three normalized sharing indexes (*var*, *rif*, *stevor*) in a dataset of 31 single-genotype genomes (B) Correlations between pairwise IBD values and the three VSAs sharing indexes for the same dataset. P-values were calculated by a mantel test.

Tables

Table 1: Genome accession numbers and metadata for 43 *P. falciparum* genome assemblies

Genome_ID	Isolate_ID	Group	Cloning dilution	Complexity of infection	Collection date	Host age	Host sex	NCBI accession number
DC01m1c	DC01m1	Dry Cohort	clone	single-genotype	24/12/2016	9	male	ERR9731534
DC02m1c	DC02m1	Dry Cohort	clone	single-genotype	24/12/2016	11	male	ERR9731535
DC03m1c	DC03m1	Dry Cohort	clone	single-genotype	22/12/2016	13	female	ERR9731539
DC04m2c	DC04m2	Dry Cohort	clone	single-genotype	26/01/2017	15	female	ERR9731540
DC07m1b	DC07m1	Dry Cohort	bulk	single-genotype	23/12/2016	23	female	ERR12263817
DC07m1c	DC07m1	Dry Cohort	clone	single-genotype	23/12/2016	23	female	ERR12263821
DC08m3b	DC08m3	Dry Cohort	bulk	poly-genotypes	13/02/2017	10	male	ERR12263802
DC08m3c	DC08m3	Dry Cohort	clone	single-genotype	13/02/2017	10	male	ERR12263818
DC08m5b	DC08m5	Dry Cohort	bulk	poly-genotypes	19/04/2017	10	male	ERR12263832
DC08m5c1	DC08m5	Dry Cohort	clone	single-genotype	19/04/2017	10	male	ERR12263826
DC08m5c2	DC08m5	Dry Cohort	clone	single-genotype	19/04/2017	10	male	ERR12263801
DC09m5b	DC09m5	Dry Cohort	bulk	poly-genotypes	20/04/2017	12	male	ERR12263810
DC09m5c1	DC09m5	Dry Cohort	clone	single-genotype	20/04/2017	12	male	ERR12263806
DC09m5c2	DC09m5	Dry Cohort	clone	single-genotype	20/04/2017	12	male	ERR12263805
DC10m1c	DC10m1	Dry Cohort	clone	single-genotype	23/12/2016	12	male	ERR9731536
DC10m2c	DC10m2	Dry Cohort	clone	single-genotype	12/01/2017	12	male	ERR9731537
DC10m3c	DC10m3	Dry Cohort	clone	single-genotype	12/02/2017	12	male	ERR9731538
WA01b	WA01	Wet Asymptomatic	bulk	single-genotype	23/11/2016	15	female	ERR12263816
WA01c	WA01	Wet Asymptomatic	clone	single-genotype	23/11/2016	15	female	ERR12263812
WA03b	WA03	Wet Asymptomatic	bulk	poly-genotypes	21/11/2016	27	female	ERR12263820
WA03c	WA03	Wet Asymptomatic	clone	single-genotype	21/11/2016	27	female	ERR12263808
WA04b	WA04	Wet Asymptomatic	bulk	single-genotype	21/11/2016	11	male	ERR12263815
WA04c	WA04	Wet Asymptomatic	clone	single-genotype	21/11/2016	11	male	ERR12263823
WA10b	WA10	Wet Asymptomatic	bulk	single-genotype	21/11/2016	8	male	ERR12263827
WA10c	WA10	Wet Asymptomatic	clone	single-genotype	21/11/2016	8	male	ERR12263807
WA11c	WA11	Wet Asymptomatic	clone	single-genotype	23/11/2016	9	male	ERR12263813

WA47b	WA47	Wet Asymptomatic	bulk	single-genotype	21/12/2016	11	male	ERR12263803
WA47c	WA47	Wet Asymptomatic	clone	single-genotype	21/12/2016	11	male	ERR12263833
WS01b	WS01	Wet Symptomatic	bulk	poly-genotypes	08/11/2016	NA	male	ERR12263825
WS03b	WS03	Wet Symptomatic	bulk	poly-genotypes	31/10/2016	13	female	ERR12263830
WS04b	WS04	Wet Symptomatic	bulk	poly-genotypes	31/10/2016	45	male	ERR12263837
WS05b	WS05	Wet Symptomatic	bulk	poly-genotypes	17/10/2016	54	female	ERR12263811
WS05c	WS05	Wet Symptomatic	clone	single-genotype	17/10/2016	54	female	ERR12263828
WS06c	WS06	Wet Symptomatic	clone	single-genotype	17/10/2016	22	female	ERR12263800
WS07c	WS07	Wet Symptomatic	clone	single-genotype	01/11/2016	10	male	ERR12263814
WS08b	WS08	Wet Symptomatic	bulk	single-genotype	07/11/2016	41	male	ERR12263804
WS09b	WS09	Wet Symptomatic	bulk	single-genotype	07/11/2016	49	female	ERR12263834
WS10b	WS10	Wet Symptomatic	bulk	poly-genotypes	08/11/2016	16	female	ERR12263838
WS11b	WS11	Wet Symptomatic	bulk	single-genotype	08/11/2016	8	male	ERR12263824
WS12b	WS12	Wet Symptomatic	bulk	poly-genotypes	08/11/2016	28	male	ERR12263831
WS12c	WS12	Wet Symptomatic	clone	single-genotype	08/11/2016	28	male	ERR12263822
WS13b	WS13	Wet Symptomatic	bulk	poly-genotypes	18/11/2016	64	female	ERR12263836
WS15b	WS15	Wet Symptomatic	bulk	poly-genotypes	18/11/2016	37	female	ERR12263835

671
672
673   Table 2: Summary statistics for 43 *P. falciparum* genome assemblies.
674

Genome_ID	Genome length	Total contigs	Telomere to telomere contigs	N50	L50	Total genes	var	LARSFADIG motifs	var2csa	rif	stevor
DC01m1c	23.4	20	3	1622224	5	5864	70	61	2	107	24
DC02m1c	23.7	28	5	1693155	5	6056	70	64	1	178	34
DC03m1c	23.6	24	7	1627210	5	6081	68	64	1	156	34
DC04m2c	23.4	24	4	1636287	5	5781	66	62	2	118	25
DC07m1b	25.07	85	6	1433249	7	6222	73	70	4	172	35
DC07m1c	26.7	187	5	1507801	6	7229	60	58	2	154	34
DC08m3b	36.34	113	4	1198427	12	8298	116	115	4	230	43
DC08m3c	24.28	64	5	1631162	5	6157	55	55	2	134	32

DC08m5b	27.88	62	5	1421197	8	6427	105	103	4	234	47
DC08m5c1	24.18	52	6	1606191	5	6085	73	71	2	133	22
DC08m5c2	25	98	4	1630462	5	6324	62	62	2	150	32
DC09m5b	27.99	164	3	1405453	9	6714	116	114	2	236	46
DC09m5c2	24.22	121	2	1661132	5	6669	84	79	1	130	28
DC09m5c1	25.61	58	6	1543257	6	6046	67	66	1	161	33
DC10m1c	23.5	23	7	1556504	5	5725	70	68	1	134	26
DC10m2c	23.7	20	9	1571854	5	5781	71	69	2	154	29
DC10m3c	23.3	27	5	1672417	5	5684	52	63	2	137	23
WA01b	25.85	127	3	1423884	7	6566	65	64	2	116	25
WA01c	23.67	25	2	1637256	5	5756	69	67	2	120	25
WA03b	24.26	49	3	1519022	5	5689	72	69	3	164	37
WA03c	23.79	46	4	1693536	5	5812	70	69	6	154	36
WA04b	24.92	81	2	1446723	7	5656	75	73	4	165	35
WA04c	23.98	44	2	1613267	5	5852	83	76	4	170	37
WA10b	24.19	36	4	1487219	6	5680	82	80	5	162	35
WA10c	26.6	182	2	1468285	6	6799	79	77	5	161	33
WA11c	23.65	24	4	1621859	5	5807	73	70	2	105	25
WA47b	29.85	328	3	1408774	8	8042	91	86	9	177	39
WA47c	23.97	48	2	1693146	5	5681	71	66	4	160	36
WS01b	26.09	104	4	1517931	6	6024	124	119	1	193	39
WS03b	24.11	38	5	1658341	5	5954	79	76	4	132	33
WS04b	30.49	144	7	1440087	8	7081	131	130	5	293	54
WS05b	25.02	56	3	1435358	6	6002	72	70	1	166	45
WS05c	23.81	42	3	1641247	5	5853	68	66	1	153	42
WS06c	24.27	65	2	1560362	5	6050	64	62	4	149	34
WS07c	24.34	73	2	1612653	5	6117	69	66	2	158	37
WS08b	24.05	51	5	1638922	5	5944	67	66	2	116	25
WS09b	24.45	54	2	1653278	5	6160	65	64	1	163	35
WS10b	32.57	123	5	1386579	7	9291	87	85	1	165	39
WS11b	25.63	108	2	1479817	6	6759	69	66	1	152	33

WS12b	33.39	113	6	1356076	10	7836	130	130	2	274	51
WS12c	24.22	45	2	1649284	5	5959	75	75	1	152	27
WS13b	24.09	40	5	1564570	5	5926	75	75	4	174	23
WS15b	24.3	41	5	1592718	5	6051	74	74	4	173	23