1

2 **Mutational divergence in local populations**

3 **of the selfing nematode *Caenorhabditis elegans***

6
7
8

9 Xu Wei[1,2], Aurélien Richaud[1], Robyn E. Tanny[3], Erik C. Andersen[3], Marie-Anne Félix[1*]

11
12 1: Institut de Biologie de l'École Normale Supérieure, Centre National de la Recherche
13 Scientifique, Paris, France
14 2: School of Life Sciences, East China Normal University, Shanghai, China
15 3: John Hopkins University, Baltimore, USA.
16
17 emails: xuweishch@outlook.com; richaud@bio.ens.psl.eu; robyn.tanny@jhu.edu;
18 erik.andersen@jhu.edu; felix@bio.ens.psl.eu
19
20 *: corresponding author: felix@bio.ens.psl.eu
21
22 Xu Wei: ORCID 0000-0002-7074-0878
23 Erik C. Andersen: ORCID 0000-0003-0229-9651

25 Data availability and funding statements are provided at the end of the manuscript as per
26 the specification of the journal. The authors declare no competing interest.

27
28
29
30 Running title: Mutational divergence in selfing populations

Abstract

Laboratory mutation accumulation experiments allow the assessment of spontaneous mutation rates and patterns with minimal selection. Here, we aimed to study the accumulation and fate of mutations in natural populations, in a spatial context. The nematode *Caenorhabditis elegans* is particularly suited for such endeavor, as it reproduces almost exclusively by selfing. We analyzed the evolution of clonal *C. elegans* genotypes along a 300-m long stream bank in the Santeuil wood (France), based on short-read whole-genome sequencing of individuals collected between 2009 and 2022. We followed along years two distinct clones (isotypes), composed of individuals only differing by recent mutations. Recombination was scarce. A temporal signal was detected: strains from earlier years were found close to inner nodes of the tree, while recent ones were found on outer tips. This signal allowed us to estimate a substitution rate of 4 to 5×10^{-8} mutations per base pair per year. Based on the spontaneous mutation rate per generation in laboratory lines, we estimated that *C. elegans* locally undergoes around 25 effective generations per year, which can be used to calibrate divergence times among and within species. Mutation densities were higher on the X chromosome, on chromosome arms, and in non-exonic regions. We detected a high transition-to-transversion ratio, not observed in *C. elegans* laboratory mutation accumulation lines. Finally, using these recent mutations, we detected a spatio-temporal pattern within the field site, indicating limited dispersal at the scale of 100 meters within 10 years.

Keywords: mutation rate, mutational bias, lifecycle, natural populations, polymorphisms, *C. elegans.*

Introduction

Genetic studies of natural populations include evolutionary processes across wide timescales. In a typical population, the genealogy of genotypes integrates numerous recombination events on chromosome fragments, each carrying polymorphisms that arose by mutation across millions of years. Often, the derived allele of these polymorphisms cannot be inferred. At the other end of the spectrum, at very short timescales, isogenized populations in the laboratory can be cultured at minimal population size for several generations of mutation accumulation, allowing to assess rate, patterns and phenotypic effects of spontaneous mutation with minimal selection (Lynch, 1985; Keightley, 1994; Keightley & Caballero, 1997; Vassilieva & Lynch, 1999; Denver *et al.*, 2009; Saxena *et al.*, 2019; Konrad *et al.*, 2019). Immediately above this laboratory timescale, we aimed here to study the mutational process in natural populations that are subject to selection, in their environment. Some organisms naturally provide isogenic populations where mutation can be easily studied in a given genomic background. However, most such species do recombine at a non-negligible rate. For example, extensive genomic surveillance has documented the accumulation of spontaneous mutations in viruses, yet re-assortment and recombination events (Holmes *et al.*, 2005; Shao *et al.*, 2017; Roemer *et al.*, 2023) complicate the inference of mutational trajectories. The crucifer *Arabidopsis thaliana* underwent a narrow initial population bottleneck during a colonization event of the American continent a few centuries ago. Using whole-genome sequencing of museum samples, (Exposito-Alonso *et al.*, 2018) leveraged this bottleneck to analyze the advent and spread of new mutations in a clonal natural lineage with rare recombination.

Studying timescales over many generations is facilitated with organisms displaying short generation times. The nematode *Caenorhabditis elegans* provides an excellent animal system where the appearance and fate of mutations can be directly studied in the wild. Given its fast lifecycle, *de novo* mutation can be detected and studied on the timescale of a few years rather than centuries. *C. elegans* reproduces mostly through selfing XX hermaphrodites (modified females that produce sperm early in adulthood) and rare X0 males arising by spontaneous non-disjunction of the X chromosome or as progeny of a cross. Outcrossing is rare in most natural populations where it was evaluated, and distinct independent clones co-exist in the same location (Haber *et al.*, 2005; Barrière & Félix, 2005; Sivasundar & Hey, 2005; Barrière & Félix, 2007; Richaud *et al.*, 2018; Crombie *et al.*, 2019; Crombie *et al.*, 2022). Based on the comparison between short-term and long-term measures of outcrossing, the cross progeny of distinct genotypes even appear to be counterselected (Barrière & Félix, 2007). This rare effective recombination in selfing *Caenorhabditis* nematodes could be explained by outbreeding depression (Dolgin *et al.*, 2007; Gimond *et al.*, 2013) and gene drive effects (Seidel *et al.*, 2008; Ben-David *et al.*, 2017; Noble *et al.*, 2021). At the species level, each group of highly similar genotypes, defined by a threshold of genomic similarity, is called an 'isotype' (Andersen *et al.*, 2012). Several of these isotypes co-exist locally (Haber *et al.*, 2005; Barrière & Félix, 2005; Sivasundar & Hey, 2005; Barrière &

Félix, 2007; Richaud *et al.*, 2018; Crombie *et al.*, 2022). Here we focus on local polymorphisms *within* an isotype, which have not been studied so far.

One main motivation in studying mutation in a natural population is to estimate the number of effective generations per year for the species, by calibrating the nucleotide substitution rate with a mutation rate per generation estimated in laboratory spontaneous mutation lines. Dating the divergence between nematode species is a challenge, because nematodes have notoriously long branches on animal phylogenies and are poorly fossilized. For example, some *Caenorhabditis* species are as distant in terms of protein sequence evolution than fish and human (Kiontke *et al.*, 2004), yet only represent a small fraction of nematode diversity. Methods that estimate *Caenorhabditis* species divergence dates require an estimate of the number of generations per year (Cutter, 2008; Picao-Osorio *et al.*, 2025; Fusca *et al.*, 2025). The lifecycle of *C. elegans* can be completed within three days in favorable conditions, but in real life development can be arrested at various points, including in a modified third larval stage called the dauer, commonly found in natural populations (Barrière & Félix, 2005; Barrière & Félix, 2007; Félix & Duveau, 2012; Richaud *et al.*, 2018). This raises a question about *C. elegans*' generation time in the wild and the effective number of generations the species undergoes per year. One aim of the present study was thus to estimate the number of mutations and of generations per year in a natural population, with implications for the species' biology as well as for inferring evolutionary divergence times.

C. elegans thrives in patchy bacteria-rich natural habitats provided by decomposing plant substrates, such as stems of herbaceous plants, fruits and flowers (Félix & Braendle, 2010; Félix & Duveau, 2012; Frézal & Félix, 2015; Schulenburg & Félix, 2017; Samuel *et al.*, 2016; Dirksen *et al.*, 2016; Crombie *et al.*, 2022). The species is found throughout the world, mostly in temperate regions, and at a higher elevation in tropical regions (Andersen *et al.*, 2012; Crombie *et al.*, 2019). To investigate within-isotype polymorphisms, we chose to focus on the Santeuil wood in France, a site for *Caenorhabditis* collection since 2009 (Félix & Duveau, 2012; Richaud *et al.*, 2018). The field site consists in a 300-meter-long North-South transect bordered by a small stream on the East side (the Viosne, running North to South), a grass field on the North end, a wood with little undercover of herbaceous plants on the West side and a stream tributary on the South end (Figure 1). We collected samples of decomposing plant material, mostly stems of herbaceous plants such as *Heracleum*, *Symphytum*, *Tussilago*, as well as some invertebrates that presumably carry *C. elegans* between bacterial-rich food sources (Barrière & Félix, 2005; Caswell-Chen *et al.*, 2005; Félix & Duveau, 2012; Petersen *et al.*, 2015; Schulenburg & Félix, 2017). Demographic and developmental stage compositions was the focus of the first study (Félix & Duveau, 2012). In the second study, three local isotypes named HS1, HS2, and HS3 were found using partial genome sequencing of animals collected from 2009 to 2014 (Richaud *et al.*, 2018). These isotypes are not merely recombinant of each other and hardly ever crossed.

The local isotype set in Santeuil provides an exemplary system to study the mutational landscape and spatial structuring of natural populations. Here we present analyses of single-nucleotide substitutions along the evolution of two isotypes in the Santeuil wood, based on short-read sequencing of the whole genomes of strains collected

146 between 2009 and 2022. We aimed to address the following questions: What are the
147 mutation rate and patterns in natural populations of *C. elegans*? How many effective
148 generations per year does *C. elegans* reproduce in the field? What are the spatiotemporal
149 distribution patterns of *C. elegans* at this local scale?

Materials and Methods

Field collection

Caenorhabditis elegans was collected along the Viosne stream in the Santeuil wood as previously described (Félix & Duveau, 2012; Richaud *et al.*, 2018). This stream border includes herbaceous plants of humid areas, such as *Symphytum officinale* (comfrey, Boraginaceae) or *Tussilago farfara* (coltsfoot, Asteraceae), as well as *Heracleum sphondylium* (common hogweed, Apiaceae) and one or several unidentified taller species in the Asteraceae family. We also collected potential invertebrate vectors such as slugs, snails, isopods, insects and annelids. We recorded at the time of collection the position of each sample along the North-South transect using GPS (on a specific hand-held device in earlier years and a smartphone in later years). In addition, we used manual notes to precisely locate samples collected at distances of less than 10 meters (Félix & Duveau, 2012; Richaud *et al.*, 2018). We collected new strains for this work in 2017, 2019, 2020 and 2022. The 2020 collection focused on collecting stems and their associated invertebrates. A list of analyzed strains is provided in Table S1.

Strains were established by selfing of single individuals and assigned a unique identifier (e.g. JU1793). The Santeuil haplotypes are called HS# for Haplotype Santeuil. The HS1, HS2 and HS3 isotypes correspond to isotypes JU1934, JU1793, and JU2600 in CaeNDR (the *Caenorhabditis* Natural Diversity Resource; (Crombie *et al.*, 2023), respectively. We sequenced and analyzed HS2 and HS3 strains from field collection spanning 2009 to 2022, usually a single strain per positive sample. Many more lines were frozen in (Félix & Duveau, 2012; Richaud *et al.*, 2018) and many genotyped by 2b-RAD sequencing in (Richaud *et al.*, 2018).

Genomic DNA sequencing

For many strains, DNA sequence alignments were extracted from the CaeNDR website version 20231213 (Crombie *et al.*, 2023). For the strains from 2022, nematode culture, harvesting, and DNA extraction were performed as in (Richaud *et al.*, 2018). Genomic DNA was sequenced with 150 bp paired-end reads using Illumina sequencing by Eurofins Genomics. The raw sequences of these strains are available at NCBI Sequence Read Archive under accession PRJNA1280162.

Read mapping and calling of single-nucleotide polymorphisms

This study focused on single-nucleotide polymorphism (SNP) in the nuclear genome. The *C. elegans* haploid nuclear genome size is 100 Mb, distributed on six chromosomes (Consortium, 1998). The pipeline is summarized in Figure S1.

New Illumina reads were trimmed and aligned to the *C. elegans* reference genome WS283 (Sternberg *et al.*, 2024) by fastp v0.20.0 (Chen *et al.*, 2018) and BWA v0.7.17 (Li & Durbin, 2009). The sequence alignments were then sorted and marked by sambamba v1.0.0 (Tarasov *et al.*, 2015) and picard v2.26.2 (Institute, 2021). Variant calling was performed using the sequence alignments (.bam files) of all strains identified at CaeNDR (version 20231213) as belonging to the isotypes JU1934 (HS1), JU1793

(HS2), JU2600 (HS3) and JU4161 (HS4) using GATK v4.2.2.0 (McKenna *et al.*, 2010).
Parametrization of read processing and variant calling referred to CaeNDR (Crombie *et*
al., 2023) (version 20231213; https://github.com/AndersenLab). We found in the 2022
samples a new Santeuil genotype HS5. We compared it to all isotypes at CaeNDR and
found it to be novel, represented for example by strain JU4370. This isotype is now
included in CaeNDR (version 20250625).
We evaluated the proportion of the different isotypes for each year of sampling
using 2b-RAD data from (Richaud *et al.*, 2018) covering years from 2009 to 2014 (as
they included many more individuals) and whole-genome sequences for the new field
collections in 2017, 2019, 2020, and 2022. The proportions of isotypes are provided in
Table S2 and plotted in Figure 1e.
Mutation rates and patterns were analyzed separately for isotypes HS2 and HS3.
All strains either found in the Santeuil wood (thereafter "local") or other locations
outside the wood ("external") in each isotype (i.e., HS2 and HS3) were employed for
accurately calling variants. We trimmed the genome (called *trimmed genome*) to
eliminate highly repetitive (masked regions in WormBase, (Sternberg *et al.*, 2024) and
hyper-divergent regions relative to the N2 reference strain (Crombie *et al.*, 2023), as well
as regions with read depths less than six to avoid miscalling. To control for potential
confounding effects of novel mutations arising in the laboratory, we removed new
mutations in the heterozygous state. SNP sets are provided in Tables S3 and S4 for HS2
and HS3, respectively.

Haplotype network construction and recombination
215 Haplotype networks were built using all known strains belonging to HS2 or HS3 by
216 SplitsTree v6.2.1-beta (Huson & Bryant, 2006). Three recombinants were identified
217 within HS2 (Figure S3). For each recombinant, the recombination breakpoints on the
218 chromosomes were inferred based on the distribution of SNPs.
219

Mutation spectra
Based on their locations in exon, intron, non-coding gene, or intergenic regions, SNPs of
local strains were categorized into four groups. Their distribution was visualized on the
trimmed genome. Trimmed genomes and category sizes are provided in Table S5. SNP
density was calculated across the trimmed genome in the following contexts: (1) per
chromosome; (2) in tips, arms, and core regions predicted by (Rockman & Kruglyak,
2009); and (3) in exon, intron, non-coding gene, and intergenic regions. We identified
nine and 27 dinucleotide variants in HS2 and HS3, respectively. For the purpose of these
density calculations, dinucleotide variants were lumped as a single site. Subsequently, a
Chi-squared goodness-of-fit test was employed to assess whether SNPs were evenly
distributed among the categories within each context. With dinucleotide variants treated
as two separate sites, we calculated the densities of all transition and transversion types
and the *Ts/Tv* ratio across exonic, intronic, non-coding gene, and intergenic regions.
233

SNP annotation

Annotation with respect to protein sequence was performed by csq function (Danecek &
McCarthy, 2017) of bcftools v1.13 (Danecek *et al.*, 2021). The SNPs located in coding
regions (CDS) were categorized into non-synonymous and synonymous mutations. A
Chi-squared test was used to assess whether the distribution of non-synonymous and
synonymous mutations differed significantly between wild strains from Santeuil and
mutation accumulation lines maintained in the laboratory (Denver *et al.*, 2009; Denver *et*
al., 2012; Konrad *et al.*, 2019).

Phylogenetic analyses and estimation of effective generations per year
To estimate local substitution rates in the *C. elegans* population from Santeuil, we
implemented a Skygrid coalescent model (Gill *et al.*, 2013) in BEAST v1.10.4
(Drummond & Rambaut, 2007). For each isotype (HS2 and HS3), all SNPs from local
non-recombinant strains were partitioned into four categories: exonic, intronic, non-
coding gene-related, and intergenic regions. A total of 10,000 invariant sites were
randomly generated and evenly distributed among these four categories to meet the base
frequency requirements of the substitution model. Tip dates were calibrated based on
sampling time. Each partition was assigned an uncorrelated relaxed clock model
(lognormal distribution) and a GTR+G+I substitution model, allowing rates to vary
across branches. To minimize the influence of singleton mutations originating from
recent laboratory mutations on substitution rate estimates, we focused on the mean
evolutionary rate along internal branches. The MCMC analysis was run for 100 million
generations, sampling 10,000 steps. A Maximum Clade Credibility (MCC) tree was
summarized using the TreeAnnotator module in BEAST for visualization.

Referring to the substitution rate of intergenic regions of both wild strains in
Santeuil ($\mu_{\text{bs.HS}}$; unit: per site per year) and mutation accumulation lines in the laboratory
($\mu_{\text{bs.MA}}$; (Konrad *et al.*, 2019); i.e., 1.82×10^{-9} per site per generation), we estimated the
number of effective generations *C. elegans* reproduced in Santeuil per year (N_{EG}; unit:
generations per year) by:

$$N_{\text{EG}} = \frac{\mu_{\text{bs.HS}}}{\mu_{\text{bs.MA}}}$$

262 A Skygrid coalescent model was re-ran for HS2 using the 47 local non-recombinant
263 strains and 6 external strains (Figure S4).

Spatio-temporal patterns
Based on the GPS coordinates and notes recorded during field sampling, we mapped the
distribution of different subclades (as defined on Figure 2) using ArcGIS v10.2 (ESRI,
2013). Euclidean distances between strains within each isotype were calculated (Tables
S9-10). For each subclade, a Mantel test with 1,000 permutations was used to evaluate
the correlation between Euclidean distance and sampling date difference among strains,
keeping a single strain per sample (for example, JU1924 for S73). For subclades with
significant Mantel test results, the slopes of regression lines can be interpreted as
estimates of the dispersal rate of *C. elegans* (unit: meters per year) (Figure S5). For this
analysis, the two remaining strains at the root of HS3.4 were conserved (JU1924 and
JU4219). Pairwise Wilcoxon rank sum tests were used to assess whether significant

spatial segregation exists among subclades, keeping a single strain per sample and not
including the HS3.4 root strains (Table S8).
278   Each strain was assigned a one-dimensional coordinate on a simplified linear axis
279   (Figure 6c), with the origin (0) set to the northernmost strain and values increasing
280   southward (Table S1). This one-dimensional location was treated as a trait and used for
281   visualizing the distribution and clustering of strains within each subclade on the MCC
282   trees (Figure S6).
283   All statistical analyses were performed in R v4.1.2 (RCoreTeam, 2023).

Results

Frequencies of co-occurring isotypes

The proportion of isotypes varied across years in our field collections (Scheirer-Hare rank test, year*isotype: $p<0.003$) (Figure 1). The HS1 isotype was predominant in 2009, then decreased and apparently disappeared in later years. The HS2 isotype became predominant, then HS3. Two new isotypes appeared in our collections in 2020 and 2022, which we named HS4 and HS5, respectively. These two isotypes are not mere recombinants of HS1-3, so immigration of new alleles likely occurred (Figure S2).

The HS2 isotype was found in several locations in France, at various distances up to 300 km away (Table S1). The HS3 isotype was only found outside the Santeuil wood a few hundred meters away in an apple (strain JU2600). The other three isotypes (HS1, HS4 and HS5) have so far not been found elsewhere.

Local mutations and rare intra-isotype recombination

After the preliminary clustering in isotypes, we focused on intra-isotype polymorphisms, using whole-genome sequencing data of strains representing HS2 (n=56 strains) and HS3 (n=99 strains). As expected, the intra-isotype polymorphism level was markedly smaller than that among the local isotypes (Figure 1d). For each isotype, we trimmed the genome of repeated and highly divergent regions to focus on genomic regions allowing high quality calls of single-nucleotide polymorphisms (SNPs). By distinguishing Santeuil wood strains ("local") from other strains of the same isotype ("external"), we called a total of 1165 SNPs for 56 HS2 strains within the 63.07 Mb of the HS2 trimmed genome, in which 600 SNPs only occurred in six external strains, and 69 SNPs only occurred in three recombinants. After excluding both external and recombinant strains, 496 SNPs were found to be polymorphic among the 47 local HS2 strains. Similarly, a total of 1214 SNPs in 66.46 Mb were called for 99 HS3 strains. After excluding the single external strain (JU2600), 1147 SNPs were found in the 98 local HS3 strains.

From these polymorphisms, we analyzed the relationships between strains of a given isotype by building haplotype networks including both local and external strains (Figures 2 and S3). The trees in Figure 2 represent 1096 SNPs for HS2 and 1214 SNPs for HS3. For HS2, the external strains from outside Santeuil appeared at the same branching point. The positions of these external strains supported the idea that the mutations among Santeuil HS2 strains may have occurred locally in recent years; in other terms, these external strains can be considered as outgroups. In addition, most polymorphisms called among strains of each isotype were not found elsewhere in the CaeNDR strain set - the exceptions are likely due to convergent mutations or sequencing/mapping errors, especially from mononucleotide runs and segmental duplications that were not removed during trimming. Specifically, 87.9 and 85.5% of the polymorphisms within HS2 and HS3, respectively, were only found in the Santeuil wood. The detected polymorphisms did not cluster in a genomic region (Figure 3a), also consistent with them originating from new mutation rather than recombination. Finally, many of these polymorphisms could be found in several Santeuil strains, indicating that

they appeared in the field and not subsequent laboratory culture. We conclude that the detected polymorphisms represent new, derived mutations occurring recently in the genomic background of this isotype. The mutational change could be polarized by inferring that the Santeuil private allele was derived.

We then wondered whether outcrossing could be more frequent *within* isotypes than that previously estimated *between* isotypes, since outbreeding depression may be less relevant in the former case. For HS2, two recombination events occurred, as revealed by the diamond shapes in the network including all strains (Figure S3a,b), and concerned a total of three strains in our dataset. Specifically, with the SNP set that we could analyse, the JU3785 and JU3786 strains displayed whole-chromosome reassortments between the inferred parental genotypes, while JU3399 displayed one recombination event on four of the six chromosomes (Figure S3c,d). Each of these two events thus appeared consistent with a single outcrossing event. Removing the three recombinant strains resulted in a tree-like pattern, where the relationships between strains could be solely explained by mutation (Figure 2a). For the sake of simplicity, we excluded recombinants to infer mutational rate and pattern below. In the HS3 isotype, no recombination was detected or if so, only concerned so few mutations that this might correspond to the resolution of two recent mutations in the heterozygous state. We thus conclude that effective outcrossing events are also rare within these local isotypes.

Mutational pattern

We analyzed the distribution of new mutations with respect to the six chromosomes, chromosome regions, gene annotation and nucleotide change. We performed these analyses on the local strains of the HS2 and HS3 genotypes separately (Figures 3 and 4). These analyses use 496 SNPs for HS2 and 1147 SNPs for HS3 and are thus more powered for the latter.

The distribution of mutations among the six chromosomes was uneven (Chi-Square Goodness of Fit Test for even rates among chromosomes: $p = 0.02$ for HS3, $p > 0.05$ for HS2). The X chromosome displayed the highest density, with 22% more substitutions per bp than autosomes for HS3; the same trend was true for HS2 (Table S5).

We then partitioned each chromosome in core, arms, and tips, following recombination regions from (Rockman & Kruglyak, 2009), where arms are defined by a higher recombination rate. The core (central) region of *C. elegans* holocentric chromosomes recombines at a lower rate and harbors fewer polymorphisms on average at the species scale, fewer repeated elements, and more conserved and essential genes (Consortium, 1998; Fraser *et al.*, 2000; Rockman & Kruglyak, 2009; Andersen *et al.*, 2012; Crombie *et al.*, 2019). In the HS3 dataset, arms carried significantly more mutations per site than other chromosomal regions (Figure 3d; Chi-Square Goodness of Fit Test for tips, arms and core: $p = 0.002$ for HS3, $p > 0.05$ for HS2, although with the same trend).

We then partitioned the SNPs in different genic regions (exons, introns, intergenic, non-coding gene). Exons displayed a significantly lower mutation density and substitution rate than the other genic regions (Chi-Square Goodness of Fit Test for

density in different genic regions: $p = 0.02$ for HS3, with the same trend in HS2) (Figure
3c, Table 1, Table S5).
We could polarize each mutation and thus classify single-nucleotide substitutions
in six chemical categories (Figure 4). GC to AT transitions were particularly frequent in
both isotypes, suggestive of mutations induced by oxidative damage (Figure 4). The
expected ratio of transitions to transversions (Ts/Tv) should be 0.5 if the mutation pattern
were random. In our dataset, transitions were found more frequently than transversions,
with a Ts/Tv of 1.20 for HS2 and 1.15 for HS3 (Table 2). At this whole-genome level,
the two isotypes did not exhibit significant differences (Chi-Square test, $p > 0.05$). In
HS3, the Ts/Tv ratio was noticeably lower in intronic regions (Table 2).
The *C. elegans* genome contains less G/C than A/T bases (36% CG), although
this trend is less strong in exons (Table S5). We weighted the six substitution rates with
the GC content of each genic region category. The mutational pattern resulted in a bias
towards AT in exons, but this bias was absent or reversed outside exons (Table S6). To
assess the effect of mutation on GC content evolution, we estimated the relative
proportion of mutations from GC to AT or TA over those from AT to GC or CG.
Weighting the number of relevant substitutions in each genic region, we found ratios of
1.75 and 0.85 for HS3 and HS2, respectively, thus a net mutational bias towards a higher
AT content in HS3. However, we note the difference between the two isotypes.
Finally, using coding gene annotations, we estimated SNP density of non-
synonymous versus synonymous mutations within coding sequences (Table S7). We
found ratios of 2.3 and 3.0 for HS2 and HS3, respectively. We did not detect a
significantly different ratio in the Santeuil sets compared to the mutation accumulation
lines (Chi-Square test, $p > 0.05$).
395
Temporal signal: mutation rate and number of generations per year
The branching pattern over time differed for strains of the two isotypes. For HS2, the
Santeuil wood lineages diverged from the external strains along two branches that could
correspond to two independent immigration events or to local mutation and selection
trimming away branches (Figure 2a). By contrast, the HS3 isotype strains showed a star-
like relationship, consistent with a single founder and lineage expansion along the years
(Figure 2b). A distinct temporal signal can be seen: the strains collected in earlier years
are on inner nodes of the phylogeny, while recently collected ones are on the outside
(Figure 2). The temporal pattern was particularly clear for HS3 and appeared weaker for
HS2, in part because of its decline.
Using sample collection dates, we inferred the most credible evolutionary
scenario using Bayesian Maximum Clade Credibility (MCC) trees, for HS2 and HS3
separately (Figure 5). Given the lack of detected recombination with this dataset, the
topology of the tree did not carry ambiguities, while the dates of nodes were the best
estimates. The two main subclades of the HS2 tree branched around year 1990 while the
HS3 tree was rooted around year 2000. When adding the external strains of HS2, a
possible isotype root was estimated to have occurred before 1960 (Figure S4).
Based on the MCC trees built using local non-recombinant strains, we estimated
the average number of substitutions per year for each isotype. In order to exclude

mutations that could have occurred in the laboratory, we removed the branches at the tree tips. In addition, we selected intergenic regions, as the rate per site per year was significantly lower in exons (see above, Figure 3c), which are more likely to be under purifying selection. In intergenic regions, mutation rates for the HS2 and HS3 natural populations were estimated to 4.13 and 4.92×10^{-8} mutations per site per year, respectively, with overlapping confidence intervals (Table 1). With a 100 Mb genome, this corresponds to 4 to 5 mutations per haploid genome per year.

To convert this substitution rate in number of generations per year, we used the spontaneous mutation rate per generation measured using laboratory mutation accumulation lines. Two recent independent studies gave similar results (Konrad *et al.*, 2019; Saxena *et al.*, 2019). Based on the rate of 1.82×10^{-9} per site per generation in intergenic regions from (Konrad *et al.*, 2019), we estimated the number of effective generations in Santeuil natural populations to 22.7 (15.6 - 30.4) and 27.0 (22.6 - 31.3) generations per year for HS2 and HS3, respectively.

Spatial structure within the Santeuil field site

We then used the new mutations detected above to study spatial structure and dynamics in the field. Because the trees in Figure 2 represent actual animal lineages across generations, the fate of mutations at the root of each clade could be followed in space - bearing in mind the spare sampling of our collection compared to the actual population size.

We first plotted the data on two-dimensional spatial maps (Figure 6), subdividing each isotype into main subclades: the HS2 isotype into two main subclades plus three recombinants, and the HS3 isotype into six main subclades, as shown in Figures 2 and S5. For HS2, the two subclades were found from North to South and were not significantly spatially segregated from each other (Wilcoxon Rank Sum Test, *p*>0.05). Strikingly, the three strains deriving from the two independent recombination events were found in a central region of the transect almost devoid of other HS2 strains - however, this corresponded to few cases. In contrast, for the six main HS3 subclades, which branched more recently than the two HS2 main subclades, we found highly significant segregation, for example of HS3.1 (North) with HS3.3 or HS3.5 (South) (Figure 6, Table S8).

We further asked whether we could detect a spatio-temporal signal within each subclade. We reasoned that if a new mutation occurred at one end (or even outside) of the transect, we may detect a spatio-temporal signal of dispersal. Mantel tests between pairwise Euclidean distance of strains and their sampling date distance found that two of eight tested HS3 subclades showed a significant signal of directional migration (Figure S5), both from South to North. The regression slopes can be interpreted as estimates of the dispersal rate of the *C. elegans* lineage carrying the mutations at the root. The subclade showing the strongest signal, HS3.4.1, gave a slope of 6 meters per year (C.I. 4.8-7.2), which if sampling was sufficient, is a minimal estimate of the maximal dispersal speed.

To represent the spatial data including sampling year and all genetic relationships, we simplified the geographical coordinates as a one-dimensional projection

on the transect line (Figure 6c) and colored the tips of the phylogenetic tree by the spatial position of the strain (Figure S6). Clustering of spatial positions on the phylogenetic tree is apparent, including across years. The broad patterns of the HS3 subclades are visible: for example, HS3.1 strains in a Northern position (smaller coordinate values) and HS3.3 and HS3.5 in a Southern position. In addition, smaller subclades within HS2 and HS3 show clustering, with the caveat of limited sampling.

Clustering of different subclades was only apparent because movement across the transect occurred. Focusing on the early years of the HS3 tree, the same mutation at the root of the sister subclades HS3.1 and HS3.5 was found in strains collected on opposite transect sides, implying movement across 200 meters at the root or early history of one or both subclades. Compared to the HS3.1 strains located in the North, early-year strains of basally branching subclades were almost all found in the Southern half of the transect, suggesting a northwards movement in the HS3.1 root, even prior to within-clade movement of Figure S6. Alternatively, this among-subclade pattern may be explained by repeated southwards movement. At a fine temporal scale, some closely related pairs of strains, for example JU4135/JU4168, JU4138/JU4166 or JU3788/JU4154 show evidence of large movement (Figure S6). These 1 mm long nematodes can move by themselves, yet faster dispersal vectors are larger invertebrate carriers (Barrière & Félix, 2005; Caswell-Chen *et al.*, 2005; Barrière & Félix, 2007; Félix & Duveau, 2012; Petersen *et al.*, 2015). We wondered whether these animals carried C. *elegans* with the same genotype as the stem on which they were collected, that is whether, in this stem, *C. elegans* may have disembarked from or embarked onto this larger invertebrate (Tables S1, S11). We found one case of concordance between the *C. elegans* genotype on the animal and the plant, six cases of discordance, and one mixed case where two different carriers gave contrasted results (Table S11). A limitation of this analysis is that a plant substrate itself - or an animal - often carries different *C. elegans* genotypes (Richaud *et al.*, 2018).

In summary, despite the fact that migration occurred, the mutated alleles at the root of these subclades did not all spread across the 300 m-long field site within 15-20 years, giving a bound on dispersal at this scale. On the other side of the coin, our observations imply cases of movement across the transect in the span of one or few years.

Discussion

Number of nucleotide substitutions and generations per year

In this study, we estimated in a *C. elegans* natural population a nucleotide substitution rate of 4 to 5×10^{-8} mutations per base pair per year. Calibrated on the laboratory spontaneous mutation rate, this rate translates into 15 to 30 effective generations per year. The estimates are somewhat lower for HS2 than for HS3 yet with overlapping credible intervals. A lineage remaining at low frequency or close to local extinction, such as HS2, could undergo fewer effective generations per year than a then thriving lineage such as HS3. Thus, surviving lineages in these populations displayed an average generation time of about two weeks, which is several-fold higher than the minimal generation time of three days in favorable laboratory conditions.

Two factors likely contribute to a longer lifecycle in natural populations. First, the duration of direct developmental cycles is strongly affected by environmental factors, such as temperature, food and pathogens. For example, development can last over a week at temperatures below 15°C, often encountered in the fall in Santeuil. Second, resources are limited, resulting in developmental arrest of individuals, including dauer diapause. In individual lineages, generations with fast direct development alternate with generations developing via a dauer stage. Given the observed census size and developmental stage composition of populations, dauer induction at high population density and low food, and the patchiness of suitable substrates, it is probable that no more than two to three generations of direct development occur before an individual lineage is induced to dauer (Félix & Duveau, 2012). Diapause can last for months when resources are scarce.

In temperate areas like Santeuil, the generation time likely varies seasonally, with two consequences on *C. elegans* generation time: 1) the availability of food patches peaks in the fall with the bulk of plant decomposition. 2) development slows down progressively during the fall when temperatures decrease. We previously found that, in an apple orchard in Orsay, near Paris, France, *C. elegans* proliferates in the fall and not summer and winter (Félix & Duveau, 2012); a fall peak of *C. elegans* was also detected in compost heaps in Germany by (Petersen *et al.*, 2014). In Santeuil, we collected *C. elegans* in the fall when decomposing stems are easier to find. We note that the Santeuil wood undergrowth is more shaded and humid than the Orsay orchard apples, thus cooler in summer, and may harbor *C. elegans* populations for a longer time. We indeed found the species in summer at the two timepoints when we collected (06 Jul 2012, 07 Aug 2013) (Richaud *et al.*, 2018) (Table S1). In summary, *C. elegans* development may occur fast when conditions are optimal in late summer/early fall, then temperatures decrease in the fall and the generation time likely slows down. Given the diapause arrests and the environmental modulation of the direct lifecycle, our estimate of 15 - 30 generations per year appears consistent with the range of expectations. Some lineages may undergo fewer generations than others in a given year, but the likelihood that they contribute to the next generations would be smaller.

A limitation concerning the conversion of mutation to generation numbers is that the mutation rate in the laboratory may not match that in the wild. First, we used

estimates from the typical laboratory environment at 20°C with direct development and
the rate may vary with the environment. However, by testing the effect of an oxidative
environment, *a priori* expected to contribute to mutation, (Rajaei *et al.*, 2021) found that
mutation rate and pattern were not strongly affected. Second, the mutation rate could
differ among genetic backgrounds. However, (Konrad *et al.*, 2019) did not report a large
variation among *Caenorhabditis* strains.

Using the present estimates, divergence between *Caenorhabditis* species was
recently dated by us and others (Picao-Osorio *et al.*, 2025; Fusca *et al.*, 2025). The
extrapolation of this local measure should in principle be taken with caution, as
demography, temperature and duration of the lifecycle will also vary with the species.
For example, on one hand tropical species may see a higher number of generations per
years; on the other hand, some species develop slower than *C. elegans*, at least in
laboratory conditions, for example *C. inopinata* (Kanzaki *et al.*, 2018) or more distant
species in the genus (Stevens *et al.*, 2019). Interestingly, Fusca et al. (2025) found low
variation among evolutionary branches of the *Caenorhabditis* genus in the substitution
rate. Thus, although further measures in different populations and species would be
welcome, using this temperate *C. elegans* population appears a useful first estimate of
the number of generations per year.

Genomic mutation pattern and detection of selection
We compared the mutation and SNP density pattern to i) a null expectation with no
mutational bias; ii) that observed in laboratory mutation lines, where selection is
minimal; ii) in some cases the species-wide pattern.

We found a higher SNP density on chromosome arms compared to centers and
tips, corresponding to regions where recombination is highest. This mutation pattern is
similar to that detected in laboratory mutation accumulation lines (Konrad *et al.*, 2019;
Saxena *et al.*, 2019) and thus consistent with meiotic events of recombination being
mutagenic. An alternative selective explanation is possible, essential genes being
enriched in central regions of *C. elegans* chromosomes (Kamath *et al.*, 2003).

More surprisingly, we detected a significantly higher SNP density on the X
chromosome compared to the autosomes, albeit of modest size (Table S5). This pattern
contrasts with 1) the pattern in laboratory mutation accumulation lines from the N2 strain
(Konrad *et al.*, 2019), where the sex chromosome had the lowest rate of substitutions; 2)
the species-wide pattern where chromosome X carries only 14.1% of the polymorphisms
(CaeNDR version 20250625, using hard-filtered VCF), whereas 23.9% of our intra-
isotype polymorphisms are on chromosome X. The different chromatin context of the X
chromosome compared to autosomes (Kelly *et al.*, 2002) may modulate the mutation rate
but cannot explain the results in these three conditions. Moreover, the GC nucleotide
content of the X chromosome is not higher than that of the autosomes (Table S5). The
fast rate of X evolution in Santeuil may be due to the genetic background: the *C. elegans*
PB306 strain showed a marginally significantly higher mutation rate on the X
chromosome compared to autosomes, whereas the N2 reference strain did not (Denver *et*
al., 2012). Alternatively, the fast rate of X evolution in the natural populations could be
due to a population genetic effect (selection), such as more frequent purifying selection

on autosomal exons. In line with this hypothesis, the proportion of nucleotides in exons is of 34.4 to 41.9% on autosomes but of 29.0% on the X chromosome, and essential genes are depleted on the latter (Kamath *et al.*, 2003). An alternative explanation for the "faster X" evolution at larger timescale is positive selection of new mutations - however, this seems unlikely to affect patterns at the scale studied here.

A possible sign of purifying selection in our data is indeed the depletion of substitutions in exons, over the whole genome. The SNP density in exons is 20 and 23% lower than in intergenic regions and introns, respectively, which cannot be explained by the GC content of exons. This pattern resembles that in laboratory mutation accumulation lines, where selection is minimal yet present (Konrad *et al.*, 2019; Saxena *et al.*, 2019). That this results from selection is tantalizing, but alternative hypotheses related to nucleotide context, chromatin structure, transcription and DNA repair mechanisms are possible. Within exons, we did not detect a difference in the ratio of non-synonymous to synonymous mutations compared to laboratory mutation accumulation lines, likely due to a lack of statistical power at this scale.

The transition-to-transversion ratio in the local natural populations is almost double to that in mutation accumulation lines in the N2 reference background, more similar to those in the PB306 background (Table 2), and close to species-wide polymorphisms (1.29; CaeNDR 20250625, using hard-filtered VCF). The higher rate of transitions may derive from the mutational pattern that favors chemical conservation of purines and pyrimidines, for example through the deamination of cytosine to thymine. Alternatively, the difference between our dataset and mutation accumulation lines could be explained by selection favoring nucleosome conservation or, in exons, synonymous or conservative amino-acid changes that are favored by transitions (Babbitt & Cotter, 2011). Finally, as the statistical difference between HS2 and HS3 mutational patterns exemplifies, the genetic background and environment contexts may influence this mutational pattern as well.

In summary, the estimated 500-generation evolutionary timescale on a selfing metapopulation may allow the detection of purifying selection on new mutations at the genome scale, yet the signal is weak if any. A particular contrast with the laboratory mutation accumulation lines is the higher mutation rate on the X chromosome (both isotypes) and the high Ts/Tv ratio for the HS3 isotype. Studies of mutation using the Santeuil genetic backgrounds in the laboratory may help in the future to determine whether this is due to the difference genetic background; alternatively, the natural environment, or selection, may influence these mutational patterns.

Spatial structure and isotype turnover
This study along a 300 m field site could detect a spatial structure of new mutations across 14 years, thus providing a picture of eco-evolutionary processes at a fine spatio-temporal scale.

C. elegans proliferates in patchy bacteria-rich decomposing plant matter. Large populations (over 100,000 individuals) always include development of young larvae into the dauer diapause stage, which forms the basis for a boom-and-bust dynamics (Félix & Duveau, 2012; Richaud *et al.*, 2018). In the Santeuil field site, herbaceous plant stems

(10 cm to 1 meter) start decomposition at their base and fall in succession, chiefly in
autumn, covering the ground in a discontinuous manner. Therefore, the density
distribution of *C. elegans* is highly heterogeneous across the site at a given timepoint.
Compared to the well-studied and larger meadow patches of the Glanville fritillary
butterfly in Finland (Hanski, 2011; Multigner *et al.*, 2025), these *C. elegans* patches are
smaller and ephemeral. Across the field site, differences are visible, with a higher density
of decomposing plants at the Northand South ends compared to the middle and a
different plant composition: for example, the coltsfoot *Tussilago farfara* was only found
on the North side across all sampled years. Finally, at a larger scale, the Viosne stream
bank is a stable habitat patch over these years, more similar to a Glanville fritillary
butterfly meadow patch. Thus, metapopulation structure may occur at several spatio-
temporal scales for *C. elegans*.

In this context, dispersal is a key parameter for population dynamics. Our
mutation tracking provides a spatio-temporal scale at which *C. elegans* does not freely
disperse. Specifically, the mutations at the root of the different HS3 subclades, estimated
to have occurred between 2002 and 2008, have not all spread through the whole 300 m
transect, which is compatible with a dispersal rate estimate of few meters per year
detected within one subclade. Field sampling of a microscopic animal is inherently
limited given the large population census size, which alters our ability for accurate
dispersal estimates. Overall, our results thus show that *C. elegans* dispersal is limited at
scales of 100 m over ten years.

The structure observed when considering new mutations within an isotype is
likely mostly mediated by limited dispersal capacity, rather than differential selection. *C.
elegans* dispersal can occur via both active self-movement and transport by animal
vectors. Self-movement can be quantified in the laboratory on a surface (Husson *et al.*,
2013) or in a volume (Kwon *et al.*, 2015). *C. elegans* adults can move at an instantaneous
speed of 0.2 mm/s on surfaces without food (Ramot *et al.*, 2008). Extrapolating this
instantaneous speed would allow movement over 20 m per day - as unlikely as Howard
Berg's imaginary molecule moving through a classroom in one second according to its
instantaneous Brown motion speed (Berg, 1993). *C. elegans* indeed uses biased random
walks, even when chemotacting towards an odour, with turns and bouts of backward
movement (Pierce-Shimomura *et al.*, 1999; Roberts *et al.*, 2016) analogous to runs and
tumbles of *E. coli* (Berg & Brown, 1972). *C. elegans* diffusion coefficient on smooth
agar surfaces is on the order of 0.01 mm^2/s (Helms *et al.*, 2019). In real life, the animals
slow down on food and, across the reproduction cycle, larvae are slower than adults;
further delays occur in the three-dimensional natural environment, with liquid volumes
where swimming is inefficient, high surface complexity, surface tension, stickiness,
bacterial biofilms affecting movement (Darby *et al.*, 2007). How far a single animal -
during its lifetime until reproduction - would move remains unclear, but likely does not
exceed a few decimeters for long-lived dauer larvae and centimeters for direct developers.
At these short distances, nematodes can move in a random or semi-directional manner
within a substrate and possibly between adjacent plant stems.

Longer-range dispersal of *C. elegans* may be mediated by physical causes such as
rain, wind, and here the waterstream along the field site. Concerning the latter, fast

movement appears to often run northwards, in opposite direction to the stream, such as mentioned for HS3.1 (root and within-clade dispersal) and also exemplified by single strains with a Northern coordinate within a Southern clade (Figure S6). Alternatively, slugs, snails, isopods and insects are able to disperse *C. elegans* individuals at medium-range distances (meters) (Barrière & Félix, 2005; Caswell-Chen *et al.*, 2005; Félix & Duveau, 2012; Petersen *et al.*, 2015; Schulenburg & Félix, 2017; Fialova & Tuf, 2025). Maximal movement speed was found in the JU4135-JU4168 pair of the HS3.4.2 lineage (Figure S6): the two strains were found at a distance of 173 meters and did not display any SNP divergence in our dataset, thus, with an average substitution rate of 5 mutations per year, diverged for less than a year. We caution that this maximal speed is sensitive to possible strain errors; however, the occurrence of several similar cases indicates occasional faster movement in an overall static population. Likely less frequent contributors to longer-range movement are vertebrates such as birds and mammals (including humans), which could move *C. elegans* within the site or further (kilometers). For example, the HS2/JU1793 isotype was found at various distances up to 300 km with a divergence date estimated to 50 years (Figure S4). Strains of the same isotype can be found on different continents, albeit rarely (Andersen *et al.*, 2012; Lee *et al.*, 2021): the "MY23" isotype appears to be a dispersal record holder, having dispersed in recent years across three continents (South America, Europe, and East Asia). The present threshold for *C. elegans* strains to be called of the same isotype at CaeNDR corresponds to about 1000 SNPs, thus 100 years of mutation on each branch (if the strains were sampled around the same years and no recombination occurred). As in the presetn study, tracking of recent mutations is a convenient approach for assessing dispersal in natural populations of microscopic organisms and will be of great value for future eco-evolutionary studies.

While the spatial structure within an isotype is largely mediated by dispersal ability, different isotypes and their spatio-temporal distribution could be affected by selection. During the 14-year period, we observed a turnover of isotypes with immigration of new alleles. One isotype (HS1) faded away early in our samples. The two HS2 local subclades branched around year 1990, which also corresponded to their divergence with external strains, and faded in the late years of our study. The local HS3 tree was rooted around year 2000 (Figure 5); given the lack of a clear outgroup, the immigration of the first individuals in Santeuil could have preceded this date. HS4 appeared in our samples in 2020, and HS5 in 2022. We cannot rule out that these isotypes were proliferating locally at low frequency in the early years, yet we find it unlikely that the larvae could subsist for several years as a "dauer bank" (by analogy with seed banks of plants). Therefore, the Santeuil wood appears to harbor a succession of selfing isotypes, some of them with a residence time of at least 20 years, and co-occurrence of a few isotypes at a given time. Neutral processes could contribute to isotype distribution and turnover dynamics, for example through limited dispersal and population bottlenecks favoring stochastic losses. Alternatively, micro-habitat heterogeneity in parameters such as microbial community, predators or physicochemical properties could result in differential selection. Laboratory competition experiments have demonstrated that diversifying selection, for example on resistance to pathogens or other

stresses, may help maintain several local isotypes on timescales of a few generations
(Richaud *et al.*, 2018). At a larger scale, how intra- and inter-annual environmental
variables, such as fluctuations in temperature, precipitation, or pathogen prevalence,
affect the persistence of local isotypes remains to be investigated.
In conclusion, the present study provides an exemplary case for eco-evolutionary
research of nematode natural populations at a fine scale. Studying metapopulations of
these small animals requires the consideration of different spatio-temporal scales,
dominated by different processes. Studies at a larger spatial scale may help understand
patterns of recombination that lead to isotype diversity and thus complement the present
study of local mutational patterns.
720

Acknowledgements
We are grateful to Gaotian Zhang and Joëlle Barido-Sottani for help and discussions on
data analysis. We thank Asher Cutter and Amanda Gibson for constructive comments on
the manuscript. We thank Buck Samuel, Robert Luallen, Patrick Phillips and members of
the Félix lab for their participation in the field. We thank WormBase. The
Caenorhabditis Natural Diversity Resource (RT, ECA) is funded by a NSF Capacity
grant (2224885). This work was funded by the Centre National de la Recherche
Scientifique and a visiting PhD scholarship #202106140037 from the China Scholarship
Council to XW.

References

Andersen E, Gerke J, Shapiro J*, et al.* (2012) Chromosome-scale selective sweeps shape *Caenorhabditis elegans* genomic diversity. *Nat. Genet.* **45**, 285-290.

Babbitt GA, Cotter CR (2011) Functional conservation of nucleosome formation selectively biases presumably neutral molecular variation in yeast genomes. *Genome Biol Evol* **3**, 15-22.

Barrière A, Félix M-A (2005) High local genetic diversity and low outcrossing rate in *Caenorhabditis elegans* natural populations. *Curr. Biol.* **15**, 1176-1184.

Barrière A, Félix M-A (2007) Temporal dynamics and linkage disequilibrium in natural *C. elegans* populations. *Genetics* **176**, 999-1011.

Ben-David E, Burga A, Kruglyak L (2017) A maternal-effect selfish genetic element in *Caenorhabditis elegans*. *Science* **356**, 1051-1055.

Berg HC (1993) *Random walks in biology*, revised edition edn. Princeton University Press.

Berg HC, Brown DA (1972) Chemotaxis in *Escherichia coli* analysed by three-dimensional tracking. *Nature* **239**, 500-504.

Broad Institute (2021) Picard Toolkit. v2.26.2. http://broadinstitute.github.io/picard/

Caswell-Chen EP, Chen J, Lewis EE*, et al.* (2005) Revising the standard wisdom of *C. elegans* natural history: ecology of longevity. *Sci. Aging Knowl. Environ.* **40**, pe30.

Chen S, Zhou Y, Chen Y, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884-i890.

Consortium, The *C. elegans* Sequencing (1998) Genome sequence of the nematode *C. elegans*: a platform for investigating biology. *Science* **282**, 2012-2018.

Crombie TA, Battlay P, Tanny RE*, et al.* (2022) Local adaptation and spatiotemporal patterns of genetic diversity revealed by repeated sampling of *Caenorhabditis elegans* across the Hawaiian Islands. *Mol Ecol* **31**, 2327-2347.

Crombie TA, McKeown R, Moya ND*, et al.* (2023) CaeNDR, the *Caenorhabditis* Natural Diversity Resource. *Nucleic Acids Res* **52**, D850-D858.

Crombie TA, Zdraljevic S, Cook DE*, et al.* (2019) Deep sampling of Hawaiian *Caenorhabditis elegans* reveals high genetic diversity and admixture with global populations. *Elife* **8**, e50465.

Cutter AD (2008) Divergence times in *Caenorhabditis* and *Drosophila*, inferred from direct estimates of the neutral mutation rate. *Mol. Biol. Evol.* **25**, 778-786.

Danecek P, Bonfield JK, Liddle J*, et al.* (2021) Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008.

Danecek P, McCarthy SA (2017) BCFtools/csq: haplotype-aware variant consequences. *Bioinformatics* **33**, 2037-2039.

Darby C, Chakraborti A, Politz SM*, et al.* (2007) *Caenorhabditis elegans* mutants resistant to attachment of Yersinia biofilms. *Genetics* **176**, 221-230.

Denver DR, Dolan PC, Wilhelm LJ*, et al.* (2009) A genome-wide view of *Caenorhabditis elegans* base-substitution mutation processes. *Proc Natl Acad Sci U S A* **106**, 16310-16314.

Denver DR, Wilhelm LJ, Howe DK, *et al.* (2012) Variation in base-substitution
mutation in experimental and natural lineages of *Caenorhabditis* nematodes.
Genome Biol Evol **4**, 513-522.
Dirksen P, Arnaud Marsh S, Braker I, *et al.* (2016) The native microbiome of the
nematode *Caenorhabditis elegans*: gateway to a new host-microbiome model.
BMC Biol **14**, 38.
Dolgin ES, Charlesworth B, Baird SE, Cutter AD (2007) Inbreeding and outbreeding
depression in *Caenorhabditis* nematodes. *Evolution* **61**, 1339-1352.
Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by
sampling trees. *BMC Evol Biol* **7**, 214.
Exposito-Alonso M, Becker C, Schuenemann VJ, *et al.* (2018) The rate and potential
relevance of new mutations in a colonizing plant lineage. *PLoS Genet* **14**,
e1007155.
Félix M-A, Braendle C (2010) The natural history of *Caenorhabditis elegans*. *Curr*
Biol **20**, R965-969.
Félix MA, Duveau F (2012) Population dynamics and habitat sharing of natural
populations of *Caenorhabditis elegans* and *C. briggsae*. *BMC Biol* **10**, 59.
Fialova R, Tuf IH (2025) Spatial activity and sheltering behaviour of terrestrial
isopods (Isopoda, Oniscidea): a field experiment. *Zookeys* **1225**, 127-139.
Fraser AG, Kamath RS, Zipperlen P, *et al.* (2000) Functional genomic analysis of *C.*
elegans chromosome I by systematic RNA interference. *Nature* **408**, 325-330.
Frézal L, Félix MA (2015) *C. elegans* outside the Petri dish. *Elife* **4**, e05849.
Fusca DD, Dall'Acqua MN, Sanchez-Ramirez S, Cutter AD (2025) Phylogenomic
timetree-calibrated speciation clocks for *Caenorhabditis* nematodes reveal
slow but disproportionate accumulation of post-zygotic reproductive isolation.
PLoS Genet **21**, e1011852.
Gill MS, Lemey P, Faria NR, *et al.* (2013) Improving Bayesian population dynamics
inference: a coalescent-based model for multiple loci. *Mol Biol Evol* **30**, 713-
724.
Gimond C, Jovelin R, Han S, *et al.* (2013) Outbreeding depression with low genetic
variation in selfing *Caenorhabditis* nematodes. *Evolution* **67**, 3087-3101.
Haber M, Schüngel M, Putz A, *et al.* (2005) Evolutionary history of *Caenorhabditis*
elegans inferred from microsatellites: evidence for spatial and temporal
genetic differentiation and the occurrence of outbreeding. *Mol. Biol. Evol.* **22**,
160-173.
Hanski IA (2011) Eco-evolutionary spatial dynamics in the Glanville fritillary
butterfly. *Proc Natl Acad Sci U S A* **108**, 14397-14404.
Helms SJ, Rozemuller WM, Costa AC, *et al.* (2019) Modelling the ballistic-to-
diffusive transition in nematode motility reveals variation in exploratory
behaviour across species. *J R Soc Interface* **16**, 20190174.
Holmes EC, Ghedin E, Miller N, *et al.* (2005) Whole-genome analysis of human
influenza A virus reveals multiple persistent lineages and reassortment among
recent H3N2 viruses. *PLoS Biol* **3**, e300.

Huson DH, Bryant D (2006) Application of phylogenetic networks in evolutionary
studies. *Mol Biol Evol* **23**, 254-267.
Husson SJ, Costa WS, Schmitt C, Gottschalk A (2013) Keeping track of worm
trackers. *WormBook* (ed. The *C. elegans* Research Community).
https://wormbook.org/chapters/www_tracking/tracking.html
Kamath RS, Fraser AG, Dong Y*, et al.* (2003) Systematic functional analysis of the
Caenorhabditis elegans genome using RNAi. *Nature* **421**, 231-237.
Kanzaki N, Tsai IJ, Tanaka R*, et al.* (2018) Biology and genome of a newly
discovered sibling species of *Caenorhabditis elegans*. *Nat Commun* **9**, 3216.
Keightley PD (1994) The distribution of mutation effects on viability in *Drosophila*
melanogaster. *Genetics* **138**, 1315-1322.
Keightley PD, Caballero A (1997) Genomic mutation rates for lifetime reproductive
output and lifespan in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* **94**,
3823-3827.
Kelly WG, Schaner CE, Dernburg AF*, et al.* (2002) X-chromosome silencing in the
germline of *C. elegans*. *Development* **129**, 479-492.
Kiontke K, Gavin NP, Raynes Y*, et al.* (2004) *Caenorhabditis* phylogeny predicts
convergence of hermaphroditism and extensive intron loss. *Proc. Natl. Acad.*
Sci. USA **101**, 9003-9008.
Konrad A, Brady MJ, Bergthorsson U, Katju V (2019) Mutational landscape of
spontaneous base substitutions and small indels in experimental
Caenorhabditis elegans populations of differing size. *Genetics* **212**, 837-854.
Kwon N, Hwang AB, You YJ, SJ VL, Je JH (2015) Dissection of *C. elegans*
behavioral genetics in 3-D environments. *Sci Rep* **5**, 9564.
Lee D, Zdraljevic S, Stevens L*, et al.* (2021) Balancing selection maintains hyper-
divergent haplotypes in *Caenorhabditis elegans*. *Nat Ecol Evol* **5**, 794-807.
Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760.
Lynch M (1985) Spontaneous mutations for life-history characters in an obligate
parthenogen. *Evolution* **39**, 804-818.
McKenna A, Hanna M, Banks E*, et al.* (2010) The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Res **20**, 1297-1303.
Multigner LF, Bras A, DiLeo MF, Saastamoinen M (2025) Relative effects of habitat
amount and fragmentation per se on genetic diversity of the Glanville fritillary
butterfly. *Mol Ecol* **34**, e70037.
Noble LM, Yuen J, Stevens L*, et al.* (2021) Selfing is the safest sex for
Caenorhabditis tropicalis. *Elife* **10**, e62587.
Petersen C, Dirksen P, Prahl S, Strathmann EA, Schulenburg H (2014) The
prevalence of *Caenorhabditis elegans* across 1.5 years in selected North
German locations: the importance of substrate type, abiotic parameters, and
Caenorhabditis competitors. *BMC Ecol* **14**, 4.
Petersen C, Hermann RJ, Barg MC*, et al.* (2015) Travelling at a slug's pace: possible
invertebrate vectors of *Caenorhabditis* nematodes. *BMC Ecol* **15**, 19.

Picao-Osorio J, Bouleau C, Gonzalez de la Rosa PM*, et al.* (2025) Evolution of developmental bias explains divergent patterns of phenotypic evolution in two nematode clades. *Proc Natl Acad Sci U S A* **122**, e2507529122.

Pierce-Shimomura JT, Morse TM, Lockery SR (1999) The fundamental role of pirouettes in *Caenorhabditis elegans* chemotaxis. *J Neurosci* **19**, 9557-9569.

Rajaei M, Saxena AS, Johnson LM*, et al.* (2021) Mutability of mononucleotide repeats, not oxidative stress, explains the discrepancy between laboratory-accumulated mutations and the natural allele-frequency spectrum in *C. elegans*. *Genome Res* **31**, 1602-1613.

Ramot D, Johnson BE, Berry TL, Jr., Carnell L, Goodman MB (2008) The Parallel Worm Tracker: a platform for measuring average speed and drug-induced paralysis in nematodes. *PLoS One* **3**, e2208.

R Core Team (2023) R: A language and environment for statistical computing. http://www.R-project.org/

Richaud A, Zhang G, Lee D, Lee J, Félix M-A (2018) The local co-existence pattern of selfing genotypes in *Caenorhabditis elegans* natural metapopulations. *Genetics* **208**, 807-821.

Roberts WM, Augustine SB, Lawton KJ*, et al.* (2016) A stochastic neuronal model predicts random search behaviors at multiple spatial scales in *C. elegans*. *Elife* **5**, e12572.

Rockman MV, Kruglyak L (2009) Recombinational landscape and population genomics of *Caenorhabditis elegans*. *PLoS Genet.* **5**, e1000419.

Roemer C, Sheward DJ, Hisner R*, et al.* (2023) SARS-CoV-2 evolution in the Omicron era. *Nat Microbiol* **8**, 1952-1959.

Samuel BS, Rowedder H, Braendle C, Félix M-A, Ruvkun G (2016) *Caenorhabditis elegans* responses to bacteria from its natural habitats. *Proc Natl Acad Sci U S A* **113**, E3941-E3949.

Saxena AS, Salomon MP, Matsuba C, Yeh SD, Baer CF (2019) Evolution of the mutational process under relaxed selection in *Caenorhabditis elegans*. *Mol Biol Evol* **36**, 239-251.

Schulenburg H, Félix M-A (2017) The natural biotic environment of *Caenorhabditis elegans*. *Genetics* **206**, 55-86.

Seidel HS, Rockman MV, Kruglyak L (2008) Widespread genetic incompatibility in *C. elegans* maintained by balancing selection. *Science* **319**, 589-594.

Shao W, Li X, Goraya MU, Wang S, Chen JL (2017) Evolution of Influenza A virus by mutation and re-assortment. *Int J Mol Sci* **18**, 1650.

Sivasundar A, Hey J (2005) Sampling from natural populations with RNAi reveals high outcrossing and population structure in *Caenorhabditis elegans*. *Curr Biol* **15**, 1598-1602.

Sternberg PW, Van Auken K, Wang Q*, et al.* (2024) WormBase 2024: status and transitioning to Alliance infrastructure. *Genetics* **227**, iyae050.

Stevens L, Félix M-A, Beltran T*, et al.* (2019) Comparative genomics of ten new *Caenorhabditis* species. *Evolution Letters* **3**, 217-236.

904   Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P (2015) Sambamba: fast
905   processing of NGS alignment formats. *Bioinformatics* **31**, 2032-2034.
906   Vassilieva LL, Lynch M (1999) The rate of spontaneous mutation for life-history
907   traits in *Caenorhabditis elegans*. *Genetics* **151**, 119-129.
908

909 **Data Accessibility statement**
910 Raw sequence reads are available at CaeNDR.org or deposited at NCBI under project
911 number PRJNA1280162.
912 Individual genotype data are available in Tables S3 and S4.
913

914 **Benefit-Sharing statement**
915 Benefits Generated: The samples were collected in mainland France where the laboratory
916 of the last author is based. Benefits from this research accrue from the sharing of our data
917 and results on public databases as described above.
918

Author contributions
MAF conceived the project. MAF performed field work and nematode isolation. RT and
ECA (CaeNDR) provided genome sequences. XW prepared genomic DNA for
sequencing, with assistance from AR. XW and MAF performed the genomic analyses.
MAF supervised the project.

Main Tables and Figures

Table 1 Substitution rates (μ_{bs}; $\times 10^{-8}$ per site per year) in exon, intron, non-coding genes, and

intergenic regions, and number of effective generations per year (N_{EG}).

	HS2		HS3	
	N_{SNP}	μ_{bs}	N_{SNP}	μ_{bs}
Exon	159	3.44 (2.34 - 4.54)	378	3.64 (3.04 - 4.26)
Intron	152	4.30 (2.98 - 5.75)	348	4.43 (3.67 - 5.20)
Non-coding genes	21	4.13 (1.74 - 6.85)	46	4.80 (3.20 - 6.40)
Intergenic†	164	4.13 (2.84 - 5.53)	375	4.92 (4.14 - 5.69)
N_{EG}	22.69 (15.60 - 30.39)		27.03 (22.58 - 31.26)	

928 N_{SNP} is the number of SNPs in each category. † The μ_{bs} of intergenic regions was employed to

929 estimate N_{EG} referring to μ_{bs} of intergenic regions of mutation accumulation lines (i.e., per site

930 per generation) from (Konrad *et al.*, 2019). Intervals in brackets represent 95% highest posterior

931 density.

Table 2 Ts/Tv ratios of different genic regions in wild isotypes sampled in the field (i.e., HS2 and HS3), and Ts/Tv ratios of the whole genome from mutation accumulation (MA) lines cultured in laboratory from [1](Denver *et al.*, 2012), [2](Denver *et al.*, 2009), [3](Konrad *et al.*, 2019), and [4](Saxena *et al.*, 2019). Cel - *Caenorhabditis elegans*; Cbr - *C. briggsae*. N_{SNP} is the number of SNPs in each category.

Isotype/Strain	region	Ts/Tv ratio	N_{SNP}
Wild HS2 (Cel)	Exon	1.69	159
	Intron	0.85	152
	Non-coding genes	1.33	21
	Intergenic	1.19	164
	Whole genome	**1.20**	**496**
Wild HS3 (Cel)	Exon	1.14	378
	Intron	1.06	348
	Non-coding genes	1.71	46
	Intergenic	1.19	375
	Whole genome	**1.15**	**1147**
MA PB306 (Cel) [1]	Whole genome	1.06	99
MA HK104 (Cbr) [1]	Whole genome	0.72	91
MA PB800 (Cbr) [1]	Whole genome	1.14	150
MA N2 (Cel) [2]	Whole genome	0.45	391
MA N2 (Cel) [1]	Whole genome	0.64	108
MA N2 (Cel) [3]	Whole genome	0.64	1112
MA N2 (Cel) [4]	Whole genome	0.74	347

Figures

Figure 1 The location of Santeuil in mainland France (a), sampling transect with surrounding landscape structure (b), substrates (e.g., rotting stems and invertebrate animals, here slug and isopod) sampled in the field (c), the number of SNPs between isotypes and within isotypes (d), and population composition of *Caenorhabditis elegans* in the sampling line across years (e). In (b), the land cover referred to ESRI (2021) and OpenStreetMap (2021). In (e), the sample sizes (N) refer to the 2b-RAD dataset in (Richaud *et al.*, 2018) (samples collected from 2009 to 2014) and whole-genome sequencing from CaeNDR (Crombie *et al.*, 2023) (2017-2020) and the laboratory (2022).

Figure 2 Haplotype networks of HS2 (a) and HS3 (b) strains. This figure includes strains from outside the Santeuil wood (i.e., external strains, in grey). In (a), to focus on mutation accumulation over time, the HS2 network is shown here without the three recombinants shown in Figure S3 of the Supplementary Materials. Subclades used for the spatial analysis are indicated by dashed-line polygons.

Figure 3 The SNP distribution along chromosomes (a) in HS2 (upper) and HS3 (lower), with SNP density expressed in $\times 10^{-5}$ per site, displayed by chromosome (b), genic region (c), and recombination domain (d). The figure uses 496 and 1147 local SNPs for HS2 strains and HS3 strains, respectively. External strains and recombinants were not included. Blank regions are those removed in the trimmed genomes. In (b)-(d), the number of SNPs in each category is annotated on the corresponding bars. In (b), the distribution is not even among chromosomes (Chi-Square Goodness of Fit Test for even rates among chromosomes, $p = 0.02$ for HS3, $p > 0.05$ for HS2), with the X chromosome displaying the highest mutation rate. In (c), exons carry significantly fewer mutations per site than introns or intergenic regions (Chi-Square Goodness of Fit Test for genic regions, $p = 0.02$ for HS3, $p > 0.05$ for HS2). In (d), the boundaries of tips, arms, and cores refer to (Rockman & Kruglyak, 2009). Arms carry significantly more mutations per site in HS3 (Chi-Square Goodness of Fit Test for tips, arms and cores, $p = 0.002$ for HS3, $p > 0.05$ for HS2). Detailed counts are in Table S5.

Figure 4 The mutational spectra in HS2 (a) and HS3 (b). The total number of mutations in each genic region category is indicated in Table 2 for each isotype. Detailed counts are in Table S6.

Figure 5 Maximum Clade Credibility trees of Bayesian Skygrid demographic model with tip-calibration for HS2 (a) and HS3 (b). Branch color represents substitution rates in intergenic regions ($\times 10^{-8}$ per site per year). Note the different scales in the two panels.

Figure 6 The overall spatial distribution of subclades within HS2 (a) and HS3 (b) along the sampling line, or along a simplified straight line (c), and spatiotemporal distribution of each subclade (e-l). In (a)-(c), the colors indicate the subclades. Detailed information about strain clustering in subclades is indicated in Figure 2. In (d)-(l), the colors indicate sampling year of each strain. The four strains in the basal position of clade HS3.4 are shown in plots of both subclades HS3.4.1 (j) and HS3.4.2 (k). The gray line represents a railway and the blue lines represent water bodies, based on data from OpenStreetMap (2021).