CDK7, CDK9, or CDK11 Inhibition Reduces Neutrophil-Driven Inflammation and Tissue Damage in Experimental Autoimmune Models

Mareile Schlotfeldt[1*], Leonie Voss[2], Adrian P. Mansini[3], Daniel Mehlberg[1], Sophia Johannisson[1], Michael Radziewitz[1], Ann-Kathrin Schneider[2], Seyed Mohammad Vahabi[3], Yulu Wang[3], Haley Gainer[3], Jing Li[3], Colin Osterloh[1], Nancy Ernst[1], Remco Visser[4], Gestur Vidarsson[4,5], Frank Petersen[6], Xinhua Yu[6], Ralf J. Ludwig[1,7], Artem Vorobyev[1,7], Kyle T. Amber[3], Anja Lux[2,8], Katja Bieber[1]

[1]Lübeck Institute of Experimental Dermatology (LIED), University of Lübeck, Lübeck, Germany

[2]Department of Biology, Institute of Genetics, Friedrich-Alexander University (FAU) Erlangen-Nürnberg, Erlangen, Germany

[3]Department of Dermatology, Rush University Medical Center, Chicago, IL, United States

[4]Immunoglobulin Research Laboratory, Sanquin Research, Amsterdam, The Netherlands

[5]Department of Biomolecular Mass Spectrometry and Proteomics, Utrecht Institute for Pharmaceutical Sciences and Bijvoet Center for Biomolecular Research, Utrecht University, Utrecht, The Netherlands

[6]Priority Area Asthma and Allergy, Research Center Borstel, Airway Research Center North, German Center for Lung Research, Borstel, Germany

[7]Department of Dermatology, University of Lübeck, Lübeck, Germany

[8]Profile Centre Immunomedicine, Friedrich-Alexander University (FAU), Erlangen, Germany

*Correspondance: Mareile Schlotfeldt, mareile.schlotfeldt@uksh.de

Abstract

Background: Cyclin-dependent kinases (CDKs) are involved in basic cellular processes like

regulation of cell-cycle progression and transcription. However, recent data also indicate a specific

role in terminally differentiated neutrophils by promoting reactive oxygen species (ROS) release,

degranulation, neutrophil extracellular trap formation, or apoptosis. Since these mechanisms are

implicated in multiple autoimmune diseases, we aimed to delineate the role of CDKs in IC-mediated

autoimmune diseases both *in vitro* and *in vivo*.

Methods: We analyzed CDK gene expression in unstimulated and immune complex (IC)-stimulated

neutrophils. Subsequently, we investigated the effect of pharmacological CDK inhibition on IC-

activated neutrophil functions. To analyze the inhibitors in a more translational approach, we

proceeded with the systemic and topical application of the effective inhibitors in a murine antibody

transfer-induced local epidermolysis bullosa acquisita (EBA) model. The most efficient inhibitor,

MC180295, was validated in two other IC-mediated models of autoimmune disease: Serum-transfer

arthritis (STA), which also, but not exclusively, depends on neutrophils and immune thrombocytopenia

(ITP), which is considered less neutrophil-dependent.

Results: We found 14 CDKs expressed in unstimulated cells, while the IC-stimulation showed an

upregulation of CDK2 and CDK4 expression. Inhibitors selectively targeting CDK1, CDK2, CDK4/6,

CDK7, CDK9, CDK11, and CDK12 showed effects on different neutrophil functions (surface

activation marker expression, ROS release, adhesion, apoptosis) *in vitro*. In the predominantly

neutrophil-driven EBA model, we observed a reduction of disease severity upon treatment with

CDK7, CDK9, or CDK11 inhibitors. Inhibiting these CDKs with topical THZ2, MC180295, or

OTS964, respectively, also improved the clinical phenotype. In line with our hypothesis, MC180295

impaired the development of STA, but not ITP.

Conclusions: CDK7, CDK11, and especially CDK9 inhibition show therapeutic potential in IC-driven

neutrophil-mediated diseases such as rheumatoid arthritis and EBA.

Keywords

cyclin-dependent kinases, neutrophils, pemphigoid disease, rheumatoid arthritis, immune thrombocytopenia, immune complexes, small-molecule inhibitors, autoimmunity, animal models, signaling

Background

Cyclin-dependent kinases (CDKs) are a group of serine-threonine kinases that were discovered for their involvement in the regulation of the eukaryotic cell cycle [1–3]. CDKs are crucially contributing to the correct replication and subsequent separation of the chromosomal DNA during mitosis [4] and are highly conserved across eukaryotes, from yeast to mammals [5]. They were termed for their critical interaction with specific regulatory protein binding partners, the cyclins [6]. The latter are expressed in a cyclic fashion during the cell cycle, while the CDKs themselves are stably expressed throughout the cell cycle [3]. There are 20 CDKs and 29 cyclins described in mammals [7], as well as other regulating proteins such as intrinsic CDK inhibitors (CKI) [8]. Mechanistically, activation of CDKs not only requires the interaction with a cyclin, also phosphorylation at a specific tyrosine residue in the activation loop is necessary [9, 10]. This phosphorylation of at least CDK1/2/4/6/9/12/13 is mediated by the CDK-activating kinase (CAK) complex [11], consisting of CDK7, cyclin H and MAT1 [12–14]. Importantly, CDK inactivation is also regulated by phosphorylation [15] or binding of CKIs [16, 17].

Beyond the well-described role of CDKs in direct cell-cycle regulation, there is rising evidence that CDKs are also involved in regulating the RNA polymerase II transcription cycle [11, 18]. It was thus proposed that there are two groups of CDKs: the cell-cycle CDKs (subfamilies CDK1, 2, 3, 4, and 6) and the transcriptional CDKs (subfamilies CDK7, 8, 9, 10, 11, 12, and 13) [7]. The remainder is referred to as atypical CDKs, although these kinases are insufficiently characterized [19]. Though this may be a helpful concept, this dichotomy is probably oversimplified as more and more studies

demonstrate roles for traditional cell-cycle CDKs in transcriptional regulation and vice versa [18, 20–

22].

Considering that CDKs are not only expressed in proliferating cells, but also in terminally

differentiated cells like neutrophils [23], it seems a reasonable assumption that their role extends

beyond mere cell cycle regulation. Neutrophils are known as the first line of defense against pathogens

[24]. Once recruited to the tissue of interest, they are activated by the binding of immune complexes

(IC) to their fragment crystallizable gamma receptors (FcγR). Downstream kinase signaling cascades

are activated, triggering effector functions such as reactive oxygen species (ROS), protease and

cytokine release to kill the invading pathogens in a controlled manner. Subsequently, the neutrophils

are cleared mostly by macrophages under inflammatory conditions [25, 26]. Notably, a role for several

CDKs in regulating neutrophil functions has been described: CDK2 is involved in migration [27],

CDK4/6 in formation of neutrophil extracellular traps (NETs) [28], CDK5 in degranulation [29], and

CDK7/9 in apoptosis [22, 30]. Still, little is known about the distinct functions of different CDKs in

direct comparison, and most studies rely on broad-range small-molecule CDK inhibitors (CDKi).

Moreover, the potential involvement of CDKs in FcγR-mediated signaling remains largely unexplored,

representing a significant gap in our understanding of CDK biology.

An increasing number of studies points towards the relevance of CDKs in various human disorders,

underlining the need for a better comprehension of these kinases. Deregulated CDK activity has been

shown to be associated not only with e.g. neurodegenerative diseases [31] and cancer [19], but also

with autoimmune diseases like rheumatoid arthritis (RA) [32] or systemic lupus erythematosus [33]. In

contrast to infectious diseases, however, there is no pathogen in autoimmunity, but autoantibodies

develop against self-antigens due to a break of tolerance [34]. To unravel the function of CDKs in the

context of neutrophil-mediated FcγR-dependent signaling, we investigated the kinase signaling

pathways upon IC stimulation in human neutrophils *in vitro*. Furthermore, we chose three well-

established murine *in vivo* models of antibody-mediated autoimmune diseases: Epidermolysis bullosa

acquisita (EBA), serum-transfer arthritis (STA) as a model for RA, and immune thrombocytopenia

(ITP). EBA belongs to the group of pemphigoid diseases (PD), which is a group of rare autoimmune

bullous skin diseases characterized by autoantibodies targeting structural proteins of the dermal-

epidermal junction. More specifically, the autoantigen in EBA is collagen VII (COL7) [35].

Neutrophils are the central effector cells in PD [36–39], causing the uncontrolled release of

inflammatory mediators. Thereby, tissue damage occurs, causing the degradation of the dermal-

epidermal adhesion complex and subepidermal split formation [38, 40]. Clinically, patients present

with pruritus, erosions, and tense, painful blisters on different sites of the body, e.g. trauma-prone

areas or the extensor skin surface [41]. Currently, the patients are most commonly treated with

systemic corticosteroids, often in combination with additional immunosuppressants, bearing a high

risk for adverse events due to general immunosuppression [42, 43].

Another more prevalent neutrophil-dependent autoimmune disease with the need for novel therapeutic

options is RA, modelled by STA [44]. In contrast to EBA, neutrophils do play a critical role in

pathogenesis [45, 46], but other cell types like macrophages [47] or mast cells [48, 49] are crucially

involved in the effector phase. Joint inflammation, bone and cartilage destruction are major hallmarks

of RA [50].

Conversely, ITP is also IC-driven, but depends less on neutrophils. In ITP, autoantibodies target

platelets leading to their phagocytosis mainly by macrophages and thus to platelet depletion [51–53].

Since kinases are generally well-druggable targets [54, 55] and considering that there are several

CDKi already in clinical use, e.g. for cancer treatment [56], a potential repurposing of the inhibitors

could accelerate the process of clinical approval. Thus, the characterization of CDK signaling cascades

in neutrophils can help to identify potential therapeutic targets for PD, but also potentially for other

neutrophil-driven autoimmune diseases like RA. This could provide targeted treatment options to the

patients and therefore drastically ameliorate the safety profile of the therapy.

Methods

Sampling of human biomaterials

Whole blood was obtained from healthy volunteers after written informed consent. All experiments

using human samples were approved by the local ethics committee (AZ #20-463, University of

Lübeck, Lübeck, Germany) and were performed in accordance with the Declaration of Helsinki.

Chemicals

All standard chemicals were supplied by Carl Roth (Karlsruhe, Germany) or Sigma-Aldrich

(Taufkirchen, Germany). CDKi were supplied by Selleck Chemicals (Houston, Texas, USA) or

MedChemExpress (Monmouth Junction, New Jersey, USA). For *in vitro* assays, 10 mM CDKi was

dissolved in dimethyl sulfoxide (DMSO) and further diluted to final concentrations of 10 µM, 1 µM,

0.1 µM, and 0.01 µM. DMSO at a final concentration of 0.1 % was included in the controls. For *in*

vivo experiments, CDKi were prepared as stated in Table 1. The respective vehicle only served as the

respective control. Additional targets, references and stage of drug development are given in

Supplemental table 1.

Table 1. CDKi used in this study. The vehicle, dosage, route and frequency for *in vivo* treatment

stated, where applicable. IC_{50}, Kd, or Ki values were adapted from Selleckchem.com. N/A, not

available.

substance	main target	vehicle	dosage [mg/kg]	route	fre-quency	IC_{50}/*Kd*/<u>Ki</u> [nM]
Ro 3306	CDK1	10 % DMSO/ 30 % PEG-300/ 10 % Tween-80	4	p.o.	daily	*20*
OTS964	CDK11B	5 % DMSO, 40 % PEG-300, 5 % Tween-80	40	p.o.	daily	<u>40</u>
Palbociclib HCl	CDK4/6	sodium lactate buffer (50 mM, pH	100	p.o.	daily	9

		4.0)				
THZ1 2HCl	CDK7	10 % DMSO/ 90 % 5 % dextrose in water	10	i.p.	twice daily	3.2
THZ2	CDK7	10 % DMSO in D5W, 5 % dextrose in water	10	i.p.	daily	13.9
MC180295	CDK9	5 % DMSO/ 40 % PEG-300/ 5 % Tween-80/ 50 % water	20	i.p.	daily	5
LY3143921 hydrate	CDC7	-	-	-	-	N/A
BMS-265246	CDK1/2	-	-	-	-	6
Purvalanol B	CDK1/2/5	-	-	-	-	6
BSJ-4-116	CDK12	-	-	-	-	107
SR-4835	CDK12/13	-	-	-	-	4.9
K03861	CDK2	-	-	-	-	9.6
AS2863619	CDK8/19	-	-	-	-	0.6
BAY1251152	CDK9	-	-	-	-	3

145 **Supplemental table 1.** Additional targets of CDKi used adopted from Selleckchem.com and

146 references for *in vivo* application and current stage of drug development.

Substance	Main target	Additional targets (IC$_{50}$/*Kd*/<u>Ki</u>)	Reference	Stage of drug development
Ro 3306	CDK1	PKCδ (<u>318 nM</u>), SGK (<u>497 nM</u>), ERK (<u>1980 nM</u>)	[57]	pre-clinical studies
OTS964	CDK11B	TOPK (28 nM)	[58]	pre-clinical studies
Palbociclib HCl	CDK4/6	CDK4-cyclin D1 (11 nM), CDK6-cyclin D2 (15 nM)	[59, 60]	FDA- and EMA- approved
THZ1 2HCl	CDK7	-	[61, 62]	pre-clinical studies
THZ2	CDK7	-	[63]	pre-clinical studies
MC180295	CDK9	CDK4-cyclin D (112 nM), CDK1-cyclin B (138 nM), CDK5-P35 (159 nM), CDK5-P25 (186 nM), CDK2-cyclin A (233 nM), CDK2-cyclin E (367 nM), CDK3-cyclin E (399 nM)	[64]	pre-clinical studies
LY3143921 hydrate	CDC7	-	[65]	clinical trial
BMS-265246	CDK1/2	CDK2-cyclin E (9 nM), CDK4-cyclin D (230 nM)	[66]	pre-clinical studies
Purvalanol B	CDK1/2/5	CDK2-cyclin A (6 nM), CDK5-P35 (6 nM), CDK2-cyclin E (9 nM)	[67]	pre-clinical studies
BSJ-4-116	CDK12	(CDK12 degrader)	[68]	clinical trial

SR-4835	CDK12/13	CDK12 (99 nM, *98 nM*)	[69]	pre-clinical studies
K03861	CDK2	CDK2 (A144C, 15.4 nM), CDK2 (C118L, 18.6 nM), CDK2 (WT, 50 nM)	[70]	pre-clinical studies
AS2863619	CDK8/19	CDK19 (4.277 nM)	[71]	pre-clinical studies
BAY1251152	CDK9	-	[72]	clinical trial

Isolation of human immune cells by magnetic activated cell sorting (MACS)

For the isolation of human neutrophils by MACS, blood from healthy young (< 30 y) donors was drawn into citrate blood collection tubes. After 1:2 dilution with 1 % (w/v) polyvinyl alcohol and sedimentation for 25 min, the supernatant (up to 40 mL) was transferred on top of a layer of 8 mL of Ficoll Paque PLUS (Cytiva, Marlborough, Massachusetts, USA). Density gradient centrifugation [73] was performed for 24 min at 850 g and room temperature (RT) without brake. The lymphocyte layer was processed for B and CD4[+] T cell isolation, whereas the granulocyte pellet was used for neutrophil purification. The lymphocyte layer was resuspended in phosphate-buffered saline (PBS) and centrifuged at 300 g for 10 min at 4 °C for platelet removal. Another washing step with PBS was performed at 400 g for 10 min at 4 °C. Next, the purified lymphocytes were subjected to MACS according to the manufacturer's instructions using the CD4[+] T cell isolation kit (130-096-533, Miltenyi Biotec, Bergisch Gladbach, Germany) and the B cell isolation kit II (130-091-151, Miltenyi Biotec). B and CD4[+] T cell purities were assessed by flow cytometry and were > 80 %.

For erythrocyte lysis, the granulocyte pellet was resuspended in distilled water, mixed for 45 s, and then neutralized with 2x PBS. Two washing steps with PBS were performed for 10 min at 300 g and RT. Subsequently, the granulocytes were subjected to eosinophil removal by MACS according to the manufacturer's instructions (130-092-010, Miltenyi Biotec). Neutrophil purity was assessed by flow cytometry and was > 98 %.

Bulk RNA sequencing

For bulk RNA sequencing (RNAseq), 100,000 MACS-purified neutrophils from healthy volunteers were sampled and frozen at -80 °C in 1 mL Trizol (Invitrogen, Waltham, Massachusetts, USA). An equal number of cells from each donor was stimulated for 6 h at 37 °C and 5 % CO_2 with immobilized IC consisting of human COL7E-F and anti-human COL7 IgG1 as previously described [74]. All samples were sent to Single Cell Discoveries (Utrecht, The Netherlands) for RNA extraction and sequencing. Data analysis was performed in RStudio (Posit Software, Boston, Massachusetts, USA; version 2024.12.1) as previously described [75]. Differentially expressed genes were identified using DESeq2 [76] on transcripts where the sum of the counts over all samples was > 10, and $p_{adj} < 0.1$ was considered significant. Gene set enrichment analysis was conducted using ClusterProfiler referring to the Gene Ontology (GO) database [77, 78]. The data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB103987 (https://www.ebi.ac.uk/ena/browser/view/PRJEB103987).

Human polymorphonuclear granulocyte (PMN) purification for real-time quantitative reverse transcription polymerase chain reaction (qRT-PCR) and functional assays

For ROS release assay, flow cytometry, adhesion assay, and qRT-PCR, PMNs were isolated from EDTA-treated human whole blood performing a PolymorphPrep™ (Serumwerk Bernburg AG, Bernburg, Germany) density gradient centrifugation as previously described [74, 75, 79]. The cells were diluted to 4×10^5 cells/mL in 1 % (w/v) bovine serum albumin (BSA) in PBS. Neutrophil purity was assessed by flow cytometry for all experiments and was higher than 90 % for all samples.

qRT-PCR

For gene expression analysis by qRT-PCR, human PMNs were stimulated with human COL7E-F anti-human COL7 IgG1 IC for 6 h at 37 °C and 5 % CO_2 as stated above. Alongside unstimulated PMNs, B cells and $CD4^+$ T cells isolated by MACS as described above, the cells were collected with a cell scraper in PBS and centrifuged for 7 min at 700 g and RT. The lysates were resuspended in 700 µL

RLT buffer (Qiagen, Hilden, Germany) and frozen at -80 °C until RNA extraction. RNA extraction

was performed using the innuPREP RNA Mini Kit 2.0 (Analytik Jena, Jena, Germany) according to

the manufacturer's instructions. Any remaining RNA was digested with DNase I (Sigma-Aldrich).

cDNA was synthesized using the RevertAid Reverse Transcriptase (Thermo Fisher Scientific,

Waltham, Massachusetts, USA) according to the manufacturer's instructions. TaqMan™ gene

expression assays (see Table 2) were purchased from Thermo Fisher Scientific and used according to

the manufacturer's instructions. qRT-PCR was performed on an Applied Biosystems 7900HT Fast

Real-Time PCR System (Thermo Fisher Scientific). The results were analyzed using the $2^{-\Delta\Delta Ct}$ method

[80] and normalized to the mean values of the housekeeping genes [81].

Table 2. TaqMan™ gene expression assays used for qRT-PCR.

	Catalogue no.	Assay ID	Gene symbol
CDKs	4453320	Hs00938777_m1	*CDK1*
	4453320	Hs01548894_m1	*CDK2*
	4448892	Hs01077186_g1	*CDK3, TEN1-CDK3*
	4453320	Hs00364847_m1	*CDK4*
	4453320	Hs00358991_g1	*CDK5*
	4453320	Hs01026371_m1	*CDK6*
	4448892	Hs00361486_m1	*CDK7*
	4448892	Hs00992501_g1	*CDK8*
	4448892	Hs00977896_g1	*CDK9*
	4448892	Hs02341397_m1	*CDK11A*
	4448892	Hs07291776_gH	*CDK11B*
	4448892	Hs00901732_m1	*CDK12*
	4448892	Hs00953418_m1	*CDK14*
	4448892	Hs01039930_g1	*CDK19*

	4453320	Hs00177487_m1	CDC7
housekeeping genes	4331182	Hs01060665_g1	ACTB
	4331182	Hs02786624_g1	GAPDH
	4331182	Hs02800695_m1	HPRT
	4331182	Hs00272002_m1	GNB2L1

ROS release assay

Plates were coated with IC consisting of human COL7 E-F at 2.5 µg/mL and anti-human COL7 IgG1

at 1.8 µg/mL as previously described [74, 82]. PMNs were added at $2x10^5$ cells per well in the

presence or absence of CDKi. Only antibody (with cells), only antigen (with cells), or only cells

served as negative controls, while IC (with cells, but without CDKi) served as positive control. DMSO

was adjusted to a final concentration of 0.01 %. All samples were measured in duplicates. Luminol

was automatically injected into each well to reach a final concentration of 90 µg/mL, and

chemiluminescence was measured for 2 h every 2 min at 37 °C in a GloMax® Discover microplate

reader (Promega, Walldorf, Germany). The area under the curve (AUC) was calculated, and the results

were normalized to IC-stimulated cells without CDKi.

Adhesion assay

PMN adhesion was investigated using impedance-based measurements as described before [83, 84]. A

96-well E-plate (Agilent, Santa Clara, California, USA) was coated with IC as described for the ROS

release assay. Inhibitor solutions were prepared and added to the plate before $2x10^5$ cells were

transferred to each well. Only antibody (with cells), only antigen (with cells), or only cells served as

negative controls, while IC (with cells, but without CDKi) served as positive control. DMSO was

adjusted to a final concentration of 0.01 %. All samples were measured in duplicates. Impedance was

measured in an xCELLigence real-time cell analysis device (Agilent) every 2 min for 120 min

expressed in arbitrary units as cell index. Data analysis was performed as for the ROS release assay.

Flow cytometry

Human PMNs were isolated as previously described and stimulated with IC, prepared in the same

manner as for the ROS release assay, for 2 h at 37 °C and 5 % CO_2. Subsequently, they were stained

for the flow cytometric analysis of the following surface markers using standard flow cytometry

procedures: CD14 (clone HCD14), CD16 (clone 3G8), CD18 (clone 1B4/CD18), CD45 (clone HI30),

CD62L (clone DREG-56), and CD193 (clone 5E8). Viability was assessed by Annexin V (B266195)

and Zombie NIR staining (B308933) [75]. Only antibody (with cells), only antigen (with cells), or

only cells served as negative controls, while IC (with cells, but without CDKi) served as positive

control. DMSO was adjusted to a final concentration of 0.01 %. All samples were measured in

duplicates. Analyses for activation markers were performed on FSC-SSC gated $CD45^+$ singlets.

Annexin V^- Zombie NIR^- cells were considered viable, Annexin V^+ Zombie NIR^- cells were considered

apoptotic. The positive control corresponds to values of up to 3 %. All dyes were purchased from

BioLegend (San Diego, California, USA). Measurements were performed on a MACSQuant®

Analyzer 10 (Miltenyi Biotec) and analysis was conducted using MACS Quantify software (version

2.13.3, Miltenyi Biotec).

Study approval

Animal experiments were approved by local authorities of the Animal Care and Use Committee

(government of Schleswig-Holstein and government of Lower Franconia) and the Institutional Animal

Care and Use Committee at Rush University. All experiments were performed by certified personnel

(AZ 108/08-15, AZ 97-11/20 and 55.2.2-2532-2-2085) following the ARRIVE guidelines.

Animal experimentation

C57BL/6J mice (Charles River, Wilmington, Massachusetts, USA) were bred in a specific pathogen–

free environment and provided standard mouse chow and acidified drinking water *ad libitum*. Mice of

both sexes, aged 8 to 14 weeks, were used for experimental EBA models with clinical examinations

performed under anaesthesia, using intraperitoneal (i.p.) administration of a ketamine (75 mg/kg,

Sigma-Aldrich) and medetomidine (1 mg/kg, Vetoquinol, Ismaning, Germany) mixture, antagonized

after ca. 30 min by atipamezole (5 mg/kg, Vetoquinol). On the final day, the mice were anaesthetized

using 15 mg/kg xylazine (Sigma-Aldrich) and 100 mg/kg ketamine. STA and ITP models were

performed with female C57BL/6J aged 12 weeks. For anaesthesia, isoflurane was used and mice were

killed by CO_2.

Local antibody-transfer induced experimental EBA

Specific anti–mouse COL7[C] IgG from rabbit serum was isolated as previously described [40, 85–87].

For systemic treatment, CDKi were administered as indicated in Table 1 at 100 µL/20 g mouse. For

topical treatment in independent experiments, THZ2, MC180295, or OTS964 were applied at

65 µg/100 µL in acetone/DMSO (20:1) per ear, beginning one day before the anti–mouse COL7[C] IgG

injection and treatment was performed daily for four days. Vehicle-treated mice served as controls in

all cases. Experimental EBA was induced by ear-base injection with 100 μg rabbit anti–mouse COL7[C]

IgG. The mice were clinically evaluated every day until day 3, and the percentage of ear thickness was

assessed using a Mitutoyo 7301 dial thickness gauge (Neuss, Germany) by a blinded person, as well as

the affected ear surface area (AESA). Ears were collected on the final day of the experiment, and half

of each ear was embedded in 4 % PBS-buffered paraffin or frozen in liquid nitrogen and stored at

-80 °C. Histological analysis of hematoxylin and eosin (H&E) staining and direct immunofluorescence

staining for murine IgG and C3 were performed as described elsewhere [87]. Scoring of H&E-stained

sections was performed to determine (i) the magnitude of epidermal thickness with values from 0 (10-

20 µm), 1 (20-40 µm), 2 (40-100 µm) to 3 (> 100 µm), (ii) the percentage of dermal-epidermal split

formation (length of split formation/length of dermal-epidermal junction), with values from 0 (no

split), 1 (< 20 %), 2 (20-50 %) to 3 (> 50 %) and (iii) the dermal infiltration with values from 0 to 3

corresponding to no, mild, moderate or severe degree, respectively. The H&E score was calculated as

the mean of these three parameters.

STA

STA was induced by injection of arthritogenic KBxN serum in female C57BL/6J mice at 10 µL/g body

weight as described before [88]. Before and after arthritis induction, mice were treated daily with 20

mg/kg body weight MC180295 or vehicle control i.p. The severity of joint swelling was monitored to

evaluate clinical disease over the course of seven days. Peripheral blood was collected retro-orbitally

on day 0 and day 7, and inflammatory cytokines were quantified in the serum by LegendPlex

multiplex assay (Mouse Inflammation 13-plex, BioLegend) as a parameter of systemic inflammation.

On day 7, mice were sacrificed, and forelegs were prepared for histological analysis. H&E staining

was performed to evaluate inflammatory infiltrates and Safranin-O staining to assess bone and

cartilage damage, as previously described [88, 89]. Stained sections were evaluated by light

microscopy using a Zeiss Axiovert 200 microscope and the AxioVision LE4.6 software (Olympus Life

Science, Waltham, Massachusetts, USA).

Immune thrombocytopenia (ITP)

Platelet depletion in the presence of kinase inhibition was assessed in female C57BL/6J mice. Platelet

depletion was induced by injection of 0.35 µg/g anti-platelet IgG2c (clone 6A6) as described [53].

Treatment with vehicle control or MC180295 (20 mg/kg body weight i.p.) was performed 24 h and 4 h

before the induction of platelet depletion. Platelet counts in peripheral blood were assessed

immediately before 6A6-IgG2c injection, and at 4 h and 24 h post-injection in peripheral blood

collected retro-orbitally. Platelet counts were determined using an ADVIA® 2120i hematocytometer

(Siemens Healthineers, Erlangen, Germany).

Statistical analysis

Data were analyzed using Prism 10.4.1 (GraphPad Software, San Diego, USA). For all *in vitro* assays,

Kruskal-Wallis test with Dunn's multiple comparisons test was performed if not indicated differently.

Dose response curves and IC_{50} values were calculated using the non-linear fit inhibitor vs. response

(three parameters) function or for EC_{50} values using the agonist vs. response (three parameters)

function where an activating response was observed. For all *in vivo* experiments, two-way ANOVA or

mixed-effects analysis (in the case of missing values) with Šidák's multiple comparisons test were

performed unless stated differently. A value of $p \leq 0.05$ was considered significant. The data are

displayed as Tukey box-and-whisker plots or else as mean ± standard deviation unless described

differently.

Results

Fourteen CDKs are expressed at mRNA level in human neutrophils and expression of CDK2 and CDK4 changes upon IC stimulation

CDKs are essential to drive cell cycle progression in proliferating cells [19]. However, at least certain CDKs seem to be functionally relevant also in terminally differentiated cells like neutrophils [22, 27–29]. Since there are conflicting data published on this topic [22, 28], we first set out to clarify which CDKs and regulating proteins are expressed at mRNA level in human neutrophils. Therefore, an RNAseq experiment on neutrophils from healthy volunteers was performed. In addition, neutrophils were stimulated with immobilized IC for 6 h and compared to unstimulated neutrophils to reveal any transcriptional changes. Fig. 1A shows the best-described CDK-cyclin complexes involved in the cell cycle and the transcriptional cycle irrespective of whether their expression changes significantly upon IC stimulation or not, whereas Fig. 1B only shows significantly altered CDKs, cyclins, and selected regulators after IC stimulation. From the 20 CDKs that are overall expressed in humans, we found 14 CDK transcripts (considering both paralogues of *CDK11*) expressed in unstimulated neutrophils and six significantly changed by IC stimulation. While *CDK13* and *CDK19* expression was downregulated after 6 h, *CDK2*, *CDK4*, *CDK7*, and *CDK12* expression was upregulated. Regarding cyclins, 16 out of 29 genes were expressed. On the one hand, the CDKs with best established roles in direct regulation of the cell cycle, namely *CDK1*, *CDK2*, *CDK4*, and *CDK6*, were all detected by RNAseq except *CDK1*. The gene expression of *CDK2* and *CDK4* was induced upon IC stimulation. In addition, their partner cyclins were also detected except cyclin E (*CCNE*), which binds to CDK2 during late G1 phase to promote G1-S transition [90, 91]. Cyclin A (*CCNA*) was unchanged after IC stimulation, whereas cyclin B (*CCNB*) and D (*CCND*) showed an increased expression. On the other hand, expression of the transcriptional kinases *CDK7*, *CDK9*, *CDK12*, *CDK13* and *CDK19* was observed, but not of *CDK18*. Their cyclin binding-partners were all detected with cyclin H (*CCNH*) and T (*CCNT*) elevated upon IC treatment. In addition, the gene expression of other CDK-regulating proteins like RB1, EP300 or MED12/13 was significantly elevated after IC stimulation (Fig. 1B). Furthermore, GO pathway

analysis of all genes significantly changed by IC stimulation revealed an activation of granulocyte-

specific pathways and a suppression of metabolic pathways (Fig. 1C).

qRT-PCR confirms the expression of 14 CDKs from RNAseq in human PMNs

To validate the results from RNAseq, an independent qRT-PCR experiment was performed on

unstimulated and IC-stimulated human PMNs. Data are provided in Additional file 1. All tested *CDKs*

were detected, although *CDK1*, *CDK3*, *CDK5*, *CDK6*, *CDK8*, *CDK11A*, and *CDC7* were expressed at

a relatively low level ($2^{-\Delta\Delta Ct} < 0.1$) under both conditions. The expression of *CDK2* and *CDK4* was

increased more than threefold or sixfold, respectively, after IC stimulation, while *CDK9* expression

decreased by half and *CDK19* expression by approximately 70 %. *CDK3* and *CDK11B* showed a trend

to be reduced (both p = 0.0556), but all other *CDKs* remained unaffected by IC stimulation (Fig. 2A).

The results support the RNAseq data which also indicate an IC-mediated induction of *CDK2* and

CDK4 transcription, but a reduction of *CDK9* and *CDK19* transcription (see section above). For

comparison with proliferating cells, B and CD4$^+$ T cells were also analyzed in qRT-PCR (Fig. 2B).

Strikingly, *CDK1*, *CDK2*, *CDK4*, *CDK5*, *CDK6*, *CDK8*, and *CDC7* were expressed at higher levels in

B and/or T cells, whereas *CDK4*, *CDK9*, *CDK11B*, *CDK14*, and *CDK19* were increased in PMNs.

Pharmacological inhibition of multiple specific CDKs partially protects *in vitro* from neutrophil

effector functions in a dose-dependent manner

To functionally evaluate the role of CDKs in neutrophil kinase signaling, we selected CDKi based on

an extensive literature review and tested them in several functional neutrophil assays, covering

multiple steps from neutrophil activation to cell death. Therefore, COL7-anti-COL7-IgG1 IC were

immobilized on plates, incubated with PMNs from healthy donors and the respective experiment was

performed. The surface expression of CD62L, an L-selectin which is shed upon neutrophil activation

[92, 93], was analyzed first. Data are provided in Additional file 2. The CDK9i MC180925, CDK7i

THZ1 2HCl (Fig. 3), and CDK1i Ro 3306 (Fig. S1) significantly reduced CD62L shedding, while the

other inhibitors tested did not (Fig. 3, S1). CDK7 inhibition with THZ2 instead significantly reduced

CD18 expression (Fig. 3), which is a β2-integrin responsible for neutrophil adhesion [94]. Similar

effects were observed for Ro 3306 and THZ1 2HCl (Fig. S2). Subsequently, adhesion measurements

demonstrate a significant reduction only for THZ2 (Fig. 3, S3). Furthermore, THZ2, MC180295,

OTS964, Ro 3306, BMS-265246, K03861, palbociclib HCl, THZ1 2HCl, and BAY1251152 inhibited

ROS release (Fig. 3, S4). Regarding apoptosis, a significant increase was measured for THZ2,

OTS964, and BSJ-4-116 (Fig. 3, S5). For BMS-265246 and AS2863619, statistical significance was

reached, however, the values did not exceed those of the negative controls (Fig. S5). Although

viability was significantly lower, especially at 10 µM, more than 99 % of the cells remained viable in

all cases (Fig. 3, S6). Taken together, CDK inhibition impairs various aspects of IgG-induced

neutrophil activation, with ROS release being significantly reduced across multiple inhibitors at doses

not substantially affecting viability. Thus, the inhibitory potential of THZ2, MC180295, and OTS964,

in particular, is highlighted.

Both systemic and topical treatment with CDK7, CDK9 or CDK11 inhibitors reduces disease

severity in the local antibody-transfer induced EBA model

Following the successful *in vitro* validation of the selected CDKi, a local antibody-transfer induced

EBA model was performed to test these compounds as systemic treatment (Fig. 4A). Data are

provided in Additional file 3. CDK9 inhibition by MC180295 completely prevented disease induction,

which is exemplified by representative images showing the ameliorated clinical presentation of the

mice and the reduced inflammation in the H&E staining despite IgG and C3 deposition along the

dermal-epidermal junction (Fig. 4B). The AESA was significantly reduced starting at day 1 and

increasing further until day 3, which is also reflected by the H&E score (Fig. 4C). CDK7 inhibition by

both THZ2 (Fig. 4D) and THZ1 2HCl (Fig. 4E) reduced the AESA, but only THZ2 treatment

significantly decreased the H&E score. Systemic CDK11 inhibition by OTS964 treatment also

ameliorated the disease severity, but the H&E score did not change significantly between the treated

and the control mice (Fig. 4F). However, treatment with Ro 3306 and palbociclib HCl had no

significant effect (Fig. S7).

The most promising inhibitors from the systemic treatment in local antibody-transfer induced EBA

were selected for topical treatment in the same model. THZ2 was chosen over THZ1 HCl since both

inhibit CDK7, but THZ2 showed better results *in vitro* and almost completely protected from disease

development *in vivo*. All three inhibitors tested prevented disease induction when applied topically

(Fig. 4G).

CDK9 inhibition with MC180295 is beneficial in neutrophil-dependent STA, but not in

neutrophil-independent ITP

The CDKi that showed the most prominent effect in the local antibody-transfer induced EBA model,

MC180295, was validated in two other IC-driven models that differ regarding the role of neutrophils

in their respective pathogenesis, namely STA and ITP. Data are provided in Additional file 4. In STA

(Fig. 5A), systemic treatment with MC180295 significantly reduced the disease score of paw swelling

(Fig. 5B). This was reflected in histological analysis showing reduced proteoglycan degradation and

thus cartilage damage as compared to control mice upon staining with Safranin-O. H&E staining

revealed pronounced immune cell infiltration in the controls, but not or to a lesser extent in the treated

mice (Fig. 5C). Out of 13 cytokines tested in the serum, two were significantly changed on the final

day between the two groups. Both IL-6 and IL-10 were reduced on day 7 in the MC180295-treated

group compared to the control group. All other measured cytokine levels were not significantly altered

over the course of the observation period of 7 days (Supplemental table 2).

Supplemental table 2. Descriptive statistics for all cytokines analysed in STA at day 0 and day 7.

	day 0			day 7		
	median ± SEM		p value	median ± SEM		p value
cyto-kine	treated	control	treated vs. control	treated	control	treated vs. control

GM-CSF	0.564 ± 0.3987	0 ± 0	0.4444	1.036 ± 0.6401	1.465 ± 0.7801	0.5556
IL-10	13.99 ± 8.837	16.63 ± 16.63	0.6825	27.06 ± 9.973	60.15 ± 9.976	0.0317
IL-12p70	1.58 ± 0.8095	0.69 ± 0.3984	0.5714	1.546 ± 1.029	3.53 ± 1.436	0.3095
IL-17A	1.144 ± 0.5896	0.44 ± 0.2	0.5556	1.53 ± 0.4803	1.47 ± 0.3853	0.9048
IL-1α	190 ± 41.09	218.5 ± 33.07	0.5556	15.69 ± 4.246	14.89 ± 5.507	0.7302
IL-1β	7.224 ± 4.069	4.81 ± 1.707	> 0.9999	3.076 ± 1.342	7.043 ± 1.508	0.1032
IL-23	29.37 ± 11.81	15.08 ± 9.157	0.4365	43.63 ± 17.1	25.55 ± 6.944	0.5476
IL-27	124.5 ± 32.39	133.7 ± 85.19	0.7302	116.6 ± 41.59	111.8 ± 18.37	0.375
IL-6	4.162 ± 1.272	2.088 ± 1.033	0.1746	2.824 ± 0.8739	34.84 ± 5.281	0.0159
IFN-β	78.04 ± 20.83	37.03 ± 27.01	0.4127	10.04 ± 6.495	29.55 ± 7.633	0.0952
IFN-γ	3.578 ± 1.287	2.833 ± 0.9399	0.7937	2.968 ± 0.6265	5.545 ± 0.9739	0.127
MCP-1	12.09 ± 2.878	7.603 ± 3.671	0.2857	9.492 ± 5.536	16.65 ± 4.481	0.3175
TNF-α	13.24 ± 2.719	11.38 ± 1.989	0.4444	10.42 ± 3.304	22.59 ± 6.41	0.0952

414 In contrast to STA, ITP (Fig. 5E), where neutrophils are not central to disease pathogenesis,

415 prophylactic treatment with MC180295 did not show any effects on platelet count 24 h after platelet

416 depletion (Fig. 5G), demonstrating the variable effect of CDKi in the different disease models.

Discussion

IC-induced FcγR-mediated kinase signaling plays a major role in intracellular signal transduction by

neutrophils under pathological conditions, e.g., in autoimmune diseases such as EBA or RA. However,

the role of CDKs in neutrophils is largely unknown. A small number of studies have reported the

expression of CDKs on protein level and suggested functional relevance in these cells [22, 28, 95]. To

the best of our knowledge, we show in the present study for the first time a comprehensive, functional

analysis of IC-triggered FcγR-dependent kinase signaling using a variety of selective CDKi. Fig. 6

summarizes our main findings.

Interestingly, we identified multiple CDKs as expressed at RNA level and found that their

transcription may change in IC-stimulated neutrophils. The expression of the two traditional cell-cycle

CDKs *CDK2* and *CDK4* increased, whereas the expression of the two transcriptional CDKs *CDK9* and

CDK19 decreased. Concerning CDK9, it has been shown that inhibition leads to the reactivation of

epigenetically silenced genes, e.g. *SYNE1*, *GLDC*, *MUC20*, *ANPEP*, *SUSD4*, and *COLEC11* [64]. All

of these genes have been linked to the immune system [96], e.g. NETosis [97–99], apoptosis [100],

complement inhibition [101], or neutrophil degranulation [102]. Therefore, CDK9 inhibition might

further influence these processes.

The trends found in RNAseq are clearly supported by our qRT-PCR analysis in an independent

dataset. Furthermore, the qRT-PCR data highlight the elevated expression of *CDK9*, *11B* and *19* in

neutrophils compared to B and CD4$^+$ T cells, making them especially promising targets in neutrophil-

dependent diseases. The tendencies concerning gene expression levels in unstimulated cells found in

our RNAseq and qRT-PCR analyses are in line with publicly available RNAseq data of unstimulated

neutrophils from six healthy donors sorted by flow cytometry [88, Human Protein Atlas

proteinatlas.org]. Several CDKs like CDK2, CDK4, and CDK11 were also detected in a kinase activity

measurement of IC-stimulated PMNs [74].

Since CDKs only become fully active after binding to their cyclin binding partners, we also examined

cyclins in our RNAseq analysis. Most prominently, we found both CDK7 and its binding partner

cyclin H significantly increased upon IC activation. Strikingly, *MNAT1* (MAT1 RNA) was not

detected, suggesting the requirement of a phosphorylation-dependent activation of CDK7 [104, 105].

Furthermore, several regulators of CDKs were differentially expressed upon IC stimulation in human

neutrophils, hence adding another layer of complexity to the regulation of CDKs in autoimmune

disease. Both *MED12* and *MED13* were transcribed more, possibly promoting the formation of a

transcriptional pre-initiator complex after association with CDK8/19 and cyclin C [106]. Moreover,

regulators of chromatin remodeling required for transcriptional activation like *RB1* [107] and *EP300*

[108, 109] were detected at higher RNA levels in stimulated cells.

In addition to the analysis focusing on CDKs and their regulators, we performed a GO pathway

analysis for the enriched gene sets from unstimulated versus IC-stimulated neutrophils. Interestingly,

our pathway analysis revealed that many metabolic pathways including ATP synthesis or catabolic

processes are suppressed, while immune pathways such as granulocyte migration, chemokine and

cytokine pathways were activated.

Thus, we hypothesized that targeting CDKs by selective small-molecule inhibitors might reduce

neutrophil effector functions in IC-stimulated neutrophils *in vitro* and consequently disease severity in

IC-driven neutrophil-dependent models of autoimmune disease *in vivo*. Although there are multiple

studies published on CDK inhibition in different contexts, they rarely focus on neutrophils and utilise

relatively unselective CDKi. However, pan-CDK inhibition is often associated with severe side effects

[110]. Here, we therefore chose more selective CDK inhibitors to investigate their role on neutrophil

functions upon IC activation.

Hence, we analyzed several functions that are critical for neutrophil recruitment as well as apoptosis.

As multiple studies suggest that ROS release might be a predictor of *in vivo* efficacy in experimental

EBA [74, 75, 79, 111], all CDKi that significantly reduced ROS release were tested as systemic

treatment in the local antibody transfer-induced EBA model (Ro 3306, palbociclib HCl, THZ1 2HCl,

THZ2, OTS964). Moreover, inhibition of CDK1, CDK2, CDK4/6, CDK7, CDK9, or CDK12

influenced other neutrophil functions such as expression of the surface activation markers CD18 and

CD62L, and adhesion, highlighting a role for CDKs in neutrophil recruitment and activation. Viability

was only slightly influenced, with values still in the range of the negative controls. However, some

inhibitors (THZ2, OTS964, BSJ-4-116) increased apoptosis, which is a highly controlled form of cell

death leaving the cell membrane intact [25]. In line with our data, a role in regulating apoptosis has

already been suggested for CDK7 in neutrophils [22], and for CDK11 in Chinese Hamster Ovary cells

and T-cell lines [112]. Concerning neutrophils, this may protect the tissue from toxic substances like

ROS which are released during alternative forms of cell death and has been described as a potential

mechanism of action of CDKi [95, 113]. Indeed, THZ2 and OTS964 increased apoptosis while

reducing ROS. However, other inhibitors reduced ROS release without affecting apoptosis (Ro 3306,

K03861, palbociclib HCl, THZ1 2HCl, BAY1251152), implicating the involvement of different

mechanisms or signaling pathways. Interestingly, CDK9 inhibition by MC180295 or BAY1251152 did

not significantly influence apoptosis, whereas CDK9 inhibition by promiscuous inhibitors such as

flavopiridol or R-roscovitine resulted in neutrophil apoptosis induction [22, 30]. Thus, this indicated

an inhibitor-specific, probably synergistic effect caused by targeting several CDKs at once.

Subsequently, the most promising CDKi were applied systemically in the local antibody-transfer

induced EBA model. Here, inhibition of CDK7 by THZ1 2HCl or THZ2, CDK9 by MC180295, or

CDK11 by OTS964 were found to significantly reduce the disease severity. Remarkably, CDK7,

CDK9, and CDK11 are all described as transcriptional kinases [3]. This is in line with our expectation

considering that neutrophils are non-proliferating cells. Inhibition of CDK1 with Ro 3306 or CDK4/6

with palbociclib HCl, in contrast, did not reduce the AESA. Although palbociclib HCl showed

promising effects *in vitro* and is licensed as treatment for breast cancer, where it improves the

neutrophil-to-lymphocyte ratio in combination with the aromatase inhibitor letrozole [114], other data

indicate an opposite role for CDK6 in regulating NET formation as CDK6$^{-/-}$ mice failed to produce

NETs upon PMA stimulation [28]. Additionally, CDK6$^{-/-}$ mice exhibited a higher amount of primary

granule proteins, suggesting CDK6 as a regulator of neutrophil granularity [115]. These data might

explain the outcome seen in EBA.

The most efficient CDKi, MC180295, was next administered in two other models to verify the

neutrophil-dependent effects: STA, which is neutrophil- and macrophage-dependent, and ITP, which is

mostly neutrophil-independent. In STA, MC180295 significantly reduced disease severity and levels

of several inflammatory cytokines. These findings complement a study describing protective effects on

human chondrocytes and cartilage explants by CDK9 inhibition with flavopiridol or small interfering

RNA. Less pro-inflammatory cytokines are released, preventing cartilage degradation [32]. Of note,

recent data show the induction of neutrophil reverse migration from the inflammation site back to the

circulation upon CDK9 inhibition in zebrafish larvae, hence contributing to the resolution of

inflammation [116].

Nevertheless, compared to the almost complete protection from disease development observed in the

local antibody transfer-induced EBA model, the MC180295-treated STA mice showed a residual mean

arthritis score of 2 on the final day, representing still 22 % of that observed in the control mice, in

contrast to only 5 % in the EBA model. This finding might reflect the different functional relevance of

neutrophils in the two distinct models. While EBA is described to be primarily neutrophil-driven [38],

neutrophils also play a critical [45], yet not exclusive, role in STA [47]. In the latter model, the

function of macrophages, NK cells, mast cells, and synovial fibroblasts during the effector phase of

the disease is also well established [44] and might explain the differences in efficacy seen on both

models. A study conducted with LPS-stimulated human monocyte-derived macrophages demonstrated

a reduction of IL-6 and TNF after treatment with the more promiscuous CDK9i AT7519 or

roscovitine, supporting a role for macrophages in the resolution of inflammation [117]. In addition,

administration of the CDK9i PC585 and PC579 in murine arthritis models led to the downregulation

of the anti-apoptotic protein MCl-1, thus favoring apoptosis [118]. As delayed apoptosis is a feature of

RA [46], CDK inhibition might be a reasonable approach in this context.

Conversely, no effects on platelet depletion or recovery were observed in ITP. Contrasting with EBA

and STA, neutrophils are considered dispensable for the effector phase of ITP. Rather other immune

cells like tissue-resident macrophages are involved [51–53]. This highlights the role for CDK9 in IC-

induced neutrophil-dependent diseases like EBA and STA.

Regarding potential limitations of this study, only one concentration of each inhibitor was used *in vivo*.

Thus, any effects which might be seen at different concentrations remain unclear. However, these

concentrations had successfully been used in mouse models before and were hence deemed safe to

apply. For example, MC180295 demonstrated an anti-tumoral effect in NOD.Cg-

Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ mice lacking mature B and T cells [64] and palbociclib is already in clinical

use for the treatment of metastatic breast cancer [119]. Furthermore, our *in vitro* data indicate a

mechanistic effect of CDKi rather than a mere cytotoxic effect.

Additionally, only one inhibitor per target (except for CDK7 since both inhibitors were equally

promising) was tested. We are aware that other inhibitors with different selectivity or bioavailability

might be more efficient, nevertheless the use of further inhibitors would have exceeded the scope of

this explorative study. The way is now paved for further studies grounding on this work.

Instead of pharmacological inhibition, genetic knockouts could be a more precise alternative to

address our research questions. Unfortunately, genetic deletions of many CDKs are embryonically

lethal, with CDK1 being indispensable for early embryogenesis [120, 121]. CDK4- and CDK6-

deficient mice display problems in embryonic development [122, 123] similar to CDK8$^{-/-}$ [124],

CDK11$^{-/-}$ [125] or CDK12$^{-/-}$ mice [126]. Therefore, we consider pharmacological inhibition to be a

practicable and clinically relevant alternative, though future studies with neutrophil-specific inducible

depletion may be informative.

542   **Conclusions**

Taken together, we have demonstrated the functional relevance of CDKs including CDK7, CDK9, and

CDK11 in neutrophil IC-induced kinase signaling both *in vitro* and *in vivo*. Our findings contribute to

the exploration of potential novel treatment options for autoimmune diseases such as PD or RA.

List of abbreviations

AUC - area under the curve

BSA - bovine serum albumin

CAK - CDK-activating kinase

CDK - cyclin-dependent kinase

CDKi - small-molecule CDK inhibitor

CKI - intrinsic CDK inhibitor

COL7 - collagen VII

DMSO - dimethyl sulfoxide

EBA - epidermolysis bullosa acquisita

ENA - European Nucleotide Archive

FcγR - fragment crystallisable gamma receptor

GO - Gene Ontology

H&E - hematoxylin and eosin

i.p. – intraperitoneal

IC - immune complex

ITP - immune thrombocytopenia

MACS - magnetic activated cell sorting

NET - neutrophil extracellular trap

PBS - phosphate-buffered saline

PD - pemphigoid disease

PMN - polymorphonuclear granulocyte

qRT-PCR - real-time quantitative reverse transcription polymerase chain reaction

RA - rheumatoid arthritis

RNAseq - bulk RNA sequencing

ROS - reactive oxygen species

RT - room temperature

STA - serum-transfer arthritis

Declarations

Ethics approval and consent to participate

Whole blood was obtained from healthy volunteers after written informed consent. All experiments

using human samples were approved by the local ethics committee (AZ #20-463, University of

Lübeck, Lübeck, Germany) and were performed in accordance with the Declaration of Helsinki.

Animal experiments were approved by local authorities of the Animal Care and Use Committee

(government of Schleswig-Holstein and government of Lower Franconia) and the Institutional Animal

Care and Use Committee at Rush University. All experiments were performed by certified personnel

(AZ 108/08-15, AZ 97-11/20 and 55.2.2-2532-2-2085) following the ARRIVE guidelines.

Consent for publication

Not applicable

Availability of data and materials

The RNAseq dataset generated and analyzed during the current study is available in the European
Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB103987
(https://www.ebi.ac.uk/ena/browser/view/PRJEB103987), all other data generated or analyzed during
this study are included in this published article and its supplementary information files.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was funded by the Cluster of Excellence "Precision Medicine in Chronic Inflammation"
(EXC 2167), the Research Training Group "Defining and Targeting Autoimmune Pre-Disease" (RTG
2633), and the Collaborative Research Center "Pathomechanisms of Antibody-mediated
Autoimmunity" (CRC 1526), all from the Deutsche Forschungsgemeinschaft and the Schleswig-
Holstein Excellence-Chair Program from the State of Schleswig Holstein.

Author's contributions

Investigation: MS, LV, APM, DM, SJ, MR, AKS, SMV, YX, HG, JL, writing (original draft): MS,
visualization: MS, LV, AKS, formal analysis: CO, NE, AV, MS, AKS, LV, KA, resources: RV, GV, FP,
XY, RJL, KTA, conceptualization: RJL, AL, KB, supervision: AL, KB, KTA, software: AV, project
administration: AL, KB, writing (review & editing): all.

Acknowledgements

We thank Kathrin Kalies, Petra Lau, Astrid Fischer, Claudia Kauderer, Alexandra Wobig, Daniela
Rieck, Norbert Reiling, Carolin Möller, and Cindy Jensen for their support.

References

1. Nurse P, Thuriaux P, Nasmyth K. Genetic control of the cell division cycle in the fission yeast

Schizosaccharomyces pombe. Mol Gen Genet MGG. 1976;146:167–78.

https://doi.org/10.1007/BF00268085.

2. Beach D, Durkacz B, Nurse P. Functionally homologous cell cycle control genes in budding and

fission yeast. Nature. 1982;300:706–9. https://doi.org/10.1038/300706a0.

3. Malumbres M. Cyclin-dependent kinases. Genome Biol. 2014;15:122.

https://doi.org/10.1186/gb4184.

4. Nurse P. A Long Twentieth Century of the Cell Cycle and Beyond. Cell. 2000;100:71–8.

https://doi.org/10.1016/S0092-8674(00)81684-0.

5. Lee MG, Nurse P. Complementation used to clone a human homologue of the fission yeast cell

cycle control gene cdc2. Nature. 1987;327:31–5. https://doi.org/10.1038/327031a0.

6. Evans T, Rosenthal ET, Youngblom J, Distel D, Hunt T. Cyclin: A protein specified by maternal

mRNA in sea urchin eggs that is destroyed at each cleavage division. Cell. 1983;33:389–96.

https://doi.org/10.1016/0092-8674(83)90420-8.

7. Cao L, Chen F, Yang X, Xu W, Xie J, Yu L. Phylogenetic analysis of CDK and cyclin proteins in

premetazoan lineages. BMC Evol Biol. 2014;14:10. https://doi.org/10.1186/1471-2148-14-10.

8. Bury M, Le Calvé B, Ferbeyre G, Blank V, Lessard F. New Insights into CDK Regulators: Novel

Opportunities for Cancer Therapy. Trends Cell Biol. 2021;31:331–44.

https://doi.org/10.1016/j.tcb.2021.01.010.

9. Draetta G, Beach D. Activation of cdc2 protein kinase during mitosis in human cells: Cell cycle-

dependent phosphorylation and subunit rearrangement. Cell. 1988;54:17–26.

https://doi.org/10.1016/0092-8674(88)90175-4.

10. Russo AA, Jeffrey PD, Pavletich NP. Structural basis of cyclin-dependent kinase activation by

phosphorylation. Nat Struct Mol Biol. 1996;3:696–700. https://doi.org/10.1038/nsb0896-696.

11. Pluta AJ, Studniarek C, Murphy S, Norbury CJ. Cyclin☐dependent kinases: Masters of the

eukaryotic universe. WIREs RNA. 2024;15:e1816. https://doi.org/10.1002/wrna.1816.

12. Tassan JP, Schultz SJ, Bartek J, Nigg EA. Cell cycle analysis of the activity, subcellular

localization, and subunit composition of human CAK (CDK-activating kinase). J Cell Biol.

1994;127:467–78. https://doi.org/10.1083/jcb.127.2.467.

13. Solomon MJ, Lee T, Kirschner MW. Role of phosphorylation in p34cdc2 activation: identification

of an activating kinase. Mol Biol Cell. 1992;3:13–27. https://doi.org/10.1091/mbc.3.1.13.

14. Fisher RP, Morgan DO. A novel cyclin associates with M015/CDK7 to form the CDK-activating

kinase. Cell. 1994;78:713–24. https://doi.org/10.1016/0092-8674(94)90535-5.

15. Parker LL, Piwnica-Worms H. Inactivation of the p34^{cdc2} -Cyclin B Complex by the Human

WEE1 Tyrosine Kinase. Science. 1992;257:1955–7. https://doi.org/10.1126/science.1384126.

16. Jeffrey PD, Tong L, Pavletich NP. Structural basis of inhibition of CDK–cyclin complexes by

INK4 inhibitors. Genes Dev. 2000;14:3115–25. https://doi.org/10.1101/gad.851100.

17. Pavletich NP. Mechanisms of cyclin-dependent kinase regulation: structures of cdks, their cyclin

activators, and cip and INK4 inhibitors. J Mol Biol. 1999;287:821–8.

https://doi.org/10.1006/jmbi.1999.2640.

18. Lim S, Kaldis P. Cdks, cyclins and CKIs: roles beyond cell cycle regulation. Development.

2013;140:3079–93. https://doi.org/10.1242/dev.091744.

19. Pellarin I, Dall'Acqua A, Favero A, Segatto I, Rossi V, Crestan N, et al. Cyclin-dependent protein

kinases and cell cycle regulation in biology and disease. Signal Transduct Target Ther. 2025;10:11.

https://doi.org/10.1038/s41392-024-02080-z.

20. Nilson KA, Guo J, Turek ME, Brogie JE, Delaney E, Luse DS, et al. THZ1 Reveals Roles for

Cdk7 in Co-transcriptional Capping and Pausing. Mol Cell. 2015;59:576–87.

https://doi.org/10.1016/j.molcel.2015.06.032.

21. Peissert S, Schlosser A, Kendel R, Kuper J, Kisker C. Structural basis for CDK7 activation by

MAT1 and Cyclin H. Proc Natl Acad Sci. 2020;117:26739–48.

https://doi.org/10.1073/pnas.2010885117.

22. Leitch AE, Lucas CD, Marwick JA, Duffin R, Haslett C, Rossi AG. Cyclin-dependent kinases 7

and 9 specifically regulate neutrophil transcription and their inhibition drives apoptosis to promote

resolution of inflammation. Cell Death Differ. 2012;19:1950–61. https://doi.org/10.1038/cdd.2012.80.

23. Syahirah R, Hsu AY, Deng Q. A curious case of cyclin-dependent kinases in neutrophils. J Leukoc

Biol. 2022;111:1057–68. https://doi.org/10.1002/JLB.2RU1021-573R.

24. Aroca-Crevillén A, Vicanolo T, Ovadia S, Hidalgo A. Neutrophils in Physiology and Pathology.

Annu Rev Pathol Mech Dis. 2024;19:227–59. https://doi.org/10.1146/annurev-pathmechdis-051222-

015009.

25. Savill JS, Wyllie AH, Henson JE, Walport MJ, Henson PM, Haslett C. Macrophage phagocytosis

of aging neutrophils in inflammation. Programmed cell death in the neutrophil leads to its recognition

by macrophages. J Clin Invest. 1989;83:865–75. https://doi.org/10.1172/JCI113970.

26. Greenlee-Wacker MC. Clearance of apoptotic neutrophils and resolution of inflammation.

Immunol Rev. 2016;273:357–70. https://doi.org/10.1111/imr.12453.

27. Hsu AY, Wang D, Liu S, Lu J, Syahirah R, Bennin DA, et al. Phenotypical microRNA screen

reveals a noncanonical role of CDK2 in regulating neutrophil migration. Proc Natl Acad Sci.

2019;116:18561–70. https://doi.org/10.1073/pnas.1905221116.

28. Amulic B, Knackstedt SL, Abu Abed U, Deigendesch N, Harbort CJ, Caffrey BE, et al. Cell-Cycle

Proteins Control Production of Neutrophil Extracellular Traps. Dev Cell. 2017;43:449-462.e5.

https://doi.org/10.1016/j.devcel.2017.10.013.

29. Rosales JL, Ernst JD, Hallows J, Lee K-Y. GTP-dependent Secretion from Neutrophils Is

Regulated by Cdk5. J Biol Chem. 2004;279:53932–6. https://doi.org/10.1074/jbc.M408467200.

30. Wang K, Hampson P, Hazeldine J, Krystof V, Strnad M, Pechan P, et al. Cyclin-Dependent Kinase

9 Activity Regulates Neutrophil Spontaneous Apoptosis. PLoS ONE. 2012;7:e30128.

https://doi.org/10.1371/journal.pone.0030128.

31. Bajaj NPS, Al-Sarraj ST, Anderson V, Kibble M, Leigh N, Miller CCJ. Cyclin-dependent kinase-5

is associated with lipofuscin in motor neurones in amyotrophic lateral sclerosis. Neurosci Lett.

1998;245:45–8. https://doi.org/10.1016/S0304-3940(98)00176-1.

32. Yik JHN, Hu Z, Kumari R, Christiansen BA, Haudenschild DR. Cyclin☐Dependent Kinase 9

Inhibition Protects Cartilage From the Catabolic Effects of Proinflammatory Cytokines. Arthritis

Rheumatol. 2014;66:1537–46. https://doi.org/10.1002/art.38378.

33. Wu L, Qin Y, Xia S, Dai M, Han X, Wu Y, et al. Identification of Cyclin☐Dependent Kinase 1 as a

Novel Regulator of Type I Interferon Signaling in Systemic Lupus Erythematosus. Arthritis

Rheumatol. 2016;68:1222–32. https://doi.org/10.1002/art.39543.

34. Sinha AA, Lopez MT, McDevitt HO. Autoimmune Diseases: The Failure of Self Tolerance.

Science. 1990;248:1380–8. https://doi.org/10.1126/science.1972595.

35. Woodley DT, Burgeson RE, Lunstrum G, Bruckner-Tuderman L, Reese MJ, Briggaman RA.

Epidermolysis bullosa acquisita antigen is the globular carboxyl terminus of type VII procollagen. J

Clin Invest. 1988;81:683–7. https://doi.org/10.1172/JCI113373.

36. Liu Z, Giudice GJ, Zhou X, Swartz SJ, Troy JL, Fairley JA, et al. A major role for neutrophils in

experimental bullous pemphigoid. J Clin Invest. 1997;100:1256–63.

https://doi.org/10.1172/JCI119639.

37. Matsumoto D, Amatya B, Tsuruta D, Hiroyasu S. The pathological function of neutrophils in

pemphigoid diseases. Dermatol Sin. 2024;42:80–8. https://doi.org/10.4103/ds.DS-D-24-00027.

38. Chiriac M, Roesler J, Sindrilaru A, Scharffetter-Kochanek K, Zillikens D, Sitaru C. NADPH

oxidase is required for neutrophil-dependent autoantibody-induced tissue damage. J Pathol.

2007;212:56–65. https://doi.org/10.1002/path.2157.

39. Sezin T, Krajewski M, Wutkowski A, Mousavi S, Chakievska L, Bieber K, et al. The Leukotriene

B4 and its Receptor BLT1 Act as Critical Drivers of Neutrophil Recruitment in Murine Bullous

Pemphigoid-Like Epidermolysis Bullosa Acquisita. J Invest Dermatol. 2017;137:1104–13.

https://doi.org/10.1016/j.jid.2016.12.021.

40. Bieber K, Witte M, Sun S, Hundt JE, Kalies K, Dräger S, et al. T cells mediate autoantibody-

induced cutaneous inflammation and blistering in epidermolysis bullosa acquisita. Sci Rep.

2016;6:38357. https://doi.org/10.1038/srep38357.

41. Schmidt E, Zillikens D. Pemphigoid diseases. The Lancet. 2013;381:320–32.

https://doi.org/10.1016/S0140-6736(12)61140-4.

42. Koga H, Prost-Squarcioni C, Iwata H, Jonkman MF, Ludwig RJ, Bieber K. Epidermolysis Bullosa

Acquisita: The 2019 Update. Front Med. 2019;5:362. https://doi.org/10.3389/fmed.2018.00362.

43. Kridin K, Kneiber D, Kowalski EH, Valdebran M, Amber KT. Epidermolysis bullosa acquisita: A

comprehensive review. Autoimmun Rev. 2019;18:786–95.

https://doi.org/10.1016/j.autrev.2019.06.007.

44. Christensen AD, Haase C, Cook AD, Hamilton JA. K/BxN Serum-Transfer Arthritis as a Model

for Human Inflammatory Arthritis. Front Immunol. 2016;7:213.

https://doi.org/10.3389/fimmu.2016.00213.

45. Wipke BT, Allen PM. Essential Role of Neutrophils in the Initiation and Progression of a Murine

Model of Rheumatoid Arthritis. J Immunol. 2001;167:1601–8.

https://doi.org/10.4049/jimmunol.167.3.1601.

46. Wright HL, Moots RJ, Edwards SW. The multifactorial role of neutrophils in rheumatoid arthritis.

Nat Rev Rheumatol. 2014;10:593–601. https://doi.org/10.1038/nrrheum.2014.80.

47. Solomon S, Rajasekaran N, Jeisy☐Walder E, Snapper SB, Illges H. A crucial role for macrophages

in the pathology of K/B × N serum☐induced arthritis. Eur J Immunol. 2005;35:3064–73.

https://doi.org/10.1002/eji.200526167.

48. Lee DM, Friend DS, Gurish MF, Benoist C, Mathis D, Brenner MB. Mast Cells: A Cellular Link

Between Autoantibodies and Inflammatory Arthritis. Science. 2002;297:1689–92.

https://doi.org/10.1126/science.1073176.

49. Nigrovic PA, Binstadt BA, Monach PA, Johnsen A, Gurish M, Iwakura Y, et al. Mast cells

contribute to initiation of autoantibody-mediated arthritis via IL-1. Proc Natl Acad Sci.

2007;104:2325–30. https://doi.org/10.1073/pnas.0610852103.

50. McInnes IB, Schett G. The Pathogenesis of Rheumatoid Arthritis. N Engl J Med. 2011;365:2205–

19. https://doi.org/10.1056/NEJMra1004965.

51. Biburger M, Aschermann S, Schwab I, Lux A, Albert H, Danzer H, et al. Monocyte Subsets

Responsible for Immunoglobulin G-Dependent Effector Functions In Vivo. Immunity. 2011;35:932–

44. https://doi.org/10.1016/j.immuni.2011.11.009.

52. Zufferey A, Kapur R, Semple J. Pathogenesis and Therapeutic Mechanisms in Immune

Thrombocytopenia (ITP). J Clin Med. 2017;6:16. https://doi.org/10.3390/jcm6020016.

53. Wöhner M, Brechtelsbauer S, Friedrich N, Vorsatz C, Bulang J, Liang C, et al. Tissue niche

occupancy determines the contribution of fetal- versus bone-marrow-derived macrophages to IgG

effector functions. Cell Rep. 2024;43:113757. https://doi.org/10.1016/j.celrep.2024.113757.

54. Schwartz PA, Murray BW. Protein kinase biochemistry and drug discovery. Bioorganic Chem.

2011;39:192–210. https://doi.org/10.1016/j.bioorg.2011.07.004.

55. Zarrin AA, Bao K, Lupardus P, Vucic D. Kinase inhibition in autoimmunity and inflammation. Nat

Rev Drug Discov. 2021;20:39–63. https://doi.org/10.1038/s41573-020-0082-8.

56. Cicenas J, Valius M. The CDK inhibitors in cancer research and therapy. J Cancer Res Clin Oncol.

2011;137:1409–18. https://doi.org/10.1007/s00432-011-1039-4.

57. Wu CX, Wang XQ, Chok SH, Man K, Tsang SHY, Chan ACY, et al. Blocking CDK1/PDK1/β-

Catenin signaling by CDK1 inhibitor RO3306 increased the efficacy of sorafenib treatment by

targeting cancer stem cells in a preclinical model of hepatocellular carcinoma. Theranostics.

2018;8:3737–50. https://doi.org/10.7150/thno.25487.

58. Matsuo Y, Park J-H, Miyamoto T, Yamamoto S, Hisada S, Alachkar H, et al. TOPK inhibitor

induces complete tumor regression in xenograft models of human cancer through inhibition of

cytokinesis. Sci Transl Med. 2014;6. https://doi.org/10.1126/scitranslmed.3010277.

59. Yang J, Luo W, Ward P, Chen S-C, Zebala J, Maeda D, et al. Combined treatment with CDK4/6,

CDK2, and CXCR1/2 inhibitors effectively halts the growth of BRAF wild-type melanoma tumors.

Front Oncol. 2025;15:1609735. https://doi.org/10.3389/fonc.2025.1609735.

60. Fry DW, Harvey PJ, Keller PR, Elliott WL, Meade M, Trachet E, et al. Specific inhibition of

cyclin-dependent kinase 4/6 by PD 0332991 and associated antitumor activity in human tumor

xenografts. Mol Cancer Ther. 2004;3:1427–38. https://doi.org/10.1158/1535-7163.1427.3.11.

61. Kwiatkowski N, Zhang T, Rahl PB, Abraham BJ, Reddy J, Ficarro SB, et al. Targeting

transcription regulation in cancer with a covalent CDK7 inhibitor. Nature. 2014;511:616–20.

https://doi.org/10.1038/nature13393.

62. Huang T, Ding X, Xu G, Chen G, Cao Y, Peng C, et al. CDK7 inhibitor THZ1 inhibits MCL1

synthesis and drives cholangiocarcinoma apoptosis in combination with BCL2/BCL-XL inhibitor

ABT-263. Cell Death Dis. 2019;10:602. https://doi.org/10.1038/s41419-019-1831-7.

63. Wang Y, Zhang T, Kwiatkowski N, Abraham BJ, Lee TI, Xie S, et al. CDK7-Dependent

Transcriptional Addiction in Triple-Negative Breast Cancer. Cell. 2015;163:174–86.

https://doi.org/10.1016/j.cell.2015.08.063.

64. Zhang H, Pandey S, Travers M, Sun H, Morton G, Madzo J, et al. Targeting CDK9 Reactivates

Epigenetically Silenced Genes in Cancer. Cell. 2018;175:1244-1258.e26.

https://doi.org/10.1016/j.cell.2018.09.051.

65. Gallagher PF, Coyle V, Evans TRJ, Plummer ER, Clive S, McGuigan L, et al. A CRUK first-in-

human phase I trial of a CDC7 Inhibitor, LY3143921 hydrate, in patients with advanced solid tumors.

J Clin Oncol. 2019;37 15_suppl:TPS3167–TPS3167.

https://doi.org/10.1200/JCO.2019.37.15_suppl.TPS3167.

66. Jiang L, Yu Y, Li Z, Gao Y, Zhang H, Zhang M, et al. BMS-265246, a Cyclin-Dependent Kinase

Inhibitor, Inhibits the Infection of Herpes Simplex Virus Type 1. Viruses. 2023;15:1642.

https://doi.org/10.3390/v15081642.

67. Ozfiliz-Kilbas P, Sarikaya B, Obakan-Yerlikaya P, Coker-Gurkan A, Arisan ED, Temizci B, et al.

Cyclin-dependent kinase inhibitors, roscovitine and purvalanol, induce apoptosis and autophagy

related to unfolded protein response in HeLa cervical cancer cells. Mol Biol Rep. 2018;45:815–28.

https://doi.org/10.1007/s11033-018-4222-8.

68. Jiang B, Gao Y, Che J, Lu W, Kaltheuner IH, Dries R, et al. Discovery and resistance mechanism

of a selective CDK12 degrader. Nat Chem Biol. 2021;17:675–83. https://doi.org/10.1038/s41589-021-

00765-y.

69. Quereda V, Bayle S, Vena F, Frydman SM, Monastyrskyi A, Roush WR, et al. Therapeutic

Targeting of CDK12/CDK13 in Triple-Negative Breast Cancer. Cancer Cell. 2019;36:545-558.e7.

https://doi.org/10.1016/j.ccell.2019.09.004.

70. Kozyrska K, Pilia G, Vishwakarma M, Wagstaff L, Goschorska M, Cirillo S, et al. p53 directs

leader cell behavior, migration, and clearance during epithelial repair. Science. 2022;375:eabl8876.

https://doi.org/10.1126/science.abl8876.

71. Akamatsu M, Mikami N, Ohkura N, Kawakami R, Kitagawa Y, Sugimoto A, et al. Conversion of

antigen-specific effector/memory T cells into Foxp3-expressing T_{reg} cells by inhibition of CDK8/19.

Sci Immunol. 2019;4:eaaw2707. https://doi.org/10.1126/sciimmunol.aaw2707.

72. Lücking U, Kosemund D, Böhnke N, Lienau P, Siemeister G, Denner K, et al. Changing for the

Better: Discovery of the Highly Potent and Selective CDK9 Inhibitor VIP152 Suitable for Once

Weekly Intravenous Dosing for the Treatment of Cancer. J Med Chem. 2021;64:11651–74.

https://doi.org/10.1021/acs.jmedchem.1c01000.

73. Skoog WA, Beck WS. Studies on the Fibrinogen, Dextran and Phytohemagglutinin Methods of

Isolating Leukocytes. Blood. 1956;11:436–54. https://doi.org/10.1182/blood.V11.5.436.436.

74. Zillikens H, Kasprick A, Osterloh C, Gross N, Radziewitz M, Hass C, et al. Topical Application of

the PI3Kβ-Selective Small Molecule Inhibitor TGX-221 Is an Effective Treatment Option for

Experimental Epidermolysis Bullosa Acquisita. Front Med. 2021;8:713312.

https://doi.org/10.3389/fmed.2021.713312.

75. Stüssel P, Schulze Dieckhoff K, Künzel S, Hartmann V, Gupta Y, Kaiser G, et al. Propranolol Is an

Effective Topical and Systemic Treatment Option for Experimental Epidermolysis Bullosa Acquisita. J

Invest Dermatol. 2020;140:2408–20. https://doi.org/10.1016/j.jid.2020.04.025.

76. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. Genome Biol. 2014;15:550. https://doi.org/10.1186/s13059-014-0550-8.

77. The Gene Ontology Consortium, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et

al. The Gene Ontology knowledgebase in 2023. GENETICS. 2023;224:iyad031.

https://doi.org/10.1093/genetics/iyad031.

78. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology: tool for

the unification of biology. Nat Genet. 2000;25:25–9. https://doi.org/10.1038/75556.

79. Ghorbanalipoor S, Matsumoto K, Gross N, Heimberg L, Krause M, Veldkamp W, et al. High

throughput screening identifies repurposable drugs for modulation of innate and acquired immune

responses. J Autoimmun. 2024;148:103302. https://doi.org/10.1016/j.jaut.2024.103302.

80. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time

Quantitative PCR and the 2−∆∆CT Method. Methods. 2001;25:402–8.

https://doi.org/10.1006/meth.2001.1262.

81. Zhang X, Ding L, Sandford AJ. Selection of reference genes for gene expression studies in human

neutrophils by real-time PCR. BMC Mol Biol. 2005;6:4. https://doi.org/10.1186/1471-2199-6-4.

82. Recke A, Trog LM, Pas HH, Vorobyev A, Abadpour A, Jonkman MF, et al. Recombinant Human

IgA1 and IgA2 Autoantibodies to Type VII Collagen Induce Subepidermal Blistering Ex Vivo. J

Immunol. 2014;193:1600–8. https://doi.org/10.4049/jimmunol.1400160.

83. Yu X, Akbarzadeh R, Pieper M, Scholzen T, Gehrig S, Schultz C, et al. Neutrophil Adhesion Is a

Prerequisite for Antibody-Mediated Proteolytic Tissue Damage in Experimental Models of

Epidermolysis Bullosa Acquisita. J Invest Dermatol. 2018;138:1990–8.

https://doi.org/10.1016/j.jid.2018.03.1499.

84. Yang K, Yin J, Yue X, Bieber K, Riemekasten G, Ludwig RJ, et al. Luteolin peracetate and

gossypolone inhibit immune complex-mediated neutrophil activation in vitro and dermal-epidermal

separation in an ex vivo model of epidermolysis bullosa acquisita. Front Immunol. 2023;14:1196116.

https://doi.org/10.3389/fimmu.2023.1196116.

85. Bieber K, Ernst AL, Tukaj S, Holtsche MM, Schmidt E, Zillikens D, et al. Analysis of serum

markers of cellular immune activation in patients with bullous pemphigoid. Exp Dermatol.

2017;26:1248–52. https://doi.org/10.1111/exd.13382.

86. Sitaru C, Mihai S, Otto C, Chiriac MT, Hausser I, Dotterweich B, et al. Induction of dermal-

epidermal separation in mice by passive transfer of antibodies specific to type VII collagen. J Clin

Invest. 2005;115:870–8. https://doi.org/10.1172/JCI200521386.

87. Kasprick A, Bieber K, Ludwig RJ. Drug Discovery for Pemphigoid Diseases. Curr Protoc

Pharmacol. 2019;84:e55. https://doi.org/10.1002/cpph.55.

88. Seeling M, Pöhnl M, Kara S, Horstmann N, Riemer C, Wöhner M, et al. Immunoglobulin G-

dependent inhibition of inflammatory bone remodeling requires pattern recognition receptor Dectin-1.

Immunity. 2023;56:1046-1063.e7. https://doi.org/10.1016/j.immuni.2023.02.019.

89. Seeling M, Hillenhoff U, David JP, Schett G, Tuckermann J, Lux A, et al. Inflammatory monocytes

and Fcγ receptor IV on osteoclasts are critical for bone destruction during inflammatory arthritis in

mice. Proc Natl Acad Sci. 2013;110:10729–34. https://doi.org/10.1073/pnas.1301001110.

90. Ohtsubo M, Roberts JM. Cyclin-Dependent Regulation of G_1 in Mammalian Fibroblasts. Science.

1993;259:1908–12. https://doi.org/10.1126/science.8384376.

91. Resnitzky D, Gossen M, Bujard H, Reed SI. Acceleration of the G_1 /S Phase Transition by

Expression of Cyclins Dl and E with an Inducible System. Mol Cell Biol. 1994;14:1669–79.

https://doi.org/10.1128/mcb.14.3.1669-1679.1994.

92. Griffin JD, Spertini O, Ernst TJ, Belvin MP, Levine HB, Kanakura Y, et al. Granulocyte-

macrophage colony-stimulating factor and other cytokines regulate surface expression of the leukocyte

adhesion molecule-1 on human neutrophils, monocytes, and their precursors. J Immunol.

1990;145:576–84. https://doi.org/10.4049/jimmunol.145.2.576.

93. Kuhns DB, Priel DAL, Gallin JI. Loss of L-Selectin (CD62L) on Human Neutrophils Following

Exudation in Vivo. Cell Immunol. 1995;164:306–10. https://doi.org/10.1006/cimm.1995.1174.

94. Arnaout MA. Biology and structure of leukocyte β2 integrins and their role in inflammation.

F1000Research. 2016;5:2433. https://doi.org/10.12688/f1000research.9415.1.

95. Rossi AG, Sawatzky DA, Walker A, Ward C, Sheldrake TA, Riley NA, et al. Cyclin-dependent

kinase inhibitors enhance the resolution of inflammation by promoting inflammatory cell apoptosis.

Nat Med. 2006;12:1056–64. https://doi.org/10.1038/nm1468.

96. Li P, Xiao J, Zhou B, Wei J, Luo J, Chen W. SYNE1 mutation may enhance the response to

immune checkpoint blockade therapy in clear cell renal cell carcinoma patients. Aging.

2020;12:19316–24. https://doi.org/10.18632/aging.103781.

97. Mercado-Evans V, Branthoover H, Chew C, Serchejian C, Saltzman AB, Mejia ME, et al. Tamm-

Horsfall protein augments neutrophil NETosis during urinary tract infection. JCI Insight.

2025;10:e180024. https://doi.org/10.1172/jci.insight.180024.

98. Feng C, Li Y, Tai Y, Zhang W, Wang H, Lian S, et al. A neutrophil extracellular traps-related

classification predicts prognosis and response to immunotherapy in colon cancer. Sci Rep.

2023;13:19297. https://doi.org/10.1038/s41598-023-45558-6.

99. Henriksen ML, Brandt J, Iyer SSC, Thielens NM, Hansen S. Characterization of the interaction

between collectin 11 (CL-11, CL-K1) and nucleic acids. Mol Immunol. 2013;56:757–67.

https://doi.org/10.1016/j.molimm.2013.07.011.

100. Li M, Feng Z, Han R, Hu B, Zhang R, Wang H. Paclitaxel promotes mTOR signaling mediated

apoptosis in esophageal cancer cells by targeting MUC20. Thorac Cancer. 2023;14:3089–96.

https://doi.org/10.1111/1759-7714.15091.

101. Holmquist E, Okroj M, Nodin B, Jirström K, Blom AM. Sushi domain□containing protein 4

(SUSD4) inhibits complement by disrupting the formation of the classical C3 convertase. FASEB J.

2013;27:2355–66. https://doi.org/10.1096/fj.12-222042.

102. Tsou P-S, Ali RA, Lu C, Sule G, Carmona-Rivera C, Lucotti S, et al. Soluble CD13 is a potential

mediator of neutrophil-induced thrombogenic inflammation in SARS-CoV-2 infection. JCI Insight.

2025;10:e184975. https://doi.org/10.1172/jci.insight.184975.

103. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, et al. Proteomics.

Tissue-based map of the human proteome. Science. 2015;347:1260419.

https://doi.org/10.1126/science.1260419.

104. Harper JW, Elledge SJ. The role of Cdk7 in CAK function, a retro-retrospective. Genes Dev.

1998;12:285–9. https://doi.org/10.1101/gad.12.3.285.

105. Fisher RP, Jin P, Chamberlin HM, Morgan DO. Alternative mechanisms of CAK assembly

require an assembly factor or an Activating Kinase. Cell. 1995;83:47–57.

https://doi.org/10.1016/0092-8674(95)90233-3.

106. Donner AJ, Ebmeier CC, Taatjes DJ, Espinosa JM. CDK8 is a positive regulator of transcriptional

elongation within the serum response network. Nat Struct Mol Biol. 2010;17:194–201.

https://doi.org/10.1038/nsmb.1752.

107. Sanidas I, Lawrence MS, Dyson NJ. Patterns in the tapestry of chromatin-bound RB. Trends Cell

Biol. 2024;34:288–98. https://doi.org/10.1016/j.tcb.2023.07.012.

108. Delvecchio M, Gaucher J, Aguilar-Gurrieri C, Ortega E, Panne D. Structure of the p300 catalytic

core and implications for chromatin targeting and HAT regulation. Nat Struct Mol Biol.

2013;20:1040–6. https://doi.org/10.1038/nsmb.2642.

109. Ogryzko VV, Schiltz RL, Russanova V, Howard BH, Nakatani Y. The Transcriptional

Coactivators p300 and CBP Are Histone Acetyltransferases. Cell. 1996;87:953–9.

https://doi.org/10.1016/S0092-8674(00)82001-2.

110. Zhang M, Zhang L, Hei R, Li X, Cai H, Wu X, et al. CDK inhibitors in cancer therapy, an

overview of recent development. Am J Cancer Res. 2021;11:1913–35.

111. Samavedam UK, Mitschker N, Kasprick A, Bieber K, Schmidt E, Laskay T, et al. Whole-Genome

Expression Profiling in Skin Reveals SYK As a Key Regulator of Inflammation in Experimental

Epidermolysis Bullosa Acquisita. Front Immunol. 2018;9:249.

https://doi.org/10.3389/fimmu.2018.00249.

112. Lahti JM, Xiang J, Heath LS, Campana D, Kidd VJ. PITSLRE protein kinase activity is

associated with apoptosis. Mol Cell Biol. 1995;15:1–11. https://doi.org/10.1128/MCB.15.1.1.

113. Leitch A, Haslett C, Rossi A. Cyclin□dependent kinase inhibitor drugs as potential novel

anti□inflammatory and pro□resolution agents. Br J Pharmacol. 2009;158:1004–16.

https://doi.org/10.1111/j.1476-5381.2009.00402.x.

114. Kim CG, Kim MH, Kim JH, Kim S-G, Kim GM, Kim TY, et al. On-treatment derived neutrophil-

to-lymphocyte ratio and survival with palbociclib and endocrine treatment: analysis of a multicenter

retrospective cohort and the PALOMA-2/3 study with immune correlates. Breast Cancer Res.

2023;25:4. https://doi.org/10.1186/s13058-022-01601-4.

115. Fleming K, Burley K, Ponce-Garcia F, Akbari P, Naveh C, Rice C, et al. Genome-wide

association analysis of neutrophil granularity identifies CDK6 as a regulator of primary granules.

iScience. 2025;28:113072. https://doi.org/10.1016/j.isci.2025.113072.

116. Kaveh A, Bruton FA, Oremek MEM, Tucker CS, Taylor JM, Mullins JJ, et al. Selective Cdk9

inhibition resolves neutrophilic inflammation and enhances cardiac regeneration in larval zebrafish.

Development. 2022;149:dev199636. https://doi.org/10.1242/dev.199636.

117. McHugh BJ, Stephen J, Robb CT, Fox S, Kipari T, Cartwright JA, et al. Inhibition of Cyclin-
Dependent Kinase 9 Downregulates Cytokine Production Without Detrimentally Affecting Human
Monocyte-Derived Macrophage Viability. Front Cell Dev Biol. 2022;10:905315.
https://doi.org/10.3389/fcell.2022.905315.

118. Hellvard A, Zeitlmann L, Heiser U, Kehlen A, Niestroj A, Demuth H-U, et al. Inhibition of CDK9
as a therapeutic strategy for inflammatory arthritis. Sci Rep. 2016;6:31441.
https://doi.org/10.1038/srep31441.

119. Beaver JA, Amiri-Kordestani L, Charlab R, Chen W, Palmby T, Tilley A, et al. FDA Approval:
Palbociclib for the Treatment of Postmenopausal Patients with Estrogen Receptor–Positive, HER2-
Negative Metastatic Breast Cancer. Clin Cancer Res. 2015;21:4760–6. https://doi.org/10.1158/1078-
0432.CCR-15-1185.

120. Satyanarayana A, Berthet C, Lopez-Molina J, Coppola V, Tessarollo L, Kaldis P. Genetic
substitution of Cdk1 by Cdk2 leads to embryonic lethality and loss of meiotic function of Cdk2.
Development. 2008;135:3389–400. https://doi.org/10.1242/dev.024919.

121. Santamaría D, Barrière C, Cerqueira A, Hunt S, Tardy C, Newton K, et al. Cdk1 is sufficient to
drive the mammalian cell cycle. Nature. 2007;448:811–5. https://doi.org/10.1038/nature06046.

122. Berthet C, Klarmann KD, Hilton MB, Suh HC, Keller JR, Kiyokawa H, et al. Combined Loss of
Cdk2 and Cdk4 Results in Embryonic Lethality and Rb Hypophosphorylation. Dev Cell.
2006;10:563–73. https://doi.org/10.1016/j.devcel.2006.03.004.

123. Malumbres M, Sotillo R, Santamaría D, Galán J, Cerezo A, Ortega S, et al. Mammalian Cells
Cycle without the D-Type Cyclin-Dependent Kinases Cdk4 and Cdk6. Cell. 2004;118:493–504.
https://doi.org/10.1016/j.cell.2004.08.002.

124. Westerling T, Kuuluvainen E, Maäkelaä TP. Cdk8 Is Essential for Preimplantation Mouse
Development. Mol Cell Biol. 2007;27:6177–82. https://doi.org/10.1128/MCB.01302-06.

125. Li T, Inoue A, Lahti JM, Kidd VJ. Failure To Proliferate and Mitotic Arrest of CDK11[p110/p58] -Null

Mutant Mice at the Blastocyst Stage of Embryonic Cell Development. Mol Cell Biol. 2004;24:3188–

97. https://doi.org/10.1128/MCB.24.8.3188-3197.2004.

126. Juan H-C, Lin Y, Chen H-R, Fann M-J. Cdk12 is essential for embryonic development and the

maintenance of genomic stability. Cell Death Differ. 2016;23:1038–48.

https://doi.org/10.1038/cdd.2015.157.

Figure Legends

Fig. 1: Cell-cycle and transcriptional *CDKs* are expressed in human neutrophils and IC-

stimulation partially alters their expression. (**A**) Human neutrophils were purified and unstimulated

cells (unstim) plus cells stimulated for 6 h with COL7 anti-COL7 IgG1-IC (6 h IC) were analyzed by

bulk RNAseq. Expression in unstimulated and stimulated neutrophils was compared for the individual

CDKs on the basis of raw counts. (**B**) Significantly differentially expressed genes among CDKs,

cyclins, and selected regulators were determined and log2 fold changes (log2 FC) are displayed for the

individual samples. (**C**) Top 20 GO activated and suppressed pathways after gene set enrichment

analysis of genes significantly differentially expressed upon IC stimulation. Adjusted p value (p.

adjust) < 0.1. Gene ratio equals the ratio of differentially expressed genes in the given GO term, count

refers to the absolute number. n = 3.

Fig. 2: *CDK2* and *CDK4* expression are induced upon IC stimulation in human PMNs. (A)

Human PMNs were isolated, stimulated for 6 h with COL7 anti-COL7 IgG1-IC and subjected to qRT-

PCR detecting 14 distinct CDKs and CDC7 using the the $2^{-\Delta\Delta Ct}$ method. (**B**) In parallel, B cells and

CD4[+] T cells were purified and analyzed in the same way for comparison with unstimulated PMNs. n

= 5. Mann-Whitney test (**A**) or Kruskal-Wallis test with Dunn's multiple comparisons test (**B**). *

p ≤ 0.05. ** p ≤ 0.01. *** p ≤ 0.001.

Fig. 3: CDK7, CDK9, and CDK11 inhibition attenuate neutrophil effector functions and trigger

apoptosis. Different neutrophil functions and characteristics were analyzed after COL7 anti-COL7

IgG1-IC stimulation of human PMNs for 2 h and inhibition with three CDKi: **(A)** CD62L shedding,

(B) CD18 expression, **(C)** adhesion, **(D)** ROS release, **(E)** apoptosis, and **(F)** viability. The results

were normalized to the positive control stimulated with IgG1-IC but without inhibitor indicated by the

red line. The dose-response curves were calculated and the IC_{50}/EC_{50} values are indicated if in the

tested concentration range. Cells only, antigen only (Ag), and antibody only (Ab) served as negative

controls. Kruskal-Wallis test with Dunn's multiple comparisons test was performed. * $p \leq 0.05$. **

$p \leq 0.01$. *** $p \leq 0.001$. For apoptosis experiment using THZ2 n = 6, rest n = 5. AUC, area under the

curve.

Fig. 4: Systemic and topical CDK7i, CDK9i, and CDK11i treatment reduces disease severity in

the local EBA . **(A)** EBA was induced by injection of anti-mCOL7[c] IgG and the mice were treated

with CDKi either systemically or topically. **(B)** Exemplary clinical and histological images are shown

with the scale bars corresponding to 100 µM. Asterisks indicate cell infiltrates and arrows indicate

binding of C3/IgG to the dermal-epidermal junction. **(C)** The affected ear surface area (AESA) and the

H&E score were evaluated in the systemic treatment group for MC180295, **(D)** THZ2, **(E)** THZ1

2HCl, and **(F)** OTS964. **(G)** For the topical model, AESA was analyzed as well. Two-way ANOVA

with Šidák's multiple comparisons test was performed for AESA evaluation, except for the systemic

treatment with THZ2 and MC180295 a mixed-effects analysis with Šidák's multiple comparisons test

was conducted. For H&E scores, Mann-Whitney test was carried out. * $p \leq 0.05$. ** $p \leq 0.01$. ****

$p \leq 0.0001$. n = 8.

1011

Fig. 5: CDK9 inhibition with MC180295 improves disease severity in STA, but not in ITP. (A) To

induce STA, mice were injected with KBxN serum and treated with MC180295 for seven days. **(B)**

The course of the disease score is depicted and **(C)** reflected by the clinical presentation of the paws

and the histological stainings. The scale bars represent 100 µm. Asterisks indicate proteoglycan

staining (Safranin-O) or cell infiltration (H&E). **(D)** Cytokine levels were measured in serum and

significantly changed cytokines are shown. **(E)** ITP was induced by anti-platelet IgG2c-induced

platelet depletion and MC180295 was applied at 24 h and 4 h before depletion. **(F)** Absolute and **(G)**

relative platelet counts were monitored for up to 24 h and values normalized to 100 % at 0 h are

indicated. For arthritis score and platelet counts, two-way ANOVA with Šidák's multiple comparisons

test was performed. For cytokine levels, Mann-Whitey test was conducted. * $p \leq 0.05$. ** $p \leq 0.01$.

**** $p \leq 0.0001$. **(A-D)** n = 5 (treatment) and n = 4 (control). **(E-G)** n = 4.

1023

Fig. 6: Successful inhibition of neutrophil functions and experimental autoimmune models using

CDKi. Out of 20 CDKs that are described in humans, 14 were detected in our gene expression

analysis. After choosing selective CDKi, different *in vitro* neutrophil functional assays were

performed. The number of efficient inhibitors is indicated. Subsequently, all CDKi that inhibited ROS

release were tested as systemic treatment in the local antibody transfer-induced EBA model, except for

BAY1251152 to avoid redundancy. Three CDKi reduced the disease severity and were thus applied

topically in the same model. Additionally, the most promising CDKi was validated in STA and ITP.

Created in https://BioRender.com.

1032

1033

Fig. S1: CDK1i Ro 3306 and CDK7i THZ1 2HCl reduce CD62L shedding in IC-stimulated

human PMNs. (A-K) Human PMNs were stimulated for 2 h with hCOL7 anti-COL7 IgG1-IC and

subsequently analyzed for CD62L shedding. Inhibitors targeting specific CDKs were tested in flow

cytometry for their ability to minimize CD62L shedding. Based on FSC-SSC gating, alive singlet

CD45$^+$ cells were considered. The results were normalized to the positive control stimulated with

IgG1-IC but without inhibitor indicated by the red line. The dose-response curves were calculated and

the IC_{50}/EC_{50} values are indicated if in the tested concentration range. Cells only, antigen only (Ag),

and antibody only (Ab) served as negative controls. Kruskal-Wallis test with Dunn's multiple

comparisons test was performed. * $p \leq 0.05$. ** $p \leq 0.01$. *** $p \leq 0.001$. n = 5.

1043

Fig. S2: CDK1i Ro 3306 or CDK7i THZ1 2HCl decrease CD18 expression on IC-activated

human PMNs. (A-K) Following stimulation of human PMNs for 2 h with hCOL7 anti-COL7 IgG1-

IC and treatment with CDKi, the cells were analyzed for their CD18 expression in flow cytometry.

Based on FSC-SSC gating. alive singlet CD45[+] cells were considered. The results were normalized to

the positive control stimulated with IgG1-IC but without inhibitor indicated by the red line. The dose-

response curves were calculated and the IC_{50} values are indicated if in the tested concentration range.

Cells only, antigen only (Ag), and antibody only (Ab) served as negative controls. Kruskal-Wallis test

with Dunn's multiple comparisons test was performed. * $p \leq 0.05$. ** $p \leq 0.01$. n = 5.

1052

Fig. S3: Adhesion of IC-activated human PMNs is not significantly reduced by the additional

CDKi used. (A-K) Human PMNs were stimulated with hCOL7 anti-COL7 IgG1-IC for 2 h.

Impedance-based measurement of adhesion was conducted using inhibitors targeting specific CDKs.

The area under the curve was determined and the results were normalized to the positive control

stimulated with IgG1-IC but without inhibitor indicated by the red line. The dose-response curves

were calculated and the IC_{50}/EC_{50} values are indicated if in the tested concentration range. Cells only,

antigen only (Ag), and antibody only (Ab) served as negative controls. Kruskal-Wallis test with

Dunn's multiple comparisons test was performed. * $p \leq 0.05$. ** $p \leq 0.01$. *** $p \leq 0.001$. n = 5.

1061

Fig. S4: Inhibition of CDK1, CDK2, CDK4/6, CDK7, or CDK9 in IC-stimulated human PMNs

minimizes ROS release. (A-K) Activation of human PMNs with hCOL7 anti-COL7 IgG1-IC for 2 h,

treatment with selective CDKi and subsequent chemiluminescence-based analysis of ROS release was

performed. The area under the curve (AUC) was calculated and the results were normalized to the

positive control stimulated with IgG1-IC but without inhibitor indicated by the red line. The dose-

response curves were calculated and the IC_{50} values are indicated if in the tested concentration range.

Cells only, antigen only (Ag), and antibody only (Ab) served as negative controls. Kruskal-Wallis test

with Dunn's multiple comparisons test was performed. * $p \leq 0.05$. ** $p \leq 0.01$. *** $p \leq 0.001$. n = 5.

1070

Fig. S5: Apoptosis in IC-stimulated human PMNs is increased by application of CDK1/2i,

CDK12i, or CDK8/19i. (A-K) Human PMNs were activated with hCOL7 anti-COL7 IgG1-IC for 2 h.

After using selective CDKi, flow cytometry was performed. Based on FSC-SSC gating, singlet CD45[+]

cells were analyzed. Annexin V[+] Zombie NIR[-] cells were considered apoptotic. The results were

normalized to the positive control stimulated with IgG1-IC but without inhibitor indicated by the red

line. The dose-response curves were calculated and the EC_{50} values are indicated if in the tested

concentration range. Cells only, antigen only (Ag), and antibody only (Ab) served as negative controls.

Kruskal-Wallis test with Dunn's multiple comparisons test was performed. * $p \leq 0.05$. ** $p \leq 0.01$. ***

$p \leq 0.001$. For apoptosis experiment using Palbociclib HCl, AS2863619, BMS-265246, and SR-4835 n

=6, rest n = 5.

1081

Fig. S6: Viability of IC-treated human PMNs is minorly affected by CDK1, CDK7, and CDK12

inhibition. (A-K) Human PMNs stimulated with hCOL7 anti-COL7 IgG1-IC and treated with CDKi

for 2 h were analyzed for toxic effects in flow cytometry. Based on FSC-SSC gating, singlet CD45[+]

cells were studied. Annexin V[-] Zombie NIR[-] cells were considered viable. The results were normalized

to the positive control stimulated with IgG1-IC but without inhibitor indicated by the red line. The

dose-response curves were calculated and the IC_{50} values are indicated if in the tested concentration

range. Cells only, antigen only (Ag), and antibody only (Ab) served as negative controls. Kruskal-

Wallis test with Dunn's multiple comparisons test was performed. * $p \leq 0.05$. n = 5.

1090

1091 **Fig. S7: CDK1i Ro 3306 or CDK4/6i palbociclib HCl do not reduce the AESA in local EBA**. **(A-**

1092 **B)** EBA was induced by injection of anti-mCOL7[c] IgG and the mice were treated systemically with

1093   three more CDKi. The affected ear surface area (AESA) was evaluated. Mixed-effects analysis with

1094   Šidák's multiple comparisons test was performed. *** $p \leq 0.001$. **(A)** n = 7, **(B)** n = 4.

1095

1096