Latent gene network expression underlies partial re-evolution of a polyphenic trait in the worker caste of ants

Angelly Vasquez-Correa[1, 2], Johanna Arnet[1], Travis Chen[1, 2], Ehab Abouheif[1, 2*]

[1] McGill University, Department of Biology, 1205 Avenue Docteur Penfield, Montreal, Quebec, H3A 1B1, Canada

[2] Center for Evolutionary & Organismal Biology, Women's Hospital, School of Medicine, Zhejiang University, Hangzhou, China

*Author for correspondence: abouheif@zju.edu.cn

Keywords: latent developmental potential, GRN, ocelli, ants, eye-antenna disc, evolutionary reversion, Formicinae, gene regulatory network, polyphenism, developmental plasticity.

Abstract

Polyphenisms—where alternative phenotypes develop from a single genome in response to environmental cues—are not only widespread in nature, but also occur at multiple levels of biological organization, from cells to individuals to societies. Polyphenism is thought to promote phenotypic diversification through the gain, loss, and re-evolution of alternative phenotypes. After the origin of a polyphenism, one of the alternative phenotypes often retains the developmental capacity to produce the ancestral trait, thereby permitting the other to evolve rapidly. Yet, little is known about the developmental processes underlying the re-evolution of polyphenic traits and how they may produce phenotypic diversification. Here we address this question by focusing on the caste polyphenism in ant societies, which produces a winged queen caste and a wingless worker caste in a single colony in response to environmental cues. We show, in a hyperdiverse group of ants, that a caste-specific trait called the ocelli (3 simple eyes on the dorsal head) is always present across queen castes but was lost and partially re-evolved multiple times, giving rise to novel patterns (1 ocelli) in the worker castes. Surprisingly, we discovered that a hidden (latent) expression of the ocelli gene regulatory network in worker castes that lost ocelli underlies the partial re-evolution of ocelli in this group. We therefore propose that latent developmental potentials may generally persist across polyphenic systems, including ant castes, and may facilitate the partial re-evolution of novel phenotypic patterns.

Introduction

Polyphenism is a form of developmental plasticity where alternative phenotypes develop from a single genome in response to environmental cues. It is phylogenetically widespread feature of plants and animals that has evolved at different levels of biological organization (Hanna et al. 2024). For example, at the population level, the mouth form polyphenism in nematode worms produces alternative big tooth (omnivorous) or small tooth (bactivorous) mouth forms that develop in response to pheromones, crowding, salt concentration, temperature, and culturing substrate (Bento et al. 2010; Bose et al. 2012; Ragsdale et al. 2013; Werner et al. 2018). At the colony-level, caste polyphenism in eusocial insects produces morphologically differentiated queen and worker castes that develop in response to temperature and nutrition (Chandra et al. 2018; Evans and Wheeler 2001; Korb 2025; Rajakumar et al. 2024). And finally, at the cellular-level, polyphenism occurs within a multicellular individual, where a single genome gives rise to differentiated cell-types during development, such as between germline and somatic cells, in response to internal cues like morphogen gradients (Brunet and King 2017; Davison and Michod 2021; Devlin et al. 2023). Polyphenism has been proposed to promote, at the macroevolutionary scale, phenotypic diversification through the gain, loss, and re-evolution of alternative phenotypes (West-Eberhard 2003). This is based on the idea that, once a polyphenic trait originates, one of the alternative morphs retains the capacity to produce the trait in the genome while the other is freer to evolve. This hypothesis has received support from a comparative study of mouth form polyphenism across 90 species of nematode worms showing a phylogenetic association between the gain and loss of alternative mouth form phenotypes and the phenotypic diversification of mouth parts (Susoy et al. 2015). Another supporting example is fat synthesis in parasitic wasps, which revealed an association between developmental plasticity and the loss and subsequent re-evolution of fat synthesis in one species (Visser et al. 2021; 2010; Peters et al. 2017). Yet, the underlying developmental and genetic processes facilitating the loss, re-evolution, and phenotypic diversification of polyphenic traits remain poorly understood. (West-Eberhard 2003; Sommer 2020).

Here we address this question by focusing on caste polyphenism in the eusocial colonies of ants, which consists of a morphological division of labor between a winged reproductive queen caste and wingless non-reproductive worker caste in almost all 16,844 described ant species (AntWeb 2024; Ward 2014). The differential expression of polyphenic traits, such as wings, that develop in queens but not workers, are called "caste-specific" traits (Abouheif 2002; Miura 2005). It has been shown that polyphenic traits, including caste polyphenism in ants, are produced during development by the differential expression of highly conserved gene regulatory networks (GRN) in response to environmental cues (Abouheif and Wray 2002; Béhague et al. 2018; Casasa et al. 2021; Davidson et al. 2023a; 2023b; Lenuzzi et al. 2023; Rajakumar et al. 2024; Vizueta et al. 2025). However, how the expression of these GRNs influences the evolution of caste-specific traits in ants remains unknown.

Here we focus on the ocelli, which are 3 small single-lens eyes on the dorsal head of most flying insects. Ocelli complement the function of the compound eyes by mediating orientation using polarised light and in the synchronization of daily activity (Buschbeck and Bok 2023; Krapp 2009). We investigate the evolution of ocelli in a hyperdiverse subfamily of ants (Formicinae), where they are universally present in the winged reproductive caste (queens and males) as 3 large ocellus that aid in mating flights and dispersal (Moser et al. 2004; Narendra et al. 2016). In contrast, ocelli in the wingless worker caste are evolutionarily labile, showing dramatic variation across species in the presence / absence or number of ocelli in the worker caste (Johnson and Rutowski 2022; Narendra et al. 2016; Narendra and Ribi 2017; Schwarz et al. 2011) (Figure 1A). In some species, adult workers completely lack ocelli, such as in *Camponotus floridanus*, while in others they are present and vary in number—there are species whose adult workers have all 3 ocelli or just 1 single ocellus, and these can be present in all or in only in a subset of workers (Figure 1A). For example, workers in *Cataglyphis bicolor* have all three ocelli, which function in light sensing and navigation, acting as a celestial compass that provides crucial directional information (Fent and Wehner 1985) (Figure 1A). In contrast, all workers of *Polyrachis bihamata* have just a single medial ocellus (Hung 1967) (Figure 1A), and in workers of *Dinomyrmex gigas*, a single medial ocellus evolved only in a subset of individuals in the worker caste called 'soldiers' with large heads but are absent in others called 'minor workers' with small heads (AntWeb 2024) (Figure 1A). How this dramatic variation in ocelli in the worker caste of formicine ants has evolved remains poorly understood.

This evolutionary lability of ocelli across the worker castes of formicine ants also provides an opportunity to understand how the GRN underlying development of ocelli influenced the evolution of this caste-specific trait. Ants are holometabolous insects, in which adult body parts develop from imaginal discs, semi-independent clusters of cells in the larvae (Held 2002; Koch and Abouheif 2019). In the fruit fly *Drosophila melanogaster*, ocelli develop from the eye-antenna imaginal disc located at the ventral region of the head capsule. The eye-antenna imaginal disc also gives rise to the head capsule, eye, antenna, and maxillary palps (Held 2002). *D. melanogaster* is the only insect where the GRN underlying ocelli development has been well characterized at the third larval stage (Blanco et al. 2009; Domínguez-Cejudo and Casares 2015; Sabat et al. 2017) (Figure 1B). The gene *orthodenticle* (*otd*) (formerly known as *ocelli-less*) is a selector gene that is necessary for specifying the ocellar region in the developing head capsule. *otd* expression in the ocellar domain, together with other genes like *hedgehog* (*hh*), initiates the development of the ocellar region and the three ocelli (two lateral ocellus and one medial ocellus) in the eye-antennal disc. The activation of these genes regulates the expression of downstream genes, such as *otd* regulating the expression of *defective proventriculus* (*dve*) (Blanco et al. 2009; Jean-Guillaume and Kumar 2022; Yorimitsu et al. 2011), whereas *hedgehog* (*hh*) activates a portion of the retinal determination network, such as *eyes absent* (*eya*), *twin of eyeless* (*toy*), *sine oculis* (*so*), and *atonal* (*ato*). These retinal determination genes have been

shown to be regulated by independent regulatory enhancers from the compound eye (Blanco et al. 2009; 2010; Jean-Guillaume and Kumar 2022) (Figure 1B). Because the eye-antennae disc and the ocelli GRN have only been well characterized in *D. melanogaster*, it remains unknown whether they are conserved in ants.

To understand how the GRN underlying ocelli development may have influenced the evolution of this caste-specific trait, we first inferred the evolutionary history of ocelli in adult workers across the Formicinae using ancestral state reconstruction. We then characterized the eye-antennae disc in ants using three genes, *eyeless* (*ey*), *distal-less* (*dll*), and *otd-1*, which are known to mark the eyes (*ey*), the antenna (*dll*), and the head capsule and ocelli (*otd-1*). This characterization allowed us to investigate the expression of five key genes in the ocelli GRN, *otd-1, hh, toy, eya,* and *so*, during development of the winged and wingless castes across two formicine species.

Results

Partial reversion to a single ocellus occurs 3 times independently within the tribe Camponotini (Formicinae)

We first performed an ancestral state reconstruction to infer the evolutionary history of worker ocelli across the subfamily Formicinae. Our ancestral state reconstruction inferred a single re-gain of worker ocelli at the base or early within the Formicinae (Figure 2). Subsequent to the gain of ocelli early in the evolution of the Formicinae, our analysis inferred a single, well-supported, loss of worker ocelli at the base of the tribe Camponotini (Figure 2). Following this single loss, we inferred three (well supported) independent and partial re-evolution to a single medial ocellus in three different genera: *Camponotus gibbinotus*, *Polyrachis bihamata*, and *Dinomyrmex gigas* Figure 2). In *P. bihamata*, the single ocellus occurs in all workers in the colony, whereas in *C. gibbinotus* and *D. gigas* the single ocellus occurs only in the large-headed soldiers (Figure 1). We therefore investigated the developmental role of the ocelli GRN underlying these independent partial reversions to a single medial ocellus in this tribe of ants.

A fate map characterizing the development of the head capsule, antennae, eyes, and ocelli within the eye-antenna disc in ants

To understand whether the GRN underlying the development of ocelli influenced the reversions of this caste-specific trait in workers, we first had to characterize the development of the head in ants. In *D. melanogaster*, the head develops from the eye-antenna disc, which is segregated into regions marked by the expression of highly conserved developmental genes (Haynie and Bryant 1986; Held 2002). In the Florida carpenter ant *C. floridanus*, we found that, similar to *D.*

melanogaster, expression of *otd-1* marks the precursor regions of the head capsule and ocelli, *ey* marks the precursor regions of the eyes, and *dll* marks the precursor regions of the antennae. During the first larval instar, expression of *otd-1* emerges primarily in the head capsule in middle part of the disc between the antenna and compound eye (Figure 3B.). In contrast, *dll* and *eya* expression delineate the precursor regions of the antenna and eye (Figure 3C, D). During the second and third larval instar, *otd-1* is expressed in the developing head capsule and ocelli in the medial region of the disc (Figure 3 G, L), while *dll* is confined to the antenna and *ey* to the compound eye region (Figure 3H, I, M, N). Finally, during the fourth (final) larval instar, a developmental threshold mediated by juvenile hormone acts as switch point to determine whether larvae will develop either into a minor worker or soldier (major worker) (MacMillan et al. 2025). Once larvae have been determined, expression patterns of *otd-1*, *dll,* and *ey* in worker-destined larvae (Figure 3 P to T) or soldier-destined larvae (Figure 3 U to Y) remain expressed in the same regions as in the second and third instars (Figure 3Q to T, V to Y). Together, our characterization shows that the eye-antennae disc and the regional identities within it, including the precursor region of the head capsule, ocelli, eyes, and antennae are conserved in ants relative to *Drosophila*.

Expression of the ocelli GRN is conserved in winged reproductive castes but is latent in species whose adult workers completely lack ocelli

We next asked whether the ocelli GRN is conserved in the winged reproductive caste (males) relative to *Drosophila* and whether it is expressed in workers that entirely lack ocelli as adults. We address these questions using two species *C. floridanus* and *Polyrachis rastellata*. We chose these two species because *C. floridanus* is closely related to *C. gibbinotus* and *P. rastellata* is closely related to *P. bihamata*, which are two of the species that our ancestral state reconstruction inferred independent partial reversions to single medial ocellus in the worker caste (Figure 2). In the winged male caste of *C. floridanus*, we found that *otd-1, eya* and *so* are expressed where the 3 ocelli will develop, while *toy* and *hh* are expressed in the inter-ocellar region (the tissue that separates the three ocellus) (see white arrowheads in Figure 4A to B' and 5A to 5C'). Because these genes are similarly expressed within the ocellar region within the eye-antennal disc of *D. melanogaster*, we infer that the ocelli GRN is conserved in the winged reproductive castes in ants.

Surprisingly, we discovered that the ocelli GRN remains latently expressed in minor worker- and soldier-destined larvae of *C. floridanus*, which completely lack ocelli as adults. In soldier-destined larvae, all five genes remain expressed in the ocellar region within the eye-antennal disc at the beginning of the last larval instar (Figure 4C, D and Figure 5D to F). By the end of this instar, *otd-1* and *hh* remain expressed (Figure 4C', D'), but *toy*, *eye*, and *so* are either down-regulated or absent relative to their expression in the compound eye region (Figure 5D' to F'). In minor worker-destined larvae, 3 of the 5 genes (*otd-1, hh, eya*) remain expressed in the ocellar

region during the early part of the last larval instar (Figure 4E, F and Figure 5 H), whereas *toy* and *so* are expressed in the compound eye region but absent (interrupted) in the ocellar region (Figure 5G, I). Furthermore, in *P. rastellata*, whose worker caste is composed of similarly sized individuals with no subcastes, we found that *otd-1*, *eya*, and *so* remain expressed in the ocellar region within the eye-antennal disc during the early part of the last larval instar (Figure 6). Together, our results show that despite the absence of ocelli in adult workers for millions of years, the expression of the ocelli GRN remains latent during larval development.

Finally, we performed Scanning Electron Microscopy (SEM) in *C. floridanus* male, soldier, and minor worker pupae. In minor worker and soldier pupae, we discovered the existence of rudimentary ocelli that appear at the beginning of pupal development, continue to be elaborated, and then are eliminated before they molt into adult workers (Figure 7). These ocelli rudiments in are highly reduced relative to the fully functional ocelli found in male pupae. Finally, the pattern and timing of development of ocelli rudiments in the minor worker and soldier pupae coincides with the spatial expression and timing of interruption of the latent expression of the ocelli GRN. In soldier-destined larvae, ocelli GRN expression is interrupted later in development than in minor worker-destined larvae, and consequently, the ocelli rudiments in soldier pupae continue to develop longer and are more elaborated relative to those in minor worker pupae (Figure 7B to D and F to H). Therefore, expression of the latent ocelli GRN in the eye-antennae disc results in the development of ocelli rudiments in worker pupae and are then eliminated in adult workers.

Discussion

Our developmental and evolutionary data provide evidence that the latent expression of genes in workers lacking ocelli as adults is part of a latently expressed ocelli GRN, which likely facilitated at least 3 independent evolutionary reversions of this trait in the worker caste of species within the Camponotini clade. The latent expression patterns of genes in the ocelli region of developing workers lacking ocelli as adults are the same as in males that will develop fully functional ocelli but are only interrupted late in larval development. Furthermore, the timing and pattern of these late interruptions coincide with the degree of development of rudimentary ocelli in minor worker and soldier pupae before they disappear in adults (Fig. 7). This indicates that, although the expression of these genes is latent, they still retain the capacity to produce rudimentary ocelli in the pupal stage before they disappear in adults. And finally, although *eya*, *toy*, and *so*, are part of both the ocelli and compound eye GRNs in *Drosophila*, the ocelli GRN has its own distinct identity and the genes within this GRN have distinct regulatory elements and are selectively regulated (Jean-Guillaume and Kumar 2022; Zimmerman et al. 2000). In ants, our data shows that the selector gene for compound eye development in insects (*eyeless* / *Pax-6*) is expressed in the compound eyes and not ocelli, and the selector gene for ocelli (*otd-1*) is expressed in the ocelli and not compound eyes. We further show that the latent expression of *toy*,

eye, and *so* are downregulated or absent in the ocellar region, but at the same time, are strongly expressed in the regions of the compound eyes. Therefore, the mutually exclusive expression of the selector genes *eyeless* / *Pax-6* in the compound eyes and *otd-1* in ocelli suggests that, like in *Drosophila*, the compound eye and ocelli GRNs have distinct identities, ultimately leading to differential expression of downstream genes and production of different cell types; compound eyes are produced from multiple imaging forming facets, while ocelli are produced from a single lense (Buschbeck and Bok 2023; Jean-Guillaume and Kumar 2022; Mishra et al. 2021). Altogether, our data show that the expression of these genes in workers lacking ocelli as adults are part of latently expressed ocelli GRN.

Several hypotheses may explain how the ocelli GRN came to be latently expressed and maintained in developing workers that lack ocelli as adults. Perhaps the most simplistic hypothesis proposes that the presence of functional ocelli in adult queen and male castes maintains the ocelli GRN intact in the genome by keeping it under positive natural selection. This hypothesis assumes that this, as a side consequence, leads to expression of the ocelli GRN in the worker castes lacking ocelli. However, caste determination between queens and workers occurs through the action of a developmental threshold or switch, where a continuous environmental cue is translated into discrete phenotypic outcomes (Abouheif 2021; MacMillan et al. 2025; Rajakumar et al. 2024; Qiu et al. 2022; Schultner et al. 2023). Once caste determination has occurred, the genome is expressed differentially during the developmental trajectories of queens and workers (Abouheif 2021; Abouheif and Wray 2002; Barkdull and Moreau 2023; Béhague et al. 2018; Khila and Abouheif 2010; Qiu et al. 2022; Chandra et al. 2018; Vizueta et al. 2025). These trajectories are decoupled, and consequently, can evolve largely independently (Abouheif 2021; Powell et al. 2020; Vong et al. 2025; Li et al. 2024a). The dramatic variation in the number and presence / absence of ocelli in worker castes across the Formicinae (see Figure 1) supports the largely independent evolution of ocelli in workers from those in queens, which always develop 3 ocellus. Furthermore, we observe similar patterns of variation in the wings and ovaries between queen and worker castes across ant species (Cronin et al. 2013; Khila et al. 2012; Monnin and Peeters 2008; Rajakumar et al. 2012; 2018; Cronin et al. 2013) . Therefore, while the presence of ocelli in males and queens maintains the ocelli GRN in the genome and creates a potential for expression of this GRN in developing workers lacking ocelli, this cannot solely explain how it this latent expression became actualized and what maintained it over millions of years.

One hypothesis for how this ocelli GRN became latently expressed and has been maintained in workers lacking ocelli is pleiotropy, which potentially results from the multiple roles that genes within the ocelli GRN play within the same imaginal disc (the eye-antennal disc). For instance, *otd-1* and *hh* also determine the regional identity of the head capsule, while *hh*, *toy*, *eya* and *so*, also play key roles in compound eye development in specifying structures such as optic lobes, cone differentiation, and rhabdomeres development (Blanco et al. 2009; Domínguez-Cejudo and

Casares 2015; Jean-Guillaume and Kumar 2022). For example, the *sonic hedgehog* (*shh*) is a highly conserved developmental regulatory gene that plays a key role in limb development across animals. The vestigialization of hindlimbs in snakes and therefore the re-evolution of hindlimbs in extinct species across the phylogeny of the group, is thought be maintained during development through pleiotropy of enhancers that drive *shh* expression. This means that the same enhancer (ZRS) that drives *shh* expression in the external genital, also drives it (pleiotropically) in the developing limb region of snakes (Leal and Cohn 2018). Future studies should attempt to explore whether expression of these conserved genes in multiple regions of the eye-antennal disc is driven by shared enhancers. Alternatively, we cannot rule out the hypothesis that ocelli GRN has been co-opted to play a novel, yet currently unknown, function during worker larval development. Recent discoveries on the evolution of the wing GRN in ants provides support for this hypothesis. The wings, another nearly universal caste-specific trait in ants, develop in the reproductive male and queen caste, but are halted in the worker caste in response to environmental cues (Abouheif and Wray 2002). The wing GRN is also found to be latently expressed in the wingless worker caste of ants and was thought to be functionless (Abouheif and Wray 2002). However, it was recently discovered that the latent expression of this wing GRN in wingless worker caste acquired a novel function to generate big-headed soldiers in the hyperdiverse ant genus *Pheidole* (Rajakumar et al. 2018; 2012).

Finally, our inference that this latent expression of the ocelli GRN in workers facilitated the partial reversion to a single medial ocellus is supported by: (1) the close phylogenetic relationship between species that lack ocelli in adult workers but retain a latent expression of the ocelli GRN (*C. floridanus* and *P. rastellata*) , and those that underwent a partial phylogenetic reversion to a single medial ocellus (*C. gibbinotus* , *Dinomyrmex gigas* and *P. bihamata*); (2) the presence of a developmental capacity or potential of the latent ocelli GRN expression to produce rudimentary ocelli in the pupal stage of *C. floridanus* workers that completely lack ocelli as adults; (3) the ability to experimentally induce only 1, only 2, or all 3 ocellus in similarly sized adult workers that normally lack them by applying high doses of Juvenile Hormone (JH) to worker-destined larvae in the ant *Monomorium pharaonis* (Rajakumar et al. 2018; 2012); and finally (4) in nature, the rare induction of a single medial ocellus by mermithid parasites in soldiers of *Pheidole pallidula* that typically lack ocelli in natural colonies (Laciny et al. 2019; Passera 1976). The ability to naturally or experimentally induce worker individuals with only a single medial ocellus in different ant species also supports the inference that the single medial ocellus can be developmentally dissociated from the other two lateral ocelli. This suggests that reversion can facilitate the appearance of novel patterns of ocelli development in the workers if selected for (Abouheif et al. 2014; Gainett et al. 2024; Klingenberg 2014; West-Eberhard 2003). In *C. floridanus* and *P. rastellata*, there is a latent ocelli GRN expression for all 3 ocelli, providing a springboard to facilitate the partial phylogenetic reversion to a single medial ocellus in *C. gibbinotus*, *P. bihamata* and *D. gigas*. In the genus *Polyrhachis*, however, some species in the same subgenus as *P. bihamata*, such as *P. bellicosa*, have three ocelli. Because the

phylogenetic relationships within this subgenus have yet to be resolved, the independent reversion of ocelli in the ancestor of this subgenus may have resulted either in a single ocellus as reflected in *P. bihamata* or in 3 ocelli as reflected in *P. bellicosa* and 2 ocelli were subsequently lost giving rise to the single medial ocellus in *P. bihamata* (Hung 1967). These possibilities further reinforce the different evolutionary pathways by which this latent potential may facilitate novelty after reversion.

More broadly, our findings raise the possibility that the ancestral and latent GRN expression (also known as ancestral developmental potential) we observed may generally underlie polyphenic systems, including caste-specific traits across ants and other eusocial organisms. We therefore propose that ancestral developmental potentials may facilitate the loss and re-evolution of polyphenic traits within lineages (West-Eberhard 2003). When ancestral developmental potential facilitates only the partial re-evolution of alternative phenotypes, novel phenotypic patterns appear. Future work should not only mechanistically test these proposals in polyphenic systems at the colony or individual level but should also be tested at the cellular-level, where a single genome produces alternative cell-types in response to internal cues, such as morphogen gradients. If alternative cell types in polyphenic or non-polyphenic organisms retain a homolog or serial homolog of the trait (Lynch 2023; Jackman et al. 2025), then this raises the possibility that ancestral and latent developmental potentials may also facilitate gain, loss, re-evolution of alternative cell types in multicellular organisms.

Data and code availability: All data reported in this paper are provided in Table S2 in the supplemental information. This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Acknowledgments
We thank Hermogenes Fernandez-Marín, who inspired this work by sharing his knowledge about ocelli in soldiers across the *Atta* species. We thank Lloyd Davis, Marc Seid, and Shelly Berger for help with collecting *Camponotus floridanus* colonies. We thank Mary Jane West-Eberhard, Friedrich Markus, Guilherme Gainett, Arjuna Rajakumar, and Rajendhran Rajakumar for discussions and/or comments on the manuscript, and Juan Carlos Penagos for input on phylogenetic analysis. We thank Erik Plante and undergraduates for help with feeding and maintaining lab colonies of *Camponotus floridanus*. Finally, we thank McGill University's Integrated Quantitative Biology Initiative (IQBI) and Advanced Bioimaging Facility (ABIF) for imaging support. This work was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant to E.A and by a Doctoral fellowship from NSERC BESS-CREATE program to A.V-C.

Author Contribution
AV-C and EA conceived the project. AV-C and JA gathered ocelli data. AV-C and JA conducted ancestral state reconstruction. A V-C conducted HCR. TC performed SEM imaging. AV-C and EA wrote the manuscript.

Declaration of interests
The authors declare no competing interests.

Figures

Figure 1. The presence and absence of ocelli in queens and workers across ants. (A) Ocelli develop in the winged reproductive castes across species of formicinae ants exemplified by *Camponotus floridanus*, *Cataglyphis bicolor, Polyrachis bihamata* and *Dinomyrmex gigas* (white dashed circles). Ocelli in the worker caste of formicine ants (white dashed circles) are evolutionarily labile, showing no ocelli (*Camponotus floridanus*), all 3 ocelli in all individuals of the worker caste (*Cataglyphis bicolor*) or only 1 ocellus in only the soldiers (*Polyrachis bihamata* and *Dinomyrmex gigas*). (B) Schematic representation of the ocelli GRN in *Drosophila melanogaster* adapted from Jean-Guillaume & Kumar (2022)(Jean-Guillaume and Kumar 2022). The genes investigated in this study are highlighted in orange. Arrowheads indicate activation, and bars indicate repression. Queens are scaled to 1mm, and workers and soldiers are scaled according to the queen of each species. Asterisks indicate that *C. floridanus* was used for gene expression studies. Photos from Antweb (AntWeb 2024)

A.

	Queen	Worker	Soldier

Camponotus floridanus*

Cataglyphis bicolor

Polyrhachis bihamata

Dinomyrmex gigas

B.

Orthodenticle

↓

Notch —— *Defective proventriculus*

↓

Hedgehog

Twin of eyeless

Ci55   *Engrailed*

↓   ↓   ↓

Eyes absent   *Patched*   *Cubitus interruptus*

↓

Sine oculis

↓

Atonal

Figure 2. Ancestral state reconstruction of ocelli reveals three well-supported reversions of ocelli in the worker caste in the tribe Camponotini. Maximum clade credibility tree of formicine ants from Blaimer et al.(Blaimer et al. 2015). Ancestral state reconstruction for the presence (blue-colored circles) and absence (red-colored circles) of ocelli based on stochastic character mapping. Each pie chart for the nodes represents the posterior probabilities, scaled by the weight of evidence for each model. Species used to analyze ocelli GRN expression are highlighted in gray, and the species that re-evolved one ocellus is indicated as partial re-evolution. Three tribes within the Formicinae are marked (Melophorini, Plagiolepidini, Lasiini, Myrmelachistini and Camponotini) by arrows.

Figure 3. Characterizing development of the eye-antenna imaginal disc in worker castes of *C. floridanus* **using** *orthodenticle-1 (otd-1), distal-less (dll)* **and** *eyeless (ey)* **gene expression to mark the developing head capsule and ocelli, antenna, and eyes.** Fluorescent images in panels A, F, K, P, U represent the development of the entire eye-antenna imaginal disc marked with the nuclear stain DAPI across all four larval stages, where the head capsule region is labelled as 'Hc', the antennal region is labelled as 'An,' and the eye region is labelled as 'Eye.' Panels B, G, L, Q, V represent the development of the head capsule (Hc) marked by the genes *orthodenticle-1 (otd-1* in magenta); Panels (C,H,M,R,W) represent the antennal region marked by *distal-less* (*dll*) in green color; and Panels (D,I,N,S,X) represent the eyes (Eye) is *eyeless (ey)* (yellow); Panels. (A-E) First instar. Note: the green or yellow staining outside of the structures highlighted by the white arrows in panels C, D and E is background noise. (F-J) second instar, (K-O) third instar, (P-T) fourth instar worker-destined larvae and, (U-Y). Fourth instar soldier-destined larvae. All images are to scale.

Figure 4. Latent expression of *otd-1* and *hh* genes in the ocelli GRN in workers and soldiers of *C. floridanus* during the 4th larval instar. Expression of *orthodenticle-1(otd-1)* is yellow, and *hedgehog (hh)* is magenta. Early 4th instar; (A, B) males (C, D) soldiers and (E, F) workers. Late 4th instar: (A,' B') males (C', D') soldiers and (E', F') workers.

Early 4th larvae instar		Late 4th larvae instar	
otd-1	*hh*	*otd-1*	*hh*

Figure 5. Latent Expression of *toy*, *eya*, and *so* within the ocelli GRN in the developing worker and soldiers of *C. floridanus* Expression of *toy* (green), *eya* (yellow)*, and so* (magenta)*.* Early 4th instar*;*(A to C) males, (D to F) soldiers and (G to I) workers. Late 4th larvae stage*;*(A' to C') males, (D' to F'), soldiers and (G' to I').

Figure 5. Latent Expression of *toy*, *eya*, and *so* within the ocelli GRN in the developing worker and soldiers of *C. floridanus* Expression of *toy* (green), *eya* (yellow)*, and so* (magenta)*.* Early 4th instar*;*(A to C) males, (D to F) soldiers and (G to I) workers. Late 4th larvae stage*;*(A' to C') males, (D' to F'), soldiers and (G' to I').

Figure 6. Latent expression of the ocelli GRN in the eye-antennal disc in worker larvae of *Polyrachis rastellata* at 4th larvae stage. Expression of selected genes *otd-1* (yellow), *eya* (yellow), and *so* (magenta) at early 4th instar larvae.

Figure 7. Development of rudimentary ocelli in worker and soldier pupae of *C. floridanus*. (A) SEM showing ocelli development in males at mid-stages of pupal development. SEM showing development of rudimentary ocelli on (B to E) soldiers and (F to I) minor workers during early (day 6-7), mid (day 16-17), and late (19-20) pupal development. These ocelli rudiments disappear prior to adult stage (E and I).

Mid (day 16-17)

Male

| | Early (day 6-7) | Mid (day 16-17) | Late (day 19-20) | Molting pupae |

Soldier

Minor worker

Methods

Ant maintenance and collection

Colonies of *C. floridanus collected at Gainesville* (*Florida,* USA) and *P. rastellata* collected at Mae Tang (Chiang Mai, Thailand), were maintained in plastic boxes with glass test tubes filled with water-constrained cotton wool. They were fed mealworms and the Bhatkar–Whitcomb diet (Bhatkar and Whitcomb 1970). Colonies were maintained at 25°C with 60% humidity in complete darkness.

Larvae fixation and *in-situ* HCR

Gene sequences were obtained from NCBI GenBank database(Sayers et al. 2022) using genome BLAST against the assembled *C. floridanus* genome: *eyeless (ey*; XM_025414466), *distal-less (dll*; XM_025412727.1), *hedgehog, (hh*; XM_011262474.3), *eye absent (eya*; XM_025414466), *sine oculis (so;* XM_011252868.3) and *twin of eyeless (toy*; XM_011268499.3) genes. For *otd*, two paralogs of the gene were found in ants (XM_020028684.2 and XM_025415314.1), which is a result of a gene duplication event that has also been reported in wasps, bees, and beetles (Lynch et al. 2006; The Honeybee Genome Sequencing Consortium 2006; Zattara et al. 2017). The two *otd* paralogs sequences were then aligned by multiple sequence alignment using all of the known *orthodenticle* related sequences in insects; *Drosophila melanogaster* (NM_001369965.1), *Apis mellifera*: *otd-1* (XM_026446161.1), *otd-2*(XM_006571236.3), *Nasonia vitripennis*: *otd-1* (XM_008212114.4), *otd-2* (XM_031926951.2) and *Tribolium castaneum otd-1* (XM_008192467.2), *otd-2* (XM_008192470.2) and *Acythosiphon pisum* (XP_008180802.1). To determine the *otd-1* paralog to *Drosophila melanogaster* (*otd-1)* a maximum likelihood gene tree was inferred using genetic distance model Hasegawa Kishino Yano (HKY) and 500 bootstrap replicates as incorporated in MEGA12 alpha (Kumar et al. 2024)(Figure S1). Probes corresponding to all genes were chosen for the hybridization chain reaction experiments using the fluorescence Hairpins (B1 546, B2 488, B3 647) synthesized by Molecular Instruments.

First, second, third and fourth larval instars of soldier, and minor worker-destined larvae and fourth instar of male destined larvae of *C. floridanus* and worker larvae of *P. rastellata* were collected and subsequently fixed in a PEM 4% formaldehyde solution for 2 hrs at room temperature. Fixed samples were then dehydrated progressively in methanol baths (25%, 50%, 75% methanol for 15 min each, and 100% overnight at 4°C) and stored in 100% methanol at −30°C until use. All gene expression analyses were conducted by In situ Hybridization Chain Reaction (HCR), following the protocol for HCR (v3.0 protocol) (Schwarzkopf et al. 2021). After the tissue was pre-hybridized in a prewarmed Probe Hybridization Buffer (Molecular Instruments) for 30 minutes at 37°C and incubated with HCR probes in a Probe Hybridization Buffer overnight at 37°C. Tissues were washed the next day in a prewarmed Probe Wash Buffer four times, 15 minutes each and washed in 5X SSCT (UltraPure 20XSSC Buffer, Invitrogen, diluted in water) three times for 5 minutes at room temperature. Tissues were pre-amplified in

Amplification Buffer (Molecular Instruments) for 30 minutes at room temperature and incubated with snap-cooled HCR hairpins in Amplification Buffer overnight at room temperature. Tissues were then washed with 5X SSCT at room temperature twice for 5 minutes, for 30 minutes, and once for 5 minutes before being mounted on glycerol-DAPI 80%.

Microscopy
Confocal imaging was used to describe gene expression using Leica SP8 confocal microscope. Fiji (Schindelin et al. 2012) was used for image processing. Scanning electron microscopy (SEM) was done on a Hitachi TM3030 Scanning Electron Microscope.

Evolution of Ocelli in the Formicine Clade
The evolution of ocelli on workers across the subfamily Formicinae was inferred using ancestral reconstruction (ASE) for discrete traits incorporated in the R package Phytools 4.3.3(Revell 2024). The ASE analysis was based on the UCE70 phylogeny for the cade Formicinae published by Blaimer *et.al.* (Blaimer et al. 2015). The species *Camponotus floridanus* and *Polyrachis bihamata* were added manually to the phylogeny. To determine the presence or absence of ocelli in workers, photographs of the studied species from the database AntWeb,Version 8.112 were used (AntWeb 2024). The observations from the database were contrasted with published information from the literature (Table S2). Ocelli were classified as present in the worker caste if individuals exhibit any of the 3 ocellus (2 lateral and 1 medial ocellus). In the case of the presence of worker polymorphism, ocelli were classified as present if any one of the 3 ocelli was present within any of the worker subcastes. Whereas the absence was the complete lack of ocelli across workers and soldiers.

Four separate models of ocelli evolution were tested for each character in phytools: Equal rate "ER", all transitions rate different "ARD", and an irreversible model allowing only transitions between presence and absence, and another irreversible model allowing only transitions between absence and presence. We compared the fit of our models by computing Akaike information criterion (AIC) and Akaike weights and conducting pairwise likelihood ratio tests. The new function incorporated in phytools 4.3.3, *simmap,* was used to generate stochastic character maps under each of the four models tested (see Table S1). The stochastic mapping that resulted from the stochastic simulation represented the frequencies that are equal to the weight of evidence supported by each model (Revell 2024)

References

Abouheif, E. 2002. "Evolution of the Gene Network Underlying Wing Polyphenism in Ants." *Science* 297 (5579): 249–52. https://doi.org/10.1126/science.1071468.

Abouheif, Ehab. 2021. "Ant Caste Evo-Devo: It's Not All about Size." *Trends in Ecology & Evolution* 36 (8): 668–70. https://doi.org/10.1016/j.tree.2021.04.002.

Abouheif, Ehab, Marie-Julie Favé, Ana Sofia Ibarrarán-Viniegra, Maryna P. Lesoway, Ab Matteen Rafiqi, and Rajendhran Rajakumar. 2014. "Eco-Evo-Devo: The Time Has Come." In *Ecological Genomics*, edited by Christian R. Landry and Nadia Aubin-Horth, vol. 781. Advances in Experimental Medicine and Biology. Springer Netherlands. https://doi.org/10.1007/978-94-007-7347-9_6.

Abouheif, Ehab, and Gregory A. Wray. 2002. "Evolution of the Gene Network Underlying Wing Polyphenism in Ants." *Science* 297 (5579): 249–52. https://doi.org/10.1126/science.1071468.

AntWeb. 2024. Version Version 8.106.1. California Academy of Science, November 13. https://www.antweb.org.

Barkdull, Megan, and Corrie S Moreau. 2023. "Worker Reproduction and Caste Polymorphism Impact Genome Evolution and Social Genes Across the Ants." *Genome Biology and Evolution* 15 (6): evad095. https://doi.org/10.1093/gbe/evad095.

Béhague, Julien, Brian L. Fisher, Romain Péronnet, Rajendhran Rajakumar, Ehab Abouheif, and Mathieu Molet. 2018. "Lack of Interruption of the Gene Network Underlying Wing Polyphenism in an Early-branching Ant Genus." *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 330 (2): 109–17. https://doi.org/10.1002/jez.b.22794.

Bento, Gilberto, Akira Ogawa, and Ralf J. Sommer. 2010. "Co-Option of the Hormone-Signalling Module Dafachronic Acid–DAF-12 in Nematode Evolution." *Nature* 466 (7305): 494–97. https://doi.org/10.1038/nature09164.

Bhatkar, A., and W. H. Whitcomb. 1970. "Artificial Diet for Rearing Various Species of Ants." *The Florida Entomologist* 53 (4): 229–32. https://doi.org/10.2307/3493193.

Blaimer, Bonnie B., Seán G. Brady, Ted R. Schultz, Michael W. Lloyd, Brian L. Fisher, and Philip S. Ward. 2015. "Phylogenomic Methods Outperform Traditional Multi-Locus Approaches in Resolving Deep Evolutionary History: A Case Study of Formicine Ants." *BMC Evolutionary Biology* 15 (1): 271. https://doi.org/10.1186/s12862-015-0552-5.

Blanco, Jorge, Tobias Pauli, Makiko Seimiya, Gerald Udolph, and Walter J. Gehring. 2010. "Genetic Interactions of Eyes Absent, Twin of Eyeless and Orthodenticle Regulate Sine Oculis Expression during Ocellar Development in Drosophila." *Developmental Biology* 344 (2): 1088–99. https://doi.org/10.1016/j.ydbio.2010.05.494.

Blanco, Jorge, Makiko Seimiya, Tobias Pauli, Heinrich Reichert, and Walter J. Gehring. 2009. "Wingless and Hedgehog Signaling Pathways Regulate Orthodenticle and Eyes Absent during Ocelli Development in Drosophila." *Developmental Biology* 329 (1): 104–15. https://doi.org/10.1016/j.ydbio.2009.02.027.

Bose, Neelanjan, Akira Ogawa, Stephan H. von Reuss, et al. 2012. "Complex Small-Molecule Architectures Regulate Phenotypic Plasticity in a Nematode." *Angewandte Chemie International Edition* 51 (50): 12438–43. https://doi.org/10.1002/anie.201206797.

Brunet, Thibaut, and Nicole King. 2017. "The Origin of Animal Multicellularity and Cell Differentiation." *Developmental Cell* 43 (2): 124–40. https://doi.org/10.1016/j.devcel.2017.09.016.

Buschbeck, Elke, and Michael Bok, eds. 2023. *Distributed Vision: From Simple Sensors to Sophisticated Combination Eyes*. Springer Series in Vision Research. Springer International Publishing. https://doi.org/10.1007/978-3-031-23216-9.

Casasa, Sofia, Joseph F Biddle, Georgios D Koutsovoulos, and Erik J Ragsdale. 2021. "Polyphenism of a Novel Trait Integrated Rapidly Evolving Genes into Ancestrally Plastic Networks." *Molecular Biology and Evolution* 38 (2): 331–43. https://doi.org/10.1093/molbev/msaa235.

Chandra, Vikram, Ingrid Fetter-Pruneda, Peter R. Oxley, et al. 2018. "Social Regulation of Insulin Signaling and the Evolution of Eusociality in Ants." *Science* 361 (6400): 398–402.

Cronin, Adam L., Mathieu Molet, Claudie Doums, Thibaud Monnin, and Christian Peeters. 2013. "Recurrent Evolution of Dependent Colony Foundation Across Eusocial Insects." *Annual Review of Entomology* 58 (1): 37–55. https://doi.org/10.1146/annurev-ento-120811-153643.

Davidson, Phillip L, Erica M Nadolski, and Armin P Moczek. 2023a. "Gene Regulatory Networks Underlying the Development and Evolution of Plasticity in Horned Beetles." *Current Opinion in Insect Science* 60 (December): 101114. https://doi.org/10.1016/j.cois.2023.101114.

Davidson, Phillip L, Erica M Nadolski, and Armin P Moczek. 2023b. "Gene Regulatory Networks Underlying the Development and Evolution of Plasticity in Horned Beetles." *Current Opinion in Insect Science* 60 (December): 101114. https://doi.org/10.1016/j.cois.2023.101114.

Davison, Dinah R., and Richard E. Michod. 2021. "Phenotypic Plasticity and Evolutionary Transitions in Individuality." In *Phenotypic Plasticity & Evolution*, 1st ed., by David W. Pfennig. CRC Press. https://doi.org/10.1201/9780429343001-13.

Devlin, Dominic K., Austen R.D. Ganley, and Nobuto Takeuchi. 2023. "A Pan-Metazoan View of Germline-Soma Distinction Challenges Our Understanding of How the Metazoan

Germline Evolves." *Current Opinion in Systems Biology* 36 (December): 100486. https://doi.org/10.1016/j.coisb.2023.100486.

Domínguez-Cejudo, M. A., and F. Casares. 2015. "Antero-Posterior Patterning of *Drosophila* Ocelli Requires an Anti-Repressor Mechanism within the *Hh-* Pathway Mediated by the Six3 Gene *Optix*." *Development*, January 1, dev.125179. https://doi.org/10.1242/dev.125179.

Evans, Jay D., and Diana E. Wheeler. 2001. "Gene Expression and the Evolution of Insect Polyphenisms†." *BioEssays* 23 (1): 62–68. https://doi.org/10.1002/1521-1878(200101)23:1%253C62::AID-BIES1008%253E3.0.CO;2-7.

Fent, Karl, and Rüdiger Wehner. 1985. "Ocelli: A Celestial Compass in the Desert Ant Cataglyphis." *Science* 228 (4696): 192–94.

Gainett, Guilherme, Benjamin C. Klementz, Pola Blaszczyk, et al. 2024. "Vestigial Organs Alter Fossil Placements in an Ancient Group of Terrestrial Chelicerates." *Current Biology* 34 (6): 1258-1270.e5. https://doi.org/10.1016/j.cub.2024.02.011.

Hanna, Lisa, Brendon E. Boudinot, Jürgen Liebig, and Ehab Abouheif. 2024. *Inferring a Role for Programmed Cell Death during the Origin and Evolution of Wing Polyphenism in Ants*. Preprint. Evolutionary Biology. https://doi.org/10.1101/2024.02.14.580404.

Haynie, John L., and Peter J. Bryant. 1986. "Development of the Eye-Antenna Imaginal Disc and Morphogenesis of the Adult Head inDrosophila Melanogaster." *Journal of Experimental Zoology* 237 (3): 293–308. https://doi.org/10.1002/jez.1402370302.

Held, Lewis, ed. 2002. "References." In *Imaginal Discs: The Genetic and Cellular Logic of Pattern Formation*. Developmental and Cell Biology Series. Cambridge University Press. Cambridge Core. https://doi.org/10.1017/CBO9780511529733.018.

Hung, Akey C. F. 1967. "A Revision of the Ant Genus Polyrhachis at the Subgeneric Level (Hymenoptera: Formicidae)." *Transactions of the American Entomological Society (1890-)* 93 (4): 395–422. https://www.jstor.org/stable/25077942.

Jackman, William R., Vincent J. Lynch, and Yann Gibert. 2025. "Challenging and Redefining Dollo's Law of Evolution: Re-Appearance of Lost Structures." *BMC Ecology and Evolution* 25 (1): 62. https://doi.org/10.1186/s12862-025-02405-8.

Jean-Guillaume, Claude Bernard, and Justin P. Kumar. 2022. "Development of the Ocellar Visual System in Drosophila Melanogaster." *The FEBS Journal* 289 (23): 7411–27. https://doi.org/10.1111/febs.16468.

Jean-Guillaume, Claude Bernard, and Justin P. Kumar. 2022. "Development of the Ocellar Visual System in *Drosophila Melanogaster*." *The FEBS Journal* 289 (23): 7411–27. https://doi.org/10.1111/febs.16468.

Johnson, Robert A., and Ronald L. Rutowski. 2022. "Color, Activity Period, and Eye Structure in Four Lineages of Ants: Pale, Nocturnal Species Have Evolved Larger Eyes and Larger Facets than Their Dark, Diurnal Congeners." *PLOS ONE* 17 (9): e0257779. https://doi.org/10.1371/journal.pone.0257779.

Khila, A., E. Abouheif, and L. Rowe. 2012. "Function, Developmental Genetics, and Fitness Consequences of a Sexually Antagonistic Trait." *Science* 336 (6081): 585–89. https://doi.org/10.1126/science.1217258.

Khila, Abderrahman, and Ehab Abouheif. 2010. "Evaluating the Role of Reproductive Constraints in Ant Social Evolution." *Philosophical Transactions of the Royal Society B: Biological Sciences* 365 (1540): 617–30. https://doi.org/10.1098/rstb.2009.0257.

Klingenberg, Christian Peter. 2014. "Studying Morphological Integration and Modularity at Multiple Levels: Concepts and Analysis." *Philosophical Transactions of the Royal Society B: Biological Sciences* 369 (1649): 20130249. https://doi.org/10.1098/rstb.2013.0249.

Koch, Sophie, and Ehab Abouheif. 2019. "The Coordination of Insect Imaginal Discs and the Regulation and Evolution of Complex Worker Caste Systems of Ants." In *Deferring Development*, 1st ed., edited by Cory D. Bishop and Brian K. Hall. CRC Press. https://doi.org/10.1201/9780429445446-10.

Korb, J. 2025. "Changes of Division of Labour along the Eusociality Spectrum in Termites, with Comparisons to Multicellularity." *Philosophical Transactions of the Royal Society B: Biological Sciences* 380 (1922): 20230268. https://doi.org/10.1098/rstb.2023.0268.

Krapp, Holger G. 2009. "Ocelli." *Current Biology: CB* 19 (11): R435-437. https://doi.org/10.1016/j.cub.2009.03.034.

Kumar, Sudhir, Glen Stecher, Michael Suleski, Maxwell Sanderford, Sudip Sharma, and Koichiro Tamura. 2024. "MEGA12: Molecular Evolutionary Genetic Analysis Version 12 for Adaptive and Green Computing." *Molecular Biology and Evolution* 41 (12): msae263.

Laciny, A., H. L. Nemeschkal, H. Zettel, B. Metscher, and I. S. Druzhinina. 2019. "Caste-Specific Morphological Modularity in the Ant Tribe Camponotini (Hymenoptera, Formicidae)." *BMC Zoology* 4 (1): 9. https://doi.org/10.1186/s40850-019-0048-7.

Leal, Francisca, and Martin J. Cohn. 2018. "Developmental, Genetic, and Genomic Insights into the Evolutionary Loss of Limbs in Snakes." *Genesis* 56 (1): e23077. https://doi.org/10.1002/dvg.23077.

Lenuzzi, Maša, Hanh Witte, Metta Riebesell, Christian Rödelsperger, Ray L. Hong, and Ralf J. Sommer. 2023. "Influence of Environmental Temperature on Mouth-form Plasticity in *Pristionchus Pacificus* Acts through *Daf-11* -dependent cGMP Signaling." *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 340 (2): 214–24. https://doi.org/10.1002/jez.b.23094.

Li, Ruyan, Xueqin Dai, Jixuan Zheng, et al. 2024a. "Juvenile Hormone as a Key Regulator for Asymmetric Caste Differentiation in Ants." *Proceedings of the National Academy of Sciences* 121 (46): e2406999121. https://doi.org/10.1073/pnas.2406999121.

Li, Ruyan, Xueqin Dai, Jixuan Zheng, et al. 2024b. "Juvenile Hormone as a Key Regulator for Asymmetric Caste Differentiation in Ants." *Proceedings of the National Academy of Sciences* 121 (46): e2406999121. https://doi.org/10.1073/pnas.2406999121.

Lynch, Jeremy A., Ava E. Brent, David S. Leaf, Mary Anne Pultz, and Claude Desplan. 2006. "Localized Maternal Orthodenticle Patterns Anterior and Posterior in the Long Germ Wasp Nasonia." *Nature* 439 (7077): 728–32. https://doi.org/10.1038/nature04445.

Lynch, Vincent J. 2023. "Is There a Loophole in Dollo's Law? A DevoEvo Perspective on Irreversibility (of Felid Dentition)." *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 340 (8): 509–17. https://doi.org/10.1002/jez.b.23163.

MacMillan, Olivia, Julia Singer, Sophia Perrakis, et al. 2025. *Juvenile Hormone Signalling Underlies the Switchpoint and Differentiation of Soldiers in Camponotus Floridanus*.

Mishra, Abhishek Kumar, Cornelia Fritsch, Roumen Voutev, Richard S. Mann, and Simon G. Sprecher. 2021. "Homothorax Controls a Binary Rhodopsin Switch in Drosophila Ocelli." *PLOS Genetics* 17 (7): e1009460. https://doi.org/10.1371/journal.pgen.1009460.

Miura, Toru. 2005. "Developmental Regulation of Caste-Specific Characters in Social-Insect Polyphenism." *Evolution & Development* 7 (2): 122–29. https://doi.org/10.1111/j.1525-142X.2005.05014.x.

Monnin, Thibaud, and Christian Peeters. 2008. "How Many Gamergates Is an Ant Queen Worth?" *Naturwissenschaften* 95 (2): 109–16. https://doi.org/10.1007/s00114-007-0297-0.

Moser, John C., John D. Reeve, José Maurício S. Bento, Terezínha M. C. Della Lucia, R. Scott Cameron, and Natalie M. Heck. 2004. "Eye Size and Behaviour of Day- and Night-Flying Leafcutting Ant Alates." *Journal of Zoology* 264 (1): 69–75. https://doi.org/10.1017/S0952836904005527.

Narendra, Ajay, Fiorella Ramirez-Esquivel, and Willi A. Ribi. 2016. "Compound Eye and Ocellar Structure for Walking and Flying Modes of Locomotion in the Australian Ant, Camponotus Consobrinus." *Scientific Reports* 6 (1): 22331. https://doi.org/10.1038/srep22331.

Narendra, Ajay, and Willi A. Ribi. 2017. "Ocellar Structure Is Driven by the Mode of Locomotion and Activity Time in *Myrmecia* Ants." *Journal of Experimental Biology* 220 (23): 4383–90. https://doi.org/10.1242/jeb.159392.

Passera, L. 1976. "Origine des intercastes dans les sociétés dePheidole pallidula (Nyl.) (Hymenoptera Formicidae) parasitées parMermis sp. (Nematoda Mermithidae)." *Insectes Sociaux* 23 (4): 559–75. https://doi.org/10.1007/BF02223482.

Peters, Ralph S., Lars Krogmann, Christoph Mayer, et al. 2017. "Evolutionary History of the Hymenoptera." *Current Biology* 27 (7): 1013–18. https://doi.org/10.1016/j.cub.2017.01.027.

Powell, Scott, Shauna L. Price, and Daniel J. C. Kronauer. 2020. "Trait Evolution Is Reversible, Repeatable, and Decoupled in the Soldier Caste of Turtle Ants." *Proceedings of the National Academy of Sciences* 117 (12): 6608–15. https://doi.org/10.1073/pnas.1913750117.

Qiu, Bitao, Xueqin Dai, Panyi Li, et al. 2022. "Canalized Gene Expression during Development Mediates Caste Differentiation in Ants." *Nature Ecology & Evolution* 6 (11): 11. https://doi.org/10.1038/s41559-022-01884-y.

Ragsdale, Erik J., Manuela R. Müller, Christian Rödelsperger, and Ralf J. Sommer. 2013. "A Developmental Switch Coupled to the Evolution of Plasticity Acts through a Sulfatase." *Cell* 155 (4): 922–33. https://doi.org/10.1016/j.cell.2013.09.054.

Rajakumar, Arjuna, Luigi Pontieri, Ruyan Li, et al. 2024. "From Egg to Adult: A Developmental Table of the Ant Monomorium Pharaonis." *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 342 (8): 557–85. https://doi.org/10.1002/jez.b.23278.

Rajakumar, Rajendhran, Sophie Koch, Mélanie Couture, et al. 2018. "Social Regulation of a Rudimentary Organ Generates Complex Worker-Caste Systems in Ants." *Nature* 562 (7728): 574–77. https://doi.org/10.1038/s41586-018-0613-1.

Rajakumar, Rajendhran, Diego San Mauro, Michiel B. Dijkstra, et al. 2012. "Ancestral Developmental Potential Facilitates Parallel Evolution in Ants." *Science* 335 (6064): 79–82. https://doi.org/10.1126/science.1211451.

Revell, Liam J. 2012. "Phytools: An R Package for Phylogenetic Comparative Biology (and Other Things)." *Methods in Ecology and Evolution* 3 (2): 217–23. https://doi.org/10.1111/j.2041-210X.2011.00169.x.

Revell, Liam J. 2024. "Phytools 2.0: An Updated R Ecosystem for Phylogenetic Comparative Methods (and Other Things)." *PeerJ* 12 (January): e16505. https://doi.org/10.7717/peerj.16505.

Sabat, Debabrat, Subhashree Priyadarsini, and Monalisa Mishra. 2017. "Understanding the Structural and Developmental Aspect of Simple Eye of Drosophila: The Ocelli." *Journal of Cell Signaling* 01 (02). https://doi.org/10.4172/2576-1471.1000109.

Sayers, Eric W, Evan E Bolton, J Rodney Brister, et al. 2022. "Database Resources of the National Center for Biotechnology Information." *Nucleic Acids Research* 50 (D1): D20–26. https://doi.org/10.1093/nar/gkab1112.

Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, et al. 2012. "Fiji: An Open-Source Platform for Biological-Image Analysis." *Nature Methods* 9 (7): 676–82. https://doi.org/10.1038/nmeth.2019.

Schultner, Eva, Tobias Wallner, Benjamin Dofka, et al. 2023. "Queens Control Caste Allocation in the Ant *Cardiocondyla Obscurior*." *Proceedings of the Royal Society B: Biological Sciences* 290 (1992): 20221784. https://doi.org/10.1098/rspb.2022.1784.

Schwarz, Sebastian, Antoine Wystrach, and Ken Cheng. 2011. "A New Navigational Mechanism Mediated by Ant Ocelli." *Biology Letters* 7 (6): 856–58. https://doi.org/10.1098/rsbl.2011.0489.

Schwarzkopf, Maayan, Mike C. Liu, Samuel J. Schulte, et al. 2021. "Hybridization Chain Reaction Enables a Unified Approach to Multiplexed, Quantitative, High-Resolution Immunohistochemistry and *in Situ* Hybridization." *Development* 148 (22): dev199847. https://doi.org/10.1242/dev.199847.

Sommer, Ralf J. 2020. "Phenotypic Plasticity: From Theory and Genetics to Current and Future Challenges." *Genetics* 215 (1): 1–13. https://doi.org/10.1534/genetics.120.303163.

Susoy, Vladislav, Erik J Ragsdale, Natsumi Kanzaki, and Ralf J Sommer. 2015. "Rapid Diversification Associated with a Macroevolutionary Pulse of Developmental Plasticity." *eLife* 4 (February): e05463. https://doi.org/10.7554/eLife.05463.

The Honeybee Genome Sequencing Consortium. 2006. "Insights into Social Insects from the Genome of the Honeybee Apis Mellifera." *Nature* 443 (7114): 931–49. https://doi.org/10.1038/nature05260.

Visser, Bertanne, Hans T. Alborn, Suzon Rondeaux, et al. 2021. "Phenotypic Plasticity Explains Apparent Reverse Evolution of Fat Synthesis in Parasitic Wasps." *Scientific Reports* 11 (1): 7751. https://doi.org/10.1038/s41598-021-86736-8.

Visser, Bertanne, Cécile Le Lann, Frank J. den Blanken, Jeffrey A. Harvey, Jacques J. M. van Alphen, and Jacintha Ellers. 2010. "Loss of Lipid Synthesis as an Evolutionary Consequence of a Parasitic Lifestyle." *Proceedings of the National Academy of Sciences of the United States of America* 107 (19): 8677–82. https://doi.org/10.1073/pnas.1001744107.

Vizueta, Joel, Zijun Xiong, Guo Ding, et al. 2025. "Adaptive Radiation and Social Evolution of the Ants." *Cell* 188 (18): 4828-4848.e25. https://doi.org/10.1016/j.cell.2025.05.030.

Vong, Erica, Shannon Parisien, Helene Orfali, and Rajendhran Rajakumar. 2025. "Evolution and Regulation of Decoupled Dimorphic Scaling Patterns in Ants." *bioRxiv*.

Ward, Philip S. 2014. "The Phylogeny and Evolution of Ants." *Annual Review of Ecology, Evolution, and Systematics* 45 (1): 23–43. https://doi.org/10.1146/annurev-ecolsys-120213-091824.

Werner, Michael S., Marc H. Claaßen, Tess Renahan, Mohannad Dardiry, and Ralf J. Sommer. 2018. "Adult Influence on Juvenile Phenotypes by Stage-Specific Pheromone Production." *iScience* 10 (December): 123–34. https://doi.org/10.1016/j.isci.2018.11.027.

West-Eberhard, Mary Jane. 2003. *Developmental Plasticity and Evolution*. Oxford University Press.

Yorimitsu, Takeshi, Naruto Kiritooshi, and Hideki Nakagoshi. 2011. "Defective Proventriculus Specifies the Ocellar Region in the Drosophila Head." *Developmental Biology* 356 (2): 598–607. https://doi.org/10.1016/j.ydbio.2011.06.015.

Zattara, Eduardo E., Anna L. M. Macagno, Hannah A. Busey, and Armin P. Moczek. 2017. "Development of Functional Ectopic Compound Eyes in Scarabaeid Beetles by Knockdown of *Orthodenticle*." *Proceedings of the National Academy of Sciences* 114 (45): 12021–26. https://doi.org/10.1073/pnas.1714895114.

Zimmerman, J E, Q T Bui, H Liu, and N M Bonini. 2000. "Molecular Genetic Analysis of Drosophila Eyes Absent Mutants Reveals an Eye Enhancer Element." *Genetics* 154 (1): 237–46. https://doi.org/10.1093/genetics/154.1.237.

Supplementary Figures

Supplementary Figure 1. Simplified Gene tree based on maximum likelihood showing relationships between *orthodenticle (otd)* orthologs in *Drosophila melanogaster*, *Apis mellifera*, *Nasonia vitripennis*, *Tribolium castaneum, Acythosiphon pisum and Camponotus floridanus.* Branch values are bootstrap support (%). Colors: *otd-2* (red), *otd-1* (blue)

Supplementary Tables.

Supplementary Table1. Model selection under maximum likelihood estimation implemented in phytools (Revell 2012). The results are ordered by decreasing Akaike Weights (w).

Model	log(L)	d.f.	AIC	weight
Equal Rates	-41.808109	1	85.6162173	0.70944317
Unequal Rates	-41.720603	2	87.441205	0.28485677
Irreversible: Absence to Presence	-46.813104	1	95.6262081	0.00475637
Irreversible: Presence to Absence	-48.430553	1	98.8611065	0.00094368

Supplementary Table 2. Database with the references used in this study for presence and number of ocellus (1) and absence (0) of ocellus.

Subfamily	Genus	Species	Ocelli Count	References
outgroup	*Acanthoponera*	*minor*	0	AntWeb.org
Formicinae	*Acropyga*	*acutiventris*	0	AntWeb.org
outgroup	*Aneuretus*	*simoni*	0	AntWeb.org
Formicinae	*Anoplolepis*	*custodiens*	0	AntWeb.org
Formicinae	*Anoplolepis*	*gracilipes*	0	AntWeb.org
Formicinae	*Aphomomyrmex*	*afer*	3	AntWeb.org
Formicinae	*Bajcaridris*	*theryi*	3	Santschi, 1936
outgroup	*Brachymyrmex*	*depilis*	0	AntWeb.org
Formicinae	*Calomyrmex*	*albertisi*	0	AntWeb.org
Formicinae	*Calomyrmex*	*laevissimus*	0	AntWeb.org
Formicinae	*Camponotus*	*floridanus*	0	AntWeb.com
Formicinae	*Camponotus*	*gibbinotus*	1	AntWeb.org
Formicinae	*Camponotus*	*hyatti*	0	MacKay & Mackay, 2002
Formicinae	*Camponotus*	*maritimus*	0	Ward, 2005
Formicinae	*Colobopsis*	*saundersi*	0	AntWeb.org
Formicinae	*Colobopsis*	*vitiensis*	0	Mann, 1920
Formicinae	*Cataglyphis*	*cursor*	3	AntWeb.org
outgroup	*Cladomyrma*	*petalae*	0	Agosti, 1991
outgroup	*Dolichoderus*	*pustulatus*	0	AntWeb.org
Formicinae	*Dinomyrmex*	*gigas*	1	AntWeb.org
Formicinae	*Echinopla*	*australis*	0	AntWeb.org
Formicinae	*Euprenolepis*	*procera*	0	Lapolla, 2009
Formicinae	*Formica*	*moki*	3	Cole Jr, 1943
Formicinae	*Formica*	*neogagates*	3	AntWeb.org
Formicinae	*Gigantiops*	*destructor*	3	Smith, 1858
Formicinae	*Iberoformica*	*subrufa*	3	Antwiki- genus
Formicinae	*Lasiophanes*	*atriventris*	3	AntWeb.org
Formicinae	*Lasius*	*californicus*	0	AntWeb.org
Formicinae	*Lasius*	*niger*	3	AntWeb.org
Formicinae	*Lepisiota*	*canescens*	3	Sharaf et al., 2020
outgroup	*Manica*	*bradleyi*	0	AntWeb.org
outgroup	*Myrmecia*	*pyriformis*	3	AntWeb.org
Formicinae	*Myrmecocystus*	*flaviceps*	3	AntWeb.org

			3 (soldiers and media, absent in minors)	
Formicinae	*Myrmecorhynchus*	*emeryi*	3 (*soldiers and media, absent in minors*)	Wheeler, 1917
outgroup	*Myrmelachista*	*flavocotea*	0	AntWeb.org
Formicinae	*Myrmoteras*	*iriodum*	3	AntWeb.org
outgroup	*Nothomyrmecia*	*macrops*	0	AntWeb.org
Formicinae	*Notoncus*	*capitatus*	3	AntWeb.org
Formicinae	*Notostigma*	*carazzii*	3	AntWeb.org; Emery, 1920
Formicinae	*Nylanderia*	*dodo*	0	Lapolla et al., 2011
Formicinae	*Nylanderia*	*hystrix*	3	Kallal & LaPolla, 2012
Formicinae	*Oecophylla*	*longinoda*	0	AntWeb.org
Formicinae	*Oecophylla*	*smaragdina*	0	Cole & Jones, 1948
Formicinae	*Opisthopsis*	*respiciens*	0	AntWeb.org
Formicinae	*Paraparatrechina*	*glabra*	3	AntWeb.org
Formicinae	*Paraparatrechina*	*oceanica*	0	AntWeb.org
Formicinae	*Paratrechina*	*antsingy*	3	LaPolla & Fisher, 2014
Formicinae	*Paratrechina*	*longicornis*	3	AntWeb.org
Formicinae	*Paratrechina*	*zanjensis*	3	LaPolla et al., 2013
Formicinae	*Petalomyrmex*	*phylax*	3	Snelling, 1979
Formicinae	*Plagiolepis*	*alluaudi*	0	AntWeb.org
Formicinae	*Polyergus*	*breviceps*	3	AntWeb.org; Smith, 1947
Formicinae	*Polyrhachis*	*decumbens*	0	Kohout, 2006
Formicinae	*Prenolepis*	*emmae*	3	AntWeb.org
Formicinae	*Prenolepis*	*imparis*	0	Williams & LaPolla, 2016
Formicinae	*Proformica*	*mongolica*	3	AntWeb.org
Formicinae	*Prolasius*	*convexus*	3	McAreavey, 1947
Formicinae	*Pseudolasius*	*australis*	0	Emery, 1925
Formicinae	*Pseudonotoncus*	*hirsutus*	3	Shattuck & O'Reilly, 2013
outgroup	*Rhytidoponera*	*chalybaea*	0	AntWeb.org
Formicinae	*Rossomyrmex*	*anatolicus*	3	AntWeb.org
Formicinae	*Santschiella*	*kohli*	3	Forel, 1916
Formicinae	*Teratomyrmex*	*greavesi*	3	Shattuck & O'Reilly, 2013
outgroup	*Tetraponera*	*rufonigra*	3	Ward, 2001
Formicinae	*Zatania*	*albimaculata*	3	AntWeb.org

Table S3 References

Agosti, D. (1991). Revision of the oriental ant genus *Cladomyrma* , with an outline of the higher
classification of the Formicinae (Hymenoptera: Formicidae). *Systematic Entomology*,
16(3), 293–310. https://doi.org/10.1111/j.1365-3113.1991.tb00690.

AntWeb.Version 8.112. California Academy of Science, online at https://www.antweb.org.
Accessed 13 November 2024.

Cole, A. C., & Jones, J. W. (1948). A Study of the Weaver Ant, Oecophylla smaragdina (Fab.)1.
The American Midland Naturalist, *39*(3), 641–651. https://doi.org/10.2307/2421529

Cole Jr, A. (1943). A New Subspecies of Formica moki Wheeler (Hymenoptera: Formicidae).
THE AMERICAN MIDLAND) NATURALIST, *29*(1), 183–184.

Emery, C. (1920). *Le genre Camponotus Mayr. Nouvel essai de sa subdivision en sous-genres.*

Emery, C. (1925). Hymenoptera, Fam. Formicidae, subfam. Formicinae. *Genera Insectorum*,
183.

Kallal, R. J., & Lapolla J. S. (2012). Monograph of Nylanderia (Hymenoptera: Formicidae) of
the World, Part II: Nylanderia in the Nearctic. *Zootaxa*, *3508*(1).
https://doi.org/10.11646/zootaxa.3508.1.1

Kohout, R. J. (2006). Review of Polyrhachis (Cyrtomyrma) Forel (Hymenoptera: Formicidae:
Formicinae) of Australia, Borneo, New Guinea and the Solomon Islands with
descriptions of new species. *Memoirs-Queensland Museum*, *52*(1), 87.

LaPolla, J., Hawkes, P., & Fisher, J. N. (2013). Taxonomic review of the ant genus Paratrechina,
with a description of a new species from Africa. *Journal of Hymenoptera Research*, *35*,
71–82. https://doi.org/10.3897/jhr.35.5628

Lapolla, J. S. (2009). Taxonomic Revision of the Southeast Asian Ant Genus Euprenolepis.
Zootaxa, *2046*(1). https://doi.org/10.11646/zootaxa.2046.1.1

Lapolla, J. S., Brady, S. G., & Shattuck, S. O. (2011). Monograph of Nylanderia (Hymenoptera:
Formicidae) of the World: An introduction to the systematics and biology of the genus.
Zootaxa, *3110*(1). https://doi.org/10.11646/zootaxa.3110.1.1

LaPolla, J. S., & Fisher, B. L. (2014). Then there were five: A reexamination of the ant genus
Paratrechina (Hymenoptera, Formicidae). *ZooKeys*, *422*, 35.
https://doi.org/10.3897/zookeys.422.7779

MacKay, W. P., & Mackay, E. (2002). *The ants of New Mexico (Hymenoptera: Formicidae)*.
Edwin Mellen Press Lewiston, NY.

Mann, W. M. (1920). Ant guests from Fiji and the British Solomon islands. *Annals of the Entomological Society of America*, *13*(1), 60–69.

McAreavey, J. (1957). Revision of the genus Stigmacros Forel. *Memoirs of the National Museum of Victoria*, *21*, 7–64.

Santschi, F. (1936). Liste et descriptions de fourmis du Maroc. *Bulletin de La Société Des Sciences Naturelles Du Maroc*, *16*(3), 198–210.

Sharaf, M. R., Aldawood, A. S., Mohamed, A. A., & Hita Garcia, F. (2020). The genus Lepisiota Santschi, 1926 of the Arabian Peninsula with the description of a new species, Lepisiota elbazi sp. Nov. From Oman, an updated species identification key, and assessment of zoogeographic affinities. *Journal of Hymenoptera Research*, *76*, 127–152. https://doi.org/10.3897/jhr.76.50193

Shattuck, S., & O'Reilly, A. (2013). Revision of the Australian endemic ant genera Pseudonotoncus and Teratomyrmex (Hymenoptera: Formicidae: Formicinae). *Zootaxa*, *3669*(3), 287–301.

Smith, F. (1858). *Catalogue of Hymenopterous Insects in the Collection of the British Museum: 6: Formicidae* (Vol. 6). order of the Trustees.

Smith, M. R. (1947). A Study of Polyergus in the United States, Based on the Workers (Hymenoptera: Formicidae). *The American Midland Naturalist*, *38*(1), 150–161. https://doi.org/10.2307/2421633

Snelling, R. R. (1979). *Aphomomyrmex and a related new genus of arboreal African ants (Hymenoptera: Formicidae)*.

Ward, P. S. (2001). Taxonomy, phylogeny and biogeography of the ant genus Tetraponera (Hymenoptera: Formicidae) in the Oriental and Australian regions. *Invertebrate Systematics*, *15*(5), 589–665.

Ward, P. S. (2005). A synoptic review of the ants of California (Hymenoptera: Formicidae). *Zootaxa*, *936*(1). https://doi.org/10.11646/zootaxa.936.1.1

Wheeler, W. (1917). The Australian ant-genus Myrmecorhynchus (Ern. André) and its position in the subfamily Camponotinae. *Transactions of the Royal Society of South Australia*, *41*, 14–19.

Williams, J. L., & Lapolla, J. S. (2016). Taxonomic revision and phylogeny of the ant genus Prenolepis (Hymenoptera: Formicidae). *Zootaxa*, *4200*(2).