Proteogenomic profiling of soft tissue leiomyosarcoma reveals distinct molecular subtypes with divergent outcomes and therapeutic vulnerabilities

Atsushi Tanaka[1,2,3]*, Makiko Ogawa[1,3], Yusuke Otani[1], Ronald C. Hendrickson[4,5,6], Zhuoning Li[7], Narasimhan P. Agaram[8], David S. Klimstra[3,9], Julia Y. Wang[10], and Michael H. A. Roehrl[1,3,11,12]*

[1]Department of Pathology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

[2]Broad Institute of MIT and Harvard, Cambridge, MA, USA

[3]Former address: Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[4]Department of Biochemistry and Molecular Biology, Miller School of Medicine, University of Miami, Miami, FL, USA

[5]Sylvester Comprehensive Cancer Center, Miller School of Medicine, University of Miami, Miami, FL, USA

[6]Former address: Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[7]Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[8]Department of Pathology and Laboratory Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[9]Department of Pathology, Yale University School of Medicine, New Haven, CT, USA

[10]Curandis, Boston, MA, USA

[11]Wyss Institute at Harvard, Boston, MA, USA

[12]Lead contact

*Correspondence to: atanaka@bidmc.harvard.edu, michael_roehrl@bidmc.harvard.edu

Running title: Proteogenomics of leimyosarcoma

Summary

Soft tissue leiomyosarcoma (STLMS) is an aggressive malignancy lacking validated molecular subclassification and effective targeted treatments. We performed comprehensive proteogenomic analysis of primary and metastatic STLMS to uncover biological traits and therapeutic weaknesses. Integrative proteomic and phosphoproteomic analyses using non-negative matrix factorization identified three subtypes. Subtype P1 shows genomic stability, low proliferation, and enrichment of FGFR2 and PDK signaling pathways. Subtype P2 exhibits chromosomal instability, inflammatory programs, activation of CDK-AURKA/B-mTOR/ERK kinome with *IGF1R*/*PDGFRA* gene alterations, and poorest survival outcomes. Subtype P3 is highly proliferative, with E2F/DNA-repair programs, elevated NCOR1, and shift towards non-homologous end joining with upregulation of PARP1. Homologous recombination deficiency (HRD) analysis distinguishes HRD-low P1 from HRD-high P2/P3. Paired analyses suggest HRD increases in metastases within P3. Immune profiling shows P2 as immunosuppressive, characterized by LGALS9 and M2 macrophages. Our proteogenomic analyses provide a molecular landscape of LMS, revealing biological insights, patient outcome stratification, and therapeutic targets.

Introduction

Soft tissue sarcomas (STS) are a diverse group of mesenchymal malignancies that include over 80 histological subtypes[1]. Leiomyosarcoma (LMS) accounts for 10-20% of all STS[2]. Soft tissue leiomyosarcoma (STLMS), i.e., extra-uterine LMS, develops mainly in the retroperitoneum, abdomen/pelvis, trunk, and extremities, with the first two being more common[3]. Clinical management of localized disease depends on anatomical site and tumor grade[4,5]. Typical STLMS shows spindle-shaped cells that are set in long intersecting fascicles, diffuse hypercellularity with high-grade histology, and the expression of one or more myogenic marker proteins. Despite multidisciplinary management, cure rates for STLMS remain low. Half of patients experience tumor relapse after surgery[6-8]. Patients with advanced or metastatic STLMS have poor outcomes and limited treatment options[9,10]. To improve outcomes, there are urgent needs to develop better classification and prognostic stratification of STLMS patients for the development of efficient therapeutics.

Several genomic and epigenomic studies of pan-STS reported enrichment of PIK3/AKT pathway alterations and deletions of tumor suppressor genes, including *TP53* (9% deep and 60% shallow deletions), *RB1* (14% deep, 78% shallow), and *PTEN* (13% deep, 68% shallow)[11-13]. Multiomic analyses showed that LMS is molecularly distinct from other STS, and that STLMS differs from uterine LMS in hormone response and DNA damage response pathways. Integrated clustering of STLMS based on WES, epigenome, and whole transcriptome data revealed two subtypes: one with short survival and frequent PI3K/AKT/MTOR pathway alterations, the other with longer survival and minimal PI3K/AKT/MTOR alterations. While MTOR inhibitors like everolimus or temsirolimus show some clinical efficacy in LMS[14,15], their efficacy is reduced by indirect AKT upregulation.

Proteins are the "machines of life" carrying out complex cellular and biochemical functions and thus serve as cancer treatment targets[16]. mRNA abundance poorly predicts protein amounts[17,18]. Proteomics complements genomic research by providing accurate assessment of tumor biology and patient stratification that is inaccessible to nucleic acid-based approaches. Two proteomic analyses of pan-STS were recently published, although the use of formalin-fixed, paraffin-embedded samples limits proteomic studies and largely precludes analyses of post-translational modifications, such as phosphorylation[19,20]. Here, we present the first integrative proteogenomic study of STLMS using freshly frozen tissue and the inclusion of phosphoproteomics to discover alterations of the kinome.

To gain a better understanding of STLMS tumor biology with focus on associations between genomic alterations and both total proteome and phosphoproteome, we analyzed 72

80 primary site and metastatic STLMS samples, including 18 paired primary-metastasis lesions.
81 Additionally, we incorporated our own independent validation cohort of 222 STLMS FFPE
82 samples as tissue microarrays.
83
Results

Unsupervised clustering of proteome identifies 3 proteomic subtypes with clinical
relevance
We obtained 72 frozen tissues of primary and metastatic STLMS (cohort 1) for liquid
chromatography-mass spectrometry (LC-MS) analysis (Figure 1A). For independent cohort 2,
we obtained FFPE blocks of 222 STLMS cases and constructed tissue microarray blocks (Figure
1A). Patient demographics are summarized in Table S1. LC-MS analysis of global proteomes
quantified 10,249 proteins in total and 7,325 proteins in at least 30% of samples (Table S2A).
For phosphoproteomics, the analysis quantified 6,542 phosphorylated proteins in total and 4,584
in at least 30% of samples (Figure S1A, Table S2B). Batch correction removed TMT multiplexing
effects from the data (Figure S1B). Using non-negative matrix factorization (NMF) on filtered
global proteome and phosphoproteome data, we identified three distinct proteome-based
subtypes (P1, P2, and P3) of STLMS (Figures 1BC, S1CD). Survival analyses showed significant
differences in overall survival time (OS) and recurrence-free survival time (RFS) between
subtypes, with P2 showing the shortest patient survival. We validated these findings using a
TCGA STLMS dataset, where P2/P3 subtypes showed significantly shorter recurrence-free
survival and overlapped with the TCGA STLMS iCluster1 subtype (Fig. S1EF). Analysis of clinical
attributes (Figure 1D) revealed no significant differences in tumor size, age, tumor status (primary
lesion or metastatic lesion), gender, TNM stage, or prior treatment between proteomic subtypes.
However, metastatic samples were enriched in P2/P3 subtypes compared to P1 (p=0.0308).
Primary tumor site distribution varied significantly among subtypes (p=0.01735), with trunk
samples predominantly in P1 and pelvis samples in P2. FNCLCC Grade 1 tumors were only seen
in P1, while FNCLCC Grade 3 tumors were only seen in P3.
107

Differences of proteogenomic landscape and transcription regulatory networks between
proteome-based subtypes
We evaluated genomic alterations in 69 samples by MSK-IMPACT targeted sequencing. P2/3
subtypes showed higher ploidy, whole genome duplication, and chromosomal instability versus
P1 (Figure 2AB, findings supported by analysis of the TCGA STLMS cohort (Figure S2AB).
Genomically, *TP53* (68%) and *RB1* (41%) were most frequently altered, followed by *NCOR1*

(35%), *MAP2K4* (33%), and *FLCN* (28%) (Figure 2C). The GENIE STLMS cohort (v15.0,

n=774)[21] showed similar frequencies. P2/P3 showed frequent genomic alterations in *NCOR1*,

MAP2K4, *FLCN*, and *IGF1R*. P2 showed commonly alterations in oncogenic drivers like *IGF1R*

or *PDGFRA* and SWI/SNIF complex genes, associated with poor outcomes[22-25]. Somatic copy

number alteration (SCNA) analysis revealed more arm-level amplifications in P2/P3 than in P1,

while P3 showed higher deletion frequencies (Figure 2DE). Each subtype showed unique focal

amplification/deletion patterns, with P1 showing fewer focal peaks (Figure 2F). The 17p12 peak,

containing *NCOR1* and *MAP2K4*, was shared between P2 and P3. *NCOR1* and *MAP2K4*

alterations occurred frequently together (p=1.03E-14, Fisher's exact test). We analyzed gene

overlap in recurrent focal peaks, finding 1,818, 1,125, and 3,556 involved genes respectively

(Figure 2G). Gene ontology analyses showed P1 focal deletion peaks enriched in copper/zinc

ion processes, P2 focal peaks in extracellular matrix processes, and P3 focal peaks in T-cell

receptor signaling (Figure S2C). KEGG analysis showed P2/P3 deletion peaks enriched in

immune-related pathways, while only P3 showed deletions involving in PI3K-Akt signaling

(Figure S2D).

Pairwise differential expression analyses between subtypes (P1 vs. P2/P3, P2 vs. P1/P3,

P3 vs. P1/P2) revealed 2,277 dysregulated proteins in P1 vs. others (207 upregulated >2-fold,

211 downregulated >2-fold), 2,503 in P2 vs. others (282 upregulated >2-fold, 286 downregulated

>2-fold), and 2,652 in P3 vs. others (106 upregulated >2-fold, 377 downregulated >2-fold) with

q-values <0.05 (Figure S2E) (Table S3ABC). KEGG and Hallmark analyses identified cancer-

related pathways in each subtype (Figure 2H). P1 showed low proliferation and enriched

metabolic processes. P2 showed enrichment of epithelial-mesenchymal transition (EMT), NF-kB

pathway, inflammatory responses, and KRAS signaling, with downregulated oxidative

phosphorylation. P3 exhibited cell cycle activation, DNA repair, low EMT, and higher muscle

differentiation. Transcriptome regulatory network analysis identified 60 dysregulated regulons

across proteome subtypes (Figure 2I). The regulons formed 3 clusters. Cluster 1 showed high

activity in the P2 subtype with poor outcome (Figure 1B), with MYC enrichment confirmed by IHC

(Figure 2J). Cluster 2 showed high activity in P3, with enriched cell cycle-related transcription

factors. DNMT1 enrichment in P3 suggests methylation differences. PAX6, a tumor progression

factor[26-28], was upregulated in P3 and confirmed by IHC (Figure 2J). Cluster 3 showed high

activity in P1 and some P3 samples. MITF, regulating immune response[29], was enriched in P1

(Figure 2J) and validated in the TCGA STLMS dataset (Figures 2J, S2F).

Impact of genomic alterations on proteomic subtypes and cell cycle-related components

We studied cis effects of copy number alteration (CNA) on mRNA and protein. Analysis found
320 and 365 cis correlations of CNA-mRNA and CNA-protein respectively (FDR <0.1) (Figure
3A), with 131 genes overlapping (Figure 3B). GOBP over-representation analysis of cis regulated
genes that showed positive CNA-protein abundance correlations revealed significant enrichment
of cell cycle-related terms (Figure 3C). Most cell cycle components showed positive correlation
with the cell cycle score at CNA, mRNA, and (phosho)protein levels (Figure 3D). The P3 subtype
showed highest cell cycle activity, Ki67 immunolabeling (Figure 3E), and AURKA protein
expression (Figure 3F), an emerging cancer treatment target[30]. E2F targets were enriched in
P2/P3 (Figure 3G), with CDK1/2 kinase activity significantly higher than in P1 (Figure 3H).
Independent TCGA STLMS data validated these findings (Figure S3A-C). CRISPR-Cas9 gene
perturbation experiments of the DepMap project showed CDK1 and AURKA inhibition greatly
affected cell survival (Figure 3I), suggesting that these proteins are potential treatment targets
for high proliferation P3 subtype STLMS. Metastatic STLMS samples showed increased cell
cycle score, proliferation rate, and CDK2 kinase activity relative to primary site STLMS (Figure
S3DE), with a trend for elevated AURKA protein expression (Figure S3F). TMA cohort 2 IHC
scores validated these findings (Figure S3G). Combined cohort analyses of recurrence-free
survival showed poor outcomes associated with increased expression of cell cycle-related
components (Figure S3H), supporting AURKA and CDKs as potential therapeutic targets against
STLMS including metastatic lesions.
167

Phosphoproteomic alterations associated with proteomic subtypes and receptor tyrosine
kinase pathways
We investigated phosphoproteomic differences between STLMS subtypes focusing on receptor
tyrosine kinase molecules (RTK-RAS pathways). Among 69 DNA-sequenced samples, 56
samples (81.2%) had at least one genomic alteration in RTK-RAS pathway genes. *MAP2K4* and
IGF1R were the most frequent events. Though alteration frequencies were higher in P2/P3 vs.
P1, differences were not statistically significant (Figure 4A). Genomic alteration frequencies of
PDGFRA (overall frequency,14%), *FGFR2* (12%), *MET* (9%), *TSC2* (7%), *ERBB3* (6%), *FGFR4*
(6%), *INPP4B* (6%) differed significantly between proteomic subtypes. All these genes except
FGFR2 showed enrichment in P2, while *FGFR2* was enriched in P1. Analyses of differential
expression at phosphoproteome level between subtypes revealed Ras protein signal
transduction enrichment in P2 and cell cycle term enrichment in P3 (Figure 4B), compatible with
RTK-RAS gene enrichment in P2 and cell cycle enrichment in P3 (as shown in Figures 2 and 3).
We profiled kinase activity using phosphopeptide expression data and the RoKAI app[31] to

generate kinase-substrate networks (Figure 4CD). CSNK1A1 and PDK1-4 were enriched in P1.
AURKA activity was downregulated in P1, matching AURKA IHC assessment (Figure 3F). CDKs
(CDK1,2,5,6), AURKB, mTOR, and MAPK1 (ERK2) were enriched in P2. CDKs (CDK2,7),
AURKB, ROCK1, and BRAF were enriched in P3. AKT1/3 showed enrichment in P3 versus
P1/P2, though not statistically significant. P1 had a distinct kinase activity profile from P2/P3,
while P2/3 shared kinase activities such as AURKB or CDK2. This difference may derive from
distinct genomic alteration patterns (Figure 4A). PTM-GSEA analyses showed similar kinase
activity differences between STLMS subtypes (Figure S4A). AURKB kinase activity scores were
higher in P2/P3 versus P1 (Figure 4E). Furthermore, AURKB IHC positivity was higher in P2/P3
compared to P1 (Figure 4F). Consistent with this observation, AURKB mRNA expression in the
independent TCGA cohort was also higher in P2/P3 versus P1 (Figure S4B). The most frequent
AURKB phosphorylation site with activating effect is T232 based on the PhosphositePlus
database (https://www.phosphosite.org/homeAction). Motif analysis of AURKB-T232 predicted
responsible kinases for phosphorylation and inferred AURKB upstream regulators (Figure 4G).
Most inferred upstream kinases showed activation in P2/P3, while AURKB-activating kinases
were downregulated in P1.
198
NCOR1 protein expression is associated with smooth muscle differentiation, cancer
stemness, and immune-suppressive phenotype
Epigenetic regulators affect gene expression via chromatin structure regulation, including histone
protein modification, DNA methylation, TF binding, and chromatin accessibility[32]. With *NCOR1*
amplification frequently observed (Figure 2C), we investigated epigenetic regulator gene
alterations in cohort 1 (Figure 5A). *NCOR1* showed highest alteration frequency (35%), followed
by *SMARCA4* (13%), *BRD4* (12%), and *CARM1* (12%). *BRD4*, *NSD1*, and *SMARCA4* alterations
differed significantly between subtypes. *NCOR1* was amplified in over 30% of P2/P3 samples
versus 14.2% in P1 (Figure 5B). *NCOR1* copy number correlated with NCOR1 protein level, with
P3 showing a trend for highest expression. P2/3 of TCGA STLMS showed significantly higher
NCOR1 amplification frequency, resulting in highest *NCOR1* mRNA expression in P3 (Figure
S5AB). As *NCOR1* was a top altered epigenomic regulator enriched in poor-outcome P3, we
investigated the impact of high vs. low NCOR1 protein expression. We divided our cohort by
median NCOR1 protein expression and performed a differential expression analysis with GSEA.
GSEA revealed muscle organ development terms enriched in the NCOR1-high group, while
immune and epithelial process terms were depleted (Figure 5E), suggesting NCOR1's role in
muscle differentiation. Smooth muscle differentiation markers trended higher in the NCOR1-high

group (Figure 5F), with TCGA STLMS data showing similar results (Figure S5C). The NCOR1-

high group showed significantly higher stemness scores (Figures 5G, S5D) and PROM1 (CD133)

and ALDH7A1 protein expression (Figure 5H). TCGA data showed higher *POU5F1* (Oct3/4),

PROM1 (CD133), and *ALCAM* mRNA in the *NCOR1*-high group (Figure S5E). Consistent with

the Hippo pathway's role in stemness, we found YAP1/TEAD expression significantly higher in

the NCOR1-high group (Figure 5IJ), confirmed by TCGA data showing YAP1/TAZ (encoded by

WWTR1)/TEAD enrichment (Figure S5F). These findings suggest a role for NCOR in stemness

maintenance and smooth muscle differentiation regulation.

Enrichment of homologous recombination deficiency and nonhomologous end joining in

poor-prognosis subtypes

We observed high chromosomal instability (CIN) in P2/P3 (Figures 2B, S2B) with whole genome

duplication and DNA repair enrichment in P3 (Figure 2H). The main repair pathway for DNA

double strand breaks (DSB) is homologous recombination repair (HRR)[33]. Inactivation leads to

genomic alterations, CIN, poor prognosis, and metastasis in tumors[34-36]. In cohort 1, we

examined 64 HRR pathway genes from the TCGA sarcoma cohort[37] and found 38 genes altered.

Using allele specific copy number and variant calls, HRR genes showed frequent loss of

heterozygosity (LOH) (Figure 6A). HRR gene events were enriched in P2/P3 (Figure 6B). The

HRD (homologous recombination deficiency) score was highest in P2 (Figure 6C) and correlated

with CIN (Figure 6D). The TCGA dataset confirmed these findings (Figure S6A-D). We

additionally assessed RAD51 protein (a key recombinase for homologous recombination[38])

expression by IHC, finding a trend for decreased RAD51 staining in P2/P3 vs. P1 (Figure 6E),

also supported by TCGA STLMS RPPA data (Figure S6E). Analysis of paired samples from the

same patients showed increased HRD scores in metastatic sites vs, primary sites for P3 STLMS

(Figure 6F), aligning with CIN-metastasis association[35,36]. We analyzed genomic alteration

patterns of the top 5 altered HRR genes in STLMS (*BRCA2*, *RPA1*, *ATM*, *MRE11*, and *FANCA*)

in cohort 1. The most frequent combination affected all 5 genes simultaneously (Figure 6G). For

single or 2-gene combinations, *BRCA2* was predominant (total events = 17). *BRCA2* LOH tumors

showed higher HRD scores than non-LOH tumors (Figure 6H). Assessment of LOH effects of

the remaining 37 HRR genes on both mRNA and protein levels revealed decreased expression

in LOH-positive STLMS (Figure 6I), a finding also supported by the external TCGA STLMS

mRNA dataset (Figure S6F). TCGA STLMS RPPA protein data confirmed reduced BRCA2

protein in HRD-high and BRCA2-LOH groups (Figure S6G). BRCA2-loss by IHC was frequent in

P2 (Figure 6J), which showed the highest HRD score among subtypes (Figure 6C). We assessed

RAD50 expression in tumor tissue by IHC since RAD50 is a core protein of DSB recognition, a
starting point of the HRR pathway. While HRD status alone showed no statistical difference in
RAD50 expression in cohort 1, P2/P3 subtypes featured decreased RAD50 expression versus
P1 (Figure 6K), a finding also supported by external TCGA data (Figure S6H). Assessment of
phosphorylated H2AX (DNA double-strand break marker) showed a positivity range shift
between subtypes (Figure 6L), suggesting impaired DSB recognition and repair in P2/P3
compared to P1. We also evaluated nonhomologous end joining (NHEJ), an alternative DSB
repair process. We assessed both classical NHEJ driven by KU70/80 (coded by XRCC6/5) and
alternative NHEJ driven by PARP1[39]. While XRCC5/6 increased in P3 compared to P1, P2
showed no significant difference to P1 (Figure 6M). PARP1 showed increased expression in the
HRD-high group and the P3 subtype (Figure 6N), with P2 showing a close-to-significant increase,
suggesting NHEJ pathway activation in P3 and possibly P2/P3. External TCGA data confirmed
these results (Figure S6I). Using RAD51 and BRCA2 IHC protein expression data from cohorts
1 and 2, we found decreased expression of both proteins significantly associated with poor
patient outcome (Figure 6O).
265
Tumor immune microenvironment profiling reveals possible mechanisms of immune
evasion
To elucidate cancer immune evasion mechanisms, we performed integrated proteogenomic
analyses with IHC assessment of immune cell infiltration. Immune microenvironment status was
determined using the ESTIMATE ImmuneScore[40], which correlates with the CD3+ T-cell count
by IHC (Figure 7A). The ImmuneScore correlates with antigen presenting machinery (APM)
components, including MHC class I pathway members and IRF1 (Figure 7A). APM-related
molecules showed good concordance between genomic copy number, mRNA, and protein
expression (Figure 7A), indicating genomic alterations affect the immune microenvironment. The
P3 subtype of STLMS showed a significantly reduced ImmuneScore (Figure 7B), a finding
corroborated by external TCGA data (Figure S7AB). Paired sample analyses showed metastatic
tumors had lower ImmuneScores (Figure 7C) and downregulated immune-related processes
(Figure S7C). Comparing immune-cold vs. immune-hot tumors revealed 356 differentially
expressed proteins, with 75 upregulated and 281 downregulated in immune-cold tumors (Figure
7D, Table S4). GSEA confirmed downregulation of antigen-presenting pathways in immune-cold
tumors (Figure 7E), also supported by TCGA STLMS data (Figure S7D). Metastatic tumors
showed similar downregulated antigen processing pathways (Figure S7E). ERAP1, TAP1/2,
MHC class I complex molecules (HLA-B/C, B2M), and proteasome components were

downregulated in immune-cold tumors (Figure 7F). IRF1, a major transcription regulator of MHC
class I and APM molecules[41], showed lower expression in immune-cold tumors (Figure 7G). We
observed positive correlations between IRF1 and MHC class I molecules (HLA-B/C, B2M),
though not statistically significant. IRF1 correlated positively with proteins in antigen processing
(including immunoproteasome components), such as TAP1, TAP2, TAPBP, ERAP1, PSMB8,
and PSMB9, while catalytic subunits of proteasome (PSMB5, PSMB7) were negatively correlated
(Figure 7H), findings supported by TCGA transcriptome data (Figure S7F). IRF1 correlated with
IFNγ-JAK-STAT pathway, which regulates its expression[42], suggesting IFNγ signaling may have
therapeutic potential against immune-cold STLMS[43-45]. Hypoxic environments may suppress host
immune responses against tumors[46-48]. However, we found no significant correlation between
hypoxia and the STLMS ImmunsScore (Figure 7A).

Next, we assessed the expression of tumor immune suppressors in immune-hot tumors,
which may abrogate an immune response. Transcriptome analysis of cohort 1 found LGALS9
was upregulated in immune-hot tumors (Figure 7I). TCGA STLMS data showed high expression
of immune suppressive molecules including LGALS9 (Figure S7G). As LGALS9 overlapped
between the two cohorts, we evaluated its protein expression by IHC in cohorts 1 and 2. IHC
confirmed tumor cells expressed LGALS9 with a trend for higher expression in the P2 (immune-
hot) subtype (Figure 7J). Interestingly, LGALS9 protein expression was not detectable by IHC in
benign smooth muscle, such as blood vessel walls or intestinal walls. The TCGA STLMS cohort
showed a similar result (Figure S7H). Given LGALS9 upregulation in immune-hot tumors, we
assessed tumor-associated macrophages via CD68 (pan-macrophage marker) and CD163 (M2
macrophage marker) IHC to evaluate their contribution to an immunosuppressive
microenvironment. M2-type macrophages were enriched in the P2 subtype (Figure 7K). Survival
analyses showed high CD68 or CD163 cell counts correlated with shorter recurrence-free
survival (Figure 7L). These results suggest that the P2 subtype evades immune surveillance
through LGALS9 expression and M2 macrophage recruitment, creating an immunosuppressive
microenvironment (despite an immune-hot background) that is associated with poor clinical
outcome.

To concisely summarize the multi-omic findings, we generated an integrative schematic
of the proteome-defined subtypes of STLMS (Figure 8). The figure summarizes, for each subtype,
the characteristic genomic and pathway alterations, kinome alterations, DNA-repair features,
tumor microenvironment, prognostic associations, and hypothesis-generating drug targets.

Discussion

In this detailed proteogenomic analysis of STLMS, we identified three proteomic subtypes (P1, P2, and P3) with notable clinical and molecular differences (Figure 8). These subtypes are associated with patient outcomes, genomic alterations, signaling pathway activations, and immune microenvironment profiles, offering insights into STLMS heterogeneity and potential targeted therapies.

Unsupervised clustering of proteomic data identified the P2 subtype as having the worst prognosis with the shortest overall survival (OS) and recurrence-free survival (RFS). Both P2 and P3 subtypes were significantly enriched in metastatic behavior and higher-grade tumors, linking proteomic profiles to tumor aggressiveness. Validation using an external TCGA STLMS dataset confirmed the clinical relevance of our proteomic subtyping. The TCGA study identified only two subtypes (iClusters 1 and 2)[12] but lacked comprehensive (phospho)proteomic data, focusing mainly on the AKT signaling pathway. Our integrated proteogenomic analyses, however, identified three molecular subtypes, providing a more detailed characterization that enhances understanding of STLMS tumor biology and heterogeneity.

Genomic analysis shows key differences in genomic instability between P1, P2, and P3 subtypes. High ploidy and whole-genome duplication in P2 and P3 indicate aggressive tumor phenotype, linked to rapid evolution and poor prognosis[49]. Enhanced chromosomal instability (CIN) in these subtypes may drive tumor heterogeneity and therapy resistance[50]. Higher alterations in oncogenic drivers like *IGF1R* and *PDGFRA* in P2 and P3 indicate activation of pathways of cell proliferation and survival[49]. Frequent alterations in SWI/SNF complex genes in P2, including *SMARCA4* and *BRD4*, suggest disrupted chromatin remodeling, causing aberrant gene expression and tumorigenesis[51]. The co-occurrence of *NCOR1* and *MAP2K4* gene amplifications in P2 and P3 is significant; NCOR1 regulates gene expression through chromatin remodeling, and MAP2K4 is vital in the MAPK/ERK pathway[49]. These amplifications contribute to tumor progression by maintaining cancer stemness and activating RTK pathways, crucial for cancer progression and metastasis[49].

Copy number alterations (CNAs) correlated with mRNA and protein expression levels, particularly in genes linked to cell cycle regulation (Figure 3). The P3 subtype showed highest cell cycle activity and elevated AURKA expression, revealing potential therapeutic targets such as CDK1/2 and AURKA against aggressive STLMS subtypes. CRISPR/Cas9 gene perturbation experiments have supported the viability of these targets, offering promising avenues for targeted therapies[52,53].

The P1 subtype shows low proliferation and metabolic process enrichment, suggesting slower-growing tumors with metabolic vulnerabilities. The P2 subtype exhibits EMT, NF-κB

signaling, immune responses, and RAS pathways, indicating an aggressive, metastatic phenotype. The P3 subtype shows cell cycle activation and DNA repair dysregulation, suggesting highly proliferative tumors susceptible to therapies targeting cell cycle checkpoints or DNA repair deficiencies (Figures 2, 3, 6, 7). Phosphoproteomic analyses have revealed subtype-specific kinome programs corresponding to biological characteristics and therapeutic hypotheses. In subtype P1, kinase activity differed from P2 and P3, favoring metabolic regulation over proliferation. CSNK1A1 and the pyruvate dehydrogenase kinase axis (PDK1-PDK4) were selectively enriched, while AURKA/B activity was diminished (Figure 4C-D). This aligns with oxidative-metabolic pathway enrichment and low proliferation in P1 and *FGFR2* alterations. These results suggest a metabolically reprogrammed, growth factor-dependent, yet cell cycle-quiescent state may be targetable with FGFR2/PDK-axis strategies (Figures 2H,4A, and 8). P2 showed an RTK-RAS/ERK-mTOR cell cycle signature. *IGF1R*/*PDGFRA* alterations were enriched, and CDK1/2/5/6, AURKB, mTOR, and MAPK1(ERK2) were highly scored, with downstream S6K/RSK families (Figure 4A-D). These signals match the EMT phenotype and poor outcomes of P2, supporting CDK1/2 or AURKB blockade with mTOR/ERK pathway inhibition (Figures 2H and 4C-D) as therapeutic targets. Subtype P3 showed a proliferative mitotic/replication kinome with CDK2/7, AURKB, ROCK1, and BRAF activity, and elevated AKT1/3 compared to P1/P2 (Figure 4C-D). Together with P3's high cell-cycle scores and AURKA protein, these data suggest sensitivity to CDK/Aurora inhibitors, while ROCK1 and BRAF enrichment implies actomyosin and MAPK dependencies (Figures 3D-H and 4C-D). AURKB activity was elevated in P2/P3 versus P1, validated by IHC, identifying AURKB as a cross-subtype mitotic driver (Figure 4E-F). AURKB-T232 motif analysis suggested upstream mitotic and checkpoint kinase activation in P2/P3, supporting AURKB-targeted therapy (Figure 4G). These kinome data define P1 as PDK-activated with metabolic/FGFR2 bias, P2 as RTK-ERK-mTOR/CDK-driven, and P3 as CDK/Aurora-centered with ROCK/MAPK inputs (Figures 4 and 8).

Transcriptional regulatory network analysis identified multiple transcription factors for each subtype (Figure 2I). These findings highlight the complexity of STLMS biology and importance of understanding subtype-specific regulatory networks for targeted therapies. Future studies should focus on validating these transcription factors as therapeutic targets and exploring their regulatory network interactions. The P2 subtype showed elevated activity of oncogenic regulators, including MYC. MYC is an oncogene that drives cell proliferation, growth, and metabolism. Its activation links to aggressive tumor phenotypes and poor prognosis[56], aligning with P2's worse outcome (Figure 1B). The P3 subtype showed enrichment for cell cycle-related transcription factors, including E2Fs, DNMT1, and PAX6, suggesting epigenetic regulation in its

pathogenesis.

NCOR1 expression in our cohort was associated with both increased smooth-muscle
marker expression and higher expression of progenitor-like stemness markers (PROM1 (CD133)
and ALDH7A1) (Figure 5). NCOR1, as a chromatin corepressor interacting with HDAC3, can
reshape transcriptional programs to favor a lineage-restricted progenitor state that co-expresses
differentiation markers while retaining self-renewal properties, which is a behavior seen in muscle
physiology and chromatin regulation. Additionally, the increased activity of YAP1/TAZ-TEAD in
tumors with high NCOR1 levels suggests a potential collaborative pathway, as YAP/TEAD
signaling encourages the proliferation of muscle progenitors and supports regenerative
processes. These combined characteristics might contribute to a differentiated yet progenitor-
like phenotype that is resistant to therapy in STLMS. To validate this hypothesis, functional
disruption and chromatin analysis will be necessary[57].

Our observations showed that HRD is common in both P2 and P3 subtypes, alongside
activation of NHEJ pathways (Figure 6)[58]. Elevated HRD scores correlated with increased
chromosomal instability. Loss of heterozygosity in HRR genes, notably *BRCA2*, was linked to
reduced HRR protein expression and unrepaired DNA double-strand breaks, as shown by higher
phosphorylated H2AX levels (Figure 6L). PARP1 upregulation suggests that PARP inhibitors
may be effective therapeutic agents in HRD-high STLMS (Figure 6N). HRD, a hallmark of
genomic instability, is characterized by inability to repair DNA double-strand breaks through
homologous recombination[59]. This deficiency leads to reliance on error-prone repair mechanisms
like NHEJ, resulting in chromosomal rearrangements and mutations. In sarcomas, particularly
uterine leiomyosarcoma, HRD is associated with aggressive tumor behavior and poor
prognosis[60]. Identification of HRD in two of our proteomic subtypes of STLMS points to the
potential for targeted therapies against STLMS exploiting this vulnerability, such as PARP
inhibitors.

Immune profiling analysis revealed that the P3 subtype and metastatic tumors exhibit
lower ImmuneScores and downregulation of antigen-presenting machinery (APM) components,
contributing to an immunosuppressive tumor microenvironment. Reduced expression of IRF1
and APM molecules in immunosuppressed tumors may facilitate immune evasion mechanisms.
Conversely, immune-hot tumors demonstrated concomitant upregulation of suppressive immune
checkpoint molecules, such as LGALS9, and M2 macrophage recruitment. These results suggest
that combinatory immunomodulatory therapies, such as LGALS9 or PD-L1 inhibition[61], could be
strategically tailored to the immune profile and proteomic subtype of STLMS.

Our study presents a large, systematic proteogenomic analysis of STLMS that also

includes a phosphoproteomic characterization of kinome alterations. By identifying subtype-
specific genomic alterations and pathway activations, the study establishes a foundational
molecular basis for precision diagnostics and therapeutic interventions. Aggressive STLMS
subtypes may benefit from therapies targeting CDK1/2 and AURKA or from exploiting HRD with
PARP inhibitors. Patients with immune-hot tumors expressing LGALS9 may respond to immune
checkpoint inhibitors. These findings highlight clinically relevant heterogeneity within STLMS and
suggest that future precision medicine testing should incorporate proteogenomic or surrogate
assays to assign molecular subtypes and guide risk stratification and treatment selection. Our
comprehensive proteogenomic dataset of primary and metastatic STLMS provides a valuable
resource for further validation of these observations and for the development of practical assays
to implement subtype–based management in routine care for patients with STLMS.
431

Materials and Methods

Clinical specimens and pathological data

We examined tissues from two cohorts of STLMS patients. Cohort 1 consists of 72 primary and
metastatic STLMS tissues from 54 patients, including 18 primary-metastasis-paired samples.
Cohort 2 consists of 222 surgically resected STLMS samples. The study was approved by the
Institutional Review Board (MSKCC #16-1683, BIDMC #2023P000933). Clinical data, such as
patient demographics, treatment history, recurrence status, clinical targeted MSK-IMPACT
sequencing results, or histologic grade were retrieved from medical records in an anonymized
fashion. Tumor content ratios for all samples were verified by board-certified pathologists.
Clinicopathologic features and prognosis information, including gender, age, pathological TNM
stage, anatomic site, are summarized in Table S1.
443

Targeted cancer gene sequencing

MSK-IMPACT targeted sequencing was conducted as previously outlined[62]. In summary,
genomic DNA was extracted from both tumor and adjacent normal tissues using the DNeasy
Tissue kit and the EZ1 Advanced XL system (Qiagen). The extracted DNA was fragmented with
the Covaris E200 device. Custom DNA probes were crafted for targeted sequencing of all exons
and selected introns across 505 genes. These probes were synthesized using the NimbleGen
SeqCap EZ library custom oligo system and were biotinylated. Sequencing libraries were
prepared following the KAPA HTP protocol (Kapa Biosystems) and the Biomek FX system
(Beckman Coulter), and sequencing was performed on the Illumina HiSeq 2500 to achieve high,
uniform coverage (>500x median coverage). Demultiplexed fastq files underwent trimming with

Trim Galore v0.2.5mod (https://github.com/FelixKrueger/TrimGalore) to remove adapters and

short reads. These files were then aligned to the human reference genome (GRCh37) using

BWA-MEM   v0.7.5a   (with   arguments   -M   -t   6)[63],   and   Picard   Tools   v2.9

(https://github.com/broadinstitute/picard) was used. AddOrReplaceReadGroups was applied to

annotate read groups, and MarkDuplicates was used to identify PCR duplicates. Genomic

regions were pinpointed using FindCoveredIntervals from the GATK Tool Kit v3.3-0[64]and

underwent indel realignment with Assembly Based ReAligner (ABRA) v2.12[65]. GATK

BaseRecalibrator was employed to identify systematic errors in base quality scores. Variant

calling was executed in paired tumor/normal mode using MuTect v1.1.4[66]for single nucleotide

variants (SNV). Pindel v0.2.5a7[67]was utilized for small insertions and deletions (indels). Vardict

v1.5.1[68]served as the variant caller to report both SNVs and indels. The vcfs generated by

MuTect, Vardict, and Pindel were subsequently combined. Additionally, copy-number variants,

including chromosomal instability (CIS) and whole-genome doubling (WGD), were identified

using   FACETS[69].   The   resulting   variants   were   annotated   using   vcf2maf   v1.6.14

(https://github.com/mskcc/vcf2maf), which employs Ensembl's Variant Effect Predictor v86. In

conclusion, we successfully conducted MSK-IMPACT sequencing on 69 out of 72 samples in

cohort 1 (LM_010P, LM_020P, LM_043P lacked sufficient DNA for sequencing). Amplifications

and   deep   deletions   were   identified   from   the   GISTIC   result   table   named

"allthresholded.bygenes.txt" (part of the GISTIC output used to determine the copy number status

of each gene in each sample). Genes with a value of +2 were classified as amplifications, while

those with a value of -2 were considered deep deletions. Sequencing results are detailed in Table

S5.

Allele-specific copynumber analysis

FACETS analyses were conducted to ascertain genome-wide allele-specific and absolute DNA

copy numbers for all samples (FACETS version 0.5.6)[69]. Allele-specific copy number data were

utilized to estimate tumor purity and ploidy. Prior to further analyses, total copy number log ratios

were adjusted for ploidy and purity. Tumors exhibiting whole-genome doubling (WGD) were

identified as those in which more than 50% of the autosomal genome possessed a major copy

number ≥2, where the major copy number is defined as the number of copies of the most

prevalent allele present in the sample[70]. Loss of heterozygosity (LOH) was defined as the loss of

one allele in cohort 1. For the TCGA STLMS cohort, allele-specific copy number data were

obtained via the R TCGAbiolink package[71], and LOH was defined using the same criteria.

HRD score calculation

The scarHRD R package was utilized to calculate the HRD score of our cohort 1 samples[72]. This metric was derived from the simple addition of three types of genetic alterations: loss of heterozygosity (LOH), large-scale transitions (LST), and telomeric allelic imbalance (TAI) based on allele-specific copy number data. LOH refers to the elimination of genomic segments exceeding 15 Mb in size, excluding entire chromosomes. LST encompasses chromosomal breakages between adjacent regions of at least 10 Mb, with breaks occurring no more than 3 Mb apart. TAI quantifies the disparities in allele sequence contributions within the telomeric regions of chromosomes. For the TCGA cohort, we downloaded precalculated HRD score from the prior study[73].

Chromosomal instability inference

Chromosomal instability (CIN) scores of our cohort 1 and the TCGA STLMS data was computed on the allele-specific copy number segments by adding the number of gains (CNV >2), losses (CNV <2), and loss of heterozygosity per sample based on data from the prior study[37].

Recurrent somatic copy number alteration detection

Recurrent somatic copy number alterations (SCNAs) were identified using the Genomic Identification of Significant Targets in Cancer (GISTIC, version 2.0.23)[74] to ascertain which SCNA regions exhibited amplification or deletion at a frequency greater than expected by chance, with a q value threshold of 0.05. The analysis employed the following parameters: Amplification Threshold = 0.3, Deletion Threshold = 0.3, Cap Values = 1.5, Confidence Level = 0.99, Join Segment Size = 4, Arm Level Peel Off = 1, and Sample Normalization Method = mean. Default values were applied for all other parameters.

RNA sequencing

For each sample, 20-30 mg of frozen tissue was homogenized in 1 mL of TRIzol Reagent (ThermoFisher catalog # 15596018), followed by phase separation induced with 200 µL of chloroform. RNA was extracted from 350 µL of the aqueous phase utilizing the miRNeasy Mini Kit (Qiagen catalog # 217004) on the QIAcube Connect (Qiagen) in accordance with the manufacturer's protocol. Samples were eluted in 34 µL of RNase-free water. Subsequent to RiboGreen quantification and quality control via the Agilent BioAnalyzer, 500 ng of total RNA with RIN values ranging from 5.9 to 9.9 underwent polyA selection and TruSeq library preparation as per the instructions provided by Illumina (TruSeq Stranded mRNA LT Kit, catalog # RS-122-

2102) with 8 cycles of PCR. Samples were barcoded and sequenced on a NovaSeq 6000 in a

PE100 run, employing the NovaSeq 6000 S1 Reagent Kit (200 Cycles) (Illumina). Read quality

was evaluated using FastQC. Trimmed raw reads were aligned to the human genome version

hg19 using STAR (v2.7.5a)[75]with default parameters, and gene annotation was incorporated

using Gencode v36, followed by the calculation of gene-level count values and TPM values via

the RSEM tool (version 1.3.1)[76].

Tissue proteome extraction

Aliquots of 5 mg of frozen tissue were lysed using a lysis buffer composed of 8 M urea and

200 mM EPPS (4-(2-Hydroxyethyl)-1-piperazinepropanesulfonic acid), at pH 8.5, supplemented

with protease inhibitor (complete mini EDTA-free, Roche) and phosphatase inhibitors (cocktails

2 and 3, Sigma). This was followed by homogenization using a Bead Ruptor. Subsequently, the

tissue mixture underwent sonication through 12 cycles of 1-minute sonication at 120 W power

(FB120, Fisher Scientific), with intermittent cooling. Post-centrifugation at 18,000 g for 10 minutes

at 4 °C, the supernatant containing all soluble proteins was collected. Protein concentrations

were quantified using BCA assays (Pierce). Pooled protein samples were prepared and stored,

with equal amounts of all samples combined to assess MS run quality.

Liquid chromatography- mass spectrometry (LC-MS) proteomic analysis

Protein samples were labeled utilizing the 16-plex TMT chemical labeling reagent (Thermo Fisher

Scientific), followed by MultiNotch MS3 LC–MS analysis employing an Orbitrap Fusion MS[77]. In

brief, 300 µg of proteins were reduced with 5 mM TCEP (tris(2-carboxyethyl) phosphine

hydrochloride), alkylated with 10 mM IAA (iodoacetamide), and quenched with 10 mM DTT

(dithiothreitol). The samples were subsequently diluted and precipitated using chloroform-

methanol[78]. The resulting pellets were resuspended in 50 µL of 200 mM EPPS buffer and

digested with Lys-C and trypsin at 37 °C overnight. Anhydrous acetonitrile was added to achieve

a final volume of 30%. TMTPro (16-plex) reagents were introduced to the peptides at a 2.8:1

(TMT reagent-to-peptide) ratio and incubated for 1 hour at room temperature. A label ratio check

was conducted to determine mixing ratios, labeling efficiency, and missed cleavages by pooling

1 µL from each sample. Based on these results, samples were mixed to achieve equal intensity

across TMT channels. The samples were dried to remove acetonitrile, desalted using C18 solid-

phase extraction (SPE) Sep-Pak (Waters), and vacuum-centrifuged. Five percent of the desalted

samples were reserved for global proteome analysis, while 95% were utilized for phosphopeptide

enrichment. Dried peptides intended for global proteomics were fractionated using the Pierce

High pH Reversed-Phase Peptide Fractionation Kit (ThermoFisher catalog # 84868) into nine fractions per the modified manufacturer's protocol. The fractions were vacuum-centrifuged and then reconstituted in 1% ACN/0.1% FA for LC–MS analysis. For optimal phosphopeptide enrichment, phosphopeptides were collected using the High-Select TiO2 Phosphopeptide Enrichment Kit (Thermo Fisher) and from the stored flow-through of Fe-NTA using the Pierce High-Select Fe-NTA Phosphopeptide Enrichment Kit (Thermo Fisher), and both were combined. Phosphopeptide samples were fractionated into nine fractions utilizing the Pierce High pH Reversed-Phase Peptide Fractionation Kit. The fractions underwent analysis via LC-MS/MS employing a nanoAQUITY UPLC (Waters) with a 50 cm (75 µm inner diameter) EASY-Spray column (PepMap RSLC, C18, 2 µm, 100 Å) maintained at 60 °C, in conjunction with an Orbitrap Fusion Lumos Tribrid Mass Spectrometer (Thermo Fisher). Peptide separation was conducted at a flow rate of 300 nL/min using a linear gradient of 1-35% acetonitrile (0.1% FA) in water (0.1% FA) over a duration of four hours, analyzed in SPS-MS3 (for global proteome) and neutral loss-triggered MS3 (phosphopeptide) modes. For global proteome analysis, MS1 scans were performed within a range of 375–1,500 m/z, with a resolution of 120 K, an AGC target of 4×1e5, and a maximum IT of 50 ms. MS2 scans were conducted on MS1 scans of charges 2-7 using 0.7 m/z isolation, collision-induced dissociation at 32%, turbo scan, and a maximum IT of 50 ms. MS3 scans employed specific precursor selection (SPS) of 10 isolation notches, an m/z range of 100–1,000, 45% CE, 50 K resolution, and an AGC target of 1e5. For phosphopeptide analysis, MS1 scans were conducted within a range of 380-1,400 m/z, with a resolution of 120 K, an AGC target of 4×1e5, and a maximum IT of 50 ms. MS2 scans were performed on MS1 scans of charges 2-7 using 0.7 m/z isolation, collision-induced dissociation at 35%, turbo scan, and a maximum IT of 50 ms. Neutral loss-triggered MS3 scans utilized a range of 100–1,000 m/z, 38% CE, 60 K resolution, an AGC target of 1e5, and a maximum IT of 118 ms. Targeted loss masses of 97.9763 and 79.9658 were employed.

TMT data analysis

Raw data files were processed utilizing Proteome Discoverer (PD) version 2.4.1.15 (Thermo Scientific). In each of the TMT experiments, raw files from all fractions were consolidated and analyzed using the SEQUEST HT search engine with a Homo sapiens UniProt protein database, downloaded on 2019/12/13 (92,249 entries). Oxidation (M), phosphorylation (STY), and deamidation (NQ) were designated as variable modifications, whereas cysteine carbamidomethylation, TMT 16plex (K), and TMT 16plex (N-terminus) were specified as fixed modifications. The precursor and fragment mass tolerances were set at 10 ppm and 0.6 Da,

respectively. A maximum of two missed trypsin cleavages were allowed. Searches employed a

reversed sequence decoy strategy to control the peptide false discovery rate (FDR), with a 1%

FDR established as the identification threshold. Phosphosite localization was determined using

the PD IMP-ptmRS node. The raw abundance of protein, phosphoprotein, and phosphopeptide

levels was normalized by sample loading normalization of each TMT channel after filtering out

protein/peptide assignments with valid values for less than 30% of samples and K-Nearest

Neighbors (KNN) imputation. Internal reference scaling based on pooled samples assigned for

each TMT plex was conducted across the entire cohort to eliminate batch effects, followed by

TMM normalization for median centralization between TMT batches[79,80]. Differential expression

analyses were performed using edgeR[81].

Unsupervised clustering of proteomic data

We used non-negative matrix factorization (NMF)-based clustering of global protein and

phosphoprotein expression data. In brief, we constructed a merged expression matrix of

normalized global protein and phosphoprotein data and selected the top 500 (phospho)proteins

with highest mean absolute deviations for unsupervised clustering to avoid noisy

(phospho)proteins in the data. The resulting matrix was then subjected to NMF analysis

leveraging the NMF R package[82]. To determine the optimal factorization rank k (number of

clusters) for the matrix, we tested a range of clusters between k = 2-7. For each k, we factorized

matrix V using 1,000 iterations with random initializations of W and H. To determine the optimal

factorization rank, we calculated two metrics for each k: (1) cophenetic correlation coefficient

(measuring how well the intrinsic structure of the data was recapitulated after clustering) and (2)

silhouette score (indicating how well each sample lies within its cluster compared to other

clusters[83]) (Figure S1CD). These metrics indicate the reproducibility of the clustering across

1,000 iterations. The cophenetic correlation coefficients for k = 2 and 3 were >0.95 and showed

a large drop at k = 4. The silhouette score at k = 2, 3 showed relatively higher scores than that

at k = 4. Based on these results and census heatmaps for each k, we chose k = 3 as the optimal

clustering number with sufficient samples in each class for downstream analysis.

Gene set enrichment analysis (GSEA)

Fold change values for gene expression between specified groups were calculated using edgeR.

We employed clusterProfiler (v2.1.2)[84]with a ranked gene list ordered by fold change value.

Enrichment tests were conducted for GOBP, Hallmark, KEGG, and Reactome gene sets

obtained from the Molecular Signatures Database (MSigDB v7.4). For the functional

characterization of each sample via single sample GSEA, we calculated normalized enrichment

scores (NES) of cancer-relevant gene sets by projecting the matrix of normalized expression

data onto GOBP, Hallmark, KEGG, and Reactome pathway gene sets. This was done using the

ssGSEA implementation available at https://github.com/broadinstitute/ssGSEA2.0 with the

following parameters: sample.norm.type = "log", weight = 0.75, statistic = "area.under.RES",

output.score.type = "NES", nperm = 1000, min.overlap = 3, correl.type = "z.score". To elucidate

gene set enrichment in each sample, we applied single sample GSEA on normalized expression

data[85]and calculated the normalized enrichment score (NES) for MSigDB terms.

633 Over-representation analysis (ORA)

634 ORA was performed to identify biological pathways and functional categories enriched in the

635 gene set of interest by using clusterProfiler (v2.1.2)[84]. Enrichment was calculated using a

636 hypergeometric test against the defined background, and p-values were adjusted for multiple

637 comparisons using the Benjamini-Hochberg method.

639 Posttranslational modification signature enrichment analysis (PTM-SEA)

To infer kinase activity based on phophosite-specific quantification data, we performed PTM-

SEA (https://github.com/broadinstitute/ssGSEA2.0), an adaptation of ssGSEA, that analyzes

site-specific signatures by scoring PTMsigDB bi-directional signature sets. PTMsigDB, sourced

from over 2,500 publications, contains modification site-specific signatures linked to

perturbations, kinase activities, and signaling pathways. Unlike other pathway databases,

PTMsigDB annotates each PTM site with its directional change upon specific perturbations or

signaling events, improving PTM-SEA's scoring. PTMsigDB's main source is the

PhosphoSitePlus database[86]. The PTM-SEA requires two main inputs: a site-centric data matrix,

denoted as m and formatted in GCT v1.3, along with the PTM signatures database. In matrix m,

each row corresponds to a distinct phosphorylation site that is precisely localized to a particular

amino acid residue, with the columns indicating the measured abundances across different

samples. When multiple phosphorylation sites are identified on the same peptide, they are

transformed into individual site-specific entities for each site.

654 Kinase enrichment analysis between subtypes

655 To assess enrichment of activated kinases and phosphatases between groups based on

656 functional information such as known protein-protein interactions or kinase-substrate

657 annotations, we input phosphopeptide abundance change values (log2(fold change)) between

658 proteome subtypes into the RoKAI application (v0.8.3)[31] and obtained kinase activity scores at

659 proteome subtype level. Default parameters were used.

Transcriptional regulatory network analysis

To deduce the transcriptional regulatory activity of STLMS based on proteome subtype, we

utilized the RTN R package[87-89], which examines the link between specific transcription factors

(TFs) or gene regulators and all possible targets, as well as the enrichment status of a network

using proteome data. In summary, we employed normalized transcriptome data (n=27) and

Lambert's human TF list[90](1,612 TFs) to initially build a potential transcriptional regulatory

network (TRN). We then eliminated non-significant associations through permutation analysis

(permutation n=1000) and bootstrapping, followed by the ARACNe algorithm[91] to preserve direct

TF-target interactions, thereby removing redundant indirect interactions. The refined TRN

dataset includes the regulator gene name, its target gene name, and adjusted p-values as a

confidence measure. For enrichment analyses over a list of regulons in the refined TRN, we first

conducted a Master Regulator Analysis (MRA)[92]over a list of regulons, with adjustments for

multiple hypothesis testing. The MRA evaluates the overlap between each regulon and the

significantly dysregulated genes between subtypes, providing adjusted p-values, which resulted

in 600 confident regulons (adjusted p-values <0.05) for further analyses. We then determined

these regulons' enrichment scores between proteome subtypes using transcriptome

expressional fold change values and the GSEA-2T (two-tailed GSEA analysis) function from the

RTN R package. To identify similarities and differences in regulatory programs among samples

in our cohort, we selected 60 statistically robust regulons (adjusted p-values <1E-8), calculated

the activity score of each regulon in each sample, and visualized the data as a heatmap.

Additionally, we incorporated CRISPR perturbation scores (sourced from the Cancer

Dependency Map (DepMap) Project[93]) of 36 sarcoma cell lines and included this data in the

heatmap.

Proteome subtype inference of TCGA STLMS samples

To externally validate differences found in cohort 1 between proteome subtypes (P1, P2, and P3)

by using the TCGA STLMS transcriptome dataset, we constructed a random forest classifier of

our proteome subtypes based on transcriptome data of cohort 1 (n=27). To select informative

genes for model construction, we first conducted differential expression analyses between

subtypes and selected only differentially expressed genes (FDR <0.01, absolute value of fold

change >2) between subtypes (668 genes). Next, we divided normalized expression data of

selected genes into training (n=19) and test (n=8) datasets, followed by random forest model

training with 10-fold cross-validation using the tidymodel R package[94]. We applied the trained

classifier to the test dataset and got a 1.0 F-means value (which means all sample subtypes

were correctly predicted). We then applied the classifier to the TCGA STLMS transcriptome

dataset (n=53) and inferred proteomic subtypes in the dataset. TCGA sample IDs and inferred

proteomic subtypes are shown in Table S6.

Assessment of immune status

ImmuneScores were derived from the normalized protein expression matrix of 72 samples

utilizing the ESTIMATE (Estimation of STromal and Immune cells in MAlignant Tumor tissues

using Expression data) package in R[40]. To evaluate immune cell infiltration, we conducted IHC

with immune cell markers (CD3, CD4, CD8, CD68, CD163) and tallied the number of positive

cells for each marker across 3 TMA cores per sample. Subsequently, we computed the average

number of positive cells per core for each sample for further analysis.

Tissue microarray (TMA) construction

TMAs were created from FFPE blocks of 67 STLMS tissues in cohort 1 and 222 STLMS tissues

in cohort 2. Three 1-mm cores were extracted from each tissue paraffin block and placed into

tissue array blocks using a TMA arrayer (3DHistech). Tumor and normal regions were chosen

after a thorough examination of individual histologic slides and electronic image-based selection

of target areas for coring.

Immunohistochemistry (IHC)

Four-μm sections were cut from FFPE tissue blocks. Paraffin was removed with xylene, and

antigens were retrieval by heat-mediated epitope retrieval. Tissue sections were stained using a

Leica BOND-MAX IHC autostainer. The antibodies used were anti-CD3 (Leica Microsystems,

NCL-L-CD3-565, 1/200 dilution), anti-CD4 (Sigma, 104R-15, 1/100 dilution), anti-CD8 (Dako,

M7103, 1/1500 dilution), anti-CD68 (Dako, M0876, 1/200 dilution), anti-CD163 (Leica

Microsystems, NCL-CD163, 1/250 dilution), anti-MITF (Abcam, ab303530, 1/100 dilution), anti-

MYC (Abcam, ab32072, 1/200 dilution), anti-PAX6 (Abcam, ab195045, 1/100 dilution), anti-

AURKA (Cell Signaling Technology, #91590, 1/100 dilution), anti-AURKB (Abcam, ab45145,

1/100 dilution), anti-CDK1 (Abcam, ab133327, 1/250 dilution), anti-CDK2 (Cell Signaling

Technology, #18048, 1/200 dilution), anti-active YAP1 (Abcam, ab205270, 1/20,000, dilution),

anti-TEAD (Abcam, ab97460, 1/400 dilution), anti-NCOR1 (Invitrogen, PA1-844A, 1/1000

dilution), anti-RAD51 (Abcam, ab133534, 1/100 dilution), anti-BRCA2 (Sigma, HPA026815, 1/50

dilution), anti-RAD50 (Cell Signaling Technology, #61915, 1/4000 dilution), anti-phosphorylated

H2AX (Cell Signaling Technology, #9718, 1/200 dilution), anti-Ki67 (Biocare, PRM325, 1/100

dilution), anti-IRF1 (Abcam, ab243895, 1/100 dilution), anti-LGALS9 (Abcam, ab227046, 1/200

dilution). For MYC, MITF, CDK1, CDK2, NCOR1, YAP1, RAD50, RAD51, PAX6, IRF1, and

LGALS9 IHC, The staining intensity of each tumor cell was rated on a scale from 0 to 3+, and the

average was taken from three separate tissue cores for each case. To determine the total

weighted IHC score (IHC H-score, theoretically ranging from 0 to 300) for a sample, the

expression intensity of individual tumor regions (rated 0-3+) was multiplied by their respective

contributions (0-100%) to the overall tumor area, and these values were then summed. Two

pathologists independently evaluated all tissue samples. If there were any discrepancies, the

cases were re-evaluated until a consensus score was achieved. For Ki67, AURKA/B, TEAD,

BRCA2, and phosphorylated H2AX IHC, we assessed the fraction of positive tumor cells. BRCA2

loss was defined as less than 50% positivity.

External datasets

The TCGA STLMS dataset (53 samples) including clinical information, transcriptome data (count

data, TPM), and gene alterations with allele-specific copy number data (data for 1 sample was

not available) was obtained via the TCGAbiolink R package (version 2.30.0)[71]. Protein

expression data from reverse phase protein array (RPPA) experiments of the TCGA cohort was

downloaded from MD Anderson Cancer Center database

(https://tcpa.drbioright.org/rppa500/main.html).

Data and code availability

Our mass spectrometry proteomics data have been deposited in Harvard Dataverse and is

available under controlled access (DOI: 10.7910/DVN/BWOCM9). Raw and processed data of

MSK-IMPACT genomic sequencing and transcriptome data were deposited to the EGA

repository with the identifiers EGAS50000000594 and EGAS50000000595, respectively. Other

data derived from this study will be shared upon reasonable request. We did not generate original

code in this study. Any additional information required to reanalyze the data reported in this paper

is available from the lead contact upon request.

Abbreviations (there are a lot more missing)

STLMS - soft tissue leiomyosarcoma. IHC - immunohistochemistry. SNV - single nucleotide

variant. FFPE - formalin fixed paraffin embedded. GSEA - gene set enrichment analysis. PTM-SEA - posttranslational modification signature enrichment analysis. CIN - chromosomal instability. CNA - copy number alteration. DSB - DNA double-strand break. EMT - epithelial–mesenchymal transition. FDR - false discovery rate. APM - antigen-presenting machinery.

Acknowledgements

MHR expresses gratitude for the grants received from the NIH/NCI (R21 CA251992, R21 CA263262, and U01 CA263986), a Cycle for Survival Equinox Innovation Grant, an Investigator Grant from the Neuroendocrine Tumor Research Foundation (NETRF), and the support from the Farmer Family Foundation. Portions of the research were funded by the MSKCC NCI Cancer Center Support Grant (P30 CA008748). The funding organizations did not participate in the study's design, nor in the data collection, analysis, or interpretation.

Author Contributions

AT designed the study, carried out experiments, performed data analyses, and wrote a draft of the manuscript. MO performed sample processing and clinical data collection. YO carried out experiments and data analysis. RH and ZL supported mass spectrometric experiments and data analyses. NA provided expert advice on sarcoma pathology. DK consulted on the study and provided partial funding. JW wrote and edited the manuscript. MR designed and supervised the study, analyzed data, wrote and edited the manuscript, and provided funding.

Declaration of Interests

Atsushi Tanaka, Makiko Ogawa, Yusuke Otani, Ronald C. Hendrickson, Zhuoning Li, and Narasimhon Agaram have stated that they have no conflicts of interest concerning this research. David S. Klimstra served as a consultant for Paige.AI, while Michael H. Roehrl is part of the Scientific Advisory Board at Universal DX. J.Y.W. is the founder of Curandis. These companies did not participate in the study's design, implementation, data analysis, or any other related activities.

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this work the authors used Paperpal in order to check English grammar. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Figure titles and legends

Figure 1. Integrative study design and clinicopathological features of proteome subtypes

(A) Summary workflow and metrics of this study

(B) Summary plot of cohort 1 including clinicopathological information and proteome subtypes. In the heatmap, proteins and phosphoproteins which show significant differences (FDR <0.05) between proteome subtypes are shown.

(C) Kaplan-Meier curves of overall survival and recurrence-free survival between proteome subtypes

(D) Differences of clinicopathological factors between proteome subtypes

Figure 2. Landscape of genomic alterations and regulatory activity across proteome subtypes

(A) A boxplot showing ploidy between subtypes. P2/P3 show significantly higher ploidy than P1.

(B) Bar chart of whole genome duplication (WGD) frequency in each subtype on the left and boxplot of chromosomal instability on the right. P2/P3 show significantly higher frequency of WGD and a higher CIN score than P1.

(C) Oncoprint of integrated genomic alterations including SNV, Indel, and CNA. Genes with 8 events across cohort 1 (>12% samples) are shown. The bar chart on the right side of the oncoprint shows the alteration frequency across cohort 1. The bar chart on the top shows the number of genomic events of each sample.

(D) Bar plot showing arm-level amplifications and deletions. In general, P1 shows lower frequency of arm-level events for each chromosome. P3 shows the highest frequency of arm-

level deletions for most chromosomes.

(E) Concordant with arm-level event frequencies for each chromosome, total event numbers of

amplifications are significantly higher in P2/P3 than in P1. For arm-level deletions, P3 shows the

highest event numbers compared with P1 or P2.

(F) Focal amplification/deletion patterns in each subtype are shown. P1 has a relatively lower

number of focal peak events. The 17p12 peak (NCOR1 and MAP2K4 gene loci) are shared

between the P2 and P3 subtypes.

(G) Venn diagram showing overlap of genes in in the recurrent focal peaks of each subtype.

(H) Balloon plot showing pathway enrichment analyses of Hallmark and KEGG terms between

subtypes. Only statistically significant results are shown.

(I) Regulatory activity profile of STLMS by subtype status based on transcriptome data.

Computed regulon activity scores are shown as a heatmap. Regulons with $p<1E-8$ from the

Master Regulator Analysis process are shown. The boxplot on the right side of the heatmap

shows DepMap CRISPER gene effect scores for 36 sarcoma cell lines.

(J) Boxplots of MYC, PAX6, MITF IHC scores by subtypes are shown. In addition, representative

IHC images are shown. Each transcription factor shows an expression profile of subtypes that is

concordant with regulon network analyses of transcriptome data. Bars, 50 μm.

Figure 3. Impact of genomic alterations on the proteome

(A) The correlations between CNA and mRNA (left) and protein expression (right) levels (y axes)

reveal both cis and trans effects. The presence of significant positive (red) and negative (green)

Spearman's correlations (FDR <0.1) between CNA and mRNA or protein is highlighted. The

numbers of significant cis and trans events corresponding to the indicated genomic loci specific

to mRNAs or proteins are represented by blue bars, while the overlap between CNA-mRNA and

CNA-protein events is depicted by black bars.

(B) Venn diagram showing cis correlation overlap between CNA-mRNA and CNA-protein.

(C) GOBP over-representatation analysis of cis correlation between CNA-protein is shown. Cell

cycle-related terms are significantly enriched.

(D) Genomic effect of cell cycle-related genes on mRNA and protein. Columns represent each

sample and are ordered by cell cycle score. The bar chart on the right side of heatmap shows

Spearman's correlation coefficients between targets and cell cycle score. "E2F targets" is based

on ssGSEA of the Hallmark E2F term. Kinase activity scores (CDK1,2 kinase activities) are from

PTM-GSE analysis. p_RB1 denotes phosphorylated protein of RB1. * denotes statistical

significance ($p<0.05$).

(E) Boxplots showing proteome subtype differences of GOBP cell cycle scores and Ki67-positive

tumor frequency.

(F) Boxplot showing AURKA protein expression (IHC score) by subtypes.

(G) Boxplot of E2F scores from ssGSEA by subtypes.

(H) Boxplot showing kinase activity scores of CDK1/2 by subtypes.

(I) Boxplot of Depmap CRISPR gene perturbation scores targeting CDKs and AURKA using 3

leiomyosarcoma cell lines. Score below -1 represents significant effect on cell survival when the

target gene is inhibited.

Figure 4. Phosphoproteomic differences between subtypes

(A) Oncoprint showing RTK-RAS pathways with 4 or more events. Amplification of MAP2K4 and

IFG1R are the top 2 most frequent alterations. * denotes statistically significant differences

(Fisher's exact test) in gene alteration frequency between subtypes.

(B) Over-representation analyses of GOBP terms. Phosphoproteins which are positively

upregulated in one subtype compared to other subtypes with FDR<0.05 are used for GOBP term

enrichment analyses.

(C) Kinase activity heatmap by subtypes. Z-scores of inferred kinase activities are plotted as a

heatmap. * denotes statistical significance. Kinase names are shown on the left side of the

heatmap.

(D) Kinase-substrate networks of the top 20 positively enriched kinases in each subtype are

shown. Note: all positively enriched kinases of P1 are shown due to its small number. Genes

shown as ellipses are kinases. Genes shown as rectangles are kinase substrates Colors of

kinase substrates denote the phosphorylation Z-score of the target's phosphorylation site. If

multiple phosphorylation sites exist in a substrate, phosphorylation averages are shown based

on color scale.

(E) Boxplot of PTM-GSEA AURKB activity score shows enrichment in P2/P3 compared to P1.

(F) Representative AURKB IHC image and boxplot of AURKB IHC score by subtype. Bar,100

μm.

(G) Networks of AURKB and inferred regulatory kinases are shown. Motif analysis of AURKB-

T232 identifies 194 kinases. Among them, kinases with significant enrichment (FDR<0.05) in

RoKAI app analysis between subtypes were selected to be shown here. Colors of kinases denote

Z-scores of kinase activity.

Figure 5. NCOR1 expression corelates with muscular differentiation, stemness, and

immune status

(A) Oncoprint of epigenetic regulators in cohort 1. NCOR1 gene is the top altered epigenetic

regulator. * denotes statistical difference of gene alteration frequency between subtypes.

(B) NCOR1 amplification shows higher frequency in P2/P3 compared to P1 in cohort 1 (butnot

statistically significant).

(C) NCOR1 copy number positively correlates with its protein level (Spearman's correlation

coefficient) in cohort 1.

(D) Boxplot of NCOR1 protein expression by subtypes in cohort 1.

(E) Top 5 positively and negatively enriched GOBP GSEA results are shown. A ranked gene list

based on fold change values between NCOR1-low and -high groups in cohort 1 was used for

GSEA.

(F) Boxplot of smooth muscle protein markers between NCOR1-low and NCOR1-high.

(G) Boxplot of GOBP_STEM_CELL_DIFFERENTIATION ssGSEA scores. In cohort 1, the

NCOR1-high group shows significantly higher scores compared to the NCOR1-low group.

(H) Boxplot of stemness markers between NCOR1-low/high groups. PROM1 (CD133) and

ALDH7A1 in the NCOR1-high group show significantly higher protein expression than the

NCOR1-low group in cohort 1.

(I) Boxplots of YAP1 IHC H-scores between NCOR1-low and -high groups in cohorts 1 and 2.

(J) Boxplots of TEAD IHC H-scores between NCOR1-low and -high groups in cohorts 1 and 2.

Figure 6. Homologous recombination repair pathway

(A) Oncoprint of HRR pathway components showing frequent LOH events in cohort 1.

(B) Boxplot of genomic alteration counts in the HRR pathway per sample shows high occurrence

in P2/P3, but not statistically significant.

(C) The HRD score is higher in P2/P3 compared to P1.

(D) The HRD score positively correlates with CIN.

(E) The RAD51 IHC scores are lower in P2/P3 compared to P1, which is concordant with the

HRD scores of these subtypes.

(F) Boxplot of HRD score comparison between paired primary and metastatic samples of P2 and

P3. Metastatic samples of P3 show a significant increase. Although overall not statistically

significant, 3 out of 4 samples of P2 show HRD score increases.

(G) UpSet plot of thetop 5 altered genes. BRCA2 is the top altered gene with multiple different

cooccurrence partners.

(H) Boxplot of HRD scores between BRCA2 LOH and non-LOH groups showing significant

increase of HRD scores in the LOH group.

(I) Boxplots of mRNA and protein expression between LOH and non-LOH groups for each HRR

gene. Among 37 HRR genes (*FANCD2* was excluded due to low LOH event number), 16 proteins

were quantified in cohort 1 and are shown here.

(J) Bar plot showing frequencies of BRCA2-retaining samples in each subtype. P2/P3 subtypes,

especially P2, show decrease of BRCA2 retention compared to P1.

(K) The RAD50 IHC score does not change significantly between HRD-low and HRD-high

groups, but it shows a downward trend in P2/P3 compared to P1.

(L) Boxplot of phosphorylated H2AX IHC scores by subtypes.

(M) Boxplot of XRCC5/6 protein expression by subtypes. P3 shows significantly higher

expression of XRCC5/6 compared to P1 or P2.

(N) Boxplot of PARP1 protein expression between HRD-low and HRD-high groups by subtypes.

(O) Kaplan-Meier analysis of overall and recurrence-free survival in combined cohorts 1 and 2,

stratified by median values of RAD51 H-score and BRCA2 tumor cell positivity.

Figure 7. Tumor microenvironment profiling shows subtype differences and immune

evasion characteristics in both immune-cold and immune-hot tumors

(A) A heatmap of immune signatures including copy number status, mRNA, and protein level. In

general, the ImmuneScore positively correlates with CD3 count, antigen presenting molecules,

and immune-related pathways. Bars on the right side of the heatmap denote Spearman's

correlation coefficients between ImmuneScores and each gene/signature.

(B) Boxplot showing the ImmuneScore by proteome subtypes for cohort 1.

(C) Boxplot showing the ImmuneScore between paired primary and metastatic tumors.

(D) Volcano plot showing differential protein expression results between immune-cold and

immune-hot. Proteins with statistical significance (q<0.05) are shown in red (log2(fold change)>1)

and blue (log2(fold change)<-1). The horizontal dashed line denotes q=0.05.

(E) GSE plot of antigen presenting pathway-related KEGG and REACTOME terms, showing

antigen presenting processes are significantly downregulated in immune-cold tumors compared

to immune-hot tumors.

(F) A bar plot of APM protein expression ratios between immune-cold and immune-hot in cohort

1. * denotes statistical significance in the differential expressional analysis shown in (D).

(G) Boxplot of IRF1 protein expression between immune-hot and immune-cold tumors

(H) A correlation plot (colored by Spearman correlation coefficient) including the ImmuneScore,

IRF1, and antigen processing machinery protein expression. *, p<0.05; **, p<0.01; ***, p<0.001.

IFNgamma_response scores are ssGSEA scores of the

"Hallmark_Interferon_Gamma_Reponse" term.

(I) Boxplot of immune-suppressive molecules between immune-cold and immune-hot tumors

showing only TIM3 is significantly upregulated in immune-hot tumors vs. immune-cold tumors.

(J) Boxplot of LGALS9 protein expression by subtypes (left side) and representative IHC image

(right side). P2 shows a trend for higher expression of LGALS9 compared to P1 or P3. Bar, 50

μm.

(K) Boxplots of CD68-positive or CD163 -sitive cell counts by subtypes.

(L) Kaplan-Meier analyses of recurrence-free survival in combined cohorts 1 and 2, stratified by

median values of CD68-positive or CD163-positive cell counts.

Figure 8. Integrated summary of proteome-defined STLMS subtypes

Schematic overview of proteome-defined STLMS subtypes (P1-P3). Major genomic alterations,

pathway alterations, kinome alterations, DNA repair alterations, tumor microenvironment

features, prognostic trends, and hypothesis-generating therapeutic targets are summarized.

APM, antigen-processing machinery; EMT, epithelial-mesenchymal transition; HRD,

homologous recombination deficiency; LOH, loss of heterozygosity; NHEJ, non-homologous

end-joining; TME, tumor microenvironment.

References

983   1.   Board, W.C.o.T.E. (2020). Soft Tissue and Bone Tumours.

984   2.   Devaud, N., Vornicova, O., Abdul Razak, A.R., Khalili, K., Demicco, E.G., Mitric, C.,
985   Bernardini, M.Q., and Gladdy, R.A. (2022). Leiomyosarcoma: Current Clinical
986   Management and Future Horizons. Surg Oncol Clin N Am *31*, 527-546.
987   10.1016/j.soc.2022.03.011.

988   3.   Serrano, C., and George, S. (2013). Leiomyosarcoma. Hematol Oncol Clin North Am *27*,
989   957-974. 10.1016/j.hoc.2013.07.002.

990   4.   Acem, I., van Houdt, W.J., Grünhagen, D.J., van der Graaf, W.T.A., Rueten-Budde, A.J.,
991   Gelderblom, H., Verhoef, C., and van de Sande, M.A.J. (2022). The role of perioperative
992   chemotherapy in primary high-grade extremity soft tissue sarcoma: a risk-stratified
993   analysis using PERSARC. European journal of cancer (Oxford, England : 1990) *165*, 71-80.
994   10.1016/j.ejca.2022.01.013.

995   5.   Gronchi, A., Miah, A.B., Dei Tos, A.P., Abecassis, N., Bajpai, J., Bauer, S., Biagini, R.,
996   Bielack, S., Blay, J.Y., Bolle, S., et al. (2021). Soft tissue and visceral sarcomas: ESMO-
997   EURACAN-GENTURIS Clinical Practice Guidelines for diagnosis, treatment and follow-
998   up(☆). Ann Oncol *32*, 1348-1365. 10.1016/j.annonc.2021.07.006.

999   6.   Pisters, P.W., Leung, D.H., Woodruff, J., Shi, W., and Brennan, M.F. (1996). Analysis of
1000   prognostic factors in 1,041 patients with localized soft tissue sarcomas of the extremities.
1001   Journal of clinical oncology : official journal of the American Society of Clinical Oncology
1002   *14*, 1679-1689. 10.1200/jco.1996.14.5.1679.

1003   7.   Lewis, J.J., Leung, D., Heslin, M., Woodruff, J.M., and Brennan, M.F. (1997). Association
1004   of local recurrence with subsequent survival in extremity soft tissue sarcoma. Journal of
1005   clinical oncology : official journal of the American Society of Clinical Oncology *15*, 646-
1006   652. 10.1200/jco.1997.15.2.646.

1007   8.   Trovik, C.S., Bauer, H.C., Alvegård, T.A., Anderson, H., Blomqvist, C., Berlin, O., Gustafson,
1008   P., Saeter, G., and Wallöe, A. (2000). Surgical margins, local recurrence and metastasis in
1009   soft tissue sarcomas: 559 surgically-treated patients from the Scandinavian Sarcoma
1010   Group Register. European journal of cancer (Oxford, England : 1990) *36*, 710-716.
1011   10.1016/s0959-8049(99)00287-7.

1012   9.   Linch, M., Miah, A.B., Thway, K., Judson, I.R., and Benson, C. (2014). Systemic treatment
1013   of soft-tissue sarcoma-gold standard and novel therapies. Nat Rev Clin Oncol *11*, 187-
1014   202. 10.1038/nrclinonc.2014.26.

1015 10.   Savina, M., Le Cesne, A., Blay, J.Y., Ray-Coquard, I., Mir, O., Toulmonde, M., Cousin, S.,
1016   Terrier, P., Ranchere-Vince, D., Meeus, P., et al. (2017). Patterns of care and outcomes of
1017   patients with METAstatic soft tissue SARComa in a real-life setting: the METASARC
1018   observational study. BMC Med *15*, 78. 10.1186/s12916-017-0831-7.

1019 11.   Chibon, F., Lagarde, P., Salas, S., Pérot, G., Brouste, V., Tirode, F., Lucchesi, C., de Reynies,
1020   A., Kauffmann, A., Bui, B., et al. (2010). Validated prediction of clinical outcome in
1021   sarcomas and multiple types of cancer on the basis of a gene expression signature related
1022   to genome complexity. Nature medicine *16*, 781-787. 10.1038/nm.2174.

1023   12.   Network, T.C.G.A.R. (2017). Comprehensive and Integrated Genomic Characterization of
1024   Adult Soft Tissue Sarcomas. Cell *171*, 950-965.e928. 10.1016/j.cell.2017.10.014.

1025   13.   Nacev, B.A., Sanchez-Vega, F., Smith, S.A., Antonescu, C.R., Rosenbaum, E., Shi, H., Tang,
1026   C., Socci, N.D., Rana, S., Gularte-Merida, R., et al. (2022). Clinical sequencing of soft tissue
1027   and bone sarcomas delineates diverse genomic landscapes and potential therapeutic
1028   targets. Nat Commun *13*, 3405. 10.1038/s41467-022-30453-x.

1029   14.   Italiano, A., Kind, M., Stoeckle, E., Jones, N., Coindre, J.M., and Bui, B. (2011).
1030   Temsirolimus in advanced leiomyosarcomas: patterns of response and correlation with
1031   the activation of the mammalian target of rapamycin pathway. Anticancer Drugs *22*, 463-
1032   467. 10.1097/CAD.0b013e3283442074.

1033   15.   Schwartz, G.K., Tap, W.D., Qin, L.X., Livingston, M.B., Undevia, S.D., Chmielowski, B.,
1034   Agulnik, M., Schuetze, S.M., Reed, D.R., Okuno, S.H., et al. (2013). Cixutumumab and
1035   temsirolimus for patients with bone and soft-tissue sarcoma: a multicentre, open-label,
1036   phase 2 trial. The Lancet. Oncology *14*, 371-382. 10.1016/s1470-2045(13)70049-4.

1037   16.   Burns, J., Wilding, C.P., R, L.J., and P, H.H. (2020). Proteomic research in sarcomas -
1038   current status and future opportunities. Semin Cancer Biol *61*, 56-70.
1039   10.1016/j.semcancer.2019.11.003.

1040   17.   Gillette, M.A., Satpathy, S., Cao, S., Dhanasekaran, S.M., Vasaikar, S.V., Krug, K., Petralia,
1041   F., Li, Y., Liang, W.W., Reva, B., et al. (2020). Proteogenomic Characterization Reveals
1042   Therapeutic Vulnerabilities in Lung Adenocarcinoma. Cell *182*, 200-225.e235.
1043   10.1016/j.cell.2020.06.013.

1044   18.   Zhang, B., Wang, J., Wang, X., Zhu, J., Liu, Q., Shi, Z., Chambers, M.C., Zimmerman, L.J.,
1045   Shaddox, K.F., Kim, S., et al. (2014). Proteogenomic characterization of human colon and
1046   rectal cancer. Nature *513*, 382-387. 10.1038/nature13438.

1047   19.   Burns, J., Wilding, C.P., Krasny, L., Zhu, X., Chadha, M., Tam, Y.B., Ps, H., Mahalingam,
1048   A.H., Lee, A.T.J., Arthur, A., et al. (2023). The proteomic landscape of soft tissue
1049   sarcomas. Nat Commun *14*, 3834. 10.1038/s41467-023-39486-2.

1050   20.   Tang, S., Wang, Y., Luo, R., Fang, R., Liu, Y., Xiang, H., Ran, P., Tong, Y., Sun, M., Tan, S., et
1051   al. (2024). Proteomic characterization identifies clinically relevant subgroups of soft
1052   tissue sarcoma. Nat Commun *15*, 1381. 10.1038/s41467-024-45306-y.

1053   21.   AACR Project GENIE: Powering Precision Medicine through an International Consortium.
1054   (2017). Cancer Discov *7*, 818-831. 10.1158/2159-8290.Cd-17-0151.

1055   22.   Ucar, D., and Lin, D.I. (2015). Amplification of the bromodomain-containing protein 4
1056   gene in ovarian high-grade serous carcinoma is associated with worse prognosis and
1057   survival. Mol Clin Oncol *3*, 1291-1294. 10.3892/mco.2015.622.

1058   23.   Brahmi, M., Lesluyes, T., Dufresne, A., Toulmonde, M., Italiano, A., Mir, O., Le Cesne, A.,
1059   Valentin, T., Chevreau, C., Bonvalot, S., et al. (2021). Expression and prognostic
1060   significance of PDGF ligands and receptors across soft tissue sarcomas. ESMO Open *6*,
1061   100037. 10.1016/j.esmoop.2020.100037.

1062   24.   Peng, L., Li, J., Wu, J., Xu, B., Wang, Z., Giamas, G., Stebbing, J., and Yu, Z. (2021). A Pan-
1063   Cancer Analysis of SMARCA4 Alterations in Human Cancers. Front Immunol *12*, 762598.
1064   10.3389/fimmu.2021.762598.

25.   Liang, J., Li, B., Yuan, L., and Ye, Z. (2015). Prognostic value of IGF-1R expression in bone
and soft tissue sarcomas: a meta-analysis. Onco Targets Ther *8*, 1949-1955.
10.2147/ott.S88293.

26.   Li, Y., Li, Y., Liu, Y., Xie, P., Li, F., and Li, G. (2014). PAX6, a novel target of microRNA-7,
promotes cellular proliferation and invasion in human colorectal cancer cells. Dig Dis Sci
59, 598-606. 10.1007/s10620-013-2929-x.

27.   Ooki, A., Dinalankara, W., Marchionni, L., Tsay, J.J., Goparaju, C., Maleki, Z., Rom, W.N.,
Pass, H.I., and Hoque, M.O. (2018). Epigenetically regulated PAX6 drives cancer cells
toward a stem-like state via GLI-SOX2 signaling axis in lung adenocarcinoma. Oncogene
37, 5967-5981. 10.1038/s41388-018-0373-2.

28.   Jing, N., Du, X., Liang, Y., Tao, Z., Bao, S., Xiao, H., Dong, B., Gao, W.Q., and Fang, Y.X.
(2024). PAX6 promotes neuroendocrine phenotypes of prostate cancer via enhancing
MET/STAT5A-mediated chromatin accessibility. J Exp Clin Cancer Res *43*, 144.
10.1186/s13046-024-03064-1.

29.   Ballotti, R., Cheli, Y., and Bertolotto, C. (2020). The complex relationship between MITF
and the immune system: a Melanoma ImmunoTherapy (response) Factor? Molecular
cancer *19*, 170. 10.1186/s12943-020-01290-7.

30.   Du, R., Huang, C., Liu, K., Li, X., and Dong, Z. (2021). Targeting AURKA in Cancer: molecular
mechanisms and opportunities for Cancer therapy. Molecular cancer *20*, 15.
10.1186/s12943-020-01305-3.

31.   Yilmaz, S., Ayati, M., Schlatzer, D., Cicek, A.E., Chance, M.R., and Koyuturk, M. (2021).
Robust inference of kinase activity using functional networks. Nat Commun *12*, 1177.
10.1038/s41467-021-21211-6.

32.   Jaenisch, R., and Bird, A. (2003). Epigenetic regulation of gene expression: how the
genome integrates intrinsic and environmental signals. Nature genetics *33 Suppl*, 245-
254. 10.1038/ng1089.

33.   Li, X., and Heyer, W.D. (2008). Homologous recombination in DNA repair and DNA
damage tolerance. Cell Res *18*, 99-113. 10.1038/cr.2008.1.

34.   Lord, C.J., and Ashworth, A. (2016). BRCAness revisited. Nature reviews. Cancer *16*, 110-
120. 10.1038/nrc.2015.21.

35.   Turajlic, S., and Swanton, C. (2016). Metastasis as an evolutionary process. Science (New
York, N.Y.) *352*, 169-175. 10.1126/science.aaf2784.

36.   Bakhoum, S.F., Ngo, B., Laughney, A.M., Cavallo, J.A., Murphy, C.J., Ly, P., Shah, P., Sriram,
R.K., Watkins, T.B.K., Taunk, N.K., et al. (2018). Chromosomal instability drives metastasis
through a cytosolic DNA response. Nature *553*, 467-472. 10.1038/nature25432.

37.   Planas-Paz, L., Pliego-Mendieta, A., Hagedorn, C., Aguilera-Garcia, D., Haberecker, M.,
Arnold, F., Herzog, M., Bankel, L., Guggenberger, R., Steiner, S., et al. (2023). Unravelling
homologous recombination repair deficiency and therapeutic opportunities in soft tissue
and bone sarcoma. EMBO Mol Med *15*, e16863. 10.15252/emmm.202216863.

38.   Baumann, P., and West, S.C. (1998). Role of the human RAD51 protein in homologous
recombination and double-stranded-break repair. Trends Biochem Sci *23*, 247-251.
10.1016/s0968-0004(98)01232-8.

39. Mansour, W.Y., Rhein, T., and Dahm-Daphi, J. (2010). The alternative end-joining
pathway for repair of DNA double-strand breaks requires PARP1 but is not dependent
upon microhomologies. Nucleic Acids Res *38*, 6065-6077. 10.1093/nar/gkq387.

40. Yoshihara, K., Shahmoradgoli, M., Martínez, E., Vegesna, R., Kim, H., Torres-Garcia, W.,
Treviño, V., Shen, H., Laird, P.W., Levine, D.A., et al. (2013). Inferring tumour purity and
stromal and immune cell admixture from expression data. Nat Commun *4*, 2612.
10.1038/ncomms3612.

41. Zhou, F. (2009). Molecular mechanisms of IFN-gamma to up-regulate MHC class I antigen
processing and presentation. Int Rev Immunol *28*, 239-260.
10.1080/08830180902978120.

42. Der, S.D., Zhou, A., Williams, B.R., and Silverman, R.H. (1998). Identification of genes
differentially regulated by interferon alpha, beta, or gamma using oligonucleotide arrays.
Proceedings of the National Academy of Sciences of the United States of America *95*,
15623-15628. 10.1073/pnas.95.26.15623.

43. van den Elsen, P.J., Holling, T.M., van der Stoep, N., and Boss, J.M. (2003). DNA
methylation and expression of major histocompatibility complex class I and class II
transactivator genes in human developmental tumor cells and in T cell malignancies.
Clinical Immunology *109*, 46-52. 10.1016/s1521-6616(03)00200-6.

44. Mora-Garcia Mde, L., Duenas-Gonzalez, A., Hernandez-Montes, J., De la Cruz-Hernandez,
E., Perez-Cardenas, E., Weiss-Steider, B., Santiago-Osorio, E., Ortiz-Navarrete, V.F.,
Rosales, V.H., Cantu, D., et al. (2006). Up-regulation of HLA class-I antigen expression and
antigen-specific CTL response in cervical cancer cells by the demethylating agent
hydralazine and the histone deacetylase inhibitor valproic acid. J Transl Med *4*, 55.
10.1186/1479-5876-4-55.

45. Angell, T.E., Lechner, M.G., Jang, J.K., LoPresti, J.S., and Epstein, A.L. (2014). MHC class I
loss is a frequent mechanism of immune escape in papillary thyroid cancer that is
reversed by interferon and selumetinib treatment in vitro. Clin Cancer Res *20*, 6034-
6044. 10.1158/1078-0432.CCR-14-0879.

46. Hu, J., Li, X., Yang, L., and Li, H. (2022). Hypoxia, a key factor in the immune
microenvironment. Biomed Pharmacother *151*, 113068. 10.1016/j.biopha.2022.113068.

47. Abou Khouzam, R., Brodaczewska, K., Filipiak, A., Zeinelabdin, N.A., Buart, S., Szczylik, C.,
Kieda, C., and Chouaib, S. (2020). Tumor Hypoxia Regulates Immune Escape/Invasion:
Influence on Angiogenesis and Potential Impact of Hypoxic Biomarkers on Cancer
Therapies. Front Immunol *11*, 613114. 10.3389/fimmu.2020.613114.

48. Pietrobon, V., and Marincola, F.M. (2021). Hypoxia and the phenomenon of immune
exclusion. J Transl Med *19*, 9. 10.1186/s12967-020-02667-4.

49. Kikutake, C., and Suyama, M. (2023). Pan-cancer analysis of whole-genome doubling and
its association with patient prognosis. BMC cancer *23*, 619. 10.1186/s12885-023-11132-
6.

50. Baker, T.M., Lai, S., Lynch, A.R., Lesluyes, T., Yan, H., Ogilvie, H.A., Verfaillie, A., Dentro,
S., Bowes, A.L., Pillay, N., et al. (2024). The History of Chromosomal Instability in Genome-
Doubled Tumors. Cancer Discov *14*, 1810-1822. 10.1158/2159-8290.Cd-23-1249.

51. Kaufmann, T.L., Petkovic, M., Watkins, T.B.K., Colliver, E.C., Laskina, S., Thapa, N., Minussi, D.C., Navin, N., Swanton, C., Van Loo, P., et al. (2022). MEDICC2: whole-genome doubling aware copy-number phylogenies for cancer evolution. Genome biology *23*, 241. 10.1186/s13059-022-02794-9.

52. Nikhil, K., and Shah, K. (2024). The significant others of aurora kinase a in cancer: combination is the key. Biomark Res *12*, 109. 10.1186/s40364-024-00651-4.

53. Jamalzadeh, S., Dai, J., Lavikka, K., Li, Y., Jiang, J., Huhtinen, K., Virtanen, A., Oikkonen, J., Hietanen, S., Hynninen, J., et al. (2024). Genome-wide quantification of copy-number aberration impact on gene expression in ovarian high-grade serous carcinoma. BMC cancer *24*, 173. 10.1186/s12885-024-11895-6.

54. Portella, G., Passaro, C., and Chieffi, P. (2011). Aurora B: a new prognostic marker and therapeutic target in cancer. Curr Med Chem *18*, 482-496. 10.2174/092986711794480203.

55. Gautschi, O., Heighway, J., Mack, P.C., Purnell, P.R., Lara, P.N., Jr., and Gandara, D.R. (2008). Aurora kinases as anticancer drug targets. Clin Cancer Res *14*, 1639-1648. 10.1158/1078-0432.Ccr-07-2179.

56. Dang, C.V. (2012). MYC on the path to cancer. Cell *149*, 22-35. 10.1016/j.cell.2012.03.003.

57. Yamamoto, H., Williams, E.G., Mouchiroud, L., Cantó, C., Fan, W., Downes, M., Héligon, C., Barish, G.D., Desvergne, B., Evans, R.M., et al. (2011). NCoR1 is a conserved physiological modulator of muscle mass and oxidative function. Cell *147*, 827-839. 10.1016/j.cell.2011.10.017.

58. Kohn, E.C., Lee, J.M., and Ivy, S.P. (2017). The HRD Decision-Which PARP Inhibitor to Use for Whom and When. Clin Cancer Res *23*, 7155-7157. 10.1158/1078-0432.Ccr-17-2186.

59. Slade, D., and Loizou, J.I. (2023). Leveraging homologous recombination repair deficiency in sarcoma. EMBO Mol Med *15*, e17453. 10.15252/emmm.202317453.

60. Dall, G., Vandenberg, C.J., Nesic, K., Ratnayake, G., Zhu, W., Vissers, J.H.A., Bedő, J., Penington, J., Wakefield, M.J., Kee, D., et al. (2023). Targeting homologous recombination deficiency in uterine leiomyosarcoma. J Exp Clin Cancer Res *42*, 112. 10.1186/s13046-023-02687-0.

61. Mimura, K., Kua, L.F., Xiao, J.F., Asuncion, B.R., Nakayama, Y., Syn, N., Fazreen, Z., Soong, R., Kono, K., and Yong, W.P. (2021). Combined inhibition of PD-1/PD-L1, Lag-3, and Tim-3 axes augments antitumor immunity in gastric cancer-T cell coculture models. Gastric cancer : official journal of the International Gastric Cancer Association and the Japanese Gastric Cancer Association *24*, 611-623. 10.1007/s10120-020-01151-8.

1184 62. Cheng, D.T., Mitchell, T.N., Zehir, A., Shah, R.H., Benayed, R., Syed, A., Chandramohan, R., Liu, Z.Y., Won, H.H., Scott, S.N., et al. (2015). Memorial Sloan Kettering-Integrated Mutation Profiling of Actionable Cancer Targets (MSK-IMPACT): A Hybridization Capture-Based Next-Generation Sequencing Clinical Assay for Solid Tumor Molecular Oncology. The Journal of molecular diagnostics : JMD *17*, 251-264. 10.1016/j.jmoldx.2014.12.006.

63. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics (Oxford, England) *25*, 1754-1760. 10.1093/bioinformatics/btp324.

64. DePristo, M.A., Banks, E., Poplin, R., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis,
A.A., del Angel, G., Rivas, M.A., Hanna, M., et al. (2011). A framework for variation
discovery and genotyping using next-generation DNA sequencing data. Nature genetics
43, 491-498. 10.1038/ng.806.

65. Mose, L.E., Wilkerson, M.D., Hayes, D.N., Perou, C.M., and Parker, J.S. (2014). ABRA:
improved coding indel detection via assembly-based realignment. Bioinformatics
(Oxford, England) *30*, 2813-2815. 10.1093/bioinformatics/btu376.

66. Cibulskis, K., Lawrence, M.S., Carter, S.L., Sivachenko, A., Jaffe, D., Sougnez, C., Gabriel,
S., Meyerson, M., Lander, E.S., and Getz, G. (2013). Sensitive detection of somatic point
mutations in impure and heterogeneous cancer samples. Nature biotechnology *31*, 213-
219. 10.1038/nbt.2514.

67. Ye, K., Schulz, M.H., Long, Q., Apweiler, R., and Ning, Z. (2009). Pindel: a pattern growth
approach to detect break points of large deletions and medium sized insertions from
paired-end short reads. Bioinformatics (Oxford, England) *25*, 2865-2871.
10.1093/bioinformatics/btp394.

68. Lai, Z., Markovets, A., Ahdesmaki, M., Chapman, B., Hofmann, O., McEwen, R., Johnson,
J., Dougherty, B., Barrett, J.C., and Dry, J.R. (2016). VarDict: a novel and versatile variant
caller for next-generation sequencing in cancer research. Nucleic Acids Res *44*, e108.
10.1093/nar/gkw227.

69. Shen, R., and Seshan, V.E. (2016). FACETS: allele-specific copy number and clonal
heterogeneity analysis tool for high-throughput DNA sequencing. Nucleic Acids Res *44*,
e131. 10.1093/nar/gkw520.

70. Bielski, C.M., Zehir, A., Penson, A.V., Donoghue, M.T.A., Chatila, W., Armenia, J., Chang,
M.T., Schram, A.M., Jonsson, P., Bandlamudi, C., et al. (2018). Genome doubling shapes
the evolution and prognosis of advanced cancers. Nature genetics *50*, 1189-1195.
10.1038/s41588-018-0165-1.

71. Colaprico, A., Silva, T.C., Olsen, C., Garofano, L., Cava, C., Garolini, D., Sabedot, T.S.,
Malta, T.M., Pagnotta, S.M., Castiglioni, I., et al. (2016). TCGAbiolinks: an R/Bioconductor
package for integrative analysis of TCGA data. Nucleic Acids Res *44*, e71.
10.1093/nar/gkv1507.

72. Sztupinszki, Z., Diossy, M., Krzystanek, M., Reiniger, L., Csabai, I., Favero, F., Birkbak, N.J.,
Eklund, A.C., Syed, A., and Szallasi, Z. (2018). Migrating the SNP array-based homologous
recombination deficiency measures to next generation sequencing data of breast cancer.
NPJ Breast Cancer *4*, 16. 10.1038/s41523-018-0066-6.

73. Shi, Z., Chen, B., Han, X., Gu, W., Liang, S., and Wu, L. (2023). Genomic and molecular
landscape of homologous recombination deficiency across multiple cancer types.
Scientific reports *13*, 8899. 10.1038/s41598-023-35092-w.

74. Mermel, C.H., Schumacher, S.E., Hill, B., Meyerson, M.L., Beroukhim, R., and Getz, G.
(2011). GISTIC2.0 facilitates sensitive and confident localization of the targets of focal
somatic copy-number alteration in human cancers. Genome Biol *12*, R41. 10.1186/gb-
2011-12-4-r41.

75.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics
(Oxford, England) *29*, 15-21. 10.1093/bioinformatics/bts635.

76.   Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq
data with or without a reference genome. BMC bioinformatics *12*, 323. 10.1186/1471-
2105-12-323.

77.   McAlister, G.C., Nusinow, D.P., Jedrychowski, M.P., Wuhr, M., Huttlin, E.L., Erickson, B.K.,
Rad, R., Haas, W., and Gygi, S.P. (2014). MultiNotch MS3 Enables Accurate, Sensitive, and
Multiplexed Detection of Differential Expression across Cancer Cell Line Proteomes.
Analytical Chemistry *86*, 7150-7158. 10.1021/ac502040v.

78.   Navarrete-Perea, J., Yu, Q., Gygi, S.P., and Paulo, J.A. (2018). Streamlined Tandem Mass
Tag (SL-TMT) Protocol: An Efficient Strategy for Quantitative (Phospho)proteome
Profiling Using Tandem Mass Tag-Synchronous Precursor Selection-MS3. J Proteome Res
17, 2226-2236. 10.1021/acs.jproteome.8b00217.

79.   Liu, H., Sadygov, R.G., and Yates, J.R., 3rd (2004). A model for random sampling and
estimation of relative protein abundance in shotgun proteomics. Analytical chemistry *76*,
4193-4201. 10.1021/ac0498563.

80.   Plubell, D.L., Wilmarth, P.A., Zhao, Y., Fenton, A.M., Minnier, J., Reddy, A.P., Klimek, J.,
Yang, X., David, L.L., and Pamir, N. (2017). Extended Multiplexing of Tandem Mass Tags
(TMT) Labeling Reveals Age and High Fat Diet Specific Proteome Changes in Mouse
Epididymal Adipose Tissue. Mol Cell Proteomics *16*, 873-890.
10.1074/mcp.M116.065524.

81.   Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. Bioinformatics
(Oxford, England) *26*, 139-140. 10.1093/bioinformatics/btp616.

82.   Gaujoux, R., and Seoighe, C. (2010). A flexible R package for nonnegative matrix
factorization. BMC bioinformatics *11*, 367. 10.1186/1471-2105-11-367.

83.   Kim, H., and Park, H. (2007). Sparse non-negative matrix factorizations via alternating
non-negativity-constrained least squares for microarray data analysis. Bioinformatics
(Oxford, England) *23*, 1495-1502. 10.1093/bioinformatics/btm134.

84.   Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for
comparing biological themes among gene clusters. Omics *16*, 284-287.
10.1089/omi.2011.0118.

85.   Barbie, D.A., Tamayo, P., Boehm, J.S., Kim, S.Y., Moody, S.E., Dunn, I.F., Schinzel, A.C.,
Sandy, P., Meylan, E., Scholl, C., et al. (2009). Systematic RNA interference reveals that
oncogenic KRAS-driven cancers require TBK1. Nature *462*, 108-112.
10.1038/nature08460.

86.   Hornbeck, P.V., Zhang, B., Murray, B., Kornhauser, J.M., Latham, V., and Skrzypek, E.
(2015). PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. Nucleic Acids Res *43*,
D512-520. 10.1093/nar/gku1267.

87.   Castro, M.A., de Santiago, I., Campbell, T.M., Vaughn, C., Hickey, T.E., Ross, E., Tilley,
W.D., Markowetz, F., Ponder, B.A., and Meyer, K.B. (2016). Regulators of genetic risk of

1275  breast cancer identified by integrative network analysis. Nature genetics *48*, 12-21.
1276  10.1038/ng.3458.

1277 88. Fletcher, M.N., Castro, M.A., Wang, X., de Santiago, I., O'Reilly, M., Chin, S.F., Rueda,
1278  O.M., Caldas, C., Ponder, B.A., Markowetz, F., and Meyer, K.B. (2013). Master regulators
1279  of FGFR2 signalling and breast cancer risk. Nat Commun *4*, 2464. 10.1038/ncomms3464.

1280 89. Robertson, A.G., Kim, J., Al-Ahmadie, H., Bellmunt, J., Guo, G., Cherniack, A.D., Hinoue,
1281  T., Laird, P.W., Hoadley, K.A., Akbani, R., et al. (2017). Comprehensive Molecular
1282  Characterization of Muscle-Invasive Bladder Cancer. Cell *171*, 540-556.e525.
1283  10.1016/j.cell.2017.09.007.

1284 90. Lambert, S.A., Jolma, A., Campitelli, L.F., Das, P.K., Yin, Y., Albu, M., Chen, X., Taipale, J.,
1285  Hughes, T.R., and Weirauch, M.T. (2018). The Human Transcription Factors. Cell *172*, 650-
1286  665. 10.1016/j.cell.2018.01.029.

1287 91. Margolin, A.A., Nemenman, I., Basso, K., Wiggins, C., Stolovitzky, G., Dalla Favera, R., and
1288  Califano, A. (2006). ARACNE: an algorithm for the reconstruction of gene regulatory
1289  networks in a mammalian cellular context. BMC bioinformatics *7 Suppl 1*, S7.
1290  10.1186/1471-2105-7-s1-s7.

1291 92. Carro, M.S., Lim, W.K., Alvarez, M.J., Bollo, R.J., Zhao, X., Snyder, E.Y., Sulman, E.P., Anne,
1292  S.L., Doetsch, F., Colman, H., et al. (2010). The transcriptional network for mesenchymal
1293  transformation of brain tumours. Nature *463*, 318-325. 10.1038/nature08712.

1294 93. Tsherniak, A., Vazquez, F., Montgomery, P.G., Weir, B.A., Kryukov, G., Cowley, G.S., Gill,
1295  S., Harrington, W.F., Pantel, S., Krill-Burger, J.M., et al. (2017). Defining a Cancer
1296  Dependency Map. Cell *170*, 564-576.e516. 10.1016/j.cell.2017.06.010.

1297 94. Kuhn M, W.H. (2020). Tidymodels: a collection of packages for modeling and machine
1298  learning using tidyverse principles.
1299

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Proteome subtypes of STLMS

	P1	P2	P3	
Genomic alterations	Genomically stable *FGFR2* amp	Genomically unstable RTK drivers (*IGF1R*, *PDGFRA*) SWI/SNF (*SMARCA4*, *BRD4*) enriched	Genomically unstable Highest arm-level deletions	
Pathway alterations	Metabolic/OXPHOS programs Low proliferation	EMT, NF-κB/inflammation High proliferation	DNA repair programs Smooth-muscle differentiation High proliferation	
Kinome alterations	CSNK, PDK high AURKA/B low	CDK1/2/5/6 high AURKA/B high mTOR, MAPK1(ERK2) high	CDK1/2/7 high AURKA/B high BRAF, AKT1/3 high	
DNA repair	HRD low	HRD high *BRCA2* LOH/protein loss frequent	HRD high Shift toward NHEJ (XRCC5/6, PARP1)	
TME	Intact APM/IRF1 axis	Immunosuppresive LGALS9 high M2 macrophages enriched	Immune-cold tendency APM down regulated	
Prognostic significance	Favorable outcome	Poor outcome	Poor outcome	
Possible drug targets	FGFR2 PDK1-4/metabolic axis	IGF1R/PDGFRA AURKA/B, CDKs, mTOR/ERK Immuno-modulation via LGALS9	CDKs, AURKA/B, PARP	

Molecular characteristics

Clinical implications

Figure 8

Figure S1. Evaluation of proteome data quality, batch effects, and robust subtype inference, related to Figure 1

(A) Bar plots showing the number of quantified proteins per sample for (top) global proteome and (bottom) phosphoproteome datasets. Samples are grouped and color-coded by TMT plex MS runs (1-6). The relatively consistent number of quantified proteins across samples and batches indicates robust and comparable proteome/phosphoproteome coverage across the TMT experiments.

(B) UMAP plots of (left) global proteome and (right) phosphoproteome profiles, colored by TMT plex MS runs (n=6). Each point represents a sample. Samples are well intermixed across TMT batches, indicating no detectable batch effect in both global proteome and phosphoproteome datasets.

(C) Consensus heatmaps for non-negative matrix factorization (NMF) clustering at ranks k=2-7, showing clustering stability across different cluster numbers. Each heatmap illustrates consensus values between sample pairs, with color bars indicating sample assignments in basis and consensus groupings. Silhouette width values are shown for each rank, reflecting clustering quality.

(D) Summary plots of NMF rank survey metrics: cophenetic coefficient (top) and silhouette width (bottom) across ranks. Rank 3 shows a relatively high silhouette score and a drop in cophenetic coefficient, suggesting it as the optimal cluster number balancing stability and separation.

(E) Association between proteome-defined subtypes (P1-P3) and TCGA iCluster LMS subtypes (iCluster1 and iCluster2) in the TCGA STLMS cohort. The distribution shows significant concordance between iCluster1 and subtype P3, and iCluster2 and subtypes P1/P2 (Fisher's exact test, p=1.6×10E-7).

(F) Kaplan-Meier analysis of recurrence-free survival in the TCGA STLMS cohort, stratified by proteome-based subtypes. Stratification by proteome-based subtypes shows significant survival differences between these subtypes. The P2/P3 subtypes show poorer recurrence-free survival compared to P1.

Figure S2. Genomic instability features and regulator profiles in the TCGA STLMS cohort and differential expression analyses of cohort 1, related to Figure 2

(A) Boxplot showing ploidy distributions by inferred proteome subtypes in the TCGA STLMS cohort (n=53).

(B) Bar chart of whole genome duplication frequency by inferred proteome subtypes in the TCGA STLMS cohort (n=53) on the left and boxplot of CIN in the TCGA STLMS cohort (n=52) on the right.

(C) Balloon plot showing GOBP over-representation analysis of genes involved in focal peak amplifications or deletions by subtype in cohort 1.

(D) Balloon plot showing KEGG over-representation analysis of genes involved in focal peak amplifications or deletions by subtype in cohort 1.

(E) Volcano plots showing differentially expressed proteins between subtypes in our cohort 1.

(F) mRNA expression profile of representative regulons by subtypes in the TCGA STLMS dataset.

Figure S3. Cell cycle feature differences in the TCGA STLMS transcriptome data and comparison of metastatic samples in cohort 1 against matched primary samples, related to Figure 3

(A) Boxplot showing cell cycle score differences by proteome subtypes in the TCGA STLMS cohort.

(B) Boxplot of "E2F targets" scores of ssGSEA by subtypes in the TCGA STLMS cohort. P2/P3 show significantly higher scores than P1.

(C) Boxplot of mRNA expression of CDK1, CDK2, and AURKA in the TCGA cohort. P2/P3 show significantly higher expressions of CDK1 and AURKA than P1. CDK2 mRNA expression in P3 is higher than in P1.

(D) Boxplot on the left side showing cell cycle score differences between 18 paired primary and metastatic samples in cohort 1. Boxplot on the right side showing Ki67-positive tumor cell frequency based on IHC by subtypes (14 paired samples) in cohort 1.

(E) Boxplot showing kinase activity scores of CDK1/2 between primary and metastatic samples. 18 paired samples were analyzed from cohort 1.

(F) Boxplot of AURKA IHC scores. 14 paired samples were available for assessment.

(G) Boxplot of Ki67, CDK1, and CDK2 IHC scores between primary and metastasis samples in cohort 2, confirming metastatic samples show significant increases of these markers.

(H) Kaplan-Meier analysis of recurrence-free survival in combined cohorts 1 and 2, stratified by Ki67, AURKA median positivities, CDK1/2 IHC H-score median values. Patients with high AURKA positivity or CDK2-high H-scores show poorer outcomes, concordant with poor survival outcome of the P2/P3 subtypes (cell cycle-enriched subtypes).

Figure S4. Phosphorylation landscape of cohort 1 and validation of findings in the independent cohort 2 and the external TCGA STLMS cohort, related to Figure 4

(A) A heatmap of PTM-GSE scores. Terms with significant differences between subtypes (ANOVA q<0.01) are shown. AURKB activity is high in P2/P3, consistent with the RoKAI results.

(B) Boxplot of AURKB mRNA expression by subtypes in the TCGA cohort. AURKB mRNA expression is significantly higher in P2/P3 compared to P1.

Figure S5. NCOR1 characteristics in the external TCGA STLMS dataset, related to Figure 5

(A) NCOR1 amplification shows higher frequency in P2/P3 compared to P1.

(B) NCOR1 mRNA expression in P3 trends higher than in other subtypes, but not statistically significant.

(C) mRNA expression of smooth muscle markers shows higher levels in the NCOR1-high group.

(D) Stem cell differentiation scores (GOBP ssGSEA scores) are significantly higher in the NCOR1-high group.

(E) mRNA expression of POU5F1, PROM1 (CD133), and ALCAM are significantly higher in the NCOR1-high group.

(F) Boxplots of YAP1/TAZ (encoded by WWTR1) related molecules. mRNA expression of YAP1, WWTR1, and TEAD1/2/3 is significantly higher in the NCOR1-high group.

Figure S6. HRR pathway in validation cohort 2 and the external TCGA cohort, related to Figure 6

(A) Oncoprint of HRR pathway coponents showing frequent LOH events in the TCGA STLMS cohort.

(B) Boxplot of genomic alteration counts in the HRR pathway per sample showing significantly higher occurrence in P2/P3 in the TCGA cohort.

(C) The HRD score is higher in P2/P3 compared to P1 in the TCGA cohort.

(D) The HRD score positively correlates with CIN in the TCGA cohort.

(E) The RAD51 protein expression is lower in the HRD-low group and P2/P3 vs. P1, but not significant in the TCGA cohort.

(F) Boxplots of mRNA expression between LOH and non-LOH groups for each HRR gene. All 38 HRR genes are shown. Most genes, including BRCA2, show lower expression in the LOH group compared to the non-LOH group. Note: all genes with statistical significance show expressional decrease in the LOH group in the TCGA cohort.

(G) Boxplots of BRCA2 protein expression between HRD-low and HRD-high groups and BRCA2 LOH and non-LOH groups. BRCA2 protein expression shows the same trend as in cohort 1.

(H) Boxplots of RAD50 protein expression between HRD-low and HRD-high groups and by proteome subtypes. RAD50 protein expression is significantly downregulated in the HRD-low group and in P2/P3 subtypes.

(I) Boxplot of PARP1 mRNA expression between HRD-low and HRD-high groups and by proteome subtypes. PARP1 mRNA expression is significantly higher in the HRD-high group and the P3 subtype.

Figure S7. Tumor microenvironment analysis of the external TCGA STLMS dataset and comparison of primary vs. metastatic lesions, related to Figure 7

(A) A heatmap with immune signatures of the TCGA dataset. Immune-cold tumors show low expression of MHC class I molecules and peptide loading molecules, confirming findings in cohort 1. Bars on the right side of the heatmap denote Spearman's correlation coefficients between ImmuneScores and each gene/signature.

(B) Boxplot showing ImmuneScores by inferred proteome subtypes. Although not statistically significant, P3 shows lower scores compared to other subtypes.

(C) Top 10 significantly downregulated GOBP terms. GSEA of differential expression between primary and metastatic tumors in cohort 1 shows significant downregulation of immune-related pathways in metastatic tumors compared to primary tumors.

(D) GSE plot of KEGG pathway analyses showing significant downregulation of antigen processing pathway coponents in immune-cold tumors compared to immune-hot tumors in the TCGA dataset.

(E) GSE plot of Reactome pathway analyses showing significant downregulation of antigen processing pathway components in metastatic tumors compared to primary tumors in cohort 1.

(F) A correlation plot (Spearman's correlation coefficients) including the ImmuneScore, IRF1, and antigen processing machinery protein expression in the TCGA cohort. *, p<0.05; **, p<0.01; ***, p<0.001. Most genes and ImmuneScore show positive correlation with each other. IFNgamma_response scores are ssGSEA scores of the "Hallmark_Interferon_Gamma_Reponse" term.

(G) mRNA expressions of immunosuppressive genes in the TCGA cohort. All genes except CD276 and VTCN1 show significantly higher expression in immune-hot tumors compared to immune-cold tumors.

(H) Boxplot of LGALS9 mRNA expression by subtypes in the TCGA cohort. P2 shows higher expression of LGALS9 compared to P1/P3, which is consistent with the findings in cohort 1.