Omicron evolution drives increased ciliated cell tropism and dysfunction in nasal epithelia

Rhys H. Parry[1,#], Helen Mostafavi[1,#], Georgina McCallum[1,#], Julian D.J. Sng[1], Merja Joensuu[2], Kirsty R. Short[1,3]*, Andrii Slonchak[1,3,4]*, Alexander A. Khromykh[1,3]*

[1.] School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD, Australia
[2.] Australian Institute for Bioengineering and Nanotechnology, The University of Queensland, Brisbane, QLD 4072, Australia
[3.] Australian Infectious Diseases Research Centre, Global Virus Network Centre of Excellence, Brisbane, QLD 4072, Australia
[4.] Systems Virology Laboratory, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia
[#]joint first authors
*joint senior and corresponding authors
Correspondence: k.short@uq.edu.au; andrii.slonchak@qimrb.edu.au; alexander.khromykh@uq.edu.au

Keywords: SARS-CoV-2, Omicron, BA.5, XBB, Nasal Epithelial Cells, Cilia, Transcriptomics, Apoptosis, Host-Virus Interaction, Pathogenesis

Abstract

SARS-CoV-2 Omicron variants demonstrate greater replication efficiency in the upper respiratory tract than both ancestral virus and Delta variant, potentially due to stronger interactions with motile cilia in human nasal epithelial cells. Despite the emergence of numerous Omicron variants, such as BA.1, BA.5, and XBB, there has been no comprehensive investigation comparing their specific interactions with ciliated cells. Here, we use differentiated primary human nasal epithelial cell (NEC) cultures to compare an ancestral isolate (QLD02) to Omicron variants BA.1, BA.5 and XBB. Transcriptomic analysis revealed that BA.5 and XBB infection, unlike BA.1 and QLD02, drives a distinct host response, defined by the profound downregulation of genes essential for cilium assembly and function at 48-hours post-infection. The BA.5 and XBB signature was also characterized by the strong upregulation of pro-apoptotic and inflammatory pathways. Immunofluorescence (IF) of nucleocapsid-positive cells in NECs showed a significant 13-fold increase for BA.5 and 9-fold increase for BA.1 relative to QLD02. Additionally, BA.1 and BA.5 exhibited greater tropism (7.6-fold for BA.1, 9-fold for BA.5) for ciliated cells compared to QLD02. Finally, while BA.5 and BA.1 infection was associated with widespread apoptosis in both infected and bystander cells, the total number of ciliated cells remained unchanged. Overall, we demonstrate a stepwise evolution at the nasal barrier: while enhanced ciliated cell tropism and widespread apoptosis is a conserved feature of Omicron infection, transcriptional ciliary dysfunction is a distinct phenotype emerging with BA.5 and preserved in the XBB variant.

Graphical Abstract

Introduction

The upper respiratory tract is the primary site of infection with SARS-CoV-2 [1]. The nasal mucosa is populated by a variety of different cells, including goblet cells, ciliated cells, mucus-producing cells, and nasal progenitor cells [2]. *In vivo,* SARS-CoV-2 displays preferential tropism for multiciliated cells of the nasal epithelium [3, 4]. For example, in nasal tissue of patients hospitalised in 2020 for COVID-19 infection, SARS-CoV-2 nucleocapsid protein (NP) expression was restricted to multiciliated cells displaying acetylated alpha tubulin and absent in goblet cells, secretory, or differentiating cells [5]. Similarly, in *in vitro* air-liquid interface cultures of human nasal epithelial cells there were significantly increased levels of SARS-CoV-2 (ancestral and alpha variant) infection in ciliated cells and secretory-ciliated cells [4]. In human challenge studies with the wildtype ancestral virus of SARS-CoV-2, single cell sequencing of nasopharyngeal swabs identified evidence of productive viral infection of both ciliated and goblet cells [6]. Additionally, this study identified a subset of ciliated cells that, whilst only constituting 4% of total infected cells, contained 67% of detected viral RNA [6].

Targeting ciliated cells has important implications for viral pathogenesis. Disruption and shedding of ciliary axonemes has been indicated to impair mucociliary clearance [7]. This impaired mucociliary clearance likely facilitates the dissemination of virus to the lower respiratory tract, which may have implications for disease severity [8, 9]. Disease severity may also be exacerbated by changes to cytokine production from infected and dying ciliated cells, resulting in a dysregulated immune response [7].

Recent evidence suggests that the extent to which SARS-CoV-2 targets ciliated epithelial cells has evolved over time [10]. The Omicron variant, which emerged in late 2021 exhibits enhanced replication efficiency in the upper respiratory tract compared to the ancestral virus and the Delta variant. This increased upper respiratory tropism is thought to be associated with increased transmissibility in Omicron variants [11]. Consistent with these data, compared to the ancestral virus

and Delta, Omicron was more likely to bind motile cilia in human nasal epithelial cells [12, 13].

Since its emergence in late 2021, Omicron has diversified substantially into a variety of different variants. The BA.1 and BA.2 variants diverged first from a common Omicron ancestor (B.1.1.529) in November of 2021. Subsequently, the BA.2 variant diverged again in early 2022, giving rise to the BA.4 and BA.5 variants, characterised by increased fusogenicity and antibody evasion, likely conferred by additional spike mutations such as L452R and F486V, respectively [14]. By mid-2022, BA.5 had become the dominant global variant, and it remained as such until approximately December 2022. Recombinant events between two BA.2 variants later gave rise to XBB, which subsequently became the predominant variant in several regions by early 2023. These different Omicron variants display marked differences in tropism and pathogenesis. For example, in human nasal epithelial cells XBB1.5 replicated more efficiently than BA.2 [15]. Whether this reflects increased targeting of ciliated cells in the upper respiratory tract remains unclear, although early studies from a limited number of nasal swabs (n=4) suggested that the later Omicron variant XBC was more effective than BA.5 at targeting cilia cells in the upper respiratory tract [16]. Here, we comprehensively examine the effect of different Omicron variants on cilia in the upper respiratory tract, to better understand variant dependent differences in pathogenesis.

Results

BA.1 and BA.5 Omicron variants of SARS-CoV-2 demonstrate higher replication efficiency compared to the ancestral virus in primary human nasal epithelial cells.

Figure 1. SARS-CoV-2 BA.1 and BA.5 Omicron variants replicate more efficiently in primary human nasal epithelial cells than the ancestral QLD02 isolate. (A) Schematic overview of experimental workflow. Primary human nasal epithelial cells (NECs) were collected from healthy donors and differentiated at air-liquid interface for 28 days to establish pseudostratified ciliated epithelium. Differentiated NECs were subsequently infected and characterised through RNA-seq and immunofluorescence staining. (B) Whole genome phylogenetic inference showing evolutionary relationships among SARS-CoV-2 variants. The Omicron variants are highlighted in red, with isolates examined in this study shown in bold (QLD02/ancestral, BA.1, BA.2, BA.5, XBB (EG.1.4)). Scale bar indicates nucleotide substitutions per site. (C) RT-qPCR analysis of viral gene expression (E, S, RdRP, and N) in primary nasal epithelial cultures infected with QLD02 SARS-CoV-2 ancestral isolate and Omicron variants BA.1 and BA.5 at 48 hours post-infection. Viral RNA loads are shown as $\log_{10}$ copies per ng of total RNA. Each data point corresponds to a NEC culture generated from an independent donor. Error bars indicate median and interquartile range. Statistical significance was determined using the Kruskal-Wallis test.

To investigate different Omicron variants at the primary site of infection, we compared the replication of three key SARS-CoV-2 isolates in primary human nasal epithelial cells (NECs) (Fig. 1A). The selected variants, as indicated by whole-genome phylogeny (Fig. 1B), include: an ancestral (Pango lineage A) isolate (QLD02, Jan 2020), an early Omicron (Pango lineage BA.1) isolate (QLD2639C, Dec 2021), and a later Omicron (Pango lineage BA.5) variant isolate (QLD-QIMR03, Jul 2022). At 48 hours post-infection, intracellular viral RNA loads for four viral genes (E, S, RdRP[nsp12], and N) were quantified by RT-qPCR (Fig. 1C). Both BA.1 and BA.5 demonstrated higher viral RNA loads than the ancestral QLD02 virus. This enhanced replication was statistically significant for the S gene in both BA.1 (p = 0.0284) and BA.5 (p = 0.0133) compared to QLD02, and for the E gene, where BA.5 replication was significantly higher than ancestral (p = 0.0339), suggesting a progressive increase in intracellular replication efficiency in NECs with Omicron variants. There were no significant differences in the levels for RNA of all 4 genes between BA.1 and BA.5 variants.

Figure 2. Infection with BA.5, but not BA.1 and ancestral isolate of SARS-CoV-2 drastically decreases levels of transcripts associated with cilia function and increases apoptotic and inflammatory responses in human NECs. (A) Differentially expressed genes (DEGs) in primary nasal epithelial cells infected with ancestral isolate (QLD02), and Omicron variants (BA.1, and BA.5) of SARS-CoV-2 relative to mock-infected cells. Significantly (FDR-corrected p-value <0.05, log$_2$FC>1) up-regulated genes are shown in red and down-regulated genes are shown in blue. The data are derived from the primary cell cultures obtained from four donors. (B) Gene Ontology (Biological Process) enrichment analysis of DEGs identified in each comparison shown in (A). Z-scores indicate

cumulative activation or suppression of biological processes. Size of the bubbles is proportional to the number of DEGs associated with the respective GO term. (C) Heatmaps representing relative expression of the genes associated with cilium assembly and function, apoptosis, inflammation, JAK/STAT signaling, and IL-17 signaling in mock- QLD02-, BA.1-, and BA.5-infected NECs. The values are the means from 4 individual donors.

Different variants of SARS-CoV-2 elicit distinct transcriptional responses in human primary nasal epithelial cultures

Having established that Omicron variants exhibit enhanced intracellular replication relative to ancestral virus (Fig. 1C), we sought to understand the corresponding impact on the host transcriptional landscape. Total RNA from differentiated NEC cultures at 48 hours post-infection (Fig. 1C) was used for bulk RNA-seq. Consistent with RT-qPCR (Fig. 1C), increased viral RNA reads were observed in BA.5 and BA.1 infected cells compared to those infected with QLD02, whilst no differences in viral RNA reads were recorded between BA.5 and BA.1 infected cells (Fig. S1A). Analysis of the host response (Fig. S1B-D) revealed distinct transcriptional responses following infection with tested SARS-CoV-2 variants At 48 hours post infection, differential expression analysis identified 189, 325, and 4,839 significantly differentially regulated genes (FDR < 0.05, $log_2 FC > 1$) in QLD02-, BA.1-, and BA.5-infected cells relative to mock-infected cells (Supplementary Tables S2-S4), respectively (Fig. 2A). Strongly upregulated genes included canonical antiviral interferon-stimulated genes (ISGs) such as IFIT1, IFIT2, IFIT3, MX1, MX2, OASL, and IFNB1, alongside pro-inflammatory chemokines (CXCL10 and CXCL11). The magnitude of ISG induction was broadly similar across variants, with CXCL10, OASL, and interferon regulatory factors consistently ranking among the most significantly upregulated transcripts. Additionally, a subset of genes associated with cell cycle progression and ciliary structure were downregulated across all infections, with the most extensive and coordinated downregulation signature observed in BA.5 infected cells.

Gene Ontology (GO) enrichment analyses of differentially regulated transcripts highlighted robust activation of innate immune pathways, including "defense response to virus," "type I interferon signaling," and "negative regulation of viral genome replication" across all variants (Fig. 2B). Genes associated with "defense response to virus" (FDR-corrected p-value 6.75E-12) remained highly significant across all SARS-CoV-2 infections. For other pathways, again, BA.5 infection elicited the most complex and distinctive transcriptional program, with GO enrichment analysis revealing a unique ciliary dysfunction signature (Fig. 2B). More specifically, the BA.5 analysis was dominated by a downregulation of ciliation terms, including "intraciliary transport" and "outer dynein arm assembly", suggesting more extensive engagement with ciliation relative to QLD02 and BA.1 analyses. BA.5 infection in NECs also showed strong downregulation of cilium-related biological processes, with the GO term "cilium assembly" (FDR-corrected p-value = 8.96E-13) emerging as the most significantly enriched term.

Heatmap visualization of the most significantly affected pathways revealed coordinated repression of cilium assembly and motility genes in both BA.1 and BA.5, with the latter exhibiting the greatest reduction (Fig. 2C). Within the cilium assembly and functions gene set, BA.5-infected cells displayed widespread downregulation of structural and motor protein-encoding genes that were relatively preserved in QLD02 and, to a lesser extent, BA.1 infections. This coordinated suppression encompassed core structural and motor protein genes, including dynein heavy and intermediate chains (DNAH5, DNAH9, DNAH11, DNAI1, DNAI2), radial spoke proteins, intra-flagellar transport genes (IFT88, IFT140, IFT172), and ciliary assembly factors such as CEP290, CFAP43, and CFAP53.

Another key difference that characterised infection across the range of SARS-CoV-2 variants related to apoptotic pathways, with variable regulation evident across all conditions. Here, both BA.1 and BA.5 infections induced a more pronounced upregulation of key pro-apoptotic factors compared to QLD02 (Fig. 2C). In particular, genes from both the intrinsic (BAX, BAK1, BCL2L11) and extrinsic (FAS) apoptotic pathways were more strongly upregulated in both Omicron variants, compared to the QLD02 virus.

Infection was also characterised by progressive upregulation of inflammation-associated genes, including chemokines (CXCL1, CXCL2, CXCL8, CCL20) and cytokines (IL1A, IL1B, IL6, TNF), from QLD02 through to BA.5. Similarly, core JAK/STAT pathway components (JAK2, STAT1, STAT2) and IL-17 signalling mediators (IL17RA, IL17RC, TRAF3IP2, NFKB1, MAPK13) showed the strongest upregulation in BA.5-infected cells, which may indicate heightened sensitivity to IL-17 signalling. Together, these data demonstrate that while all SARS-CoV-2 variants elicit canonical antiviral interferon programs in differentiated nasal epithelia, Omicron variants, particularly BA.5, induce a distinctive transcriptomic signature characterised by extensive disruption of ciliary gene expression alongside the enhanced inflammatory signalling typical of the ancestral SARS-CoV-2 virus (QLD02).

Figure 3. The BA.5 variant elicits stronger antiviral responses and causes significantly more profound disruption to cilia function compared to BA.1. (A) Differentially expressed genes (DEGs) in human primary nasal epithelial cells infected with BA.5 variant of SARS-CoV-2 in comparison to BA.1 infection at 48 hours post-infection. Significantly upregulated genes (log2FC > 1, FDR < 0.05) are shown in red and significantly downregulated genes (log2FC < -1, FDR < 0.05) are indicated in blue. Top most differentially expressed genes are labelled. (B) Gene Ontology Biological Process (GO:BP) enrichment analysis of the differentially expressed genes identified in BA.5 versus BA.1 comparison shown in (A). Bubble size corresponds to number of DEGs per pathway, colour gradient represents z-score (red = activation, blue = suppression).

BA.5 variant shows enhanced disruption of ciliary processes, and greater transcription signatures of apoptosis compared to BA.1.

To further elucidate the effect of BA.5 infection on cilia disruption we performed a differential gene expression analysis comparing BA.5 and BA.1 infections. This comparison revealed 1,513 significantly differentially expressed genes (FDR < 0.05, $|\log_2 FC| > 1$; Fig. 3A, Supplementary Table S5). Gene Ontology enrichment analysis of these DEGs (Fig. 3B) revealed key differences in the activation and suppression of pathways associated with apoptosis and ciliary processes, respectively.

First, we observed a more pronounced suppression of cilium-related biological processes in BA.5 relative to BA.1 infected cells. Multiple pathways essential for ciliary function were found to be downregulated, as indicated by highly negative z-scores. These included "cilium assembly" (z-score = -4.2), "cilium movement" (z-score = -3.8), "outer dynein arm assembly" (z-score = -3.5), and "intraciliary transport" (z-score = -3.2). This specific program of downregulation appears unique to BA.5, as these processes were also minimally affected in the BA.1 versus mock comparison (Fig. 2B).

The BA.5 and BA.1 comparison also revealed a strong positive enrichment (z-scores > 4) for pathways related to "defense response to virus" and "inflammatory response," which included a coordinated induction of apoptosis-associated genes (Supplementary Table 5). A closer analysis revealed the upregulation of numerous TNF-superfamily ligands and death-receptor–associated transcripts in BA.5 compared to BA.1, including TNFSF10 (TRAIL; logFC = 1.36) and TNFSF13B (BAFF; logFC = 2.10), along with downstream apoptotic effectors like CASP10, CASP1, and BAK1. Gene Set Enrichment Analysis (GSEA) further confirmed the significant enrichment of apoptosis-related pathways, such as KEGG Apoptosis pathway and Reactome pathway for death-receptor signalling (dynamic adj. p-values ranging from 8×10^{-3} to 3×10^{-2}).These signatures appear to represent a connected biological program: TNF-family ligands (e.g., TNFSF10) initiating death-receptor signalling upstream of caspase activation. This demonstrates that the divergent pathogenesis of BA.5 compared to BA.1 is defined by a selective suppression of ciliary gene programs and a simultaneous, robust activation of the TNF-driven extrinsic apoptosis pathway.

Figure 4. The Omicron XBB (EG.1.4) variant demonstrates comparable replication to BA.5 and results in similar transcriptional profile in primary human nasal epithelial cells. (A) Viral RNA levels in human primary nasal epithelial cells at 48 hours after infection with XBB and BA.5 SARS-CoV-2 variants. Viral RNA levels were determined by RNA-Seq. Each data point represents an independent cell culture derived from a different donor (n = 5). Bars indicate mean and interquartile range. Statistical significance determined using Mann-Whitney test. (B) Differentially expressed genes (DEGs) in XBB-infected versus mock-infected primary human nasal epithelial cells at 48 hours post-infection. (C) Gene Ontology Biological Process (GO:BP) enrichment analysis of XBB infection versus mock-infected control. Bubble size corresponds to number of DEGs per pathway, color gradient represents z-score (red = activation, blue = suppression). (D) Differentially expressed genes (DEGs) in human primary nasal epithelial cells infected with XBB SARS-CoV-2 variant in comparison to BA.5 infection. In (A) and (D) significantly upregulated genes (log2FC > 1, FDR < 0.05) are shown in red and significantly downregulated genes (log2FC < -1, FDR < 0.05) are indicated in blue. Most significantly differentially expressed genes are labelled.

XBB has a similar effect to BA.5 on host gene expression in NECs.

We next sought to investigate whether Omicron variants that emerged after BA.5 (e.g. XBB) also displayed global downregulation of cilia-related transcriptional processes. Accordingly, NECs from five independent donors were infected with the XBB and BA.5 variants. At 48 hours post-infection, total RNA was isolated from SARS-CoV-2- and mock-infected cells and analysed by RNA-Seq (Fig. 4,

Supplementary Fig. 2). Quantification of viral RNA reads in RNA-Seq data at 48
hours post-infection revealed that XBB replicated at levels comparable to those
observed for BA.5 (Fig. 4A). Transcriptional profiling of the host gene expression
followed by the multidimensional scaling (MDS) analysis demonstrated strong
separation of infected groups from mock-infected controls, while no clear separation
was observed between the groups infected with BA.5 and XBB variant
(Supplementary Fig. 2B). Although the donor-stratified MDS analysis revealed subtle
separation between BA.5 and XBB infections (Supplementary Fig. 2C), the overall
clustering patterns suggest minor transcriptomic divergence of the cells infected with
BA.5 and XBB variants.

Differential gene expression analysis comparing XBB-infected cultures to mock-
infected controls identified a robust antiviral and inflammatory transcriptional
response (Fig. 4B), characterized by strong upregulation of canonical ISGs including
IFIT1, IFIT2, IFIT3, MX1, MX2, OASL, and RSAD2, alongside elevated expression of
chemokines CXCL10, CXCL11, and CCL20. Similar induction of these genes was
observed during BA.5 infection (Fig. 2A, Supplementary Fig. 2D), demonstrating
conservation of the antiviral response between the two Omicron variants. Consistent
with what was observed in BA.5 infection, XBB-infected NECs displayed a large
number of genes with reduced expression (BA.5: 3,142 genes; XBB: 4,502 genes;
Supplementary Tables S6-S7). The GO enrichment analysis confirmed that
downregulated genes were associated with cilia-related processes, which was also
observed in BA.5 infection. Key structural and motility-associated transcripts,
including DNAH5, DNAH9, DNAI1, IFT88, and CFAP43, were consistently
downregulated, reflecting a conserved transcriptomic signature of ciliary disruption in
both XBB and BA.5 variants. Enriched upregulated processes included "defense
response to virus," "type I interferon signaling," and "negative regulation of viral
genome replication," which was consistent with the transcriptional response of the
nasal epithelial cells to all SARS-CoV-2 variants analysed in the study.

Direct comparison of XBB and BA.5 infections revealed minimal transcriptional
divergence (Fig. 4D). Only a small subset of genes satisfied the significance
threshold, and they were primarily associated with fine regulation of antiviral and
apoptotic responses rather than unique biological processes (125 DEGs in direct
comparison; Supplementary Table S8). GO enrichment analysis of these few DEGs
(data not shown) revealed no significant pathway activation or inhibition exclusive to
XBB, further indicating that XBB and BA.5 induce nearly identical host transcriptional
responses.

Collectively, these findings demonstrate that the Omicron XBB variant replicates
with equal efficiency to BA.5 in primary nasal epithelial cells and elicits a highly
conserved host response characterized by robust interferon signalling and
downregulation of ciliary gene networks. This suggests that late Omicron variants
have likely converged on a similar balance of efficient replication in nasal epithelia

and modulation of epithelial homeostasis, maintaining the phenotype of ciliary
dysfunction and antiviral activation established by BA.5.

BA.1 and BA.5 infection induces widespread apoptosis in differentiated NECs
without altering overall number of ciliated cells.

Figure 5. BA.5 infection induces elevated apoptosis and higher viral burden in primary nasal
epithelial cells. (A) Representative maximum-intensity projection immunofluorescent microscopy
images of differentiated primary NECs infected for 48 h with ancestral QLD02, Omicron BA.1, or
Omicron BA.5 variants. Cells were stained for nuclei (DAPI, blue), acetylated α-tubulin (green, cilia),
cleaved Caspase-3 (CC3; magenta, apoptotic cells), and nucleocapsid protein (NP; orange, infected
cells). Scale bars = 100 μm. (B) Maximum-intensity projections of (A) in the XZ plane. (C)
Quantification of total number of DAPI$^+$ cells, acetylated α-tubulin$^+$ cells (% ciliated cells), CC3$^+$ cells
(% apoptotic cells) and NP$^+$ cells (% infected cells) across independent donors and replicate cultures.
Each small point represents a single field of view (n = 5 per replicate). Large points represent the
means of 1-3 replicates per each donor, with 3 independent donors indicated by different coloured
data points. Statistical significance was determined using the Kruskal-Wallis test, pairwise
comparisons were made with Uncorrected Dunn's test. Error bars represent median with interquartile
range.

Our transcriptomic analyses (Figs. 2-4) consistently demonstrated that BA.5 and the
later XBB variant, induce a unique signature characterized by the robust
upregulation of pro-apoptotic programs and a simultaneous, substantial
downregulation of genes essential for ciliary function. To investigate a potential
cellular basis for these transcriptional changes, we used immunofluorescence
staining to dissect cell-specific tropism and apoptosis. We stained NECs infected
with QLD02, BA.1, or BA.5 for acetylated α-tubulin (cilia, green), cleaved caspase-3
(CC3; apoptotic cells, magenta) and viral nucleocapsid protein (NP; infected cells,
orange).

Visual inspection of the maximum-intensity projections in the XY plane (Fig. 5A) and
XZ plane (Fig. 5B) revealed clear differences between variants. In BA.1- and BA.5-
infected cultures, there was a visibly greater abundance of both infected (NP$^+$) and

apoptotic (CC3⁺) cells compared to the QLD02 and mock-infected groups. The maximum-intensity projections of the XZ plane confirmed that the cilia (green) form the apical layer of the epithelium and that this is the primary site of viral infection (orange). We also noted considerable CC3⁺ staining (magenta) in the XZ plane of the BA.1 and BA.5 infected NECs, with many apoptotic cells in proximity to, but distinct from, NP⁺ cells, suggesting a bystander apoptotic effect.

Quantification of these observations (Fig. 5C) confirmed the visual trends. Firstly, we observed no differences in total cells across all conditions, but consistent with our previous transcriptomic data, BA.5 showed a significantly higher percentage of infected (NP⁺) cells (median, 23.99%) compared to the ancestral QLD02 virus (median, 1.83%), representing a 13.11-fold increase (p = 0.0113). The 9.33-fold increase for BA.1 (median, 17.07%), relative to QLD02, while visually apparent, was not statistically significant (p = 0.1360) likely due to significant variations between samples.

Increases in the percentage of infected cells were accompanied by similarly increased percentages of apoptotic (CC3⁺) cells across BA.1 and BA.5 infection, indicated by a significantly higher percentage of apoptotic (CC3⁺) cells, overall. Specifically, BA.1 infection (median, 7.044%) resulted in a 4.20-fold increase (p = 0.0415), and BA.5 infection (median, 8.254%) drove a 4.92-fold increase (p = 0.0046), relative to mock-infected cells (median, 1.677%). However, despite the profound downregulation of ciliary genes seen in the transcriptomic data, there was no variant-dependent difference in the total number of ciliated cells at 48 hours post-infection.

BA.5 and BA.1 Omicron variants are associated with increased infection and apoptosis of ciliated cells, relative to the ancestral SARS-CoV-2 virus.

Figure 6. BA.5 and BA.1 infections of NECs are associated with increased apoptosis of infected cells and preferential ciliated cell tropism compared to QLD02. (A) Representative maximum-intensity projection immunofluorescent microscopy images of differentiated primary NECs infected for 48 h with ancestral QLD02, Omicron BA.1, or Omicron BA.5 variants. Cells were stained for nuclei (DAPI, blue), acetylated α-tubulin (green, cilia), cleaved Caspase-3 (CC3; magenta, apoptotic cells), and nucleocapsid protein (NP; orange, infected cells). Scale bars = 50 μm. (B) Quantification of the percentage of infected ciliated (NP$^+$ acetylated α-tubulin$^+$) cells (top) over total ciliated cells, all apoptotic infected (CC3$^+$ NP$^+$) cells over total CC3$^+$ cells (middle), and percentage of apoptotic ciliated (CC3$^+$ acetylated α-tubulin$^+$) cells over total CC3$^+$ cells (bottom). Each small point represents a single field of view (n = 5 per replicate). Large points represent the means of 1-3 replicates per each donor, with 3 independent donors indicated by different coloured data points. Statistical significance was determined using the Kruskal-Wallis test, pairwise comparisons were made with Uncorrected Dunn's test. Error bars represent median with interquartile range.

To further dissect the cellular interactions between infection, apoptosis, and ciliation, we used immunofluorescent staining to examine their colocalization (Fig. 6). Visual analysis of the representative images (Fig. 6A) revealed a strong overlap between infected (NP$^+$) and apoptotic (CC3$^+$) cells, particularly for BA.1 and BA.5 groups, relative to QLD02. This was accompanied by a clear and increased incidence of double-positive infected (NP$^+$) and ciliated (α-tubulin$^+$, green) cells in the Omicron variants – indicating preferential tropism for ciliated cells and prompting a quantitative analysis of these specific double-positive cell populations.

Quantitative analysis first confirmed that for Omicron variants (BA.1 and BA.5), there was an enhanced tropism for ciliated cells compared to the ancestral virus (QLD02). More specifically, BA.5-infected NECs showed a significantly higher percentage of infected ciliated cells (NP$^+$ and acetylated α-tubulin$^+$) (median, 25.73%) compared to the QLD02 strain (median, 2.87%), representing an 8.97-fold increase (Fig. 6B; p = 0.0113). The BA.1 variant also demonstrated a 7.55-fold increase in double-positive infected ciliated cells (median, 21.66%), relative to the QLD02 strain; however, this comparison was not statistically significant (p = 0.1360).

Next, we directly linked the infection status (NP$^+$) of a cell to its apoptotic status (CC3$^+$). In Omicron-infected cultures. The fraction of double-positive apoptotic and infected (CC3$^+$ and NP$^+$) cells was 3.77-fold higher for BA.5 (median, 28.34%) compared to the QLD02 group (median, 7.52%). This difference was statistically significant (Fig. 6B; p = 0.0369). The BA.1 variant also showed a 3.22-fold increase (median, 24.22%) over the QLD02 virus; however, this comparison did not reach statistical significance (p = 0.0526). There was no significant difference observed between the BA.1 and BA.5 variants in this regard (p = 0.8815).

Despite these collective data indicating widespread increases to apoptosis in infected cells and increased tropism for ciliated cells, we saw no statistically significant changes to the fraction of double-positive apoptotic ciliated (CC3$^+$ α-tubulin$^+$) cells, across all variants (Fig. 6B, % CC3+ ciliated cells/ total CC3+ cells).

Together, these data demonstrate that Omicron variants (BA.1 and BA.5) are associated with increased infection of differentiated NECs (Fig. 5C,) and exhibit an enhanced tropism for ciliated cells (Fig. 6B, top panel). Infection with these variants Omicron variants is associated with a quantifiable increase in apoptosis of infected

cells; however, despite increased tropism for ciliated cells (Fig. 6B, middle panel), there are no specific increases in apoptotic ciliated cells (Fig. 6B, bottom panel). Instead, this apparently conserved pattern of infection, tropism and cell death in Omicron variants (BA.1 and BA.5) may indicate a bystander apoptotic effect; representing a pathogenesis distinct from that observed in the ancestral QLD02 virus (Fig. 6).

Discussion

SARS-CoV-2 Omicron has evolved increased replication efficiency in the upper respiratory tract compared to the ancestral virus and Delta variant, with evidence suggesting enhanced binding to motile cilia in human nasal epithelial cells [12]. However, despite the emergence of multiple Omicron variants (e.g. BA.1, BA.5, XBB), no studies to date have systematically assessed how these variants variants differ in their interaction with and impact on, ciliated cells, leaving a gap in our understanding of disease pathogenesis associated with more recent Omicron variants.

Here, we show that whilst ancestral SARS-CoV-2 (QLD02), BA.1 and BA.5 induced numerous transcriptional changes in primary human NECs, the most striking difference between Omicron variants was a pronounced downregulation in pathways for cilia function such as 'cilium assembly' and 'cilium movement' in BA.5 infected cells, relative to BA.1 infected cells. This was accompanied by increased expression of proinflammatory genes (e.g. 'defence response to virus') and increased activation of apoptosis pathways (e.g. 'positive regulation of apoptotic processes') relative to BA.1 infected cells. It would therefore be tempting to speculate that this transcriptional phenotype in BA.5 is driven by increased death of ciliated cells and subsequent loss of cilia. However, immunofluorescence staining revealed that this was not the case. Whilst BA.5 did cause increased apoptosis (likely accounting for the increased inflammation seen with this strain by transcriptomics) this was not more pronounced in ciliated cells in any variant tested (including the ancestral virus). Similarly, there was no difference in total ciliated cell populations between the ancestral virus and Omicron variants at 48-hours post-infection. These data are broadly consistent with prior studies suggesting that both the ancestral virus [8] and Omicron variants [15] do not alter tubulin staining in the first 48-hours of infection. Our transcriptomic results are also corroborated by functional evidence that SARS-CoV-2 can decrease cilia beat frequency and coordination in the absence of widespread cilia loss [8]. Accordingly, we suggest that the observed BA.5 signature of cilia dysregulation is indicative of a loss of cilia function rather than changes to the number of ciliated cells (as estimated by acetylated α-tubulin staining). We further suggest that the increased apoptosis observed in BA.5 infected cells reflects predominantly the death of other non-ciliated cells in the differentiated epithelium (e.g. goblet and/or basal cells). Indeed, these data are consistent with previous studies showing a "bystander apoptotic effect" where necroptosis is predominately

activated in SARS-CoV-2 infected epithelial cells whilst neighbouring cells undergo
apoptosis [17].

The question therefore remains as to why the phenotype of cilia dysfunction,
increased apoptosis of non-ciliated cells and increased inflammation is more
pronounced in BA.5 and XBB, compared to earlier SARS-CoV-2 variants.
Importantly, these trends are not the result of differences in viral replication
efficiency, as BA.1, BA.5 and XBB had equivalent viral reads at 48 hours post-
infection. Instead, it is possible that these data reflect the increased fusogenicity and
syncytia formation of BA.5 compared to earlier Omicron variants [18, 19]. Syncytial
formation and the resultant altered cytoskeletal dynamics have been shown to
interfere with the coordinated orientation and beating of cilia, leading to reduced
mucociliary clearance even when cilia remain morphologically intact [8]. Syncytial
formation in ciliated cells may also account for the apoptosis of uninfected bystander
cells (and subsequent inflammation). Direct comparison of syncytial formation in
primary human NECs after BA.1, BA.5 and XBB infection thus remains an important
area for future study.

Whilst these data were generated in an *in vitro* system, it is important to consider the
in vivo implications of our observations. When ciliary motility is reduced or discordant
in vivo, mucus transport is slowed, potentially enabling pathogens such SARS-CoV-2
to penetrate beyond the conducting airways [8, 20]. However, determining whether
BA.5 or even XBB is more likely to progress to lower respiratory tract infections *in*
vivo is difficult to assess, as site specific evaluation of viral replication is confounded
by strain dependent differences in tropism for cells of the upper and lower respiratory
tract [11, 15]. Accordingly, it remains to be determined as to the precise functional
impacts of cilia dysfunction in a clinical BA.5 infection.

The present study was subject to several limitations. Firstly, due to the laborious
nature of working with primary human NECs only a limited number of SARS-CoV-2
variants were tested, and it would be informative to observe if similar transcriptional
patterns were observed with more recent Omicron variants (e.g. JN.1). Finally, future
studies would benefit from performing cilia function and motility assays to better
understand the impact of the observed phenotype associated with BA.5/XBB
infections, although we acknowledge the difficulties of performing such experiments
under biosafety level 3 containment conditions.

Nevertheless, the present study provides the first insight as to the interaction of
different Omicron variants on cilia in the nasal epithelium.

Methods

Human Ethics, Nasal Epithelial Swab Collection and Processing

488 Primary nasal epithelial cells (NECs) were collected from healthy adult volunteers as
489 described previously [21]. The study was approved by the University of Queensland
490 Human Research Ethics Committee (2017000520 and 2022002162).

Primary NECs were established as previously described [21] and cryopreserved at passage 1 or 2 in freezing medium consisting of fetal bovine serum (FBS) with 10% dimethyl sulfoxide (DMSO). Cells were expanded and maintained in PneumaCult™-Ex Plus medium (STEMCELL Technologies, Canada) and seeded at a density of 4–5 × 10⬜ cells per transwell insert (6.5-mm polyester membranes, 0.4-µm pore size; Corning Costar, USA). Cultures were maintained in PneumaCult™-Ex Plus until confluent, after which they were "air-lifted" by removing apical medium and replacing the basolateral medium with PneumaCult™-ALI differentiation medium (STEMCELL Technologies). Basolateral medium was refreshed three times per week. Cells were differentiated at the air-liquid interface (ALI) for at least three weeks, until the appearance of ciliated and mucus-producing cells and attainment of a transepithelial electrical resistance greater than 500 Ωcm². Fully differentiated NEC cultures were used for infection and downstream analyses (Table 1).

Table 1: Donor information for primary nasal epithelial cells across three experiments, grey squares indicate donors used in each experiment.

Donor	Age	Sex	Exp 1[#] – RNA seq	Exp 2[#] – RNA seq	Exp 3[#] – IF staining
YZ07	23	F	�ொ	▒	
YZ08	25	F	▒	▒	
YZ19	19	F	▒	▒	
YZ21	28	M		▒	▒
YZ03	42	M		▒	
QF08	30	F			▒
YZ10	34	F			▒
YZ05	28	F			▒

[#]Viral infection conditions per replicate include: mock (PBS, experiment 1-3), QLD02 (experiment 1 and 3), BA.1 (experiment 1 and 3), BA.5 (experiment 1-3) and XBB (experiment 2-3).

Cell Lines

VeroE6 stably expressing human TMPRSS2 (VE6-hTMPRSS2 Clone #1), as previously generated and described in [22] were cultured at 37°C and 5% CO_2 in Dulbecco modified Eagle medium (DMEM; Gibco, Thermo Fisher, USA) supplemented with 10% fetal calf serum (FCS; Bovogen; USA), 1% GlutaMAX (Cat: 35050079, Thermo Fisher, USA), and 100⬜U/mL penicillin-streptomycin (Gibco, Thermo Fisher, USA), with addition of 30⬜µg/mL of puromycin dihydrochloride (A1113803, Thermo Fisher, USA).

SARS-CoV-2 Viruses

Four low-passage SARS-CoV-2 viruses were recovered from nasopharyngeal aspirates of infected individuals provided by the Queensland Health Forensic and Scientific Services, Queensland Department of Health, and the Queensland Institute of Medical Research. These isolates included: (I) an early ancestral (PANGO v.4.2 lineage A) Australian isolate hCoV-19/Australia/QLD02/2020 (QLD02) sampled on 30/01/2020 (GISAID Accession ID; EPI_ISL_407896) [22]. (II) the Omicron BA.1,

named hCoV-19/Australia/QLD2639C/2021 collected on 14/12/2021 (GISAID
accession ID EPI_ISL_8186598) [23]; (III) the Omicron BA.5 sub-lineage BE.1
named hCoV-19/Australia/QLD-QIMR03/2022 (GISAID accession ID:
EPI_ISL_15671874.2) collected 19/07/2022 [24] and (IV) an XBB.1.9 lineage named
hCoV-19/Australia/QLD-UQ01/2023 (GISAID accession ID: EPI_ISL_17784860)
collected 11/4/2023 (Unaliased XBB.1.9.2.1.4; Pango lineage EG.1.4) [24]. All
working virus stocks were propagated and titred on VeroE6 stably expressing human
TMPRSS2 (VE6-hTMPRSS2) cells using an immuno-plaque assay [25]. All studies
with SARS-CoV-2 were performed under physical containment 3 (PC3) conditions
and were approved by the University of Queensland Biosafety Committee
(IBC/522B/SCMB/2022).

Viral infection

Differentiated NECs were infected with mock (PBS), QLD02, BA.1, BA.5, and XBB at
1.5×10^5 FFU). Specifically, 100 µL of virus or PBS was placed on the epithelial
surface in the apical compartment and incubated for 1 hour at 37 °C. Following
incubation, excess virus was removed from the transwell, and cells were incubated
at 37 °C with 5% CO_2. Every 24 hours, the basolateral media were refreshed with 1
mL of new ALI media. At predetermined time points, post-infection 100 µL of PBS
was added to the apical compartment, and cells were incubated at 37 °C with 5%
CO2 for 10 minutes. The apical supernatant was subsequently collected and stored
at –80 °C.

Immunofluorescence staining and image quantification

Infected differentiated epithelial cells grown on transwell membranes were fixed with
4% paraformaldehyde (Electron Microscopy Sciences) in PBS for 30 minutes at
room temperature and washed three times with PBS. Membranes were then excised
from transwell inserts with a scalpel, followed by permeabilization with 0.1% Triton X-
100 (Sigma-Aldrich, USA) in PBS for 15 minutes. Samples were then blocked in 1%
BSA (Sigma-Aldrich, USA) in PBS for 45 minutes at room temperature. Samples
were stained with primary antibodies diluted in 1% BSA in PBS overnight at 4°C. The
following primary antibodies were used: mouse anti–acetylated α-tubulin
(Cat#66200-1-Ig, Proteintech, USA) as a cilia marker, rabbit anti-cleaved Caspase-3
(Asp175, Cat#9661, Cell Signaling Technology), and human anti–SARS-CoV-2
nucleocapsid nanobody E2 [26].

Stained samples were washed three times in 0.5% BSA in PBS, then incubated with
secondary antibodies diluted in 1% BSA in PBS for 2.5 hours at room temperature in
the dark. Secondary antibodies included: Alexa Fluor 488 donkey anti-mouse IgG
(Cat#A32766, Invitrogen, USA), Alexa Fluor 555 donkey anti-rabbit IgG
(Cat#A31572, Invitrogen, USA), and Alexa Fluor 647 donkey anti-human IgG
(Cat#A21445, Invitrogen, USA). Nuclei were counterstained with DAPI (Invitrogen)
for 5 minutes, followed by one wash in PBS. Samples were then briefly rinsed in
Milli-Q water, and mounted between two glass coverslips using ProLong Gold
Antifade Mountant (Thermo Fisher Scientific, USA).

Z-stack images (~22 µm, 0.5 µm step size,) were acquired on a SpinSR10 spinning
disk confocal (Olympus, Japan) using a 30× silicone immersion objective. All
representative images are maximum-intensity image projections generated from Z-
stacks. All microscopy imaging was performed at the Queensland Brain Institute,
University of Queensland. Cell quantification was performed by acquiring five
random fields of view per sample at 2300 × 2300 px resolution, with 1-3 technical
replicate samples per group (specified in corresponding figure legends). Image
quantification was performed in Fiji/ImageJ (v1.54p) using custom-written scripts to
identify and count cells automatically.

Total number of cells were quantified by generating maximum-intensity projections of
Z-stacks, applying a Gaussian blur ($\sigma = 2$) and automated thresholding on the DAPI
channel. Images were converted to a binary mask and DAPI$^+$ nuclei detected using
Find Maxima (prominence = 20). Point detections were recorded as ROIs and
counted automatically using Analyze Particles (<10 µm). Total number of
nucleocapsid protein (NP$^+$), cleaved Caspase-3 and ciliated (acetylated α-tubulin)
cells were quantified using the same workflow as above but applied to their
respective fluorescence channels to create a binary mask. The Fill Hole function,
followed by Watershed separation was applied to the binary mask, and positive cells
were automatically counted using Analyze Particles (< 10 µm).

For quantification of double-positive cells, images were processed as above in the
relevant two fluorescence channels. The resulting two binary masks were then
combined using ImageJ's Image Calculator (AND) function to generate an
intersection mask representing double-positive cells. Watershed separation was
applied to the intersection mask, and double-positive cells were automatically
counted using Analyze Particles (<10 µm).

RNA sequencing and analysis

Total RNA was isolated from individual NECs using TRI Reagent (Sigma-Aldrich,
USA) as done previously [27]. RNA quality and integrity were assessed on an Agilent
TapeStation 4200 (Agilent Technologies, USA), and only samples with an RNA
integrity number (RIN) > 8 were used for downstream library preparation using the
TruSeq RNA Library Preparation Kit v2, Set A (Illumina, USA) according to the
manufacturer's protocol. Indexed cDNA libraries were pooled and sequenced on an
Illumina NextSeq 500 instrument using the NextSeq 500/550 High Output Kit v2.5
(75 cycles, Illumina, USA). Image acquisition, processing, and fastq demultiplexing
were performed using in-instrument software as described previously [28].

Processing of the raw reads was performed as described previously [29]. Raw
sequencing reads were assessed for quality using FastQC (v.0.72) and adapter and
quality trimmed using Trimmomatic (v.0.36.6) using the parameters:
ILLUMINACLIP:TruSeq3-SE:2:30:10, *LEADING:32*, *TRAILING:32*,
SLIDINGWINDOW:4:20, and *MINLEN:16*. High-quality reads were aligned to the
human reference genome assembly (GRCh38/hg38) using HISAT2 (v.2.2.1) with

default parameters. Gene-level counts were obtained using featureCounts (v2.0.1)
with the following options: *counting mode* = "Union," *strand* = "Unstranded," *feature*
type = "exon," and *ID attribute* = "gene_id".

Downstream analysis and visualization of RNA-seq data were conducted in RStudio
(v2025.09.2). Differential gene expression analysis was conducted in edgeR (v.4.2).
Low abundance reads (<1 count per million) were filtered before normalisation.
Library size and composition bias were corrected using the trimmed mean of M-
values (TMM) normalisation method. Normalized counts were analysed by
multidimensional scaling analysis under default settings and used to build quasi-
likelihood negative binomial generalized log-linear model. Contrasts were tested
using the quasi-likelihood ratio test or likelihood ratio test with correction for the false
discovery rate (FDR < 0.05).

Significantly differentially expressed genes (DEGs) were visualized as volcano plots
using ggplot2 (v3.3.2). Gene Ontology (GO) term enrichment was performed using
Database for Annotation, Visualization and Integrated Discovery (DAVID, v6.8) as
previously described [29]. Enrichment scores were integrated with expression
values, and z-scores were computed and visualised using GOplot (v.1.0.2) and
ggplot2 (v.3.3.2). Heat maps of selected GO Broad Processes (BP) terms were
generated using heatmap.2 function of gplots (v3.1.2) based on scaled expression
values (row z-scores) [30].

626 SARS-CoV-2 Gene Expression Analysis

627 For quantification of viral genomic RNA abundance, adapter- and quality-trimmed
628 RNA-seq reads as above were aligned to isolate-specific SARS-CoV-2 reference
629 genomes using Bowtie2 (v2.0.5) with default mapping parameters. Read counts
630 were obtained directly from BAM files using samtools v1.18 and normalized as
631 counts per million.

632 RNA extraction and quantitative reverse transcription PCR (qRT-PCR)

Total RNA was extracted as described above and used for reverse transcription
quantitative PCR (RT-qPCR). For qPCR, 1 μg of isolated RNA was used to generate
first-strand cDNA synthesis using the qScript cDNA SuperMix (Quantabio, USA)
following the manufacturer's instructions. Quantitative PCR was conducted on a
QuantStudio 6 Flex Real-Time PCR System (Applied Biosystems, USA) using gene-
specific primers targeting the SARS-CoV-2 E, S, RdRP, and N genes
(Supplementary Table 1). Each reaction was performed in technical triplicate for
each biological replicate. Viral RNA copy number was quantified by comparison to a
standard curve generated from serial tenfold dilutions (10^2 - 10^8 copies per reaction)
of a PCR-amplified and purified SARS-CoV-2 genomic fragment. Copy numbers
were normalized to input RNA concentration and expressed as $\log_{10}$ copies ng^{-1} total
RNA. No-template controls were included in each run.

Statistical analysis

The number of biological replicates, the sample sizes and statistical methods are indicated in the figure legends. Statistical analysis was performed using GraphPad Prism 10 software. Specific tests included the non-parametric Kruskal-Wallis test with Uncorrected Dunn's post-hoc test for multiple group comparisons and the Mann-Whitney test for two-group comparisons. RNA-seq analysis was performed using RStudio (v2025.09.2) as detailed in the 'RNA sequencing and analysis' section. Statistical significance was defined as $p < 0.05$.

Data and materials availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. RNA sequencing data generated for this study have been deposited in the NCBI Sequence Read Archive (SRA) under accession number PRJNA1347416.

Author contributions:

Conceptualization, A.A.K., K.R.S., R.H.P., and A.S.; Investigation, R.H.P., H.M., G.M., J.D.J.S., M.J., and A.S.; Visualisation, R.H.P, H.M. and A.S.; Analysis and Software, R.H.P., H.M., and A.S.; Resources, A.A.K. and K.R.S.; Administration, A.A.K., A.S., and K.R.S.; Funding acquisition, A.A.K. and K.R.S.; Writing – original draft, R.H.P., G.M., K.R.S., and A.S.; Writing – review and editing, R.H.P., H.M., G.M., J.D.J.S., M.J., K.R.S., A.S., and A.A.K.

Competing interests:

K.R.S has received speaker honoraria from Pfizer. All other authors declare no competing interests.

Acknowledgements

A.A.K. is supported by NHMRC Ideas Grant (2012883) and Australian Infectious Diseases Research Center (AID) Seed Grant. A.S. was supported by NHMRC Ideas Grant (2021272) and ARC Future Fellowship (FT230100465). The authors acknowledge the facilities and the scientific and technical assistance of the Microscopy Australia Facility at the Centre for Microscopy and Microanalysis (CMM) of The University of Queensland. We thank the Queensland Health Forensic and Scientific Services, Queensland Department of Health, and QIMR Berghofer Medical Research Institute for providing SARS-CoV-2 isolates. Next-generation sequencing was performed by ACE Sequencing (SCMB, UQ). Purchasing of NGS reagents was supported by Illumina COVID Matched Funding Scheme. We are grateful for the gift of C2/E2 SARS-CoV-2 nucleocapsid nanobody from Ariel Isaacs (SCMB, UQ). This work utilised the Australian Galaxy service (https://usegalaxy.org.au/).

References

681
1.   Schulze J, Mache C, Balázs A, Frey D, Niemeyer D, Olze H, Dommerich S,
Drosten C, Hocke AC, Mall MA, Hippenstiel S, Wolff T. Analysis of Severe Acute
Respiratory Syndrome 2 Replication in Explant Cultures of the Human Upper
Respiratory Tract Reveals Broad Tissue Tropism of Wild-Type and B.1.1.7 Variant
Viruses. *The Journal of Infectious Diseases* 2021: 224(12): 2020-2024.
2.   Davis JD, Wypych TP. Cellular and functional heterogeneity of the airway
epithelium. *Mucosal Immunology* 2021: 14(5): 978-990.
3.   Baumgardt M, Obermayer B, Balázs A, Löwa A, Wyler E, Teixeira Alves LG,
Hellwig K, Beule D, Landthaler M, Müller MA, Drosten C, Mall MA, Hippenstiel S,
Hönzke K, Hocke AC. SARS-CoV-2 Infection of Human Lung Air-Liquid Interface
Cultures Reveals Basal Cells as Relevant Targets. *The Journal of Infectious*
Diseases 2025: 231(5): 1210-1220.
4.   Chang JJY, Grimley SL, Tran BM, Deliyannis G, Tumpach C, Nguyen ANT,
Steinig E, Zhang J, Schröder J, Caly L, McAuley J, Wong SL, Waters SA, Stinear TP,
Pitt ME, Purcell D, Vincan E, Coin LJM. Uncovering strain- and age-dependent
innate immune responses to SARS-CoV-2 infection in air-liquid-interface cultured
nasal epithelia. *iScience* 2024: 27(6).
5.   Ahn JH, Kim J, Hong SP, Choi SY, Yang MJ, Ju YS, Kim YT, Kim HM,
Rahman MDT, Chung MK, Hong SD, Bae H, Lee CS, Koh GY. Nasal ciliated cells
are primary targets for SARS-CoV-2 replication in the early stage of COVID-19. *J*
Clin Invest 2021: 131(13).
6.   Lindeboom RGH, Worlock KB, Dratva LM, Yoshida M, Scobie D, Wagstaffe
HR, Richardson L, Wilbrey-Clark A, Barnes JL, Kretschmer L, Polanski K, Allen-
Hyttinen J, Mehta P, Sumanaweera D, Boccacino JM, Sungnak W, Elmentaite R,
Huang N, Mamanova L, Kapuge R, Bolt L, Prigmore E, Killingley B, Kalinova M,
Mayer M, Boyers A, Mann A, Swadling L, Woodall MNJ, Ellis S, Smith CM, Teixeira
VH, Janes SM, Chambers RC, Haniffa M, Catchpole A, Heyderman R, Noursadeghi
M, Chain B, Mayer A, Meyer KB, Chiu C, Nikolić MZ, Teichmann SA. Human SARS-
CoV-2 challenge uncovers local and systemic response dynamics. *Nature* 2024:
631(8019): 189-198.
7.   Bridges JP, Vladar EK, Huang H, Mason RJ. Respiratory epithelial cell
responses to SARS-CoV-2 in COVID-19. *Thorax* 2022: 77(2): 203-209.
8.   Robinot R, Hubert M, de Melo GD, Lazarini F, Bruel T, Smith N, Levallois S,
Larrous F, Fernandes J, Gellenoncourt S, Rigaud S, Gorgette O, Thouvenot C,
Trébeau C, Mallet A, Duménil G, Gobaa S, Etournay R, Lledo P-M, Lecuit M, Bourhy
H, Duffy D, Michel V, Schwartz O, Chakrabarti LA. SARS-CoV-2 infection induces
the dedifferentiation of multiciliated cells and impairs mucociliary clearance. *Nature*
Communications 2021: 12(1): 4354.
9.   Li Q, Vijaykumar K, Phillips SE, Hussain SS, Huynh NV, Fernandez-Petty CM,
Lever JEP, Foote JB, Ren J, Campos-Gómez J, Daya FA, Hubbs NW, Kim H,
Onuoha E, Boitet ER, Fu L, Leung HM, Yu L, Detchemendy TW, Schaefers LT,
Tipper JL, Edwards LJ, Leal SM, Jr., Harrod KS, Tearney GJ, Rowe SM. Mucociliary
transport deficiency and disease progression in Syrian hamsters with SARS-CoV-2
infection. *JCI Insight* 2023: 8(1).
10.   Shi G, Li T, Lai KK, Johnson RF, Yewdell JW, Compton AA. Omicron Spike
confers enhanced infectivity and interferon resistance to SARS-CoV-2 in human
nasal tissue. *Nature Communications* 2024: 15(1): 889.
11.   Wickenhagen A, Flagg M, Port JR, Yinda CK, Goldin K, Gallogly S, Schulz JE,
Lutterman T, Williamson BN, Kaiser F, Mukesh RK, van Tol S, Smith B, van

Doremalen N, Russell CA, de Wit E, Munster VJ. Evolution of Omicron lineage towards increased fitness in the upper respiratory tract in the absence of severe lung pathology. *Nat Commun* 2025: 16(1): 594.

12.   Wu C-T, Lidsky PV, Xiao Y, Cheng R, Lee IT, Nakayama T, Jiang S, He W, Demeter J, Knight MG, Turn RE, Rojas-Hernandez LS, Ye C, Chiem K, Shon J, Martinez-Sobrido L, Bertozzi CR, Nolan GP, Nayak JV, Milla C, Andino R, Jackson PK. SARS-CoV-2 replication in airway epithelia requires motile cilia and microvillar reprogramming. *Cell* 2023: 186(1): 112-130.e120.

13.   Fonseca BF, Robinot R, Michel V, Mendez A, Lebourgeois S, Chivé C, Jeger-Madiot R, Vaid R, Bondet V, Maloney E, Guivel-Benhassine F, Schwartz O, Duffy D, Mondal T, Gobaa S, Chakrabarti LA. Stealth replication of SARS-CoV-2 Omicron in the nasal epithelium at physiological temperature. *bioRxiv* 2025: 2025.2005.2003.652024.

14.   Pather S, Madhi SA, Cowling BJ, Moss P, Kamil JP, Ciesek S, Muik A, Türeci Ö. SARS-CoV-2 Omicron variants: burden of disease, impact on vaccine effectiveness and need for variant-adapted vaccines. *Frontiers in Immunology* 2023: Volume 14 - 2023.

15.   Hu B, Chan JF-W, Liu Y, Liu H, Chen Y-X, Shuai H, Hu Y-F, Hartnoll M, Chen L, Xia Y, Hu J-C, Yuen TT-T, Yoon C, Hou Y, Huang X, Chai Y, Zhu T, Shi J, Wang Y, He Y, Cai J-P, Zhou J, Yuan S, Zhang J, Huang J-D, Yuen K-Y, To KK-W, Zhang B-Z, Chu H. Divergent trajectory of replication and intrinsic pathogenicity of SARS-CoV-2 Omicron post-BA.2/5 subvariants in the upper and lower respiratory tract. *eBioMedicine* 2024: 99.

16.   Carolin A, Bishop CR, Grubor-Bauk B, Plummer MP, Raith E, Barry SC, Hope CM, Nguyen W, Rawle DJ, Suhrbier A. RNA-Seq of Nasopharyngeal Swabs from ICU Patients Infected with SARS-CoV-2 Omicron Sublineages in Adelaide, Australia, 2022–2023. Preprints. Preprints, 2025.

17.   Liang K, Barnett KC, Hsu M, Chou WC, Bais SS, Riebe K, Xie Y, Nguyen TT, Oguin TH, 3rd, Vannella KM, Hewitt SM, Chertow DS, Blasi M, Sempowski GD, Karlsson A, Koller BH, Lenschow DJ, Randell SH, Ting JP. Initiator cell death event induced by SARS-CoV-2 in the human airway epithelium. *Sci Immunol* 2024: 9(97): eadn0178.

18.   Qu P, Evans JP, Faraone J, Zheng YM, Carlin C, Anghelina M, Stevens P, Fernandez S, Jones D, Lozanski G, Panchal A, Saif LJ, Oltz EM, Xu K, Gumina RJ, Liu SL. Distinct Neutralizing Antibody Escape of SARS-CoV-2 Omicron Subvariants BQ.1, BQ.1.1, BA.4.6, BF.7 and BA.2.75.2. *bioRxiv* 2022.

19.   Kimura I, Yamasoba D, Tamura T, Nao N, Suzuki T, Oda Y, Mitoma S, Ito J, Nasser H, Zahradnik J, Uriu K, Fujita S, Kosugi Y, Wang L, Tsuda M, Kishimoto M, Ito H, Suzuki R, Shimizu R, Begum MSTM, Yoshimatsu K, Kimura KT, Sasaki J, Sasaki-Tabata K, Yamamoto Y, Nagamoto T, Kanamune J, Kobiyama K, Asakura H, Nagashima M, Sadamasu K, Yoshimura K, Shirakawa K, Takaori-Kondo A, Kuramochi J, Schreiber G, Ishii KJ, Hashiguchi T, Ikeda T, Saito A, Fukuhara T, Tanaka S, Matsuno K, Sato K. Virological characteristics of the SARS-CoV-2 Omicron BA.2 subvariants, including BA.4 and BA.5. *Cell* 2022: 185(21): 3992-4007.e3916.

20.   Sims AC, Baric RS, Yount B, Burkett SE, Collins PL, Pickles RJ. Severe Acute Respiratory Syndrome Coronavirus Infection of Human Ciliated Airway Epithelia: Role of Ciliated Cells in Viral Spread in the Conducting Airways of the Lungs. *Journal of Virology* 2005: 79(24): 15511-15524.

21. Zhu Y, Chew KY, Wu M, Karawita AC, McCallum G, Steele LE, Yamamoto A,
Labzin LI, Yarlagadda T, Khromykh AA, Wang X, Sng JDJ, Stocks CJ, Xia Y,
Kollmann TR, Martino D, Joensuu M, Meunier FA, Balistreri G, Bielefeldt-Ohmann H,
Bowen AC, Kicic A, Sly PD, Spann KM, Short KR. Ancestral SARS-CoV-2, but not
Omicron, replicates less efficiently in primary pediatric nasal epithelial cells. *PLoS*
Biol 2022: 20(8): e3001728.
22. Amarilla AA, Sng JDJ, Parry R, Deerain JM, Potter JR, Setoh YX, Rawle DJ,
Le TT, Modhiran N, Wang X, Peng NYG, Torres FJ, Pyke A, Harrison JJ, Freney ME,
Liang B, McMillan CLD, Cheung STM, Guevara D, Hardy JM, Bettington M, Muller
DA, Coulibaly F, Moore F, Hall RA, Young PR, Mackenzie JM, Hobson-Peters J,
Suhrbier A, Watterson D, Khromykh AA. A versatile reverse genetics platform for
SARS-CoV-2 and other positive-strand RNA viruses. *Nat Commun* 2021: 12(1):
3431.
23. Parry RH, Shang Y, Sng JDJ, McMillan CLD, Modhiran N, Amarilla AA, Isaacs
A, Liang B, Muller DA, Rawle DJ, Suhrbier A, Anderson S, Kelly S, Watterson D, Burr
LD, Khromykh AA. Vaccine-Induced and Hybrid Immunity Against SARS-CoV-2
Variants of Concern in Two Cohorts in Queensland, Australia (2021-2022). *J Med*
Virol 2025: 97(10): e70642.
24. Stewart R, Yan K, Ellis SA, Bishop CR, Dumenil T, Tang B, Nguyen W,
Larcher T, Parry R, Sng JJ, Khromykh AA, Sullivan RKP, Lor M, Meunier FA, Rawle
DJ, Suhrbier A. SARS-CoV-2 omicron BA.5 and XBB variants have increased
neurotropic potential over BA.1 in K18-hACE2 mice and human brain organoids.
Front Microbiol 2023: 14: 1320856.
25. Amarilla AA, Modhiran N, Setoh YX, Peng NYG, Sng JDJ, Liang B, McMillan
CLD, Freney ME, Cheung STM, Chappell KJ, Khromykh AA, Young PR, Watterson
D. An Optimized High-Throughput Immuno-Plaque Assay for SARS-CoV-2. *Front*
Microbiol 2021: 12: 625136.
26. Isaacs A, Amarilla AA, Aguado J, Modhiran N, Albornoz EA, Baradar AA,
McMillan CLD, Choo JJY, Idris A, Supramaniam A, McMillan NAJ, Muller DA, Young
PR, Woodruff TM, Wolvetang EJ, Chappell KJ, Watterson D. Nucleocapsid Specific
Diagnostics for the Detection of Divergent SARS-CoV-2 Variants. *Front Immunol*
2022: 13: 926262.
27. Slonchak A, Wang X, Aguado J, Sng JDJ, Chaggar H, Freney ME, Yan K,
Torres FJ, Amarilla AA, Balea R, Setoh YX, Peng N, Watterson D, Wolvetang E,
Suhrbier A, Khromykh AA. Zika virus noncoding RNA cooperates with the viral
protein NS5 to inhibit STAT1 phosphorylation and facilitate viral pathogenesis. *Sci*
Adv 2022: 8(48): eadd8095.
28. Slonchak A, Hugo LE, Freney ME, Hall-Mendelin S, Amarilla AA, Torres FJ,
Setoh YX, Peng NYG, Sng JDJ, Hall RA, van den Hurk AF, Devine GJ, Khromykh
AA. Zika virus noncoding RNA suppresses apoptosis and is required for virus
transmission by mosquitoes. *Nat Commun* 2020: 11(1): 2205.
29. Slonchak A, Chaggar H, Aguado J, Wolvetang E, Khromykh AA. Noncoding
RNA of Zika Virus Affects Interplay between Wnt-Signaling and Pro-Apoptotic
Pathways in the Developing Brain Tissue. *Viruses* 2023: 15(5).
30. Shaker MR, Slonchak A, Al-Mhanawi B, Morrison SD, Sng JDJ, Cooper-White
J, Khromykh AA, Wolvetang EJ. Choroid plexus defects in Down syndrome brain
organoids enhance neurotropism of SARS-CoV-2. *Sci Adv* 2024: 10(23): eadj4735.