Genomic evidence for co-evolution and sporadic host shifts in leishmaniaviruses

Senne Heeren[1,2,3,*], Lilian Motta Cantanhêde[4,5], Khaled Chourabi[4,5], Mayara Cristhine de Oliveira Santana[4,5], Joon Klaps[2], Alexei Y. Kostygov[6,7], Vyacheslav Yurchenko[6], Philippe Lemey[2], Jean-Claude Dujardin[1,3], Frederik Van den Broeck[1,2,*], Elisa Cupolillo[4,5,*]

[1] Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium
[2] Department of Microbiology, Immunology and Transplantation, Rega Institute for Medical Research, Katholieke Universiteit Leuven, Leuven, Belgium
[3] Department of Biomedical Sciences, University of Antwerp, Antwerp, Belgium
[4] Leishmaniasis Research Laboratory, Oswaldo Cruz Institute, Oswaldo Cruz Foundation, Rio de Janeiro 21040360, Brazil
[5] Instituto Nacional de Ciência e Tecnologia de Epidemiologia da Amazônia Ocidental, INCT EpiAmO, Porto Velho 76812100, Brazil
[6] Life Science Research Centre, Faculty of Science, University of Ostrava, Ostrava, Czechia
[7] Zoological Institute of Russian Academy of Sciences, St. Petersburg, Russia

* Correspondence: Senne Heeren (senne.heeren@kuleuven.be); Frederik Van den Broeck (fvandenbroeck@gmail.com); Elisa Cupolillo (elisa.cupolillo@ioc.fiocruz.br)

FVDB and EC contributed equally as senior author of this manuscript.

Keywords: Co-evolution; Leishmania; Leishmaniavirus; co-phylogenetics

Abstract

The evolutionary relationships between hosts and their symbionts offer valuable insights into the origins, maintenance, and consequences of biological interactions. While co-divergence and host-switching have been extensively explored in systems, such as *Wolbachia*–arthropod symbioses or viruses infecting vertebrates, similar investigations in protistan parasites remain scarce. The *Leishmania–Leishmaniavirus* (LRV) association offers a rare opportunity to study co-divergence in a medically relevant symbiotic system, in which the virus modulates the parasitic disease severity in humans. Here, we used total RNA sequencing to capture both the *Leishmania* transcriptome and the LRV genome simultaneously, enabling the first comprehensive investigation of the co-evolutionary history of *Leishmania* and LRV across different hierarchical levels (subgenus, species and population). We found significant positive correlations between the parasite and viral genetic distances at both the subgenus (R^2 = 0.89; F = 10,913.49; p < 0.001) and species level (focusing on *L. (Viannia)* spp.; R^2 = 0.59; F = 1,254; p < 0.001). This was corroborated by additional co-phylogenetic methods (global-fit and event-based), indicating a strong pattern of phylogenetic congruence between *Leishmania* and LRV up to the species level. Only at the population level, focusing on the interactions of *L. (V.) braziliensis* and *L. (V.) guyanensis* with LRV1, we found weaker co-phylogenetic signals accompanied by more instances of intraspecific host switching. Overall, our findings provide, for the first time, analytical evidence for the co-evolution between *Leishmania* and LRV, with co-speciation as the predominant process, while also shedding light on how such symbioses were maintained over larger time scales.

Introduction

Inter-species interactions are key to ecosystem functioning (e.g. pollination, dispersal, protection, parasitism, predation). In this regard, viruses also play an important role in our biosphere, where they often cause direct harm to their host's populations, sometimes leading to severe socio-economic crises on regional or global scales. Conversely, viruses may also affect populations indirectly through so-called nesting doll ("matryoshka") infections (1–6). In such cases, viruses have established persistent, sometimes mutualistic-like, relationships with organisms such as pathogenic parasites, where they may alter (and often exacerbate) the disease manifestation in humans. This may complicate treatment of the infection and ultimately hinder the disease control in endemic areas (6–9). Even though efforts are currently being made to assess the impact of such nesting-doll infections, little is known about the underlying evolutionary history between pathogenic parasites and their viruses.

To address this, we investigated the co-evolutionary history of an established virus–parasite symbiotic system comprising *Leishmania* parasites (Trypanosomatidae) and their persistent double-stranded (ds) RNA viruses of the genus *Leishmaniavirus* (LRV) (*Pseudototiviridae*, *Duplornaviricota*) (10, 11). In this system, LRV has been found to mutualistically aid its parasitic host by modulating the innate immune response of the mammalian host through an interaction with Toll-like receptor 3 (7), which, in turn, ensures the inhibition of the Nod-like receptor protein 3 inflammasome in the macrophages (12). This immunological cascade ultimately leads to increased parasite survival, disease progression, exacerbation, and an increased likelihood of treatment failure (8, 9, 12, 13). In exchange, the virus' own metabolism and replication are maintained through the help of several host-encoded proteins (e.g., kinases and proteases) (14), but the exact mechanisms involved remain unknown.

In the latest release by the International Committee on the Taxonomy of Viruses (ICTV), the genus *Leishmaniavirus* comprises six species, which are specifically associated with trypanosomatid parasites, including *Leishmania* spp. and *Blechomonas* spp. (2, 10, 11). Within *Leishmania*, different LRV species have different host ranges: *Leishmaniavirus ichi* (further refered to as "LRV1") associated with *Leishmania* (*Viannia*) spp.; *Leishmaniavirus ni* (further referred to as "LRV2") in part associated with *Leishmania* (*Leishmania*) spp. (except *L.* (*L.*) *aethiopica*) and with *Leishmania* (*Sauroleishmania*) spp.; and *Leishmaniavirus sani* associated with *L.* (*L.*) *aethiopica* (2, 11). Among the LRV species of *Blechomonas*, there is *Leishmaniavirus shi* that is associated with *B. maslovi*; *Leishmaniavirus go* with *B. wendygibsoni*; and *Leishmaniavirus roku* with *B. juanalfonzi* (11).

The current LRV classification, along with reports on the virus' host-specificity (2, 15) and the relatively low sequence identity between different LRV species (< 40% at the nucleotide level between LRV1 and LRV2; (10)), resulted in the general assumption of a long standing co-evolutionary history between the parasites and the viruses. This assumption implies that the LRV acquisition pre-dated the *Leishmania* divergence into different subgenera (at least involving *L.* (*Viannia*), *L.* (*Leishmania*), and *L.* (*Sauroleishmania*)). Widmer and Dooley (1995) were the first to propose this hypothesis based on an analysis of genetic distances between nucleotide sequences, which highlighted the mirroring patterns of the LRV1-LRV2 sequence divergence and the heterogeneity between Old and New World *Leishmania* spp. (16). This view of "strict" co-evolution, established through vertical transmission, has been contested several times by reports of horizontal transfers of LRV2 across Old World *Leishmania* spp., and between *Leishmania* and *Blechomonas* (2, 17–20). Conversely, this appears to be in stark contrast with the situation in Central and South America, where interspecific host switches between LRV1 and *Leishmania* (*Viannia*) spp. seem to

occur less frequently (10, 15, 21, 22). At the intraspecific level however, LRV1 host switching
between genetically distinct populations does generally appear to occur at a more frequent pace
(3, 23).

While numerous reports and hypotheses regarding the co-evolutionary history of
Leishmania and LRV exist in the literature (1, 10, 16, 23, 24), a comprehensive evaluation using
(co)-phylogenetic methods is lacking. This is because most studies were focused on phylogenetic
analyses of either the virus or the parasite separately; moreover, they were often based on several
genes or short genomic fragments rather than looking at a genome-wide perspective. Our study
addresses this gap, for the first time, by conducting an in-depth assessment of the virus-parasite
co-evolutionary history across different hierarchical levels (subgenus, species, and population).
The analysis integrates total RNA sequencing of 66 LRV1-positive *Leishmania (Viannia)* isolates
accompanied by the whole genome sequencing of six LRV2-positive *L. (Leishmania)* and *L.*
(Sauroleishmania) isolates, and 77 publicly available genomes of *L. (Viannia)* spp. (n = 49), LRV1
(n = 22) and LRV2 (n = 6).

Based on the current knowledge of the LRV-*Leishmania* association, we hypothesized to
find strong patterns of phylogenetic congruence (i.e. mirroring host-symbiont phylogenies; (25)) at
the subgeneric level (i.e., among *L.* (*Viannia*), *L.* (*Leishmania*) and *L.* (*Sauroleishmania*)) and
species level (i.e., within *L.* (*Viannia*)). We also expected a more general co-phylogenetic signal
(i.e. more closely related hosts tending to associate with more closely related symbionts; (25)) at
the population level of a single species. The latter would be characterized by a relatively higher
number of host transfers. Broadening our understanding of the (co-)evolutionary history of this
epidemiologically important symbiosis will shed new light on how this mutualism has stood the test
of time, while also providing insights into the interplay between viruses and (pathogenic) protists in
general.
111

Results

Expanding the known genomic diversity of LRV1 in South America

Starting from 159 *Leishmania* (*Viannia*) spp. isolates, PCR-screening revealed 66 LRV1-positive isolates (Tab. 1; S. Fig. 1; S. Tab. 1). These were subjected to total RNA sequencing to extract both the viral genomes as well as the parasite transcriptomes. Viral genomes were reconstructed based on a de novo assembly approach as implemented in a novel genome reconstruction pipeline for viral sequencing data, Viralmetagenome (26). With this pipeline we were able to reconstruct viral genomes from 49 isolates of which 42 were of high-quality for subsequent (co)-phylogenetic analyses (S. Tab. 2). These 42 isolates originated from four Amazonian states of Brazil (Acre: n = 4; Amazonas: n = 21; Pará: n = 6; Rondônia: n = 10) and one Venezuelan state (Trujillo: n = 1) which were sampled between 1985 and 2017. The remaining seven isolates were not retained for subsequent analyses as these reconstructed genomes were either too short in length (IOCL2398, IOCL3007, IOCL3515, IOCL3654), or harbored a higher number of SNPs when the sequencing reads were remapped to their respective assemblies (IOCL3562, IOCl3563, IOCL3567) (S. Tab. 2). For one isolate in particular, IOCL3567, the pipeline yielded three genomes of 4,667 – 5,185 bp with a median coverage of 370x – 372x, but with a high number of SNPs (>600). This suggests infection by at least two LRV1 variants, a phenomenon previously documented in *Leishmania* carrying that virus (3, 27). However, we were unable to further disentangle these chimeric-like assemblies and discarded this isolate from downstream analyses. The same was done for another 17 isolates for which the assembly pipeline did not yield properly reconstructed LRV1 genomes.

The 42 high-quality assemblies covered 67.5% to 99.5% of the LRV1-4 genome (5,284 bp; SRA NC_003601) serving as a reference. The percentage of sequencing reads mapping back to the assemblies ranged from 0.0002% to 0.027% (median = 0.004%). The median depths of coverage for the assemblies were between 101x and 1,093x, with an overall median of 391x (S. Tab. 2). The analysis of variants revealed a median of 2 heterozygous SNPs (average = 8; min = 0; max = 84) and no homozygous SNPs in any of the samples, demonstrating the reliability of the reconstructed genomes. All 42 genomes resembled the typical arrangement of LRV1, with two overlapping open reading frames coding for a capsid protein (CP; 2,999 bp) and an RNA-dependent RNA polymerase (RDRP; 2,637 bp) which were near-completely covered in most cases (S. Tab. 2).

Phylogenetic reconstruction and calculation of pairwise genetic distances between the 42 assembled genomes revealed 14 highly supported viral clades (Fig. 1a) exhibiting relatively low, within-clade, genetic distances (<0.125 substitutions/site; S. Tab. 3). The clades were named according to their host species with Roman numerals (i.e., LgI-V, LbI-V, LsI, LnI-II) (Fig. 1). Analysis of geographic distribution revealed that, among the studied isolates, those from the Amazonas belonged to eight clades, three of which being unique to this location (LgIII, LnI, LsI) (Fig. 1b). Other clades found in a single state included LbII and LbIV in Acre; LbI and LbIII in Rondônia; and LgVI and LnII in Pará. The remaining clades were found in two or three different neighboring states. The single LRV1 genome of Venezuelan origin belonged to the LgII-clade, which was mainly identified in Amazonas and Acre. However, the current sample size impeded our ability to infer the connection of *Leishmania* and LRV1 between Trujillo and the Western Brazilian states.

Our new panel of LRV1 genomes provides a major expansion of the known genomic diversity of the virus. This becomes clear from the phylogenetic reconstruction, combining our panel with 64 publicly available genomes. Most clades characterized above had not been identified earlier, whereas a few (LgII, LgV, LgVI, LbII, LbV) clustered with previously described ones (3, 27)

(Fig. 1c). The concomitant inference of the ancestral geographic origins (Fig. 1c, internal nodes)
along with the clear host specificity of LRV1 towards the different *Leishmania* species suggests
spatial divergence and subsequent spread of each species-specific clade. One exception to this
observed host-specificity is LRV1-Ln-IOCL3574, which clusters with viruses associated with *L.*
guyanensis (Fig. 1c). The corresponding *Leishmania* isolate was originally typed as *L. naiffi*, but
our whole transcriptome data indicated that it is a mixture both genetic material of *L. naiffi* and *L.*
guyanensis (see below).

Phylogenetic reconstructions based on the single-gene alignments (i.e. CP and RDRP)
were similar compared to the clade delimitation derived from the whole-genome data (Fig. 1b; S.
Fig. 2). Moreover, co-phylogenetic global-fit tests (Robinson-Foulds distance, ParaFit and PACo)
revealed a consistently strong co-phylogenetic signal between the different LRV1 phylogenies (i.e.
genome-wide, CP and RDRP) (S. Tab. 5; S. Fig. 2). The only striking inconsistency was the internal
position of *L. guyanensis*-associated LRV1 from French Guiana (clades: LgV, LgVI, LsI, A*, B*, D*,
E* (except M92355.1, origin: Suriname)). Based on both the CP and genome-wide phylogenies,
this clade (S. Fig. 2, clade shown in red) appears to share a common ancestor with the other *L.*
guyanensis-associated LRV1 genomes, whereas it formed a monophyletic group with the *L.*
braziliensis-associated LRV1 within the RDRP phylogeny. Nevertheless, this inconsistency does
not refute the observed LRV1 host-specificity, it merely indicates the uncertainty of the clade
divergence (i.e., low bootstrap supports) at these ancestral nodes based on the two single gene
alignments (S. Fig. 2). In addition, none of the three alignments showed any indication of
recombination, based on the pairwise homoplasy index test (whole genome: *p*-value = 1.0; CP: *p*-
value = 0.95; RDRP: *p*-value = 1.0).

Tab. 1: LRV1 prevalence across all 159 screened *Leishmania* (*Viannia*) spp.

Species	LRV1 prevalence
L. braziliensis	27.85% (22/79)
L. guyanensis	75.61% (31/41)
L. lainsoni	16.67% (1/6)
L. lindenbergi	0% (0/6)
L. naiffi	61.11% (11/18)
L. panamensis	0% (0/4)
L. shawi	33.33% (1/3)
L. utingensis	0% (0/1)
L. (*Viannia*) *spp.*	0% (0/1)
Total	**41.51% (66/159)**

Fig. 1: Lineage diversity and distribution of the LRV1 genomic diversity. a Unrooted maximum likelihood phylogenetic tree based on 42 near-whole genome LRV1 sequences. Branch values represent bootstrap support values based on 10,000 ultra-fast approximated bootstrap replicates. The scale bar depicts the number of substitutions per site. Clades were delineated based on high bootstrap support and low (within-clade) pairwise genetic distances (S. Tab. 3). Colors are set for visual purposes and do not serve any information on the ancestral states (i.e. internal nodes). The abbreviations Lb, Lg, Ln and Ls refer to the *Leishmania* hosts of the LRVs: *L. braziliensis*, *L. guyanensis*, *L. naiffi* and *L. shawi*, respectively. **b** Administrative map of South America showing the geographic origin of the 42 LRV1 genomes (dark gray area) along with geolocalized LRV1 genomes reported previously (3, 27). Administrative country- and state-level data originate from: http://www.diva-gis.org/Data. **c** Midpoint rooted maximum clade credibility tree based on the 42 newly obtained near-whole LRV1 genomes along with 64 publicly available ones. Branch support values (bold faced) represent the posterior probabilities. Nodes are colored according to their inferred geographic origin, with posterior probabilities shown adjacently in regular font. Tip labels are colored according to the viral lineages (from this study or previous ones). Bars beside the tree represent the newly or previously described viral lineages (matching the tip labels) and the associated *Leishmania* species. Single and double asterisks mark the lineages identified by Tirera et al. (2017) (27) and Heeren et al. (2023) (3), respectively.

Transcriptomic diversity of the *Leishmania* (*Viannia*) subgenus

Total RNA sequencing of the LRV1-positive *Leishmania* isolates allowed us, not only to assemble LRV1 genomes, but also to genotype the *Leishmania* isolates themselves based on the transcriptomic data. Similar to our previous whole-genome sequencing studies (28, 29), we compiled a dataset of 115 isolates containing: (i) genomic data from 66 isolates, generated for this work; and (ii) 49 publicly available sequences from various *Leishmania* spp. that served as reference strains during genotyping and species delimitation. This reference panel contained isolates from the *Leishmania* (*Viannia*) *braziliensis* clade (*Lb* L1: n = 18; *Lb* L2: n = 3; *Lb* L3: n = 2; *Lb* L4: n = 2; *L. peruviana*: n = 2), the *Leishmania* (*V.*) *guyanensis* species complex (*L. guyanensis*: n = 3; *L. panamensis*: n = 14; *L. shawi*: n = 1), *Leishmania* (*V.*) *naiffi* (n = 3), and *Leishmania* (*V.*) *lainsoni* (n = 1) (S. Tab. 6) (28).

As our data was derived from RNA sequencing, our genotyping analyses focused on the coding DNA sequence (CDS) regions. Mapping the reads against the long-read-based M2904 *L. braziliensis* reference genome (30) resulted in a median coverage of 17x (min = 1x; max = 151x) across the entire CDS for all 115 isolates. For the isolates from this study alone, the median CDS coverage was 10.5x (min = 1x; max = 43x). Considering a minimum CDS read depth of 10x, we excluded 31 isolates (30 from this study and one from a previous study) from downstream analyses and filtered out 1,184 individual CDS regions below this threshold in the retained sampels. The final genotype dataset comprised 6,299 CDS regions across 84 isolates, yielding 157,649 high-quality SNPs.

Based on the differences in the number of SNPs (S. Tab. 8) between the included *Leishmania* (*Viannia*) spp. (Chi-squared = 76.92; df = 10; *p* = 2.01e-12; pairwise Dunn tests: S. Tab. 9), all isolates, with the exception of IOCL3539 and IOCL3574, could be identified (*L. braziliensis* L1: n = 12; *L. guyanensis*: n = 18; *L. naiffi*: n = 4) (Fig. 2a). The latter two isolates harbored distinctively higher numbers of heterozygous SNPs compared to all other isolates (Fig. 2a; S. Tab. 8) and showed reticulation in the CDS-wide phylogenetic network, a pattern suggestive of recombination and potential hybrid ancestry Fig. 2b.

However, the alternate allele read depth frequency (ARDF) of these two isolates revealed uncommon profiles. In IOCL3539, the ARDF distribution was unimodal and skewed with a mode around 0.66, compared to the typical symmetric distribution around 0.5 (31) (Fig. 2c). IOCL3574, on the other hand, showed a distribution that was bimodal with modes near 0.2 and 0.8 (in contrast to the typical modes for triploid genomes at 0.33 and 0.66 (31)) (Fig. 2d). These CDS-wide shifts in the ARDF are strong indicators that the observed hybrid-like patterns are the result of either (i) extensive mosaic aneuploidy, or (ii) mixed infections by two different *Leishmania* spp., rather than true interspecific hybrids (32).

Fig. 2: Species/lineage delimitation of 36 *Leishmania* isolates based on whole transcriptome data (157,649 bi-allelic SNPs). a Quantification of homozygous (x-axis) and heterozygous (y-axis) SNPs for 36 isolates in this study and 48 reference isolates (incl. the *L. braziliensis* clade, the *L. guyanensis* species complex, *L. naiffi* and *L. lainsoni*). **b** Phylogenetic network of all 84 isolates revealing the reticulate pattern caused by IOCL3539 and IOCL3574, indicative of hybrid ancestry. **c-d** Shifted CDS-wide alternate allele frequency distribution plots of IOCL3539 (**c**) and IOCL3574 (**d**). Dashed vertical lines show the typical modalities for diploid (i.e. a unimodal distribution around 0.5) and triploid (i.e. a bimodal distribution around 0.33 and 0.66) genomes (31).

Evidence for phylogenetic congruence across different hierarchical levels of *Leishmania* and LRV

To assess the extent to which *Leishmania* and LRV have shared a similar evolutionary trajectory, we performed a matrix regression on the pairwise genetic distances of both symbionts, combined with formal co-phylogenetic tests – global-fit and event-based – to detect overall co-phylogenetic signal and evaluate true phylogenetic congruence, respectively (25). These analyses were based on a dataset including 22 Peruvian *Leishmania* (*V.*) *braziliensis* isolates with their associated LRV1 genomes (3), as well as six Old World *Leishmania* isolates (*L.* (*L.*) *major* (n = 4), *L.* (*S.*) *tarentolae* (n = 1) and *L.* (*S.*) *adleri* (n = 1)) with their respective LRV2 genomes (S. Tab. 4,6).

On the subgeneric level, we observed a highly significant, strong positive correlation between pairwise genetic distances across the 51 *Leishmania*-LRV pairs that were analyzed (MMRR: R^2 = 0.89; F = 10,913.49; p < 0.001) (Fig. 3a). In parallel, our array of global-fit analyses detected a consistently significant co-phylogenetic signal between *Leishmania* and LRV at this level (Fig. 3b; S. Tab. 10). Similarly, the event-based inference revealed strong signatures of phylogenetic congruence, suggesting that the two symbionts share a highly similar phylogenetic history (Fig. 3c). This latter analysis was conducted using nine different event-costs sets (S. Fig. 3a; S. Tab. 11), with eight of them yielding significant co-phylogenetic reconciliations, where well-supported cospeciation events predominated the divergence of Old and New World *Leishmania* subgenera with LRV2 and LRV1, respectively (S. Fig. 4).

When focusing on the species level (subgenus *L.* (*Viannia*)) and its association with LRV1, we observed a highly significant, yet more moderate positive correlation between the parasite and viral pairwise genetic distances (MMRR: R^2 = 0.59; F = 1,254; p < 0.001). Interestingly, the pronounced host specificity did not appear to be shaped by spatial separation, as viral genetic distances did not correlate with geographic distances among the isolates (MMRR: R^2 = 0.005; F = 5.28; p = 0.135). In parallel, both the global-fit and event-based methods consistently detected strong patterns of co-phylogenetic signal and phylogenetic congruence, respectively (S. Tab. 10,12; S. Fig. 5). More specifically, the event-based analysis yielded a diverse set of possible co-phylogenetic histories based under six different event-cost scenarios (S. Fig. 3b; S. Tab. 12). Strikingly, in all cases the inferred phylogenetic reconciliations were significant, and thus supporting the presence of strong phylogenetic congruence between the members of the subgenus *Viannia* and LRV1 (S. Fig. 6). Another notable observation was that co-speciation events consistently accounted for species divergence (S. Tab. 12) and were well supported, whereas duplication and transfer events appeared to more predominant within the separate species partitions (S. Fig. 6).

Finally, zooming in towards the population level of this virus-parasite co-evolutionary history, we focused on *L. (V.) braziliensis* and *L. (V.) guyanensis* with their respective LRV1 lineages (too small sample size for *L.* (*V.*) *naiffi*). This also revealed signatures of co-evolution, albeit generally weaker compared to the higher levels and occasionally lacking statistical significance. For instance, we found no significant correlation between the genetic distances of *L. braziliensis* and LRV1 (MMRR: R^2 = 0.027; F = 10.65; p = 0.215) whilst a weak, though significant, correlation was observed in the case of *L. guyanensis* (MMRR: R^2 = 0.349; F = 47.68; p = 0.002). Surprisingly, both global-fit and event-based analyses revealed significant co-phylogenetic signals and evidence of phylogenetic congruence between each *Leishmania* species and its associated LRV1 lineage (S. Fig. 3c-d, 7-9; S. Tab. 10, 13).

Fig. 3: Co-phylogenetic reconciliation of *Leishmania*-LRV symbiosis at the subgeneric level. a

Regression scatterplot of the pairwise genetic distances of *Leishmania* and LRV (51 pairs). Yellow points
represent summarized values in 200 bins along the x-axis. **b** Tanglegram of *Leishmania* and LRV phylogenies.
The *Leishmania* phylogeny was inferred from six concatenated gene sequences (S. Tab. 15; see methods).
The LRV phylogeny was inferred from concatenated sequences of CP and RDRP (ORF2 and ORF3,
respectively). Colors on the tree are set for visual purposes and do not serve any information on the ancestral
states. Branch values represent bootstrap support values based on 10,000 ultra-fast approximated bootstrap
replicates. The scale bar indicates the number of substitutions per site. **c** Event-based *Leishmania*-LRV co-
phylogeny at the subgeneric level (51 tip-pairs). The reconciliation was inferred with the following event-costs:
duplication = 1.50; transfer = 1.50; loss = 1.00 (S. Tab. 11; set 03). Support values for the inferred co-evolution
events (e.g. co-speciation, host transfer) were estimated from 100 bootstrap replicates. Colored bars adjacent
to the tip labels denote the LRV-host species as depicted in panel b.

Discussion

By harnessing total RNA sequencing of cultivated LRV-positive *Leishmania* isolates – enabling the capture of both the parasite transcriptome and (near)-complete viral genome – our study provides the first comprehensive evaluation of the virus-parasite (co)-phylogenetic history of this medically relevant symbiosis across the subgenus, species and population levels.

We first investigated the genomic diversity of each symbiont separately to position the newly sampled genomes within the known phylogenetic context of *Leishmania* and LRV, based on publicly available genomic data. For *Leishmania*, all but two isolates (IOCL3539, IOCL3574) clustered clearly with a single species – either the *L.* (*V.*) *braziliensis* clade, the *L.* (*V.*) *guyanensis* species complex, or *L.* (*V.*) *naiffi* – as judged by homozygous SNP counts and unambiguous phylogenetic placement. Conversely, the reticulate pattern in the phylogenetic network and high heterozygosity in IOCL3539 and IOCL3574 suggested either (i) interspecific hybridization (*L. guyanensis* x *L. braziliensis* L4 and *L. guyanensis* x *L. naiffi*, respectively) or (ii) a mixture of these species. Hybridization (both inter- and intraspecific) is an increasingly recognized phenomenon in parasitic protozoa (33–35) and *Leishmania* in particular (3, 28–30, 36–39). This has important epidemiological implications, making detection and reporting of new hybrids crucial within the context of genomic surveillance (3, 33, 40). Analysis of the alternate allele read depth frequencies at heterozygous sites revealed shifts from the expected uni- and bimodal patterns suggesting that the genomic signatures of these isolates are more likely the result of a mixed infection or extensive mosaic aneuploidy, rather than being true hybrid signals (31, 32). However, these isolates apeared to be devoid of distinct homozygous regions for the alternate allele and showed heterozygous streches across the entire genome – contrasting with other recently reported hybrid signatures (28, 32) – which strongly suggests that both isolates are mixtures of two *Leishmania* species rather than true genomic hybrids.

The phylogenetic reconstruction of the newly assembled LRV1 genomes substantially expanded the known genomic diversity of the virus. This included (i) an increased intra-clade diversity within groups previously described, based on whole-genome data (3, 27); and (ii) the identification of novel clades (e.g., LgI, LgIII, LgIV, LbI, LbIII, LbIV, LsI, LnI, LnII). In concordance with Santana et al. (2023) (15), we observed strict host specificity among LRV1 genomes, independent of sampling location (state/department). This was evidenced by a strong correlation of virus genetic distances with those of their host, but not with geographic distances. Such clear host specificity is already a good indicator of virus-parasite co-divergence at the species level within the *Leishmania* (*Viannia*) subgenus, with the only exceptions being the likely mixed isolates.

Although earlier studies have suggested a long-standing association between *Leishmania* and LRV, potentially involving co-divergence between and within subgenera (1, 10, 16, 23, 24), these hypotheses have not been formally tested. Most previous work focused on separate phylogenetic analyses of virus and host, often based on a few genes. Each *Leishmania* subgenus – except *L.* (*Mundinia*), which harbors a phylogenetically remote *Shilevirus* (*Leishbuviridae*, *Negarnaviricota*) (41, 42) – has its own LRV species. For LRV1, a strong pattern of host specificity and phylogenetic congruence has been observed. In contrast, LRV2 shows a more flexible pattern, with partial specificity: certain viral clusters are more frequently associated with particular *Leishmania* species, but not in a strictly co-specific manner.

The main goal of our study was to formally investigate, for the first time, the co-evolutionary
history of *Leishmania* spp. and their endosymbiotic viruses across the subgeneric, species and
population level. Overall, we revealed clear genomic evidence of virus-parasite co-divergence
across these levels. At the population level of *L.* (*V.*) *braziliensis* and *L.* (*V.*) *guyanensis*,
phylogenetic congruences decreased (e.g. due to lower signals or loss of statistical significance).
Here, we observed more instances of host-switching, as reported previously in *L. braziliensis* (3),
relative to the predominance of co-speciation events at the higher hierarchical levels. These results
align well with the seemingly mutualistic nature of the virus-parasite symbiosis, where strong
phylogenetic congruence has often been observed (43). Conversely, this view of virus-parasite co-
divergence is challenged by cases of horizontal transmission of LRV within *Leishmania* (both inter-
and intraspecific) (3, 18–23) and to other trypanosomatids, such as *Blechomonas* spp. (17). These
horizontal transfers are likely facilitated by the potential of LRV to exploit the parasite's exosomal
pathway (44–47). Nevertheless, our findings indicate that host switching is relatively rare and
occurs mainly at the intraspecific level, whereas vertical transmission remains the predominant
factor shaping the evolutionary trajectory of this symbiosis (2, 10).

This study expands our current understanding of the co-evolutionary dynamics of
Leishmania and LRV, even though it does not elucidate their full evolutionary history. A major
limitation is the discrepancy between the amount of paired genomic data between Old and New
World regions. Although our analyses revealed strong signatures of phylogenetic congruence at
the subgeneric level, these patterns warrant validation by means of larger sample sizes of Old
World isolates. Another unresolved question concerns the absence of LRV in the divergent
subgenus *Mundinia,* where it is still unknown whether LRV-presence is a plesio- (ancestral) or
apomorphic (derived) trait. In other words, LRV could have been either lost in *L.* (*Mundinia*) or
gained after its divergence from the most recent common ancestor of the other three subgenera
(*Leishmania, Viannia*, and *Sauroleishmania*) (10). Elucidating the broader virus-parasite interplay
is also becoming increasingly complex as more LRV species are described and recognized. One
additional viral species is known within *Leishmania*, *Leishmaniavirus sani* which is associated to *L.*
(*L.*) *aethiopica*, and three divergent species have been described from the flea-parasitic
trypanosomatid genus *Blechomonas*: *Leishmaniavirus shi* in *B. maslovi*, *Leishmaniavirus go* in *B.*
wendygibsoni, and *Leishmaniavirus roku* in *B. juanalfonzi* (11, 17). Moreover, LRV1 is not the sole
RNA virus infecting *L.* (*Viannia*) spp. (48). Accounting for this growing co-evolutionary and
taxonomic complexity, it is likely that additional, more cryptic, viral diversity remains undetected in
these protistan parasites. For each of these virus-parasite interactions, It will therefore be essential
to evaluate and monitor the consequences for the mammalian host (incl. humans) infected with
such virus-positive parasites.

Another point to consider is that even though our co-phylogenetic analyses focused on
LRV1-positive *Leishmania* (*Viannia*) isolates, including LRV1-negative strains could yield additional
insight into the host-virus evolutionary dynamics. The absence of LRV1 in certain parasite-lineages,
e.g., those from the Atlantic region of Brazil (49–51), may reflect historical events of viral loss,
resistance to infection, or ecological constraints limiting viral maintenance or transmission.
Incorporating these isolates into the host phylogeny could help determine whether LRV1 absence
is randomly distributed or associated with specific clades, thereby reinforcing or challenging the
current co-phylogenetic patterns. While these isolates cannot directly contribute to host–virus tree
congruence analyses due to the lack of viral sequence data, independent analyses of their
phylogenetic position and geographic distribution may shed light on the broader context of LRV1
evolution and its relationship to the diversification of *Leishmania*.

For this study, we applied a less conventional sequencing approach which proved to be a
useful and cost-efficient way to simultanuously sequence the parasite transcriptome and the LRV
genome. Even though the resulting phylogenetic patterns were consistent with those derived from
standard genome sequencing datasets for both parasite and virus, our approach had some
limitations. In particular, it was not always possible to fully reconstruct viral genomes due to the low
fraction of LRV reads. Next to that, the transcriptome read depth from the parasite data was
sometimes insufficient for making robust phylogenetic and population genomic inferences. These
issues could be mitigated either by deeper sequencing, with a potentially unjustified additional cost,
or reducing the ribosomal RNA content in the sequencing libraries. Nevertheless, for
epidemiologically relevant systems such as *Leishmania*-LRV, a strategy that captures data for both
symbionts in parallel represents a meaningful advancement for molecular surveilance.

In conclusion, our total RNA sequencing data allowed us to simultaneously perform
phylogenetic analyses on *Leishmania* isolates, via transcriptome-based genotyping, and their
respective LRV strains. This effort represents the first explicit reconstruction of the virus-parasite
co-evolutionary history, characterized by strong *Leishmania*-LRV co-divergence between and
within three parasite subgenera, with the most frequent host-switches occuring at the intraspecific
level. Although the virus-parasite interplay has not yet been fully disentangled, this work provides
fundamental biological insight in the evolution of the so-called matryoshka infections where – to
quote Hartley et al. (2012) – ultimately the host will pay the toll (52).
410

Materials and Methods

Parasite culturing & RNA sequencing

Leishmania isolates were cultured in vitro, screened for LRV1, and prepared for total RNA sequencing at the Fundação Oswaldo Cruz (Fiocruz). Parasites were grown in liquid culture medium to a concentration of 10^7–10^8 cells/ml, collected through centrifugation (on the third day after the final subculturing step), and preserved in DNA/RNA Shield (Zymo Research). Total RNA was extracted using the PureLink RNA Mini Kit (Invitrogen), following the manufacturer's instructions. All isolates were screened for the presence of LRV1 through PCR using distinct LRV1 primers: LRV F–5′- ATGCCTAAGAGTTTGGATTCG-3′ and LRV R–5′-ACAACCAGACGATTGCTGTG-3′ (53). The screening included a positive (*L. guyanensis* MHOM/BR/1975/M4147) and a negative control (*L. braziliensis* MHOM/BR/1975/M2903) (54–56). LRV1-positive isolates (n = 66) were sent for total RNA sequencing without polyA selection or rRNA depletion to make sure the LRV genome would not be lost in the sample. Samples were sequenced on either the NovaSeq (n = 56) or DNBSeq (n = 10) platform at Genewiz (Azenta Life Sciences) and BGI Genomics, respectively. The latter ten samples were part of a pilot study to test sequencing approach for capturing both the LRV genome and the parasite transcriptome. In both cases, paired-end sequences were generated to ~30,000,000 150bp reads.

Leishmaniavirus de novo assembly & phylogenetic reconstruction

Prior to the de novo assembly of viral genomes, raw reads from each of the 66 LRV-positive *L. Viannia* sequenced libraries were filtered and trimmed using fastp v.0.23.4 (57), and mapped against the *L. braziliensis* M2904 reference genome using SMALT v.0.7.6 (available at: https://www.sanger.ac.uk/tool/smalt-0/). Only unmapped reads were retained and processed with the novel bioinformatics workflow Viralmetagenome v.0.1.2 (available at: https://github.com/nf-core/viralmetagenome) for de novo reconstructions of viral genomes (26). In summary, with this workflow we combined Megahit and Spades assemblies, followed by a BLAST search to extract contigs matching publicly available LRV1 genomes (58–60). Next, all "raw" assemblies were then iteratively refined by re-mapping reads to the assembled genomes and subsequent variant and consensus calling, making use of BCFtools and iVar (61, 62). The reconstruction of hybrid consensus genomes was skipped (--skip_hybrid_consensus: true). A detailed overview of the parameter settings is provided in a supplementary table (S. Tab. 14). Viralmetagenome was able to resolve a consensus genome for 49 of the 66 LRV1-positive isolates. Additionally, we conducted a post-hoc assessment of the assembly quality which involved re-mapping of the reads against their respective assemblies, and variant calling. In total, 42 assemblies were retained based on the following criteria: length > 3,000bp, read depth > 100x, and low SNP counts (i.e., <100 heterozygous and zero homozygous SNPs). Finally, whole-genome sequences, as well as sequences for the Capsid Protein (CP) and the RNA-dependent RNA Polymerase (RDRP), were aligned using the L-INS-I algorithm, as implemented in MAFFT v.7.490 ('--localpair' option) (63).

Phylogenetic reconstructions of LRV1 were generated using IQtree2 v2.3.6 (64) over 10,000 bootstrap replicates using the ultrafast bootstrap approximation method (65). Substitution models were automaticlly selected by IQtree's ModelFinder algorithm (-m MFP) (66) which selects the best substitution model based on the Bayesian Information Criterion (BIC) after evaluating all models that are implemented in IQtree. Whole genome maximum likelihood phylogenies of the assembled LRV1 genomes were generated with and without a set of 64 publicly available genomes (S. Tab. 4) using GTR+F+I+R4 and GTR+F+I+G4 as respective substitution models. Similarly, gene specific phylogenies, including the 64 publicly available sequences, were inferred using the

GTR+F+I+R4 substitution model for both CP and RDRP. For the clade delimitation of the 42 newly
assembled genomes, we also calculated the pairwise genetic distances using the 'dist.dna' function
of the R-package ape v5.8-1 (67) (model = 'raw').

In addition to regular phylogenetic reconstructions, we also attempted to reconstruct
ancestral states across the entire LRV1 phylogeny, focusing on the virus' geographic origin and
associated *Leishmania* species. This was achieved by adopting a Bayesian phylogenetic
framework making use of the BEAST software package (v.1.10.4) (68) and the BEAGLE likelihood
calculation library (69). Consistent with previous analyses (3), no temporal signal was detected in
the data. Therefore, all sampling dates were set to zero, after which LRV1's discrete
phylogeography was inferred without time calibration, using the default strict molecular clock
model). The nucleotide substitution model was set to GTR (General Time Reversible) with
estimated base frequencies and the site heterogeneity model set to a Gamma distribution with 4
discrete rate categories and accounting for invariable sites (70–72). The tree prior model was set
to a speciation model, the Yule process (73), starting the analysis with a random starting tree
(default setting). Two discrete traits were added: the country of origin and associated *Leishmania*
species. For both traits we selected the symmetric discrete trait substitution model along with the
BSSVS option (Bayesian Stochastic Search Variable Selection procedure) in order to reconstruct
their ancstral states. Prior and operator settings were kept at their default settings. The MCMC
chain was run for ten million states, sampled every 1,000 states and providing a 10% burn-in.
Throughout the inference proces, convergence of parameter estimates was evaluated using
BEAST's auxiliary program Tracer v.1.7.2 (74). Upon completion of the runs, all effective sample
sizes (ESS) of all parameters exceeded the recommended threshold of 200, except for the C-T
transition rate in the GTR model, which reached an ESS of 110. Finally, the posterior tree
distribution was summarized as a maximum clade credibility (MCC) tree using BEAST's
TreeAnnotator.

Transcriptomic genotyping & phylogenetic reconstruction of *Leishmania* (*Viannia*) isolates

Total RNA sequencing reads of the 66 LRV1-positive *Leishmania* (*Viannia*) isolates were
mapped against the *L. braziliensis* M2904 v.65 (MHOM/BR/75/M2904) reference genome
(available at: https://tritrypdb.org/common/downloads/release-65/LbraziliensisMHOMBR75M2904
_2019) using SMALT v0.7.6 (parameters: k = 13, s = 2). Reads with a mapping quality lower than
25 were removed using SAMtools v.1.21 ('-q' option) and read depths across all positions were
extracted ('samtools depth -a') (61). This reference genome has been shown to work well for
genotyping not only *L. braziliensis* isolates, but also other *L.* (*Viannia*) species, resulting in highly
accessible genomes for downstream analyses (28-30). Because transcriptomic data were used
rather than genomic data, we discarded isolates with low median coverages (<10x) across the
coding DNA sequences (CDS) from downstream analyses. Additionally, all CDS regions, in each
sample, with a coverage below 10x were omitted, ensuring that only transcriptomic regions with
sufficient depth were included in the subsequent analyses. Variant calling (i.e., SNPs and INDELs)
was caried out using GATK v.4.1.4.1 (75). Here, the initial calling was performed on a per-sample
basis using HaplotypeCaller, after which all individual genotype vcf files were combined using
CombineGVCFs, and variants were jointly genotyped with GenotypeGVCFs. Next, SNPs and
INDELs were separated using SelectVariants after which both panels were subjected to hard
filtering with VariantFiltration. For SNPs, the following filters were applied: (i) QD < 2.0, FS > 60.0,
MQ < 40.0, MQRankSum < −12.5 and ReadPosRankSum < −8.0, following GATK's
recommendations (76); (ii) GQ < 10; and (iii) QUAL < 100. For INDELs, GATK's recommendations
were adopted: QD < 2.0, FS > 200.0, or ReadPosRankSum < −20.0 (76).

A phylogenetic network was generated using SplitsTree v.4.17.0 (77) based on the
pairwise uncorrected *p*-distances calculated across 157,649 bi-allelic SNPs. This network was built
using the NeighborNet (78) and EqualAngle (79) algorithms as implemented in SplitsTree. Forty-
nine publicly available isolates were included as reference sequences: members of the *L.* (*V.*)
braziliensis clade, the *L.* (*V.*) *guyanensis* species complex, as well as *L.* (*V.*) *naiffi* and *L.* (*V.*)
lainsoni (S. Tab. 6). In addition, the alternate allele read depth frequency (ARDF) distributions at
heterozygous sites were calculated and visualized to identify whether potential hybrid isolates were
the result from a mixed infection. This was done using a custom python script (vcf2freq.py; available
at: https://github.com/FreBio/mytools). Differences in SNP counts between the different species
were compared using a non-parametric Kruskal-Wallis (stats R-package; (80)) test, followed by
pairwise Dunn's tests with additional Benjamini-Hochberg *p*-value correction for multiple testing
(FSA R-package; (81)).

Preparing alignments and phylogenies for co-phylogenetic reconciliation

Phylogenetic reconciliations of the LRV-*Leishmania* symbiosis were performed at multiple
levels: subgenus (combining Old World and New World species), species (focusing on the South
American *Leishmania* (*Viannia*) species), population (focusing on *L.* (*V.*) *braziliensis* and *L.* (*V.*)
guyanensis with their respective LRV1 lineages) and gene level (reconciling the CP and RDRP
gene phylogenies within LRV1).

For the subgenus level analysis, we included 22 publicly available Peruvian *L.* (*V.*)
braziliensis isolates and six Old World *Leishmania* isolates positive for LRV2 (S. Tab. 4,6). Instead
of using the entire genome sequence, we concatenated the annotated CP and RDRP genes (i.e.,
ORFs 2 and 3 in the LRV genome) and aligned their sequences from LRV1 and LRV2 using the
MAFFT v.7.490 as described above (63). The LRV phylogeny was reconstructed using IQtree2
with 10,000 ultra-fast bootstrap replicates and the GTR+F+I+R3 substitution model based on the
ModelFinder algorithm (64–66). For the maximum likelihood phylogenetic reconstruction of the
Leishmania phylogeny, which included highly divergent species from different subgenera, a
concatenated alignment of six orthologous protein coding genes was used (S. Tab. 15), following
Coughlan et al. (2017) (82). The sequences of the six target genes were obtained as follows. Reads
were trimmed with fastp v.0.23.4 using similar settings used in Heeren et al. (2023): a minimum
base quality (-q) of 30; maximum percentage of unqualified bases per read (-u) to 10; quality-based
sliding window trimming front-to-tail (-5) and tail-to-front (-3) with a window size (-W) of 1 and mean
quality score (-M) of 30; right-end cutting (--cut-right) using 10 bp windows (--
cut_right_window_size) with mean quality score (--cut_right_mean_quality) below 30; and retention
of reads between 100 (-I) and 150 bp (-b) (3). Trimmed reads were then assembled de novo using
Megahit v.1.2.9 (58). Contigs with six target genes were identified using a blastn (60) search and
subsequently extracted. These sequences were combined with 222 publicly available ones
(accession numbers listed in S. Tab. 15) and aligned using MAFFT, as described above. Finally, a
phylogenetic tree was inferred using IQtree2 with 10,000 ultra-fast bootstrap replicates and
TIM2+F+G4 set as optimal substitution model (64–66).

For the reconstruction of the parasite phylogenies at the species- and population-level co-
phylogenetic analyses, we used concatenated SNP-alignments. Variants were called against the
MHOM/BR/75/M2904 reference genome, as described above, resulting in genome-wide alignment
containing 97,530 bi-allelic SNPs. From this panel, a phylogenetic tree was inferred using IQtree2
with 10,000 ultra-fast bootstrap replicates and the TVM+F+R2 as the most optimal substitution
model, correcting for ascertainment bias (+ASC) (64–66). For the population-level analyses of *L.*

braziliensis–LRV1 and *L. guyanensis*–LRV1, species specific partitions of the inferred *L.* (*Viannia*)
phylogeny were used. Finally, for the reconciliation of the LRV1 CP and RDRP genes we made
use of the reconstructed phylogenies as described above.

Genetic distance regression

551 In order to obtain an initial assessment of similarity between the viral and parasite
552 phylogenies, we calculated pairwise genetic distances from the corresponding concatenated
553 sequences using the 'dist.dna' function of the ape R-package (v.5.0) (67) (model = 'K80'). These
554 distances were converted into distance matrices, and analyzed using Multiple Matrix Regression
555 with Randomization (MMRR) (83) where *p*-values were estimated over 1,000 permutations.

Global-fit co-phylogenetic analyses

We tested for the presence of co-phylogenetic signals (i.e., testing whether closely related
hosts tend to associate with closely related symbionts) between the parasite and virus phylogenies
by making use of three global-fit methods: (i) the Robinson-Foulds distance as implemented in the
phytools R-package (84); (ii) ParaFit as implemented in ape (67); and (iii) PACo as implemented
in the paco R-package (85). For each method, both virus and parasite phylogenies were
downsampled using the 'drop.tip' function of ape (67). This way, only the isolates for which both
high quality LRV1 genomes and *L.* (*Viannia*) transcriptome-based genotypes were available. Each
approach was performed by running 1,000 permutations for p-value estimation and the Cailliez
correction method was used in ParaFit and PACo to correct for negative eigenvalues.

Event-based co-phylogenetic analysis

To test whether *Leishmania* and LRV exhibit signs of phylogenetic congruence, at one or
multiple hierarchical levels, we made use of eMPRess, an event-based phylogenetic reconciliation
tool that for reconciling phylogenies based on a set of events such as co-speciation, loss,
duplication and host-switch (25, 86–88). A key advantage of eMPRess over other event-based
reconciliation tools, is its ability to specify different sets of event-costs prior to calculating the
reconciliation, allowing the user to evaluate alternative evolutionary scenarios for obtaining more
robust inferences (86). This is important as different event-costs schemes can lead to distinct
solutions and interpretations. For each level, we visualized the cost-space (S. Fig, 7) and selected
multiple sets of event-costs covering most of the cost-space (S. Tab. 11-13). Phylogenetic
reconciliations were then calculated for each set, and a median MPR (i.e. a median value of the
MPR solution space; MPR = Maximum Parsimony reconciliation) was visualized. Significance was
assessed using 1,000 random permutations. Visualizations of the cost-spaces and reconciled
phylogenies were generated using the eMPRess GUI, while significance testing was conducted
with the CLI version ('*p*-value' function with '--n-samples 1000' option).

Acknowledgments

This work received financial support from the Directie-Generaal Ontwikkelingssamenwerking en
Humanitaire Hulp (DGD) (Belgian cooperation). F.V.d.B. and S.H. acknowledge support from the
Research Foundation Flanders (grants 1226120N and G092921N). S.H. additionally acknowledges
support from KU Leuven's Internal Funds (PDMt1/25/007). E.C. acknowledges support from
Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro: CNE -
204.165/2024; 269.891/2021; ColBio - 210.285/202; Conselho Nacional de Desenvolvimento
Científico e Tecnológico: Research Fellow - 309627/2021-4; PROEP/IOC - 441639/2024-0; INCT-
EpiAmo: 465657/2014-1; and the GGBN award: GGI-GGBN-2021-278. L.M.C. acknowledges
support from Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro,

Pós-Doutorado; Nota 10 E-26/205.730/2022 and 205.731/2022. E.C., L.M.C, M.C.d.O.S. and K.C.
also acknowledge support from the Coordenação de Aperfeiçoamento de Pessoal de Nível
Superior – Brasil (CAPES) – Finance Code 001.

Data availability

595 Sequencing data (i.e., raw sequencing reads and the the reconstructed LRV genomes), that were
596 generated for this study, will be deposited to publicly accessible databases upon publication of the
597 manuscript.

Key-words

- Co-evolution
- *Leishmania*
- *Leishmaniavirus*
- (co-)phylogenetics

References

1.	D. Grybchuk, A. Y. Kostygov, D. H. Macedo, C. M. d'Avila-Levy, V. Yurchenko, RNA viruses in trypanosomatid parasites: a historical overview. *Mem. Inst. Oswaldo Cruz* **113**, e170487 (2018).

2.	D. Klocek, *et al.*, Evolution of RNA viruses in trypanosomatids: new insights from the analysis of Sauroleishmania. *Parasitol. Res.* (2023). https://doi.org/10.1007/s00436-023-07928-x.

3.	S. Heeren, *et al.*, Diversity and dissemination of viruses in pathogenic protozoa. *Nat. Commun.* **14**, 8343 (2023).

4.	P. Gupta, *et al.*, A parasite odyssey: An RNA virus concealed in Toxoplasma gondii. *Virus Evol.* **10**, veae040 (2024).

5.	J. Charon, *et al.*, Novel RNA viruses associated with Plasmodium vivax in human malaria and Leucocytozoon parasites in avian disease. *PLoS Pathog.* **15**, e1008216 (2019).

6.	A. Ibañez-Escribano, *et al.*, Microbial matryoshka: Addressing the relationship between pathogenic flagellated protozoans and their RNA viral endosymbionts (family Totiviridae). *Vet. Sci.* **11**, 321 (2024).

7.	A. Ives, *et al.*, Leishmania RNA virus controls the severity of mucocutaneous leishmaniasis. *Science* **331**, 775–778 (2011).

8.	V. Adaui, *et al.*, Association of the Endobiont Double-Stranded RNA Virus LRV1 With Treatment Failure for Human Leishmaniasis Caused by Leishmania braziliensis in Peru and Bolivia. *J. Infect. Dis.* **213**, 112–121 (2016).

9.	E. Bourreau, *et al.*, Presence of Leishmania RNA Virus 1 in Leishmania guyanensis Increases the Risk of First-Line Treatment Failure and Symptomatic Relapse. *J. Infect. Dis.* **213**, 105–111 (2016).

10.	L. M. Cantanhêde, *et al.*, The Maze Pathway of Coevolution: A Critical Review over the Leishmania and Its Endosymbiotic History. *Genes* **12** (2021).

11.	P. Simmonds, *et al.*, Changes to virus taxonomy and the ICTV Statutes ratified by the International Committee on Taxonomy of Viruses (2024). *Arch. Virol.* **169**, 236 (2024).

12.	R. V. H. de Carvalho, *et al.*, Leishmania RNA virus exacerbates Leishmaniasis by subverting innate immunity via TLR3-mediated NLRP3 inflammasome inhibition. *Nat. Commun.* **10**, 5273 (2019).

13.	F. A. Pazmiño, *et al.*, Mucosal leishmaniasis is associated with the Leishmania RNA virus and inappropriate cutaneous leishmaniasis treatment. *PLoS One* **20**, e0317221 (2025).

14.	B. I. Hillman, A. B. Cohen, "Totiviruses (Totiviridae)" in *Encyclopedia of Virology (Fourth Edition)*, D. H. Bamford, M. Zuckerman, Eds. (Academic Press, 2021), pp. 648–657.

15.	M. C. de O. Santana, K. Chourabi, L. M. Cantanhêde, E. Cupolillo, Exploring Host-Specificity: Untangling the Relationship between Leishmania (Viannia) Species and Its Endosymbiont Leishmania RNA Virus 1. *Microorganisms* **11** (2023).

16.	G. Widmer, S. Dooley, Phylogenetic analysis of Leishmania RNA virus and Leishmania suggests ancient virus-parasite association. *Nucleic Acids Res.* **23**, 2300–2304 (1995).

17.	D. Grybchuk, *et al.*, RNA Viruses in Blechomonas (Trypanosomatidae) and Evolution of Leishmaniavirus. *MBio* **9** (2018).

18. H. Hajjaran, *et al.*, Detection and molecular identification of leishmania RNA virus (LRV)
in Iranian Leishmania species. *Arch. Virol.* **161**, 3385–3390 (2016).

19. R. Saberi, *et al.*, Presence and diversity of Leishmania RNA virus in an old zoonotic
cutaneous leishmaniasis focus, northeastern Iran: haplotype and phylogenetic based approach.
Int. J. Infect. Dis. **101**, 6–13 (2020).

20. M. Nalçacı, *et al.*, Detection of Leishmania RNA virus 2 in Leishmania species from
Turkey. *Trans. R. Soc. Trop. Med. Hyg.* **113**, 410–417 (2019).

21. R. Kariyawasam, R. Lau, B. M. Valencia, A. Llanos-Cuentas, A. K. Boggild, Novel
detection of Leishmania RNA virus-1 (LRV-1) in clinical isolates of Leishmania Viannia
panamensis. *Parasitology* **151**, 151–156 (2024).

22. A. A. Bonilla, *et al.*, Epidemiology and Genetic Characterization of Leishmania RNA Virus
in Leishmania (Viannia) spp. Isolates from Cutaneous Leishmaniasis Endemic Areas in Panama.
Microorganisms **12**, 1317 (2024).

23. A. Y. Kostygov, *et al.*, Analyses of Leishmania-LRV Co-Phylogenetic Patterns and
Evolutionary Variability of Viral Proteins. *Viruses* **13** (2021).

24. S. M. Scheffter, Y. T. Ro, I. K. Chung, J. L. Patterson, The complete sequence of
Leishmania RNA virus LRV2-1, a virus of an Old World parasite strain. *Virology* **212**, 84–90
(1995).

25. B. Perez-Lamarque, H. Morlon, Distinguishing cophylogenetic signal from phylogenetic
congruence clarifies the interplay between evolutionary history and species interactions. *Syst.*
Biol. (2024). https://doi.org/10.1093/sysbio/syae013.

26. J. Klaps, P. Lemey, nf-core community, L. E. Kafetzopoulou, Nf-core/viralmetagenome: A
novel pipeline for untargeted viral genome reconstruction. *bioRxiv* 2025.06.27.661954 (2025).

27. S. Tirera, *et al.*, Unraveling the genetic diversity and phylogeny of Leishmania RNA virus
1 strains of infected Leishmania isolates circulating in French Guiana. *PLoS Negl. Trop. Dis.* **11**,
e0005764 (2017).

28. F. Van den Broeck, *et al.*, Genome Analysis of Triploid Hybrid Leishmania Parasite from
the Neotropics. *Emerg. Infect. Dis.* **29**, 1076–1078 (2023).

29. S. Heeren, *et al.*, Evolutionary genomics of Leishmania braziliensis across the
neotropical realm. *Commun. Biol.* **7**, 1587 (2024).

30. F. Van den Broeck, *et al.*, Ecological divergence and hybridization of Neotropical
Leishmania parasites. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 25159–25168 (2020).

31. M. B. Rogers, *et al.*, Chromosome and gene copy number variation allow major structural
change between species and strains of Leishmania. *Genome Res.* **21**, 2129–2142 (2011).

32. E. Tihon, H. Imamura, J.-C. Dujardin, J. Van Den Abbeele, F. Van den Broeck, Discovery
and genomic analyses of hybridization between divergent lineages of Trypanosoma congolense,
causative agent of Animal African Trypanosomiasis. *Mol. Ecol.* **26**, 6524–6538 (2017).

33. K. C. King, R. B. Stelkens, J. P. Webster, D. F. Smith, M. A. Brockhurst, Hybridization in
Parasites: Consequences for Adaptive Evolution, Pathogenesis, and Public Health in a Changing
World. *PLoS Pathog.* **11**, e1005098 (2015).

34. W. Gibson, *et al.*, Genetic recombination between human and animal parasites creates
novel strains of human pathogen. *PLoS Negl. Trop. Dis.* **9**, e0003665 (2015).

35. A. Izeta-Alberdi, C. N. Ibarra-Cerdeña, D. A. Moo-Llanes, J. M. Ramsey, Geographical,
landscape and host associations of Trypanosoma cruzi DTUs and lineages. *Parasit. Vectors* **9**,
631 (2016).

36. N. S. Akopyants, *et al.*, Demonstration of genetic exchange during cyclical development
of Leishmania in the sand fly vector. *Science* **324**, 265–268 (2009).

37. M. B. Rogers, *et al.*, Genomic confirmation of hybridisation and recent inbreeding in a
vector-isolated Leishmania population. *PLoS Genet.* **10**, e1004092 (2014).

38. E. Inbar, *et al.*, Whole genome sequencing of experimental hybrids supports meiosis-like
sexual recombination in Leishmania. *PLoS Genet.* **15**, 1–28 (2019).

39. A. Hadermann, *et al.*, Genome diversity of Leishmania aethiopica. *Front. Cell. Infect.*
Microbiol. **13** (2023).

40. M. A. Domagalska, J.-C. Dujardin, Next-generation molecular surveillance of TriTryp
diseases. *Trends Parasitol.* **36**, 356–367 (2020).

41. D. Grybchuk, *et al.*, The First Non-LRV RNA Virus in Leishmania. *Viruses* **12** (2020).

42. H. R. Hughes, *et al.*, Summary of taxonomy changes ratified by the International
Committee on Taxonomy of Viruses (ICTV) from the Animal dsRNA and ssRNA(-) Viruses
Subcommittee, 2025. *J. Gen. Virol.* **106** (2025).

43. A. Hayward, R. Poulin, S. Nakagawa, A broadscale analysis of host-symbiont
cophylogeny reveals the drivers of phylogenetic congruence. *Ecol. Lett.* **24**, 1681–1696 (2021).

44. V. D. Atayde, *et al.*, Exploitation of the Leishmania exosomal pathway by Leishmania
RNA virus 1. *Nat Microbiol* **4**, 714–723 (2019).

45. A. Lafleur, M. Olivier, Viral endosymbiotic infection of protozoan parasites: How it
influences the development of cutaneous leishmaniasis. *PLoS Pathog.* **18**, e1010910 (2022).

46. M. Olivier, D. S. Zamboni, Leishmania Viannia guyanensis, LRV1 virus and extracellular
vesicles: a dangerous trio influencing the faith of immune response during muco-cutaneous
leishmaniasis. *Curr. Opin. Immunol.* **66**, 108–113 (2020).

47. G. Dong, V. Wagner, A. Minguez-Menendez, C. Fernandez-Prada, M. Olivier,
Extracellular vesicles and leishmaniasis: Current knowledge and promising avenues for future
development. *Mol. Immunol.* **135**, 73–83 (2021).

48. A. Y. Kostygov, *et al.*, A novel strain of Leishmania braziliensis harbors not a toti- but a
bunyavirus. *PLoS Negl. Trop. Dis.* **18**, e0012767 (2024).

49. L. Guilbride, P. J. Myler, K. Stuart, Distribution and sequence divergence of LRV1 viruses
among different Leishmania species. *Mol. Biochem. Parasitol.* **54**, 101–104 (1992).

50. L. de O. R. Pereira, *et al.*, Severity of tegumentary leishmaniasis is not exclusively
associated with Leishmania RNA virus 1 infection in Brazil. *Mem. Inst. Oswaldo Cruz* **108**, 665–
667 (2013).

51. D. H. Macedo, *et al.*, Low frequency of LRV1 in Leishmania braziliensis strains isolated
from typical and atypical lesions in the State of Minas Gerais, Brazil. *Mol. Biochem. Parasitol.*
210, 50–54 (2016).

52. M.-A. Hartley, C. Ronet, H. Zangger, S. M. Beverley, N. Fasel, Leishmania RNA virus:
when the host pays the toll. *Front. Cell. Infect. Microbiol.* **2**, 99 (2012).

53.   L. M. Cantanhêde, *et al.*, Further evidence of an association between the presence of
Leishmania RNA virus 1 and the mucosal manifestations in tegumentary leishmaniasis patients.
PLoS Negl. Trop. Dis. **9**, e0004079 (2015).

54.   A. Zakharova, *et al.*, Leishmania guyanensis M4147 as a new LRV1-bearing model
parasite: Phosphatidate phosphatase 2-like protein controls cell cycle progression and
intracellular lipid content. *PLoS Negl. Trop. Dis.* **16**, e0010510 (2022).

55.   P. I. Tarr, *et al.*, LR1: a candidate RNA virus of Leishmania. *Proc. Natl. Acad. Sci. U. S.*
A. **85**, 9572–9575 (1988).

56.   P. Ranque, "Etude morphologique et biologique de quelques Trypanosomatides récoltés
au Sénégal," Université de Aix-Marseilles. (1973).

57.   S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

58.   D. Li, C.-M. Liu, R. Luo, K. Sadakane, T.-W. Lam, MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics **31**, 1674–1676 (2015).

59.   A. Prjibelski, D. Antipov, D. Meleshko, A. Lapidus, A. Korobeynikov, Using SPAdes DE
Novo Assembler. *Curr. Protoc. Bioinformatics* **70**, e102 (2020).

60.   S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment
search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

61.   P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10** (2021).

62.   N. D. Grubaugh, *et al.*, An amplicon-based sequencing framework for accurately
measuring intrahost virus diversity using PrimalSeq and iVar. *Genome Biol.* **20**, 8 (2019).

63.   K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

64.   B. Q. Minh, *et al.*, IQ-TREE 2: New models and efficient methods for phylogenetic
inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

65.   B. Q. Minh, M. A. T. Nguyen, A. von Haeseler, Ultrafast approximation for phylogenetic
bootstrap. *Mol. Biol. Evol.* **30**, 1188–1195 (2013).

66.   S. Kalyaanamoorthy, B. Q. Minh, T. K. F. Wong, A. von Haeseler, L. S. Jermiin,
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–
589 (2017).

67.   E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526–528 (2018).

68.   M. A. Suchard, *et al.*, Bayesian phylogenetic and phylodynamic data integration using
BEAST 1.10. *Virus Evol* **4**, vey016 (2018).

69.   D. L. Ayres, *et al.*, BEAGLE: an application programming interface and high-performance
computing library for statistical phylogenetics. *Syst. Biol.* **61**, 170–173 (2012).

70.   S. Tavaré, Some probabilistic and statistical problems in the analysis of DNA sequences.
Lecture of mathematics for life science (1986).

71.   Z. Yang, Maximum likelihood phylogenetic estimation from DNA sequences with variable
rates over sites: approximate methods. *J. Mol. Evol.* **39**, 306–314 (1994).

72.   P. Lemey, A. Rambaut, A. J. Drummond, M. A. Suchard, Bayesian phylogeography finds
its roots. *PLoS Comput. Biol.* **5**, e1000520 (2009).

73.   T. Gernhard, The conditioned reconstructed process. *J. Theor. Biol.* **253**, 769–778
(2008).

74.   A. Rambaut, A. J. Drummond, D. Xie, G. Baele, M. A. Suchard, Posterior summarization
in Bayesian phylogenetics using tracer 1.7. *Syst. Biol.* **67**, 901–904 (2018).

75.   G. A. Van der Auwera, B. D. OConnor, *Genomics in the Cloud Using Docker, GATK, and*
WDL in Terra (First edition) (O'Reilly Media, 2020).

76.   G. A. Van der Auwera, *et al.*, From FastQ data to high confidence variant calls: the
Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* **43**, 11.10.1–
11.10.33 (2013).

77.   D. H. Huson, D. Bryant, Application of phylogenetic networks in evolutionary studies. *Mol.*
Biol. Evol. **23**, 254–267 (2006).

78.   D. Bryant, V. Moulton, NeighborNet: An Agglomerative Method for the Construction of
Planar Phylogenetic Networks in *Algorithms in Bioinformatics*, (Springer Berlin Heidelberg, 2002),
pp. 375–391.

79.   A. W. M. Dress, D. H. Huson, Constructing splits graphs. *IEEE/ACM Trans. Comput. Biol.*
Bioinform. **1**, 109–115 (2004).

80.   R Core Team, R: A Language and Environment for Statistical Computing. [Preprint]
(2024). Available at: https://www.R-project.org/.

81.   D. H. Ogle, J. C. Doll, A. P. Wheeler, A. Dinno, FSA: Simple fisheries stock assessment
methods. R package version 0.9. 4. [Preprint] (2023).

82.   S. Coughlan, *et al.*, The genome of Leishmania adleri from a mammalian host highlights
chromosome fission in Sauroleishmania. *Sci. Rep.* **7**, 43747 (2017).

83.   I. J. Wang, Examining the full effects of landscape heterogeneity on spatial genetic
variation: a multiple matrix regression approach for quantifying geographic and ecological
isolation. *Evolution* **67**, 3403–3411 (2013).

84.   L. J. Revell, phytools 2.0: an updated R ecosystem for phylogenetic comparative
methods (and other things). *PeerJ* [Preprint] (2024). Available at:
http://dx.doi.org/10.7717/peerj.16505.

85.   J. A. Balbuena, T. Poisot, M. Hutchinson, F. C. S. P. O. B. et al 2013
<https://doi.org/10.1371/journal.pone.0061048>, paco: Procrustes Application to Cophylogenetic
Analysis. [Preprint] (2020). Available at: https://CRAN.R-project.org/package=paco.

86.   S. Santichaivekin, *et al.*, eMPRess: a systematic cophylogeny reconciliation tool.
Bioinformatics **37**, 2481–2482 (2021).

87.   W. Dismukes, M. P. Braga, D. H. Hembry, T. A. Heath, M. J. Landis, Cophylogenetic
Methods to Untangle the Evolutionary History of Ecological Interactions. *Annu. Rev. Ecol. Evol.*
Syst. (2022). https://doi.org/10.1146/annurev-ecolsys-102320-112823.

88.   D. M. de Vienne, *et al.*, Cospeciation vs host-shift speciation: methods for testing,
evidence from natural associations and relation to coevolution. *New Phytol.* **198**, 347–385 (2013).

 Supplementary Figures

813
814 **S. Fig. 1.** Spatial distribution of LRV1-positive and negative *Leishmania* isolates. **a** Spatial distribution of all
815 isolates used in this study along with geolocalized *Leishmania* isolates from Heeren et al. (2023) (3) and Tirera
816 et al. (2017) (24). **b-d** Spatial distribution of specific *Leishmania* species: *L. braziliensis* (**b**), *L. guyanensis* (**c**)
817 and *L. naiffi* (**d**). Administrative country- and state-level data originate from: http://www.diva-gis.org/Data.

S. Fig. 2: Tanglegram of the LRV1 capsid protein (CP) and RNA-dependent RNA polymerase (RDRP) gene phylogenies. The clade colored in red occupies different positions in the CP and RDRP phylogenies. Branch values represent bootstrap support values based on 10,000 ultra-fast approximated bootstrap replicates. The scale bar depicts the number of substitutions per site. Links between the trees are colored according to the identified clades of the 42 newly assembled LRV1 genomes (Figure 1; Suppl Figure 1).

1   **S. Fig. 3:** Event-cost distributions and cost spaces from event-based co-phylogenetic analyses of the *Leishmania*–LRV symbiosis across the different hierarchical
2   levels. Per panel, the left-side plot depicts the density of the number of events that have been calculated over all the tested cost-sets; (ii) the right-side plot indicates
3   the costspace as calculated by eMPRess. Each colored zone corresponds to the same number of events and reconciliation solutions. **a** Subgenus level. **b** Species
4   level (within *L.* (*Viannia*)). Note the high number of inferred loss events in this analysis, resulting from an adjustmen of the default loss cost from 1 to 0 to enable co-
5   phylogenetic reconciliations in eMPRess. **c** Population level for *L.* (*V.*) *braziliensis*. **d** Population level for *L.* (*V.*) *guyanensis*.

2
3
4 **S. Fig. 4:** Event-based co-phylogenies of the *Leishmania* divergence, across different subgenera (*L.* (*Viannia*),
5 *L.* (*Leishmania*), *L.* (*Sauroleishmania*)), and LRV species (LRV1, LRV2). Each reconciliation was calculated
6 with different sets of event-costs (S. Fig. 3a; S. Tab. 11): **A)** set 1, **B)** set 2, **C)** set 3, **D)** set 4, **E)** set 5, **F)** set
7 6, **G)** set 7. Support values for the inferred events were based on 100 bootstrap replicates.

8
S. Fig. 5: Tanglegram of the *Leishmania Viannia* subgenus and LRV1 phylogenies based on 45 LRV1-positive
L. Viannia isolates. The *Leishmania* phylogeny was inferred from an alignment of 97,530 concatenated SNPs,
corrected for ascertainment bias. The LRV1 phylogeny was inferred from a concatenation of the two protein-
coding gene sequences, CP and RDRP (ORF2 and 3, respectively). Colors on the tree are set for visual
purposes and do not serve any information on the ancestral states (i.e. internal nodes). Branch values
represent bootstrap support values from 10,000 ultra-fast approximated bootstrap replicates. The scale bar
depicts the number of substitutions per site.
16

18
19 **S. Fig. 6:** Event-based co-phylogenies of *Leishmania Viannia* and LRV1. Each reconciliation was calculated
20 using different sets of event-costs (S. Fig. 3b;S. Tab. 12): **A)** set 1, **B)** set 2, **C)** set 3, **D)** set 4, **E)** set 5, **F)**
21 set 6. Support values for the inferred events were estimated from 100 bootstrap replicates.

22
23

S. Fig. 7: Species-specific tanglegrams of *Leishmania* (*V.*) *braziliensis* (**a**) and *L.* (*V.*) *guyanensis* (**b**) and their
associated LRV1 lineages. The *Leishmania* phylogenies represent species-specific partitions of the *L.*
(*Viannia*) SNP-based phylogeny (Suppl. Figure 5). Similarly, the LRV1 phylogenies represent specific
partitions of the LRV1 phylogeny (Suppl. Figure 5). Branch values represent bootstrap support values from
10,000 ultra-fast approximated bootstrap replicates. The scale bar depicts the number of substitutions per
site.

30
31 **S. Fig. 8:** Event-based co-phylogenies of *Leishmania* (*V.*) *braziliensis* and LRV1. Each reconciliation was
32 calculated from different sets of event-costs (S. Fig. 3c; S. Tab. 13): **A)** set 1, **B)** set 2, **C)** set 3, **D)** set 4, **E)**
33 set 5. Support values for the inferred events were estimated from 100 bootstrap replicates.

S. Fig. 9: Event-based co-phylogenies of *Leishmania* (*V.*) *guyanensis* and LRV1. Each reconciliation was calculated using different sets of event-costs (S. Fig. 3d; S. Tab. 13): **A)** set 1, **B)** set 2, **C)** set 3, **D)** set 4, **E)** set 5. Support values for the inferred events were estimated from 100 bootstrap replicates.

a

LRV pairwise genetic distance

Leishmania pairwise genetic distance

b

Leishmania spp.

Leishmaniavirus spp.

L. (S.) tarentolae
L. (S.) adleri
L. (L.) major
L. (V.) naiffi
L. (V.) braziliensis
L. (V.) guyanensis

LRV2
LRV1

c

● Cospeciation
■ Host transfer / switch
- - - Loss