**MicroRNA166-HD-ZIP III module impacts tuber shape, color and productivity**
**in potato**

Nikita Sunil Patil[1], Arati Vasav[1], Jyoti Kumari[1], Gourav Arora[1,2], Bhavani
Natarajan[1,3] and Anjan K. Banerjee[1*]

[1] Biology Division, Indian Institute of Science Education and Research (IISER) Pune,
Dr. Homi Bhabha Road, Maharashtra, India
[2] Department of Plant Developmental Biology, Max Planck Institute for Plant
Breeding Research, Cologne, Germany
[3] Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge
CB2 3EA, UK

* Corresponding author
Prof. Anjan K. Banerjee
E-mail: akb@iiserpune.ac.in
Tel: +91-20-25908057

**Summary**

MicroRNA165/166 is a highly conserved and abundant microRNA family in plants

that targets class III homeodomain leucine zipper (*HD-ZIP III*) transcription factors.

Their essential roles in plant development, organ polarity, meristem maintenance, and

vascular differentiation have been well established. However, their role in storage

organ development, particularly in potato tuberization remains unknown. Here, we

demonstrate the pleiotropic effects of *miR166* on potato (*Solanum tuberosum* spp.

*Andigena*) growth and development. Suppression of *miR166* via target mimicry

(MIM166) resulted in reduced plant height, altered vascular patterning, diminished

leaf curvature, and impaired root development. Remarkably, MIM166 lines produced

elongated, pigmented tubers with reduced yield under short-day conditions.

Transcriptomic analyses of swollen head *versus* stalk of tuberizing stolons revealed

altered expression of auxin-, cytokinin-, and gibberellin-related genes, consistent with

hormone quantification data, suggesting the key role of phytohormones in governing

tuber shape. Anthocyanin profiling indicated elevated accumulation of cyanidin 3-

glucoside, pelargonidin 3-glucoside, and delphinidin 3-glucoside in MIM166 tuber

peels, correlating with the upregulation of anthocyanin biosynthetic genes. Tuber

productivity in these lines was reduced possibly due to the decreased expression of

*SWEET11b* and distorted vascular structures in the MIM166 lines. Further,

*StREVOLUTA*, a direct miR166 target, exhibited dynamic expression during stolon

differentiation and could modulate the expression of auxin biosynthesis genes.

Collectively, our results demonstrate that the *miR166-HD-ZIP III* module is

functional in potato, plays conserved roles in regulating key developmental traits, and

is active through early stolon-to-tuber transition stages, affecting tuber shape, color

and productivity.

**Significance statement**

The microRNA166–HD-ZIP III regulatory module plays a central role in plant growth

and development. However, its function in tuber/ storage organ development remains

unexplored. Here, we show that the miR166–HD-ZIP III has distinct regulatory roles

in potato, influencing tuber shape, pigmentation, and yield. These findings reveal

previously unrecognized functions of the miR166–HD-ZIP III module in tuber crops

and broaden our understanding of miR166-mediated developmental regulation in

plants.

**Introduction**

MicroRNA 165/166 (miR165/166), a highly conserved family of microRNAs

(miRNAs), play important roles in regulating a wide range of physiological processes

in plants. They target specific transcription factors (TFs)- members of the

*Homeodomain-Leucine Zipper Class III* (*HD-ZIP III*) gene family, that are shown to

critically regulate processes such as embryo patterning, development of leaf, roots,

vasculature and apical meristems, and establishment of organ polarity, making the

*miR166-HD-ZIP III* module central to plant growth and morphogenesis (Turchi *et al.*,

2015; Ramachandran *et al.*, 2017; Roodbarkelari., 2017). *miR166* is also involved in

mediating plant responses to various abiotic stresses such as drought, salinity, and

temperature fluctuations (Yadav *et al.*, 2021; Wang *et al.*, 2025; Wei *et al.*, 2025).

Several studies have demonstrated the potential of *miR166* as an important candidate

in agricultural biotechnology, wherein its manipulation could improve crop resilience

and productivity (Boualem *et al.*, 2008; Zhang *et al.*, 2018; Zhao *et al.*, 2022).

*miR165/166* and their target genes are extensively studied in *Arabidopsis*, and

to some extent in crop plants such as rice, maize, tomato, peach, soybean, tea, and

legumes (Boualem *et al.*, 2008; Valiollahi *et al.*, 2014; Zhang *et al.,* 2015; Ding *et al.*,

2018; Zhang *et al.*, 2018; Li *et al.*, 2020; Zhao *et al.*, 2022; Zhou *et al.*, 2024). In

*Arabidopsis*, *miR165* (miR165a and miR165b) and *miR166* (miR166a-miR166g)

target five genes that encode HD-ZIP III TFs- *PHABULOSA* (*PHB*), *PHAVOLUTA*

(*PHV*), *REVOLUTA* (*REV*), *ATHB-8*, and *ATHB-15/CORONA* (*CRN*)/*INCURVATA4*

(*ICU4*). These TFs are closely related, and are known to play both overlapping and

antagonistic roles in various biological processes (Prigge *et al.*, 2005).

Overexpression of *miR165* causes a drastic reduction in all five *HD-ZIP III* genes in

*Arabidopsis*, recapitulating the phenotypes caused by loss-of-function mutations of

HD-ZIP III, such as distorted shoot apical meristem (SAM), altered organ polarity,

and defects in the development of vascular tissues and interfascicular fibers (Zhou *et*

*al.*, 2007). In tomato, HOMEOBOX15A (SlHB15A) acts as a key regulator for

preventing fruit set in the absence of fertilization. *SlHB15A* loss-of-function produces

parthenocarpic fruits under both optimal and heat stress conditions due to aberrant

ovule development (Clepet *et al.*, 2021). In soybean, knockdown of *miR166*

(*GmSTTM166*) causes increased expression of *ATHB14-LIKE*, reduces plant height, and lowers the levels of bioactive gibberellic acid (GA$_3$). ATHB14-LIKE directly represses transcription of *GmGA1* and *GmGA2* (GA biosynthesis genes), while activates the transcription of the GA catabolic gene *GmGA2ox2* (Zhao *et al.*, 2022). Thus, *miR166* plays a pivotal role in crop plants as it regulates key agronomic traits such as plant height, fruit set, and stress responses that ultimately affect yield.

Potato tuberization is a complex process governed by several molecular signals such as small RNAs, transcription factors, proteins, phytohormones, and sucrose. The mobile signals such as *StBEL5*, StSP6A, StPTB1, StPTB6 and others are systematically transported from leaves to belowground stolons under inductive conditions to initiate tuber development (Hannapel *et al.*, 2017). Overexpression of *miR156* causes development of aerial tubers from axillary meristems while overexpression of *miR172* abolishes SD-dependency of tuberization (Martin *et al.*, 2009; Bhogale *et al.*, 2014). Further, a phasiRNA *siRD29(-)* is shown to fine-tune *StGA3ox3* levels during stolon-to-tuber transitions (Malankar *et al.*, 2023). In potato, the miR166 family is amongst the most abundantly expressed miRNAs, with a total transcript per million exceeding 600,000, and exhibits tissue-specific and stolon-to-tuber stage-specific expression profiles (Lakhotia *et al.*, 2014; Kondhare *et al.*, 2018). During further analysis of the small RNA sequencing data from Kondhare *et al.* (2018), we found *miR166* to be differentially accumulated under long-day (LD; 16 h light/8 h dark; tuber non-inductive) and short-day (SD; 8 h light/16 h dark; tuber inductive) photoperiod. This observation prompted us to investigate the role of *miR166* in potato development, and tuberization in particular. Despite the diverse roles attributed to *miR166* in several plants' development, productivity, and stress responses, its functional characterization in potato and relevance in storage organ development remains unexplored. Here, we validated the functional role of the *miR166–HD-ZIP III* module in potato. Our findings show that suppression of *miR166* leads to pleiotropic effects on overall plant development, highlighting its conserved regulatory function across plants. Importantly, we reveal that *miR166* modulates key agronomic traits in potato- tuber shape, color, and productivity.

**Results**

**Identification, validation and expression analysis of miR166 in Potato**

All four precursors of microRNA166 from the miRBase (the microRNA database;
http://mirbase.org/) are predicted to form a secondary hairpin loop structure with
mature 166 in its stem region (MFold; Zuker, 2003) (Figure S1A). To validate the
presence of miR166 precursors (precursors a, b, c and d) in potato, RT-PCR was
performed and the amplified products were sequence confirmed (Figure 1A).
Precursor-b was the least abundant form, as compared to the other three precursors.
Compared to LD, the levels of precursors-a and -c decreased in the stolons under SD
photoperiod, while precursor-d increased (Figure 1B-D). The mature forms of all the
precursors are identical, except for pre-miR166b, wherein it differs by a single base
(Figure S1B). In our analysis of small RNA sequencing data from a previous study,
we found miR166 to be the most abundant microRNA in early stolon-to-tuber
transition stages and exhibited differential accumulation under short-day (SD; tuber
inductive) photoperiod (Kondhare *et al*., 2018) (Figure S2A-B). We detected the 21-
bp mature miR166 by stem-loop endpoint PCR from leaves and stolons of WT potato
plants (Figure 1E). The amplified product was verified by sequencing, thus
confirming that mature miR166 is expressed in potato.

To determine whether photoperiod affects the expression/accumulation of
miR166, plants were grown in soil under early long-day (LD; tuber non-inductive)
and SD conditions. Stem-loop quantitative reverse transcription (qRT) -PCR analysis
revealed a reduced accumulation of miR166 in leaves under SD conditions, while
stolons had comparable levels of miR166 under both the photoperiods (Figure 1F).
Stolons differentiating into tubers exhibited 10-40 percent reduction in the later stages
of stolon-to-tuber transition (Figure 1G). *In silico* analysis suggested that miR166 in
potato potentially targets six *HD-ZIP III* members, similar to *Arabidopsis*. Using
modified 5′ RLM-RACE, we mapped the cleavage site on the HD-ZIP transcripts
(Figure 1H), confirming the targets of miR166. Overall, our results validate the
existence of miR166-*HD-ZIP III* module in potato and the miR166 showed tissue-
specific accumulation under photoperiodic conditions.
164

165 **Knockdown of microRNA166 causes pleiotropic phenotypic changes**
166 Using a target mimicry approach, we generated stable transgenic lines of miR166 in *S.*
167 *tuberosum andigena ssp.* (Figure S3A). Of them, 2 independent lines (referred as KD
168 #2 and KD #3) exhibiting 50-75% suppression of mature miR166 were selected for
169 further characterization (Figure 2A-B). miR166 KD lines (MIM166) had reduced

170 shoot height but number of nodes remained unchanged, implying a decreased inter-

171 nodal distance, compared to the WT plants (Figure 2C-E). The leaf lamina from

172 miR166 KD lines appeared flat, hence we calculated the longitudinal and transverse

173 curvature indices ($CI^L$ and $CI^T$) (Liu *et al*., 2010) to quantify the curvature. A

174 curvature index of zero suggests that the leaf surface is completely flat. Both

175 curvature indices, $CI^L$ and $CI^T$ of the KD lines were significantly reduced compared to

176 WT (Figure 2F-G). The total chlorophyll content in the miR166 KD lines were

177 comparable to that of WT leaves; and leaves of KD #2 showed a slight increased

178 anthocyanin accumulation, relative to the WT (Figure S3B). The vascular structures

179 were distorted and had fewer xylem elements compared to the WT plants (Figure 2H).

180 Further, an i*n vitro* root growth assay revealed that the MIM166 lines had decreased

181 root length, shoot and root biomass, number of adventitious and lateral roots (Figure

182 S3C-H). Thus, several phenotypic characters were altered upon constitutive and

183 ubiquitous suppression of miR166 in potato.

185 **Suppression of miR166 alters tuber shape, color and productivity**

186 To examine the effect of miR166 suppression on tuber development, we subjected the

187 miR166 KD lines to tuber inductive SD photoperiod in soil conditions. Compared to

188 WT plants, the KD lines showed reduced tuber productivity in varying degrees,

189 although the number of tubers per plant did not vary significantly (Figure 3A-C).

190 Interestingly, tubers from these KD lines were elongated as opposed to the WT

191 spherical tubers. The sphericity index for these tubers was greater than 1.5, while WT

192 tubers had the sphericity index of around 1.0 (Figure 3D). However, the tuber volume

193 of KD #2 was slighty reduced when compared to WT (Figure 3E). Starch estimation

194 assay showed no significant differences in the starch accumulation in tubers of any

195 KD lines, compared to the WT tubers (Figure 3F).

196   *BEL5* and *Identity of Tuber 1* (also known as *BRC1b*) are key positive

197 regulators of tuberization (Chen *et al*., 2003; Tang *et al*., 2022). Relative

198 quantification analyses from stolons of miR166 KD lines showed a slight reduction in

199 *BEL5* and no significant changes in *IT1* levels (Figure 3G). In potato, SP6A and

200 SWEET11b are crucial for instrumenting symplastic sucrose loading under tuber-

201 inductive conditions (Abelenda *et al*., 2019). *SWEET11b* was found to be significantly

202 reduced in the stolons of both KD lines, along with varying levels of *SP6A* (Figure

203 3G). HY5 binds to the promoters to regulate the expression of *SWEET11* and

*SWEET12* in *Arabidopsis* (Chen *et al*., 2016; Sakuraba and Yanagisawa, 2018). Stolons from KD plants had reduced *HY5* levels compared to the WT (Figure 3G). Further, we found that the tuberization potential of these KD lines was greatly reduced (80-100% reduction) under *in vitro* tuber inductive conditions (Figure 3H). Overall, the KD lines exhibited reduced tuberization potential *in vivo* and *in vitro*, accompanied with down regulation of key marker genes.

**RNA-Seq analysis revealed differentially expressed genes in swollen and sub-swollen part of tuberizing stolons**

The phenotype of elongated and pigmented tuber was evident from the early stages of tuber development (Figure 4A). To identify the genes associated with the phenotype, we carried out paired-end RNA sequencing on two distinct regions—the swollen region (Head) and the sub-swelling region (SSR)—from both WT and KD #3 samples (Figure 4B; Figure S4A). The WT_Head and SSR had 37 and 376 uniquely expressed genes respectively, and KD_Head and SSR had 115 and 120 uniquely expressed genes respectively (Figure 4C). A comparison between the Head vs SSR tissue revealed 8680 DEGs (4504 up- and 4176-down regulated) in the WT; while 7846 DEGs (3830 up- and 4016-down regulated) in the KD samples (Figure S4B-C; Table S1-S2).

Using a cut-off of log2 fold change greater than 1 and p-value less than 0.05, KD_Head and SSR had 9774 and 10,209 DEGs compared to respective WT_Head and SSR tissues (Figure S4D-E; Table S3-S4). Variance stabilized transformed (VST) values were used to visualize the gene expression patterns of certain key regulators of tuberization (Figure 4D). Notably, positive regulators such as *BEL5* and *CDF1* were slightly down regulated in KD, relative to WT. *SWEET11b* levels were greatly reduced in both the KD tissues; while *SP6A* and *IT1* were specifically down regulated in the SSR region. RT-qPCR analysis of few marker genes such as *BEL5*, *SP6A* and *HY5* corroborated with the RNA-seq data (Figure S4F-H).

**Quantitative trait loci genes for tuber shape are differentially expressed in the swollen stolons of miR166 KD plants**

The developmental differences between round and elongated tuber shapes begin as early as the hook stage of the stolon. One of the major Quantitative trait locus (QTL) for tuber shape and eye depth is located on chromosome 10 (Lindqvist-Kreuze *et al*.,

2015; Fan *et al*., 2022). Gene families linked to cell structure and function—such as homeobox, expansins, extensins, and those involved in pectin biosynthesis and modification—are particularly relevant to tuber shape and eye depth. In our sequencing data, we could detect significant up regulation of plant invertase/pectin methylesterase inhibitor (PMEI) superfamily protein (Soltu.DM.10G019700). The PMEI members are key for balancing cell wall plasticity and rigidity. Further, HEAT repeat/WD domain, G-beta repeat protein (Soltu.DM.10G019120), GRAS family transcription factor (Soltu.DM.10G020490), NB-ARC domain-containing disease resistance protein (Soltu.DM.10G016590) and heat shock protein 70B (Soltu.DM.12G011130) were significantly down regulated in the KD_Head, but not in the SSR (Figure 4E; Figure S5A; Table S3-S4).

The Round (Ro) locus is a major genetic determinant of tuber shape (round vs. long/cylindrical) that maps to chromosome 10 in potato and behaves as a major-effect QTL for length:width ratio. A total of 18 genes from the Round (Ro) region were identified in a previous study (Fan *et al*., 2022), of which we could detect 16 genes in our RNAseq data. Of these, notably Leucine-rich repeat protein kinase family protein (Soltu.DM.10G018490, Soltu.DM.10G018500), Glycosyl hydrolase superfamily protein (Soltu.DM.10G018520), Aminophospholipid ATPase (Soltu.DM.10G018550), Polynucleotidyl transferase, ribonuclease H-like superfamily protein (Soltu.DM.10G018600), and HSI2-like (Soltu.DM.10G018620, Soltu.DM.10G018640) were found to be significantly down regulated in the KD_Head relative to WT_Head (Figure 4F; Table S3). Most of these genes followed a similar expression trend in the KD_SSR tissue. (Figure S5B; Table S4). Only one member belonging to the Ovate family protein (OFP)- StOFP20 (Soltu.DM.09G010420) has been characterized in potato (Ju *et al*., 2023), wherein altering its expression changes tuber shape likely through interactions that control cell division/expansion. In the KD_Head, we found multiple OFP members such as Soltu.DM.01G003200, Soltu.DM.01G031520, Soltu.DM.03G004280, Soltu.DM.05G025790, Soltu.DM.06G032770, Soltu.DM.09G010420, Soltu.DM.10G025300, Soltu.DM.10G025310, Soltu.DM.11G022190, Soltu.DM.12G024960 to be significantly up regulated; while in the KD_SSR, certain other OFP members- Soltu.DM.02G025370, Soltu.DM.09G024410 and Soltu.DM.11G006960 showed strong differential expression relative to WT (Figure S5C; Table S3-S4).

**Tuberizing stolons exhibit differential expression of phytohormone-associated genes and accumulation of hormone in swollen vs. SSR region**

A comparison between the Head and SSR tissues from WT and KD revealed significant changes in the expression patterns of hormone-related genes (Figure 5A-B; Figure S6A-B). In WT, auxin biosynthesis genes - *StTAA-like*, *StYUCCA4* and *StYUCCA8* were up regulated in the Head compared to the SSR. This spatial expression distinction was lost in the KD line. Also, *StYUCCA5a* and *StYUCCA7* exhibited reversed expression patterns in KD line relative to WT. Auxin transport genes- *StPIN2/4/8/9* displayed markedly altered expression patterns in KD tissues, suggesting perturbed auxin distribution (Figure 5A; Figure S6A). The gibberellin (GA) biosynthesis genes *StGA3ox3* and *StGA20ox2* were found to be up regulated in both the KD tissues (Figure 5A; Figure S6A). Also, the differential expression of the GA catabolic gene *StGA2ox1*, which was evident between Head and SSR in WT, was abolished in the KD line. Expression of the cytokinin (CK) biosynthesis genes *StIPT3/5* and *StCYP735A* was altered and that of *StLOG1a/3b/7c/8b* and *CKX6* in the Head *vs*. SSR tissue was reversed in the KD lines (Figure 5B; Figure S6B). Cytokinin transporter *PUP1-7/9* showed significant reversed expression in the Head *vs*. SSR tissues of respective lines. RT-qPCR analysis of selected phytohormone-related genes validated the RNA-seq results, confirming the altered expression patterns (Figure 5C; Figure S6C). In line with the transcriptomic data, auxin quantification using LC-MS/MS revealed similar auxin levels in both the Head and SSR regions of KD, whereas WT exhibited significantly lower auxin levels in the SSR (Figure 5D; Data S1.). Additionally, $GA_3$ levels in KD stolons were relatively 5.5 folds lower than those in WT, indicating an altered hormone dynamics in the KD line (Figure 5E).

**Anthocyanin-related genes are differentially expressed in the tuberizing stolons, and tuber peels show varying pigment accumulation**

miR166 suppression led to the development of pigmented tubers in KD #2 and KD #3 lines. The pigmentation was observed only in the peel of these tubers, while the flesh color was comparable to WT tubers (Figure 6A). Metabolic profiling of the tuber peels revealed that KD #2 had increased accumulation of cyanidin 3-glucoside, while KD #3 had increased accumulation of both cyanidin 3-glucoside and pelargonidin 3-glucoside, relative to WT (Figure 6B). In contrast, delphinidin 3-glucoside was

reduced in both the KD lines (Figure 6B). These shifts in anthocyanin profiles contributed to the altered pigmentation observed in KD tubers, a trait visible from early stages of tuber development (Figure 4A). Transcriptomic analysis of the Head and SSR tissues from early stages of tuberization revealed that several structural and accessory genes of the anthocyanin biosynthesis pathway were up regulated in the KD_Head as compared to the WT (Figure 6C). A weighted correlation network analysis (WGCNA) by Zhang *et al*., (2024) identified a set of TF families having the highest correlation with these anthocyanin pathway genes. Few TFs- bHLH, WRKY, GRAS, bZIP, LBD, ARF, B3, G2-like, HB-other, HD-ZIP, MYB, and MYB-related exhibited significant differential expression in the Head tissue (Figure 6D; Table S3-S4). The anthocyanin pathway genes and associated TFs were differentially expressed in the SSR tissues as well (Figure S7A-B).

**miR166 overexpression exhibits minimal effects on vasculature and tuberization**

MicroRNA166 levels are abundant in the early stolon-to-tuber transitions and decrease at the mini-tuber stage (Figure 1G). To investigate the effects of miR166 overexpression in potato, we generated stable transgenic lines overexpressing precursor miR166-d and evaluated their overall phenotype and tuberization potential under SD conditions. The overexpression (OE) lines OE #9 and #13 lines showed variable levels of mature miR166 in leaf  (Figure S8A-B). OE #13 exhibited a slight reduction in plant height and internodal distance, while the leaf curvature and overall architecture was comparable to those of WT plants (Figure S8C-F). The number of vascular cells in the OE #9 shoot seemed to be increased, relative to the WT plants (Figure S8G). Under SD conditions, the OE #9 exhibited a slight increase in tuber productivity, although the number of tubers remained largely unchanged (Figure S8H-J). RT-qPCR analysis revealed no significant difference in the levels of mature miR166 in the stolons of OE lines, which may be due to the inherently high miR166 in the WT stolon (Figure S8K). The tuber marker gene *BEL5* did not show significant variation between OE and WT stolons, while the tuber identity gene *IT1* was up regulated (Figure S8L). Additionally, auxin biosynthesis and transport-related genes were generally up regulated in the stolons of OE lines compared to WT (Figure S8M). Thus, miR166 OE lines exhibited some phenotypic alterations, comparatively less severe than miR166 KD lines.

**REVOLUTA, a miR166 target affects tuberization and influences auxin-related genes in potato**

Using modified 5'RLM-RACE, we detected the miR166 cleaved fragments of *REV* as predicted by psRNAtarget finder (https://www.zhaolab.org/psRNATarget/home) and mapped the miR166 cleavage site (Figure 7A; Figure 1H). In the WT stolons, REV transcripts were higher under long-day (LD) conditions and in shoot-forming stolons, whereas SD stolons exhibited dynamic expression across different tissues including the stolon, swollen region, SSR, and mini-tuber (Figure 7B). Stolons from miR166 KD lines had increased *REV* levels, while a slight reduction was observed in the OE stolons (Figure 7C). GUS assay of the stable transgenic lines expressing promREV::*uidA* showed promoter activity in the apical and axillary meristems, subtending regions likely corresponding to the vascular tissue, in the swollen regions of tuberizing stolons and at the apex of the mini-tuber stage (Figure 7D). Stable transgenic lines of REV anti-sense (AS) exhibited variable degrees of *REV* suppression in leaves and stolons (Figure S9A-B), however their overall growth and architecture was comparable to WT plants (Figure 9C). The stem vasculature in REV AS #6 was smaller than that of in the WT (Figure S9D). Under SD photoperiod, all three AS lines displayed significantly reduced tuber yield, although the average number of tubers per plant remained largely unchanged (Figure 7E-G). Thus, *REVOLUTA* is a spatio-temporally regulated target of miR166 in potato, and its suppression led to reduced tuber productivity. *REVOLUTA* is known to regulate the expression of auxin-related genes in *Arabidopsis* (Brandt *et al*., 2012). It is shown to bind the promoters of TAA1 and YUCCA5 through the AT[G/C]AT palindromic sequence, which was identified as the *in vitro* binding sequence for HD-ZIP III proteins (Sessa *et al*., 1998). Our *in-silico* promoter analysis of auxin biosynthesis genes in potato revealed the presence of these core binding motifs within their upstream regulatory regions (Figure 7H). Transient overexpression of *StREV* in potato leaves reduced the transcript abundance of *StYUCCA4, -7* and *-8* genes (Figure 7I), suggesting that REV might exert an inhibitory effect on the promoter activity of auxin biosynthesis genes and regulate their expression.

**Discussion**

MicroRNAs are key regulators of plant development and defense responses, and amongst them, miR166 represents a well-conserved family that targets members of

the HD-ZIP III transcription factors. Knockdown of miR166 in maize, rice,
*Arabidopsis* and tomato has been shown to affect multiple developmental traits such
as leaf rolling, vascular patterning, epidermis structure, tassel architecture, tiller angle,
yield-related traits, and abiotic stress responses (Jia *et al*., 2015; Zhang *et al*., 2018; Li
*et al*., 2020; Zhao *et al*., 2022). In potato, overexpression of miR156 led to the
development of aerial tubers under inductive SD photoperiod (Bhogale *et al*., 2014).
In another study, the authors showed that the polycomb group member- StMSI-1
alters the expression of several genes encoding histone modifiers and positively
regulates miR156 expression in potato (Kumar *et al*., 2020). Martin *et al*. (2009)
showed that overexpression of miR172 abolished the photoperiod dependency of
tuberization. A recent study (Malankar *et al*., 2023) demonstrated the role of a phasi
RNA siRD29(-) in fine-tuning the expression of *StGA3ox3* in the early stages of
stolon-to-tuber transition. These studies reveal the critical roles of microRNAs in
governing potato tuberization.

In our preliminary analysis of small RNA sequencing datasets (Lakhotia *et al*.,
2014; Kondhare *et al*., 2018), we identified miR166 as the most abundant microRNA
in potato stolons, suggesting its involvement in stolon-to-tuber transition stages.
However, to our knowledge, the functional characterization of miR166–*HD-ZIP III*
module in potato, specifically in the storage organ development, remains unknown.
Here, we demonstrate that the miR166–*HD-ZIP III* module is functional in potato,
plays conserved roles in regulating key developmental traits, and is active through
early stolon-to-tuber transition stages affecting tuber shape, color and productivity.

**The functional roles of miR166 are conserved in potato**
A constitutive and ubiquitous knockdown of miR166 in potato (MIM166) led to the
development of several pleiotropic phenotypes. The MIM166 lines had reduced plant
height, having no significant changes in the number of nodes (Figure 2C-E). In
soybean, Zhao *et al*. (2022) showed that suppression of miR166 led to decreased plant
height. ATHB14-LIKE (a HD-ZIP III target of miR166) promoted the expression of
a GA catabolic gene- *GIBBERLLIN 2 OXIDASE2*, causing decreased bioactive GA
levels in the plant affecting the height. In potato, MIM166 *in vitro* shoots and roots
had increased levels of inactive GA8 (degraded product) and elevated expression of
GA2ox1 in stolons, compared to the WT tissues (Figure S11A-B; Figure 5A),
suggesting a similar GA dynamics in potato influencing the shoot height. In

*Arabidopsis,* miR165/166 suppression led to pleiotropic effects on leaf development such as trumpet-shaped leaves, ectopic leaflets on the abaxial side, increased leaf numbers and accumulation of purple pigment (Jia *et al*., 2015). Knockdown lines of miR166 in rice exhibited a rolled-leaf phenotype, having smaller bulliform cells and abnormal sclerenchymatous cells (Zhang *et al*., 2018). In potato, MIM166 lines exhibited reduced curvature of the leaf blades, variations in the total leaflet length and number of lateral leaves (Figure 2F-G; Figure S11C). Thus, miR166 suppression causes diverse leaf phenotypes in different plants, suggesting a conserved yet species-specific role in regulating leaf development.

Two targets of miR166, ATHB15 and HB8, play antagonistic roles in the regulation of xylem development in *Arabidopsis* (Baima *et al*., 2001; Kim *et al*., 2005). *ATHB15* is exclusively expressed in vascular tissues and its suppression leads to expanded xylem tissue and inter-fascicular region, indicative of accelerated vascular cell differentiation from cambial/procambial cells (Kim *et al*., 2005). We observed that stable potato transgenic lines of miR166 (MIM and OE) had altered vascular structures in the stem (Fig 2H; Figure S8G), suggesting miR166-*HD-ZIP III* network might also influence differentiation of vascular cells in potato. Singh *et al*. (2014) showed that phytohormones dynamically regulate the expression of miR165/166 in roots of *Arabidopsis*, and control root meristem size and growth. *In vitro* root growth assay of potato MIM166 lines showed decreased root length, root biomass, number of adventitious and lateral roots (Figure S3C-H), suggesting the essential role of miR166 in root development of potato. Overall, these pleiotropic phenotypes of the potato MIM166 lines recapitulate the phenotypic traits observed in other plants, and highlight its conserved functional roles as a central regulator of growth and development.

**miR166 impacts tuber shape, color, and productivity in potato**

Phytohormone quantification, RNA sequencing of tuberizing stolons (Head *vs.* SSR), and anatomical and molecular analyses revealed crucial insights into the altered tuber phenotypes of MIM166 plants. Phytohormones such as auxin and GA influence fruit shape in tomato and apple. Stembridge and Morrell (1972) showed that spraying $GA_4$, $GA_7$, and 6-BAP increased the length/diameter ratio in apples, while Wang *et al*. (2019) showed that auxin application before anthesis produced elongated tomato fruits. $GA_3$ application promotes organ elongation and delays ripening, whereas

paclobutrazol reduces fruit shape index and shortens ripening time in tomato (Chen *et al.*, 2020). In potato stolons, high GA levels inhibit tuberization, while auxin content increases under inductive conditions. *StGA3ox3*, a GA biosynthesis gene, is active in the tuber heel and regulates stalk formation (Malankar *et al.*, 2023). In our study, the GA catabolic gene *GA2ox1* was up regulated (Figure 5A), accompanied by a 5.5-fold reduction in bioactive $GA_3$ in MIM166 stolons (Figure 5E). Genes involved in auxin biosynthesis (*StTAA-like, StYUCCA4/5a/7/8*) and transport (*StPIN2/4/8/9*) showed altered expression in KD_Head vs. WT_Head, suggesting disrupted auxin distribution. Consistently, auxin levels in KD Head vs. SSR were comparable, unlike the distinct accumulation in WT (Figure 5D). Cytokinin, which is essential for cell division and expansion during tuber bulking (Mingo-Castel *et al.*, 1976; Sattelmacher and Marschner, 1978), had reduced expression of its biosynthesis and transport genes in SSR, possibly affecting the radial cell expansion. Moreover, MIM166 tuber sprouts were stunted, pigmented, and highly branched (Figure S6D), indicating an altered auxin–CK dynamics (Muller and Leyser, 2011). We propose that disrupted auxin–GA–CK balance in differentiating stolons caused elongated tubers in MIM166 plants.

Further analysis of our transcriptomic data sets revealed that genes located within QTLs on chromosomes 5, 10, and 12—associated with tuber shape and eye depth— are differentially expressed. Several PMEI members, GRAS transcription factors, and disease resistance genes were downregulated in KD_Head relative to WT (Figure 4E; Figure S5A; Tables S3–S4). PMEIs can modulate invertase activity and sugar metabolism (Lin *et al.*, 2015; Draffehn *et al.*, 2010), while pectin methylesterase activity affects pectin structure and tuber texture (Ross *et al.*, 2010), suggesting that altered expression of these genes may have influenced cell wall remodeling and sugar partitioning in MIM166 tubers. A recent study showed that suppression of an Ovate Family Protein - *StOFP20* (Soltu.DM.09G010420) caused a shift from round to oval-shaped tubers (Ju *et al.*, 2023). In our RNA-seq data, we do not see a marked difference in the *StOFP20* gene expression; however distinct expression patterns of 19 other *OFPs* suggest the involvement of more than one *OFP* gene in influencing tuber shape (Figure S5C). Additionally, several genes from the Ro locus (Fan *et al.*, 2022) were downregulated in KD_Head compared to WT_Head (Figure 4F; Table S3). These include a Glycosyl hydrolase (Soltu.DM.10G018520) potentially involved in cell wall remodeling (Barnes and Anderson, 2018), an Aminophospholipid ATPase (Soltu.DM.10G018550) linked to membrane asymmetry

and cell expansion (Davis *et al.*, 2020), and HSI2-like genes (Soltu.DM.10G018620, Soltu.DM.10G018640), members of the B3 DNA-binding domain family that, in *Arabidopsis*, repress AGL15 via PRC2-mediated H3K27me3 deposition (Chen *et al.*, 2018). Taken together, altered phytohormone dynamics and differential expression of genes from the QTL regions, especially the Ro locus and OFPs, likely contributed to the altered tuber shape and radial expansion observed in MIM166 lines. These results raise further questions such as- what molecular factors demarcate the swelling *vs.* sub-swelling boundary during tuber development. Is there a cross talk between miR166-*HD-ZIP III*-phytohormone module and QTL genes?

Cyanidin 3-glucoside, pelargonidin 3-glucoside, and delphinidin 3-glucoside are water-soluble anthocyanins with strong antioxidant and anti-inflammatory properties (Noda *et al.*, 2002; Duarte *et al.*, 2018). In our study, MIM166 tubers showed differential accumulation of these pigments in the peel, but not in their flesh (Figure 6A-B). A similar phenotype was reported in tomato, where overexpression of a petunia chalcone isomerase (CHI) gene increased peel flavonoids by 78-fold without affecting the flesh (Muir *et al.*, 2001). In another study, Bovy *et al.* (2002) overexpressed the maize transcription factors LC and C1 in tomato that activated the flavanols pathway, but failed to produce the flavonoids, possibly due to unchanged expression of CHI and flavanone-3′-hydroxylase (F3′H) and flavanone-3′5′-hydroxylase (F3′5′H) genes. In our transcriptomic analysis, we observed an up regulation of multiple structural and accessory genes from the anthocyanin biosynthesis pathway in KD tissues compared to WT, consistent with the increased pigment accumulation (Figure 6C; Figure S7A). Transcription factor families which highly correlated with these pathway genes—bHLH, WRKY, GRAS, bZIP, LBD, ARF, B3, G2-like, HB-other, HD-ZIP, MYB, and MYB-related—also showed significant differential expression (Figure 6D; Table S3–S4; Figure S7A–B). In Apple, degradation of MdIAA121 induced by auxin treatment releases MdARF13, which acts as a negative regulator of the anthocyanin metabolic pathway (Wang *et al.*, 2018). Also, auxin inhibits the accumulation of anthocyanin in apple by degrading the MdIAA26 protein (Wang *et al.*, 2020). A tuberizing stolon is known to have high auxin at its apex for facilitating tuber development. We speculate this could have restricted pigments accumulation to the tuber peels only in the miR166 KD lines. Conversely, flavonoids may act as inhibitors of auxin transport (Gayomba *et al.*, 2016) which might have affected auxin distribution in the KD stolons. Further studies are

required to ascertain whether the auxin dynamics in tuberizing stolons and
anthocyanin accumulation are inter-linked.

Under tuber inductive SD photoperiod, SWEET11b- SP6A interacts to
mediate the symplastic loading of sucrose, and prevents its loss to the apoplastic route
(Abelenda *et al*., 2019). SWEET11b expression was drastically reduced in all the
MIM166 lines (Figure 3G and Figure 4D). Also, the vascular structures in the KD
stem were distorted (Figure 2H). The MIM166 plants had reduced tuber yield under
SD photoperiod, with no significant changes in the starch content per gram fresh
weight of the tuber (Figure 3B and 3F). Thus, the reduced productivity could be due
to the altered vascular structure of the stem, reduced symplastic loading of sucrose, or
impaired tuber-bulking at the stolon apex due to disrupted phytohormone dynamics.
Interestingly, MIM166 plants developed floral buds at the shoot apex and aerial
stolon/tubers under SD photoperiod (Figure S12A-C), denoting the formation of
ectopic sinks due to the improper sucrose distribution. As SD induction progressed,
the belowground developing tubers acted as the primary and stronger sinks, thus
limiting sucrose availability for the ectopic sinks.

**HD-ZIP III TFs are involved in tuberization**

527
Auxin plays a key role in the stolon-to-tuber transition. The stolon apical meristem
(StAM) is a major site of auxin biosynthesis, where YUC-like1 (*StYUC4*) acts as a
key contributor (Roumeliotis *et al*., 2012). In *Arabidopsis*, REVOLUTA regulates
promoters of auxin biosynthesis, transport, and signaling genes (Brandt *et* al., 2012;
Reinhart *et al*., 2013). In our study, phenotypes such as impaired root growth,
increased tuber sprout branching, and altered auxin-related gene expression suggested
a role for REV in potato (Figure 7A–C). Through GUS assay, we observed that the
promoter activity of *REV* was visible in the apex of stolons, vascular region and
subsequent tuberizing stages (Figure 7D). This was similar to the activity of an auxin
responsive promoter, (*DR5:GUS*) observed by Roumeliotis *et al*. (2012), suggesting a
putative overlap of *REV* expression and auxin accumulation. REV antisense (REV AS)
lines exhibited reduced tuber yield, suggesting a hitherto unknown role of REV in
tuberization (Figure 7E-G). The HD-ZIP III binding motif, AT[G/C]AT, was enriched
in promoters of potato auxin biosynthesis genes, implying REV-mediated regulation
(Figure 7H). Our transient overexpression of *StREV* in the leaves of potato (cv.
Désirée) suggests that REV could act as a suppressor of *StYUC4/7/8* expression

(Figure 7I) in potato, as opposed to its transcription-activator role in *Arabidopsis*. Auxin levels are dynamic during tuberization, and further studies are needed to clarify the role of REV in regulating auxin during stolon differentiation. Besides REV, other members of the HD-ZIP III family could also play a role in the stolon-to-tuber transitions (Figure S10B). For example, whether HB8 and ICU1/2 are necessary for vascular development of the tuber?

Fruit/organ shape is determined during its development. Similar to the findings in tomato and apple (Stembridge and Morrell, 1972; Wang *et al*., 2019; Chen *et al*., 2020), we hypothesize that the tuber shape in potato is governed by phytohormone dynamics (Figure 8). The miR166–HD-ZIP III module is crucial in phytohormone response pathways, and are shown to modulate the expression of each other. In potato stolons, REVOLUTA may act to repress auxin biosynthesis genes (Figure 7), while PHABULOSA, as shown in soybean, could modulate GA metabolism (Figure 8; Zhao *et al*., 2022). Together, these TFs likely coordinate auxin/GA homeostasis in the swelling and sub-swelling stolon, thereby affecting the tuber shape. Additionally, HB8 and INCURVATA are shown to exert antagonistic effects on vascular differentiation in *Arabidopsis* (Kim *et al*., 2005), and auxin positively modulates HB8 expression through ARF5/MONOPTEROS (Donner *et al*., 2009). Development of vascular tissues in differentiating stolon is crucial for delivery of nutrients and photoassimilates. We propose that the miR166–HB8/ICU1/2 likely coordinates vascular differentiation and patterning in developing stolon (Figure 8) and miR166 and its targets- the HD-ZIP III members play multifaceted roles in regulating tuber development.

To our knowledge, the role of *miR166-HD-ZIP III* module in storage organ development has not been studied. Here, we demonstrate that miR166 plays conserved roles in regulating numerous developmental processes such as plant height, vascular structure, leaf curvature, and root development, and unique roles in regulating tuber shape, color, and productivity. We also show that REVOLUTA affects tuberization and could regulate the expression of auxin related genes in potato. Further studies are required to explore the roles of miR166 and HD-ZIP III members in tuber tissue patterning and development. In conclusion, our findings establish that miR166 is an important regulatory element in potato, having significant impact on tuber development, morphology, and productivity.

## Materials and Methods

### Plant material and growth conditions

A photoperiod-responsive potato cultivar *Solanum tuberosum* ssp. *andigena* 7540 which tuberizes under SD conditions, was used in this study. *In vitro* plants (wild-type and transgenic lines) were propagated from axillary node sub-cultures on Murashige and Skoog's basal medium with 2% (w/v) sucrose (Murashige and Skoog., 1962) and were grown in a plant growth incubator (Percival Scientific, Inc.) with a light intensity of 300 μmol m−2 s−1 under LD conditions unless mentioned otherwise.

### Prediction and detection of miR166 precursors in potato

Four precursor sequences (stu-MIR166a, stu-MIR166b, stu-MIR166c and stu-MIR166d) of *miR166* were retrieved from the miRBase (http://mirbase.org/; Griffiths-Jones S *et al*., 2008). The secondary structures of these precursors were predicted using the Mfold software (http://www.unafold.org/mfold/applications/rna-folding-form.php; Zuker, 2003), and structures with lowest free energies were then further screened for having the characteristic stem-loop secondary structure with the mature miRNA sequence in the arm region of the stem-loop. All four miRNA precursors were then validated from *S. tuberosum* ssp. *andigena* 7540 by RT-PCR analysis. The sequences of all the oligos used in this study are provided in Table S5, and the IDs for all the genes are in the Table S6.

Total RNA was harvested from shoot apex, mature leaves and stolon tissues of potato using RNAiso Plus (DSS TAKARA) as per the manufacturer's instructions. cDNA was synthesized using two microgram (2μg) of total RNA, Superscript IV Reverse Transcriptase (Invitrogen), and oligo dT primers. miRNA precursors were amplified by RT-PCR using precursor-specific primer pairs. The amplicons were cloned into sub-cloning vector pGEMT-easy (Promega) and sequence verified.

### Validation of mature miR166

Total RNA was isolated using RNAiso Plus (DSS TAKARA) from mature leaves and stolons of WT potato plants. One microgram (1 μg) of total RNA was used for cDNA preparation using miR166 specific stem-loop primers (STP) followed by their end-point PCRs using miR166 specific forward and universal reverse primer as described in Varkonyi-Gasic *et al*. (2007) (Table S5).

**Tissue-specific relative abundance of miR166 under SD and LD conditions**

For investigating the influence of photoperiod on tissue-specific accumulation of mature *miR166*, *in vitro* grown wild-type *S. tuberosum* spp. *andigena* 7540 plants were transferred to soil and maintained under LD photoperiod with 300 μmol m−2 s−1 light intensity for a period of ten weeks (until they attained 10-12 leaf stages) in a growth chamber (Percival Scientific, Inc.). Later, half of the plants were transferred to tuber-inductive SD photoperiod for 7 days, while the remaining plants were maintained under LD conditions. Leaves and stages of stolon-to-tuber transitions were collected at 7 days post LD/SD induction in triplicates. Isolation of total RNA and cDNA synthesis was carried out as described before. qPCR reactions were performed on CFX96 Real-Time System (BIO-RAD) with miR166-specific forward primer and universal reverse primer, using TAKARA SYBR green master mix (Takara-Clontech). The reaction mix was incubated at 95 °C for 30 s, followed by 40 cycles at 95 °C for 5 s, gene-specific annealing temperature for 15 s and extension for 72 °C for 15 s. PCR specificity was checked by melting curve analysis, and data were analyzed using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

**Generation of constructs and potato transgenic lines**

MicroRNA166 overexpression (OE) and mimicry (MIM) constructs for overexpression and suppression of the miR166, respectively, were previously generated in our lab. Briefly, miR166 OE construct- 35S:St-pre166D-pBI121 was generated by amplifying precursor (St-pre166-d) from potato RNA using primers Pre166d-FP and Pre166d-RP (Table S5). The amplified product (108 bp) was cloned into the binary vector pBI121 downstream to the 35S CaMV constitutive promoter. For knockdown (KD) construct, we used artificial target mimicry (MIM) strategy. The KD construct (originally in pGREEN vector) was obtained from European *Arabidopsis* Stock Centre (NASC) (Todesco *et al*., 2010), and the MIM166 insert (542 bp) was re-cloned to pBI121 binary vector to generate 35S:MIM166-pBI121. Both these constructs- OE and MIM were then mobilized into *Agrobacterium tumefaciens* strain GV3101. Transgenic potato plants in *S. tuberosum* spp. *andigena* 7540 were generated as described in Banerjee *et al*., 2006. Kanamycin-resistant transgenic plants were further screened by molecular methods and selective lines were used for further experiments.

After 4 weeks of *in vitro* growth, potato plants were transferred to soil, grown

for 7-9 weeks under LD conditions, before initiating SD induction. Phenotypic

characteristics such as plant height, number of nodes and leaf curvature were recorded

just before the initiation of SD photoperiod treatment, whereas the frequency of floral

bud formation and tuber productivity parameters were measured after 1 and 8-9 weeks

of SD induction, respectively.

**Phenotypic analysis**

All the phenotypic analyses were performed on 6–7 weeks old soil-grown plants.

Shoot height was measured from the bottom-most node up to the plant apex. For

measurement of leaf curvature indices, mature terminal leaves at the fifth node were

analyzed as described in Liu *et al*., 2010. These experiments were performed at least

three times with similar results.

**Assessment of tuber productivity**

One-month-old plantlets of the transgenic lines and WT were transferred to soil in

small round pots and hardened for one week under LD conditions in a growth

chamber (Percival Scientific, Inc.) with a light intensity of 300 μmol m−2 s−1, day

temperature of 24°C, and night temperature of 22°C. This was followed by an

additional three weeks of incubation in the same pots. Subsequently, the plants were

re-potted into medium-sized pots and continued to grow under LD conditions for an

additional five weeks. When the plants reached the 10-12 leaf stage, they were

transferred to tuber inductive short-day (SD) conditions (16h dark and 8h light) with

day and night temperatures of 22°C and 20°C, respectively. A subset of these plants

was harvested after 7 days of induction, and stolons were collected for RT-qPCR

analysis. After 60 days of induction, remaining plants were evaluated for their tuber

productivity (gram fresh weight [g fr wt] per plant) and tuber numbers. This

experiment was carried out 3 times, with consistent results. For tuber sphericity index

and volume assessment, 30 tubers per line were measured for their diameters

(Diameter1, Diameter2 and height). *In vitro* tuber induction assay was performed

(experiment was repeated three times independently) as described in Kondhare *et al*.,

2021.

**Starch estimation**

Starch estimation was carried out twice, independently, as described in Kamble *et al.*
(2023), Hawkins *et al.* (2021) and Kringel *et al.* (2020), with minor modifications.
Tuber tissue was well-homogenized in cold water containing 0.35% sodium
metabisulphte. The slurry was filtered to remove the debris. Filtered solution was kept
undisturbed for 1 hour at 4 degrees. Insoluble material was collected by centrifugation,
followed by 3 washes with 80% ethanol, and then finally re-suspended in water.
Starch was digested using alpha amylase (SRL), glucoamylase (SRL) and
Amyloglucosidase (Sigma). Glucose was estimated using the DNSA method, and
compared with a standard graph prepared using Starch (Sigma), digested under
identical conditions.
690

**RNA sequencing**
For RNA-sequencing, miR166 KD #3 and WT plants were grown in soil for 10-weeks
under LD conditions and subjected to SD induction for 12 days. Swollen stolons were
harvested in triplicates (from at least 20 plants per replicate) and divided into swelling
(Head) and sub-swelling region (SSR) and snap-frozen in liquid nitrogen. The tissues
were crushed using liquid Nitrogen and stored in -80⁰ C until further use. For RNA
sequencing and RT-qPCR purposes, 100 mg of tissue from each sample was used to
isolate the total RNA using RNAiso Plus as per the manufacturer's instructions. RNA
quality and integrity were assessed, and RNA-sequencing was performed by
Novelgene Technologies Pvt. Ltd., Hyderabad. Paired-end sequencing was performed
on Illumina NovaSeq 6000, adapter sequences and low-quality reads were filtered
using FASTP to get high-quality reads on the parameter of <Q20, length < 50. Further
processing was performed as described in Kumar *et al.* 2020. Validation of selected
target genes identified in the RNA-seq analysis was done using qPCR as described
above.
706

**Detection of phytohormones and pigments by LC-MS/MS**
Tissues- stolon, swollen stolon (Head and SSR), tuber peel, tuber flesh were harvested,
crushed in liquid Nitrogen, and stored at -80⁰ C until further use. 100 mg of the
crushed powder was extracted with 1 ml of extraction solvent (For phytohormone
extraction- MS grade methanol/water/formic acid 75:20:5; having 300ng internal
standard Adonitol; For pigment extraction- 80 percent methanol in MS grade water;
having 300ng internal standard Adonitol), by 3-5 min of vigorous vortex, followed by

15 min of sonication. The solution was incubated overnight at -80°C under dark conditions. Next day, the samples were centrifuged at 4°C, maximum speed for 15 min. The supernatant was transferred into a new tube, and reduced to half of its original volume, using centrivap at 4°C, and was filtered through 0.22-micron syringe filter. The LC-MS system (Exion AD, AB Sciex) was used for detection of phytohormones and pigments. Chromatographic separation was done on AB SCIEX Exion LC AD UPLC system, using Phenomenex kinetex core shell C18 column (2.1X150 mm, 1.7µm). Separation of compounds was performed at 0.3 ml/ min flow rate with linear gradient of solvent A (milliQ + 0.1% formic acid) and solvent B (Acetonitrile + 0.1% formic acid) was performed as follows: 0 min 5% B + 95% A; 3 min, 10% B; 6 min, 50% B, 7 min, 70% B; 10 min, 95% B; 12 min, finally column was re-equilibrated to initial ratio of 5% B + 95% A for 4 min with total run time of 16 min. The m/z range for phytohormone and pigment detection was 100-400 and 100-1700, respectively. The sample injection volume was 40 µl for all the samples. Three blank runs were given every time between the biological replicates throughout run. The full scan mass spectrometry data was acquired on the X500R Q-TOF (AB Sciex, USA). The data for all the samples were acquired using electrospray ionization in both positive and negative modes. Mass calibration was performed in positive and negative ion modes before the data acquisition. Data analysis was carried out using the SCIEX OX software provided along with the machine.

**GUS assay**

*StREV* promoter transgenic lines (promREV:*uidA*) were grown *in vitro* under LD conditions for 20 d. They were also grown in soil under LD conditions for 8 weeks, followed by SD induction for 15 d. Entire *in vitro* grown plantlets, stolon and tuber samples from SD-induced soil-grown plants were subjected to GUS assay as described in Begum *et al*., (2022) and imaged using olympus MX stereozoom microscope.

**Chlorophyll and anthocyanin estimation**

Anthocyanin and chlorophyll estimation from leaves of WT and miR166 KD lines was performed as per the method in Nakata and Ohme-Takagi (2014) and Bhattacharjee and Sharma (2012), respectively. These experiments were conducted two times, with similar results.

**Histology**

For anatomical studies, stems from the middle part of soil-grown plants (6 weeks LD) were used. Free hand transverse sections were cut using a sharp razor blade, and selective sections were stained with Toluidine Blue O solution as described in Pradhan Mitra and Loqué., 2014. The sections were mounted in 70% glycerol, before imaging them on the Nikon Eclipse microscope.

**Infiltration assay**

*Agrobacterium*-mediated co-infiltration assay was performed in leaves of 6-week old WT potato (cv. *Désirée*). *35S:StREV* cloned in pCB201 construct was transformed into *A. tumefaciens* GV3101 strain. An overnight grown primary culture was used to inoculate 100 ml secondary culture (having 150 μM acetosyringone) without any antibiotics. When the secondary culture reached an OD600 of 1.00, it was centrifuged, and the cell pellet was resuspended in the infiltration buffer (10 mM $MgCl_2$, 10 mM MES-K: pH= 5.6, plus 150 μM acetosyringone), incubated at RT on a tube-roller for at least 3 hrs, followed by infiltration into the leaves of potato plants using a needleless syringe. Infiltration with only the infiltration buffer served as experimental control. Leaves were harvested 72 hrs post-infiltration, and subjected further for RNA isolation and RT-qPCR analyses. This experiment was conducted three times with similar results.

**Data availability**

The raw sequencing data were deposited in the NCBI Short Read Archive (SRA) database (http://www.ncbi.nlm.nih. gov/sra/). It can be retrieved using the SRA ID SRP630954. All other relevant data can be found within the manuscript and its supporting materials.

**Acknowledgements**

Authors acknowledge Mr. Nilesh Lohakare and Mr. Nitish Lahigude (Indian Institute of Science Education and Research, Pune) for their help in maintaining the plants in the green house facility. We thank Novelgene Technologies Pvt. Ltd. for performing RNA sequencing. We are thankful to Dr. Harpreet Singh Kalsi, Dr. Nilam Malankar, Dr. Kirtikumar Kondhare and Ms. Shikha Singh for their inputs.

**Funding**

NSP and JK are thankful to the University Grants Commission for the Senior Research Fellowship and Junior Research Fellowship, respectively. GA acknowledges funding received from the Department of Biotechnology, India. AKB acknowledges the support and generous funding from the Indian Institute of Science Education and Research (IISER) Pune.

**Authors contributions**

N.S.P. designed the experiments, generated transgenic lines, carried out phenotypic and transcriptomic data analyses, conducted experiments and wrote the manuscript draft. A.V. conducted the LC-MS/MS experiments and analyzed its data. J.K. carried out the infiltration experiment and performed its molecular analysis. G.A. and N.S.P. generated the miR166 KD and promREV::uidA transgenic lines. B.N. and N.S.P. performed RLM-RACE for target validation. A.K.B. supervised the study, obtained the funding and resources. N.S.P. and A.K.B. conceived the idea and edited the manuscript. All authors have read and approved the final manuscript.

**Conflict of Interest Statement**

The authors declare that they have no conflicts of interest associated with this work.

**Ethics Statement**

All experimental research on plants in this study complied with institutional bio-safety guidelines. The plant materials used were maintained in accordance with relevant regulations.

**Supporting Information**

**Figure S1.** Details of the precursors and mature miR166 in potato

**Figure S2.** Abundance of *miR166* in stolons under early LD vs SD stages from a photoperiod dependent tuberizing variety ssp. *andigena* 7540

**Figure S3.** Pleiotropic phenotypes of miR166 suppression on potato

**Figure S4.** PCA and Volcano plots of samples processed for RNA-sequencing

**Figure S5.** Heatmap for selective genes from the QTLs

**References**

• Abelenda JA, Bergonzi S, Oortwijn M, Sonnewald S, Du M, Visser RGF,

Sonnewald U, Bachem CWB. 2019. Source-sink regulation is mediated by

interaction of an FT homolog with a SWEET protein in potato. Curr Biol.

29:1178–1186.e6. doi:10.1016/j.cub.2019.02.018

• Baima S, Possenti M, Matteucci A, Wisman E, Altamura MM, Ruberti I,

Morelli G. 2001. The Arabidopsis ATHB-8 HD-Zip protein acts as a

differentiation-promoting transcription factor of the vascular meristems. Plant

Physiol.

• Banerjee AK, Prat S, Hannapel DJ. 2006. Efficient production of transgenic

potato (S. tuberosum L. ssp. andigena) plants via Agrobacterium tumefaciens-

mediated transformation. Plant Sci. 170:732–738.

doi:10.1016/j.plantsci.2005.11.007

• Barnes, W. J., & Anderson, C. T. (2018). Release, recycle, rebuild: Cell-wall
remodeling, autodegradation, and sugar salvage for new wall biosynthesis
during plant development. Molecular Plant, 11(1), 31–46.
https://doi.org/10.1016/j.molp.2017.08.011

• Begum S, Jing S, Yu L, Sun X, Wang E, Abu Kawochar M, Qin J, Liu J, Song
B. 2022. Modulation of JA signalling reveals the influence of StJAZ1-like on
tuber initiation and tuber bulking in potato. Plant J. 109:952–964.
doi:10.1111/tpj.15606

• Bhattacharjee S, Sharma GD. 2012. Effect of dual inoculation of arbuscular
mycorrhiza and rhizobium on the chlorophyll, nitrogen and phosphorus
contents of pigeon pea (Cajanus cajan L.). Adv Microb. 2:25945.
doi:10.4236/aim.2012.24072

• Bhogale S, Mahajan AS, Natarajan B, Rajabhoj M, Thulasiram HV, Banerjee
AK. 2014. MicroRNA156: a potential graft-transmissible microRNA that
modulates plant architecture and tuberization in Solanum tuberosum ssp.
Andigena. Plant Physiol. 164:1011–1027.

• Boualem A, Laporte P, Jovanovic M, Laffont C, Plet J, Combier JP, Niebel A,
Crespi M, Frugier F. 2008. MicroRNA166 controls root and nodule
development in Medicago truncatula. Plant J. 54:876–887. doi:10.1111/j.1365-
313X.2008.03448.x

• Brandt R, Salla-Martret M, Bou-Torrent J, Musielak T, Stahl M, Lanz C, Ott F,
Schmid M, Greb T, Schwarz M, Choi SB, Barton MK, Reinhart BJ, Liu T,
Quint M, Palauqui JC, Martínez-García JF, Wenkel S. 2012. Genome-wide
binding-site analysis of REVOLUTA reveals a link between leaf patterning
and light-mediated growth responses. Plant J. 72:31–42. doi:10.1111/j.1365-
313X.2012.05049.x

• Bovy A, Schijlen E, Hall RD (2007) Metabolic engineering of flavonoids in
tomato (Solanum lycopersicum): the potential for metabolomics.
Metabolomics 3: 399–412. https://doi.org/10.1007/s11306-007-0074-2

• Chen H, Rosin FM, Prat S, Hannapel DJ. 2003. Interacting transcription
factors from the three-amino acid loop extension superclass regulate tuber
formation. Plant Physiol. 132:1391–1404. doi:10.1104/pp.103.022434

• Chen S, Wang XJ, Tan GF, Zhou WQ, Wang GL. 2020. Gibberellin and the
plant growth retardant paclobutrazol altered fruit shape and ripening in tomato.
Protoplasma. 257:853–861. doi:10.1007/s00709-019-01471-2

• Chen X, Yao Q, Gao X, Jiang C, Harberd NP, Fu X. 2016. Shoot-to-root
mobile transcription factor HY5 coordinates plant carbon and nitrogen
acquisition. Curr Biol. 26:640–646.

• Chen, N., et al. (2018). HSI2/VAL1 silences AGL15 to regulate the
developmental transition from seed maturation to vegetative growth in
Arabidopsis. The Plant Cell, 30(3), 600–619.
https://doi.org/10.1105/tpc.17.00893

• Clepet C, Devani RS, Boumlik R, Hao Y, Morin H, Marcel F, Verdenaud M,
Mania B, Brisou G, Citerne S, Mouille G, Lepeltier JC, Koussevitzky S,
Boualem A, Bendahmane A. 2021. The miR166–SlHB15A regulatory module
controls ovule development and parthenocarpic fruit set under adverse
temperatures in tomato. Mol Plant. 14:1185–1198.
doi:10.1016/j.molp.2021.05.005

• Davis, J. A., Pares, R. B., Bernstein, T., McDowell, S. C., Brown, E., Stubrich,
J., et al. (2020). The lipid flippases ALA4 and ALA5 play critical roles in cell
expansion and plant growth. Plant Physiology, 182(4), 2111–2125.
https://doi.org/10.1104/pp.19.01332

• Ding Y, Gong S, Wang Y, Wang F, Bao H, Sun J, Cai C, Yi K, Chen Z, Zhu
C. 2018. MicroRNA166 modulates cadmium tolerance and accumulation in
rice. Plant Physiol. 177:1691–1703. doi:10.1104/pp.18.00485

• Donner, T. J., Sherr, I., & Scarpella, E. (2009). Regulation of preprocambial
cell state acquisition by auxin signaling in Arabidopsis leaves. Development,
136(19), 3235-3246. https://doi.org/10.1242/dev.037028

• Draffehn AM, Meller S, Li L, Gebhardt C. 2010. Natural diversity of potato
(Solanum tuberosum) invertases. BMC Plant Biol. 10:271. doi:10.1186/1471-
2229-10-271

• Duarte LJ, Chaves VC, Nascimento MVP dos S, Calvete E, Li M, Ciraolo E,
Ghigo A, Hirsch E, Simões CMO, Reginatto FH, Dalmarco EM. 2018.
Molecular mechanism of action of pelargonidin-3-O-glucoside, the main

anthocyanin responsible for the anti-inflammatory effect of strawberry fruits. Food Chem. 247:56–65. doi:10.1016/j.foodchem.2017.12.015

- Fan G, Wang Q, Xu J, Chen N, Zhu W, Duan S, Yang X, De Jong WS, Guo Y, Jin L, Li G. 2022. Fine mapping and candidate gene prediction of tuber shape controlling Ro locus based on integrating genetic and transcriptomic analyses in potato. Int J Mol Sci. 23:1470. doi:10.3390/ijms23031470

- Gayomba, S. R., Watkins, J. M., & Muday, G. K. (2016). Flavonols regulate plant growth and development through regulation of auxin transport and cellular redox status. In K. Yoshida, V. Cheynier, & S. Quideau (Eds.), Recent Advances in Polyphenol Research (Vol. 5, pp. 143–170). John Wiley & Sons. https://doi.org/10.1002/9781118883303.ch7

- Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ. 2008. miRBase: tools for microRNA genomics. Nucleic Acids Res. 36:D154–D158.

- Hannapel DJ, Sharma P, Lin T, Banerjee AK. 2017. The multiple signals that control tuber formation. Plant Physiol. 174:845–856.

- Hawkins E, Chen J, Watson-Lazowski A, Ahn-Jarvis J, Barclay JE, Fahy B, Hartley M, Warren FJ, Seung D. 2021. STARCH SYNTHASE 4 is required for normal starch granule initiation in amyloplasts of wheat endosperm. New Phytol. 230:2371–2386. doi:10.1111/nph.17342

- Jia X, Ding N, Fan W, Yan J, Gu Y, Tang X, Li R, Tang G. 2015. Functional plasticity of miR165/166 in plant development revealed by small tandem target mimic. Plant Sci. 233:11–21. doi:10.1016/j.plantsci.2014.12.020

- Ju A, Wang Y, Yan Y, Li C, Luo W, Ma L, Shang Y, Gao D. 2023. StOFP20 regulates tuber shape and interacts with TONNEAU1 recruiting motif proteins in potato. J Integr Agric. 22:752–761. doi:10.1016/j.jia.2022.08.069

- Kamble UN, Makhamadjonov F, Fahy B, Martins C, Saalbach G, Seung D. 2023. Initiation of B-type starch granules in wheat endosperm requires the plastidial α-glucan phosphorylase PHS1. Plant Cell. 35:4091–4110. doi:10.1093/plcell/koad217

- Kim J, Jung JH, Reyes JL, Kim YS, Kim SY, Chung KS, Kim JA, Lee M, Lee Y, Kim VN, Chua NH, Park CM. 2005. microRNA-directed cleavage of ATHB15 mRNA regulates vascular development in Arabidopsis inflorescence stems. Plant J. 42:84–94. doi:10.1111/j.1365-313X.2005.02354.x

• Kondhare KR, Kumar A, Patil NS, Malankar N, Saha K, Banerjee AK. 2021.
Development of aerial and belowground tubers in potato is governed by
photoperiod and epigenetic mechanism. Plant Physiol. 187:1071–1086.
doi:10.1093/plphys/kiab409

• Kondhare KR, Malankar NN, Devani RS, Banerjee AK. 2018. Genome-wide
transcriptome analysis reveals small RNA profiles involved in early stages of
stolon-to-tuber transitions in potato under photoperiodic conditions. 1–19.

• Kringel DH, el Halal SLM, Zavareze Eda R, Dias ARG. 2020. Methods for
the extraction of roots, tubers, pulses, pseudocereals, and other unconventional
starch    sources:    a    review.    Starch/Staerke.    72:11–12.
doi:10.1002/star.201900234

• Kumar A, Kondhare KR, Vetal PV, Banerjee AK. 2020. PcG proteins MSI1
and BMI1 function upstream of miR156 to regulate aerial tuber formation in
potato. Plant Physiol. 182:185–203. doi:10.1104/pp.19.00416

• Lakhotia N, Joshi G, Bhardwaj AR, Katiyar-Agarwal S, Agarwal M,
Jagannath A, Goel S, Kumar A. 2014. Identification and characterization of
miRNAome in root, stem, leaf and tuber developmental stages of potato
(Solanum tuberosum L.) by high-throughput sequencing. BMC Plant Biol.
14:6.

• Li N, Yang T, Guo Z, Wang Q, Chai M, Wu M, Li X, Li W, Li G, Tang J,
Tang G, Zhang Z. 2020. Maize microRNA166 inactivation confers plant
development and abiotic stress resistance. Int J Mol Sci. 21:9506.
doi:10.3390/ijms21249506

• Lin Y, Liu T, Liu J, Liu X, Ou Y, Zhang H, Li M, Sonnewald U, Song B, Xie
C. 2015. Subtle regulation of potato acid invertase activity by a protein
complex of invertase, invertase inhibitor, and sucrose nonfermenting1-related
protein kinase. Plant Physiol. 168(4):1807–1819. doi:10.1104/pp.15.00664

• Lindqvist-Kreuze H, Khan A, Salas E, Meiyalaghan S, Thomson S, Gomez R,
Bonierbale M. 2015. Tuber shape and eye depth variation in a diploid family
of Andean potatoes. BMC Genet. 16:57. doi:10.1186/s12863-015-0213-0

• Liu Z, Jia L, Mao Y, He Y. 2010. Classification and quantification of leaf
curvature. J Exp Bot. 61:2757–2767. doi:10.1093/jxb/erq111

•   Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data
using real-time quantitative PCR and the 2(-ΔΔC(T)) method. Methods.
25:402–408.

•   Malankar NN, Kondhare KR, Saha K, Mantri M, Banerjee AK. 2023. The
phased short-interfering RNA siRD29(-) regulates GIBBERELLIN 3-
OXIDASE 3 during stolon-to-tuber transitions in potato. Plant Physiol.
193:2555–2572. doi:10.1093/plphys/kiad493

•   Martin A, Adam H, Díaz-Mendoza M, Zurczak M, González-Schain ND,
Suárez-López P. 2009. Graft-transmissible induction of potato tuberization by
the microRNA miR172. Development. 136:2873–2881.

•   Mingo-Castel AM, Young RE, Smith OE. 1976. Kinetin-induced tuberization
of potato in vitro: on the mode of action of kinetin. Plant Cell Physiol. 17:557–
567.

•   Muir S, Collins G, Robinson S, et al. 2001. Overexpression of petunia
chalcone isomerase in tomato results in fruit containing increased levels of
flavonols. Nat Biotechnol. 19:470–474. doi:10.1038/88150

•   Murashige T, Skoog F. 1962. A revised medium for rapid growth and
bioassays with tobacco tissue cultures. Physiol Plant. 15:473–497.
doi:10.1111/j.1399-3054.1962.tb08052.x

•   Nakata M, Ohme-Takagi M. 2014. Quantification of anthocyanin content.
Bio-Protoc. 4:1428–1440. doi:10.21769/BioProtoc.1098

•   Noda Y, Kaneyuki T, Mori A, Packer L. 2002. Antioxidant activities of
pomegranate fruit extract and its anthocyanidins: delphinidin, cyanidin, and
pelargonidin. J Agric Food Chem. 50:166–171. doi:10.1021/jf0108765

•   Pradhan Mitra P, Loqué D. 2014. Histochemical staining of Arabidopsis
thaliana secondary cell wall elements. J Vis Exp. 87:51381.
doi:10.3791/51381

•   Prigge MJ, Otsuga D, Alonso JM, Ecker JR, Drews GN, Clark SE 2005 Class
III homeodomain-leucine zipper gene family members have overlapping,
antagonistic, and distinct roles in Arabidopsis development. Plant Cell 17: 61–
76. https://doi.org/10.1105/tpc.104.026161

• Ramachandran P, Carlsbecker A, Etchells JP. 2017. Class III HD-ZIPs govern
vascular cell fate: an HD view on patterning and differentiation. Physiol Plant.
68:55–69.

• Ross ARS, Vrebalov J, Lang L, Allan WL, Abrams SR, Hall M, Tanksley SD,
van den Berg L, McCourt P. 2010. Potato tuber pectin structure is influenced
by pectin methyl esterase activity and impacts on cooked potato texture. J Exp
Bot. 62(1):371–381. doi:10.1093/jxb/erq280

• Roodbarkelari F, Groot EP. 2017. Regulatory function of homeodomain-
leucine zipper (HD-ZIP) family proteins during embryogenesis. New Phytol.
213:95–104.

• Roumeliotis E, Kloosterman B, Oortwijn M, Kohlen W, Bouwmeester HJ,
Visser RGF, Bachem CWB. 2012. The effects of auxin and strigolactones on
tuber initiation and stolon architecture in potato. J Exp Bot. 63:4539–4547.
doi:10.1093/jxb/ers132

• Sakuraba Y, Yanagisawa S. 2018. Light signalling-induced regulation of
nutrient acquisition and utilisation in plants. Semin Cell Dev Biol. 83:123–132.
doi:10.1016/j.semcdb.2017.12.014

• Sattelmacher B, Marschner H. 1978. Cytokinin activity in stolons and tubers
of Solanum tuberosum during the period of tuberization. Z Pflanzenernähr
Bodenkd. 44.

• Sessa G, Steindler C, Morelli G, Ruberti I. 1998. The Arabidopsis Athb-8, -9
and -14 genes are members of a small gene family coding for highly related
HD-ZIP proteins. Plant Mol Biol. 38.

• Singh A, Singh S, Panigrahi KCS, Reski R, Sarkar AK. 2014. Balanced
activity of microRNA166/165 and its target transcripts from the class III
homeodomain-leucine zipper family regulates root growth in Arabidopsis
thaliana. Plant Cell Rep. 33:945–953. doi:10.1007/s00299-014-1573-z

• Stembridge GE, Morrell G. 1972. Effect of gibberellins and 6-benzyladenine
on the shape and fruit set of 'Delicious' apples. J Am Soc Hort Sci. 97:464–
467.

• Tang D, Jia Y, Zhang J, Li H, Cheng L, Wang P, Bao Z, Liu Z, Feng S, Zhu X,
et al. 2022. Genome evolution and diversity of wild and cultivated potatoes.
Nature. 606:535–541.

• Todesco M, Rubio-Somoza I, Paz-Ares J, Weigel D. 2010. A collection of
target mimics for comprehensive analysis of microRNA function in
Arabidopsis thaliana. PLoS Genet. 6:e1001031.
doi:10.1371/journal.pgen.1001031

• Turchi L, Baima S, Morelli G, Ruberti I. 2015. Interplay of HD-Zip II and III
transcription factors in auxin-regulated plant development. J Exp Bot.
66:5043–5053.

• Valiollahi E, Farsi M, Kakhki AM. 2014. Sly-miR166 and Sly-miR319 are
components of the cold stress response in Solanum lycopersicum. Plant
Biotechnol Rep. 8:349–356. doi:10.1007/s11816-014-0326-3

• Varkonyi-Gasic E, Wu R, Wood M, Walton EF, Hellens RP. 2007. Protocol: a
highly sensitive RT-PCR method for detection and quantification of
microRNAs. Plant Methods. 3:12. doi:10.1186/1746-4811-3-12

• Wang CK, Han PL, Zhao YW, Ji XL, Yu JQ, You CX, Hu DG, Hao YJ. 2020.
Auxin regulates anthocyanin biosynthesis through the auxin repressor protein
MdIAA26. Biochem Biophys Res Commun. 533:717–722.
doi:10.1016/j.bbrc.2020.09.065

• Wang D, Wang S, Zhang D, Meng Y, Qian Y, Feng S, Bai Y, Zhou Y. 2025.
Identification and expression analysis of miR166 gene family in response to
salt stress in Chrysanthemum. Horticulturae. 11:141.
doi:10.3390/horticulturae11020141

• Wang N, Xu HF, Jiang SH, Fang HC, Su MY, Zhang ZY, Zhang TL, Chen
XS. 2018. Auxin regulates anthocyanin biosynthesis through the Aux/IAA–
ARF signaling pathway in apple. Hortic Res. 5:59. doi:10.1038/s41438-018-
0068-4

• Wang Y, Clevenger JP, Illa-Berenguer E, Meulia T, van der Knaap E, Sun L.
2019. A comparison of sun, ovate, fs8.1 and auxin application on tomato fruit
shape and gene expression. Plant Cell Physiol. 60:1067–1081.
doi:10.1093/pcp/pcz024

• Wei X, Wang C, Wang Y, Zhao Y, Ma Y, Liu S, Guan S, Jiao P. 2025.
miR166e/ZmATHB14 module contributes to drought tolerance in maize root.
Int J Biol Macromol. 297:139707. doi:10.1016/j.ijbiomac.2025.139707

- Yadav A, Kumar S, Verma R, Lata C, Sanyal I, Rai SP. 2021. MicroRNA166: an evolutionarily conserved stress biomarker in land plants targeting HD-ZIP family. Physiol Mol Biol Plants. 27:2471–2485. doi:10.1007/s12298-021-01096-x

- Zhang CH, Zhang BB, Ma RJ, Yu ML, Guo SL, Guo L. 2015. Isolation and expression analysis of four HD-ZIP III family genes targeted by microRNA166 in peach. Genet Mol Res. 14:14151–14161. doi:10.4238/2015.October.29.37

- Zhang J, Zhang H, Srivastava AK, Pan Y, Bai J, Fang J, Shi H, Zhu JK. 2018. Knockdown of rice microRNA166 confers drought resistance by causing leaf rolling and altering stem xylem development. Plant Physiol. 176:2082–2094. doi:10.1104/pp.17.01432

- Zhang Y, Pu Y, Zhang Y, Li K, Bai S, Wang J, Xu M, Liu S, Zhou Z, Wu Y, Hu R, Wu Q, Kear P, Du M, Qi J. 2024. Tuber transcriptome analysis reveals a novel WRKY transcription factor StWRKY70 potentially involved in potato pigmentation. Plant Physiol Biochem. 213:108792. doi:10.1016/j.plaphy.2024.108792

- Zhao C, Ma J, Zhang Y, Yang S, Feng X, Yan J. 2022. The miR166-mediated regulatory module controls plant height by regulating gibberellic acid biosynthesis and catabolism in soybean. J Integr Plant Biol. 64:995–1006. doi:10.1111/jipb.13253

- Zhou C, Yang N, Tian C, Wen S, Zhang C, Zheng A, Hu X, Fang J, Zhang Z, Lai Z, Lin Y, Guo Y. 2024. The miR166 targets CsHDZ3 genes to negatively regulate drought tolerance in tea plant (Camellia sinensis). Int J Biol Macromol. 264:130735. doi:10.1016/j.ijbiomac.2024.130735

- Zhou GK, Kubo M, Zhong R, Demura T, Ye ZH. 2007. Overexpression of miR165 affects apical meristem formation, organ polarity establishment and vascular development in Arabidopsis. Plant Cell Physiol. 48:391–404. doi:10.1093/pcp/pcm008

- Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. Nucleic Acids Res. 31:3406–3415.

 **Figure legends**

 **Figure 1. Identification, Validation, and Expression Analysis of miR166 in Potato**

 (A) RT-PCR of miR166 precursors from shoot apex, leaves and stolons tissues of SD

 grown *Solanum tuberosum* ssp. *andigena*. (B-D) Quantification of precursor-a, -b, and

 -c levels from stolon-to-tuber transition stages (stolon, swollen stolon and mini-tuber)

 harvested from potato plants post 10 days of SD or LD induction. Relative transcript

 abundance of the gene expression in SD tissue is compared to LD stolons. Asterisks

 (one, two, three, and four) indicate significant differences at $p < 0.05$, $p < 0.01$,

 $p < 0.001$, and $p < 0.0001$, respectively, using Student's t-test. ns = not significant at

 $p > 0.05$. *StEIF3e* was used as a reference gene. Detection (E) and relative fold change

 (F) of mature miR166 from leaves and stolon tissues of SD grown (7 days post

 induction, 7 dpi) *Solanum tuberosum* ssp. *andigena* by stem-loop RT-PCR. (G)

 Relative abundance of *miR166* in stolon-to-tuber transition stages. StU6 was used for

 normalization. Error bars indicate (±) SEM of three biological replicates, each with

 three technical replicates. Asterisks (one, two, three, and four) indicate significant

 differences at $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively, using

 Student's t-test. ns = not significant at $p > 0.05$. (H) Confirmation of HD-ZIP III

 targets of miR166. Partial mRNA sequence of *ICU1*, *ICU2*, *REV*, *PHB*, *PHV*, and

 *HB8* aligned with *miR166*, numbers denote the fraction of cloned cleavage products

 that terminate at different positions (arrows).

 **Figure 2. Knockdown of microRNA166 causes pleiotropic phenotypic changes** (A)

 Expression levels of *miR166* in mimicry lines (KD #2, KD #3) compared to WT. *StU6*

 was used as a reference gene. Error bars indicate standard error of means (SEM) from

 three biological replicates (n=3). Asterisks (one, two, three, and four) indicate

 significant differences at $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively,

 using Student's t-test. ns = not significant at $p > 0.05$. (B) Plant architecture of

 miR166 KD lines along with WT. Scale bar= 5 cm. Plant height (C), number of nodes

 (D) and internodal distance (E) of miR166 KD lines in comparison with WT (n=10).

 (F) Photographs of 8-week old LD grown plants, showing reduced leaf curvature in

 KD plants (arrows). Scale bar= 2 cm. (G) The longitudinal and transverse curvature

 indices of leaves emerging from the 5th-node of 8-week old plants (n=10). Asterisks

 (one, two, three and four) indicate significant differences at $p < 0.05$, $p<0.01$, $p<0.001$

 and $p<0.0001$, respectively, using one-way ANOVA. ns= not significant at $<0.05$. (H)

1140   Transverse sections of stem from miR166 KD and WT LD grown plants. Scale bar=

1141   50 microns.

1142

**Figure 3. miR166 knockdown lines have reduced tuber productivity** (A)

Representative tuber images from miR166 KD lines compared to WT, each image

contains tubers from 6 plants per line. Scale bar= 2 cm. Tuber weight per plant (B)

and average tuber number (C) of miR166 KD lines compared to WT. Data was taken

from soil grown plants after two months of SD induction, and plotted from 14

individual plants per line. The sphericity index (D) and average tuber volume (E) of

miR166 KD tubers in comparison to WT. The data is plotted for 30 tubers from each

line. Asterisks (one, two, three and four) indicate significant differences at $p < 0.05$,

$p<0.01$, $p<0.001$ and $p<0.0001$, respectively, using one-way ANOVA. ns= not

significant at <0.05. (F) Starch content (mg/gFW) of miR166 KD tubers, relative to

WT tubers. (G) Relative fold change of key tuber marker genes in the stolons (7 days

SD) of miR166 KD, compared to the WT. Error bars indicate standard error of means

(SEM) from three biological replicates (n=3). Asterisks (one, two, three, and four)

indicate significant differences at $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$,

respectively, using Student's t-test. ns $=$ not significant at $p > 0.05$. (H) *In vitro* tuber

induction assay for miR166 KD lines along with WT.

1159

**Figure 4. RNA-Seq analysis reveals differentially expressed genes in swollen and**

**sub-swollen part of tuberizing stolons** (A) Stolon-to-tuber transition stages as seen

in the WT and miR166 KD #3. (B) A schematic illustration to depict the tissue

collection carried out for RNA-sequencing analysis. The Head and sub-swelling (SSR)

region were collected individually from WT and KD #3 stolons from plants grown

under SD photoperiod for 12 days. (C) Venn diagram shows the summary of unique

and shared genes expressed in KD #3 and WT tissues (Head vs. SSR). Heat map was

plotted for key tuberization pathway genes (D), selective genes from QTLs (E), and

16 genes belonging to the *Ro* locus (F) from WT and KD #3. Numbers inside the box

represent variance-stabilized transformed (VST) values.

1170

**Figure 5. Tuberizing stolons exhibit differential expression of phytohormone-**

**associated genes and differential accumulation of auxin and GA** (A-B) The heat

map represents differential expression of genes involved in Auxin, GA and CK

metabolism and transport. Numbers inside the box represent variance stabilized

transformed (VST) values. (C) Validation of selective auxin biosynthesis genes from

the Head vs. SSR regions. *StEIF3e* was used as reference gene. Error bars indicate

standard error of means (SEM) from three biological replicates (n=3). Asterisks (one,

two, three, and four) indicate significant differences at $p < 0.05$, $p < 0.01$, $p < 0.001$,

and $p < 0.0001$, respectively, using Student's t-test. ns = not significant at $p > 0.05$.

Relative quantification of IAA- 3 (D) and $GA_3$ (E) from the Stolon, Head and SSR

tissues from WT and miR166 KD #3.

1182

**Figure 6. Anthocyanin pathway genes are upregulated in miR166 KD** (A) Tubers

accumulate anthocyanin pigments in their peels. (B) Quantification of Cyanidin 3-

glucoside, pelargonidin 3-glucoside and delphinidin 3-glucoside from the tuber peels

of WT and miR166 KD. The mean value (n = 4) ± SEM is presented. Asterisks (one,

two, three, and four) indicate significant differences at $p < 0.05$, $p < 0.01$, $p < 0.001$,

and $p < 0.0001$, respectively, using one-way ANOVA. ns = not significant at $p > 0.05$.

The RNA-seq analysis showed up regulation of several genes encoding enzymes in

the anthocyanin pathway (C) and some of the TF families having the strong

correlation with these anthocyanin pathway genes (D) (based on Zhang *et al*., 2024)

from the tuberizing stolons of miR166 KD, compared to the WT. Numbers inside the

box represent variance stabilized transformed (VST) values.

1194

**Figure 7. REVOLUTA affects tuberization** (A) Primary PCR product from

modified 5′ RLM-RACE, to confirm *REV* as a target of *miR166* in potato. (B)

Relative fold change of *REV* in differentiating stolons from WT plants. (C) Relative

fold change of *REV* in stolons from miR166 KD and OE lines. *StEIF3e* was used as

reference gene. Error bars indicate standard error of means (SEM) from three

biological replicates (n=3). Asterisks (one, two, three, and four) indicate significant

differences at $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively, using

Student's t-test. ns = not significant at $p > 0.05$. (D) Promoter expression of *StREV* in

*promStREV:uidA* transgenic lines. (i) *in vitro* plant, (ii) axillary node, (iii) shoot-

forming stolon, (iv) swollen stolon, (v) T.S. of swollen stolon, and (vi) mini-tuber.

Scale bar= 1000 um for (i) and (vi); Scale bar= 500um for (ii)- (v). Tuber productivity

and tuber numbers (E-G) from REV AS lines, relative to the WT. Scale bar= 2 cm.

Asterisks (one, two, three and four) indicate significant differences at $p < 0.05$, $p<0.01$,

p<0.001 and p<0.0001, respectively, using one-way ANOVA. ns= not significant at <0.05. (H) Promoters of auxin biosynthesis genes harbor the AT(C/G)AT sequence, a core motif of HD-ZIP III binding site. (I) Transient infiltration of *REV* in *S. tuberosum* cv. *Désirée* led to reduced expression of *StYUC-4/-7* and *-8*, relative to the control.

**Figure 8.** A proposed model for the role(s) of miR166-HD-ZIP III module in stolon-to-tuber transition in potato. miR166 in potato targets the HD-ZIP III transcripts. REVOLUTA could suppress the expression of auxin biosynthesis genes, and PHABOLUSA might regulate GA levels, similar to that in soybean (Zhao *et al.*, 2022). Phytohormone dynamics in a developing stolon could regulate the tuber shape in potato. Auxin-ARF5-HB8 and miR166-HB8/ICU1/2 could influence vascular differentiation in a tuberizing stolon.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Proposed model for the role(s) of miR166-HD-ZIP III module in potato tuber development

Figure 8