1   Title Page

Neural Sensitivity to Word Frequency Modulated by Morphological Structure: Univariate and Multivariate fMRI Evidence from Morphologically Complex Words

Joonwoo Kim[a,b], Solbin Lee[b], Kichun Nam[a,b*]

[a] School of Psychology, Korea University, Seoul, Republic of Korea

[b] Wisdom Science Center, Korea University, Seoul, Republic of Korea

[*] Correspondence
Kichun Nam

School of Psychology, Korea University

145, Anam-ro, Seongbuk-gu, Seoul, 02739, Republic of Korea

E-mail: kichun@korea.ac.kr

Abstract

A central question in psycholinguistics in visual word recognition is whether morphologically complex words are obligatorily decomposed into stems and affixes during visual word recognition or whether whole‑word access can occur when forms are frequent and familiar. The present study investigated how morphological complexity and lexical frequency jointly shape neural responses by leveraging Korean nominal inflection, whose transparent stem–suffix structure permits a clean dissociation between base (stem) frequency and surface (whole‑word) frequency. Twenty-five native Korean speakers completed a rapid event-related fMRI lexical decision task involving simple and inflected nouns that varied parametrically in both frequency measures. Representational similarity analysis (RSA) revealed robust encoding of surface frequency—but not base frequency—in the inferior frontal gyrus (IFG) pars opercularis and supramarginal gyrus (SMG), with significantly stronger correlations for inflected than simple nouns. Univariate analyses converged with this result: surface frequency selectively increased activation for inflected nouns in inferior parietal regions, whereas base frequency showed no reliable effects in any ROI. These findings challenge models positing obligatory pre-lexical decomposition, instead supporting accounts in which morphological processing is shaped by post-lexical, usage-driven lexical statistics. Taken together, our findings shed light on a distributed perspective on morphological processing, suggesting that structural and statistical factors jointly constrain access to morphologically complex forms.

1 **INTRODUCTION**

Human language comprehension requires efficient recognition of morphologically complex

words. A central question in psycholinguistics in visual word recognition is whether the

cognitive system obligatorily decomposes these words into morphemes before meaning access or

whether whole-word representations can be accessed directly for familiar forms.

Evidence for obligatory decomposition comes primarily from masked-priming studies,

where briefly presented primes like *hunter* facilitate recognition of *hunt*, even when semantically

opaque (e.g., *corner–corn*) relative to orthographic controls (Longtin et al., 2003; Rastle et al.,

2004). Such priming, with prime durations as brief as ~50 ms, suggests rapid morpho-

orthographic segmentation that precedes semantics. ERP and MEG work corroborates this view,

showing early negativities before semantic divergence (Lavric et al., 2011; Lewis et al., 2011). It

is important to note that the empirical and theoretical frameworks reviewed here pertain

specifically to visual word recognition, where orthographic structure provides the primary cues

to morphological segmentation. Phonological processing can exhibit different morphological

dynamics, particularly in languages with rich morphophonology, and therefore lies outside the

scope of the present investigation.

In contrast, distributional perspectives argue that processing depends on lexical statistics,

transparency, and predictability (Baayen et al., 2011; Bybee, 1995; Milin et al., 2017). High-

frequency or predictable forms may be accessed holistically, whereas rare or novel forms are

decomposed. In lexical decision and naming tasks, surface frequency, the token frequency of the

whole form, often better predicts recognition than base frequency, the cumulative frequency of

words sharing a stem (Bertram et al., 2000; Kuperman et al., 2009). Morphological priming

effects are likewise modulated by transparency and familiarity (Feldman et al., 2015; Jared et al.,

2017). Connectionist models further show how distributional learning can yield decomposition-

like effects without an explicit parsing mechanism (Plaut & Gonnerman, 2000). Together, these

findings suggest morphological processing may reflect both automatic segmentation and flexible,

frequency-driven representations.

Another critical distinction in morphological processing concerns the timing of

decomposition relative to lexical access. Pre-lexical decomposition models propose that complex

words are obligatorily parsed into constituent morphemes before whole-word meaning is

accessed (Rastle et al., 2004; Taft & Forster, 1975). Under this view, recognition of *hunter*

necessarily involves early access to *hunt + -er*, regardless of whole-word frequency. In contrast**,**

post-lexical decomposition accounts propose that morphological structure is analyzed *after* initial

whole-word recognition (Giraudo & Grainger, 2000, 2001). Here, frequent derived forms like

hunter may be accessed directly as whole words, with morphological decomposition occurring

later for semantic integration or production. Critically, these accounts make different predictions

about which frequency measure should modulate neural responses: pre-lexical models predict

that base (stem) frequency should drive activation patterns because stems are accessed first,

whereas post-lexical models predict that surface (whole-word) frequency should dominate

because initial access relies on whole-word representations. Notably, surface frequency itself can

be interpreted differently across frameworks: in pre-lexical models, it reflects the frequency of

recombining morphemes, whereas in post-lexical models, it indexes the frequency of stored

whole-word forms. The present study leverages fMRI to test these competing predictions by

examining whether sustained neural activation patterns are better explained by surface or base

frequency.

Neuroimaging evidence has shown mixed results regarding whether morphological

processing reflects categorical decomposition or graded statistical effects. fMRI studies report

that complex forms engage a left-lateralized network including the inferior frontal gyrus (IFG),

inferior parietal lobule (IPL), and ventral occipitotemporal cortex. Some findings have been

interpreted as supporting categorical decomposition, which showed selective IFG activations to

inflected or transparent derivations in English and Finnish (Bozic et al., 2007; Lehtonen et al.,

2006), and distinct activation patterns for regular versus irregular inflections in the left

frontotemporal network in German and English (Beretta et al., 2003; Tyler et al., 2004).

However, other work suggests that these effects reflect graded sensitivity to phonological

complexity, semantic factors, and lexical statistics rather than a unique decomposition stage

(Devlin et al., 2004; Joanisse & Seidenberg, 2005). When phonological complexity is controlled,

categorical differences between regular and irregular forms often disappear (Desai et al., 2006),

and when differences do emerge, they are modulated by semantic variables such as semantic

transparency (Bozic et al., 2013), suggesting continuous rather than discrete processing

mechanisms. While much of this work has focused on the regular versus irregular distinction in

verb inflection, a complementary question remains: at what level—whole-word or stem—does

the brain encode lexical frequency during morphological processing?

Findings on word frequency also remain inconsistent, with some studies reporting

stronger activation for low-frequency words (Fiebach et al., 2002; Fiez et al., 1999), and others

showing divergent patterns depending on task and subregion (Carreiras et al., 2009; Graves et al.,

2010). Critically, few studies have manipulated base frequency directly, and most report limited

independent effects once surface frequency is controlled (Lehtonen et al., 2006; Pliatsikas et al.,

2014), leaving unresolved whether the brain encodes stem-level or whole-word statistics during

morphological processing.

Korean morphology encompasses derivation, compounding, and inflection, but these

subtypes differ substantially in transparency and semantic complexity. Prior work has examined

derivational or compound morphology, which can involve semantic shifts or idiosyncratic affix

behavior (Kim, Lee, et al., 2025). The present study focuses specifically on nominal inflection,

where invariant stems combine transparently with case and topic markers. Korean inflectional

nouns have transparent stem–suffix structure and invariant stems, which together provide a

useful testing ground for understanding how morphological structure and lexical statistics

contribute to visual word recognition. Previous fMRI work shows that Korean readers are

sensitive to structural aspect of inflection, with effects predominantly centered at the left IFG,

consistent with the findings in Indo-European languages. For instance, inflected verbs that

impose greater morpho-syntactic demands elicited stronger inferior frontal activation relative to

inflected nouns, and violations of morphological category recruited left IFG subregions (Kim,

Pyun, et al., 2025; Kim & Nam, 2025). Studies directly comparing regular and irregular verb

inflections also reported equivalent level of activation in left IFG (Kim et al., 2024; Yim et al.,

2006).

On the other hand, emerging evidence suggests continuous sensitivity to lexical–semantic

and distributional factors. Kim et al. (2024) observed graded increases in occipitotemporal

activity for regular inflections relative to uninflected and irregular forms, with lexical-semantic

ambiguity (i.e., homonymy) further modulating activation in both occipitotemporal and angular

gyrus regions. In the temporal domain, ERP evidence from Korean inflected verbs showed that

both form-based and lexical-semantic properties emerge rapidly during visual word recognition

(Kim et al., 2022). They showed that whole-word frequency modulates early ERP components (N100/N250) and interacts with stem length at N400. High-frequency inflected verbs were accessed more holistically, whereas lower-frequency items showed stronger stem-length effects, a pattern consistent with frequency-contingent decomposition rather than uniform pre-lexical parsing.

Together, these findings suggest that, in Korean as in Indo-European languages, readers are sensitive to morphological structure, yet sustained neural responses are primarily shaped by post-lexical, usage-driven statistics. This pattern raises an important question about the representational level at which frequency information is encoded during the processing of inflectional morphology: whether neural responses track stem-based statistics, as predicted by pre-lexical decomposition accounts, or whole-word surface frequency, as predicted by post-lexical access models. To the best of our knowledge, no fMRI study has directly contrasted surface and base frequency in Korean nominal inflection, despite the transparent agglutinative structure that allows these frequency measures to be clearly dissociated.

Recent multivariate approaches offer powerful tools for examining how linguistic information is represented in the brain. Representational similarity analysis (RSA) relates neural dissimilarities to models derived from lexical statistics (Haxby et al., 2001; Kriegeskorte, 2008), enabling examination of graded representational structure. RSA has revealed fine-grained representations of syntactic, semantic, and sublexical features across languages (Bruffaerts et al., 2013; Fischer-Baum et al., 2017; Yue & Martin, 2021), allowing us to directly examine whether neural patterns reflect surface frequency, base frequency, or morphological complexity.

The present study employed a rapid event-related fMRI design with a lexical decision task to test whether brain activation patterns track morphological structure or statistical

frequency. Participants judged simple and inflected Korean nouns varying in both measures.

Korean provides a particularly suitable test case for several reasons. Its agglutinative

morphology yields transparent stem–suffix structure, stems remain invariant across inflections,

and orthographic conventions (i.e., spacing and syllable blocking) clearly indicate morphemic

boundaries, all of which permit precise frequency estimation. Crucially, inflected forms exhibit a

natural dissociation between surface frequency (i.e., the frequency of the whole form) and base

frequency (i.e., the frequency of the bare stem). For example, the stem 까닭 /kkadak/ "reason"

can combine with different particles (e.g., 까닭은, 까닭이, 까닭을); each inflected form has its

own surface frequency, but they all share the same base frequency — the cumulative count of the

stem 까닭. By contrast, simple (monomorphemic) nouns such as 옥수수 /oksusu/ "corn" have

no inflectional suffix, so the word form and the stem are identical and the two frequency

measures necessarily converge.

We applied complementary approaches—RSA and univariate mixed-effects models—to

determine whether neural responses to inflected Korean nouns primarily reflect pre-lexical

decomposition or post-lexical morphemic activation. If morphological parsing occurs at the pre-

lexical stage, then neural activation patterns should be more strongly explained by base

frequency, which indexes the frequency of accessing the stem independent of surface form. In

contrast, if inflected nouns are accessed primarily as whole-word units, consistent with post-

lexical accounts, then surface frequency should provide a better account of both amplitude-based

responses and the representational geometry captured by RSA. The present study tests whether

neural patterns align more closely with stem-driven decomposition or whole-word-driven

statistical access by directly contrasting surface and base frequency in Korean nominal inflection.

METHODS

Participants

Twenty-five healthy adults (15 female; age: 26 ± 3.83 years) recruited from Korea University

participated in the experiment. They were all native Korean speakers, right-handed (confirmed

by the scores on the Edinburgh Handedness Test (Oldfield, 1971): $M = 7.76$, $SD = 0.28$) and had

normal or corrected-to-normal vision. Participants were informed and provided written consent

prior to the experiment and were monetarily compensated for their participation. One participant

withdrew before completing the experiment, and two others were excluded due to poor

behavioral performance (accuracy $< 60\%$). One additional participant completed the experiment

with adequate behavioral performance, but their fMRI data could not be analyzed due to file

corruption resulting from hardware failure. Behavioral data from this participant were retained.

Final sample sizes were N = 22 (11 female; age: 26.3 ± 3.97 years) for behavioral analyses and N

= 21 (11 female; age: 26.4 ± 4.07 years) for all neuroimaging analyses. The sample size was

determined based on prior fMRI studies using similar multivariate and univariate ROI-based

approaches in the language domain (Bozic et al., 2007; Carota et al., 2016; Kim et al., 2024),

which typically report medium-to-large effects in comparable tasks with sample sizes ranging

from 16 to 24 participants. Given the within-subjects design and the use of trial-wise beta

estimation, this sample size affords adequate power to detect region-level effects of morphology

and lexical frequency using both traditional and multivariate analyses. The study was conducted

in accordance with the Declaration of Helsinki and approved by the Institutional Review Board

of Korea University (protocol code KUIRB-2021-0427-01, 21 December 2021).

Stimuli

A total of 120 Korean nouns were selected from the Korean Sejong corpus (Kang & Kim, 2009),

which contains approximately 15 million words. Korean is an agglutinative language with a

transparent morphological system, in which grammatical relations are marked by attaching case

particles or topic markers to noun stems. Furthermore, the Korean writing system (Hangul)

encodes syllables both orthographically and phonologically, enabling precise control of

sublexical units. These features make Korean particularly well-suited for isolating effects of

morphological complexity, stem-level access, and syllable-level frequency during visual word

recognition.

Morphological complexity was orthogonally manipulated such that half of the stimuli were

morphologically simple nouns, and the other half were inflected nouns. Each condition included

an equal number of bi-syllabic and tri-syllabic words, ensuring that word length was fully

balanced across conditions. Simple nouns consisted of bi- or tri-syllabic stems, while inflected

nouns were composed of mono- or bi-syllabic stems followed by a monosyllabic inflectional

suffix (e.g., case or topic marker). For example, 숭어 (trout) and 뱀은 (snake + TOPIC) illustrate

the contrast between simple and inflected forms, with 옥수수 (corn) and 까닭은 (reason +

TOPIC) showing the contrast between a tri-syllabic simple noun and a tri-syllabic inflected noun

consisting of a bi-syllabic stem (까닭) with the topic particle -은. The inflected nouns were

formed using three common Korean case/topic particles. Their distribution was as follows:

accusative –을/–를 (n = 29, 48.3%), topic –은/–는 (n = 21, 35.0%), and nominative –이/–가 (n =

10, 16.7%). This distribution closely aligns with the natural frequency distribution of case and topic in the given corpus (Kang & Kim, 2009).

Word frequency was operationalized using Zipf-measure (van Heuven et al., 2014), a log-transformed word frequency per billion words in the given corpus. Stem frequency was similarly transformed using the same Zipf-measure to allow for direct comparison across lexical levels. **Fig. 1** illustrates the frequency distributions for all words (**Fig. 1A**), simple nouns (**Fig. 1B**), and inflected nouns (**Fig. 1C**), with descriptive statistics summarized in **Table 1**.

Importantly, word frequency (Zipf) was matched across simple and inflected conditions ($t(59) = 1.62$; $p = .111$), whereas stem frequency (Zipf-stem) was not explicitly controlled. This was due to the structural properties of Korean: in simple nouns, the stem and surface word are identical, yielding the same frequency value. In contrast, inflected nouns contain an additional morpheme, such that the word frequency reflects the full surface form, while the stem frequency corresponds to the base noun. Given this natural divergence, Zipf and Zipf-stem were modeled separately in all statistical analyses, allowing us to dissociate effects of surface-level lexical access and stem-level frequency. To validate the dissociation between these two measures, we examined their distributional properties. Base frequency exhibited greater dispersion than surface frequency across all stimuli (coefficient of variation [CV]: 0.141 vs. 0.081) and within inflected nouns (CV: 0.103 vs. 0.090), a difference confirmed by Levene's tests (all words: $F = 47.22$, $p < .001$; inflected only: $F = 7.23$, $p = .008$). For inflected nouns, the two measures were moderately correlated at the item level (Pearson $r = .60$, $p < .001$), yet their representational similarity structures were only weakly related (Spearman $\rho = .24$, $p < .001$), indicating that surface and stem frequency cannot be treated as interchangeable. Additional representational-similarity validation is provided in **Fig. S1**. Although stem syllable counts necessarily differ

between conditions (simple nouns: 2–3 syllable stems; inflected nouns: 1–2 syllable stems +

monosyllabic suffix), total word length was fully balanced across conditions, reducing the

likelihood that sublexical length differences at the stem level confound the frequency effects

reported here. Finally, first-syllable frequency was also matched across conditions ($t(59) = 1.16$,

$p = .251$).

The same number of pseudowords ($N = 120$) were created to serve as fillers in the lexical

decision task. All pseudowords were manually generated by recombining syllable blocks drawn

from real Korean noun stems and inflectional suffixes. Each pseudoword was constructed to be

orthographically and phonologically legal in Korean while carrying no meaning, as verified

against the Standard Korean Dictionary. These pseudowords matched the real word stimuli in

both form and length (i.e., 2–3 syllable) and were evenly divided into morphologically simple

and inflected forms. Inflected pseudowords consisted of a real suffix combined with a pseudo-

stem while simple pseudowords contained only a pseudo-stem. All pseudowords were generated

by randomly combining syllables from real word stems and suffixes, which were

orthographically and phonologically legal but none of the pseudowords carried meaning

according to the Standard Korean Dictionary.

Finally, 30 non-linguistic masks, consisting of a string of asterisks (*****), were used as

a baseline condition for a comparison with experimental conditions. These masks were visually

matched to word stimuli in font and size, but contained no linguistic or orthographic information.

Participants responded to these items using the same lexical-decision buttons, allowing us to

estimate low-level visual and motor activation while controlling for perceptual properties of

lexical stimuli.

Table 1. Descriptive statistics of the stimuli set.

Condition	Word length	# Morphemes	Surface freq	Base freq	1st syllable freq
Simple	2.5	1	3.72	3.72	2.34
(*N* = 60)	(0)	(0)	(0.27)	(0.27)	(0.45)
Inflected	2.5	2	3.63	4.58	2.25
(*N* = 60)	(0)	(0)	(0.33)	(0.48)	(0.48)
Statistical comparisons	-	-	$t(59) = 1.62$ $p = .111$	$t(59) = -12.32^{***}$ $p < .001$	$t(59) = 1.16$ $p = .251$

Note. Word length = Word length in syllable; # Morphemes = Number of morphemes;
Surface freq = Zipf-scale word frequency; Base freq = Zipf-scale stem frequency;
1st syllable freq = log-transformed first syllable frequency per million.
Values in parentheses represent standard deviations.

Figure 1. Frequency distributions across morphological conditions. (A) All words (N = 120) show overlapping distributions for surface frequency (blue) and base frequency (red). (B) Inflected nouns (N = 60) show distinct distributions for surface and base frequency measures. Histograms use 15 bins; shaded regions indicate overlapping frequencies. (C) Simple nouns (N = 60) display only surface frequency, as surface equals base by definition for monomorphemic words.

Procedure

A rapid event-related fMRI design was employed using a visual lexical decision task.

Participants were instructed to respond, as quickly and accurately as possible, whether each

presented stimulus is a real word or not. Each trial began with a fixation cross (+) displayed for

100 ms, followed by a target stimulus presented at the center of the screen for 1000 ms. After

pressing response button, blank screen was shown during the inter-trial interval (ITI), which was

jittered with an average duration of 4000 ms (range: 1000–7000 ms). Stimuli were presented in a

pseudo-random order that included words, pseudowords, and a non-linguistic baseline mask. The

trial order and ITI durations were optimized using optseq2

(https://www.nitrc.org/projects/optseq). All visual stimuli were presented in white Courier New

font (size 34 pt) on a black background, controlled using E-Prime 2.0 Professional (Psychology

Software Tools, Pittsburgh, PA). Prior to the main experiment, participants completed a practice

session consisting of 12 randomly ordered trials. Participants proceeded to the main task only if

they achieved at least 70% accuracy during practice.

Image acquisition and preprocessing

MRI data were acquired at the Korea University Brain Imaging Center using a Siemens

Magnetom Trio 3T MRI scanner (Erlangen, Germany). Functional images were

acquired using T2*-weighted gradient EPI (echo-planar imaging) sequences (TR = 2000 ms;

TE = 20 ms; flip angle = 90∘; field of view = 240 mm; slice thickness = 3 mm; no gap for

42 slices; matrix size = 80×80; and voxel size = $3 \times 3 \times 3$ mm). A high-resolution structural

image was acquired for each participant using a T1-weighted structural images were acquired

with a 3D MP-RAGE (magnetization-prepared rapid gradient echo) sequence (TR = 1900 ms;

TE = 2.52 ms; flip angle = 90∘; field of view = 256 ms; matrix size = 256×256; and voxel size =

$1 \times 1 \times 1$ mm) covering the whole head.

Preprocessing of the fMRI data was performed using SPM12 (Wellcome Centre for

Human Neuroimaging, London, UK; https://www.fil.ion.ucl.ac.uk/spm/software/spm12/) in

conjunction with custom-built MATLAB scripts. The first three volumes of each session were

discarded to allow for magnetic field stabilization. The remaining functional images were first

realigned to the first volume to correct for head motion using a six-parameter rigid-body

transformation. Slice timing correction was applied to adjust for differences in acquisition timing

between slices, using the middle slice as reference. Each participant's structural image was then

co-registered to the mean functional image and segmented. The deformation fields obtained

during segmentation were subsequently applied to normalize the functional images to the

standard MNI space, with resampling to $3 \times 3 \times 3$ mm isotropic voxels. All normalized images

were smoothed with a 6 mm full-width at half-maximum (FWHM) Gaussian kernel to improve

signal-to-noise ratio. The six motion parameters from the realignment step were included as

nuisance covariates in all first-level general linear models (GLMs).

Behavioral data analysis

Behavioral analyses were conducted on 22 participants who completed the task with adequate

performance (accuracy $\geq 60\%$) and had valid behavioral data. We analyzed reaction times (RT)

using linear mixed-effects models (LMMs) and accuracy (ACC) using generalized linear mixed-

effects models with a binomial link. All analyses reported below were conducted on word trials

only, excluding pseudowords trials. For RT, only correct trials were included, and RTs were log-

transformed to reduce skewness. Separate models were constructed for whole-word frequency

(Zipf-word) and stem frequency (Zipf-stem), each including Morphological Condition (Simple

vs. Inflected) and their interaction as fixed effects. Random intercepts for participants and items

were included to account for repeated measures (Baayen et al., 2008). Accuracy data was

analyzed using a binomial logistic regression in a general linear mixed effects model. These

analyses were performed in R (R Core Team, 2022) using the lme4 package (Bates et al., 2015),

and the significance of fixed effects was assessed using Satterthwaite's approximation for

degrees of freedom, as implemented in the lmerTest package (Kuznetsova et al., 2017).

Whole-brain analysis

All neuroimaging analyses were conducted on 21 participants who met behavioral criteria and had valid fMRI data (see Participants section for exclusion criteria). First-level analyses modeled four contrasts for each participant: Inflected > Simple, Inflected > Baseline, Simple > Baseline, and Word > Baseline. The Inflected and Simple contrasts included only real word trials from their respective morphological conditions. The Word > Baseline contrast collapsed across all real word trials (both Inflected and Simple conditions), excluding pseudowords. Events were modeled with durations of 2 s and convolved with the canonical hemodynamic response function (HRF), and six motion parameters were included as nuisance regressors. At the second level, one-sample t-tests were performed on first-level contrast images. Whole-brain results were thresholded at a voxel-wise $p < .001$ (uncorrected) with a minimum cluster size of 30 voxels, consistent with exploratory standards in prior morphological and lexical processing fMRI studies (e.g., Kim et al., 2021; Newman et al., 2010). Activation patterns were further evaluated using anatomical masks from the Harvard-Oxford Atlas.

ROI selection

Our main analyses, including univariate contrasts and representational similarity analysis (RSA), focused on six anatomically defined, left-hemisphere regions of interest (ROIs, see **Fig. 6A**). These included two subregions of the inferior frontal gyrus (pars triangularis, pars opercularis), two temporal-occipital regions including the middle temporal gyrus (MTG), fusiform gyrus (FuG), and two regions within the inferior parietal lobule (IPL): the angular gyrus (AG) and supramarginal gyrus (SMG). ROI selection was motivated by prior literature implicating these

regions in morphological and lexical-semantic processing (Binder et al., 2009; Bulut, 2022;

Vigneau et al., 2006), allowing us to examine convergent and dissociable effects across multiple

analytical frameworks. All ROIs were defined using the probabilistic Harvard-Oxford cortical

atlas (1 mm resolution). Each region was thresholded at 25% probability and binarized and

resampled to match functional image resolution (3 mm) using SPM12. All masks were visually

inspected to ensure accurate anatomical localization.

Representational Similarity Analysis (RSA)

Representational similarity analysis (RSA) was used to test whether local activation patterns

reflected graded similarity based on word or stem frequency. All representational similarity

analyses (RSA) were implemented using custom Python scripts in combination with scikit-learn

(Pedregosa et al., 2011), SciPy (Virtanen et al., 2020), and Nilearn (Abraham et al., 2014).

Analyses were performed within the six anatomically defined ROIs.

Six theoretical representational dissimilarity matrices (RDMs) were constructed based on

continuous lexical frequency measures: surface-level word frequency (Zipf) and base-form stem

frequency (Zipf-Stem). Each RDM encoded the absolute pairwise frequency difference between

trials (i.e., Euclidean distance in a 1D lexical space), producing symmetrical matrices where

larger values reflect greater dissimilarity in frequency. For each frequency measure, we created

separate model RDMs for three trial subsets: (1) all word trials (**Figs. 2A, 2D**), (2)

morphologically inflected nouns (**Figs. 2B, 2D**), and (3) only simple nouns (**Figs. 2C, 2F**). This

yielded six theoretical RDMs in total (2 frequency models $\times$ 3 morphological conditions).

Figure 2. Theoretical frequency-based representational dissimilarity matrices (RDMs). Each matrix encodes the
pairwise lexical distance that served as a model predictor in the RSA. Word-frequency RDMs (Zipf-word; panels A–
C) and stem-frequency RDMs (Zipf-Stem; panels D–F) were computed as the absolute difference (i.e., Euclidean
distance in a 1-D frequency space) in Zipf-scale frequency between two items, yielding symmetrical matrices whose
values increase with greater frequency disparity. For both frequency measures we constructed three matrices: all
nouns, only morphologically inflected nouns, and only simple nouns. Warmer colors denote larger frequency gaps;
cooler colors denote smaller gaps.

To construct neural RDMs, we extracted trial-wise beta images per participant and ROI

using NiftiMasker (standardized across voxels) and computed pairwise Pearson correlations

between activation vectors for each pair of trials. Neural dissimilarity was defined as 1 minus

Pearson r, resulting in a neural RDM for each subject, ROI, and condition (All, Simple,

Inflected). Spearman rank correlation was used to compute the correspondence between neural

and model RDMs. Correlation values were computed separately for each subject, ROI, condition,

and model.

At the group level, we assessed whether neural RDMs significantly aligned with

theoretical frequency structure using one-sample t-tests against zero (within-subject Fisher-z

transformed Spearman values). To test whether representational structure differed across

morphological conditions, we performed paired-sample *t*-tests comparing Simple vs. Inflected

correlations within each ROI and frequency model. All resulting *p*-values were corrected for

multiple comparisons using the false discovery rate (FDR) method (Benjamini & Hochberg,

1995).

Univariate analysis

To examine morphological effects and lexical frequency modulations within key language-

related regions, we conducted ROI-based univariate analyses. Trial-wise beta values were

extracted using MarsBaR (Brett et al., 2002) and submitted to linear mixed-effects models in R.

Separate models were estimated for each frequency measure (Zipf or Zipf-stem) and ROI,

with fixed effects for morphological condition (simple vs. inflected), frequency, and their

interaction. Random intercepts were included for both participants and items to account for

repeated measures and individual variability. These models assessed whether frequency effects

varied by morphological complexity and whether either measure explained additional variance in

regional activation.

RESULTS

Behavioral performance

Fig. 3 presents mixed-effects model predictions for reaction time and accuracy as

functions of surface (Zipf-word; **Fig. 3A**) and base frequency (Zipf-stem; **Fig. 3B**). In both the

Zipf-word and Zipf-stem models for RT, morphological condition did not show a significant

main effect (Zipf-word: $\beta = -0.027$, $SE = 0.036$, $t = -0.85$, $p = .399$; Zipf-stem: $\beta = -0.028$, $SE =$

0.034, $t = -0.82$, $p = .416$), nor was the interaction between morphology and frequency

significant (Zipf-word: $\beta = 0.009$, $SE = 0.01$, $t = -0.94$, $p = .347$; Zipf-stem: $\beta = 0.01$, $SE = 0.009$,

$t = 1.215$, $p = .227$). However, both frequency predictors showed significant main effects, with

faster responses to higher-frequency words. Word frequency (Zipf-word) was associated with

reduced RTs ($\beta = -0.03$, $SE = 0.01$, $t = -3.08$, $p = .003$; **Fig. 3A**), and the same pattern was

observed for stem frequency ($\beta = -0.026$, $SE = 0.009$, $t = -3.02$, $p = .003$; **Fig. 3B**).

In the accuracy model, no significant effects were found for morphological condition or

its interaction with frequency (all $ps > .83$; **Fig. 3**). Both word frequency ($\beta = 0.787$, $SE = 0.388$,

$z = 2.03$, $p = .043$; **Fig. 3A**) and stem frequency had a significant main effect ($\beta = 0.699$, $SE = $

0.34, $z = 2.07$, $p = .038$; **Fig. 3B**), suggesting faster latencies and greater accuracy for items with

higher frequencies of either the stem or the whole-form.

Figure 3. Mixed-effect model predictions for behavioral performance. Predicted reaction times (left y-axes), log-
transformed milliseconds and predicted probability of correct response (right y-axes) are shown as functions of (A)
word frequency (Zipf-word) and (B) stem frequency (Zipf-stem). Blue lines represent inflected nouns; yellow lines
represent simple nouns; shaded ribbons indicate 95% confidence intervals (CIs).

Whole-brain analysis results

Table 2 and **Fig. 4** present the results of the univariate whole-brain analysis. All contrasts were

computed at a voxel-level threshold of $p < .001$ (uncorrected) with a cluster extent threshold of k

> 30. Anatomical labels were assigned using the Harvard-Oxford cortical atlas (HOA), and all

reported clusters exceeded the statistical threshold.

When collapsed across all word conditions (*Word > Baseline*), robust activation was

observed in a broad left-lateralized network associated with visual and lexical-semantic

processing. Significant clusters were found in the left inferior frontal gyrus (pars triangularis and

pars opercularis), frontal orbital cortex, temporal occipital fusiform cortex, and lateral occipital

cortex (inferior division) bilaterally, with peak activity extending into the occipital pole.

The contrast Inflected > Rest revealed a largely overlapping activation pattern, including

the left inferior frontal gyrus, middle and posterior temporal regions, temporal occipital fusiform

cortex, frontal orbital cortex, and paracingulate gyrus. In addition, bilateral lateral occipital

cortex (inferior division) showed significant engagement, reflecting visual processing demands.

The Simple > Rest contrast similarly activated the inferior frontal gyrus, frontal orbital

cortex, and temporal occipital fusiform cortex, along with strong bilateral activation in the lateral

occipital cortex and occipital pole. However, the extent of frontal and temporal involvement was

more restricted compared to the inflected condition.

Notably, the direct contrast Inflected > Simple did not yield any suprathreshold clusters,

suggesting that the two morphological conditions engaged similar neural systems when

contrasted directly, despite observable differences in activation extent relative to baseline.

Together, these results indicate that both simple and inflected nouns recruited a left-

lateralized fronto-temporo-occipital network, with no statistically significant differences in

overall activation magnitude between conditions at the whole-brain level. However, when

examined relative to baseline, inflected nouns additionally recruited middle and posterior

temporal regions. This might reflect additional lexical-semantic processing demands associated

with morphologically complex forms, which carry richer combinatorial meaning (e.g., case or

topic information) than monomorphemic nouns.

Figure 4. Whole-brain task activation for lexical processing. Brain activation maps depict voxels significantly
engaged during word recognition in the whole-brain level (voxel-level threshold uncorrected $p < .001$, cluster extent
threshold $k \geq 30$). Red = Inflected > Baseline, Blue = Simple > baseline, and Magenta = overlap of the two
contrasts.

1   **Table 2**. MNI coordinates of significant activation identified in the whole-brain analysis.

Brain Regions		Peak MNI coordinates	Z	Cluster Size
Word > Rest				
Lateral Occipital Cortex (inferior), Occipital Pole, Occipital Fusiform Gyrus	LH	[–30, –91, –4]	8.2	2321
Lateral Occipital Cortex (inferior), Occipital Pole	RH	[33, –88, –1]	7.4	2002
Frontal Orbital Cortex	LH	[–42, 26, –16]	6.8	145
Inferior Frontal Gyrus (pars triangularis, opercularis)	LH	[–48, 26, 17]	6.5	1395
Temporal Occipital Fusiform Cortex	LH	[–43, –49, –22]	5.8	293
Inflected > Rest				
Lateral Occipital Cortex (inferior), Occipital Pole, Occipital Fusiform Gyrus	LH	[–30, –91, –4]	7.6	2293
Temporal Occipital Fusiform Cortex	LH	[–42, –46, –22]	6.5	391
Lateral Occipital Cortex (inferior)	RH	[33, –88, –1]	6.4	1421
Inferior Frontal Gyrus (pars triangularis, opercularis)	LH	[–48, 26, 17]	6.3	1759
Middle Temporal Gyrus (post)	LH	[–49, –43, 2]	5.9	430
Frontal Orbital Cortex	LH	[–40, 20, –7]	5.4	240
Paracingulate Gyrus	LH	[–3, 27, 42]	5.2	347
Simple > Rest				
Lateral Occipital Cortex (inferior), Occipital Pole, Occipital Fusiform Gyrus	LH	[–30, –91, –4]	7.7	2388
Lateral Occipital Cortex (inferior), Occipital Pole	RH	[33, –88, –1]	7.3	2732
Frontal Orbital Cortex	LH	[–42, 26, –16]	6.6	228
Inferior Frontal Gyrus (pars triangularis, opercularis)	LH	[–48, 26, 15]	6.3	1488
Temporal Occipital Fusiform Cortex	LH	[–43, –49, –22]	5.2	234

2   *Note.* Anatomical labels are from the Harvard–Oxford Cortical Atlas. All clusters survive whole-brain correction at
3   voxel-level threshold of uncorrected $p < .001$ and cluster extent threshold of $k \geq 30$. LH = left hemisphere; RH =
4   right hemisphere.
5

6   *RSA results*

7   The results of representational similarity analysis (RSA) are presented in **Fig. 5**. We assessed

8   whether local multivoxel patterns reflected lexical frequency structure by correlating neural

9   RDMs with theoretical RDMs based on either whole-word frequency (Zipf) or stem frequency

10   (Zipf-Stem). RSA was conducted within six anatomically defined ROIs (IFGoper, IFGtri, AG,

SMG, MTG, FuG). Group-level RSA effects were tested using one-sample t-tests (against zero)

for each model (i.e., Zipf and Zipf-Stem), condition (All, Inflected, Simple), and ROI. To

directly compare encoding strength across conditions, we additionally conducted paired two-

sample t-tests between Inflected and Simple nouns. All p-values were corrected using false

discovery rate (FDR) correction.

In RSA using the word frequency (Zipf-word) model (**Fig. 5A**), several ROIs showed

significant encoding for word frequency. One-sample t-tests revealed significant encoding of

Zipf frequency for inflected nouns in IFGoper ($t(20) = 3.30$, $pFDR = .025$). Notably, AG ($t(20) =$

2.50, uncorrected $p = .021$) and SMG ($t(20) = 2.22$, uncorrected $p = .038$) also showed trend-

level effects for inflected nouns, although these did not survive FDR correction. In contrast, RSA

correlations for simple nouns significantly differed from zero in IFGoper ($t(20) = -3.80$, $pFDR$

$= .020$) and SMG ($t(20) = -3.33$, $pFDR = .025$), indicating negative similarity effects, such that

items with more similar surface frequency patterns produced more dissimilar neural patterns.

Paired t-tests revealed significantly stronger encoding of word frequency for inflected compared

to simple nouns in IFGoper ($t(20) = 6.44$, $pFDR = .0001$) and SMG ($t(20) = 3.95$, $pFDR = .0096$),

suggesting that morphological complexity enhances the neural encoding of frequency

information in these regions.

In contrast, RSA using the stem frequency model (Zipf-Stem; **Fig. 5B**) showed generally

weaker effects. IFGoper ($t(20) = -3.80$, $pFDR = .020$) and SMG ($t(20) = -3.33$, $pFDR = .025$)

showed significant negative RSA correlations for simple nouns. Trend-level negative RSA

effects were observed in SMG for the All condition ($t(20) = -2.41$, uncorrected $p = .026$).

However, no ROI showed significant above-zero encoding of Zipf-Stem values in the All or

Inflected conditions, nor were significant differences found between inflected and simple nouns.

In sum, RSA revealed that whole-word frequency (Zipf) was significantly encoded in

frontoparietal regions, especially IFGoper and SMG, with stronger encoding for inflected

compared to simple nouns. Stem frequency (Zipf-Stem), in contrast, was only negatively

encoded in IFGoper and SMG for simple nouns, suggesting that morphological complexity

enhances sensitivity primarily to whole-word rather than stem-level frequency structure.

Figure 5. Representational similarity analysis (RSA) results. Mean Spearman ρ values between neural RDMs and
(A) word-frequency (Zipf) or (B) stem-frequency (Zipf-Stem) model RDMs are plotted for each ROI (blue =
Inflected; yellow = Simple); vertical bars represent ± SEM. Significance of each individual bar was tested with a
one-sample t-test against zero and is indicated directly above the bar (* $p < .05$; ** $p < .01$; *** $p < .001$; †
uncorrected $p < .05$, FDR-corrected within panel). Horizontal brackets linking the blue-yellow bar pairs denote
paired-sample t-tests that compared Inflected and Simple correlations within the same ROI, using the same symbol
conventions for significance.

Univariate analysis results

The results of the ROI analysis and illustration are presented in **Fig. 6**. We examined whether

morphologically simple versus inflected nouns elicited differential activation across six

predefined ROIs and whether this activation was modulated by surface-level (Zipf-word) (**Fig.**

6B) or stem-level (Zipf-stem) frequency (**Fig. 6C**).

Our primary analysis focused on Zipf frequency. We observed significant main effects of

word frequency in both regions in the inferior parietal lobule (IPL), including the angular gyrus

($\beta = 0.897$, *SE* = 0.39, *t* = 2.3, $p = .023$) and supramarginal gyrus ($\beta = 0.915$, *SE* = 0.397,

t = 2.3, $p = .023$), indicating increased activation for higher-frequency words. No other regions

showed significant frequency effects after FDR correction.

Importantly, a significant interaction of Condition and Zipf-word emerged in the SMG

($\beta = -1.30$, *SE* = 0.634, *t* = –2.06, $p = .042$), reflecting stronger frequency effects for

inflected compared to simple nouns. The AG also showed a marginal interaction ($\beta = -1.14$,

SE = 0.624, *t* = –1.83, $p = .07$), although this did not survive correction. No significant main

effects or interactions were found in IFG (pars opercularis/triangularis), or FuG.

We then tested the influence of stem frequency (Zipf-stem). A significant main effect was

observed in the angular gyrus ($\beta = 0.685$, *SE* = 0.27, *t* = 2.54, $p = .013$), indicating greater

activation for higher-frequency stems across conditions, though this effect did not remain

significant after correction. No other regions showed reliable stem frequency effects or

interactions with morphological condition. Importantly, SMG, which showed a robust interaction

with surface frequency, did not exhibit significant modulation by stem frequency ($\beta = -0.788$,

SE = 0.573, *t* = –1.37, $p = .172$).

Notably, as visible in **Figs. 6B–6C**, the highest frequency bin for simple nouns shows

wider confidence intervals. This pattern reflects the lexical distribution of our stimuli set. As

simple nouns equate surface and base frequency, they occupy a compressed frequency range and

only a small number of items fall into the highest frequency bin (see **Fig. 1**). The reduced item

variability in this range produces greater between-item variance and correspondingly wider

intervals, likely reflecting the combination of a small number of high-frequency items and the

inherently skewed frequency distributions observed in large lexical decision megastudies

(Brysbaert et al., 2018; Keuleers et al., 2011; Yap & Balota, 2009). Importantly, this variance

does not alter the overall pattern that surface-frequency effects are selectively and more robustly

expressed for inflected nouns.

Together, these findings suggest that surface-level frequency exerts a stronger and more

morphology-dependent influence on activation in parietal regions than stem-level frequency,

consistent with a dissociation between surface-form and base-form frequency effects in

morphological processing, in line with post-lexical accounts.

Figure 6. Regions of interest univariate analysis results. (A) Locations of anatomically defined ROIs: IFG pars opercularis (IFGoper), IFG pars triangularis (IFGtri), supramarginal gyrus (SMG), angular gyrus (AG), middle temporal gyrus (MTG), and fusiform gyrus (FuG). (B) Predicted BOLD β as a function of Zipf word frequency for inflected (blue) and simple (yellow) nouns, estimated from linear mixed-effects models. The Condition × Frequency interaction was significant in SMG ($p = .042$) and marginal in AG ($p = .07$). (C) The same analysis for Zipf stem frequency. Shaded ribbons represent 95% confidence intervals. $^{*}p < .05$; $^{\dagger}p < .10$.

DISCUSSION

The present study investigated how morphological complexity and lexical frequency jointly

shape neural responses during visual word recognition, focusing on the dissociation between

surface (whole-word) frequency and base (stem) frequency in Korean nominal inflection. Korean

provides an interesting test case because its transparent agglutinative structure allows stem-based

and surface-based statistics to be cleanly separated. By combining univariate mixed-effects

modeling with representational similarity analysis (RSA) across six left-hemisphere language

regions, we tested competing predictions of pre-lexical decomposition and post-lexical, whole-

form access accounts. Three main findings were demonstrated. First, RSA revealed robust

representational encoding of surface frequency in IFG pars opercularis (IFGoper) and

supramarginal gyrus (SMG), with encoding significantly stronger for inflected than simple nouns.

Second, base frequency did not show positive encoding in any ROI; instead, simple nouns

produced negative RSA correlations with the base-frequency RDM in IFGoper and SMG. Third,

univariate analyses in the inferior parietal lobule (IPL) revealed significant interactions of

morphology and surface frequency, with activation increasing with surface frequency for

inflected but not simple nouns. Collectively, these findings indicate that sustained neural

activation patterns primarily reflect whole-form lexical statistics rather than stem-based

morphological structure.

These results are in line with prior work in both Indo-European and Korean languages,

while extending this literature by comparing surface- and stem-based frequency within a single

inflectional system. Consistent with masked-priming and ERP evidence showing that high-

frequency complex forms can be accessed more holistically (Kim et al., 2022; Lehtonen et al.,

2007), our RSA and univariate findings demonstrate that surface frequency reliably modulates

activation patterns in frontoparietal regions. Behavioral processing advantages for high-

frequency inflected forms have been widely documented in lexical decision and naming tasks

(Bertram et al., 2000; Kuperman et al., 2009), and our data converge with these findings by

showing that surface frequency, but not stem frequency, shapes neural similarity and activation

magnitude.

Neuroimaging studies have similarly implicated graded lexical-semantic and statistical

influences on morphological processing. In languages such as English, Hebrew, Finnish, and

Italian, fMRI work has shown that putative morphological effects often reduce to or interact with

frequency, predictability, or semantic transparency (Bozic et al., 2013; Desai et al., 2006; Devlin

et al., 2004). Our results extend this observation to Korean nominal inflection; despite its

transparent morphological structure, the neural similarity in IFGoper and SMG was more

strongly associated with whole-word frequency than with stem-based factors.

These findings refine our understanding of Korean morphological processing and situate

it within broader cross-linguistic patterns. Prior fMRI research on Korean verb inflection has

consistently shown overlapping activation for regular and irregular verbs in left IFG and

temporal regions (Kim et al., 2024; Yim et al., 2006), suggesting that the processing of Korean

inflectional morphology may not solely depend on categorical dichotomy. At the same time,

these studies observed graded modulations by semantic homonymy and morphological

complexity in occipitotemporal and angular gyrus regions (Kim et al., 2024), indicating

sensitivity to experience-based properties. Our findings parallel this pattern in showing that

although morphological complexity (simple vs. inflected) modulates representational strength,

the representational structure was shaped primarily by surface-level lexical statistics.

The behavioral results complement the neural findings, further specifying how lexical

statistics shape recognition of inflected forms. Both surface and base frequency facilitated lexical

decisions, and neither interacted significantly with morphological complexity. This aligns with

large lexical-decision megastudies showing that whole-word frequency is the primary predictor

of behavioral performance (Keuleers et al., 2011; Yap & Balota, 2009; Yi et al., 2017).

Importantly, the absence of a morphology $\times$ frequency interaction behaviorally parallels the

pattern in the fusiform and MTG ROIs, which showed no reliable modulation by frequency.

The neural data, however, reveal distinctions that remain behaviorally silent. Whereas

both frequency measures facilitate decision speed, only surface frequency shapes

representational similarity in IFGoper and SMG, and only surface frequency interacts with

morphology in parietal univariate activation. This dissociation suggests that behavioral measures

and neural representational measures capture distinct aspects of word recognition; behavioral

responses index the efficiency of the recognition and decision processes at the later stage of

lexical access, whereas RSA and ROI activation reveal the internal representational architecture

supporting this process. Inflected nouns with higher surface frequency may be accessed more

efficiently behaviorally as they possess more stable or more distinct neural representations in

IFGoper and IPL. Thus, the convergence between behavioral facilitation and neural

representational structure strengthens the interpretation that surface frequency indexes a

privileged pathway for accessing inflected forms.

A central theoretical question concerns what surface frequency encoding indeed

represents neurally. In pre-lexical decomposition accounts, lexical access is assumed to proceed

via early and obligatory segmentation into morphemic units, such that base frequency —

reflecting the resting activation of stored stem representations — serves as a primary determinant

of processing efficiency (Taft & Forster, 1975). Surface frequency, in this framework, is not

expected to exert a dominant influence during initial processing stages; rather, its effects would

emerge only at later stages where decomposed morphemes are recombined and matched against

whole-word representations. In contrast, post-lexical or full-form access accounts hold that

surface frequency directly indexes the strength of stored whole-word lexical entries, with higher-

frequency forms benefiting from more robust representations and thus more efficient retrieval.

Two aspects of our findings support the latter interpretation.

First, when examined within the inflected-noun stimuli, surface and base frequency

demonstrated only moderate correlation and produced clearly distinct representational

geometries (see **Fig. S1**). If surface-frequency effects were driven by stem-level statistics, one

would expect base frequency to exhibit parallel RSA encoding, which it did not. Second,

surface-frequency encoding localized primarily to IFGoper and SMG, regions implicated in

lexical retrieval, semantic integration, and competition resolution (Binder et al., 2009; Lau et al.,

2008; Thompson-Schill et al., 1997), rather than in occipitotemporal regions, which has been

functionally linked to early morpho-orthographic segmentation (Gold & Rastle, 2007; Solomyak

& Marantz, 2010). This spatial dissociation implies that the RSA patterns reflect downstream

lexical-semantic processes tied to whole-form access rather than sublexical parsing.

Another interpretation can be derived from the Augmented Addressed Morphology

model (AAM; Caramazza et al., 1988) proposes that lexical access can proceed via both whole-

word representations and morphological decomposition, with the relative contribution of each

route depending on factors such as frequency and representation strength. Under this framework,

high-frequency inflected forms are preferentially accessed via whole-word representations,

whereas lower-frequency forms are more likely to engage decomposition processes. From this

perspective, the surface-frequency effects observed in IFGoper and SMG may reflect the degree

to which specific inflected forms have developed robust whole-word representations through

repeated exposure. Inflected nouns with high surface frequency may function as independent

lexical entries whose representational geometry is shaped by cumulative experience, consistent

with the AAM's prediction that frequency modulates the relative efficiency of the whole-word

access route. This interpretation aligns with experience-based and connectionist accounts in

which lexical representations emerge from statistical regularities in form–meaning mappings

rather than from strictly compositional morpheme-based storage (Baayen et al., 2011; Bybee,

1995; Plaut & Gonnerman, 2000).

Region-specific patterns further refine understanding of the neural architecture

supporting morphological processing. IFG pars opercularis showed robust RSA encoding of

surface frequency for inflected nouns, suggesting that this region participates in representing

graded lexical statistics rather than merely implementing competitive selection or morpho-

syntactic parsing as traditionally assumed. This finding aligns with work linking posterior IFG to

the integration of form and meaning under conditions of lexical competition or ambiguity

(Thompson-Schill et al., 1997; Whitney, Jefferies, et al., 2011).

In the IPL, both SMG and AG showed univariate interactions of morphology and surface

frequency, with increasing activation for higher-frequency inflected forms. IPL has long been

associated with lexical–semantic retrieval and the integration of distributed semantic features

(Binder et al., 2009; Price et al., 2015), as well as with controlled semantic access in tasks

requiring selection or feature weighting (Graves et al., 2010; Whitney, Jefferies, et al., 2011).

Our findings reinforce this functional characterization by indicating that IPL encodes the

accessibility or retrieval strength of whole forms rather than engaging in decomposition of stems.

In contrast, MTG showed weak or inconsistent frequency effects. Given its role in

semantic control and relational processing (Davey et al., 2016; Whitney, Kirk, et al., 2011),

MTG activation may emerge more strongly in tasks with greater semantic integration demands

such as semantic judgment rather than in lexical decision. Additionally, although whole-brain

analyses identified fusiform activation for all word stimuli, our anatomically defined fusiform

ROI did not show reliable frequency modulation. This likely reflects the broad mask used, which

spans subregions with distinct functional profiles. Only mid-fusiform cortex reliably shows

graded frequency and lexicality effects; more precise functional ROIs might reveal effects not

visible here.

The present findings offer support for whole-word access accounts over obligatory

decomposition models, which predict robust stem-frequency effects and decomposition

signatures in all morphologically complex words (Rastle et al., 2004; Taft & Forster, 1975).

Although early ERP studies support rapid form-based segmentation, such processes may be

transient and not reflected in sustained BOLD patterns. Our results instead align with flexible or

hybrid models in which whole-word access pathways are strengthened by lexical experience and

decomposition is engaged selectively when whole-form access is inefficient (Baayen et al., 2011;

Milin et al., 2017).

The results are also compatible with the broader view that morphological computation is

distributed and usage-dependent rather than categorical and rule-governed. For Korean nominal

inflection, surface frequency shapes both representational structure and activation dynamics,

suggesting that frequent inflected forms may become entrenched as distinct lexical

representations. This interpretation aligns with usage-based and connectionist perspectives in

which morphological structure emerges from learned statistical regularities (Bybee & Hopper,

2001; Plaut & Gonnerman, 2000), as well as with the AAM framework, which predicts that high-frequency inflected forms can be accessed via a direct whole-word route, effectively functioning as independent lexical entries (Caramazza et al., 1988).

Some limitations should be acknowledged. First, temporal resolution of fMRI limits direct observation of rapid pre-lexical segmentation processes occurring within the first approximately 200 ms of word processing, the processing stage indexed by masked priming and early ERP components. Early form-based segmentation may nonetheless occur transiently and be superseded by the whole-form access patterns visible in sustained BOLD responses. Multimodal recordings combining EEG/MEG with fMRI would be necessary to adjudicate whether pre- and post-lexical decomposition pathways operate sequentially during inflected word recognition.

Relatedly, stem syllable counts necessarily differ between simple and inflected conditions due to the addition of a monosyllabic suffix in inflected forms. Although total word length was matched across conditions and first-syllable frequency did not differ, we cannot entirely rule out that differences in stem length contributed to the observed morphology × frequency interactions. However, if stem length were driving the effects, one would expect base (stem) frequency—which directly indexes stem-level processing—to show stronger neural encoding for inflected nouns with shorter stems, which was not observed.

Additionally, our sample size ($N = 21$) and anatomically defined ROI approach afford adequate power for medium-to-large effects but may miss subtle regional interactions, particularly in heterogeneous regions such as MTG and mid-fusiform cortex. Similarly, the lexical-decision task emphasizes decisional speed and accuracy rather than automatic comprehension; converging evidence from speeded naming, semantic judgment, or passive reading tasks would strengthen claims about the generality of surface-frequency effects.

CONCLUSION

In conclusion, the present findings provide the first fMRI evidence comparing surface and base frequency effects on neural representational structure in morphologically complex Korean words. The results demonstrate that morphological complexity modulates neural encoding of surface rather than base frequency and that morphological condition does not yield discrete category representations in classic language areas. These findings argue against purely decompositional theories and instead support distributional or hybrid models in which both decomposition and whole-form access routes exist, with their engagement determined by lexical statistics such as surface frequency. Rather than reflecting an obligatory parsing process, morphological processing emerges from dynamic interplay between a word's structure and its statistical properties, shaped by both form-based and meaning-based constraints (Weissbart & Martin, 2024).

REFERENCES

Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language*, *59*(4), 390–412. https://doi.org/10.1016/j.jml.2007.12.005

Baayen, R. H., Milin, P., rević, D. F., Hendrix, P., & Marelli, M. (2011). An Amorphous Model for Morphological Processing in Visual Comprehension Based on Naive Discriminative Learning. *Psychological Review*, *118*(3), 438–481. https://doi.org/10.1037/a0023851

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: a Practical and Powerful Approach to Multiple Testing. In *J. R. Statist. Soc. B* (Vol. 57, Issue I). https://academic.oup.com/jrsssb/article/57/1/289/7035855

Beretta, A., Campbell, C., Carr, T. H., Huang, J., Schmitt, L. M., Christianson, K., & Cao, Y. (2003). An ER-fMRI investigation of morphological inflection in German reveals that the brain makes a distinction between regular and irregular forms. *Brain and Language*, *85*(1), 67–92. https://doi.org/10.1016/S0093-934X(02)00560-6

Bertram, R., Schreuder, R., & Baayen, R. H. (2000). The Balance of Storage and Computation in Morphological Processing: The Role of Word Formation Type, Affixal Homonymy, and Productivity. *Journal of Experimental Psychology: Learning Memory and Cognition*, *26*(2), 489–511. https://doi.org/10.1037/0278-7393.26.2.489

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral Cortex*, *19*(12), 2767–2796. https://doi.org/10.1093/cercor/bhp055

Bozic, M., Marslen-Wilson, W. D., Stamatakis, E. A., Davis, M. H., & Tyler, L. K. (2007). Differentiating morphology, form, and meaning: Neural correlates of morphological complexity. *Journal of Cognitive Neuroscience*, *19*(9), 1464–1475. https://doi.org/10.1162/jocn.2007.19.9.1464

Bozic, M., Tyler, L. K., Li, S., Wingfield, C., & Marslen-Wilson, W. D. (2013). *Neurobiological Systems for Lexical Representation and Analysis in English*. https://doi.org/10.1162/jocn_a_00420

Brett, M., Anton, J.-L., Valabregue, R., Poline, J.-B., & others. (2002). Region of interest analysis using the MarsBar toolbox for SPM 99. *Neuroimage*, *16*(2), S497.

Bruffaerts, R., Dupont, P., Peeters, R., De Deyne, S., Storms, G., & Vandenberghe, R. (2013). Similarity of fMRI activity patterns in left perirhinal cortex reflects semantic similarity between words. *Journal of Neuroscience*, *33*(47), 18597–18607. https://doi.org/10.1523/JNEUROSCI.1548-13.2013

Brysbaert, M., Mandera, P., & Keuleers, E. (2018). The Word Frequency Effect in Word Processing: An Updated Review. *Current Directions in Psychological Science*, *27*(1), 45–50. https://doi.org/10.1177/0963721417727521

Bulut, T. (2022). Neural correlates of morphological processing: An activation likelihood estimation meta-analysis. *Cortex*, *151*, 49–69. https://doi.org/10.1016/j.cortex.2022.02.010

Bybee, J. (1995). Regular morphology and the lexicon. *Language and Cognitive Processes*, *10*(5), 425–455. https://doi.org/10.1080/01690969508407111

Bybee, J., & Hopper, P. (2001). Frequency and the Emergence of Linguistic Structure. In J. L. Bybee & P. J. Hopper (Eds.), *Typological studies in languages (TSL)* (Vol. 45, Issue August). John Benjamins Publishing Company. https://doi.org/10.1075/tsl.45

Caramazza, A., Laudanna, A., & Romani, C. (1988). Lexical access and inflectional morphology. *Cognition*, *28*(3), 297–332. https://doi.org/10.1016/0010-0277(88)90017-0

Carota, F., Bozic, M., & Marslen-Wilson, W. (2016). Decompositional Representation of Morphological Complexity: Multivariate fMRI Evidence from Italian. *Journal of Cognitive Neuroscience*, *28*(12), 1878–1896. https://doi.org/10.1162/jocn_a_01009

Carreiras, M., Riba, J., Vergara, M., Heldmann, M., & Münte, T. F. (2009). Syllable congruency and word frequency effects on brain activation. *Human Brain Mapping*, *30*(9), 3079–3088. https://doi.org/10.1002/hbm.20730

Davey, J., Thompson, H. E., Hallam, G., Karapanagiotidis, T., Murphy, C., De Caso, I., Krieger-Redwood, K., Bernhardt, B. C., Smallwood, J., & Jefferies, E. (2016). Exploring the role of the posterior middle temporal gyrus in semantic cognition: Integration of anterior temporal lobe with executive processes. *NeuroImage*, *137*, 165–177. https://doi.org/10.1016/J.NEUROIMAGE.2016.05.051

Desai, R., Conant, L. L., Waldron, E., & Binder, J. R. (2006). fMRI of Past Tense Processing: The Effects of Phonological Complexity and Task Difficulty. *Journal of Cognitive Neuroscience*, *18*(2), 278–297. https://doi.org/10.1162/089892906775783633

Devlin, J. T., Jamison, H. L., Matthews, P. M., & Gonnerman, L. M. (2004). Morphology and the internal structure of words. *Proceedings of the National Academy of Sciences*, *101*(41), 14984–14988. https://doi.org/10.1073/pnas.0403766101

Feldman, L. B., Milin, P., Cho, K. W., Moscoso del Prado Martín, F., & O'Connor, P. A. (2015). Must analysis of meaning follow analysis of form? A time course analysis. *Frontiers in Human Neuroscience*, *9*. https://doi.org/10.3389/fnhum.2015.00111

Fiebach, C. J., Friederici, A. D., Müller, K., & Cramon, D. Y. von. (2002). fMRI Evidence for Dual Routes to the Mental Lexicon in Visual Word Recognition. *Journal of Cognitive Neuroscience*, *14*(1), 11–23. https://doi.org/10.1162/089892902317205285

Fiez, J. A., Balota, D. A., Raichle, M. E., & Petersen, S. E. (1999). Effects of Lexicality, Frequency, and Spelling-to-Sound Consistency on the Functional Anatomy of Reading. *Neuron*, *24*(1), 205–218. https://doi.org/10.1016/S0896-6273(00)80833-8

Fischer-Baum, S., Bruggemann, D., Gallego, I. F., Li, D. S. P., & Tamez, E. R. (2017). Decoding levels of representation in reading: A representational similarity approach. *Cortex*, *90*, 88–102. https://doi.org/10.1016/j.cortex.2017.02.017

Giraudo, H., & Grainger, J. (2000). Effects of prime word frequency and cumulative root frequency in masked morphological priming. *Language and Cognitive Processes*, *15*(4–5), 421–444. https://doi.org/10.1080/01690960050119652

Giraudo, H., & Grainger, J. (2001). Priming complex words: Evidence for supralexical representation of morphology. *Psychonomic Bulletin & Review*, *8*(1), 127–131. https://doi.org/10.3758/BF03196148

Gold, B. T., & Rastle, K. (2007). Neural Correlates of Morphological Decomposition during Visual Word Recognition. *Journal of Cognitive Neuroscience*, *19*(12), 1983–1993. https://doi.org/10.1162/jocn.2007.19.12.1983

Graves, W. W., Binder, J. R., Desai, R. H., Conant, L. L., & Seidenberg, M. S. (2010). Neural correlates of implicit and explicit combinatorial semantic processing. *NeuroImage*, *53*(2), 638–646. https://doi.org/10.1016/J.NEUROIMAGE.2010.06.055

Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001). Distributed and Overlapping Representations of Faces and Objects in Ventral Temporal Cortex. *Science*, *293*(5539), 2425–2430. https://doi.org/10.1126/science.1063736

Jared, D., Jouravlev, O., & Joanisse, M. F. (2017). The effect of semantic transparency on the processing of morphologically derived words: Evidence from decision latencies and event-related potentials. *Journal of Experimental Psychology: Learning Memory and Cognition*, *43*(3), 422–450. https://doi.org/10.1037/xlm0000316

Joanisse, M. F., & Seidenberg, M. S. (2005). Imaging the past: Neural activation in frontal and temporal regions during regular and irregular past-tense processing. *Cognitive, Affective and Behavioral Neuroscience*, *5*(3), 282–296. https://doi.org/10.3758/CABN.5.3.282

Kang, B., & Kim, H. (2009). *Frequency of Korean language: based on 15 million size Eojeol Sejong syntactic and semantic corpus*. Handkook munhwasa.

Keuleers, E., Lacey, P., Rastle, K., Brysbaert, M., Keuleers, E., Brysbaert, M., Lacey, P., & Rastle, : K. (2011). The British Lexicon Project: Lexical decision data for 28,730 monosyllabic and disyllabic English words. *Behavior Research Methods 2011 44:1*, *44*(1), 287–304. https://doi.org/10.3758/S13428-011-0118-4

Kim, J., Jung, J., & Nam, K. (2021). Neural correlates of confusability in recognition of morphologically complex Korean words. *PLOS ONE*, *16*(4), e0249111. https://doi.org/10.1371/journal.pone.0249111

Kim, J., Kang, J., Kim, J., & Nam, K. (2022). Temporal dynamics of form and meaning in morphologically complex word processing: An ERP study on Korean inflected verbs. *Journal of Neurolinguistics*, *64*, 101098. https://doi.org/10.1016/j.jneuroling.2022.101098

Kim, J., Kim, S., & Nam, K. (2024). Neural Dynamics of Processing Inflectional Morphology: An fMRI Study on Korean Inflected Verbs. *Brain Sciences*, *14*(8), 752. https://doi.org/10.3390/brainsci14080752

Kim, J., Lee, W., Song, J., Kim, J., Nam, K., & Kim, S. (2025). Korean Eojeol Representation and Lexical Processing. *Handbook on the Korean Language and Literacy: Insights into Hangul and Text Processing*, 239–276. https://doi.org/10.1007/978-3-031-88087-2_12

Kim, J., & Nam, K. (2025). Sensitivity of the inferior frontal gyrus to morphological processing through the detection of category violation. *Scientific Reports 2025 15:1*, *15*(1), 27675-. https://doi.org/10.1038/s41598-025-09921-z

Kim, J., Pyun, S. B., & Nam, K. (2025). Brain activation in noun-based- and predicate-Eojeol processing during Korean visual word recognition. *Lingua*, *326*, 104018. https://doi.org/10.1016/J.LINGUA.2025.104018

Kriegeskorte, N. (2008). Representational similarity analysis – connecting the branches of systems neuroscience. *Frontiers in Systems Neuroscience*, *2*(NOV). https://doi.org/10.3389/neuro.06.004.2008

Kuperman, V., Schreuder, R., Bertram, R., & Baayen, R. H. (2009). Reading polymorphemic Dutch compounds: Toward a multiple route model of lexical processing. *Journal of Experimental Psychology: Human Perception and Performance*, *35*(3), 876–895. https://doi.org/10.1037/a0013484

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26. https://doi.org/10.18637/JSS.V082.I13

Lau, E. F., Phillips, C., & Poeppel, D. (2008). A cortical network for semantics: (De)constructing the N400. *Nature Reviews Neuroscience*, *9*(12), 920–933. https://doi.org/10.1038/NRN2532;KWRD

Lavric, A., Rastle, K., & Clapp, A. (2011). What do fully visible primes and brain potentials reveal about morphological decomposition? *Psychophysiology*, *48*(5), 676–686. https://doi.org/10.1111/j.1469-8986.2010.01125.x

Lehtonen, M., Cunillera, T., Rodríguez-Fornells, A., Hultén, A., Tuomainen, J., & Laine, M. (2007). Recognition of morphologically complex words in Finnish: Evidence from event-related potentials. *Brain Research*, *1148*(1), 123–137. https://doi.org/10.1016/j.brainres.2007.02.026

Lehtonen, M., Vorobyev, V. A., Hugdahl, K., Tuokkola, T., & Laine, M. (2006). Neural correlates of morphological decomposition in a morphologically rich language: An fMRI study. *Brain and Language*, *98*(2), 182–193. https://doi.org/10.1016/j.bandl.2006.04.011

Lewis, G., Solomyak, O., & Marantz, A. (2011). The neural basis of obligatory decomposition of suffixed words. *Brain and Language*, *118*(3), 118–127. https://doi.org/10.1016/j.bandl.2011.04.004

Longtin, C.-M., Segui, J., & Hallé, P. A. (2003). Morphological priming without morphological relationship. *Language and Cognitive Processes*, *18*(3), 313–334. https://doi.org/10.1080/01690960244000036

Milin, P., Feldman, L. B., Ramscar, M., Hendrix, P., & Baayen, R. H. (2017). Discrimination in lexical decision. *PLoS ONE*, *12*(2). https://doi.org/10.1371/journal.pone.0171935

Newman, S. D., Ikuta, T., & Burns, T. (2010). The effect of semantic relatedness on syntactic analysis: an fMRI study. *Brain and Language*, *113*(2), 51. https://doi.org/10.1016/J.BANDL.2010.02.001

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Plaut, D. C., & Gonnerman, L. M. (2000). Are non-semantic morphological effects icompatible with a distributed connectionist approach to lexical processing? *Language and Cognitive Processes*, *15*(4–5), 445–485. https://doi.org/10.1080/01690960050119661

Pliatsikas, C., Wheeldon, L., Lahiri, A., & Hansen, P. C. (2014). Processing of zero-derived words in English: An fMRI investigation. *Neuropsychologia*, *53*(1), 47–53. https://doi.org/10.1016/j.neuropsychologia.2013.11.003

Price, A. R., Bonner, M. F., Peelle, J. E., & Grossman, M. (2015). Converging evidence for the neuroanatomic basis of combinatorial semantics in the angular gyrus. *Journal of Neuroscience*, *35*(7), 3276–3284. https://doi.org/10.1523/JNEUROSCI.3446-14.2015

R Core Team. (2022). *R: A language and environment for statistical computing* (4.3.1). R Foundation for Statistical Computing. https://www.r-project.org/

Rastle, K., Davis, M. H., & New, B. (2004). The broth in my brother's brothel: Morpho-orthographic segmentation in visual word recognition. *Psychonomic Bulletin and Review*, *11*(6), 1090–1098. https://doi.org/10.3758/BF03196742

Solomyak, O., & Marantz, A. (2010). Evidence for Early Morphological Decomposition in Visual Word Recognition. *Journal of Cognitive Neuroscience*, *22*(9), 2042–2057. https://doi.org/10.1162/jocn.2009.21296

Taft, M., & Forster, K. I. (1975). Lexical storage and retrieval of prefixed words. *Journal of Verbal Learning and Verbal Behavior*, *14*(6), 638–647. https://doi.org/10.1016/S0022-5371(75)80051-X

Thompson-Schill, S. L., D'Esposito, M., Aguirre, G. K., & Farah, M. J. (1997). Role of left inferior prefrontal cortex in retrieval of semantic knowledge: A□reevaluation. *Proceedings of the National Academy of Sciences*, *94*(26), 14792–14797. https://doi.org/10.1073/PNAS.94.26.14792

Tyler, L. K., Bright, P., Fletcher, P., & Stamatakis, E. A. (2004). Neural processing of nouns and verbs: The role of inflectional morphology. *Neuropsychologia*, *42*(4), 512–523. https://doi.org/10.1016/j.neuropsychologia.2003.10.001

van Heuven, W. J. B., Mandera, P., Keuleers, E., & Brysbaert, M. (2014). Subtlex-UK: A New and Improved Word Frequency Database for British English. *Quarterly Journal of Experimental Psychology*, *67*(6), 1176–1190. https://doi.org/10.1080/17470218.2013.850521

Vigneau, M., Beaucousin, V., Hervé, P. Y., Duffau, H., Crivello, F., Houdé, O., Mazoyer, B., & Tzourio-Mazoyer, N. (2006). Meta-analyzing left hemisphere language areas: Phonology, semantics, and sentence processing. In *NeuroImage* (Vol. 30, Issue 4, pp. 1414–1432). https://doi.org/10.1016/j.neuroimage.2005.11.002

Weissbart, H., & Martin, A. E. (2024). The structure and statistics of language jointly shape cross-frequency neural dynamics during spoken language comprehension. *Nature Communications*, *15*(1). https://doi.org/10.1038/s41467-024-53128-1

Whitney, C., Jefferies, E., & Kircher, T. (2011). Heterogeneity of the left temporal lobe in semantic representation and control: Priming multiple versus single meanings of ambiguous words. *Cerebral Cortex*, *21*(4), 831–844. https://doi.org/10.1093/cercor/bhq148

Whitney, C., Kirk, M., O'Sullivan, J., Lambon Ralph, M. A., & Jefferies, E. (2011). The Neural Organization of Semantic Control: TMS Evidence for a Distributed Network in Left Inferior

Frontal and Posterior Middle Temporal Gyrus. *Cerebral Cortex*, *21*(5), 1066–1075. https://doi.org/10.1093/CERCOR/BHQ180

Yap, M. J., & Balota, D. A. (2009). Visual word recognition of multisyllabic words. *Journal of Memory and Language*, *60*(4), 502–529. https://doi.org/10.1016/j.jml.2009.02.001

Yi, K., Koo, M.-M., Nam, K., Park, K.-N., Park, T., Bae, S., Lee, C., Lee, H., & Cho, J.-R. (2017). The Korean Lexicon Project: A Lexical Decision Study on 30,930 Korean Words and Nonwords. *Korean Journal of Cognitive and Biological Psychology*, *29*(4), 395–410. https://doi.org/10.22172/cogbio.2017.29.4.004

Yim, H., Park, C., Lim, H., & Nam, K. (2006). Mental Representation and Processing Involved in Comprehending Korean Regular and Irregular Verb Eojeols: An fMRI and Reaction Time Study. *Neural Information Processing: 13th International Conference, ICONIP 2006, Hong Kong, China, October 3-6, 2006. Proceedings, Part I 13*, *4232*, 247–254.

Yue, Q., & Martin, R. C. (2021). Maintaining verbal short-term memory representations in non-perceptual parietal regions. *Cortex*, *138*, 72–89. https://doi.org/10.1016/j.cortex.2021.01.020