Tracking morphological development in stony corals

Garrett J. Fundakowski*[1], Viviana Brambilla[2], Kyle J. A. Zawada[3], Cher F. Y. Chow[1], Emily Croasdale[2], Amelia Errington[4], Luisa Fontoura[3], Wilhelm J. Marais[5], Rachael M. Woods[6], Pim Edelaar[7], Kevin Lala[1], Joshua S. Madin[8], Maria Dornelas[1,2]

*Corresponding author: garrett.fundakowski@gmail.com

[1] Centre for Biological Diversity, School of Biology, University of St Andrews, UK

[2] MARE – Centro de Ciências do Mar e do Ambiente, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

[3] School of Natural Sciences, Macquarie University, Sydney, NSW, Australia

[4] Reefscapers, Malé, Maldives

[5] Australian Centre For Robotics (ACFR), University of Sydney, Sydney, NSW, Australia

[6] Department of Climate Change, Energy, the Environment and Water, NSW, Australia

[7] Department of Molecular Biology and Biochemical Engineering, University Pablo de Olavide, Spain

[8] Department of Biology, Lund University, Lund, Sweden

[9] Hawai'i Institute of Marine Biology, University of Hawai'i at Mānoa, O'ahu, USA

Abstract

The shape of reef-building corals largely determines how they interact with their environment and the ecosystem services they provide. However, morphology is not fixed. As corals grow and develop from singular polyps to mature adult colonies, they experience pronounced changes in their morphology. This developmental trait variation remains poorly quantified in corals, despite their importance as ecosystem engineers. To address this gap, we used photogrammetry to track the morphological development of wild coral colonies across six size-independent metrics designed to capture variation in volume compactness, surface complexity, and top-heaviness. Size-trait models and analyses of colony movement through morphological trait space revealed that small colonies shared compact morphologies and only later diverged along growth-form-specific developmental trajectories. Massive corals largely retained stable shapes as they grew, whereas non-massive colonies became less compact, more top-heavy, and in most cases, more complex. Despite these differences, there was still overlap between growth forms in their developmental trajectories through morphospace. These findings highlight that traditional categorical growth forms mask important developmental variation, particularly early in coral ontogeny. By applying a quantitative trait-based framework, this study improves understanding of coral morphological development and provides a foundation for linking individual growth trajectories to broader reef ecosystem dynamics.

Keywords

Morphological analysis, Scleractinia, 3D Photogrammetry, Traits, Growth form

Data availability

Python code to generate the 3D meshes as well as the final 3D mesh dataset are available at: https://doi.org/10.17630/b0086b3f-fec9-4765-9a0f-fe9021aa79e3. Data and code for data extraction and analysis are available at: [DOI will be generated once accepted].

Statements and Declarations

The authors of this paper declare that there is no conflict of interest.

Author contributions

Conceptualization: GJF, VB, KJAZ, JSM, MD. Funding Acquisition: VB, KL, JSM, MD. Resources: JSM, MD. Supervision: PE, KL, JSM, MD. Investigation: GJF, VB, KJAZ, CFYC, LF, WJM, RMW, MD. Methodology: GJF. Data Curation: GJF, EC, AE. Formal analysis: GJF. Writing – original draft: GJF. Writing – review and editing: all authors.

Acknowledgements

We thank the Australian Museum's Lizard Island Research Station staff for their support in facilitating this research. We would also like to thank Mariana Álvarez-Noriega, who assisted with data collection in March 2021, as well as Devynn Wulstein, Nina Schiettekatte, and Mollie Asbury, who were members of the April and November 2022 field teams.

This study was conducted under GBRMPA research permits G15/38127.1 and G22/46527.1. Funding was provided by the John Templeton Foundation (grant #60501 'Putting the Extended Evolutionary Synthesis to the Test') (KL, MD, JSM), an Ian Potter Doctoral Fellowship at Lizard Island Research Station (VB), a MASTS small grant (VB), a Leverhulme fellowship (MD), University of Hawai'i start-up funding (JSM), the Warman Foundation (MD, JSM), a National Science Foundation–Natural Environment Research Council Biological Oceanography Grant (1948946) (JSM, MD), and the European Union (CoralINT, GA 101044975) (MD, VB). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

Introduction

Morphology plays a central role in mediating how organisms interact with their environment and contribute to ecosystem processes. However, morphology is dynamic – as individuals grow and develop, ontogenetic shifts in morphological traits can alter their performance and ecological function. Trait-based ecology has emerged as a powerful framework to study these links between form and function, but it has largely focused on interspecific differences, under the assumption that traits vary more between species than within them (McGill et al. 2006). Increasingly, studies have demonstrated that within-group trait variation can be substantial (Messier et al. 2010; Siefert et al. 2015) and that traditionally neglected drivers, such as temporal and developmental change, have important consequences for broader ecosystem dynamics (Bolnick et al. 2011; Westerband et al. 2021; Cope et al. 2022). Yet the developmental component of trait variation remains understudied, particularly in marine taxa where trait-based research lags behind that in terrestrial systems (Green et al. 2022). Recently, there has been a concerted effort advocating for the use of quantitative trait-based approaches to advance coral reef ecology (Madin et al. 2016a; Madin et al. 2016b). Here, we build on that effort by using quantitative morphological traits (Zawada et al. 2019a) to track how coral morphology changes through development and capture biologically relevant variation beyond the limits of traditional growth form classifications.

Corals undergo considerable morphological changes during their lifespan and are a particularly important system for studying developmental trait variation because their morphology plays a critical role in reef functioning. Coral colonies begin their benthic life as a singular polyp, initially resembling a small hemisphere. As they replicate and the colony gets larger, their overall colony morphology differentiates and diverges across species. Some species develop into intricate, branching forms, while others form more dense, massive colonies. These ontogenetic changes in morphology influence both colony-level performance, such as vulnerability to hydrodynamic forces (Madin et al. 2014), and reef-level functioning, such as habitat provision for other organisms (Urbina-Barreto et al. 2021). Establishing links between colony size and shape is therefore important for improving our ability to understand and predict not only how individual colonies perform, but also how they collectively shape ecosystem-level processes on coral reefs.

Traditionally, Scleractinian corals are categorized into growth forms based on broad qualitative differences in adult colony morphology (Veron 2000). Growth form has been shown to be an important trait for delineating life-history strategies in corals (Darling et al. 2012), which predict their response to bleaching and fishing stressors (Darling et al. 2013). It has also been linked to differences in key demographic rates (mortality, Madin et al. 2014; fecundity, Álvarez-Noriega et

al. 2016; growth, Dornelas et al. 2017). However, such a discrete categorization does not provide

a complete description of a coral's structure (Veron 2000) nor adequately captures the variation

in colony shape both within and between species (Zawada et al. 2019a). In response to this

shortcoming, and armed with recent advances in technology, many studies have begun

quantifying the three-dimensional size and shape of coral colonies using photogrammetry, CT

scanning, and laser scanning techniques (e.g., Bythell et al. 2001; Figueira et al. 2015; Gutiérrez-

Heredia et al. 2015; Lavy et al. 2015; Reichert et al. 2016; House et al. 2018; Zawada et al. 2019a;

Lange et al. 2020; Million et al. 2021; Urbina-Barreto et al. 2021; Guendulain-García et al. 2024).

Building on these advances, Zawada et al. (2019a) proposed a suite of six size-independent,

morphological metrics that describe continuous three-dimensional shape space and position

coral morphology along three functionally relevant axes of variation – volume compactness,

surface complexity, and top-heaviness. Volume compactness, defined by sphericity and

convexity, establishes a gradient from dense, boulder-like volumes to more intricate arborescent

branching morphologies. Next, packing and fractal dimension describe surface complexity,

spanning smooth to highly convoluted surfaces. Lastly, top-heaviness, characterized by the

second moments of area and volume, describes the vertical distribution of colony surface area

and volume relative to the substrate. These quantitative morphological traits have been linked to

variation in organism performance, ecosystem function, and response to disturbance (Zawada

et al. 2019b). While Zawada et al. (2019a) also established generic size relationships using these

metrics, the corals they phenotyped were skeletons from museum collections and thus could

not capture how individual colonies change through time. Thus, there remains a gap in

understanding how these morphological traits develop through coral ontogeny, particularly in

wild populations.

In this study, we address this gap by tracking morphological development in 133 wild coral

colonies using the six quantitative metrics proposed by Zawada et al. (2019a). For each metric,

we modelled size-trait relationships across all colonies and used these to examine how colonies

move through multidimensional morphological trait space as they grow. We hypothesized that

small corals from all growth forms would initially exhibit similar, compact morphologies before

diverging along growth-form-specific developmental trajectories as they grow. Specifically, we

expected volume compactness to be high in small colonies, with non-massive growth forms

becoming less compact as they grow. We also anticipated surface complexity to be low in small

colonies, with non-massive growth forms increasing in complexity (as these growth forms

prioritize adding surface area over volume) and massive colonies maintaining their low values.

Lastly, we predicted top-heaviness to be low in small colonies, increasing across non-massive

growth forms as colonies grow, but most rapidly in tabular colonies. Overall, this study

demonstrates the utility of these quantitative metrics for tracking coral morphology and improves

our understanding of developmental trait variation in corals.

Methods

Study Site and Design

We conducted this observational study on ten reef sites around Jiigurru (also known as Lizard Island) at depths of 1-4 m in the northern Great Barrier Reef in Queensland, Australia (Fig. 1). To track morphological development in coral colonies *in situ*, we captured and quantified colony morphology of 133 coral colonies using 3D photogrammetry over the course of four years and five fieldtrips (November 2018, November 2019, March 2021, April 2022, and November 2022). We selected colonies to capture a diverse array of species and morphological growth forms, ranging in size from small, young recruits to large, adult colonies. Exact locations of colonies were noted on printed orthomosaic maps to aid in colony relocation in subsequent years, though this was not always possible.

Fig. 1 Map of study sites. The map shows the location of the 10 shallow water reef sites around Jiigurru, Australia. Moving counterclockwise from the north they are North Reef 3 (N3), Turtle Beach (TB), Resort (RS), Corner Beach (CB), Horseshoe (HS), Trimodal (TM), Lagoon 1 (L1), South Island (SI), Southeast (SE), and Gnarly Tree (GT). Spatial data for the Great Barrier Reef area and Jiigurru coastline are from Great Barrier Reef Marine Park Authority (GBRMPA) (2022)

Image capture

We took photographs of selected colonies using handheld underwater cameras in continuous shooting/interval mode with autofocus enabled. We placed three gridded scale bars (2 cm x 2 cm or 10 cm x 10 cm) on the substrate surrounding the focal colonies to provide reference points for model scaling. Snorkelers photographed each reef scene (the area immediately surrounding the selected corals) while slowly swimming in a multi-tiered circular pattern about 0.5 m away from the focal colony. Photos were taken in an upward spiral pattern with views of the focal colony at approximately 0, 30, 60, and 80 degrees, maintaining roughly 60-80% overlap between images to ensure that the complex structure of the colonies was captured as completely as possible. Additional photos were taken of any complex sections that might require additional resolution to accurately build (e.g., dark undersides of tabular colonies or dense sections of branches in arborescent colonies). The number of images taken for each reef scene ranged from 73 to 1177 (average 298), with the number required for a good quality reconstruction increasing with size, complexity, and number of colonies in the reef scene being photographed.

Model construction

We constructed three-dimensional (3D) models from the collected photosets using the proprietary photogrammetric software Agisoft Metashape Pro version 1.7.3 (Agisoft LLC 2021). The workflow used to generate the reef scene models for this study broadly follows the pipelines of similar studies that have used Metashape to construct 3D models from 2D images (e.g. - Lange et al. 2020; Million et al. 2021). However, meshes were built directly from depth maps following recommendations from the Agisoft Metashape User Manual version 1.7 (Agisoft LLC 2021). We implemented this workflow with specified parameter settings (SI Table 1) using an automated Python script adapted from Million et al. (2021).

Following automated construction, we opened the reef scene models in Metashape for visual inspection and scaling. Any models with model construction issues (e.g. major deviations in shape, holes in the focal colony, or missing scale bars) were removed from the dataset. We then scaled the remaining models using four known lengths from two gridded scale bars, with the two lengths on the same scale bar being perpendicular to each other. We checked that scale error values were less than 1 mm and manually verified scaling accuracy by measuring the unused gridded scale bar. Finally, we exported the scaled reef scene models as a Wavefront (.obj) file with

189 texture included, to allow for more precise cropping and isolation of the coral colony from the
190 surrounding reef scene during post-processing.

Mesh post-processing

We then imported each scaled reef scene model into the free mesh-editing software Autodesk
MeshMixer version 3.5 (Autodesk Inc. 2017) for post-processing to orient the reef scene, isolate
the coral colonies, create a watertight mesh, and standardize mesh resolution.

First, we manually rotated the models to align the *z*-axis with the upwards facing direction of the
reef scene *in situ* as some morphological metrics are orientation dependent. Models of the same
reef scene from different years were oriented simultaneously to ensure consistency between
years. Then we removed non-coral substrate from each reef scene model to isolate the focal
individual(s). For scenes containing multiple focal colonies, we separated each colony into its
own mesh file to allow for independent analysis.

At this point, the isolated colony meshes consisted of a continuous triangulated mesh with a hole
where the colony was attached to the reef. We smoothed the boundary of this hole, evening out
the jagged edges that were artifacts from the colony isolation process, to ensure all attachment
boundaries were of similar smoothness regardless of mesh density. From here, we generated a
new flattened attachment surface to create a watertight mesh, as this was necessary for
calculating morphological metrics related to volume. To do this, we extended the mesh from the
smoothed boundary in the direction of attachment to the reef (not always vertical). The extended
mesh was then sliced using a plane, normal to the direction of extension, at the point where the
plane intersected with the furthest extent of the smoothed attachment boundary. We filled the
resulting cut with a flat plane to make the mesh watertight.

Finally, to ensure colony meshes were comparable, we standardized mesh resolution via a
remeshing algorithm with a cell size of 2.4 mm, as this was the largest mean edge length across
all colony meshes. The remeshing algorithm first creates an approximation of the input mesh
using small cubes, or voxels, of the given cell size. The algorithm then reprojects the vertices from
the voxel approximation back to the surface of the input mesh. In doing this, meshes retained
their general shape with details smaller than the cell size being smoothed over, and resolution
was standardized across meshes. Then we ran a final inspection of the mesh to fix non-manifold
regions and fill small holes that may have been generated by the remeshing algorithm, and
exported the resulting watertight mesh in ASCII Polygon (.ply) file format. We automated the
remeshing process and export of the final mesh with a custom Python script.

Morphological metrics

To capture coral shape, we used the six morphological metrics described by Zawada et al. (2019a) – convexity, sphericity, packing, fractal dimension, and second moments of both area and volume – as they represent a quantitative, trait-based approach to describing coral morphology and have been shown to effectively partition quantitative morphological variation between qualitative growth forms. Convexity and sphericity define the morphological axis of volume compactness; packing and fractal dimension describe surface complexity; and second moments of both area and volume are top-heaviness metrics (Zawada et al. 2019a). Convexity, sphericity, packing, and fractal dimension are inherently size-independent metrics, while the second moments are not. Thus, to calculate the second moments of area and volume, meshes were first standardized by volume to remove the effect of colony size (Zawada et al. 2019a). To describe coral size for establishing size-trait relationships, we use planar area as it scales with 3D size traits (House et al. 2018) and is less biased than surface area or volume estimates to our method of generating a watertight mesh. We extracted planar area and all morphological metrics for each colony mesh using the 'habtools' package (Schiettekatte et al. 2025) in R (R Core Team 2021). Specifically, to calculate fractal dimension we used the cube-counting method, as it is the recommended method for watertight 3D meshes (Schiettekatte et al. 2025).

Colony Metadata

We identified each coral colony to species level sensu Veron (2000) and T. C. L. Bridge (personal communication). Then, we assigned growth forms for each species using the Coral Trait Database (CTD) (Madin et al. 2016b), which gives the 'typical' growth form of a species based on the morphological descriptions used in Veron (2000). Growth forms included were arborescent (referred to as open branching in the CTD), closed branching, corymbose, digitate, massive, submassive, and tabular (SI Table 2). For this study, we grouped the three colonies classified as submassives in with massives due to their similar morphologies.

The final dataset consisted of 340 watertight meshes from 133 unique colonies, with 104 colonies tracked across two or more years. The colonies spanned six growth forms, 36 species, and ranged in size from 1.03 cm^2 to 2565.28 cm^2 (see SI Table 3 for more details).

Morphological development models

To investigate how morphological metrics change as corals grow, we fit six separate Bayesian hierarchical linear models with each metric as a response variable. To explore potential differences in the size-metric relationships across growth forms, we modelled colony shape as a function of the interaction between size and growth form for each morphological metric.

Prior to model fitting, planar area was $\log_{10}$-transformed as the data spanned two orders of magnitude. Similarly, packing and both second moment metrics were also $\log_{10}$-transformed to account for the right-skewness in the data. Additionally, as sphericity and convexity are ratios bounded by zero and one, they were logit-transformed for analyses. With these transformations, response variables were approximately normal, and we modelled them with Gaussian distributions.

For each of the six models, we specified fixed effects to estimate growth-form-specific intercept ($\alpha_{m[i]}^{growth.form}$) and slope ($\beta_{m[i]}^{growth.form}$) parameters with $\log_{10}$-transformed planar area ($\log_{10}(PA_i)$) as a continuous explanatory variable. We included Colony ID as a random intercept offset ($\alpha_{c[i]}^{offset.ColonyID}$) to account for any variation between colonies within growth form. We selected weakly regularizing priors for each parameter, broadly following McElreath (2020), and visually inspected them to ensure they did not constrain the posterior distributions (SI Fig. 1). The overall model parameterization and prior structure (Table 1) we implemented is as follows:

$$morphological\ metric_i \sim Normal(\mu_i, \sigma)$$

$$\mu_i = \alpha_{m[i]}^{growth.form} + \beta_{m[i]}^{growth.form} \times \log_{10}(PA_i) + \alpha_{c[i]}^{offset.ColonyID}$$

$$\alpha_m^{growth.form} \sim Normal(\mu^x, \sigma^x)\text{ , for } m = 1, \dots, 6 \text{ growth forms}$$

$$\beta_m^{growth.form} \sim Normal(\mu^y, \sigma^y)\text{ , for } m = 1, \dots, 6 \text{ growth forms}$$

$$\alpha_c^{offset.ColonyID} \sim Normal\left(0, \sigma^{\alpha^{offset.ColonyID}}\right)\text{ , for } c = 1, \dots, 133 \text{ colony IDs}$$

$$\sigma, \sigma^{\alpha^{offset.ColonyID}} \sim Exp(\lambda^z)$$

 Table 1 Prior values used for each morphological development model

	$\alpha_m^{growth.form}$		$\beta_m^{growth.form}$		$\sigma, \sigma^{\alpha^{offset.ColonyID}}$
$morphological\ metric_i$	μ^x	σ^x	μ^y	σ^y	λ^z
$\text{logit}(convexity_i)$	2	1	0	0.6	1
$\text{logit}(sphericity_i)$	2	1	0	0.6	1
$\log_{10}(packing_i)$	−0.2	0.4	0	0.2	3
$fractal\ dimension_i$	2.2	0.2	0	0.2	3
$\log_{10}(second\ moment\ of\ area_i)$	−0.5	0.5	0	0.2	3
$\log_{10}(second\ moment\ of\ volume_i)$	0	0.5	0	0.2	3

Model fit and validation

We fit all models in R (R Core Team 2021) using stan statistical programming language (Stan Development Team 2024) via the 'brms' package (Bürkner 2017). We ran four MCMC chains of 20000 iterations per chain, with a warm-up of 5000 iterations and a thinning rate of 10 iterations, for each model. This resulted in 6000 draws in the posterior distributions. To assess model convergence, we confirmed $\hat{R}$ values were sufficiently close to 1.

To check model accuracy in reproducing the empirical data, we performed a set of posterior predictive checks for each model (Gelman et al. 1996; Gabry et al. 2019; Gelman et al. 2020). First, we produced a set of 1000 simulated datasets using the 'posterior_predict' function from the 'brms' package (Bürkner 2017). Then we calculated the mean and standard deviation of the response variable from these simulated datasets and compared them to the summary statistics from the raw data across varying grouping levels to assess model fit. We did these checks at both the growth form and site level for all size-metric models (SI Fig. 2 - SI Fig. 7).

When reporting model results, we borrow the language of evidence as proposed by Muff et al. (2022).

Developmental trajectories in morphological trait space

We used Principal Component Analysis (PCA) to construct a morphological trait space (morphospace) from the six morphological metrics derived from colony meshes. Prior to ordination, each metric was standardized to have a mean of zero and unit variance to remove the bias from varying metric scales. We conducted PCA significance tests using the 'PCAtest'

package (Camargo 2022) to determine the significant principal components (PCs) ($p < 0.05$) and
identify metrics with significant loadings on each PC.

To examine how colonies developed within this morphospace, we extracted fitted values from
each of the six morphological development models and projected them onto the PCA space. To
get colony-level trajectories, we drew 6000 samples from the expected values of the posterior
predictive distribution across the range of observed planar areas, including ColonyID as a random
effect for all 104 colonies tracked for at least two years. From these draws, we calculated colony-
level displacement vectors as the difference between the predicted positions at the minimum
and maximum sizes (standardized across growth forms), and we summarized the distribution of
vector magnitudes and directions by growth form. Finally, to compare how morphologies
occupied morphospace at different size classes, we randomly sampled 100 posterior predictive
draws per colony at four standardized planar areas (5, 50, 500, and 2500 cm2) and plotted the
density of predicted positions by growth form.

Results

Morphological development models

All growth forms except for massives exhibited strong evidence (99% credible interval below zero;
Muff et al. 2022) for negative size-compactness relationships across both convexity and
sphericity models (Fig. 2, SI Table 4). Massives had a slope estimate distribution that overlapped
with zero suggesting they maintained relatively constant volume compactness across the
observed size range. In the convexity model, tabular, digitate, and arborescent colonies had
steeper slope estimates than the closed branching and corymbose colonies. For the sphericity
model, the slope estimates for these non-massive growth forms were broadly similar, with
digitate colonies having a slightly more negative slope estimate.

Size-complexity relationships displayed distinct patterns across growth forms depending on the
surface complexity metric considered. In the packing model, arborescent, closed branching,
corymbose, and digitate colonies showed strong evidence for positive size-packing relationships
while table corals demonstrated strong evidence for a slight negative relationship across the
observed size range. Massives showed weak evidence (90% credible interval above zero; Muff et
al. 2022) for a positive size-packing relationship. On the other hand, for the fractal dimension
model, table corals were the only growth form to show strong evidence for a size-fractal

dimension relationship, which was negative. All other growth forms did not experience a change

in their fractal dimension with increasing size, as their credible intervals all overlapped zero.

Fig. 2 Scatterplots for all morphological development models. Each row corresponds to a different size-metric model (S = Sphericity, C = Convexity, P = Packing, FD = Fractal dimension, SMA = Second moment of area, SMV = Second moment of volume) faceted by growth form. Here, points represent individual meshes (n = 340) colored by growth form, with grey lines connecting the meshes from the same Colony ID through time. Bold colored lines represent linear regression lines (with 50%, 80%, and 95% credible intervals) overlaid for each growth form. Black horizontal dashed lines represent corresponding metric values for a hemisphere. (Note: a perfect hemisphere has a convexity value of 1, which on the logit scale would be positive infinity and therefore is not displayed.) Individual plot borders are colored based on whether the 95% credible interval for model slope estimates overlapped with zero for each growth form (red = negative slopes, grey = slopes overlapping zero, blue = positive slopes)

With regard to the top-heaviness models, both second moment of area (SMA) and second
moment of volume (SMV) exhibited non-zero relationships for some growth forms. In the SMA
model, arborescent, closed branching, digitate, and tabular colonies all showed strong evidence
for positive slopes, while corymbose colonies exhibited moderate evidence (95% credible
interval above zero; Muff et al. 2022). Among these, digitate and tabular colonies had the steepest
positive slopes. Massives, however, maintained a constant SMA as size increased, with the
credible interval of the slope estimate overlapping zero. The relationships were quite different for
the SMV model. Here, table corals exhibited strong evidence and digitate colonies exhibited weak
evidence for increasing SMV with increasing size. On the other hand, corymbose colonies
displayed strong evidence and massives displayed moderate evidence for a negative relationship
between size and second moment of volume. Meanwhile, arborescent and closed branching
corals had slope estimates overlapping zero, indicating no detectable change in SMV with size.

Developmental trajectories in morphological trait space

When defining the morphological trait space, the first two principal component (PC) axes were
significant ($p < 0.001$) and together captured 78.3% of the observed variation in shape (Fig. 3a).
The first PC explained 52.1% of the variation across the morphospace and was comprised of the
four volume compactness and top-heaviness metrics. The second PC captured 26.2% of the
variation and was defined by the two surface complexity metrics.

Model predictions showed that small colonies across all growth forms were concentrated in the
region of the morphospace associated with higher volume compactness (Fig. 3b). With
increasing planar area, the areas occupied by each growth form progressively diverged. At first, a
division appeared between massive and non-massive growth forms, with the latter shifting
positively along PC1, in the direction of increasing top-heaviness. Then, at larger sizes, non-
massives further differentiated along PC2, reflecting increasing variation in surface complexity.
By the largest size class, growth forms largely occupied distinct regions of the morphospace,
although a high degree of overlap remained between corymbose and closed branching colonies.

Looking at how coral colonies moved through the morphological trait space, there was a strong
pattern of growth form differences in their developmental trajectories (Fig. 3c, SI Table 6). Here,
massive colonies clearly stood out from the other growth forms with the smallest magnitude
mean trajectory vector, suggesting relatively consistent morphology with increasing planar area.
Massives were also the only group to shift on average toward increased volume compactness and

decreased top-heaviness (moving negatively along PC1). However, it is also worth noting they

exhibited the greatest colony-level variation in trajectory direction compared to other growth

forms. Table corals also had a distinctive trajectory, as the only growth form moving toward

reduced surface complexity (moving positively along PC2). The rest of the growth forms, on the

other hand, generally followed trajectories of decreasing compactness (moving positively along

PC1) and increasing surface complexity (moving negatively along PC2), though subtle differences

in magnitude and direction were evident. For instance, corymbose, closed branching, and

arborescents exhibited similar magnitudes of change, but differed slightly in direction, with

corymbose colonies shifting more strongly toward greater surface complexity (negatively along

PC2). Meanwhile, digitate colonies paralleled arborescents in trajectory direction but had a much

greater trajectory magnitude across the standardized range of planar area values.

Fig. 3 Morphological development in continuous trait space. The multidimensional space was built using the six morphological metrics (C = convexity, S = sphericity, P = packing, FD = fractal dimension, SMA = second moment of area, SMV = second moment of volume) extracted from the colony mesh models. The space is projected in two dimensions using the first two principal components (PC), which account for 78.3% of the total variance. (a) Points depict individual meshes (n = 340) colored by growth form, with grey lines connecting the meshes from the same colony through time. Black arrows represent the loading and direction of each metric with arrow lengths showing how much each trait contributes to the PCs (scaled by a factor of 4 for visualization). (b) Density contour plots representing the portion of the total morphospace occupied by individuals of each growth form at various planar area sizes. (c) Circular plot depicting the direction and magnitude of the mean trajectory (solid black line) for each growth form. Colored bars represent each growth form with their heights showing the relative magnitude of the mean trajectory and their widths depicting the 95% quantile range for the direction of individual colony trajectories. Black arrows represent each metric as it loads into the PCA

Discussion

Using a quantitative framework to describe complex three-dimensional (3D) structure, this study tracked the morphological development of coral colony growth *in situ*. We used six metrics that describe colony geometry and morphology and explicitly showed the divergence and similarities between nominal growth forms throughout development. Corals of all growth forms generally started as small semi-hemispherical colonies characterized by high volume compactness, low surface complexity, and low top-heaviness. As they grew, corals belonging to different growth forms took different developmental trajectories through morphological trait space, though there was considerable overlap between some growth forms. These findings suggest that traditional qualitative growth forms for corals may only be useful categorizations for large mature colonies, as at smaller sizes there is too much overlap in morphological space to reliably distinguish among growth forms. Furthermore, this highlights the importance of quantitative approaches for capturing the full range of morphological variation.

Overall, the morphological development observed was relatively consistent with expectations. As predicted, small corals exhibited high volume compactness, and non-massive growth forms experienced decreases in sphericity and convexity as they grew. Likewise, second moment of area increased with colony size across all non-massive growth forms. Other size-metric relationships, however, deviated from expectations and warrant closer examination.

Table corals, in particular, differed from hypothesized size-complexity relationships. While most non-massive growth forms followed the predicted positive size-packing patterns, tabular colonies exhibited a slight negative size-packing relationship. Despite this, their packing values remained close to 1 (log-packing values centered around 0) across the observed range of sizes. A packing value of 1 indicates that the surface area of the object is equal that of its convex hull, a feature obviously achieved by an entirely convex object. Although table corals are not fully convex, such packing values can also result when an object has a balanced amount of inverted and everted surfaces (Zawada et al. 2019a), and the model results suggest that table corals experience a subtle reduction in this ratio as they grow. This observed negative size-packing relationship in table corals was reinforced by an also unexpected negative relationship between size and fractal dimension. However, previous research has revealed that as table corals get larger, the shelter volume (e.g. unoccupied volume under the table coral) they provide increases exponentially (Urbina-Barreto et al. 2021), suggesting that the corals themselves occupy a progressively smaller fraction of the bounding box. It would make sense then, that for the cube-counting algorithm employed to estimate fractal dimension, that with increasing colony size,

increasingly fewer cubes will intersect with the coral surface, leading to the observed negative
relationship. Thus, although initially unexpected, both packing and fractal dimension results
converge on a consistent result – that table corals lose surface complexity as they grow.

By contrast, non-tabular morphologies maintained relatively constant fractal dimension with
size, rather than the predicted positive relationship. Notably, across all size-metric models,
fractal dimension produced some of the highest residuals suggesting that some variation was left
unexplained. This may be partly attributable to methodological constraints. To account for
different mesh resolutions resulting from variable imaging conditions, all colonies were
remeshed to a standardized resolution prior to metric extraction. While necessary to ensure
comparability between meshes, this step unavoidably introduced bias by smoothing out fine-
scale surface texture, likely reducing fractal dimension estimates and potentially obscuring
biological patterns. Though other 3D modelling technologies (i.e. laser-scanning or CT scanning)
have been shown to have higher accuracy and precision than 3D photogrammetry (Figueira et al.
2015; Guendulain-García et al. 2023) due to their higher resolution capabilities, these systems
are prohibitively expensive. At the moment, underwater photogrammetry is the most accurate
non-invasive method available for obtaining *in situ* morphological trait measurements (Figueira
et al. 2015; Gutiérrez-Heredia et al. 2015).

Unexpected results were also observed in the relationship between size and second moment of
volume (SMV). Instead, we observed that some morphologies showed no change in SMV with
increasing size, while others experienced either an increase or a decrease. One possible
explanation lies in the manual isolation step required to extract colonies from the surrounding
reef scene. By trimming the colony at the living tissue edge and extending boundaries toward the
attachment point, portions of the underlying substrate that the colony grew over (rather than
created itself) may have been inadvertently included. This effect likely varies by morphology, with
larger, broad-based colonies gaining disproportionately more volume during processing than
smaller colonies with narrow attachment points. Such methodological bias could obscure true
size-SMV relationships and likely contributed to the high residual variation observed across
morphologies as well. Future improvements, such as aligning dense point clouds between years
before colony isolation could help standardize trimming (as in Reichert et al. 2016), though at the
cost of added processing time and software requirements.

The trajectories of morphological development observed in this study have important ecological
and evolutionary implications. As coral colonies grow and change shape, their impact on the
environment and local reef community will change too. For instance, bigger corals will absorb

471 more light and cast larger shadows (Stambler and Dubinsky 2005), thus altering the light
472 availability for sessile benthic organisms in their vicinity. Additionally, as table corals grow and
473 they become larger and more top-heavy, they can offer greater shelter volume (Urbina-Barreto et
474 al. 2021), which is critical for large-bodied reef fish (Kerry and Bellwood 2011,2014). On the other
475 hand, as corals grow, their changing shape also alters how they experience their environment,
476 and in turn the selection they are exposed to. For example, larger table corals are more top-heavy
477 than smaller ones, making them more prone to dislodgement and damage from wave energy
478 (Madin and Connolly 2006; Zawada et al. 2019b). This selective impact of hydrodynamic
479 disturbances on the size structure of tabular species has potential downstream implications for
480 population recovery, as the most mechanically vulnerable tabular colonies are the most fecund
481 (Álvarez-Noriega et al. 2016). By tracking the morphological development of corals using
482 quantitative metrics, this study deepens our understanding of how coral growth might shape
483 ecological interactions in reef ecosystems.

484 While the results of this study capture some growth-form-specific size-metric relationships and
485 trajectories through continuous morphological space, they also emphasize the large amount of
486 overlap between growth forms in morphological trait space, echoing the findings of Zawada et al.
487 (2019a). However, the current results reveal even more extensive overlap in morphological traits
488 across growth forms, specifically highlighting the developmental variation of coral morphology
489 both within an individual colony and between individuals of a given growth form. This reinforces
490 the idea that these traditional growth forms lack the clear boundaries implied by their categorical
491 nature (Zawada et al. 2019a), especially at smaller colony sizes. The intra-individual and within-
492 growth-form developmental variation of coral morphology observed here once again
493 underscores the need to transition from qualitative to quantitative trait-based approaches to
494 better understand fundamental biological and ecological processes on reefs (Madin et al.
495 2016a).

496 By elucidating these size-metric relationships and individual morphological development, this
497 study lays the groundwork for future studies to more explicitly test reciprocal interactions
498 between coral morphology and reef ecosystem processes. This quantitative framework can be
499 extended to directly evaluate how morphological changes impact the surrounding environment,
500 and how environmental conditions in turn shape coral morphology, working to establish causal
501 links between the two processes (Jones et al. 1997; Odling-Smee et al. 2003). Integrating this
502 morphological data with environmental variables like light availability, temperature, and water
503 movement for example, offers a promising avenue for understanding the feedback loop between
504 coral structure and the reef environment.

References

Agisoft LLC (2021) Agisoft Metashape Professional, v1.7.3

Álvarez-Noriega M, Baird AH, Dornelas MA, Madin JS, Cumbo VR, Connolly SR (2016) Fecundity and the demographic strategies of coral morphologies. Ecology 87:3485-3493. https://doi.org/10.1002/ecy.1588

Autodesk Inc. (2017) Meshmixer, v3.5.474

Bolnick DI, Amarasekare P, Araújo MS, Bürger R, Levine JM, Novak M, Rudolf VHW, Schreiber SJ, Urban MC, Vasseur DA (2011) Why intraspecific trait variation matters in community ecology. Trends Ecol Evol 26:183-192. https://doi.org/10.1016/j.tree.2011.01.009

Bürkner P-C (2017) brms: an R package for Bayesian multilevel models using Stan. J Stat Softw 80:1 - 28. https://doi.org/10.18637/jss.v080.i01

Bythell JC, Pan P, Lee J (2001) Three-dimensional morphometric measurements of reef corals using underwater photogrammetry techniques. Coral Reefs 20:193-199. https://doi.org/10.1007/s003380100157

Camargo A (2022) PCAtest: testing the statistical significance of Principal Component Analysis in R. PeerJ 10:e12967. https://doi.org/10.7717/peerj.12967

Cope OL, Burkle LA, Croy JR, Mooney KA, Yang LH, Wetzel WC (2022) The role of timing in intraspecific trait ecology. Trends Ecol Evol 37:997-1005. https://doi.org/10.1016/j.tree.2022.07.003

Darling ES, McClanahan TR, Côté IM (2013) Life histories predict coral community disassembly under multiple stressors. Global Change Biol 19:1930-1940. https://doi.org/10.1111/gcb.12191

Darling ES, Álvarez-Filip L, Oliver TA, McClanahan TR, Côté IM (2012) Evaluating life-history strategies of reef corals from species traits. Ecol Lett 15:1378-1386. https://doi.org/10.1111/j.1461-0248.2012.01861.x

Dornelas MA, Madin JS, Baird AH, Connolly SR (2017) Allometric growth in reef-building corals. Proc R Soc B 284:20170053. https://doi.org/10.1098/rspb.2017.0053

Figueira W, Ferrari R, Weatherby E, Porter A, Hawes S, Byrne M (2015) Accuracy and precision of habitat structural complexity metrics derived from underwater photogrammetry. Remote Sens 7:16883-16900. https://doi.org/10.3390/rs71215859

Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A (2019) Visualization in Bayesian workflow. J R Stat Soc Ser A Stat Soc 182:389-402. https://doi.org/10.1111/rssa.12378

Gelman A, Meng X-L, Stern H (1996) Posterior predictive assessment of model fitness via realized discrepancies. Stat Sin 6:773-760.

Gelman A, Vehtari A, Simpson D, Margossian CC, Carpenter B, Yao Y, Kennedy L, Gabry J, Bürkner P-C, Modrák M (2020) Bayesian workflow. arXiv:1-77. https://doi.org/10.48550/arXiv.2011.01808

Great Barrier Reef Marine Park Authority (GBRMPA) (2022) Great Barrier Reef Features, Vector geospatial data. Queensland, Australia: GBRMPA. Available: https://geohub-gbrmpa.hub.arcgis.com/datasets/GBRMPA::great-barrier-reef-features-/about

Green SJ, Brookson CB, Hardy NA, Crowder LB (2022) Trait-based approaches to global change ecology: moving from description to prediction. Proc R Soc B 289:20220071. https://doi.org/10.1098/rspb.2022.0071

Guendulain-García SD, Banaszak AT, Álvarez-Filip L, Quattrini AM, López-Pérez A (2024) Three-dimensional morphological variation and physical functionality of Caribbean corals. Coral Reefs 43:405-413. https://doi.org/10.1007/s00338-024-02472-1

Guendulain-García SD, Lopez-Beltran A, Banaszak AT, Álvarez-Filip L, Ramírez-Chávez E, García-Medrano D, Sellares-Blasco R, López-Pérez A (2023) Photogrammetry for coral structural complexity: what is beyond sight? Coral Reefs 42:635-644. https://doi.org/10.1007/s00338-023-02368-6

Gutiérrez-Heredia L, D'Helft C, Reynaud EG (2015) Simple methods for interactive 3D modeling, measurements, and digital databases of coral skeletons. Limnol Oceanogr 13https://doi.org/10.1002/lom3.10017

House JE, Brambilla V, Bidaut LM, Christie AP, Pizarro O, Madin JS, Dornelas MA (2018) Moving to 3D: relationships between coral planar area, surface area and volume. PeerJ 6:e4280. https://doi.org/10.7717/peerj.4280

Jones CG, Lawton JH, Shachak M (1997) Positive and negative effects of organisms as physical ecosystem engineers. Ecology 78:1946-1957. https://doi.org/10.1890/0012-9658(1997)078[1946:Paneoo]2.0.Co;2

Kerry JT, Bellwood DR (2011) The effect of coral morphology on shelter selection by coral reef fishes. Coral Reefs 31:415-424. https://doi.org/10.1007/s00338-011-0859-7

Kerry JT, Bellwood DR (2014) Do tabular corals constitute keystone structures for fishes on coral reefs? Coral Reefs 34:41-50. https://doi.org/10.1007/s00338-014-1232-4

Lange ID, Perry CT, Cooper N (2020) A quick, easy and non-invasive method to quantify coral growth rates using photogrammetry and 3D model comparisons. Methods Ecol Evol 11:714-726. https://doi.org/10.1111/2041-210x.13388

Lavy A, Eyal G, Neal B, Keren R, Loya Y, Ilan M, McMahon S (2015) A quick, easy and non-intrusive method for underwater volume and surface area evaluation of benthic organisms by 3D computer modelling. Methods Ecol Evol 6:521-531. https://doi.org/10.1111/2041-210x.12331

Madin JS, Connolly SR (2006) Ecological consequences of major hydrodynamic disturbances on coral reefs. Nature 444:477-480. https://doi.org/10.1038/nature05328

Madin JS, Baird AH, Dornelas MA, Connolly SR (2014) Mechanical vulnerability explains size-dependent mortality of reef corals. Ecol Lett 17:1008-1015. https://doi.org/10.1111/ele.12306

Madin JS, Hoogenboom MO, Connolly SR, Darling ES, Falster DS, Huang D, Keith SA, Mizerek TL, Pandolfi JM, Putnam HM, Baird AH (2016a) A trait-based approach to advance coral reef science. Trends Ecol Evol 31:419-428. https://doi.org/10.1016/j.tree.2016.02.012

Madin JS, Anderson KD, Andreasen MH, Bridge TCL, Cairns SD, Connolly SR, Darling ES, Diaz M, Falster DS, Franklin EC, Gates RD, Harmer AMT, Hoogenboom MO, Huang D, Keith SA, Kosnik MA, Kuo C-Y, Lough JM, Lovelock CE, Luiz OJ, Martinelli J, Mizerek TL, Pandolfi JM, Pochon X, Pratchett MS, Putnam HM, Roberts TE, Stat M, Wallace CC, Widman E, Baird AH (2016b) The Coral Trait Database, a curated database of trait information for coral species from the global oceans. Sci Data 3:160017. https://doi.org/10.1038/sdata.2016.17

McElreath R (2020) Statistical Rethinking: A Bayesian Course with Examples in R and Stan. Chapman and Hall/CRC, New York, US

McGill BJ, Enquist BJ, Weiher E, Westoby M (2006) Rebuilding community ecology from functional traits. Trends Ecol Evol 21:178-185. https://doi.org/10.1016/j.tree.2006.02.002

Messier J, McGill BJ, Lechowicz MJ (2010) How do traits vary across ecological scales? A case for trait-based ecology. Ecol Lett 13:838-848. https://doi.org/10.1111/j.1461-0248.2010.01476.x

Million WC, O'Donnell S, Bartels E, Kenkel CD (2021) Colony-level 3D photogrammetry reveals that total linear extension and initial growth do not scale with complex morphological growth in the branching coral, *Acropora cervicornis*. Front Mar Sci 8:646475. https://doi.org/10.3389/fmars.2021.646475

Muff S, Nilsen EB, O'Hara RB, Nater CR (2022) Rewriting results sections in the language of evidence. Trends Ecol Evol 37:203-210. https://doi.org/10.1016/j.tree.2021.10.009

Odling-Smee FJ, Laland KN, Feldman MW (2003) Niche Construction: The Neglected Process in Evolution. Princeton University Press, Princeton, US

R Core Team (2021) R: A language and environment for statistical computing, v4.1.2. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Reichert J, Schellenberg J, Schubert P, Wilke T (2016) 3D scanning as a highly precise, reproducible, and minimally invasive method for surface area and volume measurements of scleractinian corals. Limnol Oceanogr 14:518-526. https://doi.org/10.1002/lom3.10109

Schiettekatte N, Asbury M, Chen GK, Dornelas M, Reichert J, Torres-Pulliza D, Zawada KJA, Madin JS (2025) habtools: An R package to calculate 3D metrics for surfaces and objects. Methods Ecol Evol 16:895-903. https://doi.org/10.1111/2041-210x.70027

Siefert A, Violle C, Chalmandrier L, Albert CH, Taudiere A, Fajardo A, Aarssen LW, Baraloto C, Carlucci MB, Cianciaruso MV, de L. Dantas V, de Bello F, Duarte LDS, Fonseca CR, Freschet GT, Gaucherand S, Gross N, Hikosaka K, Jackson B, Jung V, Kamiyama C, Katabuchi M, Kembel SW, Kichenin E, Kraft NJB, Lagerström A, Le Bagousse-Pinguet Y, Li Y, Mason N, Messier J, Nakashizuka T, Overton JM, Peltzer DA, Pérez-Ramos IM, Pillar VD, Prentice HC, Richardson S, Sasaki T, Schamp BS, Schöb C, Shipley B, Sundqvist M, Sykes MT, Vandewalle M, Wardle DA (2015) A global meta-analysis of the relative extent of intraspecific trait variation in plant communities. Ecol Lett 18:1406-1419. https://doi.org/10.1111/ele.12508

613   Stambler N, Dubinsky Z (2005) Corals as light collectors: an integrating sphere approach. Coral Reefs 24:1-
614   9. https://doi.org/10.1007/s00338-004-0452-4

615   Stan Development Team (2024) RStan: the R interface to Stan, v2.32.6. https://mc-stan.org/

616   Urbina-Barreto I, Chiroleu F, Pinel R, Fréchon L, Mahamadaly V, Elise S, Kulbicki M, Quod J-P, Dutrieux E,
617   Garnier R, Henrich Bruggemann J, Penin L, Adjeroud M (2021) Quantifying the shelter capacity of
618   coral reefs using photogrammetric 3D modeling: from colonies to reefscapes. Ecol Indic
619   121https://doi.org/10.1016/j.ecolind.2020.107151

620   Veron JEN (2000) Corals of the World. Vol. 1–3. Australian Institute of Marine Science, Queensland,
621   Australia

622   Westerband AC, Funk JL, Barton KE (2021) Intraspecific trait variation in plants: a renewed focus on its role
623   in ecological processes. Ann Biol 127:397-410. https://doi.org/10.1093/aob/mcab011

624   Zawada KJA, Dornelas MA, Madin JS (2019a) Quantifying coral morphology. Coral Reefs 38:1281–1292.
625   https://doi.org/10.1007/s00338-019-01842-4

626   Zawada KJA, Madin JS, Baird AH, Bridge TCL, Dornelas MA (2019b) Morphological traits can track coral reef
627   responses to the Anthropocene. Funct Ecol 33:962-975. https://doi.org/10.1111/1365-
628   2435.13358