Proline transporters balance the salicylic acid-mediated trade-off between

regeneration and immunity in plants

Dawei Xu[1], Zeinu Mussa Belew[2], Nathália Cássia Ferreira Dias[3], Li Wang[1], Andy Zhang[1],

Yun-Fan Stephanie Chen[1], Carter James Newton[1], Feng Kong[1], Yaochao Zheng[4], Yao

Yao[4], Marin Talbot Brewer[1], Paulo José Pereira Lima Teixeira[3], Hussam Hassan Nour

Eldin Auis[2], Deyang Xu[2], Li Yang[1,5*]

1. Department of Plant Pathology, College of Agricultural & Environmental Sciences,

University of Georgia, Athens 30602, USA

2. DynaMo Center, Department of Plant and Environmental Sciences, Faculty of Science,

University of Copenhagen, Frederiksberg, Denmark.

3. Department of Biology, "Luiz de Queiroz" College of Agriculture, University of São

Paulo, Sao Paulo, Brazil

4. Department of Animal and Diary Sciences, College of Agricultural & Environmental

Sciences, University of Georgia, Athens 30602, USA

5. Department of Plant Biology, Franklin College of Arts and Sciences, University of

Georgia, Athens 30602, USA

*Author for correspondence (li.yang1@uga.edu)

Abstract

A robust immune response and regenerative capacity are both essential for survival after injury. In plants, salicylic acid (SA) is essential for activating immunity but simultaneously suppresses regenerative capacity. The mechanisms coordinating the trade-off between immunity and regeneration remain poorly understood. Here, we identify proline transporters as key regulators of this balance. Mutations in two wound-induced proline transporters, ProT2 and ProT3, rescued exogenous proline-induced suppression of de novo root regeneration (DNRR) and enhanced DNRR. ProTs are required for SA–mediated suppression of regeneration, without affecting SA-dependent defense responses. Mechanistically, a ProT3–CPK1 complex modulates the dynamics of wound-induced reactive oxygen species (ROS), sustaining a ROS level that restricts DNRR after wounding. Notably, pharmacological inhibition of proline transport rescued SA-mediated suppression of regeneration and enhanced regeneration across multiple plant species. These findings establish proline transporter as a regulatory hub integrating stress-induced proline metabolism, SA signaling, and ROS homeostasis to balance immunity and regeneration, and highlight chemical inhibition of proline transport as a strategy to improve crop regeneration under biotic stresses without compromising disease resistance.

Introduction

Regeneration and immunity are two fundamental processes that enable plants to survive

wounding and other forms of tissue damage caused by pathogen invasion, herbivore

feeding, and mechanical injuries under extreme weather conditions. Upon damage,

plants initiate wound healing and, in some cases, produce new organs to restore

function [1]. During de novo root regeneration (DNRR), detached leaves or stem cuttings

undergo wound healing and callus formation, leading to adventitious root development,

which are guided exclusively by endogenous hormone signals [2]. At the same time,

multiple layers of immune responses are activated locally or systemically in response to

wound at a time scale from seconds to days to constrain microbial entry and

multiplication [3, 4]. These responses may include the activation of reactive oxygen species

(ROS) bursts, reprogramming of phytohormone signaling, strengthening of cell wall and

secretion of antimicrobial compounds [5]. Although these dual processes—defense and

regeneration—are indispensable for plant survival in natural ecosystems, regeneration is

often disrupted under high pressure of biotic stresses such as the presence of

pathogenic microbes [6].

Recent advancements have begun to unveil the molecular mechanisms through which

immune activation inhibits the regenerative potential in plants [7]. For instance, activation

of salicylic acid (SA) signaling has been repeatedly shown to impair plant regeneration.

SA accumulation after wounding inhibits callus formation in Arabidopsis and maize

tissue culture [8]. Similarly, DNRR from Arabidopsis leaf explants is strongly reduced by

exogenous SA treatment or in mutants with constitutively elevated SA levels [9]. Inside of

a callus, SA relays hypoxia stress signal to suppress shoot organogenesis [10]. Beyond

the role of SA, microbe-associated molecular patterns (MAMPs) can also block DNRR;

the bacterial elicitor flg22, a 22 amino acids peptide derived from flagellin, prevents

adventitious root initiation from wounded tissues [11]. These examples illustrate that

immune signaling—while vital for pathogen defense—can restrict regenerative

competence. Thus, fine-tuning the balance between immunity and regeneration is critical

for plant survival under combined abiotic and biotic stresses.

Wounding often induces multiple stresses simultaneously [3, 12]. Proline is a key metabolite

positioned at the interface of multiple stress responses [13, 14]. Under drought, salinity, or

oxidative stress, proline accumulates rapidly and may function as an osmoprotectant,

stabilizer of proteins and membranes, and regulator of cellular redox homeostasis [15, 16, 17,

18, 19, 20, 21]. Its levels are tightly controlled by metabolic enzymes such as P5CS (pyrroline-

5-carboxylate synthetase) for synthesis and ProDH (proline dehydrogenase) for

degradation, as well as by dedicated transporters [22, 23]. The spatial and temporal

distribution of proline in plant tissues is tightly regulated by proline transporters (ProTs),

such as the Arabidopsis ProT family (AtProT1, AtProT2, and AtProT3) [24, 25, 26].

Arabidopsis proline transporter family mediates the uptake of proline as well as γ-

aminobutyric acid (GABA), and quaternary ammonium compounds in yeast [25, 27]. ProTs

and other proline transporters such as AMINO ACID PERMEASE1 and Lysine Histidine

Transporter 1 (LHT1), link proline metabolism to plant growth, immunity, and stress

adaptation [26, 28]. Recent work shows that MAMP perception triggers LHT1-dependent

uptake of proline from the leaf apoplast, and this targeted depletion reinforces pattern-

triggered immunity and underscores nutrient restriction in apoplast as a key component

of plant defense [26].

SA and proline often act synergistically to alleviate plant stress responses. SA promotes

proline biosynthesis and catabolism under conditions such as drought, salinity, and heat

[15, 19, 29, 30, 31, 32]. Exogenous co-application of SA and proline has been shown to yield

additive or synergistic effects in multiple crops to counteract stress-induced defects [33, 34,

35, 36, 37]. In regeneration contexts, proline displays multifaceted roles. Exogenous proline

enhances callus induction and regeneration in *Brassica napus*, banana, and rice [38, 39, 40],

but in some cases, it reduces somatic embryo numbers (peanut) or inhibits cell growth

(saltgrass, *Distichlis spicata*) during regeneration [41, 42, 43]. These findings suggest that

proline influences cellular competence for regeneration in a species-, cell type- and

dose-dependent manner, although the underlying mechanism is largely unknown.

In this study, we demonstrate that the Arabidopsis proline transporters ProT2 and ProT3

play key roles in maintaining the balance between SA-mediated defense and the plant's

ability to regenerate after damage. We show that exogenous proline suppresses DNRR

in Arabidopsis, correlating with repression of the root founder cell marker *WOX11*.

Mutants of *prot2* and *prot3* exhibit enhanced regenerative capacity, whereas

overexpression of ProT3 impairs wound-induced adventitious root formation. We identify

a ProT3–CPK1 protein complex that restricts regeneration downstream of SA signaling,

likely by maintaining elevated wound-induced ROS. Importantly, this complex is

dispensable for SA-induced defense gene activation, thereby uncoupling SA-mediated

suppression of regeneration and activation of defense. Pharmacological inhibition of

proline transporter activity using the small molecule LQFM215 rescues SA-induced

regeneration defects and promotes organogenesis in Arabidopsis, tomato, and other

species. These findings offer a strategy to enhance crop regenerative capacity without

compromising immunity.

Results

Members of proline transporter family have distinct roles in regeneration

Our previous work showed that wound-induced SA suppressed DNRR, a process in

which roots regenerate from wound sites without exogenous hormones, in Arabidopsis [9].

Since SA and proline act synergistically in response to various abiotic stresses, we

sought to investigate the role of proline in DNRR. We first treated leaf explants with

various concentration of L-proline and found that L-proline reduced DNRR at 10 μM and

completely blocked adventitious root formation at 10mM (Fig. 1a). Being consistent with

the reduced DNRR, L-proline also inhibited the expression of *WOX11*, a marker of cell

fate transition from procambium cell to root founder cell during DNRR, at cutting sites

(Fig. 1b) [44]. These results suggest that endogenous proline may negatively influence

wound-induced regeneration. The analysis of genes that are induced after leaf cutting

revealed that the expression of *ProT3* was rapidly induced after cutting, resembling the

expression pattern of some SA-responsive genes (Extended Data Fig. 1a) [9, 45]. To

investigate ProT3's potential function in regeneration, we first analyzed the DNRR

phenotype of Arabidopsis *prot3-1* and *prot3-2* mutants. Both *prot3-1* and *prot3-2* alleles

exhibited enhanced adventitious root formation compared to Col-0 (Fig. 1c, d). This

enhanced regenerative capacity was restored to wild type level by a genomic *ProT3*

construct containing the *ProT3* coding region and 1919 bp upstream, confirming that the

phenotype results from a recessive loss-of-function mutation (Extended Data Fig. 1b).

We also examined wound-induced callus (WIC), another form of regeneration without

exogenous hormone application. In this system, the *prot3-2* mutant developed enlarged

calli at wound sites (Fig. 1e, f). Furthermore, two independent lines overexpressing

ProT3-YFP under the ubiquitin-10 promoter displayed significantly reduced DNRR (Fig.

1g). Similar to the application of exogenous proline, *ProT3* overexpression also

suppressed wound-induced *WOX11*, but not its expression in un-wounded tissues,

indicating that ProT3 negatively regulates root founder cell establishment during DNRR

(Fig. 1h). We also investigated the DNRR phenotype in mutants of other proline

transporter family members: *prot2* mutant enhanced DNRR similar to *prot3*, whereas

prot1 mutant had no effect, suggesting a functional divergence within the ProT family

(Extended Data Fig. 1d,e).

To further elucidate the spatial and temporal regulation of proline transporters during

DNRR, we generated promoter–GUS reporter lines for *ProT1*, *ProT2*, and *ProT3*.

Immediately after cutting (day 0, D0), GUS activity driven by the *ProT3* promoter was

detected in leaf blades but absent at cutting sites (Fig. 1i). At 1 day after cutting (1 DAC,

D1), *ProT3* was strongly induced at the wound site (Fig. 1i). With the development of

callus tissue, *proProT3::GUS* activity was excluded from callus, with activity restricted to

the peripheral zone (Fig. 1i). Similarly, *proProT2::GUS* was strongly induced at cutting

sites at 1DAC (Extended Data Fig. 1f). In contrast, *proProT1::GUS* expression was

confined to vascular tissues and was not wound-inducible at 1 DAC (Extended Data Fig.

1g). These patterns are consistent with the roles of *ProT3* and *ProT2*, but not *ProT1*, as

negative regulators of wound-induced regeneration. Collectively, these findings highlight

the divergent roles of *ProT* family members in DNRR, likely due to their distinct

expression dynamics.

ProT3 suppresses DNRR through its proline transport activity

To test the role of ProT2 and ProT3 in proline-mediated inhibition of DNRR, we first

validated proline transport activity of ProTs using *Xenopus laevis* oocyte expression

system. Oocytes expressing ProT2 or ProT3 can import ^{3}H-proline into the cells from

external buffer at pH5.0 but not at pH7.4, compared to the cells expressing GFP (***$P <$

0.001, n = 10) (Fig. 2a, b). Moreover, ProT3 showed higher ^{3}H-proline uptake than

ProT2 in 30 minutes import assay (Fig. 2a). Electrophysiological recordings revealed

robust proline-induced inward currents in ProT2- and ProT3-expressing oocytes at

pH5.0, which were absent in GFP controls (Fig. 2c). These suggested that ProTs-

mediated import of proline is driven by proton gradient across plasma membrane. When

treated with L-proline at concentrations of 10 μM and 100 μM that inhibited DNRR in

Col-0, *prot3* mutants were insensitive to such treatment, indicating that the mutant is

defective in importing extracellular proline (Fig. 2d). At 10 mM, L-proline treatment

inhibited DNRR in *prot3* mutants, probably due to toxicity or the function of other low-

affinity proline transporters (Fig. 2d). In contrast, the suppression of DNRR by mannitol

was comparable in both Col-0 and *prot3* mutants (Fig. 2e), suggesting that ProT3's role

is specifically required for proline-induced inhibition of DNRR rather than a reduced

sensitivity to osmolality. Given that ProTs are localized to the plasma membrane [25], these

findings suggest that ProT3-mediated proline import into cells at the wound site, where it

accumulates and suppresses DNRR, while the loss of ProT3 in the mutant reduces this

uptake, alleviating the inhibitory effect of proline.

We next asked whether chemical inhibition of proline transport could mimic the genetic

loss of ProT3. Previous studies have shown that LQFM215, a small-molecule inhibitor of

mammalian proline transporters (SLC6A7), blocks substrate uptake in synaptosomes [46,

47]. Inspired by this, we hypothesized that LQFM215 might also bind to Arabidopsis

ProT2 and ProT3 and inhibit their transport activity. To test this possibility, we first

predicted Arabidopsis ProT2/3 interactions with L-proline using AlphaFold v3.0.1.

Superimposition of five independent models revealed high-confidence complexes (pTM

= 0.87-0.88, ipTM=0.95-0.96 for ProT2; pTM = 0.86, ipTM=0.92-0.94 for ProT3)

(Supplementary Table 1), with L-Proline consistently positioned within a conserved

central substrate-binding cavity (Fig. 2f). We next generated ligand–protein complexes

for LQFM215. Across five independent AlphaFold predictions, LQFM215 adopted

multiple partially overlapping orientations in both ProT2 (pTM = 0.84-0.86, ipTM=0.78-

0.84) and ProT3 (pTM = 0.82-0.85, ipTM=0.69-0.8), forming a cluster of plausible poses

within the substrate pocket (Fig. 2g, Supplementary Table 1), suggesting flexibility in

inhibitor engagement with the transporters. Strikingly, nearly all residues predicted to

contact L-proline in ProT2 and ProT3 were encompassed within the broader set of

residues contacting LQFM215 (Supplementary Table 1). This overlap indicates that

LQFM215 occupies—and extends beyond—the endogenous substrate-binding site.

Because the inhibitor engages the same core residues required for proline coordination,

these structural models strongly suggest a competitive inhibition mechanism in which

LQFM215 interferes with the substrate-binding cavity and prevents proline from entering

the transport pathway.

Following this *in silico* prediction, we performed electrophysiological recordings in

Xenopus oocytes to test whether LQFM215 affects proline transport activity. Application

of 0.2 mM LQFM215 markedly reduced the proline-induced inward currents in oocytes

expressing either ProT2 or ProT3, indicating an inhibitory effect on transporter activity

(Fig. 2h). These results confirmed that LQFM215 functions as an inhibitor of Arabidopsis

ProT activity. We further examined the physiological impact of LQFM215 on DNRR in

Arabidopsis to evaluate its functional effect in planta. When LQFM215 was applied on

leaf explants, it enhanced DNRR by approximately 20% at a concentration as low as

1μM, similar as the enhanced DNRR observed in *prot3* mutants (Fig. 2i). Notably,

LQFM215 also rescued the L-proline–induced suppression of DNRR, thereby confirming

that its promotive effect on DNRR was mediated through inhibition of proline uptake (Fig.

2j). Taken together, these results demonstrate that compromising proline import, either

by knocking out endogenous ProTs function or application of inhibitors of proline

transporters, promotes the regeneration of adventitious root from leaf explants.

prot3 mutation rescues SA-mediated suppression of DNRR

We next investigated the relationship between SA and proline transporters in regulating

DNRR. Exogenous NaSA, an SA mimic, inhibited DNRR in Col-0 [9], while *prot3* and *prot2*

mutants were insensitive to this suppression (Fig. 3a,b). Similar to what we observed in

prot3 mutants, LQFM215 treatment also rescued the suppression of DNRR caused by

exogenously applied SA (Fig. 3c). To test if LQFM215 rescues DNRR defects caused by

endogenous SA, we used transgenic plants overexpressing the bifunctional SA synthase

gene *Irp9* from the pathogenic bacterium *Yersinia enterocolitica* under the control of the

35S promoter (*35S::Irp9*) [48]. *35S::Irp9* contains elevated level of SA and SA-derived

conjugates and enhances SA-mediated disease resistance [48]. DNRR was strongly

suppressed in *35S::Irp9* (Fig. 3e). However, application of 100 µM of LQFM215 partially

restored adventitious root formation from explants of *35S::Irp9* (Fig. 3e), suggesting that

interfering proline uptake rescues SA-induced suppression of regeneration. SA also

contributes to an age-dependent decline of regeneration [8, 9] with reduced DNRR capacity

in explants from 8-, 11-, and 14-day old plants. Notably, DNRR in *prot3* mutants

remained age-dependent, with the rooting ratio decreasing from 76.2% in 8-day-old

explants to 46.4% in 14-day-old explants, despite the overall higher rooting ratio

compared to Col-0 (Fig. 3f), suggesting that ProT3 only acts in a branch of SA-mediated

suppression of DNRR or other age-dependent signals inhibit DNRR in old leaves. On the

other side, L-proline-mediated inhibition of DNRR was also compromised in *NahG*

transgenic line which possess reduced SA level due to the expression of a salicylate

hydroxylase (Fig. 3g). These data suggested that SA and proline may act synergically to

suppress DNRR.

We further tested whether microbe-induced inhibition of DNRR was affected by ProT3.

Treatment with *Pseudomonas syringae pv. tomato* (*Pto*) hrcC- mutants, a compromised

bacterium defective in effector delivery but sufficient to induce PTI, inhibited DNRR in

prot3-1 and *prot3-2* mutants to a similar extent as in Col-0 (Extended Data Fig. 2a),

indicating that ProT3 does not influence PTI-mediated DNRR suppression. Similarly, 1

μM of flg22, a peptide-derived from bacterial flagellin capable of activating basal

immunity, suppressed DNRR in *prot3* as in Col-0 (Extended Data Fig. 2b). These results

are consistent with our previous observations that flg22-induced suppression of DNRR

was independent of SA [6]. Taken together, immune response activates multiple pathways

to inhibit regeneration, mutation in *prot3* only blocks the SA-mediated suppression of

DNRR.

ProT3 is dispensable for SA-mediated activation of defense genes and inhibition

of bacteria growth

SA regulates a tradeoff between regeneration capacity and immunity [8, 9, 10]. To test

whether ProT3 is also required for SA-mediated activation of immunity, we measured the

transcript levels of the SA-responsive genes *PR1* and *SARD1* following the same SA

treatment used in the SA-mediated suppression of DNRR assay (Fig. 4a,b). The

expression level of these SA responsive genes was comparable in Col-0 and *prot3*

mutants (Fig. 4a,b), indicating that ProT3 is not required for the activation of these SA

responsive genes in 11-day-old seedlings. Previous work showed that the expression of

proline transporters including ProT2 and ProT3 were activated upon flg22 treatment,

which depletes apoplastic proline as a defense strategy [26]. We also observed that *prot2*

and *prot3* single mutant showed mild enhanced susceptibility to *Pseudomonas syringae*

pv. *tomato* DC3000 (*Pto* DC3000) infection (Fig. 4c, Extended Data Fig. 3a). To

specifically test if *prot3* is required for SA-mediated suppression of bacterial growth, we

monitored bacterial load of *Pto* DC3000 in Col-0 and *prot3* mutants pre-treated with

benzothiadiazole (BTH), an SA analogue (Fig. 4c). As expected, the pretreatment of

BTH activated SA-mediated immunity and reduced bacterial growth in Col-0 (Fig. 4c).

Such reduction was also observed in *prot2* and *prot3* mutants to a similar level (Fig. 4c),

further supporting that ProT2 and 3 were dispensable for SA-dependent suppression of

bacteria multiplication.

ProT3 interacts with CPK1 to suppress regeneration

When exploring the potential function of ProT3 in regulating regeneration, we identified

calcium-dependent protein kinase 1 (CPK1) as a protein interactor of ProT3 from public

database (https://thebiogrid.org/) [49]. Bimolecular fluorescence complementation (BiFC)

and co-immunoprecipitation assays based on transiently expressed proteins in *Nicotiana*

benthamiana confirmed a physical interaction between ProT3 and CPK1 (Fig. 5a,b).

Although *CPK1* expression was not induced by cutting (Extended Data Fig. 3b), two

alleles of *cpk1* single mutants showed enhanced DNRR compared to Col-0 (Fig. 5c,d),

supporting CPK1 as a negative regulator of regeneration. Similar to *prot3* mutants, *cpk1*

mutants also displayed higher WIC formation than Col-0 (Fig. 5e,f), indicating that a

ProT3-CPK1 complex cooperatively suppress wound-induced regeneration. CPK1 was

also dispensable for SA-mediated protection against *Pto* DC3000 (Fig. 5g) or SA-

induced defense gene activation (Fig. 5h,i). However, the *cpk1* mutant did not block SA-

mediated suppression of DNRR (Extended Data Fig. 3c), indicating a signaling

divergence between ProT3 and CPK1 or redundancy between CPK1 and closely related

kinases. Taken together, these molecular and genetic evidence support a conclusion that

a ProT3-CPK1 complex is required for wound-induced regeneration but not the

activation of SA-mediated defense, decoupling the two key functions downstream of

wound-activated SA.

ProT3 regulates ROS equilibrium during DNRR

To interrogate the transcriptomic changes induced by *prot3* mutation, we performed an RNA-seq experiment comparing *prot3* and Col-0 transcriptomes at time 0 (D0) and 1 day after cutting (D1). At D0, we identified 255 and 113 genes that are differentially expressed (DEGs) in blades and petioles, respectively (Fig. 6a). At D1, there were 321 and 194 DEGs between the two genotypes in blades and petioles, respectively (Fig. 6a). It is noteworthy that only 19 DEGs in blades and 8 DEGs in petioles are shared between D0 and D1, indicating a dramatic shift of expression in response to wound (Fig. 6a). We noticed that there was no enrichment of SA-related GO terms in DEGs D0 or D1 by comparing blade or petiole samples from Col-0 and *prot3*, which further supported our observation that *prot3* did not generally contribute to SA-mediated defense (Supplementary Table 2). In addition, key genes involved in DNRR regeneration, including *YUCCA2* and *PLT7*, known to promote regeneration through auxin biosynthesis and meristem maintenance, were upregulated in *prot3* petioles at D1 (Fig. 6b,c), which was consistent with the enhanced adventitious root formation (Fig. 1c,d). To further explore the ProT3-controlled biological processes that are important for regeneration, we performed a hierarchical clustering analysis of all 714 genes that were differentially expressed between *prot3* and Col-0 at either D0 or D1 (Fig. 6b). Notably, in cluster 3, DEGs were more activated in the D1 leaf blades of *prot3* (Extended Data Fig. 4a). In addition, DEGs in cluster 4 were less activated in both blades and petioles of *prot3* (Extended Data Fig. 4b). We noticed that enriched GO terms related to oxygen metabolism such as "response to oxygen-containing compound" in cluster 3 and cluster 4 genes (Supplementary Table 3), consistent with proline's role as a ROS regulator through its metabolic cycling in cytoplasm and mitochondria. We further assessed ROS accumulation by H2DCFDA staining at cutting sites in Col-0 and *prot3 cpk1* (Fig.6d-g).

ROS signal was immediately accumulated at wound sites after cutting, no difference was

observed between Col-0 and *prot3-2cpk1-2* double mutant at D0 (Fig. 6d,e). However, at

D1, the double mutants showed reduced ROS accumulation at cutting sites compared to

Col-0 (Fig. 6f,g). Application of 10 µM LQFM215 immediately after cutting also reduced

ROS accumulation at D1 (Fig. 6h,i). Previous studies have suggested that wound-

induced ROS signal promotes DNRR [50]. Consistently, we also observed that H_2O_2

treatment at time 0 promoted DNRR in Col-0 (Fig. 6j). In contrast, application H_2O_2 at 1

day after cutting reduced rooting by ~15% in Col-0 (Fig. 6j), suggesting that a reduced

sensitivity to, or level of, ROS signal at later stage is required for full regeneration

capacity. Strikingly, *prot3* mutants were insensitive to both the promotive effect of early

H_2O_2 treatment or the inhibitory effect of H_2O_2 applied at D1 (Fig. 6j), indicating that loss

of the ProT3-CPK1 complex facilitate a permissive ROS environment at 1 day after

cutting for root regeneration. While wound-induced ROS accumulation is required to

initiate regeneration [50], subsequent ROS reduction may be necessary for DNRR

progression. Together, these results suggest that the ProT3–CPK1 complex modulates

ROS dynamics during DNRR, integrating early wound-induced stress signals into

regenerative programs.

LQFM215 promotes regeneration in crops

The *ProT* gene family is widely present and highly conserved in plants (Extended Data

Fig. 5). We further tested the effect of LQFM215 on promoting regeneration in tomato,

tobacco and rice. In tomato (*Solanum lycopersicum* cv. Money Maker), treatment of

hypocotyl explants with LQFM215 significantly increased callus size on callus-induction

medium (CIM) compared with mock treatment (Fig. 7a,b). Similarly, treatment of leaf

explants with LQFM215 resulted in a higher number of adventitious roots per explant,

with all root classes increased relative to mock (Fig. 7c,d). In *N. benthamiana*, leaf disc

explants of uniform size treated with LQFM215 exhibited higher shoot regeneration

efficiency and more shoots per explant compared with controls (Fig. 7e,f). In rice

(*Nipponbare*), scutellum-derived calli cultured on CIM were larger upon LQFM215

treatment, indicating enhanced callus proliferation (Fig. 7g,h).

Collectively, these results demonstrated that LQFM215 enhances multiple modes of

regeneration across diverse crops, extending its function beyond Arabidopsis. This

raises the possibility that LQFM215 might help overcome the trade-off between disease

resistance and regenerative capacity, providing a strategy to improve regeneration in

elite crop varieties while maintaining desirable agronomic traits.

Discussion

Our study identifies ProTs as critical regulators of DNRR in *Arabidopsis*. We uncovered

their role in balancing SA-mediated regeneration and immunity. ProTs are required for

SA-mediated suppression of regeneration but not for defense. The function of ProTs is

probably achieved via a complex with CPK1. Notably, LQFM215, an inhibitor of plant

ProTs, is sufficient to alleviate SA-mediated suppression of DNRR in multiple plant

species, suggesting that proline metabolism represents a conserved mechanism linking

biotic and abiotic stresses to regenerative capacity. These findings not only broaden the

known functions of proline transporters beyond abiotic stress tolerance but also provide

new insights into the molecular basis of the growth–defense trade-off in plants.

Wound-induced ROS signaling plays a key role in wound healing and organ

regeneration across both animals and plants [51, 52, 53]. The oxidative burst serves as an

early cue that guides wound closure, immune cell recruitment, and stem/progenitor

activation. In zebrafish tail regeneration, ROS recruit Hedgehog-expressing cells to the

wound site, thereby activating Hedgehog signaling essential for regeneration [54]. In

planarians, ROS acts upstream of the MAPK/ERK cascade and can even rescue

regeneration when ERK signaling is inhibited [55]. In amphibians such as axolotl, H_2O_2

contributes to blastema formation and growth by activating regeneration-related genes

and recruiting immune cells to the injury site [56, 57]. In plants, early ROS signals, likely

produced directly by tissue damage, play a promotive role in regeneration [50, 58, 59, 60, 61, 62].

However, evidence also indicates that excess ROS can inhibit shoot regeneration [63, 64]

and microcallus formation from protoplasts [65]. Moreover, genetic reduction of ROS levels

enhances DNRR or shoot regeneration in Arabidopsis [58, 66]. Together, these observations

suggest that while ROS act as essential early wound signals, sustained or excessive

ROS accumulation may exert inhibitory effects on regeneration beyond their initial

signaling role.

Our results showed that the effect of exogenous H_2O_2 switched from promotive to

inhibitory within one day after cutting, and that mutation in ProT3 blocked this sensitivity

(Fig. 6j), suggesting that ProT3 may play a role in regulating endogenous ROS dynamics

or cellular responses to ROS. In mitochondria, proline dehydrogenase (ProDH)

catalyzes the oxidation of proline to pyrroline-5-carboxylate (P5C), transferring electrons

to the mitochondrial electron transport chain [67, 68, 69]. This reaction generates ROS,

primarily superoxide ($O_2\bullet^-$) and hydrogen peroxide (H_2O_2), which serve as signaling

molecules that activate downstream defense and stress-response pathways [70].

Following leaf detachment, AtProDH1 was rapidly induced within one hour in explants,

exhibiting a pattern similar to that of ProT3 (Extended Data Fig. 6). These observations

suggest that proline transport and degradation may be coordinately regulated to promote

wound-induced ROS production. We speculate that excessive or prolonged ROS

accumulation derived from proline catabolism—as observed one day post-cutting at

wound sites—may inhibit the development of root founder cells, thereby constraining

DNRR. Previous work showed that ROS levels increased between 3–6 days after calli

were cultured on shoot induction medium (SIM), then declined from days 8 to 16. In

mutants of *dcc1*, a thioredoxin, and *ca2* (carbonic anhydrases 2), an interactor of *dcc1*,

ROS levels in calli remained high through day 16, correlating with a reduced shoot

regeneration phenotype [66]. In contrast, in the *prot3 cpk1* mutants, a reduced ROS level

at later stages post-wounding may create a permissive environment for root founder cell

establishment (Fig. 6f,g). The dynamics of ROS levels after wounding and their roles in

wound-induced regeneration may explain dosage- and timing dependent phenotype of

ROS impact on various forms of plant regeneration. Interestingly, *ProT3* is activated at

wound sites but later excluded from developing callus (Fig. 1i), suggesting that a low

proline, probably leading to a low ROS condition, is necessary for proceeding with the

adventitious root initiation. Thus, proline transporters such as ProT3 and ProT2 may act

as metabolic gatekeepers that couple amino acid flux to redox dynamics. Evidence from

plant immunity revealed that SA induces *ProDH* expression during the hypersensitive

response, and ProDH activity is required for full ROS accumulation and pathogen

resistance induced by SA [70]. It is possible that the loss of ProTs rescued SA-mediated

suppression of DNRR by counteracting the SA-induced ROS level. Notably, other amino

acid transporters, including LHT1 [26] and members of the AAP family [28], may also

contribute to proline homeostasis in addition to ProTs. Beyond its role as a ROS

regulator, proline may also serve as a metabolic substrate or osmo-regulator during

regeneration. Although our data show that ProT2/3 transport proline and that their

function is inhibited by LQFM215, we cannot rule out the possibility that other ProT

cargos also contribute to regeneration. How proline flux and subcellular localization are

precisely regulated to maintain redox balance remains a key question for elucidating the

mechanistic link stress response and regenerative capacity.

In plants, regeneration and immunity are tightly interconnected processes. Elite crop

varieties bred for durable disease resistance often exhibit constitutive or primed immune

activity, including elevated salicylic acid (SA) signaling, stronger microbe-associated

molecular pattern (MAMP) responses, and enhanced hypersensitive cell death. While

these traits confer resilience against pathogens, they may also compromise regenerative

competence [6]. For example, IR64, a high-yielding indica rice variety carrying resistance

genes such as Xa4 for bacterial blight, has been widely used in breeding programs but

remains one of the most recalcitrant cultivars for regeneration and Agrobacterium-

mediated transformation, requiring specialized tissue culture protocols for plant recovery

[71, 72]. Although there is no direct evidence that recalcitrance is caused by enhanced

defense responses, strategies that transiently suppress defense signaling or fine-tune

immune activation during plant tissue culture or de novo organogenesis could help

overcome these limitations. Notably, we showed that pharmacological inhibition of

proline transport by LQFM215 alleviated SA-induced suppression of regeneration in

multiple plant species, indicating that proline flux functions as a tunable regulatory node

in the antagonistic relationship between plant immunity and regeneration.

Methods

Plant growth and transformation

Arabidopsis thaliana T-DNA lines and transgenic lines used in this study are in the

Columbia-0 (Col-0) ecotype. Homozygous lines were selected by gene specific primers

LP and RP, in combination with T-DNA vector primers (Supplementary Table 4). Primers

used for vector construction are listed in Supplementary Table 4. Double knockout lines

were generated by crossing homozygous single *prot3-2* with *prot2-3* to form the *prot2-*

3prot3-2 double mutant, *prot3-2* with *cpk1-2* to form the *prot3-2cpk1-2* double mutant.

Transgenic constructs were introduced via Agrobacterium tumefaciens-mediated

transformation using the floral dip method [73]. Additionally, *Nicotiana benthamiana* plants

were cultivated under long-day conditions (16 h of light / 8 h of dark) in the growth room

at University of Georgia, maintained at 22 ℃, and 70% relative humidity, with a light

intensity of 30000 lumens/m^2.

Characterization of the mutant DNRR phenotype and wounding-induced callus

formation

DNRR phenotyping was performed as previously described [74], with minor modifications.

Arabidopsis seeds for DNRR testing were surface sterilized by 70% ethanol, germinated,

and grown on ½ MS media (Murashige & Skoog Basal Medium with Vitamins,

PhytoTechnology Laboratories) which were placed in the growth chambers (Percival,

USA) with continuous light at 22^{o}C. The 1st and 2nd true leaves from 11-day-old

seedlings were cut at the junction between the leaf blade and petiole and placed with the

abaxial side down onto Gamborg's B5 media (RPI Research Products International).

Explants were kept in continuous light condition. The adventitious roots were counted

every day under a dissecting microscope starting from 6 DAC until the 14 DAC day as

previously described [9]. Morphological assessments of whole plants were documented

using a Canon EOS Rebel T7 DSLR Camera (EF-S 18–55mm f/3.5–5.6 IS II). Leaf

explants with adventitious root formation were photographed using a Zeiss Axio

Zoom.V16 Stereo Microscope. DNRR assays performed on Phytagel plates and sand

plates (for *Pto* DC3000 hrC⁻ treatments) were performed as previously outlined [11]. The

callus size was scored 10 days after excision and callus areas were quantified using

ImageJ software.

Chemical treatments

To investigate the effects of various chemical cues on DNRR, detached Arabidopsis leaf

explants were subjected to the following treatments.

For salicylic acid (SA) treatments, sodium salicylate (NaSA) was added to B5 medium at

final concentrations of 0 µM (mock), 1 µM, 5 µM, and 25 µM. Explants were transferred

onto the respective media, and the adventitious root formation ratio was quantified as

described above.

For the regeneration response to proline, detached leaf explants were transferred onto

B5 medium containing different concentrations of L-proline (0 µM, 10 µM, 100 µM, 1 mM,

and 10 mM).

For testing the DNRR response to flg22 elicitor treatment, detached leaf explants were

pre-treated by spraying them with 1 µM flg22 for 1 hour. The explants were then placed

on B5 medium, and 10 µl of flg22 was applied directly to the wound sites after cutting.

For osmotic stress assays, D-mannitol was added to B5 medium at final concentrations

of 0 mM (mock), 10 mM, and 100 mM to impose mild and moderate osmotic stress

conditions, respectively.

For ROS treatments, freshly prepared 200 µM H_2O_2 was directly applied to the wound

sites of leaf explants at different time points after cutting to evaluate regeneration

responses under oxidative stress.

To test whether LQFM215, a small-molecule inhibitor of proline transporters, promotes

regeneration, LQFM215 was applied topically to the wound sites of leaf explants

immediately after placement on the regeneration medium. Unless otherwise indicated,

3–10 μL of 1, 10 or 100 μM LQFM215 was pipetted directly onto each cutting site. The

explants were subsequently cultured under standard regeneration conditions, and

regeneration phenotypes were evaluated at the indicated time points.

qRT-PCR analysis

For quantitative reverse transcription PCR (qRT-PCR) analysis, total RNA was extracted

from leaf blade cuttings at different time points using the E.Z.N.A Plant RNA Kit (R6827-

01, Omega BIO-TEK). First-strand cDNA was synthesized from total RNA using

GoScript™ Reverse Transcriptase (A5003, Promega). Real-time RT-PCR was

conducted with iQ SYBR Green Supermix (Bio-rad) on the Applied Biosystems

QuantStudio 5 real-time PCR system (Applied Biosystems by Thermo Fisher Scientific).

qRT-PCR procedures and data analysis were performed as previously described [9].

Primers used for qRT-PCR are listed in Supplemental Table 4. The endogenous control

Tublin 2 was not differentially expressed from time 0 to 2 days after cutting in explants [45].

It is also not affected by the genotypes [45].

GUS reporter assay for *ProT* genes expression

Plasmids *pKGWFS7-ProT1Pro::GUS*, *pKGWFS7-ProT2Pro::GUS*, *pKGWFS7-*

ProT3Pro::GUS and *pKGWFS7-CPK1Pro::GUS* were constructed to examine tissue-

specific expression of ProTs and CPK1, with the GUS marker protein driven by the 2576

bp, 2465 bp, 1919 bp and 2123 bp promoter regions of *ProT1*, *ProT2*, *ProT3*, and *CPK1*

respectively. GUS activity was visualized by staining young seedlings and leaf blade

cuttings at different time points from various transgenic lines overnight in 5-Bromo-4-

chloro-3-indolyl-b-D-glucuronide (X-Gluc), as previously described [9]. Subsequently,

tissues were cleared in 70% (v/v) ethanol. Cleared samples were observed and

photographed using a Zeiss Axio Zoom.V16 Stereo Microscope and processed using

Zeiss ZEN 3.8 and ImageJ software.

514 Bimolecular fluorescence complementation (BiFC) of ProT3 and CPK1

For BiFC analysis, the full-length coding sequences (CDS) of *ProT3* and *CPK1* were

amplified and were subcloned into the pENTR™/D-TOPO™ Cloning Kit (Thermal fisher

scientific) to form the entry vetor *pENTR-ProT3* and *pENTR-CPK1* respectively. The

resultant plasmids were then introduced to the tagging vectors *pUBN-nYFP* and *pUBN-*

cYFP by using Gateway™ LR Clonase™ II Enzyme mix Kit (Thermal fisher scientific) to

generate *ProT3-nYFP* and *CPK1-cYFP* constructs. The primers used for vector

construction are listed in Supplementary Table 4. The resulting plasmids were introduced

into *Agrobacterium tumefaciens* strain GV3101 by chemical transformation. Single

colonies were cultured overnight in LB medium containing appropriate antibiotics,

harvested by centrifugation, and resuspended in infiltration buffer (10 mM $MgCl_2$, 10 mM

MES, pH 5.6, and 150 µM acetosyringone) to an OD_{600} = 0.4. The suspensions carrying

ProT3-nYFP and *CPK1-cYFP* were mixed at a 1:1 ratio before infiltration. The

Agrobacterium mixtures were infiltrated into the abaxial side of 4–5-week-old *Nicotiana*

benthamiana leaves using a needleless syringe. After infiltration, plants were maintained

in the dark for approximately 48 h to facilitate transient expression. YFP fluorescence

signals were first detectable around 40–48 h post infiltration, and the strongest

fluorescence was typically observed at 48–60 h. Fluorescence was examined using a

Zeiss LSM880 confocal laser scanning microscope with excitation at 514 nm and

emission collected between 525–560 nm.

534 Co-immunoprecipitation of ProT3 and CPK1

For co-immunoprecipitation (Co-IP) assays, entry vetor *pENTR-ProT3* and *pENTR-*

CPK1 were cloned into the binary vectors pGWB641 (driven by the CaMV 35S promoter,

C-terminal EYFP tag) and pGWB614 (35S promoter, C-terminal 3×HA tag), respectively.

The resulting constructs (*35S:ProT3-EYFP* and *35S:CPK1-3×HA*) were introduced into

Agrobacterium tumefaciens strain GV3101 by chemical transformation. Single colonies

were cultured overnight in LB medium with appropriate antibiotics, harvested by

centrifugation, and resuspended in infiltration buffer (10 mM $MgCl_2$, 10 mM MES, 150

µM acetosyringone, pH 5.6) to an OD_{600} of 0.1. Equal volumes of Agrobacterium

suspensions carrying ProT3-EYFP and CPK1-3×HA were mixed (1:1 ratio) and infiltrated

into 4–5-week-old *Nicotiana benthamiana* leaves using a needleless syringe. Infiltrated

plants were maintained in darkness for 2 days to allow transient protein expression. To

prevent protein degradation, 50 µM MG-132 (Sigma-Aldrich) was infiltrated into the

same leaf areas 24 h post Agrobacterium infiltration, and leaves were harvested 48 h

post infiltration for protein extraction.

Total protein was extracted from infiltrated *Nicotiana benthamiana* leaves with protein

extraction buffer (50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM EDTA, 10 mM NaF, 5

mM Na3VO4 , 0.25% Triton X-100, 0.25% NP-40, 1 mM PMSF, 1× protease inhibitor

cocktail), and then incubated with 25 µl anti-GFP agarose beads (Chromotek, gta-20) for

3 h at 4 °C. The beads were washed 3 times with protein extraction buffer, and the

precipitated proteins were eluted with 2× SDS loading buffer at 95 °C for 5 min. The

samples were subjected to immunoblot analysis using anti-GFP and anti-HA through

western blot respectively.

RNA-seq analysis

Young leaf blades and petioles from Col-0 and *prot3* plants were collected at D0 and D1

(1 day after cutting), with approximately 20 leaf blades per genotype per time point for

each biological replicate. Three biological replicates were prepared for each

genotype/treatment. Samples were frozen in liquid nitrogen and milled using a

homogenizer (OMNI International). Total RNA was extracted using the E.Z.N.A Plant

RNA Kit (R6827-01, Omega BIO-TEK), and RNA concentration was measured using a

Nanodrop spectrophotometer (Thermo Scientific). RNA samples were sent to Admera

Health (https://www.admerahealth.com) for library preparation and sequencing. RNA

integrity was assessed using TapeStation or BioAnalyzer High Sensitivity RNA Assays

(Agilent Technologies), and RNA was quantified using an Infinite F Nano+ 200 Pro

(Tecan, Switzerland). Poly(A)+ RNA was purified and used for strand-specific library

construction with the NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina®,

following the manufacturer's instructions. Equimolar pooling of libraries was performed

based on QC values and sequenced on an Illumina NovaSeq X Plus platform with a

read

length configuration of 150 PE for 40 M PE reads per sample (20M in each direction).

Sequencing data processing, including read mapping, alignment, and analysis of

differentially expressed genes (DEGs) were performed as previously described [75].

Differential gene expression analysis was conducted in R using the edgeR package [76] to

compare the two genotypes (Col-0 vs *prot3*) across different tissues (leaf blade or

petiole) at each time point (D0 or D1). Genes with very low expression levels were

excluded by retaining only those with at least 1 Transcripts Per Million (TPM) in five or

more samples. The Benjamini–Hochberg correction was applied to adjust p-values for

multiple testing [77]. Genes were defined as differentially expressed when the false

discovery rate (FDR) was ≤ 0.05 and the fold-change was at least twofold. Gene

Ontology (GO) enrichment analyses were performed at the Plant GeneSet Enrichment

Analysis Toolkit (PlantGSEA) online webserver [78]. Hierarchical clustering of expression

data was conducted using the ComplexHeatmap package in R[79], applying the Euclidean

distance and the complete linkage method. Expression levels, expressed as TPM values,

were standardized to z-scores prior to clustering. Raw RNA-seq reads are deposited in

the NCBI Sequence Read Archive (SRA) with BioProject ID PRJNA1357857.

Proline transport assay in *Xenopus laevis* oocytes

The coding DNA sequences of *ProT2* and *ProT3* were cloned into *Xenopus* expression

vector pNB1u using USER cloning [80, 81] Linear DNA templates for *in vitro* transcription

were generated by PCR from the pNB1 plasmid using forward primer (5'–

AATTAACCCTCACTAAAGGGTTGTAATACGACTCACTATAGGG–3') and reverse primer

(5'–TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTATACTCAAGCTAGCCTCGAG–3')   and   a

Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific) following the

manufacturer's instructions. The PCR product was purified using E.Z.N.A. Cycle Pure Kit

(Omega BIO-TEK) following the manufacturer's instructions. Capped cRNA was in vitro

synthesized using mMessage mMachine T7 Kit (InVitrogen), following the

manufacturer's instructions. The cRNA concentration was normalized to 500 ng μl^{-1} for

oocyte injection.

Defolliculated *Xenopus laevis* oocytes at stage V–VI (Ecocyte Bioscience, Germany)

were microinjected with 50.6 nL of cRNA using the Drummond NANOJECT II

(Drummond Scientific Company, Broomall, Pennsylvania, USA), with GFP serving as a

negative control. The injected oocytes were incubated at 16°C for three days in HEPES-

based kulori buffer (90 mM NaCl, 1 mM KCl, 1 mM $MgCl_2$, 1 mM $CaCl_2$, 10 mM HEPES,

pH 7.4) supplemented with amikacin (100 $\mu g\ mL^{-1}$).

Three days after cRNA injection, oocytes were initially pre-incubated for 5 min in kulori

buffer pH 5.0 (90 mM NaCl, 1 mM KCl, 1 mM $MgCl_2$, 1 mM $CaCl_2$, 10 mM MES, pH 5) or

pH 7.4 (90 mM NaCl, 1 mM KCl, 1 mM $MgCl_2$, 1 mM $CaCl_2$, 10 mM HEPES, pH 7.4) and

then transferred to 150 μl of the same buffer (pH 5.0 or pH 7.4) containing 47 μCi/ml of

^{3}H-L-proline (93.2 Ci/mmol, Revvity) and incubated for 30 min. Afterwards, the oocytes

were washed four times with kulori buffer of the respective pH. Each oocyte was then

placed in a scintillation vial and lysed in 100 μl of 10% (w/v) SDS by vigorous vortexing.

Subsequently, 2.5 ml of EcoScint^{TM} scintillation fluid (National Diagnostics) was added to

the vial, mixed thoroughly by vortexing, and the radioactivity was measured using liquid

scintillation counting.

Two-electrode voltage-clamp electrophysiology

The two-electrode voltage-clamp (TEVC) technique was employed to determine

electrogenicity of the proline transport via ProT2 and ProT3 and further characterize

them. The TEVC recordings were carried out using the automated Roboocyte2 system

(Multichannel Systems, Reutlingen, Germany). The electrodes (resistance 280-1000 kΩ)

were filled with a solution containing 3 M potassium chlorideand 1.5 M potassium

acetate. Oocytes expressing ProT2, ProT3, or GFP (negative control) were clamped at -

60 mV membrane potential and currents were recorded under continuous perfusion with

MES-based ekulori buffer (2 mM LaCl3, 90 mM NaCl, 1 mM KCl, 1 mM MgCl2, 1 mM

CaCl2, 10 mM MES pH 5), with or without 5 mM L-proline. To test the inhibitory effect of

LQFM215, 0.2 mM LQFM215 and 5 mM L-proline were used.

Bacterial growth assay

Bacterial growth curve assays were performed as previously described [75]. Briefly,

Pseudomonas syringae pv. tomato (*Pto*) DC3000 strain was suspended in 10 mM MgCl$_2$

to a final concentration of 1 x 10□ CFU/mL and infiltrated into Arabidopsis leaves using a

1 mL syringe without a needle. After inoculation, plants were covered with transparent

lids for one hour. Day 0 samples were collected immediately after lid removal, and Day 3

samples were collected three days post-infiltration (dpi). Each sample consisted of four

leaf discs (one from each of four individual leaves) punched with a uniform-sized leaf

corer. The discs were homogenized using a homogenizer (OMNI International), and the

resulting homogenate was serially diluted (10^{-1} to $10^{-\square}$). A 10 µL aliquot of each dilution

was plated onto King's B solid medium containing rifamycin and cycloheximide and

incubated at room temperature for two days. Colony-forming units (CFUs) were

manually counted and normalized to the total leaf area sampled.

For benzothiadiazole (BTH) pretreatment, an SA analogue, plants were sprayed with 50

µM BTH solution using a sterilized sprayer one day before pathogen inoculation. After

spraying, plants were kept in the greenhouse overnight under the same growth

conditions. The following day, bacterial growth curve assays were conducted as

described above.

Reactive oxygen species (ROS) detection

Reactive oxygen species were also visualized using the fluorescent probe 2',7'-

dichlorodihydrofluorescein diacetate (H_2DCFDA; D399, Thermo Fisher Scientific) as

previously described with minor modification [82]. A 1 mM H_2DCFDA stock solution was

prepared in DMSO and stored at –20 °C in the dark. Before use, the stock was diluted

with 10 mM Tris-HCl buffer (pH 7.2) to a working concentration of 10 µM. Leaf explants

were collected at indicated time points (D0 and 1 days after cutting) and immersed in 10

µM H_2DCFDA solution, vacuum-infiltrated at 60 kPa for 5 min, and incubated at room

temperature for 10 min in the dark. After staining, samples were rinsed thoroughly with

10 mM Tris-HCl buffer to remove residual dye. Fluorescence was observed using a

Zeiss LSM 880 confocal laser scanning microscope with excitation at 488 nm and

emission at 530 nm. Green fluorescence indicated ROS accumulation, while red

autofluorescence corresponded to chlorophyll. Fluorescent areas were quantified using

ImageJ software by measuring the average signal intensity in a defined area. The area

was defined as tissues 2 mm away from the cutting edge.

Regeneration assays in crop species with LQFM215

For tomato DNRR, *Solanum lycopersicum* cv. Money Maker seeds were sterilized in 70%

ethanol for 30 s, followed by 10% bleach for 15 minutes twice, rinsed five times with

sterile water, dried on sterile filter paper, and germinated on ½ MS at 22 °C under

continuous light for ~2 weeks. The first true leaves were excised at the petiole fully using

sterile scissors and placed adaxial side up on B5 plates. Apply 3 µL of 10 µM and 100

µM of LQFM215 directly onto the wound sites. Explants were incubated at 22 °C,

continuous light, and adventitious roots (AR) were scored at 10 days post-cutting.

For hypocotyl-callus inducion, 10-day-old tomato seedlings were cut with scissors and

hypocotyl segments were incubated on CIM (MS + 3% sucrose, 2 mg/L 2,4-D, 0.1 mg/L

6-BA, 0.375% Phytagel; pH 5.8). Immediately after placement, 3 µL of 10 µM LQFM215

was applied to the cut surfaces. Plates were kept at 22 °C, continuous light, for ~2

weeks before callus phenotypes were scored. Callus images were acquired on a Zeiss

Axio Zoom.V16 stereo microscope, and projected areas were quantified in ImageJ.

For *Nicotiana tabacum* shoot induction, fully expanded healthy leaves of *Nicotiana*

tabacum were surface sterilized in 75% ethanol for 5 min, rinsed three times with sterile

water, incubated in 10% bleach for 7 min, and rinsed five times with sterile water. Sterile

leaf discs were punched with a cork borer, blotted dry, and placed abaxial side down on

regeneration medium (MS+ 3% sucrose, 1 mg/L 6-BA, 0.1 mg/L NAA, 0.375% Phytagel;

pH 5.8). Plates were incubated at 28 °C under a 16-h light/8-h dark photoperiod for ~4

weeks, with subculture to fresh medium every 14 days. Fo LQFM215 treatments, 10 µL

of 10 µM LQFM215 was applied evenly around the perimeter of each disc at day 0.

Shoot number per disc was scored at day 30.

Callus induction in *Oryza sativa* ssp. *japonica* (cv. Nipponbare) was performed as

previously described [56] with minor adaptations. Dehulled mature seeds were surface-

sterilized in 50% Clorox solution containing 5 µL Tween-20 for 30 min, rinsed three times

with sterile water, and placed on callus-induction 2N6 medium supplemented with 2.0

mg/L 2,4-D. Plates were incubated at 28 °C in darkness for 11 days. Scutellum-derived

calli were imaged by Zeiss Axio Zoom.V16 Stereo Microscope and callus area per seed

was quantified in ImageJ. Where indicated, 10 µL drop of 10 µM LQFM215 was pipetted

directly onto the seed surface immediately after plating.

Molecular docking and structure prediction

Complexes of Arabidopsis ProT2 (Q9P2962) and ProT3 (Q9SJP9) with L-proline or

LQFM215 (ProT2–L-proline, ProT2–LQFM215, ProT3–L-proline, and ProT3–LQFM215)

were predicted using AlphaFold v3.0.1[83] on an NVIDIA A100 GPU. For each protein–

ligand complex, five independent models were generated using different random seeds.

Protein–ligand contact residues of ProT2 and ProT3 with L-proline or LQFM215 were

then analyzed in these models using ChimeraX v1.10.1[84], with a standard distance cutoff

of 4 Å. Colors for different models were selected using the qualitative palette from

ColorBrewer 2.0.

Acknowledgements

We thank L. Xu (CAS Center for Excellence in Molecular Plant Sciences, China), C. H.

Khang (University of Georgia, US) for sharing materials. This project is supported by the

National Science Foundation (IOS-2039313 to L.Y.) and the National Institutes of Health

(R35GM143067 to L.Y.). Support was also provided by the Serrapilheira Institute (G-

1811-25705 to P.J.P.L.T.), by the National Council for Scientific and Technological

Development (CNPq; 308349/2022-9 to P.J.P.L.T.), and by the São Paulo Research

Foundation (FAPESP; 2018/24432-0 to P.J.P.L.T.; 2023/10167-1 to N.C.F.D.).

Computational resources were provided in part by the Georgia Advanced Computing

Resource Center, a partnership between the University of Georgia's Office of the Vice

President for Research and Office of the Vice President for Information Technology.

Author contributions

716 L.Y. and D.X. conceptualized the study. D.X., Z.M.B., A.Z., S.C., Y.Z., and F.K. performed

717 experiments. N.C.F.D., C.J.N., L.W., and M.T.B performed data analyses. D.X., L.Y.,

718 D.Y.X., H.H.N.E.A, P.J.P.L.T. and Y.Y. contributed to data interpretation. D.X. and L.Y.

719 wrote the manuscript with input from all authors. All authors reviewed and approved the

720 final manuscript.

Figure 1

Fig. 1 | ProT3 is induced by wounding and negatively regulates DNRR.

a, Ratio of adventitious root formation on Col-0 leaf explants treated with increasing concentrations of L-proline (0, 10, 100 µM, 1, and 10 mM). (Each dot represents one biological replicate with 40 explants. n=3). The insert shows an Arabidopsis seedling (left) and an explant with adventitious root (right). Red dashed line indicates cutting position.

b, Representative images of *proWOX11::GUS* staining in leaf explants treated with 100 µM L-proline at 1 day after cutting (1DAC). scale bar: 1 mm. Red arrowheads indicate the cutting sites.

c, Morphology of 11-day-old plants (top) and DNRR phenotypes (bottom) of Col-0, *prot3-1* and *prot3-2*; scale bars, 0.5 cm (top) and 1 mm (bottom).

d, Rooting ratio of Col-0, *prot3-1* and *prot3-2* explants cultured for 10 d (n > 7. 40 explants for each repeat).

e, f, *prot3* mutation enhanced wound-induced callus (WIC) (e) and the statistical analysis of callus area (f). n = 20 petioles per genotype, scale bar, 1 mm.

g. Rooting ratio in Col-0, ProT3-OE-6, and ProT3-OE-7 leaf explants. (n = 3 biological replicates, 40 explants each).

h, *proWOX11::GUS* expression in Col-0 and ProT3-OE-7 leaf explants at 1 DAC, scale
bar, 1 mm. Arrows and arrowheads indicate GUS activity at cutting sites and shoot-
hypocotyle junction, respectively.

i, *proProT3::GUS* expression in leaf explants at D0, 1DAC, 2DAC and 5DAC, scale bar,
1 mm.

Data are presented as mean ± s.d.; n denotes the number of biological replicates
Statistical significance was determined by two-tailed Student's t-test; *P < 0.05, **P
<0.01, ***P < 0.001; ns, not significant.

Figure 2

Fig. 2 | ProT3 modulates DNRR through its proline transport activity and is inhibited by a proline transporter inhibitor LQFM215.

a, b, Proline uptake assay in *Xenopus laevis* oocytes. [3]H-proline uptake was measured at pH 5.0 (a) and pH 7.4 (b). Oocytes expressing ProT2, ProT3, or GFP (negative control) were incubated in kulori buffer containing 0.5 µM [3]H-proline for 30 min (n = 10 oocytes).

c, Electrophysiological recordings of ProT2 and ProT3-mediated proline transport in *Xenopus* oocytes. Representative traces of proline-induced currents recorded from oocytes expressing ProT2, ProT3, or GFP (negative control) at a holding membrane potential of -60 mV at pH5.0.

d, Rooting ratio in Col-0 and *prot3* leaf explants treated with L-proline (0-10 mM) (n = 3 biological replicates, 40 explants each).

e, Rooting ratio in Col-0 and *prot3* leaf explants under mannitol treatment (0–100 mM) (n

= 3 biological replicates, 40 explants each).

f, g, Superimposed AlphaFold v3.0.1 models of Arabidopsis ProT2 and ProT3 in complex

with L-proline (f) and LQFM215 (g). For ProT2 and ProT3, five predicted models with

different colors are superimposed. Model 1, light blue (#a6cee3); model 2, dark blue

(#1f78b4); model 3, light green (#b2df8a); model 4, dark green (#33a02c); and model 5,

light red (#fb9a99). L-proline is shown in dark red (#e31a1c) in all models. For LQFM215,

ligand colors reflect their binding positions. In the ProT2-LQFM215 complex, ligands in

models 2 and 4 occupy a similar position and are colored purple (#984ea3), whereas

ligands in models 1, 3, and 5 are colored dark red (#e31a1c); the same coloring scheme

is applied to the ProT3-LQFM215 complex.

h, Electrophysiological analysis showing inhibition of ProT2 and ProT3 by LQFM215.

The effect of LQFM215 on ProT2- and ProT3-mediated proline-induced currents was

assessed by applying 0.2 mM LQFM215 (red bar) in the presence of 5 mM proline (black

bar) at a holding potential of -60 mV at pH5.0.

i, Rooting ratio in Col-0 by different concentration of LQFM215 treatment (n = 5

biological replicates, 40 explants each).

j, LQFM215 alleviates proline-mediated suppression of DNRR in Col-0 (n = 5 biological

replicates, 40 explants each).

Data are presented as mean ± s.d.; *n* denotes the number of biological replicates.

Statistical significance was determined by two-tailed Student's *t*-test; *$P < 0.05$, **P

<0.01, ***$P < 0.001$; ns, not significant.

Fig. 3 | Mutation of ProT3 abolishes salicylic acid (SA)-mediated suppression of DNRR.

a, b, Rooting ratio in Col-0, *prot3-1*, *prot3*-2 (a) and *prot2-3* (b) explants treated with NaSA (0–25 µM) (n = 3 biological replicates, 40 explants each).

c, Rooting ratio in Col-0 explants co-treated with 1 µM NaSA and 100 µM LQFM215 (n = 3 biological replicates, 40 explants each).

d, Representative images showing that 10 µM LQFM215 alleviates SA-mediated suppression of DNRR in Col-0 leaf explants. Scale bar: 1 mm.

e, Rooting ratio in *35S::Irp9* (high SA level) with or without 100 µM LQFM215 (n = 3 biological replicates, 40 explants each).

f, Rooting ratio in Col-0 and *prot3-2* explants from 8-, 11- and 14-day-old plants (n = 3 biological replicates, 40 explants each).

g, Rooting ratio in Col-0 and *NahG* explants treated with L-proline (10 µM–1 mM) (n = 3 biological replicates, 40 explants each).

Data presented as mean ± s.d.; n denotes the number of biological replicates. Statistical significance was determined by two-tailed Student's t-test; *P < 0.05, **P <0.01, ***P < 0.001; ns, not significant.

Fig. 4 | ProT3 does not interfere with SA-induced defense activation.

a, b, qRT-PCR analysis of *PR1* (a), *SARD1* (b) expression in Col-0 and *prot3*-2 leaf explants from 11-day old seedlings under 25 μM NaSA treatment. (n = 3 biological replicates).

c, Bacterial growth of *Pseudomonas syringae* pv. *tomato* DC3000 in seven-week-old plants of Col-0, *prot2*-3, and *prot3*-2. The left panel shows Pto DC3000 growth kinetics without chemical pretreatment. The right panel shows assessing SA-dependent resistance, where plants were spray treated with 50 μM BTH and inoculated with Pto DC3000 1 day after BTH treatment. Bacterial load was quantified 2 days post inoculation (D2). (n = 4 for D0; n = 6 for D2).

Data are presented as mean ± s.d.; Statistical significance was determined by two-tailed Student's *t*-test; *P < 0.05, **P <0.01, ***P < 0.001; ns, not significant.

Fig. 5 | ProT3 interacts with CPK1 to suppress regeneration.

a, Bimolecular fluorescence complementation (BiFC) assay showing interaction between ProT3 and CPK1 in *Nicotiana benthamiana*, scale bar, 50 µm. Fluorescent signals were detected 2 days after inoculation with Agrobacteria carrying corresponding plasmids.

b, Co-immunoprecipitation assay confirming ProT3-CPK1 interaction in *N. benthamiana*.

c, Morphology (top) and DNRR phenotype (bottom) of Col-0, *cpk1-1* and *cpk1-2*, scale bars, 0.5 cm (top) and 1 mm (bottom).

d, Rooting ratio in Col-0, *cpk1-1* and *cpk1-2* (n = 3 biological replicates, 40 explants each).

e, f, Wound-induced callus (WIC) formation in Col-0, *prot3-2* and *cpk1-1* (e) and quantification of callus area (f), n = 20 petioles per genotype, scale bar, 1 mm.

g, Bacterial growth of *Pseudomonas syringae* pv. tomato DC3000 in Col-0 and *cpk1* mutant with or without BTH pretreatment. The left panel shows DC3000 growth kinetics without chemical pretreatment. The right panel shows BTH-induced resistance; plants were spray-treated with 50 µM BTH one day before DC3000 inoculation. Bacterial loads were quantified 2 days after inoculation (n = 4 for D0; n= 6 for D2).

h,i, qRT-PCR analysis of *PR1* (h) and *SARD1* (i) expression in 11-day-old Col-0 and *cpk1-2* seedlings treated with 25 µM NaSA treatment. (n = 3 biological replicates).

Data are presented as mean ± s.d.; Statistical significance was determined by two-tailed
Student's *t*-test; *P < 0.05, **P <0.01, ***P < 0.001; ns, not significant.

Figure 6

Fig. 6 | ProT3 regulates reactive oxygen species (ROS) homeostasis during DNRR.

a. Arabidopsis leaf showing the regions used for RNA-seq (blade, bl, blue box; petiole, pt,
red box) and Venn diagram showing the overlap of differentially expressed genes (DEGs)
between *prot3-2* and Col-0 in blade and petiole tissues at D0 and 1 day after cutting (D1).

b, Hierarchical clustering of 710 genes that were differentially expressed between Col-0
and *prot3-2* at least of time point in leaf blades or petioles. Statistically significant
differentially expressed genes (FDR ≤ 0.05) are marked with green (downregulated) or
purple (upregulated) stripes. Representative auxin-related genes *YUCCA2* and *PLT7* are
highlighted.

c, Expression profiles of *PLT7* in leaf blade and petiole tissues at D0 and 1 day after
cutting (D1) in Col-0 and *prot3-2*. Box plots show normalized expression values (z-
scores) derived from RNA-seq data.

d-g, H2DCFDA fluorescence in leaf explants of Col-0 and *prot3-2/cpk1-2* mutants at D0

(d) and 1DAC (f). Quantification of H2DCFDA fluorescence was summarized below

representative images. Dotted lines outline wound sites. scale bar: 200 µm.

h,i, H2DCFDA fluorescence in leaf explants of Col-0 treated with 10 µM LQFM215

treatment at 1DAC. Quantification of fluorescence intensity is shown in (i).

j, Effect of exogenous H_2O_2 treatment on DNRR in Col-0 and *prot3-2* at different times

after cutting. H_2O_2 was applied either immediately after cutting (D0) or at D1. Rooting

ratios from each treatment are shown. (n = 3 biological replicates, each with 40 explants).

Data are presented as mean ± s.d.; Statistical significance was determined by two-tailed

Student's *t*-test; *P < 0.05, **P <0.01, and ***P < 0.001; ns, not significant.

Figure 7

Fig. 7 | LQFM215 enhances regeneration across multiple crop species.

a, b, Tomato leaf explants treated with LQFM215 (0, 10, 100 µM) showing enhanced

adventitious root formation at day 10 (a) and distribution of explants according to the

number of adventitious roots per explant, expressed as a percentage of total explants (b,

n = 20); Mann–Whitney U test; scale bar, 5 mm.

c, d, Tomato hypocotyl-derived callus formation with or without 10 μM LQFM215 (c) and

quantification (d, n = 21 explants); scale bar, 1 mm.

e, f, Shoot regeneration from *Nicotiana benthamiana* leaf discs with or without 10 μM

LQFM215 (e); shoot counts after 30 d (f, n=11); scale bar, 5 mm.

g, h, Scutellum-derived callus formation from rice (*Oryza sativa* cv. Nipponbare) seeds

with or without 10 μM LQFM215 (g) and quantification (h, n = 11 seeds); scale bars, 1

mm.

Unless otherwise indicated, data are presented as mean ± s.d.; n denotes the number of

biological replicates. Panels showing percentage distributions (Fig. 7d) represent data

from a single experiment without error bars. Statistical significance was determined by

two-tailed Student's t-test; $*P < 0.05$, $**P < 0.01$, and $***P < 0.001$; ns, not significant.

Extended Data Fig. 1 Expression and functional analysis of *ProT* genes.

a, Wound-induced expression of *ProT1*, *ProT2* and *ProT3 based on* public transcriptomic datasets [50].

b, Rooting ratio of Col-0, *prot3-2* and complementation lines (n = 3 biological replicates, 40 explants each).

c, Sequence alignment of Arabidopsis *ProTs* and homologs. Red lines indicate transmembrane domain.

d, Representative DNRR phenotype of Col-0, *prot1-1* and *prot2-3* after 10 DAC; Scale bars, 0.5 cm (top) and 1 mm (bottom).

e, Rooting ratio of Col-0, *prot1-1* and *prot2-3* (n = 3 biological replicates, 40 explants each).

f, g, GUS staining of *proProT2::GUS* (f) and *proProT1::GUS* (g) at D0 and D1 cultured on B5 medium. scale bars, 500 μm (top) and 1 mm (bottom).

Data presented as mean ± s.d.; *n* denotes the number of biological replicates. Statistical significance was determined by two-tailed Student's *t*-test; *P < 0.05, **P <0.01, ***P < 0.001; ns, not significant.

Extended Data Fig. 2 Rooting response of *prot3* mutants under biotic stress conditions.

a, Rooting inhibition rate of Col-0, *prot3-1* and *prot3-2* leaf explants upon *hrrC⁻* treatment. (n = 3 biological replicates, 40 explants each). hrcC⁻ was applied at a concentration of OD600=0.01.

b, Rooting ratio of Col-0, *prot3-1* and *prot3-2* leaf explants cultured on mock or flg22-supplemented medium for 10 days. (n = 3 biological replicates, 40 explants each).

Data presented as mean ± s.d.; *n* denotes the number of biological replicates. Statistical significance was determined by two-tailed Student's *t*-test; *P < 0.05, **P <0.01, ***P < 0.001; ns, not significant.

Extended Data Fig. 3

a

b

proCPK1::GUS

D0

D1

c

Extended Data Fig. 3 CPK1 function in DNRR and defense.

a, Summary of bacteria growth curve from multiple experiments. Each dot represents the
$\log_{10}$CFU difference between Col-0 and mutant in one independent experiment. *prot3*,
prot2 and *cpk1* mutants showed mild enhance of susceptibility to *Pto* DC3000.

b, *proCPK1::GUS* expression is not induced by wounding. GUS staining of
proCPK1::GUS Arabidopsis leaves at day 0 (D0, a) and 1 day after cutting (D1, b)
cultured on B5 medium. No detectable induction of *CPK1* promoter activity was
observed following wounding. Scale bar, 1 mm.

c, Rooting ratio of Col-0 and *cpk1-2* in response to increasing concentration of NaSA.

971 **Extended Data Fig. 4** | Expression patterns of gene clusters affected by ProT3 in
972 Arabidopsis leaf explants.
973 a, Genes in cluster 3 were more strongly induced in the leaf blades of prot3-2 at 1 day
974 after cutting (D1). b, Genes in cluster 4 displayed lower activation levels in both blades
975 and petioles of *prot3-2* compared with Col-0 at D1.

984 **Extended Data Fig. 5 Phylogenetic analysis of proline transporter (ProT)**
985 **homologs from Arabidopsis, tomato, rice, and *Nicotiana benthamiana*.**

A neighbor-joining phylogenetic tree showing the relationships among predicted proline
transporter (ProT) homologs from *Arabidopsis thaliana* (AtProT1–3), *Solanum*
lycopersicum (Solyc), *Oryza sativa* (LOC_Os), and *Nicotiana benthamiana* (Nbe). The
tree was constructed using MEGA 3.1 with 1,000 bootstrap replicates, and bootstrap
support values are indicated at each node. The scale bar represents 0.2 substitutions
per site.

1000

1001

1002

1003

1004

1005

Extended Data Fig. 6 Wound-induced expression of AtProDH1 (AT3G30775) and
AtProDH2 (AT5G38710) based on public transcriptomic datasets.

Temporal expression patterns of AtPRODH1 and AtPRODH2 following leaf cutting.
Transcript abundance (transcripts per million, TPM) was extracted from the public
wound-induced regeneration RNA-seq dataset [50]. AtPRODH1 shows a rapid induction
within 6 h after wounding, followed by a gradual decline, whereas AtPRODH2 remains at
a low basal level throughout the time course.

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

Reference

1023

1024   1.   Ikeuchi M*, et al.* Molecular mechanisms of plant regeneration. *Annual
1025   review of plant biology* **70**, 377–406 (2019).

1026
1027   2.   Xu L. De novo root regeneration from leaf explants: wounding, auxin, and
1028   cell fate transition. *Current opinion in plant biology* **41**, 39–45 (2018).

1029
1030   3.   Vega-Muñoz I, Duran-Flores D, Fernández-Fernández ÁD, Heyman J,
1031   Ritter A, Stael S. Breaking bad news: dynamic molecular mechanisms of
1032   wound response in plants. *Frontiers in plant science* **11**, 610445 (2020).

1033
1034   4.   Hilleary R, Gilroy S. Systemic signaling in response to wounding and
1035   pathogens. *Current Opinion in Plant Biology* **43**, 57–62 (2018).

1036
1037   5.   Ngou BPM, Jones JD, Ding P. Plant immune networks. *Trends in plant
1038   science* **27**, 255–273 (2022).

1039
1040   6.   Tran S, Chen Y-FS, Xu D, Ison M, Yang L. Microbial Pattern Recognition
1041   suppresses de novo organogenesis. *Development*, (2023).

1042
1043   7.   Xu D, Yang L. Regeneration and defense: unveiling the molecular
1044   interplay in plants. *New Phytologist* **246**, 2484–2494 (2025).

1045
1046   8.   Hernandez-Coronado M*, et al.* Plant glutamate receptors mediate a bet-
1047   hedging strategy between regeneration and defense. *Developmental cell*
1048   **57**, 451–465. e456 (2022).

1049
1050   9.   Tran S*, et al.* Endogenous salicylic acid suppresses de novo root
1051   regeneration from leaf explants. *PLoS Genetics* **19**, e1010636 (2023).

1052
1053   10.   Koo D, Lee HG, Bae SH, Lee K, Seo PJ. Callus proliferation-induced
1054   hypoxic microenvironment decreases shoot regeneration competence in
1055   Arabidopsis. *Molecular Plant* **17**, 395–408 (2024).

1056
1057   11.   Tran S, Stephanie Chen Y-F, Xu D, Ison M, Yang L. Microbial pattern
1058   recognition suppresses de novo organogenesis. *Development* **150**,
1059   dev201485 (2023).

1060

1061 12. Lee K, Seo PJ. Wound-induced systemic responses and their coordination
1062 by electrical signals. *Frontiers in Plant Science* **13**, 880680 (2022).

1063
1064 13. Ghosh U, Islam M, Siddiqui M, Cao X, Khan M. Proline, a multifaceted
1065 signalling molecule in plant responses to abiotic stress: understanding the
1066 physiological mechanisms. *Plant Biology* **24**, 227–239 (2022).

1067
1068 14. Hosseinifard M, Stefaniak S, Ghorbani Javid M, Soltani E, Wojtyla Ł,
1069 Garnczarska M. Contribution of exogenous proline to abiotic stresses
1070 tolerance in plants: a review. *International Journal of Molecular Sciences*
1071 **23**, 5186 (2022).

1072
1073 15. La VH, Lee B-R, Zhang Q, Park S-H, Islam MT, Kim T-H. Salicylic acid
1074 improves drought-stress tolerance by regulating the redox status and
1075 proline metabolism in Brassica rapa. *Horticulture, Environment, and
1076 Biotechnology* **60**, 31–40 (2019).

1077
1078 16. Khan P, Abdelbacki AM, Albaqami M, Jan R, Kim K-M. Proline promotes
1079 drought tolerance in maize. *Biology* **14**, 41 (2025).

1080
1081 17. Ghaffari H, Tadayon MR, Nadeem M, Cheema M, Razmjoo J. Proline-
1082 mediated changes in antioxidant enzymatic activities and the physiology of
1083 sugar beet under drought stress. *Acta physiologiae plantarum* **41**, 23
1084 (2019).

1085
1086 18. Bandurska H, Niedziela J, Pietrowska-Borek M, Nuc K, Chadzinikolau T,
1087 Radzikowska D. Regulation of proline biosynthesis and resistance to
1088 drought stress in two barley (Hordeum vulgare L.) genotypes of different
1089 origin. *Plant Physiology and Biochemistry* **118**, 427–437 (2017).

1090
1091 19. Iqbal N, Fatma M, Khan NA, Umar S. Regulatory role of proline in heat
1092 stress tolerance: modulation by salicylic acid. In: *Plant signaling
1093 molecules*). Elsevier (2019).

1094
1095 20. Tonhati R, Mello SC, Momesso P, Pedroso RM. L-proline alleviates heat
1096 stress of tomato plants grown under protected environment. *Scientia
1097 Horticulturae* **268**, 109370 (2020).

1098
1099 21. Kavi Kishor PB, Suravajhala P, Rathnagiri P, Sreenivasulu N. Intriguing
1100 role of proline in redox potential conferring high temperature stress
1101 tolerance. *Frontiers in Plant Science* **13**, 867531 (2022).

22. Lehmann S, Funck D, Szabados L, Rentsch D. Proline metabolism and transport in plant development. *Amino acids* **39**, 949–962 (2010).

23. Alvarez ME, Savouré A, Szabados L. Proline metabolism as regulatory hub. *Trends in Plant Science* **27**, 39–55 (2022).

24. Akbudak MA, Filiz E. Genome-wide investigation of proline transporter (ProT) gene family in tomato: bioinformatics and expression analyses in response to drought stress. *Plant Physiology and Biochemistry* **157**, 13–22 (2020).

25. Grallath S*, et al.* The AtProT family. Compatible solute transporters with similar substrate specificity but differential expression patterns. *Plant Physiology* **137**, 117–126 (2005).

26. Rogan CJ*, et al.* Transporter-mediated depletion of extracellular proline directly contributes to plant pattern-triggered immunity against a bacterial pathogen. *Nature Communications* **15**, 7048 (2024).

27. Breitkreuz KE, Shelp BJ, Fischer WN, Schwacke R, Rentsch D. Identification and characterization of GABA, proline and quaternary ammonium compound transporters from Arabidopsis thaliana. *FEBS letters* **450**, 280–284 (1999).

28. Wang T*, et al.* Arabidopsis AMINO ACID PERMEASE1 contributes to salt stress-induced proline uptake from exogenous sources. *Frontiers in plant science* **8**, 2182 (2017).

29. Misra N, Saxena P. Effect of salicylic acid on proline metabolism in lentil grown under salinity stress. *Plant Science* **177**, 181–189 (2009).

30. Ignatenko A, Talanova V, Repkina N, Titov A. Exogenous salicylic acid treatment induces cold tolerance in wheat through promotion of antioxidant enzyme activity and proline accumulation. *Acta Physiologiae Plantarum* **41**, 80 (2019).

31. Khan MIR, Iqbal N, Masood A, Per TS, Khan NA. Salicylic acid alleviates adverse effects of heat stress on photosynthesis through changes in proline production and ethylene formation. *Plant Signaling & Behavior* **8**, e26374 (2013).

1143

1144 32. Nazar R, Umar S, Khan N, Sareer O. Salicylic acid supplementation
1145 improves photosynthesis and growth in mustard through changes in
1146 proline accumulation and ethylene formation under drought stress. *South*
1147 *African Journal of Botany* **98**, 84–94 (2015).

1148
1149 33. Urmi TA*, et al.* Combined effect of salicylic acid and proline mitigates
1150 drought stress in rice (Oryza sativa L.) through the modulation of
1151 physiological attributes and antioxidant enzymes. *Antioxidants* **12**, 1438
1152 (2023).

1153
1154 34. Abdelaal KA*, et al.* Exogenous application of proline and salicylic acid can
1155 mitigate the injurious impacts of drought stress on barley plants
1156 associated with physiological and histological characters. *Sustainability* **12**,
1157 1736 (2020).

1158
1159 35. Sofy MR, Seleiman MF, Alhammad BA, Alharbi BM, Mohamed HI.
1160 Minimizing adverse effects of pb on maize plants by combined treatment
1161 with jasmonic, salicylic acids and proline. *Agronomy* **10**, 699 (2020).

1162
1163 36. El-Beltagi H, El-Nady M, Ismail AM, Metwaly MMS. Mitigation of salinity
1164 stress effects on Vicia faba L. growth and productivity using proline and
1165 salicylic acid foliar application. *Plant Protection Science* **61**, 222 (2025).

1166
1167 37. Agami RA, Alamri SA, Abd El-Mageed T, Abousekken M, Hashem M.
1168 Salicylic acid and proline enhance water use efficiency, antioxidant
1169 defense system and tissues' anatomy of wheat plants under field deficit
1170 irrigation stress. *J Appl Bot Food Qual* **92**, 360–370 (2019).

1171
1172 38. Pawar B, Prashant K, Bahurupe J, Jadhav A, Anil K, Pawar S. Proline and
1173 glutamine improve in vitro callus induction and subsequent shooting in rice.
1174 *Rice Science* **22**, 283–289 (2015).

1175
1176 39. Ahmadi B, Shariatpanahi ME. Proline and chitosan enhanced efficiency of
1177 microspore embryogenesis induction and plantlet regeneration in Brassica
1178 napus L. *Plant Cell, Tissue and Organ Culture (PCTOC)* **123**, 57–65
1179 (2015).

1180
1181 40. Husin N, Jalil M, Othman RY, Khalid N. Enhancement of regeneration
1182 efficiency in banana (Musa acuminata cv. Berangan) by using proline and
1183 glutamine. *Scientia Horticulturae* **168**, 33–37 (2014).

1184
1185   41.   Murch SJ, Victor JMR, Krishnaraj S, Saxena PK. The role of proline in
1186   thidiazuron-induced somatic embryogenesis of peanut. *In Vitro Cellular &*
1187   *Developmental Biology - Plant* **35**, 102–105 (1999).

1188
1189   42.   Trigiano R, Conger B. Regulation of growth and somatic embryogenesis
1190   by proline and serine in suspension cultures of Dactylis glomerata. *Journal*
1191   *of plant physiology* **130**, 49–55 (1987).

1192
1193   43.   Rodriguez MM, Heyser JW. Growth inhibition by exogenous proline and its
1194   metabolism in saltgrass (Distichlis spicata) suspension cultures. *Plant cell*
1195   *reports* **7**, 305–308 (1988).

1196
1197   44.   Liu J*, et al.* WOX11 and 12 are involved in the first-step cell fate transition
1198   during de novo root organogenesis in Arabidopsis. *The Plant Cell* **26**,
1199   1081–1093 (2014).

1200
1201   45.   Zhang G*, et al.* Jasmonate-mediated wound signalling promotes plant
1202   regeneration. *Nature plants* **5**, 491–497 (2019).

1203
1204   46.   Carvalho GA*, et al.* Novel Proline Transporter Inhibitor (LQFM215)
1205   Presents Antipsychotic Effect in Ketamine Model of Schizophrenia.
1206   *Neurochem Res* **49**, 170–183 (2024).

1207
1208   47.   Carvalho GA*, et al.* L-proline transporter inhibitor (LQFM215) promotes
1209   neuroprotection in ischemic stroke. *Pharmacol Rep* **75**, 276–292 (2023).

1210
1211   48.   Ortega MA*, et al.* Altering cold-regulated gene expression decouples the
1212   salicylic acid–growth trade-off in Arabidopsis. *The Plant Cell* **36**, 4293–
1213   4308 (2024).

1214
1215   49.   Oughtred R*, et al.* The BioGRID interaction database: 2019 update.
1216   *Nucleic acids research* **47**, D529–D541 (2019).

1217
1218   50.   Liu W*, et al.* Transcriptional landscapes of de novo root regeneration from
1219   detached Arabidopsis leaves revealed by time-lapse and single-cell RNA
1220   sequencing analyses. *Plant Communications* **3**, (2022).

1221
1222   51.   Suzuki N, Mittler R. Reactive oxygen species-dependent wound
1223   responses in animals and plants. *Free Radical Biology and Medicine* **53**,
1224   2269–2276 (2012).

1225

1226 52. Hunt M, Torres M, Bachar-Wikstrom E, Wikstrom JD. Cellular and
1227   molecular roles of reactive oxygen species in wound healing.
1228   *Communications biology* **7**, 1534 (2024).

1229

1230 53. Coffman JA, Su Y-H. Redox regulation of development and regeneration.
1231   *Current opinion in genetics & development* **57**, 9–15 (2019).

1232

1233 54. Romero MMG, McCathie G, Jankun P, Roehl HH. Damage-induced
1234   reactive oxygen species enable zebrafish tail regeneration by
1235   repositioning of Hedgehog expressing cells. *Nature communications* **9**,
1236   4010 (2018).

1237

1238 55. Jaenen V*, et al.* Reactive oxygen species rescue regeneration after
1239   silencing the MAPK–ERK signaling pathway in Schmidtea mediterranea.
1240   *Scientific reports* **11**, 881 (2021).

1241

1242 56. Carbonell M B, Zapata Cardona J, Delgado JP. Hydrogen peroxide is
1243   necessary during tail regeneration in juvenile axolotl. *Developmental*
1244   *dynamics* **251**, 1054–1076 (2022).

1245

1246 57. Carbonell-M B, Zapata Cardona J, Delgado JP. Post-amputation reactive
1247   oxygen species production is necessary for axolotls limb regeneration.
1248   *Frontiers in Cell and Developmental Biology* **10**, 921520 (2022).

1249

1250 58. Shin SY, Park S-J, Kim H-S, Jeon J-H, Lee H-J. Wound-induced signals
1251   regulate root organogenesis in Arabidopsis explants. *BMC Plant Biology*
1252   **22**, 133 (2022).

1253

1254 59. Faltin Z*, et al.* Glutathione peroxidase regulation of reactive oxygen
1255   species level is crucial for in vitro plant differentiation. *Plant and cell*
1256   *physiology* **51**, 1151–1162 (2010).

1257

1258 60. Zhang W, Rong F, Ke W. Effects of inter-culture, arabinogalactan proteins,
1259   and hydrogen peroxide on the plant regeneration of wheat immature
1260   embryos. *Journal of Integrative Agriculture* **14**, 11–19 (2015).

1261

1262 61. Szechyńska-Hebda M, Skrzypek E, Dąbrowska G, Wędzony M, van
1263   Lammeren A. The effect of endogenous hydrogen peroxide induced by
1264   cold treatment in the improvement of tissue regeneration efficiency. *Acta*
1265   *Physiologiae Plantarum* **34**, 547–560 (2012).

1266
1267   62.   Tian M, Gu Q, Zhu M. The involvement of hydrogen peroxide and
1268   antioxidant enzymes in the process of shoot organogenesis of strawberry
1269   callus. *Plant Science* **165**, 701–707 (2003).

1270
1271   63.   Kiełkowska A, Adamus A. Exogenously applied polyamines reduce
1272   reactive oxygen species, enhancing cell division and the shoot
1273   regeneration from Brassica oleracea L. var. capitata protoplasts.
1274   *Agronomy* **11**, 735 (2021).

1275
1276   64.   de Marco A, Roubelakis-Angelakis KA. Hydrogen peroxide plays a
1277   bivalent role in the regeneration of protoplasts. *Journal of plant physiology*
1278   **149**, 109–114 (1996).

1279
1280   65.   Petřivalský M*, et al.* The effects of reactive nitrogen and oxygen species
1281   on the regeneration and growth of cucumber cells from isolated
1282   protoplasts. *Plant Cell, Tissue and Organ Culture (PCTOC)* **108**, 237–249
1283   (2012).

1284
1285   66.   Zhang H*, et al.* Thioredoxin-mediated ROS homeostasis explains natural
1286   variation in plant regeneration. *Plant Physiology* **176**, 2231–2250 (2018).

1287
1288   67.   Trovato M, Forlani G, Signorelli S, Funck D. Proline metabolism and its
1289   functions in development and stress tolerance. In: *Osmoprotectant-*
1290   *mediated abiotic stress tolerance in plants: recent advances and future*
1291   *perspectives*). Springer (2019).

1292
1293   68.   Schertl P, Cabassa C, Saadallah K, Bordenave M, Savouré A, Braun HP.
1294   Biochemical characterization of proline dehydrogenase in Arabidopsis
1295   mitochondria. *The FEBS Journal* **281**, 2794–2804 (2014).

1296
1297   69.   Miller G, Honig A, Stein H, Suzuki N, Mittler R, Zilberstein A. Unraveling
1298   Δ1-pyrroline-5-carboxylate-proline cycle in plants by uncoupled expression
1299   of proline oxidation enzymes. *Journal of Biological Chemistry* **284**, 26482–
1300   26492 (2009).

1301
1302   70.   Cecchini NM, Monteoliva MI, Alvarez ME. Proline dehydrogenase
1303   contributes to pathogen defense in Arabidopsis. *Plant Physiology* **155**,
1304   1947–1959 (2011).

1305

1306 71. Mackill DJ, Khush GS. IR64: a high-quality and high-yielding mega variety.
1307 *Rice* **11**, 18 (2018).

1308
1309 72. Sahoo RK, Tuteja N. Development of Agrobacterium-mediated
1310 transformation technology for mature seed-derived callus tissues of indica
1311 rice cultivar IR64. *GM crops & food* **3**, 123–128 (2012).

1312
1313 73. Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-
1314 mediated transformation of Arabidopsis thaliana. *Plant J* **16**, 735–743
1315 (1998).

1316
1317 74. Chen X, Qu Y, Sheng L, Liu J, Huang H, Xu L. A simple method suitable to
1318 study de novo root organogenesis. *Frontiers in Plant Science* **5**, 208
1319 (2014).

1320
1321 75. Hu L, Qi P, Peper A, Kong F, Yao Y, Yang L. Distinct function of SPL genes
1322 in age-related resistance in Arabidopsis. *PLOS Pathogens* **19**, e1011218
1323 (2023).

1324
1325 76. Robinson MD, Oshlack A. A scaling normalization method for differential
1326 expression analysis of RNA-seq data. *Genome biology* **11**, R25 (2010).

1327
1328 77. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical
1329 and powerful approach to multiple testing. *Journal of the Royal statistical
1330 society: series B (Methodological)* **57**, 289–300 (1995).

1331
1332 78. Yi X, Du Z, Su Z. PlantGSEA: a gene set enrichment analysis toolkit for
1333 plant community. *Nucleic acids research* **41**, W98–W103 (2013).

1334
1335 79. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and
1336 correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–
1337 2849 (2016).

1338
1339 80. Nour-Eldin HH, Geu-Flores F, Halkier BA. USER cloning and USER fusion:
1340 the ideal cloning techniques for small and big laboratories. In: *Plant
1341 secondary metabolism engineering: methods and applications*). Springer
1342 (2010).

1343
1344 81. Nour-Eldin HH, Hansen BG, Nørholm MH, Jensen JK, Halkier BA.
1345 Advancing uracil-excision based cloning towards an ideal technique for
1346 cloning PCR fragments. *Nucleic acids research* **34**, e122–e122 (2006).

1347

1348   82.   Joo JH, Wang S, Chen JG, Jones AM, Fedoroff NV. Different signaling and
1349   cell death roles of heterotrimeric G protein alpha and beta subunits in the
1350   Arabidopsis oxidative stress response to ozone. *Plant Cell* **17**, 957–970
1351   (2005).

1352
1353   83.   Abramson J*, et al.* Accurate structure prediction of biomolecular
1354   interactions with AlphaFold 3. *Nature* **630**, 493–500 (2024).

1355
1356   84.   Meng EC*, et al.* UCSF ChimeraX: Tools for structure building and analysis.
1357   *Protein Science* **32**, e4792 (2023).

1358

1359

1360