INS-17, an insulin-receptor ligand, confers specificity

in adult IIS-regulated phenotypes in *C. elegans*

Emily J. Leptich[1,2], Priyadharshini Vijayakumar[2], Edward W. Pietryk[2,3], Meredith I. Williams[2], Rachel N. Arey[2,4*]

[1] Department of Neuroscience, Baylor College of Medicine, Houston, Texas, United States of America

[2] Center for Precision Environmental Health, Baylor College of Medicine, Houston, Texas, United States of America

[3] Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas, United States of America

[4] Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, Texas, United States of America

*Corresponding author

E-mail: rachel.arey@bcm.edu

Abstract

Insulin/IGF growth factor (IGF) signaling (IIS) is a pleiotropic signaling pathway that functions across tissues to coordinate phenotypic changes in response to nutrient status. Thus, the ubiquity of the IIS pathway hinders efforts to elucidate the mechanisms driving specific IIS-related phenotypes, Previous research in the nematode worm *C. elegans* has demonstrated that loss of function of the IGF transmembrane receptor (IR) ortholog, DAF-2, results in a doubled lifespan and enhanced learning and memory behaviors in young and aged animals. However, these findings are the result of reducing DAF-2 receptor function rather than modulating ligand-receptor interactions. In the current study, we aimed to dissect ligand-receptor interactions that may regulate associative behaviors apart from canonical IIS lifespan phenotypes in *C. elegans*. To this end, we performed targeted genetic screening of Insulin-like Peptides (ILPs) previously identified as DAF-2 antagonists to test their role in learning and memory phenotypes. We discovered that only a single uncharacterized ILP, INS-17, is required for learning and memory. We also demonstrate that INS-17 is sufficient to confer extended memory ability and can promote the maintenance of learning and memory with age. Additionally, we observe that *ins-17* regulates learning and memory ability independent of lifespan, uncoupling IIS-mutant phenotypes. We find that regulation of the *ins-17* genetic locus explains its unique requirement among ILP for learning and memory behaviors. Finally, we found that INS-17 may act to signal a state of nutrient deprivation that is required to properly process stimulus valence to promote advantageous behaviors. Our findings deepen the understanding of how IIS can regulate specific phenotypic outputs in response to changes in internal metabolic states.

Introduction

The survival of most multicellular organisms depends on the coordination of complex physiological processes between multiple, functionally discrete tissues. Remarkably, each of these tissues carries out a diverse set of physiological functions using a highly overlapping set of cell regulatory mechanisms, including ubiquitous signaling pathways (e.g. MAPK/ERK, cAMP, and ubiquitin-proteasome pathway signaling[1–4]). Major components of these pathways are commonly pleiotropic, where a single gene can control multiple, distinct phenotypes depending on the context, such as life stage, cell type, and metabolic state. Across species, one prominent pleiotropic signaling pathway is the insulin/insulin-like growth factor (IGF) receptor signaling (IIS), which regulates a wealth of phenotypes across multiple tissues, including growth and development, metabolism, fertility, aging, stress response, and survival [5–15]. Dissecting the specific signaling mechanisms by which IIS mediates such a wide range of phenotypic effects proves challenging, especially in mammalian systems. This is due to several factors, namely high costs to generate tools to examine pathways in a tissue-specific and temporally controlled manner; compensation between receptors, as a single tissue can regulate phenotypes organism-wide; and the time and resource burden to study certain IIS-regulated phenotypes, such as aging. However, invertebrate systems can help overcome these challenges, as IIS is highly conserved across species [9,10,14,16,17], and simple models allow for rapid, low-cost study of genes with remarkable temporal and tissue-specific precision [18–20].

The nematode *C. elegans* has been invaluable in elucidating the pleiotropy underlying

IIS phenotypes. In fact, foundational experiments in the worm were the first to link

insulin signaling and aging, where loss-of-function receptor mutants of the IR homolog,

daf-2, have a doubled lifespan of around 40 days compared to the 20-day lifespan of

wild-type animals (21,22). Since this initial work, studies in IIS mutants revealed the

pleiotropy of the DAF-2 receptor and its conserved downstream signaling components

(*age-1*/PI3K and transcription factors *hsf-1*/HSF1, *skn-1*/NRF2, and *daf-16*/forkhead

family (FOXO3)) in this relatively simple model (21–28), including that *daf-2* mutants

have altered development, metabolism, fertility, motility, cognitive abilities, and more

(12,26,29–39). Collectively, research indicates that DAF-*2*/IR functions across tissues

and is integral for sensing internal state along with environmental cues allowing for the

worm to adapt appropriately to improve survival.

Despite decades of study of IIS in the worm, how the DAF-2 receptor coordinates a

variety of phenotypic responses across tissues remains unclear. This is due in part to

the ubiquity of the IIS pathway across tissues, where DAF-2/IR activity in some tissues

can mask DAF-2/IR activity in other tissues. For example, it is well established that *daf-*

2 mutants have improved motility with age (30,40,41), but recent work using degron-

based approaches found that reduction of *daf-2* function in neurons and muscles has

opposing effects on maintenance of motility in aged animals (42). This work sheds light

on the remaining gap in our knowledge regarding how signaling through a single

ubiquitous receptor can impact numerous phenotypes that are often reflective of the

function of a relatively small number of cells. It is also unclear whether these distinct

phenotypes can be uncoupled, and if so by what mechanisms.

Though the DAF-2 receptor and its downstream effectors have been well-studied, less

is known about the upstream regulators of DAF-2/IR that drive specific phenotypic

outputs. The *C. elegans* genome encodes 40 insulin-like peptides (ILPs) predicted to

act as DAF-2 receptor agonists or antagonists, sometimes in a context-dependent

fashion (43–47). However, no single manipulation of an ILP fully reproduces reduced

daf-2 function phenotypes. ILPs also exhibit diverse and often non-overlapping

expression patterns and are regulated by distinct physiological contexts (48). Thus, we

hypothesize that pleiotropic phenotypes exhibited by IIS mutants may also reflect

distinct ligand-receptor interactions.

In the present study, we investigated if specific ILPs regulate *daf-2* mutant learning and

memory phenotypes. We discovered that a single ILP, the DAF-2/IR antagonist INS-17,

is required for learning and memory ability and is sufficient to promote memory. We also

found that INS-17 is specifically required for IIS regulation of behavior and does not

appear to be involved in other adult DAF-2 receptor-mediated phenotypes, including

lifespan (47,49). We find that the regulation of INS-17 is the source of this phenotypic

specificity, and that INS-17 acts as a nutrient-deprivation signal that regulates plastic

behavior in the context of changing physiological states.

Results

INS-17 is specifically required for learning and memory

ability

Decreased DAF-2 receptor function results in a 3-fold extension in short- and

intermediate associative memory (S/ITAM), as measured by positive olfactory

association assays (31). In this paradigm, animals form a positive association with the

neutral odor butanone (10% in EtOH) when paired with food, which is measured by a

training-dependent increase in butanone chemotaxis (50). A single food-butanone (1

CS-US) pairing usually results in a molecularly conserved memory that lasts

approximately 2 hours; however, young adult *daf-2* mutants maintain a food-butanone

association for up to 6 hours (Fig. 1A) (31). Percent maximum (% Max) performance

indices are shown for clarity (see methods for calculations), as *daf-2* animals have high

naïve chemotaxis to 10% butanone in multiple studies (31,34,36,37) (Fig. S1A) that

contribute to perceived learning impairments when graphed with raw performance

indices (provided in Fig. S1B). Given that partial loss-of-function *daf-2* animals display

this significant extension in S/ITAM performance, we hypothesized that insulin receptor

antagonist(s), which are predicted to reduce signaling through DAF-2/IR, may promote

learning and memory behavior (Fig. 1B). To test this hypothesis, we examined the

learning and memory ability of publicly available insulin receptor antagonist mutants

using this STAM paradigm. We tested learning and intermediate-term associative

memory (ITAM, 60 minutes post-training), as behavioral deficits are typically more

prominent at this timepoint compared to STAM (30 minutes post-training). Mutants

tested included predicted strong (*ins -17, -37, and -39)* and weak (*ins-15, -21,* and *-22)*

antagonists (47,51,52). However, we recently determined that mutants for *ins-22* have

an abnormally high preference for 10% butanone (Fig. S1I) (53). Thus, we excluded *ins-*

22 from our targeted behavioral screen, as this high naïve chemotaxis would make it

difficult to measure training-dependent increases in butanone preference. Most

antagonist mutants behaved in a manner that was indistinguishable from wild-type

animals, including baseline naïve chemotaxis (Fig. 1D-E, S1E-H); while mutant animals

for *ins-39* had significantly higher intermediate-term associative memory (ITAM)

performance compared to wild-type animals, we were screening for behavioral deficits

and focused on candidates that had impaired behavioral performance. Interestingly,

only loss-of-function mutants of *ins-17* exhibited behavioral deficits: these include

significant deficits in learning, STAM and ITAM performance (Fig 1C). These deficits

were not accompanied by impaired sensory function, as measured by naïve chemotaxis

assays to the highly attractive concentration (0.1%) of butanone (Fig. S1C-D),

suggesting that they are specific to learning and memory. INS-17 is a relatively

uncharacterized ILP; aside from its prediction as a strong antagonist, previous research

has established *ins-17*'s role in regulating dauer formation in early development (49). To

our knowledge, INS-17 is uncharacterized regarding complex behavior, and these

results (Fig. 1C-E) suggest that it is a novel learning and memory-regulating ILP.

145 **INS-17 signals through DAF-2 and promotes memory**

We next wanted to confirm that INS-17 signaling was indeed mediated by the DAF-2

receptor and tested the learning and memory behavior of *daf-2(e1370);ins-17(tm790)*

double mutants. We examined double mutant memory behavior at 3 hours after 1 CS-

US pairing, a timepoint where *daf-2* mutants exhibit extended memory (31). *daf-*

2(e1370);ins-17(tm790) worms demonstrated extended 3-hour memory

indistinguishable from *daf-2* animals (Fig. 2A, S2A-C). Thus, we infer from these results

that DAF-2/IR is epistatic to INS-17 in the regulation of learning and memory.

Next, we wanted to explore how manipulating levels of INS-17 may influence behavioral

outputs. Our behavior paradigm tests an adult-specific behavior (34), and *ins-17*

regulates dauer in development (49), so we sought to avoid confounding our results

with developmental effects of traditional transgenic approaches. Recently, we developed

a novel microbial feeding paradigm to manipulate neuropeptide signaling specifically in

adult animals (53). In these experiments, *C. elegans* are fed *E. coli* bacteria expressing

a plasmid containing the sequence encoding a neuropeptide of interest flanked by

processing enzyme cut sites (Fig. 2B). Like feeding-based RNAi approaches (54), this

method offers precise temporal control of increasing neuropeptide levels via adult-only

manipulation to avoid potential effects on early development. We have demonstrated

that this feeding method could work in an 'overexpression' manner when administered

to wild-type animals, altering sensory behavior (53). Here, we wanted to extend this

approach to assess if administering additional INS-17 to adult animals could affect

associative behaviors. We hypothesized that increasing the levels of a memory-

regulating *daf-2* antagonist would result in enhanced memory ability. Briefly, we fed L4

animals *E. coli* expressing either INS-17 or a scrambled control peptide (SCR) and

examined their behavior at Day 2 of adulthood.

To first establish the ability of this method to effectively manipulate INS-17 signaling, we

tested if INS-17 peptide feeding was sufficient to rescue learning and memory behavior

of *ins-17(tm790)* animals. SCR fed mutants still exhibited behavioral deficits, while

adult-only feeding of INS-17 to *ins-17* deficient mutants was sufficient to rescue their

behavior, similar to wild-type animals fed SCR peptide (Fig. 2C).   Notably, we

determined these results were not the result of sensory deficits in these animals (Fig.

S2D-E). We next explored if INS-17 peptide feeding was sufficient to boost wild-type

animal memory behavior past the 2-hour timepoint after 1 CS-US pairing. Remarkably,

we found that wild-type animals fed INS-17 during adulthood displayed 3-hour memory

post-training, while SCR fed animals demonstrated no memory phenotype, as expected

(Fig. 2D). We confirmed that peptide feeding to both wild-type and mutant animals did

not alter baseline chemosensation, indicating the effect is memory-specific (Fig. S2D-F).

These results suggest that increasing INS-17 is sufficient to boost cognitive

performance and phenocopy *daf-2* animals' 3-hour memory behavior.

Finally, we worked to identify the tissue where INS-17 is required to regulate memory.

Since INS-17 is highly expressed in the nervous-system, we hypothesized that neurons

may use INS-17 as a memory signal. Using a neuron-specific RNAi-sensitive line(55),

we performed an adult-only *ins-17* knockdown and observed significant S/IATM

behavioral deficits in *ins-17 RNAi* knock-down animals (Fig. 2E)—similar to what we

observed in deletion mutants (Fig. 1B)—with no detectable effects on sensory ability

(Fig. S2H-I). This demonstrates that INS-17 is a memory-promoting DAF-2/IR

antagonist, likely released from neurons.

INS-17 regulates the cognitive aging phenotypes mediated by

the DAF-2 receptor independent of lifespan

In addition to exhibiting extended STAM at young adult stages, *daf-2* mutants also

maintain the ability to learn and remember after 1 CS-US pairing with age (31). Wild-

type animals display age-related cognitive deficits as early as Days 3 and 5 of

adulthood (Fig 3A-B, S3A-C), while *daf-2* mutants maintain performance akin to Day 1

of adulthood at these timepoints. Therefore, we wanted to ask if INS-17 could also be

involved in these age-related phenotypes associated with altered *daf-2* function. As *ins-*

17 mutants already display severe behavioral defects at Day 1 of adulthood, we

examined the effect of increased INS-17 on age-related behavioral impairments in wild-

type animals, again using a peptide feeding-based approach (53).

We first assessed the behavior of animals supplemented continuously with INS-17 from

the L4 stage until Day 5 of adulthood (Fig. 3C), when worms are considered cognitively

aged (31). INS-17 fed animals demonstrated significantly better learning and memory

performance compared to wild-type SCR fed controls (Fig. 3C-D, S3D). Next, we

wondered if manipulating the DAF-2/IR pathway via INS-17 later in life could influence

cognitive aging, as previous work has shown that degradation of DAF-2/IR via an auxin

inducible degron system in older animals results in improved STAM (37). We subjected

Day 3 adult wild-type animals (which already display age-related cognitive deficits

compared to Day 1 Adults (Fig. 3A)) to SCR or INS-17 peptide feeding and tested their

learning and memory abilities at Day 5 of adulthood (Fig. 3E). Notably, we found that

Day 5 INS-17 fed animals had better learning and STAM behaviors compared to SCR

fed controls (Fig. 3E-F, S3E-F). These findings suggest that INS-17 is a DAF-2/IR ligand

that could partially contribute to the enhanced cognitive ability with age observed when

DAF-2/IR activity levels are reduced.

Since *daf-2* animals have both a doubled lifespan and enhanced memory phenotypes in

young and aged animals (21,31), we wanted to determine if manipulating INS-17*,* as a

strong DAF-2/IR antagonist, would alter lifespan in addition to associative behaviors. As

previously reported in the literature (49), we confirmed that *ins-17(tm790)* animals have

no significant difference in lifespan to wild-type animals (Fig. 3G, S3F-G). We also

tested the lifespans of wild-type animals fed SCR compared to those fed INS-17 in

adulthood, finding no significant differences between groups (Fig. 3H, S3H-I). Thus, our

findings demonstrate that either loss of *ins-17* or increased levels of INS-17 has no

effect on adult lifespan, highlighting the specific phenotypic regulation by individual

ligands underlying the pleiotropy of the DAF-2 receptor.

INS-17 engages multiple pathways downstream of DAF-2 to

regulate learning and memory

We next sought to identify the pathways downstream of DAF-2/IR that could explain the

phenotypic specificity of INS-17. The canonical downstream target of DAF-2/IR

signaling is the FOXO transcription factor homolog DAF-16(27,28), which is required for

the doubled lifespan extension of *daf-2* loss-of-function worms (26,28,56). Additionally,

daf-16 mutants exhibit severely impaired learning and memory phenotypes (Fig. 4A,

S4A (31). Since *ins-17* is required for positive associative memory behavior and there

are no adult lifespan phenotypes associated with *ins-17* (Fig 3G-H), we hypothesized

that *ins-17* activates a pathway that acts in parallel to *daf-16*. To test this, we examined

if INS-17 feeding could improve the memory impairments of *daf-16* worms. We found

that adult-only INS-17 feeding rescued the learning and memory of *daf-16(mu86)*

animals to SCR fed wild-type performance levels, while SCR fed *daf-16(mu86)* worms

still displayed behavioral deficits (Fig. 4B, S4B-C). These results support that INS-17

could act independently of DAF-16 to regulate associative behaviors.

Interestingly, recent research has uncovered evidence of pathways that can regulate

cognitive abilities downstream of DAF-2/IR in parallel to DAF-16. This includes the

discovery of a novel role for the insulin receptor substrate *ist-1*, which has been shown

to function partially in parallel to DAF-16 to regulate learning ability in an aversive

olfactory training paradigm (57). Thus, we aimed to determine if *ins-17* is functioning in

the same pathway as *ist-1*. First, we examined if *ist-1* is required in our positive

associative memory paradigm. We found that *ist-1(ok2706)* animals have impaired

positive associative STAM behavior compared to wild-type animals (Fig. 4C). Like *daf-*

16 animals, we performed adult-only INS-17 feeding with *ist-1(ok2706)* worms and

found that INS-17 peptide feeding was sufficient to boost STAM performance similar to

wild-type animals fed SCR, while *ist-1* animals fed SCR still displayed deficits (Fig. 4D).

Furthermore, we confirmed *ist-1* deficits were not the result of altered baseline

chemosensation of butanone nor peptide feeding (Fig S4D-G).

While IST-1 and DAF-16 can function in parallel, it is possible that these two pathways

can compensate for one another. Indeed, Cheng et al., (2022) identified that while *ist-1*

could regulate learning independently of *daf-16*, *daf-16* was still required for this

behavior(57). We hypothesized that INS-17 could similarly engage both *daf-16* and *ist-1*

to promote memory ability, and that partial regulation of either arm of the DAF-2

receptor signaling pathway could act as a compensatory mechanism in the mutant

backgrounds when fed INS-17. Therefore, we examined if the beneficial effects of INS-

17 administration were abolished when both genes were mutated. We generated *daf-*

16(mu86);ist-1(ok2706) double mutants and observed that they had significant learning

and memory deficits (Fig. 4E-F) compared to either single mutant alone. Because of the

striking nature of these deficits, we confirmed that double mutant animals still

maintained chemosensory ability and determined that they exhibited normal preference

for chemoattractants (Fig. SH-I). We found that INS-17 peptide administration failed to

rescue these severe behavioral defects, in contrast to the improvement observed in

single mutants (Fig. 4E-F, S4J-K). These results suggest that INS-17 feeding can

activate DAF-16 and IST-1, partially in parallel, to regulate associative behaviors.

Regulation of the *ins-17* genetic locus is required for proper

regulation of behavior

While we had identified how INS-17 might regulate distinct DAF-2/IR mediated

phenotypes, we also sought to examine what made INS-17 unique relative to other

antagonists in regard to regulating behavior. Interestingly, though the *C. elegans*

genome encodes 40 ILPs, there are only three structurally similar antagonists known as

γ-insulins; these genes are encoded across multiple genomic regions (ch. II and III) and

thus likely under the control of distinct regulatory elements (43,47,57). We therefore

hypothesized that phenotypic specificity was not necessarily due to insulin class or

structure, but instead via *ins-17*'s regulation by its surrounding genomic elements, which

likely respond to different signaling pathways and environmental cues. To test this

hypothesis, we explored if replacing *ins-17* with another strong DAF-2/IR antagonist

was sufficient to maintain normal positive associative memory ability in an *ins-17* mutant

background.

We selected *ins-37* to swap in for *ins-17* due to a key overlap in structure, as both ILPs

are predicted strong DAF-2/IR antagonists and are γ-insulins, which are defined by

three di-sulfide bonds (47). Moreover, *ins-37* mutants had no observable learning and

memory impairments (Fig. 1D, S1H) and *ins-37* is lowly expressed in neurons (58),

making it an ideal strong antagonist to test our hypothesis. Thus, we generated an *ins-*

17 CRISPR knockout in which we placed the *ins-37* coding sequence at the *ins-17*

locus (Fig. 5A). Subsequently, we examined if *ins-37* expressed at endogenous *ins-17*

levels is sufficient for normal learning and memory behavior. First, we confirmed that the

newly generated *ins-17* knockout line exhibited learning and memory impairments

similar to the predicted loss-of-function mutants (Fig 1C, 5B, S5A). Interestingly, we

found that replacing *ins-17* with *ins-37* could rescue the defective learning and memory

of *ins-17* KO animals, functionally replacing endogenous *ins-17* (Fig. 5B, S5A). These

results indicate that ILP structure is not as critical as its regulation of expression. These

results suggest that the genomic context in which *ins-17* is regulated underlies its

importance in promoting specific behavioral phenotypes.

ins-17 expression rapidly responds to nutrient deprivation

Next, we wanted to uncover the physiological conditions that regulate *ins-17* expression

and assess their role in learning and memory. We mined publicly available gene

expression datasets and discovered that *ins-17* transcription is likely regulated by

metabolic state, as it is consistently upregulated in multiple starvation paradigms and

genetic models of nutrient deprivation (S. Table 1) (49,51,59–62). Interestingly, our

appetitive behavioral paradigm involves a 1-hour food deprivation (FD) step prior to the

1 CS-US (food-butanone) pairing, which has been reported to promote a more robust

CS-US association compared to animals that do not undergo a FD period prior to

memory training (31,63). We first examined if the FD step incorporated into our

paradigm was sufficient to increase *ins-1*7. We performed qPCR analysis to measure

ins-17 levels in wild-type worms after 1 hour of FD and found that even this short time

period was sufficient to observe a significant increase in *ins-17* expression (Fig. 5C,

left). We also observed more robust upregulation of *ins-17* following 2 hours of FD (Fig.

S5B), suggesting that this locus rapidly responds to changes in nutrient availability.

INS-17 is required for properly sensing internal nutrient status to allow for processing of stimulus valence

Based on our gene expression results, we hypothesized that INS-17 could promote

memory performance by functioning as a FD signal. We first assessed the role of insulin

signaling in FD-associated improvements in associative behavior (Fig. 5C, right). We

confirmed that FD could indeed improve memory ability by comparing the behavioral

performance of wild-type animals that underwent FD prior to training to animals that

were not food-deprived (non-FD) prior to training. Similar to previous reports (31), we

found that wild-type FD animals have significantly better ITAM performance, which is

measured one hour post-training, compared to non-FD animals (Fig. 5D, S5C-D).

Interestingly, *daf-2* mutant phenotypes are strongly associated with phenotypes

paralleling those of chronically starved worms (64–67), so we hypothesized that *daf-2*

animals may not require the FD step prior to 1 CS-US pairing for robust ITAM behavior,

as they are thought to have a perceived nutrient status of chronic starvation even when

well-fed. We found that non-FD *daf-2* animals that underwent 1 CS-US pairing had

ITAM behavior nearly identical to *daf-2* animals that were FD prior to training (Fig. 5E,

S5E). Altogether, these results indicate that *daf-2* animals' improved memory phenotype

could be driven by their sensation of a chronically starved internal state.

We next tested if INS-17 could function as a FD signal. If this was the case, we

hypothesized that increased levels of INS-17 could be sufficient to promote robust ITAM

behavior in the absence of the FD step prior to 1 CS-US pairing. To test this, we

examined if wild-type animals fed INS-17 peptide required FD for enhanced ITAM. We

compared memory performance of SCR or INS-17 fed animals after training in the

absence of FD, and found that, similar to *daf-2* mutants, INS-17 fed animals exhibited

significantly more robust ITAM than SCR fed animals in the non-FD condition (Fig. 5F).

Furthermore, FD had no detectable effect on ITAM performance in INS-17 fed animals,

but significantly improved ITAM in SCR fed animals (Fig. S5F-I). These results suggest

that INS-17 acts as a FD signal to promote a stronger association between a food and

an odorant.

We next asked if INS-17 functions generally as a FD signal to promote associative

behaviors using an aversive olfactory training paradigm, where an attractive odorant

diacetyl is paired with the absence of food to form a negative association (68). If INS-17

functions as a signal of FD internal state, we hypothesized that *ins-17* mutants would

have impaired aversive learning and memory compared to wild-type animals, because

sensing starvation is required for a negative diacetyl association to be formed. Wild-type

and *ins-17* loss-of-function mutants underwent well-established aversive memory

assays using a diacetyl-starvation pairing (68). Strikingly, we found that *ins-17* is

required for normal aversive associative behaviors, as *ins-17* animals had severe

deficits in learning, STAM, and ITAM behavior compared to wild-type animals (Fig. 5G,

S5J). The aversive and appetitive olfactory conditioning paradigms we tested here

recruit distinct neuronal circuits (63,69–75). Therefore, our results suggest that *ins-17*

may be broadly required in the nervous system to signal a FD state so that the valence

of olfactory stimuli (positive when paired with food, negative when paired with

starvation) can be processed appropriately to promote advantageous behaviors (Fig.

5H). This could explain how IIS acts in the nervous system to integrate metabolic status

and regulate specific behavioral outputs, such as learning and memory.

Discussion

Across species, insulin signaling is pleiotropic. In the worm, the DAF-2 receptor

regulates a wide array of phenotypes, including those related to development, lifespan,

metabolism, fecundity, cognitive behaviors, and motility, among others. How does a

single receptor mediate such physiologically distinct phenotypes? Here we report that

some of these pleiotropic actions result from specific ligand-receptor interactions. In this

instance, we show that an olfactory memory phenotype is likely mediated by INS-17, an

antagonistic ILP with no previously reported role in behavior (to our knowledge). DAF-2

receptor partial loss-of-function mutants have a well-established extended memory

phenotype (31), so it is not surprising that a predicted antagonistic ILP regulated these

behaviors. However, we were surprised that only a single strong antagonist appeared to

be required in our paradigms. The importance of INS-17 to the extended memory

phenotype is supported by our finding that overexpression through feeding is sufficient

to recapitulate the memory phenotype of *daf-2* mutants. While we established that other

antagonists are not required for this specific behavioral phenotype, many ILPs are

diverse and act as agonists or antagonists in a context-dependent fashion, and most do

not have a described role in associative behaviors. Future studies systemically

characterizing these diverse ligands will be informative for defining DAF-2 receptor

pleiotropy across tissues.

Interestingly, several transcriptomic datasets show that *ins-17* transcripts are increased

roughly 2-fold in *daf-2* mutants and decreased in *daf-16* mutants compared to wild-type

animals (Table 1) (34,49,51,59,60,62). This indicates that INS-17 is highly regulated by

IIS signaling and is potentially acting both upstream and down-stream of DAF-2/IR. "ILP

to ILP" signaling, where the action of one ILP affects the expression of another ILP,

could be a component of *ins-17*'s role in integrating sensory cues and nutrient

availability to elicit behavioral outputs. One example of "ILP to ILP" signaling in behavior

is between INS-6 and INS-7 in aversive pathogen learning, where INS-6 released by the

ASI neuron regulates *ins-7* transcription in the URX neurons to regulate this behavior

(76). Further transcriptional analysis of *ins-17* mutants and INS-17 overexpression

models could provide insight into which ILPs may be regulating or be regulated by INS-

17.

We also report that INS-17's function in adulthood is specific to behavior, as

manipulating INS-17 does not affect lifespan (49). While *daf-2* animals' extended

lifespan is a canonical phenotype, there may exist alternate downstream IIS pathways

that are dispensable for lifespan regulation. There are reports that the sole insulin

receptor substrate homolog IST-1 may differentially activate distinct arms of the IIS

signaling pathway, as it can function upstream of—or partially in parallel to—DAF-16

and is not required for normal growth and development (77–79). Thus, it is possible the

40 known ILPs may similarly engage the IIS pathway. In the present study, we

investigated IST-1, which engages atypical IIS signaling to regulate aversive olfactory

learning (79). Here, we report that *ist-1* mutants display positive olfactory memory

deficits that were rescued by INS-17 peptide feeding; taken with our similar results in

daf-16 mutants, these findings suggest INS-17 may activate multiple arms of the IIS

pathway to regulate adult cognitive behaviors. We find that *daf-16;ist-1* double mutants

do not respond to INS-17 peptide supplementation, suggesting that, similar to other

olfactory behavioral paradigms(79), INS-17 engages multiple pathways that can act

partially in parallel to one another downstream of DAF-2 to regulate behavior. It will be

intriguing in the future to examine if other DAF-2-mediated associative behaviors

similarly engage both downstream pathways.

We also find that the genomic context in which an ILP is regulated provides specificity,

as "swapping" a structurally similar insulin, *ins-37* (47), into *the ins-17* genetic locus is

sufficient to functionally replace *ins-17* in behavior, though endogenous *ins-37* is not

required for learning phenotypes in our paradigm. This result supports that tissue-

specific gene regulation may inform DAF-2/IR pleiotropy, warranting future studies

exploring tissue-specific requirements for DAF-2/IR function.

Furthermore, we demonstrate that *ins-17* is required for both positive and aversive

associative behaviors. Both associative behavioral paradigms include a relatively short,

1-hour FD step prior to training to prime animals to form a stronger association between

food and an odorant, thus promoting a more robust behavioral response after training.

This FD step occurs within the timeframe before animals display significant changes in

sensory integration behavior due to a "lack of food sensation", which begins after 3

hours of food-deprivation (80); however, food is cleared from the alimentary system

within 2 minutes of ingestion (81,82), suggesting the need for a FD signal to inform

robust behavioral responses after only 1 hour of FD. Our gene expression data, similar

to previous studies (51,59,60), shows that the *ins-17* gene is activated in this timeframe,

suggesting that could it be involved in responding to these brief FD signals. This notion

is supported by our findings that feeding INS-17 to animals that do not undergo FD

results in normal learning and memory, bypassing the need for FD to promote memory.

Together, these results suggest that INS-17 could be released in response to acute and

prolonged FD to inform sensory integration behavior. Interestingly, the only phenotype

previously associated with INS-17 is dauer formation to protect progeny from

unfavorable conditions (49), namely starvation (83). We speculate that INS-17 may

respond to FD to coordinate phenotypic responses in different life stages that are most

advantageous; for example, while *ins-17* regulates dauer formation, which is a response

to lack of food availability and other stresses during development, it may also regulate

the ability to form associations in the context of food availability in adulthood, which is

beneficial for egg-laying and progeny survival (64,84). In the future, it will be interesting

to further examine how both the *ins-17* locus and INS-17 secretion are regulated in the

context of changing nutrient availability.

Since *ins-17* is required for both positive and aversive associative behaviors, our results

support that INS-17 may function as a broad signal for nutrient deprivation to the

nervous system. *ins-17* is widely expressed across the nervous system (58,61), but is

most highly expressed in the AIZ interneuron, which mediates a variety of behavioral

outputs in response to multiple sensory neurons and circuits (85–87). We observed that

associative behavior impairments resulting from loss of ins-17 do not stem from

baseline sensory deficits, which strongly indicates that rather sensory information may

not be integrated correctly or that sensory neurons cannot alter their responses to an

environmental stimulus in the context of altered metabolic status. We have established

that INS-17 may be essential for interpreting internal state and proper valence

processing. In mammals, neuropeptides also are important for signaling valence in

positive and aversive associative paradigms (88,89), and this is likely a theme of

peptides modulating circuits more broadly. Future studies examining how INS-17

regulates sensory circuit activity in various contexts will help further define *daf-2*

pleiotropic mechanisms.

Our results bring an attractive model for the expansion of the ILP family in *C. elegans*.

ILPs are encoded across the genome and function under the control of a variety of

regulatory elements so the animal can dynamically respond to a combination of

nutrients and environmental stimuli via DAF-2/IR to facilitate the most advantageous

phenotype. Having a diverse array of peptides that can carry out distinct outputs is

beneficial, especially if there is only one receptor to mediate such outputs. In the case of

INS-17, being able to properly make associations in the face of changing environments

can promote survival, such as remembering availability (or lack) of a food source

associated with specific odors. Together, our findings have identified INS-17 as a

physiological signal that is required for positive and aversive associative behaviors and

may integrate environmental cues and nutrient status to inform valence processing and

regulate behavioral outcomes. This research underscores the importance of future work

investigating the rich and diverse network of DAF-2/IR signaling.

Methods

General *C. elegans* maintenance

Worms were maintained at 20 °C on plates containing either 1) Standard nematode growth medium (NGM): 3 g/L NaCl, 17 g/L agar, 2.5 g/L peptone, 25 mL/L KPO4 (pH 6.0), 1 mL/L MgSO4, 1 mL/L CaCl2, and 1 mL/L cholesterol (5 mg/mL in ethanol) (90); or 2) High growth medium (HGM): 3 g/L NaCl, 30 g/L agar, 20 g/L peptone, 25 mL/L KPO4 (pH 6.0), 1 mL/L MgSO4, 1 mL/L CaCl2, and 4 mL/L cholesterol (5 mg/mL in ethanol) (50). Plates were seeded with an *E. coli* OP50 lawn for ad libitum feeding (90).

Animals were developmentally synchronized using an alkaline-bleach solution (1.5 mL 5 N NaOH, 3.0 mL 5% sodium hypochlorite, 5.5 mL water) to collect eggs from gravid adults; eggs were subsequently washed three times with M9 buffer solution (6 g/L Na2HPO4, 3 g/L KH2PO4, 5 g/L NaCl and 1 mL/L 1M MgSO4 in ultrapure water) before plating (90).

Strains

Wild-type: N2 Bristol

Mutants: *ins-17(tm790), ins-21(tm5180)*, *ins-37(tm6892)*, and *ins-39(tm6520)* were obtained from the National BioResource Project, Japan.

RB2489 (*ins-15(ok3444)*), RB2594 (*ins-22(ok3616)*), CB1370 (*daf-2(e1370)*), CF1038 (*daf-16(mu86)*), MAH677 (*sid-1(qt9) V; sqIs71[rgef-2p::GFP + rgef-1p::sid-1]*) were obtained from the Caenorhabditis Genetics Center.

CX17790 (*ist-1(ok2706)* was described previously(79) and generously gifted to us by

Dr. Cornelia Bargmann.

Strains made in collaboration with InVivo Biosystems include *ins-17* CRISPR KO

animals with *ins-37* replacement COP2848 (*ins-17(knu1330[ins-37]) III*) and *ins-17*

CRISPR KO animals COP2835 (*ins-17(knu1320) III*).

The following strains were generated by crosses: RNA34 (*daf-2(e1370)*; *ins-17(tm790))*

and RNA59 (*daf-16(mu86); ist-1(ok2706)*).

Peptide plasmid generation and transformation

The amino acid coding sequence for each peptide was identified using

www.wormbase.org (57). These sequences were codon-optimized for *C. elegans* and

cloned in pET-21a(+) cloning vector via XhoI/XhoI strategy by GenScript. The scramble

control sequence and plasmid were generated using the same cloning strategy by

GenScript. Plasmids arrived lyophilized and were re-suspended in nuclease-free H_2O to

a concentration of 100ng/µl and stored at -20 °C. Each expression clone was

transformed into competent iOP50 cells using methods previously described(91).

Briefly, iOP50 stocks were streaked out onto LB plates containing 50 µg/mL tetracycline

and grown at 37 °C overnight. Then, single colonies of iOP50 were isolated and used to

inoculate 2.0 mL LB containing 50 µg/ml tetracycline and placed in a 37 °C shaker

overnight. 0.5 mL of each culture was used to inoculate 3.0 mL of LB without antibiotics

and grown for 2 hours in a 37 °C shaker. Each culture was then aliquoted in 0.75 mL

increments into four 1.5 mL tubes and cooled for 10 minutes on ice. Next, the cells were

harvested by centrifugation at 6,000 rpm for 5 minutes at 4 °C. The resulting

supernatant was discarded, and cells were placed back on ice and re-suspended in 1.0

mL 100 mM ice-cold CaCl$_2$ solution. Cells were kept on ice for 20 minutes and then

harvested by centrifugation as in the previous step. After discarding the supernatant,

cells were re-suspended in 150µl of ice-cold CaCl$_2$ solution and placed on ice to be

immediately used for transformation.

Competent cells were transformed using the New England Biolabs protocol

(C2987H/C2987I). Single colonies of transformed cells were isolated and cultured in 3.0

mL of LB containing 50 µg/mL tetracycline and 50 µg/mL carbenicillin overnight in a 37

°C shaker. Cultures were frozen down as glycerol stocks and stored at -80 °C.

Successful transformation was verified by isolating plasmids from cultures using a

Zyppy Plasmid Miniprep Kit and sending samples for whole-plasmid sequencing

(Plasmidsaurus).

Peptide plate preparation

LB containing 50 µg/mL tetracycline and 50 µg/mL carbenicillin was inoculated with

iOP50 encoding either the peptide sequence of interest or scramble peptide sequence

and cultured for 18-20 hours in a 37 °C shaker. 100 mm NGM plates containing 50

µg/mL tetracycline, 50 µg/mL carbenicillin, and 1.0 mL/L isopropyl-β-D-thiogalactoside

(IPTG) were plated with 1.0 mL of corresponding culture. Seeded plates were left to dry

at least 24 hours at room temperature and never kept longer than 72 hours before use.

Approximately one hour before transferring worms to peptide or scramble control plates,

200 µL of 0.1 M IPTG was added to each plate and left to dry.

Peptide feeding

We adapted a previously published peptide feeding-based approach (53). Worms were

maintained on NGM agar plates seeded with OP50 *E. coli* and maintained at 20 °C until

the start of experiments. All animals were transferred to a conical tube and washed

three times with M9 buffer solution before being transferred to NGM plates containing

50 μg/mL Tetracycline, 50 μg/mL Carbenicillin, and 1.0 mM IPTG that were seeded with

1.0 mL of iOP50 expressing either the scramble peptide or the peptide of interest.

Animals were kept on the scramble or peptide-expressing lawns for at least 48 hours of

exposure time at 20 °C before performing assays. Exact peptide feeding timeline

parameters are expounded in the results section.

RNAi treatment

Worms were maintained on NGM agar plates seeded with OP50 *E. coli* at 20 °C until

the start of experiments. At the L4 stage, animals were transferred to a conical tube and

washed three times with M9 buffer solution before being transferred to NGM plates

containing 50 μg/mL Carbenicillin and 1.0 mM IPTG that were seeded with 1.0 mL of

HT115 bacteria expressing either RNAi against the gene of interest or the vector control

RNAi. Animals were kept on the RNAi lawns for 48 hours of exposure time at 20 °C

before performing assays at Day 2 of adulthood.

Naïve chemotaxis assay

Chemotaxis assays were performed using a protocol based on previously published

assays(92). Briefly, assays were performed on unseeded 100mm NGM plates. On the

back of each plate, two marks were made on opposite sides of the plate, approximately

5mm from the edge. 1µL of sodium azide (Thermo Fisher) was placed on both spots

and allowed to dry before adding 1µL of odorant (Sigma Aldrich) diluted in ethanol on

one mark and 1µL ethanol on the other. Using M9 buffer, worms were washed off their

plates and subsequently washed three times to eliminate any leftover bacteria that

could impact baseline behavior. Then, worms were placed near the bottom center of the

plate, equidistant between the two marks, and allowed to chemotax for an hour.

Chemotaxis indices were calculated as follows:

Chemotaxis Index = (# of worms butanone - # of worms ethanol) / (total # of worms)

For naïve chemotaxis assays performed to assess baseline chemosensory function, we

tested the naïve preference to a highly appetitive concentration of butanone (0.1%) as

previously reported (92).

Positive associative butanone learning and memory assay

Worms were trained and tested for learning, short- and intermediate-term memory

changes as previously described (50). Briefly, synchronized adult worms were washed

off plates with M9 buffer and allowed to settle by gravity, followed by two subsequent

washes with M9 buffer to remove leftover bacteria that could alter baseline behavior.

Next, the worms were food-deprived for one hour in M9 buffer before exposure to 1 CS-

US pairing, where food-deprived worms were transferred to 100mm NGM conditioning

plates seeded with OP50 *E. coli* bacteria (or iOP50 bacteria containing a peptide or

scramble control plasmid for peptide feeding experiments) and had 16µL of 10%

butanone (Sigma Aldrich) diluted in ethanol streaked on the lid in a '#' shape for one

hour. After conditioning, the trained population of worms were tested for chemotaxis to

10% butanone and to an ethanol control using standard, previously described

chemotaxis conditions(92). Different stages of memory were tested by measuring

chemotaxis indices of separate populations of worms at timepoints including 0 min.

(learning), 30 min. (STAM), 60 min. (ITAM) and 120 min. (forgetting) after training. For

timepoints after learning behavior, worms were transferred to hold plates seeded with

OP50 *E. coli* or iOP50 bacteria for peptide feeding experiments as stated above.

Chemotaxis indices for each timepoint were calculated as detailed in "Naïve chemotaxis

assays" section.

Performance index is the change in the chemotaxis index after training relative to the

naïve (untrained) chemotaxis index:

Performance index = Chemotaxis index$_{trained}$—Chemotaxis index$_{untrained}$

% Maximum Performance Indices are performance indices represented as a

percentage of the maximum value that can be performed relative to the naïve

chemotaxis index(31,36). This % can be more informative for visualizing performance

indices when naïve chemotaxis behavior is high, as in *daf-2(e1370)* animals. This is

calculated for each timepoint by dividing the performance index of a timepoint by the

average of the untrained chemotaxis indices of the same behavioral replicate subtracted

from 1:

% Maximum Performance Index = Performance index$_{timepoint}$/(1 - Chemotaxis

index$_{untrained}$)

Butanone enhancement assay

596   This assay was adapted from previously described methods (63). Unlike the positive

597   butanone associative memory assays, butanone enhancement assays skip the 1-hour

598   food-deprivation step prior to the conditioning step. All other methods are identical to

599   those listed in the positive butanone associative memory assay section.

Negative associative learning and memory assay

Worms were tested for learning and memory changes in response to aversive diacetyl

training, adapted from previously published methods (68). Similar to positive associative

assays, synchronized Day 1 adult worms were washed off plates with M9 buffer and

then allowed to settle by gravity and washed twice more with M9 buffer. After washing,

the worms were conditioned for one hour by food-deprivation on 100mm NGM plates

with no bacteria and 16uL of 100% diacetyl (Sigma Aldrich) streaked on the lid in a '#'

shape. The trained population of worms were then tested for chemotaxis to 1% diacetyl

and to an ethanol control using standard, previously described chemotaxis conditions

(92). Learning and memory performance indices were calculated for each timepoint as

previously stated in the positive butanone associative assay section.

Lifespan assay

Lifespan assays were performed on *ins-17(tm790)* animals and wild-type animals that

were bleached onto NGM plates seeded with OP50 *E. coli* and maintained at 20 °C; at

the L4 larval stage, worms were picked for survival analysis and transferred every other

day to freshly seeded plates. Similarly, lifespan assays with peptide feeding

administration were performed on wild-type animals that were bleached onto NGM

plates seeded with OP50 *E. coli* and maintained at 20 °C. Then, at the L4 larval stage,

worms were picked for survival analysis and transferred every other day to NGM plates

containing 50 µg/mL tetracycline, 50 µg/mL carbenicillin, and 1.0 mL/L IPTG that were

seeded with either INS-17 iOP50 or scramble iOP50 *E. coli.* Lifespans began (t = 0) at

Day 1 of adulthood and had an n = ~120 worms per strain/trial for a total of three

biological replicates per assay.

Food-deprivation behavioral experiments

For food-deprivation experiments, well-fed, synchronized adult worms were washed off

plates using M9 buffer into a 15mL conical and washed two more times to remove

bacteria that could alter baseline behavior. Next, worms were either 1) food-deprived for

one hour on a 100mm NGM plate without food or 2) non-food deprived for one hour by

being transferred to a 100mm plate seeded with a bacterial lawn (iOP50 bacteria for

peptide feeding experiments). Worms were then immediately exposed to the

conditioning step (1 CS-US pairing) and stages of learning and memory were tested as

stated in the positive associative butanone learning and memory assays section.

RNA isolation, cDNA synthesis and qRT-PCR

For qRT-PCR food-deprivation experiments, well-fed, synchronized adult worms were

washed off plates using M9 buffer into a 15mL conical and washed two more times to

remove bacteria that could alter baseline behavior. Next, worms were either 1) food-

deprived for one hour or two hours after being transferred to a 100mm plate with no

bacterial lawn or 2) non-food deprived for one hour or two hours after being transferred

to a 100mm plate seeded with a bacterial lawn.

After feeding or food deprivation, worms were crushed in liquid nitrogen using a mortar

and pestle and added to Trizol (Thermo Fisher Scientific). RNA was isolated per

manufacturer's instructions, followed by DNase treatment (Qiagen). cDNA was

synthesized with an oligo dT primer and Superscript III reverse transcriptase enzyme

(Thermo Fisher Scientific). cDNA was mixed with buffers, primers, SYBR green, and hot

start Taq polymerase in a master mix prepared by a manufacturer (Thermo Fisher

Scientific). PCR reactions were run using a Quant Studio 7 Pro Dx Real-Time PCR

System (Thermo Fisher Scientific), followed by a dissociation reaction to determine

specificity of the amplified product. Gene expression was quantified using the ΔΔCt

method using *pmp-3* as a reference gene. Primer sets were as follows:

ins-17 For: 5'- GGACACTATCCGACCACCAC -3':

ins-17 Rev: 5'- TTCACATCCCGTCTCACAGC -3':

pmp-3 For: 5'- AGTTCCGGTTGGATTGGTCC -3'

pmp-3 Rev: 5'- CCAGCACGATAGAAGGCGAT-3'

Statistical analysis and software

All statistics (excluding survival analysis) were calculated with GraphPad Prism

Software. All statistical data are reported in the main text, figures, and tables.

Significance threshold of $p < 0.05$ was used. The symbols *, **, ***, and **** refer to $p <$

0.05, 0.01, 0.001, and 0.0001. The symbols #, ##, ###, and #### refer to $p < 0.05, 0.01,$

0.001, and 0.0001. For comparing performance indices between two behavior

conditions (e.g. *vector control* vs *ins-17 RNAi*), we used a Mann-Whitney test comparing

ranks, as it does not assume normality. For comparing performance indices between

three or more groups (e.g. wild-type vs multiple insulin receptor antagonist mutants), we

performed one-way analysis of variances followed by Bonferroni post hoc tests for

multiple comparisons. Two-way ANOVAs were used for evaluating effects between

genotype and timepoint (T0, T30, T60, T120) on performance indices with a significant

interaction between factors ($p < 0.0001$) that prompted Bonferroni post-hoc analyses to

determine differences between individual groups. Sample size n represents the number

of chemotaxis assays performed for behavioral experiments, with each assay containing

approximately 50–150 worms. All experiments were performed on different days with

distinct populations to confirm reproducibility. For qPCR analysis, unpaired t-tests with

Welch's corrections were performed to determine significance between conditions.

Survival analysis statistics were generated using the log-rank (Mantel-Cox) method to

test the null hypothesis in Kaplan-Meier survival analysis and evaluated using OASIS

survival analysis software (93).

Funding

675   RNA was supported by a Glenn Foundation for Medical Research and AFAR Grant for

676   Junior Faculty, the Whitehall Foundation, and an NIH Director's New Innovator Award

677   (DP2NS132372). EWP is supported by a NIGMS NRSA, EJL is funded by a NIA NRSA.

678   Some strains were provided by the *C. elegans* Genetics Center (CGC), which is funded

679   by the NIH Office of Research Infrastructure Programs (P40 OD010440).

Acknowledgements

We thank the Bargmann lab for generously sharing the *ist-1(ok2706)* strain, and the

Caenorhabditis Genetics Center for most other strains used in this publication. We also

thank Ben Jussila and In Vivo Biosystems for strains and helpful discussion.

Additionally, we especially thank Drs. Buck Samuel and Shinya Yamamoto at Baylor

College of Medicine for helpful discussion and advice at various stages of the project.

Finally, we thank Arey lab members Katie L. Brandel-Ankrapp and Ty M. Gadberry for

feedback on the manuscript.

References

1. Cooper GM. Pathways of Intracellular Signal Transduction. The Cell: A
Molecular Approach [Internet]. 2nd ed. Sinauer Associates; 2000 [cited 2025 Sep
29]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK9870/

2. Li Y, Li S, Wu H. Ubiquitination-Proteasome System (UPS) and Autophagy Two
Main Protein Degradation Machineries in Response to Cell Stress. Cells. 2022
Mar 1;11(5).

3. Buscà R, Pouysségur J, Lenormand P. ERK1 and ERK2 map kinases: specific
roles or functional redundancy? Front Cell Dev Biol. 2016 Jun 8;4:53.

4. Lefkimmiatis K, Zaccolo M. cAMP signaling in subcellular compartments.
Pharmacol Ther. 2014 Sep;143(3):295–304.

5. Ullrich A, Bell JR, Chen EY, Herrera R, Petruzzelli LM, Dull TJ, et al. Human
insulin receptor and its relationship to the tyrosine kinase family of oncogenes.
Nature. 1985;313(6005):756–61.

6. Ullrich A, Gray A, Tam AW, Yang-Feng T, Tsubokawa M, Collins C, et al.
Insulin-like growth factor I receptor primary structure: comparison with insulin
receptor suggests structural determinants that define functional specificity. EMBO
J. 1986 Oct;5(10):2503–12.

7. White MF. The IRS-signalling system: a network of docking proteins that
mediate insulin action. Mol Cell Biochem. 1998 May;182(1–2):3–11.

8.   Taniguchi CM, Emanuelli B, Kahn CR. Critical nodes in signalling pathways:
709   insights into insulin action. Nat Rev Mol Cell Biol. 2006 Feb;7(2):85–96.

710   9.   Piñero González J, Carrillo Farnés O, Vasconcelos ATR, González Pérez A.
711   Conservation of key members in the course of the evolution of the insulin
712   signaling pathway. BioSystems. 2009 Jan;95(1):7–16.

713   10.   Kaletsky R, Murphy CT. The role of insulin/IGF-like signaling in C. elegans
714   longevity and aging. Dis Model Mech. 2010 Aug 1;3(7–8):415–9.

715   11.   Finch CE, Ruvkun G. The genetics of aging. Annu Rev Genomics Hum Genet.
716   2001;2:435–62.

717   12.   Murphy CT, Hu PJ. Insulin/insulin-like growth factor signaling in C. elegans.
718   WormBook. 2013 Dec 26;1–43.

719   13.   Semaniuk U, Piskovatska V, Strilbytska O, Strutynska T, Burdyliuk N, Vaiserman
720   A, et al. *Drosophila* insulin□like peptides: from expression to functions – a review.
721   Entomol Exp Appl. 2021 Feb;169(2):195–208.

722   14.   Biglou SG, Bendena WG, Chin-Sang I. An overview of the insulin signaling
723   pathway in model organisms Drosophila melanogaster and Caenorhabditis
724   elegans. Peptides. 2021 Nov;145:170640.

725   15.   Suzawa M, Bland ML. Insulin signaling in development. Development. 2023 Oct
726   15;150(20).

727   16.   Barbieri M, Bonafè M, Franceschi C, Paolisso G. Insulin/IGF-I-signaling
728   pathway: an evolutionarily conserved mechanism of longevity from yeast to
729   humans. Am J Physiol Endocrinol Metab. 2003 Nov;285(5):E1064-71.

730   17.   Lau HE, Chalasani SH. Divergent and convergent roles for insulin-like peptides
731   in the worm, fly and mammalian nervous systems. Invert Neurosci. 2014
732   Sep;14(2):71–8.

733   18.   Hales KG, Korey CA, Larracuente AM, Roberts DM. Genetics on the fly: A
734   primer on the drosophila model system. Genetics. 2015 Nov;201(3):815–42.

735   19.   Boulin T, Hobert O. From genes to function: the C. elegans genetic toolbox.
736   Wiley Interdiscip Rev Dev Biol. 2012 Feb;1(1):114–37.

737   20.   Simone BW, Martínez-Gálvez G, WareJoncas Z, Ekker SC. Fishing for
738   understanding: Unlocking the zebrafish gene editor's toolbox. Methods. 2018 Nov
739   1;150:3–10.

740   21.   Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A C. elegans mutant that
741   lives twice as long as wild type. Nature. 1993 Dec 2;366(6454):461–4.

742   22.   Friedman DB, Johnson TE. A mutation in the age-1 gene in Caenorhabditis
743   elegans lengthens life and reduces hermaphrodite fertility. Genetics. 1988
744   Jan;118(1):75–86.

745   23.   Hsu A-L, Murphy CT, Kenyon C. Regulation of aging and age-related disease by
746   DAF-16 and heat-shock factor. Science. 2003 May 16;300(5622):1142–5.

747   24.   An JH, Blackwell TK. SKN-1 links C. elegans mesendodermal specification to a
748   conserved oxidative stress response. Genes Dev. 2003 Aug 1;17(15):1882–93.

749   25.   Morris JZ, Tissenbaum HA, Ruvkun G. A phosphatidylinositol-3-OH kinase
750   family member regulating longevity and diapause in Caenorhabditis elegans.
751   Nature. 1996 Aug 8;382(6591):536–9.

752   26.   Kimura KD, Tissenbaum HA, Liu Y, Ruvkun G. daf-2, an insulin receptor-like
753   gene that regulates longevity and diapause in Caenorhabditis elegans. Science.
754   1997 Aug 15;277(5328):942–6.

755   27.   Lin K, Dorman JB, Rodan A, Kenyon C. daf-16: An HNF-3/forkhead family
756   member that can function to double the life-span of Caenorhabditis elegans.
757   Science. 1997 Nov 14;278(5341):1319–22.

758   28.   Ogg S, Paradis S, Gottlieb S, Patterson GI, Lee L, Tissenbaum HA, et al. The
759   Fork head transcription factor DAF-16 transduces insulin-like metabolic and
760   longevity signals in C. elegans. Nature. 1997 Oct 30;389(6654):994–9.

761   29.   Luo S, Kleemann GA, Ashraf JM, Shaw WM, Murphy CT. TGF-β and insulin
762   signaling regulate reproductive aging via oocyte and germline quality
763   maintenance. Cell. 2010 Oct 15;143(2):299–312.

764   30.   Mulcahy B, Holden-Dye L, O'Connor V. Pharmacological assays reveal age-
765   related changes in synaptic transmission at the Caenorhabditis elegans
766   neuromuscular junction that are modified by reduced insulin signalling. J Exp
767   Biol. 2013 Feb 1;216(Pt 3):492–501.

768   31.   Kauffman AL, Ashraf JM, Corces-Zimmerman MR, Landis JN, Murphy CT.
769   Insulin signaling and dietary restriction differentially influence the decline of
770   learning and memory with age. PLoS Biol. 2010 May 18;8(5):e1000372.

771   32.   Riddle DL, Swanson MM, Albert PS. Interacting genes in nematode dauer larva
772   formation. Nature. 1981 Apr 23;290(5808):668–71.

33. Gems D, Sutton AJ, Sundermeyer ML, Albert PS, King KV, Edgley ML, et al.
Two pleiotropic classes of daf-2 mutation affect larval arrest, adult behavior,
reproduction and longevity in Caenorhabditis elegans. Genetics. 1998
Sep;150(1):129–55.

34. Ange JS, Weng Y, Stevenson ME, Kaletsky R, Moore RS, Zhou S, et al. Adult
Single-nucleus Neuronal Transcriptomes of Insulin Signaling Mutants Reveal
Regulators of Behavior and Learning. BioRxiv. 2024 Feb 7;

35. Templeman NM, Luo S, Kaletsky R, Shi C, Ashraf J, Keyes W, et al. Insulin
Signaling Regulates Oocyte Quality Maintenance with Age via Cathepsin B
Activity. Curr Biol. 2018 Mar 5;28(5):753-760.e4.

36. Weng Y, Zhou S, Morillo K, Kaletsky R, Lin S, Murphy CT. The neuron-specific
IIS/FOXO transcriptome in aged animals reveals regulatory mechanisms of
cognitive aging. eLife. 2024 Jun 26;13.

37. Zhou S, Novak KE, Kaletsky R, Weng Y, Ange JS, Stevenson ME, et al. Body-
to-brain insulin and Notch signaling regulates memory through neuronal CREB
activity. Nat Aging. 2025 Jul;5(7):1232–48.

38. Hughes SE, Evason K, Xiong C, Kornfeld K. Genetic and pharmacological
factors that influence reproductive aging in nematodes. PLoS Genet. 2007 Feb
16;3(2):e25.

39. Lin K, Hsin H, Libina N, Kenyon C. Regulation of the Caenorhabditis elegans
longevity protein DAF-16 by insulin/IGF-1 and germline signaling. Nat Genet.
2001 Jun;28(2):139–45.

40. Hahm J-H, Kim S, DiLoreto R, Shi C, Lee S-JV, Murphy CT, et al. C. elegans
maximum velocity correlates with healthspan and is maintained in worms with an
insulin receptor mutation. Nat Commun. 2015 Nov 20;6:8919.

41. Bansal A, Zhu LJ, Yen K, Tissenbaum HA. Uncoupling lifespan and healthspan
in Caenorhabditis elegans longevity mutants. Proc Natl Acad Sci USA. 2015 Jan
20;112(3):E277-86.

42. Roy C, Molin L, Alcolei A, Solyga M, Bonneau B, Vachon C, et al. DAF-2/insulin
IGF-1 receptor regulates motility during aging by integrating opposite signaling
from muscle and neuronal tissues. Aging Cell. 2022 Aug;21(8):e13660.

43. Duret L, Guex N, Peitsch MC, Bairoch A. New insulin-like proteins with atypical
disulfide bond pattern characterized in Caenorhabditis elegans by comparative
sequence analysis and homology modeling. Genome Res. 1998 Apr;8(4):348–
53.

44.   Kawano T, Ito Y, Ishiguro M, Takuwa K, Nakajima T, Kimura Y. Molecular
809   cloning and characterization of a new insulin/IGF-like peptide of the nematode
810   Caenorhabditis elegans. Biochem Biophys Res Commun. 2000 Jul 5;273(2):431–
811   6.

812   45.   Pierce SB, Costa M, Wisotzkey R, Devadhar S, Homburger SA, Buchman AR, et
813   al. Regulation of DAF-2 receptor signaling by human insulin and ins-1, a member
814   of the unusually large and diverse C. elegans insulin gene family. Genes Dev.
815   2001 Mar 15;15(6):672–86.

816   46.   Li W, Kennedy SG, Ruvkun G. daf-28 encodes a C. elegans insulin superfamily
817   member that is regulated by environmental cues and acts in the DAF-2 signaling
818   pathway. Genes Dev. 2003 Apr 1;17(7):844–58.

819   47.   Zheng S, Chiu H, Boudreau J, Papanicolaou T, Bendena W, Chin-Sang I. A
820   functional study of all 40 Caenorhabditis elegans insulin-like peptides. J Biol
821   Chem. 2018 Oct 26;293(43):16912–22.

822   48.   Ritter AD, Shen Y, Fuxman Bass J, Jeyaraj S, Deplancke B, Mukhopadhyay A,
823   et al. Complex expression dynamics and robustness in C. elegans insulin
824   networks. Genome Res. 2013 Jun;23(6):954–65.

825   49.   Matsunaga Y, Nakajima K, Gengyo-Ando K, Mitani S, Iwasaki T, Kawano T. A
826   Caenorhabditis elegans insulin-like peptide, INS-17: its physiological function and
827   expression pattern. Biosci Biotechnol Biochem. 2012 Nov 7;76(11):2168–72.

828   50.   Kauffman A, Parsons L, Stein G, Wills A, Kaletsky R, Murphy C. C. elegans
829   positive butanone learning, short-term, and long-term associative memory
830   assays. J Vis Exp. 2011 Mar 11;(49).

831   51.   Chen Y, Baugh LR. Ins-4 and daf-28 function redundantly to regulate C. elegans
832   L1 arrest. Dev Biol. 2014 Oct 15;394(2):314–26.

833   52.   Zhu R, Chin-Sang ID. C. elegans insulin-like peptides. Mol Cell Endocrinol. 2024
834   May 1;585:112173.

835   53.   DiLoreto EM, Shastry S, Leptich EJ, Reilly DK, Arey RN, Srinivasan J.
836   Harnessing Microbial Tools: Escherichia coli as a Vehicle for Neuropeptide
837   Functional Analysis in Caenorhabditis elegans. Genetics. 2025 Aug 6;

838   54.   Kamath RS, Ahringer J. Genome-wide RNAi screening in Caenorhabditis
839   elegans. Methods. 2003 Aug;30(4):313–21.

840   55.   Yang Y, Arnold ML, Lange CM, Sun L-H, Broussalian M, Doroodian S, et al.
841   Autophagy protein ATG-16.2 and its WD40 domain mediate the beneficial effects

of inhibiting early-acting autophagy genes in C. elegans neurons. Nat Aging.
2024 Feb;4(2):198–212.

56.   Murphy CT, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, et al.
Genes that act downstream of DAF-16 to influence the lifespan of Caenorhabditis
elegans. Nature. 2003 Jul 17;424(6946):277–83.

57.   Sternberg PW, Van Auken K, Wang Q, Wright A, Yook K, Zarowiecki M, et al.
WormBase 2024: status and transitioning to Alliance infrastructure. Genetics.
2024 May 7;227(1).

58.   Taylor SR, Santpere G, Weinreb A, Barrett A, Reilly MB, Xu C, et al. Molecular
topography of an entire nervous system. Cell. 2021 Aug 5;184(16):4329-
4347.e23.

59.   Mueller MM, Castells-Roca L, Babu V, Ermolaeva MA, Müller R-U, Frommolt P,
et al. DAF-16/FOXO and EGL-27/GATA promote developmental growth in
response to persistent somatic DNA damage. Nat Cell Biol. 2014
Dec;16(12):1168–79.

60.   Hibshman JD, Doan AE, Moore BT, Kaplan RE, Hung A, Webster AK, et al. daf-
16/FoxO promotes gluconeogenesis and trehalose synthesis during starvation to
support survival. eLife. 2017 Oct 24;6.

61.   St Ange J, Weng Y, Kaletsky R, Stevenson ME, Moore RS, Zhou S, et al. Adult
single-nucleus neuronal transcriptomes of insulin signaling mutants reveal
regulators of behavior and learning. Cell Genomics. 2024 Dec 11;4(12):100720.

62.   Ogawa T, Isik M, Wu Z, Kurmi K, Meng J, Cho S, et al. Nutrient control of growth
and metabolism through mTORC1 regulation of mRNA splicing. Mol Cell. 2024
Dec 5;84(23):4558-4575.e8.

63.   Torayama I, Ishihara T, Katsura I. Caenorhabditis elegans integrates the signals
of butanone and food to enhance chemotaxis to butanone. J Neurosci. 2007 Jan
24;27(4):741–50.

64.   Baugh LR, Hu PJ. Starvation responses throughout the caenorhabditiselegans
life cycle. Genetics. 2020 Dec;216(4):837–78.

65.   Ben Arous J, Laffont S, Chatenay D. Molecular and sensory basis of a food
related two-state behavior in C. elegans. PLoS ONE. 2009 Oct 23;4(10):e7584.

66.   Kimura KD, Riddle DL, Ruvkun G. The C. elegans DAF-2 insulin-like receptor is
abundantly expressed in the nervous system and regulated by nutritional status.
Cold Spring Harb Symp Quant Biol. 2011 Nov 28;76:113–20.

67.   Venz R, Pekec T, Katic I, Ciosk R, Ewald CY. End-of-life targeted degradation of
877   DAF-2 insulin/IGF-1 receptor promotes longevity free from growth-related
878   pathologies. eLife. 2021 Sep 10;10.

879   68.   Stetak A, Hörndli F, Maricq AV, van den Heuvel S, Hajnal A. Neuron-specific
880   regulation of associative learning and memory by MAGI-1 in C. elegans. PLoS
881   ONE. 2009 Jun 24;4(6):e6019.

882   69.   Fadda M, De Fruyt N, Borghgraef C, Watteyne J, Peymen K, Vandewyer E, et
883   al. NPY/NPF-Related Neuropeptide FLP-34 Signals from Serotonergic Neurons
884   to Modulate Aversive Olfactory Learning in Caenorhabditis elegans. J Neurosci.
885   2020 Jul 29;40(31):6018–34.

886   70.   Choi JI, Lee HK, Kim HS, Park SY, Lee TY, Yoon K-H, et al. Odor-dependent
887   temporal dynamics in Caenorhabitis elegans adaptation and aversive learning
888   behavior. PeerJ. 2018 Jun 12;6:e4956.

889   71.   Amano H, Maruyama IN. Aversive olfactory learning and associative long-term
890   memory in Caenorhabditis elegans. Learn Mem. 2011 Oct;18(10):654–65.

891   72.   Chen Z, Hendricks M, Cornils A, Maier W, Alcedo J, Zhang Y. Two insulin-like
892   peptides antagonistically regulate aversive olfactory learning in C. elegans.
893   Neuron. 2013 Feb 6;77(3):572–85.

894   73.   Ha H, Hendricks M, Shen Y, Gabel CV, Fang-Yen C, Qin Y, et al. Functional
895   organization of a neural network for aversive olfactory learning in Caenorhabditis
896   elegans. Neuron. 2010 Dec 22;68(6):1173–86.

897   74.   O'Halloran DM, Altshuler-Keylin S, Lee JI, L'Etoile ND. Regulators of AWC-
898   mediated olfactory plasticity in Caenorhabditis elegans. PLoS Genet. 2009 Dec
899   11;5(12):e1000761.

900   75.   Chalasani SH, Kato S, Albrecht DR, Nakagawa T, Abbott LF, Bargmann CI.
901   Neuropeptide feedback modifies odor-evoked dynamics in Caenorhabditis
902   elegans olfactory neurons. Nat Neurosci. 2010 May;13(5):615–21.

903   76.   Mendler M, Riedinger C, Schlotterer A, Volk N, Fleming T, Herzig S, et al.
904   Reduction in ins-7 gene expression in non-neuronal cells of high glucose
905   exposed Caenorhabditis elegans protects from reactive metabolites, preserves
906   neuronal structure and head motility, and prolongs lifespan. J Diabetes
907   Complicat. 2017 Feb;31(2):304–10.

908   77.   Chen J, Scheiner AR, Falsztyn IB, Baugh LR. ist-1/IRS1 affects L1 starvation
909   resistance in daf-16/FoxO- dependent and independent fashion. MicroPubl Biol.
910   2025 Jun 13;2025.

911   78.   Wolkow CA, Muñoz MJ, Riddle DL, Ruvkun G. Insulin receptor substrate and
912   p55 orthologous adaptor proteins function in the Caenorhabditis elegans daf-
913   2/insulin-like signaling pathway. J Biol Chem. 2002 Dec 20;277(51):49591–7.

914   79.   Cheng D, Lee JS, Brown M, Ebert MS, McGrath PT, Tomioka M, et al.
915   Insulin/IGF signaling regulates presynaptic glutamate release in aversive
916   olfactory learning. Cell Rep. 2022 Nov 22;41(8):111685.

917   80.   Matty MA, Lau HE, Haley JA, Singh A, Chakraborty A, Kono K, et al. Intestine-
918   to-neuronal signaling alters risk-taking behaviors in food-deprived Caenorhabditis
919   elegans. PLoS Genet. 2022 May 5;18(5):e1010178.

920   81.   Viri V, Arveiler M, Lehnert T, Gijs MAM. An In Vivo Microfluidic Study of Bacterial
921   Load Dynamics and Absorption in the C. elegans Intestine. Micromachines
922   (Basel). 2021 Jul 17;12(7).

923   82.   Ghafouri S, McGhee JD. Bacterial residence time in the intestine of
924   Caenorhabditis elegans. Nematol. 2007;9(1):87–91.

925   83.   Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental
926   variant of the nematode Caenorhabditis elegans. Dev Biol. 1975 Oct;46(2):326–
927   42.

928   84.   Weinshenker D, Garriga G, Thomas JH. Genetic and pharmacological analysis
929   of neurotransmitters controlling egg laying in C. elegans. J Neurosci. 1995
930   Oct;15(10):6975–85.

931   85.   Kuhara A, Mori I. Molecular physiology of the neural circuit for calcineurin-
932   dependent associative learning in Caenorhabditis elegans. J Neurosci. 2006 Sep
933   13;26(37):9355–64.

934   86.   Tsalik EL, Hobert O. Functional mapping of neurons that control locomotory
935   behavior in Caenorhabditis elegans. J Neurobiol. 2003 Aug;56(2):178–97.

936   87.   Iino Y, Yoshida K. Parallel use of two behavioral mechanisms for chemotaxis in
937   Caenorhabditis elegans. J Neurosci. 2009 Apr 29;29(17):5370–80.

938   88.   Li H, Namburi P, Olson JM, Borio M, Lemieux ME, Beyeler A, et al. Neurotensin
939   orchestrates valence assignment in the amygdala. Nature. 2022
940   Aug;608(7923):586–92.

941   89.   Kim D-I, Kang SJ, Jhang J, Jo YS, Park S, Ye M, et al. Encoding opposing
942   valences through frequency-dependent transmitter switching in single peptidergic
943   neurons. bioRxiv. 2024 Jan 1;

90. Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974
May;77(1):71–94.

91. Neve IAA, Sowa JN, Lin C-CJ, Sivaramakrishnan P, Herman C, Ye Y, et al.
Escherichia coli Metabolite Profiling Leads to the Development of an RNA
Interference Strain for Caenorhabditis elegans. G3 (Bethesda). 2020 Jan
7;10(1):189–98.

92. Bargmann CI, Hartwieg E, Horvitz HR. Odorant-selective genes and neurons
mediate olfaction in C. elegans. Cell. 1993 Aug 13;74(3):515–27.

93. Yang J-S, Nam H-J, Seo M, Han SK, Choi Y, Nam HG, et al. OASIS: online
application for the survival analysis of lifespan assays performed in aging
research. PLoS ONE. 2011 Aug 15;6(8):e23525.

Figure legends

Fig. 1. INS-17 is a novel and specific memory-regulating insulin receptor

antagonist

(A) Partial loss-of-function *daf-2(e1370)* mutants have extended STAM that lasts 6

hours compared to 2 hours for wild-type animals after 1 CS-US food-butanone pairing.

% Max Performance reported as *daf-2(e1370)* animals have a higher baseline naïve

attraction to neutral (10%) butanone concentration (see **Fig. S1A, B**). Mean ± SEM. n =

15 per genotype. **(B)** Diagram depicting the hypothesis that insulin receptor antagonists

may inhibit DAF-2 receptor function to promote learning and memory. **(C)** *ins-17*

deletion mutants have impaired learning, STAM, and ITAM compared to wild-type

worms. Mean ± SEM. n = 15 per genotype. ***$p < 0.001$, ****$p < 0.001$; ns, not

significant ($p > 0.05$). **(D)** Weak insulin antagonist mutants for *ins-15* and *ins-21* and **(E)**

strong insulin antagonist mutants for *ins-37* and *ins-39* display no memory impairments;

ins-39 mutants have better ITAM performance compared to wild-type. Mean $\pm$ SEM. n =

15 per genotype. ***p < 0.001.

Fig. 2. INS-17 and DAF-2 are epistatic and promote memory

(A) *daf-2* and *ins-17* share an epistatic relationship. 3hrs post-training (1 CS-US

pairing), *daf-2(e1370)* mutants and *daf-2(e1370);ins-17(tm790)* double mutants display

memory behavior while *ins-17(tm790)* animals display no memory behavior. Mean $\pm$

SEM. n = 15 per genotype. ****p < 0.0001. **(B)** Diagram of microbial feeding paradigm.

iOP50 bacteria were transformed with a plasmid containing either the peptide sequence

of interest or a scramble control sequence and then fed to worms before testing

behavioral effects. **(C)** INS-17 peptide feeding rescues *ins-17(tm790)* learning, STAM

and ITAM behaviors, while SCR fed mutants display significant S/ITAM impairments

compared to SCR fed wild-type animals. Mean $\pm$ SEM. n = 15 per feeding treatment. #

is significance relative to SCR fed wild-type animal performance. ###p < 0.001, ####p

< 0.0001; ns, not significant (p > 0.05). **(D)** Wild-type animals fed INS-17 peptides

display 3-hour memory behavior after 1 CS-US pairing, similar to *daf-2(e1370)*

extended STAM performance. Mean $\pm$ SEM. n = 15 per peptide treatment. ****p <

0.0001. **(E)** Adult-only, neuron-specific RNAi knock-down of *ins-17* leads to learning,

STAM, and ITAM impairments compared to vector RNAi treated animals. Mean $\pm$ SEM.

n = 15 per RNAi treatment. ****p < 0.0001.

Fig. 3. INS-17 regulates cognitive aging phenotypes observed in *daf-2* mutants

(A) Day 5 adult *daf-2(e1370)* mutants display better learning and **(B)** memory behavior

compared to wild-type worms. Mean $\pm$ SEM. n = 10 per genotype. *p < 0.05, *** p <

0.001. **(C)** Timeline of young adult peptide treatment. Wild-type L4 animals are fed SCR

or INS-17 iOP50 bacteria until Day 5 of adulthood when S/ITAM behavior is tested. INS-

17 peptide treated worms have increased learning and **(D)** ITAM performance

compared to SCR peptide treated animals. Mean ± SEM. n = 15 per peptide treatment.

*p < 0.05. **(E)** Timeline of aged adult peptide treatment. Wild-type Day 3 adult animals

are fed SCR or INS-17 iOP50 bacteria until Day 5 of adulthood when S/ITAM behavior

is tested. INS-17 peptide treated worms have increased learning performance, but not

(F) ITAM performance, compared to SCR peptide treated animals. Mean ± SEM. n = 15

per peptide treatment. *p < 0.05, ****p < 0.0001. **(G)** Probability of survival analysis

demonstrates there is no significant difference in lifespan comparing wild-type animals

and *ins-17(tm790)* animals. One replicate depicted is representative of three replicates

(see **Fig. S3F, S3G**). Mean/median ± SEM. n ≥ 100 per replicate. ns, not significant (p

> 0.05). **(H)** Probability of survival analysis for wild-type animals treated with SCR

peptide compared to wild-type animals fed INS-17 peptide shows no significant

difference in average lifespan. One replicate depicted is representative of three

replicates (see **Fig. S3H, S3I**). Mean/median ± SEM. n ≥ 100 per replicate. ns, not

significant (p > 0.05).

Fig. 4. INS-17 likely engages parallel pathways of DAF-2 receptor signaling

(A) *daf-16(mu86)* deletion mutants have impaired learning and memory behavior

compared to wild-type animals. Mean ± SEM. n = 15 per genotype. ****p < 0.0001. **(B)**

INS-17 peptide treatment rescues *daf-16(mu86)* S/ITAM impairments. Mean ± SEM. n =

15 per genotype and peptide treatment. Wild-type animals fed INS-17 performance

indices are in **Fig. S4C**. # is significance relative to wild-type. #p < 0.05, ##p < 0.01; ns,

not significant (p > 0.05). **(C)** *ist-1(ok2706)* deletion mutants display a slight but not

significant learning deficit and impaired S/ITAM behaviors compared to wild-type worms.

Mean ± SEM. n = 15 per genotype. *p < 0.05, **p < 0.01; ns, not significant (p > 0.05).

(D) INS-17 peptide treatment rescues *ist-1(ok2706)* learning and memory impairments.

Mean ± SEM. n = 15 per genotype and peptide treatment. # is significance relative to

wild-type. #p < 0.05, ##p < 0.01, ###p < 0.001, ****p< 0.0001; ns, not significant (p >

0.05). **(E)** *daf-16(mu86);ist-1(ok2706)* animals display significant learning and **(F)**

memory impairments that are not rescued by INS-17 peptide feeding, while *daf-*

16(mu86) and *ist-1(ok2706)* deficits are rescued by INS-17 peptide administration.

Mean ± SEM. n = 15 per genotype and peptide treatment *p < 0.05, **p < 0.01, ***p <

0.001; ns, not significant (p > 0.05).

Fig. 5. INS-17 is regulated by nutrient deprivation to promote advantageous

behaviors

(A) Diagram showing *ins-17* replacement experiment design. *ins-17* and *ins-37* encode

γ -ILPs. *ins-17* was removed with CRISPR and replaced with *ins-37* at the *ins-17* locus

so *ins-37* would be expressed at endogenous *ins-17* levels. **(B)** Replacing *ins-17* with

ins-37 has no significant effect on behavior compared to wild-type animals, while *ins-17*

CRISPR KO (*ins-17(knu1320)*) animals have impaired learning and S/ITAM. Mean ±

SEM. n = 15 per genotype. # is significance relative to wild-type. ####p < 0.0001; ns,

not significant (p > 0.05). **(C)** qRT-PCR of *ins-17* mRNA levels in Day 1 adult wild-type

animals show that one hour of food-deprivation (FD) results in increased *ins-17*

transcript levels (left). For levels measured after two hours of food-deprivation, please

see **Fig. S5B.** Schematic depicting food-deprivation experiment workflow (right): well-

fed adult worms are washed three times in M9 buffer solution to remove bacteria and

placed on 100mm agar plates either seeded with a bacterial lawn (non-FD) or no lawn

(FD) prior to training and behavior testing. Mean $\pm$ SEM. n = 5 per genotype. *p < 0.05.

(D) Wild-type animals display more robust ITAM behavior if a FD step is administered

prior to 1 CS-US pairing compared to training alone (non-FD conditions). Mean $\pm$ SEM.

n = 15 per condition. *p < 0.05. **(E)** Food-deprivation prior to training (1 CS-US pairing)

does not enhance *daf-2(e1370)* ITAM performance. Mean $\pm$ SEM. n = 15 per condition.

ns, not significant (p > 0.05). **(F)** INS-17 peptide treatment in wild-type animals is

sufficient to promote robust ITAM performance in non-FD conditions prior to training

compared to non-FD SCR fed animals. See **Fig. S5I** for food-deprived SCR and INS-17

peptide fed comparisons. Mean $\pm$ SEM. n = 15 per peptide treatment. **p < 0.01. **(G)**

ins-17(tm790) mutants have significantly impaired aversive associative learning and

STAM performance compared to wild-type worms. Absolute values of learning and

memory performance indices are reported. Mean $\pm$ SEM. n = 15 per genotype. ***p <

0.001, ****p < 0.0001; ns, not significant (p > 0.05). **(H)** Putative diagram of how INS-17

is upregulated in the nervous system to signal the valence of positive and aversive

environmental conditions to inform behavioral outputs advantageous for survival.

1054

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5