INS-17, an insulin-receptor ligand, confers specificity

in adult IIS-regulated phenotypes in *C. elegans*

Emily J. Leptich[1,2], Priyadharshini Vijayakumar[2], Edward W. Pietryk[2,3], Meredith I. Williams[2], Rachel N. Arey[2,4*]

[1] Department of Neuroscience, Baylor College of Medicine, Houston, Texas, United States of America

[2] Center for Precision Environmental Health, Baylor College of Medicine, Houston, Texas, United States of America

[3] Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas, United States of America

[4] Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, Texas, United States of America

*Corresponding author

E-mail: rachel.arey@bcm.edu

Abstract

Insulin/IGF growth factor (IGF) signaling (IIS) is a pleiotropic signaling pathway that functions across tissues to coordinate phenotypic changes in response to nutrient status. Thus, the ubiquity of the IIS pathway hinders efforts to elucidate the mechanisms driving specific IIS-related phenotypes, Previous research in the nematode worm *C. elegans* has demonstrated that loss of function of the IGF transmembrane receptor (IR) ortholog, DAF-2, results in a doubled lifespan and enhanced learning and memory behaviors in young and aged animals. However, these findings are the result of reducing DAF-2 receptor function rather than modulating ligand-receptor interactions. In the current study, we aimed to dissect ligand-receptor interactions that may regulate associative behaviors apart from canonical IIS lifespan phenotypes in *C. elegans*. To this end, we performed targeted genetic screening of Insulin-like Peptides (ILPs) previously identified as DAF-2 antagonists to test their role in learning and memory phenotypes. We discovered that only a single uncharacterized ILP, INS-17, is required for learning and memory. We also demonstrate that INS-17 is sufficient to confer extended memory ability and can promote the maintenance of learning and memory with age. Additionally, we observe that *ins-17* regulates learning and memory ability independent of lifespan, uncoupling IIS-mutant phenotypes. We find that regulation of the *ins-17* genetic locus explains its unique requirement among ILP for learning and memory behaviors. Finally, we found that INS-17 may act to signal a state of nutrient deprivation that is required to properly process stimulus valence to promote advantageous behaviors. Our findings deepen the understanding of how IIS can regulate specific phenotypic outputs in response to changes in internal metabolic states.

Introduction

The survival of most multicellular organisms depends on the coordination of complex physiological processes between multiple, functionally discrete tissues. Remarkably, each of these tissues carries out a diverse set of physiological functions using a highly overlapping set of cell regulatory mechanisms, including ubiquitous signaling pathways (e.g. MAPK/ERK, cAMP, and ubiquitin-proteasome pathway signaling[1–4]). Major components of these pathways are commonly pleiotropic, where a single gene can control multiple, distinct phenotypes depending on the context, such as life stage, cell type, and metabolic state. Across species, one prominent pleiotropic signaling pathway is the insulin/insulin-like growth factor (IGF) receptor signaling (IIS), which regulates a wealth of phenotypes across multiple tissues, including growth and development, metabolism, fertility, aging, stress response, and survival [5–15]. Dissecting the specific signaling mechanisms by which IIS mediates such a wide range of phenotypic effects proves challenging, especially in mammalian systems. This is due to several factors, namely high costs to generate tools to examine pathways in a tissue-specific and temporally controlled manner; compensation between receptors, as a single tissue can regulate phenotypes organism-wide; and the time and resource burden to study certain IIS-regulated phenotypes, such as aging. However, invertebrate systems can help overcome these challenges, as IIS is highly conserved across species [9,10,14,16,17], and simple models allow for rapid, low-cost study of genes with remarkable temporal and tissue-specific precision [18–20].

The nematode *C. elegans* has been invaluable in elucidating the pleiotropy underlying IIS

phenotypes. In fact, foundational experiments in the worm were the first to link insulin

signaling and aging, where loss-of-function receptor mutants of the IR homolog, *daf-2*,

have a doubled lifespan of around 40 days compared to the 20-day lifespan of wild-type

animals (21,22). Since this initial work, studies in IIS mutants revealed the pleiotropy of

the DAF-2 receptor and its conserved downstream signaling components (*age-1*/PI3K

and transcription factors *hsf-1*/HSF1, *skn-1*/NRF2, and *daf-16*/forkhead family (FOXO3))

in this relatively simple model (21–28), including that *daf-2* mutants have altered

development, metabolism, fertility, motility, cognitive abilities, and more (12,26,29–39).

Collectively, research indicates that DAF-2/IR functions across tissues and is integral for

sensing internal state along with environmental cues allowing for the worm to adapt

appropriately to improve survival.

Despite decades of study of IIS in the worm, how the DAF-2 receptor coordinates a variety

of phenotypic responses across tissues remains unclear. This is due in part to the ubiquity

of the IIS pathway across tissues, where DAF-2/IR activity in some tissues can mask

DAF-2/IR activity in other tissues. For example, it is well established that *daf-2* mutants

have improved motility with age (30,40,41), but recent work using degron-based

approaches found that reduction of *daf-2* function in neurons and muscles has opposing

effects on maintenance of motility in aged animals (42). This work sheds light on the

remaining gap in our knowledge regarding how signaling through a single ubiquitous

receptor can impact numerous phenotypes that are often reflective of the function of a

relatively small number of cells. It is also unclear whether these distinct phenotypes can

be uncoupled, and if so by what mechanisms.

Though the DAF-2 receptor and its downstream effectors have been well-studied, less is

known about the upstream regulators of DAF-2/IR that drive specific phenotypic outputs.

The *C. elegans* genome encodes 40 insulin-like peptides (ILPs) predicted to act as DAF-

2 receptor agonists or antagonists, sometimes in a context-dependent fashion (43–47).

However, no single manipulation of an ILP fully reproduces reduced *daf-2* function

phenotypes. ILPs also exhibit diverse and often non-overlapping expression patterns and

are regulated by distinct physiological contexts (48). Thus, we hypothesize that pleiotropic

phenotypes exhibited by IIS mutants may also reflect distinct ligand-receptor interactions.

In the present study, we investigated if specific ILPs regulate *daf-2* mutant learning and

memory phenotypes. We discovered that a single ILP, the DAF-2/IR antagonist INS-17,

is required for learning and memory ability and is sufficient to promote memory. We also

found that INS-17 is specifically required for IIS regulation of behavior and does not

appear to be involved in other adult DAF-2 receptor-mediated phenotypes, including

lifespan (47,49). We find that the regulation of INS-17 is the source of this phenotypic

specificity, and that INS-17 acts as a nutrient-deprivation signal that regulates plastic

behavior in the context of changing physiological states.

Results

INS-17 is specifically required for learning and memory ability

Decreased DAF-2 receptor function results in a 3-fold extension in short- and intermediate

associative memory (S/ITAM), as measured by positive olfactory association assays (31).

In this paradigm, animals form a positive association with the neutral odor butanone (10%

in EtOH) when paired with food, which is measured by a training-dependent increase in

butanone chemotaxis (50). A single food-butanone (1 CS-US) pairing usually results in a

molecularly conserved memory that lasts approximately 2 hours; however, young adult

daf-2 mutants maintain a food-butanone association for up to 6 hours (Fig. 1A) (31).

Percent maximum (% Max) performance indices are shown for clarity (see methods for

calculations), as *daf-2* animals have high naïve chemotaxis to 10% butanone in multiple

studies (31,34,36,37) (Fig. S1A) that contribute to perceived learning impairments when

graphed with raw performance indices (provided in Fig. S1B). Given that partial loss-of-

function *daf-2* animals display this significant extension in S/ITAM performance, we

hypothesized that insulin receptor antagonist(s), which are predicted to reduce signaling

through DAF-2/IR, may promote learning and memory behavior (Fig. 1B). To test this

hypothesis, we examined the learning and memory ability of publicly available insulin

receptor antagonist mutants using this STAM paradigm. We tested learning and

intermediate-term associative memory (ITAM, 60 minutes post-training), as behavioral

deficits are typically more prominent at this timepoint compared to STAM (30 minutes

post-training). Mutants tested included predicted strong (*ins -17, -37, and -39)* and weak

(*ins-15, -21,* and *-22)* antagonists (47,51,52)*.* However, we recently determined that

mutants for *ins-22* have an abnormally high preference for 10% butanone (Fig. S1I) (53).

Thus, we excluded *ins-22* from our targeted behavioral screen, as this high naïve

chemotaxis would make it difficult to measure training-dependent increases in butanone

preference. Most antagonist mutants behaved in a manner that was indistinguishable

from wild-type animals, including baseline naïve chemotaxis (Fig. 1D-E, S1E-H); while

mutant animals for *ins-39* had significantly higher intermediate-term associative memory

(ITAM) performance compared to wild-type animals, we were screening for behavioral

deficits and focused on candidates that had impaired behavioral performance.

Interestingly, only loss-of-function mutants of *ins-17* exhibited behavioral deficits: these

include significant deficits in learning, STAM and ITAM performance (Fig 1C). These

deficits were not accompanied by impaired sensory function, as measured by naïve

chemotaxis assays to the highly attractive concentration (0.1%) of butanone (Fig. S1C-

D), suggesting that they are specific to learning and memory. INS-17 is a relatively

uncharacterized ILP; aside from its prediction as a strong antagonist, previous research

has established *ins-17*'s role in regulating dauer formation in early development (49). To

our knowledge, INS-17 is uncharacterized regarding complex behavior, and these results

(Fig. 1C-E) suggest that it is a novel learning and memory-regulating ILP.

142   INS-17 signals through DAF-2 and promotes memory

We next wanted to confirm that INS-17 signaling was indeed mediated by the DAF-2

receptor and tested the learning and memory behavior of *daf-2(e1370);ins-17(tm790)*

double mutants. We examined double mutant memory behavior at 3 hours after 1x

training, a timepoint where *daf-2* mutants exhibit extended memory (31). *daf-*

2(e1370);ins-17(tm790) worms demonstrated extended 3-hour memory indistinguishable

from *daf-2* animals (Fig. 2A, S2A-C). Thus, we infer from these results that DAF-2/IR is

epistatic to INS-17 in the regulation of learning and memory.

Next, we wanted to explore how manipulating levels of INS-17 may influence behavioral

outputs. Our behavior paradigm tests an adult-specific behavior (34), and *ins-17* regulates

dauer in development (49), so we sought to avoid confounding our results with

developmental effects of traditional transgenic approaches. Recently, we developed a

novel microbial feeding paradigm to manipulate neuropeptide signaling specifically in

adult animals (53). In these experiments, *C. elegans* are fed *E. coli* bacteria expressing

a plasmid containing the sequence encoding a neuropeptide of interest flanked by

processing enzyme cut sites (Fig. 2B). Like feeding-based RNAi approaches (54), this

method offers precise temporal control of increasing neuropeptide levels via adult-only

manipulation to avoid potential effects on early development. We have demonstrated that

this feeding method could work in an 'overexpression' manner when administered to wild-

type animals, altering sensory behavior (53). Here, we wanted to extend this approach to

assess if administering additional INS-17 to adult animals could affect associative

behaviors. We hypothesized that increasing the levels of a memory-regulating *daf-2*

antagonist would result in enhanced memory ability. Briefly, we fed L4 animals *E. coli*

expressing either INS-17 or a scrambled control peptide (SCR) and examined their

behavior at Day 2 of adulthood.

To first establish the ability of this method to effectively manipulate INS-17 signaling, we

tested if INS-17 peptide feeding was sufficient to rescue learning and memory behavior

of *ins-17(tm790)* animals. SCR fed mutants still exhibited behavioral deficits, while adult-

only feeding of INS-17 to *ins-17* deficient mutants was sufficient to rescue their behavior,

similar to wild-type animals fed SCR peptide (Fig. 2C). Notably, we determined these

results were not the result of sensory deficits in these animals (Fig. S2D-E). We next

explored if INS-17 peptide feeding was sufficient to boost wild-type animal memory

behavior past the 2-hour timepoint after 1x training. Remarkably, we found that wild-type

animals fed INS-17 during adulthood displayed 3-hour memory post-training, while SCR

fed animals demonstrated no memory phenotype, as expected (Fig. 2D). We confirmed

that peptide feeding to both wild-type and mutant animals did not alter baseline

chemosensation, indicating the effect is memory-specific (Fig. S2D-F). These results

suggest that increasing INS-17 is sufficient to boost cognitive performance and

phenocopy *daf-2* animals' 3-hour memory behavior.

Finally, we worked to identify the tissue where INS-17 is required to regulate memory.

Since INS-17 is highly expressed in the nervous-system, we hypothesized that neurons

may use INS-17 as a memory signal. Using a neuron-specific RNAi-sensitive line(55), we

performed an adult-only *ins-17* knockdown and observed significant S/IATM behavioral

deficits in *ins-17 RNAi* knock-down animals (Fig. 2E)—similar to what we observed in

deletion mutants (Fig. 1B)—with no detectable effects on sensory ability (Fig. S2H-I). This

demonstrates that INS-17 is a memory-promoting DAF-2/IR antagonist, likely released

from neurons.

INS-17 regulates the cognitive aging phenotypes mediated by

the DAF-2 receptor independent of lifespan

In addition to exhibiting extended STAM at young adult stages, *daf-2* mutants also

maintain the ability to learn and remember after 1 CS-US pairing with age (31). Wild-type

animals display age-related cognitive deficits as early as Days 3 and 5 of adulthood (Fig

3A-B, S3A-C), while *daf-2* mutants maintain performance akin to Day 1 of adulthood at

these timepoints. Therefore, we wanted to ask if INS-17 could also be involved in these

age-related phenotypes associated with altered *daf-2* function. As *ins-17* mutants already

display severe behavioral defects at Day 1 of adulthood, we examined the effect of

increased INS-17 on age-related behavioral impairments in wild-type animals, again

using a peptide feeding-based approach (53).

We first assessed the behavior of animals supplemented continuously with INS-17 from

the L4 stage until Day 5 of adulthood (Fig. 3C), when worms are considered cognitively

aged (31). INS-17 fed animals demonstrated significantly better learning and memory

performance compared to wild-type SCR fed controls (Fig. 3C-D, S3D). Next, we

wondered if manipulating the DAF-2/IR pathway via INS-17 later in life could influence

cognitive aging, as previous work has shown that degradation of DAF-2/IR via an auxin

inducible degron system in older animals results in improved STAM (37). We subjected

Day 3 adult wild-type animals (which already display age-related cognitive deficits

compared to Day 1 Adults (Fig. 3A)) to SCR or INS-17 peptide feeding and tested their

learning and memory abilities at Day 5 of adulthood (Fig. 3E). Notably, we found that Day

5 INS-17 fed animals had better learning and STAM behaviors compared to SCR fed

controls (Fig. 3E-F, S3E-F). These findings suggest that INS-17 is a DAF-2/IR ligand that

could partially contribute to the enhanced cognitive ability with age observed when DAF-

2/IR activity levels are reduced.

Since *daf-2* animals have both a doubled lifespan and enhanced memory phenotypes in

young and aged animals (21,31), we wanted to determine if manipulating INS-17, as a

strong DAF-2/IR antagonist, would alter lifespan in addition to associative behaviors. As

previously reported in the literature (49), we confirmed that *ins-17(tm790)* animals have

no significant difference in lifespan to wild-type animals (Fig. 3G, S3F-G). We also tested

the lifespans of wild-type animals fed SCR compared to those fed INS-17 in adulthood,

finding no significant differences between groups (Fig. 3H, S3H-I). Thus, our findings

demonstrate that either loss of *ins-17* or increased levels of INS-17 has no effect on adult

lifespan, highlighting the specific phenotypic regulation by individual ligands underlying

the pleiotropy of the DAF-2 receptor.

224   INS-17 engages multiple pathways downstream of DAF-2 to

225   regulate learning and memory

We next sought to identify the pathways downstream of DAF-2/IR that could explain the

phenotypic specificity of INS-17. The canonical downstream target of DAF-2/IR signaling

is the FOXO transcription factor homolog DAF-16(27,28), which is required for the

doubled lifespan extension of *daf-2* loss-of-function worms (26,28,56). Additionally, *daf-*

16 mutants exhibit severely impaired learning and memory phenotypes (Fig. 4A, S4A

(31). Since *ins-17* is required for positive associative memory behavior and there are no

adult lifespan phenotypes associated with *ins-17* (Fig 3G-H), we hypothesized that *ins-*

17 activates a pathway that acts in parallel to *daf-16*. To test this, we examined if INS-17

feeding could improve the memory impairments of *daf-16* worms. We found that adult-

only INS-17 feeding rescued the learning and memory of *daf-16(mu86)* animals to SCR

fed wild-type performance levels, while SCR fed *daf-16(mu86)* worms still displayed

behavioral deficits (Fig. 4B, S4B-C). These results support that INS-17 could act

independently of DAF-16 to regulate associative behaviors.

239   Regulation of the *ins-17* genetic locus is required for proper

240   regulation of behavior

While we had identified how INS-17 might regulate distinct DAF-2/IR mediated

phenotypes, we also sought to examine what made INS-17 unique relative to other

antagonists in regard to regulating behavior. Interestingly, though the *C. elegans* genome

encodes 40 ILPs, there are only three structurally similar antagonists known as γ-insulins;

these genes are encoded across multiple genomic regions (ch. II and III) and thus likely

under the control of distinct regulatory elements (43,47,57). We therefore hypothesized

that phenotypic specificity was not necessarily due to insulin class or structure, but

instead via *ins-17*'s regulation by its surrounding genomic elements, which likely respond

to different signaling pathways and environmental cues. To test this hypothesis, we

explored if replacing *ins-17* with another strong DAF-2/IR antagonist was sufficient to

maintain normal positive associative memory ability in an *ins-17* mutant background.

We selected *ins-37* to swap in for *ins-17* due to a key overlap in structure, as both ILPs

are predicted strong DAF-2/IR antagonists and are γ-insulins, which are defined by three

di-sulfide bonds (47). Moreover, *ins-37* mutants had no observable learning and memory

impairments (Fig. 1D, S1H) and *ins-37* is lowly expressed in neurons (58), making it an

ideal strong antagonist to test our hypothesis. Thus, we generated an *ins-17* CRISPR

knockout in which we placed the *ins-37* coding sequence at the *ins-17* locus (Fig. 5A).

Subsequently, we examined if *ins-37* expressed at endogenous *ins-17* levels is sufficient

for normal learning and memory behavior. First, we confirmed that the newly generated

ins-17 knockout line exhibited learning and memory impairments similar to the predicted

loss-of-function mutants (Fig 1C, 5B, S5A). Interestingly, we found that replacing *ins-17*

with *ins-37* could rescue the defective learning and memory of *ins-17* KO animals,

functionally replacing endogenous *ins-17* (Fig. 5B, S5A). These results indicate that ILP

structure is not as critical as its regulation of expression. These results suggest that the

genomic context in which *ins-17* is regulated underlies its importance in promoting

specific behavioral phenotypes.

267 *ins-17* expression rapidly responds to nutrient deprivation

Next, we wanted to uncover the physiological conditions that regulate *ins-17* expression

and assess their role in learning and memory. We mined publicly available gene

expression datasets and discovered that *ins-17* transcription is likely regulated by

metabolic state, as it is consistently upregulated in multiple starvation paradigms and

genetic models of nutrient deprivation (Table 1) (49,51,59–62). Interestingly, our

appetitive behavioral paradigm involves a 1-hour food deprivation (FD) step prior to the 1

CS-US (food-butanone) pairing, which has been reported to promote a more robust CS-

US association compared to animals that do not undergo a FD period prior to memory

training (31,63). We first examined if the FD step incorporated into our paradigm was

sufficient to increase *ins-17*. We performed qPCR analysis to measure *ins-17* levels in

wild-type worms after 1 hour of FD and found that even this short time period was

sufficient to observe a significant increase in *ins-17* expression (Fig. 5C, left). We also

observed more robust upregulation of *ins-17* following 2 hours of FD (Fig. S5B),

suggesting that this locus rapidly responds to changes in nutrient availability.

282 INS-17 is required for properly sensing internal nutrient status

283 to allow for processing of stimulus valence

Based on our gene expression results, we hypothesized that INS-17 could promote

memory performance by functioning as a FD signal. We first assessed the role of insulin

signaling in FD-associated improvements in associative behavior (Fig. 5C, right). We

confirmed that FD could indeed improve memory ability by comparing the behavioral

performance of wild-type animals that underwent FD prior to training to animals that were

not food-deprived (non-FD) prior to training. Similar to previous reports (31), we found

that wild-type FD animals have significantly better ITAM performance, which is measured

one hour post-training, compared to non-FD animals (Fig. 5D, S5C-D). Interestingly, *daf-*

2 mutant phenotypes are strongly associated with phenotypes paralleling those of

chronically starved worms (64–67), so we hypothesized that *daf-2* animals may not

require the FD step prior to 1 CS-US pairing for robust ITAM behavior, as they are thought

to have a perceived nutrient status of chronic starvation even when well-fed. We found

that non-FD *daf-2* animals that underwent 1 CS-US pairing had ITAM behavior nearly

identical to *daf-2* animals that were FD prior to training (Fig. 5E, S5E). Altogether, these

results indicate that *daf-2* animals' improved memory phenotype could be driven by their

sensation of a chronically starved internal state.

We next tested if INS-17 could function as a FD signal. If this was the case, we

hypothesized that increased levels of INS-17 could be sufficient to promote robust ITAM

behavior in the absence of the FD step prior to 1 CS-US pairing. To test this, we examined

if wild-type animals fed INS-17 peptide required FD for enhanced ITAM. We compared

memory performance of SCR or INS-17 fed animals after training in the absence of FD,

and found that, similar to *daf-2* mutants, INS-17 fed animals exhibited significantly more

robust ITAM than SCR fed animals in the non-FD condition (Fig. 5F). Furthermore, FD

had no detectable effect on ITAM performance in INS-17 fed animals, but significantly

improved ITAM in SCR fed animals (Fig. S5F-I). These results suggest that INS-17 acts

as a FD signal to promote a stronger association between a food and an odorant.

We next asked if INS-17 functions generally as a FD signal to promote associative

behaviors using an aversive olfactory training paradigm, where an attractive odorant

diacetyl is paired with the absence of food to form a negative association (68). If INS-17

functions as a signal of FD internal state, we hypothesized that *ins-17* mutants would

have impaired aversive learning and memory compared to wild-type animals, because

sensing starvation is required for a negative diacetyl association to be formed. Wild-type

and *ins-17* loss-of-function mutants underwent well-established aversive memory assays

using a diacetyl-starvation pairing (68). Strikingly, we found that *ins-17* is required for

normal aversive associative behaviors, as *ins-17* animals had severe deficits in learning,

STAM, and ITAM behavior compared to wild-type animals (Fig. 5G, S5J). The aversive

and appetitive olfactory conditioning paradigms we tested here recruit distinct neuronal

circuits (63,69–75). Therefore, our results suggest that *ins-17* may be broadly required in

the nervous system to signal a FD state so that the valence of olfactory stimuli (positive

when paired with food, negative when paired with starvation) can be processed

appropriately to promote advantageous behaviors (Fig. 5H). This could explain how IIS

acts in the nervous system to integrate metabolic status and regulate specific behavioral

outputs, such as learning and memory.

Discussion

Across species, insulin signaling is pleiotropic. In the worm, the DAF-2 receptor regulates

a wide array of phenotypes, including those related to development, lifespan, metabolism,

fecundity, cognitive behaviors, and motility, among others. How does a single receptor

mediate such physiologically distinct phenotypes? Here we report that some of these

pleiotropic actions result from specific ligand-receptor interactions. In this instance, we

show that an olfactory memory phenotype is likely mediated by INS-17, an antagonistic

ILP with no previously reported role in behavior (to our knowledge). DAF-2 receptor partial

loss-of-function mutants have a well-established extended memory phenotype (31), so it

is not surprising that a predicted antagonistic ILP regulated these behaviors. However,

we were surprised that only a single strong antagonist appeared to be required in our

paradigms. The importance of INS-17 to the extended memory phenotype is supported

by our finding that overexpression through feeding is sufficient to recapitulate the memory

phenotype of *daf-2* mutants. While we established that other antagonists are not required

for this specific behavioral phenotype, many ILPs are diverse and act as agonists or

antagonists in a context-dependent fashion, and most do not have a described role in

associative behaviors. Future studies systemically characterizing these diverse ligands

will be informative for defining DAF-2 receptor pleiotropy across tissues.

Interestingly, several transcriptomic datasets show that *ins-17* transcripts are increased

roughly 2-fold in *daf-2* mutants and decreased in *daf-16* mutants compared to wild-type

animals (Table 1) (34,49,51,59,60,62). This indicates that INS-17 is highly regulated by

IIS signaling and is potentially acting both upstream and down-stream of DAF-2/IR. "ILP

to ILP" signaling, where the action of one ILP affects the expression of another ILP, could

be a component of *ins-17*'s role in integrating sensory cues and nutrient availability to

elicit behavioral outputs. One example of "ILP to ILP" signaling in behavior is between

INS-6 and INS-7 in aversive pathogen learning, where INS-6 released by the ASI neuron

regulates *ins-7* transcription in the URX neurons to regulate this behavior (76). Further

transcriptional analysis of *ins-17* mutants and INS-17 overexpression models could

provide insight into which ILPs may be regulating or be regulated by INS-17.

We also report that INS-17's function in adulthood is specific to behavior, as manipulating

INS-17 does not affect lifespan (49). While *daf-2* animals' extended lifespan is a canonical

phenotype, there may exist alternate downstream IIS pathways that are dispensable for

lifespan regulation. There are reports that the sole insulin receptor substrate homolog

IST-1 may differentially activate distinct arms of the IIS signaling pathway, as it can

function upstream of—or partially in parallel to—DAF-16 and is not required for normal

growth and development (77–79). Thus, it is possible the 40 known ILPs may similarly

engage the IIS pathway. In the present study, we investigated IST-1, which engages

atypical IIS signaling to regulate aversive olfactory learning (79). Here, we report that *ist-*

1 mutants display positive olfactory memory deficits that were rescued by INS-17 peptide

feeding; taken with our similar results in *daf-16* mutants, these findings suggest INS-17

may activate multiple arms of the IIS pathway to regulate adult cognitive behaviors. We

find that *daf-16;ist-1* double mutants do not respond to INS-17 peptide supplementation,

suggesting that, similar to other olfactory behavioral paradigms(79), INS-17 engages

multiple pathways that can act partially in parallel to one another downstream of DAF-2

to regulate behavior. It will be intriguing in the future to examine if other DAF-2-mediated

associative behaviors similarly engage both downstream pathways.

We also find that the genomic context in which an ILP is regulated provides specificity, as

"swapping" a structurally similar insulin, *ins-37* (47), into *the ins-17* genetic locus is

sufficient to functionally replace *ins-17* in behavior, though endogenous *ins-37* is not

required for learning phenotypes in our paradigm. This result supports that tissue-specific

gene regulation may inform DAF-2/IR pleiotropy, warranting future studies exploring

tissue-specific requirements for DAF-2/IR function.

Furthermore, we demonstrate that *ins-17* is required for both positive and aversive

associative behaviors. Both associative behavioral paradigms include a relatively short,

1-hour FD step prior to training to prime animals to form a stronger association between

food and an odorant, thus promoting a more robust behavioral response after training.

This FD step occurs within the timeframe before animals display significant changes in

sensory integration behavior due to a "lack of food sensation", which begins after 3 hours

of food-deprivation (80); however, food is cleared from the alimentary system within 2

minutes of ingestion (81,82), suggesting the need for a FD signal to inform robust

behavioral responses after only 1 hour of FD. Our gene expression data, similar to

previous studies (51,59,60), shows that the *ins-17* gene is activated in this timeframe,

suggesting that could it be involved in responding to these brief FD signals. This notion is

supported by our findings that feeding INS-17 to animals that do not undergo FD results

in normal learning and memory, bypassing the need for FD to promote memory. Together,

these results suggest that INS-17 could be released in response to acute and prolonged

FD to inform sensory integration behavior. Interestingly, the only phenotype previously

associated with INS-17 is dauer formation to protect progeny from unfavorable conditions

(49), namely starvation (83). We speculate that INS-17 may respond to FD to coordinate

phenotypic responses in different life stages that are most advantageous; for example,

while *ins-17* regulates dauer formation, which is a response to lack of food availability and

other stresses during development, it may also regulate the ability to form associations in

the context of food availability in adulthood, which is beneficial for egg-laying and progeny

survival (64,84). In the future, it will be interesting to further examine how both the *ins-17*

locus and INS-17 secretion are regulated in the context of changing nutrient availability.

Since *ins-17* is required for both positive and aversive associative behaviors, our results

support that INS-17 may function as a broad signal for nutrient deprivation to the nervous

system. *ins-17* is widely expressed across the nervous system (58,61), but is most highly

expressed in the AIZ interneuron, which mediates a variety of behavioral outputs in

response to multiple sensory neurons and circuits (85–87). We observed that associative

behavior impairments resulting from loss of ins-17 do not stem from baseline sensory

deficits, which strongly indicates that rather sensory information may not be integrated

correctly or that sensory neurons cannot alter their responses to an environmental

stimulus in the context of altered metabolic status. We have established that INS-17 may

be essential for interpreting internal state and proper valence processing. In mammals,

neuropeptides also are important for signaling valence in positive and aversive

associative paradigms (88,89), and this is likely a theme of peptides modulating circuits

more broadly. Future studies examining how INS-17 regulates sensory circuit activity in

various contexts will help further define *daf-2* pleiotropic mechanisms.

Our results bring an attractive model for the expansion of the ILP family in *C. elegans*.

ILPs are encoded across the genome and function under the control of a variety of

regulatory elements so the animal can dynamically respond to a combination of nutrients

and environmental stimuli via DAF-2/IR to facilitate the most advantageous phenotype.

Having a diverse array of peptides that can carry out distinct outputs is beneficial,

especially if there is only one receptor to mediate such outputs. In the case of INS-17,

being able to properly make associations in the face of changing environments can

promote survival, such as remembering availability (or lack) of a food source associated

with specific odors. Together, our findings have identified INS-17 as a physiological signal

that is required for positive and aversive associative behaviors and may integrate

environmental cues and nutrient status to inform valence processing and regulate

behavioral outcomes. This research underscores the importance of future work

investigating the rich and diverse network of DAF-2/IR signaling.

Methods
429

General *C. elegans* maintenance
430

Worms were maintained at 20 °C on plates containing either 1) Standard nematode

growth medium (NGM): 3 g/L NaCl, 17 g/L agar, 2.5 g/L peptone, 25 mL/L KPO4 (pH

6.0), 1 mL/L MgSO4, 1 mL/L CaCl2, and 1 mL/L cholesterol (5 mg/mL in ethanol) (90); or

2) High growth medium (HGM): 3 g/L NaCl, 30 g/L agar, 20 g/L peptone, 25 mL/L KPO4

(pH 6.0), 1 mL/L MgSO4, 1 mL/L CaCl2, and 4 mL/L cholesterol (5 mg/mL in ethanol)

(50). Plates were seeded with an *E. coli* OP50 lawn for ad libitum feeding (90).

Animals were developmentally synchronized using an alkaline-bleach solution (1.5 mL 5

N NaOH, 3.0 mL 5% sodium hypochlorite, 5.5 mL water) to collect eggs from gravid

adults; eggs were subsequently washed three times with M9 buffer solution (6 g/L

Na2HPO4, 3 g/L KH2PO4, 5 g/L NaCl and 1 mL/L 1M MgSO4 in ultrapure water) before

plating (90).

Strains
442

Wild-type: N2 Bristol

Mutants: *ins-17(tm790), ins-21(tm5180), ins-37(tm6892)*, and *ins-39(tm6520)* were

obtained from the National BioResource Project, Japan.

RB2489 (*ins-15(ok3444)*), RB2594 (*ins-22(ok3616)*), CB1370 (*daf-2(e1370)*), CF1038

(*daf-16(mu86)*), MAH677 (*sid-1(qt9) V; sqls71[rgef-2p::GFP + rgef-1p::sid-1]*) were

obtained from the Caenorhabditis Genetics Center.

CX17790 (*ist-1(ok2706)* was described previously(79) and generously gifted to us by

Dr. Cornelia Bargmann.

Strains made in collaboration with InVivo Biosystems include *ins-17* CRISPR KO

animals with *ins-37* replacement COP2848 (*ins-17(knu1330[ins-37]) III*) and *ins-17*

CRISPR KO animals COP2835 (*ins-17(knu1320) III*).

The following strains were generated by crosses: RNA34 (*daf-2(e1370)*; *ins-17(tm790)*)

and RNA59 (*daf-16(mu86); ist-1(ok2706)*).

Peptide plasmid generation and transformation

The amino acid coding sequence for each peptide was identified using

www.wormbase.org (57). These sequences were codon-optimized for *C. elegans* and

cloned in pET-21a(+) cloning vector via XhoI/XhoI strategy by GenScript. The scramble

control sequence and plasmid were generated using the same cloning strategy by

GenScript. Plasmids arrived lyophilized and were re-suspended in nuclease-free H_2O to

a concentration of 100ng/μl and stored at -20 °C. Each expression clone was transformed

into competent iOP50 cells using methods previously described(91). Briefly, iOP50 stocks

were streaked out onto LB plates containing 50 μg/mL tetracycline and grown at 37 °C

overnight. Then, single colonies of iOP50 were isolated and used to inoculate 2.0 mL LB

containing 50 μg/ml tetracycline and placed in a 37 °C shaker overnight. 0.5 mL of each

culture was used to inoculate 3.0 mL of LB without antibiotics and grown for 2 hours in a

37 °C shaker. Each culture was then aliquoted in 0.75 mL increments into four 1.5 mL

tubes and cooled for 10 minutes on ice. Next, the cells were harvested by centrifugation

at 6,000 rpm for 5 minutes at 4 °C. The resulting supernatant was discarded, and cells

were placed back on ice and re-suspended in 1.0 mL 100 mM ice-cold $CaCl_2$ solution.

Cells were kept on ice for 20 minutes and then harvested by centrifugation as in the

previous step. After discarding the supernatant, cells were re-suspended in 150μl of ice-

cold $CaCl_2$ solution and placed on ice to be immediately used for transformation.

Competent cells were transformed using the New England Biolabs protocol

(C2987H/C2987I). Single colonies of transformed cells were isolated and cultured in 3.0

mL of LB containing 50 μg/mL tetracycline and 50 μg/mL carbenicillin overnight in a 37

°C shaker. Cultures were frozen down as glycerol stocks and stored at -80 °C. Successful

transformation was verified by isolating plasmids from cultures using a Zyppy Plasmid

Miniprep Kit and sending samples for whole-plasmid sequencing (Plasmidsaurus).

Peptide plate preparation

LB containing 50 μg/mL tetracycline and 50 μg/mL carbenicillin was inoculated with iOP50

encoding either the peptide sequence of interest or scramble peptide sequence and

cultured for 18-20 hours in a 37 °C shaker. 100 mm NGM plates containing 50 μg/mL

tetracycline, 50 μg/mL carbenicillin, and 1.0 mL/L isopropyl-β-D-thiogalactoside (IPTG)

were plated with 1.0 mL of corresponding culture. Seeded plates were left to dry at least

24 hours at room temperature and never kept longer than 72 hours before use.

Approximately one hour before transferring worms to peptide or scramble control plates,

200 µL of 0.1 M IPTG was added to each plate and left to dry.

Peptide feeding

We adapted a previously published peptide feeding-based approach (53). Worms were

maintained on NGM agar plates seeded with OP50 *E. coli* and maintained at 20 °C until

the start of experiments. All animals were transferred to a conical tube and washed three

times with M9 buffer solution before being transferred to NGM plates containing 50 µg/mL

Tetracycline, 50 µg/mL Carbenicillin, and 1.0 mM IPTG that were seeded with 1.0 mL of

iOP50 expressing either the scramble peptide or the peptide of interest. Animals were

kept on the scramble or peptide-expressing lawns for at least 48 hours of exposure time

at 20 °C before performing assays. Exact peptide feeding timeline parameters are

expounded in the results section.

RNAi treatment

Worms were maintained on NGM agar plates seeded with OP50 *E. coli* at 20 °C until the

start of experiments. At the L4 stage, animals were transferred to a conical tube and

washed three times with M9 buffer solution before being transferred to NGM plates

containing 50 µg/mL Carbenicillin and 1.0 mM IPTG that were seeded with 1.0 mL of

HT115 bacteria expressing either RNAi against the gene of interest or the vector control

RNAi. Animals were kept on the RNAi lawns for 48 hours of exposure time at 20 °C before

performing assays at Day 2 of adulthood.

Naïve chemotaxis assay

Chemotaxis assays were performed using a protocol based on previously published

assays(92). Briefly, assays were performed on unseeded 100mm NGM plates. On the

back of each plate, two marks were made on opposite sides of the plate, approximately

5mm from the edge. 1µL of sodium azide (Thermo Fisher) was placed on both spots and

allowed to dry before adding 1µL of odorant (Sigma Aldrich) diluted in ethanol on one

mark and 1µL ethanol on the other. Using M9 buffer, worms were washed off their plates

and subsequently washed three times to eliminate any leftover bacteria that could impact

baseline behavior. Then, worms were placed near the bottom center of the plate,

equidistant between the two marks, and allowed to chemotax for an hour. Chemotaxis

indices were calculated as follows:

Chemotaxis Index = (# of worms butanone - # of worms ethanol) / (total # of worms)

For naïve chemotaxis assays performed to assess baseline chemosensory function, we

tested the naïve preference to a highly appetitive concentration of butanone (0.1%) as

previously reported (92).

Positive associative butanone learning and memory assay

Worms were trained and tested for learning, short- and intermediate-term memory

changes as previously described (50). Briefly, synchronized adult worms were washed

off plates with M9 buffer and allowed to settle by gravity, followed by two subsequent

washes with M9 buffer to remove leftover bacteria that could alter baseline behavior.

Next, the worms were food-deprived for one hour in M9 buffer before exposure to 1 CS-

US pairing, where food-deprived worms were transferred to 100mm NGM conditioning

plates seeded with OP50 *E. coli* bacteria (or iOP50 bacteria containing a peptide or

scramble control plasmid for peptide feeding experiments) and had 16µL of 10%

butanone (Sigma Aldrich) diluted in ethanol streaked on the lid in a '#' shape for one hour.

After conditioning, the trained population of worms were tested for chemotaxis to 10%

butanone and to an ethanol control using standard, previously described chemotaxis

conditions(92). Different stages of memory were tested by measuring chemotaxis indices

of separate populations of worms at timepoints including 0 min. (learning), 30 min.

(STAM), 60 min. (ITAM) and 120 min. (forgetting) after training. For timepoints after

learning behavior, worms were transferred to hold plates seeded with OP50 *E. coli* or

iOP50 bacteria for peptide feeding experiments as stated above.

Chemotaxis indices for each timepoint were calculated as detailed in "Naïve chemotaxis

assays" section.

Performance index is the change in the chemotaxis index after training relative to the

naïve (untrained) chemotaxis index:

Performance index = Chemotaxis index$_{trained}$—Chemotaxis index$_{untrained}$

% Maximum Performance Indices are performance indices represented as a percentage

of the maximum value that can be performed relative to the naïve chemotaxis

index(31,36). This % can be more informative for visualizing performance indices when

naïve chemotaxis behavior is high, as in *daf-2(e1370)* animals. This is calculated for each

timepoint by dividing the performance index of a timepoint by the average of the untrained

chemotaxis indices of the same behavioral replicate subtracted from 1:

% Maximum Performance Index = Performance index$_{timepoint}$/(1 - Chemotaxis index$_{untrained}$)

Butanone enhancement assay

553 This assay was adapted from previously described methods (63). Unlike the positive

554 butanone associative memory assays, butanone enhancement assays skip the 1-hour

555 food-deprivation step prior to the conditioning step. All other methods are identical to

556 those listed in the positive butanone associative memory assay section.

Negative associative learning and memory assay

Worms were tested for learning and memory changes in response to aversive diacetyl

training, adapted from previously published methods (68). Similar to positive associative

assays, synchronized Day 1 adult worms were washed off plates with M9 buffer and then

allowed to settle by gravity and washed twice more with M9 buffer. After washing, the

worms were conditioned for one hour by food-deprivation on 100mm NGM plates with no

bacteria and 16uL of 100% diacetyl (Sigma Aldrich) streaked on the lid in a '#' shape. The

trained population of worms were then tested for chemotaxis to 1% diacetyl and to an

ethanol control using standard, previously described chemotaxis conditions (92).

Learning and memory performance indices were calculated for each timepoint as

previously stated in the positive butanone associative assay section.

Lifespan assay

Lifespan assays were performed on *ins-17(tm790)* animals and wild-type animals that

were bleached onto NGM plates seeded with OP50 *E. coli* and maintained at 20 °C; at

the L4 larval stage, worms were picked for survival analysis and transferred every other

day to freshly seeded plates. Similarly, lifespan assays with peptide feeding

administration were performed on wild-type animals that were bleached onto NGM plates

seeded with OP50 *E. coli* and maintained at 20 °C. Then, at the L4 larval stage, worms

were picked for survival analysis and transferred every other day to NGM plates

containing 50 µg/mL tetracycline, 50 µg/mL carbenicillin, and 1.0 mL/L IPTG that were

seeded with either INS-17 iOP50 or scramble iOP50 *E. coli*. Lifespans began (t = 0) at

Day 1 of adulthood and had an n = ~120 worms per strain/trial for a total of three biological

replicates per assay.

Food-deprivation behavioral experiments

For food-deprivation experiments, well-fed, synchronized adult worms were washed off

plates using M9 buffer into a 15mL conical and washed two more times to remove bacteria

that could alter baseline behavior. Next, worms were either 1) food-deprived for one hour

on a 100mm NGM plate without food or 2) non-food deprived for one hour by being

transferred to a 100mm plate seeded with a bacterial lawn (iOP50 bacteria for peptide

feeding experiments). Worms were then immediately exposed to the conditioning step (1

CS-US pairing) and stages of learning and memory were tested as stated in the positive

associative butanone learning and memory assays section.

RNA isolation, cDNA synthesis and qRT-PCR

For qRT-PCR food-deprivation experiments, well-fed, synchronized adult worms were

washed off plates using M9 buffer into a 15mL conical and washed two more times to

remove bacteria that could alter baseline behavior. Next, worms were either 1) food-

deprived for one hour or two hours after being transferred to a 100mm plate with no

bacterial lawn or 2) non-food deprived for one hour or two hours after being transferred

to a 100mm plate seeded with a bacterial lawn.

After feeding or food deprivation, worms were crushed in liquid nitrogen using a mortar

and pestle and added to Trizol (Thermo Fisher Scientific). RNA was isolated per

manufacturer's instructions, followed by DNase treatment (Qiagen). cDNA was

synthesized with an oligo dT primer and Superscript III reverse transcriptase enzyme

(Thermo Fisher Scientific). cDNA was mixed with buffers, primers, SYBR green, and hot

start Taq polymerase in a master mix prepared by a manufacturer (Thermo Fisher

Scientific). PCR reactions were run using a Quant Studio 7 Pro Dx Real-Time PCR

System (Thermo Fisher Scientific), followed by a dissociation reaction to determine

specificity of the amplified product. Gene expression was quantified using the ΔΔCt

method using *pmp-3* as a reference gene. Primer sets were as follows:

ins-17 For: 5'- GGACACTATCCGACCACCAC -3':

ins-17 Rev: 5'- TTCACATCCCGTCTCACAGC -3':

pmp-3 For: 5'- AGTTCCGGTTGGATTGGTCC -3'

pmp-3 Rev: 5'- CCAGCACGATAGAAGGCGAT-3'

Statistical analysis and software

All statistics (excluding survival analysis) were calculated with GraphPad Prism Software.

All statistical data are reported in the main text, figures, and tables. Significance threshold

of $p < 0.05$ was used. The symbols *, **, ***, and **** refer to $p < 0.05$, 0.01, 0.001, and

0.0001. The symbols #, ##, ###, and #### refer to $p < 0.05$, 0.01, 0.001, and 0.0001. For

comparing performance indices between two behavior conditions (e.g. *vector control*

vs *ins-17 RNAi*), we used a Mann-Whitney test comparing ranks, as it does not assume

normality. For comparing performance indices between three or more groups (e.g. wild-

type vs multiple insulin receptor antagonist mutants), we performed one-way analysis of

variances followed by Bonferroni post hoc tests for multiple comparisons. Two-way

ANOVAs were used for evaluating effects between genotype and timepoint (T0, T30, T60,

T120) on performance indices with a significant interaction between factors ($p<0.0001$)

that prompted Bonferroni post-hoc analyses to determine differences between individual

groups. Sample size n represents the number of chemotaxis assays performed for

behavioral experiments, with each assay containing approximately 50–150 worms. All

experiments were performed on different days with distinct populations to confirm

reproducibility. For qPCR analysis, unpaired t-tests with Welch's corrections were

performed to determine significance between conditions. Survival analysis statistics were

generated using the log-rank (Mantel-Cox) method to test the null hypothesis in Kaplan-

Meier survival analysis and evaluated using OASIS survival analysis software (93).

Funding

631 RNA was supported by a Glenn Foundation for Medical Research and AFAR Grant for

632 Junior Faculty, the Whitehall Foundation, and an NIH Director's New Innovator Award

633 (DP2NS132372). EWP is supported by a NIGMS NRSA, EJL is funded by a NIA NRSA.

634 Some strains were provided by the *C. elegans* Genetics Center (CGC), which is funded

635 by the NIH Office of Research Infrastructure Programs (P40 OD010440).

Acknowledgements

We thank the Bargmann lab for generously sharing the *ist-1(ok2706)* strain, and the

Caenorhabditis Genetics Center for most other strains used in this publication. We also

thank Ben Jussila and In Vivo Biosystems for strains and helpful discussion. Additionally,

we especially thank Drs. Buck Samuel and Shinya Yamamoto at Baylor College of

Medicine for helpful discussion and advice at various stages of the project. Finally, we

thank Arey lab members Katie L. Brandel-Ankrapp and Ty M. Gadberry for feedback on

the manuscript.

References:

1.   Cooper GM. Pathways of Intracellular Signal Transduction. The Cell: A
Molecular Approach [Internet]. 2nd ed. Sinauer Associates; 2000 [cited 2025 Sep
29]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK9870/

2.   Li Y, Li S, Wu H. Ubiquitination-Proteasome System (UPS) and Autophagy Two
Main Protein Degradation Machineries in Response to Cell Stress. Cells. 2022
Mar 1;11(5).

3.   Buscà R, Pouysségur J, Lenormand P. ERK1 and ERK2 map kinases: specific
roles or functional redundancy? Front Cell Dev Biol. 2016 Jun 8;4:53.

4.   Lefkimmiatis K, Zaccolo M. cAMP signaling in subcellular compartments.
Pharmacol Ther. 2014 Sep;143(3):295–304.

5.   Ullrich A, Bell JR, Chen EY, Herrera R, Petruzzelli LM, Dull TJ, et al. Human
insulin receptor and its relationship to the tyrosine kinase family of oncogenes.
Nature. 1985;313(6005):756–61.

6.   Ullrich A, Gray A, Tam AW, Yang-Feng T, Tsubokawa M, Collins C, et al.
Insulin-like growth factor I receptor primary structure: comparison with insulin
receptor suggests structural determinants that define functional specificity. EMBO
J. 1986 Oct;5(10):2503–12.

7.   White MF. The IRS-signalling system: a network of docking proteins that
mediate insulin action. Mol Cell Biochem. 1998 May;182(1–2):3–11.

8.   Taniguchi CM, Emanuelli B, Kahn CR. Critical nodes in signalling pathways:
insights into insulin action. Nat Rev Mol Cell Biol. 2006 Feb;7(2):85–96.

9.   Piñero González J, Carrillo Farnés O, Vasconcelos ATR, González Pérez A.
Conservation of key members in the course of the evolution of the insulin
signaling pathway. BioSystems. 2009 Jan;95(1):7–16.

10.   Kaletsky R, Murphy CT. The role of insulin/IGF-like signaling in C. elegans
longevity and aging. Dis Model Mech. 2010 Aug 1;3(7–8):415–9.

11.   Finch CE, Ruvkun G. The genetics of aging. Annu Rev Genomics Hum Genet.
2001;2:435–62.

12.   Murphy CT, Hu PJ. Insulin/insulin-like growth factor signaling in C. elegans.
WormBook. 2013 Dec 26;1–43.

13.   Semaniuk U, Piskovatska V, Strilbytska O, Strutynska T, Burdyliuk N, Vaiserman
A, et al. *Drosophila* insulin-like peptides: from expression to functions – a review.
Entomol Exp Appl. 2021 Feb;169(2):195–208.

14.   Biglou SG, Bendena WG, Chin-Sang I. An overview of the insulin signaling
pathway in model organisms Drosophila melanogaster and Caenorhabditis
elegans. Peptides. 2021 Nov;145:170640.

15.   Suzawa M, Bland ML. Insulin signaling in development. Development. 2023 Oct
15;150(20).

16.   Barbieri M, Bonafè M, Franceschi C, Paolisso G. Insulin/IGF-I-signaling
pathway: an evolutionarily conserved mechanism of longevity from yeast to
humans. Am J Physiol Endocrinol Metab. 2003 Nov;285(5):E1064-71.

17.   Lau HE, Chalasani SH. Divergent and convergent roles for insulin-like peptides
in the worm, fly and mammalian nervous systems. Invert Neurosci. 2014
Sep;14(2):71–8.

18.   Hales KG, Korey CA, Larracuente AM, Roberts DM. Genetics on the fly: A
692   primer on the drosophila model system. Genetics. 2015 Nov;201(3):815–42.

693   19.   Boulin T, Hobert O. From genes to function: the C. elegans genetic toolbox.
694   Wiley Interdiscip Rev Dev Biol. 2012 Feb;1(1):114–37.

695   20.   Simone BW, Martínez-Gálvez G, WareJoncas Z, Ekker SC. Fishing for
696   understanding: Unlocking the zebrafish gene editor's toolbox. Methods. 2018 Nov
697   1;150:3–10.

698   21.   Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A C. elegans mutant that
699   lives twice as long as wild type. Nature. 1993 Dec 2;366(6454):461–4.

700   22.   Friedman DB, Johnson TE. A mutation in the age-1 gene in Caenorhabditis
701   elegans lengthens life and reduces hermaphrodite fertility. Genetics. 1988
702   Jan;118(1):75–86.

703   23.   Hsu A-L, Murphy CT, Kenyon C. Regulation of aging and age-related disease by
704   DAF-16 and heat-shock factor. Science. 2003 May 16;300(5622):1142–5.

705   24.   An JH, Blackwell TK. SKN-1 links C. elegans mesendodermal specification to a
706   conserved oxidative stress response. Genes Dev. 2003 Aug 1;17(15):1882–93.

707   25.   Morris JZ, Tissenbaum HA, Ruvkun G. A phosphatidylinositol-3-OH kinase
708   family member regulating longevity and diapause in Caenorhabditis elegans.
709   Nature. 1996 Aug 8;382(6591):536–9.

710   26.   Kimura KD, Tissenbaum HA, Liu Y, Ruvkun G. daf-2, an insulin receptor-like
711   gene that regulates longevity and diapause in Caenorhabditis elegans. Science.
712   1997 Aug 15;277(5328):942–6.

713   27.   Lin K, Dorman JB, Rodan A, Kenyon C. daf-16: An HNF-3/forkhead family
714   member that can function to double the life-span of Caenorhabditis elegans.
715   Science. 1997 Nov 14;278(5341):1319–22.

716   28.   Ogg S, Paradis S, Gottlieb S, Patterson GI, Lee L, Tissenbaum HA, et al. The
717   Fork head transcription factor DAF-16 transduces insulin-like metabolic and
718   longevity signals in C. elegans. Nature. 1997 Oct 30;389(6654):994–9.

719   29.   Luo S, Kleemann GA, Ashraf JM, Shaw WM, Murphy CT. TGF-β and insulin
720   signaling regulate reproductive aging via oocyte and germline quality
721   maintenance. Cell. 2010 Oct 15;143(2):299–312.

722   30.   Mulcahy B, Holden-Dye L, O'Connor V. Pharmacological assays reveal age-
723   related changes in synaptic transmission at the Caenorhabditis elegans

neuromuscular junction that are modified by reduced insulin signalling. J Exp
Biol. 2013 Feb 1;216(Pt 3):492–501.

31.   Kauffman AL, Ashraf JM, Corces-Zimmerman MR, Landis JN, Murphy CT.
Insulin signaling and dietary restriction differentially influence the decline of
learning and memory with age. PLoS Biol. 2010 May 18;8(5):e1000372.

32.   Riddle DL, Swanson MM, Albert PS. Interacting genes in nematode dauer larva
formation. Nature. 1981 Apr 23;290(5808):668–71.

33.   Gems D, Sutton AJ, Sundermeyer ML, Albert PS, King KV, Edgley ML, et al.
Two pleiotropic classes of daf-2 mutation affect larval arrest, adult behavior,
reproduction and longevity in Caenorhabditis elegans. Genetics. 1998
Sep;150(1):129–55.

34.   Ange JS, Weng Y, Stevenson ME, Kaletsky R, Moore RS, Zhou S, et al. Adult
Single-nucleus Neuronal Transcriptomes of Insulin Signaling Mutants Reveal
Regulators of Behavior and Learning. BioRxiv. 2024 Feb 7;

35.   Templeman NM, Luo S, Kaletsky R, Shi C, Ashraf J, Keyes W, et al. Insulin
Signaling Regulates Oocyte Quality Maintenance with Age via Cathepsin B
Activity. Curr Biol. 2018 Mar 5;28(5):753-760.e4.

36.   Weng Y, Zhou S, Morillo K, Kaletsky R, Lin S, Murphy CT. The neuron-specific
IIS/FOXO transcriptome in aged animals reveals regulatory mechanisms of
cognitive aging. eLife. 2024 Jun 26;13.

37.   Zhou S, Novak KE, Kaletsky R, Weng Y, Ange JS, Stevenson ME, et al. Body-
to-brain insulin and Notch signaling regulates memory through neuronal CREB
activity. Nat Aging. 2025 Jul;5(7):1232–48.

38.   Hughes SE, Evason K, Xiong C, Kornfeld K. Genetic and pharmacological
factors that influence reproductive aging in nematodes. PLoS Genet. 2007 Feb
16;3(2):e25.

39.   Lin K, Hsin H, Libina N, Kenyon C. Regulation of the Caenorhabditis elegans
longevity protein DAF-16 by insulin/IGF-1 and germline signaling. Nat Genet.
2001 Jun;28(2):139–45.

40.   Hahm J-H, Kim S, DiLoreto R, Shi C, Lee S-JV, Murphy CT, et al. C. elegans
maximum velocity correlates with healthspan and is maintained in worms with an
insulin receptor mutation. Nat Commun. 2015 Nov 20;6:8919.

41.   Bansal A, Zhu LJ, Yen K, Tissenbaum HA. Uncoupling lifespan and healthspan
in Caenorhabditis elegans longevity mutants. Proc Natl Acad Sci USA. 2015 Jan
20;112(3):E277-86.

42.   Roy C, Molin L, Alcolei A, Solyga M, Bonneau B, Vachon C, et al. DAF-2/insulin
760   IGF-1 receptor regulates motility during aging by integrating opposite signaling
761   from muscle and neuronal tissues. Aging Cell. 2022 Aug;21(8):e13660.

762   43.   Duret L, Guex N, Peitsch MC, Bairoch A. New insulin-like proteins with atypical
763   disulfide bond pattern characterized in Caenorhabditis elegans by comparative
764   sequence analysis and homology modeling. Genome Res. 1998 Apr;8(4):348–
765   53.

766   44.   Kawano T, Ito Y, Ishiguro M, Takuwa K, Nakajima T, Kimura Y. Molecular
767   cloning and characterization of a new insulin/IGF-like peptide of the nematode
768   Caenorhabditis elegans. Biochem Biophys Res Commun. 2000 Jul 5;273(2):431–
769   6.

770   45.   Pierce SB, Costa M, Wisotzkey R, Devadhar S, Homburger SA, Buchman AR, et
771   al. Regulation of DAF-2 receptor signaling by human insulin and ins-1, a member
772   of the unusually large and diverse C. elegans insulin gene family. Genes Dev.
773   2001 Mar 15;15(6):672–86.

774   46.   Li W, Kennedy SG, Ruvkun G. daf-28 encodes a C. elegans insulin superfamily
775   member that is regulated by environmental cues and acts in the DAF-2 signaling
776   pathway. Genes Dev. 2003 Apr 1;17(7):844–58.

777   47.   Zheng S, Chiu H, Boudreau J, Papanicolaou T, Bendena W, Chin-Sang I. A
778   functional study of all 40 Caenorhabditis elegans insulin-like peptides. J Biol
779   Chem. 2018 Oct 26;293(43):16912–22.

780   48.   Ritter AD, Shen Y, Fuxman Bass J, Jeyaraj S, Deplancke B, Mukhopadhyay A,
781   et al. Complex expression dynamics and robustness in C. elegans insulin
782   networks. Genome Res. 2013 Jun;23(6):954–65.

783   49.   Matsunaga Y, Nakajima K, Gengyo-Ando K, Mitani S, Iwasaki T, Kawano T. A
784   Caenorhabditis elegans insulin-like peptide, INS-17: its physiological function and
785   expression pattern. Biosci Biotechnol Biochem. 2012 Nov 7;76(11):2168–72.

786   50.   Kauffman A, Parsons L, Stein G, Wills A, Kaletsky R, Murphy C. C. elegans
787   positive butanone learning, short-term, and long-term associative memory
788   assays. J Vis Exp. 2011 Mar 11;(49).

789   51.   Chen Y, Baugh LR. Ins-4 and daf-28 function redundantly to regulate C. elegans
790   L1 arrest. Dev Biol. 2014 Oct 15;394(2):314–26.

791   52.   Zhu R, Chin-Sang ID. C. elegans insulin-like peptides. Mol Cell Endocrinol. 2024
792   May 1;585:112173.

793   53.   DiLoreto EM, Shastry S, Leptich EJ, Reilly DK, Arey RN, Srinivasan J.
794   Harnessing Microbial Tools: Escherichia coli as a Vehicle for Neuropeptide
795   Functional Analysis in Caenorhabditis elegans. Genetics. 2025 Aug 6;

796   54.   Kamath RS, Ahringer J. Genome-wide RNAi screening in Caenorhabditis
797   elegans. Methods. 2003 Aug;30(4):313–21.

798   55.   Yang Y, Arnold ML, Lange CM, Sun L-H, Broussalian M, Doroodian S, et al.
799   Autophagy protein ATG-16.2 and its WD40 domain mediate the beneficial effects
800   of inhibiting early-acting autophagy genes in C. elegans neurons. Nat Aging.
801   2024 Feb;4(2):198–212.

802   56.   Murphy CT, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, et al.
803   Genes that act downstream of DAF-16 to influence the lifespan of Caenorhabditis
804   elegans. Nature. 2003 Jul 17;424(6946):277–83.

805   57.   Sternberg PW, Van Auken K, Wang Q, Wright A, Yook K, Zarowiecki M, et al.
806   WormBase 2024: status and transitioning to Alliance infrastructure. Genetics.
807   2024 May 7;227(1).

808   58.   Taylor SR, Santpere G, Weinreb A, Barrett A, Reilly MB, Xu C, et al. Molecular
809   topography of an entire nervous system. Cell. 2021 Aug 5;184(16):4329-
810   4347.e23.

811   59.   Mueller MM, Castells-Roca L, Babu V, Ermolaeva MA, Müller R-U, Frommolt P,
812   et al. DAF-16/FOXO and EGL-27/GATA promote developmental growth in
813   response to persistent somatic DNA damage. Nat Cell Biol. 2014
814   Dec;16(12):1168–79.

815   60.   Hibshman JD, Doan AE, Moore BT, Kaplan RE, Hung A, Webster AK, et al. daf-
816   16/FoxO promotes gluconeogenesis and trehalose synthesis during starvation to
817   support survival. eLife. 2017 Oct 24;6.

818   61.   St Ange J, Weng Y, Kaletsky R, Stevenson ME, Moore RS, Zhou S, et al. Adult
819   single-nucleus neuronal transcriptomes of insulin signaling mutants reveal
820   regulators of behavior and learning. Cell Genomics. 2024 Dec 11;4(12):100720.

821   62.   Ogawa T, Isik M, Wu Z, Kurmi K, Meng J, Cho S, et al. Nutrient control of growth
822   and metabolism through mTORC1 regulation of mRNA splicing. Mol Cell. 2024
823   Dec 5;84(23):4558-4575.e8.

824   63.   Torayama I, Ishihara T, Katsura I. Caenorhabditis elegans integrates the signals
825   of butanone and food to enhance chemotaxis to butanone. J Neurosci. 2007 Jan
826   24;27(4):741–50.

64.   Baugh LR, Hu PJ. Starvation responses throughout the caenorhabditiselegans
life cycle. Genetics. 2020 Dec;216(4):837–78.

65.   Ben Arous J, Laffont S, Chatenay D. Molecular and sensory basis of a food
related two-state behavior in C. elegans. PLoS ONE. 2009 Oct 23;4(10):e7584.

66.   Kimura KD, Riddle DL, Ruvkun G. The C. elegans DAF-2 insulin-like receptor is
abundantly expressed in the nervous system and regulated by nutritional status.
Cold Spring Harb Symp Quant Biol. 2011 Nov 28;76:113–20.

67.   Venz R, Pekec T, Katic I, Ciosk R, Ewald CY. End-of-life targeted degradation of
DAF-2 insulin/IGF-1 receptor promotes longevity free from growth-related
pathologies. eLife. 2021 Sep 10;10.

68.   Stetak A, Hörndli F, Maricq AV, van den Heuvel S, Hajnal A. Neuron-specific
regulation of associative learning and memory by MAGI-1 in C. elegans. PLoS
ONE. 2009 Jun 24;4(6):e6019.

69.   Fadda M, De Fruyt N, Borghgraef C, Watteyne J, Peymen K, Vandewyer E, et
al. NPY/NPF-Related Neuropeptide FLP-34 Signals from Serotonergic Neurons
to Modulate Aversive Olfactory Learning in Caenorhabditis elegans. J Neurosci.
2020 Jul 29;40(31):6018–34.

70.   Choi JI, Lee HK, Kim HS, Park SY, Lee TY, Yoon K-H, et al. Odor-dependent
temporal dynamics in Caenorhabitis elegans adaptation and aversive learning
behavior. PeerJ. 2018 Jun 12;6:e4956.

71.   Amano H, Maruyama IN. Aversive olfactory learning and associative long-term
memory in Caenorhabditis elegans. Learn Mem. 2011 Oct;18(10):654–65.

72.   Chen Z, Hendricks M, Cornils A, Maier W, Alcedo J, Zhang Y. Two insulin-like
peptides antagonistically regulate aversive olfactory learning in C. elegans.
Neuron. 2013 Feb 6;77(3):572–85.

73.   Ha H, Hendricks M, Shen Y, Gabel CV, Fang-Yen C, Qin Y, et al. Functional
organization of a neural network for aversive olfactory learning in Caenorhabditis
elegans. Neuron. 2010 Dec 22;68(6):1173–86.

74.   O'Halloran DM, Altshuler-Keylin S, Lee JI, L'Etoile ND. Regulators of AWC-
mediated olfactory plasticity in Caenorhabditis elegans. PLoS Genet. 2009 Dec
11;5(12):e1000761.

75.   Chalasani SH, Kato S, Albrecht DR, Nakagawa T, Abbott LF, Bargmann CI.
Neuropeptide feedback modifies odor-evoked dynamics in Caenorhabditis
elegans olfactory neurons. Nat Neurosci. 2010 May;13(5):615–21.

76. Mendler M, Riedinger C, Schlotterer A, Volk N, Fleming T, Herzig S, et al.
Reduction in ins-7 gene expression in non-neuronal cells of high glucose
exposed Caenorhabditis elegans protects from reactive metabolites, preserves
neuronal structure and head motility, and prolongs lifespan. J Diabetes
Complicat. 2017 Feb;31(2):304–10.

77. Chen J, Scheiner AR, Falsztyn IB, Baugh LR. ist-1/IRS1 affects L1 starvation
resistance in daf-16/FoxO- dependent and independent fashion. MicroPubl Biol.
2025 Jun 13;2025.

78. Wolkow CA, Muñoz MJ, Riddle DL, Ruvkun G. Insulin receptor substrate and
p55 orthologous adaptor proteins function in the Caenorhabditis elegans daf-
2/insulin-like signaling pathway. J Biol Chem. 2002 Dec 20;277(51):49591–7.

79. Cheng D, Lee JS, Brown M, Ebert MS, McGrath PT, Tomioka M, et al.
Insulin/IGF signaling regulates presynaptic glutamate release in aversive
olfactory learning. Cell Rep. 2022 Nov 22;41(8):111685.

80. Matty MA, Lau HE, Haley JA, Singh A, Chakraborty A, Kono K, et al. Intestine-
to-neuronal signaling alters risk-taking behaviors in food-deprived Caenorhabditis
elegans. PLoS Genet. 2022 May 5;18(5):e1010178.

81. Viri V, Arveiler M, Lehnert T, Gijs MAM. An In Vivo Microfluidic Study of Bacterial
Load Dynamics and Absorption in the C. elegans Intestine. Micromachines
(Basel). 2021 Jul 17;12(7).

82. Ghafouri S, McGhee JD. Bacterial residence time in the intestine of
Caenorhabditis elegans. Nematol. 2007;9(1):87–91.

83. Cassada RC, Russell RL. The dauerlarva, a post-embryonic developmental
variant of the nematode Caenorhabditis elegans. Dev Biol. 1975 Oct;46(2):326–
42.

84. Weinshenker D, Garriga G, Thomas JH. Genetic and pharmacological analysis
of neurotransmitters controlling egg laying in C. elegans. J Neurosci. 1995
Oct;15(10):6975–85.

85. Kuhara A, Mori I. Molecular physiology of the neural circuit for calcineurin-
dependent associative learning in Caenorhabditis elegans. J Neurosci. 2006 Sep
13;26(37):9355–64.

86. Tsalik EL, Hobert O. Functional mapping of neurons that control locomotory
behavior in Caenorhabditis elegans. J Neurobiol. 2003 Aug;56(2):178–97.

87. Iino Y, Yoshida K. Parallel use of two behavioral mechanisms for chemotaxis in
Caenorhabditis elegans. J Neurosci. 2009 Apr 29;29(17):5370–80.

88.   Li H, Namburi P, Olson JM, Borio M, Lemieux ME, Beyeler A, et al. Neurotensin
orchestrates valence assignment in the amygdala. Nature. 2022
Aug;608(7923):586–92.

89.   Kim D-I, Kang SJ, Jhang J, Jo YS, Park S, Ye M, et al. Encoding opposing
valences through frequency-dependent transmitter switching in single peptidergic
neurons. bioRxiv. 2024 Jan 1;

90.   Brenner S. The genetics of Caenorhabditis elegans. Genetics. 1974
May;77(1):71–94.

91.   Neve IAA, Sowa JN, Lin C-CJ, Sivaramakrishnan P, Herman C, Ye Y, et al.
Escherichia coli Metabolite Profiling Leads to the Development of an RNA
Interference Strain for Caenorhabditis elegans. G3 (Bethesda). 2020 Jan
7;10(1):189–98.

92.   Bargmann CI, Hartwieg E, Horvitz HR. Odorant-selective genes and neurons
mediate olfaction in C. elegans. Cell. 1993 Aug 13;74(3):515–27.

93.   Yang J-S, Nam H-J, Seo M, Han SK, Choi Y, Nam HG, et al. OASIS: online
application for the survival analysis of lifespan assays performed in aging research.
PLoS ONE. 2011 Aug 15;6(8):e23525.

Figure legends

Fig. 1. INS-17 is a novel and specific memory-regulating insulin receptor antagonist

(A) Partial loss-of-function *daf-2(e1370)* mutants have extended STAM that lasts 6 hours

compared to 2 hours for wild-type animals after 1 CS-US food-butanone pairing. % Max

Performance reported as *daf-2(e1370)* animals have a higher baseline naïve attraction to

neutral (10%) butanone concentration (see **Fig. S1A, B**). Mean $\pm$ SEM. n = 15 per

genotype. **(B)** Diagram depicting the hypothesis that insulin receptor antagonists may

inhibit DAF-2 receptor function to promote learning and memory. **(C)** *ins-17* deletion

mutants have impaired learning, STAM, and ITAM compared to wild-type worms. Mean $\pm$

SEM. n = 15 per genotype. ***$p < 0.001$, ****$p < 0.001$; ns, not significant ($p > 0.05$). **(D)**

Weak insulin antagonist mutants for *ins-15* and *ins-21* and **(E)** strong insulin antagonist

mutants for *ins-37* and *ins-39* display no memory impairments; *ins-39* mutants have better

ITAM performance compared to wild-type. Mean $\pm$ SEM. n = 15 per genotype. ***p <

0.001.

Fig. 2. INS-17 and DAF-2 are epistatic and promote memory

(A) *daf-2* and *ins-17* share an epistatic relationship. 3hrs post-training (1 CS-US pairing),

daf-2(e1370) mutants and *daf-2(e1370);ins-17(tm790)* double mutants display memory

behavior while *ins-17(tm790)* animals display no memory behavior. Mean $\pm$ SEM. n = 15

per genotype. ****p < 0.0001. **(B)** Diagram of microbial feeding paradigm. iOP50 bacteria

were transformed with a plasmid containing either the peptide sequence of interest or a

scramble control sequence and then fed to worms before testing behavioral effects. **(C)**

INS-17 peptide feeding rescues *ins-17(tm790)* learning, STAM and ITAM behaviors, while

SCR fed mutants display significant S/ITAM impairments compared to SCR fed wild-type

animals. Mean $\pm$ SEM. n = 15 per feeding treatment. # is significance relative to SCR fed

wild-type animal performance.   ###p < 0.001, ####p < 0.0001; ns, not significant (p >

0.05). **(D)** Wild-type animals fed INS-17 peptides display 3-hour memory behavior after 1

CS-US pairing, similar to *daf-2(e1370)* extended STAM performance. Mean $\pm$ SEM. n =

15 per peptide treatment. ****p < 0.0001. **(E)** Adult-only, neuron-specific RNAi knock-

down of *ins-17* leads to learning, STAM, and ITAM impairments compared to vector RNAi

treated animals. Mean $\pm$ SEM. n = 15 per RNAi treatment. ****p < 0.0001.

Fig. 3. INS-17 regulates cognitive aging phenotypes observed in *daf-2* mutants

(A) Day 5 adult *daf-2(e1370)* mutants display better learning and **(B)** memory behavior

compared to wild-type worms. Mean ± SEM. n = 10 per genotype. *p < 0.05, *** p < 0.001.

(C) Timeline of young adult peptide treatment. Wild-type L4 animals are fed SCR or INS-

17 iOP50 bacteria until Day 5 of adulthood when S/ITAM behavior is tested. INS-17

peptide treated worms have increased learning and **(D)** ITAM performance compared to

SCR peptide treated animals. Mean ± SEM. n = 15 per peptide treatment. *p < 0.05. **(E)**

Timeline of aged adult peptide treatment. Wild-type Day 3 adult animals are fed SCR or

INS-17 iOP50 bacteria until Day 5 of adulthood when S/ITAM behavior is tested. INS-17

peptide treated worms have increased learning performance, but not **(F)** ITAM

performance, compared to SCR peptide treated animals. Mean ± SEM. n = 15 per peptide

treatment. *p < 0.05, ****p < 0.0001. **(G)** Probability of survival analysis demonstrates

there is no significant difference in lifespan comparing wild-type animals and *ins-*

17(tm790) animals. One replicate depicted is representative of three replicates (see **Fig.**

S3F, S3G). Mean/median ± SEM. n ≥ 100 per replicate. ns, not significant (p > 0.05). **(H)**

Probability of survival analysis for wild-type animals treated with SCR peptide compared

to wild-type animals fed INS-17 peptide shows no significant difference in average

lifespan. One replicate depicted is representative of three replicates (see **Fig. S3H, S3I**).

Mean/median ± SEM. n ≥ 100 per replicate. ns, not significant (p > 0.05).

Fig. 4. INS-17 likely engages parallel pathways of DAF-2 receptor signaling

(A) *daf-16(mu86)* deletion mutants have impaired learning and memory behavior

compared to wild-type animals. Mean ± SEM. n = 15 per genotype. ****p < 0.0001. **(B)**

INS-17 peptide treatment rescues *daf-16(mu86)* S/ITAM impairments. Mean ± SEM. n =

15 per genotype and peptide treatment. Wild-type animals fed INS-17 performance

indices are in **Fig. S4C**. # is significance relative to wild-type. #p < 0.05, ##p < 0.01; ns,

not significant (p > 0.05). **(C)** *ist-1(ok2706)* deletion mutants display a slight but not

significant learning deficit and impaired S/ITAM behaviors compared to wild-type worms.

Mean ± SEM. n = 15 per genotype. *p < 0.05, **p < 0.01; ns, not significant (p > 0.05).

(D) INS-17 peptide treatment rescues *ist-1(ok2706)* learning and memory impairments.

Mean ± SEM. n = 15 per genotype and peptide treatment. # is significance relative to

wild-type. #p < 0.05, ##p < 0.01, ###p < 0.001, ****p< 0.0001; ns, not significant (p >

0.05). **(E)** *daf-16(mu86);ist-1(ok2706)* animals display significant learning and **(F)**

memory impairments that are not rescued by INS-17 peptide feeding, while *daf-16(mu86)*

and *ist-1(ok2706)* deficits are rescued by INS-17 peptide administration. Mean ± SEM. n

= 15 per genotype and peptide treatment *p < 0.05, **p < 0.01, ***p < 0.001; ns, not

significant (p > 0.05).

Fig. 5. INS-17 is regulated by nutrient deprivation to promote advantageous

behaviors

(A) Diagram showing *ins-17* replacement experiment design. *ins-17* and *ins-37* encode γ

-ILPs. *ins-17* was removed with CRISPR and replaced with *ins-37* at the *ins-17* locus so

ins-37 would be expressed at endogenous *ins-17* levels. **(B)** Replacing *ins-17* with *ins-*

37 has no significant effect on behavior compared to wild-type animals, while *ins-17*

CRISPR KO (*ins-17(knu1320)*) animals have impaired learning and S/ITAM. Mean ±

SEM. n = 15 per genotype. # is significance relative to wild-type. ####p < 0.0001; ns, not

significant (p > 0.05). **(C)** qRT-PCR of *ins-17* mRNA levels in Day 1 adult wild-type animals

show that one hour of food-deprivation (FD) results in increased *ins-17* transcript levels

(left). For levels measured after two hours of food-deprivation, please see **Fig. S5B.**

Schematic depicting food-deprivation experiment workflow (right): well-fed adult worms

are washed three times in M9 buffer solution to remove bacteria and placed on 100mm

agar plates either seeded with a bacterial lawn (non-FD) or no lawn (FD) prior to training

and behavior testing. Mean ± SEM. n = 5 per genotype. *$p < 0.05$. **(D)** Wild-type animals

display more robust ITAM behavior if a FD step is administered prior to 1 CS-US pairing

compared to training alone (non-FD conditions). Mean ± SEM. n = 15 per condition. *$p <$

0.05. **(E)** Food-deprivation prior to training (1 CS-US pairing) does not enhance *daf-*

2(e1370) ITAM performance. Mean ± SEM. n = 15 per condition. ns, not significant ($p >$

0.05). **(F)** INS-17 peptide treatment in wild-type animals is sufficient to promote robust

ITAM performance in non-FD conditions prior to training compared to non-FD SCR fed

animals. See **Fig. S5I** for food-deprived SCR and INS-17 peptide fed comparisons. Mean

± SEM. n = 15 per peptide treatment. **$p < 0.01$. **(G)** *ins-17(tm790)* mutants have

significantly impaired aversive associative learning and STAM performance compared to

wild-type worms. Absolute values of learning and memory performance indices are

reported. Mean ± SEM. n = 15 per genotype. ***$p < 0.001$, ****$p < 0.0001$; ns, not

significant ($p > 0.05$). **(H)** Putative diagram of how INS-17 is upregulated in the nervous

system to signal the valence of positive and aversive environmental conditions to inform

behavioral outputs advantageous for survival.

1009

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5