Modeling cetacean eDNA distribution along the Washington coast using metabarcoding from opportunistic samples and generalized additive models

Tania Valdivia-Carrillo[1]*, Megan Shaffer[2,3], Kim Parsons[2], Ally Im[3], Andrew Shelton[2], Eiren K. Jacobson[4], Abigail Wells[2], Ana Ramón-Laca[2], Krista M. Nichols[2], Ryan P. Kelly[3], Amy Van Cise[1]

[1] School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat St, Seattle, Washington, USA.

[2] Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Blvd E, Seattle, Washington, USA.

[3] School of Marine and Environmental Affairs, University of Washington, Seattle, Washington, USA.

[4] Centre for Research into Ecological and Environmental Modelling, University of St Andrews, St Andrews, Scotland, UK.

*Corresponding author: Tania Valdivia-Carrillo, School of Aquatic and Fishery Sciences, University of Washington, 1122 NE Boat St, Seattle, Washington, USA. Email: tania.valdiviac@gmail.com

Abstract

Effective cetacean conservation depends on robust monitoring, yet traditional visual and passive-acoustic surveys have constraints. We evaluated environmental DNA (eDNA) metabarcoding coupled with species distribution modeling (SDM) as a tool to study habitat use of cetaceans along the Washington State coast, USA. Seawater was collected at the surface and at a 50 m depth from 43 sites (86 samples) during the 2019 U.S.–Canada Integrated Ecosystem & Acoustic-Trawl Survey. A partial section of the mitochondrial control region was amplified with cetacean-specific primers, sequenced on an Illumina MiSeq, and taxonomically assigned with a curated reference database. Nine species were detected; we modelled the three most frequent: Pacific

white-sided dolphin (*Lagenorhynchus obliquidens*) (10 detections), humpback whale (*Megaptera*

novaeangliae) (8 detections), and Risso's dolphin (*Grampus griseus*) (6 detections). Binomial

generalized additive models related presence-absence to bathymetry, distance to shore,

longitude, slope, and sea-surface temperature; model performance was assessed with stratified

five-fold cross-validation. SDMs explained 17–51% of null deviance and presented high specificity

(≥ 0.80). The Pacific white-sided dolphin showed the highest eDNA presence probabilities

offshore, beyond the shelf break. Humpback whale eDNA presence probabilities showed hotspots

along the shelf break with secondary high-probability patches in near-shore waters. Risso's

dolphin eDNA presence probabilities were elevated in offshore zones characterized by steep

bathymetric gradients, particularly northwest of the sampled transect. These spatial patterns are

consistent with historical visual–acoustic records, suggesting that eDNA-informed SDMs can

capture cetacean habitat use. This proof of concept indicates that combining eDNA detections

with flexible SDMs could provide a cost-effective, non-invasive complement to conventional

surveys and may offer a scalable pathway for marine-mammal monitoring and spatial planning.

Keywords: eDNA metabarcoding, species distribution modeling, cetaceans, generalized additive

models

Introduction

Cetaceans play a critical role in marine ecosystems, yet their wide-ranging and often elusive

behavior makes detecting and monitoring them difficult (Kaschner et al., 2012). Understanding

and predicting their distribution is crucial for effective conservation and management in the face

of increasing anthropogenic pressures and environmental changes (Tobeña et al., 2016),

particularly for endangered or otherwise vulnerable species (Abrahms et al., 2019; Becker et al.,

2014; Breen et al., 2017; Derville et al., 2018; Salvadeo et al., 2010; Szesciorka et al., 2023).

Species distribution models (SDMs) have become valuable tools for linking species occurrences

with environmental variables, providing empirical descriptions and spatial predictions of species-

environment relationships (Elith & Leathwick, 2009). For cetaceans along the US West Coast,

SDMs based on systematic visual and acoustic surveys since 1991, have yielded estimates of

detection, abundance, or presence probabilities by incorporating relevant environmental variables

obtained from ocean models (Becker, Carretta, et al., 2020; Becker et al., 2010, 2014, 2016,

2017; Becker, Forney, et al., 2020; Fiedler et al., 2023). Within this region, the Washington State

coast, part of the California Current Ecosystem (CCE), hosts a diverse array of cetaceans,

including several species protected under the Endangered Species Act (United States, 1983).

The CCE is characterized by a broad continental shelf, shallow basins and a strong eastern-

boundary current that together create high productivity near the shelf and slope (Phillips et al.,

2023), underpinning a rich marine food web that supports these cetaceans (Rice et al., 2021).

Commercial shipping, fishing and naval exercises, however, pose risks to their populations

(Evans, 1996; Izquierdo-Serrano et al., 2022; Lewin et al., 2019; Pirotta et al., 2019; Tixier et al.,

2021). Although marine protected areas and speed-reduction zones have been implemented to

mitigate such impacts, evaluating their effectiveness for these highly mobile species requires a

thorough understanding of cetacean distribution patterns and habitat use (Bearzi, 2012; Nisi et

al., 2024; Sahri et al., 2021).

Monitoring these populations has relied primarily on visual and acoustic surveys, yet both

approaches face spatial and temporal constraints. Visual surveys provide fine spatial resolution

but depend on favorable weather conditions, specialized vessels, trained observers, and they

detect animals only when they surface within a limited viewing radius (Pikesley et al., 2012;

Vukelic et al., 2018). Passive acoustic methods can provide continuous temporal coverage but

are limited to vocally active species with described vocal repertoires (Kowarski et al., 2021) and

are often affected by ambient noise (Griffiths & Barlow, 2016). These limitations underscore the

need for alternative or complementary methods for cetacean monitoring.

Environmental DNA (eDNA) metabarcoding offers a powerful, non-invasive approach for

cetacean monitoring by detecting species-specific genetic signatures in seawater. After quality-

filtering, sequence reads are classified into amplicon sequence variants (ASVs), which capture

single-nucleotide differences comparable to haplotypes (Callahan et al., 2017; Porter &

Hajibabaei, 2020). ASVs are then matched against reference databases to assign taxonomy, a

workflow that has proved effective for a wide range of marine taxa (Poyntz-Wright et al., 2024;

Stat et al., 2018), including cetaceans (Boldrocchi et al., 2024; Merten et al., 2023; Visser et al.,

2021). Cetacean eDNA detections from depths exceeding 1000 m further demonstrate eDNA's

ability to study their habitat use throughout the water column. When coupled with SDMs, these

occurrence data link eDNA detections to environmental gradients, enabling prediction of habitat

use over unsampled areas.

Here, we combine targeted eDNA metabarcoding with generalized additive models (GAMs)-

based SDMs to map and predict the spatial distribution of cetacean eDNA along the Washington

State coast. Using seawater samples collected during a large-scale systematic fisheries survey,

we i) identify cetacean species present, ii) relate eDNA detection probabilities to environmental

covariates via GAM-based SDMs, and iii) produce spatially explicit predictions for the most

frequently detected taxa.

Material and Methods

eDNA Sampling

Water collection followed the fixed-station plan of the 2019 U.S.-Canada Integrated Ecosystem &

Acoustic-Trawl Survey for Pacific hake (de Blois, 2020). The 43 stations span 46.40° - 48.90° N

and -124.20° - -125.57° W, covering a coast-to-slope bathymetric gradient (inshore, shelf, slope)

and the latitudinal range of Wahington's continental shelf. Although opportunistic, this sampling

scheme provided systematic spatial coverage of the regions' cetacean habitats.

At each station we collected 2 samples, surface (0 m) and 50 m, for a total of 86 samples.

Sampling was carried out aboard the NOAA Ship *Bell M. Shimada* between 6 and 19 August, at

night to coincide with existing sampling protocols. For surface samples 2.5 L of seawater were

collected from the ship´s saltwater intake; 50 m samples were taken with a Niskin bottle on a

conductivity, temperature, and depth (CTD) rosette. Samples were filtered immediately through

sterile mixed cellulose ester filter membranes (1 µm pore size, 47 mm-diameter) using a vacuum

pump. Filters were stored in Longmire's buffer at room temperature and later DNA was extracted

following a modified phenol: chloroform method to increase yield (Ramón-Laca et al., 2021;

Sambrook et al., 1989). Detailed protocols for seawater collection, preservation, and eDNA

extraction are provided in Ramón-Laca et al. (2021).

Library preparation

We performed library preparation and sequencing in triplicate for each sample. We used a 2-step

PCR protocol with Dlp1.5-H and Oordlp4 primers (Baker et al., 2018) to target a 300-360 bp

segment of the cetacean mitochondrial control region. The first PCR amplification was conducted

using these primers with Illumina adapter overhang sequences, following the NEB Phusion HiFi

Taq protocol and cycling conditions (New England Biolabs). Primers and adapter sequences are

detailed in the Supplementary Material (**Table S1**). Each 18 µl PCR1 reaction included 9 µL 2X

Phusion Master Mix, 0.9 µL of each primer, 0.54 µL DMSO, 0.45 µL BSA, 4.44 µL nuclease-free

water, and 1.77 µL of eDNA. The following cycling conditions were initial denaturation of 98°C for

30 sec, followed by 45 cycles of denaturation (98°C for 10 sec), annealing (60°C for 30 sec), and

extension (72°C for 45 sec), with a final extension of 72°C for 10 min. After PCR amplification,

products were cleaned with 1X Agencourt Ampure XP Beads (Beckman Coulter), visualized on

agarose gels, and quantified using DS High Sensitivity Assay Kit (Qubit). Based on DNA quantity,

an equal sample mass of the PCR product was added to the indexing PCR reaction.

The second PCR round was performed to index samples uniquely, attaching a ten-base pairs tag

with a P5 or P7 MiSeq flow cell adapter via the Nextera overhang (**Table S1**), using IDT for

Illumina DNA/RNA UD indexes. Each 25 µl PCR2 reaction had 12.5 µL KAPA Hifi HotStart

ReadyMix, 1.25 µL of the unique IDT for Illumina DNA/RNA UD Index (one per sample), and up

to 11.25 µL of the bead-cleaned PCR1 product (volume adjusted based on normalization, with

the remaining volume filled with nuclease-free water). The cycling conditions were initial

denaturation of 95°C for 5 min, followed by eight cycles of denaturation (98°C for 20 sec),

annealing (56°C for 30 sec), and extension (72°C for 1 min), with a final extension of 72°C for 5

min. Following the index reaction, products were cleaned with 0.8X Agencourt Ampure XP beads.

The final bead-cleaned product was quantified using the DS High Sensitivity Assay Kit (Qubit)

and normalized to ensure equimolarity in the final library pool.

Libraries were sequenced on the Illumina MiSeq (v3 kit, 2x300 bp) at 7 pM with a 20% PhiX spike-

in at the University of Washington. Each of the three technical replicates was run separately. A

negative template control was included in each PCR1 to monitor contamination; all controls were

sequenced alongside field samples.

Contamination control

To minimize cross-contamination, we spatially separated pre-PCR, extraction, and post-PCR

workflows in different laboratories. Workbenches were decontaminated before and after each

session with DNA AWAY or 10% bleach solution followed by 70% ethanol. All pipettes and

consumables (plates, tips, and tubes) were exposed to UV light for 15 minutes before use. A

positive control template was processed in parallel to monitor for potential inhibition.

Bioinformatic Analysis

Raw sequence processing was performed in R and RStudio (R Core Team, 2021; RStudio Team, 2020) using custom pipeline. Primer removal was carried out using cutadapt v1.18 (Martin, 2011). Pair-end reads were trimmed, filtered, and merged using DADA2 (Callahan et al., 2016), with truncation lengths chosen by a sliding window quality (Phred ≥29 forward, ≥30 reverse) and a minimum overlap of 42 bp for merging paired-end sequences.

Taxonomy was assigned to D-loop ASVs using DADA2's naïve Bayesian classifier (*assignTaxonomy*) with a 98% bootstrap threshold (Wang et al., 2007), against a custom reference database including sequences from Baker et al. (2003).

After ASV taxonomic assignment, we collapsed counts to species-level read tallies. We then binarized detections by scoring a site as "present" when at least 1 read remained after all quality filters. We chose this ≥ 1-read rule to maximize sensitivity to rare taxa while stringent negative controls, chimera removal and high Phred scores minimizing false positives. The positive detections from three technical replicates and 2 depths were aggregated into a table for each species at each sampling location. We focused subsequent SDM analyses on the three most frequently detected cetacean species: Pacific white-sided dolphin, humpback whale, and Risso's dolphin.

Environmental predictors

Based on prior research in CCE (Becker, Carretta, et al., 2020; Becker et al., 2017; Fiedler et al., 2023), we selected five predictors: latitude, longitude, bathymetric data, distance to shore, slope, and the average seawater temperature (SST) for August 2019. Bathymetry, distance-to-shore, slope, latitude, and longitude rasters were sourced from MARSPEC (Sbrocco & Barber, 2013) via

the *sdmpredictors* R package in RStudio (Bosch & Fernandez, 2023; R Core Team, 2021;

RStudio Team, 2020). SST was obtained from Copernicus Global Ocean Physics Reanalysis

(2023). All raster layers were standardized to 0.083° grid and clipped to the Washington State

coastal polygon in QGIS (*Qgis.org*, 2024). Detailed variable descriptions can be found in

Supplementary Material **S2**.

We then built a raster brick employing the *raster* package in R and RStudio (Hijmans, 2024; R

Core Team, 2021; RStudio Team, 2020) and extracted values for each sampling site coordinate.

These covariate values were subsequently used to model the probability of eDNA occurrence of

individual cetacean species.

Cetacean species distribution models (SDMs) and cross-validation from eDNA detection

data

We modeled the probability of cetacean eDNA presence using binomial generalized additive

models (GAMs), which flexibly relate predictor variables to binary response data via smooth

functions (Wood, 2017). We fitted each smooth with thin plate regression splines (Wood, 2003)

to capture nonlinear effects of bathymetry, distance to shore, slope, and seawater temperature

(SST). The model for site *j* is:

$$Y_j \sim \text{Bernoulli}\,(\psi_j)$$

In the model, Y_j is a binary indicator of eDNA presence (1) or absence (0) at site *j*, and ψ_j is the

probability of detection. We linked ψ_j to environmental covariates using a logistic function:

$$\text{logit}(\psi_j) = \beta_0 + f_1(x_{1,j}) + f_2(x_{2,j}) + f_3(x_{3,j})$$

where β_0 is the intercept, and each f_i is a smooth thin-plate regression spline capturing potential

nonlinear effects of predictor x_i at site *j*.

Model selection and goodness-of-fit assessment

We fitted GAMs in R and RStudio (Kuhn, 2008; R Core Team, 2021; RStudio Team, 2020) using the *mgcv* and *MuMIn* packages. Each model incorporated 2-3 predictors, ensuring distance to shore and bathymetry did not co-occur (r> 0.8). We compared models using Akaike's Information Criteria (AIC) and Bozdogan's index of informational complexity (ICOMP).

Model fit was evaluated via adjusted R^2 and percent deviance explained. We used *gam.check*() function in the *mgcv* package (Wood, 2023) to inspect residuals and to verify that the basis dimension for each spline was adequate using the k-index diagnostic and residual variance tests (Wood, 2017 section 5.9). All GAMs were fitted with restricted maximum likelihood (REML) and the outer Newton optimizer, converging when changes in the REML criterion fell below a predetermined threshold. Final predictions for each species were generated from the best-supported model. The code for our modeling framework can be found on GitHub.

Cross-validation and performance evaluation

We assessed model performance with stratified k-fold cross-validation approach using the *caret* package in R and RStudio (Kuhn, 2008; R Core Team, 2021; RStudio Team, 2020). We divided the dataset into five folds (k = 5), each preserving the overall proportion of presences and absences to ensure performance estimates for the dataset. In each iteration, one-fold served as the test set and the remaining four as the training dataset. This approach was chosen because stratification better preserves the proportion of rare presences in each fold, providing more stable estimates of model performance in small, imbalanced datasets.

After obtaining probability predictions with the GAMs models for each target species, we converted them to binary presence/absence using a cut-off value of 0.3. We explored thresholds

from 0.1 to 0.5 and selected 0.3 because it maximized sensitivity while keeping false-positive rate

low, a priority given the small sample size and the low prevalence of detections. For each fold of

the stratified five-fold cross-validation, we calculated threshold-dependent metrics (i.e., accuracy,

sensitivity, specificity, and Cohen's κ) were computed via caret's confusion matrix and the

threshold-independent area under the receiver operating characteristic curve (AUC) with the

ROCR package in R and RStudio (R Core Team, 2021; RStudio Team, 2020; Sing et al., 2005).

We report the mean ± SD of each metric across the five folds as our final performance estimates.

Results

Sequencing results

We successfully amplified and sequenced 86 eDNA samples at 43 stations (0 m and 50 m),

conducting three independent MiSeq runs. We generated an average of 9,286,025 reads across

three replicate MiSeq runs (range: 8,534,845 - 9,857,378). Table **S3** in the Supplementary

Material provides detailed read counts at each step.

We identified nine cetacean species in total (<5% of the total detected ASVs; **Table 1**), with

species richness per site ranging from 0 to four species (**Fig.1**). Pacific white-sided dolphin,

humpback whale, and Risso's dolphin were the most frequently detected species (**Table 1**). In

total, 24 out of 43 sites had no detections for any of the three focal species.

Table 1. Summary of positive cetacean eDNA detections along the Washington State coast.
Counts refer to independent site-depth combinations (43 sites x 2 depths). Only taxa
detected at least once are shown.
231

Common name	Scientific name	Total detections
Pacific white-sided dolphin	*Lagenorhynchus obliquidens*	10
Humpback whale	*Megaptera novaeangliae*	8
Risso's dolphin	*Grampus griseus*	6
Dall's porpoise	*Phocoenoides dalli*	4
Northern right whale dolphin	*Lissodelphis borealis*	4
Killer whale	*Orcinus orca*	2
Baird's beaked whale	*Berardius bairdii*	1
Fin whale	*Balaenoptera physalus*	1
Harbour porpoise	*Phocoena phocoena*	1

232
Figure 1. Positive cetacean detections obtained through eDNA metabarcoding of seawater
samples. Open circles represent the sampling locations with no positive detections. The
fill of each circle represents the species detected at that sampling location; note that data
are represented as positive or negative detections for each species at each sampling
location, i.e., the proportion of the circle filled by each color does not represent the relative
sequence read count from that species in the sample. The dashed line is the 200 m isobath
and represents the shelf break.
240

Cetacean species distribution models (SDMs) from eDNA detection data

Pacific white-sided dolphin

The best-fitting GAM model retained two smooth predictors, longitude (p = 0.046) and distance to shore (p = 0.009), both significant (**Table 2**). Together they explained 47.7 % of the variation in eDNA presence on the link scale (adjusted R²) and reduced the null deviance by 50.6 %, indicating that half of the structure in the occurrence data is captured by these spatial variables. The model converged after five iterations, and k-index diagnostics confirmed adequate basis dimensions (k-index p_{lon} = 0.97; p_{dist_shore} = 0.95), providing no evidence of over- or under-smoothing.

Humpback whale

In the best-fitting GAM model for the humpback whale, eDNA bathymetry (p = 0.003) and sea-surface temperature (p = 0.026) were retained as significant smooth terms (**Table 2**). These predictors explained 28.5% of the variation in eDNA presence (adjusted R²) and reduced the null deviance by 29.2%, indicating that one-third of the spatial-environmental signal is captured. Convergence was reached after four iterations, and k-index diagnostics confirmed the adequate basis dimensions (k-index p_{bathy} = 0.52 and p_{SST} = 0.095) with no evidence of over- or under-smoothing.

Risso's dolphin

The best-fitting GAM model for Risso's dolphin retained longitude (p = 0.038) and slope (p = 0.053) as smooth terms (**Table 2**). These predictors accounted for 16.5% of the variation in eDNA presence (adjusted R²) and reduced the null deviance by 22.9%, suggesting a modest yet measurable improvement over the intercept-only model (null). The fit converged stably, and k-

index diagnostics confirmed adequate basis dimensions (k-index p_{lon} = 0.97; p_{slope} = 0.52),

suggesting no evidence of over- or under-smoothing.

Spatial patterns

Projecting these models onto the study area (**Fig.2**) revealed distinct habitat associations. Pacific

white-sided dolphin eDNA shows a higher probability of occurrence in offshore regions, beyond

the shelf break. Humpback whale eDNA showed higher probabilities along the shelf break with

localized coastal hotspots. Risso's dolphin eDNA was most probable in offshore regions with

steep bathymetric features and hotspots northwest of the sampled area.

Table 2 and **Fig.2** summarize the best GAM outputs and predicted eDNA presence probabilities,

respectively, illustrating an offshore distribution for Pacific white-sided dolphin and Risso's

dolphins, and shelf-break/coastal hotspots for humpback whales.

The Table **S4** and Figure **S5** of the Supplementary Material presents detailed information on the

modeling results.

Table 2. Best GAM models for the three more frequent cetacean species detected along the Washington coast via eDNA
metabarcoding. Models incorporate different environmental variables to predict eDNA-spatial distribution, including lat
(latitude), lon (longitude), bathy (bathymetry), dist_shore (distance to shore), slope, and seawater temperature (SST).
Statistical output parameters include R^2 (adjusted), and deviance explained. * p-value < 0.05
280

Species	(Int)	lat	lon	bathy	dist_shore	slope	SST	R^2(adj)	dev. explained
Pacific white-sided dolphin	0.003*		0.046*		0.009*			0.477	50.60
Humpback whale	0.000*			0.003*			0.026*	0.285	29.20
Risso's dolphin	0.000*		0.038*			0.053		0.165	22.90

281
282

Figure 2. eDNA presence probability derived from GAM models to evaluate eDNA species distribution for the a) Pacific white-sided dolphin, b) Humpback whale, and c) Risso's dolphin. The dashed line is the 200 m isobath and represents the shelf break.

288 **Cross-validation for eDNA-based models**

289 **Pacific white-sided dolphin**

290 The GAM for *Lagenorhynchus obliquidens* exhibited the highest predictive accuracy, with a mean

291 accuracy of 0.84 ± 0.19 and an area under the ROC curve (AUC) of 0.92 ± 0.17. Cohen's kappa

292 (κ = 0.61 ± 0.39) indicated substantial agreement beyond chance. Sensitivity, the true-positive

293 rate, averaged 0.75 ± 0.27, reflecting the model's strong ability to detect actual eDNA presences.

294 Specificity remained high at 0.87 ± 0.19, demonstrating reliable identification of absences. The

295 F_1-score of 0.72 ± 0.27 confirms a balanced trade-off between precision and recall.

296 **Humpback whale**

297 For *Megaptera novaeangliae*, the model yielded a mean accuracy of 0.80 ± 0.17 and AUC of 0.87

298 ± 0.22, indicative of good overall discrimination. Cohen's κ = 0.45 ± 0.45 denoted moderate

299 agreement beyond chance, with sensitivity of 0.70 ± 0.45 and specificity of 0.85 ± 0.18. The F_1-

300 score of 0.68 ± 0.25 illustrates that the model maintains a reasonable balance between detecting

301 true positives and minimizing false positives, despite increased variability associated with fewer

302 positive cases.

303 **Risso's dolphin**

304 The GAM for Grampus griseus faced limitations due to a small number of positive detections (n

305 = 6). It achieved a mean accuracy of 0.74 ± 0.15 but an AUC of only 0.62 ± 0.16, indicating modest

306 discriminative performance. Cohen's κ = 0.06 ± 0.32 suggests near-random agreement beyond

307 chance. Sensitivity was low (0.30 ± 0.45), while specificity remained comparatively robust at 0.80

308 ± 0.16. The F_1-score of 0.50 ± 0.24 further highlights the challenge of accurately predicting rare

309   eDNA occurrences. **Table 3** summarizes the mean ± SD of classification metrics across stratified

310   five-fold cross-validation for each species.

312 **Table 3. Summary of cross-validation results for three more frequent cetacean species detected along the Washington coast**
313 **via eDNA metabarcoding (threshold = 0.3). Values are reported as mean ± SD across folds.**
314

Species	Accuracy	Cohen's κ	Sensitivity	Specificity	F1	AUC
Pacific white-sided dolphin	0.84 ± 0.19	0.61 ± 0.39	0.75 ± 0.27	0.87 ± 0.19	0.72 ± 0.27	0.92 ± 0.17
Humpback whale	0.8 ± 0.17	0.45 ± 0.45	0.7 ± 0.45	0.85 ± 0.18	0.68 ± 0.25	0.87 ± 0.22
Risso's dolphin	0.74 ± 0.15	0.06 ± 0.32	0.3 ± 0.45	0.8 ± 0.16	0.5 ± 0.24	0.62 ± 0.16

Discussion

This study demonstrates the feasibility of integrating environmental DNA (eDNA) metabarcoding detection data into species distribution models (SDMs) to investigate cetacean spatial distributions along the Washington State coast. By applying generalized additive models (GAMs), we explored the relationship between eDNA presence and environmental variables for the three most frequently detected species: Pacific white-sided dolphin, humpback whale, and Risso's dolphin. Model performance was confirmed with stratified five-fold cross-validation, which showed consistently high values of accuracy, AUC, κ, and other diagnostics, indicating that the eDNA-based SDMs generalize well to unseen data for the Pacific white-sided dolphin and humpback whale, whereas performance for Risso's dolphin was limited by sparse detections.

It is important to clarify that the models predict the spatial distribution of eDNA detections, not the locations of individual cetaceans. Detected DNA could originate from animals that were present locally or from genetic material transported by currents, and the time elapsed since shedding is uncertain. Nevertheless, by mapping where cetacean genetic material is most likely to occur under prevailing environmental conditions, our approach offers a first step toward incorporating eDNA evidence into predictive frameworks for cetaceans, providing insights that can enhance conservation and management strategies.

Cetacean SDMs from eDNA detection data

Pacific white-sided dolphin

Pacific white-sided dolphins are pelagic species commonly found in Washington waters during the summer months. They take advantage of the productive high-latitude waters for foraging (Forney & Barlow, 1998; Rice et al., 2021). The CCE plays a vital role in shaping their distribution,

as nutrient upwelling enhances prey availability in offshore waters, particularly at the shelf break

and upper slope regions (Becker et al., 2022; Rice et al., 2021). Our model confirms these

findings, showing that the probability of detecting environmental DNA (eDNA) increases beyond

the 200-meter isobath. This aligns with previous studies identifying the continental shelf break as

a crucial habitat for dolphins (Giménez et al., 2021; Gómez-Campos et al., 2011).

Humpback whale

Humpback whales undertake long-distance migrations between tropical breeding grounds and

high-latitude summer feeding areas, including the coast of Washington (Baker et al., 1994; Barlow

et al., 2011; Darling et al., 2022; Wright et al., 2015). Their distribution is influenced by bathymetric

features and highly productive upwelling zones that concentrate prey (Derville et al., 2018;

Fleming et al., 2016; Garrigue et al., 2015; Thompson et al., 2012). Our model shows a strong

correlation between humpback whale eDNA presence and both bathymetry and SST. This pattern

is consistent with earlier studies that link bathymetry and SST to humpback foraging along shelf-

break regions (Becker, Carretta, et al., 2020; Burrows et al., 2016; Dalla Rosa et al., 2012;

Meynecke et al., 2021; Miller et al., 2022; Pelayo-González et al., 2022; Rockwood et al., 2020),

and our predicted eDNA hotspots coincide with shelf-break areas of high krill biomass (Phillips et

al., 2023), underscoring prey availability as driver of humpback whale distribution (Rockwood et

al., 2020). Consistently, both acoustic (Rice et al., 2021) and visual (Barlow & Forney, 2007;

Schorr et al., 2007) surveys report the highest humpback occurrence over the continental shelf

along the Washington coast.

Risso's dolphin

Risso's dolphins are typically found as offshore cetaceans, often observed near the outer edges

of continental shelves and upper slopes (Azzellino et al., 2008; Baumgartner, 1997; Cañadas et

al., 2002; Praca & Gannier, 2008). This habitat corresponds to areas where their primary prey,

cephalopods, are most abundant (Praca & Gannier, 2008; Visser et al., 2021). Our model

identified a significant association between the probability of detecting Risso's dolphin eDNA and

longitude, consistent with prior research highlighting the species' preference for offshore

environments (Becker, Carretta, et al., 2020; Becker, Forney, et al., 2020). Although slope was

non-significant in our model (p = 0.053), its exact role remains unclear. Higher-resolution

bathymetric data could clarify whether slope influences habitat selection by affecting prey

aggregation. Our results suggest that this study area may represent a transitional zone, where

Risso's dolphin presence declines at higher latitudes as water temperatures become more polar

(Frouin-Mouy et al., 2022; Jefferson et al., 2014).

Both Pacific white-sided dolphins and Risso's dolphins primarily forage on squid and exhibit

nocturnal activity. Our eDNA sampling at night likely coincided with an increase in surface

presence, which may enhance detection rates. This diel pattern aligns with previous studies

showing that delphinids engage in vertical migration in search of their prey (Benoit-Bird et al.,

2004; Benoit-Bird & Au, 2003).

Model robustness and limitations by cross-validation

Cross-validation serves as a critical tool for evaluating the robustness of models, particularly in

scenarios involving small datasets. The outcomes of stratified k-fold cross-validation for the three

focal species provide valuable insights into the predictive capacity and inherent limitations of our

eDNA-based SDMs. Stratified 5-fold cross-validation showed that our eDNA-based GAMs

capture meaningful spatial signal but also revealed limits imposed by a presence-rare data set.

Accuracy remained ≥ 0.74 and specificity ≥ 0.80 for all three taxa, indicating few false positives.

Metrics sensitive to the number of detections diverged: the Pacific white-sided dolphin model,

supported by the largest set of detections, showed the strongest and most reliable discrimination

(mean AUC = 0.92, κ = 0.61), whereas the humpback whale model was moderately robust (AUC

= 0.87) but variable across folds, reflecting data scarcity. With only six positive sites, the Risso's

dolphin model kept high specificity yet low sensitivity (mean 0.30) and near-random agreement

(κ = 0.06), highlighting the difficulty of modeling sparse eDNA records. Taken together, these

results suggest that eDNA-SDMs built from ≥ 10 detections provide reproducible spatial inference;

below that threshold, uncertainty increases. Also, even when presence data are limited, cross-

validation diagnostics, particularly sensitivity and κ, offer an objective benchmark for deciding

whether predictions are fit for management use or should be treated as exploratory.

Sampling-design and threshold considerations

All eDNA samples were collected opportunistically at the fixed stations of the Integrated

Ecosystem & Acoustic-Trawl Survey, which ensured a broad latitudinal and bathymetric coverage

but did under-represented shallow embayment (< 50 m) and outer-shelf waters. This sampling

may bias SDM predictions toward the environmental envelope of the survey transects.

Additionally, we adopted sensitive detection rules, both in the laboratory (≥ 1 read = presence)

and in model evaluation (0.30 probability cut-off), to minimize false negatives given the low

detection frequencies. Although contamination controls mitigate false positives, future studies

should integrate targeted eDNA stations in unsampled coastal and offshore regions and increase

biological replication to reduce spatial bias and increase detections.

Challenges of using eDNA data in SDMs

The interpretation of eDNA read counts presents several methodological and ecological

challenges that complicate their direct use in species distribution models (SDMs). While eDNA

metabarcoding enables species detection through high-throughput sequencing, read counts do

not necessarily reflect organism abundance due to various biological, technical, and

environmental factors. Given these complexities, we opted to transform our eDNA data into a

presence-absence format to improve model robustness and interpretability.

eDNA-based methods, though highly sensitive, share limitations with traditional monitoring

approaches (Fukaya et al., 2022; Mathieu et al., 2020). The amplification of specific DNA

molecules can be influenced by stochastic processes, including the limited availability of starting

material, which may lead to variability in detection outcomes (Kelly et al., 2019). Additionally,

detection probability is significantly affected by environmental conditions, sampling design, and

replication, particularly for rare or low-density species (Lahoz-Monfort et al., 2016). To minimize

underrepresentation of rare species, it is advisable to increase both biological and technical

replicates, as previous studies have demonstrated that replication enhances detection rates and

reduces bias (Alberdi et al., 2018; Buxton et al., 2021; Ficetola et al., 2015). The lack of biological

replication in this study may contribute to an underestimation of rare species' presence.

eDNA persistence and detectability in the environment vary due to multiple biological, physical,

and ecological factors (Brasseale et al., 2025). Shedding rates differ substantially across species

and are influenced by physiological and behavioral traits. Cetaceans, for example, release eDNA

through sloughed skin and feces, but the quantity shed is highly dependent on life-history traits,

body size, and health condition. Moreover, species-specific shedding rates fluctuate with

seasonality and behavior (Allan et al., 2020; Collins et al., 2018), further complicating the

interpretation of eDNA concentrations as direct indicators of abundance. Once released, eDNA

undergoes degradation at variable rates depending on environmental conditions such as

temperature, UV exposure, microbial activity, and water chemistry (Barnes et al., 2014). In cases

where degradation outpaces accumulation, the presence of a species may be underestimated,

while in other cases, eDNA may persist longer than expected, leading to false positives if not

carefully accounted for.

Another key consideration is the transport of eDNA in marine environments. Unlike terrestrial

systems, marine eDNA can be dispersed by currents, tides, and vertical mixing (Allan et al., 2021;

Andruszkiewicz et al., 2019; Brasseale et al., 2025). This movement complicates the

interpretation of read counts because high eDNA concentrations at a given location may not

necessarily indicate local species presence but rather the accumulation of transported genetic

material from upstream sources. Given this spatial complexity, a presence-absence framework

provides a more conservative and ecologically meaningful approach to SDMs.

As the scientific community improves its understanding of eDNA dynamics, we anticipate

advances in species detection accuracy, habitat modeling, and SDM precision. Further

developments, such as quantifying eDNA concentrations and linking them to species abundance

(e.g., Guri et al., 2024), could significantly enhance model performance and ecological inference.

The integration of eDNA data with comprehensive environmental datasets, including

hydrodynamic models, could allow for more precise estimations of species distributions while

accounting for eDNA transport and degradation processes.

Our study illustrates that eDNA metabarcoding can serve as a stand-alone tool for SDM

applications, particularly in scenarios where traditional survey methods (e.g., visual and acoustic

surveys) are impractical or resource intensive. While traditional observational datasets exist for

cetaceans in our study area (e.g., Becker et al., 2014), our research underscores the feasibility of

using eDNA metabarcoding to detect and model cetacean presence in a cost-effective, non-

invasive manner. However, this study serves as a proof of concept, and its statistical power is

inherently constrained by a small sample size. The continued evolution of eDNA methodologies,

combined with advancements in oceanographic modeling and environmental DNA transport

simulations, presents a clear opportunity for refining and expanding SDM applications. By

incorporating biotic and abiotic environmental predictors, as well as improved molecular

techniques, these models can be extended to a wide range of marine species, enhancing their

utility for conservation and management.

Conclusions

Our results demonstrate that environmental DNA metabarcoding, coupled with generalized

additive modeling, can generate ecologically coherent species-distribution estimates for highly

mobile cetaceans. The study was intentionally exploratory, with 43 stations and a limited number

of positive detections per species (1–10 occurrences) inherently reduce statistical power and

model generalizability. Even so, stratified k-fold cross-validation confirmed that models built from

as few as 10 positive sites can deliver reproducible spatial signal, whereas models with fewer

detections should be regarded as preliminary. The lower sensitivity observed for Risso's dolphins

underscores the challenges associated with modeling species with sparse detections,

emphasizing the need for increased sampling efforts across broader spatial and temporal scales.

A key challenge in eDNA-based SDMs is that they predict areas where cetacean DNA was

detected, rather than direct species presence. This distinction is particularly important in marine

systems, where eDNA transport via ocean currents, vertical mixing, and degradation can

influence detection locations. Future research should integrate hydrodynamic modeling to better

account for these transport processes, thereby improving spatial inferences of cetacean

distribution. Despite these challenges, the integration of eDNA detection data into SDMs presents

a significant opportunity for advancing cetacean monitoring and conservation. By expanding

sampling efforts, incorporating hydrodynamic models, and refining molecular methodologies,

future studies can enhance the accuracy and ecological relevance of eDNA-based distribution

models. This research sets the stage for the broader adoption of eDNA metabarcoding in marine

conservation, providing cost-effective, scalable, and non-invasive monitoring solutions for

cetaceans and other marine megafauna.

Data Archiving Statement

Raw fastq files will be uploaded to NCBI-Genbank and will be ready for publication. The code

used for our analysis is available on GitHub.

Author contribution

The collection of the samples was supported by the NOAA Fisheries Genomics Strategic Initiative,

led by KN. TVC, RK, AVC, and AS developed the ideas and designed the methodology for this

study. AW and ARL collected the eDNA samples and performed DNA extractions. MS conducted

laboratory experiments. AVC, AI, and TVC constructed the bioinformatic pipeline. TVC analyzed

the data obtained from bioinformatic analysis and led the manuscript writing. All authors critically

reviewed and contributed to the drafts and provided final approval for publication.

Data Archiving Statement

Raw fastq files will be uploaded to NCBI-Genbank and will be ready for publication. The code

used for our analysis is available in the GitHub repositories:

https://github.com/MMARINeDNA/metabarcoding_QAQC_pipeline and

https://github.com/MMARINeDNA/Valdiviaetal_2D-eDNA-spatial-distributions/tree/main, under

the CC0-1.0 license.

Acknowledgments

The authors gratefully acknowledge the survey expertise and support of NOAA's Pacific Hake

survey team and the NOAA Ship Bell M. Shimada personnel for supporting the collection of eDNA

samples on the West Coast Hake Survey 2019-2023.

Funding information

This material is based upon research supported by the Office of Naval Research under Award Number N00014-22-1-2719.

Conflict of interest statement

The authors declare that there is no conflict of interest.

References

Abrahms, B., Welch, H., Brodie, S., Jacox, M. G., Becker, E. A., Bograd, S. J., Irvine, L. M., Palacios, D. M., Mate, B. R., & Hazen, E. L. (2019). Dynamic ensemble models to predict distributions and anthropogenic risk exposure for highly mobile species. *Diversity & Distributions*, *25*(8), 1182–1193.

Alberdi, A., Aizpurua, O., Gilbert, M. T. P., & Bohmann, K. (2018). *Scrutinizing key steps for reliable metabarcoding of environmental samples*. *2018*(May 2017), 134–147.

Allan, E. A., DiBenedetto, M. H., Lavery, A. C., Govindarajan, A. F., & Zhang, W. G. (2021). Modeling characterization of the vertical and temporal variability of environmental DNA in the mesopelagic ocean. *Scientific Reports*, *11*(1), 21273.

Allan, E. A., Zhang, W., Lavery, A., & Govindarajan, A. F. (2020). *Environmental DNA shedding and decay rates from diverse animal forms and thermal regimes*. https://doi.org/10.1002/EDN3.141

Andruszkiewicz, E. A., Koseff, J. R., Fringer, O. B., Ouellette, N. T., Lowe, A. B., Edwards, C. A., & Boehm, A. B. (2019). Modeling Environmental DNA Transport in the Coastal Ocean Using Lagrangian Particle Tracking. *Frontiers in Marine Science*, *6*. https://doi.org/10.3389/fmars.2019.00477

Azzellino, A., Gaspari, S., Airoldi, S., & Nani, B. (2008). Habitat use and preferences of cetaceans along the continental slope and the adjacent pelagic waters in the western Ligurian Sea. *Deep Sea Research Part I: Oceanographic Research Papers*, *55*(3), 296–323.

Baker, C. S., Slade, R. W., Bannister, J. L., Abernethy, R. B., Weinrich, M. T., Lien, J., Urban, J.,

Corkeron, P., Calmabokidis, J., & Vasquez, O. (1994). Hierarchical structure of

mitochondrial DNA gene flow among humpback whales Megaptera novaeangliae, world-

wide. *Molecular Ecology*, *3*(4), 313–327.

Baker, C. S., Steel, D., Nieukirk, S., & Klinck, H. (2018). Environmental DNA (eDNA) From the

Wake of the Whales: Droplet Digital PCR for Detection and Species Identification.

Frontiers in Marine Science, *5*. https://doi.org/10.3389/fmars.2018.00133

Baker, S. C., Lavery, D. M. L., & Shane, R. H. A. (2003). www.DNA-surveillance: applied

molecular taxonomy for species conservation and discovery. *Trends in Ecology &*

Evolution, *18*(6), 271–272.

Barlow, J., Calambokidis, J., Falcone, E. A., Baker, C. S., Burdin, A. M., Clapham, P. J., Ford, J.

K. B., Gabriele, C. M., LeDuc, R., Mattila, D. K., Quinn, T. J., II, Rojas-Bracho, L., Straley,

J. M., Taylor, B. L., Urbán R., J., Wade, P., Weller, D., Witteveen, B. H., & Yamaguchi, M.

(2011). Humpback whale abundance in the North Pacific estimated by photographic

capture-recapture with bias correction from simulation studies. *Marine Mammal Science*,

27(4), 793–818.

Barlow, J., & Forney, K. (2007). *Abundance and population density of cetaceans in the California*

Current ecosystem. https://aquadocs.org/handle/1834/25509

Barnes, M. A., Turner, C. R., Jerde, C. L., Renshaw, M. A., Chadderton, W. L., & Lodge, D. M.

(2014). Environmental conditions influence eDNA persistence in aquatic systems.

Environmental Science & Technology, *48*(3), 1819–1827.

Baumgartner, M. F. (1997). The distribution of Risso's dolphin (*Grampus griseus*) with respect to

the physiography of the northern gulf of Mexico. *Marine Mammal Science*, *13*(4), 614–

638.

Bearzi, M. (2012). Cetaceans and MPAs should go hand in hand: A case study in Santa Monica

Bay, California. *Ocean & Coastal Management*, *60*, 56–59.

Becker, E. A., Carretta, J. V., Forney, K. A., Barlow, J., Brodie, S., Hoopes, R., Jacox, M. G.,

Maxwell, S. M., Redfern, J. V., Sisson, N. B., Welch, H., & Hazen, E. L. (2020).

Performance evaluation of cetacean species distribution models developed using

generalized additive models and boosted regression trees. *Ecology and Evolution*, *10*(12),

5759–5784.

Becker, E. A., Forney, K. A., Ferguson, M. C., Foley, D. G., Smith, R. C., Barlow, J., & Redfern,

J. V. (2010). Comparing California Current cetacean–habitat models developed using in

situ and remotely sensed sea surface temperature data. *Marine Ecology Progress Series*,

413, 163–183.

Becker, E. A., Forney, K. A., Fiedler, P. C., Barlow, J., Chivers, S. J., Edwards, C. A., Moore, A.

M., & Redfern, J. V. (2016). Moving Towards Dynamic Ocean Management: How Well Do

Modeled Ocean Products Predict Species Distributions? *Remote Sensing*, *8*(2), 149.

Becker, E. A., Forney, K. A., Foley, D. G., Smith, R. C., Moore, T. J., & Barlow, J. (2014).

Predicting seasonal density patterns of California cetaceans based on habitat models.

Endangered Species Research, *23*(1), 1–22.

Becker, E. A., Forney, K. A., Miller, D. L., Barlow, J., Rojas-Bracho, L., Urbán R, J., & Moore, J.

E. (2022). Dynamic Habitat Models Reflect Interannual Movement of Cetaceans Within

the California Current Ecosystem. *Frontiers in Marine Science*, *9*.

https://doi.org/10.3389/fmars.2022.829523

Becker, E. A., Forney, K. A., Miller, D. L., Fiedler, P. C., Barlow, J., & Moore, J. E. (2020). *Habitat-*

based density estimates for cetaceans in the California Current Ecosystem based on

1991-2018 survey data. https://repository.library.noaa.gov/view/noaa/27826

Becker, E. A., Forney, K. A., Thayre, B. J., Debich, A. J., Campbell, G. S., Whitaker, K., Douglas,

A. B., Gilles, A., Hoopes, R., & Hildebrand, J. A. (2017). Habitat-Based Density Models

for three Cetacean Species off Southern California Illustrate Pronounced Seasonal

Differences. *Frontiers in Marine Science*, *4*. https://doi.org/10.3389/fmars.2017.00121

Benoit-Bird, K. J., & Au, W. W. L. (2003). Prey dynamics affect foraging by a pelagic predator

(Stenella longirostris) over a range of spatial and temporal scales. *Behavioral Ecology and*

Sociobiology, *53*(6), 364–373.

Benoit-Bird, K. J., Würsig, B., & Mfadden, C. J. (2004). Dusky dolphin (*Lagenorhynchus obscurus*)

foraging in two different habitats: active acoustic detection of dolphins and their prey.

Marine Mammal Science, *20*(2), 215–231.

Boldrocchi, G., Conte, L., Galli, P., Bettinetti, R., & Valsecchi, E. (2024). Cuvier's beaked whale

(Ziphius cavirostris) detection through surface-sourced eDNA: A promising approach for

monitoring deep-diving cetaceans. *Ecological Indicators*, *161*(111966), 111966.

Bosch, S., & Fernandez, S. (2023). *sdmpredictors: Species Distribution Modelling Predictor*

Datasets (Version 0.2.15) [R package]. http://lifewatch.github.io/sdmpredictors/

Brasseale, E., Adams, N., Allan, E. A., Jacobson E. K., Kelly, R. P., Liu, O. R., Moore, S., Shaffer,

M., Xiong, J. and Parsons, K. 2025. Marine eDNA Production and Loss Mechanisms.

Journal of Geophysical Research. Oceans 130 (4): e2024JC021643.

Breen, P., Brown, S., Reid, D., & Rogan, E. (2017). Where is the risk? Integrating a spatial

distribution model and a risk assessment to identify areas of cetacean interaction with

fisheries in the northeast Atlantic. *Ocean & Coastal Management*, *136*, 148–155.

Burrows, J. A., Johnston, D. W., Straley, J. M., Chenoweth, E. M., Ware, C., Curtice, C., DeRuiter,

S. L., & Friedlaender, A. S. (2016). Prey density and depth affect the fine-scale foraging

behavior of humpback whales Megaptera novaeangliae in Sitka Sound, Alaska, USA.

Marine Ecology Progress Series, *561*, 245–260.

Buxton, A., Matechou, E., Griffin, J., Diana, A., & Griffiths, R. A. (2021). Optimising sampling and

analysis protocols in environmental DNA studies. *Scientific Reports*, *11*(1), 11637.

Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should replace

operational taxonomic units in marker-gene data analysis. *The ISME Journal*, *11*(12),

2639–2643.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*

Methods, *13*(7), 581–583.

Cañadas, A., Sagarminaga, R., & García-Tiscar, S. (2002). Cetacean distribution related with

depth and slope in the Mediterranean waters off southern Spain. *Deep Sea Research Part*

I: Oceanographic Research Papers, *49*(11), 2053–2073.

Collins, R. A., Wangensteen, O. S., Gorman, E. J. O., Mariani, S., Sims, D. W., & Genner, M. J.

(2018). Persistence of environmental DNA in marine systems. *Communications Biology*,

December. https://doi.org/10.1038/s42003-018-0192-6

Dalla Rosa, L., Ford, J. K. B., & Trites, A. W. (2012). Distribution and relative abundance of

humpback whales in relation to environmental variables in coastal British Columbia and

adjacent waters. *Continental Shelf Research*, *36*, 89–104.

Darling, J. D., Audley, K., Cheeseman, T., Goodwin, B., Lyman, E. G., & Urbán, R. J. (2022).

Humpback whales (Megaptera novaeangliae) attend both Mexico and Hawaii breeding

grounds in the same winter: mixing in the northeast Pacific. *Biology Letters*, *18*(2),

20210547.

de Blois, S. (2020). *The 2019 joint U.s.-Canada integrated ecosystem and pacific hake acoustic-*

trawl survey: Cruise report SH-19-06. Northwest Fisheries Science Center (U.S.).

https://doi.org/10.25923/P55A-1A22

Derville, S., Torres, L. G., Iovan, C., & Garrigue, C. (2018). Finding the right fit: Comparative

cetacean distribution models using multiple data sources and statistical approaches.

Diversity and Distributions, *24*(11), 1657–1673.

Elith, J., & Leathwick, J. R. (2009). Species Distribution Models: Ecological Explanation and

Prediction Across Space and Time. *Annual Review of Ecology, Evolution, and*

Systematics, *40*(1), 677–697.

Evans, P. G. H. (1996). Human disturbance of cetaceans. In *The Exploitation of Mammal*

Populations (pp. 376–394). Springer Netherlands.

Ficetola, G. F., Pansu, J., Bonin, A., Coissac, E., Giguet-Covex, C., De Barba, M., Gielly, L.,

Lopes, C. M., Boyer, F., Pompanon, F., Ray??, G., & Taberlet, P. (2015). Replication

levels, false presences and the estimation of the presence/absence from eDNA

metabarcoding data. *Molecular Ecology Resources*, *15*(3), 543–556.

Fiedler, P. C., Becker, E. A., Forney, K. A., Barlow, J., & Moore, J. E. (2023). Species distribution

modeling of deep-diving cetaceans. *Marine Mammal Science*.

https://doi.org/10.1111/mms.13057

Fleming, A. H., Clark, C. T., Calambokidis, J., & Barlow, J. (2016). Humpback whale diets respond

to variance in ocean climate and ecosystem conditions in the California Current. *Global*

Change Biology, *22*(3), 1214–1224.

Forney, K. A., & Barlow, J. (1998). Seasonal patterns in the abundance and distribution of

California cetaceans, 1991–1992. *Marine Mammal Science*, *14*(3), 460–489.

Frouin-Mouy, H., Mouy, X., Pilkington, J., Küsel, E., Nichol, L., Doniol-Valcroze, T., & Lee, L.

(2022). Acoustic and visual cetacean surveys reveal year-round spatial and temporal

distributions for multiple species in northern British Columbia, Canada. *Scientific Reports*,

12(1), 19272.

Fukaya, K., Kondo, N. I., Matsuzaki, S.-I. S., & Kadoya, T. (2022). Multispecies site occupancy

modelling and study design for spatially replicated environmental DNA metabarcoding.

Methods in Ecology and Evolution / British Ecological Society, *13*(1), 183–193.

Garrigue, C., Clapham, P. J., Geyer, Y., Kennedy, A. S., & Zerbini, A. N. (2015). Satellite tracking

reveals novel migratory patterns and the importance of seamounts for endangered South

Pacific humpback whales. *Royal Society Open Science*, *2*(11), 150489.

Giménez, J., Authier, M., Valeiras, J., Abad, E., Marçalo, A., Coll, M., Gauffier, P., Santos, M. B.,

& de Stephanis, R. (2021). Consumption rates and interaction with fisheries of

Mediterranean common dolphins in the Alboran Sea. *Regional Studies in Marine Science*,

45(101826), 101826.

Global Ocean Physics Reanalysis. (2023). [Dataset]. In *COPERNICUS*.

https://data.marine.copernicus.eu/product/GLOBAL_MULTIYEAR_PHY_001_030/descri

ption

Gómez-Campos, E., Borrell, A., Cardona, L., Forcada, J., & Aguilar, A. (2011). Overfishing of

small pelagic fishes increases trophic overlap between immature and mature striped

dolphins in the Mediterranean Sea. *PloS One*, *6*(9), e24554.

Griffiths, E. T., & Barlow, J. (2016). Cetacean acoustic detections from free-floating vertical

hydrophone arrays in the southern California Current. *The Journal of the Acoustical*

Society of America, *140*(5), EL399.

Guri, G., Shelton, A. O., Kelly R. P., Yoccoz, N., Johansen, T., Præbel, K., Hanebrekke, T., Ray,

J. L., Fall, J., and Westgaard J. I. 2024. Predicting Trawl Catches Using Environmental

DNA. *ICES Journal of Marine Science: Journal Du Conseil*, August, fsae097.

Hijmans, R. J. (2024). *raster: Geographic Data Analysis and Modeling*. https://rspatial.org/raster

Izquierdo-Serrano, M., Revuelta, O., Míguez-Lozano, R., Gozalbes, P., Ruiz-García, D., Raga, J.

A., & Tomás, J. (2022). Assessment of the interactions between cetaceans and fisheries

at the south of the Cetacean Migration Corridor and neighboring waters (Western

Mediterranean). *Frontiers in Marine Science*, *9*, 981638.

Jefferson, T. A., Weir, C. R., Anderson, R. C., Ballance, L. T., Kenney, R. D., & Kiszka, J. J.

(2014). Global distribution of Risso's dolphin Grampus griseus: a review and critical

evaluation. *Mammal Review*, *44*(1), 56–68.

Kaschner, K., Quick, N. J., Jewell, R., Williams, R., & Harris, C. M. (2012). Global coverage of

cetacean line-transect surveys: status quo, data gaps and future challenges. *PloS One*,

7(9), e44075.

Kelly, R. P., Shelton, A. O., & Gallego, R. (2019). Understanding PCR Processes to Draw

Meaningful Conclusions from Environmental DNA Studies. *Scientific Reports*, *9*(1), 12133.

Kowarski, K. A., Delarue, J. J.-Y., Gaudet, B. J., & Martin, S. B. (2021). Automatic data selection

for validation: A method to determine cetacean occurrence in large acoustic data sets.

JASA Express Letters, *1*(5), 051201.

Kuhn, M. (2008). Building Predictive Models inRUsing thecaretPackage. *Journal of Statistical*

Software, *28*(5), 1–26.

Lahoz-Monfort, J. J., Guillera-Arroita, G., & Tingley, R. (2016). Statistical approaches to account

for false-positive errors in environmental DNA samples. *Molecular Ecology Resources*,

16(3), 673–685.

Lewin, W.-C., Weltersbach, M. S., Ferter, K., Hyder, K., Mugerza, E., Prellezo, R., Radford, Z.,

Zarauz, L., & Strehlow, H. V. (2019). Potential environmental impacts of recreational

fishing on marine fish stocks and ecosystems. *Reviews in Fisheries Science &*

Aquaculture, *27*(3), 287–330.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.Journal, *17*(1), 10–12.

Mathieu, C., Hermans, S. M., Lear, G., Buckley, T. R., Lee, K. C., & Buckley, H. L. (2020). A

Systematic Review of Sources of Variability and Uncertainty in eDNA Data for

Environmental Monitoring. *Frontiers in Ecology and Evolution*, *8*.

https://doi.org/10.3389/fevo.2020.00135

Merten, V., GEOMAR Helmholtz Centre for Ocean Research Kiel, Visser, F., & Hoving, H.-J.

(2023). Deep-sea predator-prey dynamics revealed by biologging and eDNA analysis.

Oceanography (Washington, D.C.). https://doi.org/10.5670/oceanog.2023.s1.31

Merten, V. J., Visser, F., & Hoving, H.-J. T. (2023). Deep-Sea Predator-Prey Dynamics Revealed

by Biologging and eDNA Analysis. *Oceanography* , *36*(1), 96–99.

Meynecke, J.-O., de Bie, J., Barraqueta, J.-L. M., Seyboth, E., Dey, S. P., Lee, S. B., Samanta,

S., Vichi, M., Findlay, K., Roychoudhury, A., & Mackey, B. (2021). The role of

environmental drivers in humpback whale distribution, movement and behavior: A review.

707   *Frontiers in Marine Science*, *8*, 720774.

708 Miller, D. L., Becker, E. A., Forney, K. A., Roberts, J. J., Cañadas, A., & Schick, R. S. (2022).

709   Estimating uncertainty in density surface models. *PeerJ*, *10*, e13950.

710 New England Biolabs. (n.d.). *PCR Protocol for Phusion® High-Fidelity DNA Polymerase*.

711   Retrieved February 20, 2024, from https://www.neb.com/en-us/protocols/0001/01/01/pcr-

712   protocol-m0530

713 Nisi, A. C., Welch, H., Brodie, S., Leiphardt, C., Rhodes, R., Hazen, E. L., Redfern, J. V., Branch,

714   T. A., Barreto, A. S., Calambokidis, J., Clavelle, T., Dares, L., de Vos, A., Gero, S.,

715   Jackson, J. A., Kenney, R. D., Kroodsma, D., Leaper, R., McCauley, D. J., … Abrahms,

716   B. (2024). Ship collision risk threatens whales across the world's oceans. *Science (New*

717   *York, N.Y.)*, *386*(6724), 870–875.

718 Pelayo-González, L., Herra-Miranda, D., Pacheco-Polanco, J. D., Guzmán, H. M., Goodman, S.,

719   & Oviedo, L. (2022). Decreases in encounter rate of endangered Northeast Pacific

720   humpback whales in Southern Costa Rica: Possible changes in migration pattern due to

721   warming events. *Frontiers in Marine Science*, *9*.

722   https://doi.org/10.3389/fmars.2022.927276

723 Phillips, E. M., Malick, M. J., Gauthier, S., Haltuch, M. A., Hunsicker, M. E., Parker-Stetter, S. L.,

724   & Thomas, R. E. (2023). The influence of temperature on Pacific hake co-occurrence with

725   euphausiids in the California Current Ecosystem. *Fisheries Oceanography*, *32*(3), 267–

726   279.

727 Pikesley, S. K., Witt, M. J., Hardy, T., Loveridge, J., Loveridge, J., Williams, R., & Godley, B. J.

728   (2012). Cetacean sightings and strandings: evidence for spatial and temporal trends?

729   *Journal of the Marine Biological Association of the United Kingdom. Marine Biological*

730   *Association of the United Kingdom*, *92*(8), 1809–1820.

731 Pirotta, V., Grech, A., Jonsen, I. D., Laurance, W. F., & Harcourt, R. G. (2019). Consequences of

732   global shipping traffic for marine giants. *Frontiers in Ecology and the Environment*, *17*(1),

39–47.

Porter, T. M., & Hajibabaei, M. (2020). Putting COI Metabarcoding in Context: The Utility of Exact

Sequence Variants (ESVs) in Biodiversity Analysis. *Frontiers in Ecology and Evolution*, *8*.

https://doi.org/10.3389/fevo.2020.00248

Poyntz-Wright, I. P., Harrison, X. A., Pedersen, S., & Tyler, C. R. (2024). Effectiveness of eDNA

for monitoring riverine macroinvertebrates. *The Science of the Total Environment*,

941(173621), 173621.

Praca, E., & Gannier, A. (2008). Ecological niches of three teuthophageous odontocetes in the

northwestern Mediterranean Sea. *Ocean Science*, *4*(1), 49–59.

Qgis.org. (2024). QGIS Geographic Information System. Open Source Geospatial Foundation

Project. https://qgis.org/

R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Ramón-Laca, A., Wells, A., & Park, L. (2021). A workflow for the relative quantification of multiple

fish species from oceanic water samples using environmental DNA (eDNA) to support

large-scale fishery surveys. *PloS One*, *16*(9), e0257773.

Rice, A., Debich, A. J., Širović, A., Oleson, E. M., Trickey, J. S., Varga, L. M., Wiggins, S. M.,

Hildebrand, J. A., & Baumann-Pickering, S. (2021). Cetacean occurrence offshore of

Washington from long-term passive acoustic monitoring. *Marine Biology*, *168*(8), 1–22.

Rockwood, R. C., Elliott, M. L., Saenz, B., Nur, N., & Jahncke, J. (2020). Modeling predator and

prey hotspots: Management implications of baleen whale co-occurrence with krill in

Central California. *PloS One*, *15*(7), e0235603.

RStudio Team. (2020). *RStudio: Integrated Development Environment for R*. RStudio, PBC.

http://www.rstudio.com/

Sahri, A., Herwata Putra, M. I., Kusuma Mustika, P. L., Kreb, D., & Murk, A. J. (2021). Cetacean

habitat modelling to inform conservation management, marine spatial planning, and as a

basis for anthropogenic threat mitigation in Indonesia. *Ocean & Coastal Management*,

205(105555), 105555.

Salvadeo, C. J., Lluch-Belda, D., Gómez-Gallardo, A., Urbán-Ramírez, J., & MacLeod, C. D.

(2010). Climate change and a poleward shift in the distribution of the Pacific white-sided

dolphin in the northeastern Pacific. *Endangered Species Research*, *11*, 13–19.

Sambrook, J., Fritsch, E. F., & Maniatis, T. (1989). *Molecular cloning: a laboratory manual* (2d

ed.). Cold Spring Harbor Laboratory.

Sbrocco, E. J., & Barber, P. H. (2013). MARSPEC: ocean climate layers for marine spatial

ecology. *Ecology*, *94*(4), 979–979.

Schorr, G., Falcone, E., Calambokidid, J., Hildebrand, J., & Oleson, E. (2007). *Acoustic and visual*

monitoring for cetaceans along the outer Washington coast.

https://calhoun.nps.edu/bitstream/handle/10945/704/NPS-OC-09-001.pdf?sequence=3

Shelton, A. O., Gold, Z. J., Jensen, A. J., D Agnese, E., Andruszkiewicz Allan, E., Van Cise, A.,

Gallego, R., Ramón-Laca, A., Garber-Yonts, M., Parsons, K., & Kelly, R. P. (2023).

Toward quantitative metabarcoding. *Ecology*, *104*(2), e3906.

Sing, T., Sander, O., Beerenwinkel, N., & Lengauer, T. (2005). ROCR: visualizing classifier

performance in R. *Bioinformatics (Oxford, England)*, *21*(20), 3940–3941.

Stat, M., John, J., DiBattista, J. D., Newman, S. J., Bunce, M., & Harvey, E. S. (2018). Combined

use of eDNA metabarcoding and video surveillance for the assessment of fish biodiversity.

Conservation Biology: The Journal of the Society for Conservation Biology, *0*(0), 1–10.

Szesciorka, A. R., Demer, D. A., Santora, J. A., Forney, K. A., & Moore, J. E. (2023). Multiscale

relationships between humpback whales and forage species hotspots within a large

marine ecosystem. *Ecological Applications: A Publication of the Ecological Society of*

America, *33*(2), e2794.

Thompson, S. A., Sydeman, W. J., Santora, J. A., Black, B. A., Suryan, R. M., Calambokidis, J.,

Peterson, W. T., & Bograd, S. J. (2012). Linking predators to seasonality of upwelling:

Using food web indicators and path analysis to infer trophic connections. *Progress in*

Oceanography, *101*(1), 106–120.

Tixier, P., Lea, M.-A., Hindell, M. A., Welsford, D., Mazé, C., Gourguet, S., & Arnould, J. P. Y.

(2021). When large marine predators feed on fisheries catches: Global patterns of the

depredation conflict and directions for coexistence. *Fish and Fisheries (Oxford, England)*,

22(1), 31–53.

Tobeña, M., Prieto, R., Machete, M., & Silva, M. A. (2016). Modeling the potential distribution and

richness of cetaceans in the Azores from fisheries observer program data. *Frontiers in*

Marine Science, *3*, 202655.

United States. (1983). *The Endangered Species Act as Amended by Public Law 97-304 (the*

Endangered Species Act Amendments of 1982). Washington :U.S. G.P.O.

https://www.govinfo.gov/content/pkg/COMPS-3002/pdf/COMPS-3002.pdf

Visser, F., Merten, V. J., Bayer, T., Oudejans, M. G., de Jonge, D. S. W., Puebla, O., Reusch, T.

B. H., Fuss, J., & Hoving, H. J. T. (2021). Deep-sea predator niche segregation revealed

by combined cetacean biologging and eDNA analysis of cephalopod prey. *Science*

Advances, *7*(14). https://doi.org/10.1126/sciadv.abf5908

Vukelic, M., Mancini, F., Vukelic, D., & Carere, C. (2018). A cetacean monitoring system that

integrates citizen science and satellite imagery. *Rendiconti Lincei. Scienze Fisiche e*

Naturali, *29*(1), 53–59.

Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. *Applied and*

Environmental Microbiology, *73*(16), 5261–5267.

Wood, S. N. (2003). Thin plate regression splines. *Journal of the Royal Statistical Society. Series*

B, Statistical Methodology, *65*(1), 95–114.

Wood, S. N. (2008). Fast stable direct fitting and smoothness selection for generalized additive

models. *Journal of the Royal Statistical Society. Series B, Statistical Methodology*, *70*(3),

495–518.

Wood, S. N. (2017). *Generalized additive models: An introduction with R, second edition*. CRC

Press.

Wood, S. N. (2023). *mgcv: Mixed GAM Computation Vehicle with Automatic Smoothness

Estimation* (Version 1.9-1). https://cran.r-project.org/web/packages/mgcv/index.html

Wright, D. L., Witteveen, B., Wynne, K., & Horstmann-Dehn, L. (2015). Evidence of two

subaggregations of humpback whales on the Kodiak, Alaska, feeding ground revealed

from stable isotope analysis. *Marine Mammal Science*, *31*(4), 1378–1400.