Time-resolved bottlenecks reveal concurrent pathogen entry and growth during bacterial infection dynamics

Sylvain Vicente[1]* (ID), Nathanaël Boutillon[2,3,4] (ID), Raphael Forien[2] (ID), Isabelle Mila[1] (ID), Lionel Roques[2] (ID), and Nemo Peeters[1]* (ID)

[1] *Université de Toulouse, INRAE, CNRS, LIPME, Castanet-Tolosan, France*

[2] *INRAE, BioSP, Avignon 84914, France*

[3] *Aix Marseille Université, CNRS, I2M, Marseille, France*

[4] *Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, Wien, 1090, Austria*

* *Correspondence and material requests: sylvain.vicente@inrae.fr, nemo.peeters@inrae.fr*

2026/01/16

Abstract

Within-host pathogen dynamics are shaped by successive host-imposed barriers that generate bottlenecks and ultimately determine infection outcome, yet these processes remain difficult to study *in vivo*. Here, we use neutral genetic barcoding to quantify the root-entry bottleneck during infection of tomato by the bacterial soil-borne phytopathogen *Ralstonia pseudosolanacearum*. We establish an updated theoretical framework showing that the temporal structure of pathogen entry, often overlooked, is a key determinant of root colonisation. We introduce the concept of time-resolved bottleneck, capturing infection scenarios in which entry occurs continuously while the pathogen population simultaneously expands within the host. We develop quantitative tools that exploit barcode dynamics to disentangle the respective roles of motility, growth, and spatial competition, enabling direct inference of *in vivo* pathogen fluxes and growth rates and linking host susceptibility to mechanistic features of colonisation. Extending beyond plants, reanalysis of two independent animal infection datasets reveals previously unrecognised within-host development patterns. Together, these results show that time-resolved bottleneck is a general principle of bacterial infection dynamics and propose a new standard for barcoding-based studies: a host- and pathogen-agnostic framework that is directly applicable to a majority of existing STAMP(-R)-style datasets, distributed as an open, annotated R/Python notebook to facilitate community reuse.

Understanding the spatiotemporal dynamics of pathogens within their hosts is essential to explaining infection success. Host–pathogen molecular interactions and the mechanisms governing tissue colonisation are ultimately manifested in the growth, spread, and structuring of pathogen populations *in vivo*. Despite major advances in elucidating the molecular determinants of bacterial virulence, quantitatively and descriptively capturing infection dynamics within the host remains challenging, limiting our ability to link molecular mechanisms to population-level outcomes. The STAMP (Sequence Tag-Based Analysis of Microbial Population dynamics) framework[1] and its recent extensions[2,3] have been effective in monitoring the dynamics of the infecting population and identifying bottlenecks in different host-bacterial pathosystems[4]. The overarching goals of such approaches are to identify the locations of host-imposed barriers, quantify their restrictive strength against invading pathogens, and ultimately predict host survival or mortality under defined conditions.

STAMP experiments introduce neutral genetic diversity into pathogen populations by inserting short sequence tags into the genome. These tagged populations behave indistinguishably from wild-type strains, while the engineered neutral diversity acts as a heritable record of the population's genetic history during infection. A primary application of this approach is the identification and quantification of infectious bottlenecks, during which pathogen populations undergo severe reductions in size that are revealed by characteristic losses of neutral genetic diversity. The bottleneck was originally defined as *number of founding pathogen cells* that cross the barrier and expand[1]. Although many recent studies have now been published using this type of indicators (see review[4] and references within), the link between pathogen population patterns measured through STAMP and the nature of the underlying biological mechanisms so far remains unclear.

Here, we address this theoretical gap by applying STAMP to infections of tomato by the phytopathogen *Ralstonia pseudosolanacearum*. *R. pseudosolanacearum* is a soil-borne bacterium that causes bacterial wilt, a devastating disease of solanaceous crops. Infection initiates at the root, where the pathogen breaches host defences and rapidly colonises the xylem, leading to vascular occlusion, wilting, and ultimately plant death. Previous work has revealed the presence of localised host barriers along the infection route, including physical defences in the root, which impose strong constraints on pathogen entry and spread.[5,6,7,8].

We demonstrate that infection is associated with a stringent root-entry bottleneck, such that only a very small fraction of the invading pathogen population successfully establishes in the xylem. Although this bottleneck is detected by both the original STAMP framework[1] and its algorithmically refined implementation STAMPR[2], neither approach captures the fine-scale spatio-temporal patterns of root colonisation revealed by our data. By revisiting the theoretical basis of the classical definition of a bottleneck as a fixed "number of founding pathogen cells", we show that this formulation is valid only under restrictive conditions. Critically, it fails to account for infection mechanisms such as motility, *in situ* growth, and spatial competition, which are central to host colonisation. Instead, our results support a scenario in which pathogen entry, proliferation, and dispersal occur concurrently. We formalise this as a *time-resolved bottleneck model*, in which bottlenecks emerge from explicitly time-dependent

population dynamics.

To formalise this conceptual shift, we introduce a new analytical framework for neutral barcoding

data. In addition to an *instantaneous bottleneck* model corresponding to the original STAMP frame-

work[1], we develop three *time-resolved* models grounded in population genetics. Experimental datasets

were systematically evaluated against each model to identify the infection regime that best explains

the observed barcode dynamics, providing qualitative insight into the underlying mode of infection.

Quantitative metrics can then be extracted to directly characterise the processes governing infection,

including pathogen entry, growth, and competition.

Consistently, *time-resolved* models provided a markedly better fit to the plant infection data, indi-

cating that pathogen populations begin to expand immediately upon host entry and that root barriers

likely relax over the course of infection. The metrics derived from this framework explicitly capture the

temporal dimension of infection, yielding quantitative estimates of pathogen influx into the host and

in planta growth rates at the level of individual plants. Both parameters strongly correlate with host

susceptibility, demonstrating that pathogen motility and growth dynamics are integral determinants

of susceptibility rather than secondary consequences of infection.

To assess the generality of our framework, we reanalyse published animal infection datasets[9,10] and

uncover analogous time-resolved dynamics in *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* in-

fections in mice. These reanalyses serve as a stringent benchmark for the performance of our approach

and demonstrate its ability to infer pathogen influxes and *in situ* growth rates at the level of individual

organs within individual hosts. Strikingly, this quantitative dissection reveals a previously overlooked

trait in that *P. aeruginosa* colonisation is highly heterogeneous across the host, being driven by exten-

sive dispersal among systemically connected organs (lung, spleen and liver), while colonisation of the

gastrointestinal tract (gallbladder, stomach, small intestine, caecum and colon) is dominated by local

population growth.

To maximise accessibility and reproducibility, models analysis and outputs are provided along with a

fully annotated R computational document, enabling straightforward reuse, adaptation, and extension

of the methodology for the analysis of neutral barcoding experiments across diverse host–pathogen

systems.

1   Results

1.1   Physical barriers in plant roots impose a stringent pathogen-limiting bottleneck

To investigate the infection dynamics of *Ralstonia pseudosolanacearum*, we constructed a library of 119

neutral genetic tags in the reference strain GMI1000. Co-cultures of this tagged collection were used to

implement the STAMP approach[1] , enabling a quantitative analysis of the earliest stages of pathogen

colonisation within the plant host (Fig. 1a). We first asked whether root defences impose a population-

limiting barrier on *R. pseudosolanacearum*. We therefore formulated the following prediction: if a
bottleneck occurs at root entry, tag diversity should decrease between inoculum and roots then remain
stable once the pathogen has successfully colonised host tissues.

To test this hypothesis, susceptible tomato plants (*cv.* Marmande) were inoculated with the 119-
tags co-culture and sampled between 2-6 days post-inoculation (DPI). At each day, entire taproots
were harvested for colony-forming unit (CFU) quantification (Fig. 1b) and tag sequencing to compute
Shannon diversity (Fig. 1c). Bacterial loads increased steadily over time, consistent with active growth
and progressive host colonisation. In line with our prediction, the mean Shannon diversity converged
to ~ 2 by 4-6 DPI, indicating establishment and stability of pathogen diversity over time once the
population was well-established. Notably, *in planta* diversity was consistently lower than that of the
inoculum, demonstrating that a 119-tag collection reliably captured a reduction in pathogen diversity
at roots level and providing strong evidence for the existence of a root-entry bottleneck.

To assess whether physical root barriers underlie this bottleneck, we performed a second exper-
iment comparing plants with intact roots systems to plants whose roots were mechanically scarified
immediately prior to inoculation. Control Marmande tomato plants were infected as described above
with the 119-tag co-culture. For each condition, three independent replicates of $n = 8$ plants were
harvested at the onset of first disease symptoms within each experimental batch in order to collect
samples at a non-saturation phase of the infection (Supplementary Fig. 1). Root samples were pro-
cessed for CFU and diversity (Figs. 2a,2b). Control intact Marmande tomato plants with intact roots
displayed behaviour similar to that observed in the first experiment, both in terms of CFU accumu-
lation and overall infection dynamics. Intact root plants exhibit a pronounced reduction in diversity
relative to the inoculum, reflected by losses in both richness (tag extinction) and evenness (frequency
imbalance, Fig. 2c). By contrast, plants with cut roots showed significantly higher Shannon diversity
at entry (Fig. 2d), indicating enhanced pathogen ingress. Together, these results demonstrate that
the physical integrity of roots contributes to the root-entry bottleneck, in agreement with previous
qualitative reports showing that root structure impedes early infection[5,6,7,8].

1.2   Diversity profiles encode previously unexploited information about the pathogen's mode of entry

Observed diversity profiles are inconsistent with STAMP/STAMPR frameworks. As a
first approach, we quantified the root-entry bottleneck using published metrics. We computed the
standard bottleneck metric N_b from the original STAMP framework[1] and both corrected (N_r) and
simulated (N_s) bottlenecks from STAMPR[2]. Although derived differently, these metrics share the same
interpretation of a bottleneck as the "number of founding pathogen cells". To evaluate robustness, we
compared each estimate to the sample richness R, defined here as the minimum number of founders
required to generate the observed set of tags R in a given sample (Fig. 3a).

Median richness was $R \approx 30$ tags in intact plants and $R \approx 110$ in cut-root plants. Unexpectedly,

all tested bottleneck estimators exhibit some degree of inconsistency with the observed tag counts. The original STAMP metric N_b performed poorly, markedly underestimating entry given the number of detected tags (intact: median $N_b \approx 3$; cut: median $N_b \approx 14$). in contrast, the STAMPR estimators (N_r and N_s) tracked richness more closely overall, although not uniformly across conditions (intact: median $N_r \approx 12$; cut: median $N_r \approx 300$). Because N_r, appeared to be the most consistent estimator, we next tested, sample per sample, whether it could account for the observed richness (Fig. 3b). In most intact-root plants (12/17) and in a substantial subset of cut-root plants (6/24), we found $N_r < R$ indicating that systematic discrepancies are commonly. Notably, these failures clustered below a richness threshold of roughly ~ 60 tags, a regime that was not highlighted previously. Inspection of individual diversity profiles (Figs. 2) further suggest a biological origin: both N_b and N_r over-represent high-frequency tags and effectively discount low-frequency tags. This implies that standard bottleneck metrics capture only part of the colonisation process, and can miss biological meaningful complexity carried by the long tail of rarer lineages.

These discrepancies led us to re-evaluate the biological meaning of N_b-derived indicators in our infection system. Revisiting the population-genetic theory underlying these estimators[11,12] clarified that the N_b equation[1] and the derived N_r and N_s[2] are rooted in a very specific regime: barrier crossing is effectively instantaneous and decoupled from subsequent within-host growth. We refer to this class as *instantaneous bottleneck models*, which implicitly assume that a single founder event occurs over a short time window and is independent of time (Fig. 3c). A concrete representative of this class is the STAMP model ($\mathcal{M}_{\mathrm{STAMP}}$, Supplementary Fig. 3), which is sufficiently explicit to allow mathematical analysis. By contrast, N_r and N_s are computational corrections of N_b. Beyond empirical validation, they offer limited analytical tractability, making it difficult to probe their assumptions or failure modes more deeply.

Investigating time-dependent effects is necessary for a better understanding of pathogen colonisation. Unlike non-parametric summaries (*e.g.*, Shannon diversity, richness), bottleneck estimators such as N_b and N_r are only meaningful in so far as their underlying model assumptions are satisfied. Yet, for *instantaneous bottleneck models*, few practical tools exist to formally assess whether these assumptions are satisfied in a given experimental system. This motivates considering alternative mechanistic descriptions that may better reproduce the observed data. In particular, prior work suggests that *R. pseudosolanacearum* entry proceeds via a sustained influx rather than a brief, short-lived founder event[5,6,7,8]. Consistent with this view, tag frequencies within samples are not close to the homogeneous compositions expected under instantaneous entry. Instead, diversity profiles typically display pronounced gradients, with a few highly represented tags followed by a long tail of low frequency lineages (Supplementary Fig. 2), suggesting that founding and subsequent growth are intertwined over time.

If entry and growth proceed concurrently, bacteria that enter earlier should, on average, attain higher frequencies than those entering later. This motivates a class of *time-resolved bottleneck models*

in which founding overlaps with within-host expansion (Fig. 3d). rather than relying on a single summary metric, our goal is to quantify the contribution of explicit time-dependent biological processes (motility, spatial competition, growth) that cannot be interrogated within the STAMP(-R) paradigm. To challenge the instantaneous-bottleneck assumption, we introduce three alternative models grounded in distinct population genetics perspectives (detailed in Supplementary Fig. 3):

- **Constant-entry-rate model** ($\mathcal{M}_{\mathrm{CR}}$): pathogen cells enter the root at a constant rate and subsequently grow exponentially.

- **Bolthausen-Sznitman model** ($\mathcal{M}_{\mathrm{BS}}$): root colonisation is framed as a stochastic range expansion that captures diffusion and spatial occupation within the root. The resulting genealogies follow a Bolthausen-Sznitman coalescent[13].

- **Increasing-entry-rate model** ($\mathcal{M}_{\mathrm{IR}}$): a generalisation of $\mathcal{M}_{\mathrm{CR}}$ in which entry rates increase exponentially over time; while cells still grow exponentially. This model captures the progressive breaching of the host barriers as observed in $\mathcal{M}_{\mathrm{BS}}$ (Supplementary Fig. 4), while retaining a simpler, more directly interpretable mechanism.

1.3 A model-testing framework reveals new biological insights into *in planta* infection dynamics

We present a model-based procedure for analysing neutral barcoding data in which the shape of the within-host diversity profiles is treated as a diagnostic signature of the infection dynamics. The rationale is straightforward: if diversity profiles encode the mode and timing of entry, then the most plausible bottleneck mechanism can be identified by confronting observed profiles with those predicted by mechanistic models. For each sample, we simulate tag-frequency distributions under four candidate models—$\mathcal{M}_{\mathrm{STAMP}}, \mathcal{M}_{\mathrm{CR}}, \mathcal{M}_{\mathrm{BS}}, \mathcal{M}_{\mathrm{IR}}$—and estimate model parameters by optimising agreement with the empirical diversity structure (Fig. 4a). Model fit is quantified by minimising the discrepancy between a sample's Hill-numbers curve and the corresponding simulated Hill-numbers curve.

A Hill-numbers curve provides a compact and informative representation of diversity profile structure because it jointly captures tag richness and relative abundance[14]. Hill-numbers curves can be examined directly from empirical data or from the the best-fitting simulations produced by each model (Fig. 4b). Low orders q give greater weight to rare tags, whereas higher orders q increasingly emphasise dominant tags. As a result, the full Hill-numbers curve offers an essential one-to-one summary of a tag-frequency distribution, allowing diversity profiles to be compared unambiguously across samples and models.

Models' accuracy was quantified using an error score defined as the deviation between observed and simulated Hill-numbers curves, computed across all samples (Fig. 4c) or separately by condition (Supplementary Fig. 5). This model-testing framework allows us to address the nature of the bottleneck directly. $\mathcal{M}_{\mathrm{STAMP}}$ systematically failed to reproduce the contribution of low frequency tags,

indicating that an instantaneous bottleneck is not an adequate representation of the infection process. In contrast, $\mathcal{M}_{\mathrm{CR}}$, $\mathcal{M}_{\mathrm{BS}}$ and $\mathcal{M}_{\mathrm{IR}}$ achieved markedly better fits, simultaneously capturing both the near-equilibrium among dominant tags and the persistence of numerous low-frequency lineages. Together, these results support a time-resolved bottleneck as a more plausible explanation of roots infection dynamics.

Because each time-resolved bottleneck model corresponds to a distinct biological scenario, additional qualitative insight can be gained by comparing their relative performance. Notably, $\mathcal{M}_{\mathrm{BS}}$ and $\mathcal{M}_{\mathrm{IR}}$ outperformed $\mathcal{M}_{\mathrm{CR}}$ at low orders ($q \leq 2$), where the Hill-numbers are more sensitive to low-frequency tags. The key mechanistic distinction is that $\mathcal{M}_{\mathrm{CR}}$ assumes a constant entry rate, whereas both $\mathcal{M}_{\mathrm{BS}}$ and $\mathcal{M}_{\mathrm{IR}}$ allow entry to accelerate over time (Supplementary Fig. 4). This comparison refines our interpretation of the bottleneck in two ways:

- First, low-frequency tags—systematically absent in instantaneous bottlenecks models—are a natural outcome of time-resolved bottlenecks. The superior performance of $\mathcal{M}_{\mathrm{BS}}$ is consistent with its capacity to preserve rare lineages through the genealogical structure of range expansions[13] whereas $\mathcal{M}_{\mathrm{CR}}$ tends to erode this rare tail. This implies that even under highly stringent entry, minor secondary infection events can occur and remain detectable.

- Second, the improved fit of models with accelerating entry suggests that pathogen influx into the plant increases as infection progresses, consistent with a progressive weakening of host barriers. This interpretation aligns with reports of plasmolysis of cortical cells during infection[6,7,8]. By compromising tissue integrity and potentially widening access routes toward the vasculature, such damage would be expected to mechanically facilitate continued, and increasing, pathogen entry into the xylem.

1.4   Time-resolved bottlenecks quantify plant susceptibility or tolerance

To broaden inference, we extended the infection dataset across three experimental factors: *root condition* (intact *vs.* cut), *plant genotype* (susceptible *cv.* Marmande *vs.* tolerant *cv.* Hawaii7996[8,15,16]), and *temperature* ($27\,^{\circ}\mathrm{C}$ *vs.* $32\,^{\circ}\mathrm{C}$) (Fig. 5a). This factorial design yielded eight conditions, including the two reference baselines of intact or cut Marmande tomato plants infected at $27\,^{\circ}\mathrm{C}$ (*M_Intact_27*, *M_Cut_27*) from Fig. 2. Plants were sampled after symptom onset, as described in Materials and Methods.

All three factors are known to modulate disease progression. Susceptibility-enhancing modalities (Marmande tomato plants, cut roots, $32\,^{\circ}\mathrm{C}$, noted *M_Cut_32*) led to earlier symptom onset, higher bacterial load (CFU), and consequently earlier sampling (Supplementary Fig. 1). Conversely, tolerance associated modalities (Hawaii7996 tomato plants, intact roots, $27\,^{\circ}\mathrm{C}$, or *H_Intact_27*) limited colonisation, generating a continuum from highly susceptible (*M_Cut_32*) to strongly tolerant plants that remained asymptomatic and therefore could not be analysed (*H_Intact_27*).

Bottleneck stringency depends primarily on root integrity. Across conditions, the dominant driver of entry diversity was root state: intact roots admitted low diversity, whereas cut roots admitted high diversity (Fig. 5b). To compare bottleneck stringency across modems, we summarised it with a single parameter θ that can be extracted from each model ($\theta_{\mathrm{STAMP}} = N_b$; $\theta_{\mathrm{CR}} = \mu_{\mathrm{CR}}/r_{\mathrm{CR}}$; $\theta_{\mathrm{BS}} = v$; $\theta_{\mathrm{IR}} = \mu_{\mathrm{IR}}/r_{\mathrm{IR}}$; Supplementary Table 1). In time-resolved models, θ measures how rapidly new lineages enter relative to the expansion of already established lineages, making a "number of founders" interpretation inappropriate. For the models $\mathcal{M}_{\mathrm{CR}}$ and $\mathcal{M}_{\mathrm{IR}}$, the underlying entry rate μ and the population growth rate r can be recovered from θ.

In $\mathcal{M}_{\mathrm{BS}}$, θ_{BS} can be interpreted as a relative speed of barrier transversal: higher values correspond to a more synchronous entry, and therefore higher diversity at the sampling site. Because $\mathcal{M}_{\mathrm{BS}}$ provided the best overall fit (Figs. 5), we used θ_{BS} as our primary comparator across conditions (Fig. 5c). Intact roots exhibit stringent bottlenecks (median $\theta_{\mathrm{BS}} \approx 1.5$–2), whereas cut roots admitted substantially greater diversity (median $\theta_{\mathrm{BS}} \approx 4$–5). We then used a PERMANOVA to quantify how each experimental factor contributed to variation in θ_{BS} (Supplementary Fig. 6). Root condition was the only significant predictor ($R^2 = 16.3\%$) indicating that θ_{BS} is largely insensitive to susceptible differences driven by genotype or temperature. Thus, even in a time-resolved framework, a single bottleneck stringency metric θ is not sufficient to characterise infection dynamics: it summarises endpoint diversity, but does not capture the total pathogen burden within the sample.

Growth and entry rates predict susceptibility. Although $\mathcal{M}_{\mathrm{CR}}$ fit the data slightly less well than $\mathcal{M}_{\mathrm{BS}}$, it offers a key advantage: by combining diversity with CFU and sampling time, it enables estimation, at the level of individual plants, of both the pathogen growth rate r_{IR} (Fig. 5d) and a root permeability (entry rate) μ_{IR} (Fig. 5e). Unlike θ_{BS}, these two parameters reflect susceptibility across experimental conditions. PERMANOVA confirmed that all three factors (root condition, genotype, temperature) significantly influenced both r_{IR} and μ_{IR} (Supplementary Fig. 6). Genotype was the dominant driver of growth ($R^2 = 64.6\%$), with susceptible plants exhibiting an approximately twofold higher r_{IR}. Root scarification and elevated temperature had additional, smaller but significant effects on r_{IR} ($R^2 = 3.0\%$ and 3.2%, respectively). In contrast, permeability was primarily controlled by root scarification ($R^2 = 24.4\%$), with significant contributions from genotype and temperature as well ($R^2 = 5.1\%$ and 2.6%).

We also examined the role of plant size by including root samples weight as a proxy for a *plant size* factor (PERMANOVA; Supplementary Fig. 6). Plant size explained a substantial fraction of the variation of r_{IR} ($R^2 = 37.3\%$), but this effect largely reflected confounding with genotype, because tolerant plants were typically larger than susceptible plants in our setup. For μ_{IR}, only plant size, root condition and genotype remained significant (respectively $R^2 = 4.7\%$, 25.0%, and 1.2%). This pattern indicates that, independent of the other factors tested, larger plants tend to show higher estimated growth and entry rates, *i.e.* increased susceptibility, as captured by both r_{IR} and μ_{IR} overall indicating that fine susceptibility patterns are well-captured by pathogen influx parameter μ

254 and within-host growth parameter r.

1.5 Time-resolved bottleneck models extend to animal infections

To assess whether our approach generalises beyond plants, we applied the same model-testing frame-
work to a published dataset of *Pseudomonas aeruginosa* bacteraemia in mice (Fig. 2C in reference[9]),
which was previously reanalysed with STAMPR (Fig. 4 in reference[2]). Following intravenous inoc-
ulation with a tagged *P. aeruginosa* library, organ-specific bottlenecks were reported using N_b and
N_r (Fig. 6a;[2,9]). Tag-relatedness analyses further suggested distinct seeding routes: lung, spleen, and
liver primarily seeded by blood-circulating bacteria, whereas the gallbladder and gastrointestinal tract
derive from mixtures of liver-seeded and blood-circulating populations[2,9].

Model comparison identifies time-resolved bottlenecks. The original studies already argue
against an instantaneous bottleneck: the systematic underestimation of N_b relative to N_r was inter-
preted as evidence that clonal expansion occurs during organ colonisation[2]. We agree that systemic
infection reflects two concurrent processes: organ seeding by circulating bacteria (an entry rate μ) and
within-organ proliferation (a growth rate r). Our framework makes these process explicit and directly
tests the bottleneck assumption that remains implicit under STAMP(-R) metrics.
Consistent with this interpretation, error-curve comparisons show that the instantaneous model
($\mathcal{M}_{\text{STAMP}}$) provides a poor description of organ diversity profiles, whereas time-resolved models yield
substantially improved fits (Supplementary Fig. 7). Using the best-performing model ($\mathcal{M}_{\text{IR}}$), we then
extract θ_{IR}, an interpretable measure of bottleneck stringency (Fig. 6b). As in our plant system,
θ_{IR} captures whether colonisation proceeds in a largely monoclonal versus diversified manner under
continuous pathogen fluxes.

Entry and growth rates resolve organ-specific dynamics. Because our approach integrates
diversity, CFU and sampling time, a key advance is that it allows estimation of pathogen entry μ_{IR}
and growth r_{IR} at the resolution of an individual organ in an individual mouse. Estimated entry rates
μ_{IR} (Fig. 6c) indicate that lung, spleen, and liver are relatively permeable to circulating bacteria,
whereas the gallbladder and gastrointestinal tract are comparatively impermeable. These results agree
with relatedness-based inferences that the gastrointestinal tract is not directly seeded from blood[2,9],
while adding a quantitative flow-rate interpretation through μ_{IR}. As a additional, and previously
unreported insight, the inferred growth rates r_{IR} (Fig. 6d) suggest that lung and spleen share similar
dynamics (median $r_{\text{IR}} \approx 0.3$ h^{-1}), whereas liver, gallbladder, and most of the gastrointestinal tract
(with the notable exception of the stomach) display roughly twofold higher apparent growth.

Revisiting the "liver filter" interpretation in the *P. aeruginosa* infection. A previous
interpretation proposed that the liver acts as a filter for bacteria prior to entry into the gallbladder,
based on the observed reduction in N_b or N_r from liver to gallbladder[2,9]. Our time-resolved analysis

refines this view. We likewise infer a decrease in permeability between liver and gallbladder, but additionally detect a marked increase in downstream growth. Together, these results suggest an alternative mechanism: elevated growth in the gallbladder can amplify competitive exclusion, reducing effective diversity and producing an "apparent bottleneck" signature even under continued, time-resolved entry (Fig. 6e).

Model-based metrics outperform STAMPR in biological meaning and statistical power.
We next reanalysed a second dataset of systemic infection, tracking dissemination of *Klebsiella pneumoniae* following a pneumonia in mice (Fig. 1 in reference[10]). As for *P. aeruginosa*, time-resolved models provided a substantial better fit to organ diversity profiles than the instantaneous model $\mathcal{M}_{\text{STAMP}}$ (Supplementary Fig. 8). To further illustrate the added value of our framework, we compared our model-based parameters to the full set of STAMPR-derived metrics previously used to describe infection dynamics, and quantified the extent to which model-based inference improves interpretability and discriminative power (Supplementary Fig. 8).

In brief, the interpretable bottleneck metric θ_{IR} can replace N_s, avoiding the conceptual ambiguity that arises when a time-resolved process is summarised with instantaneous-bottleneck indicators. The inferred growth rate r_{IR} also provides a mechanistic counterpart to empirical proxies of expansion that are difficult to interpret, such as CFU/N_s and N_s/N_b. Likewise, the inferred entry rate μ_{IR} yields a direct estimate of pathogen transfer, an essential quantity that is not explicitly addressed in STAMPR. In terms of statistical performance, these model-based parameters generally required no log-transformation for parametric testing and more clearly separate experimental effects, suggesting both improved robustness and increased biological interpretability for analysing pathogen infection dynamics.

***K. pneumoniae* mode of dissemination is observable through within-host growth.** Two modes of *K. pneumoniae* dissemination from the lung to systemic sites have been described[10]: *metastatic dissemination*, in which systemic infection originates from a small number of dominant lineages that first expanded in the lung, and *direct dissemination*, in which a diverse, multiclonal pathogen population spreads to peripheral organs.

To strengthen this conclusion, we analysed r_{IR} across multiple organs, an inference that cannot be made from CFU alone. In *direct disseminations* cases at 24h post-infection, apparent growth rates were strikingly consistent across organs, with typical values around $r_{\text{IR}} \approx 0.45 \text{h}^{-1}$ (Supplementary Fig. 8). By contrast, *metastatic disseminations* was associated with an approximately 40% increase in r_{IR} across essentially all organs. This suggests that *metastatic dissemination* is not driven solely by enhanced replication in the lung, as originally proposed, but instead reflects a more general, host-wide increase in pathogen replication.

2 Discussion

Here, we propose to rethink the analysis of neutral barcoding experiments by treating diversity-profile structure as the result of a process combining successive infections and simultaneous growth of the pathogen population, rather than resulting from an instantaneous bottleneck event. Founder metrics (N_b, N_r, N_s) are appealingly simple, but they assume that entry and growth are decoupled. Accordingly, elevated N_s/N_b or N_r/N_b from STAMPR have been interpreted as signatures of clonal expansion during colonisation[2,10,17,18]. This interpretation is useful, yet it also highlights the limits of founder-based metrics when entry and growth co-occur.

Our results expose a counter-intuitive property of bottleneck metrics θ (i.e., N_b, N_r, N_s, θ_{CR}, θ_{BS}, θ_{IR}): Under time-resolved dynamics they do not quantify host defence or susceptibility. Instead, θ summarises the diversity among established lineages, without indicating how many bacteria (or how much flux) produced that profile. Capturing host barriers requires integrating pathogen burden and timing, via the entry and growth parameters μ and r. In our data for instance, M_Intact_32 showed rapid entry and growth with early symptoms, whereas H_Intact_32 showed slower entry and growth with delayed symptoms. Yet both display similar θ_{BS} values. They differ in susceptibility because $(\mu_{\mathrm{IR}}, r_{\mathrm{IR}})$ differ. This confusion traces back to the instantaneous STAMP assumption, where N_b conflates diversity profile structure (a θ-like quantity) with entry (μ). In time-resolved systems, μ and r cannot be recovered from N_b-derived indicators or θ_{BS}. Fully characterising host defences therefore requires explicitly estimating (μ, r), for example, using $\mathcal{M}_{\mathrm{CR}}$ or $\mathcal{M}_{\mathrm{IR}}$.

The proposed parameters (μ, r) substantially extend what can be learned from neutral barcoding experiments, enabling growth-rate inference at the level of individual hosts and even at individual organs. Critically, they require only measurements that are typically already collected: diversity profile, CFU and sampling time. Because r reflects an exponential growth process, absolute estimations of (μ, r) require CFU measurements taken during the exponential phase. In that regime, inferred r (Fig. 5e) is consistent with standard linear regression of CFU over time (Fig. 1b), while additionally providing per-plant resolution. When CFU are measured at saturation, (μ, r) can still be estimated, although they become "apparent" parameters that are reliable for relative comparisons but should not be interpreted as absolute estimations.

In *R. pseudosolanacearum* infections, we leveraged contrast in host and environment to dissect susceptibility. Relative to the susceptible cultivar Marmande, the tolerant cultivar Hawaii7996 showed reduced within-plant pathogen growth, consistent with prior work[5], whereas differences in root permeability were less clear. Root scarification increased permeability and, unexpectedly, also increased within-plant growth, suggesting that wounding decreases internal root defences.

A plausible mechanism is that damage to root-barriers, known to regulate metabolite containment and the root-associated microbiome[19,20,21], promotes leakage of exudates such as glutamine. The latter is both a chemoattractant for soil bacteria[22] and a preferred substrate for *R. pseudosolanacearum* during infection[23]. Likewise, cortical cell death reported during infection[6,7,8] could progressively widen

access routes toward the xylem, enhancing both physical entry and exudate leakage over time. Elevated temperature further increased susceptibility by boosting growth and permeability. Interestingly, r_{IR} and μ_{IR} are correlated across plants (Spearman rank correlation $R^2 = 0.40$, $p < 0.05$; Supplementary Fig. 6), suggesting a general partial coupling of the two effects during infection processes.

In *P. aeruginosa* bacteraemia in mice, our reanalysis reveals a marked within-host shift in behaviour: dissemination is strongest in blood-connected organs (lung, spleen and liver) whereas apparent growth is highest in the gastrointestinal tract (gallbladder, stomach, small intestine, caecum and colon). This pattern suggests selective pressure and competition to rapidly establish in the gastrointestinal tract, potentially because it enhances persistence and transmission through faeces shedding, or because it provides a niche for chronic infection. In line with this view, recent work links high replication potential of *P. aeruginosa* to stringent gastrointestinal tract bottlenecks[24], consistent with our growth-driven interpretation of "apparent bottlenecks".

More broadly, how common are time-resolved versus instantaneous bottlenecks? The field increasingly acknowledges that time matters in neutral barcoding analysis[2,25]. Direct evidence for truly instantaneous founding remains limited, while many frequently observed signatures, strongly heterogeneous tag frequencies, spatial and temporal declines in N_b, abundant rare-frequency tags, or $N_r/N_b > 1$, naturally arise when entry and growth overlap. This raises the possibility that instantaneous-bottleneck analysis have often been applied to systems dominated by time-dependent processes.

A notable step towards time-resolved modelling is RESTAMP for *Vibrio cholerae* in the mouse gastrointestinal tract[3,25], which emphasises transfer rates and growth across compartments. However, its high complexity (~ 30 fitted parameters[25]) and reliance on N_b to infer secondary parameters may hinder interpretability and increase the risk of over-fitting in a model that did not provide control of the fit against full tag frequency data.

Here, we take a different route: we compare a panel of mechanistic models directly against the observed diversity structures and select those best supported by the data. This strategy has three advantages: (*i*) assumptions need not be accepted *a priori*, poor models are rejected by fit, (*ii*) whether the bottleneck is instantaneous or time-resolved emerges from model performance, (*iii*) the selected models yield interpretable parameters that map onto explicit biological processes[26]. More generally, Hill-numbers curves matching provides a standardised way to test any hypothesised scenario, provided it can be formalised into a mechanistic model. Our fits are not perfect, indicating that additional mechanisms could be investigated. For instance, tissue-specific heterogeneity could affect arrival times, local densities, or growth conditions but would require spatially resolved sampling and explicit spatial models.

We show that three widely applicable time-dependant mechanisms (growth, motility and spatial competition) are intertwined in bacterial infection dynamics. Because of the generality of the addressed mechanisms, the improvements in robustness and in biological insights without modifications in current experimental protocols, our findings support a need to shift from the previous N_b-based methodology to an updated methodology accounting for time-dependant factors. We advocate for a global reanalysis

of STAMP(-R) articles under a time-resolved model approach. We foresee that underappreciated biological results may be uncovered in the following cases: (i) studies that used only N_b and may have overlooked major aspects of the infection dynamics[1,9,27,28,29], (ii) studies of colonisation across continuous space (*e.g.* gastrointestinal tract colonisation) that neglected contributions of motility and growth effects by using N_b-derived indicators[2,10,18,24,25,30,31,32], (iii) studies that empirically uncovered the importance of time dimensions using N_b-derived indicators but were limited in their analysis[17,18,30], (iv) studies that directly addressed relations between CFU dose effects and bottleneck using N_b-derived indicators but did not pair the two in an appropriate model[30,33,34].

Beyond refining earlier findings, the proposed methodology more broadly aims to connect microbial population dynamics to quantitative biological mechanisms. It opens the possibility to address questions in microbial population genetics with a unified mechanistic framework that explicitly includes time-dependant complexity. Natural extensions of this work include bringing time-resolved inference to broader microbial ecology contexts such as: tracking of complex bacterial synthetic communities[35,36], genealogies tracing of a bacterial population and inference of evolution rates[13] or mapping of complex directional pathogen fluxes at a host-wide level[37].

3 Figures

Figure 1: Roots of susceptible tomato (*cv.* Marmande) are infected by limited numbers of individual *R. pseudosolanacearum* cells. **a.** STAMP overview: 119 isogenic GMI1000 strains carrying a neutral 20 bp chromosomal barcode are generated; plants are soil-drenched with a co-culture of the tagged strains. Roots are sampled to measure bacterial load and tag diversity. Because tags are fitness-neutral, losses of diversity report bottlenecks. **b.** Root bacterial load from 2-6 DPI (log scale, $n = 8$-9 plants/day). Fit: $y = e^{a\,t+b}$ (black dotted line, equation). Blue line: local mean $\pm$ SD. **c.** Root Shannon diversity from 3-6 DPI; sequencing was feasible from 3 DPI onward when loads typically exceeded $\sim 10^4$ CFU/g. Red line: local mean $\pm$ SD. Diversity in inoculum: 4.71.

Figure 2: Tomato root-entry bottleneck depends on the physical integrity of the roots. **a and b.** Diversity profiles during colonisation of the roots of control intact Marmande plants (panel **a**, $n = 24$) or root-scarified Marmande plants (panel **b**, $n = 24$). Colours: relative frequency of the 119 tags. Bacterial load is a barplot under each diversity profile (log scale). "NA": unsuccessful sequencing due to low bacterial load. **c.** Diversity profiles in the inocula ("A27", "B27", "C27") of the three time-independent replicates of infection in panel a and b. **d.** Shannon diversity index in the roots of intact Marmande plants (*M_Intact*) or cut Marmande plants (*M_Cut*). Wilcoxon-Mann-Whitney test (two sided, **: $p = 0.0094$). Diversity in inocula: 4.70-4.72.

Figure 3: Pathogen diversity profiles better support "time-resolved" rather than "instantaneous" bottleneck models. **a.** Bottleneck estimates in intact (*M_Intact*) or root-cut Marmande plants (*M_Cut*) using richness, N_b[1] or, respectively N_r and N_s[2] (log scale). Lines link metrics from the same sample. Wilcoxon signed-rank test (two-sided, ns: $p > 0.05$; *: $p < 0.05$; ** $p < 0.01$; ****: $p < 0.0001$). **b.** N_b and N_r versus of richness (log-log scale). The diagonal $y = x$ indicates equality; points below it are lower than richness. Line links metrics from the same sample. **c.** Instantaneous bottleneck models assume a single colonisation event from the soil to the roots[11,12]. The STAMP model (N_b)[1] or the STAMPR model $(N_r$ and $N_s)$[2] follow this assumption (details in Supplementary Fig. 3). **d.** Time-resolved bottleneck models allow entry over time, enabling tests of mechanisms shaping diversity during infection (*e.g.* migration/dispersion to the xylem, early–late growth differences, and spatial competition). Examples include the CR, BS and IR models (details in Fig. *3)*

Figure 4: Time-resolved bottleneck models better explain tomato root colonisation. **a.** Diversity profiles observed in samples (intact Marmande plants) compared to the best fitting profiles under simulation of the models $\mathcal{M}_{\text{STAMP}}$, $\mathcal{M}_{\text{CR}}$, $\mathcal{M}_{\text{BS}}$ and $\mathcal{M}_{\text{IR}}$ (log scale). Colours: individual samples. **b.** Diversity profiles in samples or in simulations according to the Hill numbers ^qH from order $q = 0$ (importance of richness) to order $q = 4$ (importance of evenness). As a guide, $q = 0$ equals richness, $q = 1$ is the exponential of Shannon entropy, and $q = 2$ is the inverse Simpson index (or Simpson diversity). For a given order q, a high ^qH value in sample indicates a high diversity. Colours: individual samples. **c.** Error between simulations and real data in the range from $q = 0$ to $q = 4$, averaged over all infection conditions in Fig. 2 and Fig. 5.

Figure 5: Susceptibility to *R. pseudosolanacearum* reflects time-dependent infectious dynamics. **a.** Eight tested conditions combine three factors of plant susceptibility ($n = 24$ plants/condition). "Root condition" factor: intact plants versus root-cut plants. "Plant genotype" factor: susceptible cultivar Marmande versus tolerant cultivar Hawaii7996. "Temperature" factor: standard (27°C) versus high temperature (32°C). **b.** Hill-numbers curves between $q = 0$-6 for pathogen diversity averaged over each condition (M: Marmande, H: Hawaii7996, I: Intact, C: Cut). Condition *H_Intact_27* was tolerant and did not yield sufficient colonisation for sequencing. **c.** Estimated bottleneck parameter $\theta_{BS} = v$ obtained with $\mathcal{M}_{BS}$ (log scale). "Bottleneck stringency" θ_{BS} reflects monoclonal or diverse colonisation. Low value indicates stringent bottleneck. Kruskal-Wallis test, *post hoc* Fisher's LSD (threshold: $p < 0.01$). **d.** "Growth rate" r_{IR} in individual plant roots estimated from the parameter $\theta_{IR} = \mu_{IR}/r_{IR}$ in $\mathcal{M}_{IR}$. Kruskal-Wallis test, *post hoc* Fisher's LSD test (threshold: $p < 0.01$). **e.** "Entry rate multiplier" μ_{IR} in each individual plant roots estimated from the parameter $\theta_{IR} = \mu_{IR}/r_{IR}$ in $\mathcal{M}_{IR}$. μ_{IR} reflects the transfer rate of pathogen from soil to the plant roots. High value indicates faster entry of pathogen over time. Kruskal-Wallis test, *post hoc* Fisher's LSD test (threshold: $p < 0.01$).

Figure 6: Time-resolved models capture overlooked dynamics in systemic mouse-*Pseudomonas aeruginosa* infection (original data from Fig. 2C of article [9], $n = 10$ mice). **a.** N_b and N_r as calculated in [2,9] (log scale). Line link metrics of same sample. **b.** Bottleneck parameter θ_{IR} obtained with $\mathcal{M}_{IR}$ (log scale). Low value indicates stringent bottleneck. Kruskal-Wallis test, *post hoc* Fisher's LSD test (threshold: $p < 0.01$). **c.** Apparent growth rate r_{IR} in individual organs. Whether r_{IR} corresponds to absolute growth rate *in vivo* depends on the assumption of sampling during exponential growth. Kruskal-Wallis test, *post hoc* Fisher's LSD test (threshold: $p < 0.01$). **d.** Apparent entry rate multiplier μ_{IR} in individual organs. In this context, μ_{IR} represents the influx of systemic circulating bacteria into organs. High value indicates faster entry of pathogen over time. The accuracy of μ_{IR} is conditioned to the accuracy of r_{IR}. Kruskal-Wallis test, *post hoc* Fisher's LSD test (threshold: $p < 0.01$). **e.** Schematic hypothesis regarding mouse-*P. aeruginosa* infection. Lung, spleen and liver are colonised under a high dispersion mode (low r, high μ) with high connection to inoculation site (blood) promoting establishment of high pathogen diversity. Gastrointestinal organs are colonised under a high spatial competition mode (high r, low μ) with gallbladder being quickly colonised by the first arriving cells promoting the establishment of a homogeneous, low diversity.

4 Methods

4.1 Construction of the STAMP collection in *R. pseudosolanacearum* and co-culture of the strains

123 oligonucleotides of a 20 bp random sequence (referred as "tag", list in supplementary document "STAMP123_tags.fasta") were synthesised and hybridised with their complementary sequence to form inserts with cohesive extremities BsrGI and BglII. These fragments were inserted by restriction–ligation (BsrGI / BglII) in a pRCT-GWY[38] derived plasmid denoted pNP413. The tag is contained inside a genetic cassette of 6 kb containing a *tetR* gene, a *mCherry* gene and two flanking regions of 1 kb used for homologous recombination with *R. pseudosolanacearum* strain GMI1000 in the chromosome[38].

Produced plasmids were chemically transformed into calcium-competent *E. coli* strain DH5alpha and selected using LB medium + tetracycline 10 mg/L. Clones were verified using a wild-type discriminating PCR and Sanger sequencing of the STAMP tag (Eurofins). Validated plasmids were purified in 50 μL ultrapure water using 2 mL of stationary-phase culture with a column extraction kit (BS614, Bio Basic). 10 μL of plasmid was linearised using SfiI enzyme and mixed with 200 μL of *R. pseudosolanacearum* strain GMI1000 from a liquid culture in exponential phase, grown 1 week at 28 °C in MP medium + 2% glycerol. The mix was deposited on MP medium + 2% glycerol Petri dishes and incubated 2 days at 28 °C for natural transformation. Three days later, the bacterial lawn was suspended in distilled water and isolated on Phi medium + tetracycline 10 mg/L Petri dishes. The obtained *R. pseudosolanacearum* strains were validated for the correct insertion of the 6 kb cassette using a wild-type discriminating PCR and Sanger sequencing of the STAMP tag (Eurofins).

Individual clones were grown in 96-well microplates at 28 °C in 250 μL of Phi medium + tetracycline 10 mg/L in an Omega FLUOstar spectrophotometer (BMG Labtech) until stationary phase. An OD_{600nm}-adjusted volume (50 $AU_{OD} * \mu$L per well) was used to inoculate a secondary microplate culture in the same conditions. At stationary phase, wells were pooled 8-by-8 using OD_{600nm}-adjusted volumes (100 $AU_{OD} * \mu$L per well) for inoculation of tertiary Erlenmeyer cultures at 28 °C in 20 mL Phi medium + tetracycline 10 mg/L overnight. At stationary phase, the OD_{600nm}-adjusted volumes of each 8-tag pool (1400 $AU_{OD} * mL$) were mixed with 500 μL glycerol 60% for storage at −70 °C. The day before infection, an inoculum of 120 tags was produced by thawing the 15 tubes of 8 tags (total volume $\sim$ 25 mL) and growing them directly in an Erlenmeyer with 250 mL of Phi medium + tetracycline 10 mg/L at 28 °C overnight. Upon validation of these final co-cultures, one tag (tag007) was found missing but an otherwise almost perfect stoichiometry was obtained for the 119 other tags. The three remaining tags (denoted tag121, tag122 and tag123) were grown and kept separately for sequencing calibrations.

Used media. LB medium: Yeast extract (Duchefa Biochimie) 5 g/L; Bacto tryptone (Becton Dickinson Difco) 10 g/L; NaCl 10 g/L in distilled water. Phi medium: Bacto peptone (Gibco) 10 g/L; Casamino acid (MP) 1 g/L; Yeast extract (Duchefa Biochimie) 1 g/L in distilled water. MP medium: KH2PO4 3.4 g/L; (NH4)2, SO4 0.5 g/L; MgSO4 (7H2O) 50 mg/L; FeSO4 (7H2O) 125 µg/L in distilled

450 water. Adjust pH to 7.0 with KOH 10N.

4.2 Plant culture, plant infection and samples collection

For all experiments, tomato plants (susceptible cultivar Marmande "M" or tolerant cultivar Hawaii7996
"H") were sowed in peat-based substrate and then cultivated in 6×6 cm pots until an age of approx-
imately 3 weeks in a phytotron ($25.5\pm1\,°C$ day 12 h / $24\pm1\,°C$ night 12 h, hygrometry $60\pm15\%$). The
plants were then acclimated to the final culture chamber for 2 or 3 days either at a standard tem-
perature ($27\pm0.5\,°C$ day 12 h / $26\pm0.5\,°C$ night 12 h, hygrometry $60\pm5\%$) or at a high temperature
($32\pm1\,°C$ day and night 12 h/12 h, hygrometry $70\pm5\%$). The days of infection, two technical repli-
cates of 2 mL of the inoculum STAMP co-culture were taken for sequencing of the initial population.
The co-culture was diluted using tap water (to an $OD_{600nm} = 0.05$) and 50 mL of diluted co-culture
($\sim 2.5 \times 10^9$ CFU) was used to infect plants by watering. Right before inoculation, the condition
of "Root cut" plants was produced by scarifying roots of the plants using a scalpel by drawing an
"L"-shaped trench in the pot. "Intact" plants were otherwise watered without scarification.

For the experiments of kinetics of infection, one infection was performed (inoculum S27) on $n = 42$
plants (Conditions M_Intact_27). $n = 8$ or $n = 9$ plants were then sampled between DPI 2 to DPI 6.
The plant was unrooted and dirt removed. A root sample was collected corresponding to the whole
taproot cut at the highest lateral root and removed from all lateral roots. Samples were weighed,
surface-sterilised using 70% ethanol and cleaned using sterile distilled water. Samples were cut into the
smallest possible pieces using a scalpel and macerated in 2.5 mL sterile distilled water for approximately
3 h. The macerate was used to determine bacterial load by plating serial dilutions on Phi medium
+ tetracycline 10 mg/L Petri dishes using an automatic spreader (EasySpiral Pro, Interscience). The
remaining ~ 2 mL of macerate was centrifuged and gDNA was extracted from the bacterial pellets
using a lysis–isopropanol–ethanol protocol (Wizard Genomic DNA Purification Kit A1125, Promega)
in 50 μL final of ultrapure water.

For the experiments with single-point sampling, six temporally independent infections were per-
formed (inocula denoted A27, B27, C27, A32, B32, C32) combining a total of 192 plants (Conditions:
M_Intact_27, M_Cut_27, H_Intact_27, H_Cut_27, M_Intact_32, M_Cut_32, H_Intact_32, H_Cut_32). For
each condition three replicates were performed $n = 3 \times 8 = 24$. At the first symptoms of wilting (or at
DPI 13 or 14 if no symptoms) within a batch of 8 plants in a replicate, the whole batch was collected.
Bacterial load was quantified and DNA was extracted for sequencing as described above.

4.3 Library preparation and sequencing of the diversity profiles

481 For each purified gDNA, absorbance was measured for an $A_{260/280}$ quality ratio. Whenever possible,
482 samples were diluted to an empiric concentration between $[C] = 20$ to 100 ng/μL (corresponding
483 to a theoretical number of unique molecules of gDNA of 3×10^6 to 1.5×10^7 tags/μL) satisfying
484 a non-limiting amount of targets and an accurate measure of concentration. For samples in which

concentration was below 20 ng/μL or gave aberrant values in view of bacterial load (*e.g.* vegetal phenolic compounds), a value N_{max} defined as the *theoretical maximum number of tags* was derived from bacterial load. A 35-cycle adapter PCR1 was performed to amplify a 174 bp region containing the tag. This PCR was performed using primers with the advised 22 bp Nanopore adapter regions for PCR barcoding (SQK-LSK114, Oxford Nanopore Technologies). Additional 4 bp regions were included between primer hybridisation sequence and adapter sequence. These 4 bp represent a "run barcode" that allows filtering of cross-contamination when using the same Nanopore flow cell in several different sequencing runs. For this article, 6 different pairs of primers were sequentially used with different "run barcodes" in this fashion. For samples for which $[C] > 20$ ng/μL and $N_{max} > 2500$ tags/μL, 4 μL of purified DNA was used in a 75 μL PCR1. For samples for which $[C] < 20$ ng/μL and $N_{max} < 2500$ tags/μL or samples that did not amplify in the first condition, a larger volume of 20 μL of purified DNA was used in a 75 μL PCR1 to ensure an adequate number of targets. Samples that did not validate the two first conditions were repurified in the same volume using 2X magnetic bead-based mix in a SPRI protocol[39] before reattempting a PCR1. Samples that were validated for a ~ 226 bp band on gel electrophoresis were then purified using 1.5X magnetic bead-based mix in a SPRI protocol[39] to eliminate primers and to concentrate the PCR product twofold.

2.5 μL of reconcentrated PCR1 product was used in a 30-cycle barcoding PCR2 in a final volume of 25 μL using primers purchased from the Nanopore 96 Barcode Expansion Pack (EXP-PBC096, Oxford Nanopore Technologies). Samples that were validated for a ~ 274 bp band on gel were then quantified for DNA concentration using Quant-iT PicoGreen dsDNA Assay (P7589, Invitrogen) and pooled in equal stoichiometry before a purification with 1.2X magnetic bead-based mix in a SPRI protocol[39]. End-prep and adapter ligation steps were performed according to the PCR barcoding protocol (SQK-LSK114, Oxford Nanopore Technologies) with the exception of SPRI purifications that were adjusted to a 1.5X ratio in accordance with the length of the DNA fragments of interest.

A PromethION flow cell R10.4.1 was loaded with an adjusted quantity of 200 fmol of DNA and data acquisition was performed with a goal of minimum depth of sequencing of ~ 20k unprocessed reads for all samples. The flow cell was washed according to protocol (EXP-WSH004-XL, Oxford Nanopore Technologies) to reuse it for a total of 6 runs of ~ 96 samples. For technical simplicity, some samples went through the whole sequencing process despite not being validated in the early PCR steps. These samples were excluded from the final analysis.

4.4 Raw data processing

DNA sequences were basecalled using Dorado version 0.7.3 with the Super Accurate model v5.0.0 and a minimum Q score of 10. Briefly, sequences were first demultiplexed for the Nanopore barcode and the Nanopore barcodes were trimmed using the in-built Dorado tools. DNA sequences were then processed through a QIIME 2 bioinformatics pipeline for metagenomics. DNA sequences were size-filtered to keep only fragments between 160 and 200 bp. A first closed-reference mapping was performed to identify the "run barcode". For reads which validated the correct run barcode, a second closed-reference mapping

522 was performed against the 123 potential STAMP tags to calculate the tag abundances. Three tags
523 exogenous tags (namely tag121, tag122, tag123) were used to calibrate barcode hopping. Barcode
524 hopping was estimated by measuring the hopping between the 120 STAMP tags from samples toward
525 the three control tags. A typical order of magnitude of $\sim 1\%$ of noise reads was obtained for final
526 depths of sequencing of $> 10^4$ reads in all samples.

4.5 Validation of the 119-tags collection for STAMP experiments

All strains showed comparable growth *in vitro* and the proportions of the 119 tags remained stable
in co-culture aside from tag007 lost due to a pipetting error (Supplementary Fig. 9), indicating that
preparation steps did not introduce technical bottlenecks, that tagged strains exhibited indistinguish-
able *in vitro* growth, and that sequencing did not introduce appreciable PCR bias. To validate neutral
fitness of the collection *in planta*, we assumed that all tags should have identical probability to appear
across independent infections. Tags showed no consistent over-representation across infections (Sup-
plementary Fig. 9), validating that tagged strains have similar probabilities of establishing infection
and the observed diversity profiles reflect pathogen entry dynamics rather than tag-specific fitness
effects or technical artifacts such as sequencing bias.

4.6 N_b-derived indicators calculation and improvements of the STAMPR pipeline

For the calculation of the N_b, N_r and N_s metrics, the code from the original STAMPR pipeline[2]
was adapted. In short, the code was corrected to include the "resiliency algorithm" method and the
calculation of N_r in addition to N_s (see below). We also included additional error estimators during
the calculation of N_b and N_r.
As part of the pipeline, we set the prefiltering threshold to remove noise tags at 25 or fewer reads.
This threshold was selected based on the estimation of barcode hopping in our experimental setup.
This threshold corresponds to a threshold to filter out at least quantile 99% of tags emerging from
noise alone.). Eliminated tags typically accounted for less than 1% of tag abundance in samples and
less stringent thresholds had minimal change on N_b and N_r.
The standard bottleneck N_b in Fig. 3a, 6a was calculated based on the analytical equation formalised
in the original STAMP publication[1] using the prefiltered data:

$$N_b = \frac{g}{\left(\frac{1}{K} \sum_{i=1}^{K} \frac{(f_i - f_i^0)^2}{f_i^0 (1 - f_i^0)} - \frac{1}{S_0} - \frac{1}{S_s} \right)}, \tag{1}$$

550 K is the number of tags in the inoculum, f_i^0 and f_i are the frequencies of tag i respectively in the
551 inoculum and the sample, $g = 1$ is the number of generations, S_0 and S_s are the sequencing depths
552 respectively in the inoculum and the sample. A second way to compute N_b is proposed with θ_{STAMP}
553 based on the fit of a multinomial model (see Section 4.8)). This second way to compute standard

554 bottleneck can be approximated by a simpler analytical equation:

$$\hat{\theta}_{\text{STAMP}} = \frac{K}{\sum_{i=1}^{K} \frac{(f_i - f_i^0)^2}{f_i^0(1 - f_i^0)}},$$

Significant discrepancies between N_b and θ_{STAMP} may appear when frequencies f_i do not follow a
multinomial distribution.

The limit of detection of the N_b metric was calculated in our experiment by considering a signal-to-
noise problem. Noise was considered to be the sum of technical artefacts in tag abundances originating
from PCR noise, residual barcode hopping or misattribution of tags during bioinformatic analysis. An
artefactual N_b was calculated between technical replicate of sequencing of a same inoculum. $N_b >$
4×10^3 was obtained indicating that it is not possible to discriminate values of N_b above this threshold.

A corrected bottleneck value N_r was derived from N_b. The original code of the STAMPR pipeline[2]
was cleaned and annotated. Previously, only calculation of N_s was made available based on the
"reference resample plot" of the algorithm. Instead, we corrected the code so that both N_s and N_r
are systematically computed. Finally, an algorithm was added to estimate whether the number of
tags in the inoculum is sufficient to accurately estimate a N_b or N_r. This algorithm is similar to
the simulations that were performed in the original STAMP publication[1]. The algorithm simulates
bottleneck events under the estimated N_b or N_r and the resulting standard deviation provides the
confidence interval in the calculation of a given N_b or N_r.

4.7 Measuring diversity in samples and model comparison

571 In this section, we construct a method to compare the observed data with the data generated by
572 mechanistic models (see Section 4.8). This method allows us (i) to estimate the parameters of our
573 models and (ii) to compare the ability of our models to reproduce the observed data.

574 **Hill numbers and Hill curve.** Our comparison methods is based on the Hill numbers. The Hill
575 number of order q of a frequency vector $F^1 = (f_1, \ldots, f_K)$ is defined as:

$$^qH = \left(\sum_{i=1}^{K} (f_i)^q \right)^{\frac{1}{1-q}}.$$

Certain values of q correspond to standard diversity indices: ^0H represents species richness (number
of types); ^1H corresponds to the exponential of Shannon diversity index; ^2H is related to the inverse
Simpson diversity index, giving more weight to common tags. For all values of q, the maximum value
of ^qH is reached when $f_i = 1/K$ for all i. In this case, $^qH = K$. More generally, for each fixed q, a
lower ^qH corresponds to lower diversity, and the parameter q controls the relative emphasis placed on
common tags (large q) or rare tags (small q).

As proposed in reference[14], we describe the distribution of the frequencies through the full *diversity*
profile ^qH over a continuum of values of q. The reason for this choice is that the diversity profile fully

characterises the frequencies (proposition A22 of reference [14]), as two frequency vectors cannot lead to
the same profile unless they are equal up to a permutation.

Models and data comparison. Observing the Hill curves allows us to directly visualise the effect of
the bottleneck on the frequency distribution. These profiles also allow us to compare models and data.
Consider two frequency vectors $F^1 = (f_1^1, \ldots, f_K^1)$ and $F^2 = (f_1^2, \ldots, f_K^2)$. We define the discrepancy
between these frequencies as the average relative difference between the Hill numbers of the two vectors
over a range of q values:

$$d(F_1, F_2) = \frac{1}{\#Q} \sum_{q \in Q} \frac{|^qH(F_1) - {}^qH(F_2)|}{{}^qH(F_1) + {}^qH(F_2)},$$

where Q is a representative set of values of q, and $\#Q$ is the number of elements in this set. We took
Q to be a set of evenly spaced values ranging from 0 to 4. This range was empirically selected as
experimental data typically showed the highest variations on the interval $[0, 4]$.

Now, consider a stochastic model $\mathcal{M}(\theta)$, parametrised by $\theta \geq 0$, and describing the distribution of
the frequencies F after the bottleneck event. Given the observed experimental frequencies $\hat{F}$, for each
fixed θ, we define the performance $P[\mathcal{M}(\theta)]$ of $\mathcal{M}(\theta)$ as the 0.05 quantile of the distribution of $d(F, \hat{F})$
over replicate simulations. Then, the performance $P[\mathcal{M}]$ of the model $\mathcal{M}$ is obtained by choosing the
parameter θ that leads to the smallest value of $P[\mathcal{M}(\theta)]$. The model is more effective in reproducing $\hat{F}$
when $P[\mathcal{M}]$ is small. In other words, we let $P[\mathcal{M}(\theta)]$ be the 0.05 quantile of

$$\left\{ d(F_i[\mathcal{M}(\theta)], \hat{F}), \ i = 1, \ldots, N \right\}$$

600 where $F_i[\mathcal{M}(\theta)] \in [0, 1]^K$, $i = 1, \ldots, N$, are the final frequencies for N replicate simulations simulated
601 with model $\mathcal{M}$ and parameter θ. We then let

$$P[\mathcal{M}] := \min_\theta P[\mathcal{M}(\theta)], \qquad \theta[\mathcal{M}] := \operatorname*{argmin}_\theta P[\mathcal{M}(\theta)],$$

602 be, respectively, the performance of the model $\mathcal{M}$ and the estimated parameter for the model $\mathcal{M}$.
603 In our situation, we have several experimental frequency vectors $\hat{F}_1$, ..., $\hat{F}_N$, observed under the
604 same experimental conditions. We consider independently the vectors $\hat{F}_1$, ..., $\hat{F}_N$. For $i = 1, \ldots, N$,
605 let θ_i be the parameter that leads to the smallest value of $P[\mathcal{M}(\theta_i); F_i]$. The model performance is
606 then the average of the $P[\mathcal{M}(\theta_i); F_i]$, $i = 1, \ldots, N$, and the output parameter $\widehat{\theta}$ is the average of the θ_i.

4.8 Mechanistic Models

We propose four stochastic mechanistic models that describe different ways of crossing the barrier.
Comparing the performance of these models allows us to select the best one and infer the most
probable underlying mechanism. For each model $\mathcal{M}(\theta)$, the parameter θ determines the intensity
of the bottleneck. In all models, the smaller the value of θ, the stronger the bottleneck effect (however,
θ may scale differently according to the model).

Instantaneous bottleneck model $\mathcal{M}_{\text{STAMP}}(\theta_{\text{STAMP}})$. This is the standard model derived from the

original STAMP publication[1]. From a mechanistic perspective, the model assumes that the founding

population after the bottleneck consists of N bacteria that cross a barrier simultaneously. These

individuals are randomly selected from the initial population according to a multinomial distribution:

$$\mathcal{M}_{\text{STAMP}}(\theta_{\text{STAMP}}): \quad F \sim \text{Multinomial}(N; F_0)/N,$$

where F_0 represents the pre-bottleneck frequencies and F stands for the post-bottleneck frequencies.

The parameter of the model is the size of the founding population: $\theta_{\text{STAMP}} = N$. It can be determined

as the value θ_{STAMP}^* that maximises the performance $P[\mathcal{M}_{\text{STAMP}}(\theta_{\text{STAMP}})]$.

Constant rate model $\mathcal{M}_{\text{CR}}(\theta_{\text{CR}})$. Bacteria cross the barrier individually and stochastically at a

Poisson rate $\mu > 0$, carrying tags selected according to F_0, the pre-barrier frequencies. Once across

the barrier, bacteria reproduce deterministically following Malthusian growth at rate r. Under those

assumptions, the output $X_t = (x_1(t), \ldots, x_K(t))$ represents the number of bacteria carrying each tag

at time t. By the properties of Poisson point processes, this is equivalent to assuming that

$$x_i(t) = \sum_{j \geq 1} 1_{\{\tau_{i,j} \leq t\}} \, e^{r\tau_{i,j}}, \tag{2}$$

where, for each $i \in \{1, \ldots, K\}$, $\{\tau_{i,j}, j \geq 1\}$ are the points of a Poisson process with intensity μ/K, and

these processes are mutually independent. The corresponding frequencies depend on the observation

time and are given by $F_t = X_t/n_t$, where $n_t = \sum x_i(t)$. The parameter of the constant-rate model is

defined as $\theta_{\text{CR}} = \mu/r$. It quantifies the relative entry rate of bacteria through the bottleneck compared

to their Malthusian growth rate. We interpret θ_{CR} as a measure of bottleneck strength: high θ_{CR}

$(r \ll \mu)$ means that bacteria cross the bottleneck relatively quickly compared to their growth, making

the final population composition closer to the pre-bottleneck distribution. Conversely, low θ_{CR} $(r \gg \mu)$

means that bacteria grow much faster than they cross the bottleneck. Early crossers dominate, as they

have more time to expand before new bacteria enter, leading to a founder effect where the first few

individuals disproportionately shape the post-bottleneck population structure.

Bolthausen-Sznitman model $\mathcal{M}_{\text{BS}}(v)$. To construct the Bolthausen–Sznitman model, we assume

that the population is described by a Fisher-KPP equation. The Fisher–KPP equation is a reac-

tion–diffusion equation commonly used to model population dynamics, particularly the evolution of

neutral tags during a spatio-temporal expansion process[40] occurring at a constant propagation speed.

To account for finite-population effects, we introduce stochasticity, leading to a noisy Fisher–KPP

equation.

We see the bottleneck as an area where the population must remain very small, and outside of which

the population is large. In the experiment, n individuals are sampled shortly after the population has

crossed the barrier. *What are their genealogies?* No common ancestor can be found outside the barrier,

since the population is large in these areas. However, inside the barrier, common ancestors can be
found. The connection to our problem is that if two individuals located after the barrier are sampled,
then they will necessarily carry the same tag if they have a common ancestor that lived after the
tags were attributed. If no such common ancestor exists, the tags of the two sampled individuals are
independent. Similarly, if we know the genealogies of a sample of $n \geq 3$ individuals, then we perfectly
know the joint law of their tags. Therefore, understanding how the genealogies of individuals are
affected by the crossing of the barrier is sufficient to understand the frequencies of the different tags
that we observe at sampling time.

There exist different standard population genetics models for such genealogies. We tested two
of them, the Kingman coalescent and the Bolthausen-Sznitman coalescent, which are conjectured or
shown to arise in various forms of the noisy Fisher-KPP equations [13,41,42]. We do not show the Kingman
coalescent results in the main material, as it was unable to reproduce diversity profiles similar to the
experimental ones (Supplementary Fig. 5, 7). In Section 4.9, we provide a construction of the Kingman
coalescent and the Bolthausen-Sznitman coalescent.

The strength of the barrier in this model is measured by the time T during which the coalescent
process is run to obtain the genealogy of the individuals following the bottleneck, which can be in-
terpreted as the time taken by the population to cross the barrier. We parametrise the model with
the speed $\theta_{\mathrm{BS}} = 1/T$. A high θ_{BS} means that the population crosses the barrier rapidly, so that the
bottleneck is weak.

Increasing rate model $\mathcal{M}_{\mathrm{IR}}(\mu/r)$. Inspired by the Bolthausen–Sznitman model, the increasing rate
model offers a simpler mechanistic interpretation while still capturing a similar behaviour in terms of
arrival-time profiles (Supplementary Fig. 4). Conceptually, $\mathcal{M}_{\mathrm{IR}}$ is a generalisation of the constant-
rate model $\mathcal{M}_{\mathrm{CR}}$, in which the entry rate μ in the root is replaced by an increasing time-dependent
rate function $\lambda(t)$, leading to a non-homogeneous Poisson process. Once across the barrier, bacteria
again reproduce deterministically following Malthusian growth at rate r. Equation (2) thus remains
unchanged, but the $\{\tau_{i,j}, j \geq 1\}$ are now the points of an inhomogeneous Poisson point process on $\mathbb{R}_+$
with intensity $t \mapsto \lambda(t)$.

To maintain consistency with the observed arrival-time profiles in the Bolthausen–Sznitman model,
we define the rate as

$$\lambda(t) = \mu\, e^{krt},$$

with $k = 0.78$. This value was chosen because it yields the closest match between the arrival-time
distributions produced by the $\mathcal{M}_{\mathrm{IR}}$ model and those observed in the Bolthausen–Sznitman model, even
though the underlying parameter values differ between the two models (see Supplementary Fig. 4).
This exponential increase in the entry rate over time reflects a progressively intensifying inflow of
individuals through the barrier. We also make this choice in order to avoid introducing an additional
parameter in the model, to simplify the comparison with other models. The scaling factor rt ensures
that the frequencies produced by this model remain independent of the Malthusian growth rate r.

To clarify this point, consider the non-homogeneous Poisson process with intensity

$$\lambda_s(t) = s\,\frac{\mu}{r}\,e^{kst}.$$

681 In this process, the arrival time $t_{i,s}$ of the ith bacterium is proportional to $1/s$, that is, $t_{i,s} = t_{i,1}/s$.
682 Therefore, the relative frequency of the $(i+1)$th bacterium to the ith is

$$\begin{aligned}
\frac{x_{i+1,s}}{x_{i,s}} &= \exp\left[r(t_{i+1,s} - t_{i,s})\right] \\
&= \exp\left[\frac{r}{s}(t_{i+1,1} - t_{i,1})\right].
\end{aligned}$$

683 Thus, with the crossing intensity $\lambda_s(t)$ and Malthusian growth rate r, the relative frequencies are the
684 same as with the crossing intensity $\lambda_1(t)$ and growth rate r/s. Taking the scaling factor $s = r$, we see
685 that the model with crossing intensity $\lambda(t)$ and growth rate r is equivalent, in terms of frequencies, to
686 the model with crossing intensity

$$\lambda_1(t) = \frac{\mu}{r}\,e^{kt}$$

and growth rate 1.

Hence, as in the $\mathcal{M}_{\mathrm{CR}}$ model, only the ratio μ/r is identifiable from the observed frequencies F.
Consequently, we arbitrarily fix $r = 1$ and $t = 10$, and determine the value $\theta_{\mathrm{IR}}^* = (\mu/r)^*$ that maximises
the performance $P[\mathcal{M}_{\mathrm{IR}}(\theta_{\mathrm{IR}} = \mu/r)]$.

Summary of the models.

- In $\mathcal{M}_{\mathrm{STAMP}}(\theta_{\mathrm{STAMP}} = N_b)$, the founding population after the bottleneck consists of exactly
θ_{STAMP} bacteria that cross the barrier simultaneously, selected from the initial population ac-
cording to a multinomial distribution, leading to a stronger founder effect as θ_{STAMP} decreases.

- In $\mathcal{M}_{\mathrm{CR}}(\theta_{\mathrm{CR}} = \mu/r)$, bacteria cross the bottleneck stochastically at a Poisson rate μ and then
grow exponentially at rate r. The ratio $\theta_{\mathrm{CR}} = \mu/r$ characterises the strength of the bottleneck:
a low θ_{CR} results in strong founder effects and reduced diversity.

- In $\mathcal{M}_{\mathrm{BS}}(\theta_{\mathrm{BS}})$, bottleneck crossing is modelled as a stochastic propagation process, where the
parameter θ_{BS} corresponds to the speed at which the population crosses the bottleneck. A lower
speed θ_{BS} increases the amount of coalescence, leading to a stronger bottleneck effect.

- In $\mathcal{M}_{\mathrm{IR}}(\theta_{\mathrm{IR}} = \mu/r)$, the bottleneck is crossed at a time-inhomogeneous rate $\lambda(t) = \frac{\mu}{r}e^{kt/r}$, leading
to progressively more frequent entries as time increases. Bacteria then grow exponentially at rate
r. This variable-rate mechanism generalises the constant entry rate model $\mathcal{M}_{\mathrm{CR}}$ and reproduces
arrival profiles similar to those of the $\mathcal{M}_{\mathrm{BS}}$ model. The parameter $\theta_{\mathrm{IR}} = \mu/r$ again governs the
bottleneck strength, with smaller values indicating a stronger founder effect.

4.9 Description of the Kingman and Bolthausen-Sznitman coalescents

The Kingman coalescent. The Kingman coalescent is an object that follows the genealogy of current individuals. Let n current individuals be labelled by $1, \ldots, n$. The n-Kingman coalescent is a process going backward in time, which, at time t, contains the ancestors of the n current individuals living at time t *before* the present time. More specifically:

1. The initial state, denoted by Π_0, is the set of blocks $\{1\}, \ldots, \{n\}$. Each block corresponds to a single individual living at the present time;

2. Each $t > 0$ corresponds to a time *in the past*. Then Π_t is made of blocks of families of individuals living at the present time. For example, if $\Pi_t = \{1, 2\}, \{3\}, \{4\}$, this means that individuals 1 and 2 share a common ancestor that lived at time $-t$; and 3 and 4 do not share a common ancestor with $1, 2$ that lived at this time;

3. The dynamics of Π_t is as follows: any given pair of blocks merges at rate 1. For example, at rate 1, $\{1, 2\}$ and $\{3\}$ will try to merge to form $\{1, 2, 3\}$. A merger at time t means that the two families have found their most recent common ancestor at time t before present.

The Bolthausen-Sznitman coalescent. The Bolthausen-Sznitman coalescent, similarly to the Kingman coalescent, is a process that goes backward in time and that, at time t, regroups individuals who share a common ancestor at time t *before* the present time. However, the mechanism for the merging of blocks is different from that of the Kingman coalescent. While in the Kingman coalescent, only two blocks can merge at the same time, the Bolthausen-Sznitman coalescent allows *multiple mergers*. There are several constructions of the Bolthausen-Sznitman coalescent. It can be seen as the Λ-coalescent with $\Lambda(dx) = dx$ [43]. However, we define it here in another way, which is apter to simulations, using the construction of reference [44] (see also the description in reference [43]). First, construct a tree recursively in the following way. The root is indexed by 1. Once n vertices are positioned on the tree, the $(n+1)^{th}$ vertex chooses its parent uniformly at random among the n vertices of the tree. Each vertex of the tree corresponds to a block; initially, all blocks contain only a singleton.

The merging procedure is as follows. Each edge of the tree carries an exponential clock with rate 1. If the clock of an edge rings, the edge is "lifted"; namely, all descendants of the edge are absorbed by the parent of the edge. The procedures goes on until there is only one vertex left (see Supplementary Fig. 10).

4.10 Estimation of μ and r using the constant rate or increasing rate models

We consider here the models $\mathcal{M}_{\mathrm{CR}}$ and $\mathcal{M}_{\mathrm{IR}}$. The growth rate is estimated as follows. For each sample, we measure the bacterial population associated with the first arrival as $f_1 \times$ (bacterial load), where f_1 is the proportion of the majority tag. Assuming Malthusian growth with rate r, this population

size equals $e^{r(t-t_1)}$, where t is the observation time and t_1 is the arrival time of the first tag under the model. For the model $\mathcal{M}_{\mathrm{CR}}$, we use $t_1 \approx E[t_1] = 1/\mu = 1/(r\theta)$. For the model $\mathcal{M}_{\mathrm{IR}}$, we approximate t_1 by the median value of the first arrival time, that we compute as follows. We use the cumulative intensity function:

$$\Lambda(t) = \int_0^t \lambda(s)\,ds = \mu \cdot \frac{e^{krt} - 1}{kr}.$$

The cumulative distribution function of the first arrival time is:

$$F(t) = 1 - \exp(-\Lambda(t)).$$

The median t_{med} satisfies $F(t_{\mathrm{med}}) = 0.5$, *i.e.*,

$$\exp\left(-\mu \cdot \frac{e^{krt_{\mathrm{med}}} - 1}{kr}\right) = 0.5.$$

Thus,

$$\mu \cdot \frac{e^{krt_{\mathrm{med}}} - 1}{kr} = \ln(2).$$

and

$$t_1 \approx t_{\mathrm{med}} = \frac{1}{kr}\ln\left(1 + \frac{kr}{\mu}\ln(2)\right) = \frac{1}{kr}\ln\left(1 + \frac{k}{\theta}\ln(2)\right).$$

We then approximate r using the formula:

$$r \approx \frac{\ln(f_1 \times \text{bacterial load})}{t - t_1}.$$

For the model $\mathcal{M}_{\mathrm{CR}}$, this leads to

$$r_{\mathrm{CR}} \approx \frac{\ln(f_1 \times \text{bacterial load}) + \frac{1}{\theta}}{t}.$$

For the model $\mathcal{M}_{\mathrm{IR}}$, we get:

$$r_{\mathrm{IR}} \approx \frac{\ln(f_1 \times \text{bacterial load}) + \frac{1}{k}\ln\left(1 + \frac{k}{\theta}\ln(2)\right)}{t}.$$

References

[1] Sören Abel, Pia Abel zur Wiesch, Hsiao-Han Chang, Brigid M. Davis, Marc Lipsitch, and Matthew K. Waldor. STAMP: Sequence tag-based analysis of microbial population dynamics. *Nature methods*, 12(3):223, 2015. doi: 10.1038/nmeth.3253. Publisher: NIH Public Access.

[2] Karthik Hullahalli, Justin R Pritchard, and Matthew K Waldor. Refined quantification of infection bottlenecks and pathogen dissemination with STAMPR. *MSystems*, 6(4):10–1128, 2021.

[3] Anel Mahmutovic, Aaron Nicholas Gillman, Silje Lauksund, Natasha-Anne Robson Moe, Aime

Manzi, Merete Storflor, Pia Abel zur Wiesch, and Sören Abel. RESTAMP – Rate estimates by sequence-tag analysis of microbial populations. *Computational and Structural Biotechnology Journal*, 19:1035–1051, January 2021. ISSN 2001-0370. doi: 10.1016/j.csbj.2021.01.017.

[4] Ian W. Campbell, Karthik Hullahalli, and Matthew K. Waldor. Quantifying host-microbe interactions with bacterial lineage tracing. *Science*, 391(6780):34–40, January 2026. doi: 10.1126/science.adx5362. Publisher: American Association for the Advancement of Science.

[5] Marc Planas-Marquès, Jonathan P Kressin, Anurag Kashyap, Dilip R Panthee, Frank J Louws, Nuria S Coll, and Marc Valls. Four bottlenecks restrict colonization and invasion by the pathogen Ralstonia solanacearum in resistant tomato. *Journal of Experimental Botany*, 71(6):2157–2171, March 2020. ISSN 0022-0957. doi: 10.1093/jxb/erz562.

[6] Catherine Digonnet, Yves Martinez, Nicolas Denancé, Marine Chasseray, Patrick Dabos, Philippe Ranocha, Yves Marco, Alain Jauneau, and Deborah Goffner. Deciphering the route of Ralstonia solanacearum colonization in Arabidopsis thaliana roots during a compatible interaction: focus at the plant cell wall. *Planta*, 236(5):1419–1431, November 2012. ISSN 1432-2048. doi: 10.1007/s00425-012-1694-y.

[7] Marie Turner, Alain Jauneau, Stéphane Genin, Marie-José Tavella, Fabienne Vailleau, Laurent Gentzbittel, and Marie-Françoise Jardinaud. Dissection of Bacterial Wilt on Medicago truncatula Revealed Two Type III Secretion System Effectors Acting on Root Infection Process and Disease Development. *Plant Physiology*, 150(4):1713–1722, August 2009. ISSN 0032-0889. doi: 10.1104/pp.109.141523.

[8] Denise Caldwell, Bong-Suk Kim, and Anjali S. Iyer-Pascuzzi. Ralstonia solanacearum Differentially Colonizes Roots of Resistant and Susceptible Tomato Plants. *Phytopathology®*, 107(5):528–536, May 2017. ISSN 0031-949X. doi: 10.1094/PHYTO-09-16-0353-R. Publisher: Scientific Societies.

[9] Kelly E. R. Bachta, Jonathan P. Allen, Bettina H. Cheung, Cheng-Hsun Chiu, and Alan R. Hauser. Systemic infection facilitates transmission of Pseudomonas aeruginosa in mice. *Nature Communications*, 11(1):543, January 2020. ISSN 2041-1723. doi: 10.1038/s41467-020-14363-4. Publisher: Nature Publishing Group.

[10] Caitlyn L. Holmes, Katherine G. Dailey, Karthik Hullahalli, Alexis E. Wilcox, Sophia Mason, Bridget S. Moricz, Lavinia V. Unverdorben, George I. Balazs, Matthew K. Waldor, and Michael A. Bachman. Patterns of Klebsiella pneumoniae bacteremic dissemination from the lung. *Nature Communications*, 16(1):785, January 2025. ISSN 2041-1723. doi: 10.1038/s41467-025-56095-3.

[11] D. S. Falconer and T. F. C. Mackay. *Introduction to Quantitative Genetics*. Longmans Green, Harlow, Essex, UK, ed 4 edition, 1996.

[12] Costas B. Krimbas and Spyros Tsakas. The genetics of dacus oleae. v. changes of esterase poly-morphism in a natural population following insecticide control-selection or drift? *Evolution; International Journal of Organic Evolution*, 25(3):454–460, September 1971. ISSN 1558-5646. doi: 10.1111/j.1558-5646.1971.tb01904.x.

[13] Gabriel Birzu, Oskar Hallatschek, and Kirill S. Korolev. Genealogical structure changes as range expansions transition from pushed to pulled. *Proceedings of the National Academy of Sciences*, 118(34):e2026746118, August 2021. doi: 10.1073/pnas.2026746118. Publisher: Proceedings of the National Academy of Sciences.

[14] Tom Leinster and Christina A Cobbold. Measuring diversity: the importance of species similarity. *Ecology*, 93(3):477–489, 2012.

[15] J. F. Wang, J. Olivier, P. Thoquet, B. Mangin, L. Sauviac, and N. H. Grimsley. Resistance of tomato line Hawaii7996 to Ralstonia solanacearum Pss4 in Taiwan is controlled mainly by a major strain-specific locus. *Molecular plant-microbe interactions: MPMI*, 13(1):6–13, January 2000. ISSN 0894-0282. doi: 10.1094/MPMI.2000.13.1.6.

[16] Ai Xu, Lan Wei, Jingjing Ke, Chengfeng Peng, Pengyue Li, Changqiu Fan, Xiao Yu, and Bo Li. ETI signaling nodes are involved in resistance of Hawaii 7996 to Ralstonia solanacearum-induced bacterial wilt disease in tomato. *Plant Signaling & Behavior*, 18(1):2194747, 2023. ISSN 1559-2316. doi: 10.1080/15592324.2023.2194747.

[17] Karthik Hullahalli and Matthew K Waldor. Pathogen clonal expansion underlies multiorgan dissemination and organ-specific outcomes during murine systemic infection. *eLife*, 10:e70910, 2021. ISSN 2050-084X. doi: 10.7554/eLife.70910.

[18] Victoria Chevée, Karthik Hullahalli, Katherine G. Dailey, Leslie Güereca, Chenyu Zhang, Matthew K. Waldor, and Daniel A. Portnoy. Temporal and spatial dynamics of Listeria monocy-togenes central nervous system infection in mice. *Proceedings of the National Academy of Sciences*, 121(17):e2320311121, April 2024. doi: 10.1073/pnas.2320311121. Publisher: Proceedings of the National Academy of Sciences.

[19] Isai Salas-González, Guilhem Reyt, Paulina Flis, Valéria Custódio, David Gopaulchan, Niokhor Bakhoum, Tristan P. Dew, Kiran Suresh, Rochus Benni Franke, Jeffery L. Dangl, David E. Salt, and Gabriel Castrillo. Coordination between microbiota and root endodermis supports plant mineral nutrient homeostasis. *Science*, 371(6525):eabd0695, January 2021. doi: 10.1126/science.abd0695.

[20] Marie Barberon and Niko Geldner. Radial transport of nutrients: the plant root as a polarized epithelium. *Plant Physiology*, 166(2):528–537, October 2014. ISSN 1532-2548. doi: 10.1104/pp.114.246124.

[21] Marie Barberon, Joop Engelbertus Martinus Vermeer, Damien De Bellis, Peng Wang, Sadaf
Naseer, Tonni Grube Andersen, Bruno Martin Humbel, Christiane Nawrath, Junpei Takano,
David Edward Salt, and Niko Geldner. Adaptation of Root Function by Nutrient-Induced Plas-
ticity of Endodermal Differentiation. *Cell*, 164(3):447–459, January 2016. ISSN 1097-4172. doi:
10.1016/j.cell.2015.12.021.

[22] Huei-Hsuan Tsai, Yuanjie Tang, Lingmin Jiang, Xiaoyan Xu, Valérie Dénervaud Tendon, Jia
Pang, Yanyan Jia, Kathrin Wippel, Jordan Vacheron, Christoph Keel, Tonni Grube Andersen,
Niko Geldner, and Feng Zhou. Localized glutamine leakage drives the spatial structure of root
microbial colonization. *Science*, 390(6768):eadu4235, October 2025. doi: 10.1126/science.adu4235.
Publisher: American Association for the Advancement of Science.

[23] Caroline Baroukh, Meriem Zemouri, and Stéphane Genin. Trophic preferences of the pathogen
Ralstonia solanacearum and consequences on its growth in xylem sap. *MicrobiologyOpen*, 11(1):
e1240, February 2022. ISSN 2045-8827. doi: 10.1002/mbo3.1240.

[24] Marine Lebrun-Corbin, Bettina H. Cheung, Karthik Hullahalli, Katherine G. Dailey, Keith Bai-
ley, Matthew K. Waldor, Richard G. Wunderink, Kelly E. R. Bachta, and Alan R. Hauser. Pseu-
domonas aeruginosa population dynamics in a vancomycin-induced murine model of gastrointesti-
nal carriage. *mBio*, 16(5):e0313624, May 2025. ISSN 2150-7511. doi: 10.1128/mbio.03136-24.

[25] Aaron Nicholas Gillman, Anel Mahmutovic, Pia Abel zur Wiesch, and Sören Abel. The Infectious
Dose Shapes Vibrio cholerae Within-Host Dynamics. *mSystems*, 6(6):e00659–21, 2021. ISSN
2379-5077. doi: 10.1128/mSystems.00659-21.

[26] Myrto Vlazaki, John Huber, and Olivier Restif. Integrating mathematical models with experimen-
tal data to investigate the within-host dynamics of bacterial infections. *Pathogens and Disease*,
77(8):ftaa001, November 2019. ISSN 2049-632X. doi: 10.1093/femspd/ftaa001.

[27] Ting Zhang, Sören Abel, Pia Abel Zur Wiesch, Jumpei Sasabe, Brigid M. Davis, Darren E.
Higgins, and Matthew K. Waldor. Deciphering the landscape of host barriers to Listeria mono-
cytogenes infection. *Proceedings of the National Academy of Sciences of the United States of
America*, 114(24):6334–6339, June 2017. ISSN 1091-6490. doi: 10.1073/pnas.1702077114.

[28] Xue Liu, Jacqueline M. Kimmey, Laura Matarazzo, Vincent de Bakker, Laurye Van Maele, Jean-
Claude Sirard, Victor Nizet, and Jan-Willem Veening. Exploration of bacterial bottlenecks and
Streptococcus pneumoniae pathogenesis by CRISPRi-seq. *Cell host & microbe*, 29(1):107–120.e6,
January 2021. ISSN 1931-3128. doi: 10.1016/j.chom.2020.10.001.

[29] Sarah E Woodward, Stefanie L Vogt, Jorge Peña-Díaz, Ryan A Melnyk, Mihai Cirstea, Antonio
Serapio-Palacios, Laurel M P Neufeld, Kelsey E Huus, Madeline A Wang, Cara H Haney, and
B Brett Finlay. Gastric acid and escape to systemic circulation represent major bottlenecks to

host infection by Citrobacter rodentium. *The ISME Journal*, 17(1):36–46, January 2023. ISSN 1751-7362. doi: 10.1038/s41396-022-01321-9.

[30] Ian W. Campbell, Karthik Hullahalli, Jerrold R. Turner, and Matthew K. Waldor. Quantitative dose-response analysis untangles host bottlenecks to enteric infection. *Nature Communications*, 14:456, January 2023. ISSN 2041-1723. doi: 10.1038/s41467-023-36162-3.

[31] Julia A. Hotinger, Ian W. Campbell, Karthik Hullahalli, Akina Osaki, and Matthew K. Waldor. Quantification of Salmonella enterica serovar Typhimurium population dynamics in murine infection using a highly diverse barcoded library. *eLife*, 13:RP101388, February 2025. ISSN 2050-084X. doi: 10.7554/eLife.101388.

[32] Stefan T. Peterson, Katherine G. Dailey, Karthik Hullahalli, Daniel Sorobetea, Rina Matsuda, Jaydeen Sewell, Winslow Yost, Rosemary O'Neill, Suhas Bobba, Nicolai Apenes, Matthew E. Sherman, George I. Balazs, Charles-Antoine Assenmacher, Arin Cox, Matthew Lanza, Sunny Shin, Matthew K. Waldor, and Igor E. Brodsky. TNF signaling maintains local restriction of bacterial founder populations in intestinal and systemic sites during oral Yersinia infection. *mBio*, 0(0):e01779–25, September 2025. doi: 10.1128/mbio.01779-25. Publisher: American Society for Microbiology.

[33] Bolutife Fakoya, Karthik Hullahalli, Daniel H. F. Rubin, Deborah R. Leitner, Roma Chilengi, David A. Sack, and Matthew K. Waldor. Nontoxigenic Vibrio cholerae Challenge Strains for Evaluating Vaccine Efficacy and Inferring Mechanisms of Protection. *mBio*, 13(2):e00539–22, April 2022. doi: 10.1128/mbio.00539-22.

[34] Karthik Hullahalli, Katherine G. Dailey, and Matthew K. Waldor. Innate immune responses yield tissue-specific bottlenecks that scale with pathogen dose. *Proceedings of the National Academy of Sciences*, 120(37):e2309151120, September 2023. doi: 10.1073/pnas.2309151120. Publisher: Proceedings of the National Academy of Sciences.

[35] Jana Ordon, Julien Thouin, Ryohei Thomas Nakano, Ka-Wai Ma, Pengfan Zhang, Bruno Huettel, Ruben Garrido-Oter, and Paul Schulze-Lefert. Chromosomal barcodes for simultaneous tracking of near-isogenic bacterial strains in plant microbiota. *Nature Microbiology*, 9(4):1117–1129, April 2024. ISSN 2058-5276. doi: 10.1038/s41564-024-01619-8. Publisher: Nature Publishing Group.

[36] Benjamin B. J. Daniel, Yves Steiger, Anna Sintsova, Christopher M. Field, Bidong D. Nguyen, Christopher Schubert, Yassine Cherrak, Shinichi Sunagawa, Wolf-Dietrich Hardt, and Julia A. Vorholt. Assessing microbiome population dynamics using wild-type isogenic standardized hybrid (WISH)-tags. *Nature Microbiology*, 9(4):1103–1116, April 2024. ISSN 2058-5276. doi: 10.1038/s41564-024-01634-9. Publisher: Nature Publishing Group.

[37] Rachel T. Giorgio, My T. Le, Ting Zhang, Caitlyn L. Holmes, and Karthik Hullahalli. Accurate

interpretation of within-host dissemination using barcoded bacteria. *mSystems*, 0(0):e01460–25, December 2025. doi: 10.1128/msystems.01460-25. Publisher: American Society for Microbiology.

[38] Freddy Monteiro, Montserrat Solé, Irene van Dijk, and Marc Valls. A Chromosomal Insertion Toolbox for Promoter Probing, Mutant Complementation, and Pathogenicity Studies in Ralstonia solanacearum. *Molecular Plant-Microbe Interactions®*, 25(4):557–568, April 2012. ISSN 0894-0282. doi: 10.1094/MPMI-07-11-0201. Publisher: Scientific Societies.

[39] Nadin Rohland and David Reich. Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Research*, 22(5):939–946, May 2012. ISSN 1088-9051. doi: 10.1101/gr.128124.111.

[40] L Roques, J Garnier, F Hamel, and E K Klein. Allee effect promotes diversity in traveling waves of colonization. *Proc Natl Acad Sci USA*, 109(23):8828–8833, 2012.

[41] É. Brunet, B. Derrida, A. H. Mueller, and S. Munier. Effect of selection on ancestry: An exactly soluble case and its phenomenological generalization. *Phys. Rev. E*, 76:041104, 2007. doi: 10.1103/PhysRevE.76.041104.

[42] Alison Etheridge and Sarah Penington. Genealogies in bistable waves. *Electron. J. Probab.*, 27: Paper No. 121, 99, 2022. ISSN 1083-6489. doi: 10.1214/22-ejp845.

[43] Nathanaël Berestycki. *Recent progress in coalescent theory*, volume 16 of *Ensaios Matemáticos [Mathematical Surveys]*. Sociedade Brasileira de Matemática, Rio de Janeiro, 2009. ISBN 978-85-85818-40-1.

[44] Christina Goldschmidt and James B. Martin. Random recursive trees and the Bolthausen-Sznitman coalescent. *Electron. J. Probab.*, 10:no. 21, 718–745, 2005. ISSN 1083-6489. doi: 10.1214/EJP.v10-265.

[45] Oskar Hallatschek and David R. Nelson. Gene surfing in expanding populations. *Theoretical Population Biology*, 73(1), 2008. ISSN 00405809. doi: 10.1016/j.tpb.2007.08.008.

Data availability

All processed sequence counts and all associated metadata were kept unmodified in a all-in-one Excel data sheet available in supplementary data. Outputs from the time-resolved analysis algorithms are conserved in a semi-controlled local environment (R project "SV_bottleneck_analysis_2024") in supplementary documents. The raw sequences data and corresponding metadata for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB102723 (https://www.ebi.ac.uk/ena/browser/view/PRJEB102723).

Code availability

The code for all analyses is made available in an annotated R Markdown notebook in supplementary documents. The notebook is contained in a semi-controlled local environment inside the R project "SV_bottleneck_analysis_2024" allowing to reproduce the integrality of the presented analyses, figures and statistical tests from the original data sheet (tested environment: Windows 11, RStudio, R version 4.2.2, Python version 3.11.2). The code for the processing of raw sequences data is also provided in the file "SV_nanopore_pipeline_2024" (tested environment: Shell script in Linux Ubuntu with a Conda environment, Python version 3.11.2).

Authors contribution

S.V. and N.P. designed the plant infection experiments. S.V. and N.P. constructed and validated the biological material. S.V., I.M. and N.P. performed the plant infection experiments. S.V. sequenced and processed the sequences data. N.B., R.F. and L.R. designed and formalised the model testing approach and the time-resolved algorithms. S.V., N.B., R.F. and L.R. wrote the code of the notebook. S.V., N.B., R.F., L.R. and N.P. discussed and analysed the data. S.V. wrote an original draft. All authors corrected and contributed to the manuscript.

Acknowledgements

The authors thank Dr. Karthik Hullahalli (ID), Dr. Kelly E. R. Bachta (ID) and Dr. Michael A. Bachman (ID) for their assistance in accessing the animal infection data reanalysed in the study. The authors thank Dr. Alice Guidot (ID), Dr. Philippe Remigi and Dr. Stéphane Genin for their insightful discussions regarding the manuscript. The authors thank Baptiste Mayjonade (ID) and Sandra Moreau for their assistance and the numerous insights during the establishment of Nanopore sequencing protocols.

This work was supported by the following grants. S.V. was supported by a PhD grant from the Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation. N.P. and S.V. were supported by a seed funding "STAMP" from INRAE department Plant Health and Environment. N.B., R.F. and L.R. were supported by the ANR project ReaCh (ANR-23-CE40-0023-01). N.B. and R.F. were supported by the Chaire Modélisation Mathématique et Biodiversité (École Polytechnique, Muséum national d'Histoire naturelle, Fondation de l'École Polytechnique, VEOLIA Environnement).

Conflicts of interest statement

The authors declare no competing interests.

 # Supplementary Material

Model	Average error	θ M-Intact-27	θ M-Cut-27	θ H-Cut-27	θ M-Intact-32	θ M-Cut-32	θ H-Intact-32	θ H-Cut-32
$\mathcal{M}_{\mathrm{STAMP}}$	0.20 ± 0.01	1.00 ± 0.58	3.08 ± 1.50	2.20 ± 1.29	0.91 ± 0.54	2.65 ± 1.20	0.85 ± 0.80	2.57 ± 1.20
$\mathcal{M}_{\mathrm{CR}}$	0.12 ± 0.01	1.00 ± 0.58	2.14 ± 0.81	1.72 ± 0.77	0.92 ± 0.53	2.17 ± 0.77	0.74 ± 0.65	2.18 ± 0.79
$\mathcal{M}_{\mathrm{BS}}$	0.06 ± 0.01	1.00 ± 0.29	2.20 ± 0.90	2.35 ± 1.01	0.97 ± 0.37	1.98 ± 0.64	1.02 ± 0.39	1.91 ± 0.51
$\mathcal{M}_{\mathrm{IR}}$	0.08 ± 0.02	1.00 ± 0.74	1.92 ± 0.59	1.98 ± 0.62	1.14 ± 0.60	2.11 ± 0.51	0.75 ± 0.57	2.33 ± 0.47

Supplementary Table 1: Score of error (mean $\pm$ half-width of the 95% confidence interval) and mean normalised bottleneck parameter θ estimated for each infection condition from 2 and 5 (mean $\pm$ half-width of the 95% CI). Values of θ are normalised by the control condition M_Intact_27. Condition *H_Intact_27* was tolerant and did not yield sufficient colonisation to allow sequencing of the diversity and estimation of θ.

Supplementary figure 1

Time of sampling for all tested conditions and proportion of symptomatic plants at collection time

Bacterial load obtained at collection time for each condition

Supplementary Figure 1: Day of sampling and proportion of symptomatic plants corresponding to the all conditions in Fig. 5. Conditions *M_Intact_27* and *M_Cut_27* correspond to the data shown in Fig. 2). At the first sign of symptoms on a batch of a replicate (replicate A, B or C), the $n = 8$ plants of the batch were collected. Batches that showed no symptoms were otherwise collected at 13-14 DPI. Bacterial load was systematically quantified. Conditions that were sampled earlier showed systematically higher or similar loads compared to conditions sampled later. Ultimately, the plants that were collected earlier also corresponded to a faster colonisation in CFU but also had faster symptoms development indicating a good representation of a gradient of susceptibility to tolerance.

Supplementary figure 2

Individual assessment of the ability of Nb and Nr to correctly explain tags richness in example samples

Supplementary Figure 2: Assessment of the behaviour of N_b and N_r as a function of richness in sample (log-log scale). A reference line $y = x$ represents richness, values of N_b or N_r under the line are lower than richness in the same sample. A line connects N_b and N_r calculated from the same sample. Illustrative examples of diversity profiles are shown in three samples of intact Marmande plants (plants 40, 34 and 39) or three samples of root-cut Marmande plants (plants 51, 17, 6) (log scale). Tags are sorted in decreasing frequency. Green and red lines represent the number of tags that are successfully represented by N_b and N_r respectively.

Supplementary figure 3

Instantaneous bottleneck models:

STAMP model:
instantaneous bottleneck

STAMPR model: multiple
instantaneous bottlenecks

Time-resolved bottleneck models:

Constant entry rate
model (CR)

Bolthausen-Sznitman range
expansion model (BS)

Increasing entry rate
model (IR)

Supplementary Figure 3: Models tested in the study. **"Instantaneous bottleneck" (STAMP) model** underlying the N_b equation[1]. N_b represents a "number of founders", implicitly considering a bottleneck as a single, instantaneous founder event separating two non-overlapping populations[11,12]. Four assumptions are implied[11]: (1) neutral diversity: all tags have identical growth and virulence fitness; (2) compartmentalised space: soil and root are distinct compartments; (3) spatial homogeneity: all tags have equal access to root, tissue complexity is ignored; (4) instantaneous barrier: a single, instantaneous entry with no secondary infections. Under these assumptions, entry is quasi-instantaneous and decoupled from time-dependant effects. Assumptions (1) and (2) are supported by knowledge of the infection and the neutral tags collection. Assumptions (3) and (4) remain untested in the literature and are contradicted by the observation of motility and growth during colonisation[5,6,7,8]. **"Multiple instantaneous bottlenecks" (STAMPR) model** empirically interpreted from the N_r and N_s algorithms[2]. Colonisation occurs over k events of instantaneous entry, each contributing to $N_{b,k}$ new founders, each k subpopulation representing a proportion of the total population. Assumptions of instantaneous entry are inherited from the STAMP model. As STAMPR is an empirical correction rather than a mechanistic model, it cannot be mathematically investigated beyond empirical checks. **"Constant entry rate" (CR) model**. Plant colonisation is the combination of a constant entry process and a pure growth process. Pathogens enter the roots at a constant entry rate (or permeability) μ_{CR} while already present pathogens grow at an exponential rate r_{CR}. The bottleneck metric $\theta_{CR} = \mu_{CR}/r_{CR}$ is the ratio of entry over occupation of space by growth and informs on the quantity of unique cells entering the plant before saturation. Low θ_{CR} indicates a colonisation that is mostly governed by growth (a stringent bottleneck). **"Bolthausen-Sznitman" (BS) model**. Root barriers breaching is modelled by a range expansion in a finite population[13,45]. Pathogen cells migrate randomly within the barrier and divide at a density-dependent rate (cells in more crowded areas divide less often). Cells that manage to cross this barrier then colonise the plant. Resulting sampling of tags can be obtained using the Bolthausen-Sznitman coalescent (see Section 4.8). The bottleneck metric $\theta_{BS} = v$ is the relative speed at which pathogens can cross the root barriers and reach the final compartment. Low value indicates slow breaching of the roots barriers allowing for more branching of pathogen cells lines (a stringent bottleneck). **"Increasing entry rate" (IR) model**, a generalisation of the CR model. Colonisation of the plant is a combination of an exponentially increasing entry process and a pure growth process. As entry increases over time, an entry rate (or permeability) multiplier μ_{IR} governs the acceleration of entry of pathogen over time. Simultaneously, already present pathogens grow at an exponential rate r_{IR}. Similarly to the CR model, low $\theta_{IR} = \mu_{IR}/r_{IR}$ represents a stringent bottleneck.

Supplementary figure 4

Probability of entry of successive tags in the the four tested models

Supplementary Figure 4: Comparison of arrival time distributions under the four tested models. The first bacterium enters the root at a normalised time 0. Each curve represents the Cumulative Distribution Function (CDF) of the arrival time of the k^{th} bacterium relative to the first one. In this illustration, the parameter values are: $\mu_{\text{CR}}/r = 2$ for $\mathcal{M}_{\text{CR}}$, $v = 10$ for $\mathcal{M}_{\text{BS}}$ and $\mu_{\text{CR}}/r = 0.1$ for $\mathcal{M}_{\text{IR}}$.

Supplementary figure 5

Hill-number curves in samples and the respective best fit in each model

Mean relative error per condition

Mean relative error on all conditions

Supplementary Figure 5: Diversity profiles in samples and simulations according to the Hill numbers from order $q = 0$ (giving more importance to richness) to order $q = 4$ (giving more importance to evenness). For a given order q, a high value $^q H$ indicates a high diversity. Each colour corresponds to a sample profile from Fig. 5 including those of Fig. 2. An additional model $\mathcal{M}_K$ (Kingman model) was tested but is not presented in the main text. $\mathcal{M}_K$ provides a simpler equation to describe biological processes of $\mathcal{M}_{BS}$ but was globally unable to accurately describe the data. For each model, a score of error represents the difference between the Hill-number curve in observed data versus in best fit from a model. The error can be averaged on each condition or on all samples.

Supplementary figure 6

PERMANOVA test: contribution of all plant condition parameters to θ_{BS}

	Df	SumOfSqs	R2	parOmegaSq	F	Pr(>F)	
`Root condition`	1	2.3786	0.16279	0.155298	28.0258	0.000999	***
Genotype	1	0.0234	0.00160	-0.004948	0.2762	0.740260	
`Temperature (°C)`	1	0.0725	0.00496	-0.000997	0.8536	0.394605	
Residual	143	12.1367	0.83064				
Total	146	14.6112	1.00000				

Signif. codes: 0 `***` 0.001 `**` 0.01 `*` 0.05 `.` 0.1 ` ` 1

PERMANOVA test: contribution of all plant condition parameters to r_{IR}

	Df	SumOfSqs	R2	parOmegaSq	F	Pr(>F)	
`Root condition`	1	0.2197	0.03003	0.08524	14.698	0.000999	***
Genotype	1	4.7241	0.64577	0.68184	316.039	0.000999	***
`Temperature (°C)`	1	0.2341	0.03200	0.09069	15.661	0.000999	***
Residual	143	2.1375	0.29220				
Total	146	7.3154	1.00000				

Signif. codes: 0 `***` 0.001 `**` 0.01 `*` 0.05 `.` 0.1 ` ` 1

PERMANOVA test: contribution of collected sample weight to r_{IR}

	Df	SumOfSqs	R2	parOmegaSq	F	Pr(>F)	
`Collected fresh weight (g)`	1	2.7285	0.37298	0.55453	183.9912	0.000999	***
`Root condition`	1	0.2653	0.03627	0.10307	17.8923	0.000999	***
Genotype	1	2.1239	0.29033	0.49173	143.2182	0.000999	***
`Temperature (°C)`	1	0.0919	0.01256	0.03415	6.1979	0.005994	**
Residual	142	2.1058	0.28786				
Total	146	7.3154	1.00000				

Signif. codes: 0 `***` 0.001 `**` 0.01 `*` 0.05 `.` 0.1 ` ` 1

PERMANOVA test: contribution of all plant condition parameters to μ_{IR}

	Df	SumOfSqs	R2	parOmegaSq	F	Pr(>F)	
`Root condition`	1	6.8427	0.24463	0.256070	51.5991	0.000999	***
Genotype	1	1.4379	0.05141	0.062758	10.8432	0.000999	***
`Temperature (°C)`	1	0.7270	0.02599	0.029590	5.4823	0.008991	**
Residual	143	18.9637	0.67797				
Total	146	27.9713	1.00000				

Signif. codes: 0 `***` 0.001 `**` 0.01 `*` 0.05 `.` 0.1 ` ` 1

PERMANOVA test: contribution of collected sample weight to μ_{IR}

	Df	SumOfSqs	R2	parOmegaSq	F	Pr(>F)	
`Collected fresh weight (g)`	1	1.3196	0.04718	0.057148	9.9099	0.000999	***
`Root condition`	1	6.9936	0.25003	0.259521	52.5200	0.000999	***
Genotype	1	0.4143	0.01481	0.014162	3.1116	0.043956	*
`Temperature (°C)`	1	0.3351	0.01198	0.010213	2.5168	0.085914	.
Residual	142	18.9087	0.67600				
Total	146	27.9713	1.00000				

Signif. codes: 0 `***` 0.001 `**` 0.01 `*` 0.05 `.` 0.1 ` ` 1

Plot of growth rate r_{IR} in function of permeability μ_{IR}

Supplementary Figure 6: Statistics performed on the estimated parameters from Fig. 5c, 5d, 5e. Non-parametric PERMANOVA tests were performed to see the relative contribution of the three tested biological factors (*plant genotype*, *roots condition* and *temperature*) on θ_{BS}, μ_{IR} and r_{IR}. Another PERMANOVA test was performed on μ_{IR} and r_{IR} with the addition of root sample weight as a proxy for a *plant size* factor. As a guide, "Pr (¿F)" and "R2" respectively represents the p-value and the coefficient of determination R^2 for each tested factor. Additionally, "parOmegaSq" represents the effect size ω^2 of each factor (> 0.1: small effect; $> 0,3$: medium effect, > 0.5 large effect). Growth rate r_{IR} was plotted in function of permeability μ_{IR}. Each cross on the graph represents the average value for each condition. Values of r_{IR} and μ_{IR} correlate positively (Spearman ranks correlation, $R^2 = 0.40$, $p = 5.97 \times 10^{-7}$).

Supplementary figure 7

Mean relative error of the five tested models in each condition of the _P. aeruginosa dataset_

Supplementary Figure 7: Score of error for the models in the range $q = 0$ to $q = 4$ on the _P. aeruginosa_ infection dataset reanalysed in Fig. 6. An additional model $\mathcal{M}_{\text{K}}$ (Kingman model) was tested but is not presented in the main text. $\mathcal{M}_{\text{K}}$ provides a simpler equation to describe biological processes of $\mathcal{M}_{\text{BS}}$ but was globally unable to accurately describe the data. For each model, a score of error represents the difference between the Hill-number curve in observed data versus in best fit from a model. The error can be averaged for each organ or on all samples.

Supplementary figure 8

Mean relative error of the models over all conditions of the *K. pneumoniae* dataset

Bottleneck estimators: θ_{IR} versus N_S

Bacterial expansion estimators: r_{IR} versus CFU/Ns and Ns/Nb

Permeability estimator: no equivalents of μ_{IR}

Colonisation proxy: r_{IR} versus CFU when comparing *direct* and *metastatic* disseminations in 24h samples

Supplementary Figure 8: Time-resolved models outperform STAMPR to describe mouse-*K. pneumoniae* infection (original data from Fig. 1 of article[10] ($n = 17$ and 9 mice sampled at 24h and 48h respectively). The averaged score of error of the models over $q = 0\text{-}4$ showed a better fit of $\mathcal{M}_{\mathrm{IR}}$ over $\mathcal{M}_{\mathrm{STAMP}}$. Model-based estimators ($\theta_{\mathrm{IR}}, \mu_{\mathrm{IR}}, r_{\mathrm{IR}}$) were compared to their closest counterpart in STAMPR as follow: θ_{IR} *vs.* N_s for the bottleneck effect; r_{IR} *vs.* CFU/Ns and Ns/Nb for bacterial expansion; μ_{IR} had no equivalent in STAMPR. A comparison of r_{IR} *vs.* CFU showcases colonisation of organs at 24h between the infections characterised as *direct* or *metastatic* in article[10]. Performances of estimators were assessed using Student's two-sided t-test and Cohen's D effect size between samples collected at 24h and 48h post-infection. For STAMPR metrics ($N_s, CFU/N_s, N_s/N_b, CFU$), tests were performed after a log10 transformation (as in article[10]). Greater values of Cohen's D (positively or negatively) indicates better statistical power at discriminating 24h and 48h.

Supplementary figure 9

In vitro growth curves for the 120 tagged strains used in study and the distribution of the estimated growth rate for each strain

Diversity profiles in the co-cultures of the 120 tagged strains used for plant infection and distribution of each individual tag

Frequencies for each of the 120 tags among all samples (range 0 to 0.1 and range 0.1 to 1)

Supplementary Figure 9: **Validation of the 120 tagged strains *in vitro*.** 120 isogenic tagged strains were first grown individually in microplate cultures and monitored for their growth curve through OD_{600nm} (logarithmic scale). Growth of the 120 strain was investigated in a violin plot (thick bar: median; thin bars: 1st and 3rd quartile) and in a quantile-quantile plot against a normal distribution. No visual deviation from normality could be observed suggesting a similar growth behaviour between the strains (average growth rate $r \approx 0.176h^{-1}$). Estimation of the growth rate was also individually investigated for each strain (error bars: standard deviation on the growth rate). **Validation of the 119-tags co-cultures *in vitro*.** All co-cultures used for plant inoculation were sequenced. Visually, diversity profiles were stable across each independently prepared inoculum verifying that no technical bottlenecks were introduced at inoculum preparation. The relative frequencies at which tags appear in inocula are plotted (red bars: average across inocula; global red line: perfect frequency of 1/120). Tag007 was mistakenly forgotten from the initial pool while Tag016 was mistakenly added twice to the pool. The average frequencies did not significantly differ from the expectation of a distribution centered on a median of 1/120 (Wilcoxon-Mann-Whitney test, two-sided, $mu = 1/120$, $p = 0.74$), overall showing no indications that the 119 final tags differ in growth in co-culture. **Validation of the 119-tags collection *in planta* for the study of population dynamics.** Relative frequencies of tags in all samples pooled from Fig. 5 including samples from Fig. 2. The frequencies are observed independently of condition (total: $n = 131$ sequenced samples on 6 time-independent infections; red bars: average frequency across samples; global red line: perfect frequency of 1/120). For visual clarity, the frequencies are separated in two ranges (0 to 0.1 and 0.1 to 1) and the points within the boxplots are not shown. Under an expectation of neutral fitness, the frequencies averaged over a large number of events of infection should tend toward the initial inocula frequencies regardless of condition. The average frequencies did not significantly differ from the expectation of a distribution centred on a median of 1/120 (Wilcoxon-Mann-Whitney test, two-sided, $mu = 1/120$, $p = 0.38$), overall showing no detectable difference from the assumption of neutral fitness at a collection level nor systematic sequencing bias.

Supplementary Figure 10: The construction of the Bolthausen-Sznitman coalescent of reference[44]. Each vertex corresponds to an individual. The initial tree is a random tree where the individual $n+1$ chooses its parent uniformly at random among $1, \ldots, n$. At each step, an edge (drawn thick on the picture) is lifted.