RNA at Lipid/Water Interfaces: Molecular Insights from Coarse-Grained Simulations and Reflectivity Data Predictions

Mohd Ibrahim,[†] Jürgen Köfinger,[‡] Martin Zacharias,[†] Emanuel Schneck,[¶] and Nadine Schwierz*,[§]

[†]Center for Functional Protein Assemblies, Technical University of Munich, Ernst-Otto-Fischer-Str. 8, 85748 Garching b. München, Germany
[‡]Department of Theoretical Biophysics, Max Planck Institute of Biophysics, Max-von-Laue-Straße. 3, 60438 Frankfurt am Main, Germany
[¶]Institute for Condensed Matter Physics, TU Darmstadt, Hochschulstraße 8, 64289 Darmstadt, Germany
[§]Institute of Physics, University of Augsburg, Universitätsstraße 1, 86159 Augsburg, Germany

E-mail: Nadine.Schwierz@physik.uni-augsburg.de

Abstract

Interactions between RNA and lipids are fundamental for biological processes and are increasingly exploited for RNA delivery by lipid nanoparticles. However, RNA-lipid interactions remain challenging to characterize at the molecular level. Here, we address the modeling of RNA at lipid/water interfaces using coarse-grained (CG) simulations, experimental validation using scattering data and prediction of neutron (NR) and X-ray reflectivity (XRR) profiles from the simulations. Using neutral DOPC and cationic DOTAP bilayers, we show that lipid-RNA interactions depend strongly on RNA secondary structure, with single-stranded regions exhibiting higher interfacial affinity than double-stranded segments. We validate the CG lipid simulations, showing that while they reproduce experimental X-ray scattering data only qualitatively, the agreement improves markedly after backmapping to atomistic resolution followed by energy minimization and short all-atom molecular dynamics simulations. We further simulated distinct tRNA conformations and analyzed the influence of RNA secondary structure, concentration, solvent contrast, and lipid deuteration on NR and XRR signals identifying the conditions under which such experiments probe RNA adsorption and discriminate between different RNA conformations. Together, these results demonstrate that CG simulations combined with reflectivity data provide a powerful approach to probe RNA adsorption and structure at lipid/water interfaces and support the design and interpretation of scattering experiments.

Introduction

Understanding how RNA interacts with lipid assemblies at lipid/water interfaces is essential for elucidating biological processes such as viral entry, membrane fusion, and RNA delivery via lipid nanoparticles.[1–4] The lipid environment can influence RNA conformation, folding, and degradation, all of which are key factors in determining the efficacy of RNA-based thera-

peutics.[2,3,5–10]

Despite their biomedical significance, RNA-lipid interactions remain challenging to characterize at the molecular level under physiologically relevant conditions. XRR and NR are powerful tools for probing such interactions, providing sub-nanometer structural information normal to the interface. Both techniques have been widely applied to study supported lipid bilayers and monolayers at the air/water interface,[11–15] as well as interactions with ions and biomolecules including peptides, and nucleic acids.[16–23] Reflectivity measurements yield scattering length density (SLD) profiles across the interface encoding the spatial distributions of molecular components, such as lipid tails, head groups, and RNA, enabling inference of orientation, insertion, and adsorption geometry.[15,18,24]

However, the interpretation of reflectivity data is nontrivial. The scattering profile cannot be directly inverted into atomic positions, and traditional fitting with slab models lacks molecular details.[25] As a result, the structure of RNA at lipid/water interfaces remains poorly understood. Molecular dynamics (MD) and coarse-grained (CG) simulations address this gap by providing molecular representations of RNA-lipid systems.[26,27] MD simulations reveal lipid packing, RNA binding modes, and hydration structure, and allow the calculation of SLD profiles for direct and model free comparison with experimental scattering data.[19,20,28] For example, simulations combined with NR were used to assess the distribution of ionizable lipids in bilayers and the quality of different simulation models.[20] Integration with X-ray scattering and fluorescence enabled the determination of lipid packing and pH-dependent protonation states in lipid monolayers[19] and the structure and composition of adsorbed mRNA layers.[23] Likewise, combination of SAXS and MD simulations successfully reproduced lyotropic phases with correct mesophase symmetries, water content, and pH-induced structural transitions.[4,29,30] Similarly, CG simulations showed good agreement with NR profiles of lipid bilayers[31] and elucidated pH-dependent interaction between RNA and ionizable lipids,[32] fur-

ther highlighting the benefit of joint simulation and experimental scattering approaches. In addition to a molecular interpretation of scattering signals, simulations can also guide experimental design by enabling *a priori* prediction of reflectivity curves, optimizing parameters like deuteration and sample composition to improve signal-to-noise and reduce beam time.

In this study, we use Martini CG simulations to investigate the interactions of single-stranded (ssRNA), double-stranded RNA (dsRNA), and tRNA with neutral DOPC and cationic DOTAP lipid bilayers. Subsequently, we address the question how RNA adsorption and RNA secondary structure can be detected experimentally via surface scattering techniques including neutron and X-ray reflectivity. We compute the reflectivity profiles to assess how RNA structure, concentration, lipid composition, and solvent contrast influence the corresponding scattering signals.

Methods

System Setup and Simulation Details

We simulated DOPC, DOTAP and mixed DOPC/DOTAP bilayers in the presence of single-stranded RNA (ssRNA), double-stranded RNA (dsRNA), and transfer RNA (tRNA) molecules. For systems containing ss-RNA or dsRNA, five DOPC:DOTAP molar ratios were used: 1:0, 3:1, 1:1, 1:3, and 0:1, resulting in five distinct bilayer simulation setups. For tRNA-containing systems, only 100% DOPC and 100% DOTAP bilayers were simulated.

The lipids were described using the Martini force field (version 2.1), and RNA molecules were modeled using the corresponding Martini RNA force field.[27,33] The initial configurations were generated using the `insane.py` script.[34]

To construct a CG RNA structure, a double-stranded atomistic model was first generated using the `fd_helix.c` code from Ref.,[35] which was subsequently coarse-grained using the `martinize-nucleotide.py` script. ssRNA

was obtained by removing the complementary strand from the double-stranded structure. The RNA molecules were positioned approximately 2 nm away from the bilayer surface. All systems were solvated with CG water, including 10% anti-freeze water beads. CG sodium and chloride ions were added to achieve a physiological salt concentration of 150 mM and to ensure overall charge neutrality.

The ssRNA and dsRNA systems contained 100 lipids per monolayer, while the tRNA systems included 324 lipids per monolayer. The ssRNA sequence was a polyU chain of seven nucleotides, whereas the dsRNA comprised the sequence UCUUCUACUU and its complementary strand, which was used in a previous experimental study on RNA-lipid interactions.[36]

The initial configurations were energy-minimized using the steepest descent algorithm. This was followed by a 100 ns equilibration phase in the NPT ensemble, employing the Berendsen barostat and thermostat to maintain a pressure of 1.0 bar and a temperature of 323 K, respectively.

During the production run, the temperature was maintained at 323 K using the stochastic velocity rescaling thermostat with a time constant of 1.0 ps,[37] and the pressure was controlled semi-isotropically at 1.0 bar using the Parrinello-Rahman barostat with a time constant of 12.0 ps.[38] In all cases, the RNA molecule was restrained to the initial 3D conformation using an elastic network with a force constant of 1000 kJnm^{-2}mol^{-1}.

All simulations were performed using the GROMACS software package (versions 2018 and 2021).[39] Van-der-Waals interactions were truncated at 1.1 nm and smoothly shifted to zero. Long-range electrostatic interactions were treated using the reaction-field method with a 1.1 nm cutoff and a relative dielectric constant of 15. The equations of motion were integrated with a time step of 10 fs. Systems with ssRNA and dsRNA were simulated for 5 μs, while systems containing tRNA in various conformations were simulated for 1–3 μs each. All analyses were performed using built-in GROMACS tools or custom Python scripts based on the MD-Analysis package.[40] Visualizations were generated using VMD software.[41]

All-atom Simulations

We performed all-atom simulation of a pure DOPC bilayer with 100 lipids per leaflet using the Gromacs simulation package. The initial setup was created using the Charmm-gui webserver.[42] The bilayer was hydrated with 50 water molecules per lipid and 150 mM of NaCl was also added. The lipid molecules were described using the Amber lipid14,[43] water with TIP3P[44] and the ions with the Mamatklov-Schwierz force field[45] which was optimized for TIP3P water. The structure obtained from Charmm-gui was energy minimized with a gradient descent algorithm followed by 1.0 ns equilibration with semi-isotropic C-rescale barostat[46] with a time constant of 5.0 ps and the stochastic velocity rescaling thermostat[37] with a time constant of 1.0 ps. The temperature was maintained at 300 K and pressure at 1.0 bar. The production run was carried out for 200 ns with Parrinello-Rahman barostat[38] with 5.0 ps time constant. The Van-der-Waals interactions were cut-off and shifted to zero at 1.2 nm. Electrostatic interactions were evaluated using the particle mesh Ewald (PME) method[47] with a cut-off of 1.2 nm. The electron density profile was calculated after excluding the initial 50 ns of the trajectory for equilibration.

X-ray and Neutron SLD Profiles from CG Simulations

The neutron and X-ray SLD profiles, denoted as $\rho_n(z)$ and $\rho_x(z)$, respectively, along the bilayer normal from simulations were obtained by dividing the simulation box into small slices as:

$$
\begin{aligned}
\rho_n(z) &= \frac{\sum_i n_i(z) b_i}{\Delta V}, \\
\rho_x(z) &= r_e \frac{\sum_i n_i(z) Z_i}{\Delta V} = r_e \rho_e(z)
\end{aligned}
\tag{1}
$$

where $n_i(z)$ is the number of atoms of type i in a slice of volume ΔV centered at z, b_i is the coherent neutron scattering length, Z_i is the number of electrons of atom i, $r_e = 2.82 \times 10^{-6}$ Å

is the classical electron radius and $\rho_e(z)$ is the transverse electron density profile. We employed two distinct methods to assign the scattering length, b_i or the number of electrons, Z_i, to the CG Martini beads:

(i) Direct calculation from CG simulations: In this method, b_i or Z_i was directly assigned to the CG beads and Eqn. 1 was used to calculate $\rho_n(z)$ or $\rho_x(z)$. The b_i, Z_i values for the beads were determined using the Martini mapping file provided with the force field for each molecule. Based on this, the b_i, Z_i value of a bead was computed as the sum of the b_i or Z_i values of all atoms mapped into it, weighted by their fractional contributions. An example mapping for DOPC is shown in Table S1, where the contribution of each atom to a CG bead is specified. This method is highly efficient as it eliminates the need for backmapping and reduces disk space usage. Our scripts that automate this task are freely available at https://git.rz.uni-augsburg.de/cbio-gitpub.

(ii) Backmapping and MD Equilibration: In this method, approximately 2000 coarse-grained frames were backmapped to all-atom resolution using the `backward.py` script.[48]
Each frame was projected to all-atom resolution, followed by energy minimization and a 2.0 ps MD simulation. These backmapped and equilibrated frames were then analyzed using a custom script to compute $\rho_n(z)$ or $\rho_x(z)$ according to Eqn. 1, using known atomic b_i or Z_i values. This method is computationally more expensive and requires substantial disk storage.

Undulation Corrections

NR and XRR curves, as well as scattering form factors, are determined by the neutron or X-ray SLDs along the bilayer normal (Eqn.1). For small bilayer systems ($\sim$100 lipids/monolayer), these density profiles can be obtained by binning the system along the transverse direction, assuming the bilayer to be flat. However, in simulations of larger lipid patches, membrane undulations become significant, and simple binning under the flat-surface assumption produces a smeared-out density profile (Figure S6). Such undulations also exist in experiments at various length scales, but in supported bilayers or in monolayers at the air-water interface they may be suppressed either due to attachment to the planar substrate or the high surface tension of the interface, respectively. Therefore, to enable direct comparison between experiments and simulations, the respective undulations must be removed.[49] To correct for the undulations in the simulations, we employed the method developed by Braun et al.[49] For each frame in the trajectory, this method constructs an undulating reference surface using the Fourier coefficients of selected atom groups in the bilayer, followed by binning with respect to this dynamically defined surface. Further details and evidence of improved agreement with experimental results are provided in the Supporting Information, Section 4, and Figure S6. Our implementation is freely available at https://git.rz.uni-augsburg.de/cbio-gitpub. Note that in simulations of very large lipid bilayer systems, undulations can also arise from neighbor list update artifacts.[50] However, the systems studied here are well below the size at which such effects become significant.

NR and XRR Curves

The NR and XRR curves were obtained from $\rho_n(z)$ and $\rho_x(z)$ using the Abeles matrix formalism,[51] as implemented in the Refnx Python package.[52] The SLD profiles from simulations were discretized into layers of constant values, each 1 Å thick. NR and XRR curves were obtained for two experimental setups (i) silicon–supported lipid bilayers and (ii) lipid monolayers at the air/water interface. In both cases, we used MD simulations of free lipid bilayers in water (see section *System Setup and Simulation Details*). For supported bilayers on silicon substrates, the silicon not present in the simulations was incorporated following our previous work.[20] Specifically, the SLD values for silicon and silicon dioxide layers were set to known values, with the SiO_2 layer thickness fixed at 10 Å, which is typical for such experiments.[20]

In the second setup, the monolayer was constructed by selecting a single leaflet from the bilayer. Given the area per lipid in our simulations (~ 69.4 Å^2), this monolayer setups corresponds to experimental conditions with a surface pressure of approximately 15–20 mN/m.[53]

tRNA Secondary and Tertiary Structure Prediction

We used a tRNA molecule with the sequence `GCGGAUUUAGCUCAGUUGGGAGAGCGCCAGACUGAAGA` `UCUGGAGGUCCUGUGUUCGAUCCACAGAAUUCGCACCA`. The secondary structure of the sequence was predicted using the RNAfold package.[54] The tRNA exhibited a minimum free energy of -22.40 kcal/mol. The minimum energy structure, along with the nine closest structures within 18 kcal/mol and separated by at least 1.5 kcal/mol, were used in the simulations (Figure S7). The RNAComposer webserver was used to generate an atomistic 3D structure for each secondary structure,[55] which was then coarse-grained using the `martinize_nucleotide.py` script.

Results and Discussion

In the following sections, we discuss the interactions of ssRNA and dsRNA with lipid bilayers focusing on the influence of RNA structure and lipid charge. Subsequently, we evaluate the accuracy of the coarse-grained model in reproducing experimental X-ray scattering form factors using different backmapping strategies. Finally, we compute how RNA adsorption affects neutron and X-ray reflectivity curves of supported bilayers and monolayers at the air/water interface. We systematically analyze the effects of contrast conditions, lipid-to-nucleotide ratio, lipid composition, and RNA secondary structure.

Interactions of ssRNA and dsRNA with Lipid Layers of Increasing Charge

We simulated ssRNA and dsRNA on bilayers with increasing fractions of cationic DOTAP lipids using CG simulations. In all cases, the RNA adsorbs at the lipid/water interface and displays distinct interaction modes depending on base-pairing and membrane compositions. For ssRNA, the nucleobases are embedded near the hydrophobic lipid tails and the sugar phosphate backbone is associated with polar head groups (Figure 1A–C), consistent with previous studies.[5,56] As the cationic lipid content increases, electrostatic attraction strengthens ssRNA adsorption, resulting in a more narrow distribution close to the bilayer surface. dsRNA interacts more weakly with neutral bilayers due to base-pairing that shields hydrophobic regions (Figure 1A). On the charged bilayers, dsRNA adsorbs due to the electrostatic attraction but adopts a more diverse range of orientations, particularly at intermediate charge fractions leading to broader distributions (Figure 1E–G and S1).

These results align with experimental and all-atom simulation data showing weaker dsRNA–membrane interactions compared to ssRNA,[12,57] supporting that the CG Martini model captures the key electrostatic and hydrophobic contributions to RNA–bilayer binding. Recent all-atom MD simulations[58] also reported weaker dsRNA interactions with lipid bilayers compared to ssRNA, in agreement with our findings. In those studies, the weaker interaction was attributed to hydrogen bond formation between RNA base pairs and lipid head groups which are not represented in the current Martini CG model.

Finally, the interaction between RNA and lipids also affects the dynamics of both components. The diffusion coefficient of the lipids in the leaflet in contact with RNA is consistently lower than that of the opposite leaflet as expected. The difference between the leaflets increases as the fraction of cationic DOTAP lipids increases (Figure S2G, H). The diffusion coefficient of ssRNA and dsRNA on mixed neutral

Figure 1: **Distribution of ssRNA and dsRNA at the lipid/water interface.** Normalized probability distrivution $p(z)$ of finding a nucleobase (gray) or backbone phosphate (blue) along the membrane normal for ssRNA (top) and dsRNA (bottom). (A) $p(z)$ for ssRNA on a pure DOPC bilayer. Orange spheres indicate the DOPC choline headgroups. (B) $p(z)$ for ssRNA on a mixed DOPC/DOTAP bilayer (1:1 molar ratio). Green spheres represent the DOTAP choline headgroups. (C) $p(z)$ for ssRNA on a pure DOTAP bilayer. (D) CG model of ssRNA, with nucleobase and backbone regions colored gray and blue, respectively. (E–G) $p(z)$ for dsRNA on the bilayer with same composition as above. (H) CG structure of dsRNA, with color scheme identical to (D). Red dashed vertical lines denote the average position of the lipid choline headgroups.

and cationic bilyers is slightly lower compared to fully charged or fully neutral bilayers (Figure S2G, H).

Comparison of Different Backmapping Procedures to Reproduce Experimental Form Factors from X-ray Scattering

Next, we analyzed how well the CG model reproduces the experimental scattering data of lipid bilayers. To this end, we compare the X-ray scattering form factors calculated from DOPC bilayer simulations with experimental form factors obtained from multilamellar lipid stacks.[28] The X-ray scattering form factor ($|F(q_z)|$) is a standard quantity used to validate the structure of lipid bilayers[28,43,59] and probes the transverse electron density profile.

The form factor can be obtained directly from simulations via a Fourier transform of the electron density profiles.

The electron density profiles and corresponding form factors were computed using several CG-to-all-atom reconstruction protocols. We first applied the direct method, where form factors are computed directly from the CG electron density. This approach, while computationally efficient, leads to inaccurate bilayer core densities, particularly for long-tail lipids such as DOPC (Figure 2A). This artifact arises because the Martini force field maps lipid tails of varying length to the same number of beads (e.g., four beads for both DOPC and DPPC), concentrating more electrons in the terminal beads of longer lipids and artificially increasing the central density. However, for lipids with relatively shorter tails like DPPC, such artifacts do not arise (Figure S4).

Figure 2: **Comparison of different backmapping strategies to reproduce experimental X-ray scattering data for a DOPC bilayer.** (A) Electron density $\rho_e(z)$ obtained from all-atom MD and CG simulations with different backmapping protocols. Four CG-to-all-atom backmapping protocolls were assessed: (i) direct CG (no backmapping), (ii) backmapped to all-atom resolution (BM), (iii) backmapped followed by energy minimization (BM+EM), and (iv) backmapped with energy minimization followed by a short 2 ps MD equilibration (BM+EM+2ps MD). (B) Corresponding X-ray scattering form factors derived from each approach, compared to experimental data from Ref.[28]

To improve the agreement with experiments, we employed the standard backmapping pipeline[48] and evaluated three progressively refined backmapping strategies. In the BM (backmapped) protocol, coarse-grained frames were projected to all-atom resolution without relaxation. The resulting electron density profiles still showed central density artifacts comparable to the direct CG method. The BM+EM (backmapped with energy min-

imization) approach applied two sequential minimization, first considering bonded terms only, followed by inclusion of non-bonded interactions. This procedure improved stability but did not significantly alter the density profile. In the final BM+EM+2 ps MD protocol, which adds a short 2 ps MD equilibration with gradually increasing time steps (0.1 to 2.0 fs), the electron density profile converges to the shape and peak-to-peak spacing observed in all-atom simulations and experiments[28] (Figure 2B). The observed minor deviations at the water/lipid interface, where CG simulations show slightly elevated water density, are consistent with the known reduced water penetration in Martini models due to the larger size of CG water beads.[60]

Among all tested methods, only the BM+EM+2 ps MD approach reproduces both the bilayer thickness and the overall shape of the electron density distribution. We therefore adopt the BM+EM+2 ps MD procedure for all subsequent calculations of X-ray and neutron scattering length density profiles and reflectivity signals.

NR and XRR for tRNA at Lipid/Water Interfaces

In the following, we discuss how RNA adsorption influences NR and XRR signals from supported bilayers and monolayers at the air/water interface. We evaluate the effects of contrast conditions, lipid-to-nucleotide ratio (L/N), lipid charge, and RNA conformation. The question of RNA conformation is motivated by experimental studies showing that RNA adsorption is highly sensitive to features such as base-pairing and single-stranded regions,[5,56] as well as by previous applications of reflectivity to resolve protein and peptide conformations at bilayer surfaces.[21,22] To investigate the impact of RNA conformation we used nine distinct tRNA secondary structures (Figure S7). In the simulations, these structures adopt conformation-specific orientations on the neutral DOPC and cationic DOTAP bilayers and show distinct binding geometries and distributions (Figure 3 and Figure S10). The adsorption behavior of

Figure 3: **Simulation snapshots of two tRNA structures interacting with neutral and charged lipid bilayers after 3 μs of simulation time.** (A) tRNA6 on a neutral DOPC bilayer. (B) tRNA9 on a DOPC bilayer. (C) tRNA6 on a positively charged DOTAP bilayer. (D) tRNA9 on a DOTAP bilayer. tRNA6 and tRNA9 are the conformations which exhibited the largest differences in NR and XRR profiles. Top: tRNA secondary and tertiary structures with colors indicating membrane contacts. Red indicates high contact levels, blue low contact levels. For each structure, contact values are normalized to their respective maximum. All other tRNA structures are shown in Figure S8- S9). Bottom: Simulation snapshots of the two tRNA conformations. tRNA backbones are depicted in blue, nucleobases in gray. Choline headgroups of DOPC and DOTAP lipids are shown as orange and green spheres, respectively.

tRNA is consistent with the behavior of ssRNA and dsRNA discussed above. In most cases, tRNA adsorbs to neutral bilayer via the single-stranded regions whereas on cationic bilayers, it interacts via both single and double stranded regions (Figure 3, Figure S8- S9).

We considered two experimentally relevant setups: (i) supported bilayers on silicon substrates,[20] and (ii) monolayers at the air/water interface.[19,23] NR profiles were calculated for four different deuteration conditions: (i) non-deuterated solvent (H_2O) with regular non-deuterated lipid (hLipid), (ii) deuterated solvent (D_2O) with hLipid, (iii) H_2O with deuterated lipid (dLipid), and (iv) D_2O with dLipid. For the monolayer case, we used null contrast conditions ($\sim 92\%$ H_2O and 8.0% D_2O) instead of pure H_2O, as is typically done in such experiments.[61,62]

We begin with the NR results for supported bilayers and monolayers (Figure 4A,D). For $L/N = 1.5$, the clearest differences in the reflectivity signal were observed for supported bilayers using the hLipid/H_2O contrast and for monolayers under null contrast (Figure S11 and S18). Under these conditions, differences between the systems with and without RNA, and between the two selected tRNA structures are visible (Figure 4C,F). For bilayers, these differences are most pronounced in the q_z range 0.025–0.16 Å^{-1} (Figure 4C). For monolayers they appear for $q_z > 0.025$ Å^{-1} (Figure 4F). Notably, RNA adsorption shifts the first reflectivity minimum toward lower q_z, consistent with an increase in total thickness due to the RNA adsorption.[23]

As expected, under deuteration conditions involving D_2O and/or dLipid, such differences are

Figure 4: **NR curves of RNA interacting with supported bilayers (top) and monolayers (bottom).** (A) Solid-supported bilayer setup: Silicon/SiO$_2$ substrate, DOPC bilayer, RNA, and water. The neutron beam is incident from the left ($z < 0$). (B) Neutron SLD ρ_n for tRNA6 and tRNA9 on DOPC bilayers corresponding to setup (A). (C) NR curves for the systems shown in (B). (D) Monolayer setup: Air, DOPC monolayer, RNA, and water. (E) Corresponding neutron SLD profiles for tRNA6 and tRNA9 interacting with DOPC monolayers. (F) Monolayer systems in (E). The dashed gray lines in (C) and (F) represent the background noise level, defined as $10^{-7}q_z^4$. For an explicit noise model for reflectivity see SI Section 2. In all cases, the lipid-to-nucleotide ratio is $L/N = 1.5$. For the bilayer setup hLipid/H$_2$O contrast is used whereas for the monolayer null contrast ($\sim 92\%$ H$_2$O and 8.0% D$_2$O) is used in place of H$_2$O.

difficult to resolve (Figure S11 and S12). In these cases, the RNA contribution to the overall SLD is small compared to that of the deuterated components, resulting in reflectivity curves that are largely insensitive to RNA presence or conformation.

We now turn to X-ray reflectivity (XRR) as a complementary approach to probe RNA adsorption (Figure 5A, D). For $L/N = 1.5$, differences in reflectivity curves are observed between systems with and without RNA, and between two selected tRNA structures (Figure 5C, F) but are slightly less pronounced compared to NR. The differences appear only in the q_z range of 0.03–0.12 Å^{-1} for bilayers and up to 0.2 Å^{-1} for monolayers.

Finally, we provide an overview of the factors that influence NR and XRR profiles: lipid-to-nucleotide ratio, lipid charge, and RNA confor-

mation Figure 6. We focus on the NR signals at the supported bilayers (top) and XRR signals (bottom) for the monolayers at the air/water interface (see supporting information for all other setups).

The lipid-to-nucleotide L/N ratio is a key determinant of detectability. We investigated this by changing the L/N ratio from 4.3 to 1.5 (see supporting information section 8). Note that these ratios are similar to the ratio used in lipid nanoparticle (LNP) formulations.[63]

For NR, differences between pure lipid systems and with two different RNA conformations are clearly resolved only below $L/N < 3.0$ (Figure 6A). XRR can still detect the presence of RNA at this threshold but shows limited sensitivity to different RNA conformation (Figure 6B).

Lipid charge plays a pivotal role in RNA-lipid

Figure 5: **XRR curves of RNA interacting with solid supported bilayers (top) and monolayers (bottom).** (A) Solid-supported bilayer setup. The X-ray beam is incident from the left ($z < 0$). (B) X-ray SLD ρ_x for tRNA6 and tRNA9 on DOPC bilayers corresponding to setup (A). (C) X-ray reflectivity curves for the systems shown in (B). (D) Schematic of the monolayer setup: air, DOPC monolayer, RNA, and water. (E) Corresponding ρ_x for tRNA6 and tRNA9 interacting with DOPC monolayers. (F) X-ray reflectivity data for monolayer systems in (D). The line depicting the background noise level is omitted here since it is much lower than the signal. In all cases, the lipid-to-nucleotide ratio is $L/N = 1.5$.

interactions. In systems containing cationic DOTAP lipids, both NR and XRR profiles exhibit more pronounced differences, which can be attributed to enhanced electrostatic interactions between the cationic lipids and anionic RNA molecules (Figure 6C–F).

The impact of RNA structure is further illustrated by comparing nine different tRNA variants (Figure 6G–H). While certain conformations (e.g., tRNA6 and tRNA9) lead to distinctly different reflectivity curves, others produce nearly indistinguishable NR and XRR signals.

Collectively, these findings underscore the value of NR and XRR as mutually complementary techniques for probing RNA-lipid interactions at lipid/water interfaces. At sufficiently low lipid-to-nucleotide (L/N) ratios, both methods are sensitive enough to detect RNA adsorption at the interface. Still, forward models such as CG simulations are required for the molecular interpretation of the reflectivity curves.

Summary and Conclusion

This study addresses the modeling of RNA at lipid/water interfaces using CG simulations, validation using experimental scattering data, and prediction of NR and XRR profiles directly from the simulations. We systematically explored how ssRNA and dsRNA, as well as structured tRNA, interact with lipid DOPC/DOTAP bilayers of increasing positive charge. Our results show that RNA secondary structure is a key determinant of RNA-lipid interactions at lipid/water interfaces, with single-stranded regions exhibiting stronger interfacial affinity than double-stranded segments. ssRNA-bilayer interactions are governed by hydrophobic interactions with exposed bases and electrostatic attraction between RNA back-

Figure 6: **Influence of L/N ratio, lipid charge and RNA conformation on NR curves for solid-supported bilayers (top) and XRR curves for monolayers at the air/water interface (bottom).** Left: Influence of L/N ratio. (A) NR from a solid-supported DOPC bilayer without RNA, with tRNA6 and tRNA9. (B) XRR from a DOPC monolayer at the air/water interface with same tRNA variants. Middle: Influence of lipid charge. (C, E) NR from neutral DOPC and positively charged DOTAP bilayers without RNA, with tRNA6 and tRNA9. (D, F) XRR from neutral DOPC and positively charged monolayers with the same tRNA variants. Right: Influence of tRNA secondary structure. (G) NR from DOPC bilayers interacting with 9 different tRNA structures (Figure S7). (H) XRR from DOPC monolayers for same tRNA variants. Dashed gray lines indicate the background noise threshold, defined as $10^{-7}q_z^4$. For XRR this noise is omitted since it is much lower that the signal. It is also omitted in (A) for clarity, where all signals are well above the noise. In panels (C–H), the ratio is fixed at $L/N = 1.5$. hLipid/H$_2$O contrast was used for the neutron reflectivity curves.

bone and cationic lipid headgroups. dsRNA-bilayer interactions, constrained by base pairing, are weaker, particularly with neutral bilayers. These trends also hold for tRNA. Simulations of distinct tRNA conformations, representing different secondary structures, reveal that adsorption to neutral DOPC bilayers occurs primarily through exposed single-stranded regions, whereas in cationic DOTAP systems, adsorption is dominated by electrostatic interactions between the negatively charged tRNA backbone and the positively charged lipid headgroups.

To validate the CG bilayer model, we calculated and compared form factors from X-ray scattering. The results show that, while CG simulations reproduce experimental data qualitatively, agreement improves substantially when CG structures are backmapped to atomistic resolution and refined by energy minimization and short MD simulations.

Using this validated approach, we simulated NR and XRR curves for RNA-decorated solid-supported bilayers and monolayers at the air/water interface. We show that both NR and XRR can detect RNA adsorption at lipid in-

terfaces, especially under optimal contrast conditions and at low enough lipid-to-nucleotide ratios ($L/N < 1.5$). While reflectivity curves are sensitive to major RNA conformational changes, subtle structural differences will be challenging to resolve.

Systematic simulations can help to identify the experimental conditions best suited to resolve such subtle differences. Moreover, forward models such as the CG simulations presented here are expected to play an increasingly important role in the molecular interpretation of reflectivity data. Overall, our results demonstrate that combined simulation-scattering approaches are well suited to provide molecular insight into RNA adsorption and structure at lipid/water interfaces and to guide the design of future scattering experiments on RNA-membrane systems.

Acknowledgment

This work was supported by the German Federal Ministry of Education and Research through BMBF Project 05K18EZA within the framework of the Swedish-German research collaboration Röntgen-Ångström Cluster. The work was also supported by the Emmy Noether program of the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), number 315221747 and the research support program (Forschungspotenziale besser nutzen!) of the University of Augsburg. The authors gratefully acknowledge the scientific support and HPC resources provided by the Erlangen National High Performance Computing Center (NHR@FAU) of the Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU) under the NHR project b119ee and b253ee. We are grateful for the excellent support provided by the Max Planck Society. In particular, we are thankful to Gerhard Hummer at the Max Planck Institute for Biophysics, Frankfurt, Germany for his generous support as scientific host and for countless inspiring scientific discussions.

Conflict of Interest Statement

There are no conflicts of interest.

Supporting Information Available

Noise model for reflectivity and data for XRR and NR calculations for all the nine tRNA secondary structures for different deuteration conditions and RNA surface concentration for both bilayer and monolayer setups, undulation correction procedure and Martini bead electron and neutron scattering factor assignment procedures. The codes to obtain undulation corrected densities from CG-Martini and all-atom simulation trajectories are available at https://git.rz.uni-augsburg.de/cbio-gitpub.

References

(1) Czerniak, T.; Saenz, J. P. Lipid membranes modulate the activity of RNA through sequence-dependent interactions. Proc. Natl. Acad. Sci. U.S.A. **2022**, 119, e2119235119.

(2) Hou, X.; Zaks, T.; Langer, R.; Dong, Y. Lipid nanoparticles for mRNA delivery. Nat. Rev. Mater. **2021**, 6, 1078–1094.

(3) Schoenmaker, L.; Witzigmann, D.; Kulkarni, J. A.; Verbeke, R.; Kersten, G.; Jiskoot, W.; Crommelin, D. J. mRNA-lipid nanoparticle COVID-19 vaccines: Structure and stability. International journal of pharmaceutics **2021**, 601, 120586.

(4) Philipp, J.; Dabkowska, A.; Reiser, A.; Frank, K.; Krzysztoń, R.; Brummer, C.; Nickel, B.; Blanchet, C. E.; Sudarsan, A.; Ibrahim, M.; others pH-dependent structural transitions in cationic ionizable lipid mesophases are critical for lipid nanoparticle function. Proc. Natl. Acad. Sci. U.S.A. **2023**, 120, e2310491120.

(5) Suga, K.; Tanabe, T.; Tomita, H.; Shimanouchi, T.; Umakoshi, H. Conformational change of single-stranded RNAs induced by liposome binding. Nucleic Acids Res. **2011**, 39, 8891–8900.

(6) Stein, A.; Crothers, D. Conformational changes of transfer RNA. The role of magnesium (II). Biochemistry **1976**, 15, 160–168.

(7) Hall, M. N.; Gabay, J.; Débarbouillé, M.; Schwartz, M. A role for mRNA secondary structure in the control of translation initiation. Nature **1982**, 295, 616–618.

(8) de Smit, M. H.; van Duin, J. Control of translation by mRNA secondary structure in Escherichia coli: a quantitative analysis of literature data. J. Mol. Biol. **1994**, 244, 144–150.

(9) Chen, C.; Zhang, H.; Broitman, S. L.; Reiche, M.; Farrell, I.; Cooperman, B. S.; Goldman, Y. E. Dynamics of translation by single ribosomes through mRNA secondary structures. Nat. Struct. Mol. Biol. **2013**, 20, 582–588.

(10) Chemla, Y.; Peeri, M.; Heltberg, M. L.; Eichler, J.; Jensen, M. H.; Tuller, T.; Alfonta, L. A possible universal role for mRNA secondary structure in bacterial translation revealed using a synthetic operon. Nat. Commun. **2020**, 11, 4827.

(11) Heinrich, F.; Lösche, M. Zooming in on disordered systems: Neutron reflection studies of proteins associated with fluid membranes. Biochim Biophys Acta Biomembr **2014**, 1838, 2341–2349.

(12) Michanek, A.; Kristen, N.; Höök, F.; Nylander, T.; Sparr, E. RNA and DNA interactions with zwitterionic and charged lipid membranes—A DSC and QCM-D study. Biochim Biophys Acta Biomembr **2010**, 1798, 829–838.

(13) Nanda, H.; Datta, S. A.; Heinrich, F.; Lösche, M.; Rein, A.; Krueger, S.; Curtis, J. E. Electrostatic interactions and binding orientation of HIV-1 matrix studied by neutron reflectivity. Biophys. J. **2010**, 99, 2516–2524.

(14) Pfefferkorn, C. M.; Heinrich, F.; Sodt, A. J.; Maltsev, A. S.; Pastor, R. W.; Lee, J. C. Depth of α-synuclein in a bilayer determined by fluorescence, neutron reflectometry, and computation. Biophys. J. **2012**, 102, 613–621.

(15) Bange, L.; Mukhina, T.; Fragneto, G.; Rondelli, V.; Schneck, E. Influence of adhesion-promoting glycolipids on the structure and stability of solid-supported lipid double-bilayers. Soft Matter **2024**, 20, 2113–2125.

(16) Dabkowska, A. P.; Talbot, J. P.; Cavalcanti, L.; Webster, J. R.; Nelson, A.; Barlow, D. J.; Fragneto, G.; Lawrence, M. J. Calcium mediated interaction of calf-thymus DNA with monolayers of distearoylphosphatidylcholine: a neutron and X-ray reflectivity study. Soft Matter **2013**, 9, 7095–7105.

(17) Soranzo, T.; Martin, D. K.; Lenormand, J.-L.; Watkins, E. B. Coupling neutron reflectivity with cell-free protein synthesis to probe membrane protein structure in supported bilayers. Sci. Rep. **2017**, 7, 3399.

(18) Schneck, E.; Berts, I.; Halperin, A.; Daillant, J.; Fragneto, G. Neutron reflectometry from poly (ethylene-glycol) brushes binding anti-PEG antibodies: Evidence of ternary adsorption. Biomaterials **2015**, 46, 95–104.

(19) Grava, M.; Ibrahim, M.; Sudarsan, A.; Pusterla, J.; Philipp, J.; Rädler, J. O.; Schwierz, N.; Schneck, E. Combining molecular dynamics simulations and x-ray scattering techniques for the accurate treatment of protonation degree and packing of ionizable lipids in monolayers. J. Chem. Phys. **2023**, 159.

(20) Ibrahim, M.; Gilbert, J.; Heinz, M.; Nylander, T.; Schwierz, N. Structural Insights on Ionizable Dlin-MC3-DMA Lipids in DOPC Layers by Combining Accurate Atomistic Force Fields, Molecular Dynamics Simulations and Neutron Reflectivity. Nanoscale **2023**,

(21) Kent, M. S.; Murton, J. K.; Sasaki, D. Y.; Satija, S.; Akgun, B.; Nanda, H.; Curtis, J. E.; Majewski, J.; Morgan, C. R.; Engen, J. R. Neutron reflectometry study of the conformation of HIV Nef bound to lipid membranes. Biophys. J. **2010**, <u>99</u>, 1940–1948.

(22) Akgun, B.; Satija, S.; Nanda, H.; Pirrone, G. F.; Shi, X.; Engen, J. R.; Kent, M. S. Conformational transition of membrane-associated terminally acylated HIV-1 Nef. Structure **2013**, <u>21</u>, 1822–1833.

(23) Grava, M.; Weber, K.; Reed, J.; Weber, B.; Shen, C.; Haas, H.; Schneck, E. Lipid Monolayers Adsorbing mRNA as Models for mRNA Enclosed in Lipid Nanoparticles for Transfection. Langmuir **0**, <u>0</u>, null, PMID: 41230820.

(24) Gilbert, J.; Ermilova, I.; Fornasier, M.; Skoda, M.; Fragneto, G.; Swenson, J.; Nylander, T. On the interactions between RNA and titrateable lipid layers: implications for RNA delivery with lipid nanoparticles. Nanoscale **2024**, <u>16</u>, 777–794.

(25) Chen, P.-c.; Hub, J. S. Interpretation of solution x-ray scattering by explicit-solvent molecular dynamics. Biophys. J. **2015**, <u>108</u>, 2573–2584.

(26) Marrink, S. J.; Risselada, H. J.; Yefimov, S.; Tieleman, D. P.; De Vries, A. H. The MARTINI force field: coarse grained model for biomolecular simulations. J. Phys. Chem. B **2007**, <u>111</u>, 7812–7824.

(27) de Jong, D. H.; Singh, G.; Bennett, W. D.; Arnarez, C.; Wassenaar, T. A.; Schafer, L. V.; Periole, X.; Tieleman, D. P.; Marrink, S. J. Improved parameters for the martini coarse-grained protein force field. J. Chem. Theory Comput. **2013**, <u>9</u>, 687–697.

(28) Kučerka, N.; Nagle, J. F.; Sachs, J. N.; Feller, S. E.; Pencer, J.; Jackson, A.; Katsaras, J. Lipid bilayer structure determined by the simultaneous analysis of neutron and X-ray scattering data. Biophys. J. **2008**, <u>95</u>, 2356–2367.

(29) Carucci, C.; Philipp, J.; Muller, J. A.; Sudarsan, A.; Kostyurina, E.; Blanchet, C. E.; Schwierz, N.; Parsons, D. F.; Salis, A.; Radler, J. O. Buffer specificity of ionizable lipid nanoparticle transfection efficiency and bulk phase transition. ACS nano **2025**, <u>19</u>, 10829–10840.

(30) Philipp, J.; Sudarsan, A.; Kostyurina, E.; Meklesh, V.; Berglund, M.; Rappolt, M.; Westergren, J.; Lindfors, L.; Schwierz, N.; Raedler, J. O. Combining SAXS analysis and MD simulation to determine structure and hydration of ionizable lipid hexagonal phases. Soft Matter **2025**,

(31) Koutsioubas, A. Combined coarse-grained molecular dynamics and neutron reflectivity characterization of supported lipid membranes. J. Phys. Chem. B **2016**, <u>120</u>, 11474–11483.

(32) Settanni, G.; Brill, W.; Haas, H.; Schmid, F. pH-dependent behavior of ionizable cationic lipids in mRNA-carrying lipoplexes investigated by molecular dynamics simulations. Macromol. Rapid Commun. **2022**, <u>43</u>, 2100683.

(33) Uusitalo, J. J.; Ingólfsson, H. I.; Marrink, S. J.; Faustino, I. Martini coarse-grained force field: extension to RNA. Biophys. J. **2017**, <u>113</u>, 246–256.

(34) Wassenaar, T. A.; Ingólfsson, H. I.; Bockmann, R. A.; Tieleman, D. P.; Marrink, S. J. Computational lipidomics with insane: a versatile tool for generating custom membranes for molecular simulations. J. Chem. Theory Comput. **2015**, <u>11</u>, 2144–2155.

(35) `https://casegroup.rutgers.edu/fd_helix.c`, Accessed: 2025-04-23.

(36) Michanek, A.; Yanez, M.; Wacklin, H.; Hughes, A.; Nylander, T.; Sparr, E. RNA and DNA association to zwitterionic and charged monolayers at the air–liquid interface. Langmuir **2012**, 28, 9621–9633.

(37) Bussi, G.; Donadio, D.; Parrinello, M. Canonical sampling through velocity rescaling. J. Chem. Phys. **2007**, 126, 014101.

(38) Parrinello, M.; Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics method. J. Appl. Phys. **1981**, 52, 7182–7190.

(39) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX **2015**, 1, 19–25.

(40) Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O. MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. J. Comput. Chem. **2011**, 32, 2319–2327.

(41) Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual molecular dynamics. J. Mol. Graph. **1996**, 14, 33–38.

(42) Jo, S.; Kim, T.; Iyer, V. G.; Im, W. CHARMM-GUI: a web-based graphical user interface for CHARMM. Journal of computational chemistry **2008**, 29, 1859–1865.

(43) Dickson, C. J.; Madej, B. D.; Skjevik, Å. A.; Betz, R. M.; Teigen, K.; Gould, I. R.; Walker, R. C. Lipid14: The Amber Lipid Force Field. J. Chem. Theory Comput. **2014**, 10, 865–879.

(44) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. J. Chem. Phys. **1983**, 79, 926–935.

(45) Mamatkulov, S.; Schwierz, N. Force fields for monovalent and divalent metal cations in TIP3P water based on thermodynamic and kinetic properties. J. Chem. Phys. **2018**, 148.

(46) Bernetti, M.; Bussi, G. Pressure control using stochastic cell rescaling. J. Chem. Phys. **2020**, 153.

(47) Essmann, U.; Perera, L.; Berkowitz, M. L.; Darden, T.; Lee, H.; Pedersen, L. G. A smooth particle mesh Ewald method. J. Chem. Phys. **1995**, 103, 8577–8593.

(48) Wassenaar, T. A.; Pluhackova, K.; Bockmann, R. A.; Marrink, S. J.; Tieleman, D. P. Going backward: a flexible geometric approach to reverse transformation from coarse grained to atomistic models. J. Chem. Theory Comput. **2014**, 10, 676–690.

(49) Braun, A. R.; Brandt, E. G.; Edholm, O.; Nagle, J. F.; Sachs, J. N. Determination of electron density profiles and area from simulations of undulating membranes. Biophys. J. **2011**, 100, 2112–2120.

(50) Kim, H.; Fábián, B.; Hummer, G. Neighbor list artifacts in molecular dynamics simulations. J. Chem. Theory Comput. **2023**, 19, 8919–8929.

(51) Abelès, F.; La, F. A. théorie générale des couches minces. J. Phys. Radium **1950**, 11, 307–309.

(52) Nelson, A. R. J.; Prescott, S. W. refnx: neutron and X-ray reflectometry analysis in Python. J. Appl. Crystallogr. **2019**, 52, 193–200.

(53) Guzmán, E.; Liggieri, L.; Santini, E.; Ferrari, M.; Ravera, F. DPPC–DOPC Langmuir monolayers modified by hydrophilic silica nanoparticles: Phase behaviour, structure and rheology. Colloids Surf. A Physicochem. Eng. Asp. **2012**, 413, 174–183.

(54) Lorenz, R.; Bernhart, S. H.; Höner zu Siederdissen, C.; Tafer, H.; Flamm, C.; Stadler, P. F.; Hofacker, I. L. ViennaRNA Package 2.0. Algorithms Mol Biol **2011**, _6_, 1–14.

(55) Popenda, M.; Szachniuk, M.; Antczak, M.; Purzycka, K. J.; Lukasiak, P.; Bartol, N.; Blazewicz, J.; Adamiak, R. W. Automated 3D structure composition for large RNAs. Nucleic Acids Res. **2012**, _40_, e112–e112.

(56) Marty, R.; Nsoukpoe-Kossi, C. N.; Charbonneau, D. M.; Kreplak, L.; Tajmir-Riahi, H.-A. Structural characterization of cationic lipid–tRNA complexes. Nucleic Acids Res. **2009**, _37_, 5197–5207.

(57) Antipina, A. Y.; Gurtovenko, A. A. Toward understanding liposome-based siRNA delivery vectors: Atomic-scale insight into siRNA–lipid interactions. Langmuir **2018**, _34_, 8685–8693.

(58) Di Marco, S.; Aupic, J.; Bussi, G.; Magistrato, A. All-Atom Simulations Elucidate the Molecular Mechanism Underlying RNA–Membrane Interactions. Nano Lett. **2025**, _25_, 4628–4635.

(59) Klauda, J. B.; Venable, R. M.; Freites, J. A.; OConnor, J. W.; Tobias, D. J.; Mondragon-Ramirez, C.; Vorobyov, I.; MacKerell Jr, A. D.; Pastor, R. W. Update of the CHARMM all-atom additive force field for lipids: validation on six lipid types. J. Phys. Chem. B **2010**, _114_, 7830–7843.

(60) López, C. A.; Sovova, Z.; van Eerden, F. J.; de Vries, A. H.; Marrink, S. J. Martini force field parameters for glycolipids. J. Chem. Theory Comput. **2013**, _9_, 1694–1708.

(61) Campbell, R. A.; Kairaliyeva, T.; Santer, S.; Schneck, E.; Miller, R. Direct resolution of the interactions of a hydrocarbon gas with adsorbed surfactant monolayers at the water/air interface using neutron reflectometry. Colloids and Interfaces **2022**, _6_, 68.

(62) Gustafsson, E.; Hellsing, M. S.; Rennie, A. R.; Welbourn, R. J.; Campana, M.; Hughes, A.; Li, P.; Bowden, T. M. Understanding interactions of plasticisers with a phospholipid monolayer. Soft Matter **2024**, _20_, 2892–2899.

(63) Kulkarni, J. A.; Darjuan, M. M.; Mercer, J. E.; Chen, S.; Van Der Meel, R.; Thewalt, J. L.; Tam, Y. Y. C.; Cullis, P. R. On the formation and morphology of lipid nanoparticles containing ionizable cationic lipids and siRNA. ACS nano **2018**, _12_, 4787–4795.