A stress-dependent TRIM28–ALKBH2 feedback loop modulates chemotherapy response

Zhiming Sun, Yunfang Deng, Yue Liu, Zhaohui Liu, Jiabing Li, Lihui Wu, Liyuan Zeng, Xiaorong Feng, Lin Miao, Ying Sheng, Bei Chen, Yuming He, Ye Liu & Yu Zhao*

The National & Local Joint Engineering Laboratory of Animal Peptide Drug Development, College of Life Sciences, Hunan Normal University, Changsha, Hunan, 410081, China

* Corresponding author: Yu Zhao (smile_zhao@hotmail.com)

Running title: TRIM28–ALKBH2 modulates chemoresistance

Keywords: TRIM28, ALKBH2, MAGEA6, Ubiquitination, Chemoresistance

ABSTRACT

Chemoresistance to DNA-damaging agents, including platinum and alkylating compounds, remains a major limitation in the treatment of non-small cell lung cancer (NSCLC), often correlated with upregulated DNA repair. Here, we define a reciprocal regulatory loop between the E3 ligase TRIM28 and the alkylation repair enzyme ALKBH2. TRIM28 directly binds ALKBH2 and mediates its K48-linked polyubiquitination and proteasomal degradation, whereas ALKBH2 reciprocally enhances TRIM28 transcription, establishing a homeostatic equilibrium. Upon alkylation stress, acute MMS exposure enhances assembly of the TRIM28–ALKBH2 complex, a process facilitated by MAGEA6, and promotes ALKBH2 degradation. In contrast, prolonged MMS exposure induces K48-linked ubiquitination and degradation of TRIM28, leading to ALKBH2 stabilization and transcriptional upregulation. Clinically, ALKBH2 is elevated in lung adenocarcinoma and correlates with poor prognosis; notably, high TRIM28 expression specifically correlates with poor survival in patients receiving chemotherapy. Moreover, the combination of MMS and cisplatin enhances DNA damage and attenuates ALKBH2-mediated chemoresistance. Collectively, these findings define a stress-responsive TRIM28–ALKBH2 feedback loop that modulates alkylation responses and shapes chemotherapy sensitivity.

Introduction

Cells are continuously challenged by endogenous metabolites and environmental or therapeutic alkylating agents that generate cytotoxic and mutagenic DNA adducts, posing a constant threat to genome integrity (Fu, Calvo et al., 2012, Zhao, Majid et al., 2015). To counteract this, cells utilize three principal repair strategies: base excision repair (BER), direct reversal by O^6-methylguanine-DNA methyltransferase (MGMT), and oxidative demethylation catalyzed by the AlkB family of Fe(II)/α-ketoglutarate–dependent dioxygenases (Drabløs, Feyzi et al., 2004, Li, Feng et al., 2025, Soll, Sobol et al., 2017). These dioxygenases specifically target N-linked methyl lesions (e.g., 1-meA and 3-meC) through a coupled oxidative reaction, restoring the unmodified base without requiring nucleotide excision (Leiros, Nabong et al., 2007, Trewick, Henshaw et al., 2002). This direct reversal mechanism provides an evolutionarily conserved and critical defense against alkylation toxicity (Yi & He, 2013). In mammals, the AlkB homologues complement BER and MGMT to resolve specific lesions that escape other repair systems, thereby ensuring efficient genomic maintenance under both physiological and pathological stress (Xu & Bochtler, 2020).

Among the nine mammalian AlkB homologues, the demethylase ALKBH2 serves as the principal enzyme for repairing methylated bases in duplex DNA, a function essential for preventing alkylation-induced mutagenesis (Duncan, Trewick et al., 2002). Loss or functional impairment of ALKBH2 results in hypersensitivity to alkylating agents and the accumulation of cytotoxic base lesions, underscoring its nonredundant role in resolving adducts that escape other repair pathways (Li, Huang et al., 2016, Ringvoll, Nordstrand et al., 2006, Wang, Wu et al., 2015). Clinically, ALKBH2 is frequently upregulated in human cancers, particularly non–small cell lung cancer (NSCLC), where elevated expression correlates with increased repair capacity and chemoresistance (Johannessen, Prestegarden et al., 2013, Wang et al., 2015, Wu, Yan et al., 2021, Wu, Xu et al., 2011, Xu, Zhang et al., 2022). Given that alkylating chemotherapeutics (e.g., temozolomide) and environmental insults (e.g., cigarette smoke) generate similar methylated base lesions (Fu et al., 2012, Hegi, Diserens et al., 2005, Li, Zeng et al., 2024), the dynamic regulation of ALKBH2 constitutes a key determinant of tumor survival under sustained alkylation stress

(Johannessen et al., 2013, Wilson, Beharry et al., 2018). However, the specific post-translational mechanisms governing ALKBH2 protein stability and turnover remain poorly understood.

While the expression of AlkB family dioxygenases is modulated by diverse transcriptional and epigenetic programs (Chen, Zhou et al., 2025, Feng, Xu et al., 2022, Johannessen et al., 2013, Song, Feng et al., 2019, Wada, Naito et al., 2024), these enzymes are tightly controlled by dynamic ubiquitin-dependent regulation to ensure rapid proteostatic control. Extensive research has delineated the E3 ligase networks for other AlkB homologues, particularly FTO and ALKBH5. For instance, FTO turnover is regulated by multiple context-specific E3 ligases, including TRIM17, DTX2, MIB1, and RNF144A (Cui, Wei et al., 2024, Dai, Tian et al., 2024, Han, Tong et al., 2024, Liu, Zheng et al., 2025), while its degradation by TRIM21 is specifically primed by O-GlcNAcylation (Zhang, Liu et al., 2025). Similarly, ALKBH5 stability is controlled by the E3 ligase RNF180, or via PRMT5-mediated methylation, which primes it for recognition and ubiquitination by TRIM28 (Meng, Liu et al., 2025, Wang, Liu et al., 2024). Conversely, these degradative pathways are counterbalanced by a deubiquitinase (DUB) network, including USP7, USP14, USP18, and USP36. This regulation orchestrates dioxygenase homeostasis by stabilizing the demethylases FTO and ALKBH5 in a highly substrate-specific manner. For example, FTO is stabilized by USP7 (Lin, Zhou et al., 2023) and USP18 (Song, Yi et al., 2024), while USP14 (Zhou, Xia et al., 2025) and USP36 (Chang, Xie et al., 2023) stabilize ALKBH5. Our own previous work further extended this principle by identifying OTUD4 as a scaffolding DUB that stabilizes the demethylase ALKBH3 (Zhao et al., 2015). In stark contrast to these well-characterized homologues, the specific E3 ligase controlling ALKBH2 protein stability remains unidentified. However, a vital clue emerges from the ALKBH2-containing ASCC complex: its active recruitment of TRIM28 upon alkylation stress (MMS) (Brickner, Soll et al., 2017), thereby functionally positioning this E3 ligase within the alkylation repair machinery.

The tripartite motif (TRIM) family of RING-type E3 ligases, including TRIM24, TRIM28, and TRIM33, orchestrates ubiquitin signaling to couple chromatin remodeling with stress responses (Gu, Chen et al., 2023, Hatakeyama, 2011). As a key member of this subgroup, TRIM28

(KAP1), leverages its N-terminal RING domain and C-terminal PHD–BRD module to enforce KRAB-ZNF-dependent transcriptional silencing and maintain genomic stability (Czerwińska, Mazurek et al., 2017, Randolph, Hyder et al., 2022). While TRIM28 functions as a master corepressor and E3 ligase, its own functional state and availability are governed by a complex regulatory hierarchy. Classically, in response to DNA double-strand breaks (DSBs), TRIM28 undergoes rapid ATM/ATR-mediated phosphorylation at Ser824, a modification essential for relaxing heterochromatin to facilitate repair factor access (Goodarzi, Kurka et al., 2011, Lee, Goodarzi et al., 2012, Yalçin, Lam et al., 2024). Beyond these acute post-translational modifications, TRIM28 expression is tightly regulated at the transcriptional level by nuclear receptors such as NR3C2 (Li, Xie et al., 2025) and post-transcriptionally via NSUN2-mediated RNA methylation (m^5C) (Wang, Mierxiati et al., 2025) as well as stabilization by the lncRNA PITAR (Jana, Mondal et al., 2024). Furthermore, under stress conditions, TRIM28 protein abundance is dynamically regulated via the ubiquitin-proteasome system; specifically, its turnover is governed by the E3 ligase RNF4 during genotoxic stress and March5 under metabolic stress (Chen, Pang et al., 2025, Kuo, Li et al., 2014). Additionally, the substrate specificity of TRIM28 is often reprogrammed by melanoma-associated antigen (MAGE) proteins (Doyle, Gao et al., 2010, Kim, Lee et al., 2009, Xiao, Suh et al., 2014), such as MAGEA6, which act as substrate adaptors (Doyle et al., 2010, Li, Krone et al., 2024). However, the mechanistic interplay between TRIM28 and the alkylation repair machinery—particularly whether reciprocal feedback circuits coordinate their stability and expression under stress—remains largely unexplored.

In this study, TRIM28, together with the adaptor MAGEA6, is defined as the E3 ligase complex that regulates ALKBH2 stability through K48-linked polyubiquitination and proteasomal degradation under basal conditions. ALKBH2, in turn, enhances TRIM28 transcription, forming a homeostatic feedback loop that links E3 ligase activity to ALKBH2 turnover. Acute MMS exposure enhances assembly of the TRIM28–ALKBH2 complex and promotes ALKBH2 degradation, whereas prolonged MMS exposure induces K48-linked ubiquitination and degradation of TRIM28, resulting in ALKBH2 stabilization and transcriptional induction.

113 Together, these mechanisms reveal a stress-responsive TRIM28–MAGEA6–ALKBH2 feedback

114 loop that coordinates repair capacity with the dynamics of stress exposure and provides a

115 mechanistic basis for variations in chemotherapy response.

Results

TRIM28 directly binds ALKBH2

To identify proteins associated with TRIM28, we performed tandem affinity purification in H1299 non–small cell lung cancer (NSCLC) cells, followed by silver staining and LC–MS/MS. This analysis identified ALKBH2 as a putative interactor (Fig. 1A; Appendix Fig. S1A; Appendix Table S1). Co-immunoprecipitation (co-IP) in HEK293T cells co-expressing HA-tagged ALKBH2 and Flag-tagged TRIM28 confirmed this association, whereas the related paralog TRIM24 did not show detectable binding (Fig. 1B). Reciprocal co-IP using HA-ALKBH2 further validated the interaction (Fig. 1C). Domain mapping using TRIM28 truncation mutants demonstrated that the BCC region (B-box + coiled-coil; residues 141–400) was required for ALKBH2 binding (Fig. 1D, E). Endogenous co-IP in H1299 cells confirmed recovery of native ALKBH2 by the anti-TRIM28 antibody (Fig. 1F), and immunofluorescence analysis revealed strong nuclear co-localization of TRIM28 and ALKBH2 (Fig. 1G).

In vitro MBP pull-down assays using purified recombinant proteins demonstrated that TRIM28 directly binds ALKBH2 (Fig. 1H). Specificity profiling revealed a preferential interaction with ALKBH2 compared with the related demethylase ALKBH3 (Fig. 1I) or the direct-reversal repair enzyme MGMT (Appendix Fig. S1B). Protein–protein docking predicted a TRIM28–ALKBH2 interface overlapping the nucleic-acid-binding surface of ALKBH2, centered on the Finger (hairpin) motif and the flexible Long-loop (Yang, Yi et al., 2008), which are present in ALKBH2 but absent in ALKBH3 (Fig. 1J). Consistent with this model, single-stranded DNA (ssDNA) competitively reduced TRIM28–ALKBH2 binding in a dose-dependent manner (Fig. 1K), a strategy commonly used to probe DNA-occupied states of repair enzymes (Gu, Deng et al., 2013, Le, Ma et al., 2020). Collectively, these data reveal a direct and selective interaction between TRIM28 and ALKBH2 mediated by the TRIM28 BCC domain and indicate that DNA binding to ALKBH2 can competitively modulate this interaction.

A reciprocal regulatory loop between TRIM28 and ALKBH2 governs the stability and ubiquitination of ALKBH2

To determine whether ALKBH2 undergoes ubiquitin-dependent modification, we performed denaturing immunoprecipitation (IP) of Flag-ALKBH2 from HEK293T cells co-transfected with HA-ubiquitin. A high-molecular-weight smear corresponding to polyubiquitinated ALKBH2 was detected, and this signal markedly increased after proteasome inhibition with MG132 (Fig. 2A). Expression of a K48-only ubiquitin mutant further enhanced ALKBH2 ubiquitination, whereas the K48R mutant almost completely abolished it (Fig. 2B), indicating that ALKBH2 is predominantly modified by K48-linked polyubiquitin chains. Consistently, Halo-TUBE pull-down enriched K48-linked polyubiquitinated ALKBH2, confirming endogenous K48-linked ubiquitination (Fig. 2C).

We next examined whether TRIM28 mediates this modification. Overexpression of wild-type (WT) TRIM28 increased K48-linked ubiquitination of ALKBH2 (Fig. 2D), whereas the RING-inactive C65/68A mutant—previously shown to abolish TRIM28 E3 ligase activity (Doyle et al., 2010)—failed to promote this modification (Fig. 2F). TRIM28 depletion using two independent shRNAs markedly reduced ALKBH2 ubiquitination (Appendix Fig. S2A). Re-expression of shRNA-resistant WT TRIM28 in TRIM28-knockdown cells restored ALKBH2 ubiquitination (Fig. 2E). These results demonstrate that TRIM28 E3 ligase activity is required for efficient K48-linked ubiquitination of ALKBH2.

We then assessed how this modification affects ALKBH2 protein stability. TRIM28 knockdown in H1299 and A549 cells increased ALKBH2 protein levels without altering its mRNA expression (Fig. 2G, H), and similar results were obtained upon CRISPR/Cas9-mediated TRIM28 depletion (Appendix Fig. S2B). Cycloheximide (CHX) chase assays revealed that TRIM28 knockdown extended ALKBH2 half-life, whereas enforced TRIM28 expression shortened it in a RING-dependent manner (Appendix Fig. S2C, D). Increasing TRIM28 expression produced a dose-dependent decrease in ALKBH2 protein abundance in both H1299 and HEK293T cells (Fig. 2I). These data indicate that TRIM28 limits ALKBH2 levels through proteasome-mediated degradation rather than altered localization (Appendix Fig. S2E).

To determine whether ALKBH2 in turn regulates its E3 ligase, we depleted ALKBH2 in

H1299 cells using two independent shRNAs. TRIM28 protein levels were reduced following

ALKBH2 knockdown (Fig. 2J), and re-expression of Flag-ALKBH2 restored TRIM28 (Fig. 2K).

Overexpression of ALKBH2 further increased TRIM28 mRNA as measured by qRT-PCR (Fig.

2L). Consistent with this regulatory link, TCGA lung adenocarcinoma data revealed a positive

correlation between ALKBH2 and TRIM28 transcripts (Appendix Fig. S2F). Collectively, these

results define a reciprocal regulatory loop in which TRIM28 acts as an E3 ligase promoting K48-

linked ubiquitination and proteasomal degradation of ALKBH2, while ALKBH2 enhances

TRIM28 expression at the transcriptional level.

MAGEA6 scaffolds the TRIM28–ALKBH2 E3 ligase complex, a function impaired by

cancer-associated variants

Given that MAGE family proteins can associate with E3 ligases such as TRIM28 to modulate their

activity (Doyle et al., 2010, Feng, Gao et al., 2011), we examined the organization of the TRIM28–

ALKBH2 complex and the role of MAGEA6 in this process. Co-immunoprecipitation (co-IP) in

HEK293T cells revealed a previously unrecognized interaction between Myc-tagged MAGEA6

and the substrate ALKBH2 (Fig. 3A), which was further confirmed as direct by *in vitro* pull-down

assays using purified recombinant proteins (Fig. 3B).

We next investigated how MAGEA6 contributes to complex assembly. The MAGEA6–

ALKBH2 interaction was independent of TRIM28, as TRIM28 knockdown did not disrupt their

association (Fig. 3C), suggesting that MAGEA6 engages ALKBH2 before or independently of

TRIM28 recruitment. Overexpression of Myc-MAGEA6 enhanced TRIM28–ALKBH2 complex

formation in a dose-dependent manner (Fig. 3D; Fig. EV1A), whereas knockdown of endogenous

MAGEA3/6 attenuated this interaction (Fig. 3E). Because antibodies cannot distinguish between

the highly homologous MAGEA3 and MAGEA6 (96% sequence identity (Pineda, Ramanathan et

al., 2015)), they are referred to collectively as MAGEA3/6 throughout this study. The related

family member MAGEC2 exhibited a similar effect (Fig. EV1B), indicating a conserved scaffolding function within this subfamily.

To determine the functional consequence of this scaffolded complex, we next assessed its impact on ALKBH2 ubiquitination and stability. MAGEA6 knockdown reduced endogenous ALKBH2 ubiquitination, and this reduction was rescued by re-expression of MAGEA6 (Fig. EV1C). Conversely, MAGEA6 overexpression promoted K48-linked ubiquitination of ALKBH2, but this effect was markedly diminished upon TRIM28 knockdown, confirming that it depends on TRIM28 (Fig. 3F). Consistently, MAGEA6 accelerated ALKBH2 degradation in control cells, whereas this effect was largely abolished in TRIM28-knockout H1299 cells (Fig. 3G).

Given that MAGEA6 mutations have been reported in multiple tumor types, including lung cancer (Tsang, Wang et al., 2020), we examined whether cancer-associated variants affect this regulatory axis. Sequencing of MAGEA6 cDNA from H1299 cells identified two variants, MutA and MutB, harboring 12 and 20 amino acid substitutions, respectively (Fig. EV1D). Both mutants were individually cloned and sequence-verified for functional testing. Co-IP assays revealed that both MutA and MutB displayed weaker binding to ALKBH2, and MutB also showed reduced association with TRIM28 (Fig. 3H). As a result, these variants exhibited impaired ability to promote ALKBH2 ubiquitination and degradation compared with wild-type MAGEA6 (Fig. 3I; Fig. EV1E).

To further delineate the molecular basis of this interaction, AlphaFold3 modeling predicted a contact interface between MAGEA6 and TRIM28 (Fig. EV1F), encompassing residues D156 and S157 within the MAGE homology domain. Notably, the S157 site is found in the MutB variant and corresponds to tumor-associated mutations reported in COSMIC: S157 (MAGEA6, COSMIC ID: 1715821, melanoma) and its homologous position in MAGEA3 (S157, COSMIC ID: 3559357, detected in lung cancer, lymphoma, and melanoma). Functional assays showed that mutating these residues (D156E/S157Y) markedly reduced TRIM28 binding and impaired MAGEA6-mediated ALKBH2 degradation (Fig. 3J; Fig. EV1H). In contrast, a K211R substitution within the predicted ALKBH2-binding surface had only a modest effect on their association (Fig. 3J). Collectively,

these findings define MAGEA6 as a scaffolding cofactor that directly associates with ALKBH2

and recruits TRIM28 to promote its ubiquitination and degradation. Cancer-associated variants

that disrupt the MAGEA6–TRIM28 interface compromise this regulatory function, potentially

contributing to defective DNA repair regulation in tumors.

MMS drives a biphasic ALKBH2 response, from acute degradation to prolonged

accumulation

MMS acutely promotes ALKBH2 degradation

To investigate how MMS regulates the TRIM28–ALKBH2 axis, we analyzed their interaction

dynamics in H1299 cells. Short-term MMS exposure (3–6 h) progressively enhanced the

TRIM28–ALKBH2 association in a time- and dose-dependent manner (Fig. 4A, B). While both

proteins remained predominantly nuclear (Fig. EV2A), MAGEA3/6 accumulated in the nucleus

upon MMS treatment (Fig. 4C), suggesting that MAGEA3/6 facilitates assembly of the TRIM28–

ALKBH2 complex during genotoxic stress. Consistent with this, MMS increased K48-linked

ubiquitination of ALKBH2 in a concentration-dependent manner (Fig. 4D). To exclude

transcriptional effects, we used H1299 cells stably expressing TAP-tagged ALKBH2. MMS

exposure accelerated ALKBH2 degradation, which was further enhanced by TRIM28

overexpression (Fig. 4E). Similarly, endogenous ALKBH2 exhibited reduced stability when cells

were co-treated with MMS and cycloheximide (CHX), compared with CHX alone (Fig. EV2B, C).

These data define an acute response phase in which MMS enhances TRIM28-dependent

ubiquitination and degradation of ALKBH2.

Because alkylating agents such as MMS and temozolomide (TMZ) are known to elevate

intracellular ROS levels (Bu, Qu et al., 2021, Lombard, Kohler et al., 2020, Zhao, Zheng et al.,

2017), we next examined whether redox balance modulates the TRIM28–ALKBH2 interaction. In

co-immunoprecipitation assays, H_2O_2 enhanced complex formation, whereas single-stranded

DNA (ssDNA), ascorbate (Vc), or EDTA weakened the interaction (Fig. 4F). Reduced glutathione

(GSH) attenuated TRIM28–ALKBH2 binding, while oxidized glutathione (GSSG) had little effect.

GSH also blunted the H_2O_2-induced enhancement (Fig. EV2D). Similarly, EDTA decreased basal

interaction and prevented H_2O_2-driven binding (Fig. 4G). In agreement, oxidative stress alone was sufficient to trigger ubiquitination, as H_2O_2 increased K48-linked polyubiquitination of ALKBH2 in a dose-dependent manner (Fig. EV2E).

In contrast, the catalytically inactive ALKBH2(H171A/D173A) mutant (Lee, Jin et al., 2005, Yang et al., 2008) failed to display the ssDNA-dependent reduction or H_2O_2-induced enhancement of TRIM28 binding, suggesting that this redox control is coupled to the catalytic state of ALKBH2 (Fig. EV2F). Furthermore, the reducing agent Vc suppressed MMS-induced K48-linked ubiquitination of ALKBH2 (Fig. 4H). Together, these results identify an acute MMS response characterized by redox-sensitive, TRIM28-dependent ubiquitination leading to ALKBH2 degradation.

Prolonged MMS exposure stabilizes ALKBH2

In contrast to the early degradative phase, prolonged MMS exposure (≥8 h) led to recovery and accumulation of ALKBH2 protein. MMS treatment increased ALKBH2 abundance in both H1299 and H460 cells, with variable induction levels between lines (Fig. 4I, J). The protein levels of ALKBH3 and MGMT remained unchanged (Fig. 4I). Moreover, TRIM28 overexpression attenuated MMS-induced ALKBH2 accumulation (Fig. 4K). These data indicate that prolonged MMS treatment shifts ALKBH2 regulation from degradation to stabilization.

Prolonged MMS exposure induces transcriptional upregulation of ALKBH2

To determine whether ALKBH2 accumulation involves transcriptional activation, total RNA was isolated from H1299 cells after MMS exposure for the indicated durations and analyzed by qRT–PCR. A significant increase in ALKBH2 transcript abundance was observed at 12 h and sustained at 24 h (Fig. 4L), consistent with the delayed protein induction. Collectively, these data define a biphasic regulatory pattern of ALKBH2 under MMS exposure—an early, post-translational phase marked by TRIM28-dependent ubiquitination and degradation, followed by a late, transcriptional phase characterized by ALKBH2 stabilization and transcriptional upregulation.

MMS promotes K48-linked ubiquitination and proteasomal degradation of TRIM28 facilitated by TRIM24 and MAGEA6

To investigate how TRIM28 is regulated during alkylation stress, we examined its protein stability and post-translational modification. MMS treatment caused a time- and dose-dependent decrease in TRIM28 protein without affecting its mRNA (Fig. 5A–C), and this reduction was blocked by the proteasome inhibitor MG132, indicating proteasome-dependent degradation. Prolonged low-dose MMS exposure (100 µM, 24–72 h) led to progressive accumulation of ALKBH2, whereas other repair proteins, including ALKBH3, MGMT, and MAGEA3/6, remained unchanged (Fig. EV3A). Consistent with this relationship, pre-exposure to MMS for 12 h caused TRIM28 depletion, elevated basal ALKBH2 levels, and delayed ALKBH2 turnover (Fig. EV3B, C), supporting the notion that TRIM28 loss contributes to ALKBH2 stabilization during prolonged alkylation stress.

To define the underlying mechanism, we next analyzed TRIM28 ubiquitination. Denaturing immunoprecipitation of tagged TRIM28 showed that MMS robustly induced its polyubiquitination (Fig. 5D), which increased in a time-dependent manner in H1299 cells (Fig. 5E). Use of a K48-only ubiquitin mutant confirmed that these modifications are mainly K48-linked, consistent with proteasomal targeting (Fig. 5F). Endogenous analysis using K48-specific Halo-TUBE pulldown verified MMS-induced ubiquitination under physiological conditions (Fig. 5G). In contrast, other genotoxic agents—including TMZ, CDDP, and H_2O_2—failed to trigger either K48-linked ubiquitination or TRIM28 degradation (Fig. 5H; Fig. EV3D–F). Notably, TMZ increased ALKBH2 protein without altering TRIM28 levels (Fig. EV3D), whereas H_2O_2 decreased ALKBH2 but not TRIM28, and CDDP had little effect on either (Fig. EV3E, F). These observations indicate that TRIM28 degradation is a specific response to alkylation stress rather than a general feature of DNA damage.

To test whether SUMOylation contributes to MMS-induced ubiquitination, we examined a SUMO-acceptor-deficient TRIM28 mutant (K554/779/804R) (Ivanov, Peng et al., 2007). K48-specific Halo-TUBE pulldown revealed markedly reduced ubiquitination in the triple-lysine mutant compared with wild-type TRIM28 (Fig. 5I), suggesting that MMS-triggered ubiquitination occurs at—or is influenced by—the same lysine cluster targeted for SUMOylation.

To identify upstream regulators, we next analyzed the TRIM28 interactome (Appendix Table S2) and validated selected candidates. MMS increased TRIM28 association with the E3 ligase TRIM24 and with the proteasome subunits PSMD4 and PSMB5 in a time-dependent manner (Fig. 5J). These interactions were diminished in the SUMO-acceptor mutant (Fig. EV3G). The scaffold protein MAGEA6 enhanced formation of the TRIM28–TRIM24 complex; under MMS treatment, co-expression of MAGEA6 further strengthened TRIM28 binding to TRIM24 and to PSMD4/PSMB5, accompanied by increased K48-linked polyubiquitination (Fig. 5K). Reciprocal co-immunoprecipitation using Flag-TRIM24 confirmed the MMS-inducible TRIM28–TRIM24 interaction and revealed that TRIM24 itself underwent K48-linked ubiquitination upon MMS exposure (Fig. 5L). Overexpression of TRIM24 further amplified MMS-induced ubiquitination and loss of endogenous TRIM28, as shown by Halo-TUBE pulldown and immunoblotting (Figs. 5M and EV3H, I). Collectively, these data demonstrate that MMS specifically promotes K48-linked ubiquitination and proteasomal degradation of TRIM28 through a pathway coordinated by MAGEA6, TRIM24, and SUMO-associated regulation. This stress-specific mechanism places TRIM28 within a broader alkylation-responsive ubiquitin network distinct from other DNA-damage responses.

The TRIM28–ALKBH2 axis links clinical outcome with cellular responses to alkylation stress

To assess the clinical relevance of this pathway in lung adenocarcinoma (LUAD), we first examined ALKBH family expression. Analysis of TCGA (mRNA) and CPTAC (proteomic) datasets revealed significantly increased ALKBH2 and decreased ALKBH3, whereas MGMT levels were unchanged in tumors compared with normal lung tissue (Fig. 6A; Fig. EV4A). Immunohistochemistry confirmed stronger ALKBH2 staining in LUAD than in matched normal samples (Fig. 6B). Consistently, immunoblotting of NSCLC cell lines showed elevated ALKBH2 and reduced ALKBH3 relative to BEAS-2B bronchial epithelial cells (Fig. 6C). In survival

analyses, high ALKBH2 expression correlated with poorer overall survival, whereas high ALKBH3 correlated with better prognosis (Fig. 6D, E).

Because cigarette smoking, a leading cause of lung cancer, also generates alkylating and methylating DNA lesions (Prince, Hammerton et al., 2019), we next examined ALKBH2 expression according to smoking status. ALKBH2 transcripts were significantly higher in LUAD patients with a smoking history than in non-smokers (Fig. EV4B). Moreover, MAGEA6, a functional component of the TRIM28–ALKBH2 axis, harbored somatic mutations more frequently in smokers than in non-smokers, whereas MAGEA3 mutations showed no such enrichment (Fig. EV4C). These findings support the notion that ALKBH2 serves as a clinically relevant biomarker linked to smoking-associated alkylation stress and suggest that MAGEA6 functions as a potential co-regulator in this pathway.

To dissect how TRIM28 and ALKBH2 influence alkylation tolerance, we next employed an acute MMS exposure model. Because prolonged MMS treatment leads to TRIM28 degradation and a delayed increase in ALKBH2, we applied a brief, high-dose MMS pulse (500 µM, 1 h) that induced γ-H2AX while causing limited TRIM28 reduction and minimal change in ALKBH2 levels (Fig. EV4D). Under these conditions, ALKBH2 knockdown using two independent shRNAs reduced proliferation and markedly increased MMS sensitivity (Fig. 6F, G). While TRIM28 overexpression had no significant effect on basal proliferation (Fig. EV4E), it enhanced MMS sensitivity, an effect abolished by the RING-inactive C65/68A mutant and reversed by TRIM28 re-expression in knockdown cells (Fig. 6H, I). Co-expression of ALKBH2 restored resistance, confirming that ALKBH2 functions as the key downstream effector of TRIM28-mediated cytotoxicity (Fig. 6J).

To determine whether MAGEA6 modulates this pathway in a stress-specific manner, we next analyzed its effect on chemosensitivity. MAGEA6 overexpression markedly sensitized H1299 cells to MMS, whereas its knockdown conferred resistance; re-expression of MAGEA6 fully restored MMS sensitivity (Fig. EV4F). When TRIM28 was depleted, MAGEA6 no longer enhanced MMS cytotoxicity, and a TRIM28-binding–deficient mutant (D156E/S157Y) exhibited

greatly reduced activity (Fig. 6K, L). In contrast, MAGEA6 did not alter sensitivity to cisplatin or to the proteasome inhibitor bortezomib (Fig. EV4G, H), indicating that its regulatory role is specific to alkylation stress. Collectively, these data identify the TRIM28–ALKBH2–MAGEA6 axis as a clinically relevant regulatory module that governs cellular tolerance to alkylation damage and links the molecular stress response to chemotherapy outcome.

Disruption of the TRIM28-ALKBH2 axis enhances cisplatin cytotoxicity

Acquired chemoresistance remains a major cause of treatment failure in NSCLC. Such resistance often emerges after prolonged drug exposure, activating adaptive DNA repair responses that overlap with mechanisms of platinum resistance (Martin, Hamilton et al., 2008). Consistent with this concept, the TRIM28/*miR*-125b-5p/CREB1 axis has been reported to mediate cisplatin (CDDP) resistance in NSCLC (Tan, Ma et al., 2022). To explore the clinical relevance of TRIM28, we analyzed 771 NSCLC samples. Among treated cases, platinum-based regimens—predominantly CDDP-containing combinations—represented the main therapeutic modality (Fig. EV5A). High TRIM28 expression correlated with poorer overall survival in chemotherapy-treated patients (Fig. EV5B), but not in untreated cases (Fig. EV5C), indicating that TRIM28 functions as a treatment-dependent prognostic factor in NSCLC.

Given the clinical association of TRIM28 with chemoresistance, we examined whether sustained DNA damage responses contribute to this phenotype. To this end, we established a chronic MMS exposure model to recapitulate prolonged alkylation stress. In this system, cells were exposed to MMS for 48 h. For dose–response analysis (Fig. 7A), H1299 cells were treated with 40–120 μM MMS and analyzed immediately after exposure. For long-term assays (Fig. 7B, C), cells were treated with 100 μM MMS for 48 h, followed by medium replacement and continued culture for an additional 7 days before colony formation analysis. TRIM28 depletion rendered H1299 cells more resistant to MMS, whereas its overexpression in H460 cells increased MMS sensitivity and markedly reduced colony formation. These results demonstrate that TRIM28 enhances susceptibility to alkylation-induced cytotoxicity under chronic exposure.

To clarify the role of ALKBH2, a downstream effector of TRIM28, in drug sensitivity, we

compared MMS and CDDP responses in an ALKBH2-depleted background. Because prolonged

MMS exposure induces ALKBH2 upregulation, we used H1299 cells stably knocked down for

ALKBH2 to minimize confounding effects. Under these conditions, re-expression of Flag-

ALKBH2 increased cell viability upon MMS treatment, confirming its protective role against

alkylation damage. At 2 µM, CDDP alone caused minimal cytotoxicity; however, combined

MMS–CDDP treatment produced a concentration-dependent increase in cytotoxicity, markedly

enhancing MMS-induced cell death. Strikingly, this combination abolished the protective effect

of ALKBH2, indicating that CDDP overrides MMS resistance conferred by elevated ALKBH2

expression (Fig. 7D).

To investigate how CDDP overcomes ALKBH2-mediated MMS resistance, we examined

whether oxidative stress and cumulative DNA damage contribute to this effect. As CDDP is known

to generate reactive oxygen species, including H_2O_2 (Kim, Lee et al., 2010), we tested whether

oxidative stress influences MMS cytotoxicity. Co-treatment of H1299 cells with MMS and

increasing H_2O_2 concentrations led to a dose-dependent loss of viability (Fig. EV5D), consistent

with the enhanced cytotoxicity observed under combined MMS–CDDP exposure. Alkaline comet

assays further showed that low-dose CDDP (2 µM) or MMS (50 µM) alone caused minimal strand

breaks, whereas co-treatment markedly increased tail moments (Fig. 7E; $p < 0.001$). These

findings suggest that oxidative and alkylating stress act cooperatively to exacerbate DNA damage,

thereby counteracting ALKBH2-mediated protection.

To define subtoxic MMS concentrations suitable for combination therapy, we determined IC_{50}

values in H1299 and H460 cells, which were approximately 141.5 µM and 236.8 µM, respectively

(Fig. EV5F). Subsequent assays used MMS doses below these IC_{50} values to minimize single-

agent toxicity. Under these conditions, low-dose MMS significantly reduced the IC_{50} of CDDP in

both H1299 and H460 cells (Fig. 7F, G), demonstrating that prior alkylation stress sensitizes cells

to cisplatin. In colony formation assays, combined MMS and CDDP treatment markedly

suppressed clonogenic survival (Fig. 7H, I). In three-dimensional (3D) tumor spheroids, the

combination similarly reduced spheroid size and viability, accompanied by pronounced

morphological collapse and decreased Calcein-AM fluorescence, indicating a loss of viable cells

(Fig. 7J, K). Collectively, these findings demonstrate that combined alkylation and platinum

treatments synergistically enhance DNA damage and cytotoxicity in NSCLC cells. Furthermore,

they support a model in which persistent alkylation stress disrupts the TRIM28–ALKBH2

regulatory equilibrium, thereby increasing cellular susceptibility to platinum-based chemotherapy.

DISCUSSION

Building on our previous work showing that the deubiquitinase OTUD4 noncanonically stabilizes the alkylation repair enzymes ALKBH2 and ALKBH3 (Zhao et al., 2015), this study defines the opposing ubiquitination arm of this repair pathway. Proteomic profiling of the alkylation-responsive ASCC complex—functionally linked to ALKBH3 and also detected in association with ALKBH2 (Brickner et al., 2017)—identified TRIM28 as an ASCC2-associated component, implicating a connection between transcriptional regulators and ubiquitin signaling during alkylation repair. Consistent with this link, we identified TRIM28, in cooperation with the cancer-testis antigen MAGEA6, as an E3 ligase complex that regulates ALKBH2 stability through K48-linked ubiquitination and proteasomal degradation (Fig. 1–3). Under basal conditions, the TRIM28–MAGEA6 complex maintains steady-state ALKBH2 turnover, while ALKBH2 enhances TRIM28 transcription, forming a feedback loop that supports repair homeostasis. Upon alkylation stress, this feedback loop undergoes dynamic reprogramming. During acute MMS exposure, TRIM28–ALKBH2 binding is strengthened, leading to accelerated ALKBH2 degradation, whereas prolonged exposure redirects ubiquitination toward TRIM28, resulting in its degradation and the subsequent stabilization and transcriptional induction of ALKBH2 (Fig. 8).

This study identifies TRIM28, together with the cancer-testis antigen MAGEA6, as the E3 ligase complex responsible for ALKBH2 ubiquitination and proteasomal turnover, defining the first ubiquitin-dependent mechanism known to regulate ALKBH2 stability. This regulatory module also clarifies how substrate specificity, DNA-engagement–dependent accessibility, and reciprocal feedback regulation are integrated within the ALKBH2 control system. The AlkB family of demethylases is involved in multiple steps of nucleic acid repair (Xu & Bochtler, 2020), yet their post-translational regulation remains poorly understood. Distinct E3 ligases have been identified for other AlkB family members, including TRIM28 for ALKBH5 (Meng et al., 2025) and TRIM21 for FTO (Zhang et al., 2025); however, the mechanism regulating ALKBH2 turnover remains undefined. Notably, unlike canonical MAGE–TRIM interactions (Jin, Pan et al., 2017, Pineda & Potts, 2015), MAGEA6 functions both as a scaffolding cofactor for TRIM28 and as a

direct binding partner of ALKBH2 (Fig. 3B), suggesting a dual-binding mode that enhances substrate specificity under alkylation stress. Although ALKBH2 and ALKBH3 share conserved catalytic cores, TRIM28 preferentially associates with ALKBH2 (Fig. 1I). This selectivity reflects the unique hairpin (finger) motif and flexible long-loop region of ALKBH2—structural elements required for DNA engagement (Yang et al., 2008). Importantly, docking analysis predicts that the TRIM28–ALKBH2 contact region lies near the DNA-binding surface of ALKBH2 (Fig. 1J), and ssDNA competitively reduces their association (Fig. 1K). These observations support a model in which DNA engagement transiently restricts TRIM28 access to ALKBH2, thereby coupling catalytic activity with subsequent ubiquitin-dependent turnover once DNA dissociates. Moreover, under basal conditions, ALKBH2 enhanced TRIM28 expression at both mRNA and protein levels (Fig. 2J–L), thereby forming the transcriptional arm of this feedback loop. This regulation is consistent with previous findings that ALKBH2 modulates promoter methylation and transcription of other targets, including MGMT and BMI1 (Ke, Ye et al., 2020, Malacrida, Di Domizio et al., 2022). Together with previous evidence that the deubiquitinase OTUD4 associates with both ALKBH2 and ALKBH3 (Zhao et al., 2015), these data delineate a reciprocal ubiquitin loop in which TRIM28–MAGEA6-mediated ubiquitination counterbalances OTUD4-dependent deubiquitination, thereby maintaining ALKBH2 homeostasis.

TRIM28 is itself regulated under alkylation stress, introducing a second layer of control within the TRIM28–ALKBH2 pathway. MMS uniquely triggered K48-linked polyubiquitination and proteasomal degradation of TRIM28 (Fig. 5A–G), whereas other genotoxic agents—including temozolomide (TMZ), cisplatin (CDDP), and H_2O_2—did not induce TRIM28 modification or loss (Fig. 5H; Fig. EV3D–F). Instead, these agents produced distinct TRIM28-independent effects on ALKBH2: TMZ increased ALKBH2 protein abundance without reducing TRIM28 levels (Fig. EV3D), H_2O_2 reduced ALKBH2 (Fig. EV3F), and CDDP caused no appreciable changes (Fig. EV3E). The TMZ-driven increase in ALKBH2 aligns with previous reports of transcriptional induction in glioblastoma cells (Johannessen et al., 2013), indicating that TMZ primarily engages a transcriptional, rather than a TRIM28-dependent post-translational, regulatory mechanism.

MMS-induced ubiquitination of TRIM28 was substantially reduced in the K554/779/804R mutant

(Fig. 5I), which targets the major SUMO-acceptor lysines of TRIM28 (Ivanov et al., 2007),

suggesting that MMS-triggered ubiquitination may occur at, or be influenced by, this lysine cluster.

Conversely, interaction with TRIM24 further enhanced TRIM28 ubiquitination upon MMS

treatment, suggesting that TRIM24 facilitates this process under alkylation stress. This stress-

specific turnover contrasts with the RNF4-mediated degradation of phosphorylated and

SUMOylated TRIM28 reported in the double-strand break response (Kuo et al., 2014), indicating

that different forms of genotoxic stress engage distinct ubiquitin pathways. Consistent with this

notion, alkylation damage triggers specialized ubiquitin responses—for example, MMS promotes

HLTF degradation without affecting UV-induced PCNA monoubiquitination (Lin, Zeman et al.,

2011), and also activates the E3 ligase RNF113A to mediate recruitment of the ASCC–ALKBH3

repair complex (Tsao, Brickner et al., 2021). Collectively, these observations place TRIM28

within a broader framework of alkylation-specific ubiquitin regulation, indicating that MMS

engages a distinct ubiquitin signaling route that couples targeted protein turnover to DNA repair

adaptation. Notably, TRIM24 has been reported to cooperate with TRIM28 in transcriptional

programs across multiple cancers, where TRIM28 both stabilizes TRIM24 and promotes its

transcriptional activity (Fong, Zhao et al., 2018, Herquel, Ouararhni et al., 2011). In contrast, in

our system TRIM24 enhanced TRIM28 ubiquitination and promoted its degradation in response

to MMS treatment, indicating a stress-induced reversal of their regulatory relationship. This MMS-

selective mechanism limits TRIM28 stability and may facilitate the remodeling of repair-

associated chromatin complexes under prolonged alkylation stress.

A key mechanistic insight from this study is the discovery of a time-resolved TRIM28–

ALKBH2 feedback loop that dynamically modulates DNA repair during alkylation stress. This

loop functions as a biphasic molecular switch, acting sequentially to balance repression and

activation phases. During acute MMS exposure, TRIM28 binds ALKBH2 to promote its K48-

linked ubiquitination and proteasomal degradation, thereby transiently restraining repair capacity

(Fig. 4). As MMS exposure persists, TRIM28 undergoes ubiquitination and degradation, whereas

ALKBH2 becomes stabilized and transcriptionally induced (Fig. 4-5), thereby marking a temporal transition from repression to activation of repair capacity. This transition parallels the functional switch in demethylation repair, where ALKBH2 catalyzes the oxidative demethylation of damaged bases. This biphasic switch exemplifies a general principle of stress adaptation, reminiscent of the time-delayed feedback observed in the ATM–TRIM28 (KAP1) and p53–MDM2 circuits, which coordinate DNA repair and checkpoint recovery (Jaiswal, Benada et al., 2017, Wang, Ivanov et al., 2005). Unlike these canonical pathways, the TRIM28–ALKBH2 module integrates ubiquitin- and transcription-linked feedback loops, thus enabling the E3 ligase and its substrate to function as reciprocal regulators. Under basal conditions, ALKBH2 sustains TRIM28 expression at both mRNA and protein levels (Fig. 2J–L), thereby maintaining repair homeostasis. Prolonged alkylation stress, however, leads to TRIM28 depletion and ALKBH2 accumulation, thereby disrupting this balance and shifting the system toward enhanced repair capacity and increased tolerance to persistent damage. Such a biphasic regulatory mechanism may enable cells to fine-tune repair activity according to the duration and intensity of alkylation stress, thereby enhancing DNA repair adaptability during chemotherapy.

Clinically, the TRIM28–ALKBH2 regulatory loop emerges as a relevant determinant of NSCLC chemotherapy response. ALKBH2 is frequently upregulated in lung adenocarcinoma and correlates with poorer overall survival (Fig. 6A–D), consistent with its function in counteracting DNA alkylation damage (Duncan et al., 2002, Feng et al., 2022). By contrast, TRIM28 correlates with patient outcome only in chemotherapy-treated cohorts (Fig. EV5A-C), suggesting that its regulatory activity becomes functionally engaged under therapeutic stress. MAGEA6 mutations, which are enriched in smokers (Fig. EV4C), may link smoking-induced alkylation stress to disruption of this proteolytic module. Mechanistically, TRIM28 functions as the cognate E3 ligase for ALKBH2, with MAGEA6 serving as its scaffolding adaptor, together forming a ubiquitin-mediated checkpoint that modulates repair dynamics under chemotherapeutic stress (Figs. 1–3). Functionally, TRIM28 overexpression increased MMS sensitivity, whereas ALKBH2 re-expression reversed this effect (Fig. 6F–I), indicating that ALKBH2 acts as the primary effector

of TRIM28-dependent cytotoxicity. Combining MMS with cisplatin enhanced DNA damage, suppressed clonogenic survival, and overcame ALKBH2-mediated alkylation tolerance, resulting in synergistic cytotoxicity (Fig. 7B–K). Notably, although TRIM28 has been implicated in promoting NSCLC chemoresistance through mechanisms such as *miR*-125b-5p/CREB1 signaling, E2F1 regulation, and DNA-damage control (Ginn, Maltas et al., 2023, Liu, Xiao et al., 2017, Lundholm, Hååg et al., 2013, Saxena, Quan et al., 2025, Tan et al., 2022), it instead enhanced cytotoxicity by promoting ALKBH2 degradation in our MMS-based system. This context dependence likely reflects the mechanistic distinction between alkylation stress and platinum-induced DNA crosslinking. Prolonged MMS exposure results in TRIM28 degradation accompanied by increased ALKBH2 expression (Fig. 5A–H; Fig. EV3D–F), suggesting an adaptive enhancement of DNA repair capacity during sustained alkylation stress. Although this adaptation may confer partial tolerance to MMS, the concomitant loss of TRIM28 could sensitize cells to platinum, consistent with the synergistic cytotoxicity observed under combined treatment (Fig. 7D–K). Clinically, these findings are consistent with patient data showing that TRIM28 acquires prognostic significance during chemotherapy, where high expression correlates with poorer overall survival. This association supports a potential therapeutic rationale for combining alkylating and platinum agents in tumors with elevated TRIM28 expression.

In summary, this study delineates a stress-dependent TRIM28–ALKBH2 feedback loop that modulates alkylation repair capacity in NSCLC cells. Under basal conditions, the TRIM28–MAGEA6 complex mediates K48-linked polyubiquitination and proteasomal turnover of ALKBH2. Acute MMS exposure enhances assembly of the TRIM28–ALKBH2 complex and promotes ALKBH2 degradation, whereas prolonged MMS exposure induces K48-linked ubiquitination of TRIM28 and its proteasomal degradation, leading to ALKBH2 stabilization and transcriptional upregulation. This stress-responsive circuit integrates post-translational mechanisms with transcriptional control of ALKBH2, thereby collectively modulating cellular responses to alkylating agents. The upstream cues that initiate TRIM28 degradation and their coordination with transcriptional programs remain to be clarified. Future studies using patient-

552 derived and *in vivo* systems will provide further insight into defining these mechanisms and

553 assessing their relevance to chemotherapy response in NSCLC.

554 **Methods**

555 **Reagents and tools table**

Reagent/Resource	Reference or Source	Identifier or Catalog Number
Cell Lines		
HEK293T (*H. sapiens*)	ATCC	CRL-11268
NCI-H1299 (*H. sapiens*)	Cell Bank of the Chinese Academy of Sciences	SCSP-589
A549 (*H. sapiens*)	Cell Bank of the Chinese Academy of Sciences	SCSP-503
NCI-H460 (*H. sapiens*)	Cell Bank of the Chinese Academy of Sciences	SCSP-584
Antibodies		
Rabbit anti-TRIM28	Proteintech	Cat #15202-1-AP
Mouse anti-ALKBH2	Sigma-Aldrich	Cat #A8228
Rabbit anti-HA	Proteintech	Cat #51064-2-AP
Mouse anti-Flag	Sigma-Aldrich	Cat #F3165
Mouse anti-Myc	Millipore	Cat #05-419
Mouse anti-His	Proteintech	Cat #66005-1-Ig
Mouse anti-MBP	New England Biolabs	Cat #E8030S
Rabbit anti-MAGEA3/6	Zenbio Science	Cat #R24906
Rabbit anti-TRIM24	Zenbio Science	Cat #R27382
Rabbit anti-ALKBH3	Millipore	Cat #09-882
Rabbit anti-MGMT	Proteintech	Cat #17195-1-AP
Mouse anti-P53	Santa Cruz	Cat #sc-126
Rabbit anti-γ-H2AX	Abmart	Cat #T56572
Rabbit anti-H2AX	Cusabio	Cat #Ab-142
Rabbit anti-K48-Ubiquitin	Millipore	Cat #05-1307
Mouse anti-ubiquitin	Santa Cruz	Cat #sc-8017
Rabbit anti-PSMD4	Selleck	Cat #F1309
Rabbit anti-PSMB5	Selleck	Cat #F1421
HRP-conjugated β-actin	Genscript Biotech	Cat #A00730

HRP-conjugated β-tubulin	Earthox	Cat #E021043
HRP-conjugated anti-rabbit	Proteintech	Cat #SA00001-2
HRP-conjugated anti-mouse	Proteintech	Cat #SA00001-1
Alexa Fluor 488 goat anti-mouse	Thermo Fisher Scientific	Cat #A-11029
Alexa Fluor 594 goat anti-rabbit	Thermo Fisher Scientific	Cat #A-11037
Recombinant DNA		
pHAGE-puro-Flag-TRIM28	This study	N/A
pHAGE-puro-3xHA-TRIM28	This study	N/A
pHAGE-puro-His-TRIM28	This study	N/A
pMSCV-puro-TAP-TRIM28	This study	N/A
pHAGE-puro-Flag-ALKBH2	This study	N/A
pHAGE-blast-HA-ALKBH2	This study	N/A
pMSCV-blast-TAP-ALKBH2	This study	N/A
pCMV-Flag-TRIM24	This study	N/A
pCMV-Flag-ALKBH2	This study	N/A
pCMV-Flag-ALKBH2 H171/D173A	This study	N/A
pCMV-Flag-ALKBH3	This study	N/A
pCMV-Flag-MGMT	This study	N/A
pHAGE-puro-Flag-MAGEA6	This study	N/A
pHAGE-puro-Flag-MAGEA6 Mut	This study	N/A
pCMV-Myc-MAGEA6	This study	N/A
pLKO.1 puro	Addgene	Cat #8453
lentiCRISPR v2	Addgene	Cat #52961
pcDNA3.1-HA-ub	Zhao et al., 2015	N/A
pRK5-HA-ub K48-only	Zhao et al., 2015	N/A
pRK5-HA-ub K48R	Zhao et al., 2015	N/A
psPAX2	Zhao et al., 2015	N/A
pMD2.G	Zhao et al., 2015	N/A
pENTR-4	Zhao et al., 2015	N/A
pHAGE-CMV-Flag/HA/His	Zhao et al., 2015	N/A
Oligonucleotides and other sequence-based reagents		

ssDNA	This study	CAAAGCTTCCGAGTAC TTGCAGCTG
shTRIM28, shALKBH2, shMAGEA6, sgTRIM28	This study	Appendix Table S3
qPCR Primer	This study	Appendix Table S4

Chemicals, Enzymes and other reagents

Cell Culture & Transfection

Fetal bovine serum (FBS)	Gibco	Cat #10099-141C
Dulbecco's modified Eagle's medium (DMEM)	Gibco	Cat #C11995500BT
Penicillin-streptomycin	Gibco	Cat #15140122
0.25%Trypsin-EDTA	Gibco	Cat #C25200-072
Polybrene	Sigma-Aldrich	Cat #28728-55-4
Lipo8000™ Transfection Reagent	Beyotime	Cat #C0533

Chemicals, Inhibitors & Drugs

Cycloheximide	Cell Signaling Technology	Cat #2112
Methyl methanesulfonate	Sigma-Aldrich	Cat #129925
Cisplatin	MedChemExpress	Cat #HY-17394
Temozolomide	Selleck	Cat #S1237
Bortezomib	MedChemExpress	Cat #HY-10227
Ascorbic acid	Shanghai Hu Shi	Cat #10004014
Protease inhibitor cocktail	Beyotime	Cat #P1051

Immunoprecipitation & Affinity Purification

Anti-Flag M2-agarose beads	Sigma-Aldrich	Cat #L00425
Protein A/G Magnetic Beads	Selleck	Cat #B23202
Anti-HA magnetic beads	Selleck	Cat #B26201
HaloLink™ resin	Promega	Cat #G1915
Amylose Resin	New England Biolabs	Cat #E8021S

Molecular Biology

Gateway™ LR Clonase™ II Enzyme mix	Invitrogen	Cat #11791020
RNAeasy™ RNA Isolation Kit	Beyotime	Cat #R0026

SYBR qPCR Mix	Tsingke	Cat #TSE501
Assays & Kits		
CCK8 assay	ApexBio	Cat #K1018
Comet Assay Kit	Keygentec	Cat #KGA1302
Myco-Lumi Luminescent Mycoplasma Detection Kit	Beyotime	Cat #C0297S
Silver Staining Kit	Sangon Biotech	Cat #C500021
Staining, Imaging & Fixation		
DAPI	Beyotime	Cat #C1002
Hoechst 33342	BD Biosciences	Cat #561908
Calcein/PI Live/Dead Assay	Beyotime	Cat #C2015S
4% paraformaldehyde	Beyotime	Cat #P0099
0.1% crystal violet	Beyotime	Cat #C0121
Software		
GraphPad Prism	https://www.graphpad.com	Version 9.5.0
Pymol	https://pymol.org/	Version 2.6.0a0
RStudio	https://posit.co/	Version 4.4.2
ImageJ	https://imagej.en.softonic.com	Version 1.54K

Methods and protocols

Cell culture

Cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% heat-inactivated fetal bovine serum and penicillin–streptomycin. Cultures were maintained at 37 °C in a humidified atmosphere containing 5% CO_2. All experiments were performed using low-passage cells (passages 5–15). Mycoplasma contamination was routinely monitored using a luminescence-based detection assay.

Plasmid constructs, mutagenesis, and viral infection

Full-length human TRIM28 and ALKBH2 cDNAs were used as templates for PCR-based site-directed mutagenesis. TRIM28 deletion and point mutants (including BCC, residues 141–400, and the RING-inactive C65/68A variant) were generated by PCR, cloned into pENTR-4, and

transferred into pHAGE-CMV-Flag/HA/His through Gateway LR recombination. ALKBH2 variants were cloned into pCMV-Flag for mammalian expression and into pET28a-His-2×Flag for bacterial expression. All constructs were verified by Sanger sequencing. MAGEA6 wild-type cDNA, obtained by reverse transcription of A549 RNA, and H1299-derived MAGEA6 variants were generated analogously. Transient transfections were performed using Lipo8000.

Lentiviruses were produced in HEK293T cells by co-transfecting pLKO.1 or pHAGE vectors with the packaging plasmids psPAX2 and pMD2.G. Viral supernatants were collected at 48–72 h, clarified, and filtered through 0.45-µm membranes. Target cells were infected in the presence of polybrene (6 µg/mL) for 12 h. Stable cell lines were selected with puromycin (1 µg/mL) or blasticidin (5 µg/mL) for 3 days and maintained thereafter.

shRNA and CRISPR/Cas9 knockdown/knockout

For shRNA-mediated knockdown, annealed oligonucleotides encoding target-specific sequences were cloned into the pLKO.1 vector and packaged into lentiviral particles in HEK293T cells. Target cells were transduced and selected with puromycin for 3 days to establish stable knockdown lines. Knockdown efficiency was confirmed by western blotting (WB). A non-targeting shRNA was used as a control.

For CRISPR/Cas9-mediated knockout of TRIM28, single-guide RNAs (sgRNAs) were designed according to established criteria and cloned into the lentiCRISPRv2 vector. Lentiviral particles were produced in HEK293T cells and used to transduce target cells, followed by puromycin selection. Single-cell clones were generated by limiting dilution, and successful knockout was verified by WB. All oligonucleotide sequences are listed in Appendix Table S3.

Immunoprecipitation and western blotting

For overexpression immunoprecipitation (IP), HEK293T cells were transfected with the indicated plasmids using Lipo8000 and cultured for 2–3 days before lysis. Cells were lysed in buffer containing 150 mM NaCl, 50 mM Tris-HCl (pH 7.9), 1% Triton X-100, 10% glycerol, and 2 mM

DTT supplemented with protease and phosphatase inhibitors. Clarified lysates were incubated

overnight at 4 °C with anti-Flag M2-agarose beads or anti-HA-agarose beads. For denaturing IP,

lysates were boiled in TBS containing 1% SDS and diluted with lysis buffer to a final concentration

of 0.1% SDS before IP.

For endogenous IP from H1299 cells, lysates were prepared in the same buffer and cleared by

centrifugation (15,000 × g, 10 min, 4 °C). Supernatants were pre-cleared with Protein A/G

magnetic beads for 1 h and incubated overnight at 4 °C with anti-TRIM28 (1.5 µg). Immune

complexes were captured with Protein A/G magnetic beads, washed in lysis buffer, and eluted in

Laemmli buffer. Input and flow-through fractions were analyzed in parallel by WB to verify

expression and loading.

For ssDNA competition assays, purified Flag-ALKBH2 was added to HEK293T whole-cell

lysates supplemented with synthesized single-stranded DNA (ssDNA) oligonucleotides and

incubated overnight at 4 °C. Disruption of the TRIM28–ALKBH2 interaction was assessed by

Flag IP, followed by WB.

For WB, proteins were separated by SDS–PAGE and transferred to PVDF membranes.

Membranes were blocked with 5% non-fat milk in TBST for 1 h at room temperature, incubated

with primary antibodies overnight at 4 °C, and then with HRP-conjugated secondary antibodies

for 1–1.5 h at room temperature. Signals were detected using enhanced chemiluminescence (ECL).

In vitro MBP pull-down assay

Recombinant maltose-binding protein (MBP)–tagged TRIM28 (MBP–TRIM28), MBP–

MAGEA6, MBP alone, and His–Flag–tagged ALKBH2 (His–Flag–ALKBH2) were expressed in

E. coli Rosetta (DE3). Protein expression was induced with IPTG according to standard procedures.

Bacterial pellets were collected and lysed, and MBP fusion proteins were purified using amylose

resin.

For pull-down assays, purified MBP–TRIM28, MBP–MAGEA6, or MBP alone was

immobilized on amylose resin and washed with binding buffer. The resin was then incubated with

purified His–Flag–ALKBH2 at 4 °C with gentle rotation. After extensive washing, bound proteins

were eluted by boiling in Laemmli sample buffer and analyzed by WB using anti-His or anti-Flag

antibodies.

Immunofluorescence (IF)

Cells grown on glass coverslips were treated as indicated, fixed in 4% paraformaldehyde for 30

min, and permeabilized with 0.5% Triton X-100 in PBS for 15 min at room temperature. After

washing with PBST, samples were incubated with primary antibodies overnight at 4 °C. The

following day, coverslips were washed and incubated with fluorophore-conjugated secondary

antibodies together with Hoechst 33342 or DAPI for 1.5 h at 37 °C. Coverslips were mounted and

imaged using an Olympus fluorescence microscope. Colocalization analysis was performed using

ImageJ, and Pearson correlation coefficients were calculated when required.

Cycloheximide chase assay

For cycloheximide (CHX) chase assays, cells were treated with 150 µg/mL CHX for the indicated

durations. Cells were harvested at each time point and analyzed by WB to assess protein stability.

RNA isolation and qRT-PCR

Total RNA was isolated using the RNAeasy™ RNA Isolation Kit, following the manufacturer's

instructions. RNA concentration and purity were assessed using a NanoDrop spectrophotometer

(Thermo Fisher Scientific) by measuring absorbance at 260 nm and calculating the A260/A280

and A260/A230 ratios. First-strand cDNA synthesis was performed using the BeyoFast™ cDNA

Synthesis Kit, followed by qPCR using the BeyoFast™ SYBR Green qPCR Mix, according to the

manufacturer's guidelines. Gene expression levels were normalized to *β-actin* as an internal

control. All qRT-PCR primer sequences are listed in Appendix Table S4.

Protein–protein docking and structural modeling

The tertiary structures of human ALKBH2 (PDB ID: 3BTZ), ALKBH3 (PDB ID: 2NCU), and the

RBCC domain of TRIM28 (PDB ID: 6QAJ) were retrieved from the Protein Data Bank

(http://www.rcsb.org). Protein–protein docking simulations were conducted using the GRAMM-

X web server (http://gramm.compbio.ku.edu). Predicted complex interfaces between TRIM28 and

MAGEA6 were modeled using AlphaFold3 (https://alphafoldserver.com/). The resulting docking

complexes were visualized and analyzed using PyMOL.

Mass spectrometry

TRIM28 complexes from H1299 cells stably expressing TAP-TRIM28 were immunoaffinity-

purified using Flag M2 beads, eluted with Flag peptide, separated by SDS–PAGE, and visualized

by silver staining. Afterward, samples were analyzed by LC–MS/MS. Mass spectrometry was

performed by Shanghai Baipu Company utilizing an Easy nLC 1200 system (Thermo Fisher

Scientific) with a Dr. Maisch GmbH column (75 µm × 150 mm, 3 µm C18) for gradient separation.

Data-dependent acquisition (DDA) MS was performed on a Q-Exactive HF-X mass spectrometer

(Thermo Fisher Scientific). Peptides were matched against the UniProt Reference proteome *Homo*

sapiens (Human) [9606] database (81791-20230317.fasta), obtained from

https://www.uniprot.org/proteomes/UP000005640.

Affinity purification of K48-linked polyubiquitinated proteins (Halo-TUBE)

K48-linked ubiquitin chains and ubiquitinated proteins were isolated from cell extracts using a

tandem ubiquitin-binding entity (TUBE) containing Halo-tag (Halo-TUBE) resin under denaturing

conditions. Briefly, cells were harvested, lysed, and denatured by boiling in TBS containing 1%

SDS, then diluted with lysis buffer to reduce SDS concentration to 0.1% before affinity

purification. Extracts were incubated with 10 µL of HaloLink™ resin (Promega) pre-bound to

purified Halo-TUBE protein (10 µg per sample). The clarified extracts were incubated with Halo-

TUBE-bound resin overnight at 4 °C with rotation. Subsequently, the resin was washed five times

with washing buffer, boiled in Laemmli sample buffer, and analyzed by WB.

Cell proliferation, CDDP sensitivity, and synergy assays

For the MMS Sensitivity Assay, cells (2,000–5,000 per well, in 96-well plates) were exposed to high-concentration methyl methanesulfonate (MMS; 200–800 μM) for 1 h at 37 °C, followed by replacement with fresh medium, or low-concentration MMS (0–200 μM) continuously for 48 h. Cell viability was assessed at 48 h using the CCK-8 assay. For the Cisplatin Sensitivity Assay, cells were treated with cisplatin (CDDP; 0–40 μM) for 48 h, and cell viability was measured using the CCK-8 assay. The half-maximal inhibitory concentration (IC_{50}) was calculated using GraphPad Prism software. For the Synergy Assay, cells were treated with cisplatin (CDDP) alone or in combination with low-concentration MMS for 48 h. Cell viability was quantified using the CCK-8 assay.

Colony formation assay

To assess clonogenic survival, H460 cells were seeded in 24-well plates (400 cells per well) and incubated overnight to allow cell attachment. The following day, DMSO (vehicle control) or MMS was added to each well at the indicated concentrations. For combination treatment, the medium was replaced with fresh medium after 48 h of treatment with MMS alone, cisplatin (CDDP) alone, or their combination. After 5–10 days of incubation to allow colony formation, the medium was removed, and cells were fixed with 4% paraformaldehyde for 10 minutes at room temperature. Following fixation, cells were stained with 0.1% crystal violet according to the manufacturer's instructions. Plates were washed three times with distilled water, air-dried, and the number of colonies was counted.

Alkaline comet assay

Alkaline single-cell agarose gel electrophoresis (comet) assays were performed using a Comet Assay Kit, following the manufacturer's instructions. Comet images were visualized using an Olympus fluorescence microscope (Olympus Corporation). The percentage of DNA in the tail (Tail DNA%) and the Tail Moment were quantified using ImageJ software with the Comet Assay plugin. Tail DNA% was calculated as: Tail DNA% = (Tail DNA Intensity / (Tail DNA Intensity

703 + Head DNA Intensity)) × 100%, and the Tail Moment was calculated as: Tail Moment = Tail

704 Length × Tail DNA%.

3D tumorsphere culture

H1299 cells (~2,000 cells per drop) were cultured as hanging drops for 72 hours to facilitate

spheroid formation and then transferred to ultra-low attachment 96-well plates (LV-ULA002-

96UW, Liver Biotech). Tumor spheres were subsequently exposed to the indicated concentrations

of MMS and/or cisplatin (CDDP) for 48 hours. After drug exposure, the culture medium was

replaced, and spheroids were cultured for an extended period (up to 3–4 weeks). Tumorsphere

viability and cytotoxicity were assessed using a Calcein-AM/Propidium Iodide (PI) Live/Dead

assay kit. Fluorescence images distinguishing live (green) and dead (red) cells were acquired using

an Olympus fluorescence microscope (Olympus Corporation).

Bioinformatics and clinical data analysis

Clinical data for TCGA patients with LUAD and LUSC were obtained from the TCGA data portal

(https://tcga-data.nci.nih.gov/tcga/). Gene expression data from TCGA and protein expression data

from CPTAC were analyzed using ULCAN. Survival analysis was performed using the Kaplan–

Meier Plotter (http://kmplot.com/analysis) with default settings for overall survival, and hazard

ratios and log-rank p-values were reported. Images of ALKBH2 immunohistochemistry (IHC)

were downloaded from the Human Protein Atlas (http://www.proteinatlas.org).

MAGEA6/MAGEA3 mutations were surveyed using COSMIC (http://cancer.sanger.ac.uk/cosmic)

and cBioPortal (http://www.cbioportal.org). Cutoffs and covariates are specified in the

corresponding figure legends.

Statistical analysis

725 All graphs and statistical analyses were performed using GraphPad Prism version 9.5.0. Data are

726 presented as means ± standard deviation (SD) from at least three independent experiments ($n \geq 3$).

727 Statistical significance was determined using an unpaired, two-tailed Student's t-test. P values <

728 0.05 were considered significant (*$p < 0.05$, **$p < 0.01$).

Author contributions

Zhiming Sun: Writing – original draft, Visualization, Resources, Investigation, Data curation. **Yunfang Deng:** Formal analysis, Investigation, Data curation. **Yue Liu:** Investigation. **Zhaohui Liu:** Visualization, Formal analysis. **Jiabing Li:** Visualization, Formal analysis. **Lihui Wu:** Investigation. **Liyuan Zeng, Xiaorong Feng, Lin Miao:** Investigation, Data curation. **Ying Sheng, Bei Chen, Yuming He, Ye Liu:** Investigation. **Yu Zhao:** Writing – review & editing, Visualization, Supervision, Funding acquisition, Conceptualization.

Disclosure and competing interests statement

The authors declare that they have no competing interests.

Data availability

All data supporting the findings of this study are available within the article and its Supplementary Information.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (31971232) and the Natural Science Foundation of Hunan Province (2019JJ40185) to YZ.

References

Brickner JR, Soll JM, Lombardi PM, Vagbo CB, Mudge MC, Oyeniran C, Rabe R, Jackson J, Sullender ME, Blazosky E et al. (2017) A ubiquitin-dependent signalling axis specific for ALKBH-mediated DNA dealkylation repair. *Nature* 551: 389-393

Bu X, Qu X, Guo K, Meng X, Yang X, Huang Q, Dou W, Feng L, Wei X, Gao J et al. (2021) CD147 confers temozolomide resistance of glioma cells via the regulation of β-TrCP/Nrf2 pathway. *International journal of biological sciences* 17: 3013-3023

Chang G, Xie GS, Ma L, Li P, Li L, Richard HT (2023) USP36 promotes tumorigenesis and drug sensitivity of glioblastoma by deubiquitinating and stabilizing ALKBH5. *Neuro Oncol* 25: 841-853

Chen Y, Pang W, Ma G, Yan Y, Xiao Z, Chen Y, Ding Z, Chen L, Hou X, Cao H (2025) March5-mediated Trim28 degradation preserves islet β-cell function in mice. *Nature communications* 16: 7073

Chen Z, Zhou Y, Xue C, Zeng L, Deng S, Xu Z, Li M, Zhao H, He X, Liu S et al. (2025) Psychological stress-induced ALKBH5 deficiency promotes tumour innervation and pancreatic cancer via extracellular vesicle transfer of RNA. *Nat Cell Biol* 27: 1035-1047

Cui YH, Wei J, Fan H, Li W, Zhao L, Wilkinson E, Peterson J, Xie L, Zou Z, Yang S et al. (2024) Targeting DTX2/UFD1-mediated FTO degradation to regulate antitumor immunity. *Proceedings of the National Academy of Sciences of the United States of America* 121: e2407910121

Czerwińska P, Mazurek S, Wiznerowicz M (2017) The complexity of TRIM28 contribution to cancer. *Journal of biomedical science* 24: 63

Dai W, Tian R, Yu L, Bian S, Chen Y, Yin B, Luan Y, Chen S, Fan Z, Yan R et al. (2024) Overcoming therapeutic resistance in oncolytic herpes virotherapy by targeting IGF2BP3-induced NETosis in malignant glioma. *Nature communications* 15: 131

Doyle JM, Gao J, Wang J, Yang M, Potts PR (2010) MAGE-RING protein complexes comprise a family of E3 ubiquitin ligases. *Molecular cell* 39: 963-74

Drabløs F, Feyzi E, Aas PA, Vaagbø CB, Kavli B, Bratlie MS, Peña-Diaz J, Otterlei M, Slupphaug G, Krokan HE (2004) Alkylation damage in DNA and RNA--repair mechanisms and medical significance. *DNA Repair (Amst)* 3: 1389-407

Duncan T, Trewick SC, Koivisto P, Bates PA, Lindahl T, Sedgwick B (2002) Reversal of DNA alkylation damage by two human dioxygenases. *Proceedings of the National Academy of Sciences of the United States of America* 99: 16660-5

Feng S, Xu Z, Peng J, Zhang M (2022) The AlkB Family: Potential Prognostic Biomarkers and Therapeutic Targets in Glioblastoma. *Frontiers in oncology* 12: 847821

Feng Y, Gao J, Yang M (2011) When MAGE meets RING: insights into biological functions of MAGE proteins. *Protein Cell* 2: 7-12

Fong KW, Zhao JC, Song B, Zheng B, Yu J (2018) TRIM28 protects TRIM24 from SPOP-mediated degradation and promotes prostate cancer progression. *Nature communications* 9: 5007

Fu D, Calvo JA, Samson LD (2012) Balancing repair and tolerance of DNA damage caused by alkylating agents. *Nat Rev Cancer* 12: 104-20

Ginn L, Maltas J, Baker MJ, Chaturvedi A, Wilson L, Guilbert R, Amaral FMR, Priest L, Mole H, Blackhall F et al. (2023) A TIAM1-TRIM28 complex mediates epigenetic silencing of protocadherins to promote migration of lung cancer cells. *Proceedings of the National Academy of Sciences of the United States of America* 120: e2300489120

Goodarzi AA, Kurka T, Jeggo PA (2011) KAP-1 phosphorylation regulates CHD3 nucleosome remodeling during the
DNA double-strand break response. *Nature structural & molecular biology* 18: 831-9

Gu J, Chen J, Xiang S, Zhou X, Li J (2023) Intricate confrontation: Research progress and application potential of
TRIM family proteins in tumor immune escape. *Journal of advanced research* 54: 147-179

Gu P, Deng W, Lei M, Chang S (2013) Single strand DNA binding proteins 1 and 2 protect newly replicated telomeres.
Cell research 23: 705-19

Han T, Tong W, Xie J, Guo X, Zhang L (2024) FOXF2 suppressed esophageal squamous cell carcinoma by reducing
M2 TAMs via modulating RNF144A-FTO axis. *International immunopharmacology* 143: 113422

Hatakeyama S (2011) TRIM proteins and cancer. *Nat Rev Cancer* 11: 792-804

Hegi ME, Diserens AC, Gorlia T, Hamou MF, de Tribolet N, Weller M, Kros JM, Hainfellner JA, Mason W, Mariani
L et al. (2005) MGMT gene silencing and benefit from temozolomide in glioblastoma. *The New England journal*
of medicine 352: 997-1003

Herquel B, Ouararhni K, Khetchoumian K, Ignat M, Teletin M, Mark M, Béchade G, Van Dorsselaer A, Sanglier-
Cianférani S, Hamiche A et al. (2011) Transcription cofactors TRIM24, TRIM28, and TRIM33 associate to form
regulatory complexes that suppress murine hepatocellular carcinoma. *Proceedings of the National Academy of*
Sciences of the United States of America 108: 8212-7

Ivanov AV, Peng H, Yurchenko V, Yap KL, Negorev DG, Schultz DC, Psulkowski E, Fredericks WJ, White DE, Maul
GG et al. (2007) PHD domain-mediated E3 ligase activity directs intramolecular sumoylation of an adjacent
bromodomain required for gene silencing. *Molecular cell* 28: 823-37

Jaiswal H, Benada J, Müllers E, Akopyan K, Burdova K, Koolmeister T, Helleday T, Medema RH, Macurek L,
Lindqvist A (2017) ATM/Wip1 activities at chromatin control Plk1 re-activation to determine G2 checkpoint
duration. *The EMBO journal* 36: 2161-2176

Jana S, Mondal M, Mahale S, Gupta B, Prasasvi KR, Kandasami L, Jha N, Chowdhury A, Santosh V, Kanduri C et al.
(2024) PITAR, a DNA damage-inducible cancer/testis long noncoding RNA, inactivates p53 by binding and
stabilizing TRIM28 mRNA. *eLife* 12

Jin X, Pan Y, Wang L, Zhang L, Ravichandran R, Potts PR, Jiang J, Wu H, Huang H (2017) MAGE-TRIM28 complex
promotes the Warburg effect and hepatocellular carcinoma progression by targeting FBP1 for degradation.
Oncogenesis 6: e312

Johannessen TC, Prestegarden L, Grudic A, Hegi ME, Tysnes BB, Bjerkvig R (2013) The DNA repair protein
ALKBH2 mediates temozolomide resistance in human glioblastoma cells. *Neuro Oncol* 15: 269-78

Ke B, Ye K, Cheng S (2020) ALKBH2 inhibition alleviates malignancy in colorectal cancer by regulating BMI1-
mediated activation of NF-κB pathway. *World journal of surgical oncology* 18: 328

Kim HJ, Lee JH, Kim SJ, Oh GS, Moon HD, Kwon KB, Park C, Park BH, Lee HK, Chung SY et al. (2010) Roles of
NADPH oxidases in cisplatin-induced reactive oxygen species generation and ototoxicity. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience 30: 3933-46

Kim SH, Lee S, Lee CH, Lee MK, Kim YD, Shin DH, Choi KU, Kim JY, Park DY, Sol MY (2009) Expression of
cancer-testis antigens MAGE-A3/6 and NY-ESO-1 in non-small-cell lung carcinomas and their relationship with
immune cell infiltration. *Lung* 187: 401-11

Kuo CY, Li X, Kong XQ, Luo C, Chang CC, Chung Y, Shih HM, Li KK, Ann DK (2014) An arginine-rich motif of
ring finger protein 4 (RNF4) oversees the recruitment and degradation of the phosphorylated and SUMOylated
Krüppel-associated box domain-associated protein 1 (KAP1)/TRIM28 protein during genotoxic stress. *J Biol*

Chem 289: 20757-72

Le HP, Ma X, Vaquero J, Brinkmeyer M, Guo F, Heyer WD, Liu J (2020) DSS1 and ssDNA regulate oligomerization
of BRCA2. *Nucleic acids research* 48: 7818-7833

Lee DH, Goodarzi AA, Adelmant GO, Pan Y, Jeggo PA, Marto JA, Chowdhury D (2012) Phosphoproteomic analysis
reveals that PP4 dephosphorylates KAP-1 impacting the DNA damage response. *The EMBO journal* 31: 2403-15

Lee DH, Jin SG, Cai S, Chen Y, Pfeifer GP, O'Connor TR (2005) Repair of methylation damage in DNA and RNA by
mammalian AlkB homologues. *J Biol Chem* 280: 39448-59

Leiros I, Nabong MP, Grøsvik K, Ringvoll J, Haugland GT, Uldal L, Reite K, Olsbu IK, Knaevelsrud I, Moe E et al.
(2007) Structural basis for enzymatic excision of N1-methyladenine and N3-methylcytosine from DNA. *The*
EMBO journal 26: 2206-17

Li J, Feng X, Liu Z, Deng Y, Sun Z, Chen B, Wu L, Wang X, Miao L, Zeng L et al. (2025) USP7 promotes
temozolomide resistance by stabilizing MGMT in glioblastoma. *Cell death & disease* 16: 631

Li K, Krone MW, Butrin A, Bond MJ, Linhares BM, Crews CM (2024) Development of Ligands and Degraders
Targeting MAGE-A3. *Journal of the American Chemical Society* 146: 24884-24891

Li Q, Huang Y, Liu X, Gan J, Chen H, Yang CG (2016) Rhein Inhibits AlkB Repair Enzymes and Sensitizes Cells to
Methylated DNA Damage. *J Biol Chem* 291: 11083-93

Li X, Zeng H, Zhang L, Zhang J, Guo Y, Leng J (2024) An integrated LC-MS/MS platform for noninvasive urinary
nucleus acid adductomics: A pilot study for tobacco exposure. *Journal of hazardous materials* 474: 134780

Li Y, Xie H, Liu S, Ruan Z, Wang B (2025) Resveratrol Alleviates Ischemia-Reperfusion-Induced Neuronal Damage
by Inhibiting NR3C2-Mediated TRIM28 Expression. *Journal of integrative neuroscience* 24: 31375

Lin JR, Zeman MK, Chen JY, Yee MC, Cimprich KA (2011) SHPRH and HLTF act in a damage-specific manner to
coordinate different forms of postreplication repair and prevent mutagenesis. *Molecular cell* 42: 237-49

Lin K, Zhou E, Shi T, Zhang S, Zhang J, Zheng Z, Pan Y, Gao W, Yu Y (2023) m6A eraser FTO impairs gemcitabine
resistance in pancreatic cancer through influencing NEDD4 mRNA stability by regulating the PTEN/PI3K/AKT
pathway. *J Exp Clin Cancer Res* 42: 217

Liu L, Xiao L, Liang X, Chen L, Cheng L, Zhang L, Wu X, Xu Q, Ma C (2017) TRIM28 knockdown increases
sensitivity to etoposide by upregulating E2F1 in non-small cell lung cancer. *Oncology reports* 37: 3597-3605

Liu W, Zheng D, Huang X, Wei Z, Wei Z, Guo W (2025) TRIM17 promotes the progression of osteosarcoma by
regulating PDK1 m6A modification-mediated AKT/mTOR pathway activation through ubiquitination of FTO.
Cell death & disease 16: 767

Lombard DB, Kohler WJ, Guo AH, Gendron C, Han M, Ding W, Lyu Y, Ching TT, Wang FY, Chakraborty TS et al.
(2020) High-throughput small molecule screening reveals Nrf2-dependent and -independent pathways of cellular
stress resistance. *Science advances* 6

Lundholm L, Hååg P, Zong D, Juntti T, Mörk B, Lewensohn R, Viktorsson K (2013) Resistance to DNA-damaging
treatment in non-small cell lung cancer tumor-initiating cells involves reduced DNA-PK/ATM activation and
diminished cell cycle arrest. *Cell death & disease* 4: e478

Malacrida A, Di Domizio A, Bentivegna A, Cislaghi G, Messuti E, Tabano SM, Giussani C, Zuliani V, Rivara M,
Nicolini G (2022) MV1035 Overcomes Temozolomide Resistance in Patient-Derived Glioblastoma Stem Cell
Lines. *Biology* 11

Martin LP, Hamilton TC, Schilder RJ (2008) Platinum resistance: the role of DNA repair pathways. *Clinical cancer*
research : an official journal of the American Association for Cancer Research 14: 1291-5

Meng S, Liu H, Xu J, Deng C, Qian X, Chu S, Zhu WG, Zhu J, Yong H, Li Z et al. (2025) PRMT5-Mediated ALKBH5
Methylation Promotes Colorectal Cancer Immune Evasion via Increasing CD276 Expression. *Research*
(Washington, DC) 8: 0549

Pineda CT, Potts PR (2015) Oncogenic MAGEA-TRIM28 ubiquitin ligase downregulates autophagy by ubiquitinating
and degrading AMPK in cancer. *Autophagy* 11: 844-6

Pineda CT, Ramanathan S, Fon Tacer K, Weon JL, Potts MB, Ou YH, White MA, Potts PR (2015) Degradation of
AMPK by a cancer-specific ubiquitin ligase. *Cell* 160: 715-728

Prince C, Hammerton G, Taylor AE, Anderson EL, Timpson NJ, Davey Smith G, Munafò MR, Relton CL, Richmond
RC (2019) Investigating the impact of cigarette smoking behaviours on DNA methylation patterns in adolescence.
Hum Mol Genet 28: 155-165

Randolph K, Hyder U, D'Orso I (2022) KAP1/TRIM28: Transcriptional Activator and/or Repressor of Viral and
Cellular Programs? *Frontiers in cellular and infection microbiology* 12: 834636

Ringvoll J, Nordstrand LM, Vågbø CB, Talstad V, Reite K, Aas PA, Lauritzen KH, Liabakk NB, Bjørk A, Doughty
RW et al. (2006) Repair deficient mice reveal mABH2 as the primary oxidative demethylase for repairing 1meA
and 3meC lesions in DNA. *The EMBO journal* 25: 2189-98

Saxena T, Quan A, Chan E, Kozlova N, Matai L, Lee JD, Rupaimoole R, Beca F, Muranen T, Slack FJ (2025) EGFR-
induced lncRNA TRIDENT promotes drug resistance in non-small cell lung cancer via phospho-TRIM28-
mediated DNA damage repair. *Proceedings of the National Academy of Sciences of the United States of America*
122: e2415389122

Soll JM, Sobol RW, Mosammaparast N (2017) Regulation of DNA Alkylation Damage Repair: Lessons and
Therapeutic Opportunities. *Trends in biochemical sciences* 42: 206-218

Song H, Feng X, Zhang H, Luo Y, Huang J, Lin M, Jin J, Ding X, Wu S, Huang H et al. (2019) METTL3 and ALKBH5
oppositely regulate m(6)A modification of TFEB mRNA, which dictates the fate of hypoxia/reoxygenation-treated
cardiomyocytes. *Autophagy* 15: 1419-1437

Song M, Yi F, Zeng F, Zheng L, Huang L, Sun X, Huang Q, Deng J, Wang H, Gu W (2024) USP18 Stabilized FTO
Protein to Activate Mitophagy in Ischemic Stroke Through Repressing m6A Modification of SIRT6. *Molecular*
neurobiology 61: 6658-6674

Tan Q, Ma J, Zhang H, Wu X, Li Q, Zuo X, Jiang Y, Liu H, Yan L (2022) miR-125b-5p upregulation by TRIM28
induces cisplatin resistance in non-small cell lung cancer through CREB1 inhibition. *BMC Pulm Med* 22: 469

Trewick SC, Henshaw TF, Hausinger RP, Lindahl T, Sedgwick B (2002) Oxidative demethylation by Escherichia coli
AlkB directly reverts DNA base damage. *Nature* 419: 174-8

Tsang YH, Wang Y, Kong K, Grzeskowiak C, Zagorodna O, Dogruluk T, Lu H, Villafane N, Bhavana VH, Moreno D
et al. (2020) Differential expression of MAGEA6 toggles autophagy to promote pancreatic cancer progression.
eLife 9

Tsao N, Brickner JR, Rodell R, Ganguly A, Wood M, Oyeniran C, Ahmad T, Sun H, Bacolla A, Zhang L et al. (2021)
Aberrant RNA methylation triggers recruitment of an alkylation repair complex. *Molecular cell* 81: 4228-4242.e8

Wada Y, Naito T, Fukushima T, Saito M (2024) Evaluation of ALKBH2 and ALKBH3 gene regulation in patients with
adult T-cell leukemia/lymphoma. *Virology journal* 21: 316

Wang C, Ivanov A, Chen L, Fredericks WJ, Seto E, Rauscher FJ, 3rd, Chen J (2005) MDM2 interaction with nuclear
corepressor KAP1 contributes to p53 inactivation. *The EMBO journal* 24: 3279-90

Wang K, Liu F, Muchu B, Deng J, Peng J, Xu Y, Li F, Ouyang M (2024) E3 ubiquitin ligase RNF180 mediates the

ALKBH5/SMARCA5 axis to promote colon inflammation and Th17/Treg imbalance in ulcerative colitis mice. *Archives of pharmacal research* 47: 645-658

Wang P, Wu J, Ma S, Zhang L, Yao J, Hoadley KA, Wilkerson MD, Perou CM, Guan KL, Ye D et al. (2015) Oncometabolite D-2-Hydroxyglutarate Inhibits ALKBH DNA Repair Enzymes and Sensitizes IDH Mutant Cells to Alkylating Agents. *Cell reports* 13: 2353-2361

Wang Z, Mierxiati A, Zhu W, Li T, Xu H, Wan F, Ye D (2025) FOXA1-dependent NSUN2 facilitates the advancement of prostate cancer by preserving TRIM28 mRNA stability in a m5C-dependent manner. *NPJ precision oncology* 9: 127

Wilson DL, Beharry AA, Srivastava A, O'Connor TR, Kool ET (2018) Fluorescence Probes for ALKBH2 Allow the Measurement of DNA Alkylation Repair and Drug Resistance Responses. *Angewandte Chemie (International ed in English)* 57: 12896-12900

Wu G, Yan Y, Cai Y, Peng B, Li J, Huang J, Xu Z, Zhou J (2021) ALKBH1-8 and FTO: Potential Therapeutic Targets and Prognostic Biomarkers in Lung Adenocarcinoma Pathogenesis. *Frontiers in cell and developmental biology* 9: 633927

Wu SS, Xu W, Liu S, Chen B, Wang XL, Wang Y, Liu SF, Wu JQ (2011) Down-regulation of ALKBH2 increases cisplatin sensitivity in H1299 lung cancer cells. *Acta pharmacologica Sinica* 32: 393-8

Xiao TZ, Suh Y, Longley BJ (2014) MAGE proteins regulate KRAB zinc finger transcription factors and KAP1 E3 ligase activity. *Archives of biochemistry and biophysics* 563: 136-44

Xu GL, Bochtler M (2020) Reversal of nucleobase methylation by dioxygenases. *Nature chemical biology* 16: 1160-1169

Xu L, Zhang K, Ma X, Li Y, Jin Y, Liang C, Wang Y, Duan W, Zhang H, Zhang Z et al. (2022) Boosting cisplatin chemotherapy by nanomotor-enhanced tumor penetration and DNA adducts formation. *Journal of nanobiotechnology* 20: 429

Yalçin Z, Lam SY, Peuscher MH, van der Torre J, Zhu S, Iyengar PV, Salas-Lloret D, de Krijger I, Moatti N, van der Lugt R et al. (2024) UBE2D3 facilitates NHEJ by orchestrating ATM signalling through multi-level control of RNF168. *Nature communications* 15: 5032

Yang CG, Yi C, Duguid EM, Sullivan CT, Jian X, Rice PA, He C (2008) Crystal structures of DNA/RNA repair enzymes AlkB and ABH2 bound to dsDNA. *Nature* 452: 961-5

Yi C, He C (2013) DNA repair by reversal of DNA damage. *Cold Spring Harbor perspectives in biology* 5: a012575

Zhang L, Liu M, Xie Y, Yuan BF, Peng Z, Xiong J, Zhang XL (2025) FTO O-GlcNAcylation promotes TRIM21-mediated FTO ubiquitination degradation to sustain the negative feedback control of macrophage inflammation. *Front Immunol* 16: 1593243

Zhao W, Zheng HZ, Zhou T, Hong XS, Cui HJ, Jiang ZW, Chen HJ, Zhou ZJ, Liu XG (2017) CTT1 overexpression increases the replicative lifespan of MMS-sensitive Saccharomyces cerevisiae deficient in KSP1. *Mechanisms of ageing and development* 164: 27-36

Zhao Y, Majid MC, Soll JM, Brickner JR, Dango S, Mosammaparast N (2015) Noncanonical regulation of alkylation damage resistance by the OTUD4 deubiquitinase. *The EMBO journal* 34: 1687-703

Zhou X, Xia Q, Wang B, Li J, Liu B, Wang S, Huang M, Zhong R, Cheng SY, Wang X et al. (2025) USP14 modulates stem-like properties, tumorigenicity, and radiotherapy resistance in glioblastoma stem cells through stabilization of MST4-phosphorylated ALKBH5. *Theranostics* 15: 2293-2314

 Figure legends

Figure 1. TRIM28 directly binds ALKBH2. (A) Silver staining and LC–MS/MS identification

of ALKBH2 as a TRIM28-associated protein following affinity purification in H1299 NSCLC

cells. The TRIM28 band is indicated. **(B)** Co-immunoprecipitation (co-IP) analysis of HEK293T

cells co-expressing HA-ALKBH2 and Flag-TRIM28 or Flag-TRIM24. Anti-Flag

immunoprecipitates were analyzed by western blotting (WB). **(C)** Reciprocal co-IP using HA-

ALKBH2 and Flag-TRIM28 (or HA-GFP control) with anti-HA immunoprecipitation. **(D)**

Schematic of full-length TRIM28 and truncation mutants. **(E)** Domain mapping using Flag-IP in

HEK293T cells co-transfected with HA-ALKBH2 and Flag-TRIM28 truncation mutants. **(F)**

Endogenous ALKBH2–TRIM28 interaction in H1299 cells detected by co-IP with anti-TRIM28

antibody. **(G)** Immunofluorescence (IF) showing nuclear co-localization of endogenous ALKBH2

(green) and TRIM28 (red). Nuclei were counterstained with DAPI (blue). Scale bar, 20 μm.

Pearson correlation coefficient (PCC) = 0.896 ± 0.016 (mean ± SD, n = 42). **(H)** *In vitro* MBP

pull-down assay using recombinant His-Flag-ALKBH2 and purified MBP-TRIM28. **(I)**

Specificity profiling of endogenous TRIM28 binding in HEK293T cells transfected with Flag-

ALKBH2 or Flag-ALKBH3. **(J)** GRAMM-X docking of TRIM28 RBCC (cyan) with ALKBH2

(light red) or ALKBH3 (green). Middle: Magnified interface showing ALKBH2 Finger and Long-

loop motifs (red). Right: Superposition with ALKBH2–DNA (orange), revealing overlap of

TRIM28- and DNA-binding surfaces. **(K)** Flag-IP analysis of lysates from Flag-ALKBH2-

expressing HEK293T cells in the presence of increasing ssDNA concentrations. Data are

representative of n ≥ 3 biological replicates.

Figure 2. TRIM28 promotes K48-linked ubiquitination and proteasomal degradation of

ALKBH2. (A) *In vivo* ubiquitination assay in HEK293T cells co-transfected with Flag-ALKBH2

and HA-ubiquitin (Ub), treated with MG132 (10 μM, 6 h). Lysates were subjected to denaturing

IP (1% SDS) with anti-Flag antibody. **(B)** Denaturing Flag-IP analysis of HEK293T cells co-

transfected with Flag-ALKBH2 and HA-tagged wild-type (WT), K48-only, or K48R ubiquitin

mutants. **(C)** K48-linked polyubiquitin enrichment using Halo-TUBE beads from HEK293T cells

expressing Flag-ALKBH2 treated with MG132. **(D)** Denaturing Flag-IP of HEK293T cells co-transfected with Flag-ALKBH2, His-TRIM28, and HA-K48-only ubiquitin. **(E)** Analysis of ALKBH2 ubiquitination in HEK293T cells co-transfected with Flag-ALKBH2 and WT or RING-inactive (C65/68A) His-TRIM28. **(F)** Denaturing Flag-IP detecting K48-linked ALKBH2 ubiquitination in TRIM28-knockdown HEK293T cells reconstituted with shRNA-resistant TRIM28 (WT). **(G)** WB analysis of H1299 and A549 cells transduced with control (shCon) or TRIM28-specific shRNAs. **(H)** qRT-PCR quantification of ALKBH2 mRNA in H1299 cells following TRIM28 depletion. **(I)** WB analysis of ALKBH2 levels in H1299 and HEK293T cells overexpressing increasing doses of TRIM28. **(J)** WB analysis of H1299 cells transduced with control or ALKBH2-specific shRNAs. **(K)** WB analysis of ALKBH2-depleted cells reconstituted with Flag-ALKBH2. **(L)** qRT-PCR of TRIM28 transcript levels in H1299 cells expressing empty vector or Flag-ALKBH2. Data are representative of n ≥ 3 biological replicates. Error bars indicate mean ± SD. *$p < 0.05$, **$p < 0.01$ (unpaired, two-tailed Student's t-test). NS, not significant.

Figure 3. MAGEA6 scaffolds the TRIM28–ALKBH2 E3 ligase complex and promotes ALKBH2 degradation. (A) Anti-Flag IP analysis of the interaction between Myc-MAGEA6 and Flag-ALKBH2 (or GFP control) in HEK293T cells. **(B)** *In vitro* pulldown assay using recombinant His-Flag-ALKBH2 and purified MBP-MAGEA6. **(C)** Anti-Flag IP in HEK293T cells (± TRIM28 knockdown) co-transfected with HA-ALKBH2 and Flag-MAGEA6. **(D)** Flag-IP analysis of the ALKBH2–TRIM28 interaction in HEK293T cells transfected with Myc-MAGEA6. **(E)** HA-IP in H1299 cells stably expressing HA-ALKBH2 following MAGEA3/6 knockdown. **(F)** K48-specific Halo-TUBE pulldown in TRIM28-knockdown HEK293T cells overexpressing Myc-MAGEA6. **(G)** WB analysis of ALKBH2 stability in control or TRIM28-knockout (sgTRIM28) H1299 cells transfected with MAGEA6 and treated with cycloheximide (CHX, 150 μg/mL). **(H)** Flag-IP mapping of interactions between Flag-MAGEA6 (WT or mutants), endogenous TRIM28, and co-expressed HA-ALKBH2 in HEK293T cells. **(I)** K48-specific Halo-TUBE pulldown of ubiquitinated proteins in HEK293T cells co-expressing HA-ALKBH2 and Flag-MAGEA6 (WT

or mutants). **(J)** Anti-Flag IP of HEK293T cells transfected with HA-ALKBH2 and cancer-associated Flag-MAGEA6 mutants. Data are representative of $n \geq 3$ biological replicates.

Figure 4. MMS dynamically regulates the TRIM28–ALKBH2 pathway. (A) Anti-Flag IP from H1299 cells expressing HA-ALKBH2 and Flag-TRIM28 treated with MMS (100 µM) and MG132 (10 µM) for the indicated times. **(B)** Anti-Flag IP from H1299 cells expressing Flag-TRIM28 treated with indicated MMS concentrations (3 h, +MG132). **(C)** WB analysis of MAGEA3/6 in nuclear/cytoplasmic fractions of H1299 cells treated with MMS (100 µM, 4 h). **(D)** Denaturing anti-Flag IP of H1299 cells expressing Flag-ALKBH2 treated with MMS (0–200 µM, 3 h) plus MG132. **(E)** Degradation assay in H1299 cells expressing TAP-ALKBH2 transduced with Flag-TRIM28 or vector, treated with MMS (100 µM). **(F)** Anti-Flag IP of cell lysates incubated with ssDNA (1 µM), H_2O_2 (200 µM), Vc (500 µM), or EDTA (5 mM). **(G)** Anti-Flag IP assessing the effect of EDTA (5 mM) on H_2O_2-induced interactions in ssDNA-containing lysates. **(H)** K48-specific Halo-TUBE pulldown in H1299 cells expressing TAP-ALKBH2 treated with MG132 (10 µM), MMS (100 µM), and Vc (500 µM) for 8 h. **(I)** WB of ALKBH2 in H1299 and H460 cells treated with MMS (100 µM). **(J)** Quantification of ALKBH2 protein levels from (I). **(K)** WB analysis of H1299 cells expressing HA-TRIM28 treated with MMS (100 µM). **(L)** qRT-PCR of ALKBH2 in H1299 cells treated with MMS (100 µM). Data are representative of $n \geq 3$ biological replicates. Error bars indicate mean ± SD. *$p < 0.05$, **$p < 0.01$ (unpaired, two-tailed Student's t-test).

Figure 5. MMS induces K48-linked ubiquitination and proteasomal degradation of TRIM28. **(A, B)** WB analysis of TRIM28 levels in H1299 cells treated with MMS in a time- (A) and dose-dependent (B) manner. **(C)** qRT-PCR of TRIM28 mRNA in cells treated with MMS (100 µM). **(D)** Denaturing anti-Flag IP of HEK293T cells expressing Flag-TRIM28 and HA-Ub treated with MMS and MG132. **(E)** Anti-Flag IP and K48-specific Ub WB in H1299 cells expressing Flag-TRIM28 treated with MMS (100 µM) + MG132. **(F)** Denaturing anti-Flag IP in HEK293T cells

transfected with Flag-TRIM28 and HA-Ub (K48-only) treated with MMS (100 μM). **(G)** K48-specific Halo-TUBE pulldown of endogenous TRIM28 in H1299 cells treated with MG132 and increasing MMS. **(H)** Halo-TUBE pulldown comparing effects of MMS (200 μM), temozolomide (TMZ, 200 μM), cisplatin (CDDP, 20 μM), or H_2O_2 (300 μM) on TRIM28 ubiquitination (+MG132, 6 h). **(I)** Halo-TUBE pulldown in HEK293T cells expressing Flag-TRIM28 WT or SUMO-deficient mutant (K554/779/804R) treated with MMS. **(J)** Anti-Flag co-IP in H1299 cells expressing Flag-TRIM28 treated with MMS (100 μM) + MG132. **(K)** Anti-Flag co-IP in HEK293T cells expressing Flag-TRIM28 and Myc-MAGEA6 treated with MMS. **(L)** Anti-Flag co-IP and WB in HEK293T cells expressing Flag-TRIM24 treated with MMS + MG132. (M) Halo-TUBE enrichment of K48-linked proteins in HEK293T cells overexpressing Flag-TRIM24 treated with MMS. Data are representative of n ≥ 3 biological replicates. Error bars indicate mean ± SD. NS, not significant.

Figure 6. The TRIM28–ALKBH2 axis is clinically relevant and regulates sensitivity to alkylation damage. (A) Relative mRNA expression of ALKBH2, ALKBH3, and MGMT in LUAD vs. normal lung (TCGA dataset; Mann–Whitney test). **(B)** Representative ALKBH2 immunohistochemistry (IHC) staining in LUAD and matched normal lung (Human Protein Atlas). **(C)** WB analysis of ALKBH2, TRIM28, ALKBH3, and MGMT in NSCLC cell lines and BEAS-2B cells. **(D, E)** Kaplan–Meier overall survival for LUAD patients stratified by ALKBH2 (D) or ALKBH3 (E) mRNA expression. **(F)** CCK-8 proliferation assay in H1299 cells following ALKBH2 knockdown. **(G)** CCK-8 viability assay in H1299 cells (ALKBH2 knockdown) 48 h after a 1 h MMS pulse. **(H)** CCK-8 viability assay in H1299 cells overexpressing Flag-TRIM28 WT or RING-inactive mutant (C65/68A) after MMS pulse. **(I)** MMS sensitivity assessment in H1299 cells with TRIM28 knockdown and rescue. **(J)** Rescue of MMS hypersensitivity in H1299 cells by ALKBH2 co-expression. **(K)** Effect of MAGEA6 overexpression on MMS cytotoxicity. **(L)** Effect of TRIM28-binding-deficient MAGEA6 mutant (D156E/S157Y) on MMS sensitivity. Data are representative of n ≥ 3 biological replicates. Error bars indicate mean ± SD. *$p < 0.05$,

 **p < 0.01 (unpaired, two-tailed Student's t-test).

 Figure 7. Combining MMS with CDDP synergistically enhances cancer cell killing. (A) CCK-
8 viability assay of H1299 cells (TRIM28 knockdown) treated with continuous low-dose MMS
(48 h). **(B, C)** Colony formation assay of H460 cells overexpressing TRIM28 treated with MMS
(100 µM, 48 h). Representative images (B) and quantification (C). **(D)** CCK-8 assay of H1299
cells (ALKBH2 knockdown rescued with Flag-ALKBH2) treated with MMS ± 2 µM CDDP. **(E)**
Alkaline comet assay of H1299 cells treated with MMS (50 µM) and/or CDDP (2 µM). Tail
moments quantified (n = 70). **(F, G)** IC_{50} analysis of CDDP in H1299 (F) and H460 (G) cells with
MMS co-treatment. **(H, I)** Clonogenic survival of H460 cells treated with MMS (75 µM) and
CDDP (1 µM). **(J, K)** 3D tumor spheroid growth of H1299 cells treated with MMS (100 µM) and
CDDP (2 µM). Calcein-AM (live cells)/PI (dead cells) staining (J) and volume quantification (K).
Scale bar: 200 µm. Data are representative of n ≥ 3 biological replicates. Error bars indicate mean
± SD. *$p < 0.05$, **$p < 0.01$ (unpaired, two-tailed Student's t-test).

**Figure 8. Model of the biphasic TRIM28–ALKBH2 regulatory switch induced by alkylation
stress. (I) Basal state.** MAGEA6-assisted, TRIM28-mediated K48-linked ubiquitination
maintains low ALKBH2 levels; ALKBH2 supports TRIM28 transcription. **(II) Acute MMS:**
Enhanced assembly of the TRIM28–MAGEA6–ALKBH2 complex accelerates ALKBH2
degradation. **(III) Prolonged MMS:** Ubiquitin flux shifts toward TRIM28, leading to its
degradation and releasing ubiquitin-dependent repression of ALKBH2, together with MMS-
induced transcriptional activation that drives progressive ALKBH2 accumulation.

Expanded view figure legends

Figure EV1. MAGEA6 supports TRIM28-dependent ubiquitination of ALKBH2 (Related to Fig. 3). (A) HA-IP in HEK293T cells transfected with HA-ALKBH2, Flag-TRIM28, and increasing Myc-MAGEA6. **(B)** Anti-Flag IP in HEK293T cells co-transfected with HA-ALKBH2, Flag-TRIM28, and Myc-MAGEA6 or Myc-MAGEC2. **(C)** Halo-TUBE enrichment of K48-ubiquitinated proteins in H1299 cells (MAGEA6 knockdown) reconstituted with Flag-MAGEA6. **(D)** Sequence alignment of MAGEA6 WT and MutA/MutB variants. **(E)** WB analysis of ALKBH2 stability in H1299 cells expressing WT or mutant MAGEA6 treated with CHX. **(F)** AlphaFold3 structural model of the MAGEA6–TRIM28 complex. **(G)** ALKBH2 degradation assay in H1299 cells expressing WT or TRIM28-binding-deficient (D156E/S157Y) MAGEA6 treated with CHX. **(H)** Anti-Flag IP of HEK293T cells transfected with WT or mutant Flag-MAGEA6. Data are representative of n ≥ 3 biological replicates.

Figure EV2. Redox-modulated regulation of the TRIM28–ALKBH2 pathway under MMS exposure (Related to Fig. 4). (A) IF staining of Flag-ALKBH2 (red) and TRIM28 (green) in H1299 cells treated with MMS (100 µM, 8 h). Nuclei stained with DAPI. Scale bar = 20 µm. **(B)** WB analysis of ALKBH2 turnover in H1299 cells treated with CHX, MMS, or both. **(C)** Quantification of ALKBH2 levels from (B). **(D)** Anti-Flag IP of lysates in the presence of GSSG (1 mM), GSH (1 mM), or H_2O_2 (0.5 mM). **(E)** Anti-Flag IP in H1299 cells expressing Flag-ALKBH2 treated with H_2O_2 + MG132. **(F)** Anti-Flag IP of WT or mutant ALKBH2 (H171A/D173A) in the presence of ssDNA or H_2O_2. Data are representative of n ≥ 3 biological replicates. Error bars indicate mean ± SD. *$p < 0.05$, **$p < 0.01$.

Figure EV3. Extended analyses of MMS-induced ubiquitination and degradation of TRIM28 (Related to Fig. 5). (A) WB of H1299 cells treated with MMS (100 µM) for indicated durations. **(B)** WB monitoring ALKBH2 turnover (CHX chase) in H1299 cells pretreated with MMS (100 µM, 12 h). **(C)** Quantification of ALKBH2 levels from (B). **(D)** WB of H1299 cells treated with

TMZ (100 μM). **(E, F)** WB analysis of H1299 cells treated with CDDP (E) or H_2O_2 (F). **(G)** WB analysis of anti-Flag immunoprecipitates and whole-cell lysates from H1299 cells expressing WT or mutant Flag-TRIM28 treated with MMS. **(H)** WB of TRIM28 levels in HEK293T cells overexpressing Flag-TRIM24 treated with MMS. **(I)** Quantification of TRIM28 levels from (H). Data are representative of n ≥ 3 biological replicates. Error bars indicate mean ± SD. *$p < 0.05$, **$p < 0.01$.

Figure EV4. Clinical and functional analyses supporting the TRIM28–ALKBH2 axis in NSCLC (Related to Fig. 6). (A) CPTAC analysis (UALCAN) of ALKBH2, ALKBH3, and MGMT protein expression in LUAD vs. normal lung. **(B)** ALKBH2 mRNA expression in LUAD patients by smoking history (UALCAN/TCGA). **(C)** Frequency of MAGEA6 and MAGEA3 mutations in TCGA lung cancer samples stratified by smoking status (Fisher's exact test). **(D)** WB of TRIM28 in H1299 cells after short-term high-dose MMS (500 μM, 1 h) and release. **(E)** CCK-8 proliferation assay of H1299 cells overexpressing TRIM28. **(F)** CCK-8 survival assay in MAGEA6-knockdown H1299 cells with MAGEA6 re-expression. **(G, H)** CCK-8 viability assay of H1299 cells expressing Flag-MAGEA6 treated with CDDP (G) or bortezomib (BTZ) (H). Data are representative of n ≥ 3 biological replicates. Error bars indicate mean ± SD. *$p < 0.05$, **$p < 0.01$.

Figure EV5. Additional analyses supporting the combined effects of MMS and CDDP on cytotoxicity (Related to Fig. 7). (A) Summary of therapeutic agents for lung cancer patients (TCGA–GDC database). **(B, C)** Kaplan–Meier survival analysis of patients who received chemotherapy (B) versus those who did not receive chemotherapy (C), stratified by high vs. low TRIM28 mRNA levels. **(D)** CCK-8 viability assay of H1299 cells treated with low-dose MMS ± H_2O_2. **(E)** WB of γ-H2AX in H1299 cells treated with CDDP ± MMS. **(F)** IC_{50} determination of MMS in H1299 and H460 cells (CCK-8). Data are representative of n ≥ 3 biological replicates. Error bars indicate mean ± SD. *$p < 0.05$, **$p < 0.01$.