Novel markers for haemogenic endothelium and haematopoietic progenitors in the mouse yolk sac

Guillermo Diez-Pinel[1], Alessandro Muratore[1,2], Christiana Ruhrberg[1]*, Giovanni Canu[1]*

[1] UCL Institute of Ophthalmology, University College London, United Kingdom

[2] School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy

* Corresponding authors:

Christiana Ruhrberg: c.ruhrberg@ucl.ac.uk

Giovanni Canu: g.canu@ucl.ac.uk

KEYWORDS:

Haemogenic endothelium, erythro-myeloid progenitors, EMPs, haematopoietic development, haematopoietic differentiation, yolk sac, mouse embryo, single-cell transcriptomic, *Fxyd5*/Dysadherin, *Scarf1*, *Fcer1g*

ABSTRACT

Erythro-myeloid progenitors (EMPs) originate from the haemogenic endothelium in the yolk sac via an endothelial-to-haematopoietic transition (EHT) to generate blood and immune cells that support embryo development. Yet, the transitory nature of EHT and the limited availability of molecular markers has constrained our understanding of the origin, identity and differentiation dynamics of EMPs. Here, we have refined the annotation of yolk sac haemato-vascular populations in publicly available single-cell RNA sequencing (scRNAseq) datasets from mouse embryos to identify novel molecular markers of haemogenic endothelium and EMPs. By sub-clustering key cell populations followed by pseudotime analysis, we refined cluster annotations and then reconstructed differentiation trajectories. Subsequent differential gene expression analysis between clusters identified novel cell surface markers for haemogenic endothelial cells (*Fxyd5* and *Scarf1*) and EMPs (*Fcer1g*, *Tyrobp* and *Mctp1*). Further, we have identified candidate signalling and metabolic pathways that may regulate yolk sac haematopoietic emergence and differentiation. The specificity of FXYD5, SCARF1 and FCER1G for haemogenic endothelium and EMPs was validated by immunostaining of mouse yolk sac. These insights into the transcriptional dynamics in the yolk sac should support future investigation of EHT and haematopoietic differentiation during early mammalian development.

INTRODUCTION

Haematopoiesis in the developing embryo proceeds in three temporally and spatially overlapping waves in close proximity to vascular endothelial cells (Ivanovs *et al.*, 2017; Canu and Ruhrberg, 2021). The first wave, known as primitive haematopoiesis, arises in the yolk sac, when extra-embryonic mesoderm differentiates into vascular and haematopoietic progenitors that progressively organise into blood islands, and produces nucleated erythrocytes, megakaryocytes and macrophages (Ferkowicz and Yoder, 2005). This process occurs in the mouse from embryonic day (E) 7.0 onwards, equivalent to 2-3 post-conception weeks in human.

The second haematopoietic wave, known as pro-definitive or transient definitive haematopoiesis, also emerges in the yolk sac, when a change in cellular identity causes haematopoietic progenitors to bud off the endothelium as clusters of round cells, a process called endothelial-to-haematopoietic transition (EHT) (Frame *et al.*, 2016). In the mouse yolk sac, EHT generates mainly erythro-myeloid progenitors (EMPs) from E8.5, alongside lympho-myeloid progenitors (LMPs) from E9.5 (Böiers *et al.*, 2013; Frame, McGrath and Palis, 2013). Once the blood circulation is established by E9.5, these progenitors are transported out of the yolk sac and seed the foetal liver, from where they contribute blood and immune cells from mid-gestation until birth (Hoeffel *et al.*, 2015). Additionally, we recently showed that haematopoietic progenitors akin to EMPs can also arise from a *Pax3*-expressing lineage in the mouse and transiently seed the foetal liver by E12.5 to contribute erythroid and myeloid cells to foetal haematopoiesis (Canu *et al.*, 2024). Based on their origin from a *Pax3*-expressing progenitor, these haematopoietic cells may arise via EHT from a paraxial mesoderm-derived haemogenic endothelium, although further work is needed to address this possibility. Among the early haematopoietic populations, yolk sac-derived EMPs generate blood and immune cells important for embryogenesis (Frame, McGrath and Palis, 2013; Hoeffel *et al.*, 2015) and self-renewing tissue-resident macrophages that persist into adulthood (Moskalik, Niderla-Bielińska and Ratajska, 2021; Jiang *et al.*, 2024; Liu *et al.*, 2024).

The third haematopoietic wave, known as definitive haematopoiesis, originates intra-embryonically in the dorsal aorta, where EHT produces haematopoietic stem cells (HSCs) that enter the circulation to first colonise the foetal liver by mid-gestation and later seed the bone marrow to establish long-term, self-renewing adult haematopoiesis (Tavian *et al.*, 1996; Oberlin *et al.*, 2010; Ivanovs *et al.*, 2011; Canu and Ruhrberg, 2021; Sánchez-Lanzas, Jiménez-Pompa and Ganuza, 2024; Sugimura *et al.*, 2025). HSCs are first produced at E10.5 in the mouse, and 4-5 post-conception weeks in human.

Despite their key roles in embryonic haematopoiesis and the generation of tissue
macrophages with a role in adult homeostasis, the transitory nature of EMPs and their
overlapping markers with HSC lineages have made the study of EMP-derived cell lineages
challenging. For example, the Lin⁻ KIT⁺ CD41^{low} CD16/32⁺ cell surface profile allows EMP
isolation by flow cytometry from the early embryo, but is insufficient to unambiguously
distinguish EMPs from HSC-derived myeloid progenitors at mid-gestation (McGrath *et al.*,
2015; Van Deren *et al.*, 2022). Temporally-controlled lineage tracing via Cre-mediated
reporter gene activation has therefore been used as a complementary approach to
distinguish EMPs from HSC-derived progenitors based on the earlier production of EMPs
(Gomez Perdiguero *et al.*, 2014; Hoeffel *et al.*, 2015; McGrath, Frame and Palis, 2016).
However, this approach relies on transgenic mice expressing suitable Cre drivers and
expression reporters. Therefore, specific molecular markers are needed to allow accurate
EMP isolation and to study their formation and behaviour in the mouse embryo. Moreover,
such markers might facilitate the identification of analogous markers suitable to study foetal
haematopoiesis in human embryos, in which genetic lineage tracing is not an option.

Here, we have leveraged publicly available mouse embryonic single-cell RNA
sequencing (scRNAseq) data to investigate cell populations in the developing mouse yolk
sac to identify novel candidate markers for haemogenic endothelium and EMPs.

METHODS
Integration of publicly available E8.5 scRNAseq mouse embryo data
The file 'embryo_sce.rds' containing a processed SingleCellExperiment object including
scRNAseq data from the ArrayExpress dataset E-MTAB-6967 (Pijuan-Sala *et al.*, 2019) and
ArrayExpress dataset E-MTAB-11763 (Goh *et al.*, 2023) was downloaded from
https://marionilab.github.io/ExtendedMouseAtlas/. The cells annotated as embryonic day
E8.5 were subset for analysis in R v4.5.1, with Rstudio v2025.05.1 using the Seurat v4.5.1
R package (Satija *et al.*, 2015). The function 'as.Seurat' was used to generate a Seurat
object from the SingleCellExperiment object. For sample integration, the E-MTAB-11763
cells were subset and split into layers containing individual samples, counts were log-
normalised with the 'NormalizeData' function, the top-2000 variable features were identified
with the 'FindVariableFeatures' and scaled and centred with the 'ScaleData' function. Then,
PCA dimensionality reduction was performed with the 'RunPCA' function and Harmony
Integration (Korsunsky *et al.*, 2019) was performed using the 'IntegrateLayers' function. A
Uniform Manifold Approximation and Projection (UMAP) reduction was obtained with the
'RunUMAP' function, for which the number of principal components was selected by

examination of an elbow plot. Lastly, the layers were joined with the 'JoinLayers' function.

This same integration pipeline was used for all other data integration in the study.

Re-annotation of yolk sac cell types in E8.5 scRNAseq data

Cells with anatomy 'Yolk sac' (YS) from haemato-vascular cell types were subset and

integrated by sample. For clustering, a k-nearest neighbour (KNN) graph was calculated (k

= 20), and the Louvain algorithm was used for graph-clustering. Then, the expression of

known cell type markers in the yolk sac clusters was used alongside the anatomical origin

(yolk sac or embryo) to annotate these haemato-vascular yolk sac cells as well as the

embryo haemato-vascular cells more accurately. Afterwards, both yolk sac and embryo cells

were merged and integrated by sample. These data were then used as the reference dataset

for reference-based annotation of the E-MTAB-6967 haemato-vascular cells using the

SingleR v2.10.0 R package (Aran *et al.*, 2019), and then merged and integrated by sample

with the E-MTAB-11763 data.

Analysis of gene expression along the EHT differentiation trajectory

To identify haemogenic endothelium cells, we sub-clustered all cells annotated as

endothelial cells and EMPs in the combined dataset, including Allantois endothelium,

Venous endothelium, Embryo proper endothelium, Endocardium, YS endothelium, Vitellin

vein endothelium, EMPs and EMP-like embryo, and then examined the expression of

haemato-vascular markers.To analyse the EHT differentiation trajectory, we sub-clustered

yolk sac haemato-vascular cells, including YS endothelium, haemogenic endothelium,

EMPs, BP1, BP2 and Erythroid, and the resulting UMAP reduction was used to carry out a

pseudotime analysis to predict differentiation trajectories with the Monocle3 v1.4.26 R

package (Cao *et al.*, 2019). Genes that changed as a function of pseudotime were identified

with the 'graph_test' function and grouped according to their expression patterns into

modules with the 'find_gene_modules', which were then grouped manually into four

supermodules (downwards trend, upwards trend, peaking at EMPs and peaking at BP1).

Gene ontology and differential gene expression of haemato-vascular clusters

Enrichment analyses against the GeneOntology: Biological Pathway and WikiPathways

databases were carried out using the 'enrichGO' and 'enrichWP' functions from the

clusterProfiler v4.16.0 R package (Wu *et al.*, 2021), respectively. All genes with at least 1

count were used as the background universe. Cytoscape 3.10.3 (Shannon *et al.*, 2003) and

the RCy3 v2.28.1 R package (Gustavsen *et al.*, 2019) were used to plot the TYROBP

WikiPathway pathway. For surface marker identification, differential expression analyses were carried out with the 'FindMarkers' function, with an adjusted P-value threshold of 0.05. Then, Uniprot accession IDs were retrieved from the org.Mm.eg.db v3.21.0 R package (Carlson, 2025) using the 'select' function from the AnnotationDbi v1.70.0 R package (Pagès *et al.*, 2025). A list of curated transmembrane proteins was obtained from Uniprot (https://www.uniprot.org/) on the 30th of July 2025 using the query '(keyword:KW-0812) AND (reviewed:true)', and filtering for 'Mouse' as the relevant organism. This list was then used to filter the differential expression results for transmembrane proteins. Plotting of UMAP reductions was carried out with Seurat's 'DimPlot' and the function 'DimPlot_scCustom' from the scCustomize v3.2.0 package (Marsh *et al.*, 2025). Plotting of normalised gene expression onto UMAP reductions was carried out with Seurat's 'FeaturePlot' and scCustomize's 'FeaturePlot_scCustom' functions. Other plots were generated using the ggplot2 v4.0.0 R package (Wickham *et al.*, 2025).

Animal procedures and tissue staining

Animal procedures were performed with Animal Welfare Ethical Review Body (AWERB) and UK Home Office approval. C57BL/6J mice were timed-mated for embryo collection at E8.5 after cervical dislocation culling of the pregnant dam. Yolk sacs were fixed in 4% formaldehyde, washed in phosphate-buffered saline (PBS) and then blocked in PBS containing 2% serum-free protein block (DAKO), 2% bovine serum albumin and 0.4% Triton X-100 before staining with the following antibodies: goat anti-mouse KIT (R&D Systems #AF1356-SP, 1:50), rat anti-mouse PECAM1 (BD Pharmingen #553370, 1:50), rabbit anti-mouse FXYD5/Dysadherin (Proteintech #12166-1-AP, 1:50), rabbit anti-mouse SCARF1 (Proteintech #13702-1-AP, 1:100) or rabbit anti-mouse FCER1G (GeneTex #GTX108487s, 1:50), followed by the appropriate secondary antibodies, which were Alexa Fluor 488-conjugated donkey anti-goat Fab fragments (Stratech #705-547-003, 1:200), Cy3-conjugated donkey anti-rat Fab fragments (Stratech #712-166-150, 1:200), and Alexa Fluor 647-conjugated donkey anti-rabbit Fab fragments (Stratech #711-607-003, 1:200). Secondary-only controls were included but are not shown. DAPI-counterstained sections were imaged on a LSM710 confocal microscope (Zeiss).

RESULTS

Refined annotation of E8.5 yolk sac and embryo cell clusters

To identify the transcriptomic signature of yolk sac haemogenic endothelium and that

of its haemopoietic progeny, we re-analysed published scRNAseq data from E8.5 mouse

embryos (**Fig. 1A**), which were extracted from the ArrayExpress datasets E-MTAB-

6967 (Pijuan-Sala *et al.*, 2019) and E-MTAB-11763 (Goh *et al.*, 2023). E-MTAB-

6967 contains data from E6.5 to E8.5 mouse embryos, including yolk sac and embryo

proper, which were sequenced and analysed together to construct a molecular atlas

describing early mouse gastrulation (Pijuan-Sala *et al.*, 2019). E-MTAB-11763 instead

contains data that were generated by collecting yolk sac and embryo proper from E8.5 to

E9.5 mouse embryos and sequencing them separately. These two published datasets

overall comprise 400,000 cells from E6.5 to E9.5. A subsequent *in silico* merge of E-MTAB-

11763 data and E-MTAB-6967 embryo and yolk sac data was previously analysed to

compare the mouse versus human yolk sac contribution to metabolic and nutritional support

as well as to early erythroid and myeloid lineages (Goh *et al.*, 2023). As the goal of our

analysis was to define the transcriptome of yolk sac endothelium and its haematopoietic

progeny at the time when EMPs emerge from yolk sac haemogenic endothelium, we subset

only the E8.5 data from E-MTAB-6967 and E-MTAB-11763 (**Fig. 1A**).

First, we analysed E8.5 data from E-MTAB-11763, which contained yolk sac and

embryo proper data annotated separately, and generated a UMAP including 75 clusters,

using the cluster annotations provided by the authors of the original publication (**Fig. S1**).

Next, we selectively sub-clustered the 8,453 haemato-vascular cells that had been

sequenced from the yolk sac (**Fig. 1B,C**). To confirm the identity of the resulting clusters,

we analysed the expression of known markers of endothelial and haematopoietic progenitor

populations (**Fig. 2A-D**). The yolk sac endothelial cell (EC) cluster was identified via

expression of the pan-EC markers *Kdr* and *Cdh5* together with *Lyve1* and *Stab2* (**Fig. 2A**)

as two markers typical for yolk sac ECs (Gordon, Gale and Harvey, 2008; Shvartsman *et*

al., 2019). The EMP cluster was identified via detection of *Csf1r* and *Ptprc* as well as the

myeloid markers *Spi1 and Mpo* (**Fig. 2B**). Notably, *Kdr*, *Cdh5* and *Lyve1* were also detected

in the EMP cluster (**Fig. 2A**), agreeing with their known endothelial origin (Gomez

Perdiguero *et al.*, 2014). Clusters expressing haematopoietic progenitor markers (*Vav1*,

Myb, *Gata2*) (**Fig. 2C**) together with erythroid (*Gata1*, *Klf1*) or myeloid (*Spi1, Mpo*)

differentiation markers (**Fig. 2D**) represented intermediate progenitors at different states of

commitment and were annotated as blood progenitors (BP) 1 and BP2 (**Fig. 1C**). Finally,

high expression levels of differentiation markers (*Alas2*, *Gypa*) (**Fig. 2D**), in the absence of

endothelial (*Cdh5*, *Lyve1*), EMP (*Csf1r*, *Ptprc*) or progenitor markers (*Spi1*, *Gata2*),

identified a cluster of differentiated erythroid cells.

As expected, the detection levels of endothelial and haematopoietic genes displayed

opposite trends between the endothelial and haematopoietic clusters. Thus, *Kdr* and *Cdh5*

appeared highly expressed in ECs and EMPs but their detection declined in BP1, whilst they

were barely detectable in BP2 and absent from erythroid cells (**Fig. 2A**). Conversely,

erythroid genes such as *Klf1* and *Gypa* were not detected in ECs and EMPs, detected at

low levels in BP1, at higher levels in BP2 and at very high levels in erythroid cells (**Fig. 2D**).

Known haematopoietic progenitor markers (*e.g.*, *Myb*, *Gata2, Runx1*) displayed a gradual

decrease from EMPs to BP1 and BP2, supporting the notion that the three clusters

represented successive maturation stages (**Fig. 2C**). Myeloid markers (*e.g.*, *Csf1r, Spi1,*

Mpo) did not identify a separate cluster but were present in the EMP cluster and, at lower

levels, also in BP1 (**Fig. 2B**), agreeing with prior studies suggesting that macrophages are

only present in the yolk sac from E9.0 onwards (Hoeffel and Ginhoux, 2018).

Unexpectedly, we observed that E-MTAB-11763 contained cells that had been

isolated and sequenced from the embryo proper but nevertheless clustered with yolk sac

EMPs and BPs; accordingly, we annotated these cells as EMP-like embryo and blood

progenitors embryo, respectively (**Fig. 3A,B**). Also unexpectedly, some cells sequenced

from the embryo proper had been annotated in the original publication describing E-MTAB-

11763 as representing yolk sac-derived ECs. As the endothelium in the yolk sac vasculature

forms a continuum with the endothelium of the vitelline vein, cells within the embryo proper

that resemble yolk sac ECs likely represent vitelline vein ECs, and we re-annotated them

accordingly (**Fig. 3A,B**).

Next, we integrated E-MTAB-11763 with E-MTAB-6967 to increase the number of

cells available for downstream analysis in the haemato-vascular clusters to identify yolk sac

haemogenic endothelium. As cells derived from the yolk sac and embryo proper had not

been separately sequenced to generate E-MTAB-6967, we used a validated, reference-

based annotation method (Aran *et al.*, 2019) to harmonise cell type annotations and

distinguish cells from these sources before integrating the two datasets (**Fig. 3A**).

Specifically, this method uses gene expression information to assign the most likely cluster

from a reference dataset to a query dataset, which in our case involved applying our revised

annotation for E8.5 data in E-MTAB-11763 to the E8.5 data in E-MTAB-6967. This approach

increased the overall number of haemato-vascular cells from 8,453 to 12,351 for further

analysis (**Fig. 3B**). We also surveyed the expression of key known marker genes of the cell

populations of interest to confirm the appropriate annotation of the integrated dataset (**Fig.**

3C-F).

Identifying yolk sac haemogenic ECs via their transcriptomic signature

The haemogenic endothelium is a rare and transient population, and thus difficult to

study (Gritz and Hirschi, 2016; Scarfò *et al.*, 2024). To identify haemogenic ECs within the

yolk sac endothelial cluster, we subset EMPs and ECs from both yolk sac and embryo proper

and re-clustered them to generate a new UMAP including only these cells, which was

annotated according to our refined nomenclature. Using this approach, yolk sac ECs

clustered in an intermediate position between embryo-derived ECs and EMPs and also split

into two sub-clusters, one positioned closer to EMPs and the other closer to endocardial

ECs (**Fig. 4A**). The analysis of known marker genes showed that both sub-clusters

expressed markers of yolk sac endothelium at similar levels (*Lyve1*, *Stab2*), whereas only

the EMP-proximal sub-cluster co-expressed haematopoietic progenitor genes (*Runx1*, *Myb*,

Spi1) (**Fig. 4B**). Together, these findings indicate that a subset of yolk sac ECs have a dual

endothelial and haematopoietic cell identity, consistent with a presumed identify of

haemogenic ECs that give rise to EMPs.

Pseudotime analysis identifies transcriptomic changes accompanying EHT

Having identified a cell cluster with haemogenic endothelial identity, we next used our

refined annotation of the yolk sac populations to study signalling pathways and markers

defining the transition from haemogenic ECs into EMPs and BPs. When analysing samples

that contain cells along a continuous differentiation process, single-cell transcriptomic data

can be used to arrange cells along a pseudotime trajectory (Cannoodt, Saelens and Saeys,

2016). In essence, pseudotime analysis uses gene expression patterns to reorder cells and

arrange them based on their presumed progressive gene expression differences, thus

reconstructing trajectories along which cells may be differentiating. Thus, we used

pseudotime analysis of yolk sac cell populations to study the transition of ECs into

haematopoietic populations. In agreement with prior knowledge of EHT, pseudotime

analysis of our integrated E8.5 dataset suggested a continuum of cell differentiation from

endothelial to haemogenic endothelium and then to haematopoietic cells (**Fig. 5A**), and also

highlighted progressive erythroid maturation (**Fig. 5B**). A refined pseudotime analysis after

exclusion of erythroid cells as the predominant cell type corroborated the expected progressive transition of yolk sac ECs to haemogenic ECs, EMPs, BP1 and BP2 (**Fig. 5C**).

Next, we took advantage of a method that analyses genes that change as a function of pseudotime (Sharon *et al.*, 2019) to identify genes that showed an upward or downward trend during EHT (**Fig. 5D**, **Table S1**). We found that 2,924 genes were progressively downregulated, whereas 1,776 genes were progressively upregulated, during the predicted transition from endothelial to haematopoietic cells. We also found that 231 genes were upregulated during the predicted transition from ECs to EMPs, but downregulated during subsequent differentiation towards BP1 and then BP2 (peak transcript levels in EMPs). Moreover, 848 genes were progressively upregulated from ECs to BP1, but downregulated in BP2 (peak transcript levels in BP1) (**Fig. 5D**).

To identify potential biological processes underpinning EHT, we used gene ontology (GO) analysis on these 4 groups of co-regulated genes (**Fig. 5E-H**, **Table S2**). The GO analysis of genes with a downward expression trend during the predicted progression from endothelial to haematopoietic cells revealed enrichment in biological processes related to angiogenesis, cytoskeleton re-organisation, cell adhesion, endothelial cell migration and endothelial development (**Fig. 5E**). Genes with an upward expression trend instead were involved in mRNA processing, splicing and ribonucleoprotein biogenesis (**Fig. 5F**). Genes that peaked in EMPs showed enrichment in terms involved in leukocyte migration, myeloid proliferation and immune response (**Fig. 5G**), in agreement with EMPs being responsible for the production of foetal blood and immune cells. Finally, genes that peaked in BP1 are involved in oxidative phosphorylation, cellular respiration and ATP synthesis (**Fig. 5H**), raising the possibility that these cells have increased metabolism.

To identify genes belonging to known signalling pathways amongst each one of the 4 groups of co-regulated genes, we performed pathway analysis using the WikiPathways resource (Agrawal *et al.*, 2024) (**Table S3**). Genes that were downregulated when ECs were predicted to transition to haematopoietic cells belonged to the cholesterol biosynthesis, IL3, IL17A, PI3K/AKT/mTOR and EFGR1 signalling pathways. Genes that were upregulated during this transition are involved in RNA processing and cell cycle activation. Genes peaking in BP1 are involved in the regulation of cell metabolism, whereas genes peaking in EMPs belonged to the TYROBP and IL5 signalling pathways (**Table S3**). Notably, by comparing EMPs to other yolk sac clusters, we identified multiple genes within the TYROBP signalling pathway that, in addition to peaking in EMPs, were significantly enriched in EMPs compared to the other cells in the haemato-vascular cluster (**Fig. S2**).

Overall, this analysis provided insights into molecular changes predicted to
accompany EHT and haematopoietic differentiation in the yolk sac, and identified candidate
signalling pathways that may be studied further to understand the molecular control of these
complex cellular differentiation pathways.

Differential expression analysis identifies EMP and haemogenic endothelium markers

To identify markers for haemogenic ECs and EMPs, we used our integrated E8.5
mouse dataset to perform differential expression (DE) analysis of endothelial and
haematopoietic cell populations in both the yolk sac and embryo. First, we subset the
clusters for both cell populations and then performed DE analyses by comparing clusters in
different combinations to identify the top differentially regulated genes. To identify genes
that may encode markers useful for cell identification via immunostaining or flow cytometric
cell sorting, we filtered DE genes for those that encoded transmembrane proteins (**Table
S4**).

By comparing EMPs to haematopoietic and endothelial cells, we identified *Csf1r* and
Alox5ap as enriched in EMPs (**Fig. 6A**), agreeing with previous studies that identified them
as EMP markers (Hoeffel *et al.*, 2015; Sharon *et al.*, 2019). In addition, we identified *Fcer1g*,
Tyrobp and *Mctp1* as potential new markers for EMPs (**Fig. 6A,B**), whereas *Fxyd5* and
Scarf1 were identified as markers of yolk sac ECs and EMPs, and to a lower extent of BP1
(**Fig. 6B**). None of these seven genes were expressed in more differentiated BP2 and
erythroid cells. A comparison of genes expressed in haemogenic ECs versus EMPs showed
that acquisition of an EMP identity was associated with a loss of *Stab2* (**Fig. 6C**), as
previously described (Shvartsman *et al.*, 2019). Furthermore, we identified *Icam2*, *Cldn5*
and *Gpr182* as additional genes that were expressed in yolk sac ECs and haemogenic ECs
but downregulated in EMPs (**Fig. 6C**).

By comparing EMPs to BP1, and BP1 to BP2, we identified transcriptional changes
associated with the EMP transition to more committed progenitor states. This transition was
associated with the downregulation of the endothelial genes *Ecscr* and *Cdh5*, the EMP
marker *Alox5ap*, as well as the downregulation of our predicted novel EMP marker *Fcer1g*
and yolk sac EC and EMP markers *Fxyd5* and *Scarf1* (**Fig. 6A-D**). By contrast, the EMP
transition to more committed progenitor states was associated with the upregulation of the
megakaryocytic gene *F2r* and the erythroid gene *Smim1* (**Fig. 6D**).

To confirm that *Fxyd5*, *Scarf1* and *Fcer1g* encode EMP and haemogenic endothelial
markers, we performed immunofluorescence staining of E8.5 yolk sac. Although we had
found *Fxyd5* transcripts in yolk sac ECs and EMPs, FXYD5 staining was restricted to round

KIT$^+$ PECAM1$^+$ cells consistent with newly emerging EMPs, and there was no obvious
staining of ECs (**Figs. 6E, S3A**). Additionally, FXYD5 immunostaining identified round KIT$^-$
PECAM1^{low} cells, agreeing with our transcriptomic analysis detecting BPs with no *Kit*
transcripts and low levels of *Fxyd5* (**Fig. 6B**). Consistent with the presence of *Scarf1*
transcripts in both yolk sac ECs and EMPs, immunofluorescence staining identified SCARF1
protein in flat KIT$^-$ PECAM1$^+$ ECs and round KIT$^+$ PECAM1$^+$ emerging EMPs. Furthermore,
SCARF1 immunostaining identified flat KIT$^+$ PECAM1$^+$ cells that may be haemogenic ECs
(**Figs. 6F, S3B**). Although *Fcer1g* transcripts were detected in both EMPs and BP1,
immunofluorescence identified FCER1G protein within a subset of round KIT$^+$ cells.
Additionally, FCER1G immunostaining identified flat KIT$^-$ PECAM1$^+$ ECs, consistent with our
transcriptomic analysis detecting low levels of *Fcer1g* transcripts in yolk sac ECs (**Figs. 6G,**
S3C).

A comparison with clusters from the embryo showed that the markers identified in our
analysis as expressed in yolk sac ECs (*Fxyd5*, *Scarf1*, *Icam2*, *Cldn5*, *Gpr182*) were also
detected in the endocardium and in venous ECs (**Fig. 7A-C**). In contrast, the EMP markers
we identified (*Csf1r*, *Alox5ap*, *Fcer1g*, *Tyrobp*, *Mctp1*) were not expressed in ECs from the
embryo, confirming their specificity. Nevertheless, these markers were also expressed in
EMP-like cells from the embryo, thus corroborating their appropriate annotation.

Taken together, these findings show that novel EMP markers can be identified by
scRNAseq analysis and that they could be used to investigate both *bona fide* EMPs in the
yolk sac as well as poorly characterised EMP-like cells found in the embryo.

DISCUSSION

To capture the time when EMPs emerge from the yolk sac haemogenic endothelium
and to define the transcriptome of the cell populations involved in EHT, we re-analysed
transcriptomic data from the E8.5 mouse embryo (**Fig. 1**). Our analysis provided a refined
annotation of endothelial and haematopoietic populations in the mouse yolk sac (**Fig. 3A,B**)
and identified a subset of ECs co-expressing endothelial and haematopoietic markers,
consistent with a haemogenic endothelial phenotype (**Fig. 4A,B**). This refined annotation
also defined E8.5 yolk sac EMPs and demonstrated co-expression of myeloid, erythroid and
progenitor markers in this cluster, consistent with the known dual erythro-myeloid
differentiation potential of EMPs. Moreover, we identified EMP-like cells within the E8.5
embryo proper, which have not been highlighted in prior transcriptomic studies of early
haematopoiesis (**Fig. 3B**). Knowing that haematopoietic cells enter the embryo only from
E9.5, when the heart begins to beat to initiate blood circulation (Frame *et al.*, 2016; Kasaai

et al., 2017), we considered two possibilities to explain the existence of EMP-like cells in the

embryo. On the one hand, these cells may represent yolk sac-derived haematopoietic

progenitors that had contaminated the embryonic samples during dissection. On the other

hand, it is conceivable that these cells represent EMP-like cells originating in the embryo

rather than in the yolk sac. The latter possibility is consistent with our recent discovery of a

Pax3 lineage-derived EMP-like progenitor in the mouse embryo (Canu *et al.*, 2024).

However, further work is required to accurately describe these EMP-like cells.

Others have used scRNAseq to map potential differentiation trajectories for EMPs in

the yolk sac (Zhao *et al.*, 2023). Additionally, our analysis identified distinct BP1 and BP2

clusters as consecutive stages of EMP maturation (**Fig. 5A-C**) and provided new insight into

candidate signalling pathways controlling this transition (**Fig. 5D-H**). As primitive

erythrocytes are produced from E7.0 onwards (Palis *et al.*, 1999; Ferkowicz and Yoder,

2005), but EMPs only emerge from E8.5 onwards (Frame, McGrath and Palis, 2013; Gomez

Perdiguero *et al.*, 2014; Hoeffel *et al.*, 2015; McGrath *et al.*, 2015), the erythroid cluster in

the E8.5 dataset is expected to contain mostly primitive erythrocytes that were produced

prior to or alongside EMP emergence, with only a potential minor contribution from EMP-

derived erythrocytes. No markers of fully differentiated macrophages or other myeloid cell

types were observed, confirming previous studies showing that macrophages form in the

yolk sac from E9.0 onwards (Hoeffel and Ginhoux, 2018). We also did not detect a lymphoid

cluster at E8.5, in agreement with prior reports of LMP presence in the yolk sac only from

E9.5 onwards (Böiers *et al.*, 2013).

Previous studies established that EHT is associated with downregulation of

endothelial genes and concomitant upregulation of haematopoietic genes (Palis *et al.*, 1999;

Frame, McGrath and Palis, 2013; Gritz and Hirschi, 2016; Lilly *et al.*, 2016; Canu *et al.*,

2020). Here, pseudotime reconstruction highlighted dynamic gene expression trends during

EHT and haematopoietic maturation in the yolk sac (**Fig. 5A-D**), whereby haemogenic ECs

form a transcriptional continuum bridging the transition from endothelial to haematopoietic

cell identities, thus implying a gradual and continuous process rather than a discrete lineage

switch. Altogether, these findings align with established models of yolk sac haematopoiesis

derived from genetic lineage tracing, flow cytometry experiments and prior transcriptomic

analysis (Frame, McGrath and Palis, 2013; Hoeffel and Ginhoux, 2018; Canu and Ruhrberg,

2021; Sánchez-Lanzas, Jiménez-Pompa and Ganuza, 2024; Sugimura *et al.*, 2025).

The progressive downregulation of genes associated with angiogenesis, cytoskeletal

organisation and endothelial adhesion during the predicted EHT (**Fig. 5E**) may indicate

structural and functional changes necessary for cell detachment and transition to a

haematopoietic phenotype. Conversely, the upregulation of genes associated with RNA processing (**Fig. 5F**) suggests that haematopoietic specification is coupled to enhanced transcriptional activity. Furthermore, upregulation of genes associated to cell cycle activation and transition from the G1 to S phase (**Table S3**) agrees with previous findings that EHT requires cell cycle progression and is controlled by the cell cycle regulators CDK4/6 and CDK1 (Marcelo *et al.*, 2013; Canu *et al.*, 2020).

DE analysis identified components of the TYROBP signalling pathway as significantly enriched in EMPs (**Fig. S2**). In the adult, the activation of the TYROBP signalling pathway has been associated with the microglial response to stress (Haure-Mirande *et al.*, 2022; Zhang *et al.*, 2024). However, to the best of our knowledge, this signalling pathway has not been previously associated with EMP emergence. Further investigation should determine whether this pathway may be functionally involved in the production of EMPs or other haematopoietic progenitors (*e.g.*, HSCs from the dorsal aorta) and whether manipulating TYROBP *in vitro* during differentiation of induced pluripotent stem cells may allow to devise more efficient haematopoietic differentiation methods.

DE analysis also identified upregulation of genes associated with oxidative phosphorylation upon differentiation of EMPs to BP1 (**Fig. 5H**). Previous studies of pluripotent stem cells and neuronal progenitors showed that their differentiation is associated with metabolic reprogramming, whereby differentiating progenitors switch from glycolysis to oxidative phosphorylation (Zheng *et al.*, 2016; Dahan *et al.*, 2019; Mahmoud, 2023). Our analysis raises the possibility that an analogous process may occur when haematopoietic progenitors in the yolk sac commit to differentiation.

Our DE analysis identified both known and previously undescribed candidate markers associated with the formation of haemogenic endothelium and EMPs (**Fig. 6A-D**). Established EMP markers such as *Csf1r* and *Alox5ap* were previously described by others (Hoeffel *et al.*, 2015; Sharon *et al.*, 2019), thus validating the robustness of our transcriptomic analysis. In addition, we identified *Fcer1g*, *Tyrobp*, and *Mctp1* as potential novel EMP markers, and *Fxyd5* and *Scarf1* as shared markers between yolk sac endothelial and haematopoietic cells. The expression of *Fxyd5* and *Scarf1* in endothelial and EMP clusters, and to a lower extent in BP1, but not in more differentiated cells, suggests that these genes may serve as new markers for yolk sac ECs and emerging EMPs. Immunofluorescence staining of E8.5 yolk sac corroborated the predictions of our DE analysis: FXYD5 was present on round KIT$^+$ PECAM1$^+$ cells consistent with emerging EMPs (**Figs. 6E, S3A**); SCARF1 labelled flat KIT$^-$ PECAM1$^+$ ECs, flat KIT$^+$ PECAM1$^+$ cells consistent with haemogenic ECs and round KIT$^+$ PECAM1$^+$ cells consistent with emerging

EMPs (**Figs. 6F, S3B**); FCER1G marked flat KIT⁻ PECAM1⁺ ECs and a few round KIT⁺

PECAM1⁺ cells that may represent a subset of EMPs (**Figs. 6G, S3C**). Nevertheless, these

experiments also highlighted differences in RNA and protein expression. Thus, although

Fxyd5 RNA was detected in yolk sac ECs, haemogenic ECs and EMPs, and at lower levels

in BP1, the FXYD5 protein appeared to be prevalent in round KIT⁺ PECAM1⁺ cells consistent

with EMPs and in round KIT⁻ PECAM1ˡᵒʷ cells presumed to be myeloid-committed BP1

(**Figs. 6E, S3A**). While *Fcer1g* RNA clearly identified the EMP cluster, the FCER1G protein

was detected in only a subset of round KIT⁺ PECAM1⁺ cells, possibly marking more mature

EMPs (**Figs. 6G, S3C**). This is reminiscent of observations for *Csf1r*, a gene expressed in

EMPs at the RNA level, whereas CSF1R protein only becomes translated upon

differentiation to myeloid cells (Hoeffel *et al.*, 2015). These discrepancies between RNA and

protein expression may be due differences in expression dynamics and half-lives for mRNAs

and proteins (Vogel and Marcotte, 2012). Overall, these immunostaining results validated

our transcriptomic predictions and highlight the utility of single-cell datasets in discovering

functional cell surface markers. Previously used markers for haemogenic ECs include the

surface markers CDH5, KIT and CD41, and haematopoietic transcription factors such as

RUNX1 (Mikkola *et al.*, 2003; Nadin, Goodell and Hirschi, 2003; Lilly *et al.*, 2016), whereas

previously used markers for EMPs include KIT, CD41 and CD16/32, as well as the gene

expression reporter *Csf1r*-eGFP (McGrath *et al.*, 2011; Azzoni *et al.*, 2018; Plein *et al.*,

2018). The addition of the novel markers identified in our analysis will provide

complementary tools to define haemogenic ECs and EMPs in the mouse yolk sac.

In summary, this study refines the cellular and molecular characterisation of yolk sac

haematopoiesis, provides a detailed transcriptomic roadmap of EHT in the yolk sac and

identifies novel candidate markers for haemogenic endothelium and EMPs. The

identification of FXYD5, SCARF1 and FCER1G as markers for EMPs and haemogenic ECs

may offers valuable tools for future *in vivo* and *in vitro* studies. Finally, future studies may

wish to determine whether the novel EMP markers are present also in foetal HSCs, and

whether they are conserved in human haemopoiesis. Defining the specificity of such novel

markers may allow researchers to better define EMP identity and lineage potential as a

prerequisite for deciphering the ontogeny of the immune system and, ultimately, to harness

early haematopoietic pathways for regenerative medicine.

ACKNOWLEDGEMENTS

We thank the staff of the Biological Resources, FACS and Imaging Facilities at the UCL Institute of Ophthalmology for their technical support. This research was supported by grants from Wellcome (205099/Z/16/Z) and the British Heart Foundation (PG23/11301).

AUTHORSHIP CONTRIBUTIONS

GC, GDP and CR conceived and designed the study. GC and CR co-wrote the manuscript. GC and AM performed immunostaining experiments. GDP performed bioinformatic analyses. GC and CR supervised the project. All authors read and approved the submitted manuscript.

DISCLOSURE OF CONFLICTS OF INTEREST

The authors declare that they have no competing interests.

REFERENCES

Agrawal, A. *et al.* (2024) 'WikiPathways 2024: next generation pathway database', *Nucleic Acids Research*, 52(D1), pp. D679–D689.

Aran, D. *et al.* (2019) 'Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage', *Nature Immunology 2019 20:2*, 20(2), pp. 163–172.

Azzoni, E. *et al.* (2018) 'Kit ligand has a critical role in mouse yolk sac and aorta–gonad–mesonephros hematopoiesis', *EMBO reports*, 19(10).

Böiers, C. *et al.* (2013) 'Lymphomyeloid Contribution of an Immune-Restricted Progenitor Emerging Prior to Definitive Hematopoietic Stem Cells', *Cell Stem Cell*, 13(5), pp. 535–548.

Cannoodt, R., Saelens, W. and Saeys, Y. (2016) 'Computational methods for trajectory inference from single-cell transcriptomics', *European Journal of Immunology*, 46(11), pp. 2496–2506.

Canu, G. *et al.* (2020) 'Analysis of endothelial-to-haematopoietic transition at the single cell level identifies cell cycle regulation as a driver of differentiation', *Genome Biology*, 21(1), pp. 1–28.

Canu, G. *et al.* (2024) 'A Pax3 lineage gives rise to transient haematopoietic progenitors', *Development (Cambridge, England)*, 151(23).

Canu, G. and Ruhrberg, C. (2021) 'First blood: the endothelial origins of hematopoietic progenitors', *Angiogenesis 2021 24:2*, 24(2), pp. 199–211.

Cao, J. *et al.* (2019) 'The single-cell transcriptional landscape of mammalian organogenesis', *Nature 2019 566:7745*, 566(7745), pp. 496–502.

Carlson, M. (2025) *org.Mm.eg.db - Genome wide annotation for Mouse*, *Bioconductor*.

Dahan, P. *et al.* (2019) 'Metabolism in pluripotency: Both driver and passenger?', *Journal of Biological Chemistry*, 294(14), pp. 5420–5429.

Van Deren, D.A. *et al.* (2022) 'Defining the Hoxb8 cell lineage during murine definitive hematopoiesis', *Development (Cambridge, England)*, 149(8), p. dev200200.

Ferkowicz, M.J. and Yoder, M.C. (2005) 'Blood island formation: Longstanding observations and modern interpretations', *Experimental Hematology*, 33(9 SPEC. ISS.), pp. 1041–1047.

Frame, J.M. *et al.* (2016) 'Definitive Hematopoiesis in the Yolk Sac Emerges from Wnt-
Responsive Hemogenic Endothelium Independently of Circulation and Arterial Identity',
Stem Cells, 34(2), pp. 431–444.

Frame, J.M., McGrath, K.E. and Palis, J. (2013) 'Erythro-myeloid progenitors: "definitive"
hematopoiesis in the conceptus prior to the emergence of hematopoietic stem cells', *Blood*
cells, molecules & diseases, 51(4), pp. 220–225.

Goh, I. *et al.* (2023) 'Yolk sac cell atlas reveals multiorgan functions during human early
development', *Science*, 381(6659).

Gomez Perdiguero, E. *et al.* (2014) 'Tissue-resident macrophages originate from yolk-sac-
derived erythro-myeloid progenitors', *Nature 2014 518:7540*, 518(7540), pp. 547–551.

Gordon, E.J., Gale, N.W. and Harvey, N.L. (2008) 'Expression of the hyaluronan receptor
LYVE-1 is not restricted to the lymphatic vasculature; LYVE-1 is also expressed on
embryonic blood vessels', *Developmental Dynamics*, 237(7), pp. 1901–1909.

Gritz, E. and Hirschi, K.K. (2016) 'Specification and function of hemogenic endothelium
during embryogenesis', *Cellular and molecular life sciences : CMLS*, 73(8), pp. 1547–
1567.

Gustavsen, J.A. *et al.* (2019) 'RCy3: Network biology using Cytoscape from within R',
F1000Research 2019 8:1774, 8, p. 1774.

Haure-Mirande, J.V. *et al.* (2022) 'Microglial TYROBP/DAP12 in Alzheimer's disease:
Transduction of physiological and pathological signals across TREM2', *Molecular*
Neurodegeneration 2022 17:1, 17(1), pp. 1–15.

Hoeffel, G. *et al.* (2015) 'C-Myb+ Erythro-Myeloid Progenitor-Derived Fetal Monocytes
Give Rise to Adult Tissue-Resident Macrophages', *Immunity*, 42(4), pp. 665–678.

Hoeffel, G. and Ginhoux, F. (2018) 'Fetal monocytes and the origins of tissue-resident
macrophages', *Cellular Immunology*, 330, pp. 5–15.

Ivanovs, A. *et al.* (2011) 'Highly potent human hematopoietic stem cells first emerge in the
intraembryonic aorta-gonad-mesonephros region', *Journal of Experimental Medicine*,
208(12), pp. 2417–2427.

Ivanovs, A. *et al.* (2017) 'Human haematopoietic stem cell development: from the embryo
to the dish', *Development*, 144(13), pp. 2323–2337.

Jiang, Y. *et al.* (2024) 'Macrophages in organ fibrosis: from pathogenesis to therapeutic

targets', *Cell Death Discovery 2024 10:1*, 10(1), pp. 1–11.

Kasaai, B. *et al.* (2017) 'Erythro-myeloid progenitors can differentiate from endothelial cells
and modulate embryonic vascular remodeling', *Scientific Reports*, 7(1), pp. 1–12.

Korsunsky, I. *et al.* (2019) 'Fast, sensitive and accurate integration of single-cell data with
Harmony', *Nature Methods 2019 16:12*, 16(12), pp. 1289–1296.

Lilly, A.J. *et al.* (2016) 'Interplay between SOX7 and RUNX1 regulates hemogenic
endothelial fate in the yolk sac', *Development (Cambridge, England)*, 143(23), pp. 4341–
4351.

Liu, X. *et al.* (2024) 'Inducible CCR2+ nonclassical monocytes mediate the regression of
cancer metastasis', *The Journal of clinical investigation*, 134(22).

Mahmoud, A.I. (2023) 'Metabolic switches during development and regeneration',
Development (Cambridge), 150(20).

Marcelo, K.L. *et al.* (2013) 'Hemogenic endothelial cell specification requires c-Kit, notch
signaling, and p27-mediated cell-cycle control', *Developmental Cell*, 27(5), pp. 504–515.

Marsh, S. *et al.* (2025) *scCustomize: Custom Visualizations & Functions for Streamlined*
Analyses of Single Cell Sequencing, *CRAN: Contributed Packages*.

McGrath, K.E. *et al.* (2011) 'Definitive Erythro-Myeloid Progenitors (EMP) Emerge in the
Yolk Sac From Hemogenic Endothelium and Share Transcriptional Regulators with Adult
Hematopoiesis', *Blood*, 118(21), p. 910.

McGrath, K.E. *et al.* (2015) 'Distinct sources of hematopoietic progenitors emerge before
HSCs and provide functional blood cells in the mammalian embryo', *Cell reports*, 11(12),
p. 1892.

McGrath, K.E., Frame, J.M. and Palis, J. (2016) 'Early hematopoiesis and macrophage
development', *Seminars in immunology*, 27(6), p. 379.

Mikkola, H.K.A. *et al.* (2003) 'Expression of CD41 marks the initiation of definitive
hematopoiesis in the mouse embryo', *Blood*, 101(2), pp. 508–516.

Moskalik, A., Niderla-Bielińska, J. and Ratajska, A. (2021) 'Multiple roles of cardiac
macrophages in heart homeostasis and failure', *Heart Failure Reviews*, 27(4), p. 1413.

Nadin, B.M., Goodell, M.A. and Hirschi, K.K. (2003) 'Phenotype and hematopoietic
potential of side population cells throughout embryonic development', *Blood*, 102(7), pp.
2436–2443.

Oberlin, E. *et al.* (2010) 'Definitive human and mouse hematopoiesis originates from the
embryonic endothelium: a new class of HSCs based on VE-cadherin expression', *The*
International Journal of Developmental Biology, 54(6–7), pp. 1165–1173.

Pagès, H. *et al.* (2025) *AnnotationDbi - Manipulation of SQLite-based annotations in*
Bioconductor, Bioconductor.

Palis, J. *et al.* (1999) 'Development of erythroid and myeloid progenitors in the yolk sac
and embryo proper of the mouse', *Development*, 126(22), pp. 5073–5084.

Pijuan-Sala, B. *et al.* (2019) 'A single-cell molecular map of mouse gastrulation and early
organogenesis', *Nature 2019 566:7745*, 566(7745), pp. 490–495.

Plein, A. *et al.* (2018) 'Erythro-myeloid progenitors contribute endothelial cells to blood
vessels', *Nature 2018 562:7726*, 562(7726), pp. 223–228.

Sánchez-Lanzas, R., Jiménez-Pompa, A. and Ganuza, M. (2024) 'The evolving
hematopoietic niche during development', *Frontiers in Molecular Biosciences*, 11, p.
1488199.

Satija, R. *et al.* (2015) 'Spatial reconstruction of single-cell gene expression data', *Nature*
Biotechnology 2015 33:5, 33(5), pp. 495–502.

Scarfò, R. *et al.* (2024) 'CD32 captures committed haemogenic endothelial cells during
human embryonic development', *Nature Cell Biology 2024 26:5*, 26(5), pp. 719–730.

Shannon, P. *et al.* (2003) 'Cytoscape: A Software Environment for Integrated Models of
Biomolecular Interaction Networks', *Genome Research*, 13(11), p. 2498.

Sharon, N. *et al.* (2019) 'A Peninsular Structure Coordinates Asynchronous Differentiation
with Morphogenesis to Generate Pancreatic Islets', *Cell*, 176(4), pp. 790-804.e13.

Shvartsman, M. *et al.* (2019) 'Single-cell atlas of major haematopoietic tissues sheds light
on blood cell formation from embryonic endothelium', *bioRxiv*, p. 774547.

Sugimura, R. *et al.* (2025) 'Editorial: Plasticity of the haematopoietic niche: from embryonic
development to aging and disease', *Frontiers in Molecular Biosciences*, 12, p. 1683902.

Tavian, M. *et al.* (1996) 'Aorta-Associated CD34+ Hematopoietic Cells in the Early Human
Embryo', *Blood*, 87(1), pp. 67–72.

Vogel, C. and Marcotte, E.M. (2012) 'Insights into the regulation of protein abundance from
proteomic and transcriptomic analyses', *Nature Reviews Genetics 2012 13:4*, 13(4), pp.
227–232.

Wickham, H. *et al.* (2025) 'ggplot2: Create Elegant Data Visualisations Using the Grammar
of Graphics', *CRAN: Contributed Packages* [Preprint].

Wu, T. *et al.* (2021) 'clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data', *Innovation*, 2(3), p. 100141.

Zhang, M. *et al.* (2024) 'TYROBP serve as potential immune-related signature genes in
the acute phase of intracerebral hemorrhage', *Scientific Reports 2024 14:1*, 14(1), pp. 1–
12.

Zhao, Y.X. *et al.* (2023) 'Single-cell RNA sequencing-guided fate-mapping toolkit
delineates the contribution of yolk sac erythro-myeloid progenitors', *Cell Reports*, 42(11),
p. 113364.

Zheng, X. *et al.* (2016) 'Metabolic reprogramming during neuronal differentiation from
aerobic glycolysis to neuronal oxidative phosphorylation', *eLife*, 5(JUN2016), p. e13374.

Fig 1. Re-analysis of E8.5 mouse embryonic scRNAseq data to study yolk sac EHT.

A Workflow for the integrated re-analysis of mouse embryonic scRNAseq datasets E-MTAB-11763 and E-MTAB-6967.

B UMAP of dataset E-MTAB-11763 showing cells sequenced from the E8.5 mouse embryo and labelled according to their anatomical origin as embryo and yolk sac.

C UMAP obtained by subsetting haemato-vascular cells from E8.5 yolk sac from E-MTAB-11763 and labelled based on our refined cluster annotation.

EMP: erythro-myeloid progenitor; EC: endothelial cell; HE: haemogenic endothelium; YS: yolk sac.

Figure 2

Fig 2. Marker analysis identifies haemato-vascular cell populations in E8.5 yolk sac data in E-MTAB-11763.

UMAP representation of transcript levels for the indicated marker genes for (**A**) endothelial cells, (**B**) EMPs, (**C**) haematopoietic progenitors and (**D**) erythroid cells. The scale bar represents log-normalised transcript levels.

Figure 3

Fig 3. Refined cluster annotation of E8.5 embryo and yolk sac haemato-vascular cell clusters after integration of E-MTAB-11763 and E-MTAB-6967.

A Reference-based annotation based on gene expression was used to apply our revised annotation for E8.5 data in E-MTAB-11763 to the E8.5 data in E-MTAB-6967 before integrating the two datasets.

B UMAPs showing the E8.5 haemato-vascular clusters from both embryo and yolk sac for dataset E-MTAB-11763 before integration to dataset E-MTAB-6967 and after reference-mapping and integration of the two datasets.

C-F UMAP representation of transcript levels for the indicated marker genes for (**C**) endothelial cells, (**D**) EMPs, (**E**) haematopoietic progenitors and (**F**) erythroid cells. The scale bar represents log-normalised transcript levels.

Figure 4

A

Fig 4. Endothelial cell and EMP sub-clustering identifies haemogenic endothelium.

A UMAP obtained after subsetting and re-clustering all cells annotated as endothelial cells and EMPs after integration of E8.5 data in E-MTAB-11763 and E-MTAB-6967. The dotted box indicates an area that is zoomed on the cells previously annotated as yolk sac endothelial cells, vitellin vein endothelial cells and EMPs; note the formation of a separate cluster positioned between EMPs and the main yolk sac endothelial cluster consistent with a haemogenic endothelial identify.

B UMAP representation of key endothelial and haematopoietic markers corroborates the identification of the haemogenic endothelial cluster. The scale bars represent log-normalised transcript levels. YS: yolk sac; EMPs: erythro-myeloid progenitors.

Figure 5

Fig 5. Pseudotime analysis identifies co-regulated genes and enriched GO pathways during EHT.

A UMAP obtained after subsetting and re-clustering yolk sac haemato-vascular cells after integration of E8.5 data in E-MTAB-11763 and E-MTAB-6967, and showing our refined cluster annotation.

B,C Pseudotime analysis highlights erythroid generation and progressive maturation (**B**) and, after excluding the erythroid cluster (**C**), the progressive differentiation from yolk sac endothelial cells to haemogenic endothelial cells and then EMPs followed by blood progenitors 1 and 2. The scale bars represent pseudotime.

D Genes that changed as a function of pseudotime were identified and grouped according to their expression patterns. Heatmap shows 4 groups of co-regulated genes during differentiation from yolk sac endothelium to blood progenitors 2.

E-H Gene ontology analysis for each group of co-regulated genes.

YS: yolk sac; EMP: erythro-myeloid progenitor; BP1: blood progenitors 1.

Figure 6

Fig 6. Differential gene (DE) expression analysis and immunostaining validation identifies new markers for haemogenic endothelial cells and EMPs.

A-D UMAP representation of transcript levels for the indicated marker genes identified by DE analysis between the different yolk sac haemato-vascular clusters. The scale bars represent log-normalised transcript levels.

E-G Immunofluorescence staining of E8.5 yolk sac for the EMP marker KIT together with PECAM1 and FXYD5, SCARF1 or FCER1G. Emerging EMPs are indicated with arrows (round KIT+ PECAM1+ cells), presumed blood (myeloid) progenitors with arrowheads (round KIT- PECAM1low cells), endothelial cells with empty arrows (flat KIT- PECAM1+ cells), haemogenic endothelial cells with empty arrowheads (flat KIT+ PECAM1+ cells) and mature EMPs with asterisk (round KIT+ PECAM1+ FCER1G+ cells). Scale bars: 50 µm; n = 3 embryos.

Figure 7

A YS haemato-vascular clusters

B Embryo haemato-vascular clusters

C

Fig 7. Expression analysis in haemato-vascular cells from both yolk sac and embryo shows specificity of the new markers for haemogenic ECs and EMPs.

A,B Dot plot representation of markers identified by DE analysis in (**A**) yolk sac and (**B**) embryo haemato-vascular clusters. The dot size reflects the percentage of cells in each cluster expressing the gene. The scale bar indicates log-normalised gene expression.

C UMAP of the E8.5 haemato-vascular clusters from both embryo and yolk sac showing transcript levels for the indicated markers identified by DE analysis. The scale bars represent log-normalised transcript levels.

Supplementary Figure 1

Fig S1. Cluster annotation of E8.5 mouse scRNAseq data in E-MTAB-11763.

UMAP of the E8.5 cells in E-MTAB-11763 dataset, labelled with the cluster annotations provided by the authors of the original publication (Goh et al., 2023).

Supplementary Figure 2

TYROBP signalling pathway
EMP gene enrichment

Fig S2. The TYROBP signalling pathway is enriched in emerging EMPs.

TYROBP pathway genes enriched in EMPs compared to haemato-vascular clusters from both yolk sac and embryo. The box colour indicates the average log2 fold change. A green border indicates significant enrichment in EMPs.

Supplementary Figure 3

A E8.5 yolk sac

Fig S3. Immunostaining validation of new markers for haemogenic endothelial cells and EMPs.

A-C Immunofluorescence staining of E8.5 yolk sac for the EMP marker KIT together with PECAM1 and FXYD5, SCARF1 or FCER1G. Emerging EMPs are indicated with arrows (round KIT+ PECAM1+ cells), presumed blood (myeloid) progenitors with arrowheads (round KIT- PECAM1low cells), endothelial cells with empty arrows (flat KIT- PECAM1+ cells), haemogenic endothelial cells with empty arrowheads (flat KIT+ PECAM1+ cells) and mature EMPs with asterisk (round KIT+ PECAM1+ FCER1G+ cells). Scale bars: 50 μm; n = 3 embryos.