ApoA-I dissociated from human HDL retains its acceptor properties in ABCA1-mediated cholesterol efflux from RAW 264.7 macrophages in coronary artery disease

Veronika B. Baserova[1], Mikhail A. Popov[2], Alexander D. Dergunov[1,*]

[1]National Medical Research Center for Therapy and Preventive Medicine, Petroverigsky street 10, 101990 Moscow, Russia

[2]M.F. Vladimirsky Moscow Regional Research and Clinical Institute MONIKI, Moscow, Russia

*Correspondence: add1796@list.ru

Keywords: apoA-I dissociation; coronary artery disease; gene expression; HDL functionality; HDL structure; lipid-bound apoA-I; lipid-free apoA-I;

ABSTRACT

The significance of cholesterol efflux as a predictor of coronary artery disease (CAD) remains controversial. The major pathway of the export of intracellular cholesterol by the ABCA1 transporter includes the acceptance of effluxed cholesterol by lipid-free apolipoprotein A-I and/or high-density lipoproteins (HDL). To separate the contribution of lipid-free and lipid-bound apoA-I with different efficiencies in the total cholesterol efflux and underlying mechanism, we measured apoA-I dissociation concomitant with HDL denaturation, the expression of preselected genes, and the efficiency of cholesterol efflux. HDL preparations from plasma of 63 control and 76 CAD male patients were denatured by 4.25 M urea at the transition midpoint. ApoA-I partitioned 1.5-fold higher into the water phase for HDL from CAD patients relative to controls. The dissociation parameter D was reciprocally correlated with phospholipid and cholesterol levels and PL:apoA-I ratio in HDL from CAD patients. For control patients, the D parameter was positively correlated with *CETP* and *ABCA1* expression level. For CAD patients, the D parameter was negatively correlated with *CUBN* and *ALB* expression levels that may be associated with the increased catabolism of lipid-free apoA-I. The apoA-I functionality in ABCA1-mediated cholesterol efflux from RAW 264.7 macrophages to urea-pretreated HDL revealed by V_m, K_m, and V_m/K_m values was similar for HDL from both cohorts. The enrichment of HDL with phospholipid and cholesterol could be involved in the increased apolipoprotein dissociation in CAD. The estimates of the contribution of both lipid-free and lipid-bound apoA-I to total cholesterol efflux are required to accurately describe them as CAD predictors.

1. INTRODUCTION

The human plasma high-density lipoproteins (HDL) possess an atheroprotective effect through the reverse cholesterol transport from arterial-wall macrophages to the liver [1] that begins with cholesterol efflux via active cholesterol transport by ABCA1 and ABCG1 transporters, facilitated diffusion with SR-B1, and passive diffusion [2]. The apoA-I is a major HDL apolipoprotein that exists in lipid-bound and lipid-free forms; the latter is a primary cholesterol acceptor in the ABCA1-mediated efflux [3]. However, small HDL particles are efficient cholesterol acceptors as well [4]. The HDL heterogeneity by density and charge results in the existence of light HDL_2 and dense HDL_3 and particles with preβ-mobility and α-mobility, respectively. Lipid-free apoA-I is found only in the preβ-band [5], which also includes nascent HDL, while mature spherical HDL is localized in the α-band. The nascent HDL, as the initial product of cholesterol and phospholipid efflux by ABCA1, further accepts these lipids with the formation of disc-shaped particles. The latter are the most efficient substrate for lecithin:cholesterol acyltransferase (LCAT). The enzyme converts free cholesterol to cholesteryl ester that is thought to prevent the reabsorption of effluxed cholesterol. The subsequent steps of reverse cholesterol transport include the transfer of cholesteryl ester from HDL to larger very low-density and low-density lipoproteins (LDL) by cholesteryl ester transfer protein (CETP), the SR-B1-mediated capture of cholesteryl ester, and the uptake of a whole LDL particle by the LDL receptor in the liver [2]. The remodeling of spherical HDL by CETP and lipases results in a spherical-to-discoidal transformation with a concomitant transition between lipid-free and lipid-bound apoA-I [6]. Of note, the dissociated apoA-I [7; 8] and HDL-apoA-I exchange [9] may be potentially involved in cholesterol efflux.

A strong inverse relationship between HDL cholesterol efflux capacity (CEC) and coronary artery disease (CAD) signifies HDL functionality more than HDL-cholesterol level in the HDL atheroprotective effect [10]. HDL functionality includes HDL particle number and heterogeneity and particle ability to accept effluxed cholesterol. The significance of ABCA1-mediated efflux and HDL functionality in CAD are inconsistent [4; 11-16]. CEC from macrophages had a strong inverse association with morphological and prognostic markers of angiographic CAD [17]. The higher concentration of $preβ_1$ particles was responsible for the lower $preβ_1$ concentration-normalized ABCA1-dependent efflux capacity in CAD [18; 19]. ABCA1-dependent cholesterol efflux was positively correlated with the levels of small lipid-poor $preβ_1$ particles [18]. At variance with this, HDL efflux capacity was inversely associated with small HDL particle concentration in older adults [20]. Furthermore, the association between HDL particle size and composition and CAD risk is not conclusive, and the inclusion and magnitude of lipid-free apoA-I and HDL particles in individual steps of cholesterol efflux remain to be implicitly solved.

We describe now the relations between apoA-I dissociation, HDL functionality and the expression of selected genes controlling HDL metabolism and atherogenesis in control and CAD patients.

2. MATERIALS AND METHODS

Patient selection with widely varied HDL-C levels (63 control patients and 76 patients with CAD confirmed by coronary angiography) and laboratory tests were described previously [21]. The crude HDL preparations (pHDL) were prepared by the precipitation of apoB-containing lipoproteins with PEG 7000–9000 [7]. The concentrations of choline-containing phospholipids and total cholesterol in HDL preparations were measured by enzyme methods using Sentinel (Italy) and HUMAN (Germany) kits, respectively, and apoA-I was measured by immunonephelometry with AU 480 (Beckman Coulter, USA).

ApoA-I content in preβ- and α-fractions of pHDL was measured by agarose gel electrophoresis followed by immunodetection of apoA-I [7]. PeakFit version 4.12 (SeaSolve Software Inc.) and ImageJ software were used to determine band intensities with baseline subtraction at the conditions of linear response of each band intensity to the amount of loaded sample.

Urea-induced pHDL denaturation after incubation for 6 h at 25 °C was followed by the accumulation of lipid-free preβ-apoA-I detected in the immunoreplica of agarose gel. ApoA-I dissociation was analyzed from the fractions of preβ-signal detected after the treatment with 0 M, 4.25 M, and 7.5 M urea [7]. The degree of apolipoprotein dissociation at 4.25 M urea as a mid-transition zone that is accomplished at 7.5 M urea was characterized by dissociation parameter D (Equation 1):

$$D = (pre\beta'_{4.25} - pre\beta'_0)/(pre\beta'_{7.5} - pre\beta'_0) \tag{1},$$

where $pre\beta'_{4.25}$, $pre\beta'_0$, and $pre\beta'_{7.5}$ are preβ fractions with 4.25 M, 0 M, and 7.5 M urea, respectively. The dependence of urea-induced apoA-I release on phospholipid level in pHDL preparations was characterized by the partition coefficient K between aqueous and lipid phases adjusted for water molarity (Equation 2):

$$K = (F/B)[\text{pHDL-PL}] \tag{2},$$

where F and B are the numbers of molecules of lipid-free apoA-I in water and bound to pHDL phospholipids, respectively [22]. The F/B ratio was calculated from the D parameter measured at 4.25 M urea (Equation 3):

$F/B = D/(1 - D)$ (3).

Cholesterol efflux from murine macrophages RAW 264.7 to pHDL was measured with a fluorescent probe BODIPY-cholesterol. The basal and cAMP-stimulated effluxes were measured, and ABCA1-mediated efflux was calculated as a difference between stimulated and basal effluxes and expressed as a percent of effluxed cholesterol [7].

The expression of 65 genes involved in HDL metabolism and atherogenesis measured by us earlier by real-time PCR in peripheral blood mononuclear cells [21; 23] was compared to apoA-I dissociation from pHDL in the present study.

Statistica 13 software was used for statistical analysis. The associations between variables were analyzed by the Pearson correlation coefficient. The statistical significance limit was accepted as $p < 0.05$. Nonlinear curve fitting was done with OriginPro 9.0.0 SR1. The significance of the difference between different dataset fits was checked with the F-statistic.

3. RESULTS

3.1. The relations between apoA-I dissociation, HDL composition, and gene expression
The dissociation of apoA-I from pHDL induced by urea treatment and measured by the *D* parameter significantly decreased with the increase of absolute and apoA-I-normalized cholesterol levels in pHDL from CAD patients. The apoA-I dissociation changed quite analogously with the increase of choline-containing phospholipid (PL) (Table 1). The fit of dissociation data, passed through zero, to the linear model of apoA-I distribution between lipid-bound (B) and lipid-free (F) states (Equation 2) results in the values of the partition coefficient between aqueous and lipid phases adjusted for water molarity $K = 0.239 \pm 0.012$ ($R^2 = 0.874$) for control pHDL and $K = 0.368 \pm 0.020$ ($R^2 = 0.822$) for pHDL from CAD patients. Thus, a 1.54-fold prevalence of apoA-I distribution between water and lipid phases exists in CAD relative to controls ($p = 0.000$) with the value of the difference in the standard free energy $\Delta\Delta G^0 = \Delta G^0_{CAD} - \Delta G^0_{control} = -RT\ln(K(CAD)/K(control)) = -1.2$ kJ/mol, the difference value being positive for apolipoprotein distribution between lipid and water phases. ApoA-I in HDL from CAD patients possesses higher free energy and, thus, exhibits a higher dissociation. Of note, the increased PL:apoA-I and Chol:apoA-I ratios for pHDL from CAD patients relative to controls were measured throughout the whole range of HDL-C concentration in both cohorts (data not shown).

To reveal the associations between apoA-I dissociation and lipoprotein metabolism, the expression of 65 genes in peripheral blood mononuclear cells, sensitive to HDL metabolism and atherogenesis, was analyzed. The significant correlations between *D* and gene expression are

included in Table 1. For control patients, the positive correlation of the *D* parameter with the *CETP* expression level existed. The positive correlation of the *D* parameter with the *ABCA1* expression level may be associated with the decrease of lipid-free apoA-I with the increase of ABCA1 activity and the accompanying increase of dissociation of lipid-bound apolipoprotein. The *A2M* and *PRKACG* expression levels were negatively correlated with the *D* parameter. Remarkably, the increased expression of the *A2M* gene has been reported to be the marker of atherosclerosis progression [24]. This suggestion contrasts with the positive correlation of the dissociation parameter with the expression of the *ITGB3* and *TLR8* genes which possess the proatherogenic effect in CAD [23].

For patients with CAD, the apoA-I dissociation from pHDL is not associated with *CETP* expression level. However, the negative correlations of the *D* parameter with *CUBN* and *ALB* expression levels existed.

3.2. The relations between apoA-I and cholesterol efflux

The efflux measurements were done, firstly, with intact pHDL as cholesterol acceptors. For controls, there was no significant correlation between apoA-I dissociation and ABCA1-mediated cholesterol efflux. In CAD, efflux was less efficient relative to controls, and dissociation parameter *D* was negatively correlated with efflux (Table 1) due to a negative contribution of $pre\beta_0'$, which is a most efficient cholesterol acceptor, to the *D* parameter (Equation 1).

Due to the involvement of lipid-free and lipid-bound apoA-I in cholesterol transport by the ABCA1 transporter, the decrease of efflux efficiency in CAD may be associated with the less efficient efflux with each or both acceptors. To separate individual contributions, secondly, the functionality of lipid-free apoA-I was studied. To induce apoA-I dissociation from the HDL lipid phase to the water phase, pHDL from control and CAD patients were predenatured for 6 h at 25°C by 4.25 M urea at the middle of the HDL denaturation transition. The ABCA-mediated efflux to pHDL for 2 h was then measured, and the data were fitted to Michaelis-Menten kinetics (Table 2). The concentration of lipid-free apoA-I dissociated from mature α-HDL was calculated as a product of dissociation parameter *D* and apoA-I level in the α-band estimated by agarose gel electrophoresis and apolipoprotein immunodetection (D*α-apoA-I). The dependencies were well fitted to the equation for one substrate reaction with R^2 values 0.72-0.75 and did not differ by the F-test for control and CAD. The V_m, K_m, and V_m/K_m kinetic parameters did not differ also for pHDL from control and CAD patients by Student's *t*-test. Of note, the kinetic parameters, checked by the F-statistic, did not change significantly with pHDL predenatured with 7.5 M urea (Table 2) at the completion of the denaturation transition for pHDL from control or CAD patients, which assumes no contribution of HDL remaining at the transition midpoint into the cholesterol efflux.

Thus, the lipid-free apoA-I seems to be equally efficient as an acceptor of cholesterol effluxed by ABCA1 in CAD relative to control samples.

4. DISCUSSION

There are three major findings in the present study revealed at the comparison of compositional and functional properties of HDL from CAD and control patients. First, for HDL from CAD patients, apoA-I partitioned more into the water phase during urea-induced HDL denaturation, and distribution parameter D was inversely associated with choline-containing phospholipid, absolute and apoA-I-normalized cholesterol levels in pHDL, opposite to the lack of any association for HDL from control patients. The increased denaturation of HDL from the patients with acute coronary syndrome has been described [25]. Importantly, HDL particles from CAD patients are more enriched with PL and Chol normalized by apoA-I level throughout the whole range of HDL-C. We measured earlier the upregulated expression of *CETP*, *LPL*, and *PLTP* genes and down-regulated expression of *LCAT* gene in CAD patients [23]. The differential expression of these genes is suggested to be involved in lipid enrichment of HDL in CAD. Indeed, the enrichment of HDL with phospholipid and cholesterol molecules in CAD could occur due to a decreased phospholipid hydrolysis by LCAT which possesses phospholipase activity with a concomitant lowering of cholesteryl ester generation [26]. In turn, the free cholesterol molecules may increasingly accumulate in the vicinity of apoA-I molecules in pHDL from CAD patients with the additional perturbation of PL structural order and dynamics. The entropically favourable processes of both apoA-I exchange between discoidal HDL-bound and lipid-free pools [27] and the lipid transfer for nanodisks consisting of apoA-I and DMPC [28] with the increased activation entropy and the decrease in the standard entropy have been suggested. The exact contribution of entropy and enthalpy into apoA-I increased dissociation from mature spherical HDL in CAD remains to be determined. The different exposure of the hydrophobic amino acid residues of apoA-I to the aqueous phase in the activated state at apolipoprotein denaturation [22] may be invoved. Interestingly, cholesterol in discoidal HDL increased the number of apoA-I tryptophan residues accessible to the aqueous phase, but decreased their mean degree of hydration [29].

Second, for HDL from control patients, the D parameter was positively correlated with the *CETP* expression. CETP-induced apoA-I dissociation from HDL from healthy normolipidemic subjects has been described by Barter et al. [30;31]. Also, triglyceride enrichment of HDL core, that augments apolipoprotein dissociation from the surface [32] may underlie the positive association between the D parameter and *CETP* expression. Of note, the increased apoA-I dissociation seems to possess a proatherogenic effect in control patients due to the positive associations between the D parameter and *ITGB3* and *TLR8* gene expression. For HDL from CAD

patients, the D parameter was not associated with *CETP* expression. However, the observed negative associations of the D parameter with *CUBN* and *ALB* gene expression may be associated with the increased catabolism of lipid-free apoA-I and the accompanying decrease of the preβ fraction in CAD. Indeed, CUBN as an endocytic receptor is involved in endocytosis of apoA-I and albumin and maintenance of their blood levels [33]. Interestingly, the reciprocal associations existed between the dissociation parameter and the expression levels of two genes, i.e. negative for *PRKACG* for control patients and positive for *PRKACB* for CAD patients. The expression levels of these genes contributed reciprocally to HDL-C variability in CAD patients [23].

Third, lipid-free/lipid-poor apoA-I that may originate from the HDL-bound state both for CAD and control HDL, seems to be equally efficient in ABCA1-mediated efflux. The obtained V_m and K_m values are lower than the analogous mean values (22.3% and 0.64 µg/ml) for ABCA1-mediated efflux to lipid-free apoA-I for 6 h [4]. Of note, only 40-60% of apoA-I molecules are dissociated in water upon pHDL denaturation by 4.25 M urea (compared to 10-20% of preβ-apoA-I in intact pHDL), and V_m could not reach the maximal limiting value for lipid-free apoA-I due to the lower V_m value for lipid-bound apoA-I. However, similar efflux kinetics at two zones of denaturation transition largely exclude this possibility. The decreased ABCA1-mediated efflux efficiency of pHDL as a mixture of lipid-poor and lipid-bound apoA-I in CAD has been described [18;19]. Despite the attempts to normalize efflux to preβ$_1$-levels as an indicator of lipid-poor apolipoprotein, the conventional CEC measurements are performed with pHDL as a mixture of lipid-poor and lipid-bound apoA-I without the separation of individual efflux reactions with different acceptors [18;19;34]. However, the overall reaction velocity for a two-substrate reaction with a single enzyme depends in a non-linear manner on the concentrations of both substrates.

In conclusion, for control patients, the dissociation of apoA-I in water phase is associated with *ABCA1* and *CETP* gene expression. For CAD patients, apoA-I partitions more into the water phase due to the higher free energy of apolipoprotein in the lipid phase with the negative correlations of dissociation degree with phospholipid and cholesterol levels in HDL particle. The retained CEC of lipid-free/lipid-poor apoA-I in CAD may be masked by the lower CEC of lipid-bound apoA-I due to the lower relative apolipoprotein level in the HDL lipid phase.

Contributions. A.D. Dergunov – concept of the study; writing text of the paper; supervision of the study; M.A. Popov – patients; V.B. Baserova – conducting experiments; preparation of figures and tables. All authors read and approved final version of the manuscript.

Funding. This study was financially supported by the State Budget.

Ethics approval and consent to participate. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or

national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. The project was approved by ethics committee of the M.F. Vladimirsky Moscow Regional Research and Clinical Institute MONIKI (protocol no. 12479/2019, February 17, 2019). All involved patients provided voluntary informed consent to participate in the study.

Conflict of interest. The authors of this work declare that they have no conflicts of interest.

Declaration of generative AI and AI-assisted technologies in the writing process. During the preparation of this work the authors did not use AI in the writing process.

Table 1. Relation of dissociation parameter D with pHDL lipids, gene expression, and ABCA1-mediated cholesterol efflux.

parameter	Variable	r (p) Control	r (p) CAD
pHDL level	pHDL-C	n.s.	-0.409 (0.000)
	pHDL-PL	n.s.	-0.406 (0.000)
	pHDL-C:apoA-I	n.s.	-0.372 (0.001)
Gene expression	*ABCA1*	0.252 (0.048)	n.s.
	CETP	0.272 (0.031)	n.s.
	A2M	-0.349 (0.006)	n.s.
	PRKACG	-0.313 (0.015)	n.s.
	PRKACB	n.s.	0.303 (0.008)
	ITGB3	0.260 (0.045)	n.s.
	TLR8	0.307 (0.015)	n.s.
	CUBN	n.s.	-0.287 (0.013)
	ALB	n.s.	-0.280 (0.015)
	OLR1	n.s.	0.238 (0.040)
HDL function	Chol efflux	n.s.	-0.326 (0.004)

Pearson correlation coefficients (r) with significance level in brackets (p) are given for control (n = 63) and CAD (n = 75) patients. n.s. – nonsignificant.

Table 2. Kinetics of ABCA1-mediated efflux to pHDL predenatured by urea.

patient	Urea, M	n	Vm, %/2h	Km, µg/ml	Vm/Km
Control	4.25	29	16.6 ± 3.5	0.38 ± 0.15	44.2 ± 19.7
	4.25	5	16.8 ± 3.5	0.69 ± 0,27	24.4 ± 10.9
	7.50	5	15.0 ± 1.9	0.74 ± 0.22	20.3 ± 6.6
CAD	4.25	31	16.5 ± 3.9	0.47 ± 0.20	35.0 ± 16.9
	4.25	4	18.9 ± 1.6	0.61 ± 0.11	30.8 ± 6.0
	7.50	4	17.7 ± 1.0	0.62 ± 0.08	28.3 ± 4.1

Mean ± SE values are given. n, patient number.

References

[1] A. Rohatgi, A. Khera, J.D. Berry, E.G. Givens, C.R. Ayers, K.E. Wedin, I.J. Neeland, I.S. Yuhanna, D.R. Rader, J.A. de Lemos, P.W. Shaul, HDL cholesterol efflux capacity and incident cardiovascular events. N. Engl. J Med 371 (2014) 2383-2393. https://doi.org/10.1056/NEJMoa1409065.

[2] A.D. Dergunov, V.B. Baserova, Different Pathways of Cellular Cholesterol Efflux. Cell Biochem.Biophys. 80 (2022) 471-481. https://doi.org/10.1007/s12013-022-01081-5.

[3] A.D. Dergunov, E.A. Garaeva, E.V. Savushkin, D.Y. Litvinov, Significance of Lipid-Free and Lipid-Associated ApoA-I in Cellular Cholesterol Efflux. Curr Protein Pept. Sci 18 (2017) 92-99. https://doi.org/10.2174/1389203717666160713150223.

[4] X.M. Du, M.J. Kim, L. Hou, G.W. le, M.J. Chapman, E.M. Van, L.K. Curtiss, J.R. Burnett, S.P. Cartland, C.M. Quinn, M. Kockx, A. Kontush, K.A. Rye, L. Kritharides, W. Jessup, HDL particle size is a critical determinant of ABCA1-mediated macrophage cellular cholesterol export. Circ. Res. 116 (2015) 1133-1142. https://doi.org/10.1161/CIRCRESAHA.116.305485.

[5] A.D. Dergunov, D.Y. Litvinov, E.V. Bazaeva, V.G. Dmitrieva, E.V. Nosova, A.V. Rozhkova, L.V. Dergunova, Relation of High-Density Lipoprotein Charge Heterogeneity, Cholesterol Efflux Capacity, and the Expression of High-Density Lipoprotein-Related Genes in Mononuclear Cells to the HDL-Cholesterol Level. Lipids 53 (2018) 979-991. https://doi.org/10.1002/lipd.12104.

[6] M.N. Nanjee, E.A. Brinton, Very small apolipoprotein A-I-containing particles from human plasma: isolation and quantification by high-performance size-exclusion chromatography. Clin.Chem. 46 (2000) 207-223. https://doi.org/10.1093/clinchem/46.2.207.

[7] A.D. Dergunov, D.Y. Litvinov, A.A. Malkov, V.B. Baserova, E.V. Nosova, L.V. Dergunova, Denaturation of human plasma high-density lipoproteins by urea studied by apolipoprotein A-I dissociation. Biochim Biophys Acta Mol Cell Biol Lipids 1866 (2021) 158814. https://doi.org/10.1016/j.bbalip.2020.158814.

[8] K. Okuhira, M. Tsujita, Y. Yamauchi, S. Abe-Dohmae, K. Kato, T. Handa, S. Yokoyama, Potential involvement of dissociated apoA-I in the ABCA1-dependent cellular lipid release by HDL. J Lipid Res. 45 (2004) 645-652. https://doi.org/10.1194/jlr.M300257-JLR200.

[9] M.S. Borja, K.F. Ng, A. Irwin, J. Hong, X. Wu, D. Isquith, X.Q. Zhao, B. Prazen, V. Gildengorin, M.N. Oda, T. Vaisar, HDL-apolipoprotein A-I exchange is independently associated with cholesterol efflux capacity. J Lipid Res. 56 (2015) 2002-2009. https://doi.org/10.1194/jlr.M059865.

[10] L. Schoch, S. Alcover, T. PadrΓi, S. Ben-Aicha, G. Mendieta, L. Badimon, G. Vilahur, Update of HDL in atherosclerotic cardiovascular disease. Clin Investig. Arterioscler. 35 (2023) 297-314. https://doi.org/10.1016/j.arteri.2023.10.002.

[11] A. Gangwar, S.S. Deodhar, S. Saldanha, O. Melander, F. Abbasi, R.W. Pearce, T.S. Collier, M.J. McPhaul, J.D. Furtado, F.M. Sacks, N.J. Merrill, J.E. McDermott, J.T. Melchior, A. Rohatgi, Proteomic Determinants of Variation in Cholesterol Efflux: Observations from the Dallas Heart Study. Int. J Mol. Sci. 24 (2023) 15526. https://doi.org/10.3390/ijms242115526.

[12] T. Josefs, K. Wouters, U.J.F. Tietge, W. Annema, R.P.F. Dullaart, T. Vaisar, I.C.W. Arts, C.J.H. van der Kallen, C.D.A. Stehouwer, C.G. Schalkwijk, I.J. Goldberg, E.A. Fisher, M.M.J. van Greevenbroek, High-density lipoprotein cholesterol efflux capacity is not associated with atherosclerosis and prevalence of cardiovascular outcome: The CODAM study. J Clin Lipidol. 14 (2020) 122-132. https://doi.org/10.1016/j.jacl.2019.10.012.

[13] K. Niisuke, Z. Kuklenyik, K.V. Horvath, M.S. Gardner, C.A. Toth, B.F. Asztalos, Composition-function analysis of HDL subpopulations: influence of lipid composition on particle functionality. J Lipid Res. 61 (2020) 306-315. https://doi.org/10.1194/jlr.RA119000258.

[14] T. Paavola, S. Kuusisto, M. Jauhiainen, S. Kakko, T. Kangas-Kontio, J. Metso, P. Soininen, M. Ala-Korpela, R. Bloigu, M.L. Hannuksela, M.J. Savolainen, T. Salonurmi, Impaired HDL2-mediated cholesterol efflux is associated with metabolic syndrome in families with early onset coronary heart disease and low HDL-cholesterol level. PLoS One 12 (2017) e0171993. https://doi.org/10.1371/journal.pone.0171993.

[15] B. Shao, C. Tang, A. Sinha, P.S. Mayer, G.D. Davenport, N. Brot, M.N. Oda, X.Q. Zhao, J.W. Heinecke, Humans with atherosclerosis have impaired ABCA1 cholesterol efflux and enhanced high-density lipoprotein oxidation by myeloperoxidase. Circ.Res. 114 (2014) 1733-1742. https://doi.org/10.1161/CIRCRESAHA.114.303454.

[16] C.P.J. Talbot, J. Plat, A. Ritsch, R.P. Mensink, Determinants of cholesterol efflux capacity in humans. Prog.Lipid Res. 69 (2018) 21-32. https://doi.org/10.1016/j.plipres.2017.12.001.

[17] A.V. Khera, M. Cuchel, M. Llera-Moya, A. Rodrigues, M.F. Burke, K. Jafri, B.C. French, J.A. Phillips, M.L. Mucksavage, R.L. Wilensky, E.R. Mohler, G.H. Rothblat, D.J. Rader, Cholesterol efflux capacity, high-density lipoprotein function, and atherosclerosis. N. Engl. J Med 364 (2011) 127-135. https://doi.org/10.1056/NEJMoa1001689.

[18] B.F. Asztalos, K.V. Horvath, E.J. Schaefer, High-Density Lipoprotein Particles, Cell-Cholesterol Efflux, and Coronary Heart Disease Risk. Arterioscler. Thromb. Vasc. Biol. 38 (2018) 2007-2015. https://doi.org/10.1161/ATVBAHA.118.311117.

[19] B.F. Asztalos, T.H. Hauser, A.B. Goldfine, F.K. Welty, K.V. Horvath, E.J. Schaefer, The role of HDL- and non-HDL-related parameters in cell-cholesterol efflux capacity. Atherosclerosis 345 (2022) 1-6. https://doi.org/10.1016/j.atherosclerosis.2022.02.012.

[20] R.K. Mutharasan, C.S. Thaxton, J. Berry, M.L. Daviglus, C. Yuan, J. Sun, C. Ayers, D.M. Lloyd-Jones, J.T. Wilkins, HDL efflux capacity, HDL particle size, and high-risk carotid atherosclerosis in a cohort of asymptomatic older adults: the Chicago Healthy Aging Study. J Lipid Res. 58 (2017) 600-606. https://doi.org/10.1194/jlr.P069039.

[21] A.D. Dergunov, E.V. Nosova, A.V. Rozhkova, M.A. Vinogradina, V.B. Baserova, M.A. Popov, S.A. Limborska, L.V. Dergunova, HDL cholesterol-associated shifts in the expression of preselected genes reveal both pro-atherogenic and atheroprotective effects of HDL in coronary artery disease. Frontiers in Biosciences Landmark 29 (2024) 396. https://doi.org/10.31083/j.fbl2911396.

[22] H.J. Pownall, B.D. Hosken, B.K. Gillard, C.L. Higgins, H.Y. Lin, J.B. Massey, Speciation of human plasma high-density lipoprotein (HDL): HDL stability and apolipoprotein A-I partitioning. Biochemistry 46 (2007) 7449-7459. https://doi.org/10.1021/bi700496w.

[23] A.D. Dergunov, E.V. Nosova, A.V. Rozhkova, M.A. Vinogradina, V.B. Baserova, M.A. Popov, S.A. Limborska, L.V. Dergunova, Differential Expression of Subsets of Genes

Related to HDL Metabolism and Atherogenesis in the Peripheral Blood in Coronary Artery Disease. Curr. Issues Mol. Biol. 45 (2023) 6823-6841. https://doi.org/10.3390/cimb45080431.

[24]   A. Dziedzic, E. Miller, M. Bijak, L. Przyslo, J. Saluk-Bijak, Increased Pro-Thrombotic Platelet Activity Associated with Thrombin/PAR1-Dependent Pathway Disorder in Patients with Secondary Progressive Multiple Sclerosis. Int. J Mol. Sci. 21 (2020) 7722. https://doi.org/10.3390/ijms21207722.

[25]   V. Lopez-Olmos, M. Luna-Luna, E. CarreΓin-Torres, H. Gonzalez-Pacheco, R. Bautista-Perez, R. Posadas-Sanchez, J.M. Fragoso, G. Vargas-Alarcon, O. Perez-Mendez, Trp Fluorescence Redshift during HDL Apolipoprotein Denaturation Is Increased in Patients with Coronary Syndrome in Acute Phase: A New Assay to Evaluate HDL Stability. Int. J Mol. Sci. 22 (2021) 7819. https://doi.org/10.3390/ijms22157819.

[26]   A.D. Dergunov, Sequence-specific apolipoprotein A-I effects on lecithin:cholesterol acyltransferase activity. Mol. Cell Biochem. 378 (2013) 283-290. https://doi.org/10.1007/s11010-013-1619-5.

[27]   D. Handa, H. Kimura, T. Oka, Y. Takechi, K. Okuhira, M.C. Phillips, H. Saito, Kinetic and thermodynamic analyses of spontaneous exchange between high-density lipoprotein-bound and lipid-free apolipoprotein A-I. Biochemistry 54 (2015) 1123-1131. https://doi.org/10.1021/bi501345j.

[28]   N. Matsuzaki, T. Handa, M. Nakano, Kinetic and Thermodynamic Analysis of Cholesterol Transfer between Phospholipid Vesicles and Nanodiscs. J. Phys. Chem. B 119 (2015) 9764-9771. https://doi.org/10.1021/acs.jpcb.5b03682.

[29]   C. Talussot, G. Ponsin, Cholesterol-induced alteration of apolipoprotein A-I conformation in reassembled high density lipoprotein. Biochimie 73 (1991) 1173-1178. https://doi.org/10.1016/0300-9084(91)90001-h.

[30]   M.A. Clay, H.H. Newnham, T.M. Forte, P.I. Barter, Cholesteryl ester transfer protein and hepatic lipase activity promote shedding of apo A-I from HDL and subsequent formation of discoidal HDL. Biochim. Biophys. Acta 1124 (1992) 52-58. https://doi.org/10.1016/0005-2760(92)90125-f.

[31] H.Q. Liang, K.A. Rye, P.J. Barter, Dissociation of lipid-free apolipoprotein A-I from high density lipoproteins. J Lipid Res. 35 (1994) 1187-1199. https://doi.org/10.1016/S0022-2275(20)39962-4.

[32]   S. Jayaraman, J.L. Sanchez-Quesada, O. Gursky, Triglyceride increase in the core of high-density lipoproteins augments apolipoprotein dissociation from the surface: Potential implications for treatment of apolipoprotein deposition diseases. Biochim Biophys Acta 1863 (2017) 200-210. https://doi.org/10.1016/j.bbadis.2016.10.010.

[33]   O. Aseem, B.T. Smith, M.A. Cooley, B.A. Wilkerson, K.M. Argraves, A.T. Remaley, W.S. Argraves, Cubilin maintains blood levels of HDL and albumin. J Am. Soc. Nephrol. 25 (2014) 1028-1036. https://doi.org/10.1681/ASN.2013060671.

[34]   E.J. Schaefer, B.F. Asztalos, T. Vaisar, M.R. Diffenderfer, H.B. Brewer, Jr., E.O. Stock, J.P. Kane, High Density Lipoprotein Particle Composition, Functionality, Deficiency, and Atherosclerotic Cardiovascular Disease Risk: A Review. Curr. Atheroscler. Rep. 27 (2025) 62. https://doi.org/10.1007/s11883-025-01308-9.

Figure 1. ApoA-I distribution between water and lipid phases of pHDL predenatured by 4.25 M urea. (A) pHDL from control patients (n = 59), (B) pHDL from CAD patients (n = 73). The 95% prediction bands are given as thin dashed lines. The Y-intercepts passed through zero according to the distribution model. Slopes were 0.239 ± 0.012 ($R^2 = 0.874$) for control and 0.368 ± 0.020 ($R^2 = 0.822$) for CAD. Slope differences were significant with $p = 0.000$.

Figure 2. The kinetics of ABCA1-mediated efflux to pHDL predenatured by 4.25 M urea. (A) pHDL from control patients (n = 29), (B) pHDL from CAD patients (n = 31). The 95% prediction bands are given as thin dashed lines.