Design and analysis of synthetic carbon fixation pathways based on novel
enzymatic reactions

Vittorio Rainaldi[1,2], Sarah, D'Adamo[2], Nico J. Claassens[1]

[1]Laboratory of Microbiology, Wageningen University, Stippeneng 4, 6708 WE,
Wageningen, the Netherlands
[2]Bioprocess Engineering, Wageningen University, Droevendaalsesteeg 1, 6708 PB,
Wageningen, the Netherlands
9
10 **Corresponding authors:** NJC nico.claassens@wur.nl and VR vittorio.rainaldi@wur.nl

12 Abstract

Biological carbon fixation is currently limited to seven naturally occurring pathways,
each with its own limitations and constraints. In recent years, computational analyses of
known biochemical reaction networks have identified dozens of theoretical carbon
fixation pathways, some of which may have the potential to outperform their natural
counterparts. This mix and match approach, however, cannot account for those
reactions that have not been reported to occur in nature, which heavily limits the
possible solution space. Here, we use a bioretrosynthetic approach coupled with expert
biochemical knowledge to identify eight novel pathways that leverage enzyme
promiscuity and the latent biochemical reaction space. We analyze the thermodynamic,
stoichiometric, and kinetic parameters of these pathways and compare them to the
ubiquitous Calvin-Benson-Bassham cycle and previously proposed synthetic CO_2
fixation cycles, highlighting advantages and disadvantages. This work highlights the
need for enzyme engineering and design in the quest for efficient biological carbon
fixation.

28 Introduction

The Calvin-Benson-Bassham (CBB) cycle is responsible for over 99% of net primary
productivity on earth[1]. Despite its dominance, it is a relatively inefficient pathway,
requiring a high energy investment in terms of ATP to fix carbon. Furthermore, its key
enzyme, rubisco, is rather slow when compared to other naturally occurring
carboxylases and it presents a detrimental side activity with oxygen[2,3], which results in

additional energy costs while still leading to partial loss of carbon in a process called
photorespiration[4].

Many synthetic carbon fixation pathways have been designed so far by using a mix and
match approach that combines enzymes from disparate sources with fast and efficient
carboxylases to generate theoretical cycles that have not been described in nature[5–7].
Nevertheless, a recent quantitative analysis of natural and synthetic carbon fixation
pathways by Löwe and Kremling showed that most pathways are in fact inferior to the
CBB cycle or only marginally more efficient[7], likely making the massive engineering
efforts required to implement them unjustified.

Almost all of the current synthetic pathway designs, however, are limited by the fact that
they only use known enzymatic activities from databases such as KEGG, which vastly
limits the possible solution space. Erb, Jones, and Bar-Even have previously put
forward a proposal for categorizing pathways based on five levels, with naturally
occurring pathways in level one, and synthetic pathways based on new-to-nature
enzyme mechanisms in level five[8]. Despite recent progress, designing enzymes with
novel mechanisms remains highly challenging[9,10]. For this reason, we focused on level
four pathways, namely those that are based on unreported promiscuous enzyme
activities and reactions from the latent biochemical space. In this study, we use a
combination of bioretrosynthesis and expert biochemical knowledge to generate novel
pathway designs and analyze their thermodynamic[11], stoichiometric[12], and kinetic[13]
parameters, comparing them to the CBB cycle and highlighting advantages and
disadvantages. All these pathways require engineering of at least one enzyme to
change its substrate or cofactor specificity or other relevant parameters such as oxygen
tolerance. Oxygen tolerance is particularly important in the context of engineering CO_2
fixation in plants and algae, since these organisms produce oxygen as part of
photosynthesis, which would lead to the rapid inactivation of oxygen-sensitive enzymes.
However, oxygen tolerance is also important in hydrogen-based microbial bioproduction
from CO_2, as CO_2 fixation and bioproduction pathways require ATP, which is generated
from O_2-dependent respiration.

Overall, this paper provides a novel approach to pathway design and highlights the
need for enzyme engineering to achieve efficient carbon fixation, while offering
suggestions regarding possible starting points to engineer the proposed pathways.

Figure 1: The β-glutamate cycle. 1, succinyl-CoA synthetase; 2, succinyl-CoA reductase; 3, succinate semialdehyde dehydrogenase; 4, 4-hydroxybutyrate CoA transferase; 5, 4-hydroxybutyryl-CoA dehydratase; 6, crotonyl-CoA carboxylase; 7, glutaconate CoA transferase; 8, glutaconate ammonia lyase; 9, glutamate 2,3-aminomutase; 10, glutamate dehydrogenase; 11, 2-oxoglutarate carboxylase; 12, isocitrate dehydrogenase; 13, isocitrate lyase. In this figure and following ones the enzymes marked in red have never been reported in the literature. Bioprospecting or engineering efforts will likely be required to establish their activities.

Results

The β-glutamate cycle

We recently proposed and showed partial in vivo implementation of a new family of
carbon fixation pathways called the reductive methylaspartate cycles using the standard
mix and match approach[14]. While these cycles are promising, they suffer from two main
issues. First, the enzyme crotonyl-CoA carboxylase/reductase (CCR) is relatively
inefficient because the NADPH invested for its reductive carboxylation activity is later
only partially recovered in the form of a reduced quinone, leading to a loss of energy.
Secondly, these cycles require vitamin B_{12} as a cofactor for two enzymatic reactions.
Vitamin B_{12} is a highly complex cofactor which cannot be synthesized by most
organisms, importantly including all higher plants and many algae[15], thus limiting the
applicability of pathways containing B_{12}-dependent enzymes mostly to bacteria.

To overcome both of these issues, we designed a variant of the rMASP cycles that we
name the β-glutamate cycle based on its characteristic metabolite. According to our
computational analysis, the β-glutamate cycle shows favorable parameters when
compared to the CBB, including higher MDF, higher theoretical biomass yield, and
comparable pathway activity. These advantages increase in ambient CO_2 environments,
with a 40% increase in yield at 20% rubisco oxygenation (Table S1).

The β-glutamate cycle does not use CCR but instead depends on the promiscuous
activity of methylcrotonyl-CoA gamma carboxylase (MCC)[16,17], which we will refer to as
crotonyl-CoA gamma carboxylase (CCC). This biotin-dependent carboxylase is involved
in leucine degradation across diverse phyla including bacteria[18,19] and humans. Its
promiscuous side activity on the slightly smaller crotonyl-CoA has been reported in the
literature with about 90% of the reaction rate when compared to the native substrate.
Since the carboxylation happens in gamma, as opposed to alpha, it produces
glutaconyl-CoA instead of ethylmalonyl-CoA, thus obviating the need for a B_{12}
dependent mutase to "straighten" the carbon skeleton.

The cycle proceeds with the removal of the CoA from glutaconyl-CoA via a CoA
transferase or CoA ligase to yield glutaconate, which is further aminated to 3-
aminoglutarate, also known as β-glutamate, which gives its name to the cycle. This
amination reaction could be promiscuously performed by aspartate/methylaspartate
ammonia lyase[20], though most likely it would require engineering to expand the
substrate binding pocket to accommodate one extra carbon. An alternative to this route
could be the direct amination of glutaconyl-CoA, for example by a promiscuous β-alanyl-
CoA ammonia lyase, an enzyme found in anaerobic bacteria and used for β-alanine
fermentation[21]. The resulting beta-glutamyl-CoA would once again require a CoA
transferase or CoA ligase to remove the CoA and yield β-glutamate.

At this point, β-glutamate must undergo isomerization to produce glutamate. This
reaction is catalyzed by the naturally occurring enzyme glutamate 2,3-aminomutase[22],
which is present in bacteria and archaea, respectively for glutamate fermentation and

for the production of beta-glutamate as an osmolyte[23]. This reaction, however, is highly
challenging, requiring a radical mechanism based on S-adenosyl-methionine (SAM),
making the enzyme oxygen-sensitive. Nevertheless, there are oxygen-tolerant variants
reported to catalyze the same reaction aerobically, though using lysine as a substrate
as opposed to glutamate[24–26]. Not much is known about this enzyme class, so it is
difficult to know whether it would be easier to engineer oxygen-tolerance in a glutamate
aminomutase or change substrate specificity in a lysine aminomutase. It must be noted,
however, that while there are many examples of oxygen-tolerant radical SAM
enzymes[27,28], e.g. *bioB* and *lipA* from *E. coli*, there seems to be a tradeoff between
activity and oxygen tolerance, with oxygen-sensitive variants showing higher activity.
For this reason, it may not be possible to obtain an oxygen-tolerant aminomutase with
comparable kinetic parameters to oxygen-sensitive isoforms.

Figure 2: The 3-hydroxyglutaryl-CoA lyase cycle. 1, methylmalonyl-CoA mutase; 2-6 are as in figure 1; 7, glutaconyl-CoA hydratase; 8, 3-hydroxyglutaryl-CoA lyase; 9, malonate semialdehyde reductase; 10, propionyl-CoA synthase; 11, propionyl-CoA carboxylase; 12, methylmalonyl-CoA epimerase.

The 3-hydroxyglutaryl-CoA lyase (3HGL) cycle

In the leucine degradation pathway, the intermediate methylglutaconyl-CoA, product of methylcrotonyl-CoA gamma carboxylase (MCC), undergoes hydration to 3-hydroxy-3-methylglutaconyl-CoA and is subsequently split to acetoacetate and acetyl-CoA. We reasoned that the product of CCC, glutaconyl-CoA, can also undergo hydration rather than amination, yielding 3-hydroxyglutaryl-CoA. This promiscuous reaction is indeed reported to occur efficiently[29]. This metabolite could then be split by 3HMG-CoA lyase to acetyl-CoA and malonate semialdehyde, with the latter being reduced to propionyl-CoA and subsequently carboxylated. The resulting succinyl-CoA can then be reduced to crotonyl-CoA to restart the cycle. More pathways based on CCC and glutaconyl-CoA hydration have previously been reported[5,30], however the authors describe a hydration in C2 rather than C3, which produces 2-hydroxyglutaryl-CoA. Unfortunately, the enzyme that catalyzes this reaction is based on a ketyl radical mechanism and hence extremely oxygen sensitive[31,32]. To the best of our knowledge there has never been a report of ketyl radical enzymes operating in aerobic conditions, making these proposed routes infeasible for aerobic implementation, while their anaerobic implementation would require an external electron acceptor, making the pathways biotechnologically not relevant. The 3HGL cycle, on the other hand, does not use any oxygen-sensitive enzymes, since hydration in C3 occurs easily due to the differences in the pKa between the C2 protons and the C3 protons[33].

This pathway is comparable to the CBB cycle in terms of rate of CO_2 fixation and yield at high CO_2 concentration and it is predicted to have about 27% higher yield at ambient CO_2, with roughly double the activity (Figure 9). Moreover, the addition of acetyl-CoA carboxylase and malonyl-CoA reductase makes the cycle autocatalytic, much like the CBB cycle and reverse tricarboxylic acid (rTCA) cycle. Autocatalytic cycles are expected to be more stable and resilient to metabolic perturbations[34], potentially simplifying engineering efforts. Nevertheless, the strict B_{12} requirement for methylmalonyl-CoA mutase limits its application to bacteria and some algae[15].

Figure 3: The 3-hydroxyglutaryl-CoA mutase cycle. 1-6 are as in figure 1; 7, glutaconyl-
CoA hydratase; 8, 3-hydroxyglutaryl-CoA mutase; 9, citramalyl-CoA transferase; 10,
citramalyl-CoA lyase; 11, pyruvate carboxylase; 12, malate dehydrogenase; 13,
fumarase; 14, fumarate reductase.

The 3-hydroxyglutaryl-CoA mutase (3HGM) cycle

Alternatively, 3-hydroxyglutaryl-CoA could be isomerized in a similar reaction to the
naturally occurring interconversion of 3-hydroxybutyryl-CoA and 2-hydroxyisobutyryl-
CoA[35,36], catalyzed by a B_{12}-dependent mutase. It should be noted, however, that the
resulting molecule, namely citramalyl-CoA, would carry the CoA on C1 rather than C4,
thus requiring a specialized enzyme to swap the CoA, similarly to mesaconyl-CoA
transferase[37]. This is necessary because the subsequent citramalyl-CoA lyase reaction
is only possible if the CoA is in C4. This same enzyme was recently engineered to

accept 3-hydroxypropionyl-CoA and catalyze its interconversion to lactyl-CoA[38] as part
of the previously proposed MOG-LCM cycle[5], which we included in our comparative
computational analysis along with the MOG-BAAM cycle from the same paper (Figure
9). This engineering effort produced an enzyme with a poor k_{cat} of 0.03, which is around
an order of magnitude lower than the activity on its original substrate. Even this native
activity is rather low, and it is unclear whether it can be improved. A computational
analysis of the promiscuous activity of B_{12}-dependent glutamate mutase on the
alternative substrate 2-hydroxyglutarate (yielding threo-methylmalate) identified a
thermodynamic bottleneck in the reaction due to the formation of a glycolyl radical
instead of the usual glycyl radical which is formed in the glutamate mutase reaction. The
presence of a hydroxyl group in the glycolyl radical is thought to destabilize the radical
leading to an increase in activation energy and a proportional decrease in k_{cat} of about
two orders of magnitude (5 s^{-1} to 0.05 s^{-1})[39]. It is currently unclear whether this issue can
be solved and if it is related to glutamate mutase or general to all B_{12}-dependent
mutases. More research will be needed to improve our understanding of the chemistry
catalyzed by this complex cofactor.

The α-ketobutyrate carboxylase (ABC) cycles

While pathways based on crotonyl-CoA carboxylation use either the condensation of
two acetyl-CoA molecules or the reduction of succinyl-CoA, a different strategy can be
envisioned based on the threonine biosynthesis pathway. We recently reported the in
vivo implementation of the Serine Threonine Cycle (STC) for full formatotrophic growth

Figure 4: The *α*-ketobutyrate carboxylase-glutamate mutase cycle. 1, aspartate aminotransferase; 2, aspartate kinase; 3, aspartate semialdehyde dehydrogenase; 4, homoserine dehydrogenase; 5, homoserine kinase; 6, threonine synthase; 7, threonine deaminase; 8, *α*-ketobutyrate carboxylase; 9, methylaspartate aminotransferase; 10, glutamate mutase; 11, glutamate dehydrogenase; 12, 2-oxoglutarate carboxylase; 13, oxalosuccinate reductase; 14, aconitase; 15, ATP citrate lyase

coli[40]. The STC relies on the condensation of glycine derived from threonine cleavage with externally supplied formate and an additional carboxylation reaction based on pyruvate carboxylase to regenerate the four-carbon molecule aspartate, which is the precursor for threonine. We reasoned that threonine could instead undergo deamination to produce α-ketobutyrate, which is an α-keto acid highly similar to pyruvate and thus amenable to carboxylation via an engineered pyruvate carboxylase. While we could find

no reports of a biotin-dependent α-ketobutyrate carboxylase in the literature, there is at
least one study showing partial promiscuous activity of pyruvate carboxylase on α-
ketobutyrate[41]. This limited activity could be used as the basis for mutagenesis or
rational engineering in an approach similar to the one used to change the substrate
specificity of propionyl-CoA carboxylase to glycolyl-CoA[42].

Figure 5: The *α*-ketobutyrate carboxylase-ciramalyl-CoA lyase cycle. 1-9 are as in
figure 4. 10, methylaspartate ammonia lyase; 11, mesaconase; 12, citramalate CoA
transferase; 13, citramalyl-CoA lyase; 14, pyruvate carboxylase.

Figure 6: The *α*-ketobutyrate carboxylase-propionyl-CoA carboxylase cycle. 1-8 are as in figures 4 and 5. 9, methylmalate dehydrogenase, 10, methylmalate-CoA transferase; 11, methylmalyl-CoA lyase; 12, propionyl-CoA carboxylase; 13, methylmalonyl-CoA epimerase; 14, methylmalonyl-CoA mutase; 15, succinyl-CoA synthetase; 16, succinate dehydrogenase; 17, fumarase; 18, malate dehydrogenase.

While pyruvate carboxylation produces oxaloacetate, α-ketobutyrate carboxylation would produce 3-methyloxaloacetate (also known as 2-methyl-3-oxosuccinate or oxalopropionate). This compound can undergo reductive amination or transamination to methylaspartate[43], reduction to 3-methylmalate (erythro or threo), or condensation with acetyl-CoA to methylcitrate, though we did not investigate the latter option further. In the first case, methylaspartate can be isomerized to glutamate via a B_{12}-dependent glutamate mutase (GM) and then proceed via the same steps as the previously described rMASP cycles[14]. This variant, which we name ABC-GM cycle, compares favorably to the CBB at ambient CO_2 in terms of biomass yield, though its strict

requirement for B_{12} limits its applicability to microorganisms. To bypass this need, we

propose another variant where methylaspartate is deaminated to mesaconate and

hydrated to citramalate, which is then activated to citramalyl-CoA and split to pyruvate

and acetyl-CoA via a citramalyl-CoA lyase (CMCL). This variant, called ABC-CMCL

cycle is slightly less efficient in terms of biomass yield but has a higher predicted

pathway activity. A third option could use of the promiscuous activity of malate

dehydrogenase (MDH) for the reduction of methyloxaloacetate to erythro-methylmalate,

which can further be activated to a CoA ester and split to propionyl-CoA and glyoxylate,

however this variant known as ABC-MDH, also requires B_{12} for the isomerization of

methylmalonyl-CoA following the carboxylation of propionyl-CoA. Notably, the ABC

cycles only require one enzyme to be engineered, and their overlap with the STC shows

that they could be engineered relatively easily by applying some of the lessons learned

during the STC engineering and evolution[40]. Nevertheless, our theoretical analysis

shows that threonine biosynthesis may be limiting the overall activity of the pathway,

especially through the bottleneck enzyme homoserine dehydrogenase. It is unclear

whether faster variants of this enzyme can be found or engineered, and if the

thermodynamic constraints of this biosynthetic route can be overcome, e.g. by

optimizing enzymes and/or metabolite concentration. More research will be needed to

ascertain whether this theoretical prediction will be verified experimentally.

Figure 7: The Ribulose-5-phosphate carboxylase-Entner-Doudoroff (RED) cycle. 2PG,
2-phosphoglycerate; 3K6PG, 3-keto-6-phosphogluconate; 3PG, 3-phosphoglycerate;
6PG, 6-phosphogluconate; BPG, 1,3-bisphosphoglycerate; DHAP, dihydroxyacetone
phosphate; E4P, erythrose-4-phosphate; F6P, fructose-6-phosphate; FBP, fructose-1,6-
bisphosphate; GAP, glyceraldehyde-3-phosphate; KDPG, 2-keto-3-deoxy-
phosphogluconate, PEP, phosphoenolpyruvate; R5P, ribose-5-phosphate; Ru5P,
ribulose-5-phosphate; S7P, sedoheptulose-7-phosphate; SBP, sedoheptulose-1,7-
bisphosphate; Xu5P, xylulose-5-phosphate; 1, ribulose-5-phosphate carboxylase; 2, 6-
phosphogluconate dehydrogenase; 3, Entner-Doudoroff dehydratase; 4, Entner-
Doudoroff aldolase; 5, PEP synthetase; 6, enolase; 7, phosphoglycerate mutase, 8,

268   phosphoglycerate kinase; 9, glyceraldehyde-3-phosphate dehydrogenase; 10, triose
269   phosphate isomerase; 11, fructose bisphosphate aldolase; 12, fructose bisphosphatase;
270   13, transketolase; 14, sedoheptulose bisphosphate aldolase; 15, sedoheptulose
271   bisphosphatase; 16, ribulose-5-phosphate epimerase; 17, ribulose-5-phosphate
272   isomerase

The Ribulose-5-phosphate carboxylase-Entner-Doudoroff (RED) cycle

While all the cycles presented so far have some advantages over the CBB, it should be
noted that that they are all based on completely different architectures, thus requiring
extensive engineering efforts to establish in those organisms, including algae and plants,
that already use the CBB cycle. For this reason, we set out to establish a carbon fixation
cycle that is similar in architecture to the CBB but presents more favorable
characteristics in terms of biomass yield and pathway kinetic. The GND-Entner-
Doudoroff (GED) cycle has been previously proposed as a promising alternative to the
CBB based on natural enzymes[44], however it is severely limited by the low activity and
affinity for CO_2 of its signature enzyme, 6-phosphogluconate dehydrogenase (GND).
This enzyme normally catalyzes the oxidative decarboxylation of 6-phosphogluconate,
coupling the highly favorable decarboxylation reaction to the unfavorable generation of
NADPH. This same mechanism can be found in at least two other central metabolic
enzymes, namely the malic enzyme, which produces pyruvate and NADPH from malate,
and isocitrate dehydrogenase, which produces α-ketoglutarate and NADPH from
isocitrate.

Notably, in both of these cases there is a biotin-dependent carboxylase to overcome the
unfavorable thermodynamics and kinetics of this half-reaction, which is then coupled to
a dehydrogenase/reductase. Pyruvate is carboxylated to oxaloacetate which is then
reduced to malate, while α-ketoglutarate is carboxylated to oxalosuccinate and then
reduced to isocitrate[45–47]. A similar idea could be applied to the product of GND,
ribulose-5-phosphate. A biotin-dependent carboxylase acting on this substrate would
generate 3-keto-6-phosphogluconate, which would then be reduced to 6-
phosphogluconate. For this latter reaction GND itself could be used, provided the
residues involved in the decarboxylation half-reaction are inactivated. To the best of our
knowledge, there is no reported enzyme taking a similar substrate for the carboxylation
reaction, meaning that an extensive engineering campaign will be necessary to change
the substrate specificity. This is probably beyond what is achievable with current
strategies based on mutagenesis and selection, however it may become possible in the
future to apply novel enzyme design approaches to redesign the substrate pocket. The
RED cycle is insensitive to the presence of oxygen, does not require CoA or exotic
cofactors, it is favorable in terms of efficiency and kinetics at ambient CO_2, and it has a
high degree of overlap with the CBB cycle, making it a good candidate for
implementation in algae and plants, though it should be noted that these organisms do

not express lower glycolysis enzymes in their chloroplasts, which could increase the
engineering complexity.

Figure 8: The Malyl-CoA-Tartronyl-CoA (MCTC) cycle. 1, glycolate-CoA ligase; 2,
glycolyl-CoA carboxylase; 3, tartronyl-CoA reductase; 4, tartronate semialdehyde
reductase; 5, glycerate kinase; 6, enolase; 7, PEP carboxylase; 8, malate
dehydrogenase; 9, malate CoA transferase; 10, malyl-CoA lyase; 11, glyoxylate
reductase.

The Malyl-CoA-Tartronyl-CoA (MCTC) pathway

Scheffen et al. recently reported engineered a biotin-dependent propionyl-CoA
carboxylase to accept the smaller substrate glycolyl-CoA[42]. The resulting enzyme has
been named glycolyl-CoA carboxylase (GCC). This carboxylation reaction produces
tartronyl-CoA, which is further reduced by an engineered malonyl-CoA reductase to
tartronate semialdehyde, which is itself reduced to glycerate, a compound that can

easily be assimilated into central metabolism. The resulting pathway, known as the
tartronyl-CoA (TaCo) pathway, was conceived as a photorespiration bypass, taking 2-
phosphoglycolate derived from the rubisco oxygenation side reaction and recycling it to
3-phosphoglycerate while not only preventing loss of carbon but fixing carbon instead.
While this approach is promising, it is currently hampered by the inefficiency of glycolyl-
CoA carboxylase, which hydrolyzes more than 5 ATP per carboxylation reaction, as
opposed to 1 ATP for PCC and similar biotin-dependent enzymes. This could be
attributed to the difference between propionyl-CoA and glycolyl-CoA as substrates: the
latter is not only smaller but contains a hydroxyl group. The presence of a hydroxyl
group could destabilize the enolate intermediate required for carboxylation, leading to
futile cycles where CO_2 is released into the active site but is unable to react with the
substrate. In a subsequent work, the enzyme was further engineered by using a
machine-learning guided approach, obtaining improved variants with reduced futile ATP
hydrolysis, albeit at the expense of lower overall catalytic activity[48]. Whether this issue
can be fully solved remains an open question.

We reasoned that the TaCo pathway could also serve as the basis for a synthetic CO_2
fixation cycle (Figure 8) where the product, glycerate, goes through lower glycolysis and
anaplerosis to malate, which is activated to malyl-CoA and split to acetyl-CoA, which is
the end product of this cycle, and glyoxylate, which can be reduced to glycolate to
restart the cycle. This pathway has the advantage of being relatively compact, requiring
only eleven genes for its implementation, all of which have been tested in vitro,
potentially making it easier to engineer, especially in less genetically accessible
organisms. Assuming an improved GCC can be engineered, we calculated that this
cycle is about 20% more efficient than the CBB cycle at ambient CO_2, with comparable
pathway activity. We note that while writing this manuscript, this pathway has also been
proposed in another work, in which it has been named the LATCH pathway[49].

Coupling the CO_2 reduction (CORE) cycle to formatotrophic pathways

An interesting alternative to CO_2 fixation cycles was recently proposed by Satanowski et
al., who theorized a number of oxygen tolerant CO_2 reduction (CORE) cycles and
implemented one of them in *E. coli*[50]. The experimentally validated CORE variant relies
on the promiscuous activity of a β-keto-acid cleavage enzyme (BKACE) and yields one
molecule of formate from one molecule of bicarbonate with the input of one NADPH and
1-3 ATP. While the CORE cycle was originally proposed as a photorespiration bypass,
the resulting formate could be fully coupled to growth via a formatotrophic pathway such
as the serine cycle. We have modeled the resulting pathway to compare it with our
proposed CO_2 fixation cycles (Figure 9). Based on our results, the CORE+Serine cycle
shows a comparable yield to the CBB in high CO_2 and a roughly 20% higher yield in
ambient CO_2. Assuming the activity of the limiting enzyme, BKACE, can be improved,
the CORE+Serine cycle could be a highly relevant alternative to conventional CO_2

fixation cycles, since it is oxygen-tolerant, does not require B_{12}, and mostly relies on
reactions that are natively present in plants, or that have been shown to be functional
after heterologous expression.

Figure 9: Plot of theoretical biomass yield vs predicted pathway activity for each
pathway presented in this paper and previously proposed pathways. This figure
includes only pathways that require enzyme engineering for their implementation (other
than the CBB cycle). For additional comparisons to previously proposed mix-and-match
pathways see figure S1.

Discussion

In this paper, we proposed eight novel carbon fixation pathways based on non-natural
enzymatic reactions. We analysed the pathways based on thermodynamic driving force,
predicted biomass yield, and predicted pathway activity. This last parameter is
computed based on the measured or estimated k_{cat} and K_M values for the enzymes.
Unfortunately, the available enzymatic data in databases such as BRENDA is both
limited and noisy, since different authors perform enzymatic assays in wildly different
conditions in terms of pH, buffers, temperatures, etc. We carefully curated the chosen
values, also based on the previous work by Lowe and Kremling, however a relatively
large fraction of the enzymes was only measured in single studies, reducing their
reliability. Where multiple data points are available, it is not unusual to see a variation of
one order of magnitude or more in reported k_{cat} and K_M.

Furthermore, in vitro kinetics measurements, even when performed properly, are based
on a purified enzyme in a dilute buffer, which is a completely different scenario
compared to the densely packed cytosol of living organisms, where thousands of
enzymes and metabolites are present. Recent work has shown k_{cat} measurements in
vivo based on different growth conditions and proteomics data[51]. These results show
that while many in vitro measurements are also accurate in vivo, there are several
outliers even for well characterized enzymes in central metabolism, with a relatively
weak correlation R^2 of 0.62. All these problems are compounded when it comes to
relatively understudied enzymes, and especially their promiscuous activities. Finally, the
kinetic parameters of not yet existing enzymes are obviously impossible to obtain, and
we have simply taken representative values for similar enzymes on related substrates.

ECM measurements assume an optimal distribution of metabolites and enzymes, and
do not take into account enzyme degradation, allosteric inhibition, competitive inhibition,
and regulation. They also do not consider all the enzymes that are necessary for cell
maintenance and growth, assuming all flux is going via the pathway of interest to
generate the desired product. In practice, all these factors mean that the predicted
pathway activity is at best an upper bound for the real pathway activity in a real-life
scenario. Taking the CBB into account, measured values for CO_2 fixation rates in
exponentially growing bacteria are in the order of 3-9 nmol/min/mg (total) protein[52]. This
is a two order of magnitude difference below our computed value of around 0.5
μmol/min/mg (pathway) protein. This discrepancy is partially explained when taking into
account the fact that CBB enzymes account for about 10% of total protein [53]. The
predicted pathway activity for the CBB, then, is still one order of magnitude above
measured values, despite the fact that all of its the enzymes are well-characterized in
dozens of studies.

For the other pathways, greater uncertainty is expected due to the limited reliability of
the available data. It is unclear whether this uncertainty leads to an overestimate or an
underestimate of the total pathway activity, however it is likely that for most of the
candidate enzymes, kinetically superior isoforms could be found through a screening
process. This points to the fact that pathways other than the CBB may be
underestimated in our analysis. Further research will be needed to determine real life
(i.e. in vivo) kinetic parameters for several enzymes, leading to better estimates.

Considering all of this, our proposed pathways are especially promising for potential
future applications in terms of yield, as they can outcompete the CBB cycle, even at
high CO_2, especially the β-glutamate cycle (10% or 40% higher yield than CBB cycle at
high and ambient CO_2 respectively), and also the 3HGL, 3HGM, ABC-GM and MOG-
BAAM may outperform the CBB cycle slightly at high CO_2 and by >30% in yield at
ambient CO_2. However, none of them is predicted so far to outperform the CBB in terms
of pathway activity, at least in the high CO_2 scenario. Still, the β-glutamate cycle, 3HGL

and RED cycles are close (having 61%, 70% and 90% of predicted pathway activity of
the CBB cycle at high CO_2). Overall, this makes especially the β-glutamate and 3HGL
cycles promising to explore, given a potentially attractive yield, and reasonable
predicted pathway activity. As discussed above, pathway activity may be further
increased based on natural enzyme screening and enzyme engineering and evolution
efforts. The β-glutamate cycle especially stands out compared to 3HGL as it does not
require B_{12} and could potentially also be implemented in photosynthetic microalgae and
plants. However, given its required three new-to-nature activities and relatively large
pathway size this would be a tremendous engineering effort already in bacteria, and
definitely in microalgae and plants. The RED cycle, on the other hand, could be highly
relevant based on its overlap with the CBB and high predicted pathway activity, while
the ABC cycle strike a balance between yield, activity, and ease of implementation,
since they require only one engineered enzyme and otherwise rely on the naturally
occurring and widespread threonine biosynthesis route.

The fact that all pathways are predicted to be slower than the CBB cycle also highlights
the fact that carbon fixation cycles are not only limited by carboxylation activity and that
Rubisco, despite its relatively slow catalysis and oxygenation side activity, still leads to
relatively high pathway activity when considered in its metabolic context[54].

An open question remains for the field: why is the CBB ubiquitous, if stoichiometrically
and/or kinetically superior pathways are potentially possible? One possible explanation
for this is the high degree of overlap between the CBB cycle and the pentose phosphate
pathway, which exists in all living organisms. In fact, only Rubisco and PRK are
necessary on top of the standard pentose phosphate pathway genes, as demonstrated
by the engineering of *E. coli* to grow autotrophically via the CBB cycle, and it is common
in nature for these two genes to be expressed from a plasmid, which can easily be
horizontally transferred, thus spreading the CBB cycle across phyla. Furthermore,
several essential biomass precursors are formed as part of the cycle, which would make
a loss of activity detrimental. At least in the short term, an organism that acquired all the
necessary genes to support an alternative carbon fixation cycle pathway would have to
support both cycles at the same time, likely with competing if not outright incompatible
fluxes and metabolic profiles. The engineering approach to rewiring metabolism uses
autotrophies and bypasses to force metabolic rearrangements[55]. This strategy has
proven highly reliable in engineering a diverse array of pathways and will likely be just
as useful in the future[40,56–59].

Overall, our study expands the solution space for carbon fixation by providing realistic
and biochemically feasible avenues for enzymatic transformations based on known
enzyme mechanisms and substrate promiscuity. Notably, we identified two potential
novel carboxylases that could lead to entirely new pathway architectures once
implemented. New carbon fixation pathways with better yield and insensitivity to oxygen

could lead to improved carbon fixation in algae or plants, potentially leading to
renewable production of chemicals and increases in crop yields.

Methods

All scripts and input files are available on gitlab (https://git.wur.nl/vittorio.rainaldi/lvl4). All
pathways were computationally analyzed using flux balance analysis[60] (FBA) with
cobrapy[61] (version 0.29.1) using the *E. coli* iML1515[12] genome scale-metabolic model.
The base model was modified to change membrane-bound transhydrogenase
(TDH2pp) to translocate one proton rather than two[62]. Anaerobic reactions such as PFL,
OBTFL, and POR5 were set to zero. Non-growth associated maintenance (ATPM) was
set to zero. Additional modifications were added for each individual pathway. FBA
returns a flux value in gCDW/mol substrate. In this case, since CO_2 is freely available in
the atmosphere, it was left unconstrained, however an energy source is required to run
CO_2 fixation cycle, and this was used as a constraint for all pathways. For this purpose,
we added a dummy reaction which produces NADPH, which is the main source of
reducing power in photosynthetic organisms such as algae and plants. This results in
slightly higher values compared to using NADH as an energy source, since NADH to
NADPH conversion requires the translocation of one proton, which amounts to roughly
one-third of one ATP[62].

Max-min driving force[11] and enzyme cost[13] were computed with a custom script
modified from previous work[44,63] using equilibrator (version 0.6.0). Metabolite
concentrations were constrained to the physiologically relevant range between 10 µM
and 10 mM. ECM was computed using the following parameters: algorithm ECM,
version 3, kcat_source fwd, denominator CM, regularization volume, p_h 7.5,
ionic_strength 0.3 molar, p_mg 3, dg_confidence 0.95. Kinetics values were taken from
Brenda[64] or previous work[7]. In the case of new-to-nature enzymatic activities, the
kinetics were estimated by taking values from similar enzymes. For example, kinetic
values for α-ketobutyrate carboxylase are not available, so values for pyruvate
carboxylase were used instead. Since this enzyme catalyzes the same reaction on a
slightly smaller substrate, it is reasonable to assume that an engineered enzyme could
reach similar kinetic properties. The ECM algorithm returns a list of enzyme costs in
moles of enzyme per liter based on a total flux through the pathway for 1 mM/s of
product formed. These values were multiplied by the molecular weight of the respective
enzyme to obtain the enzyme cost in g/l. The sum of all the values is the enzyme cost
(EC) for the pathway. To obtain the pathway activity we performed the following
calculations:

$\text{Pathway activity} = \frac{\mu mol}{min*mg}$

flux $= x \frac{mM}{s} = 1000x \frac{\mu M}{s} = 60000x \frac{\mu M}{min} = 60000x \frac{\mu mol}{min*L}$

Enzyme Cost (EC) $= y \frac{g}{L} = 1000y \frac{mg}{L}$

Pathway activity $= 60000x \frac{\mu mol}{min*L} \left(\frac{1}{1000y}\right) \frac{L}{mg} = 60 \frac{x}{y} \frac{\mu mol}{min*mg} = 60 \frac{flux}{EC}$

References

1. *The Prokaryotes*. (Springer Berlin Heidelberg, Berlin, Heidelberg, 2013).

doi:10.1007/978-3-642-30123-0.

2. Prywes, N., Phillips, N. R., Tuck, O. T., Valentin-Alvarado, L. E. & Savage, D. F.

Rubisco Function, Evolution, and Engineering. *Annu. Rev. Biochem.* **92**, 385–410

(2023).

3. Tcherkez, G. G. B., Farquhar, G. D. & Andrews, T. J. Despite slow catalysis and

confused substrate specificity, all ribulose bisphosphate carboxylases may be nearly

perfectly optimized. *Proceedings of the National Academy of Sciences* **103**, 7246–

7251 (2006).

4. Claassens, N. J. *et al.* Phosphoglycolate salvage in a chemolithoautotroph using the

Calvin cycle. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 22452–22461 (2020).

5. Bar-Even, A., Noor, E., Lewis, N. E. & Milo, R. Design and analysis of synthetic

carbon fixation pathways. *Proceedings of the National Academy of Sciences* **107**,

8889–8894 (2010).

6. Schwander, T., Von Borzyskowski, L. S., Burgener, S., Cortina, N. S. & Erb, T. J. A

synthetic pathway for the fixation of carbon dioxide in vitro. *Science* **354**, 900–904

(2016).

7. Löwe, H. & Kremling, A. In-Depth Computational Analysis of Natural and Artificial

Carbon Fixation Pathways. *BioDesign Research* **2021**, 1–23 (2021).

8. Erb, T. J., Jones, P. R. & Bar-Even, A. Synthetic metabolism: metabolic engineering

meets enzyme design. *Current Opinion in Chemical Biology* **37**, 56–62 (2017).

9. Vornholt, T. *et al.* Artificial metalloenzymes. *Nat Rev Methods Primers* **4**, 78 (2024).

10. Zou, Z. *et al.* De novo design and evolution of an artificial metathase for cytoplasmic

olefin metathesis. *Nat Catal* 1–12 (2025) doi:10.1038/s41929-025-01436-0.

11. Noor, E. *et al.* Pathway Thermodynamics Highlights Kinetic Obstacles in Central

Metabolism. *PLoS Computational Biology* **10**, (2014).

12. Monk, J. M. *et al.* iML1515, a knowledgebase that computes Escherichia coli traits.

Nat Biotechnol **35**, 904–908 (2017).

13. Noor, E. *et al.* The Protein Cost of Metabolic Fluxes: Prediction from Enzymatic Rate

Laws and Cost Minimization. *PLoS Computational Biology* **12**, 1–29 (2016).

14. Rainaldi, V. *et al.* Modular in vivo engineering of the reductive methylaspartate

cycles for synthetic CO2 fixation. 2025.11.12.688011 Preprint at

https://doi.org/10.1101/2025.11.12.688011 (2025).

15. Croft, M. T., Lawrence, A. D., Raux-Deery, E., Warren, M. J. & Smith, A. G. Algae

acquire vitamin B12 through a symbiotic relationship with bacteria. *Nature* **438**, 90–

93 (2005).

16. Alban, C., Baldet, P., Axiotis, S. & Douce, R. Purification and Characterization of 3-

Methylcrotonyl-Coenzyme A Carboxylase from Higher Plant Mitochondria. *Plant*

Physiol. **102**, 957–965 (1993).

17. Huang, C. S., Ge, P., Zhou, Z. H. & Tong, L. An unanticipated architecture of the

750-kDa α6β6 holoenzyme of 3-methylcrotonyl-CoA carboxylase. *Nature* **481**, 219–

223 (2012).

18. Chavez-Avilas, M., Díaz-Pérez, A. L., Reyes-de la Cruz, H. & Campos-García, J.

The *Pseudomonas aeruginosa liuE* gene encodes the 3-hydroxy-3-methylglutaryl

coenzyme A lyase, involved in leucine and acyclic terpene catabolism. *FEMS*

Microbiology Letters **296**, 117–123 (2009).

19. Wang, H. *et al.* An essential bifunctional enzyme in *Mycobacterium tuberculosis* for

itaconate dissimilation and leucine catabolism. *Proc. Natl. Acad. Sci. U.S.A.* **116**,

15907–15913 (2019).

20. Raj, H. *et al.* Engineering methylaspartate ammonia lyase for the asymmetric

synthesis of unnatural amino acids. *Nature Chem* **4**, 478–484 (2012).

21. Herrmann, G. *et al.* Two beta-alanyl-CoA:ammonia lyases in Clostridium

propionicum. *The FEBS Journal* **272**, 813–821 (2005).

22. Ruzicka, F. J. & Frey, P. A. Glutamate 2,3-aminomutase: A new member of the

radical SAM superfamily of enzymes. *Biochimica et Biophysica Acta (BBA) -*

Proteins and Proteomics **1774**, 286–296 (2007).

23. Tunçkanat, T. *et al.* Lysine 2,3-Aminomutase and a Newly Discovered Glutamate

2,3-Aminomutase Produce β-Amino Acids Involved in Salt Tolerance in

Methanogenic Archaea. *Biochemistry* **61**, 1077–1090 (2022).

24. Chen, D., Ruzicka, F. J. & Frey, P. A. A novel lysine 2,3-aminomutase encoded by

the yodO gene of Bacillus subtilis□: characterization and the observation of organic

radical intermediates. 11 (2000).

25. Chen, D., Frey, P. A., Lepore, B. W., Ringe, D. & Ruzicka, F. J. Identification of

Structural and Catalytic Classes of Highly Conserved Amino Acid Residues in

Lysine 2,3-Aminomutase. *Biochemistry* **45**, 12647–12653 (2006).

26. Behshad, E. *et al.* Enantiomeric Free Radicals and Enzymatic Control of

Stereochemistry in a Radical Mechanism: The Case of Lysine 2,3-Aminomutases.

Biochemistry **45**, 12639–12646 (2006).

27. Sofia, H. J., Chen, G., Hetzler, B. G., Reyes-Spindola, J. F. & Miller, N. E. Radical

SAM, a novel protein superfamily linking unresolved steps in familiar biosynthetic

pathways with radical mechanisms: functional characterization using new analysis

and information visualization methods. *Nucleic Acids Res* **29**, 1097–1106 (2001).

28. Booker, S. J. & Lloyd, C. T. Twenty Years of Radical SAM! The Genesis of the

Superfamily. *ACS Bio Med Chem Au* **2**, 538–547 (2022).

29. Mack, M. *et al.* 3-Methylglutaconyl-CoA hydratase from Acinetobacter sp. *Arch*

Microbiol **185**, 297–306 (2006).

30. Liu, Y. *et al.* Systematic design and evaluation of artificial CO2 assimilation

pathways. *Synthetic and Systems Biotechnology* **10**, 1107–1118 (2025).

31. Kim, J., Hetzel, M., Boiangiu, C. D. & Buckel, W. Dehydration of (*R*)-2-hydroxyacyl-

CoA to enoyl-CoA in the fermentation of α-amino acids by anaerobic bacteria. *FEMS*

Microbiol Rev **28**, 455–468 (2004).

32. Kim, J., Darley, D. J., Buckel, W. & Pierik, A. J. An allylic ketyl radical intermediate in

clostridial amino-acid fermentation. *Nature* **452**, 239–242 (2008).

33. Bahnson, B. J., Anderson, V. E. & Petsko, G. A. Structural Mechanism of Enoyl-CoA

Hydratase: Three Atoms from a Single Water Are Added in either an E1cb Stepwise

or Concerted Fashion. *Biochemistry* **41**, 2621–2629 (2002).

34. Hordijk, W. Autocatalytic confusion clarified. *Journal of Theoretical Biology* **435**, 22–

28 (2017).

35. Kurteva-Yaneva, N. *et al.* Structural Basis of the Stereospecificity of Bacterial B12-

dependent 2-Hydroxyisobutyryl-CoA Mutase. *Journal of Biological Chemistry* **290**,

9727–9737 (2015).

36. Rohwerder, T., Rohde, M.-T., Jehmlich, N. & Purswani, J. Actinobacterial

Degradation of 2-Hydroxyisobutyric Acid Proceeds via Acetone and Formyl-CoA by

Employing a Thiamine-Dependent Lyase Reaction. *Front. Microbiol.* **11**, 691 (2020).

37. Pfister, P., Zarzycki, J. & Erb, T. J. Structural Basis for a Cork-Up Mechanism of the

Intra-Molecular Mesaconyl-CoA Transferase. *Biochemistry* **62**, 75–84 (2023).

38. Schulz-Mirbach, H. *et al.* New-to-nature CO_2-dependent acetyl-CoA assimilation

enabled by an engineered B12-dependent acyl-CoA mutase. *Nat Commun* **15**,

10235 (2024).

39. Roymoulik, I., Moon, N., Dunham, W. R., Ballou, D. P. & Marsh, E. N. G.

Rearrangement of L-2-Hydroxyglutarate to L-*threo*-3-Methylmalate Catalyzed by

Adenosylcobalamin-Dependent Glutamate Mutase. *Biochemistry* **39**, 10340–10346

(2000).

40. Wenk, S. *et al.* Evolution-assisted engineering of *E. coli* enables growth on formic

acid at ambient CO_2 via the Serine Threonine Cycle. *Metabolic Engineering* **88**, 14–

24 (2025).

41. Scrutton, M. C., Griminger, P. & Wallace, J. C. Pyruvate Carboxylase. *Journal of*

Biological Chemistry **247**, 3305–3313 (1972).

42. Scheffen, M. *et al.* A new-to-nature carboxylation module to improve natural and

synthetic CO_2 fixation. *Nature Catalysis* **4**, 105–115 (2021).

43. Rainaldi, V., Donati, S., D'Adamo, S. & Claassens, N. J. A recursive pathway for isoleucine biosynthesis arises from enzyme promiscuity. 2025.08.26.672309 Preprint at https://doi.org/10.1101/2025.08.26.672309 (2025).

44. Satanowski, A. *et al.* Awakening a latent carbon fixation cycle in Escherichia coli. *Nature Communications* **11**, 1–14 (2020).

45. Aoshima, M. & Igarashi, Y. A novel oxalosuccinate-forming enzyme involved in the reductive carboxylation of 2-oxoglutarate in Hydrogenobacter thermophilus TK-6. *Molecular Microbiology* **62**, 748–759 (2006).

46. Aoshima, M. & Igarashi, Y. Nondecarboxylating and Decarboxylating Isocitrate Dehydrogenases: Oxalosuccinate Reductase as an Ancestral Form of Isocitrate Dehydrogenase. *J Bacteriol* **190**, 2050–2055 (2008).

47. Buhrman, G. *et al.* Structure, Function, and Thermal Adaptation of the Biotin Carboxylase Domain Dimer from Hydrogenobacter thermophilus 2-Oxoglutarate Carboxylase. *Biochemistry* **60**, 324–345 (2021).

48. Marchal, D. G. *et al.* Machine Learning-Supported Enzyme Engineering toward Improved CO2-Fixation of Glycolyl-CoA Carboxylase. *ACS Synth. Biol.* **12**, 3521–3530 (2023).

49. Yang, X., Ma, Y., Yan, J., Zhao, G. & Zhang, Y. LATCH: a Linear Autocatalytic cycle Tailored for Carbon Harvesting. *Green Carbon* https://doi.org/10.1016/j.greenca.2025.10.003 (2025) doi:10.1016/j.greenca.2025.10.003.

50. Satanowski, A. *et al.* Design and implementation of aerobic and ambient CO2-

reduction as an entry-point for enhanced carbon fixation. *Nat Commun* **16**, 3134

(2025).

51. Davidi, D. *et al.* Global characterization of in vivo enzyme catalytic rates and their

correspondence to in vitro kcat measurements. *Proceedings of the National*

Academy of Sciences of the United States of America **113**, 3401–3406 (2016).

52. Gale, N. & Beck, J. Evidence for the Calvin Cycle and Hexose Monophosphate

Pathway in Thiobacillus ferrooxidans.

https://journals.asm.org/doi/epdf/10.1128/jb.94.4.1052-1059.1967 (1967)

doi:10.1128/jb.94.4.1052-1059.1967.

53. Dronsella, B. *et al. Engineered Synthetic One-Carbon Fixation Exceeds Yield of the*

Calvin Cycle. http://biorxiv.org/lookup/doi/10.1101/2022.10.19.512895 (2022)

doi:10.1101/2022.10.19.512895.

54. Bathellier, C., Tcherkez, G., Lorimer, G. H. & Farquhar, G. D. Rubisco is not really

so bad. *Plant Cell Environ* **41**, 705–716 (2018).

55. Schulz-Mirbach, H., Dronsella, B. & Erb, T. J. Escherichia coli selection strains for

growth-coupled metabolic engineering. *Trends in Biotechnology* **0**, (2025).

56. Gleizer, S. *et al.* Conversion of Escherichia coli to Generate All Biomass Carbon

from CO2. *Cell* **179**, 1255-1263.e12 (2019).

57. Kim, S. *et al.* Growth of E. coli on formate and methanol via the reductive glycine

pathway. *Nature Chemical Biology* **16**, 538–545 (2020).

58. Reiter, M. A. *et al.* A synthetic methylotrophic Escherichia coli as a chassis for

bioproduction from methanol. *Nat Catal* **7**, 560–573 (2024).

59. Chen, F. Y. H., Jung, H. W., Tsuei, C. Y. & Liao, J. C. Converting Escherichia coli to

a Synthetic Methylotroph Growing Solely on Methanol. *Cell* **182**, 933-946.e14 (2020).

60. Edwards, J. S., Ibarra, R. U. & Palsson, B. O. In silico predictions of Escherichia coli

metabolic capabilities are consistent with experimental data. *Nat Biotechnol* **19**,

125–130 (2001).

61. Ebrahim, A., Lerman, J. A., Palsson, B. O. & Hyduke, D. R. COBRApy: COnstraints-

Based Reconstruction and Analysis for Python. *BMC Systems Biology* **7**, 74 (2013).

62. Bizouarn, T., Van Boxel, G. I., Bhakta, T. & Jackson, J. B. Nucleotide binding

affinities of the intact proton-translocating transhydrogenase from Escherichia coli.

Biochimica et Biophysica Acta - Bioenergetics **1708**, 404–410 (2005).

63. Elad Noor / ged-cycle · GitLab. *GitLab* https://gitlab.com/elad.noor/ged-cycle (2024).

64. Chang, A. *et al.* BRENDA, the ELIXIR core data resource in 2021: new

developments and updates. *Nucleic Acids Res* **49**, D498–D508 (2021).