A divergent *Plasmodium* NEK4 acts as a key regulator driving the early

events of meiosis

Ryuji Yanase[1,2], Molly Hair[3], Mohammad Zeeshan[1], David J. P. Ferguson[3], Declan Brady[1],

Carla Pasquarello[4], Andrew Bottrill[5], Suhani Bhanvadia[6], Arrmund Neal[6], Eelco C. Tromer[7],

Karine G. Le Roch[6], Alexandre Hainard[4], Anthony A. Holder[8], Sue Vaughan[3], David S.

Guttery[1,2,*], Rita Tewari[1,*]

[1]School of Life Sciences, Queen's Medical Centre, University of Nottingham, Nottingham,

UK

[2]Department of Genetics, Genomics and Cancer Sciences, College of Life Sciences,

University of Leicester, Leicester, UK

[3]Oxford Brookes University, Department of Biological and Medical Sciences, Oxford, UK

[4]Proteomics Core Facility, Faculty of Medicine, University of Geneva, Switzerland

[5]School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry, UK

[6]Department of Molecular, Cell and Systems Biology, University of California, Riverside,

Riverside, United States

[7]Cell Biochemistry, Groningen Institute of Biomolecular Sciences & Biotechnology,

University of Groningen, Groningen, The Netherlands

[8]Malaria Parasitology Laboratory, The Francis Crick Institute, London, UK

*Corresponding authors:

rita.tewari@nottingham.ac.uk (RT); david.guttery@nottingham.ac.uk (DSG)

Abstract

Meiosis is a conserved yet evolutionarily varied process underpinning sexual reproduction in eukaryotes. In the malaria parasite *Plasmodium*, meiosis is unconventional: it occurs immediately after fertilisation (post-zygotic) and must be coordinated with the transformation of the zygote into a motile ookinete. The mechanisms synchronising these meiotic and morphogenetic programmes remain unknown. Here, we identify the *Plasmodium berghei* NIMA-related kinase NEK4 as a key regulator that couples meiotic initiation with zygote morphogenesis. Using ultrastructure expansion microscopy, we show that NEK4 accumulates at the microtubule-organising centre (MTOC) and the apical polar complex (APC) shortly after fertilisation, preceding the assembly of perinuclear and cortical microtubules. We reveal that *Plasmodium* zygotes undergo MTOC-associated nuclear migration, analogous to the meiotic nuclear movement in fission yeast. Deletion of the *Pbnek4* gene results in complete developmental arrest: MTOC duplication and microtubule formation are blocked, chromatin remains uncondensed, and nuclear migration and cell polarity fail to establish. Transcriptomic and phosphoproteomic analyses reveal that absence of NEK4 causes a collapse in transcriptional and phosphoregulatory networks governing meiosis and cytoskeletal organisation, leading to reduced expression and phosphorylation of important players, including HOP1, REC8, and AP2-O. These findings establish NEK4 as a key regulator driving meiotic entry and zygote maturation.

Keywords: *Plasmodium*; meiosis; NIMA-related kinase; malaria; zygote; ookinete; MTOC; microtubules; chromosome

Introduction

Meiosis is a conserved and essential mode of cell division that underpins sexual reproduction in eukaryotes. It entails a carefully choreographed sequence of events, including DNA replication, homologous recombination, pairing of homologous chromosomes (synapsis) and two rounds of chromosome segregation[1]. In mammals, it generates haploid gametes from diploid progenitors, following two successive nuclear divisions without intervening genome duplication[1]. In many well-studied organisms including yeast, plants, and mammals, meiotic progression is regulated by a network of conserved protein kinases and checkpoints, including cyclin-dependent kinases (CDKs), Polo-like kinases (PLKs), the anaphase-promoting complex/cyclosome (APC/C), ataxia-telangiectasia mutated (ATM)/ATM and Rad3-related (ATR) kinases, checkpoint kinases (CHK1/2), and members of the NIMA-related kinase (NEK) family[2, 3, 4, 5]. These regulators ensure the fidelity of chromosome segregation by integrating signals from DNA damage checkpoints, spindle dynamics, and crossover/recombination status, at the same time as maintaining synchrony with cytoskeletal changes that support cell division and differentiation.

Despite the broad conservation of its core function to facilitate obligatory chromosomal exchanges and homologous recombination between parental chromosomes, both the molecular machinery executing meiosis and its operational sequence of events display remarkable variability[6, 7]. For a wide range of unicellular eukaryotes, including parasitic protists, meiotic programmes have undergone substantial evolutionary divergence. This is evident in the malaria-causing parasite *Plasmodium*, a member of the phylum Apicomplexa[7, 8]. This haploid parasite has a complex lifecycle with proliferative asexual multiplication in the liver and blood stream of the host[9]. A subpopulation of blood-stage parasites differentiates into haploid male and female gametocytes which, following ingestion in a mosquito blood meal, start the sexual phase of the life cycle inside the gut of the vector. Gametogenesis produces haploid gametes and following fertilisation, meiosis occurs in the diploid zygote; a process known as post-zygotic meiosis. The zygote undergoes a ~24-hour developmental programme to become an elongated, motile ookinete capable of traversing the midgut epithelium[8] (Fig. 1a), where it forms an oocyst. Many haploid sporozoites then emerge from the oocyst, migrate to the mosquito salivary glands and transmit to the vertebrate host in the blood meal. *Plasmodium* infections result in over 250 million clinical cases and nearly 600,000 deaths each year, disproportionately affecting young children in

sub-Saharan Africa[10]. These outcomes are driven not only by the rapid asexual proliferation

in the human host, but also by the successful sexual reproduction and transmission through

the mosquito vector. Transmission is an obligate part of the life cycle and therefore

prevalence of the disease is critically reliant on the process of meiosis[9].

Meiosis in *Plasmodium* is divergent from the canonical process observed in model eukaryotes. In

these organisms, a diploid cell undergoes pre-meiotic DNA replication followed by prophase I

substages including leptotene, zygotene, pachytene, diplotene and diakinesis, before metaphase I and

anaphase I of the first meiotic division and then meiosis II, ultimately generating four separate germ

cells, each containing a haploid nucleus[11]. In *Plasmodium*, eukaryote mostly haploid and meiosis

begins in diploid zygote formed after fertilisation. The early meiotic events, for example DNA

replication, chromosome pairing, and formation of chromosome axes (proteinaceous scaffolds that

form along sister chromatids), are initiated within the first few hours post-fertilisation. Ultrastructural

studies have shown that *Plasmodium* zygotes progress through early prophase I stages (leptotene and

zygotene), forming distinct synaptonemal complexes (SC) that persist until metaphase I[12, 13], whereas

in canonical meiosis the SC assembles during zygotene, is fully established at pachytene, and

normally disassembles during diplotene before diakinesis and metaphase I[11]. However, in

Plasmodium later prophase I stages (including diplotene and diakinesis) and canonical meiosis II have

not been observed[14, 15]. Despite two rounds of chromosome segregation, the ookinete retains a single

nucleus with four kinetochore clusters[16, 17], indicating a divergent closed meiosis without karyokinesis

(similar to meiosis I in *S. pombe* and the closed mitosis characteristic of *Plasmodium*[17]), rather than

the formation of four separate haploid nuclei.

The cell biological divergence of meiosis is reflected at the molecular level as well.

Plasmodium species lack orthologues of several conserved meiotic regulators, including

CDC25, CDC14, PLKs, and many APC/C components[18, 19]. They do possess a reduced,

parasite-specific set of protein kinases that appear to have acquired regulatory roles normally

fulfilled by these missing elements[19, 20]. The signalling mechanisms that control *Plasmodium*

meiosis and coordinate with cytoskeletal reorganisation and establishment of cell polarity, are

poorly defined, although cGMP signalling has been suggested to initiate ookinete gliding

motility and polarization[21]. One protein kinase family of particular interest is the NIMA-

related kinases (NEKs), which are conserved across eukaryotes and regulate centrosome

function, spindle assembly, and 3D chromatin organisation[22, 23]. In metazoans, specific NEKs

such as NEK1, NEK2, and NEK9 function in meiosis and mitosis, often through regulation of

microtubule dynamics and DNA damage (NEK1) and spindle assembly (NEK2 and NEK9)

checkpoint control[22, 24]. The *Plasmodium* genome encodes four NEKs (NEK1–4), which

have, apart from *Plasmodium* NEK1 and human NEK2, no direct 1-to-1 human orthologs

(Extended Data Fig. 1a, b). Each *Plasmodium* NEK has a distinct expression pattern and

function during the parasite life cycle[25, 26, 27, 28], with NEK2 and NEK4 shown to be critical

for zygote development and parasite transmission[19, 25] (Extended Data Fig. 1c). Deletion of

the *Pbnek4* gene results in a failure of DNA replication in the zygote following fertilisation[25]

(Figure 1a), yet the underlying function and phosphosignalling network remain undefined.

Given its post-zygotic meiosis within a haploid life cycle and the lack of several canonical

meiotic regulators[8], Plasmodium provides a powerful model to investigate how minimal

signalling networks coordinate chromosome dynamics with cytoskeletal reorganisation. Here,

we show that during early zygote development where the post-zygotic meiosis commences,

Plasmodium berghei (Pb) NEK4 is located at the microtubule organising centre (MTOC) and

apical polar complex (APC). Deletion of the *Pbnek4* gene disrupts microtubule organisation,

MTOC development, and nuclear migration during meiotic S phase and prophase.

Furthermore, its absence impairs transcription and phosphorylation of key meiotic proteins,

leading to a failure in chromosome condensation and developmental arrest at the earliest

stages of meiosis. We propose that this kinase integrates the nuclear and cell differentiation

events that drive meiotic progression and ookinete development, respectively.

Results

NEK4 moves to the nucleus, MTOC, and apical polar complex during early meiotic

development

Our previous study showed that NEK4 ablation blocks zygote development[25]. To investigate

NEK4 functions during the meiotic stages of *P. berghei* development, first we determined the

spatiotemporal location of NEK4, using a transgenic parasite line expressing the protein with

a C-terminal GFP tag from the modified endogenous gene locus (PbNEK4-GFP – Extended

Data Fig. 1d-f). Live-cell imaging revealed a dynamic pattern of PbNEK4-GFP location

following female gametocyte activation, zygote differentiation and ookinete development

(Fig. 1b and Extended Data Fig. 2 and 3a). In non-activated female gametocytes, the GFP

signal was diffuse throughout the nucleoplasm and cytoplasm (Fig. 1b). Then shortly after

fertilisation and in the first two hours post-activation (hpa), the PbNEK4-GFP signal was

concentrated in two foci (Fig. 1b), one at the periphery of the nucleus near the nuclear
envelope and the other at the cell cortex. By two to six hpa, these foci were more pronounced,
suggesting a localisation of PbNEK4-GFP at both nuclear and apical regions of the cell
during this period of intense morphological transformation. The PbNEK4-GFP signal within
the nucleoplasm appeared stronger relative to the cytoplasm. In addition, a focal dot-like
PbNEK4-GFP-positive structure was observed moving within the cytoplasm (Fig. 1b; yellow
arrowhead and Supplementary Video 1 and Extended Data Fig. 3b). This structure remained
visible during the early stages of zygote-ookinete development (until ~10 hpa) but was no
longer detectable in the fully mature ookinete (Extended Data Fig. 2 and 3c). We speculate
that this focus may represent a remnant of the NEK4-associated structure that undergoes
degradation during the stages of development. The morphological transformation of the
zygote into an ookinete begins at about 4 hpa and is initiated at the site where PbNEK4-GFP
is located at the cell periphery. The onset of this transformation is accompanied by reduction
of this GFP signal and the disappearance of the nuclear PbNEK4-GFP focus, with residual
PbNEK4-GFP dispersed diffusely throughout the cytoplasm and nucleoplasm (Fig. 1b, c, and
Extended Data Fig. 3a).

To examine in more detail the location of NEK4 within the developing zygote, we used
ultrastructure expansion microscopy (U-ExM). We utilised NHS-ester staining, which labels
primary amine groups on proteins to reveal the global intracellular architecture like electron
microscopy[29]. This staining clearly revealed the nuclear MTOC and the development of the
APC at the cell periphery. Immunofluorescence labelling of the U-ExM samples with an anti-
GFP antibody confirmed that PbNEK4-GFP was located at both the MTOC and the APC
(Fig. 1d). Furthermore, at 4 hpa, the dividing and separating MTOCs and a more developed
APC were observed by NHS-ester staining. At this later time point, a distinct presence of
PbNEK4-GFP at both the MTOCs and the APC was confirmed (Fig. 1d). These observations
indicate prominent localisation of NEK4 during the early stages of zygote development (2 to
4 hpa), with a concentration at the MTOC and APC, alongside a diffuse distribution within
the nucleoplasm (Fig. 1e).

NEK4 coordinates nuclear repositioning and directional microtubule formation in the
zygote
In our study of NEK4 localisation in early meiotic zygotes, we observed that the nucleus
shows dynamic movement during early meiosis as seen across various eukaryotes[6, 30, 31]. For

example, in the fission yeast *Schizosaccharomyces pombe* (*S. pombe*), nuclear movement

during meiosis is characterised by a dramatic "horsetail" motion, where the nucleus oscillates

back and forth along the cell axis during prophase I[32, 33]. This movement is driven by

cytoplasmic dynein and microtubules anchored at the spindle pole body (SPB)[34, 35], and is

essential for efficient homologous chromosome pairing and recombination[32]. Time-lapse

imaging of PbNEK4-GFP zygotes at 2 hpa revealed that inside the zygote, PbNEK4-GFP

focus localised at the nucleus and moves together with the nucleus (magenta arrowheads in

Fig. 2a and Supplementary Video 2, 3). To examine MTOC and microtubule dynamics in

parallel during the nuclear movement, we used a transgenic *P. berghei* line expressing EB1

(End Binding protein 1) fused to GFP (PbEB1-GFP)[16], EB1 is a conserved microtubule plus-

end binding protein, but in *Plasmodium*, EB1 marks the microtubule lattice and the MTOC[16,

36]. The PbEB1-GFP focus, representing the MTOC position, was observed moving within the

zygote, leading the nucleus (magenta arrowheads in Fig. 2b). The nucleus appeared

elongated, seemingly pulled by the PbEB1-GFP focus. Additionally, PbEB1-GFP signal was

observed extending from the focus, likely representing microtubules extending from the

MTOC (Fig. 2b and Supplementary Video 4, 5). Furthermore, the fact that the localisation of

PbNEK4-GFP and PbEB1-GFP at the cell periphery remains largely stationary contrasts with

the nuclear movement (cyan arrowheads in Fig. 2a and b), indicating that this nuclear

movement is independent of whole-cell rotation or migration.

Quantification of the proportion of zygotes at different stages during development with

motile nuclei indicated that the movement peaked at between 2 and 4 hpa and then subsided

(Fig. 2c). These observations suggest that in *P. berghei* zygotes there is a form of nuclear

movement like that widely observed in *S. pombe* and other eukaryotes[6, 30, 31, 32, 37]. This

movement appears to be led by the MTOC, where both PbNEK4-GFP and PbEB1-GFP are

located (Fig. 2d). We also performed dual-colour live-cell imaging using a parasite line

expressing PbNEK4-GFP and PbEB1-mCherry. This analysis revealed a clear colocalization

of NEK4 and EB1 signals in the nucleus and at the cell periphery (Extended Data Fig. 4a).

To investigate the microtubule structures at higher resolution during the early development

stages, we performed U-ExM using an anti-α-tubulin antibody (Fig. 2e and Extended Data

Fig. 4b, c). At 2 hpa, two populations of zygotes were present: those with a single MTOC and

those with duplicated MTOCs. In cells with a single MTOC (Fig. 2e, top left panel), multiple

microtubules were visible in the perinuclear region, with one microtubule oriented as if

extending from the nuclear MTOC toward the APC. In cells with duplicated MTOCs (Fig. 2e,

top right panel), both microtubules emanating from the MTOCs and microtubule organisation

at the APC appeared more developed. Some of the microtubules running along the nuclear

periphery appeared to connect to the APC. In zygotes at 4 hpa (Fig. 2e, bottom left panel), the

two MTOCs were separating, with a microtubule structure resembling a spindle oriented

between them. Further development of microtubule structures was also evident in the

perinuclear region and at the APC. In summary, significant microtubule development occurs

at both the nuclear MTOCs and the APC, the sites of greatest PbNEK4-GFP accumulation

(Fig. 2f), suggesting that NEK4 may be involved in regulating microtubule polymerisation or

dynamics. This association of NEK4 with the MTOC and APC suggests a function as a

spatial organiser of microtubule polarity in the developing zygote. The microtubules

extending around the nucleus appear analogous to the perinuclear microtubules observed in

other eukaryotes during early meiosis, which mediate nuclear and chromosome movement[32,

38].

NEK4 is essential for nuclear migration, microtubule formation, and morphological

transformation from zygote to ookinete

To assess the importance of NEK4 during zygote/meiotic development, we used an existing

nek4 gene knockout parasite line (*Pbnek4-ko*; Extended Data Fig. 5a, b)[25]. A previous study

has shown that while *Pbnek4-ko* parasites undergo fertilisation, subsequent DNA replication

and ookinete formation are blocked[25]. Live-cell imaging over the period between 2 to 24 hpa

confirms that *Pbnek4-ko* zygotes failed to elongate or transition through the retort form into

mature ookinetes, in contrast to PbNEK4-GFP control cells that completed this

transformation within 20 hours (Fig. 3a). We then investigated the nuclear movement

observed in zygotes. Although moving nuclei were observed in a few *Pbnek4-ko* zygotes

between 2 and 4 hpa, the proportion of cells with moving nuclei was significantly reduced

compared to the PbNEK4-GFP controls (Fig. 3b).

To investigate microtubules in *Pbnek4-ko* zygotes, we used U-ExM. This revealed that the

distinctive formation of perinuclear and APC-associated microtubules in WT cells was

almost completely ablated in *Pbnek4-ko* zygotes (Fig. 3c). Although the presence of some

tubulin was detected, the formation of the microtubule-based structures at the APC was

almost completely disrupted compared to PbNEK4-GFP controls (WT; Fig. 3c).

Transmission electron microscopy (TEM) analysis also confirmed that partial formation of

the APC and the presence of microtubules at the APC and the perinuclear region were still

observable in *Pbnek4-ko* zygotes (Extended Data Fig. 5c). In non-activated female

gametocytes, a single MTOC-like structure was observed in both WT and *Pbnek4-ko*

parasites, but neither the formation of microtubules extending from it nor the assembly of the

APC structure was evident at this stage (Extended Data Fig. 5d, e). Following activation,

Pbnek4-ko zygotes exhibited partial formation of the APC and partial assembly of

microtubules at the MTOC and APC. Taken together, these results suggest that although

Pbnek4 deletion does not completely block microtubule formation, it profoundly

compromises their formation or stability. Additionally, while the formation of a nuclear

MTOC was observed, it failed to duplicate and remained as a single MTOC over the first four

hours after fertilisation. Together, these data on the consequences of *Pbnek4* gene deletion

demonstrate that NEK4 is essential for MTOC duplication and the organisation of perinuclear

and apical microtubule networks. These processes are prerequisites for DNA replication,

meiotic entry, and morphological transition from zygote to ookinete. In the absence of NEK4,

the microtubule-based cytoskeletal framework necessary for meiotic progression fails to

assemble, leading to developmental arrest prior to transition to retort form (Fig. 3d).

NEK4 is essential for establishing the ultrastructure necessary for meiotic chromatin

condensation and APC development

To further examine the ultrastructural consequences of *Pbnek4* knockout, we used TEM to

study WT and *Pbnek4-ko* zygotes (Fig. 4a). At 4 hpa, WT zygotes had a developed APC with

a well-organised microtubule structure (Fig. 4a1-2). At this stage, multiple thread-like

structures were observed within the nucleus (magenta arrowheads, Fig. 4a3-5). These

structures likely represent condensed--like chromosomes (the first stage of meiotic prophase

I, characterised by thread-like chromosome condensation), marking the entry into meiosis,

and the ends of these chromosomes appeared to be attached to the nuclear envelope (Fig.

4a5). In contrast, *Pbnek4-ko* zygotes at 4 hpa had only small APC and microtubule structures

(Fig. 4a6-7) and the condensed chromosomes seen in WT zygotes were completely absent in

Pbnek4-ko zygotes (Fig. 4a8-10).

We also utilised U-ExM coupled with NHS-ester and Hoechst 33342 DNA staining to

visualise the ultrastructure of the nucleoplasm (Fig. 4b). In WT zygotes at 4 hpa, NHS-ester

staining revealed numerous distinct thread-like structures within the nucleus that colocalised

with the Hoechst DNA signal, indicative of assembled chromosome axes and condensed

chromosomes. In contrast, these thread-like structures were absent within the nucleus in

Pbnek4-ko zygotes at 4 hpa, where the DNA signal remained widely diffuse.

To examine further the ultrastructural differences between WT and *Pbnek4-ko* zygotes, we

used Serial Block-Face Scanning Electron Microscopy (SBF-SEM). In most WT zygotes at 4

hpa, a developed APC was detected (Fig. 4c, d) and multiple condensed, thread-like

chromosomes were detected within their nuclei (Fig. 4c, magenta arrowheads and inset). In

some zygotes, these chromosomes were observed pairing with linear structures between them,

indicative of likely synaptonemal complex formation (Fig. 4c, cyan arrowheads and inset). In

contrast, in *Pbnek4-ko* zygotes at 4 hpa there was impaired APC development, and no

condensed chromosomes nor synaptonemal complexes were present within the nucleus (Fig.

4c, d). Three-dimensional models constructed from the SBF-SEM datasets illustrate the

development of the APC, the formation of condensed chromosomes (with one or both ends

appearing to attach to the nuclear envelope), and the separation of MTOCs in WT zygotes at

4 hpa. All these early meiotic processes are blocked in *Pbnek4-ko* zygotes (Fig. 4e). These

findings indicate that the deletion of the *Pbnek4* gene severely impairs the development of the

APC and its associated microtubule network in the early zygote, and also ablates the

formation of condensed chromosomes (likely leptotene chromosomes)). Given that *Pbnek4-*

ko parasites fail to undergo pre-meiotic DNA replication[25], this absence of chromosome

condensation is consistent with arrest prior to meiotic prophase.

NEK4 drives meiotic entry by coordinated control of transcription, protein synthesis

and phosphorylation

To elucidate the molecular processes governed by NEK4 during early meiosis in *P. berghei*

zygote development, we performed an integrated transcriptomic, proteomic,

phosphoproteomic, and interactomic analysis. Global transcriptomics was conducted on WT

and *Pbnek4-ko* parasites at 2 hpa and quantitative proteomics/phosphoproteomics were

conducted on non-activated and 2 hpa samples for both WT and *Pbnek4-ko* parasites. GFP-

Trap interactomics was performed using PbNEK4-GFP parasites at 2 hpa.

Quantitative real-time PCR (qRT-PCR) showed that the expression of meiosis-related genes

(e.g., *dmc1*, *hop1*, *mnd1*, and *spo11*) was significantly downregulated in *Pbnek4-ko* zygotes

at 2 hpa compared to WT zygotes at 2 hpa (Extended Data Fig. 6a). Transcriptomic profiling

via RNA-sequencing between WT and *Pbnek4-ko* parasites at 2 hpa revealed major

transcriptional changes in *Pbnek4-ko* parasites. At 2 hpa, 1,554 genes were significantly

differentially expressed (adjusted p-value ≤ 0.05, fold change ≥ 2) relative to WT parasites,

comprising 717 downregulated and 837 upregulated genes (Fig. 5a; Supplementary Table 1).

Gene Ontology enrichment of downregulated transcripts highlighted key biological

processes, including cell motility, cell differentiation, meiotic nuclear division, and

cytoskeletal organisation – hallmarks of zygote maturation (Fig. 5b; Supplementary Table 2).

Notably, transcripts encoding the meiotic recombinase *dmc1* [39] and the putative synaptonemal

complex component *hop1* (PBANKA_1407900)[40] were strongly reduced in the *Pbnek4-ko*

line.

Next, we performed GFP pulldown assays using PbNEK4-GFP and WT-GFP zygotes at 2

hpa. This analysis identified interactions of PbNEK4-GFP with putative MTOC-associate

proteins, such as a putative HEAT-repeat protein, which possesses predicted HEAT-like

repeats revealed by AlphaFold3[41] and Foldseek[42] structural prediction (PBANKA_0806700)

and a MORN repeat protein (PBANKA_1030400), as well as a regulator of chromosome

condensation (RCC; PBANKA_0905500). Furthermore, interactions with meiotic factors,

including REC8 (PBANKA_1419300)[43], DMC1, and HOP1, were also indicated (Fig. 5c;

Supplementary Table 13). These interactions may suggest that NEK4 acts as a signalling hub,

and either directly phosphorylates or stabilises other components within an integrated meiotic

complex.

To investigate the full landscape of NEK4-associated protein abundance and phosphorylation

changes during early zygote development, we performed quantitative proteomics and

phosphoproteomics analyses using non-activated (0 hpa) WT and *Pbnek4-ko* gametocytes,

and WT and *Pbnek4-ko* zygotes at 2 hpa. This design enabled us to separate activation-

associated changes (WT 2 hpa vs WT 0 hpa) from NEK4-dependent effects during early

zygote development (*Pbnek4-ko* 2 hpa vs WT 2 hpa). Initial analysis of phosphopeptide-

enriched and flow-through fractions from *Pbnek4-ko* and WT gametocytes at 0 hpa revealed

minimal differences in the abundance of identified proteins and phosphopeptides (Extended

Data Fig. 6b and d; Supplementary Table 5 and 12). These findings indicate that *Pbnek4*

deletion does not substantially reprogramme the proteome or phosphoproteome in mature

gametocytes, and instead suggest that NEK4-dependent changes are initiated upon

activation/fertilisation.

In contrast, significant differences in protein and phosphopeptide abundance were observed

when comparing gametocytes at 0 hpa with zygotes at 2 hpa, and *Pbnek4-ko* with WT

zygotes at 2 hpa (Fig. 5d-g; Extended Data Fig. 6c and f; Supplementary Table 6 and 11).

Proteomic analysis of the flow-through fractions obtained during phosphopeptide enrichment

revealed marked increases in zygote- and meiosis-associated proteins following fertilisation

in WT parasites. These included the zinc-finger protein Pb103[44], the transcription factor AP2-

Z[45], and the meiotic recombinase DMC1[39] (Fig. 5d; Supplementary Table 3). In contrast,

when *Pbnek4-ko* parasites were compared to WT controls at 2 hpa, many of these same

proteins were significantly less abundant (Fig. 5e; Supplementary Table 4), suggesting that

NEK4 activity influences their synthesis, stability, or translation during zygote development.

Phosphoproteomic profiling revealed a similar association with the presence of NEK4 for

protein phosphorylation linked to meiotic initiation. In WT parasites, fertilisation triggered

widespread phosphorylation of proteins linked to meiosis, transcription, translation, and

cytoskeletal organisation by 2 hpa. These proteins included DMC1[39], HOP1[40], REC8[43], DNA

polymerase delta subunit 3 (POLD3), AP2-Z[45, 46], AP2-O[47], Pb103[44], FD2[48], gSNF2[49],

NOT1-G[50], a putative HEAT-repeat protein (PBANKA_0806700), a serine/threonine protein

kinase putatively related to MAP/microtubule affinity-regulating kinase 1[51] and maternal

embryonic leucine zipper kinase 1[52] (MARK/MELK), zygote-ookinete specific dynein heavy

chain (DHC3)[53], and kinesin-4 (KIN4)[54] (Fig. 5f; Supplementary Table 5). In *Pbnek4-ko*

parasites phosphorylation at many sites in these proteins was significantly lower relative to

the WT control (Fig. 5g; Supplementary Table 9), Notably, some of these proteins also

showed increased overall abundance in the flow-through fraction (e.g. DMC1, Pb103, and

AP2-Z), indicating that the observed phosphorylation changes may partly reflect higher

protein levels rather than phosphorylation alone. However, the consistent reduction of

phosphorylation in *Pbnek4-ko* parasites suggests that NEK4 contribute to phosphoregulation

in addition to promoting protein accumulation. Gene Ontology (GO) analysis further

confirmed that zygote activation triggers increased phosphorylation of proteins involved in

microtubule-based movement, the cytoskeleton, and reproductive processes (Fig. 5h;

Supplementary Table 8). In contrast, *Pbnek4-ko* zygotes at 2 hpa showed reduced

phosphorylation of proteins associated with reproductive processes and meiotic nuclear

division (Fig. 5i; Supplementary Table 10). Representative proteins identified by our

proteomics and phosphoproteomic analyses, along with their abundance and putative

functions, are summarised in Figure 5j. These data support the hypothesis that NEK4 may

contributes to a phosphorylation network underpinning zygote differentiation and meiotic
progression.

Mammalian NEK kinases favour a hydrophobic residue (Leu, Met, Phe, or Trp) at the -3
position relative to the target serine or threonine, and disfavour a proline residue at the +1
position, establishing a consensus motif of [LMFW]-X-X-[S/T]-[no P][55, 56]. We observed this
consensus motif in multiple high-confidence *Plasmodium* targets ([F]-N-D-[T]-[S] in REC8,
[L]-S-V-[S]-[T] in gSNF2, and [L]-N-E-[T]-[L] in PBANKA_0806700). Importantly, all of
these motif-containing proteins were also independently identified as physical interaction
partners in the NEK4 GFP-Trap interactome (Fig. 5c and j).

Collectively, these integrated transcriptomic, interactomic, proteomic, and phosphoproteomic
datasets suggest that while NEK4 may exert some influence on transcript and protein
accumulation, its most profound and immediate impact at this early development stage (2
hpa) is on the phophoproteome. Together with the interactomic evidence, these results
identify NEK4 as a key regulator of early post-zygotic meiosis in *P. berghei*, orchestrating a
rapid rewiring of phosphosignalling networks to drive meiotic entry and zygote maturation.

Discussion

We identify *Plasmodium*NEK4 as a key kinase that links meiotic initiation and zygote
morphogenesis in *P. berghei* (Fig. 6). NEK4 localises to both the MTOC and the APC during
the earliest stages of zygote development (post-zygotic meiosis), and orchestrates MTOC
duplication, perinuclear and APC microtubule formation, and migration of the nucleus. This
directed movement resembles the "horsetail" nuclear oscillations during the meiotic prophase
in other eukaryotes, where cytoskeleton-driven repositioning of the nucleus promotes
homologous chromosome pairing and recombination[32, 37]. In *S. pombe*, this nuclear
movement occurs during prophase, where linear elements (LinEs) mediate chromosome
pairing and crossover. Similarly, in *Plasmodium*, from ~4 hpa onwards —when DNA
replication is complete, nuclear movement is evident, and morphological transformation into
the ookinete begins —the formation of condensed chromosomes reminiscent of leptotene
chromosomes and structures resembling synaptonemal complexes were observed. Although
fully condensed metaphase chromosomes seen in canonical meiosis are still elusive in
Plasmodium, this initial condensation step is clearly a prerequisite for proper meiotic

progression and this suggests that entry into meiotic prophase initiates at this stage in
Plasmodium (Fig. 6). In the absence of NEK4, the MTOC fails to duplicate, chromatin
remains diffuse, and nuclear translocation is lost, leading to arrest of cell development before
meiotic division and transformation to ookinete. These data suggest that NEK4 is a putative
central regulator linking nuclear reorganisation and cytoskeletal polarity at the onset of
meiosis. Previous studies suggest that while *Plasmodium* retains some canonical meiotic
processes well-characterised in organisms like *S. pombe*, its mechanisms and molecular
players are highly divergent, as evidenced by the evolutionary conservation analysis of
meiotic components in *Plasmodium* and by the progression through meiosis I and II without
nuclear division[8] (Fig. 6). Given this divergence, it is particularly intriguing that early meiotic
nuclear movement observed in model eukaryotes is conserved in *Plasmodium*, suggesting
that it may be important for homologous chromosome pairing and recombination in this
parasite as well.

The dual location of NEK4 and the developmental arrest that results from knocking out its
gene, place it at the intersection of nuclear and cytoskeletal organisation. Its behaviour
parallels that of NEKs in other systems, which regulate centrosome duplication, spindle
formation, and checkpoint transitions[22, 23]. In *S. pombe*, the NEK-like kinase Fin1 controls
spindle formation, specifically the maturation and inheritance of the SPB[57, 58], while NEK2
and NEK9 act at metazoan centrosomes to coordinate spindle assembly[23]. In *Plasmodium*,
which lacks canonical mitotic and meiotic regulators such as the phosphatases CDC25,
CDC14, and Polo-like kinases[18, 19], NEK4 appears to have acquired functions to integrate
both signalling and structural control within a simplified regulatory network.

Proteins identified in the PbNEK4-GFP pulldown assays, such as the putative HEAT-repeat
protein (PBANKA_0806700), MORN repeat protein (PBANKA_1030400), and Kinesin-4
(PBANKA_1208200), may interact with NEK4 at the MTOC to facilitate MTOC
development or segregation. Although the specific function of PBANKA_0806700 in
Plasmodium remains uncharacterised, HEAT-repeat proteins are universally known to form
alpha-helical solenoids that act as versatile scaffolds for protein-protein interactions[59]. Given
its massive size (4,658 aa) and multiple predicted HEAT repeats, we hypothesise that this
protein serves as a central structural scaffold at the MTOC, organising the macromolecular
complexes required for NEK4-dependent cytoskeletal remodelling during early zygote
development. Additionally, the identified regulator of chromosome condensation (RCC;

PBANKA_0905500) contains an RCC1 domain. While RCC1 domains classically mediate
interactions with chromatin or DNA[60], their functions are not restricted to chromosomal
regulation. Interestingly, human NEK8 and NEK9 possess an RCC1 domain, and this domain
is implicated in the recruitment to the centrosome[61]. Furthermore, a recent study in
Toxoplasma gondii has demonstrated that RCC-domain-containing proteins are involved in
the apical complex assembly[62]. This raises the possibility that the *Plasmodium* RCC
identified here is similarly recruited to the MTOC or the APC, where it may interact with
NEK4 to coordinate cytoskeletal and apical organisation.

Transcriptomic, proteomic and phosphoproteomic profiling revealed that fertilisation triggers
extensive reprogramming of transcription, protein abundance, and protein phosphorylation
during post-zygotic meiosis. While the apparent downregulation of many factors between 0
and 2 hpa is heavily confounded by the natural depletion of male gametocytes following
activation, our targeted comparison between WT and *Pbnek4-ko* at 2 hpa successfully
uncoupled these effects to reveal the specific NEK4-dependent networks. In WT parasites,
proteins likely essential for meiosis and zygote differentiation including protein sets relating
to meiosis (DMC1, HOP1 and REC8), DNA replication (POLD3), sexual development
(Pb103, AP2-Z, NOT1-G, FD2, AP2-O, gSNF2), and MTOC/microtubules (MARK/MELK,
DHC3, KIN4) are more phosphorylated in the zygote (Fig. 6). In *Pbnek4-ko* parasites, these
proteins were hypophosphorylated, suggesting that NEK4 contributes to phosphorylation
events required for onset of meiosis and zygote-ookinete development. Also, while the
primary effect of *Pbnek4* deletion is a sweeping reduction in phosphorylation, the
concomitant reduction in protein abundance for a subset of specific factors (e.g. DMC1,
Pb103, and AP2-Z) suggests that NEK4-dependent phosphosignalling may secondarily
influence the stability or synthesis of certain key meiotic regulators.

While distinguishing direct kinase substrates from downstream cascading effects is inherently
challenging in knockout studies, our integrated multi-omics datasets provide a powerful filter.
Analysis of the specific hypophosphorylated sites identified in our phosphoproteomic dataset
reveals that several proteins - such as REC8, gSNF2, and the putative HEAT-repeat protein
PBANKA_0806700 - possess sequence motifs that perfectly align with canonical NEK
kinase preferences. Mammalian NEK kinases favour a hydrophobic residue (Leu, Met, Phe,
or Trp) at the -3 position relative to the target serine or threonine, and disfavour a proline
residue at the +1 position, establishing a consensus motif of [LMFW]-X-X-[S/T]-[no P][55, 56].

We observed this structural signature in multiple high-confidence *Plasmodium* targets ([F]-N-

D-[T]-[S] in REC8, [L]-S-V-[S]-[T] in gSNF2, and [L]-N-E-[T]-[L] in PBANKA_0806700).

Strikingly, all of these motif-containing proteins were also independently identified as

physical interaction partners in our NEK4 GFP-Trap interactome. The convergence of

canonical motif-dependent phosphorylation and physical interaction strongly supports their

identity as likely substrates of NEK4. Although the limited number of these validated sites

currently precludes the mathematical definition of a robust, *de novo* PbNEK4 consensus

motif, these findings provide a critical molecular starting point for understanding how NEK4

directly interfaces with the early meiotic and developmental machinery.

Such coordinated phosphoregulation is consistent with broader concepts being developed for

eukaryotic meiosis. Recent phosphoproteomic studies in yeast have shown that meiotic

progression is driven by both changes in protein abundance and dynamic, stage-specific

waves of protein phosphorylation that rewire existing signalling networks[63]. These

phosphorylation cascades, directed by cyclin-dependent and Polo-like kinases, define

temporal transitions from recombination to segregation[63]. Large-scale quantitative

phosphoproteomics studies in *S. pombe* revealed that phosphorylation of thousands of sites

changes during meiotic development, for example affecting proteins involved in microtubule

organisation, recombination and chromosome cohesion[64]. Our findings are consistent with

this concept of a meiotic "phosphorewiring": the absence of NEK4 collapses a broad

phosphorylation programme encompassing transcriptional, translational and cytoskeletal

effectors, and other kinases and phosphatases, and implying that *Plasmodium* has an

analogous, temporally-ordered kinase programme despite its highly reduced kinase signalling

repertoire.

Although our study focuses on NEK4-dependent phosphorylation and protein accumulation

during early zygote development, the quantitative phosphoproteomics also identify multiple

protein kinases and phosphatases with substantial phosphorylation changes across activation

and/or in *Pbnek4-ko* parasites, and several kinase/phosphatase proteins are additionally

detected in the NEK4-GFP interactome (Supplementary Tables 13 and 14). In our curated

view of the dataset, phosphorylation of a PP2B/calcineurin catalytic subunit (CNA;

PBANKA_1227400), several PPM-family serine/threonine phosphatases (including

PPM5/6/8/11), and an NIF1-like phosphatase are dynamically modified during the 0–2 hpa

transition, with at least one PPM-family phosphatase (PPM5; PBANKA_1427200) showing

reduced phosphorylation in *Pbnek4-ko* relative to WT at 2 hpa. These observations suggest

that NEK4-associated "rewiring" is likely to involve not only kinase-driven phosphorylation

cascades but also changes in phosphatase activity, recruitment, or substrate access. The

presence of several phosphatases among the NEK4-dependent phosphosites and interactors

suggests that NEK4 participates in a broader signalling network balancing kinase and

phosphatase activities during early zygote development. Similar kinase–phosphatase

regulatory modules are known to control meiotic transitions in other eukaryotes, including

the regulation of chromosome segregation and meiotic progression by PP2A-family

phosphatases[65, 66]. Notably, conserved phosphatase control is also implicated in *Plasmodium*

atypical division programmes[18], supporting the possibility that phosphatases contribute

significantly to parasite signalling during sexual development.

Additionally, our datasets highlight the early zygote transcription factor AP2-Z as a NEK4-

associated node, with altered abundance and reduced phosphorylation in *Pbnek4-ko* parasites

at 2 hpa. Analysis of the transcriptional changes in *Pbnek4-ko* zygotes at 2 hpa showed that

~41% of annotated AP2-Z targets and ~41% of AP2-O targets, respectively are significantly

altered in the *Pbnek4-ko* line (Supplementary Tables 15 and 16), consistent with a broad

disruption of the stage-specific transcriptional outputs normally associated with these AP2

regulators during zygote/ookinete development. In contrast, when focusing specifically on

canonical meiotic genes, DMC1 was the only factor among those highlighted in our study

and is a reported AP2-Z/O target[45, 67]. This suggests that NEK4-dependent control of meiotic

initiation is not explained just by direct AP2-Z/O regulation of meiotic gene transcription but

may also involve post-transcriptional regulation and phosphosignalling-dependent changes in

protein accumulation.

Little is known about phosphorylation-dependent activation of AP2-Z or AP2-O. The

transcription of AP2-Z is regulated by upstream gametocyte factors (AP2-G and AP2-FG)

binding to the AP2-Z promoter[68]. The AP2-Z/O phosphosite(s) in the phosphoproteomic

datasets lie outside the AP2 DNA-binding domain, but this does not preclude a regulatory

role; the phosphorylation may modulate interactions with co-factors or chromatin-associated

complexes, affect stability, or determine subcellular localisation. These features may be

particularly important in the tightly staged zygote-to-ookinete transcriptional transition.

Together, these findings support a model in which NEK4 contributes to early developmental

"rewiring" that intersects with AP2-led transcriptional programmes. Defects in meiotic

progression likely reflect a combination of contributions from transcription, translation and
phosphoregulation.

It is important to note a biological limitation to our analyses: the 0 hpa parasite population is
comprised of a mixture of both male and female gametocytes. Following activation, male
gametes exflagellate and their biomass is largely depleted from the developing zygote
culture. Consequently, a substantial proportion of the proteins and phosphosites that are
diminished at 2 hpa (including known male-specific factors such as MDV1) likely reflect this
shift rather than active developmental downregulation within the female/zygote lineage.
Additional sampling earlier than at 2 hpa is likely to further enrich for primary NEK4-
dependent phosphorylation and interactions, and will be important for resolving the temporal
order of the kinase/phosphatase activation and inhibition underlying the rewiring.

Our structural observations suggest that the APC and the microtubules extending from the
nuclear MTOC are formed in a NEK4-dependent manner. Furthermore, we observed a link
between the microtubules extending from the MTOC and APC, suggesting that this
connection is associated with nuclear movement or positioning. Comparable architectures are
present in other apicomplexans: in *Toxoplasma gondii*, the APC and conoid form an
anchoring hub that coordinates cortical microtubules and organelle positioning[69, 70]. *Eimeria*
tenella and *Cryptosporidium* spp. also have similar apical scaffolds[71, 72]. Genetic modification
of key regulatory kinases has resulted in comparable defects in other apicomplexans-
specifically, the failure of MTOC/centrosome duplication and the uncoupling of nuclear
replication from cytoskeletal assembly—supporting a conserved requirement for precise
coordination between nuclear and cortical assembly. In *T. gondii*, conditional mutants of
NEK1 and the MAP kinase, MAPK2 block centrosome duplication and segregation, causing
the nucleus to replicate without successful daughter cell budding, effectively decoupling
nuclear and cytoskeletal programmes[73, 74]. *Plasmodium* NEK1 controls spindle and
kinetochore organisation during male gametogenesis[28], while loss of the phosphatase PPKL
disrupts apical complex integrity and ookinete morphogenesis[18, 75]. A putative
MARK/MELK-related serine/threonine protein kinase (PBANKA_1305200), identified in
our phosphoproteomics analysis, may be directly or indirectly activated by NEK4 to
stimulate microtubule-associated proteins, thereby promoting or stabilising microtubule
formation at the APC and MTOC in the zygote. Together, these studies highlight a recurrent

feature of Apicomplexa: kinases and phosphatases govern centrosome and apical

organisation, and their perturbation uncouples nuclear and cortical events, resulting in

developmental arrest. The function of NEK4 fits within this broader mechanistic framework,

acting as a linchpin to preserve synchrony between meiotic and morphogenetic processes.

The perinuclear microtubules observed in early zygotes may be involved in the nuclear

movement described above and intranuclear chromosome movement (Fig. 6). In *S. pombe*,

horsetail nuclear movement is driven by SPB-derived microtubules interacting with the cell

cortex via dynein[34, 35]. We speculate that a similar mechanism generates nuclear movement in

Plasmodium, through the interaction of perinuclear microtubules with dynein (e.g, DHC3,

which was identified in our phosphoproteomics). Indeed, our preliminary observation that

PbDHC3 is largely located at the nuclear periphery in early zygotes strongly supports this

hypothesis (Extended Data Fig. 7 and Supplementary Video 8 and 9). While determining the

precise biophysical mechanics and the biological necessity of this movement for meiotic

completion will require future functional studies utilising dynein inhibitors and high-

resolution tracking of microtubule dynamics, the recruitment of DHC3 provides a compelling

molecular candidate for driving this nuclear motility. Furthermore, in eukaryotes including

mice, nematodes, and yeast, it is well established that meiotic chromosomes interact with

dynein on perinuclear microtubules via the SUN (Sad-1 and UNC-84)/KASH (Klarsicht,

ANC-1, and SYNE Homology) complex on the nuclear envelope, facilitating chromosome

movement and pairing[76]. Given that SUN proteins[77] and DHC3 localise to the nuclear

periphery during the zygote-to-ookinete development in *Plasmodium*, it is plausible that

SUN proteins mediate the interaction between chromosomes and dynein on perinuclear

microtubules, thereby driving intranuclear chromosome movement to promote homologous

chromosome pairing.

In an evolutionary context, NEK4 exemplifies how *Plasmodium* has streamlined but not

abandoned a central feature of meiotic control, to ensure synchrony between nuclear and

cytoskeletal events. The parasite has a compact kinome, and a limited number of four NEKs

have diversified to perform additional discrete yet complementary roles: for example, NEK1

regulates spindle formation during male gametogenesis[28], while NEK4 governs meiotic

initiation and zygote differentiation. This division of labour underscores the adaptive

flexibility of NEKs in the absence of canonical Polo-kinases and other regulators. The NEK4

regulatory module may be a *Plasmodium*-specific version of an ancient, conserved solution to

a universal cellular problem – to use specific waves of phosphorylation to coordinate
chromosomal replication with structural reorganisation of the cell during meiosis.

In conclusion, NEK4 functions as a key regulator that co-ordinates meiotic entry in
Plasmodium, coupling microtubule organisation, MTOC duplication, nuclear migration,
DNA replication, and chromosome condensation with the establishment of apical polarity
(Fig. 6). By integrating translational and phosphoregulatory control, NEK4 contributes to the
coordination of nuclear and cytoskeletal events, which is essential to drive zygote
morphogenesis and meiosis. These findings clarify the molecular requirements of meiosis in
Apicomplexa, highlighting a conserved principle – kinase-mediated coordination of nuclear
and cellular processes – that extends across the eukaryotic kingdom. Given that NEK4 is
essential for the development of transmission-stage parasites and the absence of close
orthologs from the human host, this kinase represents a promising transmission-blocking
therapeutic target. NEK4 provides a window to investigate how the minimalist signalling
networks of divergent parasites sustain their complex developmental programmes.

Methods

Ethics statement

All animal experiments were conducted in the United Kingdom in accordance with the Animals (Scientific Procedures) Act 1986 and approved by the Home Office under Project Licence numbers PDD2D5182 and PP3589958. Protocols were reviewed and approved by the institutional Animal Welfare and Ethical Review Body (AWERB) prior to implementation. Procedures were designed to minimise animal suffering and the number of animals used, in line with the 3Rs (Replacement, Reduction, Refinement) principles. Experiments were performed on female CD1 outbred mice aged 6 to 8 weeks, housed under standard conditions with environmental enrichment and monitored daily for health and welfare.

Generation of transgenic parasites

For C-terminal GFP-tagging of NEK4 (PBANKA_0616700) by single crossover homologous recombination, a 930 bp region of *nek4* downstream of the ATG start codon was amplified using primers 5'-CCCCGGTACCGATAGCGATAAGAGAGTAAGATTGTGTG-3' and 5'-CCCCGGGCCCAACATCAACAATATCCAATAATAATG-3' (Supplementary Table 17). Genotype analysis was performed using integration PCR as illustrated in Extended Data Fig. 1d. Primer 1 (intNEK4: 5'-CCAACCATATATTACACAGAGGTTAG-3') and primer 2 (ol492)[18] were used to confirm correct integration of the *gfp* sequence at the target locus. *P. berghei* ANKA line 2.34 parasites were then transfected by electroporation[78]; confirmation of the correct protein size was performed by Western blot analysis[28]. Production of the *Pbnek4-ko* parasite line has been described previously[19, 25].

Live cell imaging

To analyse PbNEK4-GFP expression during the zygote to ookinete transition, gametocytes or zygotes activated in ookinete culture medium were purified by Nycodenz gradient[19]. Zygote and ookinete development were maintained in continuous culture for up to 24 hours post-fertilisation. At specific time points, cell aliquots were taken and immediately mixed with a Cy3-conjugated mouse monoclonal antibody 13.1 (which targets the P28 surface protein)[79] and Hoechst 33342 prior to live-cell imaging. This approach ensured that the images represent distinct, representative live cells at each time point, by avoiding phototoxicity or non-physiological effects associated with prolonged exposure to the antibody and the DNA

dye. Zygote and ookinete development were tracked for up to 24 hours post-fertilisation to

assess the dynamics of PbNEK4-GFP location during meiotic progression and

morphogenesis. Images were captured on a Zeiss Axio Imager M2 microscope equipped with

a 63× oil immersion objective and an AxioCam ICc1 digital camera (Carl Zeiss).

Fluorescence signals from GFP and Hoechst were prone to bleaching, but imaging conditions

were optimised to ensure sufficient resolution to follow protein dynamics across key

developmental transitions. Axiovision software (Rel. 4.8) was used to enhance contrast and

minimise background, ensuring threshold settings remained consistent across samples.

Representative images from at least three independent biological experiments are presented,

with protein location assessed in 30–50 individual cells per time point. For the quantification

of PbNEK4-GFP localisation, data were derived from a total of 30–50 individual cells per

time point, collected across at least three independent biological experiments. To account for

variations in the microscopic focal plane (Z-axis) which can obscure simultaneous

observation of different cellular regions, cells were independently scored for the presence of a

strong/distinct focus at the nucleus and at the cell periphery. The percentage of cells positive

for distinct focal signals at each location was calculated and plotted.

Transmission electron microscopy

For ultrastructural analysis of NEK4-deficient and wild-type parasites, zygotes were

harvested at 2 and 4 hpa and fixed in 4% glutaraldehyde in 0.1 M phosphate buffer. Samples

were processed for transmission electron microscopy following the protocol described in the

previous study[80]. Briefly, fixed cells were post-fixed in osmium tetroxide, stained en bloc

with uranyl acetate, dehydrated through a graded ethanol series, and embedded in Spurr's

epoxy resin. Ultrathin sections (approximately 70 nm) were cut using a diamond knife,

mounted on copper grids, and stained with uranyl acetate followed by lead citrate to enhance

contrast. Sections were examined using a Tecnai G2 12 BioTwin (FEI UK, UK) or a JEOL

1200EX transmission electron microscope (JEOL, Japan) operated at 100 kV. Representative

images were captured digitally for analysis.

Serial block face scanning electron microscopy (SBF-SEM) of zygotes at 4 hpa

Samples were processed for SBF-SEM following the protocol described previously[81].

Fractions enriched in WT and *Pbnek4-ko* zygotes at 4 hpa were fixed at room temperature in

4% glutaraldehyde in 0.1 M phosphate buffer. Samples were then spun and washed three

times in 0.1 M phosphate buffer, and post-fixed in 1% osmium tetroxide in 1.5% potassium

ferrocyanide in 0.1 M phosphate buffer (for 45 min at room temperature, and in the dark).

After the first osmium step, samples were washed three times in 0.1 M phosphate buffer,

incubated in 1% tannic acid in 0.1 M phosphate buffer for 30 min at room temperature, and

then subjected to a second osmium step (2% OsO_4 in ddH_2O, for 30 min, at room

temperature, and in the dark). Samples were then incubated in 2% uranyl acetate in ddH_2O

for 2 h, dehydrated in acetone (a progressive series of acetone concentrations from 20%,

40%, 90% to 100% with three changes in molecular-sieved ultradry acetone over 4 h) and

embedded in TAAB 812 Hard resin (TAAB, catalogue number T030). The tips of resin

blocks containing samples were trimmed and mounted onto aluminium pins using conductive

epoxy glue and silver dag, and then sputter coated with a layer (10–13 nm) of gold, in an

Agar Auto Sputter Coater (Agar Scientific). Before SBF-SEM imaging, ultrathin sections

(70 nm) of the block face were examined in a Jeol JEM 1400 Flash transmission electron

microscope (JEOL), to verify sample quality. Samples were then imaged in a Merlin VP

compact high resolution scanning electron microscope (Zeiss) equipped with a 3View stage

(Gatan-Ametek), and an OnPoint back-scattered electron detector (Gatan-Ametek), in

variable pressure. The following imaging conditions were used: 2 kV, 20 μm aperture, 30

pascal variable pressure, 2 nm pixel size, 5 μs pixel time, 100 nm section thickness, 100%

FCC.

Ultrastructure expansion microscopy

Ultrastructure expansion microscopy (U-ExM) was performed on PbNEK4-GFP non-

activated gametocytes or zygotes at 2 or 4 hpa, following previously published protocols with

slight modifications[82, 83]. Parasites were fixed in 4% formaldehyde in MTSB buffer (10 mM

MES, 150 mM NaCl, 5 mM EGTA, 5 mM $MgCl_2$, 5 mM glucose, pH7.0) and adhered to 10

mm poly-D-lysine–coated round coverslips for 15 min. Coverslips were incubated overnight

at 4°C in 1.4% formaldehyde (FA)/2% acrylamide (AA). Gelation was performed in

ammonium persulphate/TEMED (10% each)/monomer solution (23% sodium acrylate; 10%

AA; 0.1% BIS-AA in PBS) on ice for 5 min and at 37°C for 30 min. Gels were denatured for

15 min at 37°C and for 90 min at 95°C in denaturation buffer (200 mM SDS, 200 mM NaCl,

50 mM Tris, pH 9.0, in water). After denaturation, gels were incubated in distilled water

overnight for complete expansion. The next day, circular gel pieces with a diameter of ~13

mm were excised, and the gels were washed in PBS three times for 15 min to remove excess

water. The gels were then incubated in blocking buffer (3% BSA in PBS) at room

temperature for 30 min, incubated with rabbit polyclonal anti-GFP antibody (1:250 dilution;

A11122; Invitrogen) or mouse monoclonal anti-α-tubulin antibody (1:500 dilution: T9026;
Sigma-Aldrich) in blocking buffer at 4°C overnight, and washed three times for 15 min in
wash buffer (0.5% vol/vol Tween-20 in PBS). The gels were incubated with 8 μg/ml Atto
594 NHS-ester (Merck), 10 μg/ml Hoechst 33342 (Molecular Probes), and Alexa Fluor 488
goat anti-rabbit IgG (A11008; Invitrogen) or Alexa Fluor 488 goat anti-mouse IgG (A11001;
Invitrogen) in PBS (1:500 dilution) at 37°C for 2.5 h followed by three washes of 15 min
each in wash buffer (blocking and all antibody incubation steps were performed with gentle
shaking). The gels were then washed three times for 15 min with wash buffer and expanded
overnight in ultrapure water. The expanded gel was placed in a 35-mm glass-bottom dish
(MatTek) with the 14-mm glass coated with poly-D-lysine. High-resolution images were
acquired on a Zeiss Celldiscoverer 7 with Airyscan using a 50×/1.2 water objective. Confocal
z-stacks were acquired using line scanning and the following settings: 55×55 nm pixel size,
170-nm z-step, 2.91 μs/pixel dwell time, 850 gain, and 3.5% (405 nm), 4.5% (488 nm), and
5.0% (561 nm) laser powers. The z-stack images were processed and analysed using Fiji
(version 1.54f). Each experiment was repeated at least three times, and protein localisation
was assessed in 5–10 cells per condition. Full imaging details are provided in the figure
legends.
753

Quantitative real-time PCR (qRT-PCR) analysis
Total RNA was extracted from WT and *Pbnek4-ko* zygotes at 2 hpa (three biological
replicates per condition) using the RNA Purification Kit (Stratagene), and cDNA was
synthesised using the RNA-to-cDNA Kit (Applied Biosystems). qPCR was performed with
80 ng of RNA using SYBR Green Fast Master Mix (Applied Biosystems) on an Applied
Biosystems 7500 Fast system. Cycling conditions were: 95 °C for 20 s, followed by 40
cycles of 95 °C for 3 s and 60 °C for 30 s. Primers were designed using Primer3
(https://primer3.ut.ee/). Each gene was tested in three biological and technical replicates.
hsp70 (PBANKA_081890) and *arginyl-tRNA synthetase* (PBANKA_143420) were used as
reference genes. Primer sequences are listed in Supplementary Table 17.
764

Transcriptome study using RNA-Sequencing
RNA samples from WT and *Pbnek4-ko* zygotes at 2 hpa (three biological replicates per
condition) were vacuum-concentrated by freeze drying and transported in RNA-stabilising
tubes (Biomatrica) to ensure preservation. Strand-specific mRNA libraries were prepared
using the TruSeq Stranded mRNA Library Prep Kit LT (Illumina). Paired-end sequencing (2

× 150 bp) was performed on an Illumina HiSeq 4000 platform. Quality control of raw
sequencing reads was carried out using FASTQC
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc). Adapter trimming and removal
of low-quality sequences were performed using Trimmomatic. Cleaned reads were aligned to
the *P. berghei* ANKA reference genome (PlasmoDB release 40) using HISAT2 version 2.1.0,
applying the parameter --rna-strandness FR. Gene-level expression was quantified using
FeatureCounts. Raw read counts were converted to counts per million (CPM), and genes with
CPM <1 in all but one replicate were excluded from downstream analysis. Library sizes were
normalised using the trimmed mean of M-values (TMM) method in the EdgeR package.
Processed count data were then analysed using the voom transformation from the limma
package to model the mean–variance relationship. Differential expression analysis was
performed with DESeq2, using a false discovery rate (FDR)-adjusted p-value threshold of
0.05 (Benjamini–Hochberg correction) and a fold-change cut-off of ≥2 to define significantly
regulated genes.

Immunoprecipitation and mass spectrometry
Purified gametocytes from WT-GFP and PbNEK4-GFP parasites were activated for 2 h and
proteins were crosslinked using 1% formaldehyde for 10 min at room temperature. Excess
aldehyde was quenched by incubating samples in 0.125 M glycine for 5 min, followed by
three washes with PBS (pH 7.5). Immunoprecipitation of PbNEK4-GFP complexes was
performed using the GFP-Trap_A kit (Chromotek) in accordance with the manufacturer's
protocol. GFP-bound protein complexes were captured on beads and processed for liquid
chromatography–tandem mass spectrometry (LC-MS/MS). Beads were washed and
incubated in ammonium bicarbonate (ABC) buffer. Reduction and alkylation were carried out
using 10 mM TCEP (tris(2-carboxyethyl)phosphine hydrochloride) and 40 mM
chloroacetamide (CAA) for 5 min at 70°C. Proteins were then digested with trypsin overnight
at room temperature (1 μg trypsin per 100 μg of protein). Digestion was halted by adjusting
the pH to 3–4 with 1% trifluoroacetic acid (TFA) before mass spectrometry analysis. Tryptic
peptides were analysed via LC-MS/MS. Resulting raw data were searched using FragPipe
version 21.0 (https://fragpipe.nesvilab.org/) against the PlasmoDB-
66_PbergheiANKA_AnnotatedProteins database. Search parameters included trypsin
digestion with up to two missed cleavages, fixed carbamidomethylation of cysteines, and
variable modifications for methionine oxidation and N-terminal acetylation. Protein
identification and quantification were performed using Scaffold (version 5.3.3, Proteome

Software), and annotations were derived from PlasmoDB. Proteins were retained for analysis
if they had ≥2 unique peptides and valid quantification in at least two replicates. The analysis
was performed using two independent biological replicates.

Phosphoproteomic analysis

Protein Extraction and Digestion via SDS-FASP. Zygotes of WT and *Pbnek4-ko* lines at 0
and 2 hpa were lysed in 400 μl of buffer containing 2% SDS, 25 mM NaCl, 50 mM Tris-
HCl (pH 7.4), 2.5 mM EDTA, and 20 mM TCEP, supplemented with a protease and
phosphatase inhibitor cocktail (Halt, Thermo Fisher Scientific). Samples were vortexed and
heated at 95°C for 10 min with continuous mixing at 400 rpm. Genomic DNA was
fragmented using four 10-second sonication pulses at 50% amplitude. Following
centrifugation at 17,000 g for 30 min, supernatants were collected, and protein
concentrations were determined using the Pierce 660nm Protein Assay Kit. For alkylation the
volume corresponding to 1 mg of protein for each sample was adjusted to 300 μl and then
incubated with 48 μl of 0.5 M iodoacetamide for 1 h at room temperature in the dark.
Protein digestion was performed using the filter-aided sample preparation (FASP) method
with Amicon Ultra-4 30 kDa centrifugal filter units (Millipore). Trypsin (Promega) was
added at a 1:80 enzyme-to-protein ratio, and digestion proceeded overnight at room
temperature. Resulting peptides were desalted using Pierce Peptide Desalting Spin Columns
(Thermo Fisher Scientific) according to the manufacturer's instructions and dried using a
speed vacuum concentrator.

TMT-11plex Labelling. Peptide concentrations were measured using the Pierce Quantitative
Colorimetric Peptide Assay (Thermo Fisher Scientific). For each sample, 160 μg of peptides
were labelled with 800 μg of TMT reagents (Thermo Fisher Scientific), previously
dissolved in 220 μl of 36% acetonitrile and 200 mM EPPS buffer (pH 8.5). Labelling
reactions were carried out for 1 h at room temperature and quenched by adding
hydroxylamine to a final concentration of 0.3% (v/v). Labelled samples were pooled, desalted
using Pierce Peptide Desalting Spin Columns, and dried under vacuum.

Phosphopeptide Enrichment. Phosphopeptides were enriched using the High-Select Fe-
NTA Phosphopeptide Enrichment Kit (Thermo Fisher Scientific) following the
manufacturer's protocol. Both the enriched phosphopeptide fractions and the flow-through
were desalted using Pierce Peptide Desalting Spin Columns and dried using a speed vacuum
concentrator.

LC-MS/MS Analysis. Dried peptides were reconstituted in 5% acetonitrile with 0.1% formic

acid, and 2 µg of each sample was injected into a Vanquish NEO liquid chromatography

system (Thermo Fisher Scientific) coupled to an Orbitrap Fusion Lumos Tribrid mass

spectrometer (Thermo Fisher Scientific). Peptides were first trapped on an Acclaim PepMap

100 C18 trap column (75 µm × 20 mm, 3 µm particle size) and then separated on a

homemade analytical column (75 µm × 500 mm) packed with ReproSil-Pur C18 resin

(1.9 µm, 100 Å, Dr. Maisch GmbH). The analytical separation was run for 180 min using a

gradient of H_2O/FA 99.9%/0.1% (solvent A) and CH_3CN/H_2O/FA 80.0%/19.9%/0.1%

(solvent B). The gradient was run from 5 % B to 28 % B in 160 min, then to 40% B in 20

min, then to 99% B in 10 min with a final stay of 20 min at 99 % B. Flow rate was of 250

nL/min an total run time was of 210 min. Data-Dependent Acquisition (DDA) was

performed with MS1 full scan at a resolution of 120'000 FWHM followed by as many

subsequent MS2 scans on selected precursors as possible within 3 second maximum cycle

time. MS1 was performed in the Orbitrap with an AGC target of 4×10^5, a maximum injection

time of 50 ms and a scan range from 375 to 1500 m/z. MS2 was performed in the Orbitrap at

a resolution of 50'000 FWHM using higher-energy collisional dissociation HCD at 38%

NCE. Isolation windows was at 0.7 u with an AGC target of 5×10^4 and a maximum injection

time of 86 ms. A dynamic exclusion of parent ions of 60 s. with 10 ppm mass tolerance was

applied.

Data Processing and Analysis. Raw data files were processed using Proteome Discoverer

software version 2.4 (Thermo Fisher Scientific). Spectra were searched against the *P. berghei*

ANKA protein database (PlasmoDB.org release 68), the *mus musculus* reference proteome

database (UniProt, reviewed, release 2024_06, 17'207 entries) and an in-house database of

common contaminants using the Mascot search engine (version 2.6.2, Matrix Science).

Search parameters included trypsin as the digestion enzyme with up to one missed cleavage

allowed, a precursor mass tolerance of 10 ppm, and a fragment mass tolerance of 0.02 Da.

Carbamidomethylation of cysteine (+57.021 Da) and TMT modification of peptide N-

termini and lysine residues (+229.163 Da) were set as fixed modifications, while oxidation

of methionine (+15.995 Da) and phosphorylation of serine, threonine, and tyrosine

(+79.966 Da) were set as variable modifications. Peptide-spectrum matches (PSMs) and

peptides were validated using the Percolator algorithm, applying a false discovery rate (FDR)

of 1%. Proteins were inferred from the identified peptides and filtered to achieve an FDR of

1%. Quantitative information was extracted using reporter ion intensities from TMT tags, and

protein abundances were normalized based on the total peptide amount and scaled across all

samples. Protein ratios were calculated using the summed abundances-based approach across
groups, and statistical significance was assessed using ANOVA. For the analysis, three
independent biological replicates were used for the 0 hpa *Pbnek4-ko*, 2 hpa WT, and 2 hpa
Pbnek4-ko samples, while two independent biological replicates were used for the 0 hpa WT
sample.
876
Gene and protein annotation
Gene and protein annotations, including predicted functions and structural features, were
primarily retrieved from PlasmoDB (release 68) and UniProt (version 23). Domain
architectures and specific sequence motifs were predicted using InterPro
(https://www.ebi.ac.uk/interpro/). For the uncharacterised conserved *Plasmodium* protein
PBANKA_0806700, structural prediction was performed using the AlphaFold3 server
(https://alphafoldserver.com/)[41]. The predicted structure was subsequently used as a query in
Foldseek (https://search.foldseek.com/search)[42] to identify structurally similar proteins,
revealing the presence of putative HEAT-like repeats. Where applicable, functional
annotations were based on previously published experimental characterisations or homology-
based predictions, which are explicitly cited. A comprehensive summary of the key proteins
highlighted in this study, including their identified domains, sizes, and the basis of their
annotations, is provided in Supplementary Table 18.
890
Data availability
The mass spectrometry proteomics data have been deposited to the ProteomeXchange
Consortium via the PRIDE[84] partner repository with the dataset identifier PXD070965 and
10.6019/PXD070965 (proteomics and phosphoproteomics) and PXD070161 and
10.6019/PXD070161 (GFP immunoprecipitations). RNAseq data have been deposited to the
Gene Expression Omnibus under accession number the BioProject ID PRJNA1354107. The
authors declare that all other relevant data generated or analysed during this study are
included in the article or its supplementary information. Raw data are available for each
figure. Materials are available from the corresponding authors on reasonable request.
900
Figure Legends
Figure 1. NEK4 is located at the MTOC, APC, and nucleus during the early meiotic
phase of zygote to ookinete development. a. A schematic diagram of ookinete development
showing formation of gametocytes, gametes, and zygote, and the involvement of NEK4 in

cellular development and meiosis. The developmental stage and nuclear ploidy (from 1N to
4N) are indicated for each phase. Activation of gametocytes in the mosquito gut initiates
sexual development. In the schematic, the red blood cell membrane is depicted in red, the
parasite cytoplasm in grey, and the nucleus as a black circle. The time scale in hours post-
activation (hpa) is also indicated for the developmental stages from zygote to retort and
ookinete. **b.** Live-cell imaging of PbNEK4-GFP location at different time points (hpa). Green
arrowheads indicate the PbNEK4-GFP foci in the nucleus (Hoechst, blue) and at the cell
periphery. Insets show magnification of the cell periphery (C) and nucleus (N). The yellow
arrowhead indicates the cytoplasmic foci of PbNEK4-GFP. Images are representative of 30–
50 cells analysed across at least 3 independent biological experiments. **c.** Schematic summary
of the location of PbNEK4-GFP shown in (b), indicating the signal strength in the
nucleoplasm, nuclear focus, and at the cell periphery focus over time. **d.** Expansion
microscopy of PbNEK4-GFP zygotes at 2 hpa and 4 hpa. The cells were labelled with NHS-
ester (gray), anti-GFP antibody (green), and Hoechst (blue). The whole-cell images display
the MTOC and APC regions enclosed by superimposed magenta and cyan dashed boxes,
respectively. The corresponding magnified views of these regions are shown below. The
positions of the MTOC and APC are indicated by magenta and cyan arrows, respectively.
Representative images from three independent biological experiments (n ≥ 5 cells examined
per condition). **e.** Model illustrating the localisation of NEK4 at the MTOCs, APC and
nucleoplasm.

Figure 2. NEK4-localised MTOC leads nuclear movement, and directional microtubule
formation occurs at both the MTOC and the APC during early meiosis. a. Time-lapse
imaging of a PbNEK4-GFP expressing zygote shows nuclear movement. The PbNEK4-GFP
foci in the nucleus and at the cell periphery are indicated by magenta and cyan arrowheads,
respectively. The lower panel ("Tracing") tracks the position of the PbNEK4-GFP focus
(green dot) and the nucleus (blue) over 540 seconds. **b.** Time-lapse imaging of a zygote
expressing the microtubule plus-end/MTOC marker PbEB1-GFP (green). The PbEB1-GFP
foci in the nucleus and at the cell periphery are indicated by magenta and cyan arrowheads,
respectively. The lower panel ("Tracing") tracks the position of the PbEB1-GFP focus (green
dot) and the nucleus (blue) over 540 seconds. **c.** Quantification of the percentage of zygotes
showing nuclear movement at 0, 2, 4, and 6 hpa. The data represent the mean ± SEM from
three independent experiments (n = 3), in which 30 cells were analysed per experiment. Black

dots show the result from each experiment. **d.** Schematic model of nuclear movement,
illustrating the association of NEK4/EB1 foci with the moving nucleus at 2-4 hpa. **e.**
Expansion microscopy showing microtubule organisation in developing zygotes.
Microtubules (α-tubulin, green) are shown at 2 hpa with either one or two MTOCs, and at 4
hpa with separated MTOCs. The whole-cell images display the MTOC and APC regions
enclosed by superimposed magenta and cyan dashed boxes, respectively. The corresponding
magnified views of these regions are shown below. Representative images from three
independent biological experiments (n ≥ 5 cells examined per condition). The positions of the
MTOC and APC are indicated by magenta and cyan arrows, respectively. **f.** Model of the
microtubule organisation in a developing zygote, illustrating the formation of perinuclear and
APC microtubules and the location of PbNEK4-GFP at the MTOCs, APC and nucleoplasm.
949
Figure 3. NEK4 is essential for DNA replication, nuclear migration, microtubule
formation, and morphological transformation from zygote to ookinete. a. Live-cell
imaging of control (PbNEK4-GFP) and *Pbnek4-ko* parasites from 2 to 20 hpa. Representative
live-cell images of different cells sampled at specific time points are shown. Parasites were
labelled with Hoechst (blue) and a Cy3-conjugated 13.1 antibody (magenta), which
recognises P28 protein on the surface of zygotes and ookinetes, immediately prior to imaging.
Unlike the control which develops into an elongated ookinete, *Pbnek4-ko* parasites remain
round and fail to develop. Images are representative of 30–50 cells analysed across at least 3
independent biological experiments. **b.** Quantification of nuclear movement in control
(PbNEK4-GFP, blue) and *Pbnek4-ko* (gray) zygotes at 2- and 4 hpa. Nuclear movement is
significantly reduced in the *Pbnek4-ko* mutant both at 2- and 4 hpa. The data represent the
mean ± SEM from three independent experiments (n = 3), in which 30 cells were analysed
per experiment. Black dots show the result from each experiment. P-values were determined
by Welch's t-test. **c.** Expansion microscopy comparing microtubule organisation (α-tubulin,
green) in WT and *Pbnek4-ko* zygotes at 2 and 4 hpa. The whole-cell images display the
MTOC and APC regions enclosed by superimposed magenta and cyan dashed boxes,
respectively. The corresponding magnified views of these regions are shown below. The
positions of the MTOC and APC are indicated by magenta and cyan arrows, respectively.
Pbnek4-ko zygotes show a single MTOC and disrupted microtubule formation, in contrast to
the well-organised perinuclear and APC microtubules in WT zygotes. Representative images
from three independent biological experiments (n ≥ 5 cells examined per condition). **d.**

Schematic model summarising the defects in *Pbnek4-ko* zygotes, including the persistence of a single MTOC, disrupted microtubule formation, and reduced nuclear movement.

Figure 4. Electron microscopy shows that cytoskeletal development and condensed chromosome formation during early meiosis are blocked in *Pbnek4-ko* zygotes. a. TEM images of WT and *Pbnek4-ko* zygotes at 4 hpa. N, nucleus; APC, apical polar complex; APR, Apical polar ring; A-Mt, APC microtubule; NM, nuclear membrane. Magenta arrowheads show condensed chromosomes. Representative images from three independent biological experiments (n ≥ 20 cells examined per condition). **b.** U-ExM images of WT and *Pbnek4-ko* zygotes at 4 hpa, stained with NHS-ester (gray) and Hoechst (blue). A maximum intensity z-projection was generated using a subset of optical slices encompassing the nucleus (6 slices, 170-nm intervals) from the same datasets shown in Fig. 3c. Whole-cell images are shown at the top, and magnified views of the nuclear regions enclosed by magenta boxes are presented below. Magenta arrows indicate the positions of the MTOCs. Tracing images illustrate the thread-like structures within the WT nucleus traced with yellow lines based on the NHS-ester staining, demonstrating that these structural axes largely coincide with the Hoechst DNA signals. Representative images from three independent biological experiments (n ≥ 5 cells examined per condition). **c.** Slice images of WT and *Pbnek4-ko* zygotes at 4 hpa, obtained by SBF-SEM. The whole-cell view shows the development of the APC in the WT zygote. The magnified views of the nucleus show the presence of thread-like condensed chromosomes (magenta arrowheads) and the formation of potential synaptonemal complexes (cyan arrowheads), where these condensed chromosomes form pairs via a central linear structure. The insets show magnified views of a thread-like condensed chromosomes and synaptonemal complex. These developmental features were completely absent in *Pbnek4-ko* zygotes. **d.** Quantification of the percentage of cells with a developing APC and condensed chromosomes in WT and *Pbnek4-ko* zygotes at 4 hpa. The data represent the mean ± SEM from three independent SBF-SEM datasets (n = 3), in which 30 cells were analysed per dataset. with black dots representing the results from each dataset. **e.** 3D models of WT and *Pbnek4-ko* zygotes at 4 hpa, reconstructed from SBF-SEM datasets. The WT model illustrates the developed APC (orange) with APR (gray), the presence of condensed, thread-like chromosomes within the nucleus (blue; some of which are attached to the nuclear envelope), and the separating MTOCs (purple). In contrast, the *Pbnek4-ko* zygote model shows that none of these developmental events have occurred.

1004

Figure 5. Global transcriptomic, proteomics and phosphoproteomic analysis reveals

NEK4-dependent transcriptional and phosphorylation events during *Plasmodium*

zygote activation. a. Volcano plot showing differentially expressed transcripts in *Pbnek4-ko*

parasites compared to WT controls at 2 hpa (RNA-seq). The $\log_2$ fold change (FC) is plotted

against $-\log_{10}$(adjusted p-value). Red and blue dots indicate significantly upregulated and

downregulated genes, respectively. *NEK4* and key meiotic genes that are significantly

downregulated in *Pbnek4-ko* at 2 hpa are highlighted with yellow dots and gene names.

Dotted lines indicate the threshold for statistical significance (adjusted p-value $\leq$ 0.05 and

fold change $\geq$ 1.5). The underlying data are provided in Supplementary Table 1. **b.** Gene

Ontology (GO) analysis of differentially expressed transcripts by RNA-seq in *Pbnek4-ko*

versus WT zygotes at 2 hpa. Genes in each term are listed in Supplementary Table 2. The

analysis was performed using GO slim terms in PlasmoDB. BP = Biological Process; CC =

Cellular component; MF = Molecular Function. **c.** Table displaying selected proteins,

corresponding gene ID and representation by the number of peptides from the proteins

identified in a GFP-trap pulldown assay with precipitates of either PbNEK4-GFP or WT-GFP

zygotes at 2 hpa. **d-g.** Volcano plots displaying changes in total protein and phosphopeptide

abundance. The $\log_2$ fold change (derived from the average of three biological replicates

each) is plotted against the $-\log_{10}$ p-value. The dotted lines indicate the threshold for

statistical significance (p-value $\leq$ 0.05 and fold change $\geq$ 1.5). Significantly upregulated and

downregulated proteins/phosphopeptides are shown as red and blue dots, respectively.

Proteins/phosphopeptides with non-significant changes are shown as grey dots. **d.**

Comparison of protein abundance between WT gametocytes at 0 hpa and WT zygotes at 2

hpa. **e.** Comparison of protein abundance between *Pbnek4-ko* and WT zygotes at 2 hpa. **f.**

Comparison of phosphopeptide abundance between WT gametocytes at 0 hpa and WT

zygotes at 2 hpa. **g.** Comparison of phosphopeptide abundance between *Pbnek4-ko* and WT

zygotes at 2 hpa. Key phosphopeptides from proteins of interest that are significantly

upregulated in WT at 2 hpa (f) and downregulated in the *Pbnek4-ko* mutant at 2 hpa (g) are

highlighted with yellow dots and protein names. **h-i.** GO analysis of differentially abundant

phosphopeptides in WT zygotes at 2 hpa versus 0 hpa (h) and *Pbnek4-ko* versus WT zygotes

at 2 hpa (i). Genes in each term are listed in Supplementary Table 8 and 10. The analysis was

performed using GO slim terms in PlasmoDB. **j.** Summary table displaying selected proteins

with significant changes in phosphopeptide abundance from the phosphoproteomics screen.

Protein names highlighted with a green background indicate that these proteins were also
independently identified in the PbNEK4-GFP pulldown. The $\log_2$ fold change ($\log_2$FC)
values of identified phosphopeptides are represented by a colour gradient from red (decrease)
to blue (increase). Significant $\log_2$FC values (p-value ≤ 0.05) are shown in black text, while
non-significant values are in grey. In the phosphopeptide sequences, the regions
corresponding to the canonical NEK kinase motif ([LMFW]-X-X-S/T-[no P])[55] are
highlighted with a green background. Within these motifs, the hydrophobic residues at the -3
position are indicated in blue font, and the target phosphorylated serine/threonine residues are
indicated in orange font.

1046

Figure 6. *Plasmodium* NEK4 acts as a key regulator driving the early events of meiosis.
A model illustrating the central function of NEK4 in coordinating key developmental
processes during the zygote-to-ookinete development. The top panel shows the canonical
meiotic stages in the model organism *Schizosaccharomyces pombe* (*S. pombe*) for
comparison. The middle panel depicts the predicted steps and timing of meiosis in
Plasmodium from zygote formation, meiosis I to meiosis II, based on previous observations.
In *Pbnek4-ko* mutants, DNA replication is impaired, blocking progression to meiotic
prophase. In *S. pombe*, active nuclear movement, known as horsetail nuclear movement,
occurs during prophase, followed by chromosome pairing and crossover mediated by linear
elements (LinEs). In contrast, our observations in *Plasmodium* suggest that between 2 and 4
hpa, active nuclear movement similar to the horsetail movement occurs concurrently with the
formation of structures resembling synaptonemal complexes between condensed
chromosomes. The bottom panel proposes that NEK4 regulates four crucial modules: (1)
MTOC/Microtubule dynamics (e.g. MARK/MELK, MORN), (2) Meiosis-specific proteins
(e.g. REC8, HOP1), (3) Sexual development factors (e.g., FD2, AP2-O), (4)
DNA/chromosome functions (e.g. RCC, POLD3), (5) Uncharacterised protein (putative
HEAT-repeat protein; PBANKA_0806700). Through these pathways, NEK4 orchestrates
MTOC separation, microtubule formation at MTOC and APC, DNA replication, and
chromosome condensation, which are all essential for successful meiosis and subsequent
ookinete development. Based on the function of perinuclear microtubules in other eukaryotes
during meiosis, the perinuclear microtubules observed in *Plasmodium* zygotes are likely
associated with nuclear movement or intranuclear chromosome movement. We predict that
effectors such as cytoplasmic dynein and SUN protein mediate these functions.

1070

Extended Data Fig. legend

Extended Data Fig. 1. *Plasmodium* **NEK phylogeny, domain structure, function, and**

generation of PbNEK-GFP parasites. a. Phylogenetic tree of NIMA-related protein kinases

(NEKs) from *Plasmodium berghei* (Pb, light blue; PbNEK4, blue), *Plasmodium falciparum*

(Pf, red), *Homo sapiens* (Hs, green), including NIMA (black) from *Aspergillus nidulans*. **b.**

Domain structures of PbNEKs. **c.** Summary table of the function, expression stage, and

localisation of PbNEKs. **d.** Schematic representation of the endogenous *nek4* locus, the GFP-

tagging construct, and the recombined *nek4* locus following single homologous

recombination. Arrows indicate the position of PCR primers used to confirm successful

integration of the construct. **e.** Diagnostic PCR of *nek4* and WT-GFP parasites using the

diagnostic primers to show the correct integration. Integration of the *nek4* tagging construct

gives a band of ~1400 bp. **f.** Western blot showing the expression of endogenous PbNEK4-

GFP detected by anti-GFP antibody. The positions corresponding to the molecular weights of

GFP (~27 kDa) and PbNEK4-GFP (~74 kDa) are indicated by arrows, respectively.

1085

Extended Data Fig. 2. PbNEK4-GFP localisation dynamics during zygote-ookinete

development. Live-cell imaging of PbNEK4-GFP location at different time points (minutes

or hours post-activation, mpa or hpa). Representative live-cell images of different cells

sampled at specific time points are shown. PbNEK4-GFP (green) parasites were labelled with

Hoechst (blue) and a Cy3-conjugated 13.1 antibody (magenta), which recognises P28 protein

on the surface of zygotes and ookinetes, immediately prior to imaging. Images are

representative of 30–50 cells analysed across at least 3 independent biological experiments.

1093

Extended Data Fig. 3. Analyses of PbNEK4-GFP localisation dynamics. a. Line graph

showing the percentage of cells exhibiting a distinct PbNEK4-GFP focus in the nucleus (blue

line) and at the cell surface (orange line) from pre-activation (0 hpa) to 24 hpa.

Representative live-cell images of a 1 hpa zygote (PbNEK4-GFP alone and a GFP/Hoechst

merge) are shown adjacent to the graph to illustrate the scoring. Orange and blue arrowheads

indicate the distinct PbNEK4-GFP foci at the cell surface and within the nucleus,

respectively. The y-axis represents the percentage of cells positive for the focal signal at each

location. Scoring for the two locations was performed independently. Note that due to the

spherical nature of the early zygote, simultaneously capturing both the nucleus and the cell

surface in sharp microscopic focus (Z-plane) is technically challenging. Consequently, cells

lacking an optimal focal plane for either region may have been scored as negative, meaning
the percentages presented here might represent a slight underestimation. PbNEK4-GFP
localisation was assessed in 30–50 individual cells per time point from at least three
independent parasite preparation. **b.** Time-lapse imaging of a focal dot-like PbNEK4-GFP-
positive structure in the Hoechst (blue)-stained PbNEK4-GFP (green) zygotes at 5 hpa. The
focal dot-like PbNEK4-GFP-positive structure and the PbNEK4-GFP foci in the nucleus and
at the cell periphery are indicated by yellow, magenta, and cyan arrowheads, respectively.
Images are representative of 30–50 cells analysed across at least 3 independent biological
experiments. **c.** Live-cell imaging of a focal dot-like PbNEK4-GFP-positive structure in the
Hoechst (blue)-stained PbNEK4-GFP (green) developing ookinete at 10 hpa. The yellow
arrowhead indicates the focal dot-like PbNEK4-GFP-posivite structure. Images are
representative of 30–50 cells analysed across at least 3 independent biological experiments.
1116
Extended Data Fig. 4. Live-cell imaging and expansion microscopy of early zygotes. a.
PbNEK4-GFP and PbEB1-mCherry zygotes at 2 hpa. PbNEK4-GFP, PbEB1-mch, and
colocalisation foci were indicated with green, magenta, and white arrow heads, respectively.
Images are representative of at least 30 cells analysed across at least 3 independent biological
experiments. **b.** Expansion microscopy of the zygote at 2 hpa displaying single MTOC with
microtubule structures extending from it. Representative images from three independent
biological experiments (n ≥ 5 cells examined). **c.** Zygotes activated for 2 hours displayed the
development of the APC and MTOCs, along with microtubule structures extending form
them. Representative images from three independent biological experiments (n ≥ 5 cells
examined).
1127
Extended Data Fig. 5. Generation and genotypic and phenotypic analysis of *Pbnek4-ko*
parasites. a. Schematic representation of the endogenous *nek4* locus, the targeting knockout
construct and the recombined *nek4* locus following double homologous crossover
recombination. Arrows P1 and P2 indicate PCR primers used to confirm successful
integration in the *nek4* locus following recombination, and arrows P3, 4, 5 and 6 indicate
PCR primers used to show deletion of the *Pbnek4* gene. **b.** Integration PCR of the *nek4* locus
in WT and *Pbnek4-ko* parasites using the primers, P1-6. **c.** TEM images of WT and *Pbnek4-*
ko zygotes at 2 hpa. N, nucleus; APC, apical polar complex; A-Mt, APC microtubule; Pn-Mt,
perinuclear microtubule; NM, nuclear membrane. Magenta box show condensed
chromosomes. Representative images from three independent biological experiments (n ≥ 20

cells examined per condition). **d, e.** NHS-ester staining (gray) revealed that the apical polar
complex (APC) had not yet formed in both WT (c) and *Pbnek4-ko* (d) female gametocytes
before activation. A structure presumed to be the MTOC was also visible (magenta
arrowhead), although it appeared smaller compared to that in zygotes at 2 hpa. Furthermore,
anti α-tubulin antibody (green) and Hoechst (blue) staining revealed that no microtubule
formation was observed around the potential MTOC and nucleus. Representative images
from three independent biological experiments (n ≥ 5 cells examined per condition).

1145

Extended Data Fig. 6. qRT-PCR analysis of *Pbnek4-ko* and WT zygotes at 2 hpa and
proteomic and phosphoproteomic analysis of *Pbnek4-ko* and WT zygotes at 0 hpa and
***Pbnek4-ko* gametocytes at 0 hpa and *Pbnek4-ko* zygotes at 2 hpa. c.** Expression level
validation of *nek4* and meiotic related genes using qRT-PCR in *Pbnek4-ko* and WT zygotes
at 2 hpa. Shown is mean ± SEM; n = 3 independent experiments. Multiple comparisons *t* test,
with post hoc test of Holm–Sidak showed significant differences in relative expression.
*adjusted p-value $<$ 0.05, **adjusted p-value $<$ 0.01. **b-e.** Volcano plots displaying
changes in total protein and phosphopeptide abundance between *Pbnek4-ko* and WT zygotes
at 0 hpa and between *Pbnek4-ko* gametocytes at 0 hpa and *Pbnek4-ko* zygotes at 2 hpa. The
$\log_2$ fold change (derived from the average of three biological replicates each) is plotted
against the -$\log_{10}$ p-value. The dotted lines indicate the threshold for statistical significance
(p-value ≤ 0.05 and fold change ≥ 1.5). Significantly upregulated and downregulated
proteins/phosphopeptides are shown as red and blue dots, respectively.
Proteins/phosphopeptides with non-significant changes are shown as grey dots. **b.**
Comparison of protein abundance between *Pbnek4-ko* and WT zygotes at 0 hpa. **c.**
Comparison of protein abundance between *Pbnek4-ko* gametocytes at 0 hpa and *Pbnek4-ko*
zygotes at 2 hpa. **d.** Comparison of phosphopeptide abundance between *Pbnek4-ko* and WT
zygotes at 0 hpa. **e.** Comparison of phosphopeptide abundance between *Pbnek4-ko*
gametocytes at 0 hpa and *Pbnek4-ko* zygotes at 2 hpa. Key phosphopeptides from proteins of
interest that are significantly upregulated in WT at 2 hpa (f) and downregulated in the
Pbnek4-ko mutant at 2 hpa (g) are highlighted with yellow dots and protein names.

1167

1168 **Extended Data Fig. 7. PbDHC3-GFP localisation dynamics during zygote-ookinete**
1169 **development.** Live-cell imaging of PbDHC3-GFP location at different time points.
1170 Representative live-cell images of different cells sampled at specific time points are shown.

PbDHC3-GFP (green) parasites were labelled with Hoechst (blue) and a Cy3-conjugated 13.1 antibody (magenta), which recognises P28 protein on the surface of zygotes and ookinetes, immediately prior to imaging. Magenta and cyan arrowheads indicate PbDHC3-GFP foci around the nucleus and PbDHC3-GFP localisation on the cortical region, respectively. Images are representative of 30–50 cells analysed across at least 3 independent biological experiments.

Supplementary Table 1: RNA-Seq data analysis of *Pbnek4-ko* vs WT parasite 2 hpa

Supplementary Table 2: Gene Ontology Analysis of RNA-Seq data

Supplementary Table 3: Quantitative proteomics of WT 2 hpa versus WT 0 hpa parasites

Supplementary Table 4: Quantitative proteomics of *Pbnek4-ko* 2 hpa versus WT 2 hpa parasites

Supplementary Table 5: Quantitative proteomics of *Pbnek4-ko* 0 hpa versus WT 0 hpa parasites

Supplementary Table 6: Quantitative proteomics of *Pbnek4-ko* 2 hpa versus *Pbnek4-ko* 0 hpa parasites

Supplementary Table 7: Quantitative phosphoproteomics of WT 2 hpa versus WT 0 hpa parasites

Supplementary Table 8: Gene Ontology Analysis of phosphoproteomic data for WT 2 hpa versus WT 0 hpa parasites

Supplementary Table 9: Quantitative phosphoproteomics of *Pbnek4-ko* 2 hpa versus WT 2 hpa parasites

Supplementary Table 10: Gene Ontology Analysis of phosphoproteomic data for *Pbnek4-ko* 2 hpa versus WT 2 hpa parasites

Supplementary Table 11: Quantitative phosphoproteomics of *Pbnek4-ko* 2 hpa versus *Pbnek4-ko* 0 hpa parasites

Supp. Table 12: Quantitative phosphoproteomics of *Pbnek4-ko* 0 hpa versus WT 0 hpa parasites

Supplementary Table 13: GFP-Trap downs of PbNEK4-GFP parasites 2 hpa

Supplementary Table 14: Phosphatases and kinases identified in the *Pbnek4-ko* phosphoproteomics

Supplementary Table 15: AP2-Z target genes significantly altered in the *Pbnek4-ko* RNA-seq data

1205 **Supplementary Table 16: AP2-O target genes significantly altered in the *Pbnek4-ko***
1206 **RNA-seq data**
1207 **Supplementary Table 17: Primers used in this study**
1208 **Supplementary Table 18: Summary of key proteins highlighted in this study and their**
1209 **functional annotations**
1210

1211 **Supplementary Video 1**
1212 Time-lapse video microscopy of a focal dot-like PbNEK4-GFP-positive structure in the
1213 Hoechst (blue)-stained PbNEK4-GFP (green) zygotes at 5 hpa. Video is played back at 20×
1214 speed. Scale bar = 2 μm.
1215

1216 **Supplementary Video 2**
1217 Time-lapse video microscopy of nuclear movement in the Hoechst (blue)-stained PbNEK4-
1218 GFP (green) zygotes at 2 hpa. Video is played back at 50× speed. Scale bar = 5 μm.
1219

1220 **Supplementary Video 3**
1221 Time-lapse video microscopy of nuclear movement in the single zygote from Supplementary
1222 Video 1 (blue; Hoechst, green; PbNEK4-GFP). Video is played back at 50× speed. Scale bar
1223 = 5 μm.
1224

1225 **Supplementary Video 4**
1226 Time-lapse video microscopy of nuclear movement in the Hoechst (blue)-stained PbEB1-
1227 GFP (green) zygotes at 2 hpa. Video is played back at 50× speed. Scale bar = 5 μm.
1228

Supplementary Video 5
Time-lapse video microscopy of nuclear movement in the single zygote from Supplementary
Video 3 (blue; Hoechst, green; PbEB1-GFP). Video is played back at 50× speed. Scale bar =
5 μm.
Supplementary Video 6
Time-lapse video microscopy of moving nuclei in the Hoechst (blue)-stained PbNEK4-GFP
zygotes at 4 hpa. Video is played back at 20× speed. Scale bar = 5 μm.
1236

1237 **Supplementary Video 7**

Time-lapse video microscopy of static nuclei in the Hoechst (blue)-stained *Pbnek4-ko* zygotes at 4 hpa. Video is played back at 20× speed. Scale bar = 5 μm.

Supplementary Video 8

Time-lapse video microscopy of nuclear movement in the Hoechst (blue)-stained PbDHC3-GFP (green) zygotes at 3 hpa. Video is played back at 50× speed. Scale bar = 5 μm.

Supplementary Video 9

Time-lapse video microscopy of nuclear movement in the Hoechst (blue)-stained PbDHC3-GFP (green) zygotes at 5 hpa. Video is played back at 50× speed. Scale bar = 5 μm.

Contributions

RT and DSG conceived the project. RY, RT, MZ, DB performed the live imaging, genotypic and phenotype analysis; RY, MH, DJF, SV performed TEM and SBFSEM analysis; SB,AN, KGLR performed RNA seq analysis; RY, DB, ARB and ECT performed proteomic analysis; CP, AH performed phosphoproteomics, RY , MZ performed U-ExM observation; RY, MH, MZ, DJPF, CP, AN, ECT, KGLR, AAH, SV, DSG, RT performed formal analysis; RT, DSG, KGLR, ECT acquired funding; RT, DSG and AAH supervised the project. RY, DSG, RT wrote the manuscript draft; All authors reviewed and edited the manuscript.

Acknowledgements

RT is supported by an ERC advance grant funded by UKRI Frontier Science (EP/X024776/1), MRC UK (MR/K011782/1), and BBSRC (BB/L013827/1, BB/X014681/1). MZ, MH and DB were supported as research fellows and senior technician (EP/X024776/1). RY and DSG are supported by the BBSRC (BB/X014681/1). AAH is supported by the Francis Crick Institute (FC001097), which receives core funding from the Cancer Research UK (FC001097), the UK Medical Research Council (FC001097), and the Wellcome Trust (FC001097). KGLR is supported by the NIH/NIAID (R01 AI136511) and the University of California, Riverside NIFA-Hatch-225935. ECT was supported by a personal fellowship from the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), the Netherlands (grant no. VI. Veni.202.223). We thank the School of Life Sciences Imaging

(SLIM) for providing access to confocal and SIM microscopy and Dr Robert Markus for

technical assistance. We thank the Nanoscale and Microscale Research Centre (nmRC) for

providing access to instrumentation and Dr Michael W. Fey, Dr Julie Watts and Ms Nicola J.

Weston for technical assistance. For Open Access, the authors have applied a CC BY public

copyright licence to any Author Accepted Manuscript version arising from this submission.

We thank Cleidiane Zampronio at Warwick University for mass spectrometry methods and

Bio Support Unit, University of Nottingham for maintenance of mice used in this study.

1278

1279

References

1.   Bolcun-Filas E, Handel MA. Meiosis: the chromosomal foundation of reproduction. *Biol*
Reprod **99**, 112-126 (2018).

2.   Marston AL, Amon A. Meiosis: cell-cycle controls shuffle and deal. *Nat Rev Mol Cell Biol* **5**,
983-997 (2004).

3.   Ohkura H. Meiosis: an overview of key differences from mitosis. *Cold Spring Harb Perspect*
Biol **7**, (2015).

4.   Pesin JA, Orr-Weaver TL. Regulation of APC/C activators in mitosis and meiosis. *Annu Rev*
Cell Dev Biol **24**, 475-499 (2008).

5.   Pereira C, Smolka MB, Weiss RS, Brieno-Enriquez MA. ATR signaling in mammalian
meiosis: From upstream scaffolds to downstream signaling. *Environ Mol Mutagen* **61**, 752-
766 (2020).

6.   Loidl J. Conservation and Variability of Meiosis Across the Eukaryotes. *Annu Rev Genet* **50**,
293-316 (2016).

7.   Mirzaghaderi G, Horandl E. The evolution of meiotic sex and its alternatives. *Proc Biol Sci*
283, (2016).

8.   Guttery DS, Zeeshan M, Holder AA, Tromer EC, Tewari R. Meiosis in Plasmodium: how
does it work? *Trends Parasitol* **39**, 812-821 (2023).

9.   Guttery DS, Zeeshan M, Ferguson DJP, Holder AA, Tewari R. Division and Transmission:
Malaria Parasite Development in the Mosquito. *Annu Rev Microbiol* **76**, 113-134 (2022).

10. WHO. World malaria report 2024. *World Health Organisation* **ISBN 978-992-974-010444-**
010440, (2024).

1311

1312   11.   Zickler D, Kleckner N. Meiosis: Dances Between Homologs. *Annu Rev Genet* **57**, 1-63
1313   (2023).

1314
1315   12.   Sinden RE, Hartley RH, Winger L. The development of Plasmodium ookinetes in vitro: an
1316   ultrastructural study including a description of meiotic division. *Parasitology* **91 (Pt 2)**, 227-
1317   244 (1985).

1318
1319   13.   Sinden RE, Hartley RH. Identification of the meiotic division of malarial parasites. *J*
1320   *Protozool* **32**, 742-744 (1985).

1321
1322   14.   Sinden RE. Gametocytogenesis in Plasmodium spp., and observations on the meiotic division.
1323   *Ann Soc Belg Med Trop* **65 Suppl 2**, 21-33 (1985).

1324
1325   15.   Sinden RE, Canning EU, Spain B. Gametogenesis and fertilization in Plasmodium yoelii
1326   nigeriensis: a transmission electron microscope study. *Proc R Soc Lond B Biol Sci* **193**, 55-76
1327   (1976).

1328
1329   16.   Zeeshan M, *et al.* Plasmodium ARK2 and EB1 drive unconventional spindle dynamics,
1330   during chromosome segregation in sexual transmission stages. *Nat Commun* **14**, 5652 (2023).

1331
1332   17.   Zeeshan M, *et al.* Real-time dynamics of Plasmodium NDC80 reveals unusual modes of
1333   chromosome segregation during parasite proliferation. *J Cell Sci* **134**, (2020).

1334
1335   18.   Guttery DS, *et al.* Genome-wide functional analysis of Plasmodium protein phosphatases
1336   reveals key regulators of parasite development and differentiation. *Cell Host Microbe* **16**,
1337   128-140 (2014).

1338
1339   19.   Tewari R, *et al.* The systematic functional analysis of Plasmodium protein kinases identifies
1340   essential regulators of mosquito transmission. *Cell Host Microbe* **8**, 377-387 (2010).

1341
1342   20.   Ward P, Equinet L, Packer J, Doerig C. Protein kinases of the human malaria parasite
1343   Plasmodium falciparum: the kinome of a divergent eukaryote. *BMC Genomics* **5**, 79 (2004).

1344
1345   21.   Gao H, *et al.* ISP1-Anchored Polarization of GCbeta/CDC50A Complex Initiates Malaria
1346   Ookinete Gliding Motility. *Curr Biol* **28**, 2763-2776 e2766 (2018).

1347
1348   22.   Fry AM, Bayliss R, Roig J. Mitotic Regulation by NEK Kinase Networks. *Front Cell Dev*
1349   *Biol* **5**, 102 (2017).

1350
1351   23.   Fry AM, O'Regan L, Sabir SR, Bayliss R. Cell cycle regulation by the NEK family of protein
1352   kinases. *J Cell Sci* **125**, 4423-4433 (2012).

1353
1354   24.   Brieno-Enriquez MA, Moak SL, Holloway JK, Cohen PE. NIMA-related kinase 1 (NEK1)
1355   regulates meiosis I spindle assembly by altering the balance between alpha-Adducin and
1356   Myosin X. *PLoS One* **12**, e0185780 (2017).

1357

1358 25. Reininger L, *et al.* A NIMA-related protein kinase is essential for completion of the sexual
1359   cycle of malaria parasites. *J Biol Chem* **280**, 31957-31964 (2005).

1360
1361 26. Reininger L, Garcia M, Tomlins A, Muller S, Doerig C. The Plasmodium falciparum, Nima-
1362   related kinase Pfnek-4: a marker for asexual parasites committed to sexual differentiation.
1363   *Malar J* **11**, 250 (2012).

1364
1365 27. Reininger L, *et al.* An essential role for the Plasmodium Nek-2 Nima-related protein kinase in
1366   the sexual development of malaria parasites. *J Biol Chem* **284**, 20858-20868 (2009).

1367
1368 28. Zeeshan M, *et al.* Plasmodium NEK1 coordinates MTOC organisation and kinetochore
1369   attachment during rapid mitosis in male gamete formation. *PLoS Biol* **22**, e3002802 (2024).

1370
1371 29. M'Saad O, Bewersdorf J. Light microscopy of proteins in their ultrastructural context. *Nat
1372   Commun* **11**, 3850 (2020).

1373
1374 30. Cromer L, *et al.* Rapid meiotic prophase chromosome movements in Arabidopsis thaliana are
1375   linked to essential reorganization at the nuclear envelope. *Nat Commun* **15**, 5964 (2024).

1376
1377 31. Christophorou N, Rubin T, Bonnet I, Piolot T, Arnaud M, Huynh JR. Microtubule-driven
1378   nuclear rotations promote meiotic chromosome dynamics. *Nat Cell Biol* **17**, 1388-1400
1379   (2015).

1380
1381 32. Saito TT, Tougan T, Okuzaki D, Kasama T, Nojima H. Mcp6, a meiosis-specific coiled-coil
1382   protein of Schizosaccharomyces pombe, localizes to the spindle pole body and is required for
1383   horsetail movement and recombination. *J Cell Sci* **118**, 447-459 (2005).

1384
1385 33. Ding DQ, Chikashige Y, Haraguchi T, Hiraoka Y. Oscillatory nuclear movement in fission
1386   yeast meiotic prophase is driven by astral microtubules, as revealed by continuous
1387   observation of chromosomes and microtubules in living cells. *J Cell Sci* **111 (Pt 6)**, 701-712
1388   (1998).

1389
1390 34. Yamamoto A, West RR, McIntosh JR, Hiraoka Y. A cytoplasmic dynein heavy chain is
1391   required for oscillatory nuclear movement of meiotic prophase and efficient meiotic
1392   recombination in fission yeast. *J Cell Biol* **145**, 1233-1249 (1999).

1393
1394 35. Vogel SK, Pavin N, Maghelli N, Julicher F, Tolic-Norrelykke IM. Self-organization of dynein
1395   motors generates meiotic nuclear oscillations. *PLoS Biol* **7**, e1000087 (2009).

1396
1397 36. Yang S, *et al.* EB1 decoration of microtubule lattice facilitates spindle-kinetochore lateral
1398   attachment in Plasmodium male gametogenesis. *Nat Commun* **14**, 2864 (2023).

1399
1400 37. Chikashige Y, *et al.* Telomere-led premeiotic chromosome movement in fission yeast.
1401   *Science* **264**, 270-273 (1994).

1402

1403   38.   Labella S, Woglar A, Jantsch V, Zetka M. Polo kinases establish links between meiotic
1404   chromosomes and cytoskeletal forces essential for homolog pairing. *Dev Cell* **21**, 948-958
1405   (2011).

1406
1407   39.   Mlambo G, Coppens I, Kumar N. Aberrant sporogonic development of Dmc1 (a meiotic
1408   recombinase) deficient Plasmodium berghei parasites. *PLoS One* **7**, e52480 (2012).

1409
1410   40.   Tromer EC, Wemyss TA, Ludzia P, Waller RF, Akiyoshi B. Repurposing of synaptonemal
1411   complex proteins for kinetochores in Kinetoplastida. *Open Biol* **11**, 210049 (2021).

1412
1413   41.   Abramson J*, et al.* Accurate structure prediction of biomolecular interactions with AlphaFold
1414   3. *Nature* **630**, 493-500 (2024).

1415
1416   42.   van Kempen M*, et al.* Fast and accurate protein structure search with Foldseek. *Nat*
1417   *Biotechnol* **42**, 243-246 (2024).

1418
1419   43.   Sayers C*, et al.* Systematic screens for fertility genes essential for malaria parasite
1420   transmission reveal conserved aspects of sex in a divergent eukaryote. *Cell Syst* **15**, 1075-
1421   1091 e1076 (2024).

1422
1423   44.   Hirai M, Maeta A, Mori T, Mita T. Pb103 Regulates Zygote/Ookinete Development in
1424   Plasmodium berghei via Double Zinc Finger Domains. *Pathogens* **10**, (2021).

1425
1426   45.   Nishi T, Kaneko I, Iwanaga S, Yuda M. Identification of a novel AP2 transcription factor in
1427   zygotes with an essential role in Plasmodium ookinete development. *PLoS Pathog* **18**,
1428   e1010510 (2022).

1429
1430   46.   Yuda M, Kaneko I, Murata Y, Iwanaga S, Nishi T. Mechanisms of triggering malaria
1431   gametocytogenesis by AP2-G. *Parasitol Int* **84**, 102403 (2021).

1432
1433   47.   Modrzynska K*, et al.* A Knockout Screen of ApiAP2 Genes Reveals Networks of Interacting
1434   Transcriptional Regulators Controlling the Plasmodium Life Cycle. *Cell Host Microbe* **21**,
1435   11-22 (2017).

1436
1437   48.   Russell AJC*, et al.* Regulators of male and female sexual development are critical for the
1438   transmission of a malaria parasite. *Cell Host Microbe* **31**, 305-319 e310 (2023).

1439
1440   49.   Kaneko I, Nishi T, Iwanaga S, Yuda M. Differentiation of Plasmodium male gametocytes is
1441   initiated by the recruitment of a chromatin remodeler to a male-specific cis-element. *Proc*
1442   *Natl Acad Sci U S A* **120**, e2303432120 (2023).

1443
1444   50.   Hart KJ, Power BJ, Rios KT, Sebastian A, Lindner SE. The Plasmodium NOT1-G paralogue
1445   is an essential regulator of sexual stage maturation and parasite transmission. *PLoS Biol* **19**,
1446   e3001434 (2021).

1447

1448 51. Drewes G, Ebneth A, Preuss U, Mandelkow EM, Mandelkow E. MARK, a novel family of
1449 protein kinases that phosphorylate microtubule-associated proteins and trigger microtubule
1450 disruption. *Cell* **89**, 297-308 (1997).

1451
1452 52. Cao LS, Wang J, Chen Y, Deng H, Wang ZX, Wu JW. Structural basis for the regulation of
1453 maternal embryonic leucine zipper kinase. *PLoS One* **8**, e70031 (2013).

1454
1455 53. Liu B, Liu C, Li Z, Liu W, Cui H, Yuan J. A subpellicular microtubule dynein transport
1456 machinery regulates ookinete morphogenesis for mosquito transmission of Plasmodium
1457 yoelii. *Nat Commun* **15**, 8590 (2024).

1458
1459 54. Depoix D*, et al.* Vital role for Plasmodium berghei Kinesin8B in axoneme assembly during
1460 male gamete formation and mosquito transmission. *Cell Microbiol* **22**, e13121 (2020).

1461
1462 55. van de Kooij B*, et al.* Comprehensive substrate specificity profiling of the human Nek
1463 kinome reveals unexpected signaling outputs. *Elife* **8**, (2019).

1464
1465 56. Alexander J*, et al.* Spatial exclusivity combined with positive and negative selection of
1466 phosphorylation motifs is the basis for context-dependent mitotic signaling. *Sci Signal* **4**, ra42
1467 (2011).

1468
1469 57. Grallert A, Hagan IM. Schizosaccharomyces pombe NIMA-related kinase, Fin1, regulates
1470 spindle formation and an affinity of Polo for the SPB. *EMBO J* **21**, 3096-3107 (2002).

1471
1472 58. Grallert A, Krapp A, Bagley S, Simanis V, Hagan IM. Recruitment of NIMA kinase shows
1473 that maturation of the S. pombe spindle-pole body occurs over consecutive cell cycles and
1474 reveals a role for NIMA in modulating SIN activity. *Genes Dev* **18**, 1007-1021 (2004).

1475
1476 59. Yoshimura SH, Hirano T. HEAT repeats - versatile arrays of amphiphilic helices working in
1477 crowded environments? *J Cell Sci* **129**, 3963-3970 (2016).

1478
1479 60. Ji DD, Sultan AA, Chakrabarti D, Horrocks P, Doerig C, Arnot DE. An RCC1-type guanidine
1480 exchange factor for the Ran G protein is found in the Plasmodium falciparum nucleus. *Mol
1481 Biochem Parasitol* **95**, 165-170 (1998).

1482
1483 61. Zalli D, Bayliss R, Fry AM. The Nek8 protein kinase, mutated in the human cystic kidney
1484 disease nephronophthisis, is both activated and degraded during ciliogenesis. *Hum Mol Genet*
1485 **21**, 1155-1171 (2012).

1486
1487 62. Burette M*, et al.* Two anchoring proteins control daughter apical complex assembly in
1488 Toxoplasma gondii. *bioRxiv*, (2026).

1489
1490 63. Koch LB, Spanos C, Kelly V, Ly T, Marston AL. Rewiring of the phosphoproteome executes
1491 two meiotic divisions in budding yeast. *EMBO J* **43**, 1351-1383 (2024).

1492

1493 64.   Sivakova B*, et al.* Quantitative proteomics and phosphoproteomics profiling of meiotic
1494   divisions in the fission yeast Schizosaccharomyces pombe. *Sci Rep* **14**, 23105 (2024).

1495
1496 65.   Swartz SZ, Nguyen HT, McEwan BC, Adamo ME, Cheeseman IM, Kettenbach AN.
1497   Selective dephosphorylation by PP2A-B55 directs the meiosis I-meiosis II transition in
1498   oocytes. *Elife* **10**, (2021).

1499
1500 66.   Okumura E, Morita A, Wakai M, Mochida S, Hara M, Kishimoto T. Cyclin B-Cdk1 inhibits
1501   protein phosphatase PP2A-B55 via a Greatwall kinase-independent mechanism. *J Cell Biol*
1502   **204**, 881-889 (2014).

1503
1504 67.   Kaneko I, Iwanaga S, Kato T, Kobayashi I, Yuda M. Genome-Wide Identification of the
1505   Target Genes of AP2-O, a Plasmodium AP2-Family Transcription Factor. *PLoS Pathog* **11**,
1506   e1004905 (2015).

1507
1508 68.   Murata Y, Nishi T, Kaneko I, Iwanaga S, Yuda M. Coordinated regulation of gene expression
1509   in Plasmodium female gametocytes by two transcription factors. *Elife* **12**, (2024).

1510
1511 69.   Anderson-White B, Beck JR, Chen CT, Meissner M, Bradley PJ, Gubbels MJ. Cytoskeleton
1512   assembly in Toxoplasma gondii cell division. *Int Rev Cell Mol Biol* **298**, 1-31 (2012).

1513
1514 70.   Tosetti N*, et al.* Essential function of the alveolin network in the subpellicular microtubules
1515   and conoid assembly in Toxoplasma gondii. *Elife* **9**, (2020).

1516
1517 71.   Harding CR, Frischknecht F. The Riveting Cellular Structures of Apicomplexan Parasites.
1518   *Trends Parasitol* **36**, 979-991 (2020).

1519
1520 72.   Burrell A*, et al.* Cellular electron tomography of the apical complex in the apicomplexan
1521   parasite Eimeria tenella shows a highly organised gateway for regulated secretion. *PLoS*
1522   *Pathog* **18**, e1010666 (2022).

1523
1524 73.   Chen CT, Gubbels MJ. TgCep250 is dynamically processed through the division cycle and is
1525   essential for structural integrity of the Toxoplasma centrosome. *Mol Biol Cell* **30**, 1160-1169
1526   (2019).

1527
1528 74.   Hu X, O'Shaughnessy WJ, Beraki TG, Reese ML. Loss of the Conserved Alveolate Kinase
1529   MAPK2 Decouples Toxoplasma Cell Growth from Cell Division. *mBio* **11**, (2020).

1530
1531 75.   Guttery DS*, et al.* A unique protein phosphatase with kelch-like domains (PPKL) in
1532   Plasmodium modulates ookinete differentiation, motility and invasion. *PLoS Pathog* **8**,
1533   e1002948 (2012).

1534
1535 76.   Mytlis A, Levy K, Elkouby YM. The many faces of the bouquet centrosome MTOC in
1536   meiosis and germ cell development. *Curr Opin Cell Biol* **81**, 102158 (2023).

1537

1538 77. Zeeshan M*, et al.* A novel SUN1-ALLAN complex coordinates segregation of the bipartite
1539 MTOC across the nuclear envelope during rapid closed mitosis in Plasmodium berghei. *Elife*
1540 **14**, (2025).

1541
1542 78. Janse CJ*, et al.* High efficiency transfection of Plasmodium berghei facilitates novel selection
1543 procedures. *Mol Biochem Parasitol* **145**, 60-70 (2006).

1544
1545 79. Winger LA, Tirawanchai N, Nicholas J, Carter HE, Smith JE, Sinden RE. Ookinete antigens
1546 of Plasmodium berghei. Appearance on the zygote surface of an Mr 21 kD determinant
1547 identified by transmission-blocking monoclonal antibodies. *Parasite Immunol* **10**, 193-207
1548 (1988).

1549
1550 80. Ferguson DJ*, et al.* Maternal inheritance and stage-specific variation of the apicoplast in
1551 Toxoplasma gondii during development in the intermediate and definitive host. *Eukaryot Cell*
1552 **4**, 814-826 (2005).

1553
1554 81. Hair M, Moreira-Leite F, Ferguson DJP, Zeeshan M, Tewari R, Vaughan S. Atypical flagella
1555 assembly and haploid genome coiling during male gamete formation in Plasmodium. *Nat*
1556 *Commun* **14**, 8263 (2023).

1557
1558 82. Yanase R*, et al.* Divergent Plasmodium kinases drive MTOC, kinetochore and axoneme
1559 organisation in male gametogenesis. *Life Sci Alliance* **8**, (2025).

1560
1561 83. Liffner B, Silva T, Vega-Rodriguez J, Absalon S. Mosquito Tissue Ultrastructure-Expansion
1562 Microscopy (MoTissU-ExM) enables ultrastructural and anatomical analysis of malaria
1563 parasites and their mosquito. *BMC Mehods* **1**, (2024).

1564
1565 84. Perez-Riverol Y*, et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res* **53**,
1566 D543-D553 (2025).

1567
1568

Figure 1

Figure 2

Figure 3

Figure 4

a

4 h activated WT zygotes

1 | 2 — APC, APR, A-Mt

3 | 4 (N) | 5 (N, NM)

4 h activated *Pbnek4-ko* zygotes

6 (N) | 7 — APR, A-Mt, APC

8 (N) | 9 (N) | 10 (N, NM)

b

| 4 h activated WT zygote | 4 h activated *Pbnek4-ko* zygote |

NHS-ester/Hochest

NHS-ester — Tracing

Hoechst

NHS-ester/Hochest

c

Whole cell | Nucleus

WT — APC, N

WT — APC, N

Pbnek4-ko — APC, N

d

Developing APC
p = 0.0002
% of cells
WT   *Pbnek4-ko*

Condensed chromosomes
p = 0.0004
% of cells
WT   *Pbnek4-ko*

e

WT
Nucleus, APR, APC, MTOCs, Condensed chromosomes

Pbnek4-ko
Nucleus, APR, APC, MTOC, Uncondensed chromosomes

Figure 5

Figure 6

Meiosis in *S. pombe*

Cell membrane Nucleus ● SPB/MTOC ● Kinetochore — Microtubule

Chromosomes — Sister chromatids — LinEs formation Crossover

| Meiotic entry (2N) | Meiotic S phase (2-4N) | Prophase (4N) | Metaphase I (4N) | Anaphase I (4N) | Metaphase II (2×2N) | Anaphase II (4×1N) |

Horsetail nuclear movement

Meiosis in *Plasmodium*

NEK4 KO

SCs formation Crossover?

| 0-1 hpa Zygote (2N) | 2-4 hpa Meiotic S phase (2-4N) | 4-6 hpa Prophase (4N) | 6-8 hpa Metaphase I (?) (4N) | 8-10 hpa Anaphase I (?) (4N) | 10-12 hpa Meiosis II start (?) (4N) | 12-24 hpa Meiosis II (?) (4N) |

Nuclear movement

Meiosis
Rec8
DMC1 HOP1

DNA Chromosomes
RCC (0905500)
POLD3

Uncharacterised
Putative HEAT-repeat protein (0806700)

NEK4

MTOC Microtubules
MARK/MELK
DHC3 KIN4
MORN (1030400)

Sexual development
Pb103 AP2-Z
FD1/2 AP2-O
NOT1-G gSNF2

NEK4

APC microtubule formation

Nuclear/chromosome movement?
Dynein? Microtubules Dynein? SUN?
MTOC
Nucleus Chromosomes Nuclear membrane

MTOC separation

Perinuclear microtubule formation

DNA replication Chromosome condensation

Meiosis initiation Ookinete development