TXNDC5 Governs Extracellular Matrix Homeostasis in Pulmonary Hypertension

Zhen Chen[1*]; Fanhao Kong[1*]; Lixin Huang[1*]; Wenze Wu[1*]; Ziping Wang[1]; Hanbin Chen[1]; Junting Zhang[1]; Lin Deng[6]; Chengrui Cao[1]; Xiaoyan Zhang[1]; Zixin Liu[2]; Kai-Chien Yang[3#]; Wei-Ting Chang[4, 5#]; Jin-Song Bian[1#]; Xiaowei Nie[2#];

Department of Pharmacology, Joint Laboratory of Guangdong-Hong Kong Universities for Vascular Homeostasis and Diseases, SUSTech Homeostatic Medicine Institute, School of Medicine, Southern University of Science and Technology, Shenzhen 518055, China.

Department of Human Cell Biology and Genetics, SUSTech Homeostatic Medicine Institute, School of Medicine, Southern University of Science and Technology, Shenzhen 518055, China.

Department and Graduate Institute of Pharmacology, National Taiwan University College of Medicine, Taipei, Taiwan

School of Medicine, College of Medicine, National Sun Yat-sen University, Kaohsiung, Taiwan

Division of Cardiology, Department of Internal Medicine, Chi Mei Medical Center, Tainan

Department of Cardiology, The Eighth Affiliated Hospital, Sun Yat-sen University, Shenzhen, China

[*]These authors contributed equally to this work.

[#]Correspondence to:

Xiaowei Nie, PhD, E-mail niexw@sustech.edu.cn.

Jin-Song Bian, PhD, E-mail bianjs@sustech.edu.cn.

Wei-Ting Chang, PhD, E-mail cmcvecho2@gmail.com.

Kai-Chien Yang, PhD, E-mail allenmy0920@gmail.com.

ABSTRACT

BACKGROUND: Pulmonary hypertension (PH) is characterized by vascular remodeling without effective treatments. Thioredoxin domain containing 5 (TXNDC5), a member of the protein disulfide isomerases (PDI) family, regulates protein folding and vascular homeostasis, yet its role in PH remains unknown.

METHODS: Label-free proteomics was used to profile PDI expression in hypoxic PH patients. TXNDC5 expression and localization were examined by multiplex immunofluorescence and Western blot. Gain- and loss-of-function approaches, including adeno-associated virus-mediated lung endothelial overexpression, global knockout, and endothelial ablation of TXNDC5, were used in Sugen5416/hypoxia (SuHx)-induced rodent PH models. *In vitro* endothelial cell assays, RNA sequencing, protein-protein interaction analysis, and pharmacological intervention were conducted to elucidate underlying mechanisms and therapeutic potential.

RESULTS: TXNDC5 was significantly upregulated in the lungs of both PH patients and experimental PH models, predominantly in endothelial cells of remodeled distal pulmonary arteries. Endothelial TXNDC5 overexpression exacerbated pulmonary vascular remodeling, elevated right ventricular systolic pressure, and promoted right ventricular hypertrophy, whereas global or endothelial-specific TXNDC5 deficiency conferred protection against SuHx-induced PH. RNA sequencing and protein-protein interaction analysis revealed that TXNDC5 regulated extracellular matrix homeostasis through biglycan (BGN), which was required for TXNDC5-mediated inhibition of HIF-$1\alpha/2\alpha$ ubiquitination, leading to pathological HIF accumulation. Pharmacological inhibition of TXNDC5 with E64FC26 and endothelial-targeted TXNDC5 gene therapy significantly attenuated PH severity in SuHx rats.

CONCLUSIONS: Our study reveals that TXNDC5 is a main modulator to regulate ECM homeostasis and may serve as a promising target for the treatment of PH.

Key Words: pulmonary hypertension**,** extracellular matrix, TXNDC5, BGN, HIF signaling

INTRODUCTION

Pulmonary hypertension (PH) is a serious cardiovascular disease characterized by elevated pulmonary artery pressure and right heart failure, with high morbidity and mortality[1]. Hypoxia triggers PH, with Group III being common clinically[2]. Dysfunction of pulmonary artery endothelial cells (PAECs) under hypoxia initiates vasoconstriction[3], while pulmonary artery smooth muscle cells (PASMCs) become hyperproliferative and apoptosis-resistant phenotype, leading to vascular remodeling[4]. However, there remains no curative clinical treatment option for reversing remodeled vessels against the disease[5,6]. It highlights the need for further exploration of PH pathogenesis and new therapeutic targets.

Protein disulfide isomerase (PDI) is an endoplasmic reticulum enzyme family that mediates disulfide bond formation, protein folding, and regulates endothelial function to maintain vascular homeostasis[7,8]. PDIA1 controls HIF-1α expression and redox state via lysosomal degradation and enhances vascular endothelial growth factor receptor 2 signaling, promoting endothelial migration, proliferation, and angiogenesis[9,10]. TXNDC5, mainly expressed in endothelial cells (ECs) and also called endothelial PDI, is linked to fibrosis[11], diabetes[12], rheumatoid arthritis[13], and cancers[14]. Inhibition of TXNDC5 restores endothelial dysfunction to mitigate vascular calcification and atherosclerosis[15,16]. However, its role in pulmonary vascular remodeling in PH remains unclear.

Extracellular matrix (ECM) is a complex network of proteins and polysaccharides secreted by cells, providing structural and communicative support to adjacent cells[17]. Remodeling of pulmonary arterial ECM occurs in the early stage of PH pathogenesis, before increased wall thickness and pulmonary artery pressure. ECM remodeling is the cause rather than a consequence of pulmonary vascular remodeling in PH[18]. Recently, we revealed that the ECM proteins in PAECs accelerated pulmonary vascular remodeling through interaction with HIF-1α in a positive feedback loop manner, providing evidence for early intervention in ECM remodeling to prevent PH

progression[19,20].

In this study, using 4D label-free proteomics, we identified TXNDC5 as a

markedly upregulated PDI protein in lungs of patients with hypoxic PH. Global or

endothelial-specific deficiency of TXNDC5 alleviated Sugen5416/hypoxia (SuHx)-

induced PH, whereas TXNDC5 overexpression aggravated PH. TXNDC5 silencing

preserved ECM homeostasis, reduced PA remodeling, and restored endothelial function.

In addition, TXNDC5 directly interacted with and assisted its the folding of biglycan

(BGN), which was required to regulate the HIF-1α/2α ubiquitination. Pharmacological

inhibition or pulmonary endothelium-targeted silencing of TXNDC5 effectively

mitigated PH in rats. The present study therefore demonstrated for the first time that

TXNDC5 may be a key protein to control ECM homeostasis and a promising

therapeutic target for PH.

METHODS

Detailed expanded methods and materials are available in **Supplemental Material 1**.

Data Availability Statement

Source data and original blots or gel images for this study are available from the

corresponding author upon reasonable request.

RESULTS

TXNDC5 Expression is Increased in PH Patients and Rodent Models

We performed 4D label-free proteomics on lung samples from 5 patients with

hypoxic PH and 5 non-PH donors (**Figure S1A**). Among the human PDI family proteins,

TXNDC5 was the most markedly dysregulated member in PH patients compared with

donors (fold change=3.45, *P*=0.0048) (**Figure 1A**). Increased TXNDC5 expression in

PH lungs was confirmed at both protein and mRNA levels (**Figure 1B-C**), and similarly

observed in multiple rat and mouse PH models (**Figure 1D** and **Figure S1B**). In distal

pulmonary arteries (PAs), multiplex immunofluorescence (mIF) showed pronounced co-localization of TXNDC5 with the endothelial marker CD31, α-SMA (a marker for smooth muscle cells), and COL1A1 (a marker for fibroblasts) (**Figure 1E-G** and **Figure S1C**). Furthermore, TXNDC5 was elevated in the intima of remodeled PAs from interstitial lung disease (ILD)-PH and chronic obstructive pulmonary disease (COPD)-PH patients (**Figure 1H**). Similar results were observed in the remodeled PAs of SuHx- or monocrotaline (MCT)-induced PH rat models (**Figure 1I-K** and **Figure S1D**). Western blot analysis of primary lung ECs corroborated these findings (**Figure S1E-F**). Notably, intimal TXNDC5 levels positively correlated with right ventricular systolic pressure (RVSP) and Fulton index (right ventricle weight to left ventricle plus interventricular septum weight, ie, RV/(LV+S)), and negatively with tricuspid annular plane systolic excursion (TAPSE) (**Figure 1L-N**). Taken together, these findings suggest that the TXNDC5 may be a critical dysregulated factor in ECs within the pulmonary vasculature that contributes to PH pathogenesis.

TXNDC5 Mediates the Development of SuHx-Induced PH

To investigate the role of TXNDC5 *in vivo* in the development of experimental PH, TXNDC5 knockout (TXNDC5$^{-/-}$) mice were challenged with or without SuHx (**Figure 2A**). TXNDC5 deletion alone had no significant effects but markedly reduced RVSP (**Figure 2B**) and RV/(LV+S) (**Figure S2A**) induced by SuHx. Echocardiography indicated improved right ventricular (RV) function, with increased PA acceleration time/ PA ejection time ratio (PAT/PET), TAPSE, and right ventricular fractional area change (RVFAC) (**Figure S2B-D**), and reduced right ventricular free wall thickness (RVFWT) (**Figure S2E**). In addition, mIF and Victoria blue/Sirius red (VB/SR) showed reduced distal PA muscularization, medial wall thickness, and collagen deposition under SuHx (**Figure 2C-F**). Consistent with the PA changes, wheat germ agglutinin (WGA) staining proved that the RV cardiomyocyte hypertrophy was markedly inhibited in SuHx-treated TXNDC5$^{-/-}$ mice when compared with their control littermates (**Figure 2G**).

With a gain-of-function approach, an adeno-associated virus serotype engineered

to specifically target lung endothelial cells (AAV-TXN) was delivered via tail vein to overexpress TXNDC5 specifically in lung ECs (**Figure 2H**). *Ex vivo* imaging confirmed lung targeting (**Figure 2I**), and co-immunofluorescence and Western blot verified EC-specific overexpression (**Figure S2F-G**). Hemodynamic measurement revealed that mice administered with AAV-TXN increased the RVSP compared with those administered with control AAV (AAV-NC) even under normoxic conditions. In response to the SuHx challenge, AAV-TXN mice further increased in both RVSP and RV/(LV+S) (**Figure 2J** and **Figure S2H**). Echocardiography analysis showed reduced PAT/PET, TAPSE RVFAC, and increased RVFWT under SuHx (**Figure S2I-L**). In addition, mIF and VB/SR staining confirmed exacerbated distal PA remodeling with increased muscularization, medial thickness, and collagen deposition (**Figure 2K-N**). Conversely, endothelial TXNDC5 overexpression was found to exacerbate SuHx-induced PH.

Endothelial Cell-Specific Deletion of TXNDC5 Ameliorates SuHx-Induced PH

To examine the role of endothelial TXNDC5 in PH, endothelial cell-specific TXNDC5 knockout mice (TXNDC5$^{\text{ECKO}}$) were generated, validated (**Figure S2M**), and challenged by SuHx (**Figure 3A**). Deletion of endothelial TXNDC5 significantly attenuated SuHx-induced increases in RVSP (**Figure 3B**) and RV/(LV+S) (**Figure 3C**) and improved RV function (**Figure S2N-Q**). Under SuHx, TXNDC5$^{\text{ECKO}}$ mice exhibited reduced distal PA muscularization, medial wall thickness, and collagen deposition (**Figure 3D-G**). Lung angiography confirmed better intrapulmonary contrast filling in TXNDC5$^{\text{ECKO}}$ mice compared with TXNDC5$^{\text{f/f}}$ controls, indicating preserved pulmonary vasculature (**Figure 3H** and **Figure S2R**). In addition, SuHx-induced RV hypertrophy was also markedly attenuated, with decreased RV wall thickness and cardiomyocyte cross-sectional area (**Figure 3I-J**). Together, these findings underscore the crucial role of endothelial TXNDC5 in the development of PH and highlight its potential as a sex-independent therapeutic target.

TXNDC5 Promotes Hypoxia-Induced Dysfunction in Pulmonary Arterial Cells

Exposure to hypoxia significantly increased the TXNDC5 protein and mRNA

levels in hPAECs, predominantly in the cytoplasm (**Figure S3A-C**). Silencing

TXNDC5 with small-interfering RNA oligonucleotides (siTXNDC5) suppressed

proliferation and migration of hPAECs under both normoxic and hypoxic conditions

(**Figure S3D-E**), and prevented G1-to-S phase transition (**Figure S3F**). Conversely,

TXNDC5 overexpression via lentivirus (LV-TXNDC5) disrupted cytoskeletal

architecture, inducing prominent stress fibers and diffuse paxillin distribution (**Figure**

S3G-H). Taken together, TXNDC5 regulates many important functional abnormalities

associated with PH pathogenesis, with disruption of cytoskeletal architecture in

hPAECs.

The hyper-proliferation and resistance to apoptosis of PASMCs directly accelerate

the remodeling of PAs in PH[21]. Thus, we investigated whether endothelial TXNDC5

affects the function of hPASMCs via paracrine signaling. For this purpose, conditioned

medium from siTXNDC5-treated hPAECs (siTXN-CM) was used to treat hPASMCs.

Western blot showed that siTXN-CM significantly downregulated cyclin A and

proliferating cell nuclear antigen (PCNA) expression under both normoxic and hypoxic

conditions (**Figure S3I**). Consistently, EdU incorporation and Cell Counting Kit-8

(CCK-8) assays further confirmed that siTXN-CM suppressed the proliferation and

viability of hPASMCs (**Figure S4A-B**). In addition, siTXN-CM enhanced hPASMCs

apoptosis, as indicated by increased cleaved caspase-3 expression (**Figure S3J**).

Collectively, our data demonstrate that hPAECs TXNDC5 plays a critical role in

promoting hPASMCs proliferation, survival, and resistance to apoptosis.

TXNDC5 Regulates ECM Expression and Homeostasis

To explore the underlying downstream molecular mechanisms of TXNDC5, we

performed a bulk RNA sequencing (RNA-seq) analysis on hPAECs transfected with

siRNA targeting TXNDC5 under hypoxia exposure. Gene ontology analysis indicated

that the silencing of TXNDC5 resulted in altered expression of genes mainly enriched

in "collagen-containing ECM", "ECM", "ECM organization", "ECM structural

constituent", "response to hypoxia", and "cellular response to hypoxia" (**Figure 4A**).

We selected the differentially expressed ECM genes and then constructed a protein-

protein interaction (PPI) network (**Figure 4B**). The network revealed that TXNDC5-

affected ECM proteins were clustered into different functional subgroups, including

"proteoglycans and glycoproteins", "collagens", "matrix modulators and adhesion",

"metallopeptidase activity", "basement membrane", "growth factors and signaling

regulators", and "angiogenesis". Immunofluorescence staining confirmed that periostin

(POSTN), biglycan (BGN), lysyl oxidase (LOX), and transglutaminase 2 (TGM2) were

excessively deposited in the PAs from TXNDC5$^{f/f}$ mice but not in TXNDC5^{ECKO} mice

under SuHx conditions (**Figure 4C-F** and **Figure S4C**). Western blot analysis also

confirmed that TXNDC5 silencing reversed the abnormal expression of ECM proteins

(COL1A1, TGM2, FN1, POSTN, LOX) induced by hypoxia treatment in hPAECs

(**Figure 4G**).

To further investigate whether TXNDC5 regulates ECM secretion, we collected

the supernatants from hPAECs. Dot blot analysis revealed that the silence of TXNDC5

markedly reduced the secretion of BGN, POSTN, and LOX **(Figure S4D-G)**. These

results provide a mechanistic explanation for why the conditional medium from

TXNDC5-silenced hPAECs suppresses proliferation and viability while promoting

apoptosis of hPASMCs. Collectively, these data highlight the key role TXNDC5 played

in regulating ECM homeostasis.

BGN Is a Promising Biomarker and Therapeutic Target for Hypoxic Pulmonary

Hypertension

We next explored TXNDC5-targeted ECM proteins critical for EC function. We

performed a computational analysis of the PPI network between the ECM proteins

dataset and the ECs function-related proteins dataset. By integrating these datasets, we

first constructed a comprehensive PPI network, which was then refined into a core

network using degree-based filtering (**Figure 5A**). Through the intersection of ECM

datasets, core ECM in the above PPI network, TXNDC5 silencing in hPAECs, and

hypoxia-treated hPAECs (GSE160255), we finally identified biglycan (BGN) as a key

candidate (**Figure S4H**).

BGN protein levels were increased in lung biopsies from patients with hypoxic

PH or donors (**Figure S4I**). A GEO dataset (GSE254617) confirmed elevated BGN

levels in idiopathic pulmonary arterial hypertension (IPAH) lungs (**Figure 5B**).

Similarly, BGN was upregulated in rodent PH models (**Figure S4J-K**). We also

investigated whether BGN can reflect the severity of PH by analyzing BGN levels in

serum. Surprisingly, a markedly increased BGN level was found in the serum of 27

patients with hypoxic PH compared with the serum of 27 donors (**Figure 5C**). Serum

BGN levels exhibited a significant correlation with systolic pulmonary artery pressure

(sPAP) and mean pulmonary artery pressure (mPAP) (**Figure 5D-E**), indicating its

potential as a biomarker. As a secreted protein, we further investigated the biological

function of circulating BGN *in vitro*. Functionally, recombinant human BGN (rhBGN)

enhanced hPASMC proliferation (**Figure S4L**) and viability (**Figure S4M**) compared

with the negative control (bovine serum albumin, BSA). In parallel, rhBGN induced an

apoptosis-resistant phenotype in hPASMCs, as evidenced by reduced cleaved caspase-

3 expression (**Figure S4N**) and decreased apoptotic cell populations detected by flow

cytometry (**Figure S4O**).

To evaluate the specific role of BGN in PH generation, we generated endothelial-

specific BGN knockout mice (BGN^{ECKO}) (**Figure S5A**). Since BGN is located on the

X chromosome, male mice carry a single floxed allele (BGN$^{f/0}$) and female mice carry

two floxed alleles (BGN$^{f/f}$), and these mice are collectively referred to as BGN^{flox} mice.

Notably, SuHx-induced increases in RVSP and RV/(LV+S) were significantly

attenuated in BGN^{ECKO} mice compared with BGN^{flox} controls (**Figure 5F-G**).

Consistently, BGN^{ECKO} mice showed reduced PAs remodeling (**Figure 5H-K**). Lung

angiogram also revealed the better-preserved lung vasculature in BGN^{ECKO} mice

(**Figure 5L**). Histological assessment of the RV showed a significant reduction in

hypertrophy in BGN^{ECKO} mice treated with SuHx (**Figure 5M**). Echocardiography

further demonstrated that SuHx-induced decreases in PAT/PET, TAPSE, RVFAC, and

increases in RVFWT were less pronounced in BGN^{ECKO} mice than in BGN^{flox} mice

(**Figure S5B-E**). Together, our data suggest that endothelial BGN plays an important

role in PH development.

TXNDC5 Assists BGN Protein Folding to Drive Pulmonary Hypertension

To elucidate how TXNDC5 regulates BGN, we first performed Western blots to validate the positive regulation of TXNDC5 on BGN both *in vitro* and *in vivo*. The silencing of TXNDC5 reversed the overexpression of BGN induced by hypoxia stimulation in hPAECs (**Figure S5F**). Reduced BGN expression was also observed in the lungs of TXNDC5-deficient mice (**Figure S5G**), which was consistent with the findings in primary lung ECs isolated from TXNDC5[ECKO] mice (**Figure S5H**). In contrast, BGN expression was elevated in hPAECs following lentiviral-mediated TXNDC5 overexpression (**Figure S5I**) and in primary lung ECs isolated from AAV-TXN-treated mice (**Figure S5J**).

To further assess the direct interaction between TXNDC5 and BGN, we investigated the subcellular location at which TXNDC5 binds to BGN. Immunostaining assays showed that TXNDC5 colocalized with BGN (**Figure 6A**). We also performed co-IP analysis and found that TXNDC5 bound to BGN, and the binding was enhanced upon hypoxia exposure (**Figure 6B** and **Figure S5K**). To identify the protein domain of TXNDC5 that was involved in the interaction with BGN, we performed molecular docking analysis and revealed that multiple sites of TXNDC5 (Cys89, Ser125, Gly351, and His352) might be involved in the interaction with BGN (**Figure 6C**). Thus, we constructed four TXNDC5 mutants (C89A, S125A, G351A, and H352A) to identify the residue that plays a key role in this process. Co-IP analysis showed that C89A and G351A could partially abolish TXNDC5 binding with BGN (**Figure 6D**), suggesting that these amino acid residues might be essential for the binding of TXNDC5 and BGN.

TXNDC5, an endoplasmic reticulum resident protein with three thioredoxin domains containing CGHC motifs, catalyzes disulfide bond formation[14]. Thus, we hypothesized that TXNDC5 facilitates the folding of BGN, enabling its maturation, signaling, and secretion. To test this, we generated a PDI-inactive TXNDC5 mutant (TXNDC5 Mut) with cysteine-to-alanine substitutions in all three domains (**Figure S5L**). Fluorescence resonance energy transfer (FRET)-based protein folding assays (**Figure S5M**) showed that LV-TXNDC5 increased BGN folding efficiency, whereas

TXNDC5 Mut did not (**Figure 6E**). Moreover, cycloheximide (CHX) chase assays further demonstrated that overexpression of TXNDC5 markedly prolonged the half-life of BGN (**Figure S5N**), whereas TXNDC5 silencing accelerated BGN degradation (**Figure S5O**). These results indicate that TXNDC5 relies on its PDI activity to promote BGN folding, stability, and extracellular secretion from hPAECs.

To assess whether BGN mediates the endothelial TXNDC5 *in vivo*, we overexpressed TXNDC5 in lung ECs of BGN^ECKO and BGN^flox mice (**Figure 6F**). In BGN^flox mice under SuHx conditions, AAV-TXN exacerbated distal PA muscularization, medial wall thickness, and collagen deposition in BGN^flox mice, whereas these effects were largely abolished in BGN^ECKO mice (**Figure 6G-J**). Consistently, AAV-TXN markedly increased RVSP (**Figure 6K**) and aggravated RV hypertrophy in BGN^flox mice, whereas both changes were significantly attenuated in BGN^ECKO mice (**Figure 6L**). Echocardiographic assessment revealed that AAV-TXN led to pronounced deterioration of RV performance, while these adverse functional changes were largely blunted in BGN^ECKO mice (**Figure S5P-S**). Together, these results indicate that endothelial BGN is required for TXNDC5-mediated pulmonary vascular remodeling and PH development, confirming BGN as a key downstream effector of TXNDC5.

A Positive Feedback Loop Exists Between TXNDC5 and HIF Signaling

HIF α subunits are the dominant effector activated by hypoxia, suggesting that HIF signaling may regulate TXNDC5 transcription. qPCR analysis showed that silencing of HIF-2α, but not HIF-1α, markedly downregulated TXNDC5 mRNA levels under both normoxia and hypoxia conditions (**Figure S6A-B**). Western blot analysis also demonstrated that the protein level of TXNDC5 was significantly decreased when HIF-2α was knocked down (**Figure S6C**). Next, two putative HIF-2α binding sites (BS1 and BS2) were identified in the TXNDC5 promoter (**Figure S6D**). Chromatin immunoprecipitation (ChIP) assays confirmed HIF-2α binding to both sites (**Figure S6E**), and dual-luciferase reporter assays showed that mutation of either site reduced promoter activity, with mutation of both causing a further decrease (**Figure 7A**). These results indicate that BS1 and BS2 synergistically mediate HIF-2α-driven TXNDC5

transcription. Moreover, we found that silencing TXNDC5 markedly reduced HIF-

1α/2α protein levels under hypoxia (**Figure S6F**), consistent with observations in the

lungs of TXNDC5-deficient mice (**Figure 7B**). Nuclear localization of HIF-1α/2α was

also decreased upon TXNDC5 knockdown, as confirmed by Western blot and

immunofluorescence (**Figure S6G-H**). These results indicate a positive feedback loop

between TXNDC5 and HIF signaling during hypoxia.

To assess whether HIF signaling mediates the pathogenic effects of endothelial

TXNDC5 *in vivo*, mice overexpressing TXNDC5 via AAV-TXN were treated with

selective HIF-1α (PX-478) or HIF-2α (PT-2385) inhibitors under normoxia or SuHx

(**Figure S6I**). As expected, AAV-TXN markedly exacerbated PH phenotypes, including

distal PA remodeling, increased RVSP, RV hypertrophy, and impaired RV function.

Importantly, HIF inhibitors significantly attenuated these TXNDC5-induced vascular

and cardiac abnormalities, as shown by histology (**Figure 7C-F**), hemodynamics

(**Figure S6J**), evaluation of RV hypertrophy (**Figure S6K**), and echocardiographic

analysis (**Figure S6L-O**). Collectively, these results indicate that TXNDC5 promotes

PH via a HIF-dependent positive feedback loop.

TXNDC5 Requires BGN to Inhibit the Ubiquitination of HIF-1α/2α

Interestingly, we found in the present study that the HIF-1α/2α protein levels were

significantly decreased by the silencing of BGN upon hypoxia exposure in hPAECs

(**Figure S7A**). Since TXNDC5 directly interacted with BGN, we hypothesized that the

regulation of HIF-1α/2α by TXNDC5 required the presence of BGN. Western blots

revealed that LV-TXNDC5 increased the expression of HIF-1α/2α both under normoxic

and hypoxic conditions in hPAECs (**Figure S7B**). More importantly, knockdown of

BGN abolished the upregulation of HIF-1α/2α induced by LV-TXNDC5 (**Figure 7G**).

These data suggest that TXNDC5 requires BGN to regulate the HIF signaling pathway.

Under normoxia, HIF-α is rapidly ubiquitinated by E3 ligase von Hippel-Lindau

(VHL) and degraded[22]. Notably, TXNDC5 overexpression increased the HIF-1α/2α

levels even under normoxic conditions, which was also observed in primary lung ECs

from AAV-TXN-treated mice (**Figure S7C**). CHX chase assays showed that TXNDC5

prolonged the half-life of HIF-1α/2α in hPAECs (**Figure 7H** and **Figure S7D**). Mechanistically, TXNDC5 silencing increased VHL expression, while concurrent knockdown of VHL rescued the reduction of HIF-1α/2α caused by TXNDC5 depletion under hypoxia (**Figure 7I**). *In vivo* ubiquitination assays further revealed that TXNDC5 overexpression suppressed HIF-1α/2α ubiquitination, whereas BGN silencing enhanced ubiquitination by upregulating VHL. Importantly, BGN knockdown abolished the TXNDC5-induced inhibition of HIF-1α/2α ubiquitination and VHL downregulation (**Figure 7J** and **Figure S7E**). Collectively, these results demonstrate that TXNDC5 stabilizes HIF-1α/2α by inhibiting VHL-mediated ubiquitination in a BGN-dependent manner.

E64FC26 Exhibits Therapeutic Effects Against PH in Rats

E64FC26 is a potent PDI pan-inhibitor, with IC_{50} of 16.3 μM against TXNDC5[23]. To evaluate the toxicity and safety of E64FC26, we performed histological analyses on the major organs of rats continuously treated with E64FC26 for 5 weeks under normoxic conditions. In the kidneys, the renal tubules and glomeruli maintained their integrity, with no signs of tubular dilation, epithelial cell injury, necrosis, or noticeable inflammatory cell infiltration in the kidneys. Similarly, hepatocytes were well-organized with clear cell boundaries, and there were no indications of necrosis, cell swelling, or inflammatory cell infiltration in the liver. Histological examination of the spleen revealed intact white and red pulp structures, without evidence of lymphoid depletion, congestion, or inflammatory infiltration (**Figure S8A**). Furthermore, body weight measurements showed no significant differences after E64FC26 treatment compared with the vehicle (**Figure S8B**). These findings collectively indicate that E64FC26 exhibits a favorable safety profile without causing overt organ or systemic toxicity.

Then, to examine whether pharmacological inhibition of TXNDC5 activity produces a therapeutic effect on PH, E64FC26 was administered intraperitoneally in SuHx-induced PH rats (**Figure 8A**). Hemodynamic measurements and morphometric analysis demonstrated that E64FC26 administration attenuated the elevation of RVSP

(**Figure 8B**) and the increase in RV/(LV+S) ratio (**Figure 8C**) in SuHx-induced PH rats compared with those with vehicle treatment. Histological analyses revealed that E64FC26 reversed SuHx-induced PA remodeling compared with vehicle-treated rats, as evidenced by reduced distal PA muscularization, decreased medial wall thickness, and attenuated collagen deposition area (**Figure 8D** and **Figure S8C-E**). Echocardiographic assessment further indicated that E64FC26 alleviated RV dysfunction in SuHx-induced PH rats (**Figure S8F**). Consistent with the echocardiographic improvements, histological analyses revealed that E64FC26 administration significantly reversed the RV hypertrophy (**Figure 8E**) and reduced the RV fibrosis area (**Figure 8F**) in SuHx-induced PH rats compared with those treated with vehicle.

To assess whether the therapeutic effects of E64FC26 in PH *in vivo* are primarily mediated through TXNDC5 rather than off-target inhibition of other PDI family members, we treated TXNDC5^{ECKO} mice with E64FC26 under SuHx conditions (**Figure 8G**). Notably, E64FC26 treatment also exhibited therapeutic effects in TXNDC5$^{f/f}$ mice against SuHx challenged, consistent with the results observed in rats. More importantly, in TXNDC5 endothelial-deficient mice, administration of E64FC26 did not confer additional protection against SuHx-induced distal PA remodeling, as indicated by unchanged medial wall thickness and collagen deposition area compared with vehicle-treated TXNDC5^{ECKO} mice (**Figure 8H** and **Figure S8G-H**). Similarly, lung angiography showed that E64FC26 administration failed to improve distal pulmonary perfusion in SuHx-treated TXNDC5^{ECKO} mice (**Figure 8I**). Consistently, E64FC26 also did not attenuate RVSP elevation (**Figure S8I**), RV hypertrophy (**Figure S8J**), and RV functional impairment (**Figure S8K**) in these mice. These findings indicated that the therapeutic effect of E64FC26 in PH would be largely dependent on the targeting of endothelial TXNDC5 rather than off-target inhibition of other PDI family members. In a word, these results indicate that pharmacological inhibition of TXNDC5 activity may be a promising strategy to alleviate the progression of PH.

Pulmonary Endothelium-Targeted TXNDC5 Gene Therapy Ameliorates PH

We further established a pulmonary endothelium-targeted gene therapy approach
to enhance organ specificity. Using a self-complementary adeno-associated virus
(scAAV) vector encoding shRNA against TXNDC5 (shTXN), we achieve rapid and
endothelial-specific gene silencing *in vivo*. The vector was administered via the tail vein
to SuHx-induced PH rats to selectively suppress TXNDC5 in lung ECs and evaluate its
therapeutic efficacy (**Figure 8J**). *Ex vivo* imaging confirmed efficient lung targeting of
the AAV (**Figure 8K**), and Western blotting demonstrated marked suppression of
TXNDC5 expression in primary rat lung ECs (**Figure S8L**).

Treatment with shTXN significantly attenuated SuHx-induced PA remodeling
compared with rats receiving the control vector (shNC), as evidenced by reduced distal
PA muscularization, decreased medial wall thickness, and lower collagen deposition
area (**Figure 8L** and **Figure S8M-O**). Hemodynamic measurements further
demonstrated that shTXN administration attenuated the elevation of RVSP in SuHx-
induced PH rats (**Figure 8M**). Moreover, morphological and histological analyses
showed that shTXN decreased the RV/(LV+S) ratio (**Figure S8P**) and alleviated RV
wall thickening (**Figure 8N**). Consistently, echocardiographic assessment revealed
improved RV function, reflected by increased PAT/PET, TAPSE, RVFAC, and reduced
RVFWT (**Figure S8Q**). These findings demonstrate that endothelial-targeted TXNDC5
silencing effectively mitigates pulmonary vascular remodeling and right heart
dysfunction in PH. Collectively, these findings demonstrate that endothelial-targeted
TXNDC5 silencing effectively mitigates pulmonary vascular remodeling and right
heart dysfunction in PH, highlighting the therapeutic potential of TXNDC5 inhibition.
439

DISCUSSION

PDI family catalyzes disulfide bond formation and rearrangement in the
endoplasmic reticulum. Using 4D label-free proteomics, we identified TXNDC5 as the
most significantly dysregulated PDI family member in the lungs of patients with
hypoxic PH. TXNDC5, dominantly expressed in endothelial cells, was also known as
endothelial PDI (Endo-PDI)[24]. We found that the silencing of TXNDC5 inhibited the

proliferation and migration of hPAECs, whereas its overexpression led to changes in the cytoskeletal architecture of hPAECs, implying a critical role of TXNDC5 in endothelial dysfunction in PH. *In vivo*, global or endothelial-specific TXNDC5 deletion attenuated, while endothelial TXNDC5 overexpression exacerbated SuHx-induced PH. Pharmacological inhibition with E64FC26, a pan-PDI inhibitor with an IC50 of 16.3 μM against TXNDC5, alleviated pulmonary arterial remodeling and right ventricular hypertrophy, thereby preventing PH progression in rats. These findings identify TXNDC5 as a promising therapeutic target for PH.

We next examined the molecular mechanisms of the critical role of TXNDC5 in PH development. Accumulating evidence indicates that ECM remodeling, which precedes pulmonary vascular remodeling, is a key driver of PH progression by altering vascular stiffness and resistance[25]. In this study, we identified TXNDC5 as a central regulator of ECM homeostasis in hypoxic PH. TXNDC5 broadly modulated ECM components, including proteoglycans and glycoproteins that govern matrix stiffness and elasticity, collagens that provide structural support, matrix modulators and adhesion proteins involved in ECM crosslinking, and metallopeptidases responsible for ECM turnover. TXNDC5 also influenced basement membrane proteins and angiogenesis-related factors, thereby affecting endothelial barrier function and vascular homeostasis. Notably, excessive deposition of POSTN, BGN, LOX, and TGM2 was observed in the PAs of SuHx-treated TXNDC5[f/f] mice but was markedly reduced in TXNDC5[ECKO] mice. As key ECM regulators, POSTN and BGN promote matrix organization and mechanotransduction[26–28], whereas LOX and TGM2 facilitate ECM crosslinking and collagen stabilization[29,30], collectively driving vascular stiffening and remodeling. These findings imply that TXNDC5 inhibition shifts the ECM equilibrium toward a less fibrotic and more compliant phenotype, mitigating PH progression by preserving vascular ECM homeostasis.

BGN has been implicated in inflammation, metabolic, musculoskeletal, and oncogenic diseases[31]. In the present study, we identified BGN as a direct binding partner of TXNDC5, with enhanced interaction with hypoxia exposure. Mutations at Cys89

and Gly351 within the PDI active domains of TXNDC5 markedly disrupted TXNDC5–BGN binding, indicating that PDI activity is essential for this interaction. Consistently, WT-TXNDC5, but not the PDI-inactive mutant, significantly increased FRET efficiency in the BGN folding assay, demonstrating that TXNDC5 facilitates BGN folding in a PDI-dependent manner. This process is critical for BGN maturation, stability, and secretion. Further investigation revealed that BGN was upregulated in the lungs of PH patients and rodent models. Endothelial cell-specific deficiency of BGN significantly alleviated PH progression in mice, highlighting endothelial-derived BGN as a key contributor to disease pathogenesis. Importantly, BGN deficiency largely abolished TXNDC5 overexpression-induced distal PA remodeling and RV hypertrophy, establishing BGN as a critical downstream effector of TXNDC5-driven PH. Notably, serum BGN levels were significantly elevated in PH patients and strongly correlated with disease severity, suggesting that BGN may serve as a potential diagnostic biomarker for PH.

HIF signaling is a central driver of hypoxic PH, with HIF-α subunits rapidly degraded under normoxia via the VHL E3 ligase[32,33]. In this study, we showed that TXNDC5-induced upregulation of HIF-1α/2α depends on BGN, as endothelial BGN silencing restored VHL-mediated ubiquitination and degradation of HIF-1α/2α. These findings uncover a previously unrecognized mechanism linking ECM remodeling to hypoxia-driven signaling in PH and suggest that TXNDC5, BGN, and HIF form a coordinated regulatory axis during disease progression. Excessive systemic inhibition of HIF-1α or HIF-2α has been associated with organ dysfunction, developmental defects, and embryonic lethality in mice, highlighting safety concerns for direct HIF targeting.[34–36]. We found that global TXNDC5 deletion prevented SuHx-induced HIF activation without causing adverse effects on growth or organ development. These findings suggest that targeting TXNDC5 offers a more tissue-selective strategy to modulate pathological HIF signaling while minimizing systemic toxicity, providing a promising therapeutic alternative for PH.

In summary, we identified for the first time the novel role of PDI protein TXNDC5

in the pathogenesis of hypoxic PH. TXNDC5 facilitates ECM protein (e.g., BGN) folding and reduces HIF-1α/2α ubiquitination, which contributes to endothelial dysfunction. TXNDC5 promotes a positive feedback loop between BGN and HIF signaling, driving the progression of pulmonary vascular remodeling in PH. This study highlights TXNDC5 as a promising therapeutic target for hypoxia-induced PH and lays the groundwork for future clinical investigations.

Acknowledgments

We thank Professor John Shyy from the University of California, San Diego, United States, for critical reading and advice on this article.

Sources of Funding

This work was supported by the Key Program of National Natural Science Foundation of China (NSFC, grant No. 82430004 to Jin-Song Bian), Shenzhen Science Fund for Distinguished Young Scholars (grant No. RCJC20210706091946002 to Xiaowei Nie), Shenzhen Science and Technology Program (grant No. GJHZ20240218111401002 to Jin-Song Bian), the National Natural Science Foundation of China (NSFC, grant No. 82241021 to Xiaowei Nie), and the National Natural Science Foundation of China (NSFC, grant No. 82400063 to Junting Zhang).

References

1. Johnson S, Sommer N, Cox-Flaherty K, Weissmann N, Ventetuolo CE, Maron BA. Pulmonary Hypertension: A Contemporary Review. *Am J Respir Crit Care Med*. 2023;208:528–548.

2. Shlobin OA, Adir Y, Barbera JA, Cottin V, Harari S, Jutant E-M, Pepke-Zaba J, Ghofrani H-A, Channick R. Pulmonary hypertension associated with lung diseases. *Eur Respir J*. 2024;64:2401200.

3. Evans CE, Cober ND, Dai Z, Stewart DJ, Zhao Y-Y. Endothelial cells in the pathogenesis of pulmonary arterial hypertension. *Eur Respir J*. 2021;58:2003957.

4. Ma B, Cao Y, Qin J, Chen Z, Hu G, Li Q. Pulmonary artery smooth muscle cell

phenotypic switching: A key event in the early stage of pulmonary artery
hypertension. *Drug Discov Today*. 2023;28:103559.

5. Mandras SA, Mehta HS, Vaidya A. Pulmonary Hypertension: A Brief Guide for
Clinicians. *Mayo Clin Proc*. 2020;95:1978–1988.

6. Olsson KM, Corte TJ, Kamp JC, Montani D, Nathan SD, Neubert L, Price LC,
Kiely DG. Pulmonary hypertension associated with lung disease: new insights
into pathomechanisms, diagnosis, and management. *Lancet Respir Med*.
2023;11:820–835.

7. Kanemura S, Matsusaki M, Inaba K, Okumura M. PDI Family Members as Guides
for Client Folding and Assembly. *Int J Mol Sci*. 2020;21:9351.

8. Kuramochi T, Yamashita Y, Arai K, Kanemura S, Muraoka T, Okumura M.
Boosting the enzymatic activity of CxxC motif-containing PDI family members.
Chem Commun (Camb). 2024;60:6134–6137.

9. Nagarkoti S, Kim Y-M, Ash D, Das A, Vitriol E, Read T-A, Youn S-W, Sudhahar
V, McMenamin M, Hou Y, et al. Protein disulfide isomerase A1 as a novel redox
sensor in VEGFR2 signaling and angiogenesis. *Angiogenesis*. 2023;26:77–96.

10. Kobayashi Y, Oguro A, Hirata Y, Imaoka S. The regulation of Hypoxia-Inducible
Factor-1 (HIF-1alpha) expression by Protein Disulfide Isomerase (PDI). *PLoS*
One. 2021;16:e0246531.

11. Hung C-T, Tsai Y-W, Wu Y-S, Yeh C-F, Yang K-C. The novel role of ER protein
TXNDC5 in the pathogenesis of organ fibrosis: mechanistic insights and
therapeutic implications. *J Biomed Sci*. 2022;29:63.

12. Jin Y, Sharma A, Bai S, Davis C, Liu H, Hopkins D, Barriga K, Rewers M, She J-
X. Risk of type 1 diabetes progression in islet autoantibody-positive children can
be further stratified using expression patterns of multiple genes implicated in
peripheral blood lymphocyte activation and function. *Diabetes*. 2014;63:2506–
2515.

13. Wang L, Dong H, Song G, Zhang R, Pan J, Han J. TXNDC5 synergizes with
HSC70 to exacerbate the inflammatory phenotype of synovial fibroblasts in
rheumatoid arthritis through NF-κB signaling. *Cell Mol Immunol*. 2018;15:685–
696.

14. Wang X, Li H, Chang X. The role and mechanism of TXNDC5 in diseases. *Eur J*
Med Res. 2022;27:145.

15. Yeh C-F, Cheng S-H, Lin Y-S, Shentu T-P, Huang R-T, Zhu J, Chen Y-T, Kumar
S, Lin M-S, Kao H-L, et al. Targeting mechanosensitive endothelial TXNDC5 to
stabilize eNOS and reduce atherosclerosis in vivo. *Sci Adv*. 2022;8:eabl8096.

16. Wu S, Luo X, Chen Y, Wang Z, Liu X, Sun N, Zhao J, Luo W, Zhang J, Tong X, et al. Sodium-glucose cotransporter 2 inhibitors attenuate vascular calcification by suppressing endoplasmic reticulum protein thioredoxin domain containing 5 dependent osteogenic reprogramming. *Redox Biol*. 2024;73:103183.

17. Fu Y, Zhou Y, Wang K, Li Z, Kong W. Extracellular Matrix Interactome in Modulating Vascular Homeostasis and Remodeling. *Circ Res*. 2024;134:931–949.

18. Thenappan T, Chan SY, Weir EK. Role of extracellular matrix in the pathogenesis of pulmonary arterial hypertension. *Am J Physiol Heart Circ Physiol*. 2018;315:H1322–H1331.

19. Nie X, Shen C, Tan J, Wu Z, Wang W, Chen Y, Dai Y, Yang X, Ye S, Chen J, et al. Periostin: A Potential Therapeutic Target For Pulmonary Hypertension? *Circ Res*. 2020;127:1138–1152.

20. Bian J-S, Chen J, Zhang J, Tan J, Chen Y, Yang X, Li Y, Deng L, Chen R, Nie X. ErbB3 Governs Endothelial Dysfunction in Hypoxia-Induced Pulmonary Hypertension. *Circulation*. 2024;150:1533–1553.

21. Zhang Y, Li X, Li S, Zhou Y, Zhang T, Sun L. Immunotherapy for Pulmonary Arterial Hypertension: From the Pathogenesis to Clinical Management. *Int J Mol Sci*. 2024;25:8427.

22. Luo Z, Tian M, Yang G, Tan Q, Chen Y, Li G, Zhang Q, Li Y, Wan P, Wu J. Hypoxia signaling in human health and diseases: implications and prospects for therapeutics. *Signal Transduct Target Ther*. 2022;7:218.

23. Robinson RM, Reyes L, Duncan RM, Bian H, Reitz AB, Manevich Y, McClure JJ, Champion MM, Chou CJ, Sharik ME, et al. Inhibitors of the protein disulfide isomerase family for the treatment of multiple myeloma. *Leukemia*. 2019;33:1011–1022.

24. Sullivan DC, Huminiecki L, Moore JW, Boyle JJ, Poulsom R, Creamer D, Barker J, Bicknell R. EndoPDI, a novel protein-disulfide isomerase-like protein that is preferentially expressed in endothelial cells acts as a stress survival factor. *J Biol Chem*. 2003;278:47079–47088.

25. Jandl K, Radic N, Zeder K, Kovacs G, Kwapiszewska G. Pulmonary vascular fibrosis in pulmonary hypertension - The role of the extracellular matrix as a therapeutic target. *Pharmacol Ther*. 2023;247:108438.

26. Sun X-J, Ma W-Q, Zhu Y, Liu N-F. POSTN promotes diabetic vascular calcification by interfering with autophagic flux. *Cell Signal*. 2021;83:109983.

27. Nikoloudaki G, Snider P, Simmons O, Conway SJ, Hamilton DW. Periostin and matrix stiffness combine to regulate myofibroblast differentiation and fibronectin

synthesis during palatal healing. *Matrix Biol*. 2020;94:31–56.

28.   Yu M, He X, Song X, Gao J, Pan J, Zhou T, Wang Q, Zhu W, Ma H, Zeng H, et al.
Biglycan promotes hepatic fibrosis through activating heat shock protein 47. *Liver*
Int. 2023;43:500–512.

29.   Wang Z-J, Guan Q-W, Zhou H-H, Mao X-Y, Chen F-H. Mechanistic insight into
lysyl oxidase in vascular remodeling and angiogenesis. *Genes Dis*. 2023;10:771–
785.

30.   Zhang M, Li L, Zhang W, Li M, Yan G, Tang C. TG2 participates in pulmonary
vascular remodelling by regulating the senescence of pulmonary artery smooth
muscle cells. *Cell Signal*. 2024;121:111296.

31.   Appunni S, Rubens M, Ramamoorthy V, Anand V, Khandelwal M, Sharma A.
Biglycan: an emerging small leucine-rich proteoglycan (SLRP) marker and its
clinicopathological significance. *Mol Cell Biochem*. 2021;476:3935–3950.

32.   Tarade D, Ohh M. The HIF and other quandaries in VHL disease. *Oncogene*.
2018;37:139–147.

33.   Pullamsetti SS, Mamazhakypov A, Weissmann N, Seeger W, Savai R. Hypoxia-
inducible factor signaling in pulmonary hypertension. *J Clin Invest*.
2020;130:5638–5651.

34.   Kotch LE, Iyer NV, Laughner E, Semenza GL. Defective vascularization of HIF-
1alpha-null embryos is not associated with VEGF deficiency but with
mesenchymal cell death. *Dev Biol*. 1999;209:254–267.

35.   Chen H, Houshmand G, Mishra S, Fong G-H, Gittes GK, Esni F. Impaired
pancreatic development in Hif2-alpha deficient mice. *Biochem Biophys Res*
Commun. 2010;399:440–445.

36.   Peng Y-J, Nanduri J, Khan SA, Yuan G, Wang N, Kinsman B, Vaddi DR, Kumar
GK, Garcia JA, Semenza GL, et al. Hypoxia-inducible factor 2α (HIF-2α)
heterozygous-null mice exhibit exaggerated carotid body sensitivity to hypoxia,
breathing instability, and hypertension. *Proc Natl Acad Sci U S A*. 2011;108:3065–
3070.

Figure 1. TXNDC5 was upregulated in PH.

A, Heatmap of 4D label-free proteomics showing PDI family expression in lung tissues

from 5 non-PH donors and 5 patients with hypoxic PH. **B**, Representative Western blots and quantification of TXNDC5 and Vinculin in the lungs of control donors and PH patients (n=7). **C**, qPCR analysis of TXNDC5 mRNA levels in lung homogenates from donors and patients with PH (n=7). **D**, Western blots showing the TXNDC5 protein levels in the lungs collected from MCT (n=7) or SuHx-treated rats (n=6) or rats under normoxia. **E**, **F**, and **G**, Representative confocal images of multiplex immunofluorescence (mIF) staining for TXNDC5 (red), CD31 (green), α-SMA (white), and COL1A1 (purple) in distal pulmonary arteries (PAs) from donors, ILD-PH, and COPD-PH patients. Nuclei were counterstained with DAPI (blue). White arrows indicate the colocalization of TXNDC5 with CD31 (yellow). White dashed lines indicate the outer layer of smooth muscle cells. Scale bars=25 μm. **H**, Quantitative analysis showing elevated TXNDC5 mean fluorescence intensity (MFI) in the $CD31^+$ intima of distal PAs from patients with ILD-PH and COPD-PH compared with non-PH donors. (n=5). **I**, **J**, and **K**, Representative images of mIF staining for TXNDC5 (red), CD31 (green), α-SMA (cyan), and COL1A1 (purple) in distal PAs from SuHx- or MCT-treated rats or rats under normoxia. Nuclei were counterstained with DAPI (blue). White arrows indicate the colocalization of TXNDC5 with CD31 (yellow). White dashed lines indicate the outer layer of smooth muscle cells. Scale bars=20 μm. **L**, **M**, and **N**, Correlation analyses of intimal TXNDC5 MFI and right ventricular systolic pressure (RVSP), Fulton index [RV/(LV+S)], and tricuspid annular plane systolic excursion (TAPSE) in rats. Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**B-D**) or 2-way ANOVA (**H**), followed by Tukey honestly significant difference test for multiple comparisons and the Pearson correlation test (**L-N**). PDI indicates protein disulfide isomerase; PH, pulmonary hypertension; TXNDC5, thioredoxin domain containing 5; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; DAPI, 4′,6-diamidino-2-phenylindole; Nor, normoxia; SuHx, Sugen5416/hypoxia; and MCT, monocrotaline.

Figure 2. TXNDC5 contributed to SuHx-induced pulmonary vascular remodeling
A, Schematic protocol for the animal treatments in the SuHx-induced PH mice. **B**,
Representative tracings and quantification of right ventricular systolic pressure (RVSP)
in WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=8). **C**, Representative

multiplex immunofluorescence (mIF) for CD31 (green), α-SMA (white), and COL1A1 (red), and Victoria blue/Sirius red (VB/SR) staining for elastic lamina (dark blue), collagens (red), and muscle fiber (yellow) in distal pulmonary arteries (PAs) from WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx. Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **D**, Quantification of distal PA muscularization based on α-SMA positive area in WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=8). **E**, Measurement of medial wall thickness in distal PAs, defined by the elastic lamina visualized with VB staining (n=8). **F**, Quantification of collagen deposition area in distal PAs based on SR staining, indicating decreased collagen deposition in TXNDC5-deficient mice (n=8). **G**, Immunofluorescence staining of WGA (green) in right ventricles from WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=8). Nuclei were counterstained with DAPI (blue). Scale bars=20 μm. **H**, Schematic diagram illustrating the *in vivo* pulmonary endothelial-specific overexpression of TXNDC5 in mice. Mice received a single tail vein injection of AAV-NC or AAV-TXN at the beginning of the 2 weeks, followed by SuHx exposure. **I**, Representative images of major organs from saline-treated control or AAV-TXN-treated mice, showing GFP fluorescence delivered by AAV detected by IVIS imaging system. **J**, Hemodynamic measurements of RVSP in mice administered with AAV-NC or AAV-TXN under normoxia or SuHx situations (n=8). **K**, Representative mIF of distal PAs from AAV-NC or AAV-TXN-treated mice under SuHx or normoxia, showing CD31 (green), α-SMA (white), and COL1A1 (red), together with VB/SR staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **L**, **M,** and **N**, Quantitative analysis of distal PA remodeling in AAV-NC or AAV-TXN-treated mice under normoxic or SuHx conditions (n=8). PA muscularization, medial wall thickness, and collagen deposition area were assessed using α-SMA, VB, and SR staining, respectively. Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by Tukey honestly significant difference test for multiple comparisons. DAPI indicates 4′,6-diamidino-2-phenylindole; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; WGA, wheat germ agglutinin;

Figure 3. Endothelial cell-specific deletion of TXNDC5 ameliorated SuHx-induced PH.

A, Schematic protocol for the endothelial cell-specific TXNDC5 knockout mice assays.

Mice were intraperitoneally injected with 75 mg/kg tamoxifen once daily for 5 days

before being exposed to SuHx. **B**, Hemodynamic measurements of right ventricular

systolic pressure (RVSP) in TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice exposed to normoxia

or SuHx (n=8). **C**, Fulton index [RV/(LV+S)] in TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice

(n=8). **D**, Representative multiplex immunofluorescence (mIF) of distal pulmonary

arteries (PAs) from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice under normoxia or SuHx,

showing CD31 (green), α-SMA (white), and COL1A1 (red), together with Victoria

blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens

(red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L

indicates lumen. Scale bars=25 μm. **E**, Quantification of distal PA muscularization

based on α-SMA positive area (n=8). **F**, Medial wall thickness of distal PAs, measured

across the elastic lamina visualized by VB staining (n=8). **G**, Perivascular collagen

deposition quantified using SR staining (n=8). **H**, Representative lung angiogram of the

left lungs from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with SuHx or under

normoxic conditions. Scale bars=2 mm. **I**, Histological analysis of the cross-sectioned

heart at the mid-RV (red arrows) plane from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice with

or without SuHx exposure (n=8). Scale bars=1 mm. **J**, Immunofluorescence staining of

WGA (green) in RV from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice exposed to normoxia or

SuHx (n=8). Nuclei were counterstained with DAPI (blue). Scale bars=20 μm. Data

represent the mean±SD. Comparisons of parameters were performed with 2-way

ANOVA, followed by Tukey honestly significant difference test for multiple

comparisons. DAPI indicates 4′,6-diamidino-2-phenylindole; H&E, hematoxylin-eosin;

α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; WGA, wheat germ

agglutinin; f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; and RV, right

ventricle.

Figure 4. TXNDC5 regulated ECM protein expression and distribution.

A, HPAECs were transfected with siTXNDC5 or siNC under hypoxic conditions, followed by RNA sequencing analysis (n=3). Bubble chart showing the Gene Ontology enrichment using DAVID for the differentially expressed genes. **B**, Differentially

expressed ECM proteins were selected and visualized by heatmap (left). The PPI

network showed that TXNDC5-targeted ECM proteins could be classified into different

functional subgroups (right). **C**, **D**, **E**, and **F**, Representative images of

immunofluorescence staining of α-SMA (red), CD31 (green), and POSTN, BGN, LOX,

or TGM2 (white) in PAs from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice under normoxia and

SuHx conditions. Nuclei were counterstained with DAPI (blue). L indicates lumen.

Scale bars=20 μm. **G**, Western blots showing the protein levels of COL1A1, TGM2,

FN1, POSTN, and LOX in hPAECs transferred with siTXNDC5 or siNC in normoxia

or hypoxia conditions (n=6). Data represent the mean±SD. Comparisons of parameters

were performed with the 2-way ANOVA, followed by the Tukey honestly significant

difference test for multiple comparisons. ECM indicates extracellular matrix; PPI,

protein-protein interaction; Nor, normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia;

si, small interfering; α-SMA, α-smooth muscle actin; POSTN, periostin; BGN, biglycan;

LOX, lysyl oxidase; TGM2, transglutaminase 2; FN1, fibronectin; and COL1A1,

collagen alpha-1 chain.

Figure 5. BGN served as a biomarker and therapeutic target of PH.

A, The progressive screening of the PPI network between ECM genes and ECs function-related genes. Each new PPI network is composed of core genes in the previous network. **B,** Box-whisker plot showing the relative mRNA levels of TXNDC5

in lung tissues from PH patients (n=96) and healthy donors (n=52) (GSE254617). **C**,

Enzyme-linked immunosorbent assay showing the BGN level in the serum of non-PH

donors (n=27) and patients with PH (n=27). **D-E**, Linear regression analysis showing

the correlation between serum BGN concentrations with sPAP and mPAP of PH patients.

F, Quantification of right ventricular systolic pressure (RVSP) in BGN^{flox} and BGN^{ECKO}

mice exposed to normoxia or SuHx (n=8). **G**, Fulton index [RV/(LV+S)] in BGN^{flox}

and BGN^{ECKO} mice exposed to normoxia or SuHx (n=8). **H**, Representative multiplex

immunofluorescence (mIF) of distal pulmonary arteries (PAs) from BGN^{flox} and

BGN^{ECKO} mice under normoxia or SuHx, showing CD31 (green), α-SMA (white), and

COL1A1 (red), together with Victoria blue/Sirius red (VB/SR) staining of elastic

lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were

counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **I**,

Quantification of distal PA muscularization based on α-SMA positive area (n=8). **J**,

Medial wall thickness of distal PAs, measured across the elastic lamina visualized by

VB staining (n=8). **K**, Quantification of collagen deposition of distal PAs from BGN^{flox}

and BGN^{ECKO} mice under normoxia or SuHx(n=8). **L**, Representative lung angiogram

of the left lungs from BGN^{flox} and BGN^{ECKO} mice treated with SuHx or under normoxic

conditions. Scale bars=2 mm. **M**, Histological analysis of heart cross-sections at the

mid-RV plane (top) and highlighted right ventricular wall (bottom) of the BGN^{flox} and

BGN^{ECKO} mice exposed to normoxia or SuHx (n=8). Scale bars=1 mm. Data represent

the mean±SD. Comparisons of parameters were performed with the 2-tailed Mann-

Whitney *U* test (**B** and **C**), Pearson correlation test (**D** and **E**), or 2-way ANOVA (**F-G**,

I-K, and **M**), followed by Tukey honestly significant difference test for multiple

comparisons. H&E indicates hematoxylin-eosin; α-SMA, α-smooth muscle actin;

COL1A1, collagen type I alpha 1; DAPI, 4′,6-diamidino-2-phenylindole; Nor,

normoxia; SuHx, Sugen5416/hypoxia; RV, right ventricle; sPAP, systolic pulmonary

arterial pressure; and mPAP, mean pulmonary arterial pressure.

Figure 6. TXNDC5 assisted BGN protein folding to promote PH.

A, Immunofluorescence staining showing the localization of TXNDC5 (green) and BGN (red) in hPAECs. Nuclei (blue) were stained with DAPI. White arrows in the enlarged view indicate the colocalization of TXNDC5 with BGN (yellow). Scale

bars=10 μm. **B**, Coimmunoprecipitation analysis showing the interaction between TXNDC5 and BGN in hPAECs under both normoxic and hypoxic situations. IP: TXNDC5, immunoprecipitation with TXNDC5 antibody. IgG for the negative control. **C**, Molecular docking showing the binding domains of human TXNDC5 and BGN. **D**, Immunoprecipitation with anti-HA beads in HEK293T cells transfected with FLAG-BGN and HA-TXNDC5 mutant plasmids. **E**, FRET acceptor photobleaching of the hPAECs transfected with CFP-BGN-YFP plasmid. CFP and YFP images were taken both before and after acceptor photobleaching. FRET efficiency is displayed as a pseudocolor representation and qualification by ImageJ (n=5). Scale=10 μm. **F**, Schematic of the *in vivo* rescue experiments: adeno-associated virus encoding TXNDC5 (AAV-TXN) or control AAV (AAV-NC) was delivered to lung ECs in BGN^{flox} and BGN^{ECKO} mice under normoxia or SuHx conditions. **G**, Representative multiplex immunofluorescence (mIF) images of distal pulmonary arteries (PAs) from BGN^{flox} and BGN^{ECKO} mice treated with AAV-NC or AAV-TXN, showing CD31 (green), α-SMA (white), and COL1A1 (red), with Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei stained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **H**, Quantification of distal PA muscularization in BGN^{flox} and BGN^{ECKO} mice with AAV-NC or AAV-TXN administration (n=6). **I**, Medial wall thickness of distal PAs measured across the elastic lamina, visualized by VB staining (n=6). **J**, Quantification of collagen deposition area in distal PAs (n=6). **K**, Right ventricular systolic pressure (RVSP) in BGN^{flox} and BGN^{ECKO} mice treated with AAV-NC or AAV-TXN under SuHx or normoxia conditions (n=6). **L**, Fulton index [RV/(LV+S)] in BGN^{flox} and BGN^{ECKO} mice with AAV-NC or AAV-TXN treatment (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the 1-way (**E**) or 2-way (**H-L**) ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. ECs, endothelial cells; Nor, normoxia; Hyp, hypoxia; IgG, immunoglobulin G; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; FRET, fluorescence resonance energy transfer; CFP, cyan fluorescent protein; and YFP, yellow fluorescent protein.

Figure 7. Endothelial TXNDC5 Driven PH via BGN-dependent stabilization of

HIF-1α/2α.

A, Dual-luciferase assay showing that luciferase activity in constructs with BS1

mutation (Mut-1), BS2 mutation (Mut-2), or double mutation of BS1 and BS2 (dMut)

(n=6). **B**, Western blots showing the decreased protein levels of HIF-1α and HIF-2α in the lungs from TXNDC5 deficiency mice compared with WT exposed to SuHx (n=6). **C**, Representative distal pulmonary arteries (PAs) images from AAV-NC or AAV-TXN-treated mice exposed to SuHx or normoxia, with HIF-1α inhibitor (PX-478) or HIF-2α inhibitor (PT-2385), or vehicle administration. Multiplex immunohistochemical (mIF) showing CD31 (green), α-SMA (white), and COL1A1 (red) and Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **D**, **E**, and **F**, Quantitative analysis of distal PA remodeling, including PA muscularization assessed by α-SMA-positive area (n=6), medial wall thickness measured across the elastic lamina visualized by VB staining (n=6), and perivascular collagen area quantified using SR staining (n=6). **G**, Representative Western blots showing that the LV-TXNDC5-induced upregulation of HIF-1/2α was abolished by siBGN in hPAECs both under normoxia and hypoxia conditions (n=6). **H**, Western blot showing that the protein stabilization of HIF-1/2α was enhanced by LV-TXNDC5 treatment in hPAECs under normoxic conditions (n=5). **I**, Representative Western blots of HIF-1α, HIF-2α, and VHL in hPAECs after treatment with TXNDC5 knockdown only or double knockdown (dKO) of TXNDC5 and VHL (n=6). **J**, Western blot analysis indicating the HIF-2α ubiquitylation profile was reduced by LV-TXNDC5 but increased by siBGN treatment. hPAECs were treated with 10 μM MG132 for 6 h before harvest. Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. BS indicates binding site; VHL, von Hippel-Lindau; Ub, ubiquitin; Nor, normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypixa; IgG, immunoglobulin G; LV, lentivirus; α-SMA, α-smooth muscle actin; and COL1A1, collagen type I alpha 1.

Figure 8. Pharmacological and Pulmonary Endothelium-Targeted Inhibition of TXNDC5 Ameliorated PH.

A, Schematic protocol for E64FC26 administration for SuHx-induced PH rats. Rats were treated with SuHx for 3 weeks and then intraperitoneal injection with 2 mg/kg

E64FC26 daily for 2 weeks under normoxic conditions. **B**, Quantification of right

ventricular systolic pressure (RVSP) in rats administered with E64FC26 or vehicle

exposed to normoxia or SuHx (n=6). **C**, Fulton index [RV/(LV+S)] of rats with

E64FC26 or vehicle treatment under normoxia or SuHx conditions (n=6). **D**,

Representative multiplex immunofluorescence (mIF) of distal pulmonary arteries (PAs)

from rats treated with vehicle or E64FC26 under normoxia or SuHx conditions,

showing CD31 (green), α-SMA (white), and COL1A1 (red), together with Victoria

blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens

(red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L

indicates lumen. Scale bars = 25 μm. **E**, Histological and quantification analysis of the

cross-sectioned heart at the mid-RV plane (top) and the highlighted right ventricular

wall (bottom) of rats treated with E64FC26 or vehicle under normoxia or SuHx

conditions (n=6). Scale bars=2 mm. **F**, Masson's trichrome staining and quantification

of the fibrotic area by collagen volume fraction measured by Image J of RV from rats

treated with E64FC26 or vehicle exposed to normoxia or SuHx (n=6). Scale bars=100

μm. **G**, Schematic of vehicle or E64FC26 treatment in TXNDC5$^{f/f}$ and TXNDC5^{ECKO}

mice under SuHx conditions to investigate whether the effects of E64FC26 are

TXNDC5-dependent. **H**, Representative distal PAs images from TXNDC5$^{f/f}$ and

TXNDC5^{ECKO} mice treated with vehicle or E64FC26 under SuHx or normoxia

conditions. Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue),

perivascular collagens (red), and muscle fiber (yellow). L indicates lumen. Scale

bars=25 μm. **I**, Representative lung angiogram of SuHx-treated TXNDC5$^{f/f}$ and

TXNDC5^{ECKO} mice with or without E64FC26 administration, showing distal

pulmonary perfusion. Scale bars=2 mm. **J**, Schematic diagram of the pulmonary

endothelium-targeted gene therapy strategy. Rats received a tail vein injection of self-

complementary adeno-associated virus (scAAV) encoding shRNA against TXNDC5

(shTXN) or control shRNA (shNC). **K**, Representative IVIS images of major organs

from rats treated with Saline or scAAV-shTXN, showing GFP fluorescence delivered

by the scAAV vector. **L**, Representative mIF for CD31 (green), α-SMA (white), and

COL1A1 (red), and VB/SR staining for elastic lamina (dark blue), collagens (red), and

muscle fiber (yellow) in distal PAs from rats receiving shNC or shTXN exposed to normoxia or SuHx. Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **M**, Hemodynamic measurement of RVSP in SuHx-induced PH rats treated with shNC or shTXN (n=6). **N**, Histological sections of the heart at the mid-RV plane and quantification of right ventricular wall thickness from SuHx-induced PH rats treated with shNC or shTXN (n=6). Scale bars = 2 mm. Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by Tukey honestly significant difference test for multiple comparisons. H&E indicates hematoxylin-eosin; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; shRNA, short hairpin RNA; f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; and RV, right ventricle.

Graphic Abstract. The role of TXNDC5 in regulating ECM homeostasis during PH. In the healthy state, BGN folding is properly maintained and HIF-1α/2α is rapidly degraded. During PH, TXNDC5 is upregulated, leading to excessive accumulation of BGN and impaired ubiquitination-mediated degradation of HIF-1α/2α, resulting in an imbalance in ECM homeostasis and contributing to vascular remodeling. Pharmacological or genetic inhibition of TXNDC5 restores BGN folding capacity and facilitates HIF-1α/2α degradation, thereby reestablishing ECM homeostasis and attenuating PH progression.

Supplemental Figures

Figure S1. Upregulation of TXNDC5 in the lungs and lung endothelial cells of PH

rodent models.

A, Schematic illustration showing the 4D label-free proteomics of lungs from non-PH

donors and patients with hypoxic PH. **B**, Western blot analysis and quantification of TXNDC5 and Vinculin in the lungs of mice exposed to SuHx or hypoxia or under normoxic situations (n=6). **C**, Quantitative analysis of mIF showing TXNDC5 was predominantly co-localized with CD31 in distal PAs. (n=5). **D**, Quantification of mean fluorescence intensity (MFI) of TXNDC5 in remodeled distal PAs from SuHx- and MCT-induced rat PH models compared with rats under normoxic conditions. (n=6). **E**, Western blot of TXNDC5 in primary lung ECs from SuHx-induced mice or normoxic controls, showing increased protein levels in ECs. **F**, Western blot of TXNDC5 in lung ECs isolated from SuHx-induced PH rats, demonstrating elevated expression compared with normoxic controls. Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student t-test (**B**, **E**, and **F**) or 2-way ANOVA (**D**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; and ECs, endothelial cells.

Figure S2. Role of TXNDC5 in PH development.

A, Fulton index [RV/(LV+S)] in WT and TXNDC5$^{-/-}$ mice exposed to normoxia or

SuHx (n=8). **B**, **C**, **D**, and **E**, Echocardiography analysis of PAT/PET, TAPSE, RVFAC,

and RVFWT in WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=8). **F,**

Representative confocal images showing GFP-TXNDC5 (green), CD31 (red), and α-SMA (white) in lung sections from AAV-TXN-treated mice. Nuclei were counterstained with DAPI (blue). L indicates lumen. A indicates airway. Scale bars=50 μm. **G**, Western blot analysis of TXNDC5 protein levels in primary lung ECs isolated from AAV-TXN- or AAV-NC-treated mice (n=6). **H**, RV/(LV+S) in mice administered with AAV-NC or AAV-TXN exposed to normoxia or SuHx (n=8). **I**, **J**, **K**, and **L**, Echocardiography analysis of PAT/PET, TAPSE, RVFAC, and RVFWT in mice administered with AAV-NC or AAV-TXN exposed to normoxia or SuHx (n=8). **M**, Representative Western blots and quantification of TXNDC5 protein levels in primary lung ECs isolated from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice (n=6). **N**, **O**, **P**, and **Q**, Echocardiographic assessment of right ventricular function in TXNDC5$^{f/f}$ and TXNDC5^{ECKO} under normoxia or SuHx, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=8). **R**, Enlarged lung angiogram images of the left lungs from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with SuHx or under normoxic conditions. Scale bars=2 mm. Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**G** and **M**) or 2-way ANOVA (**A-E**, **H-L,** and **N-Q**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; SuHx, Sugen5416/hypoxia; PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio; TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional area change; RVFWT, right ventricular free wall thickness; DAPI indicates 4′,6-diamidino-2-phenylindole; α-SMA, α-smooth muscle actin; and f/f, flox/flox.

Figure S3. TXNDC5 regulated cell function in pulmonary artery.

A and **B**, Quantification of TXNDC5 protein levels by Western blots and TXNDC5

mRNA levels by qPCR analysis in hPAECs in both normoxia and hypoxia situations

(n=6). **C**, Immunofluorescence staining analysis showing increased protein levels of

TXNDC5 exposed to hypoxia. Scale bars=10 μm. **D**, Representative EdU (red) staining image and group data showing the effects of TXNDC5 silencing on the proliferative activity of hPAECs (n=6). Nuclei were counterstained with DAPI (blue). Scale bars=200 μm. **E**, Representative images and group data showing the migration of hPAECs treated with hPAECs siTXNDC5 or siNC in normoxia or hypoxia conditions (n=6). Scale bars=200 μm. **F**, Flow cytometry showing the effect of siTXNDC5 on the cell cycle of hPAECs exposed to normoxia or hypoxia. The representative images and quantification of percentages of cells in G1, S, and G2 were shown (n=6). X and Y axes, respectively, indicate the denoted DNA content stained with propidium iodide (PI) and cell counts. **G** and **H**, Confocal microscopy images showing the effect of LV-TXNDC5 on the cytoskeletal architecture of hPAECs. Cells were stained with phalloidin (green, scale bars=20 μm) or paxillin (red, scale bars=10 μm). Nuclei were counterstained with DAPI (blue). **I** and **J**, Representative Western blots and quantification showing the protein levels of Cyclin A, proliferating cell nuclear antigen (PCNA), and cleaved-caspase 3 of hPASMCs treated with conditioned medium from hPAECs transfected with siTXNDC5 or siNC in normoxia or hypoxia conditions (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**A** and **B**) or 2-way ANOVA (**D**, **E**, **I**, and **J**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; LV, lentivirus; si, small interfering; CM, conditioned medium; and EdU, 5-ethynyl-20-deoxyuridine.

Figure S4. TXNDC5-driven endothelial paracrine signaling and ECM remodeling.

A, Representative EdU staining image and group data showing the proliferative activity

of hPASMCs treated with conditioned medium from hPAECs transfected with

siTXNDC5 (siTXN-CM) or siNC (siNC-CM) in normoxia or hypoxia conditions (n=6).

Nuclei were counterstained with DAPI (blue). Scale bars=200 μm. **B**, CCK-8 assays demonstrating decreased cell viability of hPASMCs after treatment with siTXN-CM or siNC-CM (n=6). **C**, Quantification of POSTN, BGN, LOX, and TGM2 deposition in distal PAs from SuHx-treated TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice, assessed by mean fluorescence intensity (MFI) (n=6). **D**, **E**, and **F**, Representative dot blot analysis of BGN, POSTN, and LOX in the supernatants of hPAECs transfected with siTXNDC5 or control siRNA (siNC). **G**, Quantification of dot blot intensities for BGN, POSTN, and LOX, showing significantly reduced secretion upon TXNDC5 knockdown in normoxia or hypoxia conditions (n=5). **H**, Venn diagram showing that BGN exists in the overlapping of the core ECM genes in the PPI network, the differentially expressed ECM proteins in TXNDC5 silencing hPAECs, and the differentially expressed ECM genes in hypoxia-treated hPAECs (GSE160255). **I**, Representative Western blots and quantification of BGN and vinculin in the lungs of donors and PH patients (n=7). **J**, Representative Western blots and quantification of BGN and β-actin in the lungs from mice exposed to SuHx or under normoxic conditions (n=6). **K**, Western blots showing the BGN protein levels in the lungs collected from MCT or SuHx-treated rats or rats under normoxia (n=7). **L**, Representative EdU staining image and group data showing 10 μg/ml recombinant human biglycan (rhBGN) treatment for 24 h enhanced the proliferative activity of hPASMCs (n=5). Scale bars=200 μm. **M**, CCK-8 assay confirming increased cell viability of hPASMCs after rhBGN treatment (n=6). **N**, Western blots showing the protein levels of cleaved-caspase 3 of hPASMCs treated with 10 μg/ml rhBGN or the negative control (bovine serum albumin, BSA) for 24 h (n=6). **O**, Flow cytometry analysis of Annexin V-FITC/PI staining demonstrating decreased apoptotic cell populations in hPASMCs treated with rhBGN (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**I-O**) or 2-way ANOVA (**A-C** and **G**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; MCT, monocrotaline; si, small interfering; DAPI, 4′,6-diamidino-2-phenylindole; f/f, flox/flox; POSTN, periostin; BGN, biglycan; and LOX, lysyl oxidase; TGM2, transglutaminase 2.

1012

Figure S5. TXNDC5 stabilized BGN to regulate its protein expression.

A, Representative Western blots and quantification of BGN protein levels in primary

lung ECs isolated from BGN^{flox} and BGN^{ECKO} mice (n=6). **B**, **C**, **D**, and **E**,

Echocardiographic assessment of right ventricular function in BGN^{flox} and BGN^{ECKO}

under normoxia or SuHx, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=8). **F**, Representative Western blots and quantification of TXNDC5 and BGN in hPAECs treated with siNC or siTXNDC5 under normoxia or hypoxia conditions (n=6). **G**, Western blot analysis showing the downregulation of BGN in the lungs from TXNDC5 deficiency mice compared with WT mice (n=6). **H**, Representative Western blots and quantification of TXNDC5 and BGN in lung ECs isolated from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice under normoxia or SuHx conditions (n=6). **I**, Representative Western blots showing that the LV-TXNDC5-induced upregulation of BGN in hPAECs both under normoxia and hypoxia conditions (n=6). **J**, Western blot analysis showing the upregulation of BGN in the lung ECs isolated from endothelial-specific AAV-NC or AAV-TXN-treated mice exposed to SuHx or under normoxic conditions (n=6). **K**, Coimmunoprecipitation analysis showing the interaction between TXNDC5 and BGN was enhanced by hypoxia stimulation in hPAECs. IP: BGN, immunoprecipitation with BGN antibody. IgG for the negative control. **L**, Scheme of WT (TXNDC5 WT) and mutant TXNDC5 (TXNDC5 Mut) sequences with Cysteine (C)-to-Alanine (A) mutations in all three thioredoxin domains. **M**, Schematic illustration for using FRET to determine the extent of BGN protein folding in hPAECs. When the BGN protein folds correctly (bottom), the distance between CFP and YFP will be less than 10 nm, resulting in FRET occurring. **N**, Western blot showing that TXNDC5 overexpression prolonged the BGN protein half-life in hPAECs (n=6). **O**, Western blot analysis demonstrating that TXNDC5 silencing destabilized BGN in hPAECs, resulting in accelerated protein degradation (n=6). **P**, **Q**, **R**, and **S**, Echocardiographic evaluation of right ventricular function in BGN$^{f/0(f/f)}$ and BGN^{ECKO} mice with AAV-NC or AAV-TXN treatment (n=6), including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; IgG, immunoglobulin G; LV, lentivirus; WT, wild-type; Mut, mutation; CHX, cycloheximide; FRET, fluorescence resonance energy transfer; CFP, cyan fluorescent protein; YFP, yellow fluorescent protein; PAT/PET, pulmonary artery acceleration time/pulmonary

1047 artery ejection time ratio; TAPSE, tricuspid annular plane systolic excursion; RVFAC,

1048 right ventricular fractional area change; and RVFWT, right ventricular free wall

1049 thickness.

1050

1051 **Figure S6. HIF signaling mediated TXNDC5-driven PH.**

A and **B**, qPCR analysis showing the effects of HIF-1α (A) or HIF-2α (B) silencing on the mRNA expression of TXNDC5. Average expression values are normalized to the β-actin mRNA level. **C**, Western blot showing that siHIF-2α reduced the protein levels of TXNDC5 in hPAECs under hypoxic conditions (n=6). **D**, Schematic illustration showing 2 sites in HIF-2α predicted to bind to the TXNDC5 promoter region. **E**, Chromatin immunoprecipitation (ChIP) followed by polymerase chain reaction analysis showing that the BS1 and BS2 regions are both for HIF-2α binding. **F**, Western blot analysis showing that silencing of TXNDC5 reduced the protein levels of HIF-1α and HIF-2α in hPAECs under hypoxic conditions (n=6). **G**, Western blots showing that siTXNDC5 reduced the protein levels of HIF-1α and HIF-2α in the nucleus exposed to hypoxia in hPAECs (n=6). **H**, Immunofluorescent staining analysis showing that siTXNDC5 reversed the increased levels of HIF-1α (green, left) or HIF-2α (green, right) induced by hypoxia in hPAECs. Nuclei were counterstained with DAPI (blue). Scale bars=50 μm. **I**. Schematic diagram of the pharmacological rescue experiment. AAV-NC or AAV-TXN was delivered via the tail vein to overexpress TXNDC5 in lung endothelial cells of mice, followed by normoxia or SuHx exposure, and then with HIF-1α inhibitor (PX-478) or HIF-2α inhibitor (PT-2385), or vehicle treatment. **J**. Hemodynamic measurements showing right ventricular systolic pressure (RVSP) in mice treated as described in (I) (n=6). **K**, Fulton index [RV/(LV+S)] of mice, assessing right ventricular hypertrophy (n=6). **L**, **M**, **N,** and **O**, Echocardiographic evaluation of right ventricular function, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**G**) or 2-way ANOVA (**A-C**, **F,** and **J-O**), followed by Tukey honestly significant difference test for multiple comparisons. BS indicates binding site; Nor, normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypixa; ChIP, chromatin immune precipitation; PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio; TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional area change; and RVFWT, right ventricular free wall thickness.

1080

Figure S7. TXNDC5 inhibited the ubiquitination of HIF-1α/2α in a BGN-dependent manner.

A, Representative Western blots and quantification of BGN, and HIF-1/2α in hPAECs treated with siNC or siBGN under normoxia or hypoxia conditions (n=6). **B**, Western

blot showing that the LV-TXNDC5-induced upregulation of HIF-1/2α in hPAECs both under normoxic and hypoxic situations (n=6). **C**, Representative Western blots and quantification of TXNDC5, HIF-1α, and HIF-2α in the lung ECs isolated from AAV-TXN or AAV-NC treated mice exposed to SuHx or under normoxic conditions (n=6). **D**, Quantification analysis showing that the protein half-life of endogenous HIF-1/2α was prolonged by LV-TXNDC5 treatment in hPAECs under normoxic conditions (n=5). **E**, Western blot analysis indicated that the HIF-1α ubiquitylation profile was reduced by LV-TXNDC5 but increased by siBGN treatment. hPAECs were treated with 10 μM MG132 for 6 h before harvest. Data represent the mean±SD. Comparisons of parameters were performed with the 2-way ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; Ub, ubiquitin; IgG, immunoglobulin G; AAV, adeno-associated viruses; LV, lentivirus; and CHX, cycloheximide.

Figure S8. TXNDC5 inhibited the ubiquitination of HIF-1α/2α in a BGN-dependent manner.

A, Representative H&E staining of major kidney, liver, and spleen from rats maintained under normoxia for 5 weeks and continuously treated with E64FC26 or vehicle,

showing preserved tissue morphology without apparent injury or inflammation. Scale bars=100 μm. **B**, Gain of body weight of female (top panel) and male (bottom panel) rats maintained under normoxia for 5 weeks with E64FC26 or vehicle treatment (n=6). **C**, **D**, and **E**, Quantification of distal pulmonary arteries (PAs) remodeling in SuHx-induced PH rats treated with vehicle or E64FC26, including distal PA muscularization, medial wall thickness, and collagen deposition area (n=6). **F**, Echocardiographic assessment of right ventricular function in rats with vehicle or E64FC26 treatment, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). **G** and **H**, Quantification of distal PA medial wall thickness and collagen deposition area SuHx-treated TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice with or without E64FC26 administration under SuHx or normoxia conditions (n=6). **I**, Right ventricular systolic pressure (RVSP) in TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with vehicle or E64FC26 (n=6). **J**, Fulton index [RV/(LV+S)] of TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice with vehicle or E64FC26 treatment (n=6). **K**, Echocardiographic assessment of right ventricular function in TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with vehicle or E64FC26 exposed to SuHx or normoxia, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). **L**, Western blot showing efficient knockdown of TXNDC5 in lung ECs isolated from rats receiving pulmonary endothelium-targeted self-complementary adeno-associated virus (scAAV) carrying shRNA against TXNDC5 (shTXN) compared with control shRNA (shNC) (n=6). **M**, **N**, and **O**, Quantification of distal PA remodeling in SuHx-induced PH rats treated with shNC or shTXN, including distal PA muscularization, medial wall thickness, and perivascular collagen deposition area (n=6). **P**, Fulton index [RV/(LV+S)] in rats treated with shNC or shTXN under normoxia or SuHx conditions (n=6). **Q**, Echocardiographic assessment of right ventricular function in rats receiving shNC or shTXN, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**L**) or 2-way ANOVA (**B-K**, and **M-Q**), followed by Tukey honestly significant difference test for multiple comparisons. H&E indicates hematoxylin-eosin; f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; shRNA, short hairpin RNA; PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio;

1133　TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional

1134　area change; and RVFWT, right ventricular free wall thickness.