TXNDC5 Governs Extracellular Matrix Homeostasis in Pulmonary Hypertension

Zhen Chen[1][*]; Fanhao Kong[1][*]; Lixin Huang[1][*]; Wenze Wu[1][*]; Ziping Wang[1]; Hanbin Chen[1]; Junting Zhang[1]; Lin Deng[6]; Chengrui Cao[1]; Xiaoyan Zhang[1]; Zixin Liu[2]; Kai-Chien Yang[3][#]; Wei-Ting Chang[4, 5][#]; Jin-Song Bian[1][#]; Xiaowei Nie[2][#];

Department of Pharmacology, Joint Laboratory of Guangdong-Hong Kong Universities for Vascular Homeostasis and Diseases, SUSTech Homeostatic Medicine Institute, School of Medicine, Southern University of Science and Technology, Shenzhen 518055, China.

Department of Human Cell Biology and Genetics, SUSTech Homeostatic Medicine Institute, School of Medicine, Southern University of Science and Technology, Shenzhen 518055, China.

Department and Graduate Institute of Pharmacology, National Taiwan University College of Medicine, Taipei, Taiwan

School of Medicine, College of Medicine, National Sun Yat-sen University, Kaohsiung, Taiwan

Division of Cardiology, Department of Internal Medicine, Chi Mei Medical Center, Tainan

Department of Cardiology, The Eighth Affiliated Hospital, Sun Yat-sen University, Shenzhen, China

[*]These authors contributed equally to this work.

[#]Correspondence to:

Xiaowei Nie, PhD, E-mail niexw@sustech.edu.cn.

Jin-Song Bian, PhD, E-mail bianjs@sustech.edu.cn.

Wei-Ting Chang, PhD, E-mail cmcvecho2@gmail.com.

Kai-Chien Yang, PhD, E-mail allenmy0920@gmail.com.

ABSTRACT

BACKGROUND: Pulmonary hypertension (PH) is characterized by vascular remodeling without effective treatments. The initiation and progression of PH are heavily influenced by the dysfunction of pulmonary artery endothelial cells. Thioredoxin domain containing 5 (TXNDC5), a member of the protein disulfide isomerases (PDI) family, catalyzes the formation and rearrangement of disulfide bonds in proteins, crucial for endothelial cell function and maintaining vascular homeostasis in cardiovascular diseases, yet its role in PH remains unclear.

METHODS: Label-free proteomics was used to reveal the PDI protein expression profile in hypoxic PH patients. Multiplex immunofluorescence and Western blot analyses were performed to investigate the pathological role of TXNDC5. Mice with adeno-associated virus-mediated lung endothelial TXNDC5 overexpression, global knockout, and endothelial ablation of TXNDC5 were used to study the role of TXNDC5 in rodent PH models. *In vitro* experiments with cultured pulmonary arterial endothelial cells and bioinformatics assays were used to elucidate the underlying mechanisms. Pharmacological inhibition and endothelial-targeted TXNDC5 gene therapy were evaluated for their effects in PH.

RESULTS: TXNDC5 was significantly upregulated in the lungs of both PH patients and rodent models, with prominent expression in the endothelial layer of remodeled distal pulmonary arteries. Overexpression of TXNDC5 exacerbated Sugen5416/hypoxia (SuHx)-induced pulmonary artery remodeling, increased right ventricular systolic pressure, and right ventricular hypertrophy. Global or endothelial-specific deficiency of TXNDC5 exerted protective efficacy against the SuHx challenge. RNA sequencing and protein-protein interaction analysis revealed TXNDC5 as a potential regulator of extracellular matrix (ECM) homeostasis, with biglycan (BGN) identified as a critical downstream effector. TXNDC5 required BGN to regulate the ubiquitination of HIF-1α/2α, resulting in its abnormal accumulation to drive PH development. E64FC26, a TXNDC5 inhibitor, and endothelial-targeted TXNDC5 gene therapy exhibited therapeutic efficacy in SuHx-induced PH rats.

CONCLUSIONS: Our study reveals that TXNDC5 is a main modulator to regulate

ECM homeostasis and may serve as a promising target for the treatment of PH.

Key Words: pulmonary hypertension, extracellular matrix, TXNDC5, BGN, HIF

signaling

INTRODUCTION

Pulmonary hypertension (PH) is a serious cardiovascular disease characterized by

elevated pulmonary artery pressure and right heart failure, with high morbidity and

mortality[1]. Hypoxia is a significant trigger for PH, with hypoxia-induced and/or lung

disease-related PH (Group III) being one of the most common types clinically[2].

Hypoxia-induced dysfunction in pulmonary artery endothelial cells (PAECs) initiates

pulmonary vasoconstriction[3]. Subsequently, pulmonary artery smooth muscle cells

(PASMCs) switch to a hyperproliferative and apoptosis-resistant phenotype, leading to

vascular remodeling[4]. However, there remains no curative clinical treatment option for

reversing remodeled vessels against the disease[5,6]. It highlights the need for further

exploration of PH pathogenesis and new therapeutic targets.

Protein disulfide isomerase (PDI) is a family of enzymes located in the

endoplasmic reticulum that facilitates the disulfide bond formation and rearrangement,

aiding protein proper folding[7,8]. This family comprises 21 members, each containing

thioredoxin-like domains involved in redox reactions, particularly under hypoxic

conditions[9]. PDI proteins are implicated in the modulation of endothelial cell function

to maintain vascular homeostasis. PDIA1 regulated the expression and redox state of

hypoxia-inducible factor (HIF)-1α by promoting its lysosomal degradation by

influencing the formation of disulfide bonds[10]. It also enhances vascular endothelial

growth factor receptor 2 (VEGFR2) signaling, thereby promoting endothelial cell

migration, proliferation, and angiogenic responses[11]. As a member of the PDI family,

TXNDC5 is mainly expressed in endothelial cells (ECs) and is also known as

endothelial PDI (Endo-PDI). Its abnormal expression is associated with multiple

disorders such as organ fibrosis[12], diabetes[13], rheumatoid arthritis[14], and cancers[15].

Additionally, inhibition of TXNDC5 restores endothelial dysfunction to mitigate

vascular calcification and atherosclerosis[16,17]. However, the role of PDI proteins in

pulmonary vascular remodeling in PH remains unclear.

Extracellular matrix (ECM) is a complex network of proteins and polysaccharides

secreted by cells, providing structural and communicative support to adjacent cells[18].

Remodeling of pulmonary arterial ECM occurs in the early stage of PH pathogenesis,

before increased wall thickness and pulmonary artery pressure. ECM remodeling is the

cause rather than a consequence of pulmonary vascular remodeling in PH[19]. Recently,

we revealed that the ECM proteins in PAECs accelerated pulmonary vascular

remodeling through interaction with HIF-1α in a positive feedback loop manner,

providing evidence for early intervention in ECM remodeling to prevent PH

progression[20,21].

In this study, we performed 4D label-free proteomics to reveal the expression

profile of PDI proteins from the lungs of patients with hypoxic PH. We found that

TXNDC5 was one of the most significantly upregulated proteins. Global or endothelial

cell-specific deficiency of TXNDC5 alleviated Sugen5416/hypoxia (SuHx)-induced

PH, whereas TXNDC5 overexpression aggravated PH. Importantly, the silencing of

TXNDC5 preserved ECM homeostasis to alleviate PA remodeling and restore

endothelial function. In addition, TXNDC5 directly interacted with biglycan (BGN)

and assisted its protein folding to drive PH development. Furthermore, TXNDC5

required BGN to regulate the HIF-1α/2α ubiquitination, resulting in its abnormal

aggregation. Importantly, pharmacological inhibition of TXNDC5 with E64FC26 or

pulmonary endothelium-targeted TXNDC5 silencing effectively mitigated PH in rats.

The present study therefore demonstrated for the first time that TXNDC5 may be a key

protein to control ECM homeostasis and a promising therapeutic target for PH.

METHODS

Detailed expanded methods and materials are available in **Supplemental Material 1**.

Study Approval

All human samples were obtained from the Shenzhen People's Hospital (China) and Wuxi People's Hospital (China) with the informed consent of the patients, which was approved by the Ethics Committee for the Use of Human Subjects following The Code of Ethics of the Helsinki Declaration of the World Medical Association. All animal studies followed the guidelines of the Committee on Animal Research and Ethics of the Southern University of Science and Technology (China).

Data Availability Statement

Source data and original blots or gel images for this study are available from the corresponding author upon reasonable request.

Statistical Analysis

Data were analyzed with GraphPad Prism 8.0 software and were expressed as the mean±standard deviation (mean±SD). For comparisons between two groups with a normal distribution, an unpaired 2-tailed Student *t*-test was performed. When the data did not follow a normal distribution, the non-parametric Mann-Whitney U test was applied. For data involving more than two groups, one-way ANOVA was used, followed by Tukey's honestly significant difference test for multiple comparisons. For two variables, two-way ANOVA with Tukey's honestly significant difference test for multiple comparisons was applied. Statistical significance was defined as $P<0.05$.

RESULTS

TXNDC5 Expression is Increased in PH Patients and Rodent Models

We performed 4D label-free proteomics on lung samples from 5 patients with hypoxic PH and 5 non-PH donors (**Figure S1A**). Among the human PDI family proteins, TXNDC5 was the most markedly dysregulated member in PH patients compared with donors (fold change=3.45, *P*=0.0048) (**Figure 1A**). The involvement of TXNDC5 in PH was then confirmed by assessing its protein and mRNA expressions in the lungs of

PH patients as compared with non-PH donors (**Figure 1B-C**). Western blotting further showed that TXNDC5 expression was also significantly increased in multiple rat PH models and mouse PH models compared with controls (**Figure 1D** and **Figure S1B**). In distal pulmonary arteries (PAs), multiplex immunofluorescence (mIF) assays demonstrated pronounced co-localization of TXNDC5 with the endothelial marker CD31. By contrast, co-localization of TXNDC5 with α-SMA (a marker for smooth muscle cells) and COL1A1 (a marker for fibroblasts) was marginal (**Figure 1E-G** and **Figure S1C**). Furthermore, TXNDC5 expression was significantly elevated in the CD31$^+$ intima of severely remodeled PAs from patients with interstitial lung disease (ILD)-PH and chronic obstructive pulmonary disease (COPD)-PH patients, compared with non-PH donors (**Figure 1H**). Similar results were observed in the remodeled PAs of SuHx- or monocrotaline (MCT)-induced PH rat models, indicating that TXNDC5 upregulation is a conserved feature of pulmonary artery (PA) remodeling (**Figure 1I-K** and **Figure S1D**). Moreover, Western blot analysis of primary lung ECs isolated from PH rodent models confirmed the TXNDC5 upregulation (**Figure S1E-F**). Notably, correlations between intimal TXNDC5 mean fluorescence intensity (MFI) and PH severity were observed in rats, with positive correlations with right ventricular systolic pressure (RVSP) and Fulton index (right ventricle weight to left ventricle plus interventricular septum weight, ie, RV/(LV+S)), and a negative correlation with tricuspid annular plane systolic excursion (TAPSE) (**Figure 1L-N**). Taken together, these findings suggest that the TXNDC5 may be a critical dysregulated factor in ECs within the pulmonary vasculature that contributes to PH pathogenesis.

TXNDC5 Mediates the Development of SuHx-Induced PH

To investigate the role of TXNDC5 *in vivo* in the development of experimental PH, we created TXNDC5 knockout (TXNDC5$^{-/-}$) mice, which were then challenged with or without SuHx (**Figure 2A**). We found that deletion of TXNDC5 alone had no significant effects but markedly reduced both RVSP (**Figure 2B**) and RV/(LV+S) (**Figure S2A**) in male mice. Echocardiography analysis also showed that TXNDC5 deficiency significantly improved the right ventricular (RV) function, as indicated by

increased PA acceleration time/ PA ejection time ratio (PAT/PET), TAPSE, and right

ventricular fractional area change (RVFAC) (**Figure S2B-D**), along with reduced right

ventricular free wall thickness (RVFWT) (**Figure S2E**). In addition, using mIF and

Victoria blue/Sirius red (VB/SR) staining, we evaluated distal PA remodeling in mice

(**Figure 2C**). Under normal conditions, no significant abnormal histological change

was found in TXNDC5$^{-/-}$ mice compared with wild-type (WT) littermates. Notably,

TXNDC5 deficiency markedly reduced PA muscularization, decreased medial wall

thickness, and diminished collagen deposition area compared with WT littermates

(**Figure 2D-F**). Consistent with the PA changes, wheat germ agglutinin (WGA) staining

also proved that the RV cardiomyocyte hypertrophy was markedly inhibited in SuHx-

treated TXNDC5$^{-/-}$ mice when compared with their control littermates (**Figure 2G**).

Notably, these protective effects of TXNDC5 deficiency were also confirmed in female

mice. In SuHx-treated female TXNDC5$^{-/-}$ mice, we observed similar reductions in

RVSP and RV/(LV+S) (**Figure S2F-G**), improvements in echocardiographic

parameters (**Figure S2H-K**), as well as attenuated distal PA remodeling compared with

female WT controls (**Figure S2L-O**). These data indicate that TXNDC5 deletion

protects against SuHx-induced PH.

With a gain-of-function approach, we next examined whether overexpression of

TXNDC5 can influence the development of PH in SuHx-treated mice. TXNDC5 was

exogenously delivered via tail vein injection using an adeno-associated virus serotype

engineered to specifically target lung endothelial cells (AAV-TXN) (**Figure 2H**). *Ex*

vivo imaging of various organs confirmed that preferential accumulation of AAV-TXN

in the lungs (**Figure 2I**). Consistently, co-immunofluorescence staining further

demonstrated strong colocalization of GFP-TXNDC5 with anti-CD31 in pulmonary

arteries (**Figure S2P**). Furthermore, Western blot analysis of primary lung ECs further

verified the EC-specific TXNDC5 overexpression (**Figure S2Q**). Hemodynamic

measurement revealed that both male and female mice administered with AAV-TXN

increased the RVSP compared with those administered with control AAV (AAV-NC)

even under normoxic conditions. In response to the SuHx challenge, AAV-TXN mice

further increased in both RVSP and RV/(LV+S) compared with AAV-NC mice (**Figure**

2J and **Figure S2R**). Echocardiography analysis also showed that AAV-TXN reduced PAT/PET, TAPSE RVFAC, and increased RVFWT compared with AAV-NC mice under SuHx, although no significant differences were observed under normoxic conditions (**Figure S2S-V**). In addition, mIF and VB/SR staining confirmed exacerbated remodeling of distal PAs in AAV-TXN mice compared with AAV-NC mice upon SuHx treatment, as indicated by increased PA muscularization, medial wall thickness, and collagen deposition area (**Figure 2K-N**). Conversely, endothelial TXNDC5 overexpression was found to exacerbate SuHx-induced PH.

Endothelial Cell-Specific Deletion of TXNDC5 Ameliorates SuHx-Induced PH

To examine the role of endothelial TXNDC5 in the development of PH, we generated endothelial cell-specific TXNDC5 knockout mice (TXNDC5^{ECKO}) (**Figure 3A** and **Figure S3A**) and validated them with Western blots (**Figure S3B**). SuHx-induced RVSP and RV/(LV+S) elevation (**Figure 3B-C**), as well as impaired RV function (**Figure S3C-F**), were significantly improved by the deletion of endothelial TXNDC5. The morphology of PAs in normoxic TXNDC5^{ECKO} mice did not differ from that of control littermates (TXNDC5$^{f/f}$ mice) under normoxia. In line with the hemodynamic and echocardiographic changes, TXNDC5^{ECKO} mice exhibited reduced muscularized PAs, medial wall thickness, and collagen deposition induced by SuHx compared with TXNDC5$^{f/f}$ mice (**Figure 3D-G**). Using a standardized lung angiography, we confirmed that SuHx treatment led to distal PA remodeling, manifested as restricted intrapulmonary contrast agent distribution in TXNDC5$^{f/f}$ mice. In contrast, TXNDC5^{ECKO} mice displayed markedly improved angiographic filling, indicative of better-preserved pulmonary vasculature (**Figure 3H** and **Figure S3G**). In addition to vascular changes, SuHx-induced RV hypertrophy was also markedly attenuated in TXNDC5^{ECKO} mice, as evidenced by reduced RV wall thickness and decreased cardiomyocyte cross-sectional area (**Figure 3I-J**). Importantly, comparable protective effects were also observed in female TXNDC5^{ECKO} mice, as evidenced by reduced RVSP and RV hypertrophy (**Figure S3H-I**), improved RV function on echocardiography (**Figure S3J-M**), and attenuated distal PA remodeling (**Figure S3N-**

Q). Together, these findings demonstrate that endothelial cell-specific deletion of TXNDC5 ameliorates SuHx-induced PH in mice, with consistent benefits across sexes. These data underscore the crucial role of endothelial TXNDC5 in the development of PH and highlight its potential as a sex-independent therapeutic target.

TXNDC5 Promotes Hypoxia-Induced Dysfunction in Pulmonary Arterial Cells

Exposure to hypoxia significantly increased the protein and mRNA levels of TXNDC5 in hPAECs (**Figure S4A-B**). Immunofluorescent staining further confirmed that TXNDC5 expression was upregulated upon hypoxia treatment and was mainly located in the cytoplasm (**Figure S4C**). Targeting TXNDC5 small-interfering RNA oligonucleotides (siTXNDC5) was used to silence TXNDC5 expression in hPAECs. EdU staining showed that knocking down TXNDC5 expression suppressed the proliferation of hPAECs under both normoxic and hypoxic conditions (**Figure S4D**). In addition, siTXNDC5 treatment also abated the hypoxia-induced migration ability of hPAECs (**Figure S4E**). To further evaluate the role of increased TXNDC5 in hPAECs, we overexpressed TXNDC5 in hPAECs by transducing with lentivirus (LV-TXNDC5). TXNDC5 overexpression visibly altered cytoskeletal architecture, which displayed prominent stress fibers and a more diffuse pattern of paxillin distribution (**Figure S4F-G**). Furthermore, flow cytometry showed that the silencing of TXNDC5 prevented the G1 to S phase transition (**Figure S4H**) of hPAECs under both normoxic and hypoxic conditions. Taken together, TXNDC5 regulates many important functional abnormalities associated with PH pathogenesis, with disruption of cytoskeletal architecture in hPAECs.

The hyper-proliferation and resistance to apoptosis of PASMCs directly accelerate the remodeling of PAs in PH[22]. Thus, we also investigated whether endothelial TXNDC5 affects the function of hPASMCs via influencing paracrine patterns. For this purpose, hPAECs were transfected with siTXNDC5 under hypoxia or normoxia conditions, and their conditioned medium (siTXN-CM) was collected to treat hPASMCs. Assessment of cell proliferation showed that siTXN-CM significantly downregulated cyclin A and proliferating cell nuclear antigen (PCNA) expression under

both normoxic and hypoxic conditions (**Figure S4I**). Consistently, EdU incorporation

assays further confirmed that siTXN-CM suppressed the proliferation of hPASMCs

(**Figure S5A**). Cell Counting Kit-8 (CCK-8) assays further demonstrated that siTXN-

CM decreased the viability of hPASMCs (**Figure S5B**). In addition, siTXN-CM

enhanced hPASMCs apoptosis, as indicated by increased cleaved caspase-3 expression

(**Figure S4J**). Collectively, our data demonstrate that hPAECs TXNDC5 plays a critical

role in promoting hPASMCs proliferation, survival, and resistance to apoptosis.

TXNDC5 Regulates ECM Expression and Homeostasis

To explore the underlying downstream molecular mechanisms of TXNDC5, we

performed a bulk RNA sequencing (RNA-seq) analysis on hPAECs transfected with

siRNA targeting TXNDC5 under hypoxia exposure. Gene ontology analysis indicated

that the silencing of TXNDC5 resulted in altered expression of genes mainly enriched

in "collagen-containing ECM", "ECM", "ECM organization", "ECM structural

constituent", "response to hypoxia", and "cellular response to hypoxia" (**Figure 4A**).

We selected the differentially expressed ECM genes and then constructed a protein-

protein interaction (PPI) network (**Figure 4B**). The network revealed that TXNDC5-

affected ECM proteins were clustered into different functional subgroups, including

"proteoglycans and glycoproteins", "collagens", "matrix modulators and adhesion",

"metallopeptidase activity", "basement membrane", "growth factors and signaling

regulators", and "angiogenesis". Immunofluorescence staining confirmed that periostin

(POSTN), biglycan (BGN), lysyl oxidase (LOX), and transglutaminase 2 (TGM2) were

excessively deposited in the PAs from TXNDC5$^{f/f}$ mice but not in TXNDC5^{ECKO} mice

under SuHx conditions (**Figure 4C-F** and **Figure S5C**). Western blot analysis also

confirmed that TXNDC5 silencing reversed the abnormal expression of ECM proteins

(COL1A1, TGM2, FN1, POSTN, LOX) induced by hypoxia treatment in hPAECs

(**Figure 4G**).

To further investigate whether TXNDC5 regulates ECM secretion, we collected

the supernatants from hPAECs. Dot blot analysis revealed that the silence of TXNDC5

markedly reduced the secretion of BGN, POSTN, and LOX (**Figure S5D-G**). These

results provide a mechanistic explanation for why the conditional medium from TXNDC5-silenced hPAECs suppresses proliferation and viability while promoting apoptosis of hPASMCs. Collectively, these data highlight the key role TXNDC5 played in regulating ECM homeostasis.

BGN Is a Promising Biomarker and Therapeutic Target for Hypoxic Pulmonary Hypertension

We further explore the key TXNDC5-targeted ECM proteins that are responsible for the biological function of ECs. We performed a computational analysis of the PPI network between the ECM proteins dataset and the ECs function-related proteins dataset. By combining these two datasets, we constructed a huge original PPI network. A core network was finally obtained by progressively simplifying through degree values (**Figure 5A**). Through the intersection of 3 ECM datasets, core ECM in the above PPI network, TXNDC5 silencing in hPAECs, and hypoxia-treated hPAECs (GSE160255), we finally identified biglycan (BGN) (**Figure S5H**).

We first examined the level of BGN in lung biopsies from patients with hypoxic PH or healthy donors. Protein levels of BGN were increased in the lungs of PH patients as compared with non-PH donors (**Figure S5I**). A recently published GEO dataset (GSE254617) also showed markedly upregulated mRNA levels of BGN in the lungs of patients with idiopathic pulmonary arterial hypertension (IPAH) (**Figure 5B**). Consistent with human specimens, protein levels of BGN were also increased in rodent PH models, including SuHx-treated mice and rats and MCT-treated rats, as compared with controls (**Figure S5J-K**). We also investigated whether BGN can reflect the severity of PH by analyzing BGN levels in serum. Surprisingly, a markedly increased BGN level was found in the serum of 27 patients with hypoxic PH compared with the serum of 27 non-PH donors (**Figure 5C**). Additionally, BGN expression in serum exhibited a significant correlation with systolic pulmonary artery pressure (sPAP) (**Figure 5D**), suggesting that BGN may serve as a biomarker associated with PH. As a secreted protein, we further investigated the biological function of circulating BGN *in vitro*. EdU staining showed that treatment with recombinant human BGN (rhBGN)

significantly enhanced the proliferative capacity of hPASMCs compared with the negative control (bovine serum albumin, BSA) (**Figure S5L**). Consistently, CCK-8 assays demonstrated increased cell viability following rhBGN treatment (**Figure S5M**). In parallel, rhBGN induced an apoptosis-resistant phenotype in hPASMCs, as evidenced by reduced cleaved caspase-3 expression (**Figure S5N**) and decreased apoptotic cell populations detected by flow cytometry (**Figure S5O**).

To evaluate the specific role of BGN in PH generation, we then developed an endothelial cell-specific BGN knockout mouse (BGN^{ECKO}) (**Figure S6A**), and the absence of BGN in lung ECs was validated (**Figure S6B**). As BGN is an X chromosome-located gene, male mice carry a single floxed allele (BGN$^{f/0}$), whereas female mice carry double floxed alleles (BGN$^{f/f}$). Firstly, RVSP and RV/(LV+S) were elevated in male mice with SuHx treatment, while endothelial cell-specific BGN ablation ameliorated these compared with their control littermates (BGN$^{f/0}$) (**Figure 5E-F**). Consistent with changes in hemodynamics, BGN^{ECKO} mice showed abrogated PAs remodeling, as shown by the mIF and VB/SR staining (**Figure 5G-I** and **Figure S6C**). Lung angiogram also revealed the better-preserved lung vasculature in BGN^{ECKO} mice, further indicating that endothelial BGN deletion effectively protects against distal PA remodeling (**Figure 5J**). Histological assessment of the RV also revealed an increase in RV wall thickness in the SuHx-BGN$^{f/0}$ group, which was significantly reduced in BGN^{ECKO} mice treated with SuHx (**Figure 5K**). Moreover, echocardiography analysis also showed that SuHx-induced reductions in PAT/PET, TAPSE, RVFAC, and increases in RVFWT were greater in BGN$^{f/0}$ mice than in BGN^{ECKO} mice (**Figure S6D-G**). Notably, similar protective effects were observed in female BGN^{ECKO} mice, with comparable improvements in RVSP (**Figure S6H**), RV hypertrophy and function (**Figure S6I-M**), and distal PA pathological remodeling compared with BGN$^{f/f}$ female littermates (**Figure S6N-Q**). Together, our data suggest that endothelial BGN plays an important role in PH development.

TXNDC5 Assists BGN Protein Folding to Drive Pulmonary Hypertension

To elucidate how TXNDC5 regulates BGN, we first performed Western blots to

validate the positive regulation of TXNDC5 on BGN both *in vitro* and *in vivo*. The

silencing of TXNDC5 reversed the overexpression of BGN induced by hypoxia

stimulation in hPAECs (**Figure S7A**). Reduced BGN expression was also observed in

the lungs of TXNDC5-deficient mice (**Figure S7B**), which was consistent with the

findings in primary lung ECs isolated from TXNDC5[ECKO] mice (**Figure S7C**). In

contrast, BGN expression was elevated in hPAECs following lentiviral-mediated

TXNDC5 overexpression (**Figure S7D**) and in primary lung ECs isolated from AAV-

TXN-treated mice (**Figure S7E**).

To further assess the direct interaction between TXNDC5 and BGN, we

investigated the subcellular location at which TXNDC5 binds to BGN. Immunostaining

assays showed that TXNDC5 colocalized with BGN (**Figure 6A**). We also performed

co-IP analysis and found that TXNDC5 bound to BGN, and the binding was enhanced

upon hypoxia exposure (**Figure 6B** and **Figure S7F**). To identify the protein domain of

TXNDC5 that was involved in the interaction with BGN, we performed molecular

docking analysis and revealed that multiple sites of TXNDC5 (Cys89, Ser125, Gly351,

and His352) might be involved in the interaction with BGN (**Figure 6C**). Thus, we

constructed four TXNDC5 mutants (C89A, S125A, G351A, and H352A) to identify

the residue that plays a key role in this process. Co-IP analysis showed that C89A and

G351A could partially abolish TXNDC5 binding with BGN (**Figure 6D**), suggesting

that these amino acid residues might be essential for the binding of TXNDC5 and BGN.

As an endoplasmic reticulum resident protein with 3 thioredoxin domains,

TXNDC5 catalyzes the formation or rearrangement of substrate disulfide bonds. Each

thioredoxin domain contains a highly conserved CGHC motif, which serves as the

active site for PDI activity[15]. Thus, we hypothesized that TXNDC5 functions to

facilitate the folding of BGN, allowing it to mature and subsequently exert signaling

transduction and travel through the secretory pathway. To confirm the requirement for

the PDI activity of TXNDC5 in assisting BGN folding, we generated a TXNDC5

mutant (TXNDC5 Mut) that lacks PDI activity by introducing cysteine-to-alanine

mutations in each of its 3 thioredoxin domains (**Figure S7G**). As described previously[23],

the extent of BGN protein folding can be examined using a fluorescence resonance

energy transfer (FRET)-based protein folding assay (**Figure S7H**). Application of a standard acceptor photobleaching FRET protocol in hPAECs revealed that LV-TXNDC5 significantly increased FRET efficiency while TXNDC5 Mut overexpression did not, suggesting that TXNDC5 assists BGN protein folding dependent on its PDI activity in hPAECs (**Figure 6E**). Moreover, a cycloheximide (CHX) protein chase assay was conducted to evaluate the stability of BGN in hPAECs. Overexpression of TXNDC5 markedly prolonged the half-life of BGN protein compared with LV-NC (**Figure S7I**), whereas TXNDC5 silencing accelerated BGN degradation, exerting the opposite effect (**Figure S7J**). Taken together, these results indicate that TXNDC5 depends on its PDI activity to contribute to the proper folding of BGN, maintaining its protein stability and subsequently secreting extracellular from hPAECs.

To assess whether BGN mediates the pathogenic effects of endothelial TXNDC5 *in vivo*, we performed a rescue experiment in endothelial BGN-deficient mice by specifically overexpressing TXNDC5 in lung ECs (**Figure 6F**). As shown by mIF and VB/SR staining, AAV-TXN exacerbated distal PA muscularization, medial wall thickness, and collagen deposition in $BGN^{f/f(f/0)}$ mice under SuHx conditions, whereas these effects were largely abolished in BGN^{ECKO} mice (**Figure 6G-J**). Consistently, AAV-TXN markedly increased RVSP (**Figure 6K**) and also aggravated RV hypertrophy in $BGN^{f/f(f/0)}$ mice, whereas both changes were significantly attenuated in BGN^{ECKO} mice (**Figure 6L**). Furthermore, echocardiographic assessment revealed that AAV-TXN led to pronounced deterioration of RV performance, as reflected by reduced PAT/PET, TAPSE, RVFAC, and elevated RVFWT. In contrast, these adverse functional changes were largely blunted in BGN^{ECKO} mice (**Figure S7K-N**). Together, these results indicate that endothelial BGN is required for TXNDC5-mediated pulmonary vascular remodeling and PH development, confirming BGN as a key downstream effector of TXNDC5.

A Positive Feedback Loop Exists Between TXNDC5 and HIF Signaling

Given that the HIF α subunits are the dominant effector activated by hypoxia, we hypothesize that HIF signaling regulates the transcription of TXNDC5. qPCR analysis

showed that silencing of HIF-2α, but not HIF-1α, markedly downregulated TXNDC5

mRNA levels under both normoxia and hypoxia conditions (**Figure S8A-B**). Western

blot analysis also demonstrated that the protein level of TXNDC5 was significantly

decreased when HIF-2α was knocked down (**Figure S8C**). Next, we identified two

putative HIF-2α binding sites (BS) in the TXNDC5 promoter region by utilizing the

available transcriptional factors database (hTFtarget and AnimalTFDB 4.0) (**Figure**

S8D). To study how HIF-2α transcriptionally regulates TXNDC5, we performed a

chromatin immunoprecipitation (ChIP) assay to examine whether HIF-2α binds to these

BS. We designed primers that span the sequence from two BS to amplify from HIF-2α

ChIP samples. Gel electrophoresis results showed that both primers could be amplified,

indicating that HIF-2α binds to these regions of the TXNDC5 promoter (**Figure S8E**).

Dual-luciferase reporter assays further revealed that mutation of each putative HIF-2α

binding site within the TXNDC5 promoter individually reduced luciferase activity

compared with the wild-type construct. Mutation of both sites caused a further decrease

(**Figure 7A**). These results indicate that both BS1 and BS2 contribute synergistically to

HIF-2α-mediated transcriptional activation of TXNDC5.

Following validation, we found that the HIF-1α/2α protein levels were

significantly decreased by silencing of TXNDC5 under the hypoxic situation (**Figure**

S8F), and similar results were also observed in the lungs of TXNDC5-deficient mice

(**Figure 7B**). We also examined the nuclear localization of HIF-1α/2α. Western blot

analysis and immunofluorescence staining consistently confirmed that TXNDC5

silencing decreased HIF-1α/2α levels in the nucleus under hypoxia exposure (**Figure**

S8G-H). Together, these results suggest that a positive feedback loop exists between

TXNDC5 and HIF signaling during hypoxia.

To further investigate whether HIF signaling is required for the pathogenic effects

of endothelial TXNDC5 *in vivo*, we performed a pharmacological rescue experiment

using selective inhibitors of HIF-1α (PX-478) or HIF-2α (PT-2385). Mice were

administered AAV-TXN to overexpress endothelial TXNDC5 and subsequently

subjected to normoxia or SuHx conditions, with or without HIF inhibitor treatment

(**Figure S8I**). As expected, AAV-TXN markedly exacerbated PH phenotypes, including

distal PA remodeling, increased RVSP, RV hypertrophy, and impaired RV function

under SuHx conditions. Importantly, treatment with HIF-1α or HIF-2α inhibitors

significantly mitigates these TXNDC5-induced vascular and cardiac abnormalities, as

evidenced by histological staining (**Figure 7C-F**), hemodynamic measurements

(**Figure S8J**), assessment of RV hypertrophy (**Figure S8K**), and echocardiographic

analysis (**Figure S8L-O**). Collectively, these results indicate that TXNDC5 drives PH

progression in a HIF signaling-dependent manner, establishing a functional TXNDC-

HIF positive feedback loop.

TXNDC5 Requires BGN to Inhibit the Ubiquitination of HIF-1α/2α

Interestingly, we found in the present study that the HIF-1α/2α protein levels were

significantly decreased by the silencing of BGN upon hypoxia exposure in hPAECs

(**Figure S9A**). Since TXNDC5 directly interacted with BGN, we hypothesized that the

regulation of HIF-1α/2α by TXNDC5 required the presence of BGN. Western blots

revealed that LV-TXNDC5 increased the expression of HIF-1α/2α both under normoxic

and hypoxic conditions in hPAECs (**Figure S9B**). More importantly, knockdown of

BGN abolished the upregulation of HIF-1α/2α induced by LV-TXNDC5 (**Figure 7G**).

These data suggest that TXNDC5 requires BGN to regulate the HIF signaling pathway.

Under normoxia, HIF-α is rapidly ubiquitinated by E3 ligase von Hippel-Lindau

(VHL) and leads to proteasomal degradation[24]. Interestingly, we found that TXNDC5

overexpression increased the HIF-1α/2α levels even under normoxic conditions, and

similar results were also observed in primary lung ECs isolated from AAV-TXN-treated

mice (**Figure S9C**). The protein half-life of HIF-1α/2α examined using CHX chase

assays showed that LV-TXNDC5-treated hPAECs displayed a prolonged half-life of

HIF-1α/2α as compared with control (**Figure 7H** and **Figure S9D**). We then further

determined whether HIF-1α/2α accumulation by TXNDC5 is VHL-dependent. It was

found that TXNDC5 silencing increased VHL expression, and simultaneous

knockdown of VHL and TXNDC5 significantly reversed the decreased HIF-1α/2α

protein levels caused by TXNDC5 silencing in hPAECs under hypoxic conditions

(**Figure 7I**). These data suggest that TXNDC5 modulates HIF-1α/2α to interfere with

VHL-mediated HIF-1α/2α degradation. To investigate whether TXNDC5 inhibits HIF-

1α/2α stability dependent on BGN, we performed *in vivo* ubiquitination assays. Results

showed that LV-TXNDC5 significantly suppressed the ubiquitination of HIF-1α/2α,

while siBGN increased it by upregulating the expression of VHL. More importantly,

the silencing of BGN eliminated the decrease in HIF-1α/2α ubiquitination and the

increase in VHL expression caused by TXNDC5 overexpression (**Figure 7J** and **Figure**

S9E). Taken together, our findings suggest that BGN is required for TXNDC5 to inhibit

the ubiquitination of HIF-1α/2α in a VHL-dependent manner.

E64FC26 Exhibits Therapeutic Effects Against PH in Rats

E64FC26 is a potent PDI pan-inhibitor, with IC$_{50}$ of 16.3 μM against TXNDC5[25].

To evaluate the toxicity and safety of E64FC26, we performed histological analyses on

the major organs of rats continuously treated with E64FC26 for 5 weeks under

normoxic conditions. In the kidneys, the renal tubules and glomeruli maintained their

integrity, with no signs of tubular dilation, epithelial cell injury, necrosis, or noticeable

inflammatory cell infiltration in the kidneys. Similarly, hepatocytes were well-

organized with clear cell boundaries, and there were no indications of necrosis, cell

swelling, or inflammatory cell infiltration in the liver. Histological examination of the

spleen revealed intact white and red pulp structures, without evidence of lymphoid

depletion, congestion, or inflammatory infiltration (**Figure S10A**). Furthermore, body

weight measurements showed no significant differences in both male and female rats

after E64FC26 treatment compared with the vehicle (**Figure S10B**). These findings

collectively indicate that E64FC26 exhibits a favorable safety profile without causing

overt organ or systemic toxicity.

Then, to examine whether pharmacological inhibition of TXNDC5 activity

produces a therapeutic effect on PH, E64FC26 was administered intraperitoneally in

SuHx-induced PH rats (**Figure 8A**). Hemodynamic measurements and morphometric

analysis demonstrated that E64FC26 administration attenuated the elevation of RVSP

(**Figure 8B**) and the increase in RV/(LV+S) ratio (**Figure 8C**) in SuHx-induced PH rats

compared with those with vehicle treatment. Histological analyses revealed that E64FC26 reversed SuHx-induced PA remodeling compared with vehicle-treated rats, as evidenced by reduced distal PA muscularization, decreased medial wall thickness, and attenuated collagen deposition area (**Figure 8D** and **Figure S10C-E**). Echocardiographic assessment further indicated that E64FC26 alleviated RV dysfunction in SuHx-induced PH rats, as reflected by increased PAT/PET and TAPSE, improved RVFAC, and reduced RVFWT (**Figure S10F**). Consistent with the echocardiographic improvements, histological analyses revealed that E64FC26 administration significantly reversed the RV hypertrophy (**Figure 8E**) and reduced the RV fibrosis area (**Figure 8F**) in SuHx-induced PH rats compared with those treated with vehicle.

To assess whether the therapeutic effects of E64FC26 in PH *in vivo* are primarily mediated through TXNDC5 rather than off-target inhibition of other PDI family members, we treated TXNDC5^{ECKO} mice with E64FC26 under SuHx conditions (**Figure 8G**). Notably, E64FC26 treatment also exhibited therapeutic effects in TXNDC5$^{f/f}$ mice against SuHx challenged, consistent with the results observed in rats. More importantly, in TXNDC5 endothelial-deficient mice, administration of E64FC26 did not confer additional protection against SuHx-induced distal PA remodeling, as indicated by unchanged medial wall thickness and collagen deposition area compared with vehicle-treated TXNDC5^{ECKO} mice (**Figure 8H** and **Figure S10G-H**). Similarly, lung angiography showed that E64FC26 administration failed to improve distal pulmonary perfusion in SuHx-treated TXNDC5^{ECKO} mice (**Figure 8I**). Consistently, E64FC26 also did not attenuate RVSP elevation (**Figure S10I**), RV hypertrophy (**Figure S10J**), and RV functional impairment (**Figure S10K**) in these mice. These findings indicated that the therapeutic effect of E64FC26 in PH would be largely dependent on the targeting of endothelial TXNDC5 rather than off-target inhibition of other PDI family members. In a word, these results indicate that pharmacological inhibition of TXNDC5 activity may be a promising strategy to alleviate the progression of PH.

Pulmonary Endothelium-Targeted TXNDC5 Gene Therapy Ameliorates PH

We further established a pulmonary endothelium-targeted gene therapy approach to enhance organ specificity. Using a self-complementary adeno-associated virus (scAAV) vector encoding shRNA against TXNDC5 (shTXN), we achieve rapid and endothelial-specific gene silencing *in vivo*. The vector was administered via the tail vein to SuHx-induced PH rats to selectively suppress TXNDC5 in lung ECs and evaluate its therapeutic efficacy (**Figure 8J**). *Ex vivo* imaging confirmed efficient lung targeting of the AAV (**Figure 8K**), and Western blotting demonstrated marked suppression of TXNDC5 expression in primary rat lung ECs (**Figure S10L**).

Treatment with shTXN significantly attenuated SuHx-induced PA remodeling compared with rats receiving the control vector (shNC), as evidenced by reduced distal PA muscularization, decreased medial wall thickness, and lower collagen deposition area (**Figure 8L** and **Figure S10M-O**). Hemodynamic measurements further demonstrated that shTXN administration attenuated the elevation of RVSP in SuHx-induced PH rats (**Figure 8M**). Moreover, morphological and histological analyses showed that shTXN decreased the RV/(LV+S) ratio (**Figure S10P**) and alleviated RV wall thickening (**Figure 8N**). Consistently, echocardiographic assessment revealed improved RV function, reflected by increased PAT/PET, TAPSE, RVFAC, and reduced RVFWT (**Figure S10Q**). These findings demonstrate that endothelial-targeted TXNDC5 silencing effectively mitigates pulmonary vascular remodeling and right heart dysfunction in PH. Collectively, these findings demonstrate that endothelial-targeted TXNDC5 silencing effectively mitigates pulmonary vascular remodeling and right heart dysfunction in PH, highlighting the therapeutic potential of TXNDC5 inhibition.

DISCUSSION

PDI family catalyzes the formation and rearrangement of disulfide bonds of proteins in the endoplasmic reticulum. They play crucial roles in protein folding and disulfide bond formation. In the present study, we utilized 4D label-free proteomics to

profile the PDI proteins in the lungs of patients with hypoxic PH, identifying TXNDC5

as the most significantly altered member in the PDI family. TXNDC5, dominantly

expressed in endothelial cells, is also known as endothelial PDI (Endo-PDI)[26]. We

found that the silencing of TXNDC5 inhibited the proliferation and migration of

hPAECs, whereas its overexpression led to changes in the cytoskeletal architecture of

hPAECs, implying a critical role of TXNDC5 in endothelial dysfunction in PH. In *in*

vivo studies, we found that global or endothelial-specific TXNDC5 deficiency

attenuated, whereas pulmonary endothelial-specific TXNDC5 overexpression

aggravated SuHx-induced PH. E64FC26, a pan-PDI inhibitor with IC50 of 16.3 μM

against TXNDC5, effectively delayed the progression of multiple myeloma *in vivo* in

the previous study[25]. We demonstrated in the present study that E64FC26

administration successfully alleviated the PA remodeling and RV hypertrophy,

effectively preventing the development and progression of SuHx-induced PH in rats.

Notably, administration of E64FC26 did not confer additional protection in endothelial

TXNDC5-deficient mice, indicating that its therapeutic effects are largely dependent

on endothelial TXNDC5. Collectively, these findings indicate that targeting TXNDC5

could be a promising strategy to treat PH.

We next examined the molecular mechanisms of the critical role of TXNDC5 in

PH development. Recent studies have demonstrated that alterations in ECM

composition and stiffness contribute to the pathological changes observed in PH,

influencing vascular resistance and exacerbating disease progression[27]. More

importantly, ECM remodeling is the cause rather than a consequence of pulmonary

vascular remodeling in PH[19]. We identified in the present study that TXNDC5 might

play a central role in orchestrating ECM homeostasis in hypoxic PH. Specifically,

TXNDC5 modulated proteoglycans and glycoproteins, which determine ECM stiffness

and elasticity, as well as collagens that define ECM structural integrity. Additionally,

TXNDC5 regulated matrix modulators and adhesion proteins involved in ECM

crosslinking, along with metallopeptidases responsible for ECM degradation and

turnover. Furthermore, TXNDC5 influenced basement membrane proteins,

angiogenesis, and growth factor-associated proteins, contributing to endothelial barrier

function and vascular homeostasis. More importantly, we demonstrated that POSTN, BGN, LOX, and TGM2 were excessively deposited in the PAs of TXNDC5[f/f] mice but were significantly reduced in TXNDC5[ECKO] mice under SuHx conditions. As key proteoglycans, POSTN and BGN regulate ECM organization and mechanotransduction, contributing to vascular stiffening[28–30]. Meanwhile, LOX and TGM2, categorized under matrix modulators, facilitate ECM crosslinking and collagen stabilization, further exacerbating pulmonary vascular remodeling[31,32]. These findings imply that TXNDC5 inhibition shifts the ECM equilibrium toward a less fibrotic and more compliant phenotype, mitigating PH progression by preserving vascular ECM homeostasis.

BGN has been demonstrated to play a significant role in clinical conditions related to inflammation, metabolic disturbances, musculoskeletal irregularities, and tumorigenesis[33]. We found in the present study that BGN was a direct binding partner of TXNDC5, with enhanced interaction with hypoxia stimulation. Mutations at Cys89 and Gly351 of TXNDC5 significantly disrupted the association between TXNDC5 and BGN. Notably, both amino acids were within the PDI active domain of TXNDC5, implying that its PDI activity is directly involved in mediating BGN binding and stabilization. Additionally, WT-TXNDC5, but not PDI-inactive TXNDC5 mutant, markedly increased the FRET efficiency in the BGN folding assay. These findings suggest that TXNDC5 facilitates BGN folding dependent on PDI activity, which is essential for the maturation and secretion of BGN as well as many other ECM proteins. Further investigation revealed that BGN was upregulated in the lungs of PH patients and rodent models, particularly in the intima of remodeled pulmonary arteries. Endothelial cell-specific deficiency of BGN alleviated PH progression in mice, implying that targeting endothelial-derived BGN may be a potential therapeutic strategy for PH. Importantly, endothelial-specific deficiency of BGN restored distal PA remodeling and RV hypertrophy induced by TXNDC5 overexpression in mice, indicating that BGN is a critical downstream effector mediating TXNDC5-driven PH. Additionally, rhBGN promoted the proliferation and apoptosis resistance of hPASMCs, suggesting that extracellular BGN also contributes to vascular remodeling. More importantly, we found that serum BGN level was significantly increased in PH patients

and was perfectly correlated with the disease severity. Our data, therefore, highlighted a potential for BGN to be a diagnostic biomarker of PH.

HIF signaling is one of the most dominant pathways involved in the development of hypoxic PH[34]. HIF-α subunits are rapidly degraded under normoxic conditions by the VHL E3 ligase[35]. In the present study, we demonstrated that silencing of BGN effectively abolished the upregulation of HIF-1α/2α induced by TXNDC5 overexpression and restored HIF-1α/2α ubiquitination by relieving VHL inhibition. These findings provide new insights into how ECM remodeling is coordinated with hypoxia-driven cellular responses in PH. Given that TXNDC5 potentially regulates ECM homeostasis and the established interplay between ECM and the HIF pathway, we speculate that factors form a complex regulatory network that contributes to the PH progression. Whether other ECM components are involved in the TXNDC5-HIF axis warrants further investigation. Taken together, targeting TXNDC5 may offer therapeutic strategies for regulating both ECM dynamics and hypoxic signaling.

Evidence has shown that excessive depletion of HIF-1α or HIF-2α may lead to organ dysfunction, developmental defects, and even embryonic lethality in mice[36–38]. This highlights the potential risks of systemic HIF inhibition. Interestingly, although clinical trial data are not yet available, preclinical studies have shown that various small-molecule HIF inhibitors exhibit promising therapeutic effects in PH models[34]. In the present study, we demonstrated that TXNDC5 is primarily expressed and upregulated in endothelial cells. Targeting TXNDC5 may offer a more tissue-specific approach to inhibit HIF signaling, thereby potentially reducing or avoiding the systemic side effects. Notably, global TXNDC5 knockout prevented SuHx-induced HIF signaling activation, without causing any adverse effects on growth and organ development in mice. These findings suggest that targeting TXNDC5 blocks HIF-driven pathological processes in PH, offering a promising alternative to HIF inhibition with better tissue specificity and fewer adverse effects.

In summary, we identified for the first time the novel role of PDI protein TXNDC5 in the pathogenesis of hypoxic PH. TXNDC5 facilitates ECM protein (e.g., BGN) folding and reduces HIF-1α/2α ubiquitination, which contributes to endothelial

dysfunction. TXNDC5 promotes a positive feedback loop between BGN and HIF signaling, driving the progression of pulmonary vascular remodeling in PH. This study highlights TXNDC5 as a promising therapeutic target for hypoxia-induced PH and lays the groundwork for future clinical investigations.

Acknowledgments

We thank Professor John Shyy from the University of California, San Diego, United States, for critical reading and advice on this article.

Sources of Funding

This work was supported by the Key Program of National Natural Science Foundation of China (NSFC, grant No. 82430004 to Jin-Song Bian), Shenzhen Science Fund for Distinguished Young Scholars (grant No. RCJC20210706091946002 to Xiaowei Nie), Shenzhen Science and Technology Program (grant No. GJHZ20240218111401002 to Jin-Song Bian), and the National Natural Science Foundation of China (NSFC, grant No. 82241021 to Xiaowei Nie).

References

1. Johnson S, Sommer N, Cox-Flaherty K, Weissmann N, Ventetuolo CE, Maron BA. Pulmonary Hypertension: A Contemporary Review. *Am J Respir Crit Care Med*. 2023;208:528–548. doi: 10.1164/rccm.202302-0327SO

2. Shlobin OA, Adir Y, Barbera JA, Cottin V, Harari S, Jutant E-M, Pepke-Zaba J, Ghofrani H-A, Channick R. Pulmonary hypertension associated with lung diseases. *Eur Respir J*. 2024;64:2401200. doi: 10.1183/13993003.01200-2024

3. Evans CE, Cober ND, Dai Z, Stewart DJ, Zhao Y-Y. Endothelial cells in the pathogenesis of pulmonary arterial hypertension. *Eur Respir J*. 2021;58:2003957. doi: 10.1183/13993003.03957-2020

4. Ma B, Cao Y, Qin J, Chen Z, Hu G, Li Q. Pulmonary artery smooth muscle cell phenotypic switching: A key event in the early stage of pulmonary artery hypertension. *Drug Discov Today*. 2023;28:103559. doi: 10.1016/j.drudis.2023.103559

5.   Mandras SA, Mehta HS, Vaidya A. Pulmonary Hypertension: A Brief Guide for
Clinicians. *Mayo Clin Proc*. 2020;95:1978–1988. doi:
10.1016/j.mayocp.2020.04.039

6.   Olsson KM, Corte TJ, Kamp JC, Montani D, Nathan SD, Neubert L, Price LC,
Kiely DG. Pulmonary hypertension associated with lung disease: new insights
into pathomechanisms, diagnosis, and management. *Lancet Respir Med*.
2023;11:820–835. doi: 10.1016/S2213-2600(23)00259-X

7.   Kanemura S, Matsusaki M, Inaba K, Okumura M. PDI Family Members as Guides
for Client Folding and Assembly. *Int J Mol Sci*. 2020;21:9351. doi:
10.3390/ijms21249351

8.   Kuramochi T, Yamashita Y, Arai K, Kanemura S, Muraoka T, Okumura M.
Boosting the enzymatic activity of CxxC motif-containing PDI family members.
Chem Commun (Camb). 2024;60:6134–6137. doi: 10.1039/d4cc01712a

9.   Rahman NSA, Zahari S, Syafruddin SE, Firdaus-Raih M, Low TY, Mohtar MA.
Functions and mechanisms of protein disulfide isomerase family in cancer
emergence. *Cell Biosci*. 2022;12:129. doi: 10.1186/s13578-022-00868-6

10.   Kobayashi Y, Oguro A, Hirata Y, Imaoka S. The regulation of Hypoxia-Inducible
Factor-1 (HIF-1alpha) expression by Protein Disulfide Isomerase (PDI). *PLoS*
One. 2021;16:e0246531. doi: 10.1371/journal.pone.0246531

11.   Nagarkoti S, Kim Y-M, Ash D, Das A, Vitriol E, Read T-A, Youn S-W, Sudhahar
V, McMenamin M, Hou Y, et al. Protein disulfide isomerase A1 as a novel redox
sensor in VEGFR2 signaling and angiogenesis. *Angiogenesis*. 2023;26:77–96. doi:
10.1007/s10456-022-09852-7

12.   Hung C-T, Tsai Y-W, Wu Y-S, Yeh C-F, Yang K-C. The novel role of ER protein
TXNDC5 in the pathogenesis of organ fibrosis: mechanistic insights and
therapeutic implications. *J Biomed Sci*. 2022;29:63. doi: 10.1186/s12929-022-
00850-x

13.   Jin Y, Sharma A, Bai S, Davis C, Liu H, Hopkins D, Barriga K, Rewers M, She J-
X. Risk of type 1 diabetes progression in islet autoantibody-positive children can
be further stratified using expression patterns of multiple genes implicated in
peripheral blood lymphocyte activation and function. *Diabetes*. 2014;63:2506–
2515. doi: 10.2337/db13-1716

14.   Wang L, Dong H, Song G, Zhang R, Pan J, Han J. TXNDC5 synergizes with
HSC70 to exacerbate the inflammatory phenotype of synovial fibroblasts in
rheumatoid arthritis through NF-κB signaling. *Cell Mol Immunol*. 2018;15:685–
696. doi: 10.1038/cmi.2017.20

15.   Wang X, Li H, Chang X. The role and mechanism of TXNDC5 in diseases. *Eur J*

Med Res. 2022;27:145. doi: 10.1186/s40001-022-00770-4

16. Yeh C-F, Cheng S-H, Lin Y-S, Shentu T-P, Huang R-T, Zhu J, Chen Y-T, Kumar S, Lin M-S, Kao H-L, et al. Targeting mechanosensitive endothelial TXNDC5 to stabilize eNOS and reduce atherosclerosis in vivo. *Sci Adv*. 2022;8:eabl8096. doi: 10.1126/sciadv.abl8096

17. Wu S, Luo X, Chen Y, Wang Z, Liu X, Sun N, Zhao J, Luo W, Zhang J, Tong X, et al. Sodium-glucose cotransporter 2 inhibitors attenuate vascular calcification by suppressing endoplasmic reticulum protein thioredoxin domain containing 5 dependent osteogenic reprogramming. *Redox Biol*. 2024;73:103183. doi: 10.1016/j.redox.2024.103183

18. Fu Y, Zhou Y, Wang K, Li Z, Kong W. Extracellular Matrix Interactome in Modulating Vascular Homeostasis and Remodeling. *Circ Res*. 2024;134:931–949. doi: 10.1161/CIRCRESAHA.123.324055

19. Thenappan T, Chan SY, Weir EK. Role of extracellular matrix in the pathogenesis of pulmonary arterial hypertension. *Am J Physiol Heart Circ Physiol*. 2018;315:H1322–H1331. doi: 10.1152/ajpheart.00136.2018

20. Nie X, Shen C, Tan J, Wu Z, Wang W, Chen Y, Dai Y, Yang X, Ye S, Chen J, et al. Periostin: A Potential Therapeutic Target For Pulmonary Hypertension? *Circ Res*. 2020;127:1138–1152. doi: 10.1161/CIRCRESAHA.120.316943

21. Bian J-S, Chen J, Zhang J, Tan J, Chen Y, Yang X, Li Y, Deng L, Chen R, Nie X. ErbB3 Governs Endothelial Dysfunction in Hypoxia-Induced Pulmonary Hypertension. *Circulation*. 2024;150:1533–1553. doi: 10.1161/CIRCULATIONAHA.123.067005

22. Zhang Y, Li X, Li S, Zhou Y, Zhang T, Sun L. Immunotherapy for Pulmonary Arterial Hypertension: From the Pathogenesis to Clinical Management. *Int J Mol Sci*. 2024;25:8427. doi: 10.3390/ijms25158427

23. Shih Y-C, Chen C-L, Zhang Y, Mellor RL, Kanter EM, Fang Y, Wang H-C, Hung C-T, Nong J-Y, Chen H-J, et al. Endoplasmic Reticulum Protein TXNDC5 Augments Myocardial Fibrosis by Facilitating Extracellular Matrix Protein Folding and Redox-Sensitive Cardiac Fibroblast Activation. *Circ Res*. 2018;122:1052–1068. doi: 10.1161/CIRCRESAHA.117.312130

24. Luo Z, Tian M, Yang G, Tan Q, Chen Y, Li G, Zhang Q, Li Y, Wan P, Wu J. Hypoxia signaling in human health and diseases: implications and prospects for therapeutics. *Signal Transduct Target Ther*. 2022;7:218. doi: 10.1038/s41392-022-01080-1

25. Robinson RM, Reyes L, Duncan RM, Bian H, Reitz AB, Manevich Y, McClure JJ, Champion MM, Chou CJ, Sharik ME, et al. Inhibitors of the protein disulfide

isomerase family for the treatment of multiple myeloma. *Leukemia*. 2019;33:1011–1022. doi: 10.1038/s41375-018-0263-1

26. Sullivan DC, Huminiecki L, Moore JW, Boyle JJ, Poulsom R, Creamer D, Barker J, Bicknell R. EndoPDI, a novel protein-disulfide isomerase-like protein that is preferentially expressed in endothelial cells acts as a stress survival factor. *J Biol Chem*. 2003;278:47079–47088. doi: 10.1074/jbc.M308124200

27. Jandl K, Radic N, Zeder K, Kovacs G, Kwapiszewska G. Pulmonary vascular fibrosis in pulmonary hypertension - The role of the extracellular matrix as a therapeutic target. *Pharmacol Ther*. 2023;247:108438. doi: 10.1016/j.pharmthera.2023.108438

28. Sun X-J, Ma W-Q, Zhu Y, Liu N-F. POSTN promotes diabetic vascular calcification by interfering with autophagic flux. *Cell Signal*. 2021;83:109983. doi: 10.1016/j.cellsig.2021.109983

29. Nikoloudaki G, Snider P, Simmons O, Conway SJ, Hamilton DW. Periostin and matrix stiffness combine to regulate myofibroblast differentiation and fibronectin synthesis during palatal healing. *Matrix Biol*. 2020;94:31–56. doi: 10.1016/j.matbio.2020.07.002

30. Yu M, He X, Song X, Gao J, Pan J, Zhou T, Wang Q, Zhu W, Ma H, Zeng H, et al. Biglycan promotes hepatic fibrosis through activating heat shock protein 47. *Liver Int*. 2023;43:500–512. doi: 10.1111/liv.15477

31. Wang Z-J, Guan Q-W, Zhou H-H, Mao X-Y, Chen F-H. Mechanistic insight into lysyl oxidase in vascular remodeling and angiogenesis. *Genes Dis*. 2023;10:771–785. doi: 10.1016/j.gendis.2022.05.011

32. Zhang M, Li L, Zhang W, Li M, Yan G, Tang C. TG2 participates in pulmonary vascular remodelling by regulating the senescence of pulmonary artery smooth muscle cells. *Cell Signal*. 2024;121:111296. doi: 10.1016/j.cellsig.2024.111296

33. Appunni S, Rubens M, Ramamoorthy V, Anand V, Khandelwal M, Sharma A. Biglycan: an emerging small leucine-rich proteoglycan (SLRP) marker and its clinicopathological significance. *Mol Cell Biochem*. 2021;476:3935–3950. doi: 10.1007/s11010-021-04216-z

34. Pullamsetti SS, Mamazhakypov A, Weissmann N, Seeger W, Savai R. Hypoxia-inducible factor signaling in pulmonary hypertension. *J Clin Invest*. 2020;130:5638–5651. doi: 10.1172/JCI137558

35. Tarade D, Ohh M. The HIF and other quandaries in VHL disease. *Oncogene*. 2018;37:139–147. doi: 10.1038/onc.2017.338

36. Kotch LE, Iyer NV, Laughner E, Semenza GL. Defective vascularization of HIF-

1alpha-null embryos is not associated with VEGF deficiency but with
mesenchymal cell death. *Dev Biol*. 1999;209:254–267. doi:
10.1006/dbio.1999.9253

37. Chen H, Houshmand G, Mishra S, Fong G-H, Gittes GK, Esni F. Impaired
pancreatic development in Hif2-alpha deficient mice. *Biochem Biophys Res*
Commun. 2010;399:440–445. doi: 10.1016/j.bbrc.2010.07.111

38. Peng Y-J, Nanduri J, Khan SA, Yuan G, Wang N, Kinsman B, Vaddi DR, Kumar
GK, Garcia JA, Semenza GL, et al. Hypoxia-inducible factor 2α (HIF-2α)
heterozygous-null mice exhibit exaggerated carotid body sensitivity to hypoxia,
breathing instability, and hypertension. *Proc Natl Acad Sci U S A*. 2011;108:3065–
3070. doi: 10.1073/pnas.1100064108

Figure legends

Figure 1. TXNDC5 was upregulated in PH.

A, Heatmap of 4D label-free proteomics showing PDI family expression in lung tissues

from 5 non-PH donors and 5 patients with hypoxic PH. **B**, Representative Western blots

and quantification of TXNDC5 and Vinculin in the lungs of control donors and PH

patients (n=7). **C**, qPCR analysis of TXNDC5 mRNA levels in lung homogenates from donors and patients with PH (n=7). **D**, Western blots showing the TXNDC5 protein levels in the lungs collected from MCT (n=7) or SuHx-treated rats (n=6) or rats under normoxia. **E**, **F**, and **G**, Representative confocal images of multiplex immunofluorescence (mIF) staining for TXNDC5 (red), CD31 (green), α-SMA (white), and COL1A1 (purple) in distal pulmonary arteries (PAs) from donors, ILD-PH, and COPD-PH patients. Nuclei were counterstained with DAPI (blue). White arrows indicate the colocalization of TXNDC5 with CD31 (yellow). White dashed lines indicate the outer layer of smooth muscle cells. Scale bars=25 μm. **H**, Quantitative analysis showing elevated TXNDC5 mean fluorescence intensity (MFI) in the CD31$^+$ intima of distal PAs from patients with ILD-PH and COPD-PH compared with non-PH donors. (n=5). **I**, **J**, and **K**, Representative images of mIF staining for TXNDC5 (red), CD31 (green), α-SMA (cyan), and COL1A1 (purple) in distal PAs from SuHx- or MCT-treated rats or rats under normoxia. Nuclei were counterstained with DAPI (blue). White arrows indicate the colocalization of TXNDC5 with CD31 (yellow). White dashed lines indicate the outer layer of smooth muscle cells. Scale bars=20 μm. **L**, **M**, and **N**, Correlation analyses of intimal TXNDC5 MFI and right ventricular systolic pressure (RVSP), Fulton index [RV/(LV+S)], and tricuspid annular plane systolic excursion (TAPSE) in rats. Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**B-D**) or 2-way ANOVA (**H**), followed by Tukey honestly significant difference test for multiple comparisons and the Pearson correlation test (**L-N**). PDI indicates protein disulfide isomerase; PH, pulmonary hypertension; TXNDC5, thioredoxin domain containing 5; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; DAPI, 4′,6-diamidino-2-phenylindole; Nor, normoxia; SuHx, Sugen5416/hypoxia; and MCT, monocrotaline.

Figure 2. TXNDC5 contributed to SuHx-induced pulmonary vascular remodeling

A, Schematic protocol for the animal treatments in the SuHx-induced PH mice. **B**, Representative tracings and quantification of right ventricular systolic pressure (RVSP) in male WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=10). **C**,

Representative multiplex immunofluorescence (mIF) for CD31 (green), α-SMA (white), and COL1A1 (red), and Victoria blue/Sirius red (VB/SR) staining for elastic lamina (dark blue), collagens (red), and muscle fiber (yellow) in distal pulmonary arteries (PAs) from male WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx. Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **D**, Quantification of distal PA muscularization based on α-SMA positive area in WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=8). **E**, Measurement of medial wall thickness in distal PAs, defined by the elastic lamina visualized with VB staining (n=8). **F**, Quantification of collagen deposition area in distal PAs based on SR staining, indicating decreased collagen deposition in TXNDC5-deficient mice (n=8). **G**, Immunofluorescence staining of WGA (green) in right ventricles from male WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=6). Nuclei were counterstained with DAPI (blue). Scale bars=20 μm. **H**, Schematic diagram illustrating the *in vivo* pulmonary endothelial-specific overexpression of TXNDC5 in both male and female mice. Mice received a single tail vein injection of AAV-NC or AAV-TXN at the beginning of the 2 weeks, followed by SuHx exposure. **I**, Representative images of major organs from saline-treated control or AAV-TXN-treated mice, showing GFP fluorescence delivered by AAV detected by IVIS imaging system. **J**, Hemodynamic measurements of RVSP in male and female mice administered with AAV-NC or AAV-TXN under normoxia or SuHx situations (n=8). **K**, Representative mIF of distal PAs from AAV-NC or AAV-TXN-treated male and female mice under SuHx or normoxia, showing CD31 (green), α-SMA (white), and COL1A1 (red), together with VB/SR staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **L**, **M,** and **N**, Quantitative analysis of distal PA remodeling in AAV-NC or AAV-TXN-treated mice under normoxic or SuHx conditions (n=8). PA muscularization, medial wall thickness, and collagen deposition area were assessed using α-SMA, VB, and SR staining, respectively. Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by Tukey honestly significant difference test for multiple comparisons. DAPI indicates 4′,6-diamidino-2-

phenylindole; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; WGA,
wheat germ agglutinin; GFP, green fluorescent protein; Nor, normoxia; SuHx, and
Sugen5416/hypoxia.

Figure 3. Endothelial cell-specific deletion of TXNDC5 ameliorated SuHx-induced

PH.

A, Schematic protocol for the endothelial cell-specific TXNDC5 knockout mice assays. Mice were intraperitoneally injected with 75 mg/kg tamoxifen once daily for 5 days before being exposed to SuHx. **B**, Hemodynamic measurements of right ventricular systolic pressure (RVSP) in male TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice exposed to normoxia or SuHx (n=8). **C**, Fulton index [RV/(LV+S)] in male TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice (n=8). **D**, Representative multiplex immunofluorescence (mIF) of distal pulmonary arteries (PAs) from male TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice under normoxia or SuHx, showing CD31 (green), α-SMA (white), and COL1A1 (red), together with Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **E**, Quantification of distal PA muscularization based on α-SMA positive area (n=8). **F**, Medial wall thickness of distal PAs, measured across the elastic lamina visualized by VB staining (n=8). **G**, Perivascular collagen deposition quantified using SR staining (n=8). **H**, Representative lung angiogram of the left lungs from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with SuHx or under normoxic conditions. Scale bars=2 mm. **I**, Histological analysis of the cross-sectioned heart at the mid-RV (red arrows) plane from male TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice with or without SuHx exposure (n=6). Scale bars=1 mm. **J**, Immunofluorescence staining of WGA (green) in RV from male TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice exposed to normoxia or SuHx (n=6). Nuclei were counterstained with DAPI (blue). Scale bars=20 μm. Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by Tukey honestly significant difference test for multiple comparisons. DAPI indicates 4′,6-diamidino-2-phenylindole; H&E, hematoxylin-eosin; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; WGA, wheat germ agglutinin; f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; and RV, right ventricle.

Figure 4. TXNDC5 regulated ECM protein expression and distribution.

A, HPAECs were transfected with siTXNDC5 or siNC under hypoxic conditions, followed by RNA sequencing analysis (n=3). Bubble chart showing the Gene Ontology enrichment using DAVID for the differentially expressed genes. **B**, Differentially

expressed ECM proteins were selected and visualized by heatmap (left). The PPI network showed that TXNDC5-targeted ECM proteins could be classified into different functional subgroups (right). **C**, **D**, **E**, and **F**, Representative images of immunofluorescence staining of α-SMA (red), CD31 (green), and POSTN, BGN, LOX, or TGM2 (white) in PAs from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice under normoxia and SuHx conditions. Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=20 μm. **G**, Western blots showing the protein levels of COL1A1, TGM2, FN1, POSTN, and LOX in hPAECs transferred with siTXNDC5 or siNC in normoxia or hypoxia conditions (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the 2-way ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. ECM indicates extracellular matrix; PPI, protein-protein interaction; Nor, normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; si, small interfering; α-SMA, α-smooth muscle actin; POSTN, periostin; BGN, biglycan; LOX, lysyl oxidase; TGM2, transglutaminase 2; FN1, fibronectin; and COL1A1, collagen alpha-1 chain.

Figure 5. BGN served as a biomarker and therapeutic target of PH.

A, The progressive screening of the PPI network between ECM genes and ECs function-related genes. Each new PPI network is composed of core genes in the previous network. **B**, Box-whisker plot showing the relative mRNA levels of TXNDC5

in lung tissues from PH patients (n=96) and healthy donors (n=52) (GSE254617). **C**, Enzyme-linked immunosorbent assay showing the BGN level in the serum of non-PH donors (n=27) and patients with PH (n=27). **D**, Linear regression analysis showing the correlation between serum BGN concentrations with sPAP of PH patients (n=27). **E**, Quantification of right ventricular systolic pressure (RVSP) in male $BGN^{f/0}$ and BGN^{ECKO} mice exposed to normoxia or SuHx (n=8). **F**, Fulton index [RV/(LV+S)] in $BGN^{f/0}$ and BGN^{ECKO} mice exposed to normoxia or SuHx (n=8). **G**, Representative multiplex immunofluorescence (mIF) of distal pulmonary arteries (PAs) from $BGN^{f/0}$ and BGN^{ECKO} mice under normoxia or SuHx, showing CD31 (green), α-SMA (white), and COL1A1 (red), together with Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **H**, Quantification of distal PA muscularization based on α-SMA positive area (n=8). **I**, Medial wall thickness of distal PAs, measured across the elastic lamina visualized by VB staining (n=8). **J**, Representative lung angiogram of the left lungs from $BGN^{f/0}$ and BGN^{ECKO} mice treated with SuHx or under normoxic conditions. Scale bars=2 mm. **K**, Histological analysis of heart cross-sections at the mid-RV plane (top) and highlighted right ventricular wall (bottom) of the male $BGN^{f/0}$ and BGN^{ECKO} mice exposed to normoxia or SuHx (n=6). Scale bars=1 mm. Data represent the mean±SD. Comparisons of parameters were performed with the 2-tailed Mann-Whitney *U* test (**B** and **C**), Pearson correlation test (**D**), or 2-way ANOVA (**E-F**, **H-I,** and **K**), followed by Tukey honestly significant difference test for multiple comparisons. H&E indicates hematoxylin-eosin; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; DAPI, 4′,6-diamidino-2-phenylindole; f/0, flox/0; Nor, normoxia; SuHx, Sugen5416/hypoxia; RV, right ventricle; and sPAP, systolic pulmonary arterial pressure.

Figure 6. TXNDC5 assisted BGN protein folding to promote PH.

A, Immunofluorescence staining showing the localization of TXNDC5 (green) and BGN (red) in hPAECs. Nuclei (blue) were stained with DAPI. White arrows in the enlarged view indicate the colocalization of TXNDC5 with BGN (yellow). Scale

bars=10 μm. **B**, Coimmunoprecipitation analysis showing the interaction between TXNDC5 and BGN in hPAECs under both normoxic and hypoxic situations. IP: TXNDC5, immunoprecipitation with TXNDC5 antibody. IgG for the negative control. **C**, Molecular docking showing the binding domains of human TXNDC5 and BGN. **D**, Immunoprecipitation with anti-HA beads in HEK293T cells transfected with FLAG-BGN and HA-TXNDC5 mutant plasmids. **E**, FRET acceptor photobleaching of the hPAECs transfected with CFP-BGN-YFP plasmid. CFP and YFP images were taken both before and after acceptor photobleaching. FRET efficiency is displayed as a pseudocolor representation and qualification by ImageJ (n=5). Scale=10 μm. **F**, Schematic of the *in vivo* rescue experiments: adeno-associated virus encoding TXNDC5 (AAV-TXN) or control AAV (AAV-NC) was delivered to lung ECs in $BGN^{f/0(f/f)}$ and BGN^{ECKO} mice under normoxia or SuHx conditions. **G**, Representative multiplex immunofluorescence (mIF) images of distal pulmonary arteries (PAs) from $BGN^{f/0(f/f)}$ and BGN^{ECKO} mice treated with AAV-NC or AAV-TXN, showing CD31 (green), α-SMA (white), and COL1A1 (red), with Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei stained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **H**, Quantification of distal PA muscularization in $BGN^{f/0(f/f)}$ and BGN^{ECKO} mice with AAV-NC or AAV-TXN administration (n=6). **I**, Medial wall thickness of distal PAs measured across the elastic lamina, visualized by VB staining (n=6). **J**, Quantification of collagen deposition area in distal PAs (n=6). **K**, Right ventricular systolic pressure (RVSP) in $BGN^{f/0(f/f)}$ and BGN^{ECKO} mice treated with AAV-NC or AAV-TXN under SuHx or normoxia conditions (n=6). **L**, Fulton index [RV/(LV+S)] in $BGN^{f/0(f/f)}$ and BGN^{ECKO} mice with AAV-NC or AAV-TXN treatment (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the 1-way (**E**) or 2-way (**H-L**) ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. ECs, endothelial cells; Nor, normoxia; Hyp, hypoxia; IgG, immunoglobulin G; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; FRET, fluorescence resonance energy transfer; CFP, cyan fluorescent protein; and YFP, yellow fluorescent protein.

Figure 7. Endothelial TXNDC5 Driven PH via BGN-dependent stabilization of

HIF-1α/2α.

A, Dual-luciferase assay showing that luciferase activity in constructs with BS1

mutation (Mut-1), BS2 mutation (Mut-2), or double mutation of BS1 and BS2 (dMut)

(n=6). **B**, Western blots showing the decreased protein levels of HIF-1α and HIF-2α in the lungs from TXNDC5 deficiency mice compared with WT exposed to SuHx (n=6). **C**, Representative distal pulmonary arteries (PAs) images from AAV-NC or AAV-TXN-treated mice exposed to SuHx or normoxia, with HIF-1α inhibitor (PX-478) or HIF-2α inhibitor (PT-2385), or vehicle administration. Multiplex immunohistochemical (mIF) showing CD31 (green), α-SMA (white), and COL1A1 (red) and Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **D**, **E**, and **F**, Quantitative analysis of distal PA remodeling, including PA muscularization assessed by α-SMA-positive area (n=6), medial wall thickness measured across the elastic lamina visualized by VB staining (n=6), and perivascular collagen area quantified using SR staining (n=6). **G**, Representative Western blots showing that the LV-TXNDC5-induced upregulation of HIF-1/2α was abolished by siBGN in hPAECs both under normoxia and hypoxia conditions (n=6). **H**, Western blot showing that the protein stabilization of HIF-1/2α was enhanced by LV-TXNDC5 treatment in hPAECs under normoxic conditions (n=5). **I**, Representative Western blots of HIF-1α, HIF-2α, and VHL in hPAECs after treatment with TXNDC5 knockdown only or double knockdown (dKO) of TXNDC5 and VHL (n=6). **J**, Western blot analysis indicating the HIF-2α ubiquitylation profile was reduced by LV-TXNDC5 but increased by siBGN treatment. hPAECs were treated with 10 μM MG132 for 6 h before harvest. Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. BS indicates binding site; VHL, von Hippel-Lindau; Ub, ubiquitin; Nor, normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypixa; IgG, immunoglobulin G; LV, lentivirus; α-SMA, α-smooth muscle actin; and COL1A1, collagen type I alpha 1.

1023

Figure 8. Pharmacological and Pulmonary Endothelium-Targeted Inhibition of

TXNDC5 Ameliorated PH.

A, Schematic protocol for E64FC26 administration for SuHx-induced PH rats. Rats

were treated with SuHx for 3 weeks and then intraperitoneal injection with 2 mg/kg

E64FC26 daily for 2 weeks under normoxic conditions. **B**, Quantification of right

ventricular systolic pressure (RVSP) in rats administered with E64FC26 or vehicle

exposed to normoxia or SuHx (n=6). **C**, Fulton index [RV/(LV+S)] of rats with

E64FC26 or vehicle treatment under normoxia or SuHx conditions (n=6). **D**,

Representative multiplex immunofluorescence (mIF) of distal pulmonary arteries (PAs)

from rats treated with vehicle or E64FC26 under normoxia or SuHx conditions,

showing CD31 (green), α-SMA (white), and COL1A1 (red), together with Victoria

blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens

(red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L

indicates lumen. Scale bars = 25 μm. **E**, Histological and quantification analysis of the

cross-sectioned heart at the mid-RV plane (top) and the highlighted right ventricular

wall (bottom) of rats treated with E64FC26 or vehicle under normoxia or SuHx

conditions (n=6). Scale bars=2 mm. **F**, Masson's trichrome staining and quantification

of the fibrotic area by collagen volume fraction measured by Image J of RV from rats

treated with E64FC26 or vehicle exposed to normoxia or SuHx (n=6). Scale bars=100

μm. **G**, Schematic of vehicle or E64FC26 treatment in TXNDC5$^{f/f}$ and TXNDC5^{ECKO}

mice under SuHx conditions to investigate whether the effects of E64FC26 are

TXNDC5-dependent. **H**, Representative distal PAs images from TXNDC5$^{f/f}$ and

TXNDC5^{ECKO} mice treated with vehicle or E64FC26 under SuHx or normoxia

conditions. Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue),

perivascular collagens (red), and muscle fiber (yellow). L indicates lumen. Scale

bars=25 μm. **I**, Representative lung angiogram of SuHx-treated TXNDC5$^{f/f}$ and

TXNDC5^{ECKO} mice with or without E64FC26 administration, showing distal

pulmonary perfusion. Scale bars=2 mm. **J**, Schematic diagram of the pulmonary

endothelium-targeted gene therapy strategy. Rats received a tail vein injection of self-

complementary adeno-associated virus (scAAV) encoding shRNA against TXNDC5

(shTXN) or control shRNA (shNC). **K**, Representative IVIS images of major organs

from rats treated with Saline or scAAV-shTXN, showing GFP fluorescence delivered

by the scAAV vector. **L**, Representative mIF for CD31 (green), α-SMA (white), and

COL1A1 (red), and VB/SR staining for elastic lamina (dark blue), collagens (red), and

muscle fiber (yellow) in distal PAs from rats receiving shNC or shTXN exposed to normoxia or SuHx. Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **M**, Hemodynamic measurement of RVSP in SuHx-induced PH rats treated with shNC or shTXN (n=6). **N**, Histological sections of the heart at the mid-RV plane and quantification of right ventricular wall thickness from SuHx-induced PH rats treated with shNC or shTXN (n=6). Scale bars = 2 mm. Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by Tukey honestly significant difference test for multiple comparisons. H&E indicates hematoxylin-eosin; α-SMA, α-smooth muscle actin; COL1A1, collagen type I alpha 1; shRNA, short hairpin RNA; f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; and RV, right ventricle.

Graphic Abstract. The role of TXNDC5 in regulating ECM homeostasis during PH. In the healthy state, BGN folding is properly maintained and HIF-1α/2α is rapidly degraded. During PH, TXNDC5 is upregulated, leading to excessive accumulation of BGN and impaired ubiquitination-mediated degradation of HIF-1α/2α, resulting in an imbalance in ECM homeostasis and contributing to vascular remodeling. Pharmacological or genetic inhibition of TXNDC5 restores BGN folding capacity and facilitates HIF-1α/2α degradation, thereby reestablishing ECM homeostasis and

1078   attenuating PH progression.

1079

Supplemental Figures

1081

Figure S1. Upregulation of TXNDC5 in the lungs and lung endothelial cells of PH rodent models.

A, Schematic illustration showing the 4D label-free proteomics of lungs from non-PH donors and patients with hypoxic PH. **B**, Western blot analysis and quantification of TXNDC5 and Vinculin in the lungs of mice exposed to SuHx or hypoxia or under normoxic situations (n=6). **C**, Quantitative analysis of mIF showing TXNDC5 was predominantly co-localized with CD31 in distal PAs. (n=5). **D**, Quantification of mean fluorescence intensity (MFI) of TXNDC5 in remodeled distal PAs from SuHx- and MCT-induced rat PH models compared with rats under normoxic conditions. (n=6). **E**, Western blot of TXNDC5 in primary lung ECs from SuHx-induced mice or normoxic controls, showing increased protein levels in ECs. **F**, Western blot of TXNDC5 in lung ECs isolated from SuHx-induced PH rats, demonstrating elevated expression compared with normoxic controls. Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**B**, **E**, and **F**) or 2-way ANOVA (**D**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; and ECs, endothelial cells.

1099

Figure S2. Role of TXNDC5 in PH development.

A, Fulton index [RV/(LV+S)] in male WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=8). **B**, **C**, **D**, and **E**, Echocardiography analysis of PAT/PET, TAPSE, RVFAC, and RVFWT in male WT and TXNDC5$^{-/-}$ mice exposed to normoxia or SuHx (n=10). **F**, Right ventricular systolic pressure (RVSP) in female WT and TXNDC5$^{-/-}$

mice under normoxia or SuHx (n=6). **G**, RV/(LV+S) in female WT and TXNDC5$^{-/-}$ mice (n=6). **H**, **I**, **J**, and **K**, Echocardiographic assessment of right ventricular function of female WT and TXNDC5$^{-/-}$ mice, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). **L**, Representative images of distal PAs from female WT and TXNDC5$^{-/-}$ mice under normoxia or SuHx conditions, showing α-SMA immunofluorescence (green) and Victoria blue/Sirius red (VB/SR) staining of elastic lamina (dark blue), perivascular collagens (red), and muscle fiber (yellow). Nuclei were counterstained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **M**, **N**, and **O**, Quantification of distal PA remodeling in female WT and TXNDC5$^{-/-}$ mice, including PA muscularization, medial wall thickness, and collagen deposition area (n=6). **P**, Representative confocal images showing GFP-TXNDC5 (green), CD31 (red), and α-SMA (white) in lung sections from AAV-TXN-treated mice. Nuclei were counterstained with DAPI (blue). L indicates lumen. A indicates airway. Scale bars=50 μm. **Q**, Western blot analysis of TXNDC5 protein levels in primary lung ECs isolated from AAV-TXN- or AAV-NC-treated mice (n=6). **R**, RV/(LV+S) in male and female mice administered with AAV-NC or AAV-TXN exposed to normoxia or SuHx (n=8). **S**, **T**, **U**, and **V**, Echocardiography analysis of PAT/PET, TAPSE, RVFAC, and RVFWT in both male and female mice administered with AAV-NC or AAV-TXN exposed to normoxia or SuHx (n=8). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student t-test (**Q**) or 2-way ANOVA (**A-K**, **M-O**, and **R-V**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; SuHx, Sugen5416/hypoxia; PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio; TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional area change; and RVFWT, right ventricular free wall thickness.

1131

Figure S3. Protective effects of endothelial cell-specific TXNDC5 deletion on SuHx-induced PH.

A, Schematic illustrating the generation of endothelial cell-specific TXNDC5 knockout (TXNDC5^{ECKO}) mice. **B**, Representative Western blots and quantification of TXNDC5 protein levels in primary lung ECs isolated from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice

(n=6). **C**, **D**, **E**, and **F**, Echocardiographic assessment of right ventricular function in male TXNDC5$^{f/f}$ and TXNDC5^{ECKO} under normoxia or SuHx, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). **G**, Enlarged lung angiogram images of the left lungs from TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with SuHx or under normoxic conditions. Scale bars=2 mm. **H**, Right ventricular systolic pressure (RVSP) in female TXNDC5$^{f/f}$ and TXNDC5^{ECKO} under normoxia or SuHx (n=6). **I**, Fulton index [RV/(LV+S)] in female TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice (n=6). **J**, **K**, **L**, and **M**, Echocardiographic evaluation of right ventricular function in female TXNDC5^{ECKO} and littermate control mice, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). **N**, Representative distal PA images from female TXNDC5$^{f/f}$ and TXNDC5^{ECKO} exposed to normoxia or SuHx, showing α-SMA (green) and VB/SR staining of elastic lamina (dark blue), collagens (red), and muscle fiber (yellow). Nuclei stained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **O**, **P**, and **Q**, Quantitative analysis of distal PA remodeling in female TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice, including PA muscularization, medial wall thickness, and collagen deposition (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**B**) or 2-way ANOVA (**C-F**, **H-M**, and **O-Q**), followed by Tukey honestly significant difference test for multiple comparisons. DAPI indicates 4′,6-diamidino-2-phenylindole; α-SMA, α-smooth muscle actin; f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio; TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional area change; and RVFWT, right ventricular free wall thickness.

1159

Figure S4. TXNDC5 regulated cell function in pulmonary artery.

A and **B**, Quantification of TXNDC5 protein levels by Western blots and TXNDC5

mRNA levels by qPCR analysis in hPAECs in both normoxia and hypoxia situations

(n=6). **C**, Immunofluorescence staining analysis showing increased protein levels of

TXNDC5 exposed to hypoxia. Scale bars=10 μm. **D**, Representative EdU (red) staining image and group data showing the effects of TXNDC5 silencing on the proliferative activity of hPAECs (n=6). Nuclei were counterstained with DAPI (blue). Scale bars=200 μm. **E**, Representative images and group data showing the migration of hPAECs treated with hPAECs siTXNDC5 or siNC in normoxia or hypoxia conditions (n=6). Scale bars=200 μm. **F** and **G**, Confocal microscopy images showing the effect of LV-TXNDC5 on the cytoskeletal architecture of hPAECs. Cells were stained with phalloidin (green, scale bars=20 μm) or paxillin (red, scale bars=10 μm). Nuclei were counterstained with DAPI (blue). **H**, Flow cytometry showing the effect of siTXNDC5 on the cell cycle of hPAECs exposed to normoxia or hypoxia. The representative images and quantification of percentages of cells in G1, S, and G2 were shown (n=6). X and Y axes, respectively, indicate the denoted DNA content stained with propidium iodide (PI) and cell counts. **I** and **J**, Representative Western blots and quantification showing the protein levels of Cyclin A, proliferating cell nuclear antigen (PCNA), and cleaved-caspase 3 of hPASMCs treated with conditioned medium from hPAECs transfected with siTXNDC5 or siNC in normoxia or hypoxia conditions (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**A** and **B**) or 2-way ANOVA (**D**, **E**, **I**, and **J**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; LV, lentivirus; si, small interfering; CM, conditioned medium; and EdU, 5-ethynyl-20-deoxyuridine.

1185

Figure S5. TXNDC5-driven endothelial paracrine signaling and ECM remodeling.

1186

A, Representative EdU staining image and group data showing the proliferative activity

1188

of hPASMCs treated with conditioned medium from hPAECs transfected with

1189

siTXNDC5 (siTXN-CM) or siNC (siNC-CM) in normoxia or hypoxia conditions (n=6).

Nuclei were counterstained with DAPI (blue). Scale bars=200 μm. **B**, CCK-8 assays demonstrating decreased cell viability of hPASMCs after treatment with siTXN-CM or siNC-CM (n=6). **C**, Quantification of POSTN, BGN, LOX, and TGM2 deposition in distal PAs from SuHx-treated TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice, assessed by mean fluorescence intensity (MFI) (n=6). **D**, **E**, and **F**, Representative dot blot analysis of BGN, POSTN, and LOX in the supernatants of hPAECs transfected with siTXNDC5 or control siRNA (siNC). **G**, Quantification of dot blot intensities for BGN, POSTN, and LOX, showing significantly reduced secretion upon TXNDC5 knockdown in normoxia or hypoxia conditions (n=5). **H**, Venn diagram showing that BGN exists in the overlapping of the core ECM genes in the PPI network, the differentially expressed ECM proteins in TXNDC5 silencing hPAECs, and the differentially expressed ECM genes in hypoxia-treated hPAECs (GSE160255). **I**, Representative Western blots and quantification of BGN and vinculin in the lungs of donors and PH patients (n=7). **J**, Representative Western blots and quantification of BGN and β-actin in the lungs from mice exposed to SuHx or under normoxic conditions (n=6). **K**, Western blots showing the BGN protein levels in the lungs collected from MCT or SuHx-treated rats or rats under normoxia (n=7). **L**, Representative EdU staining image and group data showing 10 μg/ml recombinant human biglycan (rhBGN) treatment for 24 h enhanced the proliferative activity of hPASMCs (n=5). Scale bars=200 μm. **M**, CCK-8 assay confirming increased cell viability of hPASMCs after rhBGN treatment (n=6). **N**, Western blots showing the protein levels of cleaved-caspase 3 of hPASMCs treated with 10 μg/ml rhBGN or the negative control (bovine serum albumin, BSA) for 24 h (n=6). **O**, Flow cytometry analysis of Annexin V-FITC/PI staining demonstrating decreased apoptotic cell populations in hPASMCs treated with rhBGN (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**I-O**) or 2-way ANOVA (**A-C** and **G**), followed by Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; MCT, monocrotaline; si, small interfering; DAPI, 4′,6-diamidino-2-phenylindole; f/f, flox/flox; POSTN, periostin; BGN, biglycan; and LOX, lysyl oxidase; TGM2, transglutaminase 2.

1220

Figure S6. Endothelial BGN deficiency protected against pulmonary vascular remodeling and RV failure in PH.

A, Schematic illustrating the generation of endothelial cell-specific BGN knockout (BGN^{ECKO}) mice. As BGN is an X chromosome-located gene, male mice carry a single floxed allele (BGN$^{f/0}$), whereas female mice carry double floxed alleles (BGN$^{f/f}$). **B**, Representative Western blots and quantification of BGN protein levels in primary lung ECs isolated from BGN$^{f/0(f/f)}$ and BGN^{ECKO} mice (n=6). **C**, Quantification of collagen

deposition of distal pulmonary arteries (PAs) from male BGN$^{f/0}$ and BGN^{ECKO} mice under normoxia or SuHx(n=8). **D**, **E**, **F**, and **G**, Echocardiographic assessment of right ventricular function in male BGN$^{f/0}$ and BGN^{ECKO} under normoxia or SuHx, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=8). **H**, Right ventricular systolic pressure (RVSP) in BGN$^{f/f}$ and female BGN^{ECKO} under normoxia or SuHx (n=6). **I**, Fulton index [RV/(LV+S)] in BGN$^{f/f}$ and BGN^{ECKO} mice (n=6). **J**, **K**, **L,** and **M**, Echocardiographic evaluation of PAT/PET, TAPSE, RVFAC, and RVFWT in female BGN^{ECKO} and littermate control mice (n=6). **N**, Representative distal PA images from BGN$^{f/f}$ and BGN^{ECKO} exposed to normoxia or SuHx, showing α-SMA (green) and VB/SR staining of elastic lamina (dark blue), collagens (red), and muscle fiber (yellow). Nuclei stained with DAPI (blue). L indicates lumen. Scale bars=25 μm. **O**, **P**, and **Q**, Quantitative analysis of distal PA remodeling in BGN$^{f/f}$ and female BGN^{ECKO} mice, including PA muscularization, medial wall thickness, and collagen deposition (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**B**) or 2-way ANOVA (**C-M,** and **O-Q**), followed by Tukey honestly significant difference test for multiple comparisons. DAPI indicates 4′,6-diamidino-2-phenylindole; α-SMA, α-smooth muscle actin; f/0, flox/0; f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio; TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional area change; and RVFWT, right ventricular free wall thickness.

1249

Figure S7. TXNDC5 stabilized BGN to regulate its protein expression.

A, Representative Western blots and quantification of TXNDC5 and BGN in hPAECs

treated with siNC or siTXNDC5 under normoxia or hypoxia conditions (n=6). **B**,

Western blot analysis showing the downregulation of BGN in the lungs from TXNDC5

deficiency mice compared with WT mice (n=6). **C**, Representative Western blots and

quantification of TXNDC5 and BGN in lung ECs isolated from TXNDC5$^{f/f}$ and

TXNDC5^{ECKO} mice under normoxia or SuHx conditions (n=6). **D**, Representative

Western blots showing that the LV-TXNDC5-induced upregulation of BGN in hPAECs

both under normoxia and hypoxia conditions (n=6). **E**, Western blot analysis showing the upregulation of BGN in the lung ECs isolated from endothelial-specific AAV-NC or AAV-TXN-treated mice exposed to SuHx or under normoxic conditions (n=6). **F**, Coimmunoprecipitation analysis showing the interaction between TXNDC5 and BGN was enhanced by hypoxia stimulation in hPAECs. IP: BGN, immunoprecipitation with BGN antibody. IgG for the negative control. **G**, Scheme of WT (TXNDC5 WT) and mutant TXNDC5 (TXNDC5 Mut) sequences with Cysteine (C)-to-Alanine (A) mutations in all three thioredoxin domains. **H**, Schematic illustration for using FRET to determine the extent of BGN protein folding in hPAECs. When the BGN protein folds correctly (bottom), the distance between CFP and YFP will be less than 10 nm, resulting in FRET occurring. **I**, Western blot showing that TXNDC5 overexpression prolonged the BGN protein half-life in hPAECs (n=6). **J**, Western blot analysis demonstrating that TXNDC5 silencing destabilized BGN in hPAECs, resulting in accelerated protein degradation (n=6). **K**, **L**, **M**, and **N**, Echocardiographic evaluation of right ventricular function in $BGN^{f/0(f/f)}$ and BGN^{ECKO} mice with AAV-NC or AAV-TXN treatment (n=6), including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). Data represent the mean±SD. Comparisons of parameters were performed with 2-way ANOVA, followed by the Tukey honestly significant difference test for multiple comparisons. Nor indicates normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypoxia; IgG, immunoglobulin G; LV, lentivirus; WT, wild-type; Mut, mutation; CHX, cycloheximide; FRET, fluorescence resonance energy transfer; CFP, cyan fluorescent protein; and YFP, yellow fluorescent protein.

1280

Figure S8. HIF signaling mediated TXNDC5-driven PH.

A and **B**, qPCR analysis showing the effects of HIF-1α (A) or HIF-2α (B) silencing on

the mRNA expression of TXNDC5. Average expression values are normalized to the β-

actin mRNA level. **C**, Western blot showing that siHIF-2α reduced the protein levels of

TXNDC5 in hPAECs under hypoxic conditions (n=6). **D**, Schematic illustration showing 2 sites in HIF-2α predicted to bind to the TXNDC5 promoter region. **E**, Chromatin immunoprecipitation (ChIP) followed by polymerase chain reaction analysis showing that the BS1 and BS2 regions are both for HIF-2α binding. **F**, Western blot analysis showing that silencing of TXNDC5 reduced the protein levels of HIF-1α and HIF-2α in hPAECs under hypoxic conditions (n=6). **G**, Western blots showing that siTXNDC5 reduced the protein levels of HIF-1α and HIF-2α in the nucleus exposed to hypoxia in hPAECs (n=6). **H**, Immunofluorescent staining analysis showing that siTXNDC5 reversed the increased levels of HIF-1α (green, left) or HIF-2α (green, right) induced by hypoxia in hPAECs. Nuclei were counterstained with DAPI (blue). Scale bars=50 μm. **I**. Schematic diagram of the pharmacological rescue experiment. AAV-NC or AAV-TXN was delivered via the tail vein to overexpress TXNDC5 in lung endothelial cells of mice, followed by normoxia or SuHx exposure, and then with HIF-1α inhibitor (PX-478) or HIF-2α inhibitor (PT-2385), or vehicle treatment. **J**. Hemodynamic measurements showing right ventricular systolic pressure (RVSP) in mice treated as described in (I) (n=6). **K**, Fulton index [RV/(LV+S)] of mice, assessing right ventricular hypertrophy (n=6). **L**, **M**, **N,** and **O**, Echocardiographic evaluation of right ventricular function, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**G**) or 2-way ANOVA (**A-C**, **F,** and **J-O**), followed by Tukey honestly significant difference test for multiple comparisons. BS indicates binding site; Nor, normoxia; Hyp, hypoxia; SuHx, Sugen5416/hypixa; ChIP, chromatin immune precipitation; PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio; TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional area change; and RVFWT, right ventricular free wall thickness.

1310

Figure S9. TXNDC5 inhibited the ubiquitination of HIF-1α/2α in a BGN-dependent manner.

A, Representative Western blots and quantification of BGN, and HIF-1/2α in hPAECs treated with siNC or siBGN under normoxia or hypoxia conditions (n=6). **B**, Western blot showing that the LV-TXNDC5-induced upregulation of HIF-1/2α in hPAECs both under normoxic and hypoxic situations (n=6). **C**, Representative Western blots and quantification of TXNDC5, HIF-1α, and HIF-2α in the lung ECs isolated from AAV-TXN or AAV-NC treated mice exposed to SuHx or under normoxic conditions (n=6). **D**, Quantification analysis showing that the protein half-life of endogenous HIF-1/2α was prolonged by LV-TXNDC5 treatment in hPAECs under normoxic conditions (n=5). **E**, Western blot analysis indicated that the HIF-1α ubiquitylation profile was reduced

Figure S10. TXNDC5 inhibited the ubiquitination of HIF-1α/2α in a BGN-dependent manner.

A, Representative H&E staining of major kidney, liver, and spleen from rats maintained under normoxia for 5 weeks and continuously treated with E64FC26 or vehicle, showing preserved tissue morphology without apparent injury or inflammation. Scale bars=100 μm. **B**, Gain of body weight of female (top panel) and male (bottom panel) rats maintained under normoxia for 5 weeks with E64FC26 or vehicle treatment (n=6). **C**, **D**, and **E**, Quantification of distal pulmonary arteries (PAs) remodeling in SuHx-induced PH rats treated with vehicle or E64FC26, including distal PA muscularization, medial wall thickness, and collagen deposition area (n=6). **F**, Echocardiographic assessment of right ventricular function in rats with vehicle or E64FC26 treatment, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). **G** and **H**, Quantification of distal PA medial wall thickness and collagen deposition area SuHx-treated TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice with or without E64FC26 administration under SuHx or normoxia conditions (n=6). **I**, Right ventricular systolic pressure (RVSP) in TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with vehicle or E64FC26 (n=6). **J**, Fulton index [RV/(LV+S)] of TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice with vehicle or E64FC26 treatment (n=6). **K**, Echocardiographic assessment of right ventricular function in TXNDC5$^{f/f}$ and TXNDC5^{ECKO} mice treated with vehicle or E64FC26 exposed to SuHx or normoxia, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). **L**, Western blot showing efficient knockdown of TXNDC5 in lung ECs isolated from rats receiving pulmonary endothelium-targeted self-complementary adeno-associated virus (scAAV) carrying shRNA against TXNDC5 (shTXN) compared with control shRNA (shNC) (n=6). **M**, **N**, and **O**, Quantification of distal PA remodeling in SuHx-induced PH rats treated with shNC or shTXN, including distal PA muscularization, medial wall thickness, and perivascular collagen deposition area (n=6). **P**, Fulton index [RV/(LV+S)] in rats treated with shNC or shTXN under normoxia or SuHx conditions (n=6). **Q**, Echocardiographic assessment of right ventricular function in rats receiving shNC or shTXN, including PAT/PET, TAPSE, RVFAC, and RVFWT (n=6). Data represent the mean±SD. Comparisons of parameters were performed with the unpaired 2-tailed Student *t*-test (**L**) or 2-way ANOVA (**B-K**, and **M-Q**), followed by Tukey honestly significant difference test for multiple comparisons. H&E indicates hematoxylin-eosin;

1361   f/f, flox/flox; Nor, normoxia; SuHx, Sugen5416/hypoxia; shRNA, short hairpin RNA;

1362   PAT/PET, pulmonary artery acceleration time/pulmonary artery ejection time ratio;

1363   TAPSE, tricuspid annular plane systolic excursion; RVFAC, right ventricular fractional

1364   area change; and RVFWT, right ventricular free wall thickness.