Pangenome Analysis of *Proteus mirabilis* Reveals Lineage-Specific Antimicrobial

Resistance Profiles and Discordant Genotype-Phenotype Correlations

Authors: Namrata Deka[1], Aimee L. Brauer[1], Katherine Connerton[1,2], Blake M.

Hanson[3,4], Jennifer N. Walker[5,3,4], Chelsie E. Armbruster[1*]

*address correspondence to Chelsie E. Armbruster, chelsiea@buffalo.edu

[1]Department of Microbiology and Immunology, Jacobs School of Medicine and

Biomedical Sciences, Buffalo, NY, 14203, NY, USA. [2]School of Nursing, Niagara

University, Buffalo, NY, USA. [3]Department of Epidemiology, UTHealth Houston School

of Public Health; [4]Center for Infectious Diseases, UTHealth Houston School of Public

Health; [5]Department of Microbiology and Molecular Genetics, McGovern Medical

School, The University of Texas Health Science Center at Houston, Texas, USA.

Author Contributions: ND and CEA designed experiments. ND, KC, and ALB
conducted MIC experiments, ND and CEA analyzed results. ND performed genome
sequencing, pangenome analysis, and AMR detection. BH and JW provided guidance
on pangenome analysis, AMR detection, and data visualization. ND visualized the data
and prepared the first draft of the manuscript. CEA provided project administration. All
authors critically reviewed and revised the manuscript.
22

23 **Conflict of Interest:** All authors declare no conflicts of interest.

24 **KEYWORDS:** *Proteus mirabilis*; antimicrobial resistance; pangenome; antimicrobial
25 susceptibility testing; multi-locus sequence typing; trimethoprim; sulfamethoxazole;
26 chloramphenicol; urinary tract infection

ABSTRACT (<250 words)

Urinary tract infections (UTIs) impose a large healthcare burden, with escalating

antimicrobial resistance (AMR) and treatment failure. *Proteus mirabilis* is an under-

characterized and challenging UTI pathogen due to intrinsic resistances and biofilm

formation. To understand *P. mirabilis* population genomics, we combined pangenome

analysis, *in silico* AMR predication, and phenotypic antimicrobial susceptibility testing

(AST) across 1,027 *P. mirabilis* genomes derived from human urine specimens. This

revealed a mosaic pangenome driven by extensive accessory genome plasticity.

Multilocus sequence typing (MLST) identified 213 MLSTs with only 7% having ≥10

genomes, highlighting strain diversity. AMR gene profiles were largely lineage specific,

with 25% of genomes harboring resistances for >6 antimicrobial subclasses. ST135 was

identified as a highly MDR lineage, with 95% of genomes carrying ≥16 resistance

genes. Mobile genetic element (MGE) analysis of 22 clinical isolates with complete,

reference level genomes revealed that Tn7 transposons, IS26-mediated genomic

islands, and class 1 integrons act as vehicles for high AMR gene dissemination,

including IS26-mediated gene stacking within a *Proteus mirabilis* Genomic Resistance

Island 1 (*PmGRI1*) in ST135 isolates. While presence of genes like *aph(3')-la* reliably

predicted kanamycin resistance, discordance for antibiotics such as trimethoprim-

sulfamethoxazole and chloramphenicol revealed that AMR gene stacking, regulatory

context, and intrinsic mechanisms, like efflux pumps, modulate phenotypic outcomes. In

summary, our study provides a comprehensive and phenotypic resolution of *P.*

mirabilis AMR, establishing that resistance architecture is lineage structured, MGE-

driven, and phenotypically non-deterministic. We emphasize the need to shift towards

50   standardized, genome-informed surveillance framework to translate into diagnostic and

51   therapeutic strategies.

INTRODUCTION

Urinary tract infections (UTIs) are among the most common bacterial infections, with an estimated healthcare cost of >$3 billion in the United States [1-4]. They are a leading cause of antimicrobial prescriptions in both outpatient and inpatient settings, with 50-60% of women experiencing at least one UTI in their lifetime [5]. UTI can be classified as complicated UTI (cUTI) or uncomplicated UTI (uUTIs) depending on the presence and absence of structural or functional abnormalities of the urinary tract, catheterization, or immunocompromised states [5]. Recent epidemiological data estimate approximately 174,000 cases of uUTI and 2.38 million cases of cUTI annually [6, 7]. Despite standard antibiotic therapy, treatment failure rates for UTIs remain prevalent, occurring in 16.7% of cases in adult female outpatients with cUTI, 27% of elderly men, and 15% of elderly women, often leading to recurrence, secondary infection, progression to pyelonephritis, or urosepsis [8, 9].

Among uropathogens, *Proteus mirabilis* is an important Gram-negative bacterium and a major cause of UTIs, contributing to approximately 4.2% of uUTIs and 4.7% of cUTIs [5]. Notably, *P. mirabilis* accounts for up to 45% of catheter-associated UTIs (CAUTIs) and 13-21% of secondary bacteremia, making it a key pathogen in medical device-associated infections [5, 10, 11]. *P. mirabilis* has a strong propensity for biofilm formation and causes encrustation of the catheter by deposition of struvite and apatite crystals, leading to persistent infections and catheter blockage [10, 12-15]. These urinary stones can remain in the bladder as crystalline deposits despite multiple doses of antibiotic treatment, catheter changes, and even periods without catheterization [15-17]. Infection with *P. mirabilis* can become fatal when it progresses to symptomatic

CAUTI, acute bacteremia, and urosepsis due to its ability to disseminate from the

bladder to other organs [14, 18].

Antimicrobial resistance (AMR) in UTI pathogens is an escalating global concern,

with increasing reports of multidrug-resistant (MDR) strains complicating treatment [19].

Treatment failure for *P. mirabilis*-associated UTIs has been linked to MDR, leading to

prolonged hospitalization, increased healthcare costs, and higher morbidity [15, 19, 20].

While antibiotic therapy remains the primary treatment approach, its efficacy is

increasingly compromised by both intrinsic and acquired resistance mechanisms that

limit therapeutic options [14, 21-23]. The resistance to first-line UTI antibiotics such as

trimethoprim-sulfamethoxazole (TMP-SMX), fluoroquinolones, and β-lactams increases

and the prevalence of ESBL genes like *blaTEM* and *blaCTX-M* further exacerbates

resistance. Among the key drivers of AMR dissemination in *P. mirabilis* are mobile

genetic elements (MGEs) such as SXT/R391 integrative and conjugative elements

(ICEs) [19, 24, 25]. However, characterization of MGEs in *P. mirabilis* UTI clinical

isolates remains limited. Understanding the MGE landscape is essential for deciphering

the mechanisms underlying horizontal gene transfer, the evolution and spread of MDR,

and the discordance between genotypic predictions and phenotypic resistance.

Numerous studies have characterized the model *P. mirabilis* strain HI4320,

originally isolated from the female urinary tract, and its pathogenic potential [10, 26].

However, a major gap remains in our understanding of the diversity of strains isolated

from recent clinical urine samples, either during infection or in the presence of a

catheter. Multilocus sequence typing (MLST) provides an unambiguous method for

characterizing bacterial isolates based on the sequences of internal fragments of

typically 6-7 housekeeping genes. While previous studies in other bacterial species

suggest that certain lineages correlate with varying levels of resistance and biofilm

production, such associations in *P. mirabilis* remain largely unexplored. For example, in

methicillin-resistant *Staphylococcus aureus* (MRSA), expression of the *ica* genes—

critical for biofilm formation—varies among different STs, with certain STs showing

higher *ica* gene expression, particularly in strains isolated from wound and catheter

samples [27]. If similar correlations exist in *P. mirabilis*, routine sequence typing of

clinical isolates could serve as a valuable tool to predict AMR potential and

pathogenicity.

Several typing schemes have been developed to characterize *P. mirabilis* strains,

but no standard scheme is followed for clinical classification. Core genome MLST

(cgMLST), a high-resolution genomic method, has identified 205 clonal groups (CGs)

linked to severe UTIs and AMR [28]. Historically, *P. mirabilis* strain differentiation has

relied on methods such as bacteriophage typing, serotyping, and Dienes typing, but

genomic approaches now offer superior discriminatory power [29-33]. Even with the

established role of *P. mirabilis* in CAUTI, the publicly available genome database is

limited compared to other UTI pathogens [32, 34, 35]. Furthermore, there remains a

lack of integrated data linking sequence types (ST) to AMR gene carriage in urinary

isolates using a standard classification framework. To address this gap, we focused

specifically on genomes isolated from human urine, enabling a targeted analysis of

AMR profiles and providing insights that may inform risk stratification for treatment

failure and severe disease.

There are numerous ways to predict AMR, including the standardized clinical

laboratory practices of testing resistance phenotypes to specific antibiotics, as well as

genomic or machine learning methods to predict resistant gene expression [36].

Genomic prediction tools can be useful for quick detection of AMR in clinical practice,

and there may be a correlation between AMR gene prediction and phenotypic

expression, but the nature of the correlation can vary depending on specific genes and

the bacterial species. For instance, *Escherichia coli* frequently harbors resistance genes

for β-lactams, sulfonamides, trimethoprim, and chloramphenicol on mobile genetic

elements, leading to a stronger correlation between genotype and phenotype [37-39]. In

contrast, according to some studies, *Proteus mirabilis* carrying β-lactam and

sulfonamide (*sul1* and *sul2*) genes exhibits this correlation less consistently, showing

weaker alignment between genomic predictions and observed resistance profiles [37,

40-42].

Tools like AMRFinder and CARD are accurate *in silico* prediction tools;

AMRFinder was able to validate the presence of genes pertaining to AMR phenotypes

for *Salmonella enterica, Campylobacter spp.*, and *E. coli* with 98.4% consistent

predictions [43]. Further, a 99% concordance was found between phenotypic

antimicrobial susceptibility testing and *in silico* prediction of AMR genes using whole-

genome sequencing (WGS) in 1,321 isolates of *Salmonella enterica* from human

infections in Canada [20]. While WGS-based methods provide a promising avenue to

predict AMR in other species, no equivalent variation exist for *P. mirabilis*. Hence,

integrating genotypic AMR prediction and phenotypic MIC testing in diverse strains may

offer a more comprehensive approach to predicting antimicrobial resistance in this

species.

In this study, we conducted a comparative genomic analysis of *P. mirabilis* using

1,001 publicly available human urine isolate sequences to characterize the pangenome,

AMR gene profiling, and *in-silico* MLST typing. We benchmarked this against 26 clinical

isolates collected from patients with long-term indwelling catheters [15, 17]. Our

analysis revealed that AMR gene profiles can link to specific lineages rather than

isolation source, with isolates of the same MLST clustering together and exhibiting

related resistant genes. We found that ST135 lineage is a hotspot for resistance gene

acquisition in our dataset, driven by the integration of MGEs such as IS26 and the

PmGRI1 genomic island with multiple overlapping resistance genes. We further

identified a high rate of discordance between genotype-based susceptibility prediction

and phenotypic resistance. This discrepancy can be partially attributed to

uncharacterized MGE's and efflux pump mechanisms, highlighting the limitations of

relying on gene detection only for resistance profiling. Our findings underscore the need

for integrating lineage-specific MGE surveillance with phenotyping testing to guide

treatment options and reduce the overuse of broad-spectrum antibiotics in managing *P.*

mirabilis CAUTI infections.

MATERIALS AND METHODS

Data availability statement

Sequencing data have been deposited in NCBI BioProject PRJNA1367153: *Proteus mirabilis* genome sequences from human urine sources in the United States. All other data are available within the article and its supplemental material. All basic commands and databases used are available in the supplementary appendix.

Ethics approval

This body of work includes sequencing of *P. mirabilis* strains that were previously isolated from human subjects in two different studies, for which human subject protocols were approved by the University of Buffalo (IRB #STUDY00002526) and Washington University School of Medicine (IRB #201410058).

Patient population and Sample collection

To investigate whether *P. mirabilis* MLSTs correlate with AMR patterns, we assessed *P. mirabilis* genomes from three separate cohorts: publicly available genomes from NCBI (**Table S1**), nursing home residents with long-term catheters (CEA isolates), and community-dwelling catheterized patients (HUC isolates).

For comparison of *P. mirabilis* genomes, we sought to query all publicly available high-quality genomes of *P. mirabilis* isolates obtained from urine samples in hospital environments in the United States. Using advanced search parameters, publicly available raw genome data were retrieved from the NCBI database. The initial search,

conducted in October 2024, yielded 1,337 hits using the terms "*Proteus mirabilis*,"

isolation source "urine," and host "*Homo sapiens.*" Filtering the results to include only

isolates originating from the "United States" reduced the dataset to 1,152 genomes. The

dataset was further refined by only including whole-genome sequencing data. These

filters resulted in a final dataset of 1,064 (**Table S1**) genomes, which were subsequently

downloaded from NCBI. Following FastQC v0.11.9-Java-11 analysis and adapter

trimming using trimmomatic v0.39-Java-11.0.16. [44, 45], 1,026 genomes were retained

for downstream analyses. Samples were excluded if they exhibited poor quality, had

high GC content, high read duplication rates, or if paired end read files were missing (46

samples). Raw paired end read sequences were assembled using spades v3.15.5 [46].

After genome assembly, we assessed quality and completeness using QUAST (**Table

S2**), selecting 1,001 high-quality genomes with >95% completeness and fewer than

1,000 contigs which were annotated with prokka v1.14.5 to facilitate downstream

analyses [47, 48].

Isolates designated 101–210 were cultured from urine specimens collected

during a prospective cohort study on asymptomatic catheter-associated bacteriuria at

two nursing homes in Buffalo, New York, USA, between July 2019 and March 2020 [15].

Inclusion criteria required participants to be 21 years or older at the time of consent and

have an indwelling urinary catheter in place for at least 12 months at the start of the

study. Study participants underwent an initial enrollment visit, followed by weekly study

visits for up to seven months. Most of the study participants were White (79%) and male

(79%), an average age of 65 years, and 68% had suprapubic catheters [15].

Participants exhibited a high degree of functional dependence in activities of daily living,

with the most common comorbidities including neurogenic bladder (74%), hemiplegia

(42%), diabetes (32%), renal disease (32%), multiple sclerosis (26%), and chronic heart

failure (26%) [15]. Urine specimens were cultured on MacConkey agar and Columbia

Nutrient Agar, and Gram-negative bacteria were identified to the species level using

API-20E test strips (BioMérieux). *P. mirabilis* species designation was further confirmed

based on characteristic swarming motility on blood agar plates (Hardy Diagnostics) and

whole genome sequencing (**Table S3**).

Additional *P. mirabilis* isolates with designations of "HUC" were obtained from the

lab of Scott J. Hultgren, Washington University School of Medicine [17]. These isolates

were derived from urine samples collected from community-dwelling catheterized

patients during routine catheter removals in the urology clinic as part of their standard

care. Most of the participants were male (64%) with an average age of 64 years.

Inclusion criteria required participants to be 18 years or older at the time of consent and

scheduled for the removal of an indwelling urinary device, including urethral and

suprapubic catheters or ureteral stents. The primary indications for catheterization

included bladder outlet obstruction (33%), peripheral or central neuropathy (40%),

rectovesical or rectourethral fistula (7%), stress urinary incontinence (7%), atonic

bladder (11%), and temporary post-operative complications (2%) [17]. Bacterial colonies

were distinguished based on morphology, size, and color, and 1–9 single colonies per

sample were selected for DNA extraction. *P. mirabilis* isolates were identified via 16S

rRNA sequencing and confirmed by whole genome sequencing (**Table S3**).

Whole Genome Sequencing of Clinical Isolates

Freezer stocks were streak-plated on MacConkey agar to isolate single colonies of

Proteus mirabilis. 5 ml of Low Salt Luria Bertani (LSLB; 0.1 g/L NaCl, 10 g/L Tryptone, 5

g/ yeast extract) broth was inoculated with a single colony and incubated for 16 hours at

37°C with shaking. Genomic DNA was extracted using the DNeasy Blood and Tissue

Kit (Qiagen, Germantown, Maryland, USA) according to the manufacturer's instructions

with specific modifications to ensure high purity of DNA; samples were incubated in

proteinase K for 2 hours instead of 10 minutes, and three washes of AW2 buffer were

conducted to ensure high-quality DNA free of salts. Samples were sent to SeqCoast

Genomics for whole genome sequencing (WGS). All samples were prepared using the

Illumina DNA Prep tagmentation kit and IDT For Illumina Unique Dual Indexes.

Sequencing was performed on the Illumina NextSeq2000 platform using a 300-cycle

flow cell kit to produce 2x150bp paired reads, as previously described [49]. 1-2% PhiX

control was spiked into the run to support optimal base calling. The sequencing data

has been uploaded in BioProject PRJNA1367153: *Proteus mirabilis* genome sequences

from human urine sources in the United States. Read trimming and run analytics were

performed using trimmomatic v0.39.0 [45]. For long read sequencing, Oxford Nanopore

Technologies Native Barcoding Kit (#SQK-NBD114) and long fragment buffer was used

to promote longer read lengths of DNA. DNA was sequenced on the PromethION 2

Solo platform using a FLO-PRO114M Flow Cell (vR10.4.1). Sequencing was performed

with a translocation speed of 400bps. Base calling was done on the GridION using the

super-accurate basecalling model with barcode trimming enabled.

Assembly of short reads from the clinical isolates and the genomes acquired

from the NCBI database was performed using spades v3.15.5, while hybrid assembly of

long and short reads from clinical isolates was performed using Unicycler v0.5.0 [46].

Here, long reads were used to assemble the genome structure and the short reads

were used to polish the genome. The assemblies were quality-checked using QUAST

v5.2.0 selecting genomes with < 500 contigs for only short read assembled genomes

and <3 contigs for hybrid (**Table S3**) If sequences were not of good quality, they were

further polished using pilon/1.23-Java-11.0.16 [50]. Genomes that passed quality

filtering were annotated using prokka v1.14.5 [48].

Multi Locus Sequence Typing (MLST) and Phylogenetic trees

Multilocus sequence typing (MLST) analysis was conducted to identify the

sequence types (STs) present in the dataset using mlst software, which incorporates

the PubMLST database [51, 52]. The annotated genomes were then fed into roary

v3.13.0 using a core threshold cluster of 99%, to identify their pangenome profile and

develop a gene presence/absence matrix [53]. The filtered core genome alignment was

constructed using FastTree v2.1.11. The resulting newick file was visualized as a mid-

rooted phylogenetic tree using the iTOL website and annotated with source, and

antibiotic metadata files [54, 55].

Drug Resistance Gene Prediction

AMRFinderPlus (using the Bacterial Antimicrobial Resistance Reference Gene

Database) and CARD (The Comprehensive Antibiotic Resistance Database) were used

to predict AMR genes from the coding sequence of the *P. mirabilis* genomes including

the 26 CEA and HUC strains and clinical reference strain HI4320 [36, 43].

Mobile genetic elements were characterized in the hybrid-sequenced clinical

isolates using multiple approaches. Insertion sequence (IS) elements were identified

using ISEScan v1.7.3 [56]. Prophage regions were predicted using PHASTER (PHAge

Search Tool Enhanced Release) [57-59] and classified as intact (score >90),

questionable (70-90), or incomplete (<70). Plasmid replicons were detected using

ABRicate v1.0.1 with the PlasmidFinder database (≥80% identity, ≥60% coverage) [60,

61]. Integrative and conjugative elements (ICEs) were identified via ICEberg 3.0 [62].

Spearman's rank correlation in R v4.3.0 assessed associations between IS abundance

and AMR gene content and virulence factors with significance set at $p < 0.05$.

Antibiotic Susceptibility Testing (AST)

The clinical strains were grown overnight in 5 mL LSLB broth at 37°C with shaking.

Antibiotic solutions: Tetracycline HCl (Research Products International (RPI), CAS: 64-

75-5), Chloramphenicol (RPI, CAS: 56-75-7), Streptomycin (RPI, CAS: 3810-74-0),

Streptothricin (Goldbio, CAS: 96736-11-7), Sulfonamide (sulfamethoxazole, Cayman

chemical company, CAS: 723-46-6), Trimethoprim (Cayman chemical company, CAS:

738-70-5), Ampicillin (RPI, CAS: 69-53-4), and Kanamycin (RPI, CAS: 8063-07-8), were

prepared in Mueller-Hinton Broth (MHB) at a concentration range encompassing the

sensitive-to-resistant spectrum according to Clinical Laboratory Standards Institute

(CLSI) guidelines for *Enterobacteriales* [63]. Stock solutions of all the antibiotics were

prepared and stored at –20°C, and working dilutions were made fresh from the stock

immediately prior to testing.

1 ml of the overnight cultures were centrifuged at 1,000g to pellet the cells. The

supernatant was removed, and the pellets were resuspended in 0.9% saline to wash

away residual LB. The inoculum was adjusted to 4×10^5 CFU/ml in MHB. One antibiotic

with increasing concentrations was tested per 96-well plate, with four technical and

duplicate biological replicates. The plates were incubated at 37°C with double-orbital

shaking in Biotech Synergy H1, and the OD_{600} was measured every 15 minutes for 20

hours to monitor bacterial growth with OD_{600} threshold set at 0.15.

Statistical analysis

Logistic regression models were used to explore AMR gene carriage as a

function of sequence type, geographic location of isolation, and year of isolation. Data

were analyzed using Stata/IC version 15.1 (StataCorp LP, College Station, TX).

RESULTS

To assess the population structure and genetic diversity of *Proteus mirabilis* in the urinary tract, we analyzed 1,001 publicly available genomes isolated from human urine specimens (**Table S1**). The genomes downloaded from NCBI were predominantly derived from clinical settings, focusing on medical surveillance and characterization of urinary samples.

Key contributing sources include research from Washington University School of Medicine (WUSM), the University of Pittsburgh Enhanced Detection System for Healthcare-Associated Transmission (EDS-HAT) study, and national antimicrobial surveillance programs such as the Center of Disease Control's Healthcare-Associated Infection Sequencing (CDC HAI-Seq) Gram-Negative Bacteria Study and studies on the human urinary microbiome. Among the genomes, contig counts ranged from 60 to 973, with an average of 258 and a median of 180 contigs. The GC content averaged 38.76%, with a median of 38.69%, making them suitable for downstream analyses (**Table 1 and S2**). While all sequences were confirmed to be derived from urine samples, the clinical context of these specimens—whether they were associated with asymptomatic bacteriuria or infection—is not known.

Phylogenetic analyses (mid-rooted tree) revealed a monophyletic, star-like phylogeny, with a central ancestral node from which several peripheral branches radiate (**Fig 1a**), which may indicate recent diversification events. When considering branching, a subset of genomes branched out creating three clades, *clade 1* (964 genomes, dominant), *clade 2* (5 genomes), and *clade 3* (32 genomes) (**Fig S1**). This breakdown is consistent with previously reported intra-species core genome variation in *P. mirabilis*

[32, 33, 35]. For example, Potter *et. al.* studied a cohort of 2,060 *P. mirabilis* genomes

collected worldwide from multiple anatomical sources and observed that the essential

core genes remained relatively conserved with three divergent sub-species. [32].

Pan-genomic analysis of *P. mirabilis* reveals a mosaic structure

To characterize the pangenome of *P. mirabilis* from human urine samples we

assessed its structural composition, including core, shell, and cloud gene categories. In

the 1,001 genomes analyzed, a total of 41,121 gene families were identified. Of these,

the core genome (genes conserved in >99% of the samples analyzed) is small with

1,937 genes indicating a high degree of conservation. Considering that a typical *P.*

mirabilis genome harbors roughly 4,600-4,900 coding sequences (CDS), the core

genome accounts for approximately 40-45% of the average gene content. This indicates

a highly conserved genomic backbone with 4.7% of it supporting essential cellular

functions, coexisting with an expansive and dynamic accessory genome (**Fig S1**). The

shell of the 1,001 genomes contained 1,749 genes (4.08%), and 531 soft core genes

shared among a subset of strains that likely contribute to niche-specific adaptations.

The majority of genes (37,204 genes, 90%) were classified as cloud genes, making the

accessory component the largest in the pangenome. Having a small, stable core and

large, variable accessory genome likely represents strain-specific differences driven by

horizontal gene transfer.

The pangenome exhibited an open conformation representing a mosaic pattern

(**Fig S1**), with the size expanding as additional genomes were analyzed (**Fig 1b**). This

expansion reflects the continuous acquisition of novel gene families, highlighting the

dynamic nature of the *P. mirabilis* genome and its capacity for adaptation through

horizontal gene transfer (**Fig 1c**). The stability of core genes is crucial for the consistent

expression of virulence factors, while the expanding accessory genes may help adapt to

polymicrobial environments and/or antibiotic pressures. These findings indicate the

genetic plasticity of *P. mirabilis* isolated from urine and provide a foundation for future

studies on its pathogenic potential and AMR mechanisms in the urinary tract.

Publicly available *P. mirabilis* genomes comprise 213 defined and 118 undefined

MLSTs

The specific genes used in MLST of *P. mirabilis* are a combination of

housekeeping genes *atpA, dnaJ, mdh, pyrC, recA,* and *rpoD* [51, 52]. We identified a

total of 213 defined STs and approximately 118 novel or undefined genomes, reflecting

the diversity of *P. mirabilis* human urine isolates within the dataset (**Fig 1**). The untyped

MLST have been submitted to the PubMLST database to update the schema in *P.*

mirabilis.

A detailed analysis of ST prevalence within the dataset revealed that certain STs

were more frequently observed. The distribution of MLSTs across the dataset was

highly skewed (**Fig 2a, Table S4 and S5**), with 46.26% of the STs having only one

representative genome in the dataset while ~7% of the STs had ≥10 genomes (**Fig 2a**).

Many genomes in the same ST came from the same study/location but it was not the

rule as the top 15 overrepresented ST's show mixed isolation sources (**Fig 1a**). The

uneven ST distribution is further visualized in a bar chart of the top 20 MLSTs,

highlighting the dominance of a few clonal lineages across sampled populations (**Fig

2b). ST96 was the most abundant (49 genomes; 4.90%), followed by ST185 (42

genomes; 4.20%), ST99 (38 genomes; 3.80%), ST135 (38 genomes; 3.80%), and ST97

(37 genomes; 3.70%). As expected, genomes from each of these STs clustered

together by lineages, even if isolated from different geographical locations and reported

in different studies (**Fig 1a**).

Phylogenetic analysis reveals shared lineages and novel variants across clinical

***P. mirabilis* isolates**

To further examine the clinical relevance of *P. mirabilis* MLSTs, we next

sequenced and analyzed a collection of 25 clinical isolates from two distinct cohorts

(CEA isolates collected from the urine of nursing home residents with long-term

indwelling catheters, and HUC isolates collected from the urine of catheterized

community-dwelling individuals during routine catheter changes), as well as the

reference strain HI4320 that was previously isolated from the urine of a catheterized

nursing home resident (**Fig 3a**). The analysis revealed a population structure forming

two primary clades, termed *clade 1* (23 isolates*)* and *clade 2* (3 isolates) (**Fig 3b**). Lack

of the third clade identified in the 1,001 genomes was expected considering its low

frequency (0.5%) and the small sample size of the clinical isolate collection.

Phylogenetic clustering was in accordance with the core genome and gene

content irrespective of the clinical origin (HUC vs CEA, **Fig 3a**). Across the 26 isolates,

we identified 15 distinct STs; 10 STs were identified in the HUC cohort, 4 in the CEA

cohort, and 1 represented solely by HI4320. Six of the 10 overrepresented STs in the

NCBI genome dataset were also represented among the clinical isolates; namely STs

96, 97, 135, 145, 146, and 185. ST185 was observed in both the CEA and HUC

cohorts (H120-04 and 102-00, 206-16), demonstrating genetic overlap across different

geographic locations. The two most prevalent STs in the NCBI dataset were also the

most prevalent in the clinical isolates: ST185 (3 isolates, 12%) and ST96 (3 isolates,

12%). Their presence confirms that our clinical cohort captured high-impact,

epidemiologically relevant STs rather than rare or sporadically enriched variants even

with a limited sample size. The full summary of isolated sources and STs are provided

in **Table 2 and Table S3.**

Proteus mirabilis **urine isolates possess resistance factors to multiple subclasses**

of antibiotics

To understand how AMR complicates *P. mirabilis* UTI treatment and contributes

to the global public health burden, we examined the AMR profiles of the 1,001 publicly

available *P. mirabilis* genomes. A total of 24 antibiotic resistance subclasses were

identified, reflecting the diverse mechanisms by which this pathogen can evade

antibiotics. The most prevalent genes were those conferring resistance to tetracycline

and chloramphenicol, observed in nearly all genomes (**Table S6)**.

P. mirabilis is known to have intrinsic resistance to tetracycline, so it was not

surprising that tetracycline resistance genes were detected in 991 (99.0%) of the 1,001

genomes, with *tetJ* specifically encoded by 985 of those 991 (99.5%) (**Table 3**). Several

other tetracycline resistance determinants, including *tetA, tetB, tetC*, and *tetD*, were

detected at lower frequencies. Interestingly, 934/991 (94.2%) of the genomes carried

only a single *tet* gene, while 48 (4.8%) carried 2, and 9 (0.9%) carried 3 *tet* genes (**Fig**

4a). Notably, a small number of genomes (11, 1.1%) had no detectable tetracycline

resistance genes. This could represent a true absence of the genes, or it could be a

result of fragmented assemblies or sequence dissimilarity not found in the AMR

prediction database.

Chloramphenicol resistance was detected in 952 (95%) of the genomes, with 748

harboring the *catA* gene. Additionally, 69 genomes harbored both *catA* and *catA1,* 43

harbored *catA* and *floR,* and the presence of all three genes was seen in 15 genomes

(**Fig 4b, Table S6**). These findings highlight the prevalence of phenicol resistance in *P.*

mirabilis genomes despite limited use of chloramphenicol in modern clinical settings.

Resistance to aminoglycosides, a critical class of antibiotics for treating severe

Gram-negative infections, was predicted in 716 (71%) of the genomes. Notably, we

observed streptomycin resistance genes in 352 genomes, suggesting the persistence of

resistance despite limited clinical use (**Table 3**). The presence of multiple

aminoglycoside resistance determinants indicates that *P. mirabilis* maintains a robust

arsenal against this class of antimicrobials, including amikacin, gentamycin, kanamycin,

streptomycin and tobramycin.

Resistance to trimethoprim (TMP), mediated by *dfrA* alleles, was detected in 290

(29%) genomes. The most prevalent allele was *dfrA1* (n=248), followed by *dfrA17*

(n=44), and *dfrA32* (n=17). 235 (81.02%) genomes had a single *dfrA* gene, while 50

(17.2%) carried 2, and 5 (1.72%) carried 3 *dfrA* genes (**Fig 4c**). The diversity of *dfrA*

genes highlights the adaptive potential of *P. mirabilis* to acquire and maintain resistance

determinants. Resistance to sulfonamides (SMX) was similarly predicted in 286 (28%)

genomes, with *sul1* detected in 180, *sul2* in 187, and 81 genomes harboring both *sul1*

and *sul2* genes (**Fig 4d, Table S6**). Importantly, 223 of the 363 genomes encoding

resistance genes for either TMP or SMX had resistance determinants for both (**Fig 4e**).

Since TMP and SMX are often used in combination, the co-occurrence of these genes

would likely complicate efficacy of combined treatment. Additionally, we detected 185

genomes (18%) with Extended Spectrum β-Lactamase (ESBL)-associated genes. Of

these, 148 harbored the *blaTEM-1* gene, expected to confer resistance to ampicillin

(**Table 3**).

Finally, the number of resistance subclasses detected per *P. mirabilis* genome

ranged from 1 to 24, with an average of 4.61 AMR subclasses per isolate. A total of 71

genomes (7%) exhibited resistance to only a single antibiotic subclass, most commonly

tetracycline. In contrast, 931 (93%) had resistance to two or more subclasses, and a

substantial subset of 246 genomes (25%) harbored more than six antibiotic subclasses,

indicating a high prevalence of multidrug resistance in this sample set. 176 genomes

(18%) had at least one resistance determinant from all five of the above resistance gene

groups, while 56 (6%) had combinations of tetracycline, chloramphenicol, and β-lactam

resistance genes. Additionally, phylogenetic analyses (**Fig 1a**) determined that certain

clusters consistently encoded resistance genes to nearly all antibiotic classes shown,

such as genomes belonging to ST135, while others exhibited a complete absence of

resistance genes other than tetracycline, independent of study source.

Antibiotic resistance subclasses are lineage-associated but not strictly

determined by ST

To evaluate the relationship between ST and AMR burden in *P. mirabilis*, we

quantified the number of predicted AMR subclasses per genome. Logistic regression

confirmed that MLST was not a significant driver of high AMR carriage overall (OR

0.0997, 95% CI 0.992-1.002, p=0.304). However, striking lineage-specific enrichment

was observed with stratification by resistance burden. Hence, to enable robust

comparisons, we focused on a subset of 422 genomes representing the 16 STs with

≥10 genomes each. This accounted for 41% of the dataset (**Table S7, S8**). Within this

subset, the mean AMR subclass count was 4.4 per genome, with most genomes

carrying modest resistomes (80th percentile ~6 subclasses).

Genomes encoding resistances to ≥6 distinct subclasses were enriched in 5 STs

(121/142 [85%]): ST135 (37/37 genomes [100%]), ST185 (33/47 [80%]), ST279 (10/12

[83%]), ST97 (25/36 [69%]), and ST145 (16/32 [50%]) (Pearson χ^2 p<0.001). Among

these 142 genomes, 41 (29%) carried resistance genes to ≥12 subclasses, 33 of which

(80%) belonged to ST135. In contrast, ST145, ST93, ST33, and ST202 typically carried

2-3 resistance subclasses with some outliers, highlighting inter lineage heterogeneity

(**Table S8**). We acknowledge that sampling bias is possible as many surveillance

studies only pick up on hospitalized or severe infections, but our dataset also includes

general isolate submission from the CDC surveillance study, and we are most

interested to identify clinically relevant strains.

ST135 emerged as an exceptionally high-risk clone; 95% of ST135 isolates

(35/37) harbored ≥16 AMR genes, and 100% carried resistance to ≥6 subclasses. We

next examined whether geographic origin confounded this association and created a

bias. Within the 422-genome subset, 325 genomes had associated location data (US

states/cities) and 167 were from St. Louis. Although a slightly higher proportion of St.

Louis isolates exhibited high AMR carriage (≥10 subclasses) compared to other

locations (3.4% vs. 0.9%; Pearson χ² p=0.037), isolation source was not an

independent predictor in multivariable logistic regression controlling for MLST (OR 1.31,

95% CI 0.88–1.97, p=0.185). The strong enrichment of high-level AMR in ST135

regardless of geographical location points to lineage-intrinsic genetic factors as the

primary drivers.

The distribution of resistance gene burden is consistent across the dataset

To contextualize our 26 clinical isolates and HI4320 within the broader *P.*

mirabilis population, we compare their AMR gene content against the 1,001 NCBI

genomes. In total, resistance genes representing 14 different antibiotic subclasses were

identified in the clinical isolates compared to 24 from the NCBI dataset. The 26 clinical

isolates and Reference strain HI4320 (total 27) exhibited substantial variation in

resistance gene burden, ranging from 1 gene (minimum, observed in ST765 and ST591

isolates carrying only a tetracycline resistance determinant) to 15 genes (maximum, in

ST135 isolate H115-00, H115-29, and H127-00), with an average of 3.8 resistance

genes per isolate (**Table 3 and Table S9**).

Despite these differences in dataset size and diversity, several resistance traits

were highly conserved across both datasets. Tetracycline resistance approached

fixation; *tetJ* was present in all isolates predicted to be tetracycline resistant from both

datasets, and co-occurrence of *tetJ* with *tetA and tetC* (11%, 3/26 clinical) was also

observed. Similarly, chloramphenicol resistance genes demonstrated strong cross-

517 reservoir conservation, with *catA* (type-4) detected in 89% (23/26) of clinical isolates

518 compared to 95% of NCBI genomes, and 26% of clinical isolates (6/23) co-

519 harbored *catA1* while 11% (3/26) harbored no chloramphenicol resistance genes (**Table**

520 **S9**). ST135 strains also exhibited a rare combination of *floR* + *catB3*.

521  Fewer of the clinical isolates had resistance genes for cephalosporins or

522 quaternary ammonium in comparison to the NCBI dataset (**Table 3**). However, clinical

523 isolates showed enrichment of resistance genes under UTI treatment pressure including

524 TMP-SMX (22%, *sul1/sul2*; 37%, *dfr*) and aminoglycosides (streptomycin (37%, most

525 common: *aadA1*), streptothricin (37%, *sat2*), and kanamycin (26%, *aph(3')-Ia*) (**Table**

526 **3**). In terms of extended β-lactam ecology, clinical isolates predominantly

527 carried *blaTEM-1* (18%) while NCBI genomes exhibited broader cephalosporin

528 resistance genes (**Table 3, Table S6**). Collectively, these comparisons reveal

529 a conserved genetic backbone of core resistance determinants (tetracycline,

530 chloramphenicol) across *P. mirabilis* populations, upon which lineage-specific variation

531 is superimposed.

533 **The mobile genetic element (MGE) landscape of *P. mirabilis* reveals an IS26-**

534 **mediated major gene cluster in ST135.**

535  To determine if MGE's are associated with AMR gene dissemination in *P.*

536 *mirabilis*, we characterized 22 complete, polished reference quality genomes from the

537 clinical cohort. We carried out MGE analysis only using only complete, polished

538 genomes for high accuracy. MGE was not done for short read assembled genomes of

539 the NCBI database since incomplete genomes it is not feasible to get an accurate

picture from segmented genomes. Among the 22 isolates, a median of 22 IS elements

per genome was detected (mean=23.68; range 16-36) that were distributed across 17

IS families (**Fig 5a, Table S7**). The most abundant family was IS200/IS605 (48.5%),

followed by IS3 (16.31%), and ISNCY (7.2%). The three isolates with the highest total

IS count (36 each) were ST135 (H115-00, H115-29, and H127-00).

The high AMR gene count observed in the three ST135 isolates is positively co-

related with IS element count (**Fig. 5b**). A large AMR gene cluster comprising 14 genes

(*aadA2–ere(A)–dfrA32–sul1–arr-3–catB3–blaOXA-1–aac(6')-Ib-D181Y–tet(C)–sul2–*

aph(3'')-Ib–aph(6)-Id–tet(A)–floR) was associated with flanking IS6 (also known as

IS26) (**Fig. 5c, 5d**). This cluster spanned ~72.6 kb of contiguous AMR gene content

within an IS-bounded region of ~180 kb. IS26 elements (3 copies, **Fig 5d**) were

positioned within and flanking this cluster, with the closest IS26–AMR gene distances of

88–956 bp for six clinically important resistance genes including *aac(6')-Ib-D181Y* (92

bp), *aph(3')-Ia* (88 bp), and *sul1* (112 bp). This organization closely matches the

PmGRI1 genomic resistance island previously characterized in *P. mirabilis* Cluster-1

strains [64]. In contrast, non-ST135 isolates with moderate to low AMR burdens (e.g.,

ST185 with 11 genes, ST145 with 5–11 genes) lacked IS26 entirely (**Fig 5a**). IS

elements flanking AMR genes in these isolates belonged primarily to the unclassified

"new" family, IS3, and IS5 (**Fig 5d**).

A conserved Tn7-associated gene cassette (*aadA1-sat2-dfrA1*) was identified in

9/22 isolates (40.9%), three of which were ST135 (**Table S10**). This cassette displayed

an identical 1,938 bp span and gene order in all positive isolates regardless of

phylogenetic lineage, confirming the clonal stability of this segment. Interestingly, the

three ST135 isolates also harbored a second AMR gene cluster (*catA1-blaTEM-1-*

aph(3')-Ia-dfrA17-aadA5-sul1) spanning ~14.3 kb with identical gene content and order

between genomes.

The MGE landscape revealed Type IV secretion-system-type integrative and

conjugative elements (ICE) in *P. mirabilis* genomes

Genome mining using ICEberg 3.0 [62] identified 1-2 Type IV secretion system

(T4SS)-type ICEs in the 22 complete, polished chromosomes of *P. mirabilis*. The GC

content of these elements ranged from 44.86%-46.19%, they integrated near the

tRNA-Phe gene, flanked by identical 52-bp direct repeats (attL/attR), and carried a

mobilization protein family H (MOBH) relaxase and a type G mating pair formation

system. ST135 isolates (H115-00, H115-29, H127-00) carried one identical ICE

harboring 13 AMR genes along with a copy of the IS26-flanked cluster of 5 genes

(*aacA4-blaTEM-catA1-emrE-folP1*) (**Fig 6a**). In contrast, ST185 isolates (102-00, H120-

04) had 2 regions with T4SS-type ICEs. Annotation of the ICE identified a range of

virulence, housekeeping, and metabolic genes (e.g., permease, PikA5); **Fig. 6b**). The

other isolates carried only one ICE, harboring similar virulence-associated genes as the

region 1 ICE in ST185 (**Fig. 6c**).

Prophages and plasmids have a minimal contribution to AMR gene carriage in *P.*

mirabilis

PHASTER analysis of the 22 clinical isolate genomes identified 189 prophage

regions (mean: 8.2/genome), including 64 intact (33.8%), 20 questionable (10.5%), and

104 incomplete (55%) (**Table S10**). Coordinate-overlap analysis revealed 14 AMR gene

instances within prophage regions across 3/11 isolates (27%), none of which were

ST135. These AMR genes in these prophage regions included 10 genes spanning six

resistance classes (aminoglycosides, β-lactams, chloramphenicol, sulfonamides,

trimethoprim, and disinfectants).

PlasmidFinder detected replicons in only 2/26 isolates (6.7%): IncN4

(conjugative) plus Col3M and ColE10 in H94-00 (ST696), and a single ColE10 in H81-

01 (ST765) (**Table S10**). The only confirmed plasmid-borne AMR gene across all 30

isolates was *qnrD1* (quinolone resistance) on the Col3M plasmid in H94-00 (ST696).

This indicates that the overwhelming majority of AMR in this *P. mirabilis* collection are

chromosomally encoded.

Hierarchical MGE overlaps reveal a "Russian doll" architecture driving multidrug

resistance in ST135

Systematic analysis of MGE overlaps within the 22 clinical isolates revealed a

nested hierarchical architecture most pronounced in ST135 isolates. Class 1 integrons

with their 3'-CS (*qacEΔ1-sul1*) are embedded near IS26 sites. Each integron carries

distinct aminoglycoside and trimethoprim resistance cassettes (*aadA, dfrA*). These

modules are themselves clustered within *PmGRI1*-like genomic islands containing IS

elements. In ST135 isolates, IS26 serves as the structural organizer of *PmGRI1*: Region

1 contains 6 IS elements (3× IS26, 2× ISL3, 1× IS3) interspersed among 16 AMR genes

(**Fig 7a**); Region 2 contains 8 IS elements (3× IS26, 3× IS5, IS66, IS200/IS605) among

10 AMR genes (**Fig 7b**). IS26 may act as a boundary element for sub-modules within

PmGRI1, enabling modular rearrangement and stepwise accumulation of resistance. In

the same arrangement, Prophage region 10 (35.1 kb, intact, score 130, with *att* sites)

simultaneously overlaps region 2. This represents a four-way overlap of all element

types within a single 35 kb region, nested within *PmGRI1*.

Experimental AMR phenotyping of *P. mirabilis* clinical isolates reveals variable

genotype-phenotype concordance

AMRFinderPlus reports gene presence/absence and point mutation but does not

infer phenotypic resistance due to factors including gene expression and regulation.

Thus, many of the resistance genes detected by established computational databases

may not be relevant for clinical management or antimicrobial surveillance. To determine

concordance between *in-silico* AMR predication and phenotypic resistance in *P.*

mirabilis, we determined the AMR of all 27 *P. mirabilis* clinical isolates by standardized

Antimicrobial Susceptibility Tests (AST). We tested growth kinetics of strains with

specific resistance genes at 15-minute intervals over a 20-hour time course to evaluate

the minimum inhibitory concentrations as well as more nuanced dose-dependent effects

on the growth rates and doubling times of the *P. mirabilis* strains (**Fig 8 and Fig 9**).

Antibiotic concentrations were selected based on the Clinical and Laboratory Standards

Institute (CLSI) guidelines for *Enterobacterales* and were designed to span the

thresholds for sensitive (lowest) to resistant (highest). The genotypic and phenotypic

resistance profiles are summarized in **Table S9 and S11.**

Tetracycline, Kanamycin, Streptomycin, Streptothricin, and Ampicillin

For tetracycline, 100% of clinical isolates (26/26) exhibited resistance with MIC

values >20 µg/mL, which is consistent with the presence of *tetJ*. Growth kinetics

demonstrated uniformly robust growth across all concentrations (**Fig 9a**). Thus, a

perfect genotype-phenotype concordance was observed. Carriage of any additional

MGE-acquired *tet* genes (*tetA* or *tetC*) are phenotypically redundant.

Kanamycin resistance genes (*aac(6')-Ib-cr5, aph(3')-Ia*) were detected in 7/26

clinical isolates and genotype-phenotype concordance was observed. All 7 isolates

encoding the resistance genes had MIC >34 ug/mL, while 106-15, H81-01, H94-04, and

HI4320 were predicted to lack all kanamycin resistance determinants and were

susceptible, but some exhibited MICs ranging from 14 – 20.5 ug/ml (**Fig 9b**).

Different combinations of streptomycin resistance genes (*aadA1, aph(6)-Id,*

aph(3")-Ib, aadA5, aadA2) were observed in 11/26 clinical isolates (**Fig 8a, Table S10**),

and all provided resistance to streptomycin when tested in our AST assay (**Fig 9c**).

Unexpectedly, resistance was also observed in 102-24, 106-15, H81-01, H94-00, and

reference strain HI4320 which lacks canonical streptomycin resistance genes. However,

these strains all exhibited a dose-dependent decrease in growth when incubated with

streptomycin. It is also notable that clinical isolate 210-04 (1 predicted gene, *aadA1*)

exhibited a similar dose-dependent decrease in growth even though other strains (206-

05, H97-00, and H97-11) with a singular *aadA1* gene with 100% amino acid sequence

similarity to that of 210-04 did not show dose dependency. It is possible that regulation

of *aadA1* differs in 210-04, leading to a less efficient or inducible mechanism of

resistance compared to the other isolates. The presence of multiple, stacked

aminoglycoside resistance genes on MGEs like *PmGRI1* in H115-00

(e.g., *aadA1, aph(6)-Id, aph(3'')-Ib)* can confer broader and potentially higher-level

resistance compared to strains carrying only a single gene, explaining quantitative

differences in MICs beyond simple gene presence.

An ampicillin resistance gene (*blaTEM*) was identified in 6/26 clinical isolates and

all exhibited MICs >34 ug/mL. The resistance phenotype was also observed in HI4320,

which lacks any known β-lactamase genes. However, strains that did not encode

blaTEM, including HI4320, exhibited dose-dependent growth inhibition with increasing

concentrations of ampicillin, despite displaying resistance at CLSI breakpoints (**Fig 9d**).

The dose-dependent growth inhibition observed in some strains with *bla*TEM, and the

low-level resistance in strains lacking it, may be due to IS-proximal regulatory effects

influencing gene expression or the presence of alternative, non-canonical resistance

mechanisms located within genomic islands.

Streptothricin resistance gene *sat2* was observed in 11/26 clinical isolates (**Fig

8a, Table S9**), yet the AST assay revealed a mix of intermediate to resistant

phenotypes with MICs ranging from 8 to >22 ug/mL (CLSI breakpoints are ≤8 for

susceptible and ≥16 for resistant). Since streptothricin resistance is conferred by *sat1,*

sat2, and *sat4* genes, the presence of only one gene may confer only partial resistance

with dose-dependency increasing with gene stacking. The genotypically-positive

isolates 102-00, 206-16 H97-11, H115-00, H115-29, H120-00, and H120-04 exhibited

dose-dependent decreases in growth with increasing streptothricin concentrations.

However, other genotypically positive strains 206-05 and 210-04 were sensitive (**Fig

9e). HI4320 exhibited robust growth regardless of streptothricin concentration, yet this

strain does not encode any known streptothricin resistance genes (**Fig 9e).**

Sulfonamide and Trimethoprim

Sulfonamide resistance genes (*sul1* and *sul2)* were both observed in the 26

clinical isolates, with *sul1* present in 3 isolates (H115-0, H115-05, H120-00), *sul2* in 5

isolates (102-00, 206-16, H120-04, H115-00, H115-05), and both genes co-occurring in

3 isolates (H115-00, H115-05. H127-00) (**Table S9**). AST with sulfamethoxazole

revealed high level resistance (MIC >514 ug/ml) in all isolates, irrespective of *sul* gene

carriage (single or double), indicating that the presence of an additional *sul* gene did not

further elevate resistance. Unexpectedly, even strains that did not encode *sul* (102-24,

106-15, 210-04, H81-01, H94-00, H97-11, and HI4320) had MICs >514 ug/ml, but

partial growth inhibition was clearly present even at the lowest tested concentration (250

ug/ml, **Fig. 10a**). The uniform MIC >514 µg/mL across all strains implies clinical

resistance per CLSI guidelines, as it far exceeds the susceptible breakpoint (typically

≤38 µg/mL). However, the partial growth inhibition of strains that do not encode *sul*

highlights the complexity of interpreting MICs near the resistance threshold. These data

suggest that, while canonical *sul*-mediated resistance mechanisms were absent in

these isolates, IS elements located near AMR genes or regulatory regions may

contribute to the observed resistance (above threshold) phenotype.

Trimethoprim resistance genes were also detected in 10/26 clinical isolates, with

MICs of >15 µg/mL (exceeding clinical breakpoints) regardless of which resistance gene

(*dfrA1*, *dfrA17*, or *dfrA32*) was present **(Fig 10b)**. However, one isolate (H97-00)

harboring *dfrA1* exhibited dose-dependent growth inhibition, contrasting with other

strains harboring *dfrA1* with 100% amino acid similarity that showed no dose-

dependency. This implies strain-specific phenotypic variability (e.g., differential gene

expression, compensatory mutations) despite shared genetic determinants. The *dfrA*-

negative reference strain H81-01 also displayed full resistance, while strain HI4320 that

did not carry the *dfrA* gene or gene mutation displayed intermediate resistance and

mirrored the dose-dependent growth inhibition of strain H97-00. These observations

support the existence of non-*dfrA* resistance mechanisms in HI4320 and H81-01, such

as altered folate metabolism or reduced drug uptake. The discordance between genetic

markers and phenotypic outcomes also underscores a key challenge: genetic testing

alone cannot always predict antibiotic response. However, *dfr* negative strains 102-24,

106-15, and H94-00 were sensitive to trimethoprim, in concordance with the genotype.

Since the genotype-negative strains were all phenotypically resistant to

sulfamethoxazole (SMX), albeit with reduced growth rate, we further tested the impact

of combinatory treatment (trimethoprim-sulfamethoxazole, TMP-SMX) on *P. mirabilis*

growth. This is an important parameter since, historically, TMP-SMX was the first-line

empirical treatment for acute uncomplicated UTIs. In the strains exhibiting 100%

genotype-phenotype concordance for both SMX and TMP resistance (i.e. true resistant

genotypes correlated with resistant phenotypes), all isolates (102-00, 206-16, H115-00,

H115-05, H120-00, and H120-04), exhibited complete resistance to the combinatory

drug without any dose-dependent decreases in growth (**Fig 10c**). In contrast, strains

such as 206-05, 210-04, and H97-11 that were genotypically SMX susceptible TMP

resistant but phenotypically resistant to both antibiotics when used alone exhibited a

dose-dependent resistance phenotype when treated with TMP-SMX. Additionally,

strains that were genotypically predicted to be sensitive to both antibiotics but were

phenotypically resistant demonstrated intermediate to resistant phenotypes for

combination treatment: H81-01 displayed an MIC of 8/152 ug/mL (resistant), while 102-

24 displayed an MIC >2/38 ug/mL (intermediate). This is clinically concerning as growth

was observed at breakpoint concentrations. Other discordant strains that had lower

growth yields under single SMX exposure (106-15, H94-00, H97-00, HI4320) were

found to be sensitive to the TMP-SMX combinations (**Fig 10c**).

Chloramphenicol

Chloramphenicol resistance displayed genotype-phenotype concordance, with all

isolates possessing a *catA* gene exhibiting resistance and those lacking *cat* genes

exhibiting susceptibility (**Fig 11**). MIC values ranged from 10-14 ug/ml in isolates lacking

a *cat* gene but were >26 ug/ml in isolates with *catA* alone and >34 ug/ml in isolates

harboring both *catA* and *catA1* (**Table S9**). Interestingly, many isolates with a single

catA gene (16/17) exhibited dose-dependent growth inhibition (**Fig 11**), while isolates

possessing both *catA* and *catA1* demonstrated hyper-resistance (hR) with no dose

dependency (**Fig 11**). The exception was HI4320, which only encodes *catA* but

exhibited robust growth at all but the highest three concentrations of chloramphenicol.

No strains were predicted to have *catA1* alone, so we were not able to determine

whether *catA1* provides a greater level of resistance than *catA,* but our data suggests

that the presence of two *catA* genes enhances functional resistance.

We next compared the sequences of the *catA* and *catA1* genes in our *P.*

mirabilis clinical isolates that harbored both genes as *catA1* has not previously been

reported in *P. mirabilis*. The two genes share 76% amino acid sequence identity (with

representative strain 102-00; **Fig S2a**), and *catA* from strain 102-00 showed 100%

identity to the *catA* gene in reference strain *P. mirabilis* HI4320 (**Fig S2b**). The

nucleotide and amino acid sequences of *catA1* from our isolates share 100% amino

acid identity with *catA1* from *E. coli* and ~98% amino acid identity with *catA1*

from *Photobacterium damselae* and *Shigella flexneri* (**Fig S2c**).

Even though we had a limited sample size, within-lineage discordance was

observed in ST145 and can be attributed to differential MGE acquisition. 102-24 carries

11 AMR genes (including IS5-flanked accessory cluster with *aph(3″)-Ib, aph(3′)-Ia*,

aph(6)-Id, blaTEM-1, catA1, sul2) while 210-04 carries only 5 genes. The 6 additional

genes in 102-24 are MGE-acquired (flanked by IS5 elements), conferring predicted

resistance to 3+ additional drug classes. This means two isolates of the same sequence

type would have different AST profiles, a difference that is possibly explained by

differential MGE acquisition.

DISCUSSION

CAUTIs caused by *P. mirabilis* remain clinically challenging due to crystalline biofilm formation, recurrent infections, and persistence despite antibiotic treatment and catheter changes [15]. This comprehensive genomic analysis of *P. mirabilis* from human urinary sources expands our understanding of *P. mirabilis* genetic diversity, AMR profiles, and the accessory genome. We analyzed 1,027 publicly available and clinical isolates and determined that urinary *P. mirabilis* isolates possess an open pangenome with more than 41,121 gene families and a core genome of 1,937 genes. Open pangenomes are characteristics of sympatric species with a high capacity of horizontal gene transfer (HGT) mediated by MGEs, highlighting genomic plasticity. This may allow *P. mirabilis* to rapidly adapt to the urinary tract under antibiotic exposure, host immune responses, and competition in the polymicrobial catheterized environment [65-67]. This dynamic gain, loss, and exchange of MGEs with accessory genomes comprising ~90% (37,918 genes) emphasizes the importance of flexible genomic elements in driving diversity in the urinary tract.

The presence of 213 defined STs and ~118 previously undefined genomes and a star-like phylogeny further supports ongoing diversification. Almost half of the STs were represented by single isolates (46%), showcasing heterogeneity. Yet, most genomes were distributed across a small subset of STs (ST96, ST185, ST99, ST135, ST97), resulting in a skewed distribution. While MLST remains a widely used typing method due to its reproducibility and interpretability [51, 68], we want to highlight its limitations for epidemiological investigations [68, 69]. The substantial proportion of unidentified STs

and limited sample size constrain ST-specific characterization in the entire 1,027

genomes but highlights the need for continuously updated typing schemes.

The AMR landscape was complex and dynamic, with variable genotype-

phenotype concordance. Overall, 93% of the isolates were resistant to ≥2 antibiotic

subclasses and 25% resistant to >6 subclasses posing a challenge for effective

treatment of *P. mirabilis* UTI's. Although MLST was not a significant predictor of high

AMR carriage in the full dataset of 1,001 genomes, ST135 emerged as a high-risk clone

is the subset of 422/1,001 with >10 genomes. 95% of ST135 genomes had ≥16

resistance genes and 100% had resistance to >6 antimicrobial subclasses (**Fig 1a**).

This near-uniform high-level resistance within ST135 is attributed to the IS26-mediated

MGEs that are rarely observed in other lineages. IS6/IS26-encoding genomes carry

3.2× more AMR genes (20.8 vs 6.5) and 2.8× more drug subclasses (13.9 vs 4.9), with

a significant dose-response relationship as IS6 copy number increases. MLST ST135

had the highest IS diversity in our dataset (37/422 genomes + 3 clinical isolates),

highest IS6/IS26 prevalence, and AMR gene burden with 100% MDR (>3 antibiotic

subclass).

Geographic confounding was excluded by our analysis; although ST135 was

predominantly isolated from St. Louis, isolation source alone was not an independent

driver of high AMR carriage (p=0.051) pointing to lineage-intrinsic factors rather than

regional antibiotic selection pressures. The absence of geographical associations,

together with the hierarchical MGE architecture we identified (**Fig 7**) implicates lineage-

specific MGE carriage as the mechanistic basis for extreme resistance phenotypes.

This is known for other species as well—*S. aureus* USA300 (ST8/5) carry the mecA

cassette and are seen globally [70-72]. The overlapping MGE architecture (H115-00;

Fig 7) reveals a multi-level system for AMR propagation. Class 1 integrons capture

individual resistance cassettes, IS26 organizes these into modular translocatable units,

PmGRI1 provides stable chromosomal integration, and prophages enable horizontal

transfer of complete AMR modules. This nested hierarchy has clinical implications

because redundant dissemination pathways can ensure robust AMR spread. Clustering

of multiple resistance genes within shared elements means selection for any single

antibiotic co-selects the entire MDR module, and chromosomal integration ensures

stable inheritance while IS26 and phage mechanisms retain mobility. The exclusive

association of this architecture with ST135, which carries significantly more AMR genes

than IS26-negative lineages (mean 25.0 vs. 4.1, p=0.0019), implicates these

overlapping MGEs as key drivers of this lineage's epidemiological success.

A recent study by *Zhang et. al. (2026)* identified the same high-risk lineage which

they assigned Cluster-1 genomes comprising 233 globally distributed isolates [64]. Our

analysis confirmed that Cluster-1, which mostly carries ST135 (159/233), had

significantly elevated AMR gene burden. The high AMR gene counts across isolates

from different locations and hosts that was reported by *Zhang et. al.* further supports

ST135 as a lineage of major clinical concern [64]. The T4SS ICE and IS26-mediated

AMR gene clusters help to stack AMR genes within *PmGRI1*, enabling vertical

transmission and potentially facilitating further resistance acquisition. Such

lineage-specific AMR enrichment has critical implications for infection control and

genomic surveillance, as clonal expansion of ST135 could rapidly disseminate multidrug

resistance across healthcare networks like that or Methicillin resistant *S. aureus*

(MRSA), Vancomycin resistance *E. faecalis* (VRE) [70, 72-78]. Prospective surveillance

should therefore prioritize tracking this high-risk lineage, and future functional studies

should dissect the precise MGEs responsible for its resistance burden and why is it

limited to this specific lineage. In contrast, the broader distribution of common

resistance subclasses across multiple STs reflects ongoing horizontal gene transfer.

These findings highlight the critical need for genomic surveillance targeting high-risk

lineages, as clonal expansion of ST135 could rapidly disseminate multidrug resistance

across healthcare networks.

The overwhelming prevalence of tetracycline resistance genes (98.9%),

particularly *tetJ*, aligns with intrinsic resistance of *P. mirabilis.* The dominance

of *tetJ* and the frequent carriage of only a single *tet* gene (94.3%) suggest this

determinant is often sufficient for the observed resistance phenotype. Tetracycline

resistance also exemplifies how MGE-acquired genes can be phenotypically redundant:

tet(J) alone confers complete resistance (MIC >20 µg/mL), yet some genomes such as

the ST135 isolates carry three genes (*tet*(J), *tet*(A), *tet*(C)) acquired on the PmGRI1

genomic island via IS26. Thus, the additional MGE-borne genes inflate genotypic

counts without meaningfully expanding the phenotype.

Chloramphenicol resistance genes (*catA*, 95%) were prevalent despite limited

clinical use, possibly reflecting co-selection with other antibiotics or acquisition due to

extensive use in agricultural and environmental settings [79, 80]. The co-occurrence

of *catA* and *catA1* was detected in 19.2% of *cat*-positive isolates, with rare

floR and *catB3* gene stacking. Strains with only a single *cat* gene exhibited dose-

dependent growth inhibition, while additional *catA1* resulted in a hyper-resistant

phenotype (no growth inhibition at all at any tested concentrations), enhancing the

resistance phenotype. This observation suggests that the combined expression of

multiple acetyltransferases, driven by IS26 elements positioned within 10 kb,

quantitatively enhances resistance. Detection of *catA1* in *P. mirabilis* is a significant

finding as this gene has not been well reported and characterized in *P. mirabilis* but

match closely to related bacterial species [81, 82].

Additionally, we observed resistance to aminoglycosides (71%) with streptomycin

resistance often co-occurring with other determinants of carbapenems, cephalosporins,

fluoroquinolones and trimethoprim-sulfonamides (**Fig 1**). This persistence despite

reduction in clinical use may reflect MGE mediated clonal spread of resistant lineages.

Trimethoprim (TMP, *dfr*, 29%) and sulfonamide (SMX, *sul*, 28%) resistance

determinants also co-occurred in 22% of genomes, threatening the efficacy of this

combination therapy for *P. mirabilis* for UTI treatment. While ESBL-associated genes

were less common (18.5% and primarily *blaTEM-1*), their presence is concerning for

broader β-lactam resistance development. Resistance for quinolones (*qnr aac(6')-Ib-cr,*

gyrA/parC mutations) and the chloramphenicol/florfenicol (*cat, floR*) combination were

rare in both datasets (~7%) but noteworthy as potential molecular fingerprints of human

driven selection linked to agriculture use and MGE gene transfer from *P. mirabilis*

lineages found in animal sources [35, 64, 83-86].

Our comprehensive AMR analysis shows a complex genotype-to-phenotype

landscape with both expected concordance and clinically critical discordance.

Concordance for antibiotics like tetracycline was expected and kanamycin resistance

reliably predicted by the presence of an *aph(3')-Ia* gene, validated the robustness of our

genomic approach for identifying known resistance determinants [75, 87, 88]. However,

discordances provide compelling insights into AMR dissemination. For example,

resistance to streptomycin and streptothricin could be due to novel acetyltransferases.

Similarly, the ampicillin resistance in strains 102-24, 106-15, H94-00, H97-11, and

HI4320 that lack a detectable β-lactamase gene may reflect permeability or efflux

mechanisms, which is a hallmark of concentration-dependent antibiotics [89, 90], where

higher concentrations or adjuvants can eventually overcome cellular defenses.

A critical finding was the discordance between genotypic SMX susceptibility and

TMP-SMX phenotypic response (**Fig. 8**). Strains that lacked the canonical *sul* genes yet

exhibited variable resistance phenotypes could possibly be explained by MGE-related

mechanisms: 1) IS elements positioned ≤100 bp from AMR genes in seven isolates may

drive upregulation via outward-facing promoters; 2) ST135 carries three *dfr* genes

(*dfrA1*, *dfrA17*, *dfrA32*) accumulated via IS26 on *PmGRI1*, while some isolates carry

only *dfrA1* via non-IS26 Tn7 transposon. Thus, strains like 206-05 and 210-04 (MIC

>16/305 μg/mL) likely possess MGE-driven mechanisms robust enough to contribute to

resistance, albeit less efficiently than truly resistant strains, while the sensitivity of

strains like HI4320 have mechanisms still overcome by trimethoprim. Such prediction

errors can be dangerous for treatment outcomes.

The genotype-phenotype discordance also highlights a significant limitation of

current genomic prediction models: their reliance on curated databases of known

resistance genes and mutations. Our study confirms that WGS-based prediction is

highly reliable for well-characterized resistance determinants in *P. mirabilis* but can fail

to predict phenotypes that are likely governed by cryptic mutations, efflux, and MGEs

that impact gene expression, copy number, and functional redundancy. Not much has

been reported on the mechanism of resistance involving efflux systems and membrane

permeability [10, 83, 91]. Efflux systems such as *AcrAB-TolC* reported in many *P.*

mirabilis strains can be overwhelmed by high drug concentrations, rather than the

efficient enzymatic hydrolysis provided by a beta-lactamase and is linked to a very high

multidrug-resistance rates [92]. Genotype based predictions are also prone to

sequencing, assembly, and annotation errors, ideally needing long-read or reference-

based sequencing to create hybrid assemblies. However, this can become impractical

and costly for diagnostic laboratories.

Several limitations should be considered when interpreting our findings. First, the

publicly available genomes were only derived from human hosts in the US, potentially

biasing global population inferences. Second, our MGE analysis was restricted to the 22

clinical hybrid-assembled genomes to ensure accuracy, limiting the sample size for

these comparisons but short reads are not an accurate predictor for deep MGE

analysis. Third, while we identified compelling associations between MGE architecture

in ST135 and AMR burden, it was done only in the sample set of 26 clinical isolate

genomes and HI4320 with short-read assembly. Fourth, functional validation of IS26-

mediated gene stacking and prophage transduction is necessary to confirm the

mechanistic roles proposed here. Finally, our genotype-phenotype discordance

analysis, though comprehensive, was necessarily limited by the number of clinical

isolates with paired AST data.

In conclusion, while WGS has the potential to be used widely in clinical context

for AMR predictions, our findings reinforce that assessing gene presence/absence does

not replace phenotypic AST of *P. mirabilis* isolates for accurate clinical decision-making

and surveillance. Accurate resistance prediction requires larger sampling size, database

curation, resistance gene detection, and comprehensive characterization of the MGE

landscape in *P. mirabilis*. These findings highlight the need for genomic surveillance

targeting both high-risk lineages and the mobile elements that enable resistance gene

flux across phylogenetic boundaries.

Acknowledgements

This work was supported by an American Heart Association (AHA) pre-doctoral

fellowship 24PRE1195538 to ND, and by the National Institute of Diabetes and

Digestive and Kidney Diseases under awards R01 DK123158 and R01 DK140371 to

CEA.

References

1. Yang, X., et al., *Disease burden and long-term trends of urinary tract infections: A worldwide report.* Front Public Health, 2022. **10**: p. 888205.
2. Zeng, Z., et al., *Global, regional, and national burden of urinary tract infections from 1990 to 2019: an analysis of the global burden of disease study 2019.* World J Urol, 2022. **40**(3): p. 755-763.
3. Biondo, C., *New Insights into the Pathogenesis and Treatment of Urinary Tract Infections.* Pathogens, 2023. **12**(10).
4. Schaffer, J.N. and M.M. Pearson, *Proteus mirabilis and Urinary Tract Infections.* Microbiol Spectr, 2015. **3**(5).
5. Gaston, J.R., et al., *Polymicrobial interactions in the urinary tract: is the enemy of my enemy my friend?* Infect Immun, 2021.
6. Zilberberg, M.D., et al., *Descriptive epidemiology and outcomes of emergency department visits with complicated urinary tract infections in the United States, 2016-2018.* J Am Coll Emerg Physicians Open, 2022. **3**(2): p. e12694.
7. Ku, J.H., et al., *Multidrug Resistance of Escherichia coli From Outpatient Uncomplicated Urinary Tract Infections in a Large United States Integrated Healthcare Organization.* Open Forum Infect Dis, 2023. **10**(7): p. ofad287.
8. Fromer, D.L., et al., *Risk Factors for Empiric Treatment Failure in US Female Outpatients with Uncomplicated Urinary Tract Infection: an Observational Study.* J Gen Intern Med, 2025. **40**(4): p. 862-870.
9. Wolterink, I., et al., *Nitrofurantoin Failure in Elderly Men: A Retrospective Observational Study.* Antibiotics (Basel), 2020. **9**(5).
10. Armbruster, C.E., H.L.T. Mobley, and M.M. Pearson, *Pathogenesis of Proteus mirabilis Infection.* EcoSal Plus, 2018. **8**(1).
11. Yuan, F., et al., *Pathogenesis of Proteus mirabilis in Catheter-Associated Urinary Tract Infections.* Urol Int, 2021. **105**(5-6): p. 354-361.
12. Brauer, A.L., et al., *Preferential catabolism of L- vs D-serine by Proteus mirabilis contributes to pathogenesis and catheter-associated urinary tract infection.* Molecular microbiology, 2022. **118**(3): p. 125–144.
13. Armbruster, C.E., et al., *Genome-wide transposon mutagenesis of Proteus mirabilis: Essential genes, fitness factors for catheter-associated urinary tract infection, and the impact of polymicrobial infection on fitness requirements.* PLoS Pathog, 2017. **13**(6): p. e1006434.
14. White, A.N., et al., *Catalase Activity is Critical for Proteus mirabilis Biofilm Development, Extracellular Polymeric Substance Composition, and Dissemination during Catheter-Associated Urinary Tract Infection.* Infect Immun, 2021. **89**(10): p. e0017721.
15. Armbruster, C.E., et al., *Prospective assessment of catheter-associated bacteriuria clinical presentation, epidemiology, and colonization dynamics in nursing home residents.* JCI Insight, 2021. **6**(19).
16. Sabbuba, N.A., et al., *Genotyping demonstrates that the strains of Proteus mirabilis from bladder stones and catheter encrustations of patients undergoing long-term bladder catheterization are identical.* J Urol, 2004. **171**(5): p. 1925-8.

17. Nye, T.M., et al., *Microbial co-occurrences on catheters from long-term*
catheterized patients. Nat Commun, 2024. **15**(1): p. 61.
18. Armbruster, C.E., et al., *The Pathogenic Potential of Proteus mirabilis Is*
Enhanced by Other Uropathogens during Polymicrobial Urinary Tract Infection.
Infect Immun, 2017. **85**(2).
19. Ramatla, T., et al., *Antibiotic resistance and virulence profiles of Proteus mirabilis*
isolated from broiler chickens at abattoir in South Africa. Vet Med Sci, 2024.
10(2): p. e1371.
20. Bharat, A., et al., *Correlation between Phenotypic and In Silico Detection of*
Antimicrobial Resistance in Salmonella enterica in Canada Using Staramr.
Microorganisms, 2022. **10**(2).
21. Gaston, J.R., et al., *Enterococcus faecalis Polymicrobial Interactions Facilitate*
Biofilm Formation, Antibiotic Recalcitrance, and Persistent Colonization of the
Catheterized Urinary Tract. Pathogens, 2020. **9**(10).
22. Luzzaro, F., et al., *Prevalence and epidemiology of microbial pathogens causing*
bloodstream infections: results of the OASIS multicenter study. Diagn Microbiol
Infect Dis, 2011. **69**(4): p. 363-9.
23. Kanayama, A., I. Kobayashi, and K. Shibuya, *Distribution and antimicrobial*
susceptibility profile of extended-spectrum beta-lactamase-producing Proteus
mirabilis strains recently isolated in Japan. Int J Antimicrob Agents, 2015. **45**(2):
p. 113-8.
24. Filipiak, A., et al., *Pathogenic Factors Correlate With Antimicrobial Resistance*
Among Clinical Proteus mirabilis Strains. Front Microbiol, 2020. **11**: p. 579389.
25. Li, Y., et al., *Genetic analysis of resistance and virulence characteristics of*
clinical multidrug-resistant Proteus mirabilis isolates. Front Cell Infect Microbiol,
2023. **13**: p. 1229194.
26. Nielubowicz, G.R. and H.L. Mobley, *Host-pathogen interactions in urinary tract*
infection. Nat Rev Urol, 2010. **7**(8): p. 430-41.
27. Tahmasebi H, et al., *Relationship between biofilm gene expression with*
antimicrobial resistance pattern and clinical specimen type based on sequence
types (STs) of methicillin-resistant S. aureus. . Molecular Biology Reports 2020.
47(2): p. 1309-1320.
28. Chen, S.L., et al., *A Core Genome Multilocus Sequence Typing Scheme for*
Proteus mirabilis. Biomed Environ Sci, 2023. **36**(4): p. 343-352.
29. Hickman FW and F.J. 3rd, *Differentiation of Proteus mirabilis by bacteriophage*
typing and the Dienes reaction. Journal of Clinical Microbiology, 1976. **3**(3): p.
350-8.
30. William C. Schmidt and C.D. Jeffries, *Bacteriophage Typing of Proteus mirabilis,*
Proteus vulgaris, and Proteus morganii. Applied and Environmental Microbiology,
1974.
31. Pfaller MA, et al., *Evaluation of the Discriminatory Powers of the Dienes Test and*
Ribotyping as Typing Methods for Proteus mirabilis. Journal of Clinical
Microbiology, 2000. **38**(3): p. 1077-80.
32. Potter RF, et al., *Uncharacterized and lineage-specific accessory genes within*
the Proteus mirabilis pan-genome landscape. mSystems, 2023. **84**(4): p. 00159-
23.

33. Lian S, et al., *Genomic insights on cgMLST markers, drug resistance, and urease cluster of Proteus mirabilis strains.* 2024. **13**(1).
34. Yin, Z., et al., *Insights into novel diagnostic assay development, antimicrobial resistance, and pathogenicity in Proteus mirabilis through pan-genome analysis.* Appl Environ Microbiol, 2026: p. e0189825.
35. Zhang, S., et al., *Genomic analysis of Proteus mirabilis: Unraveling global epidemiology and antimicrobial resistance dissemination - emerging challenges for public health and biosecurity.* Environ Int, 2025. **199**: p. 109487.
36. Feldgarden, M., et al., *AMRFinderPlus and the Reference Gene Catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence.* Sci Rep, 2021. **11**(1): p. 12728.
37. Sung, K., et al., *Comprehensive Genomic Analysis of Uropathogenic E. coli: Virulence Factors, Antimicrobial Resistance, and Mobile Genetic Elements.* Pathogens, 2024. **13**(9).
38. Chin, J.J., et al., *High Carriage of tetA, sul1, sul2 and bla (TEM) Resistance Genes among the Multidrug-resistant Uropathogenic Escherichia coli (UPEC) Strains from Malaysian Patients.* Trop Life Sci Res, 2024. **35**(2): p. 211-225.
39. Kandi, V., et al., *Molecular Characterization of Escherichia coli Causing Urinary Tract Infections Through Next-Generation Sequencing: A Comprehensive Analysis of Serotypes, Sequence Types, and Antimicrobial and Virulence Genes.* Cureus, 2024. **16**(3): p. e55556.
40. He, X.T., et al., *The risk factors, antimicrobial resistance patterns, and outcomes associated with extended-spectrum beta-lactamases-Producing pathogens in pediatric urinary tract infection.* Pediatr Neonatol, 2024. **65**(3): p. 242-248.
41. Hassuna, N.A., et al., *Characterization of antimicrobial resistance among Proteus mirabilis isolates from catheter-associated urinary tract infections and non-catheter-associated urinary tract infections in Egypt.* BMC Infect Dis, 2025. **25**(1): p. 767.
42. Mirzaei, A., et al., *From the Urinary Catheter to the Prevalence of Three Classes of Integrons, beta-Lactamase Genes, and Differences in Antimicrobial Susceptibility of Proteus mirabilis and Clonal Relatedness with Rep-PCR.* Biomed Res Int, 2021. **2021**: p. 9952769.
43. Feldgarden, M., et al., *Validating the AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates.* Antimicrob Agents Chemother, 2019. **63**(11).
44. Simon, A. *FastQC: A quality control tool for high throughput sequence data.* 2010 Available from:https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.
45. Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina sequence data.* Bioinformatics, 2014. **30**(15): p. 2114-20.
46. Bankevich, A., et al., *SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing.* J Comput Biol, 2012. **19**(5): p. 455-77.
47. Gurevich, A., et al., *QUAST: quality assessment tool for genome assemblies.* Bioinformatics, 2013. **29**(8): p. 1072-5.
48. Seemann, T., *Prokka: rapid prokaryotic genome annotation.* Bioinformatics, 2014. **30**(14): p. 2068-9.

49.   Hunt. BC, Brix. V, and G.L. Vath. J, Taddei. SM, Deka. N, Learman. BS, Brauer.
AL, Shen. S, Qu. J, Armbruster. CE. , *Metabolic interplay between Proteus*
mirabilis and Enterococcus faecalis facilitates polymicrobial biofilm formation and
invasive disease. mBio 15:e02164-24.
. mBio, 2024. **15**(12).
50.   Walker, B.J., et al., *Pilon: an integrated tool for comprehensive microbial variant*
detection and genome assembly improvement. PLoS One, 2014. **9**(11): p.
e112963.
51.   Jolley, K.A., J.E. Bray, and M.C.J. Maiden, *Open-access bacterial population*
genomics: BIGSdb software, the PubMLST.org website and their applications.
Wellcome Open Res, 2018. **3**: p. 124.
52.   Torsten, S. *mlst: Scan contig files against PubMLST typing scheme.* [cited 2025;
Available from: https://github.com/tseemann/mlst.
53.   Page, A.J., et al., *Roary: rapid large-scale prokaryote pan genome analysis.*
Bioinformatics, 2015. **31**(22): p. 3691-3.
54.   Letunic, I. and P. Bork, *Interactive Tree Of Life (iTOL) v5: an online tool for*
phylogenetic tree display and annotation. Nucleic Acids Res, 2021. **49**(W1): p.
W293-W296.
55.   Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree: computing large minimum*
evolution trees with profiles instead of a distance matrix. Mol Biol Evol, 2009.
26(7): p. 1641-50.
56.   Xie, Z. and H. Tang, *ISEScan: automated identification of insertion sequence*
elements in prokaryotic genomes. Bioinformatics, 2017. **33**(21): p. 3340-3347.
57.   Arndt, D., et al., *PHASTER: a better, faster version of the PHAST phage search*
tool. Nucleic Acids Res, 2016. **44**(W1): p. W16-21.
58.   Arndt, D., et al., *PHAST, PHASTER and PHASTEST: Tools for finding prophage*
in bacterial genomes. Brief Bioinform, 2019. **20**(4): p. 1560-1567.
59.   Wishart, D.S., et al., *PHASTEST: faster than PHASTER, better than PHAST.*
Nucleic Acids Res, 2023. **51**(W1): p. W443-W450.
60.   Carattoli, A., et al., *In silico detection and typing of plasmids using PlasmidFinder*
and plasmid multilocus sequence typing. Antimicrob Agents Chemother, 2014.
58(7): p. 3895-903.
61.   Seemann, T. *Abricate.* Available from: https://github.com/tseemann/abricate.
62.   Wang, M., et al., *ICEberg 3.0: functional categorization and analysis of the*
integrative and conjugative elements in bacteria. Nucleic Acids Res, 2024.
52(D1): p. D732-D737.
63.   *Performance Standard for Antimicrobial Susceptibility Testing*, CLSI, Editor.
2023.
64.   Zhang, T., et al., *Genomic survey uncovers the emergence of a multidrug-*
resistant dominant lineage in Proteus mirabilis populations. NPJ Antimicrob
Resist, 2026. **4**(1).
65.   Friedman, J., E.J. Alm, and B.J. Shapiro, *Sympatric speciation: when is it*
possible in bacteria? PLoS One, 2013. **8**(1): p. e53539.
66.   Zhao, L., et al., *Evolution alters ecological mechanisms of coexistence in*
experimental microcosms. Functional Ecology, 2015. **30**(8): p. 1440-1446.

67.   Mukherjee, S., et al., *Incipient Sympatric Speciation and Evolution of Soil*
1112   *Bacteria Revealed by Metagenomic and Structured Non-Coding RNAs Analysis.*
1113   Biology (Basel), 2022. **11**(8).
1114   68.   Maiden, M.C., *Multilocus sequence typing of bacteria.* Annu Rev Microbiol, 2006.
1115   **60**: p. 561-88.
1116   69.   Perez-Losada, M., et al., *Population genetics of microbial pathogens estimated*
1117   *from multilocus sequence typing (MLST) data.* Infect Genet Evol, 2006. **6**(2): p.
1118   97-112.
1119   70.   Campbell, S.J., et al., *Genotypic characteristics of Staphylococcus aureus*
1120   *isolates from a multinational trial of complicated skin and skin structure*
1121   *infections.* J Clin Microbiol, 2008. **46**(2): p. 678-84.
1122   71.   Phumthanakorn, N., et al., *Genetic characteristics and SCCmec variations in pet-*
1123   *associated MRSA CC398 and ST9 strains.* Vet J, 2026. **315**: p. 106532.
1124   72.   Chanchaithong, P., et al., *Chromosomal and plasmid localization of ileS2 in high-*
1125   *level mupirocin-resistant Staphylococcus pseudintermedius and Staphylococcus*
1126   *aureus isolated from canine and feline origins.* J Antimicrob Chemother, 2024.
1127   **79**(8): p. 1856-1864.
1128   73.   Anahtar, M.N., et al., *Prediction of Antimicrobial Resistance in Clinical*
1129   *Enterococcus faecium Isolates Using a Rules-Based Analysis of Whole-Genome*
1130   *Sequences.* Antimicrob Agents Chemother, 2022. **66**(1): p. e0119621.
1131   74.   Osman, M., et al., *Critical genomic insights into vancomycin-resistant*
1132   *Enterococcus faecium in Lebanon.* Microbiol Spectr, 2025. **13**(9): p. e0017125.
1133   75.   El-Baz, A.M., et al., *The Link between Occurrence of Class I Integron and*
1134   *Acquired Aminoglycoside Resistance in Clinical MRSA Isolates.* Antibiotics
1135   (Basel), 2021. **10**(5).
1136   76.   Gonzalez, M.J., et al., *Phylogenomic insights into LA-MRSA from Argentine pig*
1137   *farm environments: novel OptrA variant and regional emergence of an ST9*
1138   *lineage co-circulating with international CC398 lineages.* Front Microbiol, 2025.
1139   **16**: p. 1662779.
1140   77.   Habib, I., et al., *Exploring human-to-food contamination potential: Genome-based*
1141   *analysis of diversity, toxin repertoire, and antimicrobial resistance of methicillin-*
1142   *resistant Staphylococcus aureus in United Arab Emirates retail meat.* One
1143   Health, 2025. **21**: p. 101216.
1144   78.   Kim, D.H., et al., *Comparative Genomic and Epidemiologic Analysis of*
1145   *Methicillin-Resistant Staphylococcus aureus Isolates in Republic of Korea.*
1146   Antibiotics (Basel), 2026. **15**(3).
1147   79.   Graf, F.E., et al., *Molecular mechanisms of re-emerging chloramphenicol*
1148   *susceptibility in extended-spectrum beta-lactamase-producing Enterobacterales.*
1149   Nat Commun, 2024. **15**(1): p. 9019.
1150   80.   Puangseree, J., et al., *Molecular basis of the persistence of chloramphenicol*
1151   *resistance among Escherichia coli and Salmonella spp. from pigs, pork and*
1152   *humans in Thailand.* PLoS One, 2024. **19**(5): p. e0304250.
1153   81.   Pitout, J.D.D. and L. Chen, *The Significance of Epidemic Plasmids in the*
1154   *Success of Multidrug-Resistant Drug Pandemic Extraintestinal Pathogenic*
1155   *Escherichia coli.* Infect Dis Ther, 2023. **12**(4): p. 1029-1041.

82. Marcoux, P.E., et al., *Host Dependent-Transposon for a Plasmid Found in*
Aeromonas salmonicida subsp. salmonicida That Bears a catB3 Gene for
Chloramphenicol Resistance. Antibiotics (Basel), 2023. **12**(2).
83. ElTaweel, M., H.S. Said, and R. Barwa, *Emergence of extensive drug resistance*
and high prevalence of multidrug resistance among clinical Proteus mirabilis
isolates in Egypt. Ann Clin Microbiol Antimicrob, 2024. **23**(1): p. 46.
84. Piekarska, K., et al., *The molecular mechanisms of fluoroquinolone resistance*
found in rectal swab isolates of Enterobacterales from men undergoing a
transrectal prostate biopsy: the rationale for targeted prophylaxis. Ann Clin
Microbiol Antimicrob, 2021. **20**(1): p. 81.
85. Qu, X., et al., *Genomic Investigation of Proteus mirabilis Isolates Recovered*
From Pig Farms in Zhejiang Province, China. Front Microbiol, 2022. **13**: p.
952982.
86. Thaotumpitak, V., et al., *Emergence of colistin resistance and characterization of*
antimicrobial resistance and virulence factors of Aeromonas hydrophila,
Salmonella spp., and Vibrio cholerae isolated from hybrid red tilapia cage culture.
PeerJ, 2023. **11**: p. e14896.
87. Quirke, J.C.K., et al., *Structure-Activity Relationships for 5'' Modifications of 4,5-*
Aminoglycoside Antibiotics. ChemMedChem, 2022. **17**(13): p. e202200120.
88. Thacharodi, A. and I.L. Lamont, *Aminoglycoside-Modifying Enzymes Are*
Sufficient to Make Pseudomonas aeruginosa Clinically Resistant to Key
Antibiotics. Antibiotics (Basel), 2022. **11**(7).
89. Geng, B., et al., *Identification and Characterization of Genes Related to*
Ampicillin Antibiotic Resistance in Zymomonas mobilis. Antibiotics, 2022. **11**(11).
90. Darby, E.M., et al., *Molecular mechanisms of antibiotic resistance revisited.* Nat
Rev Microbiol, 2023. **21**(5): p. 280-295.
91. Dulanto Chiang, A. and J.P. Dekker, *Efflux pump-mediated resistance to new*
beta lactam antibiotics in multidrug-resistant gram-negative bacteria. Commun
Med (Lond), 2024. **4**(1): p. 170.
92. Al-Kaim, H.W. and H.O.M. Al-Dahmoshi, *Prevalence of AcrAB-TolC efflux pump*
among Proteus mirabilis isolated from cystitis patients in Hilla City, Iraq. Medical
Journal of Babylon, 2023. **20**(3): p. 503-510.
1188
1189

1190   **Table 1: Summary statistics of *P. mirabilis* genome sequences**

Publicly available *P. mirabilis* genomes isolated from urine				
Metric	**Average**	**Max**	**Min**	**Median**
# Contigs	258.43	973	60	180
Total Length	4020761.51	8382181	3667799	3992140
N50	174294.28	1994055	4981	174571
GC Content	38.76	43.24	37.57	38.69
Clinical *P. mirabilis* genomes isolated from catheterized individuals				
Metric	**Average**	**Max**	**Min**	**Median**
# Contigs	66.52	98	2	69
Total Length	4015048.48	4165371	3795264	4015495
N50	317358.93	4063606	119871	182416
GC Content	38.74	38.98	38.38	38.78

1191

1192   **Table 2: *P. mirabilis* clinical isolates used in this study**

Isolate ID	Study	Species	MLST	Patient designation in cited publication	Isolation week
102-00	CEA	*P. mirabilis*	185	Participant A [15]	Visit 0
102-24	CEA	*P. mirabilis*	145	Participant A [15]	Visit 24
102-25	CEA	*P. mirabilis*	258	Participant A [15]	Visit 25
104-00	CEA	*P. mirabilis*	258	Participant C [15]	Visit 0
106-15	CEA	*P. mirabilis*	552	Participant H [15]	Visit 15
201-07	CEA	*P. mirabilis*	258	Participant G [15]	Visit 7
203-01	CEA	*P. mirabilis*	615	Participant F [15]	Visit 1
206-05	CEA	*P. mirabilis*	185	Participant D [15]	Visit 5
206-16	CEA	*P. mirabilis*	185	Participant D [15]	Visit 16
210-04	CEA	*P. mirabilis*	145	Participant J [15]	Visit 4
H81-01	HUC	*P. mirabilis*	765	Patient 81-urine [17]	Visit 1
H85-10	HUC	*P. mirabilis*	330	Patient 85-urine [17]	Visit 10
H94-00	HUC	*P. mirabilis*	696	Patient 94-urine [17]	Visit 0
H97-00	HUC	*P. mirabilis*	157	Patient 97-urine [17]	Visit 0
H97-11	HUC	*P. mirabilis*	157	Patient 97-urine [17]	Visit 11
H105-00	HUC	*P. mirabilis*	591	Patient 105-urine [17]	Visit 0
H105-06	HUC	*P. mirabilis*	765	Patient 105-urine [17]	Visit 6
H108-00	HUC	*P. mirabilis*	96	Patient 108-urine [17]	Visit 0
H115-00	HUC	*P. mirabilis*	135	Patient 115-urine [17]	Visit 0
H115-05	HUC	*P. mirabilis*	135	Patient 115-urine [17]	Visit 5
H119-03	HUC	*P. mirabilis*	146	Patient 119-urine [17]	Visit 3
H120-00	HUC	*P. mirabilis*	97	Patient 120-urine [17]	Visit 0
H120-04	HUC	*P. mirabilis*	185	Patient 120-urine [17]	Visit 4
H125-06	HUC	*P. mirabilis*	565	Patient 125-urine [17]	Visit 6
H127-00	HUC	*P. mirabilis*	96	Patient 127-urine [17]	Visit 0
H132-05	HUC	*P. mirabilis*	96	Patient 132-urine [17]	Visit 5
HI4320	Reference	*P. mirabilis*	90	Refseq: GCA_000069965.1_ASM6996v1_genomic	-

1193

1194

Table 3: AMR subclass count and percentage in *P. mirabilis* genomes

Resistance subclass	Count (NCBI)	Genes detected (NCBI)	Percentage (1001 NCBI)	Count (clinical)	Genes detected (clinical)	Percentage (26 clinical)	NCBI/Clinical Comparison
AMIKACIN/KANAMYCIN / QUINOLONE / TOBRAMYCIN	11	*aac(6')-Ib, aph(3')-IIIa, aph(3')-VI, aph(3')-VIb, aph(3')-XV*	7.19%	1	*aac(6')-Ib-cr5*	3.70%	Lower in clinical
BETA-LACTAM	185	*blaCARB-2, blaOXA, blaOXA-1033, blaOXA-2, blaOXA-9, blaTEM, blaTEM-1, blaTEM-2, blaTEM-30, hugA*	18.46%	5	*blaTEM-1*	18.52%	Similar in prevalence
CEPHALOSPORIN	90	*ampC, blaCMY-2, blaCMY-6, blaCTX-M-14, blaCTX-M-15, blaCTX-M-2, blaCTX-M-3, blaCTX-M-55, blaCTX-M-65, blaCTX-M-91, blaDHA, blaDHA-1, blaFOX-5, blaOXA-1, blaOXA-10, blaPDC-3, blaPER, blaPER-1, blaSHV-12, blaSHV-31, blaTEM-10, blaTEM-155, blaVEB-6*	8.98%	1	*blaOXA-1*	3.70%	Lower in clinical
CHLORAMPHENICOL	891	*catA, catA1, catA2, catA3, catB11, catB2, catB3, catB7, catB8, cmlA, cmlA5, cmlA6*	88.92%	23	*catA, catA1, catB3, floR*	85.19%	Similar in prevalence
ERYTHROMYCIN	36	*ere(A), ere(B) mph(E)*	3.50%	1	*ere(A)*	3.70%	Similar in prevalence

KANAMYCIN	137	*aph(3')-IIb, aph(3')-Ia*	13.67%	7	*aph(3')-Ia*	25.93%	Higher in clinical
QUATERNARY AMMONIUM	192	*qacE, qacEdelta1*	19.16%	3		11.11%	Lower in clinical
QUINOLONE	68	*qnrA, qnrA1, qnrB19, qnrB2, qnrB4, qnrD, qnrD1, qnrD2, qnrS1, qnrVC1*	6.79%	2	*qnrD*	7.41%	Similar in prevalence
RIFAMYCIN	28	*arr-2, arr-3*	2.79%	1	*arr-3*	3.70%	Similar low prevalence
STREPTOMYCIN	352	*aadA1, aadA10, aadA16, aadA2, aadA3, aadA31, aadA5, ant(6)-Ia, aph(3'')-Ib, aph(6)-Id*	35.13%	10	*aadA1, aph(6)-Id, aph(3'')-Ib, aadA5, aadA2*	37.04%	Similar in prevalence
STREPTOTHRICIN	282	*sat2*	28.14%	10	*sat2*	37.04%	Higher in clinical
SULFONAMIDE	286	*sul1, sul2*	28.54%	6	*sul1, sul2*	22.22%	Slightly lower in clinical
TETRACYCLINE	991	*tet(59), tet(A), tet(B), tet(C), tet(D), tet(G), tet(H), tet(J), tet(L), tet(M), tet(Y)*	98.90%	26	*tet(J), tet(A), tet(C)*	100%	Nearly universal presence in both
TRIMETHOPRIM	290	*dfrA1, dfrA10, dfrA12, dfrA14, dfrA16, dfrA17, dfrA19, dfrA27, dfrA32, dfrA42, dfrA46, dfrA5, dfrA7*	28.94%	10	*dfrA1, dfrA17, dfrA32*	37.04%	Higher in clinical
AZITHROMYCIN / CLARITHROMYCIN / CLINDAMYCIN / ERYTHROMYCIN / TELITHROMYCIN	1	*erm(A)*	0.10%	NA	NA	NA	Only in NCBI
AZITHROMYCIN / ERYTHROMYCIN / SPIRAMYCIN / TELITHROMYCIN	15	*mph(A)*	1.5	NA	NA	NA	Only in NCBI

AZITHROMYCIN / ERYTHROMYCIN / STREPTOGRAMIN	26	*msr(E)*	2.59%	NA	NA	NA	Only in NCBI
BLEOMYCIN	52	*ble, bleO*	5.19%	NA	NA	NA	Only in NCBI
CARBAPENEM	171	*blaIMP, blaIMP-27, blaKPC-2, blaKPC-3, blaNDM-1, blaNDM-4, blaNDM-5, blaNDM-50, blaNDM-6, blaNDM-7, blaOXA, blaOXA-23, blaOXA-48, blaVIM-1, blaVIM-78*	17.07%	NA	NA	NA	Only in NCBI
FLORFENICOL	61	*floR, floR2*	6.09%	NA	NA	NA	Only in NCBI
FOSFOMYCIN	14	*fosA, fosA3*	1.40%	NA	NA	NA	Only in NCBI
GENTAMICIN	54	*aac(3)-IId, aac(3)-IIe, aac(3)-VIa, aac(6')-Ib, aac(6')-Ib4, armA*	5.39%	NA	NA	NA	Only in NCBI
GENTAMICIN / KANAMYCIN / TOBRAMYCIN	72	*ant(2")-Ia*	7.19%	NA	NA	NA	Only in NCBI
HYGROMYCIN	13	*aph(4)-Ia*	1.30%	NA	NA	NA	Only in NCBI
KASUGAMYCIN	1	*aac(2')-IIa*	0.10%	NA	NA	NA	Only in NCBI
LINCOSAMIDE	42	*lnu(F), lnu(G)*	4.19%	NA	NA	NA	Only in NCBI
SPECTINOMYCIN / STREPTOMYCIN	1	*aadA27*	0.10%	NA	NA	NA	Only in NCBI

Figure Legends

Fig 1. Phylogenetic tree and pangenome analysis of publicly-available *Proteus mirabilis* genomes from human urine samples. (**1a**) maximum-likelihood phylogenetic tree of 1,001 *P. mirabilis* human urine genomes with metadata arranged as concentric bands. The first two bands display multi-locus sequence type (ST), and the third band indicates the study source in the NCBI database associated with the Bioproject. Bands 4-8 indicate presence (colored bars) or absence (grey) of resistance genes across different antibiotic classes. Distinct clusters were observed that either had resistance genes for nearly all classes of antibiotics or lacked them entirely, which was dependent on ST and independent of geographical source. (**1b**) Roary collector curve showing conserved genes (~1,700 core genome) present in all of the genomes analyzed as well as the number of total genes, which increases as new genomes are added, indicating an open pangenome. (**1c**) Roary collector curve showing unique vs new gene accumulation across all *P. mirabilis* genomes. Here, the total unique genes is >15,000 indicating massive accessory gene diversity and an open pangenome. Early genomes introduce a lot of new genes before dropping out. However, after addition of >250 genomes, new genes (vertical lines) are still accumulated, indicating horizontal gene transfer (HGT), mobile genetic elements, or prophage insertions.

Fig 2. Sequence type distribution of publicly-available *Proteus mirabilis* genomes from human urine samples. (**2a**) Pie-chart illustrating the distribution of MLSTs

1218   grouped by their sample count frequency. (**2b**) Bar plot showing the top 20 MLSTs

1219   based on sample count.

1220

Fig 3: Phylogenetic tree and pangenome analysis of *Proteus mirabilis* strains

isolated from individuals with indwelling urinary catheters. (**3a**) A mid-rooted

maximum likelihood phylogenetic tree of clinical *P. mirabilis* isolates collected from

patients with long-term urinary catheters. The phylogenetic tree displays the participant

ID, the cohort ID (strip1), and multi-locus sequence type (ST, strip 2). Many of the

isolates cluster by ST, but not by geographical location. (**3b**). Mosaic pangenome with

presence/absence matrix of core-genome phylogenetic tree for 26 *P. mirabilis* clinical

isolates collected from patients with long-term urinary catheters.

1229

Fig 4: AMR gene carriage in publicly-available *P. mirabilis* genomes. (**4a**) Number

of *P. mirabilis* strains in the NCBI dataset carrying 1, 2, and 3 tetracycline resistance

(*tet)* genes. (**4b**) Three-way Venn diagram exhibiting the shared chloramphenicol

resistance genes (*catA, catA1,* and *floR*) by all the *P. mirabilis* isolates in the NCBI

dataset. (**4c**) Number of *P. mirabilis* strains in the NCBI dataset carrying 1, 2, and 3

trimethoprim resistance (*dfrA)* genes. (**4d**) Venn diagram of shared sulfonamide

resistance genes (*sul1* and *sul2*). (**4e**) Venn diagram of shared trimethoprim (TMP) and

sulfonamide (SMX) resistance genes.

1238

Fig 5. IS Element landscape overview of clinical isolates. **(5a)** Stacked bar chart of IS family distribution across STs. **(5b)** Scatter plot of total IS count vs. AMR gene count with Spearman correlation line (ρ=0.560, p=0.008). **(5c)** Proximity heatmap showing IS26 distance to each AMR gene in all isolates. **(5d)** Co-occurrence of IS family-AMR genes in isolates with IS within 10kb of AMR genes.

Fig 6. Genetic organization of the T4SS integrative and conjugative elements (ICEs) identified in *Proteus mirabilis*. (**6a**) Genetic structure of the ICE identified in strain H115-00**.** The element includes the integrase gene *intS* and regulatory gene *umuC*, followed by insertion sequence elements (IS3000**,** IS8 family) and multiple accessory genes. A cluster of antibiotic resistance genes is present, including genes such as *bcr**, tetR*, *tetB**, dfrA**, aac,** aph*, and *erm* family members, interspersed with transposition- and recombination-related genes (*tnpA**, xerC).* **(6b)** 2 T4SS-ICEs detected in strain 102-00. Genes are shown as arrows indicating the direction of transcription. The element contains recombination genes *recC* and *uvrC*, followed by multiple accessory and housekeeping genes. Conjugation-associated genes including *tra* components are present toward the right side of the element together with additional metabolic genes. **(6c)** T4SS-ICE detected in strain 106-00. This is a representative for all the non-ST135 and ST145 in our dataset. The ICE is similar to the first ICE observed in ST145 in 6b.

Fig 7: Comprehensive map of the *Proteus mirabilis* Genomic Resistance Island 1 (PmGRI1)-like region in H115-00 (ST135): (7a) Putative PmGRI1 region 1 (3.29 Mb-

1261 3.37 Mb), and **(7b)** Putative PmGRI1 region 2 (3.885 Mb- 4.0 Mb). PmGRI1 is a

1262 chromosomally integrated, modular resistance island identified by tyrosine-family site-

1263 specific recombinase/integrase (~394 aa), insertion at the 3' end of tRNA-Sec, and 20bp

1264 direct repeats flanking the element.

1265

Fig 8: Overview of antibiotic resistance and MIC. (**7a**) Overview of antibiotic

resistance phenotype found in the clinical CAUTI isolates. Color coding denotes

resistance status: red = resistant, yellow = intermediate resistance, and green =

susceptible strains. The presence or absence of resistance-associated genes is

indicated by filled and empty circles, respectively. (**7b**) Heatmap representing the MIC

for each antibiotic tested across clinical *P. mirabilis* CAUTI isolates. Color intensity

corresponds to MIC values, indicating variation in resistance levels.

1273

Fig 9: Dose response curves and MIC. Mean normalized OD600 values at

experimental endpoint (20 hours) are plotted against antibiotic concentrations in ug/ml.

OD is normalized to a no drug control set to 1.0 (maximum growth). The red vertical line

indicates the resistance breakpoint (CLSI) for *Enterobacterales.* The MIC was

determined as the lowest antibiotic concentration for which growth was restricted below

a threshold of a normalized OD of 0.15 (grey horizontal dashed line). (**8a**) Tetracycline

(MIC range tested: 2-20 ug/ml), (**8b**) Kanamycin (MIC range tested: 6-34 ug/ml), (**8c**)

Streptomycin (MIC range tested: 4-22 ug/ml), (**8d**) Ampicillin (MIC range tested: 6-34

ug/ml), (**8e**) Streptothricin (MIC range tested: 4-22 ug/ml).

1283

Fig 10: Dose response curves and MIC. Mean normalized OD600 values at

experimental endpoint (20 hours) are plotted against antibiotic concentrations in ug/ml.

OD is normalized to a no drug control set to 1.0 (maximum growth). The red vertical line

indicates the resistance breakpoint (CLSI) for *Enterobacterales.* The MIC was

determined as the lowest antibiotic concentration for which growth was restricted below

a threshold of a normalized OD of 0.15 (grey horizontal dashed line). (**9a**)

Sulfamethoxazole (MIC range tested: 250-512 ug/ml), (**9b**) Trimethoprim (MIC range

tested: 4-22 ug/ml), (**9c**) TMP-SMX (MIC range tested: 0.5/9.5-16/305 ug/ml).

1292

Fig 11: Dose response curves and MIC of Chloramphenicol. Mean normalized

OD600 values at experimental endpoint (20 hours) are plotted against antibiotic

concentrations in ug/ml. OD is normalized to a no drug control set to 1.0 (maximum

growth). The red vertical line indicates the resistance breakpoint (CLSI) for

Enterobacterales. The MIC was determined as the lowest antibiotic concentration for

which growth was restricted below a threshold of a normalized OD of 0.15 (grey

horizontal dashed line). Chloramphenicol concentrations tested ranged from 6-34 ug/ml.

a

ST135 *

TMP-SMX
Fluoroquinolones
Cephalosporins/Penicilins
Carbapenems
Aminoglycosides
Isolation_Source
MLST-top15
MLST_Status

Band 1: MLST Classification
- Known MLST
- Novel/Untyped

Band 2: Top 15 MLST
- ST96
- ST185
- ST99
- ST135
- ST97
- ST145
- ST133
- ST93
- ST202
- ST146
- ST157
- ST92
- ST196
- ST495
- ST279

Band 3: Study/Isolation Region
- University of Pittsburgh (EDS-HAT)
- CDC (Routine reports)
- Female Urinary Microbiota study
- Rush University (Gut UPEC)
- WUSM (multiple regions)
- Human Bladder Bacteria study
- Antimicrobial Surveillance, Brigham & Women
- Nosocomial infections study
- Bacterial Surveillance for LDH study
- New York City Hospital
- Urine: UTI with AMR study

Bands 4-8: Antibiotic Subclasses
- Aminoglycosides
- Carbapanemes
- Cephalosporins/Penicilins
- Fluoroquinolones
- TMP-SMX

b

Key
— Conserved genes
···· Total genes

c

Key
— New genes
···· Unique genes

a

Tree scale: 0.01 ⊢━━━┤

▷ Strip 1: MLST

Clinical cohorts

- ■ CEA cohort
- ■ HUC cohort
- ■ Reference HI4320

Strips 3-12: Antibiotic Class

- ■ Tetracycline
- ■ Chloramphenicol
- ■ Aminoglycosides
- ■ Carbapenems
- ■ Cephalosporins-Penicillins
- ■ TMP-SMX
- ■ Streptothricin
- ■ Erythromycin + Rifamycin
- ■ Quinolone
- ■ Biocide

b

Clade 1

Clade 2

a

Class 1 integron

Incomplete prophage

AMR genes

aadA2 ereA dfrB2 aar3 catB3 APH(3) APH(6) floR
sul1 OXA1 sul2 tet(A)
qacEΔ1 AAC(6)
tet(C)

IS elements

ISL3 IS26 IS26 IS26 ISL3 IS3

Putative PmGRI1-like genomic island (~138 kb)

3285   3330   3375

Genomic position (kb)

b

Prophage region #10 (Intact, score=130)

Class 1 integron
(dfrA17-aadA5-qacEΔ1-sul1)

AMR genes

catA1

blaTEM-1 APH(3')-Ia dfrA1 aadA5 qacEΔ1 sul1

IS elements

IS66 IS5 IS26 IS26 IS26 IS5 IS5 new

Within PmGRI1-like genomic island (~115 kb)

3885   3900   3915

Genomic position (kb)

Chloramphenicol dose–response

STRAIN ID

102-00	H97-11
102-24	H105-00
102-25	H105-06
104-00	H108-00
106-15	H115-00
201-07	H115-29
203-01	H119-03
206-05	H120-00
206-16	H120-04
210-04	H125-06
H81-01	H127-00
H85-10	H132-05
H94-00	HI4320
H97-00	

Normalized OD (endpoint mean) vs Concentration (µg/mL)

catA+ catA1+

catA+ catA1-

catA+ catA1-

catA- catA1-