Pangenome Analysis of *Proteus mirabilis* Reveals Lineage-Specific Antimicrobial

Resistance Profiles and Discordant Genotype-Phenotype Correlations

Authors: Namrata Deka[1], Aimee L. Brauer[1], Katherine Connerton[1,2], Blake Hanson[3,4],

Jennifer N. Walker[5,3,4], Chelsie E. Armbruster[1*]

*address correspondence to Chelsie E. Armbruster, chelsiea@buffalo.edu

[1]Department of Microbiology and Immunology, Jacobs School of Medicine and

Biomedical Sciences, Buffalo, NY, 14203, NY, USA. [2]School of Nursing, Niagara

University, Buffalo, NY, USA. [3]Department of Epidemiology, UTHealth Houston School

of Public Health; [4]Center for Infectious Diseases, UTHealth Houston School of Public

Health; [5]Department of Microbiology and Molecular Genetics, McGovern Medical

School, The University of Texas Health Science Center at Houston, Texas, USA.

Author Contributions: ND and CEA designed experiments. ND, KC, and ALB
conducted MIC experiments, ND and CEA analyzed results. ND performed genome
sequencing, pangenome analysis, and AMR detection. BH and JW provided guidance
on pangenome analysis, AMR detection, and data visualization. ND visualized the data
and prepared the first draft of the manuscript. CEA provided project administration. All
authors critically reviewed and revised the manuscript.
22

23 **Conflict of Interest:** All authors declare no conflicts of interest.

24 **KEYWORDS:** *Proteus mirabilis*; antimicrobial resistance; pangenome; antimicrobial
25 susceptibility testing; multi-locus sequence typing; trimethoprim; sulfamethoxazole;
26 chloramphenicol; urinary tract infection

ABSTRACT (<250 words)

Urinary tract infections (UTIs) impose a substantial health care burden with increase

antimicrobial resistance and treatment failure rates. *Proteus mirabilis* is a challenging

UTI pathogen due to intrinsic resistances coupled with formation of crystalline biofilms.

We combined pangenome analysis, antimicrobial resistance gene (AMR) predication,

and antimicrobial susceptibility testing (AST) to determine whether highly multidrug-

resistant (MDR) isolates of *P. mirabilis* arise from distinct lineages and explored the

clinical utility of multilocus sequence typing (MLST). The pangenome of 1,002 *P.

mirabilis* genomes from human urine revealed an open conformation driven by strain

diversity and the accessory genome. A total of 213 sequence types (STs) were

identified and only 7% had ≥11 genomes, highlighting strain diversity. 93% of the *P.

mirabilis* genomes harbored resistance genes for ≥2 antibiotic subclasses, and 25%

were predicted to be resistant to >6 subclasses, confirming a high MDR burden. By

focusing on the 15 most prevalent STs, we observed that AMR subclasses were largely

lineage-specific. However, isolates with very high resistance gene counts (>20) were

distributed across multiple STs, indicating that extreme resistance gene accumulation is

not restricted to ST. Comprehensive AST of 27 *P. mirabilis* clinical isolates further

revealed variable genotype-phenotype concordance, indicating unresolved mechanisms

of resistance not captured in current AMR prediction databases. In summary, our study

emphasizes the critical need to expand genomic surveillance of *P. mirabilis* to fully

understand the complex landscape of AMR in this species and translate these insights

into improved diagnostic and therapeutic strategies.

IMPORTANCE (<150 words)

Proteus mirabilis is a clinically-challenging cause of urinary tract infections due to multidrug resistance and its ability to form crystalline biofilms that provide further antibiotic protection. In this study, we sought to determine how well sequence typing and antimicrobial resistance gene prediction correlate with laboratory-based antimicrobial susceptibility testing. By analyzing more than 1,000 *P. mirabilis* genomes from human urine samples, we found that some resistance patterns were sequence type-specific. However, the genome structure of this species suggests frequent horizontal gene transfer, and the most highly-resistant strains did not cluster by lineage. Importantly, many isolates that appeared "susceptible" based on their genomes were in fact resistant upon laboratory testing, revealing hidden or uncharacterized resistance mechanisms. These findings show that current gene-based prediction tools can miss clinically relevant resistance, underscoring the need for further study to guide effective treatment of *P. mirabilis* infections.

INTRODUCTION

Urinary tract infections (UTIs) are among the most common bacterial infections, with an estimated healthcare cost of >$3 billion in the United States [1-4]. They are a leading cause of antimicrobial prescriptions in both outpatient and inpatient settings, with 50-60% of women experiencing at least one UTI in their lifetime [5]. UTI can be classified as complicated UTI (cUTI) or uncomplicated UTI (uUTIs) depending on the presence and absence of structural or functional abnormalities of the urinary tract, catheterization, or immunocompromised states [5]. Recent epidemiological data estimate approximately 174,000 cases of uUTI and 2.38 million cases of cUTI annually [6, 7]. Despite standard antibiotic therapy, treatment failure rates for UTIs remain prevalent, occurring in 16.7% of cases in adult female outpatients with cUTI, 27% of elderly men, and 15% of elderly women, often leading to recurrence, secondary infection, progression to pyelonephritis, or urosepsis [8, 9].

Among uropathogens, *Proteus mirabilis* is an important Gram-negative bacterium and a major cause of UTIs, contributing to approximately 4.2% of uUTIs and 4.7% of cUTIs [5]. Notably, *P. mirabilis* accounts for up to 45% of catheter-associated UTIs (CAUTIs) and 13-21% of secondary bacteremias, making it a key pathogen in medical device-associated infections [5, 10, 11]. *P. mirabilis* has a strong propensity for biofilm formation and causes encrustation of the catheter by deposition of struvite and apatite crystals, leading to persistent infections and catheter blockage. Struvite and apatite crystals form due to the production of a potent urease enzyme by *P. mirabilis* that increases urine pH, promoting ion precipitation, crystal deposition, and urinary stone formation [10, 12-15]. It can remain in bladder crystalline deposits despite multiple

doses of antibiotic treatment, catheter changes, and even periods without

catheterization [15-17]. Infection with *P. mirabilis* can become fatal when it progresses

to symptomatic CAUTI, acute bacteremia, and urosepsis due to its ability to disseminate

from the bladder to other organs [14, 18].

Antimicrobial resistance (AMR) in UTI pathogens is an escalating global concern,

with increasing reports of multidrug-resistant (MDR) strains complicating treatment [19].

Treatment failure for *P. mirabilis*-associated UTIs has been linked to MDR, leading to

prolonged hospitalization, increased healthcare costs, and higher morbidity rates [15,

19, 20]. Treatment typically involves antibiotics but becomes challenging due to intrinsic

and acquired resistance mechanisms, including AmpC β-lactamase production, efflux

pumps, and extended-spectrum β-lactamase (ESBL) expression [14, 21-23]. These

factors contribute to resistance to first-line UTI antibiotics such as trimethoprim-

sulfamethoxazole (TMP-SMX), fluoroquinolones, and β-lactams. The prevalence of

ESBL genes like *blaTEM* and *blaCTX-M* further exacerbates resistance, while mobile

genetic elements such as SXT/R391 integrative and conjugative elements (ICEs) play a

crucial role in the spread of AMR genes in *P. mirabilis* strains [19, 24, 25].

Numerous studies have characterized the model *P. mirabilis* strain HI4320,

originally isolated from the female urinary tract, including virulence factors such as

adhesins for attachment and biofilm formation, flagella for motility and dissemination,

urease enzyme activity leading to renal calculi, and iron-binding siderophores [10, 26].

However, a major gap remains in our understanding of the diversity of strains isolated

from clinical urine samples, either during infection or in the presence of a catheter.

Multilocus sequence typing (MLST) provides an unambiguous method for characterizing

bacterial isolates based on the sequences of internal fragments of typically 6-7

housekeeping genes. An open question in the field is whether specific sequence types

(STs) in *P. mirabilis* are associated with the accumulation of similar AMR and virulence

genes. While previous studies in other bacterial species suggest that certain lineages

correlate with varying levels of resistance and biofilm production, such associations in

P. mirabilis remain largely unexplored. For example, in methicillin-resistant

Staphylococcus aureus (MRSA), expression of the *ica* genes—critical for biofilm

formation—varies among different STs, with certain STs showing higher *ica* gene

expression, particularly in strains isolated from wound and catheter samples [27]. If

similar correlations exist in *P. mirabilis*, routine sequence typing of clinical isolates could

serve as a valuable tool to predict AMR potential and pathogenicity.

Despite advances in next-generation sequencing (NGS), *P. mirabilis* has not

been typed extensively at the genomic level. Several typing schemes have been

developed to characterize *P. mirabilis* strains, but no standard scheme is followed for

clinical classification. Core genome MLST (cgMLST), a high-resolution genomic

method, has identified 205 clonal groups (CGs) linked to severe UTIs and AMR [28].

Some other older methods include proticine production and sensitivity typing, which

differentiates strains based on bacteriocin activity, and bacteriophage typing, which

classifies strains by susceptibility to phage lysis [29, 30]. Additionally, Dienes typing,

based on swarming behavior, provides a simple means of strain differentiation, while

serotyping classifies strains based on O-antigen structures, revealing a diverse

antigenic profile among clinical isolates [31]. Recent comprehensive genomic studies by

Potter *et. al.* and Lian *et. al.* have given insights into some key genomic features of *P.*

mirabilis populations and distribution [32, 33]. However, there remains a lack of data on

the prevalence of *P. mirabilis* isolates through a single classification system and the link

between ST to AMR genes in urine isolates. This gap in knowledge highlights the need

to investigate whether specific STs are predictive of resistance patterns or increased

virulence within the urinary tract. By focusing on genomes isolated from human urine

samples, we aim to gain critical insights into AMR and virulence of *P. mirabilis* in urine

to have better indicators for risk of treatment failure and severe disease.

There are numerous ways to predict AMR, including the standardized clinical

laboratory practices of testing resistance phenotypes to specific antibiotics, as well as

genomic or machine learning methods to predict resistant gene expression [34].

Genomic prediction tools can be useful for quick detection of AMR in clinical practice,

and there may be a correlation between AMR gene prediction and phenotypic

expression, but the nature of the correlation can vary depending on specific genes and

the bacterial species. For instance, *Escherichia coli* frequently harbors resistance genes

for β-lactams, sulfonamides, trimethoprim, and chloramphenicol on mobile genetic

elements, leading to a stronger correlation between genotype and phenotype [35-37]. In

contrast, according to some studies, *Proteus mirabilis* carrying β-lactam and

sulfonamide (*sul1* and *sul2*) genes exhibits this correlation less consistently, showing

weaker alignment between genomic predictions and observed resistance profiles [35,

38-40].

Tools like AMRFinder and CARD are accurate *in silico* prediction tools;

AMRFinder was able to validate the presence of genes pertaining to AMR phenotypes

for *Salmonella enterica, Campylobacter spp.*, and *Escherichia coli* with 98.4%

consistent predictions [41]. Further, a 99% concordance was found between phenotypic

antimicrobial susceptibility testing and *in silico* prediction of AMR genes using whole-

genome sequencing (WGS) in 1,321 isolates of *Salmonella enterica* from human

infections in Canada [20]. Similarly, a machine-learning approach was used to predict

the Minimum Inhibitory Concentrations (MIC) of ciprofloxacin in *Escherichia coli* where

MIC was predicted accurately in 65%-93% of the samples within a 2-to-4-fold dilution

range [42]. While genetic data provides a promising avenue to predict AMR, it is

important to understand the limitations and variability in gene expression and resistance

phenotypes in different bacteria. There has been no such correlation established for *P.*

mirabilis, and it would be particularly important to examine for urine isolates. Hence,

integrating genotypic AMR prediction and phenotypic MIC testing in diverse strains may

offer a more comprehensive approach to predicting antimicrobial resistance in this

species.

In this study, we conducted a comparative genomic analysis of *P. mirabilis* using

1,001 publicly available human urine isolate sequences to characterize the pangenome,

prevalent MLST in clinical isolates, and AMR profiles. Additionally, we compare these

findings with 26 clinical isolates collected from patients with long-term indwelling

catheters [15, 17]. Given the increasing challenges associated with AMR and persistent

infections, we sought to identify predictive factors that correlate genomic data with

resistance mechanisms in a clinical setting. By evaluating urine isolates from both

community and healthcare settings, we aimed to provide a comprehensive

understanding of *P. mirabilis* antimicrobial susceptibility patterns and resistance

determinants. We found that *P. mirabilis* isolates from urine exhibit a heterogenous

AMR gene profile that is independent of collection source. However, similar MLSTs

clustered together regardless of isolation source and showcased related AMR profiles.

Importantly, the intrinsic resistance of *P. mirabilis* to tetracycline and the high incidence

of TMP-SMX resistance eliminates the use of two primary UTI therapy agents. This

necessitates reliance on more expansive broad-spectrum alternatives, fueling the AMR

crisis. Our study also demonstrates that numerous *P. mirabilis* CAUTI isolates that

would be classified as antibiotic susceptible by AMR gene prediction were resistant

upon phenotypic testing, indicating that additional as-yet unidentified resistance

mechanisms may be prevalent in urine isolates of this species and likely to influence

treatment efficacy in clinical settings.

MATERIALS AND METHODS

Data availability statement

Sequencing data have been deposited in NCBI BioProject PRJNA1367153: *Proteus mirabilis* genome sequences from human urine sources in the United States. All other data are available within the article and its supplemental material.

Ethics approval

This body of work includes sequencing of *P. mirabilis* strains that were previously isolated from human subjects in two different studies, for which human subject protocols were approved by the University of Buffalo (IRB #STUDY00002526) and Washington University School of Medicine (IRB #201410058).

Patient population and Sample collection

To investigate whether *P. mirabilis* MLSTs correlate with AMR patterns, we assessed *P. mirabilis* genomes from three separate cohorts: publicly available genomes from NCBI (**Table S1**), nursing home residents with long-term catheters (CEA isolates), and community-dwelling catheterized patients (HUC isolates).

For comparison of *P. mirabilis* genomes, we sought to query all publicly available high-quality genomes of *P. mirabilis* isolates obtained from urine samples in hospital environments in the United States. Using advanced search parameters, publicly available raw genome data were retrieved from the NCBI database. The initial search, conducted in October 2024, yielded 1,337 hits using the terms "*Proteus mirabilis*,"

isolation source "urine," and host "*Homo sapiens*." Filtering the results to include only

isolates originating from the "United States" reduced the dataset to 1,152 genomes. The

dataset was further refined by only including whole-genome sequencing data. These

filters resulted in a final dataset of 1,064 (**Table S1**) genomes, which were subsequently

downloaded from NCBI. Following FastQC v0.11.9-Java-11 analysis and adapter

trimming using trimmomatic v0.39-Java-11.0.16. [43, 44], 1,026 genomes were retained

for downstream analyses. Samples were excluded if they exhibited poor quality, had

high GC content, high read duplication rates, or if paired end read files were missing (46

samples). Raw paired end read sequences were assembled using spades v3.15.5 [45].

After genome assembly, we assessed quality and completeness using QUAST (**Table**

S2), selecting 1,001 high-quality genomes with >95% completeness and fewer than

1,000 contigs which were annotated with prokka v1.14.5 to facilitate downstream

analyses [46, 47].

Isolates designated 101–210 were cultured from urine specimens collected

during a prospective cohort study on asymptomatic catheter-associated bacteriuria at

two nursing homes in Buffalo, New York, USA, between July 2019 and March 2020 [15].

Inclusion criteria required participants to be 21 years or older at the time of consent and

have an indwelling urinary catheter in place for at least 12 months at the start of the

study. Study participants underwent an initial enrollment visit, followed by weekly study

visits for up to seven months. Most of the study participants were White (79%) and male

(79%), an average age of 65 years, and 68% had suprapubic catheters [15].

Participants exhibited a high degree of functional dependence in activities of daily living,

with the most common comorbidities including neurogenic bladder (74%), hemiplegia

(42%), diabetes (32%), renal disease (32%), multiple sclerosis (26%), and chronic heart

failure (26%) [15]. Urine specimens were cultured on MacConkey agar and Columbia

Nutrient Agar, and Gram-negative bacteria were identified to the species level using

API-20E test strips (BioMérieux). *P. mirabilis* species designation was further confirmed

based on characteristic swarming motility on blood agar plates (Hardy Diagnostics) and

whole genome sequencing (**Table S3**).

Additional *P. mirabilis* isolates with designations of "HUC" were obtained from the

lab of Scott J. Hultgren, Washington University School of Medicine [17]. These isolates

were derived from urine samples collected from community-dwelling catheterized

patients during routine catheter removals in the urology clinic as part of their standard

care. Most of the participants were male (64%) with an average age of 64 years.

Inclusion criteria required participants to be 18 years or older at the time of consent and

scheduled for the removal of an indwelling urinary device, including urethral and

suprapubic catheters or ureteral stents. The primary indications for catheterization

included bladder outlet obstruction (33%), peripheral or central neuropathy (40%),

rectovesical or rectourethral fistula (7%), stress urinary incontinence (7%), atonic

bladder (11%), and temporary post-operative complications (2%) [17]. Bacterial colonies

were distinguished based on morphology, size, and color, and 1–9 single colonies per

sample were selected for DNA extraction. *P. mirabilis* isolates were identified via 16S

rRNA sequencing and confirmed by whole genome sequencing (**Table S3**).

Whole Genome Sequencing of Clinical Isolates

Freezer stocks were streak-plated on MacConkey agar to isolate single colonies of

Proteus mirabilis. 5 ml of Low Salt Luria Bertani (LSLB; 0.1 g/L NaCl, 10 g/L Tryptone, 5

g/ yeast extract) broth was inoculated with a single colony and incubated for 16 hours at

37°C with shaking. Genomic DNA was extracted using the DNeasy Blood and Tissue

Kit (Qiagen, Germantown, Maryland, USA) according to the manufacturer's instructions

with specific modifications to ensure high purity of DNA; samples were incubated in

proteinase K for 2 hours instead of 10 minutes, and three washes of AW2 buffer were

conducted to ensure high-quality DNA free of salts. Samples were sent to SeqCoast

Genomics for whole genome sequencing (WGS). All samples were prepared using the

Illumina DNA Prep tagmentation kit and IDT For Illumina Unique Dual Indexes.

Sequencing was performed on the Illumina NextSeq2000 platform using a 300-cycle

flow cell kit to produce 2x150bp paired reads, as previously described [48]. 1-2% PhiX

control was spiked into the run to support optimal base calling. The sequencing data

has been uploaded in BioProject PRJNA1367153: *Proteus mirabilis* genome sequences

from human urine sources in the United States.

Read trimming and run analytics were performed using trimmomatic v0.39.0 [44].

The processed reads were assembled using spades v3.15.5 [45] . The genome

assemblies were quality-checked using QUAST v5.2.0 selecting genomes with fewer

than 500 contigs (**Table S3**). Genomes that passed quality filtering were annotated

using prokka v1.14.5 [47].

Multi Locus Sequence Typing (MLST) and Phylogenetic trees

Multilocus sequence typing (MLST) analysis was conducted to identify the

sequence types (STs) present in the dataset using mlst software, which incorporates

the PubMLST database [49, 50]. The annotated genomes were then fed into roary

v3.13.0 using a core threshold cluster of 95%, to identify their pangenome profile and

develop a gene presence/absence matrix [51]. The filtered core genome alignment was

constructed using FastTree v2.1.11. The resulting newick file was visualized as a mid-

rooted phylogenetic tree using the iTOL website and annotated with source, and

antibiotic metadata files [52, 53].

Drug Resistance Gene Prediction

We used AMRFinderPlus, which uses the Bacterial Antimicrobial Resistance

Reference Gene Database to predict AMR genes from the coding sequence of the *P.*

mirabilis genomes including the 26 CEA and HUC strains and clinical reference strain

HI4320 [34, 41].

Antibiotic Susceptibility Testing (AST)

The clinical strains were grown overnight in 5 mL LSLB broth at 37°C with shaking.

Antibiotic solutions: Tetracycline HCl (Research Products International (RPI), CAS: 64-

75-5), Chloramphenicol (RPI, CAS: 56-75-7), Streptomycin (RPI, CAS: 3810-74-0),

Streptothricin (Goldbio, CAS: 96736-11-7), Sulfonamide (sulfamethoxazole, Cayman

chemical company, CAS: 723-46-6), Trimethoprim (Cayman chemical company, CAS:

738-70-5), Ampicillin (RPI, CAS: 69-53-4), and Kanamycin (RPI, CAS: 8063-07-8), were

prepared in Mueller-Hinton Broth (MHB) at a concentration range encompassing the

sensitive-to-resistant spectrum according to Clinical Laboratory Standards Institute

(CLSI) guidelines for *Enterobacteriales* [54]. Stock solutions of all the antibiotics were

prepared and stored at −20°C, and working dilutions were made fresh from the stock

immediately prior to testing.

1ml of the overnight cultures were centrifuged at 1,000g to pellet the cells. The

supernatant was removed, and the pellets were resuspended in 0.9% saline to wash

away residual LB. The inoculum was adjusted to 4×10^5 CFU/ml in MHB. One antibiotic

with increasing concentrations was tested per 96-well plate, with four technical and

duplicate biological replicates. The plates were incubated at 37°C with double-orbital

shaking in Biotech Synergy H1, and the OD_{600} was measured every 15 minutes for 20

hours to monitor bacterial growth with OD_{600} threshold set at 0.15.

RESULTS

To assess the population structure and genetic diversity of *Proteus mirabilis* in the urinary tract, we analyzed 1,001 publicly available genomes isolated from human urine specimens (**Table S1**). The genomes downloaded from NCBI that were used in this analysis were predominantly derived from clinical settings, focusing on medical surveillance and characterization of urinary samples.

Key contributing sources include research from Washington University School of Medicine (WUSM), the University of Pittsburgh Enhanced Detection System for Healthcare-Associated Transmission (EDS-HAT) study, and national antimicrobial surveillance programs such as the Center of Disease Control's Healthcare-Associated Infection Sequencing (CDC HAI-Seq) Gram-Negative Bacteria Study and studies on the human urinary microbiome. Among the genomes, contig counts ranged from 60 to 973, with an average of 258 and a median of 180 contigs. The GC content averaged 38.76%, with a median of 38.69%, making them suitable for downstream analyses (**Table 1 and S2**). While all sequences were confirmed to be derived from urine samples, the clinical context of these specimens—whether they were associated with asymptomatic bacteriuria or infection—is not known.

Phylogenetic analyses (mid-rooted tree) revealed a monophyletic, star-like phylogeny, with a central ancestral node from which several peripheral branches radiate (**Fig 1**), which may indicate recent diversification events. When considering branching, a subset of genomes branched out creating three clades, *clade 1* (964 genomes, dominant), *clade 2* (5 genomes), and *clade 3* (32 genomes) (**Fig S1**). This breakdown is consistent with previously reported intra-species core genome variation in *P. mirabilis*

[32, 33, 55]. For example, Potter *et. al.* studied a cohort of 2,060 *P. mirabilis* genomes

collected worldwide from multiple anatomical sources and observed that the essential

core genes remained relatively conserved with three divergent sub-species. [33].

Pan-genomic analysis of *P. mirabilis* reveals a mosaic structure

To characterize the pangenome of *P. mirabilis* from human urine samples we

assessed its structural composition, including core, shell, and cloud gene categories. In

the 1,001 genomes analyzed, a total of 41,121 gene families were identified. Of these,

the core genome (genes conserved in >99% of the samples analyzed) is small with

1,937 genes indicating a high degree of conservation. Considering that a typical *P.*

mirabilis genome harbors roughly 4,600-4,900 coding sequences (CDS), the core

genome accounts for approximately 40-45% of the average gene content. This indicates

a high conserved genomic backbone with 4.7% of it supporting essential cellular

functions, coexisting with an expansive and dynamic accessory genome (**Fig S1**). The

shell of the 1,001 genomes contained 1,749 genes (4.08%), and 531 soft core genes

shared among a subset of strains that likely contribute to niche-specific adaptations.

The majority of genes (37,204 genes, 90%) were classified as cloud genes, making the

accessory component the largest in the pangenome. Having a small, stable core and

large, variable accessory genome likely represents strain-specific differences driven by

horizontal gene transfer.

The pangenome exhibited an open conformation representing a mosaic pattern

(**Fig S1**), with the size expanding as additional genomes were analyzed (**Fig 1b**). This

expansion reflects the continuous acquisition of novel gene families, highlighting the

dynamic nature of the *P. mirabilis* genome and its capacity for adaptation through

horizontal gene transfer (**Fig 1c**). The stability of core genes is crucial for the consistent

expression of virulence factors, while the expanding accessory genes may help adapt to

polymicrobial environments and antibiotic pressures. These findings indicate the genetic

plasticity of *P. mirabilis* isolated from urine and provide a foundation for future studies

on its pathogenic potential and AMR mechanisms in the urinary tract.

Publicly available *P. mirabilis* genomes comprise 213 defined and 118 undefined

MLSTs

The specific genes used in MLST of *P. mirabilis* are a combination of

housekeeping genes *atpA, dnaJ, mdh, pyrC, recA,* and *rpoD* [49, 50]. We identified a

total of 213 defined STs and approximately 118 novel or undefined genomes, reflecting

the diversity of *P. mirabilis* human urine isolates within the dataset (**Fig 1**). The untyped

MLST have been submitted to the PubMLST database to update the schema in *P.*

mirabilis.

A detailed analysis of ST prevalence within the dataset revealed that certain STs

were more frequently observed. The distribution of MLSTs across the dataset was

highly skewed (**Fig 2a, Sup Table S4 and S5**), with 46.26% of the STs having only one

representative genome in the dataset while ~7% of the STs had ≥11 genomes (**Fig 2a**).

Many genomes in the same ST came from the same study/location but it was not the

rule as the top 15 overrepresented ST's show mixed isolation sources (**Fig 1a**). The

uneven ST distribution is further visualized in a bar chart of the top 20 MLSTs,

highlighting the dominance of a few clonal lineages across sampled populations (**Fig**

2b). ST96 was the most abundant (49 genomes; 4.90%), followed by ST185 (42

genomes; 4.20%), ST99 (38 genomes; 3.80%), ST135 (38 genomes; 3.80%), and ST97

(37 genomes; 3.70%). Notably, genomes from each of these STs clustered together in

distinct lineages, even if isolated from different geographical locations and reported in

different studies (**Fig 1a**). This pattern supports the core genome phylogenetic tree

clustering and also that accessory gene variation in the whole genome in *P. mirabilis*

are closely linked to clonal lineages, meaning that the non-essential adaptive genes of

P. mirabilis are not random but are co-related to its evolutionary history linking multiple

sources [33].

Phylogenetic analysis reveals shared lineages and novel variants across clinical

P. mirabilis **isolates**

To begin further examining the clinical relevance of *P. mirabilis* MLSTs, we next

sequenced and analyzed a collection of 25 clinical isolates from two distinct cohorts

(CEA isolates collected from the urine of nursing home residents with long-term

indwelling catheters, and HUC isolates collected from the urine of catheterized

community-dwelling individuals during routine catheter changes), as well as the

reference strain HI4320 that was previously isolated from the urine of a catheterized

nursing home resident (**Fig 3a**). The analysis revealed a population structure forming

two primary clades, termed *clade 1* (23 isolates*)* and *clade 2* (3 isolates) (**Fig 3b**). Lack

of the third clade identified in the 1,001 genomes was expected considering its low

frequency (0.5%) and the small sample size of the clinical isolate collection.

Phylogenetic clustering was in accordance with the core genome and gene content

irrespective of the clinical origin (HUC vs CEA, **Fig 3a**). Across the 26 isolates, we

identified 15 distinct STs; 10 STs were identified in the HUC cohort, 4 in the CEA cohort

(includes a novel ST), and 1 represented solely by HI4320. Six of the 10

overrepresented STs in the NCBI genome dataset were also represented among the

clinical isolates; namely STs 96, 97, 135, 145, 146, and 185. ST185 was observed in

both the CEA and HUC cohorts (H120-04 and 102-00, 206-16), demonstrating genetic

overlap across different geographic locations. The two most prevalent STs in the NCBI

dataset were also the most prevalent in the clinical isolates: ST185 (3 isolates, 12%)

and ST96 (3 isolates, 12%). This concordance with the NCBI dataset suggests these

lineages are not only geographically widespread but also clinically prevalent, likely due

to conserved traits (e.g., AMR, virulence, or niche adaptation) that enhance their

transmission and success across diverse environments. Their presence also confirms

that our clinical cohort captured high-impact, epidemiologically relevant STs rather than

rare or sporadically enriched variants even with a limited sample size.

Phylogenetic analyses demonstrated moderate genetic diversity overall, with

branch lengths suggesting varied degrees of relatedness. The pangenome of the

isolates further highlights the heterogenicity within the clinical isolates (**Fig. 3b**),

supporting the mosaic lineage of these isolates. Three isolates (H94-00, H81-01, H105-

06) exhibited genetic divergence from the rest of the *P. mirabilis* strains (clade 2),

possibly due to horizontal gene transfer or independent evolution. The full summary of

isolated sources and STs are provided in **Table 2 and Table S3.**

428   ***Proteus mirabilis* urine isolates possess resistance factors to multiple subclasses**

429   **of antibiotics**

430   Understanding AMR profiles in *P. mirabilis* is crucial to understand its ability to

431   resist commonly used antibiotics including β-lactams, aminoglycosides, sulfonamides,

432   and trimethoprim, since AMR complicates treatment strategies during UTI and

433   contributes to the global public health burden. Analyzing the AMR genes in publicly

434   available *P. mirabilis* genomes can also provide insights into the dissemination of

435   resistance determinants, guiding surveillance and control measures. We first examined

436   the AMR profiles of the 1,001 publicly available *P. mirabilis* genomes. A total of 24

437   antibiotic resistance subclasses were identified, reflecting the diverse mechanisms by

438   which this pathogen can evade antibiotics. The most prevalent genes were those

439   conferring resistance to tetracycline and chloramphenicol, observed in nearly all

440   genomes (**Table S6**).

441   *P. mirabilis* is known to have intrinsic resistance to tetracycline, so it was not

442   surprising that tetracycline resistance genes were detected in 991 (99.0%) of the 1,001

443   genomes, with *tetJ* specifically encoded by 985 of those 991 (99.5%) (**Table 3**). Several

444   other tetracycline resistance determinants, including *tetA, tetB, tetC*, and *tetD*, were

445   detected at lower frequencies. Interestingly, 934/991 (94.2%) of the genomes carried

446   only a single *tet* gene, while 48 (4.8%) carried 2, and 9 (0.9%) carried 3 *tet* genes (**Fig**

447   **4a**). In cases of multiple tetracycline resistance determinants, the genes may be

448   functionally redundant or may have been acquired during co-selection of other

449   antibiotics. Notably, a small number of genomes (11, 1.1%) had no detectable

450   tetracycline resistance genes. This could represent a true absence of the genes, or it

could be a result of fragmented assemblies or sequence dissimilarity not found in the

AMR prediction database.

Chloramphenicol resistance was detected in 952 (95%) of the genomes, with 748

harboring the *catA* gene. Additionally, 69 genomes harbored both *catA* and *catA1,* 43

harbored *catA* and *floR,* and the presence of all three genes was seen in 15 genomes

(**Fig 4b, Table S6**). These findings highlight the prevalence of phenicol resistance in *P.*

mirabilis genomes despite limited use of chloramphenicol in modern clinical settings,

suggesting that selective pressures from other environments or co-selection with other

resistance genes may play a role.

Resistance to aminoglycosides, a critical class of antibiotics for treating severe

Gram-negative infections, was predicted in 716 (71%) of the genomes. Notably, we

observed streptomycin resistance genes in 352 genomes, suggesting the persistence of

resistance despite limited clinical use (**Table 3**). The presence of multiple

aminoglycoside resistance determinants indicates that *P. mirabilis* maintains a robust

arsenal against this class of antimicrobials, including amikacin, gentamycin, kanamycin,

streptomycin and tobramycin.

Resistance to trimethoprim, mediated by *dfrA* alleles, was detected in 290 (29%)

genomes. The most prevalent allele was *dfrA1* (n=248), followed by *dfrA17* (n=44),

dfrA32 (n=17), *dfrA5* (n=12), *dfrA19* (n=7), *dfrA14* (n=7), *dfrA7* and *dfrA16* (n=4), *dfr12*

and *dfrA27* (n=2), and *dfrA10, dfrA42*, and *dfrA46* (n=1). 235 (81.02%) genomes had a

single *dfrA* gene, while 50 (17.2%) carried 2, and 5 (1.72%) carried 3 *dfrA* genes (**Fig**

4c). The diversity of *dfrA* genes highlights the adaptive potential of *P. mirabilis* to

acquire and maintain resistance determinants. Resistance to sulfonamides was similarly

predicted in 286 (28%) genomes, with *sul1* detected in 180, *sul2* in 187, and 81

genomes harboring both *sul1* and *sul2* genes (**Fig 4d, Table S6**). Importantly, 223

(22%) genomes were predicted to encode resistance determinants for both trimethoprim

and sulfonamide (**Fig 4e**). Since trimethoprim and sulfamethoxazole are often used in

combination, the co-occurrence of these genes, potentially mediated by mobile genetic

elements, would likely complicate efficacy of combined treatment. Additionally, we

detected 185 genomes (18%) with Extended Spectrum β-Lactamase (ESBL)-associated

genes. Of these, 148 harbored the *blaTEM-1* gene, expected to confer resistance to

ampicillin (**Table 3**).

Finally, the number of resistance subclasses detected per *P. mirabilis* genome

ranged from 1 to 24, with an average of 4.61 AMR subclasses per isolate. A total of 71

genomes (7%) exhibited resistance to only a single antibiotic subclass, most commonly

tetracycline. In contrast, 931 (93%) had resistance to two or more subclasses, and a

substantial subset of 246 genomes (25%) harbored more than six antibiotic subclasses,

indicating a high prevalence of multidrug resistance in this sample set. 176 genomes

(18%) had at least one resistance determinant from all five of the above resistance gene

groups, while 56 (6%) had combinations of tetracycline, chloramphenicol, and β-lactam

resistance genes. Additionally, phylogenetic analyses (**Fig 1a**) determined that certain

clusters consistently had resistance genes to nearly all antibiotic classes shown, while

others exhibited a complete absence of resistance genes, independent of study source.

Antibiotic resistance subclasses have a common distribution within lineages

To evaluate the relationship between ST and AMR burden in *P. mirabilis*, we

quantified the number of predicted AMR subclasses per genome and summarized these

across 215 distinct STs. Analysis of AMR gene burden across the STs revealed

significant lineage-associated differences in resistome size. Several STs exhibited AMR

subclass counts substantially exceeding the dataset-wide mean of 4.4 subclasses per

genome. Focusing on the STs with ≥3 genomes, ST431, ST425, ST485, ST450, ST421,

ST284, ST269, ST 417, and ST 335 demonstrated the highest mean resistance

subclass burdens, each surpassing twice the overall mean (**Fig 5a**). For example,

ST485 maintained a high AMR burden (subclass range:11-12 subclasses) with minimal

intra-ST variability, suggesting a clonally stable and highly resistant lineage. In contrast,

ST431 (range: 9-20 subclasses), ST425 (range: 5-24 subclasses) and ST417 (range: 4-

15 subclasses) exhibited broad resistance distributions, indicating greater intra-lineage

heterogeneity. This diversity may reflect horizontal gene acquisition or ongoing

divergent evolutionary trajectories and underscores the complex dynamics of AMR

dissemination within clonal backgrounds.

At the population level, the geometric mean resistome size was 3.6 genes,

reflecting the typical genome. However, the arithmetic mean was higher at 5.5 genes

due to a small subset of highly resistant genomes. 93.8% of our dataset harbored

resistance genes for ☐10 subclasses and 99.8% of our dataset had ☐20 subclasses,

with the 80[th] percentile at ~6 subclasses (**Fig 5b**). More than 450 genomes had

approximately 5 resistance genes, while 35 exhibited elevated AMR gene counts that

extend beyond 20 resistance genes; 4 genomes had 30 gene counts, and 1 extreme

case had 36 gene counts that skewed the arithmetic mean. Together, these data

indicate that the majority of *P. mirabilis* genomes from this collection encode low to

moderate numbers of resistance genes, with only a small fraction exhibiting very high

AMR gene carriage.

To further explore AMR-ST associations, we analyzed the distribution of

resistance subclasses and number of genes across the 10 most prevalent STs where

notable inter-lineage variation was observed (**Fig 5c, 5d**). ST96 (n=49 isolates), the

most prevalent lineage, exhibited the highest frequency and diversity of resistance

genes (range:1-17, mean: 3.7). These genes conferred resistance to aminoglycosides

(e.g., amikacin, tobramycin, apramycin, gentamicin), macrolides (azithromycin),

streptogramins, and fosfomycin, in addition to tetracycline and chloramphenicol. 14

genomes from ST96 exhibited resistance to >6 antibiotic subclasses, highlighting

extensive genetic adaptation, and this was also observed in 7 genomes from ST97 (n=

37 genomes, range: 1-16, mean: 3.27). Similarly, ST185 (n=42 genomes, range: 1-10,

mean: 3.78), ST99 (n=38 genomes, range: 1-12, mean: 3.36) and ST135 (n= 38

genomes, range:1-8, mean: 3.05) frequently displayed resistance to >4 subclasses (**Fig**

5c, 5d). Most genomes carried modest resistomes, with those possessing more than 18

genes being uncommon (80[th] percentile); 93.5 **%** of genomes had fewer than 30

resistance genes, and 98.3 **%** had 50 or fewer genes (**Fig. 5e**). The AMR burden was

dispersed across multiple STs rather than confined to a single lineage. Although most

lineages carried few resistance gene, ST 96, ST135 and 185 exhibited higher

resistance burden overall and may be linked to more extensive AMR profiles both in the

urine genomes and our clinical isolates. A notable conservation of AMR profiles was

observed within ST, irrespective of their source of isolation. Genomes in STs 97, 185,

495, and 279 –even though isolated in different studies— had similar resistance profiles

within the same ST (**Fig 1a**). Similarly, ST135 exhibited similar AMR profiles acquiring

all important subclasses highlighted in **Fig 1a**.

In contrast, ST145 (n=34 genomes, range: 1-8, mean: 2.41), ST33 (n=32

genomes, range: 1-9, mean 2.81), ST93 (n=29 genomes, range: 1-8, mean: 3.41) and

ST 202 (n=28 genomes, range: 1-10, mean: 3.00) typically harbored resistance to just

2-3 subclasses with a few outliers (**Fig 5c, 5d**). Again, within STs 92, 93, 133, 145, 196,

and 202 we observe only sporadic accumulation of AMR genes, a pattern that was

consistent across all genomes within these ST's (**Fig 1a**). Collectively, these findings

highlight the varied and ST-specific resistance gene profiles found in *P. mirabilis* urine

isolates. Some STs exhibit consistently high resistance burdens suggesting a shared

capability to acquire or retain the same genes, while others show significant variability,

indicating that a combination of clonal expansion and horizontal gene transfer likely play

an important role in the spread of AMR.

While some resistome subclass profiles correspond to ST, gene composition was

not always determined by ST. Ordination analysis using Jaccard distances show mixing

of the ST (**Fig. S2a**). Some clustering is seen for ST185 (red circle) and ST135 (green

circle). However, some subclasses as well as genes did not co-relate (marked with red

and green arrows for ST185 and ST135) with the genomes captured at the time of

isolation. Highlighting the top 3 most prevalent STs, we observe a 95% covariance

ellipse (**Fig S2b**) without corresponding to specific AMR gene compositions.

Additionally, a comparison of Jaccard distances of the resistomes within ST was similar

to the differences between different STs (**Sup S2c**), probably driven by the skewed

sampling size. However, we see a small number of STs (ST201, ST114, and ST130 in

PCoA and ST201, ST279, and ST93 in NMDS) that showcase compact clusters in the

top 20 genomes (**Fig S2d and S2e**). This may suggest that common resistome

subclasses (ex. Tetracycline, chloramphenicol, aminoglycosides, carbapenemes,

penicillins, cephalosporins, TMP, SMX) are ST-specific, but some less common

antibiotic subclasses (ex. erythromycin, clindamycin) are spread across STs depicting a

dynamic horizontal gene transfer network.

The distribution of resistance gene burden reveals reservoir-specific selection

Comparing the 26 clinical isolates and HI4320 against the 1,001 NCBI genomes

revealed reservoir-specific selection pressures. In total, resistance genes representing

14 different antibiotic subclasses were identified in the clinical isolates compared to 24

from the NCBI genomes. The 26 clinical isolates exhibited substantial variation in

resistance gene burden, ranging from 1 gene (minimum, observed in ST765 and ST591

isolates carrying only a tetracycline resistance determinant) to 15 genes (maximum, in

ST135 isolate H115-00), with an average of 3.8 resistance genes per isolate. This

diversity included resistance to tetracycline, chloramphenicol, streptomycin,

trimethoprim, kanamycin, erythromycin, sulfonamide, β-lactams, and other subclasses

(**Table 3 and Table S8**). Tetracycline resistance gene carriage was nearly ubiquitous in

both datasets, and the majority of genomes from both datasets also harbored genes for

chloramphenicol resistance, demonstrating a strong correlation despite the small size of

the clinical dataset. Tetracycline resistance approached fixation; *tetJ* was present in all

isolates predicted to be tetracycline resistant from both datasets, and co-occurrence of

tetJ with *tetA* (8%, 2/26 clinical) and *tetC* (4%, 1/26 clinical) was also observed.

Similarly, chloramphenicol resistance genes demonstrated strong cross-reservoir

conservation, with *catA* (type-4) detected in 89% (22/26) of clinical isolates compared to

95% of NCBI genomes, and 23% of clinical isolates (5/22) co-harbored *catA1* while 11%

(3/26) harbored no chloramphenicol resistance genes (**Table S8**). Strikingly, one clinical

strain (H115-05) exhibited a rare combination of *floR* + *catB3*; *floR* was seen at a

frequency of 1.2% in the 1,001 genomes while *catB3* was seen at a frequency of 0.34%

(**Fig. 4b**).

Fewer of the clinical isolates had resistance genes for cephalosporins or

quaternary ammonium in comparison to the NCBI dataset (**Table 3**). However, clinical

isolates showed enrichment of resistance genes under UTI treatment pressure including

TMP-SMX (22%, *sul1/sul2*; 37%, *dfr*) and aminoglycosides (streptomycin (37%, most

common: *aadA1*), streptothricin (37%, *sat2*), and kanamycin (26%, *aph(3')-Ia*)) (**Table**

3). In terms of extended β-lactam ecology, clinical isolates predominantly

carried *blaTEM-1* (18%) while NCBI genomes exhibited broader cephalosporin

resistance genes (**Table 3, Table S6**). These patterns demonstrate how reservoir-

specific selection pressures like clinical antibiotic exposure may shape distinct

resistance landscapes from a conserved genetic backbone.

Experimental AMR phenotyping of *P. mirabilis* clinical isolates reveals variable

genotype-phenotype concordance

AMRFinderPlus reports gene presence/absence and point mutation but does not

infer phenotypic resistance due to factors including gene expression and regulation.

Thus, many of the resistance genes detected by established computational databases

may not be relevant for clinical management or antimicrobial surveillance. To determine

concordance between *in-silico* AMR predication and phenotypic resistance in *P.*

mirabilis, we determined the AMR of *P. mirabilis* clinical isolates by standardized

Antimicrobial Susceptibility Tests (AST). We tested growth kinetics of strains with

specific resistance genes at 15-minuite intervals over a 20-hour time course to evaluate

the minimum inhibitory concentrations as well as more nuanced dose-dependent effects

on the growth rates and doubling times of the *P. mirabilis* strains (**Fig 6 and Fig 7**).

Antibiotic concentrations were selected based on the Clinical and Laboratory Standards

Institute (CLSI) guidelines for *Enterobacterales* and were designed to span the

thresholds for sensitive (lowest) to resistant (highest). The genotypic and phenotypic

resistance profiles are summarized in **Table S7 and S9.**

Tetracycline, Kanamycin, Streptomycin, Streptothricin, and Ampicillin

For tetracycline, 100% of clinical isolates (26/26) exhibited resistance with MIC

values >20 µg/mL, which is consistent with the presence of *tetJ*. Growth kinetics

demonstrated uniformly robust growth across all concentrations (**Fig 7a**). Thus, a

perfect genotype-phenotype concordance was observed. Carriage of any additional *tet*

genes (*tetA* or *tetC*) did not alter MIC or growth kinetics within the tested concentration

range.

Kanamycin resistance genes (*aac(6′)-Ib-cr5, aph(3′)-Ia*) were detected in 7/26

clinical isolates and genotype-phenotype concordance was again observed. All 7

isolates encoding the resistance genes had MIC >34 ug/mL, while 102-24, 106-15, H81-

01, H94-04, and HI4320 were predicted to lack all kanamycin resistance determinants

and were susceptible, but some exhibited MICs ranging from 14 – 20.5 ug/ml (**Fig 7b**).

Different combinations of streptomycin resistance genes (*aadA1, aph(6)-Id,*

aph(3")-Ib, aadA5, aadA2) were observed in 10/26 clinical isolates (**Fig 6a, Table S8**),

and all provided resistance to streptomycin when tested in our AST assay (**Fig 7c**).

Unexpectedly, resistance was also observed in 102-24, 106-15, H81-01, H94-00, and

reference strain HI4320 which lacks canonical streptomycin resistance genes,

suggesting the presence of an alternative resistance mechanism. However, these

strains all exhibited a dose-dependent decrease in growth when incubated with

streptomycin. It is also notable that clinical isolate 210-04 (1 predicted gene, *aadA1*)

exhibited a similar dose-dependent decrease in growth even though other strains (206-

05, H97-00, and H97-11) with a singular *aadA1* gene with 100% amino acid sequence

similarity to that of 210-04 did not show dose dependency. It is possible that regulation

of *aadA1* differs in 210-04, leading to a less efficient or inducible mechanism of

resistance compared to the other isolates.

An ampicillin resistance gene (*blaTEM*) was identified in 5/26 clinical isolates and

all exhibited MICs >34 ug/mL. The resistance phenotype was also observed in HI4320,

which lacks any known β-lactamase genes. However, strains that did not encode

BlaTEM, including HI4320, exhibited dose-dependent growth inhibition with increasing

concentrations of ampicillin, despite displaying resistance at CLSI breakpoints (**Fig 7d**).

Streptothricin resistance gene *sat2* was observed in 10/26 clinical isolates (**Fig**

6a, S8), yet the AST assay revealed a mix of intermediate to resistant phenotypes with

MICs ranging from 8 to >22 ug/mL (CLSI breakpoints are ≤8 for susceptible and ≥16 for

resistant). Since streptothricin resistance is conferred by *sat1, sat2*, and *sat4* genes, the

presence of only one gene may confer only partial resistance. The genotypically-

positive isolates 102-00, 206-16 H97-11, H115-00, H115029, H120-00, and H120-04

exhibited dose-dependent decreases in growth with increasing streptothricin

concentrations. However, other genotypically positive strains 206-05 and 210-04 were

sensitive (**Fig 7e**). HI4320 exhibited robust growth regardless of streptothricin

concentration, yet this strain does not encode any known streptothricin resistance

genes (**Fig 7eF**). This discordance suggests that HI4320 may possess a novel, non-

canonical and potentially constitutive mechanism of high-level streptothricin resistance.

Sulfonamide and Trimethoprim

Sulfonamide resistance genes (*sul1* and *sul2)* were both observed in the clinical

isolates, with *sul1* present in 3 isolates (H115-0, H115-05, H120-00), *sul2* in 5 isolates

(102-00, 206-16, H120-04, H115-00, H115-05), and both genes co-occurring in 2

isolates (H115-00, H115-05) (**Table S7**). AST with sulfamethoxazole revealed high level

resistance (MIC >514 ug/ml) in all isolates, irrespective of *sul* gene carriage (single or

double), indicating that the presence of an additional *sul* gene did not further elevate

resistance. Unexpectedly, even strains that did not encode *sul* (102-24, 106-15, 210-04,

H81-01, H94-00, H97-11, and HI4320) had MICs >514 ug/ml, but partial growth

inhibition was clearly present even at the lowest tested concentration (250 ug/ml, **Fig.**

8a). The uniform MIC >514 µg/mL across all strains implies clinical resistance per CLSI

guidelines, as it far exceeds the susceptible breakpoint (typically ≤38 µg/mL). However,

the partial growth inhibition of strains that do not encode *sul* highlights the complexity of

interpreting MICs near the resistance threshold. These data suggest that, while

canonical *sul*-mediated resistance mechanisms were absent in these isolates, non-

genetic or alternative pathways (e.g. efflux pumps, metabolic adaptations) may

contribute to the observed resistance (above threshold) phenotype.

Trimethoprim resistance genes were also detected in 10 clinical isolates, with

MICs of >15 µg/mL (exceeding clinical breakpoints) regardless of which resistance gene

(*dfrA1*, *dfrA17*, or *dfrA32*) was present (**Fig 8b)**. However, one isolate (H97-00)

harboring *dfrA1* exhibited dose-dependent growth inhibition, contrasting with other

strains harboring *dfrA1* with 100% amino acid similarity that showed no dose-

dependency. This implies strain-specific phenotypic variability (e.g., differential gene

expression, compensatory mutations) despite shared genetic determinants. The *dfrA*-

negative reference strain H81-01 also displayed full resistance, while strain HI4320 that

did not carry the *dfrA* gene or gene mutation displayed intermediate resistance and

mirrored the dose-dependent growth inhibition of strain H97-00. These observations

support the existence of non-*dfrA* resistance mechanisms in HI4320 and H81-01, such

as altered folate metabolism or reduced drug uptake. The discordance between genetic

markers and phenotypic outcomes also underscores a key challenge: genetic testing

alone cannot always predict antibiotic response. However, *dfr* negative strains 102-24,

106-15, and H94-00 were sensitive to trimethoprim, in concordance with the genotype.

Since the genotype-negative strains were all phenotypically resistant to

sulfamethoxazole (SMX), albeit with reduced growth rate, we further tested the impact

of combinatory treatment (trimethoprim-sulfamethoxazole, TMP-SMX) on *P. mirabilis*

growth. This is an important parameter since, historically, TMP-SMX was the first-line

empirical treatment for acute uncomplicated UTIs. In the strains exhibiting 100%

genotype-phenotype concordance for both SMX and TMP resistance (i.e. true resistant

genotypes correlated with resistant phenotypes), all isolates (102-00, 206-16, H115-00,

H115-05, H120-00, and H120-04), exhibited complete resistance to the combinatory

drug without any dose-dependent decreases in growth (**Fig 8c**). In contrast, strains

such as 206-05, 210-04, and H97-11 that were genotypically SMX susceptible TMP

resistant but phenotypically resistant to both antibiotics when used alone exhibited a

dose-dependent resistance phenotype when treated with TMP-SMX. Additionally,

strains that were genotypically predicted to be sensitive to both antibiotics but were

phenotypically resistant demonstrated intermediate to resistant phenotypes for

combination treatment: H81-01 displayed an MIC of 8/152 ug/mL (resistant), while 102-

24 displayed an MIC >2/38 ug/mL (intermediate). This is clinically concerning as growth

was observed at breakpoint concentrations. Other discordant strains that had lower

growth yields under single SMX exposure (106-15, H94-00, H97-00, HI4320) were

found to be sensitive to the TMP-SMX combinations (**Fig 8c**).

Chloramphenicol

Chloramphenicol resistance displayed genotype-phenotype concordance, with all

isolates possessing a *catA* gene exhibiting resistance and those lacking *cat* genes

exhibiting susceptibility (**Fig 9a**). MIC values ranged from 10-14 ug/ml in isolates lacking

a *cat* gene but were >26 ug/ml in isolates with *catA* alone and >34 ug/ml in isolates

harboring both *catA* and *catA1* (**Table S9**). Interestingly, the majority of isolates with a

single *catA* gene (16/17) exhibited dose-dependent growth inhibition (**Fig 9**), while

isolates possessing both *catA* and *catA1* demonstrated hyper-resistance (hR) with no

dose dependency (**Fig 9**). The exception was HI4320, which only encodes *catA* but

exhibited robust growth at all but the highest three concentrations of chloramphenicol.

No strains were predicted to have *catA1* alone, so we were not able to determine

whether *catA1* provides a greater level of resistance than *catA,* but our data suggests

that the presence of two *catA* genes enhances functional resistance.

732   We next compared the sequences of the *catA* and *catA1* genes in our *P.*

mirabilis clinical isolates that harbored both genes as *catA1* has not previously been

reported in *P. mirabilis*. The two genes share 76% amino acid sequence identity (with

representative strain 102-00; **Sup Fig 3a**), and *catA* from strain 102-00 showed 100%

identity to the *catA* gene in reference strain *P. mirabilis* HI4320 (**Sup Fig 3b**). The

nucleotide and amino acid sequences of *catA1* from our isolates share 100% amino

acid identity with *catA1* from *E. coli* and ~98% amino acid identity with *catA1*

from *Photobacterium damselae* and *Shigella flexneri* (**Fig S3c)**. These findings indicate

that the coexistence of multiple resistance genes can significantly alter phenotypic

resistance, highlighting the complexity of chloramphenicol resistance in *P.*

mirabilis clinical isolates.

DISCUSSION

CAUTIs caused by *P. mirabilis* remain clinically challenging due to crystalline

biofilm formation, AMR, recurrent infections, and persistence even after antibiotic

treatment and catheter changes [15]. This comprehensive genomic analysis of *P.*

mirabilis from human urinary sources expands our understanding of *P. mirabilis* genetic

diversity, AMR profiles, and MLST prevalence. We studied the population structure and

genomic diversity of a dataset of 1,001 publicly available genomes from human urinary

sources supplemented by phylogenetic analysis of 27 clinical isolates, through which we

determined that urinary *P. mirabilis* isolates possess an open pangenome with more

than 41,121 gene families and a core genome of 1,937 genes for basic cellular

functions, colonization, and survival.

In the context of bacterial genomics, open pangenomes are characteristics of

sympatric species with a high capacity of horizontal gene transfer (HGT), highlighting

genomic plasticity that may allow *P. mirabilis* to rapidly adapt to the urinary tract in the

presence of antibiotic exposure, host immune responses, and competition in the

polymicrobial catheterized environment [56-58]. *P. mirabilis* likely integrates genetic

material from diverse sources, explaining the accessory genes that comprise the vast

majority of the pangenome (37,918 genes, ~90%). This dynamic gain, loss, and

exchange of genetic material emphasizes the importance of flexible genomic elements

in driving diversity in the urinary tract.

The presence of 213 defined STs and approximately 118 previously undefined

genomes and a star-like phylogeny further supports ongoing diversification. Almost half

of the STs in the NCBI and clinical genomes were represented by single isolates (46%),

showcasing heterogeneity. The majority of genomes were distributed across a small

subset of STs (ST96, ST185, ST99, ST135, ST97), resulting in a skewed distribution.

However, the highest AMR burdens were found in rarer STs (e.g., ST431, ST425,

ST279). These strains carried numerous AMR gene subclasses, but we cannot

comment if it is epidemiologically relevant due to low sampling (**Table 3**).

In this study we used MLST to identify lineages as it remains a widely adopted

method of bacterial typing due to its reproducibility and ease of interpretation [49, 59].

However, for *P. mirabilis*, our results highlight limitations in the discriminatory power of

MLST when applied alone for epidemiological and outbreak investigations [59, 60].

While MLST reliably identifies clonal expansions of prevalent lineages like ST96 and

ST185, the large number of undefined STs in our analysis indicates gaps in current

MLST schema and highlights the need for a continuously updated and standardized

typing scheme, which we aim to do in the future.

Overall, the AMR landscape of the 1,001 urine genomes and 27 clinical

isolates was complex and dynamic, with variable genotype-phenotype concordance. An

important observation was the high prevalence of MDR, with 93% of the isolates

resistant to ≥2 antibiotic subclasses and 25% resistant to >6 subclasses. This MDR

burden poses a challenge for effective treatment of *P. mirabilis* urinary tract infections.

The most important finding was the identification of similar antibiotic subclasses in same

MLST even if they originated from different studies/geographical locations. We

observed a lineage-specific AMR burden across STs. However, a widespread

distribution of isolates with very high resistance gene counts (>20) across multiple STs

indicate that extreme resistance gene accumulation is not restricted to ST but is a

function of divergent genetic and environmental context. We can identify if a specific

antibiotic subclass will be effective therapeutically by ST, at least in our top 15 isolates.

However, this approach would not be useful for the >46% (**Fig 2a**) genomes with only 1-

2 genomes per ST. Extending our analysis to more genomes for each rare ST will be

necessary to verify the AMR-ST link of many *P. mirabilis* lineages.

The overwhelming prevalence of tetracycline resistance genes (98.9%),

particularly *tetJ*, aligns with the well-documented intrinsic resistance of *P. mirabilis.* The

dominance of *tetJ* and the frequent carriage of only a single *tet* gene (94.3%) suggest

this determinant is often sufficient for the observed resistance phenotype. However, the

prevalence of chloramphenicol resistance genes (*catA*, 95%) despite limited clinical use

of chloramphenicol is striking. This phenomenon has been observed in other

Enterobacterales and may be linked to co-selection with other antibiotics or acquisition

due to extensive use in agricultural and environmental settings [61, 62]. *P. mirabilis* also

exhibit the co-occurrence of *catA* and *catA1* in 19.2% of *cat*-positive isolates, with

additional *floR* and *catB3* genes in a few isolates. Strains with only a single *cat* gene

exhibited dose-dependent growth inhibition by chloramphenicol while the additional

presence of *catA1* resulted in a hyper-resistant phenotype (no growth inhibition at all at

any tested concentrations), enhancing the resistance phenotype. Detection of *catA1* in

P. mirabilis is a significant finding as this gene has not been well reported and

characterized in *P. mirabilis*. Since the sequences of both genes match those from

closely related bacterial species, this may indicate HGT events via Mobile Genetic

Elements (MGE) that carry *catA* [62] [63, 64].

Additionally, we observed resistance to important classes of antibiotics like

aminoglycosides (71%) with common streptomycin resistance present within lineages

that carried resistance determinants to other antibiotic classes as well, including

carbapenems, cephalosporins, fluoroquinolones and trimethoprim-sulfonamides (**Fig 1**).

The continued presence of resistance determinants for antibiotics such as

aminoglycosides despite reduction in their clinical use may indicate either environmental

influence or co-selection with other antibiotics in *P. mirabilis* urine isolates.

Trimethoprim (TMP, *dfr* genes, 29%) and sulfonamide (SMX, *sul* genes, 28%)

resistance determinants were present at similar frequencies and co-occurred in 22% of

genomes, which is likely due to the common use of TMP-SMX in UTI treatment and

threatens the efficacy of this combination therapy for *P. mirabilis*. While ESBL-

associated genes were less common (18.5% and primarily *blaTEM-1*), their presence is

concerning for broader β-lactam resistance development. Resistance for quinolones

(*qnr aac(6')-Ib-cr, gyrA/parC* mutations) and the chloramphenicol/florfenicol (*cat, floR*)

combination were less common overall in both datasets (~7%) but worth noting because

they act as molecular fingerprints of human driven selection under antimicrobial

pressures due to their prevalence in agriculture [65-68].

Our comprehensive AMR analysis shows a complex landscape of genotype-to-

phenotype relationships, which is characterized by both expected concordance and

clinically critical discordance. The high degree of concordance observed for antibiotics

like tetracycline was expected. Additionally, concordance observed for kanamycin,

where presence of an *aph(3')-la* gene predicted a resistant phenotype, validated the

robustness of the genomic approach for identifying known resistance determinants. It

has been observed that enzymes like aminoglycoside-modifying enzyme like APH(3')-la

confer high-level, constitutive resistance impacting clinical efficacy of these drugs

independent of other mechanisms like efflux pumps [69-71]. Hence, we see a clear

concordance where strains with *aph(3')-la* were resistant while the negative strains had

sensitive MIC. Hence, the presence of AMR determinants are a good indication of

resistance, but the lack of AMR determinants result in less clear resistance predictions.

Our phenotypic analysis also shows that genotyping alone cannot always predict

resistance phenotypes and can indicate multiple mechanisms at play. The observed

discordances provide compelling insights into the evolving and multifaceted nature of

AMR. For example, resistance to streptomycin and streptothricin could be due to novel

resistance mechanisms, such as alternate acetyltransferases with low sequence

homology to known enzymes. Similarly, the ampicillin resistance in strains 102-24, 106-

15, H94-00, H97-11, and HI4320 that lack a detectable β-lactamase gene could be due

to non-enzymatic mechanisms or plasmids. In general, resistance mediated by

permeability or efflux is a hallmark of concentration-dependent antibiotics, where higher

antibiotic concentrations can eventually overcome cellular defenses in concentration-

dependent antibiotics [72, 73]. This is an important consideration in clinical settings, as

treating strains that exhibit dose-dependent resistance with a higher doses or adjuvant

may still be effective, whereas strains with enzymatic resistances lacking dose-

dependent inhibition may require a different approach altogether. Additionally, the

genomic AMR prediction is limited by the existing databases for less well studied

organisms like *P. mirabilis,* lowering the detection of AMR genes and mutations. Hence,

this study highlights the need to characterize AMR mechanisms as well as update

databases, given the clinical importance of this pathogen in UTI.

One of the most critical findings from our genotype-phenotype studies was the

observation that genotypic susceptibility to SMX did not always correspond to

phenotypic susceptibility to TMP-SMX combination treatment (**Fig. 8**). We identified a

spectrum of strains that lacked SMX resistance factors that were either sensitive,

resistant but inhibited in a dose-dependent manner, or completely resistant to

combination treatment, suggesting the presence of unknown SMX resistance

mechanisms that vary in their efficacy. This kind of prediction error can be very

dangerous in a clinical setting since it could lead to treatment failure. Strains like 206-05

and 210-04 (with MIC >16/305 ug/mL) possess a mechanism that is robust enough to

contribute to TMP-SMX resistance, albeit less efficiently than the truly resistant strains.

In contrast, the sensitivity of strains like HI4320, H97-00, and 106-15 to the TMP-SMX

combination treatment implies that the yet unidentified SMX resistance mechanisms can

still be overcome by trimethoprim.

The chloramphenicol resistance profile further illustrates the nuanced relationship

between genotype and phenotype. Isolates with a single *catA* gene exhibited a clear

dose-dependent response, where increasing antibiotic concentrations led to

progressively greater growth inhibition. In contrast, strains possessing both *catA* and

catA1 genes demonstrated a constitutive high-level resistance phenotype with robust

growth maintained even at the highest concentrations tested. This may suggest a gene

dosage effect, where combined expression of two acetyltransferases enhances

resistance. Overall, these data demonstrate how genetic complexity within a strain can

directly dictate the clinical level of resistance.

The genotype-phenotype discordance observed in our small sample study

highlights a significant limitation of current genomic prediction models: their reliance on

curated databases of known resistance genes and mutations. Our study confirms that

WGS-based prediction is highly reliable for well-characterized resistance determinants

in *P. mirabilis* but can fail to predict phenotypes that are likely governed by cryptic

mutations or mechanisms involving efflux and permeability. Not much has been

reported on the mechanism of resistance involving efflux systems and membrane

permeability [10, 65, 74]. Efflux systems such as *AcrAB-TolC* are reported in many *P.*

mirabilis strains can be overwhelmed by high drug concentrations, rather than the

efficient enzymatic hydrolysis provided by a beta-lactamase and is linked to a very high

multidrug-resistance rate of 96 % [75].

Predictions based on genotype are also prone to sequencing and

assembly/annotation errors and ideally need to be reinforced with long-read or

reference-based sequencing to create hybrid assemblies. However, this can become

impractical and costly for diagnostic laboratories. In conclusion, while WGS has the

potential to be used widely in clinical context for AMR predictions, our findings reinforce

that assessing gene presence/absence does not remove the necessity for phenotypic

AST of *P. mirabilis* isolates for accurate clinical decision-making and surveillance.

Further exploration of isolates that were predicted to be sensitive but were either fully

resistant or exhibited dose-dependent growth inhibition may yield discovery of new

resistance mechanisms in this problematic bacterial species.

Acknowledgements

This work was supported by an American Heart Association (AHA) pre-doctoral

fellowship 24PRE1195538 to ND, and by the National Institute of Diabetes and

Digestive and Kidney Diseases under awards R01 DK123158 and R01 DK140371 to

CEA.

References

1. Yang, X., et al., *Disease burden and long-term trends of urinary tract infections: A worldwide report.* Front Public Health, 2022. **10**: p. 888205.
2. Zeng, Z., et al., *Global, regional, and national burden of urinary tract infections from 1990 to 2019: an analysis of the global burden of disease study 2019.* World J Urol, 2022. **40**(3): p. 755-763.
3. Biondo, C., *New Insights into the Pathogenesis and Treatment of Urinary Tract Infections.* Pathogens, 2023. **12**(10).
4. Schaffer, J.N. and M.M. Pearson, *Proteus mirabilis and Urinary Tract Infections.* Microbiol Spectr, 2015. **3**(5).
5. Gaston, J.R., et al., *Polymicrobial interactions in the urinary tract: is the enemy of my enemy my friend?* Infect Immun, 2021.
6. Zilberberg, M.D., et al., *Descriptive epidemiology and outcomes of emergency department visits with complicated urinary tract infections in the United States, 2016-2018.* J Am Coll Emerg Physicians Open, 2022. **3**(2): p. e12694.
7. Ku, J.H., et al., *Multidrug Resistance of Escherichia coli From Outpatient Uncomplicated Urinary Tract Infections in a Large United States Integrated Healthcare Organization.* Open Forum Infect Dis, 2023. **10**(7): p. ofad287.
8. Fromer, D.L., et al., *Risk Factors for Empiric Treatment Failure in US Female Outpatients with Uncomplicated Urinary Tract Infection: an Observational Study.* J Gen Intern Med, 2025. **40**(4): p. 862-870.
9. Wolterink, I., et al., *Nitrofurantoin Failure in Elderly Men: A Retrospective Observational Study.* Antibiotics (Basel), 2020. **9**(5).
10. Armbruster, C.E., H.L.T. Mobley, and M.M. Pearson, *Pathogenesis of Proteus mirabilis Infection.* EcoSal Plus, 2018. **8**(1).
11. Yuan, F., et al., *Pathogenesis of Proteus mirabilis in Catheter-Associated Urinary Tract Infections.* Urol Int, 2021. **105**(5-6): p. 354-361.
12. Brauer, A.L., et al., *Preferential catabolism of L- vs D-serine by Proteus mirabilis contributes to pathogenesis and catheter-associated urinary tract infection.* Molecular microbiology, 2022. **118**(3): p. 125–144.
13. Armbruster, C.E., et al., *Genome-wide transposon mutagenesis of Proteus mirabilis: Essential genes, fitness factors for catheter-associated urinary tract infection, and the impact of polymicrobial infection on fitness requirements.* PLoS Pathog, 2017. **13**(6): p. e1006434.
14. White, A.N., et al., *Catalase Activity is Critical for Proteus mirabilis Biofilm Development, Extracellular Polymeric Substance Composition, and Dissemination during Catheter-Associated Urinary Tract Infection.* Infect Immun, 2021. **89**(10): p. e0017721.
15. Armbruster, C.E., et al., *Prospective assessment of catheter-associated bacteriuria clinical presentation, epidemiology, and colonization dynamics in nursing home residents.* JCI Insight, 2021. **6**(19).
16. Sabbuba, N.A., et al., *Genotyping demonstrates that the strains of Proteus mirabilis from bladder stones and catheter encrustations of patients undergoing long-term bladder catheterization are identical.* J Urol, 2004. **171**(5): p. 1925-8.

17. Nye, T.M., et al., *Microbial co-occurrences on catheters from long-term*
catheterized patients. Nat Commun, 2024. **15**(1): p. 61.
18. Armbruster, C.E., et al., *The Pathogenic Potential of Proteus mirabilis Is*
Enhanced by Other Uropathogens during Polymicrobial Urinary Tract Infection.
Infect Immun, 2017. **85**(2).
19. Ramatla, T., et al., *Antibiotic resistance and virulence profiles of Proteus mirabilis*
isolated from broiler chickens at abattoir in South Africa. Vet Med Sci, 2024.
10(2): p. e1371.
20. Bharat, A., et al., *Correlation between Phenotypic and In Silico Detection of*
Antimicrobial Resistance in Salmonella enterica in Canada Using Staramr.
Microorganisms, 2022. **10**(2).
21. Gaston, J.R., et al., *Enterococcus faecalis Polymicrobial Interactions Facilitate*
Biofilm Formation, Antibiotic Recalcitrance, and Persistent Colonization of the
Catheterized Urinary Tract. Pathogens, 2020. **9**(10).
22. Luzzaro, F., et al., *Prevalence and epidemiology of microbial pathogens causing*
bloodstream infections: results of the OASIS multicenter study. Diagn Microbiol
Infect Dis, 2011. **69**(4): p. 363-9.
23. Kanayama, A., I. Kobayashi, and K. Shibuya, *Distribution and antimicrobial*
susceptibility profile of extended-spectrum beta-lactamase-producing Proteus
mirabilis strains recently isolated in Japan. Int J Antimicrob Agents, 2015. **45**(2):
p. 113-8.
24. Filipiak, A., et al., *Pathogenic Factors Correlate With Antimicrobial Resistance*
Among Clinical Proteus mirabilis Strains. Front Microbiol, 2020. **11**: p. 579389.
25. Li, Y., et al., *Genetic analysis of resistance and virulence characteristics of*
clinical multidrug-resistant Proteus mirabilis isolates. Front Cell Infect Microbiol,
2023. **13**: p. 1229194.
26. Nielubowicz, G.R. and H.L. Mobley, *Host-pathogen interactions in urinary tract*
infection. Nat Rev Urol, 2010. **7**(8): p. 430-41.
27. Tahmasebi H, et al., *Relationship between biofilm gene expression with*
antimicrobial resistance pattern and clinical specimen type based on sequence
types (STs) of methicillin-resistant S. aureus. . Molecular Biology Reports 2020.
47(2): p. 1309-1320.
28. Chen, S.L., et al., *A Core Genome Multilocus Sequence Typing Scheme for*
Proteus mirabilis. Biomed Environ Sci, 2023. **36**(4): p. 343-352.
29. Hickman FW and F.J. 3rd, *Differentiation of Proteus mirabilis by bacteriophage*
typing and the Dienes reaction. Journal of Clinical Microbiology, 1976. **3**(3): p.
350-8.
30. William C. Schmidt and C.D. Jeffries, *Bacteriophage Typing of Proteus mirabilis,*
Proteus vulgaris, and Proteus morganii. Applied and Environmental Microbiology,
1974.
31. Pfaller MA, et al., *Evaluation of the Discriminatory Powers of the Dienes Test and*
Ribotyping as Typing Methods for Proteus mirabilis. Journal of Clinical
Microbiology, 2000. **38**(3): p. 1077-80.
32. Lian S, et al., *Genomic insights on cgMLST markers, drug resistance, and*
urease cluster of Proteus mirabilis strains. 2024. **13**(1).

33.   Potter RF, et al., *Uncharacterized and lineage-specific accessory genes within the Proteus mirabilis pan-genome landscape.* mSystems, 2023. **84**(4): p. 00159-23.

34.   Feldgarden, M., et al., *AMRFinderPlus and the Reference Gene Catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence.* Sci Rep, 2021. **11**(1): p. 12728.

35.   Sung, K., et al., *Comprehensive Genomic Analysis of Uropathogenic E. coli: Virulence Factors, Antimicrobial Resistance, and Mobile Genetic Elements.* Pathogens, 2024. **13**(9).

36.   Chin, J.J., et al., *High Carriage of tetA, sul1, sul2 and bla (TEM) Resistance Genes among the Multidrug-resistant Uropathogenic Escherichia coli (UPEC) Strains from Malaysian Patients.* Trop Life Sci Res, 2024. **35**(2): p. 211-225.

37.   Kandi, V., et al., *Molecular Characterization of Escherichia coli Causing Urinary Tract Infections Through Next-Generation Sequencing: A Comprehensive Analysis of Serotypes, Sequence Types, and Antimicrobial and Virulence Genes.* Cureus, 2024. **16**(3): p. e55556.

38.   He, X.T., et al., *The risk factors, antimicrobial resistance patterns, and outcomes associated with extended-spectrum beta-lactamases-Producing pathogens in pediatric urinary tract infection.* Pediatr Neonatol, 2024. **65**(3): p. 242-248.

39.   Hassuna, N.A., et al., *Characterization of antimicrobial resistance among Proteus mirabilis isolates from catheter-associated urinary tract infections and non-catheter-associated urinary tract infections in Egypt.* BMC Infect Dis, 2025. **25**(1): p. 767.

40.   Mirzaei, A., et al., *From the Urinary Catheter to the Prevalence of Three Classes of Integrons, beta-Lactamase Genes, and Differences in Antimicrobial Susceptibility of Proteus mirabilis and Clonal Relatedness with Rep-PCR.* Biomed Res Int, 2021. **2021**: p. 9952769.

41.   Feldgarden, M., et al., *Validating the AMRFinder Tool and Resistance Gene Database by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection of Isolates.* Antimicrob Agents Chemother, 2019. **63**(11).

42.   Pataki, B.A., et al., *Understanding and predicting ciprofloxacin minimum inhibitory concentration in Escherichia coli with machine learning.* Sci Rep, 2020. **10**(1): p. 15026.

43.   Simon, A. *FastQC: A quality control tool for high throughput sequence data*. 2010 [cited 2025; Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

44.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina sequence data.* Bioinformatics, 2014. **30**(15): p. 2114-20.

45.   Bankevich, A., et al., *SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing.* J Comput Biol, 2012. **19**(5): p. 455-77.

46.   Gurevich, A., et al., *QUAST: quality assessment tool for genome assemblies.* Bioinformatics, 2013. **29**(8): p. 1072-5.

47.   Seemann, T., *Prokka: rapid prokaryotic genome annotation.* Bioinformatics, 2014. **30**(14): p. 2068-9.

48.   Hunt. BC, Brix. V, and G.L. Vath. J, Taddei. SM, Deka. N, Learman. BS, Brauer. AL, Shen. S, Qu. J, Armbruster. CE. , *Metabolic interplay between Proteus*

mirabilis and Enterococcus faecalis facilitates polymicrobial biofilm formation and
invasive disease. mBio 15:e02164-24.
. mBio, 2024. **15**(12).
49.   Jolley, K.A., J.E. Bray, and M.C.J. Maiden, *Open-access bacterial population*
genomics: BIGSdb software, the PubMLST.org website and their applications.
Wellcome Open Res, 2018. **3**: p. 124.
50.   Torsten, S. *mlst: Scan contig files against PubMLST typing scheme.* [cited 2025;
Available from: https://github.com/tseemann/mlst.
51.   Page, A.J., et al., *Roary: rapid large-scale prokaryote pan genome analysis.*
Bioinformatics, 2015. **31**(22): p. 3691-3.
52.   Letunic, I. and P. Bork, *Interactive Tree Of Life (iTOL) v5: an online tool for*
phylogenetic tree display and annotation. Nucleic Acids Res, 2021. **49**(W1): p.
W293-W296.
53.   Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree: computing large minimum*
evolution trees with profiles instead of a distance matrix. Mol Biol Evol, 2009.
26(7): p. 1641-50.
54.   *Performance Standard for Antimicrobial Susceptibility Testing*, CLSI, Editor.
2023.
55.   Zhang, S., et al., *Genomic analysis of Proteus mirabilis: Unraveling global*
epidemiology and antimicrobial resistance dissemination - emerging challenges
for public health and biosecurity. Environ Int, 2025. **199**: p. 109487.
56.   Friedman, J., E.J. Alm, and B.J. Shapiro, *Sympatric speciation: when is it*
possible in bacteria? PLoS One, 2013. **8**(1): p. e53539.
57.   Zhao, L., et al., *Evolution alters ecological mechanisms of coexistence in*
experimental microcosms. Functional Ecology, 2015. **30**(8): p. 1440-1446.
58.   Mukherjee, S., et al., *Incipient Sympatric Speciation and Evolution of Soil*
Bacteria Revealed by Metagenomic and Structured Non-Coding RNAs Analysis.
Biology (Basel), 2022. **11**(8).
59.   Maiden, M.C., *Multilocus sequence typing of bacteria.* Annu Rev Microbiol, 2006.
60: p. 561-88.
60.   Perez-Losada, M., et al., *Population genetics of microbial pathogens estimated*
from multilocus sequence typing (MLST) data. Infect Genet Evol, 2006. **6**(2): p.
97-112.
61.   Graf, F.E., et al., *Molecular mechanisms of re-emerging chloramphenicol*
susceptibility in extended-spectrum beta-lactamase-producing Enterobacterales.
Nat Commun, 2024. **15**(1): p. 9019.
62.   Puangseree, J., et al., *Molecular basis of the persistence of chloramphenicol*
resistance among Escherichia coli and Salmonella spp. from pigs, pork and
humans in Thailand. PLoS One, 2024. **19**(5): p. e0304250.
63.   Pitout, J.D.D. and L. Chen, *The Significance of Epidemic Plasmids in the*
Success of Multidrug-Resistant Drug Pandemic Extraintestinal Pathogenic
Escherichia coli. Infect Dis Ther, 2023. **12**(4): p. 1029-1041.
64.   Marcoux, P.E., et al., *Host Dependent-Transposon for a Plasmid Found in*
Aeromonas salmonicida subsp. salmonicida That Bears a catB3 Gene for
Chloramphenicol Resistance. Antibiotics (Basel), 2023. **12**(2).

65. ElTaweel, M., H.S. Said, and R. Barwa, *Emergence of extensive drug resistance*
and high prevalence of multidrug resistance among clinical Proteus mirabilis
isolates in Egypt. Ann Clin Microbiol Antimicrob, 2024. **23**(1): p. 46.
66. Piekarska, K., et al., *The molecular mechanisms of fluoroquinolone resistance*
found in rectal swab isolates of Enterobacterales from men undergoing a
transrectal prostate biopsy: the rationale for targeted prophylaxis. Ann Clin
Microbiol Antimicrob, 2021. **20**(1): p. 81.
67. Qu, X., et al., *Genomic Investigation of Proteus mirabilis Isolates Recovered*
From Pig Farms in Zhejiang Province, China. Front Microbiol, 2022. **13**: p.
952982.
68. Thaotumpitak, V., et al., *Emergence of colistin resistance and characterization of*
antimicrobial resistance and virulence factors of Aeromonas hydrophila,
Salmonella spp., and Vibrio cholerae isolated from hybrid red tilapia cage culture.
PeerJ, 2023. **11**: p. e14896.
69. El-Baz, A.M., et al., *The Link between Occurrence of Class I Integron and*
Acquired Aminoglycoside Resistance in Clinical MRSA Isolates. Antibiotics
(Basel), 2021. **10**(5).
70. Quirke, J.C.K., et al., *Structure-Activity Relationships for 5" Modifications of 4,5-*
Aminoglycoside Antibiotics. ChemMedChem, 2022. **17**(13): p. e202200120.
71. Thacharodi, A. and I.L. Lamont, *Aminoglycoside-Modifying Enzymes Are*
Sufficient to Make Pseudomonas aeruginosa Clinically Resistant to Key
Antibiotics. Antibiotics (Basel), 2022. **11**(7).
72. Geng, B., et al., *Identification and Characterization of Genes Related to*
Ampicillin Antibiotic Resistance in Zymomonas mobilis. Antibiotics, 2022. **11**(11).
73. Darby, E.M., et al., *Molecular mechanisms of antibiotic resistance revisited.* Nat
Rev Microbiol, 2023. **21**(5): p. 280-295.
74. Dulanto Chiang, A. and J.P. Dekker, *Efflux pump-mediated resistance to new*
beta lactam antibiotics in multidrug-resistant gram-negative bacteria. Commun
Med (Lond), 2024. **4**(1): p. 170.
75. Al-Kaim, H.W. and H.O.M. Al-Dahmoshi, *Prevalence of AcrAB-TolC efflux pump*
among Proteus mirabilis isolated from cystitis patients in Hilla City, Iraq. Medical
Journal of Babylon, 2023. **20**(3): p. 503-510.
1118
1119

1120 **Table 1: Summary statistics of *P. mirabilis* genome sequences**

Publicly available *P. mirabilis* genomes isolated from urine				
Metric	**Average**	**Max**	**Min**	**Median**
# Contigs	258.43	973	60	180
Total Length	4020761.51	8382181	3667799	3992140
N50	174294.28	1994055	4981	174571
GC Content	38.76	43.24	37.57	38.69
Clinical *P. mirabilis* genomes isolated from catheterized individuals				
Metric	**Average**	**Max**	**Min**	**Median**
# Contigs	66.52	98	2	69
Total Length	4015048.48	4165371	3795264	4015495
N50	317358.93	4063606	119871	182416
GC Content	38.74	38.98	38.38	38.78

1121

1122 **Table 2: *P. mirabilis* clinical isolates used in this study**

Isolate ID	Study	Species	MLST	Patient designation in cited publication	Isolation week
102-00	CEA	*P. mirabilis*	185	Participant A [15]	Visit 0
102-24	CEA	*P. mirabilis*	145	Participant A [15]	Visit 24
102-25	CEA	*P. mirabilis*	UD	Participant A [15]	Visit 25
104-00	CEA	*P. mirabilis*	UD	Participant C [15]	Visit 0
106-15	CEA	*P. mirabilis*	552	Participant H [15]	Visit 15
201-07	CEA	*P. mirabilis*	UD	Participant G [15]	Visit 7
203-01	CEA	*P. mirabilis*	615	Participant F [15]	Visit 1
206-05	CEA	*P. mirabilis*	185	Participant D [15]	Visit 5
206-16	CEA	*P. mirabilis*	185	Participant D [15]	Visit 16
210-04	CEA	*P. mirabilis*	145	Participant J [15]	Visit 4
H81-01	HUC	*P. mirabilis*	765	Patient 81-urine [17]	Visit 1
H85-10	HUC	*P. mirabilis*	330	Patient 85-urine [17]	Visit 10
H94-00	HUC	*P. mirabilis*	696	Patient 94-urine [17]	Visit 0
H97-00	HUC	*P. mirabilis*	157	Patient 97-urine [17]	Visit 0
H97-11	HUC	*P. mirabilis*	157	Patient 97-urine [17]	Visit 11
H105-00	HUC	*P. mirabilis*	591	Patient 105-urine [17]	Visit 0
H105-06	HUC	*P. mirabilis*	765	Patient 105-urine [17]	Visit 6
H108-00	HUC	*P. mirabilis*	96	Patient 108-urine [17]	Visit 0
H115-00	HUC	*P. mirabilis*	135	Patient 115-urine [17]	Visit 0
H115-05	HUC	*P. mirabilis*	135	Patient 115-urine [17]	Visit 5
H119-03	HUC	*P. mirabilis*	146	Patient 119-urine [17]	Visit 3
H120-00	HUC	*P. mirabilis*	97	Patient 120-urine [17]	Visit 0
H120-04	HUC	*P. mirabilis*	185	Patient 120-urine [17]	Visit 4
H125-06	HUC	*P. mirabilis*	565	Patient 125-urine [17]	Visit 6
H127-00	HUC	*P. mirabilis*	96	Patient 127-urine [17]	Visit 0
H132-05	HUC	*P. mirabilis*	96	Patient 132-urine [17]	Visit 5
HI4320	Reference	*P. mirabilis*	90	Refseq: GCA_000069965.1_ASM6996v1_genomic	-

1123

1124

Table 3: AMR subclass count and percentage in *P. mirabilis* genomes

Resistance subclass	Count (NCBI)	Genes detected (NCBI)	Percentage (1001 NCBI)	Count (clinical)	Genes detected (clinical)	Percentage (26 clinical)	NCBI/Clinical Comparison
AMIKACIN/KANAMYCIN / QUINOLONE / TOBRAMYCIN	11	*aac(6')-Ib, aph(3')-IIIa, aph(3')-VI, aph(3')-VIb, aph(3')-XV*	7.19%	1	*aac(6')-Ib-cr5*	3.70%	Lower in clinical
BETA-LACTAM	185	*blaCARB-2, blaOXA, blaOXA-1033, blaOXA-2, blaOXA-9, blaTEM, blaTEM-1, blaTEM-2, blaTEM-30, hugA*	18.46%	5	*blaTEM-1*	18.52%	Similar in prevalence
CEPHALOSPORIN	90	*ampC, blaCMY-2, blaCMY-6, blaCTX-M-14, blaCTX-M-15, blaCTX-M-2, blaCTX-M-3, blaCTX-M-55, blaCTX-M-65, blaCTX-M-91, blaDHA, blaDHA-1, blaFOX-5, blaOXA-1, blaOXA-10, blaPDC-3, blaPER, blaPER-1, blaSHV-12, blaSHV-31, blaTEM-10, blaTEM-155, blaVEB-6*	8.98%	1	*blaOXA-1*	3.70%	Lower in clinical
CHLORAMPHENICOL	891	*catA, catA1, catA2, catA3, catB11, catB2, catB3, catB7, catB8, cmlA, cmlA5, cmlA6*	88.92%	23	*catA, catA1, catB3, floR*	85.19%	Similar in prevalence
ERYTHROMYCIN	36	*ere(A), ere(B) mph(E)*	3.50%	1	*ere(A)*	3.70%	Similar in prevalence
KANAMYCIN	137	*aph(3')-IIb, aph(3')-Ia*	13.67%	7	*aph(3')-Ia*	25.93%	Higher in clinical
QUATERNARY AMMONIUM	192	*qacE, qacEdelta1*	19.16%	3		11.11%	Lower in clinical
QUINOLONE	68	*qnrA, qnrA1, qnrB19, qnrB2, qnrB4, qnrD, qnrD1, qnrD2, qnrS1, qnrVC1*	6.79%	2	*qnrD*	7.41%	Similar in prevalence
RIFAMYCIN	28	*arr-2, arr-3*	2.79%	1	*arr-3*	3.70%	Similar low prevalence
STREPTOMYCIN	352	*aadA1, aadA10, aadA16, aadA2, aadA3, aadA31, aadA5, ant(6)-Ia, aph(3'')-Ib, aph(6)-Id*	35.13%	10	*aadA1, aph(6)-Id, aph(3'')-Ib, aadA5, aadA2*	37.04%	Similar in prevalence
STREPTOTHRICIN	282	*sat2*	28.14%	10	*sat2*	37.04%	Higher in clinical
SULFONAMIDE	286	*sul1, sul2*	28.54%	6	*sul1, sul2*	22.22%	Slightly lower in clinical

TETRACYCLINE	991	*tet(59), tet(A), tet(B), tet(C), tet(D), tet(G), tet(H), tet(J), tet(L), tet(M), tet(Y)*	98.90%	26	*tet(J), tet(A), tet(C)*	100%	Nearly universal presence in both
TRIMETHOPRIM	290	*dfrA1, dfrA10, dfrA12, dfrA14, dfrA16, dfrA17, dfrA19, dfrA27, dfrA32, dfrA42, dfrA46, dfrA5, dfrA7*	28.94%	10	*dfrA1, dfrA17, dfrA32*	37.04%	Higher in clinical
AZITHROMYCIN / CLARITHROMYCIN / CLINDAMYCIN / ERYTHROMYCIN / TELITHROMYCIN	1	*erm(A)*	0.10%	NA	NA	NA	Only in NCBI
AZITHROMYCIN / ERYTHROMYCIN / SPIRAMYCIN / TELITHROMYCIN	15	*mph(A)*	1.5	NA	NA	NA	Only in NCBI
AZITHROMYCIN / ERYTHROMYCIN / STREPTOGRAMIN	26	*msr(E)*	2.59%	NA	NA	NA	Only in NCBI
BLEOMYCIN	52	*ble, bleO*	5.19%	NA	NA	NA	Only in NCBI
CARBAPENEM	171	*blaIMP, blaIMP-27, blaKPC-2, blaKPC-3, blaNDM-1, blaNDM-4, blaNDM-5, blaNDM-50, blaNDM-6, blaNDM-7, blaOXA, blaOXA-23, blaOXA-48, blaVIM-1, blaVIM-78*	17.07%	NA	NA	NA	Only in NCBI
FLORFENICOL	61	*floR, floR2*	6.09%	NA	NA	NA	Only in NCBI
FOSFOMYCIN	14	*fosA, fosA3*	1.40%	NA	NA	NA	Only in NCBI
GENTAMICIN	54	*aac(3)-IId, aac(3)-IIe, aac(3)-VIa, aac(6')-Ib, aac(6')-Ib4, armA*	5.39%	NA	NA	NA	Only in NCBI
GENTAMICIN / KANAMYCIN / TOBRAMYCIN	72	*ant(2'')-Ia*	7.19%	NA	NA	NA	Only in NCBI
HYGROMYCIN	13	*aph(4)-Ia*	1.30%	NA	NA	NA	Only in NCBI
KASUGAMYCIN	1	*aac(2')-IIa*	0.10%	NA	NA	NA	Only in NCBI
LINCOSAMIDE	42	*lnu(F), lnu(G)*	4.19%	NA	NA	NA	Only in NCBI
SPECTINOMYCIN / STREPTOMYCIN	1	*aadA27*	0.10%	NA	NA	NA	Only in NCBI

Figure Legends

Fig 1. Phylogenetic tree and pangenome analysis of publicly-available *Proteus mirabilis* genomes from human urine samples. (**1a**) maximum-likelihood phylogenetic tree of 1,001 *P. mirabilis* human urine genomes with metadata arranged as concentric bands. The first two bands display multi-locus sequence type (ST), and the third band indicates the study source in the NCBI database associated with the Bioproject. Bands 4-8 indicate presence (colored bars) or absence (grey) of resistance genes across different antibiotic classes. Distinct clusters were observed that either had resistance genes for nearly all classes of antibiotic or lacked them entirely, which was dependent on ST and independent of geographical source. (**1b**) Roary collector curve showing conserved genes (~1,700 core genome) present in all of the genomes analyzed as well as the number of total genes, which increases as new genomes are added, indicating an open pangenome. (**1c**) Roary collector curve showing unique vs new gene accumulation across all *P. mirabilis* genomes. Here, the total unique genes is >15,000 indicating massive accessory gene diversity and an open pangenome. Early genomes introduce a lot of new genes before dropping out. However, after addition of >250 genomes, new genes (vertical lines) are still accumulated, indicating horizontal gene transfer (HGT), mobile genetic elements, or prophage insertions.

Fig 2. Sequence type distribution of publicly-available *Proteus mirabilis* genomes from human urine samples. (**2a**) Pie-chart illustrating the distribution of MLSTs

1148   grouped by their sample count frequency. (**2b**) Bar plot showing the top 20 MLSTs

1149   based on sample count.

1150

Fig 3: Phylogenetic tree and pangenome analysis of *Proteus mirabilis* strains

isolated from individuals with indwelling urinary catheters. (**3a**) A mid-rooted

maximum likelihood phylogenetic tree of clinical *P. mirabilis* isolates collected from

patients with long-term urinary catheters. The phylogenetic tree displays the participant

ID, the cohort ID (strip1), and multi-locus sequence type (ST, strip 2). Many of the

isolates cluster by ST, but not by geographical location. (**3b**). Mosaic pan genome with

presence/absence matrix of core-genome phylogenetic tree for 26 *P. mirabilis* clinical

isolates collected from patients with long-term urinary catheters.

1159

Fig 4: AMR gene carriage in publicly-available *P. mirabilis* genomes. (**4a**) Number

of *P. mirabilis* strains in the NCBI dataset carrying 1, 2, and 3 tetracycline resistance

(*tet)* genes. (**4b**) Three-way Venn diagram exhibiting the shared chloramphenicol

resistance genes (*catA, catA1,* and *floR*) by all the *P. mirabilis* isolates in the NCBI

dataset. (**4c**) Number of *P. mirabilis* strains in the NCBI dataset carrying 1, 2, and 3

trimethoprim resistance (*dfrA)* genes. (**4d**) Venn diagram of shared sulfonamide

resistance genes (*sul1* and *sul2*). (**4e**) Venn diagram of shared trimethoprim (TMP) and

sulfonamide (SMX) resistance genes.

1168

Fig 5: Distribution of AMR subclass carriage in isolates from the most common

STs. (**5a**) AMR subclass count for the top MLSTs that each had >3 representative

genomes. Each black point represents the AMR count for one isolate within the

indicated ST. The overall dataset-wide subclass mean is indicated by the red line. (**5b**)

Empirical cumulative distribution function (ECDF) showing the cumulative proportion of

P. mirabilis genomes as a function of the number of AMR subclasses detected per

genome, with 80^{th} percentile noted at 6 subclasses. (**5c**) Mean AMR subclass count for

the 10 most prevalent multi-locus sequence types (ST), ordered by prevalence. Bars

summarize the average burden of AMR subclasses per isolate in each ST. (**5d**)

Distribution of AMR subclass count per isolate within each ST among the 10 most

prevalent ST, with each black point representing an isolate. (**5e**) Empirical cumulative

distribution function (ECDF) showing the cumulative proportion of isolates as a function

of the number of AMR genes detected per isolate, with 80^{th} percentile noted at 18

genes.

1183

Fig 6: Overview of antibiotic resistance and MIC. (**6a**) Overview of antibiotic

resistance phenotype found in the clinical CAUTI isolates. Color coding denotes

resistance status: red = resistant, yellow = intermediate resistance, and green =

susceptible strains. The presence or absence of resistance-associated genes is

indicated by filled and empty circles, respectively. (**6b**) Heatmap representing the MIC

for each antibiotic tested across clinical *P. mirabilis* CAUTI isolates. Color intensity

corresponds to MIC values, indicating variation in resistance levels.

1191

Fig 7: Dose response curves and MIC: Mean normalized OD600 values at

experimental endpoint (20 hours) are plotted against antibiotic concentrations in ug/ml.

OD is normalized to a no drug control set to 1.0 (maximum growth). The red vertical line

indicates the resistance breakpoint (CLSI) for *Enterobacterales.* The MIC was

determined as the lowest antibiotic concentration for which growth was restricted below

a threshold of a normalized OD of 0.15 (grey horizontal dashed line). (**7a**) Tetracycline

(MIC range tested: 2-20 ug/ml), (**7b**) Kanamycin (MIC range tested: 6-34 ug/ml), (**7c**)

Streptomycin (MIC range tested: 4-22 ug/ml), (**7d**) Ampicillin (MIC range tested: 6-34

ug/ml), (**7e**) Streptothricin (MIC range tested: 4-22 ug/ml).

1201

Fig 8: Dose response curves and MIC: Mean normalized OD600 values at

experimental endpoint (20 hours) are plotted against antibiotic concentrations in ug/ml.

OD is normalized to a no drug control set to 1.0 (maximum growth). The red vertical line

indicates the resistance breakpoint (CLSI) for *Enterobacterales.* The MIC was

determined as the lowest antibiotic concentration for which growth was restricted below

a threshold of a normalized OD of 0.15 (grey horizontal dashed line). (**8a**)

Sulfamethoxazole (MIC range tested: 250-512 ug/ml), (**8b**) Trimethoprim (MIC range

tested: 4-22 ug/ml), (**8c**) TMP-SMX (MIC range tested: 0.5/9.5-16/305 ug/ml).

1210

Fig 9: Dose response curves and MIC of Chloramphenicol. Mean normalized

OD600 values at experimental endpoint (20 hours) are plotted against antibiotic

concentrations in ug/ml. OD is normalized to a no drug control set to 1.0 (maximum

growth). The red vertical line indicates the resistance breakpoint (CLSI) for

Enterobacterales. The MIC was determined as the lowest antibiotic concentration for

which growth was restricted below a threshold of a normalized OD of 0.15 (grey

horizontal dashed line). Chloramphenicol concentrations tested ranged from 6-34 ug/ml.

a

Band 1: MLST Classification
- ■ Known MLST
- ■ Novel/Untyped

Band 2: Top 15 MLST
- ■ ST96
- ■ ST185
- ■ ST99
- ■ ST135
- ■ ST97
- ■ ST145
- ■ ST133
- ■ ST93
- ■ ST202
- ■ ST146
- ■ ST157
- ■ ST92
- ■ ST196
- ■ ST495
- ■ ST279

Band 3: Study/Isolation Region
- ■ University of Pittsburgh (EDS-HAT)
- ■ CDC (Routine reports)
- ■ Female Urinary Microbiota study
- ■ Rush University (Gut UPEC)
- ■ WUSM (multiple regions)
- ■ Human Bladder Bacteria study
- ■ Antimicrobial Surveillance, Brigham & Women
- ■ Nosocomial infections study
- ■ Bacterial Surveillance for LDH study
- ■ New York City Hospital
- ■ Urine: UTI with AMR study

Bands 4-8: Antibiotic Subclasses
- ■ Aminoglycosides
- ■ Carbapanemes
- ■ Cephalosporins/Penicilins
- ■ Fluoroquinolones
- ■ TMP-SMX

b

c

Total MLST = 214

a

Number of representative genomes

■ 1 □ 2 □ 3-5 ■ 6-10 □ 11-50

b

a

Tree scale: 0.01 ⊢——⊣

Band 1: MLST

Band 2: Clinical cohorts
- CEA cohort
- HUC cohort
- Reference HI4320

Bands 3-12: Antibiotic Class
- Tetracycline
- Chloramphenicol
- Aminoglycosides
- Carbapenems
- Cephalosporins-Penicillins
- TMP-SMX
- Streptothricin
- Erythromycin + Rifamycin
- Quinolone
- Biocide

b

Clade 1

Clade 2

Chloramphenicol dose–response

Normalized OD (endpoint mean)

Concentration (μg/mL)

catA+ catA1+

catA+ catA1-

catA+ catA1-

catA- catA1-

STRAIN ID

102-00	H97-11
102-24	H105-00
102-25	H105-06
104-00	H108-00
106-15	H115-00
201-07	H115-29
203-01	H119-03
206-05	H120-00
206-16	H120-04
210-04	H125-06
H81-01	H127-00
H85-10	H132-05
H94-00	HI4320
H97-00	

Fig S1. Mosaic pan genome with presence/absence matrix of core-genome phylogenetic tree for 1,001 *P. mirabilis* genomes. Three clades with different branch lengths (*sb1, sb2,* and *sb3*) are represented in the phylogenetic tree, where *sb2* has a clear difference in core genome.

Fig. S2. a. Classical PCoA (principal component analysis/metric multidimensional scaling) of Jaccard distances computed from AMR gene presence–absence in isolates of the 20 most prevalent MLSTs. Each point is an isolate, colored by MLST. Axes are labeled with the percentage of distance variation (PCo1 ≈ 42.2%, PCo2 ≈ 10.6%). **b.** PCoA with 95% covariance ellipse highlighting the top 3 MLST by count, where similar distributions and overlapping ellipses indicate limited separation by ST. **c.** Within vs between MLST Jaccard distances of the 20 most prevalent 20 MLST, depicting similar distances **d.** PCoA highlighting the top 3 MLST by tightness. **e.** Non-metric multidimensional scaling (NMDS) highlighting the top 3 MLST by tightness. In **d and e,** only the three most compact MLSTs by 95% ellipse area are highlighted. These groups are cohesive but still overlap with the main cloud (grey). NMDS stress is ~0.19, indicating weak clustering by AMR.

a

b

c

Fig S3. Comparative sequence analysis of *catA* and *catA1* genes in *P. mirabilis* clinical isolates. a. Amino acid equence comparison between *catA* and *catA1* demonstrates 78% amino acid sequence identity. **b.** Sequence comparison between the *catA* gene from strain 102-00 to the *catA* gene of reference strain *P. mirabilis* HI4320 shows 100% amino acid sequence identity. **c**. Sequence comparison between the *catA1* gene in different species of *Enterobacterales*.