Trans-acting mutations reveal non-nuclear modulators of both intrinsic and extrinsic gene expression noise in a eukaryote.

Nena Martin[1], Julie Kleine-Schultjann[1], Quentin Boussau[1], Agnès Dumont[1], Hélène Duplus-Bottin[1], Laurent Modolo[1], Gaël Yvert[1], Patricia J. Wittkopp[2,3] and Fabien Duveau[1,3]

Affiliations:

[1]Laboratory of Biology and Modeling of the Cell, Ecole Normale Supérieure de Lyon, CNRS, Université Claude Bernard Lyon, Université de Lyon, France

[2]Department of Molecular, Cellular, and Developmental Biology, University of Michigan, USA

[3]Department of Ecology and Evolutionary Biology, University of Michigan, USA

Corresponding author:

Fabien Duveau, fabien.duveau@ens-lyon.fr

LBMC, ENS de Lyon, CNRS, UMR5239

46 allée d'Italie

69007 Lyon

France

ABSTRACT

Gene expression regulation is a stochastic process that can be modified by mutations not only in a

deterministic manner but also in a probabilistic way, for example by changing the extent of cell-to-cell

variability in gene expression (also called "gene expression noise") without necessarily changing the

expected (mean) expression level. Such mutations can act either in *cis* (perturbing expression noise at

their own locus) or in *trans* (perturbing expression noise of a gene located elsewhere in the genome).

Although systematic studies have successfully analyzed the properties of *cis*-acting mutations

modulating gene expression noise, less is known about the type of genetic changes that may alter

expression noise in *trans*. Here, we applied genetic mapping on yeast strains (*Saccharomyces*

cerevisiae) generated by random mutagenesis and identified three mutations (in *chs1*, *yme2* and *msh1*

genes) that changed the expression noise of a reporter gene in the nuclear genome regulated by the

yeast *TDH3* promoter. Each of these mutations affected either extrinsic noise (variability due to cell-

specific factors), intrinsic noise (inherent variability within each cell), or both. Surprisingly, all three

mutations targeted proteins located outside of the nucleus: Yme2 and Msh1 being involved in the

maintenance of mitochondrial genome integrity, and Chs1 being necessary for the repair of cell-wall

defects in freshly-born daughter cells. Our results reveal that mitochondrial state can modulate the

extent of intrinsic expression noise of eukaryotic nuclear genes.

INTRODUCTION

Gene expression levels vary among cells even in isogenic populations originating from a common cell type and maintained in a constant environment. This variability - commonly named *gene expression noise* - results from random fluctuations that inevitably occur along molecular and cellular processes impacting gene expression (reviewed in Pal & Dhar, 2024; Sanchez & Golding, 2013; Verhagen & Hansen, 2025). Expression noise can be modified by genetic changes, leading to differences in cell-to-cell variability among genotypes that can be either detrimental, neutral or beneficial depending on the gene and on the environmental condition; expression noise is therefore a substrate for evolution (M. Richard and Yvert, 2014). As a result, gene expression noise can take diverse forms depending on the shape of the distribution of expression levels observed among cells and how this distribution is modified by genetic changes. In extreme cases, a bimodal (or multimodal) distribution has been observed with two (or more) groups of isogenic cells displaying distinct expression levels of a gene. In certain cases, bimodal expression was shown to be part of a bet-hedging strategy when the two groups of cells were adapted to different environmental conditions (Baptista et al., 2025; Gamba et al., 2015; Levy et al., 2012). Even when the distribution of expression among cells is unimodal, which is the case for most genes (Newman et al., 2006; Taniguchi et al., 2010), genetic variants changing the variance of this distribution can impact population fitness (Duveau et al., 2018; Schmiedel et al., 2019) and therefore be targeted by selection (Metzger et al., 2015). Importantly, a global relationship is known to exist between mean expression and noise, whether among different genes (Newman et al., 2006; Taniguchi et al., 2010), among different genetic variants of a promoter (Carey et al., 2013; Hornung et al., 2012) or among environmental conditions (Bar-Even et al., 2006). Although the confounding effect of mean expression has complicated the study of the genetic sources and functional consequences of variation in expression noise, statistical and experimental approaches have been specifically designed to uncouple variation in these two metrics.

Several studies have shown that total noise in gene expression has two components, *extrinsic* and *intrinsic*, each attributable to distinct sources (Elowitz et al., 2002; Raser & O'Shea, 2004). Technically, extrinsic and intrinsic noise can be measured in cells where transcriptional activities of two copies of the same promoter are distinguishable (for instance using two fluorescent reporter genes of different colors; but see Stewart-Ornstein et al. (2012) for an alternative approach). In this setup, extrinsic noise represents expression variation among isogenic cells (or over time in a cell) that is correlated between the two copies of the promoter, while intrinsic noise represents independent

variation of expression between the two copies. Biologically, e*xtrinsic noise* results from sources
generating heterogeneity in global state among cells, with consequences on how individual cells
express the gene. In contrast, *intrinsic noise* corresponds to fluctuations that remain in cells after
accounting for their state due to the inherent stochasticity of regulatory processes. Experimental and
theoretical work have previously studied the properties of these two components and how specific steps
of gene regulation contribute to them. The predominant source of intrinsic noise was shown to be
transcription, which occurs stochastically in bursts (Blake et al., 2006; Harper et al., 2011; Pedraza &
Paulsson, 2008; Quarton et al., 2020; Suter et al., 2011). The extent to which transcription contributes
to intrinsic noise depends on chromatin remodeling (Raser & O'Shea, 2004), on nucleosome
occupancy (Tirosh & Barkai, 2008), on the presence of TATA boxes in regulatory sequences (Blake et
al., 2006; Choi & Kim, 2009; Hornung et al., 2012; Sun & Zhang, 2020) and on the number and type of
transcription factors binding to the promoter (Parab et al., 2022; Sharon et al., 2014). Extrinsic noise is
known to result from various cell-to-cell differences such as their pool of RNA polymerase (Yang et
al., 2014), micro-environment (Snijder et al., 2009), age (P. Liu et al., 2017; Martinez-Jimenez et al.,
2017), cell cycle stage (Sukys & Grima, 2025; Zopf et al., 2013), global transcription rate (Sherman et
al., 2015), mitochondrial content (Guantes et al., 2015) and proliferation rate (Keren et al., 2015).
Extrinsic noise was also shown to be reduced for genes regulated by miRNAs (Siciliano et al., 2013).

All factors modulating gene expression noise are possible targets for mutations and therefore
provide a potential substrate for the natural evolution of cell-to-cell variability. As for any trait, two
driving forces may influence the evolution of gene expression noise: the effects of random mutations
on noise and the effects of selection on the fate of these mutations in natural populations. Regarding the
action of selection on expression noise, several important observations have been reported. First, gene
expression noise can be evolved experimentally through artificial selection (Wolf et al., 2015; You et
al., 2019). Second, several studies provided evidence of negative selection acting against elevated
expression noise in natural populations, suggesting that the detrimental effect of elevated noise could
be counter-selected separately from any effect of a genetic variant on mean expression (Barroso et al.,
2018; Metzger et al., 2015; Puzović et al., 2023; Schor et al., 2017; Vlková & Silander, 2022).
Reciprocally, other studies showed that gene expression noise could be positively selected in certain
cases, particularly in fluctuating environments (Acar et al., 2008; Eldar & Elowitz, 2010; Rocabert et
al., 2020; Schmutzer & Wagner, 2020). When cells are maintained in a steady environment, increasing
expression noise of a gene can be either beneficial or detrimental depending on the mean expression
level and on the shape of the relationship that exists between the expression level of the gene and the

rate of cell proliferation (Duveau et al., 2018; Schmiedel et al., 2019). In yeast, transcriptome-wide

analyses revealed a particularly high level of expression noise for plasma-membrane transporters,

which play a critical role in cellular responses to environmental changes (Zhang et al., 2009).

Experimental studies in bacteria (Wolf et al., 2015) and yeast (Bódi et al., 2017) showed that elevated

expression noise could facilitate the evolution of gene regulation. By generating fitness variation

among individuals, expression noise can also potentially reduce the efficacy of selection and therefore

affect phenotypic evolution (Mineta et al., 2015; Wang & Zhang, 2011). In multicellular organisms,

expression noise was associated with cell fate decision (Balázsi et al., 2011; Coomer et al., 2022;

Ohnishi et al., 2014; Pal & Dhar, 2024; A. Richard et al., 2016; Wernet et al., 2006) and in the

resistance of cancer cells to chemotherapy (Shaffer et al., 2017). Finding what genetic variants and

molecular mechanisms can possibly modulate expression noise is therefore essential for understanding

many fundamental biological processes.

In this regard, important discoveries were made about the type of genetic changes that can tune

gene expression noise and their mechanisms of action. First, genetic variants segregating in natural

populations were found to generate different levels of gene expression noise among individuals. In

other words, variability among isogenic cells can itself vary among genetically-diverse individuals.

This was first shown in yeast (Ansel et al., 2008; Fehrmann et al., 2013; J. Liu et al., 2015; Metzger et

al., 2015) and later in bacteria (Vlková & Silander, 2022), flies (Huang et al., 2015) and human cells

(Lu et al., 2016; Morgan et al., 2020; Triqueneaux et al., 2020). Second, as previously demonstrated for

changes in mean expression (Wittkopp et al., 2004), a mutation may modify the expression noise of a

focal gene in two different ways (Raser & O'Shea, 2004): the mutation may act in *cis* (if the mutation

is at the locus of the regulated gene) or in *trans* (if the mutation is at a locus physically distant from the

regulated gene). Importantly, a single mutation may act in *cis* on the expression of a nearby gene and in

trans on the expression of another more distant gene, for example if the first gene is a regulator of the

second gene. *Cis* effects of mutations are better accessible for investigations than *trans* effects because

of the close proximity in the genome between *cis*-acting mutations and the gene with altered

expression. Several studies were therefore conducted on specific genes to understand how *cis*-located

mutations modify the expression noise of these genes. A seminal study focusing on the *GAL1* promoter

in yeast showed how mutations in the TATA box could modulate noise by altering the dynamics of

transcriptional bursts (Blake et al., 2006). In a later study, a more comprehensive random mutagenesis

of 22 yeast promoters revealed a mathematical relationship between mean expression and expression

noise, showing that mutations primarily altered the frequency of transcriptional bursts while burst size

differed among promoters, producing promoter-specific relationships between mean and noise (Hornung et al., 2012). Using an ingenious high-throughput approach to quantify mean expression and expression noise from thousands of designed synthetic promoters, another yeast study showed that both the number of functional transcription-factor binding sites and the frequency of poly(dT/dA) motifs - which favor nucleosome positioning – were key determinants of expression noise (Sharon et al., 2014). A fourth study compared the effects of 236 artificial mutations and 45 natural polymorphisms in the promoter of the yeast gene *TDH3*, providing evidence of negative selection against *cis*-acting mutations that increased noise of *TDH3* promoter activity (Metzger et al., 2015).

Although there are only few known examples of mutations affecting gene expression noise in *trans*, their effects on noise may differ from those of *cis*-acting mutations. For example, in the case of the yeast *TDH3* promoter, *trans*-acting mutations obtained via random mutagenesis of the genome were found to have a broader distribution of effects on expression noise than *cis*-acting mutations located in the *TDH3* promoter; in addition, random *trans*-acting mutations were as likely to increase as to decrease expression noise, whereas *cis*-acting mutations were more likely to increase expression noise (Metzger et al., 2016). Consistently, a study quantifying expression noise of surface proteins in human cells detected more *trans*-acting than *cis*-acting genetic regulators, suggesting that natural variation modulating gene expression noise was more likely to occur in *trans* (Morgan et al., 2020). Intuitively, another possible difference between *cis* and *trans* effects of mutations could be their impact on intrinsic and extrinsic noise. Indeed, theoretical models predict that *cis*-acting mutations in a regulatory sequence may modulate intrinsic noise by altering the frequency and magnitude of transcriptional bursts (Pedraza & Paulsson, 2008; Zenklusen et al., 2008). It is more difficult to imagine a scenario where a *cis*-acting mutation would modulate extrinsic noise. Mutations affecting expression noise in *trans*, however, could modulate either intrinsic or extrinsic noise. For instance, while mutations in a transcription factor may affect intrinsic noise of its target genes, mutations in a cell cycle regulator may impact extrinsic noise of other genes by generating heterogeneity of states among cells. Given these differences, it is necessary to further characterize what type of mutations can modulate gene expression noise in *trans* and what their mechanisms of action are to understand how expression noise is controlled genetically and evolves. One way to do so is by introducing target mutations in molecular complexes participating to gene expression and determine how they modify cell-to-cell variability in protein levels, as was done for mutations targeting chromatin remodeling (Raser & O'Shea, 2004) or transcription elongation (Ansel et al., 2008; Trinh et al., 2023). Such targeted approach, however, is driven by existing knowledge on gene regulation and does not interrogate the possible roles of

unexpected players. An alternative approach is to identify *trans*-regulators segregating in natural

populations using adapted GWAS or QTL mapping methods (Chuffart et al., 2016; Fehrmann et al.,

2013; Lu et al., 2016; Morgan et al., 2020; Sarkar et al., 2019; Yvert, 2014). Although this approach is

well-suited to characterize genetic variants that have been filtered by selection, it does not capture all

possible genetic sources of variation in expression noise. Identifying random mutations that modulate

expression noise of a gene in *trans* - regardless of their fitness consequences - is an important

complement to understand how the evolution of expression noise is shaped by the mutational process

and natural selection (Hill et al., 2021).

Here we used a random mutagenesis screen followed by genetic mapping and functional assays

to identify mutations that changed in *trans* the expression noise of a focal gene in yeast. We took

advantage of a collection of 1241 mutant strains obtained by chemical mutagenesis of a parental strain

expressing a fluorescent reporter gene controlled by the *TDH3* promoter (P_{TDH3}-*YFP*). In its native

context, the *TDH3* promoter drives strong constitutive expression of a glyceraldehyde-3-phosphate

dehydrogenase involved in glycolysis and gluconeogenesis (Partow et al., 2010). Using flow

cytometry, we identified five mutants from the collection that showed altered P_{TDH3}-*YFP* expression

noise relative to the parental strain without significant change in mean expression. Via genetic

mapping, we identified six candidate causal mutations in these mutants and we validated the effects of

three of them using strains engineered to each contain only one of these three candidate mutation

(single-site mutants). We then characterized the contribution of these mutations to extrinsic and

intrinsic noise using two-color reporters. The three mutations altered the coding sequences of *YME2*,

MSH1 and *CHS1* genes, none of which were previously associated with the control of noise or mean

gene expression. In particular, our results suggest a potentially central and unexpected role of

mitochondrial genome integrity in the modulation of intrinsic expression noise of nuclear genes.

RESULTS

Mapping mutations with *trans*-acting effects on expression noise

We previously identified mutations that affected the median expression level of a focal gene

(P_{TDH3}-*YFP*, where expression of a *YFP* reporter gene is controlled by the *TDH3* promoter) from a yeast

collection obtained by ethyl methanesulfonate (EMS) mutagenesis of a reference strain (Duveau et al.,

2021; Metzger et al., 2016). Here, we took advantage of this collection to identify and characterize

mutations that altered expression noise of the focal gene in *trans* (*i.e.* mutations outside of the *TDH3*

promoter and causing increased or decreased cell-to-cell variability of fluorescence level). To this end,

we first quantified median expression and expression noise (coefficient of variation of normalized

fluorescence; see Methods) of P_{TDH3}-*YFP* in 1241 EMS mutants from the collection and in the reference

strain by measuring fluorescence of thousands of cells for each strain using flow cytometry (**Figure

1a**). This initial screen revealed a wide variation of expression noise among mutants that was largely –

but not completely – independent from variation in median expression. Next, we selected 254 mutants

that showed largest changes of expression noise relative to the reference strain and we quantified their

fluorescence a second time in an independent assay (**Figure 1 – figure supplement 1**). Among these

254 mutants, we chose five mutants for further analyses that showed significant changes of expression

noise relative to the reference strain in both assays (*t*-test, adjusted $P < 0.01$), with no significant

change in median expression in the first assay (*t*-test, adjusted $P \geq 0.05$). Three of these mutants had an

increased level of P_{TDH3}-*YFP* expression noise and two mutants had a decreased level of noise (**Figure

1a**, **Table 1**). This procedure ensured that the five selected mutants showed reproducible changes in

expression noise that were not explained by variation of median expression levels. By whole-genome

re-sequencing (median coverage of 46x), we found between 17 and 47 mutations in the genomes of

these five mutants (**Table 1**), which was consistent with the range (14 to 41) of mutations previously

identified in other mutants from the same collection (Duveau et al., 2021). Given that the vast majority

of mutants in the collection showed no significant change in expression noise, we expected that only

one or a few mutations were responsible for the altered expression noise in each of the five selected

mutants. We therefore applied a genetic mapping strategy specifically designed to identify these

presumably-rare mutations (**Figure 1b**). We used a bulk segregant analysis approach where each

haploid mutant was crossed to a wild-type strain also expressing P_{TDH3}-*YFP* and random segregants

obtained from the meiosis of resulting diploid cells were sorted by FACS based on their fluorescence

levels (see Methods). Among all segregants, the subset of cells that inherited a mutation responsible for

a change in expression noise of the fluorescent reporter (a causative mutation) should show either a broader or a narrower distribution of fluorescence levels than the subset of cells inheriting the wild-type allele. For this reason, we sorted random segregants in three bulks that captured distinct ranges of fluorescence levels: one bulk corresponding to the 2% of cells with lowest fluorescence (low bulk), one bulk corresponding to the 2% of cells with highest fluorescence (high bulk) and one bulk corresponding to the 10% of cells with fluorescence closest to the median fluorescence of all segregants (mid bulk). We reasoned that a causative mutation should be found at different frequencies in the mid bulk than in the low and high bulks. To further increase this difference, we repeated the fluorescence selection twice for each bulk with a few generations of growth between each selection step (**Figure 1b**). For each cross, frequencies of all segregating mutations were then quantified by deep sequencing of genomic DNA extracted from each bulk. Likelihood-ratio tests (*G*-tests) were used to determine if the frequency of each mutation was significantly different between the mid bulk and the combined low and high bulks (**Figure 1 – figure supplement 2**), in which case the mutation was considered causative (or genetically linked to a causative mutation). Using this approach, we found one candidate causative mutations in two out of five mutants analyzed and two candidate mutations in another mutant (**Table 2**). For mutant YPW2162, although no mutation was significantly associated with expression noise, a single mutation was found at very low frequency in all fluorescence bulks (*m2* in **Table 2**). This indicated that the mutation was strongly counter-selected during the procedure used to establish fluorescent bulks, which greatly reduced the statistical power to detect potential association with expression noise. For this reason, we considered mutation *m2* to be possibly causative and further investigated its impact on expression noise in the functional assays described below. For the last mutant (YPW2143), we noticed that the frequency of all 26 mutations was either near zero or near one in the low bulk of segregants. This unexpected observation suggested that a single haplotype was nearly fixed in the low bulk, either due to a strong selective advantage of this haplotype or more likely due to a strong bottleneck in population size (random genetic drift) when this bulk was established. For this reason, we only compared allele frequencies between mid and high bulks, but no mutation was found at a significantly different frequency between these two bulks (**Figure 1 – figure supplement 2c**). Therefore, we used another strategy based on linkage analysis and tetrad dissection to determine which mutation in mutant YPW2143 was associated with variation of expression noise. The candidate mutation we tested first was mutation *m3* (**Table 2**) because it was found at a slightly higher frequency in the high bulk (consistent with the increased noise observed in YPW2143) and this mutation appeared to be counter-selected in both bulks, which reduced the statistical power to detect different frequencies

of the mutation between bulks. After meiosis of diploid cells from the cross between YPW2143 and the

mapping wild-type strain, we measured P_{TDH3}-*YFP* expression noise in 84 segregant clones issued from

21 dissected tetrads. We observed a very strong association between expression noise and the presence

or absence of mutation *m3* in each clone (**Figure 1 – figure supplement 3**; ANOVA, *F* value = 792.31,

P = 1.47 x 10^{-82}).

Functional validation of effects of candidate mutations on expression noise

Statistical associations described above may be explained by a causal effect of candidate

mutations on P_{TDH3}-*YFP* expression noise, but they could also result from genetic linkage with a

causative mutation or from unsuspected selection for another trait. To demonstrate the causal impact of

each candidate mutation (six mutations in **Table 2**) on expression noise, we used site-directed

mutagenesis to introduce each mutation in a common genetic background (reference strain) that does

not harbor any other mutations found in EMS mutants and that contains two distinct reporter genes

inserted in opposite orientations at the *ho* locus (**Figure 1c**). As further described in the next sections,

the use of two reporter genes (*YFP* and *RFP* coding sequences transcribed under control of distinct

copies of the *TDH3* promoter) allowed us to distinguish two sources of expression noise. We then used

flow cytometry to quantify median expression levels and expression noise of all site-directed mutants,

the reference strain and EMS mutants from which candidate mutations were identified (note that EMS

mutants do not express the *RFP* reporter gene). In almost all experiments reported below, we included

two independent clones for each genotype (*i.e.* two strains established from two colonies from the same

transformation) to detect cases where a secondary mutation not shared by all transformed cells would

contribute to phenotypic variation (*i.e.* cases where the two independent clones would show statistically

different phenotypes).

For three candidate mutations (*m1*, *m2* and *m3*), we observed a significant difference of

expression noise between site-directed mutants and the reference strain (*t*-test, adjusted *P* < 0.01), both

for YFP (**Figure 2a-c**) and for RFP (**Figure 2 – Figure supplement 2g-i**). In all three cases, the

mutation caused an increased in expression noise relative to the allele of the reference strain. In

addition, no significant difference of YFP expression noise was observed between each site-directed

mutant and the EMS mutant carrying the same mutation (**Figure 2a-c**), indicating that the three

candidate mutations were responsible for most, if not all, of the variation of expression noise observed

in EMS mutants from which they were identified. As expected, we observed no significant difference

of median expression between the reference strain and site-directed or EMS mutants carrying mutations

m1 and *m2* (*t*-test, adjusted $P \geqslant 0.01$), both for YFP (**Figure 2 – figure supplement 1a-b**) and for RFP

(**Figure 2 – figure supplement 2a-b**). However, all strains carrying mutation *m3* showed a significant

increase of YFP (**Figure 2 – figure supplement 1c**) and RFP (**Figure 2 – figure supplement 2c**)

median expression levels relative to the reference strain (*t*-test, adjusted $P < 0.01$), which was

unexpected based on the initial screen (**Figure 1a**) where the EMS mutant carrying mutation *m3* did

not show an increased median expression level. We further investigated the cause of this apparent

inconsistency below (**Figure 6 – figure supplement 2**).

For the other three candidate mutations tested (*m4, m5* and *m6*), no significant change in

expression noise was observed between site-directed mutants and the reference strain for YFP (**Figure

2d-f**) and RFP (**Figure 2 – figure supplement 2j-l**); except for one of the two tested clones carrying

mutation *m5* and only for RFP, which we considered to be a false positive. These mutations also did

not significantly impact the median expression level of YFP (**Figure 2 – figure supplement 1d-f**) and

RFP (**Figure 2 – figure supplement 2d-f**). Because mutations *m5* and *m6* were identified from the

same EMS mutant, we hypothesized that these mutations may be required together to significantly

impact expression noise. We therefore constructed a double mutant carrying both *m5* and *m6*, but this

mutant did not show any significant change of expression noise relative to the reference strain (**Figure

2 – figure supplement 3**). In these experiments, the two EMS mutants from which *m4, m5* and *m6*

mutations were identified showed a significant change of expression noise relative to the reference

strain, as expected. Several hypotheses may explain why we did not find a causative mutation for these

two EMS mutants (see Discussion).

A missense mutation in *YME2* gene increases intrinsic noise of P_{TDH3}-*YFP* expression.

To further investigate potential mechanisms of action for each causative mutation, we next

measured how these mutations affected two distinct sources of noise: intrinsic and extrinsic noise. Even

though intrinsic and extrinsic noise both contribute to the total expression noise of a gene, they are

confounded when expression is measured from a single allele. However, these sources of noise can be

distinguished when measuring expression from two reporter genes (in our case P_{TDH3}-*YFP* and P_{TDH3}-

RFP) in individual cells (**Figure 3a**), as previously demonstrated (Elowitz et al., 2002). Indeed, while

intrinsic noise creates differences of expression levels between the two reporters within a cell, extrinsic

noise creates expression variation among cells that is well correlated between the two reporters (**Figure

1c**). Intrinsic noise is expected to result from stochastic processes involved in the independent

transcription, translation and post-translational regulation of each reporter gene, while extrinsic noise

results from more global differences among cells such as differences in physiological states or in the

amount of transcription factors regulating both reporters (Raser & O'Shea, 2005). We calculated

intrinsic and extrinsic noise of mutant and reference strains from YFP and RFP fluorescence levels of

individual cells as described in Fu & Pachter (2016) (see Methods).

We first focused on mutation *m1* corresponding to a missense mutation located in the coding

sequence of the *YME2* gene (*G1234a* substitution; **Table 3**). Site-directed mutants carrying the mutant

allele *yme2(G1234a)* showed an increased intrinsic noise relative to the reference strain (**Figure 3b**; *t*-

test, adjusted $P < 0.001$ for both clones). This effect is visible through the lower correlation observed

between YFP and RFP expression levels of mutant cells than wild-type cells (**Figure 3 – figure**

supplement 1). The impact of *yme2(G1234a)* mutation on extrinsic noise was less clear as only one of

the two tested clones showed a significant increase in extrinsic noise (**Figure 3c**; *t*-test, adjusted $P <$

0.001 for clone 1 and adjusted $P \geqslant 0.05$ for clone 2). Overall, the impact of *yme2(G1234a)* mutation on

intrinsic noise explained most of the variation in total expression noise of the two reporter genes

(**Figure 3d**). This is particularly surprising since YME2p is an integral protein of the mitochondria

inner membrane with known functions linked to fusion and fission of mitochondrial nucleoids and

integrity of the mitochondrial genome (Hanekamp & Thorsness, 1996; Park et al., 2006). Indeed,

YME2p is not known to directly regulate the expression of nuclear genes. An indirect effect of

mitochondrial activity on transcriptional or post-transcriptional regulation of nuclear genes would be

expected to affect the expression of both reporter genes in a coordinated fashion, resulting in variation

of extrinsic noise, which is not what we observed. Therefore, we tested whether potential experimental

artifacts may explain the elevated intrinsic noise of *yme2(G1234a)* mutant. In particular, the detection

of intrinsic noise may not only by caused by biological noise, but also by technical noise in the

measurements of YFP an RFP fluorescence using flow cytometry. To exclude this possibility, we fused

the coding sequences of *YFP* and *RFP* downstream of the *TDH3* promoter sequence (P_{TDH3}-*YFP-RFP*)

and inserted the resulting transgene at the *ho* locus in genetic backgrounds carrying either the wild-type

YME2 allele or the *yme2(G1234a)* mutation. We reasoned that expressing a YFP-RFP protein fusion

would lead to the same amount of yellow and red chromophores within a cell (variation can still exist

among cells), which would completely prevent the detection of biological intrinsic noise (**Figure 3e**).

Therefore, any difference of intrinsic noise detected in this context could be attributed to technical

noise affecting the detection of YFP and RFP signals. We observed no significant effect of

yme2(G1234a) mutation on intrinsic noise detected in strains expressing P_{TDH3}-*YFP-RFP* (**Figure 3f**),

while the mutation significantly increased levels of extrinsic noise (**Figure 3g**) and total noise (**Figure**

3h). This result confirmed that the impact of *yme2(G1234a)* mutation on intrinsic noise detected with
two reporter genes was caused by actual differences in the amount of YFP and RFP molecules within
each cell.

We noticed in flow cytometry data a small but statistically significant decrease of median cell
size (estimated from forward scatter signal) in *yme2(G1234a)* mutants relative to the reference strain
(**Figure 4a**). When clustering cells in five quantiles based on cell size for each genotype, we observed
that the increase of intrinsic noise associated with *yme2(G1234a)* mutation was more pronounced in
smaller cells (**Figure 4b**). The extrinsic noise of wild-type and mutant cells was indistinguishable
regardless of their size, even though a negative relationship was globally observed between extrinsic
noise and the size of genetically identical cells (**Figure 4c**). In budding yeast with asymmetric cell
division, smallest cells in non-synchronized populations often correspond to daughter cells just after
division. Although variation of intrinsic noise is typically thought to result from alteration in processes
regulating gene expression (Hypothesis 1 in **Figure 4d**), an alternative hypothesis would be that the
effect of *yme2(G1234a)* mutation on intrinsic noise was caused by altered partitioning of YFP and RFP
molecules in daughter cells after division, potentially because of their reduced size compared to wild-
type daughter cells (Hypothesis 2 in **Figure 4d**). To test these hypotheses, we constructed a P_{TDH3}-*YFP*-
RFP gene fusion where coding sequences of *YFP* and *RFP* were separated by a short sequence
encoding the cleavage site for the tobacco etch virus (TEV) protease (ENLYFQS peptide sequence).
This construct was inserted at *ho* locus in strains that constitutively express the wild-type TEV or the
improved uTEV3 protease from Sanchez and Ting (2020), with either the wild-type or *G1234a* mutant
allele of *YME2*. We reasoned that sources of noise acting on expression at any step from transcription
to translation would be undetectable in these strains because YFP and RFP are expressed from the same
coding sequence. However, if the increased intrinsic noise was caused by altered partitioning of YFP
and RFP molecules, we should still be able to detect it in these strains because YFP-RFP proteins are
cleaved into YFP and RFP peptides that are free to segregate independently in mother and daughter
cells after division (**Figure 4d**). Western blotting with an antibody against RFP showed that YFP-RFP
was fully cleaved in strains expressing the improved uTEV3 protease (undetectable level of YFP-RFP
fusion protein), but only partially cleaved in strains expressing the wild-type TEV (**Figure 4e**).
Therefore, we measured expression noise in strains expressing uTEV3. We observed no detectable
effect of mutation *yme2(G1234a)* on intrinsic noise in strains producing YFP and RFP peptides from
cleavage of YFP-RFP fusion proteins (**Figure 4f**; *t*-test, adjusted $P > 0.05$), regardless of cell size
(**Figure 4h**). However, a significant increase of extrinsic noise was observed for *yme2(G1234a)* strains

relative to wild-type strains (**Figure 4g**; *t*-test, adjusted $P < 0.001$ for clone 1, adjusted $P < 0.01$ for clone 2). These results are not consistent with the hypothesis that *yme2(G1234a)* mutation affects expression noise via altered partitioning of proteins during cell division. Instead, the mutation must cause an increased intrinsic noise by acting on a process regulating gene expression at some point from transcription to protein degradation, via an unknown mechanism.

A nonsense mutation in *CHS1* increases extrinsic noise via impaired chitin septum reparation in daughter cells.

We next quantified the impact of mutation *m2* on extrinsic and intrinsic noise using site-directed mutants that expressed both P_{TDH3}-*YFP* and P_{TDH3}-*RFP*. *m2* is a nonsense mutation in the *CHS1* gene (*G1752a* substitution) that introduces an early stop codon reducing the open reading frame by 48.4% (**Table 3**). We found that *chs1(G1752a)* allele had no clear impact on intrinsic noise (**Figure 5a**; *t*-test, adjusted $P \geqslant 0.05$ for clone 1 and $P < 0.01$ for clone 2 with decreased intrinsic noise). However, it significantly increased extrinsic noise (**Figure 5b**; *t*-test, adjusted $P < 0.01$ for clone 1 and $P < 0.001$ for clone 2) and total noise (**Figure 5c**; *t*-test, adjusted $P < 0.01$ for clone 1 and $P < 0.001$ for clone 2) relative to the reference allele. Consistent with this increase in extrinsic noise, *chs1(G1752a)* cells showed stronger correlation of YFP and RFP expression levels than wild-type cells, with a higher proportion of mutant cells showing either low or high expression of both reporters (**Figure 5 – figure supplement 1**). As expected, we also detected an increased extrinsic noise in *chs1(G1752a)* mutant cells that expressed a YFP-RFP fusion protein (**Figure 5 – figure supplement 2**).

We next investigated what mechanisms could explain the effect of *chs1(G1752a)* mutation on expression noise based on the known function of *CHS1*. *CHS1* encodes a chitin synthase specifically involved in the repair of the chitin septum after cell division in daughter cells (Cabib et al., 1989, 1992), which are the smaller cells resulting from asymmetric division in budding yeast. Detachment between mother and daughter cells involves partial degradation of the septum on the daughter side, which can sometimes damage the cell wall of daughter cells and lead to their lysis in the absence of repair. *chs1(G1752a)* allele generates an early stop codon that leads to a truncated protein lacking several residues and domains that are essential for *chs1* function (Chen et al., 2023); therefore, it is likely to be a null allele of *chs1*. Flow cytometry data showed an enrichment of cells with a smaller size for the *chs1(G1752a)* site-directed mutant relative to the reference strain (**Figure 5d**), consistent with an accumulation of daughter cells with damaged or lysed cell wall in the mutant. When splitting wild-type and *chs1(G1752a)* samples in different intervals of cell size, we observed that the median

expression level of P_{TDH3}-YFP did not depend on cell size (**Figure 5e**). Therefore, the increased expression noise of *chs1(G1752a)* strain is not explained by a difference of median expression level between damaged daughter cells and larger cells in this mutant. Instead, we observed a large increase of extrinsic noise mostly in the smallest cells of the *chs1(G1752a)* mutant: larger *chs1(G1752a)* cells and wild-type cells of all sizes showed much lower extrinsic noise (**Figure 5f**). This observation is consistent with an increased extrinsic noise in damaged daughter cells that accumulate in the *chs1(G1752a)* mutant.

To further show that the effect of *chs1(G1752a)* mutation on extrinsic noise was due to impaired chitin septum repair, we measured expression noise of wild-type and mutant cells grown in presence of glucosamine in the culture medium. Supplementing the culture medium with glucosamine is known to promote chitin synthesis via the activity of CHS3p (Bulik *et al.,* 2003), preventing damage to the chitin septum in daughter cells. Therefore, any consequence of *chs1(G1752a)* mutation caused by impaired chitin septum repair is expected to be suppressed in presence of glucosamine. We indeed observed that the extrinsic noise of the smallest *chs1(G1752a)* mutant cells decreased with increasing doses of glucosamine in the culture medium (**Figure 5g**). At 15 mM glucosamine, there was almost no difference of extrinsic noise and of cell size between wild-type and *chs1(G1752a)* cells. When considering all cell sizes (**Figure 5 – figure supplement 3c**), the increase of extrinsic noise in *chs1(G1752a)* cells relative to wild-type cells varied from 136% in absence of glucosamine (*t*-test, $P = 3.35 \times 10^{-7}$) to only 12% in presence of 15 mM glucosamine in the culture medium (*t*-test, $P = 5.8 \times 10^{-3}$). We also observed a gradual decrease of median cell size with increasing dose of glucosamine, a pattern that was more pronounced for wild-type cells than for *chs1(G1752a)* cells (**Figure 5 – figure supplement 3a**). This is consistent with the previous finding that yeast cells grown in presence of glucosamine retain less water, for unknown reasons (Bulik *et al.,* 2003). Importantly, the doses of glucosamine used did not significantly impact cell growth rates during the first phase of growth after glucosamine was added to the medium (**Figure 5 – figure supplement 4**). Altogether, these observations strongly suggest that *chs1(G1752a)* mutation increases extrinsic noise of P_{TDH3} activity exclusively in daughter cells with damaged chitin septum, a phenomenon that is suppressed in presence of glucosamine. The mechanism linking chitin septum repair to extrinsic noise of gene expression remains to be characterized.

Density-dependent effect of a missense mutation in *MSH1* gene on intrinsic and extrinsic noise.

Mutation *m3* is a missense mutation located in the coding sequence of the *MSH1* gene (*G1262a*

substitution; **Table 3**). As mentioned above, we observed considerable variation of P_{TDH3}-*YFP* median

expression and expression noise in the EMS mutant carrying mutation *m3* (YPW2143) between

independent experiments. In the initial mutagenesis screen (**Figure 1a**), mutant YPW2143 showed a

modest decrease of median expression (-0.9%) and a large increase of expression noise (+72.8%)

relative to the reference strain. However, in a subsequent experiment (**Figure 2d**, **Figure 2 – figure**

supplement 1c), the same mutant showed a substantial increase of median expression (+6.5%) and a

much lower increase of expression noise (+6.3%). Since these two experiments were performed in

different laboratories, the observed phenotypic variation could be caused by uncontrolled variation in

experimental conditions. We hypothesized that the initial density of cell cultures – a parameter difficult

to control exactly when handling a large number of samples in parallel – might influence the effect of

msh1(G1262a) mutation on P_{TDH3}-*YFP* expression. To test this hypothesis, we initiated cell cultures

from four different dilutions of a preculture and incubated these cultures for the same duration (20

hours) at 30°C before quantifying absorbance as a proxy for final cell density and expression using

flow cytometry. This experimental procedure was applied to haploid strains expressing both P_{TDH3}-*YFP*

and P_{TDH3}-*RFP* reporters with either the wild type allele or the *G1234a* allele of *MSH1*. We found a

strong association between the initial dilution of cell cultures and the effects of *msh1(G1234a)*

mutation on expression noise of either P_{TDH3}-*YFP* (**Figure 6a**) or P_{TDH3}-*RFP* (**Figure 6b**). Expression

noise increased with initial cell density both for the reference strain and for the *msh1(G1262a)* mutant,

but this trend was stronger for the *msh1(G1262a)* mutant, resulting in higher expression noise for the

mutant relative to the reference strain at the highest initial density tested (*i.e.* 16:500 dilution). At the

lowest initial cell density (*i.e.* 1:500 dilution), we observed a negligible increase of expression noise for

P_{TDH3}-*YFP* (+4.2%; *t*-test, adjusted P = 0.41) and P_{TDH3}-*RFP* (+6.4%; *t*-test, adjusted P = 0.035) in the

msh1(G1262a) mutant relative to the reference strain (**Figure 6 – figure supplement 2**). In contrast, at

the highest initial cell density (*i.e.* 16:500 dilution), expression noise was increased by 45.7% for P_{TDH3}-

YFP (*t*-test, adjusted P = 3.0 x 10^{-4}) and by 34% for P_{TDH3}-*RFP* (*t*-test, adjusted P = 9.9 x 10^{-4}) in

msh1(G1262a) mutant cells (**Figure 6 – figure supplement 2**). We also observed a complex

relationship between initial cell density and median expression of P_{TDH3}-*YFP* and P_{TDH3}-*RFP* (**figure 6 –**

figure supplement 2): median expression remained constant among the different initial densities tested

for the reference strain; however, for *msh1(G1262a)* mutant cells, median expression changed

significantly depending on initial cell density and this variation was not the same for P_{TDH3}-*YFP* and

P_{TDH3}-*RFP* median expression. The fact that initial cell density altered the effect of *msh1* mutation on

expression noise and median expression can explain the variability we previously observed between

independent experiments. However, variation of initial cell density is not necessarily the direct cause of

expression variation. A more direct cause could be a variation of final cell densities after growth of

cultures initiated at different cell densities. This seems unlikely, though, as we observed significant

differences of intrinsic (*t*-test, adjusted $P = 1.98 \times 10^{-3}$) and extrinsic (*t*-test, adjusted $P = 4.42 \times 10^{-2}$)

noise for *msh1* mutant cells diluted at 2:500 and at 4:500 (**Figure 6 – figure supplement 2**) despite no

significant difference of final cell densities (**Figure 6 – figure supplement 2b**; *t*-test, adjusted $P =$

0.24). Cultures initiated at different cell densities are expected to reach different phases of growth after

20 h of incubation at 30°C, which could explain why we observed density-dependent effects of

msh1(G1262a) mutation on median expression and expression noise.

We found that *msh1(G1262a)* mutation impacted both extrinsic and intrinsic noise (**Figure 6c**),

but only at the highest initial cell density tested (*i.e.* 16:500 pre-culture dilution) and the effect on

intrinsic noise was greater that the effect on extrinsic noise (**Figure 6 – figure supplement 2**). No

significant difference of intrinsic noise (*t*-test, adjusted $P = 0.69$) and extrinsic noise (*t*-test, adjusted P

$= 0.16$) was observed between *msh1(G1262a)* mutant cells and wild-type cells for a 1:500 pre-culture

dilution, while a 244% increased intrinsic noise (*t*-test, adjusted $P = 2.8 \times 10^{-3}$) and a 41% increased

extrinsic noise (*t*-test, adjusted $P = 0.058$) was observed in mutant relative to wild-type cells for a

16:500 pre-culture dilution. The effect of *msh1* mutation on intrinsic noise is visible at the single cell

level via the lower correlation observed between YFP and RFP expression levels of mutant cells

relative to wild-type cells (**Figure 6 – figure supplement 1**).

MSH1 encodes a DNA-binding protein of the mitochondria involved in the repair of

mitochondrial DNA. Disruption of this gene is known to cause a high frequency of respiratory-

deficient cells (*ρ*- petite cells), which may generate cell-to-cell heterogeneity and could provide a

simple explanation for the increased expression noise observed in the *msh1(G1262a)* mutant. We

compared the frequencies of *ρ*- cells in wild-type and *msh1(G1262a)* mutant strains after growth in a

rich medium (YPD) containing a fermentable carbon source (see Methods) and using two different pre-

culture dilutions (1:500 and 8:500). Both *ρ*+ and *ρ*- cells can grow in this medium. As expected, we

observed a much lower frequency of *ρ*- cells in the wild-type strain (1.3% and 0.3% *ρ*- cells for pre-

culture dilutions of 1:500 and 8:500; **Figure 6d**) than in the *msh1(G1262a)* mutant strain (94.5% and

88.6% *ρ*- cells for pre-culture dilutions of 1:500 and 8:500; **Figure 6d**). We isolated a *ρ*- colony from

the *msh1(G1262a)* mutant strain and noticed, as expected, that it produced 100% of *ρ*- cells after

growth in YPD medium. We reasoned that if the impact of *msh1(G1262a)* mutation on expression

noise was explained by the co-existence of $\rho+$ and $\rho-$ cells in the mutant strain, then expression noise
should be reduced in the *msh1(G1262a)* strain with 100% of $\rho-$ cells, and its noise should no longer
depend on initial cell density. However, fixing the frequency of $\rho-$ cells to 100% did not reduce the
effect of *msh1(G1262a)* mutation on intrinsic and extrinsic noise (**Figure 6d**; **Figure 6 - figure**
supplement 3). In addition, the expression noise of *msh1(G1262a)* strain was higher in cultures
initiated at high cell densities than in cultures initiated at low cell density regardless of the frequency of
$\rho-$ cells in the population (**Figure 6d**; **Figure 6 - figure supplement 3**). These observations indicate
that the effect of *msh1(G1262a)* mutation on expression noise is not directly explained by cell-to-cell
heterogeneity in respiratory capacity caused by the mutation.

DISCUSSION

In this study, we used random mutagenesis followed by genetic mapping to identify three
trans-acting mutations that increased cell-to-cell variability of expression of a yeast reporter gene. This
unbiased approach made it possible to discover any mutation that could possibly alter the expression
noise of the focal gene. Indeed, all three mutations identified here were located in the coding sequence
of genes that were not known regulators of *TDH3* promoter activity and were therefore not expected to
regulate P_{TDH3}-*YFP* expression noise. Importantly, these three mutations had very little impact on the
median expression of P_{TDH3}-*YFP*. Their action on expression noise can therefore not be explained by
the well-known scaling of expression noise with expression mean (Newman et al., 2006). The three
mutations targeted proteins with different biochemical functions involved in different – but not
necessarily unrelated – cellular processes. *chs1(G1752a)* non-sense mutation increased extrinsic noise
exclusively in the smallest cells of yeast populations, which was likely caused by damaged cell walls of
daughter cells improperly repaired in the absence of chitin synthase I activity. The second mutation
targeted the inner mitochondrial membrane protein Yme2 and increased intrinsic noise of P_{TDH3}
expression. The third mutation, targeting the Msh1 ATPase involved in mitochondrial DNA repair, also
increased P_{TDH3} intrinsic noise and this effect varied with the growth phase of cells. Intriguingly,
although *yme2(G1234a)* and *msh1(G1262a)* mutations impacted genes with distinct functions, both
genes are important for the maintenance of mitochondrial genome integrity, leaving open the
possibility that these mutations modulate intrinsic expression noise via a common mechanism. After
describing the potential limits of our screen-based approach, we discuss this possibility, as well as the
main implications of these results, below.

We decided to identify genetic modulators of expression noise by performing a random
mutagenesis screen rather than testing candidate genes because it allowed the discovery of unknown
mechanisms without ascertainment biases. Although EMS mutagenesis mostly induces G:C to A:T
substitutions, which could slightly bias the screen towards GC-rich genes and coding sequences
(relative to regulatory sequences), we do not expect this potential bias to significantly impact our
conclusions. Indeed, *YME2*, *CHS1* and *MSH1* coding sequences are not particularly enriched in G or C
nucleotides (GC content is 39.7%, 36.8% and 37.0% in these three genes relative to 39.6% among all
ORFs) and, in the yeast genome, GC content is not much lower in intergenic sequences (34.4%) than in
ORFs (39.6%). We note that bulk-segregant analysis (BSA-seq), which is generally very powerful to
map causal mutations in yeast (Duveau et al., 2014; Ehrenreich et al., 2010; Magwene et al., 2011), was
only successful here for two of the five mutants that we analyzed (we identified the third causal
mutation by another approach based on tetrads). Several possibilities may explain the limited power of
BSA-seq in the present study. First, the statistical power to map a variant impacting expression noise is
expected to be lower than to the power to map a variant impacting mean expression (Sarkar et al.,
2019). Second, based on previous simulations (Duveau et al., 2014), our procedure included three
cycles of growth and FACS selection. This is predicted to increase detection power, but only for
mutations that do not impair fitness during the growth phase of each cycle. It is therefore possible that
some mutations causing elevated expression noise escaped BSA-seq detection because they also
impacted growth. Another possibility is that the effect of these mutations may be particularly sensitivite
to microenvironmental conditions, which could have differed between the characterization of the
mutants and the conditions of BSA-seq selection. In the present mapping study where the genetic
diversity between the two parental strains was low, tetrad analysis proved to be more powerful than
BSA-seq to map mutations altering gene expression noise.

Few studies previously characterized *trans*-acting genetic variants impacting expression noise.
Besides pioneering work showing the effect of chromatin remodelers on noise using gene deletions
(Raser & O'Shea, 2004), expression-noise modulators were also identified as natural polymorphisms in
genes involved in cell-sensing and responses to environmental variation. The first examples discovered
were the *MUP1* and *ERC1* genes, which regulate the import of extracellular methionine in yeast cells,
and where natural polymorphisms affected noise in expression of the methionine-sensitive reporter
P_{MET17}-*GFP* (Fehrmann et al., 2013). Another example was later identified in humans, where a disease-
associated variant in the *PDE4D* phosphodiesterase - controlling cAMP signalling - was associated
with noise in expression of the HLA-DR receptor at the surface of CD8+ T cells and of the CD38

receptor at the surface of CD4+CD45RA+ naive T cells (Lu et al., 2016). Therefore, we could have

expected our screen to reveal mutations perturbing cell sensing, but this was not the case. Our screen

also did not reveal mutations in transcription factors known to regulate *TDH3* promoter activity, or in

the TATA box binding protein (TBP), which were obvious candidates because mutations in

transcription factor binding sites and in the TATA box of the *TDH3* promoter are known to sometimes

alter expression noise independently from mean expression (Duveau et al., 2018; Hornung et al., 2012;

Loell et al., 2022; Metzger et al., 2015). We note that the number of mutations we mapped (3) is small

in comparison to the size of the mutational target represented by these genes in the genome (~0.05% of

the genome) and it is possible that a larger screen could have revealed such genes. It is also possible

that most mutations altering these candidate genes are lethal – TBP and many transcription factors are

essential proteins – and thus could not be recovered from our mutagenesis screen. Conversely, it is not

necessarily surprising that the classes of genes we identified here have not been previously associated

with *trans*-modulation of gene expression noise. Indeed, previous reports focused on natural variants

which, unlike the artificial random mutations studied here, had been exposed to the filter of natural

selection. The mutations we identified here would probably be counter-selected in nature since they

cause dysfunctions in cell wall repair and mitochondrial genome integrity. The classes of genes we

identified as *trans*-acting noise modulators therefore illustrate the complementarity between the

analysis of natural variants and the analysis of laboratory mutations.

In a previous study, we screened the same collection of EMS mutants to identify 69 *trans*-acting

mutations that changed the median expression of P_{TDH3}-*YFP* (Duveau et al., 2021). These mutations

were located in genes with different functions (iron metabolism, purine synthesis, transcription) from

the three genes we identified here, indicating that different cellular processes are involved in the

regulation of median expression level and expression noise. However, all of these mutations (those

impacting noise and those impacting median expression) were non-synonymous coding mutations

outside annotated regulatory sequences. Therefore, when not exposed to natural selection, random

mutations in coding sequences are more likely to alter gene expression (either median level or noise) of

another gene in *trans* than mutations in regulatory sequences. To draw more general conclusions about

the genetic modulation of expression noise variation, more mutations will need to be identified, in

different systems, cell types and organisms, and for the regulation of multiple genes.

The identity of two of the three mutations we identified (*yme2(G1234a)* and *msh1(G1262a)*)

suggests that the integrity of the mitochondrial genome is needed to limit expression noise of the P_{TDH3}

nuclear gene promoter. Previous studies linked mitochondrial variability among clonal yeast or human

cells to the variability of various cellular traits, including proliferation and transcription rates (Dhar et
al., 2019; Guantes et al., 2015; Neves et al., 2010). In these past studies, cell-to-cell variability in
mitochondrial content or activity acted as an extrinsic source of expression noise: cells inherited
different numbers of mitochondria after division, causing correlated variation in the expression of many
genes among cells (Johnston et al., 2012). The situation we observe here is different: the *yme2* and
msh1 mutations increased the variability of expression of two reporter genes of P_{TDH3} activity within the
same cell (intrinsic noise), but not their co-variation among cells (extrinsic noise). This is a very
surprising observation because it cannot be explained by the classic extrinsic model, where the
mutation would generate variable mitochondrial defects among cells that would in turn cause
differences of gene expression among these cells. Instead, our results suggest a more complex model:
the mutation amplifies stochastic variability within cells without necessarily generating diverse
mitochondrial states among cells. In this model of increased intrinsic noise, the same mitochondrial
defect in different cells can lead to different expression levels. How may this happen? It is known that
intrinsic noise can be altered by changing the frequency and magnitude of transcriptional or
translational bursts. A factor decreasing burst frequency while increasing burst size could increase
intrinsic noise without altering mean expression (Pedraza & Paulsson, 2008; Tan & van Oudenaarden,
2010; Zenklusen et al., 2008), as we observed in *yme2(G1234a)* and *msh1(G1262a)* mutants. Thus, we
can hypothesize that a signal is activated in response to mitochondrial defects and transmitted to the
molecular complexes involved in transcription or translation. A promising candidate for such a signal
would be the retrograde response. This pathway is activated in response to mitochondria dysfunction
and enables cells to compensate for the lack of mitochondrial activity via the regulation of nuclear
genes (Butow & Avadhani, 2004; Z. Liu & Butow, 2006). The retrograde response was best
characterized in yeast and was also described in other eukaryotes, including human cells, which
revealed that its mechanisms and functions are partially conserved across species (Jazwinski & Kriete,
2012). We hypothesize that *yme2(G1234a)* and *msh1(G1262a)* mutations may both activate the
retrograde response by causing mitochondrial defects, which could then lead to alteration of
transcriptional bursting at the *TDH3* promoter, generating intrinsic noise. The known binding site
(GGTCAC) of the retrograde transcription factor (Rtg1-Rtg3 heterodimer; Jia et al., 1997; X. Liao &
Butow, 1993) is absent in the *TDH3* promoter, suggesting it is not a direct target of the pathway.
However, previous findings suggest that the expression level of *CIT2* – the best-characterized target
gene of the retrograde pathway (X. S. Liao et al., 1991) – may impact *TDH3* promoter activity, thereby
an indirect effect of the retrograde response is possible. Indeed, we previously showed that mutations in

the Cyc8-Tup1 complex, a direct regulator of *CIT2* transcription (Conlan et al., 1999), altered *TDH3*

promoter activity (Duveau et al., 2021). Further work is necessary to determine whether activation of

the retrograde response can indeed increase intrinsic noise of the *TDH3* promoter, and whether

yme2(G1234a) and *msh1(G1262a)* mutations act via this mechanism. If this is the case, it will then

become interesting to investigate whether mitochondrial retrograde signaling has a similar effect on

expression noise of other genes and in other species.

What can be the consequences of an increase in expression noise? Previous studies showed that

varying the expression noise of *TDH3* or other genes could have a small but detectable impact on

fitness (Duveau et al., 2018), highlighting how genetic modifiers of noise could be subjected to natural

selection. The mutations we identified here compromise the integrity of important cellular functions

(mitochondria and cell wall) and this direct effect is likely their major - negative - impact on fitness.

The fact that they also modulate noise in gene expression could also cause marginal fitness changes,

especially if this modulation is not restricted to the expression of *TDH3* but affects expression noise of

other genes involved in other biological processes.

In conclusion, the present study provides examples of *trans*-acting mutations affecting gene

expression noise, revealing an unexpected link between mitochondrial defects and intrinsic variability

in expression of a nuclear gene. In the future, identifying more *trans*-acting mutations, and quantifying

their pleiotropic effects on expression noise at the transcriptomic scale and on fitness (as done in Vande

Zande et al., (2022) for mutations impacting median expression levels) will be essential to understand

how random mutations contribute to natural variation of expression noise.

MATERIALS AND METHODS

Biological resources

All yeast strains, oligonucleotides and plasmids used in this study are listed in Supplementary

File 1. A genealogy of yeast strains with a brief summary of their genotypes can be found in

Supplementary File 2. Laboratory stocks were traced using MyLabStocks web application (Chuffart &

Yvert, (2014); https://gitbio.ens-lyon.fr/LBMC/yvertlab/mylabstocks).

Reference strain & EMS mutants

The collection of 1241 mutant strains obtained by ethyl methanesulfonate (EMS) mutagenesis
of the reference strain YPW1139 was described in Metzger et al. (2016) and Duveau et al. (2021).
YPW1139 was itself obtained from YPW978, a *MATα* haploid strain with S288c genetic background
carrying alleles of *RME1* and *TAO3* from SK1 strain that conferred increased sporulation efficiency
(Deutschbauer & Davis, 2005) and alleles of *SAL1*, *CAT5* and *MIP1* from RM11 strain that conferred
decreased frequency of respiratory deficient (petites) cells (Dimitrov et al., 2009). YPW1139 was
obtained by introducing at the *ho* locus a reporter gene that contained the *TDH3* promoter upstream of
the *Venus YFP* coding sequence and the *CYC1* terminator along with a *KanMX4* drug resistance gene.
As shown in Supplementary Files 1 & 2, all strains constructed in this study were derived from
YPW1139.

Strain constructions

We describe below different procedures that were applied to generate yeast strains used in this study.
For each genotype, we systematically constructed and used in experiments two strains that originated
from two independent colonies obtained after the same transformation. Comparing phenotypes of these
independent transformants allowed us to distinguish true effects of a targeted genetic change from
undesired effects of potential spontaneous secondary mutations.

Reference strains with two fluorescent reporters

To be able to distinguish intrinsic and extrinsic sources of expression noise, we constructed
strains GY2458 and GY2459 by inserting a P_{TDH3}-*RFP*-T_{CYC1} construct at *ho* locus in the genome of
strain YPW1139. The resulting genotype is *ho*::P_{TDH3}-*YFP*-T_{CYC1}-*KanMX4*-T_{CYC1}-*RFP*-P_{TDH3} with the
two reporter genes in opposite orientation. We used the *mCherry* variant of *RFP*. The genomic
insertion was done using a CRISPR/Cas9 approach adapted to yeast (Laughery et al., 2015). First, we
used overlap extension PCR to generate a P_{TDH3}-*RFP* fragment with homologies to the insertion site at
5' and 3' ends. P_{TDH3} was amplified by PCR from genomic DNA of strain YPW1139 using
oligonucleotides 1Q65 and 1S07. *RFP*-T_{CYC1} was amplified by PCR from genomic DNA of strain
GY2433 using oligonucleotides 1S04 and 1S05. Equimolar amounts of both amplicons were fused by
overlap extension PCR, and 70 bp of homology to the insertion site was added by a final PCR with
oligonucleotides 1S06 and 1S08. In parallel, we generated plasmid pGY639 by cloning hybridized
oligonucleotides 1R98 and 1R99 between BclI and SwaI restriction sites of plasmid pML104 (encoding
guide RNA and Cas9). The repair fragment and pGY639 were transformed in YPW1139 using a classic
LiAc/polyethylene glycol and heatshock protocol (Gietz & Schiestl, 2007). Transformants were

selected on SD -Ura agar plate (synthetic defined medium lacking uracil). Successful transformation
was verified by PCR and sequencing of the insert. Two positive transformants were transferred to a
YPG plate to counter select petite cells and then transferred to a SD + 5-FOA plate (0.9 g/L 5-
Fluoroorotic acid) to counter select cells carrying the pGY639 plasmid. The two strains were stored at -
70°C in 15% glycerol.

Single-site mutants

Point mutations identified via genetic mapping were inserted individually in the genome of
strain GY2458. *yme2(G1234a)*, *chs1(G1752a)*, *ald4(C-344t)*, *nop19(C-575t)* and *rsn1(G86a)*
mutations were inserted using a similar CRISPR/Cas9 approach as described above and in Duveau *et*
al. (2021). For each mutation, a specific recognition site of the guide RNA was cloned in plasmid
pML104 corresponding to a sequence adjacent to a PAM motif and near the mutation site in the
genome. The repair fragment was obtained by hybridization of two complementary 90-bases
oligonucleotides containing the target mutation and flanking genomic sequences (Supplementary File
1). *msh1(G1262a)* mutation was obtained using the *delitto perfetto* approach (Stuckey et al., 2011),
because a suitable recognition site for the guide RNA could not be found near the mutation site. First,
we inserted a cassette containing *Ura3* and *hphMX4* selection markers (amplified by PCR from plasmid
pCORE-UH) at the mutation site in *msh1*. Transformants were selected consecutively on SD medium
lacking uracil and on YPD medium complemented with 300 mg/L of hygromycin B (Sigma-Aldrich
H3274-1G). Correct insertion of the *Ura3-hphMX4* cassette without mutation was verified by PCR and
Sanger sequencing. Second, the *Ura3-hphMX4* cassette inserted at *msh1* locus was replaced by a repair
fragment that contained *msh1(G1262a)* mutation. This time transformants were selected on SD
medium complemented with 0.9 mg/L of 5-FOA. Correct insertion of the mutation was validated by
Sanger sequencing. However, resulting strains were unable to grow on a non-fermentable carbone
source (petites), a phenotype frequently observed in *msh1* mutant. We restored respiratory efficiency
by crossing these strains with YPW1240 (*MAT***a** strain with same genetic background as YPW1139)
and by selecting *MATα*; *msh1(G1262a)*; *ho::P$_{TDH3}$-YFP-T$_{CYC1}$-KanMX4-T$_{CYC1}$-RFP-P$_{TDH3}$* segregating
cells after meiosis of the diploid. For each single-site mutant, two independent clones were stored at -
80C in 15% glycerol.

YFP-RFP gene fusion

We edited the genome of strains YPW1139 (*MATα*) and YPW1240 (*MAT***a**) using CRISPR/Cas9 to
fuse the *RFP* coding sequence at the 3' end of the *YFP* coding sequence with addition of a 30-bp linker
sequence between the two reporters. The repair fragment was obtained by PCR amplification of *RFP*

from GY2433 genomic DNA using oligonucleotides 1U75 and 1U76. CRISPR plasmid pGY666 was

obtained by cloning hybridized oligonucleotides 1U79 and 1U80 in pML104. Correct insertion was

verified by PCR and Sanger sequencing. We then inserted separately mutations *yme2(G1234a)* and

chs1(G1752a) in the genome of the *MATα* strain that expressed the YFP-RFP fusion protein. The

yme2(G1234a) single-site mutant was crossed to the *MAT***a** strain with *YFP-RFP* gene fusion. We then

performed tetrad dissection after meiosis of the resulting diploid, and selected segregant clones with

MATα; yme2(G1234a); *ho::P$_{TDH3}$-YFP-RFP* genotype. *chs1(G1752a)* mutation was introduced using

the CRISPR/Cas9 approach described above.

YFP-RFP *gene fusion with uTEV cleaving site*

We also constructed strains with a *P$_{TDH3}$-YFP-RFP* gene fusion at *ho* where coding sequences of

YFP and RFP were separated by a short sequence encoding a cleavage site (ENLYFQS polypeptide)

for the tobacco etch virus protease uTEV (Kapust et al., 2001). Using CRISPR/Cas9, we first inserted

the uTEV cleavage site between YFP and RFP coding sequences in the genome of strains described

above carrying a *YFP-RFP* gene fusion and with or without *yme2(G1234a)* mutation. CRISPR plasmid

pGY682 was obtained by cloning hybridized oligonucleotides 1W21 and 1W22 in pML104. The repair

fragment containing uTEV cleavage site and homology to 3' end of *YFP* and 5' end of *RFP* coding

sequences was obtained by PCR amplification of oligonucleotides 1W23 and 1W24. Correct insertion

was verified by PCR and Sanger sequencing. Second, we inserted a *P$_{TDH3}$-uTEV* construct at *LYS2* locus

in these strains using CRISPR/Cas9 gene editing. The CRISPR plasmid targeting a guide RNA at *LYS2*

was obtained by cloning hybridized oligonucleotides and in pML104. The repair fragment was

obtained by overlap extension PCR. *P$_{TDH3}$* was amplified by PCR from genomic DNA of strain

YPW1139 using oligonucleotides 1R33 and 1W26. *uTEV* coding sequence was amplified from plasmid

pRK792 (obtained from Addgene) using oligonucleotides 1W27 and 1W28. Equimolar amounts of

both amplicons were fused by overlap extension PCR, and 70 bp of homology to the insertion site was

added by a final PCR with oligonucleotides 1W29 and 1W30. Transformants were selected as

described above for reference strains. Correct insertion was verified by PCR and Sanger sequencing

using oligonucleotides 1W31, 1W32 and 1W33.

Sequencing genomes of EMS mutants

The genome of the parental strain YPW1139 was sequenced in a previous study (Duveau et al.,

2021). Here, we sequenced the genomes of EMS mutant strains YPW2088, YPW2162, YPW2143,

YPW2210 and YPW2153 to identify all point mutations that were present in each strain. After growth of each strain in 5 mL of YPD medium for 16 hours at 30°C, genomic DNA was extracted from ~2 x 10^8 cells per strain using a Wizard genomic DNA purification kit (Promega). Illumina sequencing libraries were then prepared using a DNA Library Prep kit from ActiveMotif. Samples were sequenced on a MiSeq instrument by ProfileXpert, using a MiSeq standard V2 2 x 250 bp sequencing kit. The sequencing depth was ~40x per genome (15 M paired-end reads). Sequencing data were analyzed with a Nextflow pipeline. Sequences corresponding to P5 and P7 Illumina adapters were removed using *cutadapt*, then trimmed reads were aligned to S288c reference genome using *bowtie2*. Variant calling was performed using *freebayes*, and false positive calls were filtered out based on mapping quality MQ. We also removed variants shared by strain YPW1139 and mutant strains, as well as mutations found in at least four mutant strains. All scripts use for analyzing whole-genome sequencing data can be found at: https://gitbio.ens-lyon.fr/LBMC/yvertlab/mapping-noise.

Mapping mutations altering expression noise

BSA-seq approach

We previously used a BSA-seq approach to identify mutations associated with variation of P_{TDH3}-*YFP* median expression (Duveau et al., 2021). Here, we adapted this approach for mapping mutations associated with variation of P_{TDH3}-*YFP* expression noise. First, each haploid mutant strain (resistant to G418) was crossed to the mapping strain YPW1240 (resistant to Nourseothricin) on YPG agar plates. Hybrids were selected after replication on YPD agar complemented with 60 mg/L of Nourseothricin and 200 mg/L of G418. Then,meiosis was induced by incubation for 5 days on YPA agar plates (10 g/L Bactopeptone, 10 g/L Bacto Yeast extract, 2% Kac). Random spores were isolated after incubation in 0.3 µg/L zymolyase 100T for 1 hour at 30°C as previously described (Duveau et al., 2021). Briefly, spores and vegetative cells were pelleted by centrifugation in a microcentrifuge tube, resuspend in 100 µL H_2O and then vigorously vortexed for 2 minutes. At this stage, spores adhered to tube walls, allowing elimination of vegetative cells and undigested tetrad via several washes with 1 mL of sterile water. Spores were then resuspended in 0.02% Triton-X using brief sonication. Next, 10^6 spores that did not express RFP (*MATα*) were sorted for each sample by FACS (BD FACS Aria II) at the University of Michigan Flow Cytometry Core. RFP fluorescence was quantified using a 561 nm laser for excitation and a 582/15 nm optical filter for emission. Sorted cells were cultivated in 1 mL of YPD for 24 hours at 30°C. We next sorted each sample into three bulks of 10^5 cells based on cell

fluorescence levels using FACS. YFP fluorescence was quantified using a 488 nm laser for excitation and a 530/30 nm optical filter for emission. The low fluorescence bulk corresponded to cells sorted among the 2% cells in the population with the lowest YFP fluorescence. The high fluorescence bulk corresponded to cells sorted among the 2% cells in the population with the highest YFP fluorescence. The mid fluorescence bulk corresponded to cells sorted among the 10% cells in the population with a fluorescence closest to the median YFP fluorescence. When establishing gates for cell sorting, we were particularly careful to keep the median cell size (area of forward scatter) as similar as possible among the three bulks. Sorted cells were then cultivated for 24 hours in 1.5 ml of YPD. Next, we proceeded with a second round of cell sorting based on YFP fluorescence. 10^5 cells were sorted by FACS for each fluorescence bulk. For each fluorescent bulk, we only sorted cells with YFP levels from the same portion of the population as in the previous round of FACS. After growth for 24 hours in 1.5 ml of YPD, a third round of cell sorting and growth was performed using the same procedure. Next, genomic DNA was extracted from each sample using a Gentra Puregene yeast/Bact kit (Qiagen). DNA libraries were prepared from 1 ng of genomic DNA using a Nextera XT DNA Library Prep kit (Illumina). Deep sequencing of DNA libraries was performed on a HiSeq 4000 instrument (Illumina) at the University of Michigan Sequencing Core Facility, using a 2 x 150 bp sequencing kit. Demultiplexing of sequencing reads were performed using *bcl2fastq2* v2.17. Sequencing data were next analyzed using a Nextflow pipeline on clusters of the PSMN at ENS de Lyon. The first steps of the pipeline were the same as described above for the sequencing of EMS mutants. Variant calling was performed using *freebayes*, which provided the number of reads supporting each allele at each variant site. We then used custom *R* scripts to only retain variants that were identified in the genome of the corresponding EMS mutant for each sample. Finally, likelihood ratio tests (*G*-tests) were used to determine for each variant site whether the proportion of wild-type and mutant alleles was statistically different in the mid fluorescence bulk and in the combined low and high fluorescence bulks. We reasoned that a mutation associated with an increased expression noise should be found at a lower frequency in the mid bulk than in combined data from low and high bulks. We expected to observe the opposite pattern for a mutation associated with a decreased expression noise, and no significant difference among bulks for a mutation without effect on expression noise and not linked to a causative mutation. All scripts use for analyzing BSA-seq data can be found at: https://gitbio.ens-lyon.fr/LBMC/yvertlab/mapping-noise.

Tetrad analysis approach

Mutation *msh1(G1262a)* was the best candidate to explain the increased expression noise of mutant
YPW2143 after using BSA-seq, but no significant association could be established due to unexpected
allele frequencies in one of the three segregant bulks. To statistically test the association between
expression noise and *msh1* genotype, we used linkage analysis based on tetrad dissection. First, we
induced meiosis in the diploid strain YPW2318 (obtained by crossing YPW2143 mutant with
YPW1240 mapping strain) by incubation for 5 days in YPA medium (10 g/L Bactopeptone, 10 g/L
Bacto Yeast extract, 2% KAc). After sporulation, tetrads were digested in 0.5 mg/mL zymolyase 20T
(Seikagaku) and 1M sorbitol for 10 minutes at 30°C. 32 tetrads were dissected with a platinum needle
mounted on a inverted microscope, resulting in the isolation of the 4 spores from each tetrad on YPD
agar plates with 1M sorbitol. After 48 h of incubation at 30°C, colonies were replicated with sterile
velvets on selective media to determine how three markers segregated among the 4 spores in each
tetrad: *NatMX4* was selected on Nourseothricin (60 mg/L), *KanMX4* was selected on G418 (200 mg/L)
and *HphMX* was selected on Hygromycin B (200 mg/L). In addition, the segregation of a red
fluorescent reporter inserted at the *MAT* locus (*mata2::yEmRFP-HphMX)* was characterized using a
fluorescence stereomicroscope (Nikon SMZ800N). 22 tetrads showed a 2:2 segregation of all markers
with a co-segregation of red fluorescence and Hygromycin B resistance, as expected. We then
quantified P_{TDH3}-*YFP* median expression and expression noise for the 88 segregants from these 22
tetrads using flow cytometry as described below. In parallel, we used Sanger sequencing of PCR
amplicons obtain with oligonucleotides 1T05 and 1T06 to determine the genotype at *MSH1* locus in the
88 segregants. Statistical association between *MSH1*genotype and expression noise was determined
using a linear model with expression noise treated as a continuous response variable and genotypes at
MSH1, MAT and *NatMX4* treated as categorical explanatory variables.

Quantification of median expression and expression noise using flow cytometry

Measurements of median expression and expression noise of the P_{TDH3}-*YFP* reporter gene in the
1241 EMS mutants and control strains YPW1139 and YPW978 were performed at the University of
Michigan as previously described (Duveau et al., 2021; Metzger et al., 2016), using a BD Accuri C6
flow cytometer (BD Biosciences). Subsequent experiments to measure P_{TDH3}-*YFP* expression in single-
site mutants were performed at ENS Lyon following a similar procedure to prepare samples, but using
a MACS Quant VYB flow cytometer (Miltenyi Biotec). For each experiment, yeast strains were
revived from -80°C glycerol stocks on YPG plates (20 g/L yeast extract, 10 g/L peptone, 5% glycerol
v/v and 20 g/L agar) at least 3 days and no more than 2 weeks before starting the experiment. After 2
days of incubation at 30°C, YPG plates were kept at room temperature. Then, a small amount of cells

($\sim$$10^5$ cells) was inoculated in each well of a 96-deep well plate containing 0.5 mL of filter-sterilized

YPD medium (20 g/L yeast extract, 10 g/L peptone, 20 g/L dextrose from Fisher bioreagents) and one

3-mm glass bead per well according to a randomized plate design. We observed that changing the

brand of reagents used to make YPD medium could lead to inconsistent measures of expression noise.

Culture plates were incubated for 24 h at 30°C with 250 rpm orbital shaking. Cells were maintained in

suspension both because of the shaking and the presence of a single glass bead in each well. After this

first phase of growth, 2 µL of culture from each well ($\sim$ 2 x 10^5 cells) were transferred to another plate

containing 0.5 mL of fresh YPD per well and a second growth phase was carried on by incubating new

plates at 30°C for 18 h with 250 rpm orbital shaking. For the experiment where the second culture was

initiated at different cell densities (Figure 6), we transferred 1, 2, 4 or 16 µL of culture in different

wells and we estimated cell density after growth by measuring the optical density at 660 nm of 0.2 mL

of each culture using a Sunrise (Tecan) plate reader. Then, for all experiments, 200 µL from each

sample were transferred to 96-well　plates with 1.2 µm filters (Millipore MSBVS1210). Culture

medium was aspirated with vacuum, then cells were washed twice in 200 µL of phosphate-buffer saline

(PBS) before fixation in 200 µL of 2% paraformaldehyde (PFA) for 8 minutes. After aspiration, cells

were resuspended in 200 µl of PBS containing 0.1 M Glycine for 12 minutes, and washed twice in 200

µL of PBS. A final 1/20 dilution was applied by transferring 5 µL of each sample into 195 µL of PBS

in a flat bottom 96-well plate (Falcon). Plates were stored up to 24 hours at 4°C prior to flow cytometry

acquisition.

Forward scatter and fluorescence were recorded for 10,000 events per sample with the MACS

Quant VYB. A 488-nm laser was used for excitation and a 525/50 optical filter was used for acquisition

of YFP signal, while a 561-nm laser was used for excitation and a 615/20 optical filter was used for

acquisition of RFP signal. Flow cytometry data were analyzed using custom R scripts (Supplementary

File 3) with a similar approach as in Duveau et al. (2018), except for certain points mentioned below.

First, we filtered out artifactual events and doublets (events corresponding to multiple cells) based on

values of the area and height of forward scatter (FSC.A and FSC.H) as previously described. Then, we

excluded cells that were clear outliers in each fluorescence channel using flowClust package. Raw

fluorescence values of cells are strongly correlated with forward scatter (a proxy for cell size). We

therefore computed normalized fluorescence values that were independent from cell size, as previously

described (Duveau et al., 2018). For each sample, we then computed the median cell size among cells,

as well as the median fluorescence and the coefficient of variation (CV) of fluorescence for each

fluorescence channel (YFP and RFP) as measures of expression noise. Extrinsic and intrinsic noise

were calculated using formula from Fu and Pachter (2016). First, we applied quantile normalization to YFP and RFP fluorescence values of cells (after cell size normalization) using *normalize.quantiles* function from R package *preprocessCore*. Then, intrinsic noise was calculated as:

$$intrinsic = \frac{\frac{1}{n}\left(\sum_{i=1}^{n}\left(\left(RFP_i - YFP_i\right)^2\right)\right)}{2 \cdot \overline{RFP} \cdot \overline{YFP}},$$

and extrinsic noise was calculated as:

$$extrinsic = \frac{\frac{1}{n}\left(\sum_{i=1}^{n}\left(RFP_i \cdot YFP_i\right)\right) - 2 \cdot \overline{RFP} \cdot \overline{YFP}}{2 \cdot \overline{RFP} \cdot \overline{YFP}},$$

where n is the number of cells in a sample, RFP_i and YFP_i are quantile-normalized fluorescence levels of each cell and $\overline{RFP}$ and $\overline{YFP}$ are means of quantile-normalized fluorescence levels among cells. For each sample, median cell size, median fluorescence, CV of fluorescence, intrinsic noise and extrinsic noise were also calculated for five subsets of cells defined based on five quantiles (quintiles) of FSC.A values (area of forward scatter). Statistical comparisons between genotypes or conditions were performed via pairwise *t*-tests using data from replicate samples (samples sharing the same genotype and condition). Benjamini-Hochberg procedure (FDR) was used to adjust *P* values for multiple testing.

Western blot

The protease from tobacco etch virus (TEV) cleaves the specific site ENLYFQS in proteins (Kapust et al., 2001; Parks et al., 1994), resulting into two residues when the target sequence is inserted in a protein sequence. A western blot on whole-cell protein extract was performed to determine how complete was the cleavage of YFP-ENLYFQS-RFP fusion protein by the classic TEV and improved uTEV protease both in wild-type and *yme2(G1234a)* cells. Control genotypes that expressed either no RFP, RFP alone or YFP-RFP fusion protein were also included (seven strains in total). YFP-RFP and RFP are detected using a primary antibody targeted against mCherry and distinguished based on band sizes after electrophoresis. GAPDH constitutive control were detected with anti-GAPDH antibody for all proteins extract.

Yeast strains were thawed from glycerol stocks kept at –80°C onto YPG agar plates 3 days before experiment. After 2 days of incubation at 30°C, YPG plates were placed at room temperature. The day before the experiment, a small amount of cells (~10^5 cells) was inoculated in 4 mL of YPD medium and incubated overnight at 30°C with 250 rpm orbital shaking. 1 mL of cell culture was then

transferred to 50 mL of YPD medium and incubated for another 4 hr at 30°C to reach the exponential

phase of growth. All cells were pelleted by centrifugation and resuspended in 1 mL of sterile water.

After another centrifugation at 10,000 g for 1 minute, cell pellets were resuspended on ice in 300 µL of

fresh lysis buffer (0,1 % w/v sodium deoxycholate, 1 mM EDTA pH8, 50 mM HEPES-KOH pH 7.5,

140 mM NaCl, 1 % w/v Triton W-100 and proteinase inhibitors (Sigma P8215). Reagents were mixed

by pipetting up and down. To disrupt cell walls and membranes, we added 500 µL of cold acid-washed

glass beads (Sigma G8772) and the tubes were placed on a PreCellys apparatus. Tubes were shaken at

6,800 rpm through 2 cycles, each cycle consisting of 10 s ON, 10 s OFF, 10 s ON. Tubes were placed

for 5 min on ice between the two cycles. Lysates were collected in a microtube and stored at -80°C.

Protein concentration was measured using the Bio-Rad Protein Assay (#500-0006) from optical density

at 595 nm (Bradford) based on linear regression curve made with standards. The samples were diluted

in lysis buffer to load 50 µg of proteins and denatured 5 min at 95°C.

Proteins were separated on a 12 % SDS-PAGE gel electrophoresis at 100 V for 2 hours and

then transferred to nitrocellulose membrane at 0,2 µM by semi-dry transfer. The membrane was

blocked 1 hour at room temperature with 5 % non-fat dry milk in Tris-buffered saline (TBS) containing

0,02 % Tween. The membrane was incubated with primary antibodies against mCherry in TBS with

0,02% Tween and 0,3% milk (1:1000; Invitrogen, PA5-34974) overnight at 4°C. The membrane was

washed three times during 10 minutes in TBS with 0,02% Tween and incubated with the secondary

antibodies against rabbit in TBS with 0,02% Tween and 0,3% milk (1:10000; Odyssey; 926-32210) for

1 hour at room temperature. The membrane was washed as before and the protein band was visualized

by a ChemiDoc MP imaging system (Biorad) with ECL kit (Thermo Scientific) and analyzed using

ImageJ. YFP-RFP fusion protein had an expected size of 55 kDa and the RFP residue alone had an

expected size of 26 kDa. The membrane was washed one last time and incubated with GAPDH-HRP

conjugate antibody in TBS with 0,02% Tween and 0,3% milk [1:2000; AbCam; ab85760] for 1 hour at

room temperature. The GAPDH protein bands were revealed as before, with an expected size of 36

kDa.

Figure 1. Identification of mutations affecting cell-to-cell variability of expression of a target gene. (a) A mutagenesis screen was performed to identify 5 strains (red dots) with altered expression noise and unchanged median expression of the P_{TDH3}-*YFP* reporter. These five strains were identified among a collection of 1241 strains obtained by ethyl methanesulfonate (EMS) mutagenesis (Duveau *et al.* 2021); their altered expression noise was further validated in another independent experiment (**Figure supplement 1**). Each dot represents the mean expression value among four replicate samples relative to the mean expression of the unmutagenized reference strain. Error bars show 95% confidence intervals. **(b)** Scheme of the bulked segregant analysis and whole-genome sequencing mapping strategy (BSA-seq) used to determine which mutations were statistically associated with expression noise. All mutations present in the 5 selected mutant strains were identified by whole-genome sequencing. In parallel, each mutant strains was crossed to a reference strain to

obtain a large population of segregating cells after meiosis of the resulting diploids. Segregating cells

were sorted by FACS in three bulks based on their fluorescence levels. Each bulk was further enriched

for a particular fluorescence level by two additional rounds of growth and cell sorting. Whole-genome

sequencing was then performed at ~100x coverage to quantify frequencies of segregating mutations in

the three bulks. A mutation that does not impact *YFP* expression noise is expected to be found at

similar frequencies in the three bulks, while a mutation impacting expression noise or genetically

linked to a causative mutation is expected to be found at different frequencies in the bulk of cells with

average fluorescence levels and in the two bulks of cells with extreme fluorescence levels. **(c)** Strategy

used to confirm the effects of mapped mutations on expression noise. Mapped mutation were

individually introduced in the genome of a strain expressing two different reporter genes (*RFP* and

YFP), each under control of a copy of the *TDH3* promoter. These single-site mutants allowed us to

quantify the effects of each mutation not only on expression noise of the two fluorescent reporters, but

also on intrinsic and extrinsic noise (from the covariation of fluorescence of the two reporters, see

Methods).

Figure 1 – figure supplement 1. Second fluorescence screen used to select mutant strains
with reproducible changes of expression noise for genetic mapping. Median expression and
expression noise of YFP were quantified in 254 mutant strains that showed significant changes of YFP
expression (median and/or noise) relative to the reference strain in the initial screen (**Figure 1a**). Each
dot represents the mean expression value among four replicate samples relative to the mean expression
of the unmutagenized reference strain. The five strains selected for genetic mapping are shown in red.
Error bars show 95% confidence intervals.

1000

1001

Figure 1 – figure supplement 2. Statistical comparisons of allele frequencies between
different fluorescent bulks at each mutation site in the genome of segregants from five crosses.
For each mutation site (each vertical bar) in **(a)**, **(b)**, **(d)**, and **(e)**, the *P*-value was computed using a *G*-
test (as described in Methods) to compare read counts with mutant and wild-type alleles between i) low
and high fluorescence bulks combined and ii) mid fluorescence bulk. For the cross involving mutant
YPW2143 **(c)**, we only compared high and mid fluorescence bulks because a single haplotype was

nearly fixed in the low fluorescence bulk. *P*-values were adjusted using the Benjamini-Hochberg

procedure to control the false-discovery rate. To visualize directions of effects, we represented

$-\log_{10}(P)$ values with a positive sign if a mutation was found at higher frequency in the low and high

bulks than in the medium bulk (*i.e.* mutation associated with increased expression noise) and a negative

sign if a mutation was found at lower frequency in the low and high bulks than in the medium bulk (*i.e.*

mutation associated with decreased expression noise). Vertical dotted lines show boundaries between

chromosomes. Horizontal red dotted lines show *P*-value thresholds of 0.01 for mutations with positive

and negative effects. The six mutations for which we attempted to validate the effect on expression

noise in **Figure 2** are represented in red.

1018

1019

1020

1021

1022

a

b

Analysis of variance table

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MSH1	1	4.39E-02	4.39E-02	7.92E+02	1.47E-82
TETRAD	19	1.19E-03	6.27E-05	1.13E+00	3.20E-01
PLATE	3	4.08E-03	1.36E-03	2.45E+01	4.54E-14
MAT	1	1.15E-03	1.15E-03	2.06E+01	8.31E-06
MARKER	1	1.52E-04	1.52E-04	2.74E+00	9.89E-02
Residuals	272	1.51E-02	5.55E-05		

Figure 1 – figure supplement 3. Linkage analysis showing a strong association between

expression noise and segregation of *m3* mutation (*msh1(G1262a)*) in F2 hybrids. P_{TDH3}-*YFP*

expression noise was quantified by flow cytometry and *msh1* genotype was determined by Sanger

sequencing in 80 F2 clones isolated from 20 tetrads. F1 diploid cells were obtained by crossing mutant

YPW2143 with the mapping strain YPW1240. **(a)** Each dot shows the mean expression noise measured

from 6094 to 8799 cells in 3 or 4 replicate populations for each F2 clone. Lines connect the four clones

issued from the same tetrad. **(b)** Results of an ANOVA used to determine the statistical significance of

the additive effects of the following explanatory variables on expression noise: i) *MSH1* genotype, ii)

the tetrad of origin of each clone, iii) the culture plate (each replicate was grown in a different plate),

iv) the mating type and v) the drug resistance marker segregating in F2 clones. The adjusted *R*-squared

of the linear model used in the ANOVA is 0.7489 and the associated *F*-statistic is 36.43 (*P*-value = 3.30 x 10^{-72}).

Figure 2. Testing the effects of six mapped mutations on YFP expression noise using single-site mutants. Thick lines, boxes and whiskers represent the median, interquartile range (IQR) and 1.5 IQR of expression noise measured among 7 to 26 samples, with colors corresponding to different genotypes as indicated in the legend of panel **(a)**. Dots represent the expression noise of individual samples measured from 2102 to 4707 cells. For the reference genotype, circles and triangles represent data collected for two independent clones obtained from the same transformation experiment. For genotypes carrying single mutations, data collected for independent clones from the same transformation are shown in two orange boxes. Pairwise t-tests were performed to compare the mean expression noise among replicates between pairs of genotypes, with p-values adjusted using Benjamini-Hochberg correction. For each panel, the mean expression noise is statistically different between two genotypes (adjusted p-value < 0.05) if no letter is shared above the boxes of the two genotypes. Above each plot, positions of mutations are indicated relative to the start codon of the gene with positive and

1051 negative numbers if mutations are respectively downstream (*i.e.* in coding sequence) and upstream of
1052 the start codon. Capital and lowercase letters indicate respectively reference and alternate nucleotides at
1053 each mutation site.

1054

Figure 2 – figure supplement 1. Testing the effects of six mapped mutations on YFP
median expression using single-site mutants. Thick lines, boxes and whiskers represent the median,
interquartile range (IQR) and 1.5 IQR of median expression measured among 7 to 26 samples, with
colors corresponding to different genotypes as indicated in the legend of panel **(c)**. Dots represent the
median expression of individual samples measured from 2102 to 4707 cells. For the reference
genotype, circles and triangles represent data collected for two independent clones obtained from the
same transformation experiment. For genotypes carrying single mutations, data collected for
independent clones from the same transformation are shown in two orange boxes. Pairwise *t*-tests were
performed to compare the mean of median expression levels among replicates between pairs of
genotypes, with *p*-values adjusted using Benjamini-Hochberg correction. For each panel, the median
expression is statistically different between two genotypes (adjusted *p*-value < 0.05) if no letter is
shared above the boxes of the two genotypes. Above each plot, positions of mutations are indicated

 relative to the start codon of the gene with positive and negative numbers if mutations are respectively

1069 downstream (*i.e.* in coding sequence) and upstream of the start codon. Capital and lowercase letters

1070 indicate respectively reference and alternate nucleotides at each mutation site.

1071

1072

Figure 2 – figure supplement 2. Testing the effects of six mapped mutations on RFP

median expression and expression noise using single-site mutants. Thick lines, boxes and whiskers

represent the median, interquartile range (IQR) and 1.5 IQR of median expression **(a-f)** or expression

noise **(g-l)** measured among 7 to 26 samples, with colors corresponding to different genotypes as

indicated in the legend of panel **(c)**. Dots represent the median expression or expression noise of

individual samples measured from 2102 to 4707 cells. For the reference genotype, circles and triangles

represent data collected for two independent clones obtained from the same transformation experiment.

For genotypes carrying single mutations, data collected for independent clones from the same

transformation are shown in two orange boxes. Pairwise *t*-tests were performed to compare the mean of

median expression levels (or expression noise) among replicates between pairs of genotypes, with *p*-

values adjusted using Benjamini-Hochberg correction. For each panel, the median expression (or

expression noise) is statistically different between two genotypes (adjusted *p*-value < 0.01) if no letter is shared above the boxes of the two genotypes. Above each plot, positions of mutations are indicated relative to the start codon of the gene with positive and negative numbers if mutations are respectively downstream (*i.e.* in coding sequence) and upstream of the start codon. Capital and lowercase letters indicate respectively reference and alternate nucleotides at each mutation site.

Figure 2 – figure supplement 3. Combined effects of *rsn1(G86a)* and *nop19(C-575t)* mutations on YFP and RFP median expression and expression noise. Thick lines, boxes and whiskers represent the median, interquartile range (IQR) and 1.5 IQR of **(a)** YFP median expression,

(b) RFP median expression, **(c)** YFP expression noise and **(d)** RFP expression noise measured among 6 to 14 samples. Colors corresponding to different genotypes as indicated in the legend of panel **(a)**. Dots represent expression values of individual samples measured from 4226 to 4551 cells. For the reference genotype, circles and triangles represent data collected for two independent clones obtained from the same transformation experiment. For genotypes carrying single mutations, data collected for independent clones from the same transformation are shown in two orange boxes. Pairwise *t*-tests were performed to compare the mean of median expression levels among replicates between pairs of genotypes, with *p*-values adjusted using Benjamini-Hochberg correction. For each panel, the median expression is statistically different between two genotypes (adjusted *p*-value < 0.05) if no letter is shared above the boxes of the two genotypes. Above each plot, positions of mutations are indicated relative to the start codon of the gene with positive and negative numbers if mutations are respectively downstream (*i.e.* in coding sequence) and upstream of the start codon. Capital and lowercase letters indicate respectively reference and alternate nucleotides at each mutation site.

Figure 3. Quantifying the effects of a non-synonymous mutation in *yme2* on intrinsic and

extrinsic noise using a dual reporter system. (a) A construct consisting of P_{TDH3}-*RFP* and P_{TDH3}-*YFP*

reporter genes transcribed from opposite strands and separated by a *KanMX4* drug resistance cassette

was inserted at the *ho* locus in genetic backgrounds with wild-type or mutant (*G1234a*) alleles of *yme2*.

By quantifying both YFP and RFP fluorescence of individual cells using flow cytometry, it is possible

to distinguish the effects of *yme2* mutation on intrinsic and extrinsic noise of P_{TDH3} activity. **(b)**

Intrinsic, **(c)** extrinsic and **(d)** total noise of P_{TDH3} activity in strains carrying the dual reporter system

with wild-type (light and dark gray bars) or mutant (light and dark orange bars) alleles of *yme2*. Bars

show mean values among 7 to 26 replicate populations. Error bars are 95% confidence intervals of

means. Dots show values of individual samples obtained from 4312 to 4689 cells. **(e)** A construct

consisting of fused *YFP* and *RFP* coding sequences expressed under control of a single P_{TDH3} copy was

inserted at *ho* locus in genetic backgrounds with wild-type or mutant (*G1234a*) alleles of *yme2*. This

system was used to test whether the change in intrinsic noise detected in **(b)** using the dual reporter

system was a technical artifact or not. Indeed, the mutation should not have detectable impact on

intrinsic noise when the two fluorescent proteins are fused, because their expression is forced to

covary. **(f)** Detected intrinsic noise, **(g)** detected extrinsic noise and **(h)** detected total noise in strains

expressing YFP-RFP fusion proteins and with wild-type (light and dark gray bars) or mutant (light and

dark orange bars) alleles of *yme2*. Bars show mean values among 8 to 17 replicate populations. Error

bars are 95% confidence intervals of means. Dots show values of individual samples obtained from

4424 to 4714 cells. In **(d,h)**, light and dark bars indicate respectively the contributions of intrinsic and

extrinsic noise to total noise. **(b-d,f-h)** For the reference genotype, circles and triangles represent data

collected for two independent clones obtained from the same transformation experiment. For *yme2*

mutants, data collected for independent clones from the same transformation are shown in two different

colored bars. Pairwise *t*-tests were performed to compare mean values between bars: ns $P \geq 0.05$, *

$0.05 > P \geq 0.01$, ** $0.001 > P \geq 0.001$, *** $P < 0.001$.

1134

1135

Reference
strain

yme2(G1234a)
clone1

Figure 3 – figure supplement 1. YFP and RFP expression levels of individual cells from

wild-type and *yme2(G1234a)* strains expressing P_{TDH3}-*YFP* and P_{TDH3}-*RFP* reporter genes.

Expression of the two fluorescent reporters in **(a-c)** three replicate populations of strain GY2458

carrying a wild-type allele of *YME2* and **(d-f)** three replicate populations of strain GY2634 carrying

mutation *G1234a* in *YME2* coding sequence (4312 to 4689 cells per sample). Histograms show the

distribution of data on each axis. Correlations between YFP and RFP expression levels are visually

lower for *yme2(G1234a)* cells than wild-type cells, consistent with the increased intrinsic noise

quantified from mutant cells.

1146

1148

Figure 4 – *yme2(G1234a)* mutation increases intrinsic noise via altered gene expression

and not via altered protein partitioning at cell division. (a) Median size of wild-type and

yme2(G1234a) mutant cells quantified from forward scatter area (FSC.A) in flow cytometry data.

Thick lines, boxes and whiskers represent the median, interquartile range (IQR) and 1.5 IQR of median

cell size measured among 7 to 26 samples. Dots represent the median cell size of individual samples

measured from 4312 to 4689 cells. For the reference genotype, circles and triangles represent data

collected for two independent clones obtained from the same transformation experiment. For genotypes

carrying single mutations, data collected for independent clones from the same transformation are

shown in two green boxes. Pairwise *t*-tests were performed to compare median cell sizes between

boxes: ns $P \geq 0.05$, * $0.05 > P \geq 0.01$, ** $0.001 > P \geq 0.001$, *** $P < 0.001$. **(b)** Intrinsic noise and **(c)**

extrinsic noise of five groups of cells with different cell sizes for strains with wild-type (gray) and

G1234a (orange) alleles of *yme2*. Cell groups correspond to five fractions of the population of cells

delimited by 5-quantiles of cell sizes (same number of cells in each group but different cell sizes). Dots

indicate the median cell size (x-axis) and intrinsic **(b)** or extrinsic **(c)** noise (y-axis) of each cell group

(906 cells per group in 24 replicate samples for wild-type and 909 cells per group in 18 replicate

samples for *yme2* mutant). Shaded areas represent 95% confidence intervals. **(d)** Schemes representing

i) two alternative hypotheses to explain how *yme2(G1234a)* mutation can lead to higher intrinsic noise

detection and ii) a strategy to test these hypotheses. In hypothesis 1, higher intrinsic noise is explained

by higher variability of expression of the two fluorescent reporters (yellow and red hexagons), either

via altered regulation of transcription or translation. In hypothesis 2, higher detection of intrinsic noise

is not caused by altered regulation of expression, but instead by altered partitioning of fluorescent

proteins at cell division. If fewer proteins are inherited by daughter cells (because of smaller bud size or

other mechanisms), this could lead to higher variability of colors detected by fluorescence in daughter

cells and therefore higher intrinsic noise detected. A strategy to test these hypotheses is to express a

YFP-RFP fusion protein that can be cleaved by a protease in two separate YFP and RFP fluorescent

proteins after translation. With this system, we expect *yme2* mutation to have no detectable impact on

intrinsic noise if it acts at the expression level (hypothesis 1) because the expression of the two

reporters is forced to covary. However, we expect to detect increased intrinsic noise if *yme2* mutation

acts on protein partitioning (hypothesis 2) because cleaved proteins can segregate independently at cell

division. **(e)** Western blot used to detect the presence of YFP-RFP fusion protein, RFP single protein

and GAPDH constitutive control in seven genotypes (as indicated below the plot). YFP-RFP and RFP

are detected using an antibody against mCherry and distinguished based on the size of bands after

electrophoresis. GAPDH is detected with anti-GPADH antibody. The last two lanes on the right show

complete cleavage of YFP-ENLYFQS-RFP fusion protein by uTEV3 protease both in wild-type and

yme2(G1234a) cells. ENLYFQS is the peptide sequence targeted by the protease. **(f)** Detected intrinsic

noise and **(g)** extrinsic noise in strains expressing YFP-ENLYFQS-RFP fusion protein cleaved by

uTEV3 and with wild-type (light and dark gray bars) or mutant (light and dark orange bars) alleles of

yme2. Bars show mean values among 9 to 16 replicate populations. Error bars are 95% confidence

intervals of means. Dots show values of individual samples obtained from 4061 to 4673 cells. For the

reference genotype, circles and triangles represent data collected for two independent clones obtained

from the same transformation experiment. For *yme2(G1234a)* strains, data collected for independent

clones from the same transformation are shown in two different colored bars. Pairwise *t*-tests were

performed to compare mean values between bars: ns $P \geq 0.05$, * $0.05 > P \geq 0.01$, ** $0.001 > P \geq 0.001$,

*** $P < 0.001$. **(h)** Intrinsic noise of five groups with different cell sizes for strains expressing YFP-

ENLYFQS-RFP fusion protein cleaved by uTEV3 with wild-type (gray) and mutant *G1234a* (orange)

alleles of *yme2*. Cell groups correspond to five fractions of the population of cells delimited by 5-

quantiles of cell sizes. Dots indicate the median cell size (x-axis) and intrinsic noise (y-axis) of each

cell group (902 cells per group in 17 replicate samples per group for wild-type and 906 cells per group

in 18 replicate samples per group for *yme2(G1234a)*). Shaded areas represent 95% confidence intervals

of intrinsic noise.

1199

Figure 5 – *chs1(G1752a)* **mutation increases extrinsic noise exclusively in small cells in absence of glucosamine precursor for chitin synthesis. (a)** Intrinsic, **(b)** extrinsic and **(c)** total noise of P_{TDH3} activity in strains carrying the dual reporter system (P_{TDH3}-*YFP* and P_{TDH3}-*RFP*) with wild-type (light and dark gray bars) or mutant (light and dark orange bars) alleles of *chs1*. Bars show mean values among 7 to 16 replicate populations. Error bars are 95% confidence intervals of means. Dots show values of individual samples obtained from 3684 to 4657 cells. **(d)** Cumulative distribution of cell sizes for wild-type (gray lines) and *chs1(G1752a)* cells (orange lines) expressing the dual reporter system,

shown for individual samples (thin lines) or for all samples sharing the same genotype (thick lines).

Cell sizes were quantified by flow cytometry (FSC.A: area of forward scatter). The number of samples

and cells analyzed is the same as in **(a-c)** as data are from the same experiment. **(e)** YFP median

expression and **(f)** extrinsic noise of five groups of cells with different cells sizes for strains with wild-

type (gray) and *G1752a* (orange) alleles of *chs1*. Cell groups correspond to five fractions of the

population of cells delimited by 5-quantiles of cell sizes (same number of cells in each group but

different cell sizes). Dots indicate the median cell size (x-axis) and YFP median expression **(e)** or

extrinsic **(c)** noise (y-axis) of each cell group (1790 cells per group in 16 replicate samples for wild-

type and 1603 cells per group in 15 replicate samples for *chs1* mutant). **(g)** Extrinsic noise of cells of

different sizes measured after growth in media with different concentrations of glucosamine (0, 1, 5 or

15 mM) for strains with wild-type (gray) and *G1752a* (orange) alleles of *chs1*. Cells were grouped

based on median cell size as explained for **(e-f)**. Dots indicate the median cell size (x-axis) and

extrinsic noise (y-axis) for 1660 to 1878 cells per group measured in 5 to 16 replicates. **(e-g)** Shaded

areas represent 95% confidence intervals.

1222

Figure 5 – figure supplement 1. YFP and RFP expression levels of individual cells from
wild-type and *chs1(G1752a)* strains expressing P_{TDH3}-YFP and P_{TDH3}-RFP reporter genes.
Expression of the two fluorescent reporters in **(a-c)** three replicate populations of strain GY2458
carrying a wild-type allele of *CHS1* and **(d-f)** three replicate populations of strain GY2534 carrying
mutation *G1752a* in *CHS1* coding sequence (3917 to 4627 cells per sample). Histograms show the
distribution of data on each axis. The variability and covariation of YFP and RFP expression levels are
both visually higher for *chs1(G1752a)* cells than for wild-type cells, consistent with the increased
extrinsic noise quantified from mutant cells.

1232

Figure 5 – figure supplement 2. Fusion of YFP and RFP fluorescent proteins does not
change the detected effect of *chs1(G1752a)* mutation on extrinsic noise. (a) Detected intrinsic noise,
(b) detected extrinsic noise and **(c)** detected total noise in strains expressing YFP-RFP fusion proteins
and with wild-type (light and dark gray bars) or mutant (light and dark orange bars) alleles of *chs1*.
Bars show mean values among 8 to 16 replicate populations. Error bars are 95% confidence intervals of
means. Dots show values of individual samples obtained from 4196 to 4899 cells. Pairwise *t*-tests were
performed to compare mean values between bars: ns $P \geq 0.05$, * $0.05 > P \geq 0.01$, ** $0.001 > P \geq 0.001$,
*** $P < 0.001$.

1242

1243　　　　　**Figure 5 – figure supplement 3. Impact of glucosamine concentration on (a) median cell**
1244　　**size, (b) intrinsic noise, (c) extrinsic noise and (d) total noise.** These traits were quantified in wild-
1245　　type (gray dots) and *chs1(G1752a)* (orange dots) strains expressing P_{TDH3}-*YFP* and P_{TDH3}-*RFP* reporter
1246　　genes. Error bars show 95% confidence intervals of means measured among 5 to 16 replicate samples.
1247

Figure 5 – figure supplement 4. Growth of wild-type cell populations exposed to different

concentrations of glucosamine. Densities of cell cultures were quantified at different time points

(dots) over two cycles of growth using a spectrophotometer. At $t = 0$, cells were transferred to different

media containing the indicated concentrations of glucosamine (colors), then grown for 18.5 hours,

diluted to fresh media containing the same concentrations of glucosamine and grown again for 30

hours.

1255

Figure 6. The effect of *msh1(G1262a)* mutation on expression noise depends on cell density. Expression noise was quantified either from YFP fluorescence **(a)**, RFP fluorescence **(b)** or both **(c)** in strains carrying the dual reporter system (P_{TDH3}-*YFP* and P_{TDH3}-*RFP*) with wild-type (circles) or mutant (triangles) alleles of *msh1*. **(d)** Intrinsic noise plotted against the proportion of respiratory deficient cells (petites ρ-) for a strain with wild-type allele of *msh1* (circle), a strain derived from a ρ+ cell with *msh1(G1262a)* mutation (triangle) and a strain derived from a ρ- cell with *msh1(G1262a)* mutation (diamond). The proportion of ρ- cell was quantified for each strain as described in Materials and methods. **(a-d)** Dots indicate mean values among 4 replicates and error bars are 95% confidence intervals of the mean. Colors indicate the average density of cell cultures measured after growth and just before fixing cells to quantify their fluorescence levels, with color legend shown on top left for **(a-c)** and at the bottom for **(d)**.

Reference
strain

msh1(G1262a)

1270

Figure 6 – figure supplement 1. YFP and RFP expression levels of individual cells from

wild-type and *msh1(G1262a)* strains expressing P_{TDH3}-*YFP* and P_{TDH3}-*RFP* reporter genes.

Expression of the two fluorescent reporters in **(a-c)** three replicate populations of strain GY2458

carrying a wild-type allele of *MSH1* and **(d-f)** three replicate populations of strain GY2534 carrying

mutation *G1262a* in *MSH1* coding sequence (7885 to 8394 cells per sample). Histograms show the

distribution of data on each axis.

1277

1279

Figure 6 – figure supplement 2. Effect of *msh1(G1262a)* mutation on 8 quantitative traits

depending on cell culture dilutions. (a) Median cell size, **(b)** cell density, **(c)** YFP median expression,

(d) RFP median expression, **(e)** YFP expression noise, **(f)** RFP expression noise, **(g)** intrinsic noise and

(h) extrinsic noise were quantified in strains with the dual reporter system carrying wild-type (gray) or

G1262a (orange) alleles of *msh1*. **(a-h)** Thick lines, boxes and whiskers represent the median,

interquartile range (IQR) and 1.5 IQR of traits indicated on y-axis measured among 4 replicate

samples. Dots represent mean values for individual samples measured from 7203 to 9218 cells. Starting

from a culture at saturation (~10^8 cells/ml), cells were diluted to 500 µl of fresh SD medium at 4

different dilution ratios as indicated on x-axis and incubated for another 18 hours at 30°C before

fixation and phenotyping by flow cytometry. As intended, the different dilution ratios resulted in

different cell densities after incubation as shown in **(b)**. Pairwise *t*-tests were performed to compare

mean phenotypic values among replicates between wild-type and *msh1(G1262a)* strains for each

dilution ratio, with p-values adjusted using Benjamini-Hochberg correction: ns $P \geq 0.05$, * $0.05 > P \geq$

0.01, ** $0.001 > P \geq 0.001$, *** $P < 0.001$.

1294

Figure 6 – figure supplement 3. Cell density changes the effect of *msh1(G1262a)* on

expression noise independently from an increased proportion of respiratory deficient cells. (a)

Extrinsic noise, **(b)** YFP expression noise and **(c)** RFP expression noise plotted against the proportion

of respiratory deficient cells (petites ρ-) for a strain with wild-type allele of *msh1* (circle), a strain

derived from a ρ+ cell with *msh1(G1262a)* mutation (triangle) and a strain derived from a ρ- cell with

msh1(G1262a) mutation (diamond). The proportion of ρ- cell was quantified for each strain as

described in Materials and methods. Dots indicate mean values among 4 replicates and error bars are

95% confidence intervals of the mean. Colors indicate the average density of cell cultures measured

after growth and just before fixing cells to quantify their fluorescence levels, with color legend shown

on top left.

TABLES

Strain name	Type	YFP median expr. (relative) Assay 1	YFP noise (relative) Assay 1	YFP noise (absolute) Assay 1	YFP median expr. (relative) Assay 2	YFP noise (relative) Assay 2	YFP noise (absolute) Assay 2
YPW1139	Reference	1	1	0.0152	1	1	0.0165
YPW2088	EMS mutant	0.994	1.329	0.0203	0.996	1.255	0.0208
YPW2162	EMS mutant	0.995	1.123	0.0172	0.994	1.140	0.0188
YPW2143	EMS mutant	0.991	1.728	0.0261	0.989	1.904	0.0309
YPW2210	EMS mutant	0.990	0.792	0.0121	1.008	0.870	0.0146
YPW2153	EMS mutant	0.997	0.791	0.0120	1.013	0.824	0.0134

1307   **Table 1. Properties of the reference strain and of the 5 mutants with altered P_{TDH3}-YFP**
1308   **expression noise included in the genetic mapping experiments.**

1309

Mutation	Chr.	Position (nt)	Nucleotide change	EMS mutant	Mutant noise	Frequency low bulk	Frequency middle bulk	Frequency high bulk	P-value
m1	XIII	871,392	C→T	YPW2088	Increase	0.66	0.24	0.36	4 x 10^{-9}
m2	XIV	278,253	G→A	YPW2162	Increase	0.04	0.01	0.01	0.41
m3	VIII	350,835	G→A	YPW2143	Increase	NA	0.11	0.03	NA
m4	XV	1,039,495	C→T	YPW2210	Decrease	0.37	0.61	0.50	0.007
m5	VII	995,063	C→T	YPW2153	Decrease	0.37	0.56	0.38	0.002
m6	XIII	798,603	G→A	YPW2153	Decrease	0.44	0.57	0.39	0.004

1310   **Table 2. Candidate mutations showing statistical association with variation of P_{TDH3}-YFP**
1311   **expression noise or low frequency in all fluorescence bulks.**

Mutation	Gene	Function	Position in protein	Amino acid change
m1	YME2	Integral inner mitochondrial membrane protein; role in maintaining mitochondrial nucleoid structure and number; mutants exhibit an increased rate of mitochondrial DNA escape; shows some sequence similarity to exonucleases	412	Asp→Asn
m2	CHS1	Chitin synthase I; requires activation from zymogenic form in order to catalyze the transfer of N-acetylglucosamine (GlcNAc) to chitin; required for repairing the chitin septum during cytokinesis; transcription activated by mating factor	584	Trp→Stop
m3	MSH1	DNA-binding protein of the mitochondria; involved in repair of mitochondrial DNA; has ATPase activity and binds to DNA mismatches; has homology to E. coli MutS; transcription is induced during meiosis	428	Arg→His

1312   **Table 3. Mutations with confirmed effect on P_{TDH3}-YFP expression noise.**

AUTHORS CONTRIBUTION

FD and PJW conceived the study; FD and LM supervised the work; NM, FD, QB, AD and HDB performed experiments; FD, JKS and QB analyzed data; FD, GY and PJW interpreted results; FD wrote the manuscript; GY reviewed and edited the manuscript. These contributions may change prior to publication.

ACKNOWLEDGEMENTS

We thank Gérard Triqueneaux for sharing protocols, Taslima Haque for comments on the manuscript, the SFR Biosciences Gerland-Lyon Sud (UAR3444/US8) and University of Michigan Flow Cytometry Core for access to flow cytometers and technical assistance, the University of Michigan Sequencing Facility and ProfileXpert for Illumina sequencing, the Pôle Scientifique de Modélisation Numérique for computing resources. We also thank developers of R, Bioconductor, Ubuntu and Git.

FUNDING

Agence Nationale de la Recherche – JCJC VORTEX (ANR-20-CE12-0022)
- Fabien Duveau

Fonds de Recherche ENS Lyon – Projet Emergent 2020-2022 (R02/B-R02-FR-S60)
- Fabien Duveau

Horizon Europe, European Research Council Consolidator eGRIDE (101126053_ERC-2023-COG)
- Fabien Duveau

National Institutes of Health (R01GM108826)
- Patricia Wittkopp

National Institutes of Health (R35GM118073)
- Patricia Wittkopp

National Science Foundation (MCB-1929737)
- Patricia Wittkopp

European Molecular Biology Organization – Long-Term Fellowship (1114-2012)
- Fabien Duveau

1344   The funders had no role in study design, data collection and interpretation, or the decision to
1345   submit the work for publication.

1346

DATA AVAILABILITY

Illumina sequencing data have been deposited on the European Nucleotide Archive of the EBI
under Study PRJEB104107 (https://www.ebi.ac.uk/ena/browser/home). This includes data for genome
sequencing of the five EMS mutant strains described in the study, as well as data for deep genome
sequencing of three bulks of segregants sorted based on fluorescence (BSA-seq data) for each of the
five mutants crossed to the mapping strain YPW1240.

1353

1354

1355

1356   REFERENCES

Acar, M., Mettetal, J. T., & van Oudenaarden, A. (2008). Stochastic switching as a survival strategy in fluctuating environments. *Nature Genetics*, *40*(4), 471–475. https://doi.org/10.1038/ng.110

Ansel, J., Bottin, H., Rodriguez-Beltran, C., Damon, C., Nagarajan, M., Fehrmann, S., François, J., & Yvert, G. (2008). Cell-to-cell stochastic variation in gene expression is a complex genetic trait. *PLoS Genetics*, *4*(4), e1000049. https://doi.org/10.1371/journal.pgen.1000049

Balázsi, G., van Oudenaarden, A., & Collins, J. J. (2011). Cellular decision making and biological noise: From microbes to mammals. *Cell*, *144*(6), 910–925. https://doi.org/10.1016/j.cell.2011.01.030

Baptista, I. S. C., Dash, S., Arsh, A. M., Kandavalli, V., Scandolo, C. M., Sanders, B. C., & Ribeiro, A. S. (2025). Bimodality in *E. coli* gene expression: Sources and robustness to genome-wide stresses. *PLOS Computational Biology*, *21*(2), e1012817. https://doi.org/10.1371/journal.pcbi.1012817

Bar-Even, A., Paulsson, J., Maheshri, N., Carmi, M., O'Shea, E., Pilpel, Y., & Barkai, N. (2006). Noise in protein expression scales with natural protein abundance. *Nature Genetics*, *38*(6), 636–643. https://doi.org/10.1038/ng1807

Barroso, G. V., Puzovic, N., & Dutheil, J. Y. (2018). The Evolution of Gene-Specific Transcriptional Noise is driven by selection at the pathway level. *Genetics*, *208*(1), 173–189. https://doi.org/10.1534/genetics.117.300467

Blake, W. J., Balázsi, G., Kohanski, M. A., Isaacs, F. J., Murphy, K. F., Kuang, Y., Cantor, C. R., Walt, D. R., & Collins, J. J. (2006). Phenotypic consequences of promoter-mediated transcriptional noise. *Molecular Cell*, *24*(6), 853–865. https://doi.org/10.1016/j.molcel.2006.11.003

Bódi, Z., Farkas, Z., Nevozhay, D., Kalapis, D., Lázár, V., Csörgő, B., Nyerges, Á., Szamecz, B., Fekete, G., Papp, B., Araújo, H., Oliveira, J. L., Moura, G., Santos, M. A. S., Jr, T. S., Balázsi, G., & Pál, C. (2017). Phenotypic heterogeneity promotes adaptive evolution. *PLOS Biology*, *15*(5), e2000644. https://doi.org/10.1371/journal.pbio.2000644

Butow, R. A., & Avadhani, N. G. (2004). Mitochondrial Signaling: The retrograde response. *Molecular Cell*, *14*(1), 1–15. https://doi.org/10.1016/S1097-2765(04)00179-0

Cabib, E., Sburlati, A., Bowers, B., & Silverman, S. J. (1989). Chitin synthase 1, an auxiliary enzyme for chitin synthesis in *Saccharomyces cerevisiae*. *The Journal of Cell Biology*, *108*(5), 1665–1672. https://doi.org/10.1083/jcb.108.5.1665

Cabib, E., Silverman, S. J., & Shaw, J. A. (1992). Chitinase and chitin synthase 1: Counterbalancing activities in cell separation of *Saccharomyces cerevisiae*. *Journal of General Microbiology*, *138*(1), 97–102. https://doi.org/10.1099/00221287-138-1-97

Carey, L. B., van Dijk, D., Sloot, P. M. A., Kaandorp, J. A., & Segal, E. (2013). Promoter sequence determines the relationship between expression level and noise. *PLoS Biology*, *11*(4), e1001528. https://doi.org/10.1371/journal.pbio.1001528

Chen, D.-D., Wang, Z.-B., Wang, L.-X., Zhao, P., Yun, C.-H., & Bai, L. (2023). Structure, catalysis, chitin transport, and selective inhibition of chitin synthase. *Nature Communications*, *14*(1), 4776. https://doi.org/10.1038/s41467-023-40479-4

Choi, J. K., & Kim, Y.-J. (2009). Intrinsic variability of gene expression encoded in nucleosome positioning sequences. *Nature Genetics*, *41*(4), 498–503. https://doi.org/10.1038/ng.319

Chuffart, F., Richard, M., Jost, D., Burny, C., Duplus-Bottin, H., Ohya, Y., & Yvert, G. (2016). Exploiting single-cell quantitative data to map genetic variants having probabilistic effects. *PLoS Genetics*, *12*(8), e1006213. https://doi.org/10.1371/journal.pgen.1006213

Chuffart, F., & Yvert, G. (2014). MyLabStocks: A web-application to manage molecular biology materials. *Yeast (Chichester, England)*, *31*(5), 179–184. https://doi.org/10.1002/yea.3008

Conlan, R. S., Gounalaki, N., Hatzis, P., & Tzamarias, D. (1999). The Tup1-Cyc8 protein complex can shift from a transcriptional co-repressor to a transcriptional co-activator. *The Journal of Biological Chemistry*, *274*(1), 205–210. https://doi.org/10.1074/jbc.274.1.205

Coomer, M. A., Ham, L., & Stumpf, M. P. H. (2022). Noise distorts the epigenetic landscape and shapes cell-fate decisions. *Cell Systems*, *13*(1), 83-102.e6. https://doi.org/10.1016/j.cels.2021.09.002

Deutschbauer, A. M., & Davis, R. W. (2005). Quantitative trait loci mapped to single-nucleotide resolution in yeast. *Nature Genetics*, *37*(12), Article 12. https://doi.org/10.1038/ng1674

Dhar, R., Missarova, A. M., Lehner, B., & Carey, L. B. (2019). Single cell functional genomics reveals the importance of mitochondria in cell-to-cell phenotypic variation. *eLife*, *8*, e38904. https://doi.org/10.7554/eLife.38904

Dimitrov, L. N., Brem, R. B., Kruglyak, L., & Gottschling, D. E. (2009). Polymorphisms in multiple genes contribute to the spontaneous mitochondrial genome instability of *Saccharomyces cerevisiae* S288C Strains. *Genetics*, *183*(1), 365–383. https://doi.org/10.1534/genetics.109.104497

Duveau, F., Hodgins-Davis, A., Metzger, B. P., Yang, B., Tryban, S., Walker, E. A., Lybrook, T., & Wittkopp, P. J. (2018). Fitness effects of altering gene expression noise in *Saccharomyces cerevisiae*. *eLife*, *7*, e37272. https://doi.org/10.7554/eLife.37272

Duveau, F., Metzger, B. P. H., Gruber, J. D., Mack, K., Sood, N., Brooks, T. E., & Wittkopp, P. J. (2014). Mapping small effect mutations in *Saccharomyces cerevisiae*: Impacts of experimental design and mutational properties. *G3 (Bethesda, Md.)*, *4*(7), 1205–1216. https://doi.org/10.1534/g3.114.011783

Duveau, F., Vande Zande, P., Metzger, B. P., Diaz, C. J., Walker, E. A., Tryban, S., Siddiq, M. A., Yang, B., & Wittkopp, P. J. (2021). Mutational sources of *trans*-regulatory variation affecting gene expression in *Saccharomyces cerevisiae*. *eLife*, *10*, e67806. https://doi.org/10.7554/eLife.67806

Ehrenreich, I. M., Torabi, N., Jia, Y., Kent, J., Martis, S., Shapiro, J. A., Gresham, D., Caudy, A. A., & Kruglyak, L. (2010). Dissection of genetically complex traits with extremely large pools of yeast segregants. *Nature*, *464*(7291), 1039–1042. https://doi.org/10.1038/nature08923

Eldar, A., & Elowitz, M. B. (2010). Functional roles for noise in genetic circuits. *Nature*, *467*(7312), 167–173. https://doi.org/10.1038/nature09326

Elowitz, M. B., Levine, A. J., Siggia, E. D., & Swain, P. S. (2002). Stochastic gene expression in a single cell. *Science (New York, N.Y.)*, *297*(5584), 1183–1186. https://doi.org/10.1126/science.1070919

Fehrmann, S., Bottin-Duplus, H., Leonidou, A., Mollereau, E., Barthelaix, A., Wei, W., Steinmetz, L. M., & Yvert, G. (2013). Natural sequence variants of yeast environmental sensors confer cell-to-cell expression variability. *Molecular Systems Biology*, *9*(1). https://doi.org/10.1038/msb.2013.53

Fu, A. Q., & Pachter, L. (2016). Estimating intrinsic and extrinsic noise from single-cell gene expression measurements. *Statistical Applications in Genetics and Molecular Biology*, *15*(6), 447–471. https://doi.org/10.1515/sagmb-2016-0002

Gamba, P., Jonker, M. J., & Hamoen, L. W. (2015). A novel feedback loop that controls bimodal expression of genetic competence. *PLoS Genetics*, *11*(6), e1005047. https://doi.org/10.1371/journal.pgen.1005047

Gietz, R. D., & Schiestl, R. H. (2007). High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nature Protocols*, *2*(1), Article 1. https://doi.org/10.1038/nprot.2007.13

Guantes, R., Rastrojo, A., Neves, R., Lima, A., Aguado, B., & Iborra, F. J. (2015). Global variability in gene expression and alternative splicing is modulated by mitochondrial content. *Genome Research*, *25*(5), 633–644. https://doi.org/10.1101/gr.178426.114

Hanekamp, T., & Thorsness, P. E. (1996). Inactivation of YME2/RNA12, which encodes an integral inner mitochondrial membrane protein, causes increased escape of DNA from mitochondria to the nucleus in *Saccharomyces cerevisiae*. *Molecular and Cellular Biology*, *16*(6), 2764–2771. https://doi.org/10.1128/MCB.16.6.2764

Harper, C. V., Finkenstädt, B., Woodcock, D. J., Friedrichsen, S., Semprini, S., Ashall, L., Spiller, D. G., Mullins, J. J., Rand, D. A., Davis, J. R. E., & White, M. R. H. (2011). Dynamic analysis of stochastic transcription cycles. *PLOS Biology*, *9*(4), e1000607. https://doi.org/10.1371/journal.pbio.1000607

Hill, M. S., Vande Zande, P., & Wittkopp, P. J. (2021). Molecular and evolutionary processes generating variation in gene expression. *Nature Reviews. Genetics*, *22*(4), 203–215. https://doi.org/10.1038/s41576-020-00304-w

Hornung, G., Bar-Ziv, R., Rosin, D., Tokuriki, N., Tawfik, D. S., Oren, M., & Barkai, N. (2012). Noise–mean relationship in mutated promoters. *Genome Research*, *22*(12), 2409. https://doi.org/10.1101/gr.139378.112

Huang, W., Carbone, M. A., Magwire, M. M., Peiffer, J. A., Lyman, R. F., Stone, E. A., Anholt, R. R. H., & Mackay, T. F. C. (2015). Genetic basis of transcriptome diversity in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*, *112*(44), E6010–E6019. https://doi.org/10.1073/pnas.1519159112

Jazwinski, S. M., & Kriete, A. (2012). The yeast retrograde response as a model of intracellular signaling of mitochondrial dysfunction. *Frontiers in Physiology*, *3*. https://doi.org/10.3389/fphys.2012.00139

Jia, Y., Rothermel, B., Thornton, J., & Butow, R. A. (1997). A basic helix-loop-helix-leucine zipper transcription complex in yeast functions in a signaling pathway from mitochondria to the nucleus. *Molecular and Cellular Biology*, *17*(3), 1110–1117. https://doi.org/10.1128/MCB.17.3.1110

Johnston, I. G., Gaal, B., Neves, R. P. das, Enver, T., Iborra, F. J., & Jones, N. S. (2012). Mitochondrial variability as a dource of extrinsic cellular noise. *PLOS Computational Biology*, *8*(3), e1002416. https://doi.org/10.1371/journal.pcbi.1002416

Kapust, R. B., Tözsér, J., Fox, J. D., Anderson, D. E., Cherry, S., Copeland, T. D., & Waugh, D. S. (2001). Tobacco etch virus protease: Mechanism of autolysis and rational design of stable mutants with wild-type catalytic proficiency. *Protein Engineering*, *14*(12), 993–1000. https://doi.org/10.1093/protein/14.12.993

Keren, L., van Dijk, D., Weingarten-Gabbay, S., Davidi, D., Jona, G., Weinberger, A., Milo, R., & Segal, E. (2015). Noise in gene expression is coupled to growth rate. *Genome Research*, *25*(12), 1893–1902. https://doi.org/10.1101/gr.191635.115

Laughery, M. F., Hunter, T., Brown, A., Hoopes, J., Ostbye, T., Shumaker, T., & Wyrick, J. J. (2015). New vectors for simple and streamlined CRISPR-Cas9 genome editing in *Saccharomyces cerevisiae*. *Yeast (Chichester, England)*, *32*(12), 711–720. https://doi.org/10.1002/yea.3098

Levy, S. F., Ziv, N., & Siegal, M. L. (2012). Bet hedging in yeast by heterogeneous, age-correlated expression of a stress protectant. *PLoS Biology*, *10*(5), e1001325. https://doi.org/10.1371/journal.pbio.1001325

Liao, X., & Butow, R. A. (1993). *RTG1* and *RTG2*: Two yeast genes required for a novel path of communication from mitochondria to the nucleus. *Cell*, *72*(1), 61–71. https://doi.org/10.1016/0092-8674(93)90050-z

Liao, X. S., Small, W. C., Srere, P. A., & Butow, R. A. (1991). Intramitochondrial functions regulate nonmitochondrial citrate synthase (CIT2) expression in *Saccharomyces cerevisiae*. *Molecular and Cellular Biology*, *11*(1), 38–46. https://doi.org/10.1128/mcb.11.1.38-46.1991

Liu, J., Martin-Yken, H., Bigey, F., Dequin, S., François, J.-M., & Capp, J.-P. (2015). Natural yeast promoter variants reveal epistasis in the generation of transcriptional-mediated noise and its potential benefit in stressful conditions. *Genome Biology and Evolution*, *7*(4), 969–984. https://doi.org/10.1093/gbe/evv047

Liu, P., Song, R., Elison, G. L., Peng, W., & Acar, M. (2017). Noise reduction as an emergent property of single-cell aging. *Nature Communications*, *8*(1), 680. https://doi.org/10.1038/s41467-017-00752-9

Liu, Z., & Butow, R. A. (2006). Mitochondrial retrograde signaling. *Annual Review of Genetics*, *40*, 159–185. https://doi.org/10.1146/annurev.genet.40.110405.090613

Loell, K., Wu, Y., Staller, M. V., & Cohen, B. (2022). Activation domains can decouple the mean and noise of gene expression. *Cell Reports*, *40*(3), 111118. https://doi.org/10.1016/j.celrep.2022.111118

Lu, Y., Biancotto, A., Cheung, F., Remmers, E., Shah, N., McCoy, J. P., & Tsang, J. S. (2016). Systematic analysis of cell-to-cell expression variation of T lymphocytes in a human cohort identifies aging and genetic associations. *Immunity*, *45*(5), 1162–1175. https://doi.org/10.1016/j.immuni.2016.10.025

Magwene, P. M., Willis, J. H., & Kelly, J. K. (2011). The statistics of bulk segregant analysis using next generation sequencing. *PLoS Computational Biology*, *7*(11), e1002255. https://doi.org/10.1371/journal.pcbi.1002255

Martinez-Jimenez, C. P., Eling, N., Chen, H.-C., Vallejos, C. A., Kolodziejczyk, A. A., Connor, F., Stojic, L., Rayner, T. F., Stubbington, M. J. T., Teichmann, S. A., de la Roche, M., Marioni, J. C., & Odom, D. T. (2017). Aging increases cell-to-cell transcriptional variability upon immune stimulation. *Science (New York, N.Y.)*, *355*(6332), 1433–1436. https://doi.org/10.1126/science.aah4115

Metzger, B. P. H., Duveau, F., Yuan, D. C., Tryban, S., Yang, B., & Wittkopp, P. J. (2016). Contrasting frequencies and effects of *cis*- and *trans*-regulatory mutations affecting gene expression. *Molecular Biology and Evolution*, *33*(5), 1131–1146. https://doi.org/10.1093/molbev/msw011

Metzger, B. P. H., Yuan, D. C., Gruber, J. D., Duveau, F., & Wittkopp, P. J. (2015). Selection on noise constrains variation in a eukaryotic promoter. *Nature*, *521*(7552), 344–347. https://doi.org/10.1038/nature14244

Mineta, K., Matsumoto, T., Osada, N., & Araki, H. (2015). Population genetics of non-genetic traits: Evolutionary roles of stochasticity in gene expression. *Gene*, *562*, 16–21.

Morgan, M. D., Patin, E., Jagla, B., Hasan, M., Quintana-Murci, L., & Marioni, J. C. (2020). Quantitative genetic analysis deciphers the impact of *cis* and *trans* regulation on cell-to-cell variability in protein expression levels. *PLoS Genetics*, *16*(3), e1008686. https://doi.org/10.1371/journal.pgen.1008686

Neves, R. P. das, Jones, N. S., Andreu, L., Gupta, R., Enver, T., & Iborra, F. J. (2010). Connecting variability in global transcription rate to mitochondrial variability. *PLOS Biology*, *8*(12), e1000560. https://doi.org/10.1371/journal.pbio.1000560

Newman, J. R. S., Ghaemmaghami, S., Ihmels, J., Breslow, D. K., Noble, M., DeRisi, J. L., & Weissman, J. S. (2006). Single-cell proteomic analysis of *S. cerevisiae* reveals the architecture of biological noise. *Nature*, *441*(7095), 840–846. https://doi.org/10.1038/nature04785

Ohnishi, Y., Huber, W., Tsumura, A., Kang, M., Xenopoulos, P., Kurimoto, K., Oleś, A. K., Araúzo-Bravo, M. J., Saitou, M., Hadjantonakis, A.-K., & Hiiragi, T. (2014). Cell-to-cell expression variability followed by signal reinforcement progressively segregates early mouse lineages. *Nature Cell Biology*, *16*(1), 27–37. https://doi.org/10.1038/ncb2881

Pal, S., & Dhar, R. (2024). Living in a noisy world – Origins of gene expression noise and its impact on cellular decision-making. *FEBS Letters*, *598*(14), 1673–1691. https://doi.org/10.1002/1873-3468.14898

Parab, L., Pal, S., & Dhar, R. (2022). Transcription factor binding process is the primary driver of noise in gene expression. *PLOS Genetics*, *18*(12), e1010535. https://doi.org/10.1371/journal.pgen.1010535

Park, S., Hanekamp, T., Thorsness, M. K., & Thorsness, P. E. (2006). Yme2p is a mediator of nucleoid structure and number in mitochondria of the yeast *Saccharomyces cerevisiae*. *Current Genetics*, *50*(3), 173–182. https://doi.org/10.1007/s00294-006-0087-9

Parks, T. D., Leuther, K. K., Howard, E. D., Johnston, S. A., & Dougherty, W. G. (1994). Release of proteins and peptides from fusion proteins using a recombinant plant virus proteinase. *Analytical Biochemistry*, *216*(2), 413–417. https://doi.org/10.1006/abio.1994.1060

Partow, S., Siewers, V., Bjørn, S., Nielsen, J., & Maury, J. (2010). Characterization of different promoters for designing a new expression vector in *Saccharomyces cerevisiae*. *Yeast (Chichester, England)*, *27*(11), 955–964. https://doi.org/10.1002/yea.1806

Pedraza, J. M., & Paulsson, J. (2008). Effects of molecular memory and bursting on fluctuations in gene expression. *Science (New York, N.Y.)*, *319*(5861), 339–343. https://doi.org/10.1126/science.1144331

Puzović, N., Madaan, T., & Dutheil, J. Y. (2023). Being noisy in a crowd: Differential selective pressure on gene expression noise in model gene regulatory networks. *PLoS Computational Biology*, *19*(4), e1010982. https://doi.org/10.1371/journal.pcbi.1010982

Quarton, T., Kang, T., Papakis, V., Nguyen, K., Nowak, C., Li, Y., & Bleris, L. (2020). Uncoupling gene expression noise along the central dogma using genome engineered human cell lines. *Nucleic Acids Research*, *48*(16), 9406–9413. https://doi.org/10.1093/nar/gkaa668

Raser, J. M., & O'Shea, E. K. (2004). Control of stochasticity in eukaryotic gene expression. *Science (New York, N.Y.), 304*(5678), 1811–1814. https://doi.org/10.1126/science.1098641

Raser, J. M., & O'Shea, E. K. (2005). Noise in gene expression: Origins, consequences, and control. *Science (New York, N.Y.), 309*(5743), 2010–2013. https://doi.org/10.1126/science.1105891

Richard, A., Boullu, L., Herbach, U., Bonnafoux, A., Morin, V., Vallin, E., Guillemin, A., Gao, N. P., Gunawan, R., Cosette, J., Arnaud, O., Kupiec, J.-J., Espinasse, T., Gonin-Giraud, S., & Gandrillon, O. (2016). Single-cell-based analysis highlights a surge in cell-to-cell molecular variability preceding irreversible commitment in a differentiation process. *PLOS Biology, 14*(12), e1002585. https://doi.org/10.1371/journal.pbio.1002585

Richard, M., & Yvert, G. (2014). How does evolution tune biological noise? *Frontiers in Genetics, 5,* 374. https://doi.org/10.3389/fgene.2014.00374

Rocabert, C., Beslon, G., Knibbe, C., & Bernard, S. (2020). Phenotypic noise and the cost of complexity. *Evolution, 74*(10), 2221–2237. https://doi.org/10.1111/evo.14083

Sanchez, A., & Golding, I. (2013). Genetic determinants and cellular constraints in noisy gene expression. *Science (New York, N.Y.), 342*(6163), 1188–1193. https://doi.org/10.1126/science.1242975

Sarkar, A. K., Tung, P.-Y., Blischak, J. D., Burnett, J. E., Li, Y. I., Stephens, M., & Gilad, Y. (2019). Discovery and characterization of variance QTLs in human induced pluripotent stem cells. *PLOS Genetics, 15*(4), e1008045. https://doi.org/10.1371/journal.pgen.1008045

Schmiedel, J. M., Carey, L. B., & Lehner, B. (2019). Empirical mean-noise fitness landscapes reveal the fitness impact of gene expression noise. *Nature Communications, 10*(1), 3180. https://doi.org/10.1038/s41467-019-11116-w

Schmutzer, M., & Wagner, A. (2020). Gene expression noise can promote the fixation of beneficial mutations in fluctuating environments. *PLOS Computational Biology, 16*(10), e1007727. https://doi.org/10.1371/journal.pcbi.1007727

Schor, I. E., Degner, J. F., Harnett, D., Cannavò, E., Casale, F. P., Shim, H., Garfield, D. A., Birney, E., Stephens, M., Stegle, O., & Furlong, E. E. M. (2017). Promoter shape varies across populations and affects promoter evolution and expression noise. *Nature Genetics, 49*(4), 550–558. https://doi.org/10.1038/ng.3791

Shaffer, S. M., Dunagin, M. C., Torborg, S. R., Torre, E. A., Emert, B., Krepler, C., Beqiri, M., Sproesser, K., Brafford, P. A., Xiao, M., Eggan, E., Anastopoulos, I. N., Vargas-Garcia, C. A., Singh, A., Nathanson, K. L., Herlyn, M., & Raj, A. (2017). Rare cell variability and drug-induced reprogramming as a mode of cancer drug resistance. *Nature, 546*(7658), 431–435. https://doi.org/10.1038/nature22794

Sharon, E., van Dijk, D., Kalma, Y., Keren, L., Manor, O., Yakhini, Z., & Segal, E. (2014). Probing the effect of promoters on noise in gene expression using thousands of designed sequences. *Genome Research, 24*(10), 1698–1706. https://doi.org/10.1101/gr.168773.113

Sherman, M. S., Lorenz, K., Lanier, M. H., & Cohen, B. A. (2015). Cell-to-cell variability in the propensity to transcribe explains correlated fluctuations in gene expression. *Cell Systems, 1*(5), 315–325. https://doi.org/10.1016/j.cels.2015.10.011

Siciliano, V., Garzilli, I., Fracassi, C., Criscuolo, S., Ventre, S., & di Bernardo, D. (2013). miRNAs confer phenotypic robustness to gene networks by suppressing biological noise. *Nature Communications, 4*(1), 2364. https://doi.org/10.1038/ncomms3364

Snijder, B., Sacher, R., Rämö, P., Damm, E.-M., Liberali, P., & Pelkmans, L. (2009). Population context determines cell-to-cell variability in endocytosis and virus infection. *Nature, 461*(7263), 520–523. https://doi.org/10.1038/nature08282

Stewart-Ornstein, J., Weissman, J. S., & El-Samad, H. (2012). Cellular noise regulons underlie fluctuations in *Saccharomyces cerevisiae*. *Molecular Cell, 45*(4), 483–493. https://doi.org/10.1016/j.molcel.2011.11.035

Stuckey, S., Mukherjee, K., & Storici, F. (2011). *In vivo* site-specific mutagenesis and gene collage using the delitto perfetto system in yeast *Saccharomyces cerevisiae*. In H. Tsubouchi (Ed.), *DNA Recombination* (Vol. 745, pp. 173–191). Humana Press. https://doi.org/10.1007/978-1-61779-129-1_11

Sukys, A., & Grima, R. (2025). Cell-cycle dependence of bursty gene expression: Insights from fitting mechanistic models to single-cell RNA-seq data. *Nucleic Acids Research, 53*(7), gkaf295. https://doi.org/10.1093/nar/gkaf295

Sun, M., & Zhang, J. (2020). Allele-specific single-cell RNA sequencing reveals different architectures of intrinsic and extrinsic gene expression noises. *Nucleic Acids Research, 48*(2), 533–547. https://doi.org/10.1093/nar/gkz1134

Suter, D. M., Molina, N., Gatfield, D., Schneider, K., Schibler, U., & Naef, F. (2011). Mammalian genes are transcribed with widely different bursting kinetics. *Science (New York, N.Y.), 332*(6028), 472–474. https://doi.org/10.1126/science.1198817

Tan, R. Z., & van Oudenaarden, A. (2010). Transcript counting in single cells reveals dynamics of rDNA transcription. *Molecular Systems Biology, 6*(1), 358. https://doi.org/10.1038/msb.2010.14

Taniguchi, Y., Choi, P. J., Li, G.-W., Chen, H., Babu, M., Hearn, J., Emili, A., & Xie, X. S. (2010). Quantifying *E. coli* proteome and transcriptome with single-molecule sensitivity in single cells. *Science (New York, N.Y.), 329*(5991), 533–538. https://doi.org/10.1126/science.1188308

Tirosh, I., & Barkai, N. (2008). Two strategies for gene regulation by promoter nucleosomes. *Genome Research, 18*(7), 1084–1091. https://doi.org/10.1101/gr.076059.108

Trinh, D.-C., Martin, M., Bald, L., Maizel, A., Trehin, C., & Hamant, O. (2023). Increased gene expression variability hinders the formation of regional mechanical conflicts leading to reduced organ shape robustness. *Proceedings of the National Academy of Sciences of the United States of America, 120*(30), e2302441120. https://doi.org/10.1073/pnas.2302441120

Triqueneaux, G., Burny, C., Symmons, O., Janczarski, S., Gruffat, H., & Yvert, G. (2020). Cell-to-cell expression dispersion of B-cell surface proteins is linked to genetic variants in humans. *Communications Biology, 3*(1), 346. https://doi.org/10.1038/s42003-020-1075-1

Vande Zande, P., Hill, M. S., & Wittkopp, P. J. (2022). Pleiotropic effects of *trans*-regulatory mutations on fitness and gene expression. *Science, 377*(6601), 105–109. https://doi.org/10.1126/science.abj7185

Verhagen, P. G. A., & Hansen, M. M. K. (2025). Exploring the central dogma through the lens of gene expression noise. *Journal of Molecular Biology*, 169202. https://doi.org/10.1016/j.jmb.2025.169202

Vlková, M., & Silander, O. K. (2022). Gene regulation in *Escherichia coli* is commonly selected for both high plasticity and low noise. *Nature Ecology & Evolution, 6*(8), 1165–1179. https://doi.org/10.1038/s41559-022-01783-2

Wang, Z., & Zhang, J. (2011). Impact of gene expression noise on organismal fitness and the efficacy of natural selection. *Proceedings of the National Academy of Sciences, 108*(16), E67–E76. https://doi.org/10.1073/pnas.1100059108

Wernet, M. F., Mazzoni, E. O., Çelik, A., Duncan, D. M., Duncan, I., & Desplan, C. (2006). Stochastic spineless expression creates the retinal mosaic for colour vision. *Nature, 440*(7081), 174–180. https://doi.org/10.1038/nature04615

Wittkopp, P. J., Haerum, B. K., & Clark, A. G. (2004). Evolutionary changes in *cis* and *trans* gene regulation. *Nature, 430*(6995), 85–88. https://doi.org/10.1038/nature02698

Wolf, L., Silander, O. K., & van Nimwegen, E. (2015). Expression noise facilitates the evolution of gene regulation. *eLife, 4*, e05856. https://doi.org/10.7554/eLife.05856

Yang, S., Kim, S., Rim Lim, Y., Kim, C., An, H. J., Kim, J.-H., Sung, J., & Lee, N. K. (2014). Contribution of RNA polymerase concentration variation to protein expression noise. *Nature Communications, 5*(1), 4761. https://doi.org/10.1038/ncomms5761

You, S.-T., Jhou, Y.-T., Kao, C.-F., & Leu, J.-Y. (2019). Experimental evolution reveals a general role for the methyltransferase Hmt1 in noise buffering. *PLoS Biology, 17*(10), e3000433. https://doi.org/10.1371/journal.pbio.3000433

Yvert, G. (2014). 'Particle genetics': Treating every cell as unique. *Trends in Genetics: TIG, 30*(2), 49–56. https://doi.org/10.1016/j.tig.2013.11.002

Zenklusen, D., Larson, D. R., & Singer, R. H. (2008). Single-RNA counting reveals alternative modes of gene expression in yeast. *Nature Structural & Molecular Biology, 15*(12), 1263–1271. https://doi.org/10.1038/nsmb.1514

Zhang, Z., Qian, W., & Zhang, J. (2009). Positive selection for elevated gene expression noise in yeast. *Molecular Systems Biology, 5*, 299. https://doi.org/10.1038/msb.2009.58

Zopf, C. J., Quinn, K., Zeidman, J., & Maheshri, N. (2013). Cell-cycle dependence of transcription dominates noise in gene expression. *PLOS Computational Biology, 9*(7), e1003161. https://doi.org/10.1371/journal.pcbi.1003161

1357