Broad Effects of Activation of Alcohol Dehydrogenase 1 on Healthspan Extension

Avery Sukienik[1,3], Samuel Berhanu[1,5,6], Abbas Ghaddar[1,4,6], Elizabeth MacPhail[1], and Eyleen J. O'Rourke[1,2,7]

1. Department of Biology, College of Arts and Sciences, University of Virginia, Charlottesville, VA 22903, USA
3. Department of Cell Biology, School of Medicine, University of Virginia, Charlottesville, VA 22903, USA
4. Current address: Department of Bioengineering, George R. Brown School of Engineering and Computing, Rice University, Houston, Texas 77005, United States
5. Current address: Emory University School of Medicine, 2015 Uppergate Dr, Atlanta, GA 30307
6. Equal contribution
7. Correspondence to ejorourke@virginia.edu

Summary

Nutritional, genetic, and pharmacological interventions can extend lifespan; however, fewer have been shown to extend healthspan—the period of life free from chronic, debilitating diseases. In line with this, the molecular effectors that drive healthspan are even less understood than those responsible for lifespan extension. We recently reported that activation of Alcohol Dehydrogenase 1 (ADH-1) extends lifespan in yeast and *C. elegans*. In addition, *adh-1* is transcriptionally activated in yeast, worms, mice, and humans in response to caloric restriction—an intervention that extends not only lifespan but also healthspan. Therefore, we investigated whether activating *adh-1* could also extend healthspan. We demonstrate here that *adh-1* activation has broad and robust effects on health, including resistance to age-related obesity, delayed sarcopenia, and attenuated neurodegeneration. Mechanistically, ADH-1-driven healthspan extension is associated with improved proteostasis. These findings position ADH-1 as a promising target for future research aimed at promoting healthy aging.

Keywords

Alcohol Dehydrogenase 1, ADH-1, Healthspan, Lifespan, Aging, Longevity, Age-related disease, Sarcopenia, Neurodegeneration, Proteostasis, ER stress, Oxidative stress

Introduction

Healthspan is the length of life spent free from serious illness or disability. Although human life expectancy has increased by approximately 30 years over the past century(Crimmins, 2015), healthspan extension has not kept pace—a phenomenon referred to as the *healthspan gap* (Rowe & Kahn, 1987). This gap is also observed in laboratory conditions following genetic, nutritional, and pharmacological interventions that extend lifespan. For instance, *Caenorhabditis elegans* carrying loss-of-function mutations in *clk-1*, a gene encoding a protein required for coenzyme Q (a.k.a. ubiquinone-9, a key component of the electron transport chain) biosynthesis, exhibit increased longevity but also severe developmental and motility defects that are partially diet-dependent(Branicky et al., 2000; Hihi et al., 2002). Even the best-characterized long-lived *C. elegans* mutant, *daf-2*—which encodes the worm's sole insulin/IGF-1 receptor, shows pronounced developmental defects(Gems et al., 1998). Vertebrate longevity models also commonly display healthspan gaps. For example, mice carrying point mutations in codon 72 of the tumor suppressor p53 are long-lived(Migliaccio et al., 1999) but exhibit increased cancer susceptibility(Zhao et al., 2018). Similarly, methionine-restricted mice have extended lifespan but reduced bone density(Ables & Johnson, 2017), and the most studied mouse models of longevity, Ames and Dwarf(Bartke et al., 2001), show strong developmental and fertility defects(Kano, 2013). Thus, across diverse organisms, lifespan-extending interventions do not necessarily lead to proportional healthspan gains.

We recently reported that elevated Alcohol Dehydrogenase I (ADH-1) activity is necessary for lifespan extension driven by caloric restriction, as well as by mTOR and insulin signaling inhibition in *C. elegans* (Ghaddar, Mony, et al., 2023). (Ghaddar et al., 2023). Importantly, constitutive activation of ADH-1 was also sufficient to extend lifespan, not only in worms but also in yeast. Furthermore, we found that calorically restricted mice, rats, pigs, Rhesus monkeys, and humans exhibit elevated *Adh1*/ADH1 expression (Ghaddar, Mony, et al., 2023), suggesting evolutionary conservation of this response. In that study, and in the present one, we used three independent transgenic *C. elegans* lines that constitutively overexpress *adh-1* (hereafter referred to as ADH-1[OE]). These lines were validated to exhibit increased *adh-1* mRNA and protein expression and, critically,

chronically elevated alcohol dehydrogenase activity(Ghaddar et al., 2023). The absence of obvious detrimental phenotypes in these long-lived animals (Ghaddar, Mony, et al., 2023; Molotkov et al., 2002) supports the notion that ADH-1 activation may represent a viable strategy to promote longevity. However, whether ADH-1 activation also extends healthspan—delaying the onset or severity of age-associated functional decline—remains unknown.

Here, we demonstrate that the sole overexpression of ADH-1 (ADH-1^{OE}) is sufficient to robustly extend healthspan in *C. elegans* feeding *ad libitum*. ADH-1^{OE} animals are resistant to both age-related and diet-induced obesity and exhibit reduced sarcopenia and neurodegeneration. Mechanistically, ADH-1^{OE} animals show transcriptional activation of a proteostatic program, accompanied by increased resistance to protein aggregation. These findings establish ADH-1 as a targetable driver of healthspan extension, supporting its potential as a therapeutic intervention against age-related decline.

Results

ADH-1^{OE} *C. elegans* are Resistant to Age-related and Diet-induced Obesity

Adiposity increases with age and is associated with the overall decline in health observed in aging animals (Bischof & Park, 2015; Palmer & Jensen, 2022; Palmer & Kirkland, 2016). Across organisms, life-extending interventions have diverse effects on adiposity, ranging in *C. elegans* from leanness under caloric restriction (Klapper et al., 2011) to obesity in long-lived notch (*glp-1*) and insulin receptor (*daf-2*) mutants (O'Rourke et al., 2009). To establish whether ADH-1 activity influences age-driven changes in adiposity, we first tested whether age-related obesity can be captured in *C. elegans* using the fat-specific dye Oil Red O (ORO) (Ke et al., 2018; Wählby et al., 2014a). For this, we stained young (day 1) and aged (day 10) wild-type animals. As anticipated, we observed higher ORO signal in aged worms (Fig. 1B), confirming that age-related fat accumulation can be effectively modeled and assessed in *C. elegans*.

C. elegans overexpressing ADH-1 (ADH-1^{OE}) exhibit approximately 5-fold higher *adh-1* expression than wild-type animals (Ghaddar et al., 2023)–a level comparable to the induction observed under caloric restriction, a condition known to mobilize fat stores in *C. elegans* and other organisms (O'Rourke et al., 2009; Ghaddar et al., 2023). This parallel led us to test whether sustained ADH-1 activation could be sufficient to promote fat mobilization and counteract age-associated obesity. Indeed, ADH-1^{OE} worms markedly reduced lipid stores by day 10 of adulthood (Fig. 1A, B), suggesting improved metabolic health with age. Interestingly, even at day 1, ADH-1^{OE} worms were slightly leaner than wild-type controls (Fig. 1C, DataS1A, B).

To rule out the possibility that the observed reduction in Oil Red O (ORO) staining reflected altered cuticle permeability rather than true fat depletion, we tested staining across different incubation times. The rationale was that the cuticle becomes more permeable with age, and if ADH-1^{OE} animals remained more "youthful," their cuticles might take longer to absorb the dye. However, we found that the lean phenotype of aged ADH-1^{OE} animals was consistent regardless of staining duration and was evident even after ORO saturation of ADH-1^{OE} animals had been reached (DataS2A), supporting a genuine reduction in fat content.

The experiments we describe were performed using the PHX2888 strain, one of three independently generated ADH-1^{OE} lines. In our previous work (Ghaddar et al., 2023), we characterized all three lines and found them to exhibit indistinguishable phenotypes across multiple biological scales, including *adh-1* expression levels, enzymatic activity, subcellular localization, and organismal phenotypes such as lifespan extension. Likewise, the lean phenotype was consistent across all three independent ADH-1^{OE} strains (DataS1D-F). We selected the PHX2888 line—hereafter referred to simply as ADH-1^{OE}— for the detailed healthspan analyses presented in this study, as it showed an intermediate degree of both leanness and lifespan extension.

Because aging also increases susceptibility to diet-induced metabolic dysregulation, we next tested whether ADH-1 activity could counter the obesogenic effects of fructose feeding in aged *C. elegans* (Ke et al., 2021). Consistent with this hypothesis, ADH-1^{OE} animals fed a fructose-enriched diet accumulated less fat than wild-type controls at both days 1 and 10 (Fig. 1C, DataS1C).

Because reduced fat stores in aged animals could reflect impaired feeding or abnormal food perception rather than improved metabolic health, we directly tested these possibilities at day 1 and 10. First, we examined pharyngeal function and found that both young and aged ADH-1^{OE} worms displayed normal contractility with aged ADH-1^{OE} animals displaying greater contractility relative to wild type (Video 1, Video 2; Fig. 1D), indicating enhanced rather than diminished feeding capacity. We then assessed whether reduced adiposity might arise from defects in food detection or active avoidance. Both young and aged ADH-1^{OE} animals responded normally to food cues and did not avoid food with aged ADH-1^{OE} performing better in the food cue response relative to WT (Fig. 1E). Together, these controls support the conclusion that reduced fat stores in aged ADH-1^{OE} animals reflect improved metabolic homeostasis, rather than feeding or sensory defects. The results also suggest better muscle function (pharynx) and locomotory and sensory capacity (chemotaxis) in ADH-1^{OE} *C. elegans*.

Figure 1. ADH-1 overexpression causes resistance to age-driven and diet-induced obesity without compromising feeding capacity. (A) Representative images of wild-type (WT) and ADH-1[OE] *C. elegans* stained with the neutral-fat dye Oil Red O (ORO) at day 10 of adulthood (n=3 biological replicates). (B) Quantification of rORO intensity in WT and ADH-1[OE] animals fed a standard *E. coli* OP50 diet at day 1 and day 10 of adulthood (n=3 biological replicates). (C) Quantification of ORO intensity in WT and ADH-1[OE] animals fed an obesogenic High Fructose Diet (HFrD) (n=3 replicates, refer to DataS1C). (D) Quantification of pharyngeal pumping in WT and ADH-1[OE] animals at day 1 and day 8 of adulthood (n=3 biological replicates). (E) Chemotactic index (CI) of WT and ADH-1[OE] animals at day 1 and day 10 of adulthood (n=3 biological replicates). Statistical testing was conducted using Welch's t-test after outlier correction or Wilcoxon when populations were non-normal (Panels D, E, F, and G). Error bars denote SEM. ns= not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, **MPI** = Mean Pixel Intensity.

ADH-1[OE] *C. elegans* are Resistant to Oxidative Stress

Fats stored in lipid droplets, like those stained by ORO, can act as sinks for reactive oxygen species (ROS) that would otherwise cause broad-spectrum cellular damage (Bailey et al., 2015). For instance, insulin-insensitive *daf-2* mutant *C. elegans*—which have larger-than-wild-type fat stores—are more resistant to oxidative stress (Honda & Honda, 1999). Given that overexpressing ADH-1 reduces fat stores, we considered the possibility that this might compromise the animals' resistance to oxidative stress.

To test whether reduced lipid buffering might increase ROS toxicity in ADH-1[OE] animals, we exposed them to paraquat and measured survival, as previously described (Ishii et al., 1990; Mony et al., 2021; Senchuk et al., 2017). Surprisingly, ADH-1[OE] animals exhibited enhanced survival compared to wild-type controls (Fig. 2A and B, DataS3 B-D), suggesting that fat depletion in these animals does not impair, but instead improves oxidative stress resilience.

Consistent with this enhanced resistance, ADH-1[OE] worms exhibit constitutively elevated expression of antioxidant genes, including *ctl-2, gpx-5, gst-4, gst-5, and gst-16* (Back et al., 2012a; Hu et al., 2018). Notably, *ctl-2* has been directly implicated in protection from ROS-induced damage(Musa et al., 2022; Petriv & Rachubinski, 2004). Additional redox-

responsive genes such as *ctl-3* and *sod-1* also trended toward upregulation in ADH-1[OE] worms (Data S3A), consistent with a broader activation of antioxidant defenses (Blackwell et al., 2015).

Mild elevations in ROS levels have been documented in other long-lived *C. elegans* mutants, such as *clk-1* and *sod-2*, and in some cases are essential for their extended lifespan(Raamsdonk & Hekimi, 2009; Yang & Hekimi, 2010b). Similarly, low-dose oxidant exposure (e.g., juglone or paraquat) has been shown to increase lifespan via hormesis—whereby transient oxidative stress triggers protective responses that increase long-term survival(Heidler et al., 2010; Lee et al., 2010; Yang & Hekimi, 2010a). This model suggests that improved survival following mild oxidative stress may arise not from targeted adaptation to each specific insult that an animal could encounter throughout its lifespan, but from a generalized enhancement of stress resilience. Such broad-spectrum protection could defend against diverse ROS-linked threats encountered throughout life, including fatty acid oxidation, pathogen attack and its associated immune responses(Back et al., 2012b; Ristow & Schmeisser, 2011; Schulz et al., 2007), and is often referred to as hormesis.

To determine whether ADH-1[OE] animals exhibit elevated endogenous ROS that might trigger such a hormetic response, we measured hydrogen peroxide (H_2O_2) levels using the HyPer biosensor—a genetically encoded, ratiometric probe combining a circularly permuted YFP with the H_2O_2-sensitive regulatory domain of *E. coli* OxyR (Belousov et al., 2006). ADH-1[OE] day 1 adults displayed significantly elevated H_2O_2 levels compared to wild-type animals (Fig. 2D, E), consistent with a state of mild endogenous oxidative stress. Notably, despite this increase, ADH-1[OE] animals not only survived paraquat exposure better than controls but also showed upregulation of multiple antioxidant defense genes. These findings support a model in which enhanced ADH-1 activity initiates a preconditioning response that boosts redox detoxification capacity. Thus, rather than sensitizing animals to oxidative damage, fat depletion via ADH-1 overexpression appears to enhance oxidative stress resistance, potentially through hormetic activation of stress response pathways.

Figure 2. ADH-1 overexpression increases resistance to oxidative stress. (A) Day-1 ADH-1[OE] *C. elegans* adults show enhanced survival to 100mM paraquat when compared to WT (representative of 2 biological replicates, see also DataS1C). (B) Day-8 ADH-1[OE] *C. elegans* adults show enhanced survival to 100mM paraquat when compared to WT (representative of 3 biological replicates, see also DataS1D and E). (C) Day-1 ADH-1[OE] adults show constitutive activation of an anti-oxidant transcriptional program (n=4 biological replicates), (D) ADH-1[OE] animals show heightened H_2O_2 levels at day 1 of age as indicated by the HyPer::YFP H_2O_2 sensor, (E) Quantification of HyPer::YFP signal showing ADH-1[OE] animals have increased ROS levels when compared to WT (n=3 biological replicates). Statistical testing was conducted using Welch's t-test after outlier

correction or Wilcoxon when populations were non-normal (Panel E). Statistical testing of survival conducted by Kaplan-Meier Log-Rank Fold test. Error bars denote SEM. ns= not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

ADH-1^{OE} Enhances ER Stress Response and Promotes Metabolic Homeostasis

Lean phenotypes and resistance to oxidative stress have been associated with enhanced endoplasmic reticulum (ER) function and improved protein homeostasis (proteostasis) in multiple model systems, including *C. elegans* and mammals(Ben-Zvi et al., 2009; Henis-Korenblit et al., 2010; Taylor & Dillin, 2013a). This relationship is particularly relevant to aging and metabolic disease, as chronic or excessive ER stress—when the protein-folding capacity of the ER is overwhelmed—contributes to metabolic dysfunction, insulin resistance, and obesity(Hotamisligil, 2010; Özcan et al., 2004; Zhang & Kaufman, 2008). Paradoxically, transient or mild activation of the ER stress response, particularly the unfolded protein response (UPR), can be protective. A growing body of evidence suggests that low-level UPR activation can enhance the cellular capacity to manage misfolded proteins and oxidative stress, promoting metabolic resilience and extending organismal healthspan (Calfon et al., 2002a; Hetz & Papa, 2018; Wang & Kaufman, 2016). In this context, the UPR acts not as a sign of dysfunction but as an adaptive, preemptive defense mechanism.

To explore whether the improved metabolic and stress resilience of ADH-1^{OE} animals might stem from enhanced ER homeostasis, we examined the expression of *hsp-4*, a key component of the unfolded protein response and the *C. elegans* ortholog of mammalian BiP/GRP78. Using a *hsp-4*P::GFP reporter, we found that *hsp-4* expression is modestly but consistently elevated in ADH-1^{OE} worms under non-stressed conditions, particularly in the intestine and tail regions—tissues central to metabolic regulation (Fig. 3A and B). This trend was also reflected, albeit to a lesser extent, in *hsp-4* transcript levels (DataS4C).

Importantly, while *hsp-4*P::GFP levels are elevated at baseline in ADH-1^{OE} animals, this increase is far more modest than the strong induction observed following exposure to

5 µg/mL tunicamycin, a well-established ER stressor(Calfon et al., 2002b) (Fig. 3C and D). This distinction suggests that the elevation in *hsp-4* expression in ADH-1[OE] animals likely reflects a mild, hormetic activation of the UPR rather than a pathological stress response. In turn, this preconditioning may enhance the animal's capacity to cope with future ER stress, potentially contributing to its improved metabolic health and oxidative stress resistance.

A

Day 1
*hsp-4*P::GFP Expression

WT

ADH-1^OE

B

Day 1 Expression of *hsp-4*P::GFP

****

WT   ADH-1^OE

C

Day 1 Tunicamycin
*hsp-4*P::GFP Expression

WT

ADH-1^OE

D

Day 1 log10(Expression) of *hsp-4*P::GFP

****

WT   ADH-1^OE

Figure 3. Activating ADH-1 causes mild activation of ER stress response. (A) Representative images showing that ADH-1[OE] animals show enhanced expression of the gene encoding the ER-specific chaperone HSP-4 in the intestine–mainly in the anterior and posterior intestine, which accumulate most fat stores in *C. elegans*. (B) Quantification of *hsp-4*P::GFP intensity in unstressed wild-type and ADH-1[OE] *C. elegans* as shown in panel A (n=3, see also DataS4A-B). (C) Quantification of *hsp-4*P::GFP intensity in wild-type *C. elegans* stressed with 5µg/µl of Tunicamycin, a treatment that induces pathological ER stress (n=2). Error bars denote SEM. Statistical testing was conducted using Welch's t-test after outlier correction or Wilcoxon when populations were non-normal (Panels B and D). ns= not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

ADH-1[OE] Delays Age-dependent Loss of Locomotory Capacity

The improved pharyngeal pumping observed in aged *C. elegans* overexpressing ADH-1 (Fig. 1D) suggests that ADH-1[OE] enhances the function of aging striated autonomic muscles. We next examined whether skeletal muscle integrity is similarly preserved in these animals. To assess this, we measured locomotory function using the thrashing assay, which quantifies the number of body bends per second (bbps) as worms respond to liquid immersion. This behavior declines with age and is a widely used proxy for neuromuscular health in *C. elegans*(Dillin et al., 2002; Glenn et al., 2004; Herndon et al., 2002; Hosono, 1978; Hosono et al., 1980; Hsu et al., 2009; C. Huang et al., 2004; Johnson et al., 1988; Liu et al., 2013 At day 1 of adulthood, locomotory activity was similar between wild-type and ADH-1[OE] animals (Fig. 4A). However, by day 10, ADH-1[OE] worms maintained significantly higher bbps compared to age-matched controls (Fig. 4A; DataS5A-B), indicating that the locomotory benefits of ADH-1[OE] emerge with age.

Age-dependent locomotory decline is typically accompanied by progressive deterioration of skeletal muscle fibers and motor neurons (Chow et al., 2006; Cruz-Jentoft et al., 2019; Dhondt et al., 2021; Giunti et al., 2024; X. Huang et al., 2002; Pan et al., 2011; Tank et

al., 2011). Thus, the improved locomotory activity observed in aged ADH-1^{OE} worms suggests that elevated alcohol dehydrogenase activity would help preserve motor neuron and/or skeletal muscle function.

GABAergic motor neurons are particularly prone to early age-dependent decline, experiencing functional decline before the onset of sarcopenia (Glenn et al., 2004; Herndon et al., 2002). This decline is characterized most prominently by late-life structural abnormalities in the commissures—the axons that project from the ventral and dorsal cord neurons of *C. elegans* GABAergic motor neurons (Fig. 4C). Common abnormalities include bulges, beading, and fasciculation(Pan et al., 2011; Tank et al., 2011), collectively referred to as "stippling", which serve as markers of overall neuronal health(Giunti et al., 2024). To determine whether the enhanced locomotory function in ADH-1^{OE} worms is associated with improved motor neuron maintenance, we examined commissure structure in aging worms(Gjorgjieva et al., 2014). In line with improved motor neuron maintenance, 12-day-old ADH-1^{OE} worms exhibited less commissure stippling than their wild-type counterparts (Fig. 4B & C). Notably, while WT control animals showed examples of extreme neurodegeneration (DataS5C), ADH-1^{OE} animals were not noted to have any such cases at day 12.

Having established that ADH-1^{OE} preserves motor neuron integrity with age, we next asked whether skeletal muscle structure might also be protected. Age-related locomotory decline is strongly associated with sarcopenia—the progressive degeneration of body wall muscles—which manifests cytologically as disorganized and degraded muscle fibers (Cruz-Jentoft et al., 2019, 2019; Dhondt et al., 2021). To assess whether the improved locomotory capacity in ADH-1^{OE} worms was also associated with preserved muscle architecture, we used a strain expressing a GFP::MYO-3 fusion protein(Campagnola et al., 2002), which allows visualization of the body wall muscle sarcomeres through green fluorescence. Sarcopenia was assessed by binary scoring of GFP::MYO-3 in the midbody: worms with well-structured sarcomeres received a score of 1, while those with severe sarcomere disorganization received a score of 0 (Fig. 4D, E). As expected, muscle fibers in both wild-type and ADH-1^{OE} worms exhibited parallel, well-organized sarcomere layers—most scoring as grade 1—in young day 1 adults (Fig. 4D, E). While sarcomere deterioration occurred with age in both genotypes, 10-day-old ADH-

1^{OE} worms showed significantly better preservation of striated muscle structure (Fig. 4F). Interestingly, the proportion of intermediately damaged fibers was similar between groups (DataS5E, H), suggesting that muscle aging may proceed in discrete steps rather than as a gradual decline. This pattern is consistent with an age-dependent increase in severely damaged fibers in wild-type worms, a trajectory from which ADH-1^{OE} animals appear partially protected (DataS5D–I).

Together, these results suggest that the improved locomotory function observed in aged ADH-1^{OE} worms stems from the combined structural preservation of both GABAergic motor neurons and body wall muscles.

Figure 4. ADH-1 Overexpression Preserves Motor Neuron and Skeletal Muscle Structure During Aging. (A) Quantification of locomotion capacity through the thrashing assay shows no baseline advantage of ADH-1^{OE} animals at day 1 (n = 3 biological replicates), but at day 12, ADH-1^{OE} animals exhibit more BBPS than WT animals, indicative of improved locomotory capacity relative to same-age WT *C. elegans* (representative of 3 biological replicates; see also DataS5). (B) Representative images of the axonal commissures of GABAergic motor neurons in WT (left) and ADH-1^{OE} (right) animals at day 12 (n = 2 biological replicates). (C) Quantification of GABAergic motor neurodegeneration showing that ADH-1^{OE} animals have less axonal stippling events per commissure than same-age WT worms (n = 2 biological replicates). (D) Schematic showing the muscle health scoring system used to assess sarcopenia status. (E) At day 1, WT and ADH-1^{OE} animals display similarly well-structured sarcomeres. (F) However, at day 10, the muscle health scores of ADH-1^{OE} animals are significantly higher than those of WT (n = 2 biological replicates). Statistical testing was conducted using Welch's t-test

after outlier correction, or Wilcoxon test when populations were non-normal (Panel A and F). ns = not significant; *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001.

ADH-1^{OE} Improves Muscle Proteostasis and Late-Age Locomotion

Impaired proteostasis is a fundamental driver of age-related decline in muscle structure and function across species, contributing to sarcopenia and reduced physical performance. In *C. elegans*, as in mammals, the progressive accumulation of misfolded and aggregated proteins in muscle cells underlies functional deterioration with age.(Hipp et al., 2019; Labbadia & Morimoto, 2014; Meller & Shalgi, 2021). Given that ADH-1 overexpression preserves muscle integrity and locomotory performance during aging, we tested whether it also improves muscle proteostasis. To address this, we employed two well-established *C. elegans* models of muscle-specific proteotoxic stress.

Most proteins fold optimally within a narrow temperature range, and deviations from this range activate the unfolded protein response (UPR) and recruit chaperones to assist in protein refolding. Thus, one approach to challenging the global proteostasis network is to expose animals to heat stress. Chronic exposure to 27°C is sufficient to induce proteostatic stress in *C. elegans*, providing a straightforward assay for the organism's capacity to maintain protein homeostasis as it ages. Consistent with improved proteostasis, ADH-1^{OE} animals survived longer than wild-type controls under chronic heat stress (Fig. 5A, DataS3E), indicating enhanced resilience to protein misfolding stress.

While heat stress resistance is a widely used proxy for organismal proteostasis, it reflects the integrated response of multiple cellular pathways and stress axes, making it an informative but indirect readout of proteostatic capacity. To directly assess muscle-specific proteostasis, we next used a well-established model of muscle proteotoxicity in *C. elegans*: the AM140 strain, which carries the integrated transgene *rmIs132* (*unc-54p::Q35::YFP*). This construct expresses a polyglutamine (polyQ) stretch fused to YFP under control of the *unc-54* muscle-specific promoter, leading to age-dependent aggregation of the fusion protein in body wall muscle cells. In young AM140 animals, the

YFP signal is diffuse and outlines the muscle fibers, producing a smooth contour along the body wall. As the animals age, the YFP signal becomes increasingly punctate due to protein aggregation, and the fiber-defined contour is lost—reflecting a decline in muscle proteostasis. (Ellis et al., 2023; Morley et al., 2002).

To test whether ADH-1 overexpression enhances muscle proteostasis, we crossed the *Q35::YFP* transgene into the ADH-1^{OE} background. Longitudinal analysis revealed that ADH-1^{OE} animals exhibited longer and more continuous stretches of diffuse YFP signal compared to wild-type animals, beginning as early as day 4 of adulthood (Fig. 5B-C). This observation suggests reduced aggregation and supports the conclusion that ADH-1^{OE} extends the period during which muscle proteostasis is maintained.

To investigate the molecular basis of this enhanced proteostasis, we measured the expression of several heat shock proteins encoding genes (*hsp*), which are critical components of the proteostatic network. Quantitative PCR analysis showed that ADH-1^{OE} animals exhibit elevated expression of *hsp-16.1*, *hsp-16.2*, and *hsp-60* (Fig. 5D).

To investigate the molecular basis of the enhanced proteostasis observed in ADH-1^{OE} animals, we measured the expression of several heat shock protein (*hsp*) genes, which are key components of the proteostasis network. Quantitative PCR analysis revealed that ADH-1^{OE} worms exhibit elevated expression of *hsp-16.1*, *hsp-16.2*, and *hsp-60* (Fig. 5D).

To test whether transcriptional activation of heat shock proteins plays a role in the locomotory benefits of ADH-1^{OE}, we knocked down *hsf-1*, the master regulator of the heat shock response (Hajdu-Cronin et al., 2004; Morley & Morimoto, 2004). Although *hsf-1* RNAi had no effect on locomotion in day 1 adults (Fig. 5E), it suppressed the improved locomotory capacity of aged ADH-1^{OE} animals (Fig. 5F), suggesting that *hsf-1*-dependent transcriptional programs are required for the maintenance of locomotion in these animals.

Together, these findings suggest that ADH-1^{OE} enhances muscle proteostasis at multiple levels, including both tissue-specific reductions in protein aggregation and systemic improvements in heat stress resilience, likely mediated through *hsf-1*-dependent upregulation of HSPs. These results support a model in which ADH-1^{OE} extends

healthspan not only by improving metabolic homeostasis, but also by maintaining proteostatic capacity in aging muscle through the activation of *hsf-1*–driven stress

response genes.

Figure 5. ADH-1 overexpression results in resistance to proteostatic stress. (A) Representative survival curve showing that ADH-1^{OE} animals better survive chronic heat stress (n=2 replicates, see DataS3E) (B) Representative images of day-4 animals expressing polyQ::YFP (depicted green) with masks highlighting the thresholded aggregates (depicted yellow to white according to their increasing signal intensity) (C) Quantification of total surface area of aggregates per animal shows that ADH-1^{OE} animals have less polyQ::YFP aggregation than same-age WT worms (n=3 biological replicates), (D) ADH-1^{OE} animals show increased expression of key proteostatic genes (n=4 biological replicates), (E) *hsf-1i* has no impact on locomotory capacity at day 1, (F) *hsf-1i* suppresses ADH-1^{OE} dependent locomotory capacity improvements at day 8 (n=1 biological replicate). Statistical testing conducted by Welch's t-test after outlier correction or by Wilcoxon when populations were non-normal (Panels E and F). Error bars denote SEM. ns= not significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

ADH-1 May Act Through a Non-Cell-Autonomous Mechanism

Our findings indicate that overexpression of *adh-1* preserves the function of multiple tissues in aging *C. elegans*, including GABAergic motor neurons (Fig. 3D). However, single-cell RNA sequencing data (Fig. 6A; Ghaddar, Armingol, et al., 2023) and confocal imaging of endogenously tagged (CRISPR-edited) ADH-1 (Fig. 6B) reveal a more restricted expression pattern for this enzyme. Specifically, *adh-1* is predominantly expressed in body wall muscle, pharyngeal muscle, marginal cells, and the intestine— but not in GABAergic motor neurons (Fig. 6C–F). Similarly, we observed no detectable *adh-1* expression in sensory neurons (Fig. 6D), which—although not directly examined here—are likely better maintained in ADH-1^{OE} animals, based on their improved chemotactic performance (Fig. 1F, G).

Together, these findings suggest that the beneficial effects of ADH-1 overexpression extend beyond its sites of expression, supporting a cell non-autonomous mechanism of action. While the precise inter-tissue signals mediating these effects remain to be elucidated, our results establish that ADH-1 activation enhances healthspan across multiple systems in *C. elegans*. This observation sets the stage for future studies aimed

at identifying systemic signals—such as metabolites, redox-active molecules, myokines, or other intercellular messengers—that may underlie the broad physiological improvements associated with ADH-1 activity.

Figure 6. ADH-1 may act through a non-cell autonomous mechanism. (A) Transcripts per million (TPMs) of *adh-1* in tissues of interest as extracted from wormseq.org(Ghaddar, Armingol, et al., 2023). "Proportion" indicates percentage of cells in the cluster that express *adh-1* (B) Schematic of endogenous CRISPR tagging with wrmScarlet of the *adh-1* open reading frame (edited genotype: *adh-1*p::ADH-1::wrmScarlet::*adh-1* 3'UTR). (C) ADH-1 is strongly expressed in the intestine (outlined in yellow) and the pharynx. (D)

ADH-1 is strongly expressed in the pharynx (yellow arrow) but not in AWB neurons, as indicated by lack of colocalization with *str-1*p::GFP (green arrow). (E) ADH-1 is expressed in body wall muscle (BWM), as defined by colocalization with *myo-3p::GFP*. (F) ADH-1 is not detectable in GABAergic motor neurons, as indicated by lack of colocalization with *unc-25*p::GFP. Muscle expression of *adh-1*::wrmScarlet is outlined in yellow for comparison.

Discussion

Many established geroprotective interventions extend lifespan, but they often do so unevenly across organ systems, sometimes improving select features of healthspan while neglecting or impairing others. For example, loss-of-function in *clk-1* (CoQ7) extends lifespan but causes delayed development and reduced locomotion(Cristina et al., 2009). As another illustration of trade-offs, methionine restriction, while extending lifespan in mice, has been associated with reduced bone density(Ables et al., 2012). As another illustration of trade-offs, methionine restriction, while extending lifespan in mice, has been associated with reduced bone density(Ables et al., 2016). By contrast, ADH-1 overexpression yields coordinated health benefits across distinct physiological domains—including proteostasis, stress resilience, locomotion, neuronal maintenance, and leanness. This breadth of benefit invites consideration of how aging processes across tissues might be modulated by a common upstream factor.

At a mechanistic level, our findings suggest that ADH-1 promotes healthspan through metabolic rewiring that subtly alters cellular redox balance (Summary Figure). The modest but consistent increase in reactive oxygen species (ROS) observed in ADH-1^{OE} animals is reminiscent of mitohormesis, wherein low-level oxidative stress activates protective transcriptional programs that confer stress resistance and longevity(Ristow & Zarse, 2010; Yun & Finkel, 2014). Such hormetic ROS signaling has been implicated in longevity-promoting effects of metformin (De Haes et al., 2014), mitochondrial uncoupling (Cristina et al., 2009), and proline catabolism (Zarse et al., 2012).

Given ADH-1's biochemical function in alcohol oxidation and its impact on NAD$^+$/NADH balance, its overexpression may stimulate NAD$^+$ cycling and enhance mitochondrial metabolism via β-oxidation and glycerol catabolism—both of which can influence ROS production and systemic energy homeostasis. Indeed, our previous work showed that ADH-1 promotes leanness under nutrient-rich conditions (Ghaddar & Mony et al., 2023), positioning it as a modulator of intracellular flux and metabolic flexibility.

This redox shift appears to prime cytoprotective pathways. ADH-1^{OE} animals exhibit hallmarks of enhanced proteostasis, including upregulation of multiple hsp genes and activation of the ER unfolded protein response (UPR), as shown by both transcriptomic analyses and *hsp-4*p::GFP reporter expression. While the UPR is classically associated with cellular stress, moderate activation enhances protein folding capacity and extends lifespan in *C. elegans* and other models (Taylor & Dillin, 2013b). Disruptions in protein glycosylation or folding, potentially resulting from glucose limitation via glycerol diversion, could underlie the ER stress observed in ADH-1^{OE} animals(Cruz-Rodríguez et al., 2025) . ROS can also engage UPR sensors (e.g., IRE-1 and ATF6) through redox-sensitive modifications (Cao & Kaufman, 2014; Malhotra & Kaufman, 2007), providing a molecular basis for redox–UPR coupling. Importantly, our genetic data indicate that such coupling is not uniformly required for all ADH-1 benefits: while UPR priming may contribute to proteostasis and heat-stress resilience, RNAi against UPR branches did not eliminate the leanness phenotype (with only partial trends), suggesting that metabolic outcomes can be decoupled from IRE-1/ATF-6–dependent signaling in this context.

Muscle function is another critical determinant of healthspan, as the maintenance of muscle integrity is essential for preserving locomotory capacity in aging animals. The improved sarcomere organization and muscle fiber integrity observed in ADH-1^{OE} animals are consistent with the observed upregulation of small heat shock proteins, including *hsp-16.1*, *hsp-16.2*, and *hsp-70*, which are critical for maintaining muscle proteostasis(Kutzner et al., 2024; Shemesh et al., 2013; van Oosten-Hawle et al., 2013). RNAi against *hsf-1*— the master regulator of the heat-shock response—suppressed the locomotory

improvements, indicating that this transcriptional axis is required for extended muscle performance in ADH-1^{OE} animals.

Intriguingly, several tissues that benefit from ADH-1 overexpression do not express the gene endogenously. We observed no detectable ADH-1 expression in GABAergic or olfactory neurons, yet ADH-1^{OE} animals maintain better axonal structure and chemotaxis later in life. This disconnect between expression and benefit supports a non-cell-autonomous mode of action, in which ADH-1 activity in muscle, pharynx, or intestine initiates systemic protective programs (Summary Figure). Prior work has established that transcription factors such as HSF-1, DAF-16, and SKN-1 can orchestrate broad organismal responses, including cross-tissue proteostasis signaling (Prahlad et al., 2008; Taylor & Dillin, 2013c). In contrast to these transcriptional master regulators, ADH-1 is a metabolic enzyme, raising the possibility that enzymatic remodeling of metabolites and redox tone constitutes a distinct, under-appreciated mechanism of tissue crosstalk that warrants further investigation.

Rather than contrasting ADH-1 with "isolated" longevity axes, we emphasize potential mechanistic overlap with canonical pathways. In (Ghaddar, Mony, et al., 2023), *adh-1* activity was required for the longevity of *daf-2* mutants, *eat-2* (DR) models, and mTOR RNAi worms, suggesting that ADH-1 intersects with, or is permissive for, these pathways. This positioning is consistent with ADH-1 acting at a metabolic–proteostatic convergence point that integrates redox signaling, chaperone capacity, and inter-organ communication. Candidate distal effectors most consistent with our data include metabolites (e.g., acetate; Miskelly et al., 2019), changes in NAD$^+$/NADH ratios, and redox-active small molecules. While additional mediators cannot be excluded, our current data more strongly support metabolic and redox intermediates than peptide hormones as primary drivers.

From a translational perspective, these non-cell-autonomous benefits are especially compelling. In mammals, ADH1C—the ortholog of *C. elegans adh-1*—is secreted in liver-derived exosomes and has been detected in systemic circulation(Wang et al., 2022). Complementing this, human ADH1C has been detected in liver-derived exosomes and in

circulation(Yoshida et al., 2020), raising the prospect that ADH1 activation in one tissue could influence others via vesicle-associated or freely circulating metabolites in mammals.

In sum, ADH-1 overexpression confers broad protection across cellular systems by combining redox adaptation, proteostatic reinforcement, and inter-tissue exchange of beneficial effectors or signals. This integration positions ADH-1 as a regulator of healthspan architecture—capable not only of delaying decline in individual tissues but also of orchestrating organism-wide resilience. Future work should aim to map the upstream triggers of *adh-1* induction, define its metabolic and, possibly, secreted effectors, and determine whether its benefits can be mobilized pharmacologically or genetically in other species. These investigations will clarify whether ADH-1 represents a conserved axis for systemic geroprotection, and whether it can be leveraged to promote healthy aging in humans.

Summary Figure

Methods

C. elegans Strains and Maintenance

The full list of *C. elegans* strains used in this study is provided in the Key Resources Table.

For age-related phenotype assessments, *C. elegans* were synchronized by L1 arrest: hatchlings were incubated overnight (18–24 h) in incomplete S-buffer. Approximately 200 synchronized L1s were then dispensed onto 6-cm NGM plates seeded with 200 µL of freshly harvested OP50 and incubated at 20 °C until the young adult stage. At that point, worms were transferred to fresh NGM plates supplemented with 100 µg/mL FUdR to prevent progeny production. Phenotypic assessments were performed on day 1, 2, 8, 10, and/or 12 of adulthood, depending on the assay.

Fat Staining

Wild-type (N2) and ADH-1^{OE} (GMW20–22; see Key Resources Table) worms were grown and synchronized as described above. Oil Red O (ORO) staining was performed as previously described (Ke et al., 2018, 2021; Wählby et al., 2014). Briefly, day 1 and day 10 adults were washed four times in incomplete S-buffer, concentrated, and fixed by resuspension in 60% isopropanol. Fixed worms were stained with freshly prepared, filtered 60% ORO solution and incubated in a humid chamber at 25 °C overnight. Staining duration was empirically tested at 5, 10, and 20 h at 25 °C in a humid chamber.

Imaging was conducted on Nikon Eclipse Ti inverted microscope using a 10x Objective. Image acquisition was carried out with a DS-Ri2 Nikon camera and NIS software. Images were exported as RGB and processed in ImageJ/Fiji (fiji.sc). The blue channel was isolated; wild-type images were inverted and thresholded to capture worm bodies, and the resulting threshold parameters were then applied identically to all other conditions. Binary masks generated from this thresholding were analyzed with Analyze Particles to extract mean pixel intensity (MPI) per worm as the ORO signal. At least three independent biological replicates were performed.

RT-qPCR

Approximately 1,500 synchronized L1 worms were seeded per 10-cm RNAi plate containing control *E. coli* XU363. Upon reaching the young adult stage, worms were transferred to control or experimental RNAi plates supplemented with 100 µg/mL FUdR. On day 8 of adulthood, animals were harvested using a mesh to remove dead eggs, flash-frozen in liquid nitrogen, and stored at −80 °C until processing.

Total RNA was extracted from frozen worms with TRI Reagent (MRC, USA) according to the manufacturer's instructions. RNA purity and concentration were assessed by NanoDrop spectrophotometry. Three micrograms of RNA (3 µg) were used to synthesize complementary DNA (cDNA). Quantitative PCR was performed using cDNA, SYBR Green (Bio-Rad), and gene-specific primers (Table S1) on a real-time PCR thermal cycler (Bio-Rad, USA). Fold changes were calculated using the Pfaffl method (Pfaffl, 2001) , and statistical significance relative to the WT control was determined using an unpaired Student's t-test. At least three independent biological replicates were performed.

Pharyngeal Pumping Assay

Wild-type (N2) and ADH-1[OE] (GMW20; see Key Resources Table) worms were grown and synchronized as described in *C. elegans* Strains and Maintenance. Pharyngeal pumping was assessed on day 1 and day 8 of adulthood.

Sixty-second videos of the head region were acquired on a Zeiss Axio Zoom.V16 stereomicroscope (PlanNeoFluar Z 2.3×/0.57 FWD objective; ~112× total magnification) equipped with a digital camera. Videos were played back at 10 fps, and pumps were counted within multiple non-overlapping 30-s windows per animal. Because animals could transiently drift out of focus during a 60-s recording, pumps per minute were calculated by averaging the counts from the available 30-s windows and extrapolating to per-minute rates. At least three independent biological replicates were performed.

Chemoattractant assay

Wild-type (N2) and ADH-1[OE] (GMW20; see Key Resources Table) worms were grown and synchronized as described in *C. elegans* Strains and Maintenance. Chemotaxis assays were performed on unseeded NGM plates (10 cm).

Each plate was divided into four quadrants, a 0.5-cm diameter circle was marked at the center, and a single "solution point" was designated in each quadrant, positioned equidistant from one another and ≥2 cm from the center. Solutions of attractant (diacetyl in incomplete S-buffer) and control (incomplete S-buffer) were prepared.

On day 1 and day 10 of adulthood, worms were harvested by washing plates with incomplete S-buffer, washed four times to remove residual bacteria, and concentrated. Approximately 150 worms were deposited at the center of each plate, after which control and attractant solutions were pipetted onto their respective solution points (T0). After 1hr, , assays were analyzed following the protocol detailed in (Margie et al., 2013). At least three independent biological replicates were performed.

Paraquat Challenge

Wild-type (N2) and ADH-1^{OE} (GMW20; see Key Resources Table) worms were grown and synchronized as described in *C. elegans* Strains and Maintenance. Oxidative stress assays were performed as previously described (Mony et al., 2021; Senchuk et al., 2017)with minor modifications. Six-centimeter NGM plates were seeded with fresh OP50. Paraquat (methyl viologen) was added to a final concentration of 100 mM; plates were placed on an orbital shaker for 1 h to ensure even distribution and then dried in a biosafety hood for 20 min.

Day 1 and day 8 adults were transferred to paraquat plates by picking. Unless oth) and noted, ~20–30 worms were placed per plate, with ≥3 plates per condition (experimental replicates), and incubated at 20 °C. Survival was scored at regular intervals until at least 50% mortality was reached. Animals that crawled off the agar, ruptured, or displayed internal hatching were censored. At least three independent biological replicates were performed.

Thrashing Assay

Wild-type (N2) and ADH-1^{OE} (GMW20–22; see Key Resources Table) worms were grown and synchronized as described in *C. elegans* Strains and Maintenance. Thrashing was assessed in day 1 and day 12 adults.

Twenty-four–well plates were pre-filled with 2 mL S-buffer per well. Using a wire pick, individual worms were placed into separate wells directly into the buffer. After a 2-4 min acclimation (kept ≤5 min uniformly across animals), 60-s videos of each well were acquired sequentially on a ZEISS Axio Zoom.V16 stereomicroscope (PlanNeoFluar Z 2.3×/0.57 FWD objective; ~112× total magnification) using ZEISS ZEN software (RRID:SCR_013672). Movies were analyzed in ImageJ/Fiji (RRID:SCR_002285) (Schindelin et al., 2012) and the WrmTrck plugin(Nussbaum-Krammer et al., 2015) to quantify body bends per second (BBPS). A "body bend/thrash" was defined as one complete change in lateral bending of the mid-body (left→right→left or right→left→right). When acquisition frame rates differed between runs, bend counts were normalized to elapsed time to report BBPS. At least three independent biological replicates were performed.

Fluorescent Imaging and Processing

Imaging was performed on either (1) a Nikon Eclipse Ti inverted microscope equipped with a Yokogawa CSU-X1 spinning disk, a Hamamatsu ORCA-Flash 4.0 sCMOS camera, and 405/488/561/640-nm lasers using a Plan Apo 10x, or (2) a Leica Stellaris 8 Fluorescent Spinning Disk Confocal with a 63×/1.4 NA oil-immersion objective. GABAergic motor neurons and body-wall muscles were imaged on the Leica system. PolyQ aggregates, reference images of ADH-1 tissue localization, HyPer::YFP images, and *hsp-4*p::GFP images were acquired on the Nikon system.

For all imaging, worms were immobilized in 10 mM levamisole on 2% agarose pads.. For experiments requiring fluorescent imaging, Z-stacks were collected. Equal laser intensity and exposure time was used for each replicate of a respective experiment. Image processing was solely conducted in ImageJ/Fiji (fiji.sc) software. Maximum projections were generated in almost all cases. In some cases, excessive worm movements made maximum projections not possible. In those instances, the brightest single slices were chosen from each respective group. Images then had rolling background subtraction applied (radius of 50) except in cases where thresholding was used (eg. PolyQ imaging). Per worm data was collected via manual worm tracing in Fiji. Worm ROIs were drawn

from the first two intestinal cells to the tails of the animal This was done to minimize any false positive signal from the green pharynxes of ADH-1^{OE} animals. Animals were excluded if they were determined to be much younger than the majority of worms in the image (eg. smaller total area discrepancy and/or lack of presence of eggs). Due to the green pharynx ADH-1^{OE} have, blinding was not possible in most cases. Blinding was conducted in the muscle scoring analysis as muscle images were taken from worm midbodies (pharynx not visible) and the scoring system was more subjective so blinding was heavily prioritized. Then, properties of each ROI were measured using the Fiji "measure" function. Data analyses was then conducted in R (see Statistical Testing).

Muscle Integrity Assay

RW1596 and GMW031 (see Key Resources Table) *C. elegans* were grown and synchronized as described in *C. elegans* Strains and Maintenance. The mid-body region was imaged to avoid distortions near the vulva, head, and tail. Z-stacks were acquired on a Leica Stellaris 8 fluorescent confocal microscope using a 40× objective with 2 μm optical steps. Images were collected at days 1, 6, 8, and 10; data shown are from day-10 adults. Image acquisition and processing followed the workflow described in "Fluorescent Imaging and Processing". Images were blinded using Blinder (Cothren et al., 2018) prior to scoring, then categorized into three classes: healthy, intermediately damaged, or severely damaged as exemplified in Fig. 4D-F. Two independent biological replicates were performed.

Neuronal integrity Assay

CZ13799 and GMW028 (see Key Resources Table) worms were grown and synchronized as described in *C. elegans* Strains and Maintenance. Animals were immobilized in 10 mM levamisole on 2% agarose pads, and VD/DD GABAergic axonal commissures were imaged from head to tail on a Leica Stellaris 8 fluorescent spinning-disk confocal microscope using a 63×/1.4 NA oil-immersion objective. Z-stacks were acquired at 1 μm optical steps. For image processing, the workflow described in "Fluorescent Imaging and Processing" was followed. Specific to this assay, commissural axons were cropped away from surrounding tissue to prevent spurious detections. Neurodegeneration was

quantified in ImageJ/Fiji (fiji.sc) using the FindFoci plugin (Herbert et al., 2014). Axonal "stippling" events were operationally defined as punctate GFP spikes above the commissural background. FindFoci was initialized by establishing a baseline detection threshold on a manually curated training set of healthy commissures; this threshold and all analysis parameters were then fixed and applied uniformly to all images and conditions. For each worm, spike counts were computed per commissure and then averaged across all commissures to yield a single value (mean stippling events per commissure per worm). At least two independent biological replicates were performed.

Polyglutamine aggregate assessment

AM140 and GMW0024(see Key Resources Table) worms were grown and synchronized as described in *C. elegans* Strains and Maintenance. For imaging, animals were immobilized in 1 mM levamisole on 2% agarose pads. Z-stacks were acquired at .9 µm optical steps with a 20× objective on the Nikon spinning-disk system (Eclipse Ti + Yokogawa CSU-X1; Hamamatsu ORCA-Flash 4.0). Whole-animals were tiled and stitched in NIS-Elements (RRID:SCR_014329) using linear blending with 50% overlap.

Image analysis was performed in ImageJ/Fiji (RRID:SCR_002285)(Schindelin et al., 2012). Worm outlines were cropped to generate per-animal ROIs. Aggregates were enhanced for detection by local automatic thresholding (Otsu, default settings) benchmarked against WT animals and then applied to ADH-1^{OE} animals. The same threshold was applied to all images of within that replicate. Particles were then segmented with Analyze Particles (size: 2–500; other parameters at default). This size window was chosen to exclude GFP-positive pharynxes in ADH-1^{OE} animals and to avoid counting large diffuse GFP signals as large aggregates. All processing parameters were applied identically across conditions. Two independent biological replicates were performed.

Statistical Testing

All analyses and visualizations were performed in R (R Foundation) using RStudio (Posit). Core packages included: Tidyverse(*Tidyverse*, n.d.), ggplot2(*Authors and Citation*, n.d.), ggpubr(*Ggplot2 Based Publication Ready Plots*, n.d.), patchwork(*Getting Started*, n.d.), RColorBrewer(Neuwirth, 2022), survival(Therneau et al., 2024), ggsurvfit(Sjoberg et al.,

2024), dplyr(Wickham, François, et al., 2023), readxl(Wickham et al., 2025), infer(Bray et al., 2025), svglite(Wickham, Henry, et al., 2023), survminer(Kassambara et al., 2024), glue(Hester et al., 2024), ggtext(Wilke & Wiernik(@bmwiernik), 2022) , rstatix(Kassambara, 2023), knitr(Xie [aut et al., 2025) , stringr(Wickham, Software, et al., 2023), and extrafont(Chang, 2023). Data were assessed for normality (Shapiro–Wilk test). When assumptions were met, two-sided t-tests (Welch's correction when variances were unequal). When assumptions were violated, Wilcoxon/Mann–Whitney or Kruskal–Wallis tests were applied with appropriate multiple-comparison procedures. Survival data were analyzed by Kaplan–Meier estimates and log-rank tests. The specific test used for each comparison is stated in the corresponding figure legend.

Declaration of interests

The authors declare no competing interests.

References

Ables, G. P., Hens, J. R., & Nichenametla, S. N. (2016). Methionine restriction beyond life-span

extension. *Annals of the New York Academy of Sciences*, *1363*(1), 68–79.

https://doi.org/10.1111/nyas.13014

Ables, G. P., & Johnson, J. E. (2017). Pleiotropic responses to methionine restriction.

Experimental Gerontology, *94*, 83–88. https://doi.org/10.1016/j.exger.2017.01.012

Ables, G. P., Perrone, C. E., Orentreich, D., & Orentreich, N. (2012). Methionine-Restricted

C57BL/6J Mice Are Resistant to Diet-Induced Obesity and Insulin Resistance but Have

Low Bone Density. *PLoS ONE*, *7*(12), e51357.

https://doi.org/10.1371/journal.pone.0051357

Authors and Citation. (n.d.). Retrieved April 24, 2025, from

https://ggplot2.tidyverse.org/authors.html#citation

Back, P., Braeckman, B. P., & Matthijssens, F. (2012a). ROS in Aging Caenorhabditis elegans:

Damage or Signaling? *Oxidative Medicine and Cellular Longevity*, *2012*(1), 608478.

https://doi.org/10.1155/2012/608478

Back, P., Braeckman, B. P., & Matthijssens, F. (2012b). ROS in Aging Caenorhabditis elegans:

Damage or Signaling? *Oxidative Medicine and Cellular Longevity*, *2012*, 608478.

https://doi.org/10.1155/2012/608478

Bailey, A. P., Koster, G., Guillermier, C., Hirst, E. M. A., MacRae, J. I., Lechene, C. P., Postle,

A. D., & Gould, A. P. (2015). Antioxidant Role for Lipid Droplets in a Stem Cell Niche of

Drosophila. *Cell*, *163*(2), 340–353. https://doi.org/10.1016/j.cell.2015.09.020

Bartke, A., Wright, J. C., Mattison, J. A., Ingram, D. K., Miller, R. A., & Roth, G. S. (2001).

Extending the lifespan of long-lived mice. *Nature*, *414*(6862), 412–412.

https://doi.org/10.1038/35106646

Belousov, V. V., Fradkov, A. F., Lukyanov, K. A., Staroverov, D. B., Shakhbazov, K. S.,

Terskikh, A. V., & Lukyanov, S. (2006). Genetically encoded fluorescent indicator for

intracellular hydrogen peroxide. *Nature Methods*, *3*(4), 281–286.

https://doi.org/10.1038/nmeth866

Ben-Zvi, A., Miller, E. A., & Morimoto, R. I. (2009). Collapse of proteostasis represents an early

molecular event in Caenorhabditis elegans aging. *Proceedings of the National Academy*

of Sciences of the United States of America, *106*(35), 14914–14919.

https://doi.org/10.1073/pnas.0902882106

Blackwell, T. K., Steinbaugh, M. J., Hourihan, J. M., Ewald, C. Y., & Isik, M. (2015). SKN-1/Nrf,

stress responses, and aging in Caenorhabditis elegans. *Free Radical Biology &*

Medicine, *88*(Pt B), 290–301. https://doi.org/10.1016/j.freeradbiomed.2015.06.008

Branicky, R., Bénard, C., & Hekimi, S. (2000). Clk-1, mitochondria, and physiological rates.

BioEssays, *22*(1), 48–56. https://doi.org/10.1002/(SICI)1521-

1878(200001)22:1%253C48::AID-BIES9%253E3.0.CO;2-F

Bray, A., Ismay, C., Chasnovski, E., Couch [aut, S., cre, Baumer, B., Cetinkaya-Rundel, M.,

Laderas, T., Solomon, N., Hardin, J., Kim, A. Y., Fultz, N., Friedman, D., Cotton, R., &

Fannin, B. (2025). *infer: Tidy Statistical Inference* (Version 1.0.8) [Computer software].

https://cran.r-project.org/web/packages/infer/index.html

Calfon, M., Zeng, H., Urano, F., Till, J. H., Hubbard, S. R., Harding, H. P., Clark, S. G., & Ron,

D. (2002a). IRE1 couples endoplasmic reticulum load to secretory capacity by

processing the XBP-1 mRNA. *Nature*, *415*(6867), 92–96.

https://doi.org/10.1038/415092a

Calfon, M., Zeng, H., Urano, F., Till, J. H., Hubbard, S. R., Harding, H. P., Clark, S. G., & Ron,

D. (2002b). IRE1 couples endoplasmic reticulum load to secretory capacity by

processing the XBP-1 mRNA. *Nature*, *415*(6867), 92–96.

https://doi.org/10.1038/415092a

Campagnola, P. J., Millard, A. C., Terasaki, M., Hoppe, P. E., Malone, C. J., & Mohler, W. A.

(2002). Three-Dimensional High-Resolution Second-Harmonic Generation Imaging of

Endogenous Structural Proteins in Biological Tissues. *Biophysical Journal*, *82*(1), 493–

508. https://doi.org/10.1016/S0006-3495(02)75414-3

Cao, S. S., & Kaufman, R. J. (2014). Endoplasmic Reticulum Stress and Oxidative Stress in Cell

Fate Decision and Human Disease. *Antioxidants & Redox Signaling*, *21*(3), 396–413.

https://doi.org/10.1089/ars.2014.5851

Chang, W. (2023). *extrafont: Tools for Using Fonts* (Version 0.19) [Computer software].

https://cran.r-project.org/web/packages/extrafont/index.html

Chow, D. K., Glenn, C. F., Johnston, J. L., Goldberg, I. G., & Wolkow, C. A. (2006). Sarcopenia

in the Caenorhabditis elegans pharynx correlates with muscle contraction rate over

lifespan. *Experimental Gerontology*, *41*(3), 252–260.

https://doi.org/10.1016/j.exger.2005.12.004

Cothren, S. D., Meyer, J. N., & Hartman, J. H. (2018). Blinded Visual Scoring of Images Using

the Freely-available Software Blinder. *Bio-Protocol*, *8*(23), e3103.

https://doi.org/10.21769/BioProtoc.3103

Crimmins, E. M. (2015). Lifespan and Healthspan: Past, Present, and Promise. *The

Gerontologist*, *55*(6), 901–911. https://doi.org/10.1093/geront/gnv130

Cristina, D., Cary, M., Lunceford, A., Clarke, C., & Kenyon, C. (2009). A Regulated Response to

Impaired Respiration Slows Behavioral Rates and Increases Lifespan in Caenorhabditis

elegans. *PLOS Genetics*, *5*(4), e1000450. https://doi.org/10.1371/journal.pgen.1000450

Cruz-Jentoft, A. J., Bahat, G., Bauer, J., Boirie, Y., Bruyère, O., Cederholm, T., Cooper, C.,

Landi, F., Rolland, Y., Sayer, A. A., Schneider, S. M., Sieber, C. C., Topinkova, E.,

Vandewoude, M., Visser, M., & Zamboni, M. (2019). Sarcopenia: Revised European

consensus on definition and diagnosis. *Age and Ageing*, *48*(1), 16–31.

https://doi.org/10.1093/ageing/afy169

Cruz-Rodríguez, M., Chevet, E., & Muñoz-Pinedo, C. (2025). Glucose sensing and the unfolded

protein response. *The FEBS Journal*, *292*(14), 3581–3595.

https://doi.org/10.1111/febs.70113

De Haes, W., Frooninckx, L., Van Assche, R., Smolders, A., Depuydt, G., Billen, J., Braeckman,

B. P., Schoofs, L., & Temmerman, L. (2014). Metformin promotes lifespan through

mitohormesis via the peroxiredoxin PRDX-2. *Proceedings of the National Academy of

Sciences of the United States of America*, *111*(24), E2501–E2509.

https://doi.org/10.1073/pnas.1321776111

Dhondt, I., Verschuuren, C., Zečić, A., Loier, T., Braeckman, B. P., & De Vos, W. H. (2021).

Prediction of biological age by morphological staging of sarcopenia in Caenorhabditis

elegans. *Disease Models & Mechanisms*, *14*(11), dmm049169.

https://doi.org/10.1242/dmm.049169

Dillin, A., Hsu, A.-L., Arantes-Oliveira, N., Lehrer-Graiwer, J., Hsin, H., Fraser, A. G., Kamath, R.

S., Ahringer, J., & Kenyon, C. (2002). Rates of behavior and aging specified by

mitochondrial function during development. *Science (New York, N.Y.)*, *298*(5602), 2398–

2401. https://doi.org/10.1126/science.1077780

Ellis, Z. L., Siriwardhana, C. C., & Legleiter, J. A. (2023). Probing the morphological and

mechanical changes in a C. elegans model of Huntington's disease. *Biophysical Journal*,

122(3), 349a–350a. https://doi.org/10.1016/j.bpj.2022.11.1940

Gems, D., Sutton, A. J., Sundermeyer, M. L., Albert, P. S., King, K. V., Edgley, M. L., Larsen, P.

L., & Riddle, D. L. (1998). Two pleiotropic classes of daf-2 mutation affect larval arrest,

adult behavior, reproduction and longevity in Caenorhabditis elegans. *Genetics*, *150*(1),

129–155. https://doi.org/10.1093/genetics/150.1.129

Getting Started. (n.d.). Retrieved April 24, 2025, from https://patchwork.data-

imaginist.com/articles/patchwork.html

Ggplot2 Based Publication Ready Plots. (n.d.). Retrieved April 24, 2025, from

>   https://rpkgs.datanovia.com/ggpubr/

Ghaddar, A., Armingol, E., Huynh, C., Gevirtzman, L., Lewis, N. E., Waterston, R., & O'Rourke,

>   E. J. (2023). Whole-body gene expression atlas of an adult metazoan. *Science*

>   *Advances*, *9*(25), eadg0506. https://doi.org/10.1126/sciadv.adg0506

Ghaddar, A., Mony, V. K., Mishra, S., Berhanu, S., Johnson, J. C., Enriquez-Hesles, E.,

>   Harrison, E., Patel, A., Horak, M. K., Smith, J. S., & O'Rourke, E. J. (2023). Increased

>   Alcohol Dehydrogenase 1 activity promotes longevity. *Current Biology : CB*, *33*(6), 1036-

>   1046.e6. https://doi.org/10.1016/j.cub.2023.01.059

Giunti, S., Blanco, M. G., De Rosa, M. J., & Rayes, D. (n.d.). The ketone body β-

>   hydroxybutyrate ameliorates neurodevelopmental deficits in the GABAergic system of

>   daf-18/PTEN Caenorhabditis elegans mutants. *eLife*, *13*, RP94520.

>   https://doi.org/10.7554/eLife.94520

Giunti, S., Blanco, M. G., De Rosa, M. J., & Rayes, D. (2024). The ketone body β-

>   hydroxybutyrate ameliorates neurodevelopmental deficits in the GABAergic system of

>   daf-18/PTEN Caenorhabditis elegans mutants. *eLife*, *13*, RP94520.

>   https://doi.org/10.7554/eLife.94520

Gjorgjieva, J., Biron, D., & Haspel, G. (2014). Neurobiology of Caenorhabditis elegans

>   Locomotion: Where Do We Stand? *Bioscience*, *64*(6), 476–486.

>   https://doi.org/10.1093/biosci/biu058

Glenn, C. F., Chow, D. K., David, L., Cooke, C. A., Gami, M. S., Iser, W. B., Hanselman, K. B.,

>   Goldberg, I. G., & Wolkow, C. A. (2004). Behavioral deficits during early stages of aging

>   in Caenorhabditis elegans result from locomotory deficits possibly linked to muscle

>   frailty. *The Journals of Gerontology. Series A, Biological Sciences and Medical*

>   *Sciences*, *59*(12), 1251–1260. https://doi.org/10.1093/gerona/59.12.1251

Hajdu-Cronin, Y. M., Chen, W. J., & Sternberg, P. W. (2004). The L-Type Cyclin CYL-1 and the

 Heat-Shock-Factor HSF-1 Are Required for Heat-Shock-Induced Protein Expression in

 Caenorhabditis elegans. *Genetics*, *168*(4), 1937–1949.

 https://doi.org/10.1534/genetics.104.028423

Heidler, T., Hartwig, K., Daniel, H., & Wenzel, U. (2010). Caenorhabditiselegans lifespan

 extension caused by treatment with an orally active ROS-generator is dependent on

 DAF-16 and SIR-2.1. *Biogerontology*, *11*(2), 183–195. https://doi.org/10.1007/s10522-

 009-9239-x

Henis-Korenblit, S., Zhang, P., Hansen, M., McCormick, M., Lee, S.-J., Cary, M., & Kenyon, C.

 (2010). Insulin/IGF-1 signaling mutants reprogram ER stress response regulators to

 promote longevity. *Proceedings of the National Academy of Sciences*, *107*(21), 9730–

 9735. https://doi.org/10.1073/pnas.1002575107

Herbert, A. D., Carr, A. M., & Hoffmann, E. (2014). FindFoci: A Focus Detection Algorithm with

 Automated Parameter Training That Closely Matches Human Assignments, Reduces

 Human Inconsistencies and Increases Speed of Analysis. *PLOS ONE*, *9*(12), e114749.

 https://doi.org/10.1371/journal.pone.0114749

Herndon, L. A., Schmeissner, P. J., Dudaronek, J. M., Brown, P. A., Listner, K. M., Sakano, Y.,

 Paupard, M. C., Hall, D. H., & Driscoll, M. (2002). Stochastic and genetic factors

 influence tissue-specific decline in ageing C. elegans. *Nature*, *419*(6909), 808–814.

 https://doi.org/10.1038/nature01135

Hester, J., Bryan, J., Software, P., & PBC. (2024). *glue: Interpreted String Literals* (Version

 1.8.0) [Computer software]. https://cran.r-project.org/web/packages/glue/index.html

Hetz, C., & Papa, F. R. (2018). The Unfolded Protein Response and Cell Fate Control.

 Molecular Cell, *69*(2), 169–181. https://doi.org/10.1016/j.molcel.2017.06.017

Hihi, A. K., Gao, Y., & Hekimi, S. (2002). Ubiquinone Is Necessary for Caenorhabditis

    elegansDevelopment at Mitochondrial and Non-mitochondrial Sites *. *Journal of*

    Biological Chemistry, *277*(3), 2202–2206. https://doi.org/10.1074/jbc.M109034200

Hipp, M. S., Kasturi, P., & Hartl, F. U. (2019). The proteostasis network and its decline in

    ageing. *Nature Reviews Molecular Cell Biology*, *20*(7), 421–435.

    https://doi.org/10.1038/s41580-019-0101-y

Honda, Y., & Honda, S. (1999). The daf-2 gene network for longevity regulates oxidative stress

    resistance and Mn-superoxide dismutase gene expression in Caenorhabditis elegans.

    The FASEB Journal, *13*(11), 1385–1393. https://doi.org/10.1096/fasebj.13.11.1385

Hosono, R. (1978). Age dependent changes in the behavior of Caenorhabditis elegans on

    attraction to Escherichia coli. *Experimental Gerontology*, *13*(1–2), 31–36.

    https://doi.org/10.1016/0531-5565(78)90027-x

Hosono, R., Sato, Y., Aizawa, S. I., & Mitsui, Y. (1980). Age-dependent changes in mobility and

    separation of the nematode Caenorhabditis elegans. *Experimental Gerontology*, *15*(4),

    285–289. https://doi.org/10.1016/0531-5565(80)90032-7

Hotamisligil, G. S. (2010). Endoplasmic Reticulum Stress and the Inflammatory Basis of

    Metabolic Disease. *Cell*, *140*(6), 900–917. https://doi.org/10.1016/j.cell.2010.02.034

Hsu, A.-L., Feng, Z., Hsieh, M.-Y., & Xu, X. Z. S. (2009). Identification by machine vision of the

    rate of motor activity decline as a lifespan predictor in C. elegans. *Neurobiology of*

    Aging, *30*(9), 1498–1503. https://doi.org/10.1016/j.neurobiolaging.2007.12.007

Hu, Q., D'Amora, D. R., MacNeil, L. T., Walhout, A. J. M., & Kubiseski, T. J. (2018). The

    Caenorhabditis elegans Oxidative Stress Response Requires the NHR-49 Transcription

    Factor. *G3: Genes|Genomes|Genetics*, *8*(12), 3857–3863.

    https://doi.org/10.1534/g3.118.200727

Huang, C., Xiong, C., & Kornfeld, K. (2004). Measurements of age-related changes of

    physiological processes that predict lifespan of Caenorhabditis elegans. *Proceedings of*

the *National Academy of Sciences of the United States of America*, *101*(21), 8084–8089. https://doi.org/10.1073/pnas.0400848101

Huang, X., Cheng, H.-J., Tessier-Lavigne, M., & Jin, Y. (2002). MAX-1, a Novel PH/MyTH4/FERM Domain Cytoplasmic Protein Implicated in Netrin-Mediated Axon Repulsion. *Neuron*, *34*(4), 563–576. https://doi.org/10.1016/S0896-6273(02)00672-4

Ishii, N., Takahashi, K., Tomita, S., Keino, T., Honda, S., Yoshino, K., & Suzuki, K. (1990). A methyl viologen-sensitive mutant of the nematode Caenorhabditis elegans. *Mutation Research*, *237*(3–4), 165–171. https://doi.org/10.1016/0921-8734(90)90022-j

Johnson, T. E., Conley, W. L., & Keller, M. L. (1988). Long-lived lines of Caenorhabditis elegans can be used to establish predictive biomarkers of aging. *Experimental Gerontology*, *23*(4–5), 281–295. https://doi.org/10.1016/0531-5565(88)90031-9

Kano, K. (2013). Dwarfism, Mouse. In S. Maloy & K. Hughes (Eds.), *Brenner's Encyclopedia of Genetics (Second Edition)* (pp. 435–437). Academic Press. https://doi.org/10.1016/B978-0-12-374984-0.00455-1

Kassambara, A. (2023). *rstatix: Pipe-Friendly Framework for Basic Statistical Tests* (Version 0.7.2) [Computer software]. https://cran.r-project.org/web/packages/rstatix/index.html

Kassambara, A., Kosinski, M., Biecek, P., & Fabian, S. (2024). *survminer: Drawing Survival Curves using "ggplot2"* (Version 0.5.0) [Computer software]. https://cran.r-project.org/web/packages/survminer/index.html

Kutzner, C. E., Bauer, K. C., Lackmann, J.-W., Acton, R. J., Sarkar, A., Pokrzywa, W., & Hoppe, T. (2024). Optogenetic induction of mechanical muscle stress identifies myosin regulatory ubiquitin ligase NHL-1 in C. elegans. *Nature Communications*, *15*, 6879. https://doi.org/10.1038/s41467-024-51069-3

Labbadia, J., & Morimoto, R. I. (2014). Proteostasis and longevity: When does aging really begin? *F1000Prime Reports*, *6*, 7. https://doi.org/10.12703/P6-7

Lee, S.-J., Hwang, A. B., & Kenyon, C. (2010). Inhibition of respiration extends C. elegans'

lifespan via reactive oxygen species that increase HIF-1 activity. *Current Biology : CB*,

20(23), 2131–2136. https://doi.org/10.1016/j.cub.2010.10.057

Liu, J., Zhang, B., Lei, H., Feng, Z., Liu, J., Hsu, A.-L., & Xu, X. Z. S. (2013). Functional Aging in

the Nervous System Contributes to Age-Dependent Motor Activity Decline in C. elegans.

Cell Metabolism, *18*(3), 392–402. https://doi.org/10.1016/j.cmet.2013.08.007

Malhotra, J. D., & Kaufman, R. J. (2007). Endoplasmic Reticulum Stress and Oxidative Stress:

A Vicious Cycle or a Double-Edged Sword? *Antioxidants & Redox Signaling*, *9*(12),

2277–2294. https://doi.org/10.1089/ars.2007.1782

Margie, O., Palmer, C., & Chin-Sang, I. (2013). C. elegans Chemotaxis Assay. *Journal of

Visualized Experiments : JoVE*, *74*, 50069. https://doi.org/10.3791/50069

Meller, A., & Shalgi, R. (2021). The aging proteostasis decline: From nematode to human.

Experimental Cell Research, *399*(2), 112474.

https://doi.org/10.1016/j.yexcr.2021.112474

Migliaccio, E., Giorgio, M., Mele, S., Pelicci, G., Reboldi, P., Pandolfi, P. P., Lanfrancone, L., &

Pelicci, P. G. (1999). The p66shc adaptor protein controls oxidative stress response and

life span in mammals. *Nature*, *402*(6759), 309–313. https://doi.org/10.1038/46311

Molotkov, A., Fan, X., Deltour, L., Foglio, M. H., Martras, S., Farrés, J., Parés, X., & Duester, G.

(2002). Stimulation of retinoic acid production and growth by ubiquitously expressed

alcohol dehydrogenase Adh3. *Proceedings of the National Academy of Sciences of the

United States of America*, *99*(8), 5337–5342. https://doi.org/10.1073/pnas.082093299

Mony, V. K., Drangowska-Way, A., Albert, R., Harrison, E., Ghaddar, A., Horak, M. K., Ke, W., &

O'Rourke, E. J. (2021). Context-specific regulation of lysosomal lipolysis through

network-level diverting of transcription factor interactions. *Proceedings of the National

Academy of Sciences of the United States of America*, *118*(41), e2104832118.

https://doi.org/10.1073/pnas.2104832118

Morley, J. F., Brignull, H. R., Weyers, J. J., & Morimoto, R. I. (2002). The threshold for polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in Caenorhabditis elegans. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(16), 10417–10422. https://doi.org/10.1073/pnas.152161099

Morley, J. F., & Morimoto, R. I. (2004). Regulation of longevity in Caenorhabditis elegans by heat shock factor and molecular chaperones. *Molecular Biology of the Cell*, *15*(2), 657–664. https://doi.org/10.1091/mbc.e03-07-0532

Musa, M., Dionisio, P. A., Casqueiro, R., Milosevic, I., Raimundo, N., & Krisko, A. (2022). Lack of peroxisomal catalase affects heat shock response in Caenorhabditis elegans. *Life Science Alliance*, *6*(1), e202201737. https://doi.org/10.26508/lsa.202201737

Neuwirth, E. (2022). *RColorBrewer: ColorBrewer Palettes* (Version 1.1-3) [Computer software]. https://cran.r-project.org/web/packages/RColorBrewer/index.html

Nussbaum-Krammer, C. I., Neto, M. F., Brielmann, R. M., Pedersen, J. S., & Morimoto, R. I. (2015). Investigating the Spreading and Toxicity of Prion-like Proteins Using the Metazoan Model Organism C. elegans. *Journal of Visualized Experiments : JoVE*, *95*, 52321. https://doi.org/10.3791/52321

O'Rourke, E. J., Soukas, A. A., Carr, C. E., & Ruvkun, G. (2009). C. elegans major fats are stored in vesicles distinct from lysosome-related organelles. *Cell Metabolism*, *10*(5), 430–435. https://doi.org/10.1016/j.cmet.2009.10.002

Özcan, U., Cao, Q., Yilmaz, E., Lee, A.-H., Iwakoshi, N. N., Özdelen, E., Tuncman, G., Görgün, C., Glimcher, L. H., & Hotamisligil, G. S. (2004). Endoplasmic Reticulum Stress Links Obesity, Insulin Action, and Type 2 Diabetes. *Science*, *306*(5695), 457–461. https://doi.org/10.1126/science.1103160

Pan, C.-L., Peng, C.-Y., Chen, C.-H., & McIntire, S. (2011). Genetic analysis of age-dependent defects of the Caenorhabditis elegans touch receptor neurons. *Proceedings of the*

National Academy of Sciences, *108*(22), 9274–9279.

https://doi.org/10.1073/pnas.1011711108

Petriv, O. I., & Rachubinski, R. A. (2004). Lack of peroxisomal catalase causes a progeric

phenotype in Caenorhabditis elegans. *The Journal of Biological Chemistry*, *279*(19),

19996–20001. https://doi.org/10.1074/jbc.M400207200

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT–PCR.

Nucleic Acids Research, *29*(9), e45. https://doi.org/10.1093/nar/29.9.e45

Prahlad, V., Cornelius, T., & Morimoto, R. I. (2008). Regulation of the Cellular Heat Shock

Response in Caenorhabditis elegans by Thermosensory Neurons. *Science*, *320*(5877),

811–814. https://doi.org/10.1126/science.1156093

Raamsdonk, J. M. V., & Hekimi, S. (2009). Deletion of the Mitochondrial Superoxide Dismutase

sod-2 Extends Lifespan in Caenorhabditis elegans. *PLOS Genetics*, *5*(2), e1000361.

https://doi.org/10.1371/journal.pgen.1000361

Ristow, M., & Schmeisser, S. (2011). Extending life span by increasing oxidative stress. *Free

Radical Biology and Medicine*, *51*(2), 327–336.

https://doi.org/10.1016/j.freeradbiomed.2011.05.010

Ristow, M., & Zarse, K. (2010). How increased oxidative stress promotes longevity and

metabolic health: The concept of mitochondrial hormesis (mitohormesis). *Experimental

Gerontology*, *45*(6), 410–418. https://doi.org/10.1016/j.exger.2010.03.014

Rowe, J. W., & Kahn, R. L. (1987). Human aging: Usual and successful. *Science (New York,

N.Y.)*, *237*(4811), 143–149. https://doi.org/10.1126/science.3299702

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V.,

Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji—An Open Source platform for

biological image analysis. *Nature Methods*, *9*(7), 10.1038/nmeth.2019.

https://doi.org/10.1038/nmeth.2019

Schulz, T. J., Zarse, K., Voigt, A., Urban, N., Birringer, M., & Ristow, M. (2007). Glucose

    Restriction Extends *Caenorhabditis elegans* Life Span by Inducing Mitochondrial

    Respiration and Increasing Oxidative Stress. *Cell Metabolism*, *6*(4), 280–293.

    https://doi.org/10.1016/j.cmet.2007.08.011

Senchuk, M. M., Dues, D. J., & Van Raamsdonk, J. M. (2017). Measuring Oxidative Stress in

    Caenorhabditis elegans:Paraquat and Juglone Sensitivity Assays. *Bio-Protocol*, *7*(1),

    e2086. https://doi.org/10.21769/BioProtoc.2086

Shemesh, N., Shai, N., & Ben-Zvi, A. (2013). Germline stem cell arrest inhibits the collapse of

    somatic proteostasis early in Caenorhabditis elegans adulthood. *Aging Cell*, *12*(5), 814–

    822. https://doi.org/10.1111/acel.12110

Sjoberg, D. D., Baillie, M., Fruechtenicht, C., Haesendonckx, S., & Treis, T. (2024). *ggsurvfit:*

    Flexible Time-to-Event Figures (Version 1.1.0) [Computer software]. https://cran.r-

    project.org/web/packages/ggsurvfit/index.html

Tank, E. M. H., Rodgers, K. E., & Kenyon, C. (2011). Spontaneous Age-Related Neurite

    Branching in Caenorhabditis elegans. *Journal of Neuroscience*, *31*(25), 9279–9288.

    https://doi.org/10.1523/JNEUROSCI.6606-10.2011

Taylor, R. C., & Dillin, A. (2013a). XBP-1 Is a Cell-Nonautonomous Regulator of Stress

    Resistance and Longevity. *Cell*, *153*(7), 1435–1447.

    https://doi.org/10.1016/j.cell.2013.05.042

Taylor, R. C., & Dillin, A. (2013b). XBP-1 Is a Cell-Nonautonomous Regulator of Stress

    Resistance and Longevity. *Cell*, *153*(7), 1435–1447.

    https://doi.org/10.1016/j.cell.2013.05.042

Taylor, R. C., & Dillin, A. (2013c). XBP-1 Is a Cell-Nonautonomous Regulator of Stress

    Resistance and Longevity. *Cell*, *153*(7), 1435–1447.

    https://doi.org/10.1016/j.cell.2013.05.042

Therneau, T. M., until 2009), T. L. (original S.->R port and R. maintainer, Elizabeth, A., &

Cynthia, C. (2024). *survival: Survival Analysis* (Version 3.8-3) [Computer software].

https://cran.r-project.org/web/packages/survival/index.html

Tidyverse. (n.d.). Retrieved April 24, 2025, from https://www.tidyverse.org/

van Oosten-Hawle, P., Porter, R. S., & Morimoto, R. I. (2013). Regulation of Organismal

Proteostasis by Transcellular Chaperone Signaling. *Cell*, *153*(6), 1366–1378.

https://doi.org/10.1016/j.cell.2013.05.015

Wang, M., & Kaufman, R. J. (2016). Protein misfolding in the endoplasmic reticulum as a

conduit to human disease. *Nature*, *529*(7586), 326–335.

https://doi.org/10.1038/nature17041

Wickham, H., Bryan, J., Posit, attribution), P. (Copyright holder of all R. code and all C. code

without explicit copyright, code), M. K. (Author of included R., code), K. V. (Author of

included libxls, code), C. L. (Author of included libxls, code), B. C. (Author of included

libxls, code), D. H. (Author of included libxls, & code), E. M. (Author of included libxls.

(2025). *readxl: Read Excel Files* (Version 1.4.5) [Computer software]. https://cran.r-

project.org/web/packages/readxl/index.html

Wickham, H., François, R., Henry, L., Müller, K., Vaughan, D., Software, P., & PBC. (2023).

dplyr: A Grammar of Data Manipulation (Version 1.1.4) [Computer software].

https://cran.r-project.org/web/packages/dplyr/index.html

Wickham, H., Henry, L., Pedersen, T. L., Luciani, T. J., Decorde, M., Lise, V., code), T. P. (Early

line dashing, code), D. G. (Line dashing code and early raster, implementation), Y. Q.

(Improved styles; polypath, code), H. M. (Opacity, Posit, & PBC. (2023). *svglite: An*

"SVG" Graphics Device (Version 2.1.3) [Computer software]. https://cran.r-

project.org/web/packages/svglite/index.html

Wickham, H., Software, P., & PBC. (2023). *stringr: Simple, Consistent Wrappers for Common String Operations* (Version 1.5.1) [Computer software]. https://cran.r-project.org/web/packages/stringr/index.html

Wilke, C. O., & Wiernik (@bmwiernik), B. M. (2022). *ggtext: Improved Text Rendering Support for "ggplot2"* (Version 0.1.2) [Computer software]. https://cran.r-project.org/web/packages/ggtext/index.html

Xie [aut, Y., cre] (https://yihui.org), Sarma, A., Vogt, A., Andrew, A., Zvoleff, A., Al-Zubaidi, A., http://www.andre-simon.de), A. S. (the C. files under inst/themes/ were derived from the H. package, Atkins, A., Wolen, A., Manton, A., Yasumoto, A., Baumer, B., Diggs, B., Zhang, B., Yapparov, B., Pereira, C., Dervieux, C., Hall, D., … PBC. (2025). *knitr: A General-Purpose Package for Dynamic Report Generation in R* (Version 1.50) [Computer software]. https://cran.r-project.org/web/packages/knitr/index.html

Yang, W., & Hekimi, S. (2010a). A Mitochondrial Superoxide Signal Triggers Increased Longevity in Caenorhabditis elegans. *PLoS Biology*, *8*(12), e1000556. https://doi.org/10.1371/journal.pbio.1000556

Yang, W., & Hekimi, S. (2010b). Two modes of mitochondrial dysfunction lead independently to lifespan extension in Caenorhabditis elegans. *Aging Cell*, *9*(3), 433–447. https://doi.org/10.1111/j.1474-9726.2010.00571.x

Yun, J., & Finkel, T. (2014). Mitohormesis. *Cell Metabolism*, *19*(5), 757–766. https://doi.org/10.1016/j.cmet.2014.01.011

Zarse, K., Schmeisser, S., Groth, M., Priebe, S., Beuster, G., Kuhlow, D., Guthke, R., Platzer, M., Kahn, C. R., & Ristow, M. (2012). Impaired Insulin/IGF1 Signaling Extends Life Span by Promoting Mitochondrial L-Proline Catabolism to Induce a Transient ROS Signal. *Cell Metabolism*, *15*(4), 451–465. https://doi.org/10.1016/j.cmet.2012.02.013

Zhang, K., & Kaufman, R. J. (2008). From endoplasmic-reticulum stress to the inflammatory response. *Nature*, *454*(7203), 455–462. https://doi.org/10.1038/nature07203

Zhao, Y., Wu, L., Yue, X., Zhang, C., Wang, J., Li, J., Sun, X., Zhu, Y., Feng, Z., & Hu, W.

(2018). A polymorphism in the tumor suppressor p53 affects aging and longevity in

mouse models. *eLife, 7*, e34701. https://doi.org/10.7554/eLife.34701