Bacteriophage genome-wide transposon mutagenesis

Alex Chan[1,+], Wearn-Xin Yee[1,+], Deepto Mozumdar[1,#], Claire Kokontis[1,#], Matias Rojas-Montero[1], Li Yuping[1,X], Ying Yang[1], Joseph Bondy-Denomy[1,*]

Affiliations
[1]Department of Microbiology & Immunology, University of California, San Francisco
+ Co-first authors
# Co-second authors
x Current affiliation: Biozentrum, University of Basel
*Correspondence: joseph.bondy-denomy@ucsf.edu

Abstract

Bacteriophages have genomes that span a wide size range, are densely packed with coding sequences, and frequently encode genes of unknown function. Classical forward genetics has defined essential genes for phage replication in a few model systems but remains laborious and non-scalable. Unbiased functional genomics approaches are therefore needed for phages, particularly for large lytic phages. Here, we develop a phage transposon sequencing (TnSeq) platform that uses the *mariner* transposase to insert an anti-CRISPR selectable marker into phage genomes. CRISPR-Cas13a–based enrichment of transposed phages followed by pooled sequencing identifies both fitness-conferring and dispensable genes. Using the *Pseudomonas aeruginosa*-infecting nucleus-forming jumbo phage ΦKZ (280,334 bp; 371 predicted genes) as a model, we show that ~110 genes are fitness-conferring via phage TnSeq. These include conserved essential genes involved in phage nucleus formation, protein trafficking, transcription, DNA replication, and virion assembly. We also isolate hundreds of individual phages with insertions in non-essential genes and reveal conditionally essential genes that are specifically required in clinical isolates, at environmental temperature, or in the presence of a defensive nuclease. Phage TnSeq is a facile, scalable technology that can define essential phage genes and generate knockouts in all non-essential genes in a single experiment, enabling conditional genetic screens in phages and providing a broadly applicable resource for phage functional genomics.

Introduction

Bacteriophage genomes are rich in genes of unknown function and exhibit extensive genetic and proteomic diversity. Many fundamental discoveries likely remain hidden in uncharacterized phage genes[1]. However, few tools exist to enable unbiased genome-wide studies to bring phages into the functional genomics era. Most challenging perhaps are large lytic phages (e.g. jumbo phages with genomes >200 kb), where most genes are of unknown function[2]. For most of these phages, even the basic inventory of genes essential for replication is unknown.

Here we focus on ΦKZ, a model member of the *Chimalliviridae* family with distinctive infection biology[3,4]. ΦKZ has many notable features, including two distinct multi-subunit RNA polymerase complexes[5–8], a lipid-based early phage infection vesicle[9–11], and a protein-based phage "nucleus"[12,13] that enables resistance to nucleases[14]. It remains unclear which genes are strictly required for this core biology, and which are dispensable but conditionally important, for example in specific hosts or environments. A method that could both rapidly identify all essential genes and simultaneously generate a library of knockouts in all other genes would greatly accelerate mechanistic dissection of such phages.

Targeted phage genome manipulation tools such as homology-directed recombination (HDR) and CRISPR-Cas counter-selection have been widely used[15] albeit at low throughput. Recombineering is also effective and has recently been adapted to genome-wide mutagenesis[16] and *in vitro* genome manipulation and rebooting have expanded the toolkit further[17]. RNA-targeting Cas13 has been particularly useful for selecting for spontaneous[14,18] or HDR-derived phage mutants[19]. To enable facile and scalable bacteriophage genome modification with the ease afforded to bacteria by transposon insertion sequencing (TnSeq), we sought to develop an analogous genome-wide insertional mutagenesis method for phages. Here, we deploy *mariner* transposition in the nucleus-forming jumbo phage ΦKZ, using an anti-CRISPR (*acrVIA1*) selectable marker and CRISPR-Cas13a–mediated enrichment to isolate thousands of independently transposed phages. High-throughput mapping of insertion sites reveals essential genes at genome scale and yields a large collection of phage mutants for conditional phenotypic screens.

Genome-wide knockout approaches that generate permanent changes in phage genomes, such as the one presented here, are complementary to recently developed protein knockdown approaches based on antisense oligomers (ASOs) and dCas13d-mediated translation interference (CRISPRi-ART)[20,21]. These perturbation tools are powerful for studying essential genes that cannot be deleted outright. By contrast, phage TnSeq requires no large-scale oligo or guide RNA design, *mariner* transposase is active in many bacterial hosts[22], and inserts transposons semi-randomly[23,24]. Together, these features should enable broad adoption of phage TnSeq as an easy-to-implement, low-cost genome-wide mutagenesis platform which can, in principle, be executed with any combination of a strong phage defense system and cognate anti-defense molecule.

Results
Establishing transposition in a nucleus-forming jumbo phage

ΦKZ protects its genome within a protein-based nucleus-like structure during infection and selectively imports proteins. To enable transposition inside this compartment, we fused the

Mariner transposase to an sfCherry fluorescent reporter and to the nucleus-localized phage
protein Nlp1[12,25] (gp152) (Figure 1A). Upon induction of this fusion and infection with ΦKZ,
fluorescence microscopy revealed co-localization of sfCherry–Mariner–Nlp1 with DAPI-
stained phage DNA, indicating import of the transposase into the phage nucleus (Figure 1B).

The initial transposon donor (V1) carried a selectable *acrVIA1* cassette flanked by a ΦKZ
consensus early promoter[8,26] and a terminator, to drive early *acrVIA1* expression (Table S1).
For selection, we co-expressed *Listeria seeligeri* Cas13a (LseCas13a)[19,27] with two crRNAs:
one targeting *orf120* (major capsid) and one targeting *orf55* (a nvRNAP subunit). Each
crRNA alone strongly restricted phage infection but was individually escapable (Figure 1C).
We therefore designed a single repeat–spacer–repeat–spacer array expressing both
crRNAs, which efficiently blocked infections by previously isolated escaper phages (Figure
S1A). No escaper mutants were detected (limit of detection (LOD) <2.5 x 10^{-10} pfu/mL), so
this dual crRNA array was used for all subsequent selections (Figure 1C, Figure S1B).

To generate a transposed phage population, we infected a log-phase culture expressing
Mariner–Nlp1 with ΦKZ for 16 h and then selected output phages on the dual-crRNA
Cas13a strain using full-plate infections. Single plaques were isolated, and *acrVIA1* was
detected by PCR (Figure S1C). Plaque formation required both the transposase and the
transposon donor (Figure 1D–E, Figure S1D). Whole-genome sequencing of four
independently isolated phages confirmed bona fide transposon insertions in *orf303*, *orf208*,
orf241, and *orf252*. In each case, two inverted repeats flanked an insertion at a TA
dinucleotide in the ΦKZ genome (schematized in Figure S1C).

Transposition using the V1 donor occurred at a rate of ~1 in 3 × 10□ phages, demonstrating
that transposition is inefficient but feasible in a large population (Figure 1E). No plaques
were detected when the donor plasmid was present without transposase (LOD: 1.3 x 10^{-10}
pfu/mL (Figure 1E)). A second donor (V2) lacking the early promoter yielded slightly lower
efficiency (1 in 6.7 × 10□ phages; Figure 1E), consistent with *acrVIA1* expression relying on
endogenous promoters. To facilitate compatibility with bacterial Tn libraries, we engineered a
third donor (V3, derived from V1) that altered the transposon inverted repeat junctions to
introduce orthogonal primer binding sites. Transposition efficiency with V3 (1 in 7 × 10□
phages) was similar to V1 and V2 and remained strictly dependent on the Nlp1–Mariner
fusion (LOD: 3.9 x 10^{-11} pfu/mL). Transposition was detected across a range of input phage
concentrations and was most efficient at a relatively low input (~2 × 10□ pfu/mL; Figure
S1E). We therefore used the V3 donor to generate a pool of mutants for insertion
sequencing.

A ΦKZ phage TnSeq library with genome-wide coverage
To build independent pools of transposon mutants, we performed ten biological replicate
transposition infections (Figure S2). Mutants from each reaction were enriched on the
Cas13a counter-selection host, and fractions from all ten pools were combined to generate a
single mutant library. We isolated DNA, prepared sequencing libraries, and amplified
transposon–genome junctions for Illumina sequencing (Methods). We obtained ~3.4 million
high-quality reads, 90% of which contained transposon sequence. Of these, 55.7% mapped
to the ΦKZ genome and the remainder to the V3 donor plasmid. Across the ΦKZ genome,
6,667 unique insertion sites were identified (Figure 2A), corresponding to a mean of one
insertion every 42 bp over the 280,334 bp genome.

To define confident gene insertions, we first examined genes expected to be essential. Many
insertion sites yielded low sequencing depth, suggesting that insertion occurred but rendered
phages unfit. We therefore required >4 reads per insertion for downstream analysis. In
addition, insertions within the first or last 10% of a gene were excluded to avoid
misclassifying overlapping genes or insertions that might not fully disrupt gene function.
Using these criteria, ~30% of ΦKZ genes lacked disruptive insertions, whereas ~70%
tolerated insertions (Figure S3).

ΦKZ phage TnSeq identifies essential genes
As an initial accuracy check, we focused on large genes, which should, by chance,
accumulate many insertions unless they are essential. Tailed phage genomes often encode
large essential structural and polymerase proteins. In our ΦKZ TnSeq dataset, all 16 genes
>2,154 bp and 30 of 34 genes >1,560 bp lacked disruptive insertions (Figure 2B), strongly
indicating essentiality given the genome-wide insertion frequency (one unique insertion
every 42 bp). Below, we highlight examples of essential genes in tail, capsid, and conserved
non-virion modules with all features reported in Figure 3A

Tail genes. The largest gene in ΦKZ, *orf181*, encodes a lysozyme-containing tail protein
likely serving as the tape measure[28]. *orf181* and its neighbor *orf182* (encoding a baseplate
protein) both lack transposon insertions (Figure 2B). Given the 6,714-bp size of *orf181*, its
complete lack of insertions makes it a high-confidence essential gene. Other tail-associated
genes lacking Tn insertions include putative tail fibers (gp145, gp146), a putative baseplate
wedge protein (gp87), a paralogous baseplate protein family (gp131–135)[28], the tail
sheath[29,30] (gp29), and the tail tube (gp30) (Figure 2B). Virion-associated proteins of
unknown function such as gp27 (DUF7193) and a low-copy virion protein gp128 also lack
insertions. Virion localization and functional annotations were supported by previously
published mass spectrometry datasets[31–33] and HHpred/PSI-BLAST analyses (Figure 3A).

Head genes. Several large genes that appear essential by phage TnSeq encode for capsid
assembly, DNA packaging, or injected proteins. These include the major capsid protein
(gp120)[32,34], the portal (gp129)[32,35], and the large terminase (gp25)[32] (Figure 2B). Mass
spectrometry of a tail-less ΦKZ mutant revealed >50 distinct polypeptides incorporated into
the head, many of which are proteolytically processed by an essential (per phage TnSeq)
head protease, gp175[32,36]. Many head proteins (gp92, 93, 94, 95, 96, 97, 153, 162, 163;
PF12699.12 (C-terminus)[10,32]) are members of a paralogous family[10,33] and some of them are
injected during infection[10]. Interestingly, several paralogous genes (e.g., *orf92, 93, 96, 162,*
163) readily tolerate insertions, whereas genes encoding other high-copy inner body
proteins[32] such as *orf89* and *orf90*, along with genes encoding the vRNAP (*orf178, 149, 180,*
80)[8,37] appear essential by phage TnSeq.

To validate a subset of phage TnSeq-based essentiality calls, we used Cas13a to target
transcripts encoding head proteins and isolated escape mutants. Strong targeting yielded
phages with large deletions in non-essential genes or point mutations in essential genes.
Deletions were obtained in several genes predicted to be non-essential by Phage TnSeq
(*orf93, 94, 162*, *163, 203, 303*), while targeting of genes predicted to be essential (*orf89*, *90*,
99, *119*, *177*) did not result in deletions (Figure S4A-C), corroborating their essentiality.
Notably, *orf97*, *orf157*, and *orf85–86* tolerated deletions despite lacking transposon

insertions, likely due to polar effects resulting from insertions (described below). *orf95* is
dispensable by Cas13a targeting but incurs a fitness cost in competition (Figure S5).
174
Conserved *Chimalliviridae* Proteins: In addition to gene size, evolutionary conservation is
expected to be correlated strongly with essentiality. A previous study defined 20 ΦKZ genes
that are conserved across Chimalliviridae yet unique to this phage family[38]. Of the "Prichard
20", 19 appear essential or strongly fitness conferring by phage TnSeq (15 with zero
insertions; 4 with only 1–3 insertions; Figure 3A, Supplementary Data 1). The lone exception
is the paralogous protein family encoded in multiple copies (*orf93/162/163*). As an example
of an essential conserved gene, Chimallin A (*orf54*), the phage nucleus shell protein[39,40], was
previously shown to be refractory to knockout in ΦKZ[19] and essential upon translational
repression in the related *E. coli* phage Goslar[9]. No Tn insertions were identified in this gene.
Other conserved genes encoding protein importers (*imp1*/*orf69*; *imp2*/*orf47*; *imp3*/*orf59*;
imp4/*orf48*), non-virion RNA polymerase (nvRNAP) subunits (*orf123, orf74, orf68, orf71–73,*
orf55–56.1), and the nucleus-associated *imp6/chmC/orf67*[25,41] also lack disruptive insertions
(Figures 2B, 3A). Additional conserved non-virion genes, such as *orf165* (a putative SbcCD-
like ABC ATPase with SMC homology), also lacked insertions. A subset of these genes were
also queried for essentiality previously via ASO-based translational repression[20], with
generally consistent results and some exceptions. Figure 3A summarizes the concordance
between phage TnSeq fitness scores and ASO knockdown.
192
Polar Effects and Transcriptional Architecture
Insertional approaches in bacteria and phages are susceptible to polar effects, where an
insertion disrupts expression of downstream genes in the same transcriptional unit. Polar
effects are more severe when a small number of operons are driven by few transcription
start sites (TSSs), and less problematic when multiple TSSs distribute transcription across
densely packed loci. Although phages are sometimes depicted as having few promoters,
recent short-read RNA sequencing identified 149 TSSs across the ΦKZ genome[26], in line
with earlier predictions of 134 operons[8].
201
To ask whether abundant TSSs mitigate polar effects in ΦKZ, we again examined the *chmA*
(*orf54*) gene. *orf53* and *chmA* appear in a putative operon with a TSS driving *orf53-chmA*,
yet multiple transposon insertions are present in *orf53* while none occur in *chmA* (Figure 3B).
This indicates that Tn insertion in *orf53* does not abrogate *chmA* transcription. An
independent TSS is found directly upstream of *chmA,* likely sustaining *chmA* transcription
even when *orf53* is disrupted. *orf55* (nvRNAP subunit) also appears essential, while *orf56–*
57 are non-essential and *orf58–59* appear essential (Figure 3B). A TSS upstream of *orf59*
(*imp3*) supports the interpretation that the absence of insertions in *orf58* reflects essentiality
of gp58, rather than a polar effect on *imp3*. This pattern in a tightly packed locus
underscores how multiple TSSs can insulate neighboring essential genes from polar effects.

Another example of polar effects being mitigated arises in the lysis cassette. Biochemical
studies show that gp144 is a highly active peptidoglycan hydrolase and the likely
endolysin[42,43]. A recent preprint further identified gp142 as the holin and gp143 as a lysis
regulator[44]. By phage TnSeq, *orf142* and *orf144* lack insertions and appear essential,
whereas *orf143* is disrupted by five insertions in its 534 bp coding sequence. Multiple TSSs
in this region are again identified, upstream of *orf139*, *orf143*, and *orf144*[26] (Figure 3C).
Thus, many potential polar effects are mitigated by the underlying transcriptional

architecture. Importantly, abundant TSSs are not unique to ΦKZ: several smaller
Pseudomonas phages profiled by the same method also exhibit numerous TSSs (e.g., 14–1,
50 TSSs; LUZ24, 16; LUZ19, 9; PAK_P3, 75; YuA, 21)[26].

By contrast, polar effects are expected when genes are expressed from a single
polycistronic transcript without internal TSSs. Two clear examples are the ΦKZ intron-
encoded homing endonucleases gp72 and gp179, both lying within RNA polymerase
operons. ASO knockdown data suggested that gp72 is essential while gp179 is not[20], while
both genes appear essential by phage TnSeq (no insertions). Moreover, a gp179 homolog in
ΦPA3 is non-essential but implicated in phage–phage competition[45]. To resolve this
discrepancy, we used Cas13a to target each gene and readily isolated deletion mutants in
orf72 and *orf179*, demonstrating that neither gene is strictly essential for ΦKZ replication
(Figure S4C). These loci thus exemplify polar effects: transposon insertion in *orf72* or *orf179*
likely terminates transcriptional readthrough, blocking expression of the downstream
essential genes (*orf73* or *orf180*). Similar cases likely exist elsewhere in the ΦKZ genome.
Cas13-based targeting of individual genes coupled to sequencing of escape mutants
provides a straightforward way to discriminate true essential genes from polar-effect artifacts
uncovered by phage TnSeq.
236
Non-essential genes revealed by dense insertion coverage
Genes with abundant, evenly distributed insertions can be confidently classified as non-
essential. *orf39* encodes the PhuZ tubulin required for nucleus positioning[12,13,46,47] contains
many insertions across its length and has been previously deleted[19]. ChmB (gp2), a protein
proposed to cooperate with PhuZ to facilitate capsid treadmilling and docking at the phage
nucleus[48], likewise appears non-essential, suggesting that core aspects of DNA packaging
remain to be fully explained.

Other genes with many Tn insertions and known non-essential functions encode: a ribosome
binding protein (gp14)[49], an injected protein[19] (gp93), Thoeris anti-defense protein 1[50] (Tad1,
gp184), a cluster uncharacterized non-essential proteins gp206-216[19], the activator of Juk
defense (gp241)[51], and proteins encoded nearby (gp237gp240, 242)[51]. Dip (gp37), a
previously studied RNA degradosome inhibitor[52], also appears dispensable. Interestingly,
most virion proteins in ΦKZ appear essential, but notable exceptions include a helicase
gp203 and gp303 (unknown function), both of which are proteolytically processed and
packaged into the head[32]. Gp203 and gp93 co-pellet with phage genomic DNA after capsid
disruption[32], indicating tight association with packaged DNA despite being non-essential
under the conditions tested
254
Small genes were particularly likely to tolerate insertions. Of 210 genes shorter than 500 bp
(arbitrarily defined as "small"), 185 contain at least one insertion, while just 25 lack
insertions. Because small genes are more likely to be misclassified as essential by chance,
this likely underestimates the true number of non-essential small genes. Even so, our
insertion data strongly indicate that approximately half of ΦKZ genes are both small and
non-essential under the laboratory conditions used here (37 °C, LB, aeration). Across the full
genome, roughly 70% of genes appear non-essential under these conditions. Many of these
non-essential genes may be conditionally essential in specific hosts or environments
(examples are presented in the next section)—phenotypes that phage Tn mutagenesis is
uniquely poised to uncover.

Minor capsid proteins are required at environmental temperature

P. aeruginosa is frequently isolated from soil, freshwater, and human-associated environments[53,54]. We therefore queried whether any phage genes are dispensable at 37 °C but required at a lower environmental temperature (18 °C). We subjected a randomly chosen set of 90 transposed mutants to infection at 18 °C. Four mutants that grew poorly at 18 °C were identified and sequenced alongside a control mutant that grew normally (Figure 3D). Insertions mapped to *orf162*, *orf199* (two redundant insertions), and *orf244*. All three genes encode proteins packaged in the capsid[32]. gp244 is notable as it appears to decorate the outside of the phage capsid[34]. The *orf244::Tn* mutant completely failed to replicate at 18 °C, a defect rescued by expressing *orf244*[33] in trans (Figure 3E).

To understand the role of minor capsid proteins in this phage, we next checked what phage TnSeq and a recent cryo-EM structure of the capsid reveal[34]. The cryo-EM structure revealed ~10 "minor capsid" proteins whose N-termini weave into major capsid hexamers[34]. Most of the genes encoding these minor capsid proteins, including *orf35*, *orf93*, *orf162*, *orf184* (*tad1*), and *orf244*, appear non-essential by phage TnSeq, at 37 °C. Interestingly, only three minor capsid genes appear essential at 37 °C: *orf28*, *orf91*, and *orf119*. The structural data provide an explanation for this, revealing a junction between vertex-binding complexes in which gp28 from one complex contacts gp91 and gp119 from the adjacent complex[34]. Cas13a targeting of these genes generated escapers lacking deletions, confirming essentiality and supporting a model where they form an interlocking complex critical for capsid assembly (Figure S4B). These findings therefore suggest multiple roles for minor capsid proteins: i) three minor capsid proteins that form an essential (at 37 °C) complex in the capsid, ii) additional minor capsid proteins, including a capsid decorating protein gp244, that are dispensable at 37 °C, but essential at 18 °C, and iii) non-essential proteins whose N-termini are wedged into the capsid and the C-termini that are liberated by a head protease[32] are injected during the next round of infection (i.e. gp93, gp184/Tad1)[33,50]. The transposed mutant library and pooled TnSeq can thus be used to rapidly identify genes whose contributions emerge only in ecologically relevant conditions—a likely major evolutionary pressure for *P. aeruginosa* phages[55].

gp298 is a conditionally essential tail protein for adsorption to wild isolates

Phage Tn mutagenesis provides a straightforward route to isolate knockouts for every non-essential gene, enabling systematic screens for strain- or condition-specific phenotypes. To explore host-range determinants, we arrayed ~1,500 transposed ΦKZ mutants in 96-well format and plated the collection on 18 wild *P. aeruginosa* strains. Host-specific phage "dropouts" were readily detected (Figure S6).

One striking example is *orf298*, which has 28 insertions across its 1,008-bp coding region after propagation in PAO1, yet an *orf298::Tn* mutant failed to plaque on clinical isolate PA1032 (from a human acute infection, San Francisco) and on Env201 (a soil isolate from Toronto). gp298 contains a Kelch domain and a predicted N-acetylneuraminate epimerase domain and is present at low abundance in the virion[33]. In infection time courses, wild-type ΦKZ efficiently adsorbed to all three hosts (as indicated by decreased free phage titers within 15 min), whereas the *orf298::Tn* mutant specifically failed to adsorb to PA1032 and Env201 but adsorbed normally to PAO1 (Figure 3F). These data support gp298 as a previously unrecognized tail-associated factor required for adsorption and/or receptor access on a subset of wild *P. aeruginosa* strains. Notably, *orf298* is not located near canonical tail genes; instead, it clusters with other non-essential genes of unknown function.

Nlp1 is a fitness-conferring recombinase required for NucC resistance

To identify phage genes that are conditionally essential for DNA replication or genome stability, we screened the 1,500 arrayed ΦKZ mutants against two nucleases: EcoRI and the Type III-C CBASS system with a NucC effector. Wild-type ΦKZ naturally resists both[14,25]. We hypothesized that mutants with increased nuclear permeability might become sensitive to EcoRI, a ~30 kDa protein that cannot normally access the phage genome but cleaves ΦKZ DNA if granted access to it *in vivo* and *in vitro*[14]. However, none of the mutants were restricted by EcoRI, indicating that genomic segregation from the cytoplasm—and thus resistance to both EcoRI and the endogenous PAO1 Type I RM system[14,56]—is robustly maintained across all tested mutants.

By contrast, one ΦKZ mutant completely lost replication on strains expressing a Type III-C CBASS system (Figure 4A). This system senses unknown phage factors, produces cAAA signals, and activates NucC, a nuclease effector previously implicated in blocking a ΦKZ-like phage in *Serratia*[57]. In that context, degradation of the host genome outside the phage nucleus was proposed to halt phage progression. We were therefore particularly interested in mutants that become sensitive to NucC, because ΦKZ-like phages degrade the host genome[46,47].

The NucC-sensitive mutant carried a Tn insertion in *nlp1* (nucleus-localized protein 1), encoding a protein homologous to phage T4 UvsX[58]. UvsX is a recombinase that is not strictly essential in T4, but its loss confers sensitivity to DNA damage[58]. To confirm this result, we deleted *nlp1* by replacing it with *acrVIA1* (ΦKZ *nlp1::acrVIA1*, hereafter referred to as *Δnlp1*) under complementation with wild-type *nlp1 in trans*. Like the Tn mutant, the *Δnlp1* phage was sensitive to Type III-C CBASS NucC, and this sensitivity was rescued by Nlp1 expression in trans (Figure 4A). Sensitivity was dependent on canonical CBASS activity because Acb2, a cyclic-nucleotide sponge that sequesters cAAA[59], rescued the *Δnlp1* phage. In contrast, Nlp1 expression did not rescue CBASS NucC sensitivity of an unrelated phage (JBD67Δ*acb2*), indicating that Nlp1 is not a general CBASS inhibitor (Figure 4A).

During infection, NucC activity modestly reduced WT ΦKZ DNA levels yet completely abolished DNA replication in the *Δnlp1* mutant (Figure 4B). Attempts to visualize NucC localization were unsuccessful because fluorescent tagging rendered NucC inactive. Nonetheless, the genetic data support a model in which Nlp1's recombinase activity repairs NucC-induced phage DNA damage, analogous to UvsX-mediated recombination of fragmented T4 genomes after UV or Cas9/Cas12 cleavage[60]. While NucC is often viewed as anti-phage through host genome destruction, our data suggest that NucC can directly damage phage DNA and that ΦKZ relies on Nlp1 to tolerate this assault. Consistent with a DNA repair defect specific to NucC, *Δnlp1* phage remains fully resistant to EcoRI (Figure S7).

The absence of Tn insertions in *nlp1* in the pooled phage TnSeq library suggested that it is a strongly fitness-conferring gene. *Δnlp1* phages replicated slower (Figure 4B), formed smaller plaques than wild-type ΦKZ, and plaque size was restored by Nlp1 complementation (Figure 4C), a phenotype reminiscent of T4 *uvsX* mutants[58]. Tn insertions in *nlp1* were likely present at low frequency in the initial transposed population but were outcompeted during propagation. In a direct competition assay over 4 h in liquid culture, control acr insertions in *orf2/chmB*, *orf184/tad1*, *orf39/phuZ*, or *orf241* exhibited competitive indices of ~0.35–2.0 (Figure S5), while *Δnlp1* showed a competitive index of 0.05, confirming a strong fitness

357 defect. *nlp1* thus exemplifies a gene for which full deletion is possible, but the mutant drops
358 out during phage competition or in the presence of CBASS NucC.
359
360 Together, these findings highlight the value of pairing pooled phage TnSeq to map
361 essential/fitness genes with an arrayed mutant library to interrogate conditional phenotypes
362 and recover mutants that are underrepresented in pooled selections.

Non-essential gene clusters and defense-related functions

363
Many non-essential ΦKZ genes cluster within a broad region of largely contiguous non-
essential genes spanning from ~*orf184* through *orf304* and then *orf1–orf20* on the left arm
(Figure 2B). ΦKZ is circularly permuted and its genome ends are arbitrary. This region
roughly corresponds to the portion absent from a reduced size *P. aeruginosa* jumbo phage,
ΦEL[35,61]. This observation suggests that this region encodes host-interacting or defense-
related adaptations rather than core replication modules.

It is tempting to speculate that these non-essential regions constitute "anti-defense loci", as
described for smaller phages[62], although this remains to be systematically tested. Our
arrayed mutant screens offer initial support. When we plated the 1,500 mutants on *P.*
aeruginosa strain Env201, we identified a mutant with an insertion in *orf302* that failed to
form plaques. This phage adsorbed normally but failed to produce progeny, consistent with
an Env201-specific intracellular defense that is absent in PAO1 (Figure S8). In a separate
screen against the PfJuk[51] (jumbo phage killer) defense system, which restricts ΦKZ, we
observed robust restriction of most mutants but identified one that grew well. Sequencing
revealed a Tn insertion in *orf241*, encoding the previously described Juk activator[51]. Thus,
the transposed mutant collection can rapidly uncover both inhibitors and activators of diverse
defense systems across wild *P. aeruginosa* isolates.

Discussion

381
Phage functional genomics is beginning to match the scale and sophistication of bacterial
genetics, but tools remain limited, particularly for large and complex phage genomes. Here,
we establish a generalizable phage transposon mutagenesis and phage TnSeq platform that
(i) maps essential and fitness-conferring genes genome-wide, (ii) generates a library of
knockouts in nearly all non-essential genes, and (iii) identifies safe harbor loci suitable for
insertion and expression of transgenes.

By directing *mariner* transposition into the ΦKZ nucleus and selecting for insertions using an
anti-CRISPR and CRISPR-Cas13a, we demonstrate a rapid and scalable workflow for
phage genome mutagenesis. Because *mariner* is widely used in bacteria, requires no host-
specific factors, and inserts semi-randomly, we anticipate that this approach will be broadly
adaptable across bacteria–phage pairs. The transposon donor can be easily reconfigured—
for example, by removing or modifying promoters—to profile therapeutic phages and identify
optimal genomic contexts for stable expression of payload genes such as anti-defense
factors or therapeutic cargo.

Phage TnSeq is highly complementary to recently developed ASO and CRISPRi-ART
platforms[20,21], which are particularly suited to titrating expression of essential genes. In
contrast, TnSeq requires no guide RNA or oligo design, can be deployed at genome scale in
a single experiment, and yields hundreds to thousands of individual insertion mutants that
can be archived and screened under diverse conditions. Tn insertions, however, are not
ideal for fine-tuned modulation of essential gene expression and are inherently limited by

issues such as polar effects. Moving forward, we anticipate that the recent explosion in
defense and anti-defense system discovery means that one can have the ability to
transpose, and select for transposed plaque formation, with any phage of interest.

One of the most striking outcomes of this work is the revelation that a majority of ΦKZ genes
are dispensable under standard laboratory conditions. Some of these non-essential genes
encode tail (e.g. gp298) or capsid-associated proteins (e.g. gp244) with host- or
temperature-specific roles, inhibitors (e.g. gp302) or activators (e.g. gp241) of bacterial
defense systems, and a recombinase (Nlp1/gp152) that becomes critical in the presence of
a specific nuclease. The ease with which transposon mutagenesis can interrogate this
abundant pool of non-essential genes sets the stage for systematic dissection of phage
accessory and anti-defense functions.

Smaller phages likely encode a lower fraction of non-essential genes than ΦKZ, yet the total
number of non-essential genes per phage may still be modest enough that a complete
collection of knockout mutants could be arrayed in a single or a few 96-well plates. This
makes it feasible to screen for defense inhibitors, host-range factors, and conditionally
essential genes across multiple hosts, temperatures, and selective pressures. As interest
grows in phage diversity, basic biology, genome engineering, and therapeutic applications,
we anticipate that phage Tn mutagenesis will provide a broadly useful resource and
methodology. The essentiality maps, mutant libraries, and design principles described here
should be transferable to other phage systems, enabling the community to move from
sequence catalogs to functional insight at scale.

Limitations
Phage TnSeq, as implemented here, requires a robust defense system that is difficult for
phages to escape and a cognate anti-defense factor that can be encoded on the transposon.
These requirements may constrain applicability for some phage–host systems.
Nevertheless, the programmability of CRISPR-Cas systems and the growing catalog of anti-
CRISPR proteins[63] provide a flexible toolbox for adapting selection schemes.

A second, inherent limitation of insertional mutagenesis in prokaryotes and phages is polar
effects: insertions can appear to reveal essential genes when they instead disrupt
expression of neighboring essential genes. We show that the high density of TSSs in ΦKZ
alleviates many potential polar effects, but we also identify clear cases where phage TnSeq
misclassifies non-essential homing endonucleases as essential due to operon structure. In
some regions, we identified insertion direction biases suggesting that perhaps read through
transcription can alleviate or induce confounders. Using Tn donors that lack transcriptional
terminators or that incorporate unidirectional terminators could reduce some polar effects.
Consequently, putative essential genes of interest should be addressed by complementary
approaches such as targeted knockdowns or targeted deletions.

Acknowledgements
J.B.-D. is supported by the National Institutes of Health (nos. R01 AI171041
and R01 AI167412). C.K. received support from the UCSF Discovery Fellowship and the
NIH award 2T32AI060537-21. D.M. received support from the NIH Ruth L. Kirschstein
National Research Service (NRSA) award 1F32GM149125-01/-02. We thank Bondy-
Denomy laboratory members for input into this work.

445
Author contributions

A.W.C. - Conceptualization, Formal analysis, Investigation, Methodology, Validation,
Visualization, Software, Writing - original draft,
W.X.Y. - Investigation, Methodology, Validation, Visualization, Writing - original draft, Writing
- review and editing, Writing - original draft, Writing - review and editing
D.M. - Investigation, Visualization, Writing - original draft, Writing - review and editing
C.K. - Investigation, Visualization, Writing - original draft, Writing - review and editing
M.R.M. - Investigation
Y.L. - Conceptualization, Resources
Y.Y. - Resources
J.B.D. - Conceptualization, Funding acquisition, Investigation, Writing - original draft, Writing
- review and editing
458
459   **Declaration of interests** J.B.-D. is a scientific advisory board member of SNIPR Biome and
460   Excision Biotherapeutics, a consultant to LeapFrog Bio and a scientific advisory board
461   member and cofounder of Acrigen Biosciences and ePhective Therapeutics. The remaining
462   authors declare no competing interests. The Bondy-Denomy laboratory received
463   past research support from Felix Biotechnology.

Main Figure Legends

Figure 1. Transposon mutagenesis of a nucleus-forming jumbo phage
(A) Schematic illustrating the ΦKZ transposon mutagenesis strategy. The Nlp1–Mariner transposase fusion localizes to the phage nucleus, where a transposon encoding *acrVIA1* inserts into the phage genome, generating insertional mutants. Output phages are subsequently subjected to Cas13-based counter-selection.
(B) Fluorescence microscopy of *P. aeruginosa* infected with ΦKZ expressing sfCherry-tagged transposase (red) and stained with DAPI (blue). White arrows indicate co-localization of transposase with the phage nucleus in infected cells.
(C) Quantification of ΦKZ replication efficiency in the presence of single versus dual Cas13a crRNAs.
(D) Representative plaque phenotypes of transposed ΦKZ forming on dual Cas13a targeting strains.
(E) Efficiency of plaquing (EOP) of ΦKZ on dual Cas13a strains following transposition with variants V1 (*acrVIA1* with a phage early promoter), V2 (*acrVIA1* lacking promoter), or V3 (derived from V1 with modified primer binding sites). All dark grey bars are at the limit of detection, where a pseudocount of 1 was used for calculating detection limits.

Figure 2. Genome-wide map of ΦKZ gene essentiality
(A) Genome-wide distribution of transposon (Tn) insertion reads across the ΦKZ genome. All forward reads are shown without filtering, with a pseudocount added to sites with single reads for log-scale visualization. The ΦKZ genome map is displayed below, with genes in the forward direction in dark gray and reverse in light gray.
(B) Schematic summary of gene essentiality across the ΦKZ genome, as determined by insertion sequencing (TnSeq) analysis.

Figure 3. Discovery of novel ΦKZ gene functions using Tn mutagenesis
(A) Heat map showing gene fitness scores compared with antisense oligonucleotide (ASO) assay results. Gene conservation and virion localization are indicated. Genes are rank ordered by essentiality.
(B–C) Examples of Tn insertion read density across representative genomic regions (*orf 53–59* and *141–144*).
(D) Plaque assays showing environmental temperature-sensitive mutants at 37 °C and 18 °C. Wild-type ΦKZ and a transposed control phage are included.
(E) Adsorption efficiency of wild-type and *orf298::Tn* ΦKZ phages in *P. aeruginosa* PAO1 and clinical isolates.

Figure 4. Nlp1, a ΦKZ-specific UvsX homolog, protects against CBASS defense
(A) Efficiency of plating (EOP) of wild-type and *Δnlp1* ΦKZ in the presence of CBASS. Control phages JBD67 and JBD67Δ*acb2* confirm CBASS targeting specificity, while Acb2 expression neutralizes CBASS activity.

(B) Quantitative PCR analysis of ΦKZ wild-type and ΦKZ *nlp1::acr* ΦKZ and replication in the presence and absence of CBASS.

(C) Full-plate infections comparing ΦKZ and ΦKZ *nlp1::acrVIA1* with or without *nlp1* complementation.

Supplemental Figure Legends

Figure S1. Construction and validation of Cas13 dual-guide counter-selection system
(A) Plaque assays of wild-type ΦKZ, ΦKZ *phuZ*::*acrVIA1*, and single-guide escapers on targeting and non-targeting Cas13 strains.
(B) Escape frequencies of single versus dual Cas13a crRNAs targeting ΦKZ.
(C) Agarose gel PCR verification of *acrVIA1* insertions in transposed plaques. Low-transposition control pools are shown for comparison.
(D) Plaque assays of transposition reactions across targeting and non-targeting hosts.
(E) Quantification of transposition efficiency as a function of input ΦKZ titer (pfu/mL).

Figure S2. Characterization of individual transposition reactions before and after enrichment
(A) Plaque assays comparing ten independent transposition reactions across selection conditions: no selection, dual Cas13a targeting, and JukAB. Wild-type ΦKZ is shown as control.
(B) Full-plate infections (20 µL input) of each transposition reaction.
(C) Post-enrichment plaque assays confirming selective recovery of transposed ΦKZ under Cas13a counter-selection.

Figure S3. Rank-ordered distribution of ΦKZ gene fitness scores
Cumulative ranking of ΦKZ genes based on transposon insertion frequency and essentiality scoring.

Figure S4. Cas13 escaper phages acquire large genomic deletions
(A) Schematic overview of genes targeted by Cas13. Protein function, essentiality, and conservation across jumbo phages and *Chimalliviridae* are indicated.
(B) Plaque assays showing targeting of wild-type ΦKZ and ΦKZ *orf105*::*acrVIA1* (*acrVIA1* control) with crRNAs directed genes encoding injected structural proteins. The Cas13a dual guide construct targeting *orf55* and *120* are included as control.
(C) Schematic representation of deletion events identified in Cas13 escaper genomes. The annotations start (s), dual (d) and cleavage (c) refer to the position of guide targeting. Numbers shown indicate the base number within the *orf* that had mutations. Top: virion, bottom: non-virion.

543 ──

Figure S5. Competitive index of ΦKZ mutants

Competitive fitness of ΦKZ mutants carrying *acrVIA1* insertions in different genes, compared with wild-type phage.

547 ──

Figure S6. Representative plaques selected for downstream phenotypic follow-up

Example images of plaques from arrayed ΦKZ mutants pinned onto *P. aeruginosa* lawns.

550 ──

Figure S7. *In vivo* EcoRI targeting of ΦKZ and ΦKZ *nlp1::acrVIA1*

Quantification of *in-vivo* EcoRI restriction of wild-type and ΦKZ *nlp1::acrVIA1*. F8 phage is included as an EcoRI-sensitive control unrelated to ΦKZ.

554 ──

Figure S8. Adsorption and replication kinetics of ΦKZ *orf302::Tn*

Adsorption and 2-hour replication profile of *orf302::Tn* compared to wild-type ΦKZ.

Supplemental Table Legends

Table S1. List of oligonucleotides used in this paper

Table S2. Raw TraDIS output for gene essentiality after trimming

Table S3. Raw TraDIS output for all insertions across the ΦKZ genome

References

1. Hatfull, G.F. (2015). Dark matter of the biosphere: The amazing world of bacteriophage
diversity. J. Virol. *89*, 8107–8110.

2. Mesyanzhinov, V.V., Robben, J., Grymonprez, B., Kostyuchenko, V.A., Bourkaltseva,
M.V., Sykilinda, N.N., Krylov, V.N., and Volckaert, G. (2002). The genome of
bacteriophage phiKZ of Pseudomonas aeruginosa. J. Mol. Biol. *317*, 1–19.

3. Prichard, A., and Pogliano, J. (2024). The intricate organizational strategy of nucleus-
forming phages. Curr. Opin. Microbiol. *79*, 102457.

4. Mozumdar, D., A Agard, D., and Bondy-Denomy, J. (2025). The complex developmental
mechanisms of nucleus-forming jumbo phages. Curr. Opin. Microbiol. *88*, 102676.

5. deYMartín Garrido, N., Orekhova, M., Lai Wan Loong, Y.T.E., Litvinova, A., Ramlaul, K.,
Artamonova, T., Melnikov, A.S., Serdobintsev, P., Aylett, C.H.S., and Yakunina, M.
(2021). Structure of the bacteriophage PhiKZ non-virion RNA polymerase. Nucleic Acids
Res. *49*, 7732–7739.

6. Yakunina, M., Artamonova, T., Borukhov, S., Makarova, K.S., Severinov, K., and
Minakhin, L. (2015). A non-canonical multisubunit RNA polymerase encoded by a giant
bacteriophage. Nucleic Acids Res. *43*, 10411–10420.

7. Antonova, D., Belousova, V.V., Zhivkoplias, E., Sobinina, M., Artamonova, T.,
Vishnyakov, I.E., Kurdyumova, I., Arseniev, A., Morozova, N., Severinov, K., et al.
(2023). The dynamics of synthesis and localization of jumbo phage RNA polymerases
inside infected cells. Viruses *15*. https://doi.org/10.3390/v15102096.

8. Ceyssens, P.-J., Minakhin, L., Van den Bossche, A., Yakunina, M., Klimuk, E., Blasdel,
B., De Smet, J., Noben, J.-P., Bläsi, U., Severinov, K., et al. (2014). Development of
giant bacteriophage KZ is independent of the host transcription apparatus. J. Virol. *88*,
10501–10510.

9. Armbruster, E.G., Rani, P., Lee, J., Klusch, N., Hutchings, J., Hoffman, L.Y.,
Buschkaemper, H., Enustun, E., Adler, B.A., Inlow, K., et al. (2025). Sequential
membrane- and protein-bound organelles compartmentalize genomes during phage
infection. Cell Host Microbe *33*, 484–497.e6.

10. Mozumdar, D., Fossati, A., Stevenson, E., Guan, J., Nieweglowska, E., Rao, S., Agard,
D., Swaney, D.L., and Bondy-Denomy, J. (2024). Characterization of a lipid-based
jumbo phage compartment as a hub for early phage infection. Cell Host Microbe *32*,
1050–1058.e7.

11. Antonova, D., Nichiporenko, A., Sobinina, M., Wang, Y., Vishnyakov, I.E., Moiseenko,
A., Kurdyumova, I., Chesnokov, Y.M., Stepanchikova, E., Bourkaltseva, M., et al.
(2024). Genomic transfer via membrane vesicle: a strategy of giant phage phiKZ for
early infection. J. Virol. *98*, e0020524.

12. Chaikeeratisak, V., Nguyen, K., Khanna, K., Brilot, A.F., Erb, M.L., Coker, J.K.C.,
Vavilina, A., Newton, G.L., Buschauer, R., Pogliano, K., et al. (2017). Assembly of a

nucleus-like structure during viral replication in bacteria. Science *355*, 194–197.

13. Chaikeeratisak, V., Nguyen, K., Egan, M.E., Erb, M.L., Vavilina, A., and Pogliano, J.
(2017). The Phage Nucleus and Tubulin Spindle Are Conserved among Large
Pseudomonas Phages. Cell Rep. *20*, 1563–1571.

14. Mendoza, S.D., Nieweglowska, E.S., Govindarajan, S., Leon, L.M., Berry, J.D., Tiwari,
A., Chaikeeratisak, V., Pogliano, J., Agard, D.A., and Bondy-Denomy, J. (2020). A
bacteriophage nucleus-like compartment shields DNA from CRISPR nucleases. Nature
577, 244–248.

15. Mahler, M., Costa, A.R., van Beljouw, S.P.B., Fineran, P.C., and Brouns, S.J.J. (2023).
Approaches for bacteriophage genome engineering. Trends Biotechnol. *41*, 669–685.

16. Chen, J., Nilsen, E.D., Chitboonthavisuk, C., Corban, J.E., Yang, M., Mo, C.Y., and
Raman, S. (2025). Systematic, high-throughput characterization of bacteriophage gene
essentiality on diverse hosts. Cell Host Microbe *33*, 1363–1380.e11.

17. Kilcher, S., Studer, P., Muessner, C., Klumpp, J., and Loessner, M.J. (2018). Cross-
genus rebooting of custom-made, synthetic bacteriophage genomes in L-form bacteria.
Proc. Natl. Acad. Sci. U. S. A. *115*, 567–572.

18. Adler, B.A., Hessler, T., Cress, B.F., Lahiri, A., Mutalik, V.K., Barrangou, R., Banfield, J.,
and Doudna, J.A. (2022). Broad-spectrum CRISPR-Cas13a enables efficient phage
genome editing. Nat. Microbiol. *7*, 1967–1979.

19. Guan, J., Oromí-Bosch, A., Mendoza, S.D., Karambelkar, S., Berry, J.D., and Bondy-
Denomy, J. (2022). Bacteriophage genome engineering with CRISPR-Cas13a. Nat.
Microbiol. *7*, 1956–1966.

20. Gerovac, M., Buhlmann, L., Zhu, Y., Đurica-Mitić, S., Rech, V., Carien, S., Gräfenhan,
T., Popella, L., and Vogel, J. (2025). Programmable antisense oligomers for phage
functional genomics. Nature. https://doi.org/10.1038/s41586-025-09499-6.

21. Adler, B.A., Al-Shimary, M.J., Patel, J.R., Armbruster, E.G., Colognori, D., Charles, E.J.,
Miller, K.V., Lahiri, A., Cui, M.L., Oromí-Bosch, A., et al. (2025). CRISPRi-ART enables
functional genomics of diverse bacteriophages using RNA-binding dCas13d. Nat.
Microbiol. *10*, 694–709.

22. Rubin, E.J., Akerley, B.J., Novik, V.N., Lampe, D.J., Husson, R.N., and Mekalanos, J.J.
(1999). In vivo transposition of mariner-based elements in enteric bacteria and
mycobacteria. Proc. Natl. Acad. Sci. U. S. A. *96*, 1645–1650.

23. Zhang, L., Dawson, A., and Finnegan, D.J. (2001). DNA-binding activity and subunit
interaction of the mariner transposase. Nucleic Acids Res. *29*, 3566–3575.

24. Lampe, D.J., Churchill, M.E., and Robertson, H.M. (1996). A purified mariner
transposase is sufficient to mediate transposition in vitro. EMBO J. *15*, 5470–5479.

25. Kokontis, C., Klein, T.A., Silas, S., and Bondy-Denomy, J. (2025). Multi-interface
licensing of protein import into a phage nucleus. Nature *639*, 456–462.

26. Putzeys, L., Wicke, L., Boon, M., van Noort, V., Vogel, J., and Lavigne, R. (2024).
Refining the transcriptional landscapes for distinct clades of virulent phages infecting
Pseudomonas aeruginosa. Microlife *5*, uqae002.

27. Meeske, A.J., Nakandakari-Higa, S., and Marraffini, L.A. (2019). Cas13-induced cellular
dormancy prevents the rise of CRISPR-resistant bacteriophage. Nature *570*, 241–245.

28. Sycheva, L.V., Shneider, M.M., Sykilinda, N.N., Ivanova, M.A., Miroshnikov, K.A., and
Leiman, P.G. (2012). Crystal structure and location of gp131 in the bacteriophage phiKZ
virion. Virology *434*, 257–264.

29. Aksyuk, A.A., Kurochkina, L.P., Fokine, A., Forouhar, F., Mesyanzhinov, V.V., Tong, L.,
and Rossmann, M.G. (2011). Structural conservation of the myoviridae phage tail
sheath protein fold. Structure *19*, 1885–1894.

30. Kurochkina, L.P., Aksyuk, A.A., Sachkova, M.Y., Sykilinda, N.N., and Mesyanzhinov,
V.V. (2009). Characterization of tail sheath protein of giant bacteriophage phiKZ
Pseudomonas aeruginosa. Virology *395*, 312–317.

31. Lecoutere, E., Ceyssens, P.-J., Miroshnikov, K.A., Mesyanzhinov, V.V., Krylov, V.N.,
Noben, J.-P., Robben, J., Hertveldt, K., Volckaert, G., and Lavigne, R. (2009).
Identification and comparative analysis of the structural proteomes of phiKZ and EL, two
giant Pseudomonas aeruginosa bacteriophages. Proteomics *9*, 3215–3219.

32. Thomas, J.A., Weintraub, S.T., Wu, W., Winkler, D.C., Cheng, N., Steven, A.C., and
Black, L.W. (2012). Extensive proteolysis of head and inner body proteins by a
morphogenetic protease in the giant Pseudomonas aeruginosa phage φKZ. Mol.
Microbiol. *84*, 324–339.

33. Fossati, A., Mozumdar, D., Kokontis, C., Mèndez-Moran, M., Nieweglowska, E., Pelin,
A., Li, Y., Guo, B., Krogan, N.J., Agard, D.A., et al. (2023). Next-generation proteomics
for quantitative Jumbophage-bacteria interaction mapping. Nat. Commun. *14*, 5156.

34. Yang, Y., Shao, Q., Guo, M., Han, L., Zhao, X., Wang, A., Li, X., Wang, B., Pan, J.-A.,
Chen, Z., et al. (2024). Capsid structure of bacteriophage ΦKZ provides insights into
assembly and stabilization of jumbo phages. Nat. Commun. *15*, 6551.

35. Cornelissen, A., Hardies, S.C., Shaburova, O.V., Krylov, V.N., Mattheus, W., Kropinski,
A.M., and Lavigne, R. (2012). Complete genome sequence of the giant virus OBP and
comparative genome analysis of the diverse ΦKZ-related phages. J. Virol. *86*, 1844–
1852.

36. Thomas, J.A., and Black, L.W. (2013). Mutational analysis of the Pseudomonas
aeruginosa myovirus KZ morphogenetic protease gp175. J. Virol. *87*, 8713–8725.

37. Thomas, J.A., Rolando, M.R., Carroll, C.A., Shen, P.S., Belnap, D.M., Weintraub, S.T.,
Serwer, P., and Hardies, S.C. (2008). Characterization of Pseudomonas chlororaphis
myovirus 201varphi2-1 via genomic sequencing, mass spectrometry, and electron
microscopy. Virology *376*, 330–338.

38. Prichard, A., Lee, J., Laughlin, T.G., Lee, A., Thomas, K.P., Sy, A.E., Spencer, T.,

Asavavimol, A., Cafferata, A., Cameron, M., et al. (2023). Identifying the core genome of
the nucleus-forming bacteriophage family and characterization of Erwinia phage RAY.
Cell Rep. *42*, 112432.

39. Nieweglowska, E.S., Brilot, A.F., Méndez-Moran, M., Kokontis, C., Baek, M., Li, J.,
Cheng, Y., Baker, D., Bondy-Denomy, J., and Agard, D.A. (2023). The ☐PA3 phage
nucleus is enclosed by a self-assembling 2D crystalline lattice. Nat. Commun. *14*, 927.

40. Laughlin, T.G., Deep, A., Prichard, A.M., Seitz, C., Gu, Y., Enustun, E., Suslov, S.,
Khanna, K., Birkholz, E.A., Armbruster, E., et al. (2022). Architecture and self-assembly
of the jumbo bacteriophage nuclear shell. Nature *608*, 429–435.

41. Enustun, E., Armbruster, E.G., Lee, J., Zhang, S., Yee, B.A., Malukhina, K., Gu, Y.,
Deep, A., Naritomi, J.T., Liang, Q., et al. (2024). A phage nucleus-associated RNA-
binding protein is required for jumbo phage infection. Nucleic Acids Res. *52*, 4440–
4455.

42. Briers, Y., Volckaert, G., Cornelissen, A., Lagaert, S., Michiels, C.W., Hertveldt, K., and
Lavigne, R. (2007). Muralytic activity and modular structure of the endolysins of
Pseudomonas aeruginosa bacteriophages phiKZ and EL. Mol. Microbiol. *65,* 1334–
1344.

43. Fokine, A., Miroshnikov, K.A., Shneider, M.M., Mesyanzhinov, V.V., and Rossmann,
M.G. (2008). Structure of the bacteriophage phi KZ lytic transglycosylase gp144. J. Biol.
Chem. *283*, 7242–7250.

44. Manohar, P., Wan, J., Ganser, G., Sharp, K., and Young, R. (2025). The Lysis Cassette
of Jumbophage phiKZ. bioRxiv. https://doi.org/10.1101/2025.11.11.687635.

45. Birkholz, E.A., Morgan, C.J., Laughlin, T.G., Lau, R.K., Prichard, A., Rangarajan, S.,
Meza, G.N., Lee, J., Armbruster, E., Suslov, S., et al. (2024). An intron endonuclease
facilitates interference competition between coinfecting viruses. Science *385*, 105–112.

46. Kraemer, J.A., Erb, M.L., Waddling, C.A., Montabana, E.A., Zehr, E.A., Wang, H.,
Nguyen, K., Pham, D.S.L., Agard, D.A., and Pogliano, J. (2012). A phage tubulin
assembles dynamic filaments by an atypical mechanism to center viral DNA within the
host cell. Cell *149*, 1488–1499.

47. Erb, M.L., Kraemer, J.A., Coker, J.K.C., Chaikeeratisak, V., Nonejuie, P., Agard, D.A.,
and Pogliano, J. (2014). A bacteriophage tubulin harnesses dynamic instability to center
DNA in infected cells. Elife *3*. https://doi.org/10.7554/eLife.03197.

48. Enustun, E., Deep, A., Gu, Y., Nguyen, K.T., Chaikeeratisak, V., Armbruster, E.,
Ghassemian, M., Villa, E., Pogliano, J., and Corbett, K.D. (2023). Identification of the
bacteriophage nucleus protein interaction network. Nat. Struct. Mol. Biol. *30*, 1653–
1662.

49. Gerovac, M., Chihara, K., Wicke, L., Böttcher, B., Lavigne, R., and Vogel, J. (2024).
Phage proteins target and co-opt host ribosomes immediately upon infection. Nat.
Microbiol. *9*, 787–800.

50. Li, D., Xiao, Y., Fedorova, I., Xiong, W., Wang, Y., Liu, X., Huiting, E., Ren, J., Gao, Z.,
Zhao, X., et al. (2024). Single phage proteins sequester signals from TIR and cGAS-like
enzymes. Nature *635*, 719–727.

51. Yuping, L., Guan, L., Becher, I., Makarova, K.S., Cao, X., Hareendranath, S., Guan, J.,
Stein, F., Yang, S., Boergel, A., et al. (2025). Jumbo phage killer immune system
targets early infection of nucleus-forming phages. Cell *188*, 2127–2140.e21.

52. Van den Bossche, A., Hardwick, S.W., Ceyssens, P.-J., Hendrix, H., Voet, M.,
Dendooven, T., Bandyra, K.J., De Maeyer, M., Aertsen, A., Noben, J.-P., et al. (2016).
Structural elucidation of a novel mechanism for the bacteriophage-based inhibition of
the RNA degradosome. Elife *5*, e16413.

53. Weimann, A., Dinan, A.M., Ruis, C., Bernut, A., Pont, S., Brown, K., Ryan, J., Santos,
L., Ellison, L., Ukor, E., et al. (2024). Evolution and host-specific adaptation of
Pseudomonas aeruginosa. Science *385*, eadi0908.

54. Crone, S., Vives-Flórez, M., Kvich, L., Saunders, A.M., Malone, M., Nicolaisen, M.H.,
Martínez-García, E., Rojas-Acosta, C., Catalina Gomez-Puerto, M., Calum, H., et al.
(2020). The environmental occurrence of Pseudomonas aeruginosa. APMIS *128*, 220–
231.

55. Greenrod, S.T.E., Hector, T.E., Blazanin, M., Cazares, D., and King, K.C. (2025).
Temperature as a driver of phage ecology and evolution. Annu. Rev. Microbiol. *79*, 497–
522.

56. Shmidov, E., Villani, A., Mendoza, S.D., Avihu, E., Lebenthal-Loinger, I., Karako-
Lampert, S., Shoshani, S., Ye, C., Wang, Y., Yan, H., et al. (2025). Multigenerational
proteolytic inactivation of restriction upon subtle genomic hypomethylation in
Pseudomonas aeruginosa. Nat. Microbiol. *10*, 2498–2510.

57. Malone, L.M., Warring, S.L., Jackson, S.A., Warnecke, C., Gardner, P.P., Gumy, L.F.,
and Fineran, P.C. (2020). A jumbo phage that forms a nucleus-like structure evades
CRISPR-Cas DNA targeting but is vulnerable to type III RNA-based immunity. Nat
Microbiol *5*, 48–55.

58. Hamlett, N.V., and Berger, H. (1975). Mutations altering genetic recombination and
repair of DNA in bacteriophage T4. Virology *63*, 539–567.

59. Cao, X., Xiao, Y., Huiting, E., Cao, X., Li, D., Ren, J., Fedorova, I., Wang, H., Guan, L.,
Wang, Y., et al. (2024). Phage anti-CBASS protein simultaneously sequesters cyclic
trinucleotides and dinucleotides. Mol. Cell *84*, 375–385.e7.

60. Wu, X., Zhu, J., Tao, P., and Rao, V.B. (2021). Bacteriophage T4 escapes CRISPR
attack by minihomology recombination and repair. MBio *12*, e0136121.

61. Hertveldt, K., Lavigne, R., Pleteneva, E., Sernova, N., Kurochkina, L., Korchevskii, R.,
Robben, J., Mesyanzhinov, V., Krylov, V.N., and Volckaert, G. (2005). Genome
comparison of Pseudomonas aeruginosa large phages. J. Mol. Biol. *354*, 536–545.

62. Pinilla-Redondo, R., Shehreen, S., Marino, N.D., Fagerlund, R.D., Brown, C.M.,

Sørensen, S.J., Fineran, P.C., and Bondy-Denomy, J. (2020). Discovery of multiple anti-
CRISPRs uncovers anti-defense gene clustering in mobile genetic elements. bioRxiv.
https://doi.org/10.1101/2020.05.22.110304.

63. Bondy-Denomy, J., Davidson, A.R., Doudna, J.A., Fineran, P.C., Maxwell, K.L.,
Moineau, S., Peng, X., Sontheimer, E.J., and Wiedenheft, B. (2018). A Unified
Resource for Tracking Anti-CRISPR Names. CRISPR J *1*, 304–305.
763

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, **Joseph Bondy-Denomy** (joseph.bondy-denomy@ucsf.edu).

Materials availability

Plasmids and strains generated in this study are available from the lead contact upon reasonable request and, where possible, will be deposited with a public repository (e.g., Addgene).

Data and code availability

- Insertion sequencing data and ΦKZ whole-genome sequencing data will be deposited in a public repository and accession numbers will be provided upon publication.

- Custom analysis scripts are available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Bacterial strains

Unless otherwise indicated, experiments were performed in *Pseudomonas aeruginosa* PAO1 and PAO1 derivatives. Bacteria were grown in LB broth at 37 °C with aeration. Where appropriate, media were supplemented with gentamicin (50 µg/mL) or carbenicillin (250 µg/mL).

Phages

The jumbo phage ΦKZ and derivative mutants generated in this study were propagated on *P. aeruginosa* PAO1 or specified host strains. For host-range and conditional screens, a

panel of wild *P. aeruginosa* isolates, including clinical isolate PA1032 and environmental
isolate Env201, were used as indicated.

____

METHOD DETAILS

Construction of mariner transposase expression strains

The mariner transposase gene was amplified from pBTK30 (kindly provided by the Lory lab)
and assembled into a Tn7 plasmid carrying sfCherry–gp152. For construction of FLAG-
tagged transposase, plasmid p30T expressing sfCherry-Mariner–Nlp1 was linearized with
primers oAWC088/oAWC021, and the resulting backbone was ligated with gBlock oAWC093
using NEBuilder HiFi DNA Assembly (New England Biolabs).

All plasmids were sequence-verified (Quintara Biosciences) and introduced into *P.*
aeruginosa PAO1 by electroporation together with pTNS3 for integration at the Tn7 site[1].
Transformants were selected on LB agar containing 50 µg/mL gentamicin. The Flp
recombinase plasmid pFLP was then introduced into these strains to excise the gentamicin
resistance cassette. Colonies were counter-selected on sucrose, and loss of gentamicin
resistance was confirmed by replica plating.

Construction of *acrVIA1* transposon donors (V1–V3)

For construction of *acrVIA1* transposon donors V1 and V2, p30HERDT (abbreviated p30T)
was linearized with primers oAWC047/oAWC048. gBlocks encoding distinct constructs
(oAWC122 and oAWC123 for donors V1 and V2, respectively) were ligated into the
linearized backbone using HiFi Assembly and transformed into *E. coli* XL1-Blue.

For the V3 donor used in pooled TnSeq experiments, the p20T plasmid was linearized
instead of p30T and ligated with gBlocks oAWC146 and oAWC147 using HiFi Assembly. All
constructs were confirmed by Sanger sequencing and subsequently transformed into PAO1
strains carrying the mariner transposase, as described above.

Construction of pHERD30T expression plasmids

To generate gene inserts in p30T, the plasmid backbone was first amplified with primers
WX_016 and WX_017. Genes of interest were amplified with the appropriate primer pairs

and assembled into the backbone using HiFi Assembly. Constructs were cloned into XL1-

Blue, sequence-verified, and introduced into PAO1 by electroporation.

Construction of CRISPR-Cas13 guide plasmids

For cloning of Cas13 guide arrays for single or dual targeting, the plasmid backbone was

amplified with primers oAWC004/oAWC024. Complementary oligonucleotides (Table S1)

encoding the desired crRNAs (100 µM each) were annealed in Duplex Buffer A and ligated

into the backbone using HiFi Assembly.

For Cas13 guide plasmids used to target candidate essential or non-essential genes in ΦKZ

(to isolate escaper mutants), the Cas13 backbone plasmid was amplified with primers that

incorporated the crRNA sequence directly. Products were assembled with HiFi and cloned

into XL1-Blue. All guide plasmids were confirmed by sequencing and transformed into PAO1

or PAO1 *attTn7::cas13aLse* as appropriate.

————————————————————————————————————

Generation of transposed ΦKZ arrays

Transposed ΦKZ phages were selected on a Cas13 strain expressing dual crRNAs targeting

orf120 and *orf55*. Individual plaques were picked into SM buffer. In total, 1,800 plaques were

collected from 20 plates in a 96-well format, with plate corners reserved for wild-type ΦKZ

controls and two central wells containing no phage. The phage array was replicated onto

lawns of different *P. aeruginosa* isolates and incubated overnight at 37 °C to assess host-

range and conditional phenotypes.

Cas13a-mediated isolation of ΦKZ deletion mutants

839   High-titer ΦKZ stocks were plated on induced PAO1 *attTn7::cas13aLse* strains carrying

840   targeting guides as described above. Escaper plaques were picked and propagated on

841   PAO1 *attTn7::cas13aLse* expressing the same guides. Regions spanning the putative

842   deletion were amplified using primers flanking the targeted locus and sequenced to confirm

843   the nature of the deletions or point mutations.

Deletion of individual ΦKZ genes by *acrVIA1* replacement

Homology-directed constructs for gene replacement with *acrVIA1* were synthesized (Twist

Bioscience) with compatible overhangs and cloned into NheI-digested p30T using NEB HiFi

Gibson assembly. Constructs were selected with gentamicin and confirmed by sequencing.

Phage recombinants were generated and selected similarly to a previous report[2]. PAO1

strains harboring these plasmids were grown overnight, subcultured 1:100 into fresh LB, and

grown at 37 °C to an OD_{600} of 0.6. Cultures were diluted 1:100, and ~10^3 PFU of ΦKZ were

added for overnight infection. The following day, cultures were centrifuged at 10,000 × g for

2 min, and chloroform was added to the supernatant at a 1:8 (v/v) ratio. After ≥15 min at

room temperature, lysates were centrifuged again to clarify.

For selection of recombinants, 50 µL of lysate were mixed with 300 µL of PAO1

attTn7::cas13aLse + p30T crRNA[55/120] in the presence of 0.3% arabinose and 1 mM IPTG

and plated in top agar. Resistant plaques were picked and propagated on the same strain

under the same induction conditions. Candidate recombinant phages were screened by PCR

using primers outside the homology arms to confirm *acrVIA1* insertion into the phage

genome and to exclude plasmid contamination. An internal PCR targeting the wild-type locus

was used to monitor residual WT phage. Serial passages and screening were repeated until

no WT amplicon was detectable.

Transposition infections for pooled TnSeq

Transposition strains and control strains were grown overnight from glycerol stocks in LB

supplemented with either 50 µg/mL gentamicin (V1, V2 donors) or 250 µg/mL carbenicillin

(V3 donor) at 37 °C with shaking. Overnight cultures were subcultured 1:100 into LB

containing 1 mM IPTG (to induce transposase expression), the appropriate antibiotic, and 10

mM Mg^{2+} and grown for 2 h at 37 °C with shaking.

Log-phase cultures were adjusted to OD_{600} = 0.5 in 3 mL volumes. Bacteria were infected

with 10 µL of 2 × 10^8 PFU/mL ΦKZ at an MOI of approximately 2.22 × 10^{-3} and incubated

for 16 h at 37 °C with shaking. The next day, 2 mL of each culture were centrifuged at

16,000 × g for 2 min to pellet unlysed cells. Supernatants were transferred to new tubes,

mixed with 100 µL chloroform, and incubated at room temperature with shaking for 10 min.

After centrifugation at 10,000 × g for 2 min, supernatants were transferred to fresh tubes,

extracted once more with 100 µL chloroform, clarified, and stored at 4 °C.

For the genome-wide pooled TnSeq library, 10 independent transposition reactions done

using PAO1 *attTn7:mariner-sfCherry-nlp1* + p20T-V3 were performed and processed as

described above.

Cas13a counter-selection of transposed phages

Transposition output phages were first titered by serial dilution spot assays on Cas13 non-

targeting and dual-targeting strains, as well as in the presence of *jukAB*, to verify identity and

exclude contaminants. Typical output titers were ~4 × 10$^{1\square}$ PFU/mL.

To enrich for transposed phages, PAO1 *attTn7::cas13aLse* + p30T^{nlp1} crRNA[55/120] strain was

grown to OD$\square\square\square$ ~1.8 in 3 mL LB and infected at an approximate MOI of 5 using 400 μL of

the ~1 × 10$^{1\square}$ PFU/mL transposition output. Counter-selection cultures contained 1 mM

IPTG, 0.3% arabinose, 50 μg/mL gentamicin, and 10 mM Mg$^{2\square}$ and were incubated

overnight at 37 °C with shaking. Phages were harvested the following day by chloroform

treatment as described above and stored at 4 °C. Serial dilution spot assays and full-plate

infections on the counter-selection strain confirmed enrichment and Cas13 resistance, and

titers were typically ~2 × 10$^{1\square}$ PFU/mL.

For pooled TnSeq, enriched phages from the ten reactions were combined in equal PFU

amounts.

―――――――――――――――――――――――――――――――――――

Library preparation for transposon insertion sequencing

Phage DNA isolation

Genomic DNA was purified from enriched phage lysates using SDS/proteinase K treatment

followed by column purification. Briefly, 50–200 μL of high-titer phage lysate were mixed 1:1

with lysis buffer (10 mM Tris-HCl pH 8.0, 10 mM EDTA, 100 μg/mL proteinase K, 100 μg/mL

RNase A, 0.5% SDS), incubated at 37 °C for 30 min and at 55 °C for 30 min, and then

processed with the DNA Clean & Concentrator kit (Zymo Research). DNA concentration was

measured by spectrophotometry.

Illumina library construction and nested PCR

Transposon junction libraries were prepared using the NEBNext Ultra II FS DNA Library

Prep Kit for Illumina with modifications to enrich for transposon–genome junctions.

To generate indexed i7 adaptors, oAWC190 was pre-annealed separately with oAWC191,
192, 194, or 195 in 10 mM NaCl at 200 µM. For each pair, 12 µL of each primer were mixed,
heated to 95 °C for 5 min, and cooled to 12 °C at 0.1 °C/sec. Annealed oligos were diluted to
15 µM and stored at −20 °C.

For fragmentation, 1,000 ng of phage DNA in 26 µL 10 mM Tris-HCl pH 8.0 were mixed with
7 µL Ultra II FS Reaction Buffer and 2 µL Ultra II FS Enzyme Mix, vortexed, and incubated at
37 °C for 24 min to generate 100–250 bp fragments, followed by 65 °C for 30 min and hold
at 4 °C. I7 adapters were ligated by adding 30 µL Ultra II Ligation Master Mix, 1 µL Ligation
Enhancer, and 2.5 µL of 15 µM pre-annealed adaptor, followed by incubation at 20 °C for 15
min.

For size selection, the reaction was brought to 100 µL with 10 mM Tris-HCl pH 8.0, and 40
µL AMPure XP beads (Beckman Coulter) were added. After 10 min at room temperature and
5 min on a magnetic rack, the supernatant was transferred to a new tube. An additional 20
µL AMPure XP beads were added, incubated 10 min, and the beads washed twice with 200
µL 80% ethanol. Beads were air-dried for 3–5 min and eluted in 16 µL water. Fifteen
microliters of eluate containing size-selected DNA were carried forward.

To amplify transposon junctions, 15 µL of size-selected DNA were mixed with 25 µL
NEBNext Ultra II Q5 Master Mix , 5 µL 10 µM oAWC196, and 5 µL 10 µM oAWC224 (total
50 µL). PCR was performed with: 98 °C for 30 s; 12 cycles of 98 °C for 10 s, 65 °C for 75 s;
65°C for 5 min; hold at 12 °C. PCR products were cleaned with AMPure XP beads (45 µL
beads per 50 µL reaction), washed twice with 80% ethanol, and eluted in 16 µL water (15 µL
carried forward).

A second PCR to further enrich junctions and attach i5 adaptors was performed using 15 µL
eluate and 35 µL of master mix (25 µL Q5 Master Mix, 5 µL 10 µM oAWC199, and 5 µL of
10 µM oAWC200, 201, 202, or 203). Products were cleaned with AMPure XP beads as
above, eluted in 33 µL water, and 30 µL were retained.

Library quality was assessed using the Qubit 1× dsDNA HS Assay (expected 20–50 ng/µL)
and Agilent D1000 ScreenTape (expected size ~300–350 bp).

____

Illumina sequencing

Libraries were sequenced on an Illumina MiSeq i100 instrument using the MiSeq i100 Series

25M Reagent Kit (100-cycle; 20126567). Indexed libraries were pooled and diluted with

water to ~4 nM (Qubit-based). A 10× loading concentration of 0.8 nM in 100 µL

Resuspension Buffer was prepared according to the manufacturer's instructions.

A separate indexed sample for ΦKZ whole-genome sequencing was prepared at 0.8 nM and

included as a diversity spike-in. The loading mix consisted of 75 µL pooled library and 25 µL

spike-in. Runs were configured with 8 cycles for index 1, 8 cycles for index 2, 100 cycles for

read 1, and 0 cycles for read 2. Quality control and demultiplexing were performed in

Illumina BaseSpace, and FASTQ files for each index were exported for downstream

analysis.

____

Insertion sequencing analysis

Demultiplexed FASTQ reads were filtered for those containing the transposon inverted

repeat sequence marking the insertion junction

(GACCGGGGACTTATCAGCCAACCTGTTA). Retained reads were trimmed to keep the last

10 nt of the inverted repeat (CAACCTGTTA) plus 35 nt of downstream genomic sequence.

Trimmed reads were mapped to the reverse complement of the ΦKZ reference genome

(NC_004629) using the TraDIS toolkit (bacteria_tradis). Insertion sites supported by ≤2

reads were removed before essentiality analysis. Gene-level insertion statistics were

obtained using tradis_gene_insert_sites on both raw and filtered insertion datasets; for

filtered data, insertions within the first and last 10% of each annotated ORF were excluded.

Fitness-conferring genes were annotated using the tradis_essentiality.R script. TraDIS

outputs were further processed and visualized in R.

____

Plaque assays

Unless otherwise noted, plaque assays were performed at 37 °C. Strains were grown

overnight in LB at 37 °C (EcoRI experiments were typically ~13 h). For spot assays, 100 µL

of overnight culture were mixed with 3–5 mL of 0.4% LB top agar supplemented with Mg²⬚

and poured onto LB + Mg plates. After solidification, ΦKZ or derivative phages were serially

diluted and 2 µL of each dilution were spotted on the lawn. Plates were incubated overnight

at 37 °C.

For complementation assays with the ΦKZ *nlp1::acrVIA1* phage, 150 μL of overnight culture
were mixed with 10 μL of phage at the desired dilution and 3.5 mL of 0.35% top agar with
Mg²⁺, poured on LB + Mg plates containing 50 μg/mL gentamicin, and incubated overnight
at 30 °C.

For cold-sensitivity assays, plates were incubated for 48 h at 18 °C. A water-filled plate was
placed on top of assay plates to provide weight and limit evaporation.

───

qPCR for CBASS experiments

Strains were grown overnight at 37 °C and diluted 1:100 into fresh LB the next day. Cultures
were grown for 6 h, and ΦKZ was added at MOI ~0.5. At each time point, 100 μL of culture
were harvested, washed once with 1× PBS, and pellets were snap-frozen on dry ice and
stored at −20 °C.

Genomic DNA was extracted using the Zymo Genomic DNA kit. Approximately 1 ng of DNA
was used as input for qPCR with primer pairs 15F/15R (phage) and rpoD_F/rpoD_R
(bacterial chromosomal control). Phage DNA levels were first normalized to bacterial DNA
(ΔCt), and relative phage DNA abundance was then calculated relative to t = 0.

───

Adsorption and replication assays

Overnight cultures were diluted 1:100 into fresh LB and grown at 37 °C for ~2.5 h to $OD_{600} \approx$
0.6. ΦKZ (10^4 PFU) was added to each culture. For adsorption measurements, 100 μL of
culture were sampled at t = 15 min, mixed with chloroform to lyse cells, and titrated to
determine unadsorbed phage. For replication assays, 100 μL of culture were collected at t =
2 h, lysed with chloroform, and titrated on PAO1 to quantify total phage yield.

───

Competition assays

Mutant phages carrying *acrVIA1* were first verified to form plaques on both a non-targeting
strain (PAO1 *attTn7::cas13aLse* + p30T^empty) and a Cas13 single-guide strain PAO1
attTn7::cas13aLse + p30T crRNA[120], while wild-type ΦKZ formed plaques only on PAO1

attTn7::cas13aLse + p30T^{empty}. Full-plate infections of mutant and wild-type phage on PAO1

attTn7::cas13aLse + p30T^{empty} were used to generate starting stocks.

Overnight cultures of PAO1 *attTn7::cas13aLse* with and without guide RNA and were grown

at 37 °C. For competition, PAO1 *attTn7::cas13aLse* + p30T^{empty} was subcultured 1:100 and

grown to $OD_{600} \approx 0.6$. Mutant and wild-type phage were mixed at an approximate 1:1 ratio

and added to PAO1 *attTn7::cas13aLse* + p30T^{empty} at a total MOI of 1 in 150 μL volumes in

96-well plates. Infection was monitored over 4 h in a plate reader.

The initial ratio of mutant to wild-type phage was determined at t = 0 by full-plate infection of

PAO1 *attTn7::cas13aLse* + p30T^{empty} (total phage) and PAO1 *attTn7::cas13aLse* + p30T

crRNA[120](mutant-only) at appropriate dilutions for plaque counting. After 4 h, chloroform was

added to each well and incubated for ~15 min, and lysates were clarified by centrifugation.

The final ratio of mutant to wild-type phage was determined by titration on PAO1

attTn7::cas13aLse + p30T^{empty} and PAO1 *attTn7::cas13aLse* + p30T crRNA[120].

The **Competitive Index (CI)** was calculated as:

$$Competitive\ Index\ (CI) = \frac{\left(\frac{Mutant\ phage\ PFU\ at\ t = 0}{Total\ phage\ PFU\ at\ t = 0}\right)}{\left(\frac{Mutant\ phage\ PFU\ at\ t = 4h}{Total\ phage\ PFU\ at\ t = 4h}\right)}$$

1008

A CI of 1 indicates no change in mutant frequency during the assay (mutant fitness

equivalent to wild type), CI < 1 indicates a competitive disadvantage, and CI > 1 indicates a

competitive advantage.

____

QUANTIFICATION AND STATISTICAL ANALYSIS

For TnSeq analysis, insertions supported by ≤2 reads were excluded and the first and last

10% of each ORF were masked prior to essentiality calling (results in Table S2). Essential

and fitness-conferring genes were identified using the TraDIS essentiality pipeline

(tradis_essentiality.R). For calculation of total insertions in *orf*s, all reads were considered

(results in Table S3). Graphical analysis related to TnSeq were all performed in R.

Plaque assays, adsorption/replication assays, qPCR experiments, and competition assays

were typically performed with at least three biological replicates unless otherwise indicated in

figure legends. Statistical analysis (e.g., calculation of mean, standard deviation, and, where

appropriate, significance testing) was performed in GraphPad Prism; details of tests are

indicated in the corresponding figure legends.

1024

1025 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Bacterial and virus strains		
PAO1 p30T[empty]	Qiu *et al.*, 2008[3]	
PAO1 p30T[244-FLAG]	Fossati, Mozumdar *et al.*, 2023[4]	
PAO1 *attTn7::empty*	This paper	
PAO1 *attTn7::jukAB*	Yuping *et al.*, 2025[5]	
PAO1 *attTn7::CBASS*	Huiting *et al.*, 2023[6]	
PAO1 *attTn7::CBASS* + p30T[empty]	Huiting *et al.*, 2023	
PAO1 *attTn7::CBASS* + p30T[nlp1]	This paper	
PAO1 *attTn7::CBASS* + p30T[acb2]	Huiting *et al.*, 2023	
PAO1 *attTn7::empty* + p30T[empty]	This paper	
PAO1 *attTn7::empty* + p30T[nlp1]	Huiting *et al.*, 2023	
PAO1 *attTn7::empty* + p30T[acb2]	Huiting *et al.*, 2023	
PAO1 p30T[Gb135]	This paper	
PAO1 p30T[Gb136]	This paper	
PAO1 p30T[Gb144]	This paper	
PAO1 *attTn7::mariner-sfCherry-nlp1*	This paper	
PAO1 *attTn7::mariner-sfCherry-nlp1* + p30T-V1	This paper	
PAO1 + p30T-V1	This paper	
PAO1 *attTn7::mariner-sfCherry-nlp1* + p30T-V2	This paper	
PAO1 + p30T-V2	This paper	
PAO1 *attTn7::mariner-FLAG* + p20T-V3	This paper	
PAO1 *attTn7::mariner-sfCherry-nlp1* + p20T-V3	This paper	
PAO1 + p20T-V3	This paper	
PAO1 *attTn7::cas13aLse* + p30T[nlp1] crRNA[55/120]	This paper	
PAO1 *attTn7::cas13aLse* + p30T[empty]	Guan *et al.*, 2022[2]	
PAO1 *attTn7::cas13aLse* + p30T crRNA[120]	Guan *et al.*, 2022	
PAO1 *attTn7::cas13aLse* + p30T crRNA[55]	Guan *et al.*, 2022	
PAO1 *attTn7::cas13aLse* + p30T crRNA[55/120]	This paper	

PAO1 *attTn7::cas13aLse* + p30T crRNA[86start]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[95cleav]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[97start]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[97cleav]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[99start]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[119start]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[162dual]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[163dual]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[177start]	This paper	
PAO1 *attTn7::cas13aLse* + p30T crRNA[303start]	This paper	
Env201	This paper	
PA1032	This paper	
Phages		
ΦKZ	Alan Davidson Lab	AF399011
ΦKZ *orf298::Tn*	This paper	
ΦKZ *orf302::Tn*	This paper	
ΦKZ_*orf152::Tn*	This paper	
ΦKZ *nlp1::acr*	This paper	
ΦKZ *orf2::acr*	This paper	
ΦKZ *orf95::acr*	This paper	
ΦKZ *orf93::acr*	Guan *et al.*, 2022	
ΦKZ *orf105::acr*	This paper	
ΦKZ *orf244::acr*	This paper	
ΦKZ*orf184::acr*	Li, Xiao, Federova… *et al.*, 2024[7]	
ΦKZ*orf241::acr*	Yuping *et al.*, 2025	
JBD67	Alan Davidson Lab	NC_042135.1
JBD67Δ*acb2*	Huiting *et al.*, 2023	
Biological samples		
Chemicals, peptides, and recombinant proteins		
Gentamicin	RPI	
Carbenicillin	BPBio	
IPTG	Fisher	
Chloroform	Alta Aesar	

Proteinase K	Thermofisher Scientific	
RNaseA	Omega	
Critical commercial assays		
Luna qPCR master mix	New England Biolabs	M3003X
NEBNext Ultra II FS DNA Library Prep Kit	New England Biolabs	E7805
NEBuilder HiFi DNA Assembly Master mix	New England Biolabs	E2621L
DNA Clean and Concentrator Kit	Zymo Research	D4034
AMPure XP Beads	Beckman Coulter	A63880
Qubit 1x dsDNA HS Assay Kit	Thermofisher Scientific	Q33230
Agilent D1000 ScreenTape and Reagents	Agilent Technologies	5067-5582 5067-5583 5067-5602 5067-5586
Deposited data		
Experimental models: Cell lines		
Experimental models: Organisms/strains		
Pseudomonas aeruginosa strain PAO1		
Oligonucleotides		
See Table S1 for details		
Recombinant DNA		
p30T^{empty}	Qiu *et al.*, 2008	
p30T crRNA^{120}	Guan *et al.*, 2022	
p30T crRNA^{55}	Guan *et al.*, 2022	
p30T crRNA$^{55/120}$	This paper	
p30T crRNA^{86start}	This paper	
p30T crRNA^{95cleav}	This paper	
p30T crRNA^{97start}	This paper	
p30T crRNA^{97cleav}	This paper	
p30T crRNA^{99start}	This paper	

p30T crRNA[119start]	This paper	
p30T crRNA[162dual]	This paper	
p30T crRNA[163dual]	This paper	
p30T crRNA[177start]	This paper	
p30T crRNA[303start]	This paper	
p30T V1	This paper	
p30T V2	This paper	
p20T V3	This paper	
p30T[244-FLAG]	Fossati, Mozumdar *et al.*, 2023	
p30T*nlp1*	Huiting *et al.*, 2023	
p30T*acb2*	Huiting *et al.*, 2023	
p30T[Gb135]	This paper	
p30T[Gb136]	This paper	
p30T[Gb144]	This paper	
attTn7::mariner-sfCherry-nlp1	This paper	
attTn7::mariner-FLAG	This paper	
TNS3	Choi et al., 2007	
Software and algorithms		
GraphPad PRISM	GraphPad Software	
RStudio	R Foundation	
TraDIS Toolkit	Barquist *et al.*, 2016[8]	
Other		

1026

1027

References

1. Choi, K.-H., and Schweizer, H.P. (2006). mini-Tn7 insertion in bacteria with single
attTn7 sites: example Pseudomonas aeruginosa. Nat. Protoc. *1*, 153–161.

2. Guan, J., Oromí-Bosch, A., Mendoza, S.D., Karambelkar, S., Berry, J.D., and Bondy-
Denomy, J. (2022). Bacteriophage genome engineering with CRISPR-Cas13a. Nat.
Microbiol. *7*, 1956–1966.

3. Qiu, D., Damron, F.H., Mima, T., Schweizer, H.P., and Yu, H.D. (2008). PBAD-based
shuttle vectors for functional analysis of toxic and highly regulated genes in
Pseudomonas and Burkholderia spp. and other bacteria. Appl. Environ. Microbiol. *74*,
7422–7426.

4. Fossati, A., Mozumdar, D., Kokontis, C., Mèndez-Moran, M., Nieweglowska, E., Pelin,
A., Li, Y., Guo, B., Krogan, N.J., Agard, D.A., et al. (2023). Next-generation proteomics
for quantitative Jumbophage-bacteria interaction mapping. Nat. Commun. *14*, 5156.

5. Yuping, L., Guan, L., Becher, I., Makarova, K.S., Cao, X., Hareendranath, S., Guan, J.,
Stein, F., Yang, S., Boergel, A., et al. (2025). Jumbo phage killer immune system
targets early infection of nucleus-forming phages. Cell *188*, 2127–2140.e21.

6. Huiting, E., Cao, X., Ren, J., Athukoralage, J.S., Luo, Z., Silas, S., An, N., Carion, H.,
Zhou, Y., Fraser, J.S., et al. (2023). Bacteriophages inhibit and evade cGAS-like
immune function in bacteria. Cell *186*, 864–876.e21.

7. Li, D., Xiao, Y., Fedorova, I., Xiong, W., Wang, Y., Liu, X., Huiting, E., Ren, J., Gao, Z.,
Zhao, X., et al. (2024). Single phage proteins sequester signals from TIR and cGAS-like
enzymes. Nature *635*, 719–727.

8. Barquist, L., Mayho, M., Cummins, C., Cain, A.K., Boinett, C.J., Page, A.J., Langridge,
G.C., Quail, M.A., Keane, J.A., and Parkhill, J. (2016). The TraDIS toolkit: sequencing
and analysis for dense transposon mutant libraries. Bioinformatics *32*, 1109–1111.

1053
1054

Figure 1. Transposon mutagenesis of a nucleus-forming jumbo phage

(A) Schematic illustrating the ΦKZ transposon mutagenesis strategy. The Nlp1–mariner transposase fusion localizes to the phage nucleus, where a transposon encoding *acrVIA1* inserts into the phage genome, generating insertional mutants. Output phages are subsequently subjected to Cas13-based counter-selection.

(B) Fluorescence microscopy of *P. aeruginosa* infected with ΦKZ expressing SfCherry-tagged transposase (red) and stained with DAPI (blue). White arrows indicate co-localization of transposase with the phage nucleus in infected cells.

(C) Quantification of ΦKZ replication efficiency in the presence of single versus dual Cas13a crRNAs.

(D) Representative plaque phenotypes of transposed ΦKZ forming on dual Cas13a targeting strains.

(E) Efficiency of plating (EOP) of ΦKZ on dual Cas13a strains following transposition with variants V1 (*acrVIA1* with a phage early promoter), V2 (*acrVIA1* lacking promoter), or V3 (derived from V1 with modified primer binding sites). All dark grey bars are at the limit of detection, where a pseudocount of 1 was used for calculating detection limits.

Figure 2. Genome-wide map of ΦKZ gene essentiality

(A) Genome-wide distribution of transposon (Tn) insertion reads across the ΦKZ genome. All forward reads are shown without filtering, with a pseudocount added to sites with single reads for log-scale visualization. The ΦKZ genome map is displayed below, with forward *orf*s in dark gray and reverse *orf*s in light gray.

(B) Schematic summary of gene essentiality across the ΦKZ genome, as determined by insertion sequencing (TnSeq) analysis.

Figure 3. Discovery of novel ΦKZ gene functions using Tn mutagenesis

(A) Heat map showing gene fitness scores compared with antisense oligonucleotide (ASO) assay results. Gene conservation and virion localization are indicated. Genes are rank-ordered by essentiality.

(B–C) Examples of Tn insertion read density across representative genomic regions (*orfs 53–59* and *141–144*).

(D) Plaque assays showing environmental temperature-sensitive mutants at 37 °C and 18 °C. Wild-type ΦKZ and a transposed control phage are included.

(E) Adsorption efficiency of wild-type and *orf298::Tn* ΦKZ phages in *P. aeruginosa* PAO1 and clinical isolates.

Figure 4. Nlp1, a ΦKZ-specific UvsX homolog, protects against CBASS defense

(A) Efficiency of plating (EOP) of ΦKZ wild type and Δ*nlp1* in the presence of CBASS. Control phages JBD67 and JBD67Δ*acb2* confirm CBASS targeting specificity, while Acb2 expression neutralizes CBASS activity.
(B) Quantitative PCR analysis of ΦKZ wild type and ΦKZ *nlp1::acrVIA1* replication in the presence and absence of CBASS.
(C) Full-plate infections comparing ΦKZ and ΦKZ *nlp1::acrVIA1* with or without *nlp1* complementation.

Figure S1. Construction and validation of Cas13 dual-guide counter-selection system

(A) Plaque assays of wild-type ΦKZ, ΦKZ *phuZ::acrVIA1*, and single-guide escapers on targeting and non-targeting Cas13 strains.
(B) Escape frequencies of single versus dual Cas13a crRNAs targeting ΦKZ.
(C) Agarose gel PCR verification of *acrVIA1* insertions in transposed plaques. Low-transposition control pools are shown for comparison. Schematic of an example of *acrVIA1* insertion is as shown.
(D) Plaque assays of transposition reactions across targeting and non-targeting hosts.
(E) Quantification of transposition efficiency as a function of input ΦKZ titer (pfu/mL).

Figure S2. Characterization of individual transposition reactions before and after enrichment

(A) Plaque assays comparing ten independent transposition reactions across selection conditions: no selection, dual Cas13a targeting, and JukAB. Wild-type ΦKZ is shown as control.
(B) Full-plate infections (20 μL input) of each transposition reaction.
(C) Post-enrichment plaque assays confirming selective recovery of transposed ΦKZ under Cas13a counter-selection.

Figure S3. Rank-ordered distribution of ΦKZ gene fitness scores

Cumulative ranking of ΦKZ genes based on transposon insertion frequency and essentiality scoring.

Figure S4. Cas13 escaper phages acquire large genomic deletions

(A) Schematic overview of genes targeted by Cas13. Protein function, essentiality, and conservation across jumbo phages and *Chimalliviridae* are indicated.

(B) Plaque assays showing targeting of wild-type ΦKZ and ΦKZ *orf105::acrVIA1* (*acrVIA1* control) with crRNAs directed genes encoding injected structural proteins. The Cas13a dual guide construct targeting *orf*s *55* and *120* are included as control.

(C) Schematic representation of deletion events identified in Cas13 escaper genomes. The annotations start (s), dual (d) and cleavage (c) refer to the position of guide targeting. Numbers shown indicate the base number within the *orf* that had mutations. Top: virion, bottom: non-virion.

Figure S5. Competitive index of ΦKZ mutants

Competitive fitness of ΦKZ mutants carrying *acrVIA1* insertions in different genes, compared with wild-type phage.

non targeting plate    Clinical isolate

Plaque of interest ⟶ Picked ⟶ Propagated

Figure S6. Representation of plaque selection for downstream phenotypic follow-up

Example images of plaques from arrayed ΦKZ mutants pinned onto *P. aeruginosa* lawns.

Figure S7. *In vivo* **EcoRI targeting of ΦKZ and ΦKZ** *nlp1::acrVIA1*

Quantification of *in-vivo* EcoRI restriction of wild-type and ΦKZ *nlp1::acrVIA1*. F8 phage is included as an EcoRI-sensitive control unrelated to ΦKZ.

Figure S8. Adsorption and replication kinetics of ΦKZ *orf302::Tn*

Adsorption and 2-hour replication profile of *orf302::Tn* compared to wild-type ΦKZ.