Systematic yeast two-hybrid screening identifies novel functions for SET1C/COMPASS

Short title : SET1C methylates the ARTSTRGR motif of Snf2 AT-hook

Pierre Luciano[1, ¶], Kihyun Park[2, ¶], Stéphane Audebert[1], Luc Camoin[1], Carlos A. Niño[1], Da Kyeong Park[3], Isabella E. Maudlin[4], Marion Dubarry[1], Lara Lee[1], Marlene Oeffinger[5,6,7], Jean D. Beggs[4], Young Hye Kim[3], Jaehoon Kim[2,*], Bernhard Dichtl[8, *], Vincent Géli[1,*].

[1]Marseille Cancer Research Center (CRCM), U1068 Inserm, UMR7258 CNRS, Aix Marseille University, Institut Paoli-Calmettes, France. Equipe labellisée Ligue.

[2]Department of Biological Sciences, Korea Advanced Institute of Science and Technology, Daejeon, 34141, South Korea.

[3]Ochang Institute of Biological and Environmental Science, Korea Basic Science Institute, Cheongju 28119, South Korea

[4]Wellcome Centre for Cell Biology and Institute of Cell Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3BF, UK.

[5] Institut de recherches cliniques de Montréal, Center for Genetic and Neurological Diseases, 110 avenue des Pins Ouest, Montréal, Québec, H2W 1R7, Canada

[6] Département de biochimie et médecine moléculaire, Faculté de Médicine, Université de Montréal, Québec, H3C 3J7, Canada

[7] Division of Experimental Medicine, Faculty of Medicine, McGill University, Montréal, Québec, H4A 3J1, Canada

[8] School of Life and Environmental Sciences, Deakin University, Geelong, Australia.

* To whom correspondence should be addressed
vincent.geli@inserm.fr
kimjaehoon@kaist.edu
bernhard.dichtl@deakin.edu.au

¶Joint authors

Present Addresses:

Isabella E. Maudlin, Sir William Dunn School of Pathology, University of Oxford, Oxford, OX1 3RE, UK.

Marion Dubarry, Univ Lyon, Université Claude Bernard Lyon 1, INSA-Lyon, CNRS, UMR5240, Microbiologie, Adaptation et Pathogénie, 10 rue Raphaël Dubois, F-69622, Villeurbanne, France

Carlos A. Niño, IFOM ETS, The AIRC Institute of Molecular Oncology, Milan, Italy.

Vincent Géli, Institute for Research on Cancer and Aging, Nice (IRCAN), Nice, France

ABSTRACT

Set1 is the catalytic subunit of SET1C or COMPASS, which methylates histone H3K4 and serves as a scaffold for the association of seven tightly bound polypeptides. We have employed yeast two-hybrid screenings to determine the interactome of Set1 and each subunit, providing a unique resource for exploring known and novel roles of the complex. Our screenings identified a multitude of interactors involved in chromatin regulation, DNA replication, meiotic breaks, and Ty transposition, processes previously associated with SET1C. Consistent with Set1 being an RNA-binding protein, the screens link SET1C to multiple aspects of RNA biogenesis, including pre-mRNA splicing and polyadenylation. The results reveal that Set1 interacts with several importins and with RGG motif-containing proteins, providing insights into the mechanisms by which Set1 moves between cytoplasmic and nuclear compartments. We demonstrate that the transcriptional corepressor Nrm1 is methylated by SET1C *in vitro* suggesting that H3K4-like domains may represent a class of non-histone substrates for SET1C. We further reveal that reconstituted SET1C interacts with the AT hook domain of the chromatin remodeler Snf2 and methylates multiple arginines within this domain. *In vivo*, we report that the ARTSTRGR AT-hook motif is methylated in a Set1-dependent manner revealing new interplay between lysine and arginine methylation.

INTRODUCTION

Post-translational modifications of histones shape the chromatin landscape and provide essential mechanisms of regulating DNA accessibility, thereby controlling gene expression, genome maintenance and transmission. Multiprotein complexes of the SET/MLL family methylate histone H3 on lysine 4 (H3K4), which modulates multiple aspects of genome biology (Ruthenburg *et al*, 2007). Each complex contains additional factors specifying their recruitment to explicit chromatin domains and their specific biological effects (Cenik & Shilatifard, 2021). The budding yeast Set1 complex called COMPASS (for Complex of Proteins Associated with Set1) or SET1C has proved to be an excellent model to study the SET1/MLL family complexes. In *Saccharomyces cerevisiae*, all H3K4 methylation is carried out by SET1C that is composed of Set1, the catalytic subunit, acting as a scaffold for seven other components (Swd1 [RbBP5], Swd2 [WDR82], Swd3 [WDR5], Bre2 [ASHL2], Sdc1 [DPY30], Spp1 [CFP1] and Shg1 [BOD1]) (Nagy *et al*, 2002; Briggs *et al*, 2001; Roguev *et al*, 2001; Miller *et al*, 2001). Swd1, Swd3, Bre2 and Sdc1 copurify with the SET domain of Set1, yet none of these proteins can interact alone with the SET domain, suggesting that interactions between subunits are required for catalytic domain formation (Kim *et al*, 2013). The crystal structure of the SET domain of Set1 associated with Swd1, Swd3, Bre2, Sdc1 shows that the catalytic module is organized by Swd1, whose C-terminal tail nucleates Swd3 and a sub-complex formed by Bre2-Sdc1 (Qu *et al*, 2018; Hsu *et al*, 2018). For its side, Spp1 associates with the N-SET domain of Set1, while Shg1 binds to the central region of Set1 and Swd2 contacts the N-terminal region of Set1 (Kim *et al*, 2013; Roguev *et al*, 2001; Dehe *et al*, 2006; Halbach *et al*, 2009). Swd1 contacts Spp1 and also interacts with the N-terminal region of Set1, suggesting that the C- and N-terminal regions of Set1 may interact with each other (Qu *et al*, 2018; Acquaviva *et al*, 2013b; Jeon *et al*, 2018). Such SET1C organization was

confirmed by cryo-electron microscopy and cross-linking experiments performed on the full-
length complex (Wang *et al*, 2018). Interestingly, SET1C has a remarkable mode of assembly
that is initiated in the cytoplasm while the nascent Set1 polypeptide emerges from the
ribosome. Set1 is initially bound during its translation, by Shg1, Spp1 and Swd1, then Swd2,
Swd3, Bre2 and Sdc1 associate with the initial pre-complex to form the full SET1C. This
explains why Set1 is associated with its own mRNA (Luciano *et al*, 2017; Halbach *et al*,
2009).
Swd2, the only essential SET1C subunit, is also part of the APT complex (for Pta1-
associated), a subcomplex of the Cleavage Polyadenylation Factor (CPF) involved in the
transcriptional termination of mRNA and snoRNA suggesting a functional link between
SET1C and 3' end formation/termination (Cheng *et al*, 2004; Nedea *et al*, 2003; Dichtl,
2004). Set1 has also been linked genetically to the Nrd1-Nab3-Sen1 (NNS) complex (Kim *et*
al, 2006; Arigo *et al*, 2006), but intriguingly, it was reported to have two seemingly
contradictory effects on ncRNA termination: on the one hand Set1 was suggested to promote
efficient snoRNA termination mediated by Nrd1 (Terzi *et al*, 2011), and on the other hand it
appeared to interfere with the early termination of a large number of ncRNAs (Castelnuovo *et*
al, 2014; Margaritis *et al*, 2012). Interestingly, various lysines distributed between Nrd1,
Nab3 and Sen1, are methylated, notably lysine 363 in the RNA Recognition Motif (RRM) of
Nab3 whose mono methylation depends on Set1 and Set3 (Lee *et al*, 2020).
Set1 interacts co-transcriptionally with the RNA polymerase II Carboxy Terminal
Domain (PolII CTD) phosphorylated on Ser5 of the heptad repeats, producing an H3K4
methylation gradient that starts at nucleosome +1 and fades away from the promoter (Soares
et al, 2017; Ng *et al*, 2003). It was recently shown that the N-terminal region of Set1 and
Swd2 interact cooperatively with the CTD (Carboxy Terminal Domain) of Rbp1 (RNA
polymerase II large subunit) promoting SET1C recruitment to transcription elongation
complexes at the 5′ ends of genes (Bae *et al*, 2020). Deletion of residues 200-210 of Set1
abolishes interaction with the Rpb1-CTD and Swd2, while deletion of the first 200 amino
acids of Set1 strongly reduces trimethylation of H3K4 (H3K4me3) at the 5' end of transcribed
genes (Bae *et al*, 2020). Interestingly, Swd2 is ubiquitinated by Rad6/Bre1 and preventing
Swd2 ubiquitination affects H3K4 trimethylation with a concomitant reduction in Spp1
recruitment to chromatin suggesting a cross-talk between Swd2 and Spp1(Vitaliano-Prunier *et*
al, 2008). In a similar vein, di- and tri-methylation of H3K4 by Set1, which depends on prior
ubiquitination of histone H2B by Rad6/Bre1 (Dover *et al*, 2002; Sun & Allis, 2002), requires
a contact between Spp1 and Swd1 (Hsu *et al*, 2019; Jeon *et al*, 2018). Other modes of SET1C
recruitment to chromatin must exist in the absence of the Set1 N-terminal region, probably via
direct interactions between SET1C's catalytic domain and the nucleosome (Jeon *et al*, 2018;
Dehé & Géli, 2006; Thornton *et al*, 2014).
The central region of Set1 contains two RRMs positioned in tandem (Trésaugues *et al*,
2006) through which Set1 binds directly to RNA. This binding contributes to retain Set1 in
the 5' region of genes thus facilitating their H3K4 trimethylation (Luciano *et al*, 2017;
Battaglia *et al*, 2017; Sayou *et al*, 2017). Both the N-SET domain and Spp1 also contribute to
Set1 binding to RNA. Set1 associates post-transcriptionally with transcripts produced by
specific classes of genes including snRNA, Ty1 and a number of genes involved in adaptive
responses (Luciano *et al*, 2017). The dRRM itself is flanked by an autoinhibitory region that

negatively regulates H3K4me3 (Schlichter & Cairns, 2005). It is possible that in the context of full-length Set1, the alternative mode of SET1C recruitment may be inhibited by the autoinhibitory domain.

In *S. cerevisiae*, H3K4me3 methylation is counteracted by the demethylase Jhd2, a conserved JARID1 family protein (Huang *et al*, 2010; Liang *et al*, 2007). The erasure of ancestral histone methylation states results from both active enzymatic demethylation by Jhd2 and passive dilution of parental histones during replication (Radman-Livaja *et al*, 2010). SET1C is a dimer and its dimerization depends on the Sdc1 subunit (Choudhury *et al*, 2019). It has been proposed that the symmetrical methylation of H3K4 on nucleosomes is a consequence of the dimeric nature of SET1C and that Jhd2 preferentially demethylates asymmetrical H3K4me3 (Choudhury *et al*, 2019). Interestingly, H3K4me3 at environmental stress genes depends on H3-P16 isomerisation, a process that controls K4me3 by balancing the actions of Jhd2 and Spp1 (Howe *et al*, 2014).

The complexity of the *S. cerevisiae* SET1C, the specific arrangement of subunits within the complex, and the specific roles of certain subunits, notably Spp1 and Swd2 (Acquaviva *et al*, 2013a), raise the question of the contribution of SET1C and of its subunits to multiple processes. Overall, H3K4 methylation, SET1C and its individual subunits have been involved in multiple processes in *S. cerevisiae* such as meiotic recombination (He *et al*, 2019; Sollier *et al*, 2004; Borde *et al*, 2008; Acquaviva *et al*, 2013b; Sommermeyer *et al*, 2013; Karányi *et al*, 2018; Adam *et al*, 2018), stress response and epigenetic transcriptional memory (D'Urso *et al*, 2016; Kim & Buratowski, 2009; Weiner *et al*, 2012), DNA repair (Faucher & Wellinger, 2010), telomere and rDNA silencing (Jezek *et al*, 2023; Corda *et al*, 1999; Briggs *et al*, 2001; Bryk *et al*, 2002; Santos-Rosa *et al*, 2004; Nislow *et al*, 1997), cell wall biogenesis (Nislow *et al*, 1997), chromosome segregation (Beilharz *et al*, 2017; Zhang *et al*, 2000), antisense transcription (Murray *et al*, 2015; van Dijk *et al*, 2011; Margaritis *et al*, 2012; Castelnuovo *et al*, 2014), transcription termination (Kaczmarek Michaels *et al*, 2020; Nedea *et al*, 2003; Dichtl, 2004; Cheng *et al*, 2004; Nedea *et al*, 2008; Terzi *et al*, 2011; Soares & Buratowski, 2012; Castelnuovo *et al*, 2013; Lee & Wang, 2018), Ty silencing (Luciano *et al*, 2017; Berretta *et al*, 2008), chronological aging (Gong *et al*, 2023; Walter *et al*, 2014; Mei *et al*, 2019), ergosterol homeostasis (South *et al*, 2013), stress response (Deshpande *et al*, 2022; Nadal-Ribelles *et al*, 2015), lipid homeostasis (Giaever *et al*, 2019), and DNA replication (Ghaddar *et al*, 2023; Sollier *et al*, 2004; Rizzardi *et al*, 2012; Chong *et al*, 2020; Delamarre *et al*, 2020; Santos-Rosa *et al*, 2021; de La Roche Saint-André & Géli, 2021; Serra-Cardona *et al*, 2022).

The multiple roles of Set1 and its subunits led us to perform global two-hybrid screening to identify interactors either of Set1-full length, or of its N- and C-terminal regions and of each of the individual subunits (Swd2, Shg1, Spp1, Swd1, Swd2, Sdc1, Bre2). The identification of interactors is discussed not only for each bait, but also as a whole, revealing new potential functions for the complex. In addition, we demonstrate that the transcriptional corepressor Nrm1 and the Snf2 AT-hook are both methylated by SET1C *in vitro*. In vivo, deleting *SET1* abrogates arginine methylation within the ARTSTRGR motif of Snf2 AT-hook. These results suggest that H3K4-like sequences may represent a class of non-histone substrates for SET1C and reveal new interplay between Lysine and arginine methylation. This work is an invaluable

177 resource for further exploring known and unsuspected roles of the SET1C complex and its
178 subunits.
179
RESULTS
Identification of Set1 and Set1 subunits interactors
We produced a total of ten yeast two-hybrid (Y2H) screens (Hybrigenics, See Methods). For
the Set1 protein, we performed a total of three screens using as bait the full-length Set1
protein (Set1 FL), the amino-terminal region including amino acids 1-754 or a C-terminal
fragment including amino acids 754-1081 (Fig. 1A). We have chosen to separate Set1 into
these 2 regions because they have been described as having well-defined properties. The Set1
1-754 fragment includes the domain involved in Set1 recruitment to chromatin, the double
RRM (Trésaugues *et al*, 2006), and the central self-inhibitory domain (Schlichter & Cairns,
2005) that all have regulatory roles (Dehé & Géli, 2006). The Set1 region 754-1081 contains
the N-SET domain, the SET domain and the post-SET domain: the latter is capable of
methylating H3K4 on its own (Thornton *et al*, 2014) (Fig. S1). For each of the subunits (SU)
of the complex whose organization is described in Fig. 1A we used the whole protein as bait.
Each gene encoding Set1 (and its fragments) or its subunits was cloned downstream the Gal4-
BD except for Swd2 that was placed upstream the Gal4-BD. These screens have proven their
power and effectiveness. In particular, they identified Mer2 as an interactor of Spp1
(Acquaviva *et al*, 2013b), and the CTD of Rpb1 as an interactor of the N-terminal region of
Set1 (Bae *et al*, 2020) (Fig. S1). The results of the ten Y2H screens are presented in Fig. 1 and
Table S2. Each interactor is characterised with a confidence score based on several criteria,
including the frequency with which a prey protein is found for a particular screen and the
presence of overlapping fragments, which allow the delineation of the interaction domain
involved (see Methods). Very high confidence interactors (indicated by a color code) are
likely to interact directly with their bait. For Set1 and its N- and C-terminal fragments, the
high confidence Y2H interactors are shown in Fig. 1B. All interactors are shown in Table S2.
The predicted protein-protein interactions within the Set1 interactors reveal functional
information that is detailed in the following paragraphs. The high confident interactors of the
seven SET1C subunits are shown in Fig. 1C-E. We found that Spp1, Shg1 and Swd2 interact
alone with Set1 (Fig. 1C). The minimum Set1 region for which an interaction is found for
each of these 3 subunits is shown in Fig. 1C. The high confidence interactors of the seven
SET1C subunits are shown in Fig. 1C-E. We found that Spp1, Shg1 and Swd2 display Y2H
interactions with Set1 (Fig. 1C). The high confidence interactors of Spp1, Shg1 and Swd2 are
indicated in Fig. 1D (see also Table S2). In contrast, we did not find any interaction between
Swd1, Swd3, Bre2 and Sdc1 (SET-c components) and Set1 in the various Y2H screens,
suggesting that these subunits must cooperate to bind to Set1. Only Bre2 was found to interact
with Sdc1 in both Sdc1 and Bre2 Y2H screens (Fig. 1E). These results are consistent with the
structure of the extended SET1C catalytic module and full-length SET1C (Wang *et al*, 2018;
Hsu *et al*, 2018; Qu *et al*, 2018).
217
218 **Set1 1-754 interacts with the importin Kap104**
219 We found that Set1 1-754 interacted with very high confidence with the importin Kap104,
220 suggesting a direct interaction (Fig. 2A). Kap104 has been reported to recognize specific

cargos containing a specific class of NLS termed PY-NLS (Soniat *et al*, 2013; Xu *et al*, 2010). This PY-NLS includes a N-terminal or central hydrophobic or basic motif, which often contains hydrophobic (R/H/K) residues and which can include RGG repeats (arginine-glycine-glycine motifs) and a C-terminal Proline-Tyrosine (PY) dipeptide near the C-terminus. Such cargos include the mRNA export factor Nab2, the subunit of the THO/TREX complex Hrp1, and the transcription factor Tfg2 (Lee & Aitchison, 1999; Süel *et al*, 2008). Interestingly, Nab2 is a confident interactor of Sdc1 (Fig. 2A). The Set1 interacting domain (SID) of Kap104 extends from residue 359 to 621 and includes the HEAT-like repeat (Yoshimura & Hirano, 2016) (Fig. 2B). Consistent with this result, we identified two PY-NLS in the N-terminal region of Set1 that are likely recognized by Kap104 (Fig. 2C). Along the same line, Shg1, which binds RRM2, interacted with very high confidence with Mog1 (Oliete-Calvo *et al*, 2018). Mog1 has been involved in the modulation of the nucleotide state of Ran-GTP in the nucleus and of Ran-GDP in the cytoplasm, thereby conferring directionality to the nuclear import pathway (Baker *et al*, 2001). Mog1 has been reported to interact directly with the Ran homologue Gsp1 (Oki & Nishimoto, 1998) (Fig. 2A). We have also identified, Kap123, Msn5 (Kap142), Nup82 and Mtr10 as interactors of Set1 1-754 that are also involved in protein import/export (Fig. 2A). Kap123 is a major karyopherin that recognizes NLS of cytoplasmic H3 and H4 (An *et al*, 2017) while Msn5/Kap142 was shown to mediate the import into the nucleus of the subunits of RPA (Yoshida & Blobel, 2001). Nup82 for its part belongs to a module at the cytoplasmic face of the NPC and interacts with karyopherins (Beck & Hurt, 2017) (Fig. 2A). Finally, Mtr10 has been implicated in the nuclear import of the mRNA-binding protein Npl3 (Senger *et al*, 1998; Pemberton *et al*, 1997). Collectively, these multiple interactions reveal insights for understanding the nuclear import of Set1, which was reported to bind co-translationally with Shg1, Swd1, and Spp1 (Halbach *et al*, 2009).

Set1 1-754 interacts with Snf2 and RGG motif proteins

Set1 1-754 and Set1 FL interacted with very high confidence with Snf2, the catalytic subunit of the SWI/SNF chromatin remodeling complex (Côté *et al*, 1994) (Fig. 1B). Interestingly, two additional Snf2 complex components, Snf5 and Swi1, have been identified in the Set1FL screen (Table S2). Together these observations strongly support the idea that the suggested interaction of Set1 and SWI/SNF is of biological significance (Hirschhorn *et al*, 1992). The various Y2H screens revealed Set1 and its subunits interacted with a number of proteins involved in chromatin structure regulation although the relevance of these interactions remains to be demonstrated (Fig. 3). Interestingly, the SID region of Snf2 is juxtaposed to an RGG motif composed of several RGG/RG repeats that is often found in RNA-binding proteins (Thandapani *et al*, 2013) and may be the substrate for arginine methyltranferases (McBride *et al*, 2005) (Fig. 2A, 2D). RGG repeats are multifunctional motifs that mediate RNA and DNA binding and nuclear import. They have been involved in RNA metabolism and chromatin dynamics possibly via arginine methylation, which modulates RNA affinity and nuclear localization (108). Along the same line, Set1 1-754 also interacted with the mRNA export factor Gbp2 (Poornima *et al*, 2021) and the nucleolar protein Nop1, both of which contain an RGG motif (Fig. 2A, 2D). Other RGG proteins such as polyadenylated RNA-binding protein Nab2 and the RNA-helicase Dbp1 interacted with

Set1 754-1081 (and Sdc1) and Spp1, respectively (Fig. 2A, 2D). For each of these proteins,
the putative SID includes the RGG motif. Of note, both Nop1 and Nab2 are methylated by the
Arginine methyltransferase Hmt1 (Smith *et al*, 2020; Green *et al*, 2002).
Our screens indicate that the minimal region of Snf2 interacting with Set1 1-754 was located
between Snf2 residues 1430-1499. We further refined this interaction by Y2H testing of five
Set1 fragments covering the entire Set1 length against the Snf2 1349-1650 region. The two-
hybrid analysis revealed that Snf2 (1349-1650) interacted with the N-terminal domain (Set1
1-230) and the N-SET region (Set1 762-938) (Fig. S2). To validate these interactions we
obtained recombinant protein where GST was fused to the C-terminal domain of Snf2
(Snf2C), the AT-hook (Snf2C-AT-hook) and the Bromo domain (Snf2C-Bromo) (Kim *et al*,
2010) (Fig. 4A). We performed pull-down assays to test the interaction of the GST-fusion
proteins with reconstituted SET1C or with the SET1C762 complex in which Set1 has a N-
terminal truncation of the first 761 residues, respectively (See METHODS) (Fig. S3 A-E). We
found that SET1C as well as the SET1C762 complex were both associated with Snf2C and
Snf2C-AT-hook but not to Snf2C-Bromo (Fig. 4 B, C), indicating that AT-hook is a key
region for interaction with SET1C. We then delineated the minimal AT-hook region required
for interaction with SET1C and found that residues 1461-1547 are essential for binding to
SET1C (Fig. 4 D-E). Collectively, these results indicate that SET1C interacts with Snf2C-AT-
hook region suggesting that the Set1 and Snf2 complexes cooperate to modify chromatin
structure.
285
SET1C is involved in multiple aspects of RNA biogenesis
Set1 was previously shown to bind RNA nascent transcripts through its dRRM contributing to
position Set1 and H3K4me3 predominantly to the 5′ regions of genes. Of note, Set1 showed a
higher occupancy within introns, at transcripts from ribosomal DNA (rDNA), and tRNAs
(Luciano *et al*, 2017; Battaglia *et al*, 2017; Sayou *et al*, 2017). Moreover, Set1 also binds
post-transcriptionally to Ty1 retrotransposon transcripts and mRNA encoding specific
transcription factors genes (Luciano *et al*, 2017). The SET1C Y2H interactome revealed that
many proteins involved in RNA biogenesis were interacting directly or indirectly with Set1 or
its subunits (Fig. 5). We find a strong connection between SET1C and pre-mRNA processing,
as highlighted by the identification of multiple spliceosome subunits (Fig. 5). Some
spliceosome subunits were found in several screens (Fig. 1B, D; Table S2). For instance, the
Prp8 subunit of the U5 snRNP that function in critical molecular rearrangements during the
splicing process (Grainger & Beggs, 2005) interacted with Set1 FL, Spp1, and Swd1.
Similarly, Prp22 interacts with Set1FL and Set1 754-1080. Prp22 is a DEAH-box helicase
that associates with newly spliced mRNA and promotes its release from the spliceosome (Will
& Lührmann, 2011). We have refined the interaction region between Set1, Prp8 and Prp22,
showing that Prp8 and Prp22 interact strongly with Set1-F4. (n-SET). Prp22 interacts in
addition with Set1-F1 (Fig. S2). We have further strengthened the link between Set1 and
Prp22 by showing that Set1 is co-immunoprecipitated with Prp22 *in vivo* in an RNA-
independent manner. (Fig S4A, B). The preferential binding of Set1 to genes with introns
(Luciano *et al*, 2017) and its interaction with splicing factors, in particular Prp22, suggest that
Set1 may be involved in late splicing events. Alternatively, Prp22 and/or other splicing factors
could be involved in H3K4 methylation.

We also found a number of factors involved in rRNA processing that could be related to the
binding of Set1 to ncRNA transcripts derived from the rDNA intergenic spacer regions
(Sayou *et al*, 2017) (Fig. 5). Next, and consistent with the observation that SET1C regulates
the choice of the polyadenylation site and the recruitment of the cleavage/polyadenylation
complex (Kaczmarek Michaels *et al*, 2020), the Y2H screens revealed multiple potential
interactions between SET1C and factors involved in RNA polyadenylation (Fig. 5). Some of
these interactions are known, such as the interactions between Swd2 with Pta1 and Ref2, all
three proteins belonging to the APT complex (Nedea *et al*, 2008). Interestingly, we found that
Set1 1-754 interacts with Not1 and Not4 that belongs to the ubiquitin-protein ligase CCR4-
NOT (Liu *et al*, 2001) (Fig. 5). This complex was shown to be involved in the regulation of
H3K4me3 via a ubiquitin-dependent pathway (Mulder *et al*, 2007; Laribee *et al*, 2007) that
was subsequently linked to Jhd2 degradation (Huang *et al*, 2010; Mersman *et al*, 2009). These
results raise the question of which of the Y2H interactions described in this study are linked to
the regulation of H3K4 methylation states.
Finaly and somewhat surprisingly, we found that many Y2H interactors of Set1 1-754 were
involved in tRNA nuclear transport, modification, and synthesis (Fig. 5). Swd2, which
interacts with the Set1 N-terminus, exhibit a 2H interaction with a high degree of confidence
with the cytoplasmic asparaginyl-tRNA synthetase Ded81 and the prolyl-tRNA synthetase
Yhr020 while Sdc1interacted with the glycyl-tRNA synthase Grs1 and Bre2 with the
glutamine tRNA synthetase Gln4. Of note, Trm1 (Liu *et al*, 1998) and Trm732 (Guy *et al*,
2012) are both directly or indirectly involved in tRNA methylation and form part of what has
been defined as the cell's global methytransferome (Giaever *et al*, 2019).
331
SET1C Y2H interactors are involved in several aspects of DNA transactions
As mentioned in the introduction, it has been shown that SET1C, and in particular Spp1,
regulate the selection of meiotic breaks, the progression of the replication fork, DNA repair,
chromosome segregation, and transposition of Ty elements (Deshpande & Bryk, 2023). We
have classified all Set1 and subunit interactors according to these SET1C roles (Fig. S5). We
found a number of interactors involved in meiosis. In particular, Spp1 not only interacts with
Mer2, but also with the meiosis specific DNA helicase Mer3 (Nakagawa & Kolodner, 2002)
that interacts also with Swd1 (Fig. S5). We further found Set1 interactions with the
kinetochore proteins Spc25 and Cbf2 suggesting that Set1 could be transiently localized at the
spindle pole body. Regarding the biology of retrotransposons, we previously reported that
Set1 binds post-transcriptionally to Ty1 mRNA and represses Ty1 mobility (Luciano *et al*,
2017). Interestingly, the two-hybrid screens reveal that Set1 1-754 interacted with Gag
capsid-like proteins of Ty1 (Fig. S5) raising the possibility that Set1 binding to Ty1 mRNA is
linked to the interaction of Set1 1-754 with Gag. Concerning the role of Set1 and subunits in
DNA replication and repair, we report multiple potential interactions (Fig. S5). In particular,
Swd2, Swd1, and Spp1interacted with high confidence with Orc6, Nrm1, and Mcm2,
respectively (Fig. 1 and Fig. S5). Orc2 was previously described to interact physically with
Spp1 (Kan *et al*, 2008). We recently reported that Spp1 is recruited at replication forks stalled
at the Tus/Ter barrier independently of its interaction with Set1 (Ghaddar *et al*, 2023).
Interestingly, in the Y2H screen Spp1 interacted with the extreme C-terminal region of Mcm2
(791-867) (Fig. S6A) that corresponds to a non-conserved accessible alpha-helix within the

MCM complex (Li *et al*, 2015b). We fused Spp1 and Mcm2 to GST and performed GST pull-
down experiments. We confirmed *in vitro* in both directions a weak interaction between Spp1
and Mcm2 (Fig. S6B). Whether Spp1 is recruited by Mcm2 at stalled replication fork remains
to be determined. On its side Nrm1 inactivates MBF, a major regulator of the G1/S
transcription (de Bruin *et al*, 2006). During replication stress, Nrm1 phosphorylation by the
checkpoint kinase prevents its binding to MBF target promoters leading to the activation of
G1/S transcription (Travesa *et al*, 2012). An exciting idea is that Swd1 recruits Nrm1 to
stalled forks by promoting its phosphorylation by Rad53. Swd1 would play a role in linking
replication stress and transcriptional regulation via Nrm1. Of note, Nrm1 was identified as a
gene required for the cell-cycle pattern of H3K79me2 during early S phase (Schulze *et al*,
2009). The interaction between Swd1 and Nrm1 is described in more detail below.
Finally, Y2H screening indicated that Set1 and its subunits interacted with a number of
proteins involved in protein SUMOylation (Fig. S5). The proteins were either involved in
SUMO conjugation or SUMO-dependent degradation. Remarkably, the C-terminus of Nis1
(360-407) that contains a potential SUMO-binding site (Hannich *et al*, 2005) was identified a
high-confidence interactor of Spp1, Shg1, and Sdc1 (Fig. 1). Nis1 is localized at the bud neck
(Iwase & Toh-e, 2001) at the vicinity of the septin collar containing several highly
SUMOylated proteins (Shs1, Cdc11) (Wykoff & O'Shea, 2005) and has been implicated in
preventing bud recovery at the site of division (Meitinger *et al*, 2014). We thus sought to
confirm biochemically the interaction of Nis1 with the three SET1C subunits. We fused the
GST to Spp1, Sdc1, and Sgh1 to perform pull-down experiments with *in vitro* translated Nis1.
We confirmed the interaction of Nis1 with Spp1, and Sdc1, but not with Sgh1 (Fig. S7).
However, mass spectrometry analyses on TAP-Nis1 did not reveal the presence of SET1C
subunits (Table S3) suggesting that interaction between Nis1 and Spp1/Sdc1 might be
transient. The relevance of the Nis1 and other putative interactors remains unclear, especially
as many of those proteins are not known to be located in the nucleus. Interestingly, Nis1 has
been reported to shuttle from the bud neck to the nucleus when overexpressed (Perez &
Thorner, 2019). In Fig. S8, we have identified a number of interactors, including Sdc1 and
Spp1, showing changes in localization following hypoxia (Henke *et al*, 2011). Of note, one of
the two human SET1C homolog SET1B has a cytoplasmic location with functions unrelated
to H3K4 methylation (Wang *et al*, 2017). Combined, these observations raise the question of
the minor or transient localization of Set1 and its subunits outside the nucleus, or conversely
of the transient localization of interactors within the nucleus.
386
Set1 is SUMOylated
Global analyses of SUMOylated proteins in fission yeast revealed that Set1 and Spf1 (Spp1)
were SUMOylated (Shin *et al*, 2005; Nie *et al*, 2015). As Set1 had two-hybrid interactions
with Slx5 and Wss1, both of which act on SUMOylated proteins as they undergo protein
degradation (Mullen *et al*, 2010), we tested whether Set1 can be SUMOylated. Cells
expressing Myc-Set1 (Dehe *et al*, 2006) or transformed with pPB66-SET1 expressing GBD-
Set1-FL (GAL4 binding domain) were transformed with a plasmid encoding His_6-SUMO or a
control plasmid. Proteins were purified on Ni-NTA agarose beads and detected by Western
blot either using anti-MYC or anti-GAL4 antibodies. The results revealed that Set1 can be
mono-SUMOylated (Fig. 6A) or di-SUMOylated (Fig. 6B). This difference may be due to the

fact that GBD-SET1-FL is under the control of the ADH promoter in plasmid pGB66 and thus
overexpressed.

We sought to refine the SUMOylated Set1 region. We transformed the plasmid encoding
His6-SUMO and the control plasmid into cells expressing Set1 fragments F1, F2, F3, F4,
F3+F4, and F5 (Fig. S9A). We expressed F3+F4 instead of the isolated F3 and F4 fragments
in order to preserve the K769-centered motif predicted to be highly SUMOylated (Fig. S9B).
F1, F2, F3, and F4 were not SUMOylated (not shown) while the F3+F4 fragment was clearly
SUMOylated (Fig. 6C). In order to identify whether K769 is important for F3+F4
SUMOylation, we mutated K769 to R. We show that introduction of K769R into F3+F4
abolishes its SUMOylation (Fig. 6C), indicating that K769 is likely to be the SUMOylated
lysine within the F3+F4 fragment. We then found that F5 was also SUMOylated (Fig. 6D). In
this case, the substitution K1055R, K1060R, or the double substitution, does not affect the
SUMOylation of F5 indicating that this motif also predicted to be SUMOylated with a high
score is not the SUMOylated motif (Fig. 6D). Taken together, these experiments indicate that
Set1 can be SUMOylated in the N-SET and SET domains, in the interaction region of Spp1
and Swd1-Swd2-Bre2-Sdc1, respectively. This raises the possibility that SUMOylation
regulates the interaction of Set1 with its subunits, in particular with Spp1, which interacts
dynamically with SET1C (Ghaddar *et al*, 2023; D'Urso *et al*, 2016; Karányi *et al*, 2018;
Serra-Cardona *et al*, 2022). Interestingly, in mammals, the SUMO peptidase SENP3 (the
ortholog of Ulp1 in budding yeast) interacts with MLL1 and MLL2, catalyzing the
deSUMOylation of RbBP5 (Swd1). This process regulates the association of specific subunits
of MLL1/MLL2, such as menin and Ash2L (Bre2), with the DLX3 gene, which plays a role in
osteogenic differentiation (Nayak *et al*, 2014).

421   The transcriptional corepressor Nrm1 interacts with SET1C

Screening with *SWD1* yielded a total of forty clones, fourteen of which contained fragments
of the *NRM1/YNR009w* gene, which encodes a basic protein of 249 amino-acids (Fig. 7A). Its
amino-terminus carries a putative D-box motif that was found to destabilize the protein (de
Bruin *et al*, 2006). The Y2H results furthermore delineated a central domain (amino-acids 58
to 195) mediating the interaction with Swd1. A sequence motif within this domain displays
strong similarity to the amino-terminus of histone H3; lysine 118 within Nrm1 (Nrm1-K118)
aligns with lysine 4 of histone H3, the substrate of SET1C. We refer to this sequence as
H3K4-like domain. The similarity of the H3K4 like domain and the H3K4 modification site
suggested that Nrm1 may represent a non-histone substrate for SET1C. To test this idea, we
initially performed *in vitro* methylation experiments with partially purified SET1C, S-
adenosyl (methyl-^{3}H) methionine and peptide substrates. We found that a H3K4 peptide was
efficiently methylated, but a peptide carrying a H3K4A substitution was not (Fig. 7B). Also,
no methylation was observed with a peptide encompassing K118 of Nrm1. The possibility
remained that this peptide lacked parts of Nrm1 that were required for methylation. For this
purpose, we obtained recombinant protein where the N-terminus of Nrm1 was fused to the 42
kDa maltose binding protein (MBP). Interestingly, we observed *in vitro* methylation of this
protein by SET1C but not with MBP only (Fig. 7C). Note, however, that histone methylation
occurred much more efficiently under these reaction conditions. Next, we asked whether
K116 and K118 of Nrm1 were required for the observed methylation activity. Fig. 7D shows

that wild-type and mutant MBP-fusion proteins were found to be methylated suggesting that
the H3K4-like domain is not required for the observed SET1C methylation activity.
The H3K4-like domain in Nrm1 raised our attention to other yeast proteins that carry such
sequences. We used the scansite search algorithm (http://scansite.mit.edu) to systematically
identify sequence motifs that are related to the SET1C modification site in histone H3. Search
parameters included four to six identical residues of the ARTKQT sequence that is found at
the H3K4 modification site. In addition, we did allow for biochemically equivalent amino
acid changes. Fig. 7E shows eight candidate proteins based on their function and the sequence
context of the H3K4-like domain. Importantly, some of these gene products are involved in
cellular pathways that functionally overlap with SET1C, e.g. transcription (Not5), rDNA
silencing (Irs4) and cell-cycle control (Dbf2 and Dbf20), increasing the possibility of
occurring physical and functional interactions.
453
SET1C Y2H interactors regulate metabolism and stress responses
Set1 was reported to regulate ergosterol levels (South *et al*, 2013). We identified Erg9, Ugt51,
Vhr1, Vhr2 and Ste20, all involved in ergosterol metabolism (Daicho *et al*, 2020; Lees *et al*,
1995; Warnecke & Heinz, 1994). Erg9 and Ste20 interacted with Set1 while Ugt51 and
Vhr1/Vhr2 were high confidence interactors of Shg1 and Sdc1, respectively (Fig. S10).
Along the same line, the various Y2H screens revealed numerous genes involved in
phosphatidyl inositol metabolism interacting with Set1 and its subunits. In particular Vip1 is
a high-confidence interactor of Set1 754-1081 (Fig. 1B). Vip1 is a bifunctional inositol
pyrophosphate kinase and phosphatase that regulates IP7 levels in the inositol pyrophosphate
(PP-IP) synthesis pathway (Lee *et al*, 2007; Mulugu *et al*, 2007). Interestingly, Vip1 was
reported to regulate the environmental stress response (ESR) through IP7 that activates the
HDAC Rdp3 (Worley *et al*, 2013) suggesting potential new avenues to explain ESR
regulation by Set1 (Weiner *et al*, 2012). Along the same line, Swd1 and Swd3 were identified
in a screen aimed to identify genes that negatively regulate the PHO pathway in a Vip1-
dependent manner (Choi *et al*, 2017). How the putative interaction between Set1 754-1081,
which also interacts with Pho23 subunit of the Rpd3L complex, and Vip1 fits into these
processes remains to be discovered. Of note, Vip1 has been shown in *Arabidopsis thaliana* to
change localization upon hypo-osmotic stress from the cytosol to the nucleus (Takeo & Ito,
2017). On their side, Y2H Interactors identified as stress-responsive genes interact either with
Set1 1-754 or Swd2 (Fig. S10). Swd2 interacted with high confidence with Nar1, an essential
Fe/S protein required for the assembly of cytosolic Fe/S proteins (Balk *et al*, 2004) and the
calcineurin phosphatase Cmp2 activated in response to ER stress (Mizuno *et al*, 2018). The
Swd2/Nar1 interaction can be linked to the fact that Set1 754-1081 interacts with relatively
good confidence with TyW1 according to hybrigenics criteria (see Methods), an iron-sulfur
protein required for synthesis of wybutosine modified tRNA (Noma *et al*, 2006). In contrast
to the stress genes mentioned above, many interactors involved in glucose metabolism
(repression) interact only with the Set1 1-1081 or with members of the nSET module (Fig.
S10).
482
Reconstituted Set1C methylates the Snf2 RG motif within the AT-hook *in vitro*

We have shown above that SET1C interacts with Snf2C-AT-hook region (Figure 4). We thus
tested whether the Snf2C-AT-hook could be methylated by Set1C. Set1C and truncated
Set1C were reconstituted and affinity purified from Sf9 cells coinfected with baculoviruses
that express FLAG-Set1 (or truncated Set1) and the seven other untagged subunits (Kim *et al*,
2013)(Fig. S3). We incubated reconstituted Set1C with the Snf2C, AT-hook and Bromo
fragments in the presence of radioactive S-adenosylmethionine (^{3}H-SAM). The results
indicate that Set1C methylates *in vitro* purified Snf2C and Snf2C-AT-hook, but not Bromo
(Fig. 8A, B) and that Snf2C-AT-hook methylation requires Set1 FL (Fig. 8C). Collectively
these results show that the Snf2C-AT-hook (1384-1547) interacts and is methylated by Set1
FL *in vitro*. Lys 1494 (K1494) and Lys 1498 (K1498) located between the AT-hook domains
of Snf2 were previously shown to be acetylated by Gcn5 (Kim *et al*, 2010). We individually
mutated all the Lys of the Snf2-AT-hook into Arg and tested the methylation of the mutated
fragments. None of the substitutions abolished or decreased the methylation of the AT-hook
indicating that he AT-hook could be methylated on multiple sites or on other type of residues
(Fig. S11). We thus sought to identify the minimal region within the Snf2-AT-hook that is
methylated by Set1C *in vitro*. The Snf2-AT-hook domain was divided in 6 regions that were
fused to the GST (Fig. 8D). *In vitro* methylation assays indicated that the Snf2-B1, B2, B3
were methylated by the reconstituted Set1C (Fig. 8E). As the Snf2-B3 fragment that contains
the last four lysines (K1488, K1494, K1498, K1526) of Snf2-AT-hook domain was still
methylated, it suggested that one of several of these lysines are methylated by Set1C. We then
examined which of these 4 lysine residues are methylated by Set1C *in vitro*. To our surprise,
mutating all the lysine residues of the Snf3-B3 fragment did not abolish the methylation of
Snf3-B3 (Fig. 8F). These results led us to think that Set1C could methylate arginine residues,
in particular those contained in the RGG motif of the Snf2-B3 fragment. We thus deleted the
RG repeats of Snf2-B3 (Fig. 8G) and purified the *Snf2-B3ΔRG*. We found that deletion of the
Snf2-B3 RG repeats abolished the interaction between Snf2-B3 and Set1 and methylation of
Snf2-B3 (Fig. 8H, I). The fact that both, interaction and methylation are lost upon deletion of
the RG motif argues in favor that Set1C, and not a potential contaminant from the
reconstituted Set1C, is responsible for the methylation of the Snf2-B3 RG repeats. Individual
deletion of pairs of arginine residues in the RG motif did not suppress methylation of the
Snf2-B3 fragment, suggesting flexibility in Set1C's ability to methylate Snf2-B3 (Fig. S12).
To further confirm that Snf2-B3 is methylated on arginine, we incubated a mutant Snf2-B3 (in
which lysines were mutated in alanine to limit protease digestion prior mass spectroscopy
analysis) with reconstituted Set1C and SAM and further purified the Snf2-B3 mutant on a Ni-
NTA resin (Fig. 9A-D). We then analyzed by mass spectroscopy arginine methylation in
Snf2-B3. We found that R1490, R1501, R1505, R1507 and R1517 were mono- and di-
methylated while R1509, R1513, and R1519 were monomethylated (Fig. 9E). We thus
confirmed that reconstituted Set1C has the ability to methylate in vitro multiple arginines in
the Snf2B3 region.
523
Snf2 is methylated *in vivo* on its AR**TST**R**G**R **AT-hook motif in a Set1-dependent way**
Our results show that an activity associated with reconstituted Set1C is capable of methylating
arginines in the vicinity and in the RG repeats of Snf2C-AT-hook. The fact that only Set1FL
is capable of methylating the AT-hook is in favor of this activity being associated with Set1C.

Nevertheless, one cannot formally rule out that methylation the AT-hook is due to a
contaminant from insect cells from which Set1C has been reconstituted.
We thus investigated the interaction between Snf2 and Set1 *in vivo* and whether Snf2 is
methylated by Set1C. We found that Set1 was coimmunoprecipitated with Snf2 tagged with a
C-terminal Myc epitope (Fig. 10A). In contrast to what we observed *in vitro*, deletion of the
RG motif of Snf2 did not affect the interaction between Snf2 and Set1, which is consistent
with the fact that the SID of Snf2 is located upstream of the RG repeats (Fig. 10A).
To further investigate Snf2 methylation by Set1C *in vivo*, we purified Snf2-GFP and
its subunits with GFP nanobodies in the presence or absence of Set1 (Fig. 10B). The protein
composition of the complex was determined by mass spectrometry in the presence of absence
of Set1 (Figure 10B, Table S5). We recovered in both cases all the known components of the
Snf2 complex (Snf2, Swi1, Swi3, Snf5, Snf12, Swp82, Arp9, Arp7, Snf6, Rtt102, Snf11,
Taf14) (Table S5, Fig. S13A). Interestingly, we found that the arginine methyltransferase
Rmt2 was found specifically enriched with the Snf2-GFP complex in *set1Δ* cells (Fig. S13).
We then excised the Snf2-GFP gel band and subjected it to mass spectrometry. We
examine the post-translational modifications (PTM) in the Snf2 (1485-1549) region in either
WT or *set1Δ* strains (Table S6). We were unable to purify a peptide containing the RG repeats
but we could identify the peptides flanking the RG motif (Fig. 10C). Fig. 10C represents the
combined results of two independent experiments. In WT cells, we found that the region
between K1494 and R1505 was reproducibly subjected to multiple PTM (Fig. 10C, upper
panel). K1494 and K1498 that were previously found acetylated by Gcn5 (Kim *et al*, 2010)
were found acetylated, however both lysines were also found to be methylated (Fig. 10 C). Of
note H3K4 that is methylated by Set1C has been reported to be also acetylated by Gcn5
(Guillemette *et al*, 2011). T1995, S1996, S1999, T1502, S1503, and T1504 were found
phosphorylated. Interestingly, R1501 was found mono-methylated while R1505 and R1507
were found di-methylated in the two independent Snf2-GFP purifications. These 3 arginines
belong to a ARTSTRGR motif that lies in the Snf2-B3 fragment. They were also found to be
methylated *in vitro* by the reconstituted Set1C (Fig. 9E). We then examine the
posttranslational modifications of the peptides flanking the RG motif in the *set1Δ* strain (Fig.
10C, lower panel). Phosphorylation of serines and threonines was not modified in the *set1Δ*
strain except for T1502. In the *set1Δ* strain, K1494 and K1498 remained methylated
indicating that methylation of both lysines does not depend on Set1C. Strikingly, deleting
SET1 abolished the mono-methylation of R1501 and the di-methylation of R1505, R1507
(Fig. 10C, lower panel). The results thus show that SET1C is required for the methylation of
the three arginines of the ARTSTRGR motif, in agreement with the *in vitro* results.
563
564
DISCUSSION
We have exploited the power of Y2H screening technology to establish an extended
interaction network for the histone H3K4 methylase complex SET1C. Comprehensive data
integration unveiled many potential functions of either the whole complex or individual
subunits. This study is partly validated by the fact that we found interactors in known
functions of SET1C, or putative functions for which the mechanism is unknown. We have
also provided a number of validations by confirming the interaction through biochemical

approaches. It seems likely that some proteins interact transiently with Set1 and its subunits,
or under specific conditions of stress or nutrient limitations, explaining why these interactors
have not previously been found associated with SET1C in biochemical approaches. The most
illustrative case for this is the high-affinity interaction between Spp1 and Mer2 that takes
place during the first meiotic division, which we had identified from the Spp1 screen and
which has been extensively characterized (Acquaviva *et al*, 2013b). Nevertheless, considering
the multitude of interactions identified, it seems unlikely that they all represent direct
contacts, especially where multiple subunits of complexes or multiple components of a given
metabolic pathways interact with Set1C. One caveat is that Y2H interactions can be mediated
by endogenous proteins that bridge the interactions. But this does not make them less
functionally significant. We anticipate that this systematic screen will be an invaluable
resource for further investigation of the role of SET1C in known and novel processes revealed
in this study. Processes discussed here include Set1's export/import mechanism through the
nucleus, interaction with RGG proteins some of which are targeted by arginine
methyltransferase, cooperation of SET1C with chromatin remodeling factors in particular
Snf2C, interaction with splicing factors notably with Prp8 and Prp22, coupling replication
with histone deposition (Mcm2/Spp1), SET1C plasticity regulated by SUMOylation, and Ty
element transposition. A recurrent question related to the identification of potential SET1C
interactors is whether they could be methylated by SET1C. Based on the identification
of domains that closely resembles the histone H3 modification site, we have shown that the
Swd1 interactor Nrm1 is methylated by SET1C *in vitro* suggesting that proteins with H3K4-
like domains represent a class of non-histone substrates that may be regulated by SET1C.

We show that Set1C interacts both *in vitro* and *in vivo* with Snf2. Strikingly,
reconstituted Set1C from insect cells is able to methylate *in vitro* the three arginines located
within the A**R**TST**R**G**R** motif of the AT-hook of Snf2. The *in vivo* results confirm that Set1 is
required for the methylation of these arginines within the A**R**TST**R**G**R** motif. It is interesting
to note that this motif resembles the A**R**TKQTA**R** motif of the N-terminal H3. In yeast, H3R2
is mono- and di-methylated and its asymmetric di-methylation prevents H3K4me3 (Kirmizis
et al, 2009, 2007) . In contrast symmetric di-methylation of H3R2 correlates with H3K4me3
and requires Set1 *in vivo* but not the classical arginine methyltransferases: Hmt1, Hmt2,
Rkm2, Rkm3, Rkm4, Hsl7, and Efm1 (Li *et al*, 2015a; Yuan *et al*, 2012). It has been
proposed that Set1C could be responsible for both H3R2me2s and H3K4me3 or that
H3R2me2s deposited by a yet unidentified arginine methyltransferase requires the prior
deposition of H3K4me3 (Yuan *et al*, 2012).

Interestingly, previous tandem affinity protein purification of the mono- and
asymmetric arginine methyltransferase Hmt1 (also termed Rmt1) identified 108 proteins
associated with it (Jackson *et al*, 2012), a number of which were shown to be methylated.
Snf2 was found to be copurified with Hmt1 *in vivo* and methylated *in vitro* by Hmt1 (Jackson
et al, 2012) making Hmt1 a potential candidate for methylating Snf2. *In vitro*, purified Hmt1
catalyzes mono-methylation and asymmetric di-methylation of H3R2, however,
deleting *HMT1* does not affect the asymmetric di-methylation of H3R2 *in vivo*, which
suggests that several methyltransferases could act redundantly on H3R2 (Li *et al*, 2015a). In
mammals, H3R2 and H3R8 are asymmetrically di-methylated by PRMT6 (Hamey *et al*, 2021;
Guccione *et al*, 2007; Hyllus *et al*, 2007). It has been also reported that PRMT5 can be found

associated with hSWI/SNF and has the ability to methylate H3R8 (Pal *et al*, 2004). We did
not identify Rmt1 in our purification but found that the arginine mono-methyl transferase
Rmt2 enriched with Snf2-GFP in *set1Δ* cells. Rmt2 is a type IV methyl transferase that was
reported to methylate the ribosomal protein L12 (Low & Wilkins, 2012; Chern *et al*, 2002).
We could envision the possibility the possibility that Set1C could cooperate with distinct
protein arginine methyl transferases to promote the mono-methylation of R1501 and the di-
methylation R1505 and R1507 within the A**R**TST**R**G**R** motif. However, it was previously
reported that the combined inactivation of Rmt1, Rmt2 and Hsl7 did not affect the mono-
methylation of H3R2, weakening this hypothesis (Kirmizis *et al*, 2009). It is possible that in
yeast there is a redundancy of enzymes capable of methylating the Snf2 A**R**TST**R**G**R** motif,
but that for each of them their activity on this motif depends on the presence of Set1.
Alternatively, Set1C could directly methylate the A**R**TST**R**G**R** motif as discussed for the
A**R**TKQTA**R** H3 motif (Yuan *et al*, 2012). The fact that Set1C interacts with Snf2, Gbp2,
Nop1, Nab2, Dbb1 (Fig. 2), all of which have RG motifs and are mostly Hmt1 substrates,
raises the possibility of a general interplay between methylarginine and methyllysine.
Interestingly, a very recent article shows that PRMT1 binds to the N-terminal region
of MLL2 and methylates multiple arginine residues within its RGG/RG motifs (An *et al*,
2025).
634
635 **MATERIALS and METHODS**
636 **Strain construction**
637 All strains and plasmids used in this study are listed in Table S1. To obtain gene deletions and
638 expression of tagged proteins, we amplified by PCR a disruption or a tagging cassette
639 containing the appropriate marker as described (Janke *et al*, 2004).
640
Yeast two-hybrid analysis
Yeast two-hybrid screening was performed by Hybrigenics Services, S.A.S., Evry, France
(73). The coding sequence for Set1-FL, Set1 1-754, Set1 754-1081 and Bre2 were cloned into
pB66 (GAL4-bait) as C-terminal fusions to the Gal4 DNA-binding domain while those of
Swd1, Swd3, Sdc1, Spp1, and Shg1 were cloned into pB27 as a C-terminal fusion to LexA
(LEXA-bait). *SWD2* was cloned into pB43 as a N-terminal fusion (SWD2-GAL4). Individual
bait cloning was performed by using specific primers for PCR for every bait. Every PCR
fragment subcloned as bait is entirely sequenced to avoid eventual mismatches in the coding
sequence. The constructs were used as baits to screen a genomic *S. cerevisiae* library
constructed into pP6 based on the pGADGH plasmid (Wilson *et al*, 1993). pB6, pB66 and
pB43 are derived from the original pAS2Δ vector (Fromont-Racine *et al*, 1997). The library
was submitted to very strict quality controls. Each protein is represented by several fragments
(domains) and the library has been screened with a set of 6 reference baits before any other
bait proteins are screened.
The Gal4 constructs were screened using a mating approach with YHGX13 (Y187 *ade2-*
101::loxP-kanMX-loxP, MATalpha) and CG1945 (*MATa*) yeast strains. The LexA constructs
were screened using a mating approach with YHGX13 (Y187 *ade2-101::loxP-kanMX-loxP,*
mat alpha) and L40ΔGal4 (*MATa*) yeast strains as previously described (Fromont-Racine *et*
al, 1997). Positive clones were selected on a medium lacking tryptophan, leucine and

histidine and supplemented with 3-aminotriazole (3AT) if necessary to handle bait auto-
activation. The prey fragments of the positive clones were amplified by PCR and sequenced at
their 5' and 3' junctions. The resulting sequences were used to identify the corresponding
interacting proteins in the GenBank database (NCBI) using a fully automated procedure.
A confidence score (PBS, for Predicted Biological Score) was attributed to each interaction as
previously described (Formstecher *et al*, 2005). The confidence score relies on two different
levels of analysis. Firstly, a local score takes into account the redundancy and independency
of prey fragments, as well as the distribution of reading frames and stop codons in
overlapping fragments. Secondly, a global score takes into account the interactions found in
all the screens performed at Hybrigenics using the same library. This global score represents
the probability of an interaction being nonspecific. For practical use, the scores were divided
into four categories, from A (highest confidence) to D (lowest confidence). A fifth category
(E) specifically flags interactions involving highly connected prey domains previously found
several times in screens performed on libraries derived from the same organism. Finally,
several of these highly connected domains have been confirmed as false-positives of the
technique. The PBS scores have been shown to positively correlate with the biological
significance of interactions. When possible, the bait interacting domain of each prey is
provided. The PBS scores have been shown to positively correlate with the
biological significance of interactions (Rain *et al*, 2001). e-values for the interactions are
available in the Hybrigenics database.
680
681

Protein expression and protein interaction assays
Protein expression was done in *Escherichia coli* BL21 cells and purification of GST-fusion
proteins were as described (Dichtl *et al*, 2002). MBP-fusion proteins were purified according
to manufacturer`s instructions (New England BioLabs, Beverly, MA, USA). For His-tagged
proteins expressed in bacteria, cDNAs were inserted into the pET28 vector (Novagen),
expressed in *Escherichia coli*, and purified using Ni-NTA beads (Qiagen) following the
previously described procedure (Kim & Roeder, 2011). GST pull-down assays were done as
described (Dichtl *et al*, 2002).
A baculovirus expression system was used to express and reconstitute Set1 complexes
containing either FLAG-Set1 or FLAG-Set1-C762. cDNAs were subcloned into
pFASTBAC1, with or without an epitope tag, and baculoviruses were produced following the
manufacturer's instructions (Gibco-Invitrogen). Sf9 cells were infected with various
combinations of baculoviruses, and the complexes were purified by affinity chromatography
using M2 agarose, as previously described (Kim & Roeder, 2011).
696

TAP-Nis1 affinity purification and mass spectrometry analysis
TAP-Nis1 and control cells (W303a) were grown in YPD, harvested in logarithmic phase
(OD600 0.6-0.85) and cryo-lysed as previously described (Trahan & Oeffinger, 2022).
Affinity purification was performed in RNP100 (20 mM HEPES-KOH pH 7.4, 100 mM
NH$_4$OAc, 0.5% Triton X-100, 0.1% Tween 20, 1:100 solution P, 1:5000 antifoam A, 100 mM
NaCl) as described in (Trahan & Oeffinger, 2022). Following washes, the samples were on-
bead trypsin digested in a volume of 50µl (f. c. 20 µg/mL trypsin (Sigma, proteomics grade)

in 20 mM Tris-HCl (pH 8.0), 37 °C, 900 rpm, 16-20 h; stopped with 2% formic acid) and

analyzed by tandem mass spectrometry as described in (Trahan & Oeffinger, 2022) using a

70-minute gradient on a LTQ Orbitrap Velos is a hybrid mass spectrometer (ThermoFisher

Scientific) in Data-dependent mode. Data were processed with Thermo Excalibur to generate

a raw file. Mascot search server (53 Version 2.3.02) was used with a parent tolerance of 10

ppm for precursor ions, 0.52Da for fragments, and only considering 1 possible missed

cleavage as well as a mass change of +16 for methionine oxidations in the mass calculation.

Data were searched against *S. cerevisiae* NCBI database and analyzed in Scaffold (version

3.6.4). The threshold and false discovery rates (FDR) were set to 80% and 0.37% respectively

for peptides, and to 95% (1 peptide minimum) and 1.9% for respectively for proteins.

Exclusive spectrum counts (ESC) were used for analysis and for each prey, the highest values

obtained in the controls were removed from those of the samples during analysis.

717   *In vitro* methylation

In vitro methylation reactions were done essentially as described (Roguev *et al*, 2001). 30 ml

reactions typically contained 2 to 4 ml of partially purified SET1C complex, 2 to 4 mg

substrate (either peptide, or core histone mixture, or recombinant H3) and 2 ml S-adenosyl

(methyl-^{3}H) methionine in MTA buffer (50 mM Tris 8.5, 20 mM KCl, 10 mM $MgCl_2$, 250

mM sucrose). Reactions with core histones and recombinant H3 were resolved on 4-20%

NuPAGE gels and subjected to fluorography. For Snf2 methylation assays, reaction mixtures

containing purified SET1C (with 30 ng of the Bre2 subunit) and 200 ng of Snf2 fragments in

20 μl of reaction buffer (25 mM HEPES [pH 7.6], 50 mM KCl, 5 mM MgCl2, 0.1 mM

EDTA, and 10% glycerol) were supplemented with 1 μCi of S-adenosyl (methyl-^{3}H)

methionine (PerkinElmer) and incubated at 30°C for 1 hour. Proteins were resolved by SDS-

PAGE and subjected to fluorography. For fluorography, gels were fixed for 30 min in 40%

methanol, 10% acetic acid, treated with EN^{3}HANCE solution (Perkin Elmer) for 60 min,

washed in cold dH_20 for 30 min, dried and exposed for five to fourteen days to photographic

film.

733   Mass spectrometry-based identification of Arg methylation in Snf2-B3

Mutant Snf2-B3 underwent a methylation reaction with SET1C and SAM, followed by

purification via Ni-NTA affinity chromatography and separation by SDS-PAGE. The gel

bands corresponding to the mutant Snf2-B3 were excised and subjected to in-gel digestion

with AspN enzyme, followed by peptide extraction. The resulting peptide fractions were

analyzed using an Easy-nLC 1200 coupled to an Orbitrap Fusion Lumos mass spectrometer

(Thermo Fisher Scientific, MA, USA) at the Korea Basic Science Institute (Ochang). Peptides

were separated on a C18 column using a 150-minute gradient, and data were acquired in data-

dependent acquisition (DDA) mode. Full MS scans were acquired in the Orbitrap at a

resolution of 60,000 over an m/z range of 350–2,000. Precursors for MS/MS analysis were

selected for higher-energy collisional dissociation (HCD) fragmentation at a normalized

collision energy of 30%. Raw MS data were processed using Proteome Discoverer with the

SEQUEST search engine. The data were searched against the sequence of Snf2-B3 with a

precursor mass tolerance of 10 ppm and a fragment mass tolerance of 0.02 Da. AspN was set

as the digestion enzyme, allowing for up to two missed cleavages. Methylation,

dimethylation, and methionine oxidation were specified as variable modifications. The false
discovery rate (FDR) was set to 1% at the peptide level.
750
Y2H interaction of Set1 fragments with selected preys
The Set1 fragments were amplified from the pB66-Set1-FL plasmid and cloned into the SfiI
site of pB66. The selected preys (Snf2, Prp8, and Prp22) into p6 (Hybrigenics) were extracted
from the screen. Plasmid pB66 contains the Gal4 DNA binding domain and the *TRP1* marker
while pP6 expresses the Gal4 activating domain and the *LEU2* marker. TOTO cells were
transformed with the different pB66-Set1-Fragments and the pP6-interactors and incubated 3
days at 30°C on SD-LEU-TRP. The transformant colonies were then streaked on SD-LEU-
TRP-HIS containing either 5 mM or 20 mM of 3AT. To visualize interaction between Set1
fragments and the interactors, yeast cells were incubated 2 days at 30°C and cell growth was
examined.
761
CO-IP of Prp22-FLAG with Myc-Set1.
Co-immunoprecipitation experiment was performed in W303 expressing chromosomally
encoded Myc-Set1 (Dehe *et al*, 2006) and Prp22AID-FLAG (Mendoza-Ochoa *et al*, 2019).
250 ml of culture at an OD600 of 0.8 was harvested by centrifugation at 1000 x *g* and washed
twice in ice-cold 1 X PBS. The cell pellet was re-suspended in 900 µl lysis buffer (50 mM
Tris-HCl pH 7.5, 2 mM Mg2Cl2, 150 mM NaCl, 0.2% NP-40 and one complete EDTA-free
proteinase inhibitor tablet (Roche #11836145001) and 400 µl zirconia beads. Cells were lysed
using a Mini-Beadbeater-24 (BioSpec Products) twice at 2000 rpm for 2 min followed by 2
min on ice. The sample was centrifuged at 1000 x *g* for 2 min, the supernatant was collected
and additionally centrifuged at 20,000 x *g* for 30 min at 4°C and used for
immunoprecipitation. The concentration of protein was measured using the Bradford assay,
and 1 mg of protein used per IP. Prior to immunoprecipitation, extract was pre-cleared by
adding ½ volume unconjugated Protein A/G Dynabeads (Life Technologies
##10001D/10003D). 50 µl Protein A/G Dynabeads conjugated to antibody were incubated
with the appropriate volume of extract on a rotating wheel overnight at 4°C. The next day,
beads were washed 8 times in lysis buffer (non-bound fraction kept for analysis). 20 µl of
loading buffer was added to the beads, input and non-bound samples, which were boiled for
10 min before loading on a NuPAGE 4-12% Bis-Tris gel Bis-Tris (Invitrogen) and western
blotting was performed.
781
Set1 SUMOylation analysis
Cells were transformed with a plasmid encoding 6His-SUMO under the *CUP1* promoter
(YEp352-6His-SUMO) or the corresponding empty vector (Niño *et al*, 2016). The
transformed cells were grown on selective medium and stimulated overnight with 0.1 mM
$CuSO_4$. 200 OD_{600} of cells were collected and lysed with glass beads in a 20% TCA solution,
the final TCA concentration was adjusted to 12%. Cell lysates were incubated at 4°C during
45 min, and precipitated proteins were collected by centrifugation. Proteins were resuspended
in loading buffer (6 M guanidinium-HCl, 100 mM KH_2PO_4, 20 mM Tris-HCl, pH 8,0, 100
mM NaCl, 0,1% Triton X-100, and 10 mM imidazole). 6His-SUMOylated proteins were
purified using Ni-NTA agarose beads (Qiagen), pre-equilibrated with loading buffer, and

incubated for 1 h at room temperature. After incubation the beads were collected by centrifugation (3000 rpm, 2min) and washed twice with wash buffer (8 M urea, 100 mM Na$_2$HPO$_4$/NaH$_2$PO$_4$ pH 6,4, 10 mM Tris–HCl, pH 6.4, 10 mM Imidazole, 10 mM β-mercaptoethanol, and 0.1% Triton X-100). 6His-SUMOylated proteins were eluted with 50 ml 2X laemmli buffer (95°C, 5min). The proteins in the eluted fraction were analyzed by Western blot using anti-MYC (for endogenous Myc-Set1), anti-GAL4 (for GBD-Set1-full length and fragments) and polyclonal rabbit anti-Smt3.

Interaction Snf2-AT-hook with SET1C and SET1C-762

GST-tagged Snf2 fragments (final concentration 250 nM) were combined with either the full-length SET1C or C762SET1C (final concentration 25 nM), along with 15 µl of glutathione-Sepharose 4B resin and BSA (final concentration 0.2 mg/ml) in a binding buffer containing 20 mM Tris-Cl (pH 7.9), 150 mM NaCl, 0.2 mM EDTA, 20% glycerol, 0.1% NP-40, and 1 mM PMSF, making a total volume of 300 µl. The mixtures were rotated at 4°C for 3 hours, followed by four washes with the binding buffer. The proteins bound to the resin were then eluted, separated by SDS-PAGE, and analyzed by immunoblotting.

CO-IP of Snf2-Myc and Snf2ΔRG-Myc with Set1

WT cells or expressing *Snf2-Myc* and *Snf2ΔRG-Myc* (200 ml) were grown at 30°C in YPD to and O.D$_{600}$ = 0.8. Cells were collected by centrifugation, washed once with 10 mM Tris-HCL, pH 8.0 and snap-frozen in liquid nitrogen. Cells were lysed using Retsch with the following parameters :2 times 2 min at 30 m/S. The powder was resuspended in 3 ml of TMG 50 (10 mM Tris-HCL, pH 8.0, 0,1 mM MgCl$_2$, 10% (V/V) glycerol, 50 mM NaCl, 0.1 mM EDTA, 0.1 mM DTT) containing protease inhibitors, 10 mM MG-132 and 1 mM PMSF. Cell lysates were clarified by centrifugation (13 300 rpm, 15', 4°C) and 1.4 ml of supernatant were recovered. Protein concentration was determined using nanodrop and samples were adjusted to the same concentration by adding lysis buffer. Immunoprecipitation was performed overnight with 5 ml of 9E10 antibody (Santa Cruz Biotechnology) following by an incubation for 3 hours with 25 ml of pre-equilibrated protein-G dynabeads (Invitrogen). Immunoprecipitations were washed 3X with TMG 50 buffer and eluted using 25 ml of 1X Laemmli loading Buffer. Samples were resolved on a 7.5% Acrylamide gel, transferred on a nitrocellulose membrane and reveal with either anti-Myc 9E10 (Santa Cruz Biotechnology) and anti-Set1 antibodies.

Snf2-GFP Complex purification

800 ml of *Snf2-GFP* and *Snf2-GFP set1Δ* cells were grown at 30°C in YPD to O.D$_{600}$ = 0.8. Cell pellets were treated as described above except that supernatants were incubated with 25 ml of GFP nanobody coated Dynabeads (Gift from M. Modesti, CRCM Marseille) for 250 min, wash 3X with TMG 50 buffer and eluted in 30 ml of 2X Laemmli Loading buffer containing 50 mM DTT. For Western blotting, 1 ml was resolved on a NuPAGE 4-12% Bis-Tris Gel (Invitrogen), transferred on a nitrocellulose membrane and revealed using an anti-GFP antibody (A-11122-Invitrogen). For the Silver Staining, 2,5 ml were resolved on a NuPAGE 4-12% Bis-Tris Gel (Invitrogen) and stained using Pierce^{TM} Silver Stain kit (Thermo Scientific).

836

837 Mass spectrometry-based identification of post-translational modifications in Snf2-GFP

Purified Snf2-GFP (n=3, biological replicates) from WT and *set1Δ* strains were loaded on
NuPAGE™ 4–12% Bis–tris acrylamide gels according to the manufacturer's instructions
(Life Technologies). Running of protein was stopped as soon as proteins stacked in a single
band. Protein containing bands were stained with Imperial Blue (Pierce), cut from the gel and
digested with high sequencing grade trypsin (Promega) before mass spectrometry analysis.
Briefly, gel pieces were washed and destained using few steps of 100 mM NH_4HCO_3.
Destained gel pieces were shrunk with 100 mM ammonium bicarbonate in 50% acetonitrile
and dried at RT. Protein spots were then rehydrated using 10mM DTT in 25 mM ammonium
bicarbonate pH 8.0 for 45 min at 56°C. This solution was replaced by 55 mM iodoacetamide
in 25 mM ammonium bicarbonate pH 8.0 and the gel pieces were incubated for 30 min at
room temperature in the dark. They were then washed twice in 25 mM ammonium
bicarbonate and finally shrunk by incubation for 5 min with 25 mM ammonium bicarbonate
in 50% acetonitrile. The resulting alkylated gel pieces were dried at room temperature. The
dried gel pieces were reswollen by incubation in 25 mM ammonium bicarbonate pH 8.0
supplemented with 12.5 ng/µl trypsin (Promega) for 1h at 4°C and then incubated overnight at
37°C. Peptides were harvested by collecting the initial digestion solution and carrying out two
extractions; first in 5% formic acid and then in 5% formic acid in 60% acetonitrile. Pooled
extracts were dried down in a centrifugal vacuum system. Samples were reconstituted in 0.1%
TFA 4% acetonitrile before mass spectrometry using an Orbitrap Fusion Lumos Tribrid Mass
Spectrometer (ThermoFisher Scientific, San Jose, CA) online with an Ultimate
3000RSLCnano chromatography system (ThermoFisher Scientific, Sunnyvale, CA). Peptides
were separated at 40°C using a two steps linear gradient (4–20% acetonitrile/H2O; 0.1%
formic acid for 110 min and 20-32% acetonitrile/H2O; 0.1% formic acid for 10 min). An
EASY-Spray nanosource was used for peptide ionization (2,200 V, 275°C). MS was
conducted using a data-dependent acquisition mode (DDA). The Orbitrap Lumos was used in
data dependent mode to switch consistently between MS and MS/MS. Time between Masters
Scans was set to 3 seconds. MS spectra were acquired with the Orbitrap in the range of m/z
400-1600 at a FWHM resolution of 120 000 measured at 400 m/z. AGC target was set at
4.0e5 with a 50 ms Maximum Injection Time. For internal mass calibration the 445.120025
ion was used as lock mass. The more abundant precursor ions were selected and collision-
induced dissociation fragmentation was performed in the ion trap to have maximum
sensitivity and yield a maximum amount of MS/MS data. Number of precursor ions was
automatically defined along run in 3s windows using the "Inject Ions for All Available
parallelizable time option" with a maximum injection time of 300 ms. The signal threshold for
an MS/MS event was set to 5000 counts. Charge state screening was enabled to exclude
precursors with 0 and 1 charge states. Dynamic exclusion was enabled with a repeat count of
1 and duration of 60 s.
For identification of PTMs the region corresponding to Snf2-GFP was excised from the gel
and subjected to enzymatic digestion under the same conditions as above. Raw mass
spectrometry files were analysed using MaxQuant software (version 1.6.3.4) as above with
the exception of the Max Missed cleavage set at 5, the FDR at peptide and proteins levels at
5% and with the following variable modifications activated; Lysine acetylation (+42.0106);

Serine, threonine and tyrosine phosphorylation (+79.966); Lysine and arginine methylation (+14.0157), dimethylation (+28.0313) and trimethylation (+42.0470).

Data Processing Protocol

Relative intensity-based label-free quantification (LFQ) was processed using the MaxLFQ algorithm from the freely available MaxQuant computational proteomics platform, version 1.6.2.1 (Cox et al. 2014, Cox et al. 2008). The acquired raw LC Orbitrap MS data were first processed using the integrated Andromeda search engine (Cox et 2011). Spectra were searched against the Saccharomyces cerevisiae (organism ID 4932) extracted from UniProt containing 7904 entries. The following parameters were used for searches: (i) trypsin allowing cleavage before proline; (ii) two missed cleavage was allowed; (iii) cysteine carbamidomethylation (+57.02146) as a fixed modification and methionine oxidation (+15.99491) and N-terminal acetylation (+42.0106) as variable modifications. The match between runs option was enabled. The false discovery rate (FDR) at the peptide and protein levels were set to 1% and determined by searching a reverse database. For protein grouping, all proteins that cannot be distinguished based on their identified peptides were assembled into a single entry according to the MaxQuant rules. The statistical analysis was done with Perseus program (version 1.6.1.3) (Tyanova and Cox 2018) from the MaxQuant environment (www.maxquant.org). Quantifiable proteins were defined as those detected in above 70% of samples in one condition or more. Protein LFQ normalized intensities were base 2 logarithmized to obtain a normal distribution. Missing values were replaced using data imputation by randomly selecting from a normal distribution centered on the lower edge of the intensity values that simulates signals of low abundant proteins using default parameters (a downshift of 1.8 standard deviation and a width of 0.3 of the original distribution). The protein composition of the complex was determined using, a two-sample t-test using permutation-based FDR-controlled at 0.01 and employing 250 permutations and a scaling factor s0 with a value of 3. All proteins passing these criteria with a positive fold change are considered to be the potential interactome of Snf2.

Antibodies

For IP: Mouse anti-FLAG (Sigma M2 #F1804); 10 μl/IP with Dynabeads and Protein G.

For westerns - primary antibodies: Mouse anti-c-MYC (Santa Cruz #SC-40x) 1:1000; Rat anti-FLAG (Agilent #200474) 1:1000; Mouse anti-Gal4 Binding Domain (Euromedex) 1:5000; Rabbit anti-Smt3 is a gift from B. Palancade (Institut Jacques Monod, Paris). Anti-Bre2 is a gift from Peter Nagy. The rabbit anti-Spp1 antibody was developed in V. Géli's lab. The anti-Spp1 antibody is showing significant cross-reactivity with other cellular proteins but did allow the identification of Spp1 in purified and partially purified fractions. Secondary antibodies: Goat anti-mouse IRDye680RD (LI-COR #926-68070) 1:10,000, Goat anti-rat IRDye800RD (LI-COR #926-32219) 1:10,000.

DATA AVAILABILITY

Additional data and reagents are available upon request to the corresponding authors. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium

924 (www.proteomexchange.org) via the PRIDE partner repository
925 (https://www.ebi.ac.uk/pride/login): Snf2-B3 data (accession number PXD061448), Snf2-
926 GFP complex (accession number PXD061496), Snf2 PTM (accession number PXD061531).

928 .

ACKNOWLEDGEMENTS
930
931 We are grateful to Sandra Holbein and Andre Halbach for help with MBP-Nrm1 fusion
932 proteins. We thank Benoit Palancade for his help in the interpretation of the Y2H results and
933 reagents.

FUNDING
937
This work is supported by the "Ligue Nationale Contre le Cancer" (LNCC) (Equipe
labellisée, GELI/2019). Funding to B.D. was provided by grants from the Swiss National
Science Foundation (PP00A--102941/1 and 31003A_1327010). I.E.M. was funded by
Wellcome Trust PhD Studentship (105256) and work in the Wellcome Centre for Cell
Biology was supported by Wellcome core funding (092076). Funding to M. O. was provided
by the Natural Sciences and Engineering Research Council of Canada to M.O. (RGPIN-2020-
06924). Funding to J. K was provided by the National Research Foundation of Korea (NRF-
2022R1A2C3012803). Y. H. K is supported by the National Research Foundation of Korea
(NRF) grant funded by the Korea government (MSIT) (No. RS-2024-00454407). Proteomics
analyses using the mass spectrometry facility of Marseille Proteomics (marseille-
proteomique.univ-amu.fr) are supported by IBISA, the Cancéropôle PACA, the Provence-
Alpes-Côte d'Azur Region, the Institut Paoli-Calmettes, and Fonds Européens de
Développement Regional (FEDER).

CONFLICT OF INTEREST
953
954 The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS
956
B.D. J. K and V.G. designed the study and secured the funding. All the Y2H screens and prey
identification were performed by Hybrigenics in close interaction with B.D. and V. G. P.L.
performed molecular cloning, contributed to validation experiments, and purified Snf2-GFP
for subsequent analyses. SET1C and SET1C-C762 purification and interaction assays with
Snf2-AT-hook were performed by J. K. and K.P. Set1 SUMOylation experiments were
performed by C. N. M. D and B. D. organized the results shown in Table S2. L. L
constructed the F1-F5 fragments and contributed to validation experiments. Pull-down
experiments and Nrm1 methylation assay were performed by B. D. TAP-Nis1 data were
provided by M.O. I.M. and J. B. performed and discuss the Set1/Prp22 interaction. D.K. P
and H. H. K performed mass spectrometry analysis of the Snf2-B3. S.A. and L.C. performed

mass spectrometry analysis of the Snf2-GFP. V.G. and B. D. wrote the manuscript with inputs
of the authors.
969
970
971 **REFERENCES**
972

Acquaviva L, Drogat J, Dehé P-M, de La Roche Saint-André C & Géli V (2013a) Spp1 at the
crossroads of H3K4me3 regulation and meiotic recombination. *Epigenetics* 8:
355–360

Acquaviva L, Szekvolgyi L, Dichtl B, Dichtl BS, Saint Andre C d LR, Nicolas A & Geli V
(2013b) The COMPASS Subunit Spp1 Links Histone Methylation to Initiation of
Meiotic Recombination. *Science* 339: 215–218

Adam C, Guérois R, Citarella A, Verardi L, Adolphe F, Béneut C, Sommermeyer V, Ramus
C, Govin J, Couté Y, *et al* (2018) The PHD finger protein Spp1 has distinct
functions in the Set1 and the meiotic DSB formation complexes. *PLoS Genet* 14:
e1007223

An D, Kim J, Moon B, Kim H, Nguyen H, Park S, Lee JE, Kim J-A & Kim J (2025) PRMT1-
mediated methylation regulates MLL2 stability and gene expression. *Nucleic*
Acids Res 53

An S, Yoon J, Kim H, Song J-J & Cho U-S (2017) Structure-based nuclear import
mechanism of histones H3 and H4 mediated by Kap123. *eLife* 6: e30244

Arigo JT, Carroll KL, Ames JM & Corden JL (2006) Regulation of yeast NRD1 expression
by premature transcription termination. *Mol Cell* 21: 641–651

Bae HJ, Dubarry M, Jeon J, Soares LM, Dargemont C, Kim J, Geli V & Buratowski S (2020)
The Set1 N-terminal domain and Swd2 interact with RNA polymerase II CTD to
recruit COMPASS. *Nat Commun* 11: 2181

Baker RP, Harreman MT, Eccleston JF, Corbett AH & Stewart M (2001) Interaction
between Ran and Mog1 is required for efficient nuclear protein import. *J Biol*
Chem 276: 41255–41262

Balk J, Pierik AJ, Netz DJA, Mühlenhoff U & Lill R (2004) The hydrogenase-like Nar1p is
essential for maturation of cytosolic and nuclear iron-sulphur proteins. *EMBO J*
23: 2105–2115

Battaglia S, Lidschreiber M, Baejen C, Torkler P, Vos SM & Cramer P (2017) RNA-
dependent chromatin association of transcription elongation factors and Pol II
CTD kinases. *eLife* 6: e25637

Beck M & Hurt E (2017) The nuclear pore complex: understanding its function through
structural insight. *Nat Rev Mol Cell Biol* 18: 73–89

Beilharz TH, Harrison PF, Miles DM, See MM, Le UMM, Kalanon M, Curtis MJ, Hasan Q,
Saksouk J, Margaritis T, *et al* (2017) Coordination of Cell Cycle Progression and

Mitotic Spindle Assembly Involves Histone H3 Lysine 4 Methylation by
Set1/COMPASS. *Genetics* 205: 185–199

Berretta J, Pinskaya M & Morillon A (2008) A cryptic unstable transcript mediates
transcriptional trans-silencing of the Ty1 retrotransposon in S. cerevisiae. *Genes*
Amp Dev 22: 615–626

Borde V, Robine N, Lin W, Bonfils S, Géli V & Nicolas A (2008) Histone H3 lysine 4
trimethylation marks meiotic recombination initiation sites. *EMBO J* 28: 99–111

Briggs SD, Bryk M, Strahl BD, Cheung WL, Davie JK, Dent SY, Winston F & Allis CD (2001)
Histone H3 lysine 4 methylation is mediated by Set1 and required for cell growth
and rDNA silencing in Saccharomyces cerevisiae. *Genes Amp Dev* 15: 3286–3295

de Bruin RAM, Kalashnikova TI, Chahwan C, McDonald WH, Wohlschlegel J, Yates J,
Russell P & Wittenberg C (2006) Constraining G1-specific transcription to late G1
phase: the MBF-associated corepressor Nrm1 acts via negative feedback. *Mol Cell*
23: 483–496

Bryk M, Briggs SD, Strahl BD, Curcio MJ, Allis CD & Winston F (2002) Evidence that Set1,
a factor required for methylation of histone H3, regulates rDNA silencing in S.
cerevisiae by a Sir2-independent mechanism. *Curr Biol CB* 12: 165–170

Castelnuovo M, Rahman S, Guffanti E, Infantino V, Stutz F & Zenklusen D (2013) Bimodal
expression of PHO84 is modulated by early termination of antisense
transcription. *Nat Struct Mol Biol* 20: 851–858

Castelnuovo M, Zaugg JB, Guffanti E, Maffioletti A, Camblong J, Xu Z, Clauder-Münster S,
Steinmetz LM, Luscombe NM & Stutz F (2014) Role of histone modifications and
early termination in pervasive transcription and antisense-mediated gene
silencing in yeast. *Nucleic Acids Res* 42: 4348–4362

Cenik BK & Shilatifard A (2021) COMPASS and SWI/SNF complexes in development and
disease. *Nat Rev Genet* 22: 38–58

Cheng H, He X & Moore C (2004) The Essential WD Repeat Protein Swd2 Has Dual
Functions in RNA Polymerase II Transcription Termination and Lysine 4
Methylation of Histone H3. *Mol Cell Biol* 24: 2932–2943

Chern M-K, Chang K-N, Liu L-F, Tam T-CS, Liu Y-C, Liang Y-L & Tam MF (2002) Yeast
Ribosomal Protein L12 Is a Substrate of Protein-arginine Methyltransferase 2. *J*
Biol Chem 277: 15345–15353

Choi J, Rajagopal A, Xu Y-F, Rabinowitz JD & O'Shea EK (2017) A systematic genetic
screen for genes involved in sensing inorganic phosphate availability in
Saccharomyces cerevisiae. *PloS One* 12: e0176085

Chong SY, Cutler S, Lin J-J, Tsai C-H, Tsai H-K, Biggins S, Tsukiyama T, Lo Y-C & Kao C-F
(2020) H3K4 methylation at active genes mitigates transcription-replication
conflicts during replication stress. *Nat Commun* 11: 809

Choudhury R, Singh S, Arumugam S, Roguev A & Stewart AF (2019) The Set1 complex is dimeric and acts with Jhd2 demethylation to convey symmetrical H3K4 trimethylation. *Genes Dev* 33: 550–564

Corda Y, Schramke V, Longhese MP, Smokvina T, Paciotti V, Brevet V, Gilson E & Geli V (1999) Interaction between Set1p and checkpoint protein Mec3p in DNA repair and telomere functions. *Nat Genet* 21: 204–208

Côté J, Quinn J, Workman JL & Peterson CL (1994) Stimulation of GAL4 derivative binding to nucleosomal DNA by the yeast SWI/SNF complex. *Science* 265: 53–60

Daicho K, Koike N, Ott RG, Daum G & Ushimaru T (2020) TORC1 ensures membrane trafficking of Tat2 tryptophan permease via a novel transcriptional activator Vhr2 in budding yeast. *Cell Signal* 68: 109542

Dehe PM, Dichtl B, Schaft D, Roguev A, Pamblanco M, Lebrun R, Rodriguez-Gil A, Mkandawire M, Landsberg K, Shevchenko A, *et al* (2006) Protein Interactions within the Set1 Complex and Their Roles in the Regulation of Histone 3 Lysine 4 Methylation. *J Biol Chem* 281: 35404–35412

Dehé P-M & Géli V (2006) The multiple faces of Set1. *Biochem Cell Biol Biochim Biol Cell* 84: 536–548

Delamarre A, Barthe A, de la Roche Saint-André C, Luciano P, Forey R, Padioleau I, Skrzypczak M, Ginalski K, Géli V, Pasero P, *et al* (2020) MRX Increases Chromatin Accessibility at Stalled Replication Forks to Promote Nascent DNA Resection and Cohesin Loading. *Mol Cell* 77: 395-410.e3

Deshpande N & Bryk M (2023) Diverse and dynamic forms of gene regulation by the S. cerevisiae histone methyltransferase Set1. *Curr Genet* 69: 91–114

Deshpande N, Jordan R, Henderson Pozzi M & Bryk M (2022) Histone 3 lysine 4 monomethylation supports activation of transcription in S. cerevisiae during nutrient stress. *Curr Genet* 68: 181–194

Dichtl B (2004) Functions for S. cerevisiae Swd2p in 3' end formation of specific mRNAs and snoRNAs and global histone 3 lysine 4 methylation. *RNA* 10: 965–977

Dichtl B, Blank D, Sadowski M, Hübner W, Weiser S & Keller W (2002) Yhh1p/Cft1p directly links poly(A) site recognition and RNA polymerase II transcription termination. *EMBO J* 21: 4125–4135

van Dijk EL, Chen CL, d'Aubenton-Carafa Y, Gourvennec S, Kwapisz M, Roche V, Bertrand C, Silvain M, Legoix-Né P, Loeillet S, *et al* (2011) XUTs are a class of Xrn1-sensitive antisense regulatory non-coding RNA in yeast. *Nature* 475: 114–117

Dover J, Schneider J, Tawiah-Boateng MA, Wood A, Dean K, Johnston M & Shilatifard A (2002) Methylation of histone H3 by COMPASS requires ubiquitination of histone H2B by Rad6. *J Biol Chem* 277: 28368–28371

D'Urso A, Takahashi Y-H, Xiong B, Marone J, Coukos R, Randise-Hinchliff C, Wang J-P,
Shilatifard A & Brickner JH (2016) Set1/COMPASS and Mediator are repurposed
to promote epigenetic transcriptional memory. *eLife* 5: 111

Faucher D & Wellinger RJ (2010) Methylated H3K4, a Transcription-Associated Histone
Modification, Is Involved in the DNA Damage Response Pathway. *PLoS Genet* 6:
e1001082

Formstecher E, Aresta S, Collura V, Hamburger A, Meil A, Trehin A, Reverdy C, Betin V,
Maire S, Brun C, *et al* (2005) Protein interaction mapping: a Drosophila case
study. *Genome Res* 15: 376–384

Fromont-Racine M, Rain JC & Legrain P (1997) Toward a functional analysis of the yeast
genome through exhaustive two-hybrid screens. *Nat Genet* 16: 277–282

Ghaddar N, Corda Y, Luciano P, Galli M, Doksani Y & Géli V (2023) The COMPASS subunit
Spp1 protects nascent DNA at the Tus/Ter replication fork barrier by limiting
DNA availability to nucleases. *Nat Commun* 14: 5430

Giaever G, Lissina E & Nislow C (2019) Network dynamics of the yeast
methyltransferome. *Microb Cell Graz Austria* 6: 356–369

Gong X, Wang S, Yu Q, Wang M, Ge F, Li S & Yu X (2023) Cla4 phosphorylates histone
methyltransferase Set1 to prevent its degradation by the APC/CCdh1 complex. *Sci
Adv* 9: eadi7238

Grainger RJ & Beggs JD (2005) Prp8 protein: at the heart of the spliceosome. *RNA N Y N*
11: 533–557

Green DM, Marfatia KA, Crafton EB, Zhang X, Cheng X & Corbett AH (2002) Nab2p is
required for poly(A) RNA export in Saccharomyces cerevisiae and is regulated by
arginine methylation via Hmt1p. *J Biol Chem* 277: 7752–7760

Guccione E, Bassi C, Casadio F, Martinato F, Cesaroni M, Schuchlautz H, Lüscher B &
Amati B (2007) Methylation of histone H3R2 by PRMT6 and H3K4 by an MLL
complex are mutually exclusive. *Nature* 449: 933–937

Guillemette B, Drogaris P, Lin H-HS, Armstrong H, Hiragami-Hamada K, Imhof A, Bonneil
É, Thibault P, Verreault A & Festenstein RJ (2011) H3 Lysine 4 Is Acetylated at
Active Gene Promoters and Is Regulated by H3 Lysine 4 Methylation. *PLOS Genet*
7: 1–16

Guy MP, Podyma BM, Preston MA, Shaheen HH, Krivos KL, Limbach PA, Hopper AK &
Phizicky EM (2012) Yeast Trm7 interacts with distinct proteins for critical
modifications of the tRNAPhe anticodon loop. *RNA N Y N* 18: 1921–1933

Halbach A, Zhang H, Wengi A, Jablonska Z, Gruber IML, Halbeisen RE, Dehé P-M,
Kemmeren P, Holstege F, Géli V, *et al* (2009) Cotranslational assembly of the
yeast SET1C histone methyltransferase complex. *EMBO J* 28: 2959–2970

Hamey JJ, Rakow S, Bouchard C, Senst JM, Kolb P, Bauer U, Wilkins MR & Hart-Smith G
(2021) Systematic investigation of PRMT6 substrate recognition reveals broad
specificity with a preference for an RG motif or basic and bulky residues. *FEBS J*
288: 5668–5691

Hannich JT, Lewis A, Kroetz MB, Li S-J, Heide H, Emili A & Hochstrasser M (2005)
Defining the SUMO-modified proteome by multiple approaches in Saccharomyces
cerevisiae. *J Biol Chem* 280: 4102–4110

He C, Liu N, Xie D, Liu Y, Xiao Y & Li F (2019) Structural basis for histone H3K4me3
recognition by the N-terminal domain of the PHD finger protein Spp1. *Biochem J*
476: 1957–1973

Henke RM, Dastidar RG, Shah A, Cadinu D, Yao X, Hooda J & Zhang L (2011) Hypoxia
elicits broad and systematic changes in protein subcellular localization. *Am J*
Physiol Cell Physiol 301: C913-928

Hirschhorn JN, Brown SA, Clark CD & Winston F (1992) Evidence that SNF2/SWI2 and
SNF5 activate transcription in yeast by altering chromatin structure. *Genes Dev* 6:
2288–2298

Howe FS, Boubriak I, Sale MJ, Nair A, Clynes D, Grijzenhout A, Murray SC, Woloszczuk R
& Mellor J (2014) Lysine acetylation controls local protein conformation by
influencing proline isomerization. *Mol Cell* 55: 733–744

Hsu PL, Li H, Lau H-T, Leonen C, Dhall A, Ong S-E, Chatterjee C & Zheng N (2018) Crystal
Structure of the COMPASS H3K4 Methyltransferase Catalytic Module. *Cell* 174:
1106-1116.e9

Hsu PL, Shi H, Leonen C, Kang J, Chatterjee C & Zheng N (2019) Structural Basis of H2B
Ubiquitination-Dependent H3K4 Methylation by COMPASS. *Mol Cell* 76: 712-
723.e4

Huang F, Chandrasekharan MB, Chen Y-C, Bhaskara S, Hiebert SW & Sun Z-W (2010) The
JmjN domain of Jhd2 is important for its protein stability, and the plant
homeodomain (PHD) finger mediates its chromatin association independent of
H3K4 methylation. *J Biol Chem* 285: 24548–24561

Hyllus D, Stein C, Schnabel K, Schiltz E, Imhof A, Dou Y, Hsieh J & Bauer U-M (2007)
PRMT6-mediated methylation of R2 in histone H3 antagonizes H3 K4
trimethylation. *Genes Dev* 21: 3369–3380

Iwase M & Toh-e A (2001) Nis1 encoded by YNL078W: a new neck protein of
Saccharomyces cerevisiae. *Genes Genet Syst* 76: 335–343

Jackson CA, Yadav N, Min S, Li J, Milliman EJ, Qu J, Chen YC & Yu MC (2012) Proteomic
analysis of interactors for yeast protein arginine methyltransferase Hmt1 reveals
novel substrate and insights into additional biological roles. *PROTEOMICS* 12:
3304–3314

Janke C, Magiera MM, Rathfelder N, Taxis C, Reber S, Maekawa H, Moreno-Borchart A,
Doenges G, Schwob E, Schiebel E, *et al* (2004) A versatile toolbox for PCR-based
tagging of yeast genes: new fluorescent proteins, more markers and promoter
substitution cassettes. *Yeast* 21: 947–962

Jeon J, McGinty RK, Muir TW, Kim J-A & Kim J (2018) Crosstalk among Set1 complex
subunits involved in H2B ubiquitylation-dependent H3K4 methylation. *Nucleic
Acids Res* 46: 11129–11143

Jezek M, Sun W, Negesse MY, Smith ZM, Orosz A & Green EM (2023) Set1 regulates
telomere function via H3K4 methylation-dependent and -independent pathways
and calibrates the abundance of telomere maintenance factors. *Mol Biol Cell* 34:
ar6

Kaczmarek Michaels K, Mohd Mostafa S, Ruiz Capella J & Moore CL (2020) Regulation of
alternative polyadenylation in the yeast Saccharomyces cerevisiae by histone
H3K4 and H3K36 methyltransferases. *Nucleic Acids Res* 48: 5407–5425

Kan J, Zou L, Zhang J, Wu R, Wang Z & Liang C (2008) Origin Recognition Complex (ORC)
Mediates Histone 3 Lysine 4 Methylation through Cooperation with Spp1 in
Saccharomyces cerevisiae. *J Biol Chem* 283: 33803–33807

Karányi Z, Halász L, Acquaviva L, Jónás D, Hetey S, Boros-Oláh B, Peng F, Chen D, Klein F,
Géli V, *et al* (2018) Nuclear dynamics of the Set1C subunit Spp1 prepares meiotic
recombination sites for break formation. *J Cell Biol* 217: 3398–3415

Kim J, Kim J-A, McGinty RK, Nguyen UTT, Muir TW, Allis CD & Roeder RG (2013) The n-
SET Domain of Set1 Regulates H2B Ubiquitylation-Dependent H3K4 Methylation.
Mol Cell 49: 1121–1133

Kim J & Roeder RG (2011) Nucleosomal H2B ubiquitylation with purified factors.
Methods San Diego Calif 54: 331–338

Kim J-H, Saraf A, Florens L, Washburn M & Workman JL (2010) Gcn5 regulates the
dissociation of SWI/SNF from chromatin by acetylation of Swi2/Snf2. *Genes Dev*
24: 2766–2771

Kim M, Vasiljeva L, Rando OJ, Zhelkovsky A, Moore C & Buratowski S (2006) Distinct
pathways for snoRNA and mRNA termination. *Mol Cell* 24: 723–734

Kim T & Buratowski S (2009) Dimethylation of H3K4 by Set1 recruits the Set3 histone
deacetylase complex to 5' transcribed regions. *Cell* 137: 259–272

Kirmizis A, Santos-Rosa H, Penkett CJ, Singer MA, Green RD & Kouzarides T (2009)
Distinct transcriptional outputs associated with mono- and dimethylated histone
H3 arginine 2. *Nat Struct Mol Biol* 16: 449–451

Kirmizis A, Santos-Rosa H, Penkett CJ, Singer MA, Vermeulen M, Mann M, Bähler J, Green
RD & Kouzarides T (2007) Arginine methylation at histone H3R2 controls
deposition of H3K4 trimethylation. *Nature* 449: 928–932

de La Roche Saint-André C & Géli V (2021) Set1-dependent H3K4 methylation becomes
critical for limiting DNA damage in response to changes in S-phase dynamics in
Saccharomyces cerevisiae. *DNA Repair* 105: 103159

Laribee RN, Shibata Y, Mersman DP, Collins SR, Kemmeren P, Roguev A, Weissman JS,
Briggs SD, Krogan NJ & Strahl BD (2007) CCR4/NOT complex associates with the
proteasome and regulates histone methylation. *Proc Natl Acad Sci U S A* 104:
5836–5841

Lee DC & Aitchison JD (1999) Kap104p-mediated nuclear import. Nuclear localization
signals in mRNA-binding proteins and the role of Ran and Rna. *J Biol Chem* 274:
29031–29037

Lee K & Wang K (2018) Small Noncoding RNAs in Agrobacterium tumefaciens. *Curr Top*
Microbiol Immunol 418: 195–213

Lee KY, Chopra A, Burke GL, Chen Z, Greenblatt JF, Biggar KK & Meneghini MD (2020) A
crucial RNA-binding lysine residue in the Nab3 RRM domain undergoes SET1 and
SET3-responsive methylation. 1–15

Lee Y-S, Mulugu S, York JD & O'Shea EK (2007) Regulation of a cyclin-CDK-CDK inhibitor
complex by inositol pyrophosphates. *Science* 316: 109–112

Lees ND, Skaggs B, Kirsch DR & Bard M (1995) Cloning of the late genes in the ergosterol
biosynthetic pathway of Saccharomyces cerevisiae--a review. *Lipids* 30: 221–226

Li H-T, Gong T, Zhou Z, Liu Y-T, Cao X, He Y, Chen CD & Zhou J-Q (2015a) Yeast Hmt1
catalyses asymmetric dimethylation of histone H3 arginine 2 *in vitro*. *Biochem J*
467: 507–515

Li N, Zhai Y, Zhang Y, Li W, Yang M, Lei J, Tye B-K & Gao N (2015b) Structure of the
eukaryotic MCM complex at 3.8 Å. *Nature* 524: 186–191

Liang G, Klose RJ, Gardner KE & Zhang Y (2007) Yeast Jhd2p is a histone H3 Lys4
trimethyl demethylase. *Nat Struct Mol Biol* 14: 243–245

Liu HY, Chiang YC, Pan J, Chen J, Salvadore C, Audino DC, Badarinarayana V,
Palaniswamy V, Anderson B & Denis CL (2001) Characterization of CAF4 and
CAF16 reveals a functional connection between the CCR4-NOT complex and a
subset of SRB proteins of the RNA polymerase II holoenzyme. *J Biol Chem* 276:
7541–7548

Liu J, Liu J & Stråby KB (1998) Point and deletion mutations eliminate one or both
methyl group transfers catalysed by the yeast TRM1 encoded tRNA
(m22G26)dimethyltransferase. *Nucleic Acids Res* 26: 5102–5108

Low JKK & Wilkins MR (2012) Protein arginine methylation in *Saccharomyces cerevisiae*.
FEBS J 279: 4423–4443

Luciano P, Jeon J, El-Kaoutari A, Challal D, Bonnet A, Barucco M, Candelli T, Jourquin F,
Lesage P, Kim J, *et al* (2017) Binding to RNA regulates Set1 function. *Cell Discov* 3:
17040

Margaritis T, Oreal V, Brabers N, Maestroni L, Vitaliano-Prunier A, Benschop JJ, van Hooff
S, van Leenen D, Dargemont C, Géli V, *et al* (2012) Two distinct repressive
mechanisms for histone 3 lysine 4 methylation through promoting 3'-end
antisense transcription. *PLoS Genet* 8: e1002952

McBride AE, Cook JT, Stemmler EA, Rutledge KL, McGrath KA & Rubens JA (2005)
Arginine methylation of yeast mRNA-binding protein Npl3 directly affects its
function, nuclear export, and intranuclear protein interactions. *J Biol Chem* 280:
30888–30898

Mei Q, Xu C, Gogol M, Tang J, Chen W, Yu X, Workman JL & Li S (2019) Set1-catalyzed
H3K4 trimethylation antagonizes the HIR/Asf1/Rtt106 repressor complex to
promote histone gene expression and chronological life span. *Nucleic Acids Res*
47: 3434–3449

Meitinger F, Khmelinskii A, Morlot S, Kurtulmus B, Palani S, Andres-Pons A, Hub B, Knop
M, Charvin G & Pereira G (2014) A memory system of negative polarity cues
prevents replicative aging. *Cell* 159: 1056–1069

Mendoza-Ochoa GI, Barrass JD, Maudlin IE & Beggs JD (2019) Blocking late stages of
splicing quickly limits pre-spliceosome assembly in vivo. *RNA Biol* 16: 1775–1784

Mersman DP, Du H-N, Fingerman IM, South PF & Briggs SD (2009) Polyubiquitination of
the demethylase Jhd2 controls histone methylation and gene expression. *Genes*
Dev 23: 951–962

Miller T, Krogan NJ, Dover J, Erdjument-Bromage H, Tempst P, Johnston M, Greenblatt JF
& Shilatifard A (2001) COMPASS: a complex of proteins associated with a
trithorax-related SET domain protein. *Proc Natl Acad Sci U S A* 98: 12902–12907

Mizuno T, Nakamura M & Irie K (2018) Induction of Ptp2 and Cmp2 protein
phosphatases is crucial for the adaptive response to ER stress in Saccharomyces
cerevisiae. *Sci Rep* 8: 13078

Mulder KW, Brenkman AB, Inagaki A, van den Broek NJF & Timmers HTM (2007)
Regulation of histone H3K4 tri-methylation and PAF complex recruitment by the
Ccr4-Not complex. *Nucleic Acids Res* 35: 2428–2439

Mullen JR, Chen C-F & Brill SJ (2010) Wss1 is a SUMO-dependent isopeptidase that
interacts genetically with the Slx5-Slx8 SUMO-targeted ubiquitin ligase. *Mol Cell*
Biol 30: 3737–3748

Mulugu S, Bai W, Fridy PC, Bastidas RJ, Otto JC, Dollins DE, Haystead TA, Ribeiro AA &
York JD (2007) A conserved family of enzymes that phosphorylate inositol
hexakisphosphate. *Science* 316: 106–109

Murray SC, Haenni S, Howe FS, Fischl H, Chocian K, Nair A & Mellor J (2015) Sense and
antisense transcription are associated with distinct chromatin architectures
across genes. *Nucleic Acids Res* 43: 7823–7837

Nadal-Ribelles M, Mas G, Millán-Zambrano G, Solé C, Ammerer G, Chávez S, Posas F & de
Nadal E (2015) H3K4 monomethylation dictates nucleosome dynamics and
chromatin remodeling at stress-responsive genes. *Nucleic Acids Res* 43: 4937–
4949

Nagy PL, Griesenbeck J, Kornberg RD & Cleary ML (2002) A trithorax-group complex
purified from Saccharomyces cerevisiae is required for methylation of histone
H3. *Proc Natl Acad Sci U S A* 99: 90–94

Nakagawa T & Kolodner RD (2002) The MER3 DNA helicase catalyzes the unwinding of
holliday junctions. *J Biol Chem* 277: 28019–28024

Nayak A, Viale-Bouroncle S, Morsczeck C & Muller S (2014) The SUMO-specific
isopeptidase SENP3 regulates MLL1/MLL2 methyltransferase complexes and
controls osteogenic differentiation. *Mol Cell* 55: 47–58

Nedea E, He X, Kim M, Pootoolal J, Zhong G, Canadien V, Hughes T, Buratowski S, Moore
CL & Greenblatt J (2003) Organization and Function of APT, a Subcomplex of the
Yeast Cleavage and Polyadenylation Factor Involved in the Formation of mRNA
and Small Nucleolar RNA 3'-Ends. *J Biol Chem* 278: 33000–33010

Nedea E, Nalbant D, Xia D, Theoharis NT, Suter B, Richardson CJ, Tatchell K, Kislinger T,
Greenblatt JF & Nagy PL (2008) The Glc7 Phosphatase Subunit of the Cleavage
and Polyadenylation Factor Is Essential for Transcription Termination on
snoRNA Genes. *Mol Cell* 29: 577–587

Ng HH, Robert F, Young RA & Struhl K (2003) Targeted recruitment of Set1 histone
methylase by elongating Pol II provides a localized mark and memory of recent
transcriptional activity. *Mol Cell* 11: 709–719

Nie M, Vashisht AA, Wohlschlegel JA & Boddy MN (2015) High Confidence Fission Yeast
SUMO Conjugates Identified by Tandem Denaturing Affinity Purification. *Sci Rep*
5: 14389

Niño CA, Guet D, Gay A, Brutus S, Jourquin F, Mendiratta S, Salamero J, Géli V &
Dargemont C (2016) Posttranslational marks control architectural and functional
plasticity of the nuclear pore complex basket. *J Cell Biol* 212: 167–180

Nislow C, Ray E & Pillus L (1997) SET1, a yeast member of the trithorax family, functions
in transcriptional silencing and diverse cellular processes. *Mol Biol Cell* 8: 2421–
2436

Noma A, Kirino Y, Ikeuchi Y & Suzuki T (2006) Biosynthesis of wybutosine, a hyper-
modified nucleoside in eukaryotic phenylalanine tRNA. *EMBO J* 25: 2142–2154

Oki M & Nishimoto T (1998) A protein required for nuclear-protein import, Mog1p,
directly interacts with GTP-Gsp1p, the Saccharomyces cerevisiae ran homologue.
Proc Natl Acad Sci U S A 95: 15388–15393

Oliete-Calvo P, Serrano-Quílez J, Nuño-Cabanes C, Pérez-Martínez ME, Soares LM, Dichtl
B, Buratowski S, Pérez-Ortín JE & Rodríguez-Navarro S (2018) A role for Mog1 in
H2Bub1 and H3K4me3 regulation affecting RNAPII transcription and mRNA
export. *EMBO Rep* 19: e45992

Pal S, Vishwanath SN, Erdjument-Bromage H, Tempst P & Sif S (2004) Human SWI/SNF-
Associated PRMT5 Methylates Histone H3 Arginine 8 and Negatively Regulates
Expression of *ST7* and *NM23* Tumor Suppressor Genes. *Mol Cell Biol* 24: 9630–
9645

Pemberton LF, Rosenblum JS & Blobel G (1997) A distinct and parallel pathway for the
nuclear import of an mRNA-binding protein. *J Cell Biol* 139: 1645–1653

Perez AM & Thorner J (2019) Septin-associated proteins Aim44 and Nis1 traffic between
the bud neck and the nucleus in the yeast Saccharomyces cerevisiae. *Cytoskelet
Hoboken NJ* 76: 15–32

Poornima G, Srivastava G, Roy B, Kuttanda IA, Kurbah I & Rajyaguru PI (2021) RGG-
motif containing mRNA export factor Gbp2 acts as a translation repressor. *RNA
Biol* 18: 2342–2353

Qu Q, Takahashi Y-H, Yang Y, Hu H, Zhang Y, Brunzelle JS, Couture J-F, Shilatifard A &
Skiniotis G (2018) Structure and Conformational Dynamics of a COMPASS
Histone H3K4 Methyltransferase Complex. *Cell* 174: 1117-1126.e12

Radman-Livaja M, Liu CL, Friedman N, Schreiber SL & Rando OJ (2010) Replication and
Active Demethylation Represent Partially Overlapping Mechanisms for Erasure of
H3K4me3 in Budding Yeast. *PLoS Genet* 6: e1000837

Rain JC, Selig L, De Reuse H, Battaglia V, Reverdy C, Simon S, Lenzen G, Petel F, Wojcik J,
Schächter V, *et al* (2001) The protein-protein interaction map of Helicobacter
pylori. *Nature* 409: 211–215

Rizzardi LF, Dorn ES, Strahl BD & Cook JG (2012) DNA replication origin function is
promoted by H3K4 di-methylation in Saccharomyces cerevisiae. *Genetics* 192:
371–384

Roguev A, Schaft D, Shevchenko A, Pijnappel WW, Wilm M, Aasland R & Stewart AF
(2001) The Saccharomyces cerevisiae Set1 complex includes an Ash2 homologue
and methylates histone 3 lysine 4. *EMBO J* 20: 7137–7148

Ruthenburg AJ, Allis CD & Wysocka J (2007) Methylation of lysine 4 on histone H3:
intricacy of writing and reading a single epigenetic mark. *Mol Cell* 25: 15–30

Santos-Rosa H, Bannister AJ, Dehe PM, Géli V & Kouzarides T (2004) Methylation of H3
lysine 4 at euchromatin promotes Sir3p association with heterochromatin. *J Biol
Chem* 279: 47506–47512

Santos-Rosa H, Millán-Zambrano G, Han N, Leonardi T, Klimontova M, Nasiscionyte S, Pandolfini L, Tzelepis K, Bartke T & Kouzarides T (2021) Methylation of histone H3 at lysine 37 by Set1 and Set2 prevents spurious DNA replication. *Mol Cell* 81: 2793-2807.e8

Sayou C, Millán-Zambrano G, Santos-Rosa H, Petfalski E, Robson S, Houseley J, Kouzarides T & Tollervey D (2017) RNA Binding by Histone Methyltransferases Set1 and Set2. *Mol Cell Biol* 37: e00165-17

Schlichter A & Cairns BR (2005) Histone trimethylation by Set1 is coordinated by the RRM, autoinhibitory, and catalytic domains. *EMBO J* 24: 1222–1231

Schulze JM, Jackson J, Nakanishi S, Gardner JM, Hentrich T, Haug J, Johnston M, Jaspersen SL, Kobor MS & Shilatifard A (2009) Linking cell cycle to histone modifications: SBF and H2B monoubiquitination machinery and cell-cycle regulation of H3K79 dimethylation. *Mol Cell* 35: 626–641

Senger B, Simos G, Bischoff FR, Podtelejnikov A, Mann M & Hurt E (1998) Mtr10p functions as a nuclear import receptor for the mRNA-binding protein Npl3p. *EMBO J* 17: 2196–2207

Serra-Cardona A, Duan S, Yu C & Zhang Z (2022) H3K4me3 recognition by the COMPASS complex facilitates the restoration of this histone mark following DNA replication. *Sci Adv* 8: eabm6246

Shin JA, Choi ES, Kim HS, Ho JCY, Watts FZ, Park SD & Jang YK (2005) SUMO modification is involved in the maintenance of heterochromatin stability in fission yeast. *Mol Cell* 19: 817–828

Smith D-L, Erce MA, Lai Y-W, Tomasetig F, Hart-Smith G, Hamey JJ & Wilkins MR (2020) Crosstalk of Phosphorylation and Arginine Methylation in Disordered SRGG Repeats of Saccharomycescerevisiae Fibrillarin and Its Association with Nucleolar Localization. *J Mol Biol* 432: 448–466

Soares LM & Buratowski S (2012) Yeast Swd2 Is Essential Because of Antagonism between Set1 Histone Methyltransferase Complex and APT (Associated with Pta1) Termination Factor. *J Biol Chem* 287: 15219–15231

Soares LM, He PC, Chun Y, Suh H, Kim T & Buratowski S (2017) Determinants of Histone H3K4 Methylation Patterns. *Mol Cell* 68: 773-785.e6

Sollier J, Lin W, Soustelle C, Suhre K, Nicolas A, Géli V & de La Roche Saint-André C (2004) Set1 is required for meiotic S-phase onset, double-strand break formation and middle gene expression. *EMBO J* 23: 1957–1967

Sommermeyer V, Béneut C, Chaplais E, Serrentino ME & Borde V (2013) Spp1, a member of the Set1 Complex, promotes meiotic DSB formation in promoters by tethering histone H3K4 methylation sites to chromosome axes. *Mol Cell* 49: 43–54

Soniat M, Sampathkumar P, Collett G, Gizzi AS, Banu RN, Bhosle RC, Chamala S, Chowdhury S, Fiser A, Glenn AS, *et al* (2013) Crystal structure of human

Karyopherin β2 bound to the PY-NLS of Saccharomyces cerevisiae Nab2. *J Struct*
Funct Genomics 14: 31–35

South PF, Harmeyer KM, Serratore ND & Briggs SD (2013) H3K4 methyltransferase Set1
is involved in maintenance of ergosterol homeostasis and resistance to Brefeldin
A. *Proc Natl Acad Sci U S A* 110: E1016-25

Süel KE, Gu H & Chook YM (2008) Modular organization and combinatorial energetics of
proline-tyrosine nuclear localization signals. *PLoS Biol* 6: e137

Sun Z-W & Allis CD (2002) Ubiquitination of histone H2B regulates H3 methylation and
gene silencing in yeast. *Nature* 418: 104–108

Takeo K & Ito T (2017) Subcellular localization of VIP1 is regulated by phosphorylation
and 14-3-3 proteins. *FEBS Lett* 591: 1972–1981

Terzi N, Churchman LS, Vasiljeva L, Weissman J & Buratowski S (2011) H3K4
Trimethylation by Set1 Promotes Efficient Termination by the Nrd1-Nab3-Sen1
Pathway. *Mol Cell Biol* 31: 3569–3583

Thandapani P, O'Connor TR, Bailey TL & Richard S (2013) Defining the RGG/RG motif.
Mol Cell 50: 613–623

Thornton JL, Westfield GH, Takahashi Y-H, Cook M, Gao X, Woodfin AR, Lee J-S, Morgan
MA, Jackson J, Smith ER, *et al* (2014) Context dependency of Set1/COMPASS-
mediated histone H3 Lys4 trimethylation. *Genes Dev* 28: 115–120

Trahan C & Oeffinger M (2022) Single-Step Affinity Purification (ssAP) and Mass
Spectrometry of Macromolecular Complexes in the Yeast S. cerevisiae. *Methods*
Mol Biol Clifton NJ 2477: 195–223

Travesa A, Kuo D, de Bruin RAM, Kalashnikova TI, Guaderrama M, Thai K, Aslanian A,
Smolka MB, Yates JR, Ideker T, *et al* (2012) DNA replication stress differentially
regulates G1/S genes via Rad53-dependent inactivation of Nrm1. *EMBO J* 31:
1811–1822

Trésaugues L, Dehé P-M, Guérois R, Rodriguez-Gil A, Varlet I, Salah P, Pamblanco M,
LUCIANO pierre, Quevillon-Cheruel S, Sollier J, *et al* (2006) Structural
Characterization of Set1 RNA Recognition Motifs and their Role in Histone H3
Lysine 4 Methylation. *J Mol Biol* 359: 1170–1181

Vitaliano-Prunier A, Menant A, Hobeika M, Géli V, Gwizdek C & Dargemont C (2008)
Ubiquitylation of the COMPASS component Swd2 links H2B ubiquitylation to
H3K4 trimethylation. *Nat Cell Biol* 10: 1365–1371

Walter D, Matter A & Fahrenkrog B (2014) Loss of histone H3 methylation at lysine 4
triggers apoptosis in Saccharomyces cerevisiae. *PLoS Genet* 10: e1004095

Wang L, Collings CK, Zhao Z, Cozzolino KA, Ma Q, Liang K, Marshall SA, Sze CC,
Hashizume R, Savas JN, *et al* (2017) A cytoplasmic COMPASS is necessary for cell

1420  survival and triple-negative breast cancer pathogenesis by regulating
1421  metabolism. *Genes Dev* 31: 2056–2066

1422 Wang Y, Ding Z, Liu X, Bao Y, Huang M, Wong CCL, Hong X & Cong Y (2018) Architecture
1423  and subunit arrangement of the complete Saccharomyces cerevisiae COMPASS
1424  complex. *Sci Rep* 8: 17405

1425 Warnecke DC & Heinz E (1994) Purification of a Membrane-Bound UDP-Glucose:Sterol
1426  [beta]-D-Glucosyltransferase Based on Its Solubility in Diethyl Ether. *Plant*
1427  *Physiol* 105: 1067–1073

1428 Weiner A, Chen HV, Liu CL, Rahat A, Klien A, Soares L, Gudipati M, Pfeffner J, Regev A,
1429  Buratowski S, *et al* (2012) Systematic Dissection of Roles for Chromatin
1430  Regulators in a Yeast Stress Response. *PLoS Biol* 10: e1001369

1431 Will CL & Lührmann R (2011) Spliceosome structure and function. *Cold Spring Harb*
1432  *Perspect Biol* 3: a003707

1433 Wilson TE, Padgett KA, Johnston M & Milbrandt J (1993) A genetic method for defining
1434  DNA-binding domains: application to the nuclear receptor NGFI-B. *Proc Natl Acad*
1435  *Sci U S A* 90: 9186–9190

1436 Worley J, Luo X & Capaldi AP (2013) Inositol pyrophosphates regulate cell growth and
1437  the environmental stress response by activating the HDAC Rpd3L. *Cell Rep* 3:
1438  1476–1482

1439 Wykoff DD & O'Shea EK (2005) Identification of sumoylated proteins by systematic
1440  immunoprecipitation of the budding yeast proteome. *Mol Cell Proteomics MCP* 4:
1441  73–83

1442 Xu D, Farmer A & Chook YM (2010) Recognition of nuclear targeting signals by
1443  Karyopherin-β proteins. *Curr Opin Struct Biol* 20: 782–790

1444 Yoshida K & Blobel G (2001) The karyopherin Kap142p/Msn5p mediates nuclear import
1445  and nuclear export of different cargo proteins. *J Cell Biol* 152: 729–740

1446 Yoshimura SH & Hirano T (2016) HEAT repeats - versatile arrays of amphiphilic helices
1447  working in crowded environments? *J Cell Sci* 129: 3963–3970

1448 Yuan C-C, Matthews AGW, Jin Y, Chen CF, Chapman BA, Ohsumi TK, Glass KC, Kutateladze
1449  TG, Borowsky ML, Struhl K, *et al* (2012) Histone H3R2 Symmetric Dimethylation
1450  and Histone H3K4 Trimethylation Are Tightly Correlated in Eukaryotic Genomes.
1451  *Cell Rep* 1: 83–90

1452 Zhang Z, Shibahara K & Stillman B (2000) PCNA connects DNA replication to epigenetic
1453  inheritance in yeast. *Nature* 408: 221–225

1454 **FIGURES AND LEGENDS**

1455

1456 **FIGURE 1**

1457
1458
1459
Figure 1. Schematic representation of the Set1 and subunit major interactors identified
in the systematic yeast two-hybrid screens.
A) Schematic representation of Set1 FL and Set1 fragments 1-754 and 754-1081. **B**) Set1 FL,
1-754 and 754-1081 major Y2H interactors). **C**) Interacting regions of Set1 with Swd2, Shg1
and Spp1. **D**) Swd2, Spp1, Shg1 major Y2H interactors. **E**) Swd1, Swd3, Bre2, Sdc1 major
Y2H interactors. The term "interactor" is used to mean a high confidence two-hybrid
interaction, with the limitations that this entails. The color reflects the Predicted Biological
Score (see METHODS). Red, highest confidence; Blue, high confidence; Green, good
confidence.
1469
1470
1471
1472

1473 **FIGURE 2**

1474
1475

1476
1477

Figure 2. Set1 1-754 interacts with RGG proteins and the importin Kap104.

A) RGG proteins and import/export proteins interacting with Set1 1-754, Set1 754-1081 and Spp1, Shg1, and Sdc1. **B**) Set1 interacting domain (SID) (blue) within Kap104. Heat like repeat 9 and 10 are represented in purple **C**) PY-NLS in the N-terminal region of Set1. **D**) SID (blue) and RGG motif (green) within the RGG proteins. The interaction domains indicated represent the minimal overlapping DNA sequence present in multiple independent Y2H interacting clones of the same gene. Each genomic fragment of a Y2H clone was analyzed, and the shared overlapping region for given gene was determined to be the only common element among all interacting clones. As such, this region represents the minimal sequence required for interaction.

1488   **FIGURE 3**

1489
1490

1491
1492
1493   **Figure 3. Chromatin regulators identified in all the two-hybrid screens.** The lines indicate
1494   the individual proteins involved in the Y2H interaction. Red line refers to a very high
1495   confidence Y2H interaction. All Y2H interactors are described in Table S2. Interactors are
1496   grouped according to the complex to which they belong.

 FIGURE 4

1503
1504

1505
1506
1507

Figure 4. SET1C interacts *in vitro* with Snf2C-AT-hook.
A) A schematic diagram depicting the domains of Snf2 and the Snf2 fragments used in this study, along with the SDS-PAGE/Coomassie staining of the purified GST-tagged Snf2 fragments. **B** and **C**) GST pull-down assay using purified GST-tagged Snf2 fragments. The purified SET1C (**B**) or SET1C-C762 complex (**C**) was mixed with GST-tagged Snf2 fragments, followed by GST pull-down, and the bound proteins were analyzed by immunoblotting. **D**) A schematic diagram illustrating Snf2 fragments with a more detailed breakdown of the AT-hook domain, along with the SDS-PAGE/Coomassie Blue staining of the purified Snf2 fragments. The lysines present in the AT-hook are represented by the letter K. **E**) GST pull-down assay using purified GST-tagged Snf2 fragments and SET1C.

1518
1519
1520
1521

FIGURE 5

1524
1525
1526
1527 **Figure 5. The SET1C Y2H interactome identifies proteins involved in RNA biogenesis.**
1528 All Y2H interactors are described in Table S2. The processes linked to RNA metabolism in
1529 which the different interactors are involved are shown in the figure. The green lines linking
1530 Prp22 to Set1FL/Set1 754-1080 and Prp8 to Set1 FL and Spp1 indicate interactions with a
1531 high degree of confidence.

 FIGURE 6

Figure 6. Set1 is SUMOylated.
6His-SUMO–conjugated proteins were purified from cells transformed (+) or not transformed
(−) with a plasmid encoding 6His-SUMO under control of the *CUP1* promoter. Cell lysates
(Input) and Ni-purified material (Elutes) were analyzed by Western blotting with an anti-
MYC antibody (**A**) or and anti-GAL4 antibody (**B-D**). Analysis of 6His-SUMO -conjugated
forms of (**A**) genomically MYC-tagged Set1 or (**B**) GB-Set1 transformed cells was performed
(left panels), in both the cases SUMO expression and efficiency of purification were
controlled using an anti-SUMO antibody (right panels). **C**) SUMOylation analysis of Set1
fragment F3+F4 (aa. 351-956) WT and the K769R mutant. **D**) SUMOylation analysis of Set1
fragment F5 (aa. 956-1080) WT, single mutants K1055R and K1060R, and the double mutant
K1055R/K1060R mutant.

1562 **FIGURE 7**

1563
1564
1565
Figure 7. Nrm1 is methylated *in vitro* by SET1C.
A) Schematic representation of the *S. cerevisiae* Nrm1 protein. Nrm1 carries a D-box sequence at its
amino-terminus and a central domain that mediated interaction with the Swd1 subunit of SET1C in a
yeast two-hybrid screen. A H3K4 like domain that closely resembles the modification site of SET1C
in histone H3 is contained within the Swd1 interaction domain. Identical and similar amino-acid
positions are shown in italic. Lysine 118 (K118) of Nrm1 aligns with lysine 4 of histone H3; also
indicated are K116R and K118A mutant sequences. **B**) *In vitro* methylation reactions using partially
purified SET1C, [3]H-SAM and the indicated peptides. Radioactive reaction products retained on
Whatman P81 filter paper following extensive washes were measured by scintillation counting. Each
reaction was done in triplicate and error bars indicate the standard deviation. **C**) *In vitro* methylation
reactions as in (B), however, core histone (2, 4 and 6 mg) and recombinant MBP and MBP-Nrm1
fusion protein (0.5, 1 and 2 mg) were tested as substrates for methylation. Reaction products were
separated on 4 to 12% NuPAGE gels and analyzed by fluorography. **D**) *In vitro* methylation reactions
as in (C), however, core histone (2 mg) and recombinant MBP-Nrm1 wild-type and K118A and
K116R fusion protein (0.5, 1 and 2 mg) were tested as substrates for methylation. (upper panel). Note
that the molecular mass marker proteins (MM) have been resolved together with the core histone
methylation reaction in the same lane. The asterisk indicates migration of a non-specific background
signal of unclear origin. The substrate proteins included in the reactions were analyzed on a parallel
gel and stained with colloidal Coomassie G-250 (lower panel). **E**) H3K4-like proteins. Shown are
selected proteins containing sequences similar to the modification site found in histone H3. An
exhaustive list can be found in Table S4.
1587

1588
1589
1590
1591

    FIGURE 8

1593
1594

1595
1596
Figure 8. Snf2 is methylated in tandem AT-hook domain by reconstituted Set1C.

A) *In vitro* methyltransferase assay using purified SET1C and Snf2 fragments. ³H-SAM was used as a methyl-donor and methylated proteins were detected by autoradiography. The band marked with a red star is a degradation product of Snf2C. **B)** *In vitro* methyltransferase assay using SET1C and two Snf2-AT-hook fragments with two different tags. **C)** Schematic diagram of N-terminal truncated SET1 complexes (left) and *in vitro* methyltransferase assay with GST-Snf2-AT-hook and truncated SET1 complexes. **D)** Schematic diagram showing the positions of all lysines in Snf2-AT-hook and the further cleaved fragments of Snf2-AT-hook. The red box indicates the two lysines that are acetylated by Gcn5. **E)** Coomassie staining of purified Snf2 fragments (lower) and an *in vitro* methyltransferase assay using these fragments with SET1C (upper). **F)** *In vitro* methyltransferase assay by SET1C when each of the four lysines in the C-terminal region of the Snf2-AT-hook is substituted with arginine or when both lysines known to be acetylated by Gcn5 are substituted. **G)** A schematic diagram showing the position of the RG-repeat region and the design of Snf2-AT-hook with RG-repeat truncation. H and I) GST pull-down assay (**H**) and *in vitro* methyltransferase assay (**I**) using purified SET1C and GST-Snf2-AT-hook with or without RG-repeats.

1613
1614

1615 **FIGURE 9**

1616
1617
Figure 9. The arginines in the RG-repeat of Snf2 are methylated by reconstituted
SET1C.
A) A diagram showing the WT Snf2-B3 fragment and the Mut Snf2-B3 with all four lysines
substituted with alanine. B) Coomassie staining of purified WT and Mut Snf2-B3. C) Mass-
spectrometry experiment design to identify the methylation sites of Snf2-B3. D) Coomassie
staining of Mut Snf2-B3 after methylation reaction and an additional purification step using
Ni-NTA. The band corresponding to Mut Snf2-B3 (~18 KDa) was excised and used for mass
spectrometry analysis. The ~37 KDa band observed in lanes 5–8 appears to be a SET1C
subunit that binds non-specifically to Ni-NTA, likely SWD2 based on its size. E) Mass
spectrometry analysis result of Snf2-B3 methylation sites revealed that multiple arginines in
the RG-repeat were methylated. Amino acid sequence of Snf2 highlighting the arginines
methylated by SET1. The sequence of the B3 fragment is shown in bold, and the arginines
methylated by SET1 are marked in red. Methylation and demethylation are denoted as M and
DM, respectively.
1632

1633   **FIGURE 10**

1634

1635
1636

Figure 10. The arginines of the motif ARTSTRGR within the AT hook are methylated *in*
***vivo* in a Set1-dependent way.**
A) Set1 interacts *in vivo* with Snf2 and Snf2-ΔRG. Myc-tagged Snf2 and Snf2ΔRG were
immunoprecipitated with 9E10 Myc antibodies (see Methods) and revealed with either
9E10 (Upper panel) or Set1 antibodies (lower panel). **B**) Snf2C complex was purified
from WT and *set1Δ* strains, separated on a 4-12% Bis-Tris Gel and Silver Stained (*Left*);
the presence of Snf2-GFP is detected by Western-blotting with anti-GFP antibody
(*Right*). The area corresponding to Snf2-GFP was excised from the gel and used for mass
spectrometry analysis. Peptides flanking the RG repeats (1485-1549) with their PTM are
shown in Fig. S14. **C**) Panel C show a focus on the amino sequence flanking the RG
repeats. The positions of residues from D_{1485} to V_{1549} are indicated on the figure. Peptides
identified after digestion of Snf2-GFP are indicated in color with their identified PTM
indicated by the color code shown in the top of the panel. The small numbers above the
amino acids indicate the probability of localisation according to the MS2 peaks. It should
be noted that for wild-type K1488 and R1490, a peptide with dimethylation is detected,
but no discriminating MS2 peak allows us to conclude whether K or R are dimethylated.
This is also the case for dimethylation on R1528 and R1535. The results presented
represent the observed PTMs from two independent experiments, each containing 3
replicates (Table S6).