Absence of Local Retinotopy in the Mouse Optic Tract

Authors: Matteo Tripodi[1], Hiroki Asari*[1,2]

Affiliations:

1. Epigenetics and Neurobiology Unit, EMBL Rome, European Molecular Biology Laboratory, Monterotondo, 00015, Italy

2. Current address: Scuola Internazionale Superiore di Studi Avanzati (SISSA), Trieste, 34136, Italy

*Corresponding author: Hiroki Asari, E-mail: asari@embl.it

Author contributions: M.T. collected the experimental data; H.A. designed the study, analyzed the data, and wrote and revised the manuscript.

Competing interests: The authors declare no competing financial interests.

Key words: retinotopy; optic tract; retinal ganglion cells; lateral geniculate nucleus; mouse

Acronyms: AP, anterior-posterior; DV, dorsal-ventral; IR, infrared; LED, light emitting diode; LGN, lateral geniculate nucleus; ML, medial-lateral; OT, optic tract; PFA, paraformaldehyde; RF, receptive field; RGC, retinal ganglion cell; STA, spike-triggered average; UV, ultraviolet.

Acknowledgements: This work was supported by research grants from EMBL (H.A.). The EMBL Light Imaging Facility is acknowledged for support in histological image acquisition; EMBL IT Support for provision of computer and data storage servers; and the LAR facility for taking care of animals. We thank all the Asari lab members for useful discussions.

Abstract

Retinotopy is a fundamental organizational principle of the visual system, where neighboring neurons represent adjacent points in visual space. This spatial relationship is established by precise anatomical wiring across successive areas, e.g., from the retina to the lateral geniculate nucleus (LGN) to the visual cortex. To examine the precision of this topographic arrangement within the long-range projection axons themselves, we recorded retinal ganglion cell (RGC) axons in the mouse optic tract (OT) and mapped their receptive fields (RFs). As expected for a retinotopically organized area, we found that nearby LGN cell pairs had significantly smaller RF distances than distant pairs. In contrast, no such relationship was observed among RGC axons in the OT. Modelling analyses further confirmed that the observed RF distances in the OT were incompatible with any locally retinotopic arrangement. Instead, the OT retained only coarse topography, with ~18° RF deviations or ~40 µm axonal displacements from an ideal retinotopic organization. These results demonstrate that the mouse OT lacks fine-scale retinotopy and maintains only broad topographic structure.

Introduction

The vertebrate visual system maintains spatial relationships from the retina to the downstream visual areas via topographic projections (Huberman et al., 2008; Seabrook et al., 2017; Cang et al., 2018). This retinotopic organization emerges early in development (Debski and Cline, 2002; McLaughlin and O'Leary, 2005; Cang and Feldheim, 2013), and provides a fundamental platform for visual processing, ensuring that nearby points in visual space are represented by neighboring neuronal populations. In mice, retinal ganglion cells (RGCs) project to >40 brain areas (Morin and Studholme 2014; Martersteck et al, 2017), with nearly all axons passing through the same pathway – the optic nerve, optic chiasm, and optic tract (OT) – before diverging to targets, such as the superior colliculus and the lateral geniculate nucleus (LGN). While retinotopy in these downstream areas has been well characterized (McLaughlin, et al., 2003; Piscopo et al., 2013; Molotkov et al., 2023), it remains elusive to what extent retinotopy is maintained in the long-range RGC axons themselves.

Previous anatomical and tracing studies have demonstrated that retinotopy is largely

preserved in the optic nerve (Bunt and Horder, 1983; Reese, 2011; but see Horton et al. 1979).

Axons arising from specific retinal quadrants occupy consistent positions within the optic nerve,

forming a reliable mapping of the retinal geometry as they exit the eye. This topographic

organization is, however, substantially disrupted in the optic chiasm (Marcus and Mason, 1995;

Colerro and Guillery, 1998), where axons from the two eyes intersect and distribute themselves

into either ipsilateral or contralateral side. The optic chiasm has thus been considered as a sorting

hub, where retinotopy is relaxed to permit reorganization based on molecular cues, midline

crossing decision, and eventual target-specific routing (Jeffery, 2001). After passing through the

optic chiasm, RGC axons reorganize and partially recover retinotopic organization in the OT via

pre-target sorting (Simon and O'Leary, 1991; Plas et al., 2005; Sitko et al., 2018). This indicates

the presence of coarse retinotopy in the OT before reaching the target; however, questions remain

on the precision and functional consequences of this reorganization.

Using in vivo electrophysiological recordings, here we mapped visual receptive fields

(RFs) of individual RGC axons in the mouse OT. We quantified the spatial organization of these

RFs, and took a data-driven modelling approach to evaluate how faithfully visual space is

represented in the OT. Our results revealed a moderate degree of retinotopy, but not a fine-

grained one. This highlights a robustness of the visual system to imprecision in long-range wiring,

with refinement of the topographic organization taking place in each target area.

Results

In a brain region with retinotopy, by definition, neighboring cells have adjacent receptive fields

(RFs), representing nearby points in visual space (Cang and Feldheim, 2013; Seabrook et al.,

2017). To examine if the optic tract (OT) has a retinotopic organization, we thus performed in vivo

extracellular recordings of retinal ganglion cell (RGC) axons in the mouse OT, and mapped their

RFs using white-noise stimuli and reverse correlation (e.g., **Fig. 1A-E**). As a positive control,

recordings were also made from the dorsal lateral geniculate nucleus (LGN; e.g., **Fig. 1F-J**), a

major retinorecipient area well-known to have a retinotopy (Piscopo, et al., 2013).

Figure 1: Receptive fields (RFs) of retinal ganglion cell (RGC) axons simultaneously recorded in the mouse optic tract (OT) are more widely spread across the visual field than those of lateral geniculate nucleus (LGN) cells. A: Schematic of OT recordings in awake head-fixed mice and a representative histological image showing the electrode position (DiI, red). B–D: Spatial RF (top) and temporal kernel (bottom) of three representative single-units that had the largest signals on the same recording site from a representative OT recording (left, average spike waveform in each recording site). Orange ellipses show a two-dimensional Gaussian envelope (at 1.5σ) fitted to the spatial RF profile of each unit. E: RF center locations of all simultaneously recorded single-units from the representative OT recording (crosses; N=20). Those with the largest signals on the same recording site are shown with the Gaussian envelopes of the same color. F–J: Corresponding data for a representative LGN recording (F, schematic and histological image; G–I, spatial RF, temporal kernel, and spike waveforms of three representative units that had the largest signals on the same recording site; J, RF center locations of all simultaneously recorded single-units, N=40).

Using multichannel silicon probes (4 shanks, each with 8-16 recording sites; **Fig. 1**), we

simultaneously recorded 6±4 single-units from the OT (mean ± standard deviation, 33 animals)

and 17±16 single-units from the LGN (17 animals). We found that the measured RFs of those

single-units from the OT recordings were widely distributed across the visual stimulation area (73°

and 44° in azimuth and elevation, respectively; RF size range, 1.3°–13.1°; e.g., **Fig. 1E**). This was

also the case even for those units that had the largest signals on the same recording site, hence

were supposedly located near each other in close proximity to the recording site (e.g., **Fig. 1B-D**).

In contrast, the measured RFs of LGN cells were spatially more clustered (e.g., **Fig. 1J**) and

demonstrated substantial overlap, especially among neighboring cells (e.g., **Fig. 1G-I**) as

expected from known retinotopy (Huberman et al., 2008; Seabrook et al., 2017). Such qualitative

comparison already suggested that the mouse OT lacks retinotopy.

For quantitative population-level data analysis, we examined how the RF distance relates

to the physical distance between simultaneously recorded units (**Fig. 2**). We first used the

electrode distance as a proxy of the distance between the units, and found a significant positive

correlation for the LGN (Pearson correlation coefficient R=0.42, p<0.001, N=4491 pairs; **Fig. 2C**),

but not for the OT (R=-0.01, p=0.8, N=756 pairs; **Fig. 2A**). The RF size differences between the

pairs were small (LGN, 1.4°±0.9°; OT, 1.1°±0.6°; median ± median absolute deviation), indicating

that consistent populations were sampled in each group. We next made a comparison between

nearby and distant cell pairs, where nearby cell pairs were defined as those single-units that had

the largest signals on the same recording site (i.e., zero electrode distance), whereas distant pairs

as those with the largest signals on different recordings sites (i.e., non-zero electrode distance).

In the LGN, the measured RF distance was significantly shorter for nearby cell pairs (4.8°±2.3°,

median ± median absolute deviation; N=150 from 11 mice) than for distant cell pairs (14.0°±5.8°;

N=4341 from 17 mice; p(U-test)<0.001; **Fig. 2D**). In the OT, however, RF distances did not differ

between nearby (21.3°±15.1°, N=39 from 18 mice) and distant pairs (29.0°±12.7°, N=717 from 33

mice, **Fig. 2B**; p(U-test)=0.10). Thus, neighboring RGC axons in the mouse OT do not necessarily

represent adjacent visual locations.

97
Figure 2: The mouse optic tract (OT) lacks local topographic organization, while
maintaining global retinotopy. A,B: The RF center distance between RGC axons in the mouse
OT as a function of the distance between the recording sites (A), and the probability distribution
of the RF center distance (B: light red, nearby pairs with the largest signals on the same
recording site, N=39 from 18 animals; dark red, distant pairs with the largest signals on different
recording sites, N=717 from 33 animals; gray, expected distribution for nearby pairs by Monte
Carlo simulation): circle, median; bar, the first and third quartiles. C,D: Corresponding data for
LGN cells (light blue, N=150 nearby pairs from 11 animals; dark blue, N=4341 distant pairs from
17 animals; gray, Monte Carlo simulation). E,F: Simulated RF distance between nearby RGC
axons at different retinotopy levels. Additive Gaussian noise (at different jitter levels, σ; E) was
introduced to RF locations to achieve different spatial monotonicity levels (Kendall's τ) of their
topographic organization (F): black line, median; gray shade, 25 and 75 percentile range; black
dotted line, expected σ and τ values given the measured data (red). G,H: Simulated topographic
map (color-coded) of the mouse OT (G, precise retinotopy; H, moderate retinotopy
corresponding to experimental data).

To further evaluate this, we modelled the expected RF distance for nearby cell pairs under

the assumption of retinotopy (see Methods for details). For LGN, Monte Carlo simulations yielded

an estimated RF distance between 4.2° and 4.9° (95% confidence interval; median, 4.6°; **Fig. 2D**),

given the retinotopy gradient of 0.11°/μm (Piscopo et al., 2013) and a reliable single-unit recording

range of 40 μm from an extracellular electrode (Anastassiou et al., 2015). This is consistent with

the measured RF distance (median, 4.8°), validating our modelling approach. Applying the same

logic to the mouse OT (diameter, ~300 μm; visual field extending ~135° in both azimuth and

elevation; Paxinos and Franklin, 2001), our model predicted the median RF distance of 7.9°–10.7°

for nearby pairs (median, 9.3°; **Fig. 2B**) if RGC axons were arranged retinotopically. This

disagreed with the measured RF distance (median, 21.3°), hence rejecting the presence of

retinotopy imposed as a key premise of the model.

To what extent does the mouse OT retain topographic organization? To address this

question, we extended our model by introducing additive Gaussian noise to the RF location of the

simulated RGC axons (see Methods for details), where σ=1 corresponds to the represented visual

field (135°). We then used Kendall's τ as a measure of spatial monotonicity to quantify the degree

of retinotopy, where τ=1 and 0 represent precise and no retinotopy, respectively. As expected, the

retinotopy level τ decreased with increasing noise level σ, and the estimated RF distance of nearby

RGC axons expanded (**Fig. 2E,F**). From the intersection between the measured and simulated

RF distances, we then identified that the mouse OT had a moderate level of retinotopy (τ=0.49)

with σ=0.13, corresponding to a deviation of RF location from a retinotopically ideal position by

~18°, or equivalently, a displacement of RGC axons by ~40 μm (**Fig. 2G,H**). This indicates that

the mouse OT lacks local retinotopy, while maintaining topographic organization at a global level.

Discussion

Here we provide in vivo electrophysiological evidence for the absence of local retinotopy in the

mouse optic tract (OT). Both population-level analyses and data-driven models failed to identify a

fine-scale topographic organization in the OT that would make neighboring retinal ganglion cell

(RGC) axons encode adjacent locations in the visual field. In contrast, corresponding analyses on

the dorsal lateral geniculate nucleus (LGN) supported the presence of retinotopy as expected

from previous studies (Huberman et al., 2008; Seabrook et al., 2017), thereby serving as a control

for our functional circuit characterization.

We acknowledge several caveats in our results. First, the OT is predominantly comprised

of RGC axons, but also contains axons from other sources, such as the parabigeminal nucleus

(Reinhard et al., 2019; Tokuoka et al., 2020) and the superior colliculus (Gale and Murphy, 2014).

Our physiological criteria (i.e., robust visual responses with short latencies and well-defined,

spatially-confined linear RFs) may not be sufficient to fully exclude those nontargets. Second,

while the OT is largely intermingled to form a broad retinotopy, there is partial and graded

segregation of RGC axons by the cell type and projection target (Erskine and Herrera, 2014;

Robles et al., 2014). Our models did not incorporate such typology, and thus may underestimate

the actual degree of retinotopy in OT. Another caveat is that the nodes of Ranvier from adjacent

RGC axons can be offset by up to ~70 μm (Butt et al., 1994); hence, they may not be detected on

the same recording site. Nevertheless, the units we defined as "nearby" should still be in close

proximity to a given recording site, and thus near each other if not the closest. This guarantees

the validity and robustness of our data/model analyses.

Our finding that RGC axons in the mouse OT retain only coarse retinotopy implies a

fundamental constraint of early visual pathway organization: i.e., pre-target axon sorting cannot

establish precise spatial relationships over long distances (Plas et al., 2005). This is consistent

with known limits of axon guidance mechanisms, such as Eph/ephrin signaling and axon

fasciculation, that establish broad topographic gradients but lack the resolution required for single-

axon precision (McLaughlin and O'Leary, 2005; Cang and Feldheim, 2013). Thus, fine retinotopy

must be re-established within each target via activity-dependent mechanisms—including retinal

waves and Hebbian plasticity—to impose local precision on coarse inputs (Debski and Cline,

2002; McLaughlin et al., 2003).

Such a two-step architecture likely reflects an adaptive circuit organization principle: by

relaxing long-range wiring constraints, the system reduces developmental cost while increasing

robustness. Small positional deviations in the OT have then minimal impact because final

mappings are refined locally in individual targets. Moreover, each area remains free to optimize

the retinotopic map for its own computational needs (Knapen, 2021), rather than relying on the

geometric arrangement of incoming axons themselves. Testing these functional implications in

the visual system—and examining if they extend across the central nervous system—will be an

important direction for future studies.

Materials and Methods

174 No statistical method was used to predetermine the sample size. The significance level was 0.05

175 in all analyses unless otherwise noted. All experiments were performed under the license

176 233/2017-PR and 220/2024-PR from the Italian Ministry of Health, following protocols approved

177 by the Institutional Animal Care and Use Committee at European Molecular Biology Laboratory.

178 The data analyses were done in Python and Matlab.

Animals

180 Animals were housed on a 12h light-dark cycle, with ad libitum access to water and food. In total,

181 50 female wild-type mice (C57BL6/J; RRID:IMSR_JAX:000664), 4-19 weeks old (median, 9.2

182 weeks old) at the time of surgery, were used for in vivo electrophysiology (optic tract, 33 animals;

183 lateral geniculate nucleus, 17 animals).

In vivo electrophysiology

In vivo electrophysiology was performed as described previously (Tripodi and Asari, 2025). Briefly,

we first implanted a head-plate to animals for fixing their head during in vivo electrophysiological

recordings. Before the surgery, animals were injected with Carprofen (5 mg/kg) and then

anaesthetized with isoflurane (4% for induction, 1% for maintenance in O_2). During the surgery,

the animals were placed in a stereotaxic frame (Stoelting 51625) with a heating pad (Supertech

Physiological) to keep their temperature stable at 37°C; and ointment (VitA-Pos, Ursapharm) was

applied on both eyes to prevent them from drying. A portion of the scalp was removed to expose

the skull, and the periosteum was scraped away with a round scalpel to increase adherence of

the dental cement. A titanium head-plate with a hole (diameter, 8mm) was then cemented on the

skull with a mixture of cyanoacrylate (Loctite 401, Henkel) and dental cement (Paladur, Kulzer).

The skull surface was then glazed with a thick layer of cyanoacrylate to support the skull with

mechanical, atmospheric and biological protection, while still allowing for visual identification of

reference points (bregma and lambda). After the surgery, the animals were placed on a heating

pad for recovery, and then housed in individual cages. During the following seven days, the mice

were administered with analgesia (Carprofen; 50 mg/mL) diluted in drinking water. The animals

were then rehoused together to reduce post-surgical isolation.

After recovery from the surgery, the animals were habituated to head fixation on the

experimental apparatus for one week (twice a day for up to two hours). On the day of the recording,

we first placed the subject animal in the recording set up with its head fixed, and determined the

electrode penetration path to the target area (optic tract, [-1.34, +1.87, +4.74], [-1.70, +1.87, +4.74],

or [-1.82, +2.35, +4.07] in [anterior-posterior (AP), medial-lateral (ML), dorsal-ventral (DV)]

coordinates; LGN, [-2.3, +2.3, +2.8]) using the robotic stereotaxic system (StereoDrive,

NeuroStar). The animal was then briefly anesthetized with isoflurane for about 5 minutes, and a

hole was drilled around the electrode entry point on the skull. After the removal of the anesthesia,

an acute silicon probe (P2, Cambridge Neurotech, **Fig. 1A**; or Buzsaki32L, Neuronexus, **Fig. 1F**)

coated with a fluorescent dye (DiI stain, Invitrogen, D282) was lowered at 5 µm/s using the robotic

arm until visual responses were found in the target area. A battery of visual stimuli (see below for

details) were then presented for recordings.

After the recording session, the electrode position was verified histologically. After

retracting the silicon probe, the mice were anesthetized (2.5% Avertin, 16 µL/g, intraperitoneal

injection) and perfused with paraformaldehyde (PFA; 4% in phosphate buffer solution), followed

by brain tissue harvest and overnight post-fixation in 4% PFA at 4°C. Coronal sections of the brain

tissue (thickness, 100-150 µm) were then examined under a fluorescence microscope (Leica,

LMD7000 with N2.1 filter cube) to visualize the trace left by the DiI stain on the probe (**Fig. 1A,F**).

Visual stimulation

Visual stimuli were presented as described before (Boissonnet et al., 2023). Briefly, visual stimuli

were projected to a spherical screen (radius, 20 cm) placed ~20 cm from the animal's left eye,

covering about a quarter of its visual field (±36.5° and ±22° in azimuth and elevation, respectively).

A gamma-corrected digital light processing device (Texas Instruments, DLPDLCR3010EVM-LC)

was used as a light source, where the green and the red light-emitting diodes (LEDs) were

replaced with ultraviolet (UV; 365nm, LZ1-00UV00, LED Engine) and infrared (IR; 950nm, SFH

4725S, Osram) LEDs, respectively. The UV and remaining blue channels were used for visual

stimulation (frame rate, 60 Hz; maximum intensity, 31 mW/m^2), while the IR signals were recorded

with a photodiode (PDA100A2, Thorlabs) for data synchronization. For receptive field (RF)

mapping, a black-and-white binary noise stimulus was presented for 15 minutes, consisting of a

32-by-18 pixels checkerboard patterns, where each pixel randomly and independently flickered at

60 Hz while keeping the overall luminance of each frame constant at the mean intensity.

232 Data analysis

We adapted previously established methods of spike sorting and data analysis (Boissonnet et al.,

2023; Tripodi and Asari, 2025). In brief, we used Kilosort 2.0 to sort spikes with a set of default

parameters, except for the spike detection threshold to be 6 during optimization. Single-units were

then identified by clustering in principal component space using Phy for visualization and manual

data curation. Only those units that maintained the average spike waveforms and

autocorrelograms with a minimal refractory period of 1 ms were kept for subsequent analyses.

For the visual RF analysis, we employed stimulus ensemble statistical techniques (reverse

correlation methods; 500 ms window; $\Delta t = 1/60$ s bin size). Specifically, we first obtained the linear

spatiotemporal RF of each recorded cell by calculating a spike-triggered average (STA) of the

"checkerboard" stimuli with ±1 being "white" and "black" for each pixel, respectively. As a quality

measure, *p*-value was computed for each voxel against a null hypothesis that the STA follows a

normal distribution $N(0, 1/C)$, where C is the total number of spikes. For those with $p < 10^{-5}$ at any

voxel, we ran a singular value decomposition to obtain temporal and spatial filters, respectively,

assuming the separability of RGC/LGN spatiotemporal RFs. We then fitted a two-dimensional

Gaussian envelope to those spatial filters with a single, distinct and localized feature; and

considered the center of the Gaussian as the RF center location (**Fig. 1**). Single-units with little or

no visual responses were excluded as nontarget cells, such as the axons from the parabigeminal

nucleus in the OT (Reinhard et al., 2019; Tokuoka et al., 2020). For the OT recordings, we also

applied an RF size threshold of 15° to exclude axons from wide-field cells in the superior colliculus

(Gale and Murphy, 2014; Relota et al., 2025). In total, we obtained 60 ON RGCs (peak latency of

the temporal filter, 46±5 ms, median ± median absolute deviation; RF size, 3.6°±0.9°, estimated

as the mean of the fitted Gaussian envelope short- and long-axis diameters at 1 σ), 133 OFF

RGCs (51±8 ms; 3.5°±0.9°), 137 ON LGN cells (46±4 ms; 4.7°±1.2°), and 155 OFF LGN cells

(46±5 ms; 4.3°±1.0°).

To a first approximation, at short distances, the amplitude of extracellular signals is

inversely proportional to the distance between the cell and the probe (Anastassiou et al., 2015).

We thus considered that the recorded units were physically located nearby if they both had the

largest spike waveforms on the same recording site on the probe, whereas distant if they had the

largest spike waveforms on different recording sites. Using these criteria, we identified pairs of

nearby and distant single-units that were recorded simultaneously (optic tract, 39 and 717 pairs

in total, respectively; LGN, 150 and 4341 pairs in total, respectively), and compared the distance

between their RF center locations at the population level using Mann-Whitney U-test (**Fig. 2**).

Model analysis

We used Monte Carlo methods to obtain an expected RF distance distribution of nearby cell pairs,

given a retinotopy gradient *g*. In particular, we first randomly sampled two points in a sphere,

where the radius *r* represents the maximum distance from an extracellular electrode for reliable

single-unit recording. If the distance between the two points, *D*, was larger than an exclusion

threshold *d* due to a physical mass of neurons, we then estimated the RF distance between two

neurons located at those two sampled points as *D×g*. We repeated the procedure $1000n$ times to

compute the probability distribution function of the estimated RF distance, as well as a 95%

confidence interval of the median value (of sample size *n*). For the mouse OT (*n* = 42; **Fig. 2A**),

we used *r* = 20 μm for recording range, *d* = 2 μm for RGC axon diameter, and *g* = (represented

visual field) / (optic tract diameter) = 135° / 300 μm (Paxinos and Franklin, 2001). For the mouse

LGN (*n* = 150; **Fig. 2C**), we used *r* = 40 μm for recording range, *d* = 10 μm for cell body size, and

g = 0.11°/μm (Piscopo et al., 2013).

To run simulations at different retinotopy levels for the mouse OT (**Fig. 2E-H**), we added

noise to the simulated RF locations, drawn from a Gaussian distribution $N(0, \sigma^2)$ where $\sigma=1$

corresponds to the full range of the represented visual field (135°), or equivalently, the full width

of the mouse OT (300 µm; Paxinos and Franklin, 2001). Simulations were done with $\sigma=[0, 0.5]$ in

steps of 0.05, as described above except that we randomly sampled two points within the distance

d from a randomly chosen probe position within the visual stimulation area (73° and 44° in azimuth

and elevation, respectively), and eliminated a trial if any cell on those two sampled points had its

RF outside the visual stimulation area due to the added jitter. With increasing jitter, the simulated

RF distance increased and the topographic map exhibited greater disorganization. To quantify the

level of retinotopy, we used Kendall's τ as a measure of spatial monotonicity, where $\tau=0$ indicates

a random mapping with no retinotopy; and $\tau=1$ represents a perfect retinotopy. The intersection

between the measured and the simulated RF distances was then used to estimate the level of

retinotopic organization in the mouse OT.