Tug-of-war between cortical and cytoplasmic forces shaping planar 4-cell stage embryos

Silvia Caballero-Mancebo[1], Daniel Gonzalez Suarez[1], Janet Chenevert[1], Sameh Ben-Aicha[2], Lydia Besnardeau[1], Alex McDougall[1] and Rémi Dumollard[1*]

[1]Laboratoire de Biologie du Développement de Villefranche-sur-Mer, Institut de la Mer de Villefranche-sur-Mer, Sorbonne Université, CNRS, 06230 Villefranche-sur-Mer, France.
[2]Institut de Biologie de Valrose, Université Côte-d'Azur, CNRS, 03108 Nice, France.

* Corresponding author: remi.dumollard@imev-mer.fr

ABSTRACT
Early embryonic cleavages often follow conserved geometric rules, resulting in species-specific cleavage patterns. How these rules are mechanistically implemented, however, varies widely across species and remains poorly understood. Here, using quantitative 3D live imaging, mechanical and biochemical manipulations, we dissect the mechanisms governing centrosomal complex (CC) migration and spindle orientation in ascidian 2-cell stage embryos to generate the characteristic planar, square 4-cell stage. We show that following the first mitotic division, CCs in the two blastomeres form at variable orientations relative to the mother spindle and progressively achieve parallel and coplanar alignment during interphase of the 2-cell stage. Final spindle orientation is established prior to nuclear envelope breakdown, contrasting with dynamic spindle reorientation reported in other systems.
Our analyses reveal that CC rotation is driven by forces acting on long astral microtubules and guided by an anisotropic, ER-rich cytoplasmic domain that surrounds the CCs. This ER domain correlates with a dense astral microtubule network, enabling length-dependent microtubule pulling that orients the CCs independently of cell shape. In parallel, dynein-mediated cortical pulling refines spindle tilt and maintains coplanarity. Together, these findings uncover a cell-cycle-regulated tug-of-war between ER-mediated cytoplasmic forces and anisotropic cortical forces that ensures robust planar cleavage in ascidian early embryos.

INTRODUCTION

Species-specific patterns of early cleavage divisions have fascinated developmental biologists for more than a century. Despite the extraordinary diversity of embryo sizes and shapes, early cleavage planes typically follow two conserved geometric principles: Hertwig's rule, which states that the mitotic spindle aligns along the longest axis of the cell, highlighting the role of cell shape in dictating the orientation of the division plane[1,2] and the Sach's (or orthogonal) rule, where division occurs at right angles to the previous one[3], due to either centrosome chirality or cell shape. These rules collectively explain the widespread emergence of the planar, square arrangement of blastomeres at the 4-cell stage, a configuration observed across a broad spectrum of animal taxa, including cnidarians, echinoderms, ascidians, cephalochordates, and amphibians[4–6]. Yet, despite the apparent universality of these geometric rules, the mechanisms by which they are implemented are species-dependent and remain the subject of active investigation and debate.

In many species, the physical environment of the embryo—such as the presence of a rigid eggshell—can impose significant constraints on blastomere arrangement and cleavage orientation. For example, nematode embryos, encased in a rigid eggshell, display a striking diversity of 4-cell stage morphologies, ranging from square to linear configurations, depending on the mechanical properties and the aspect ratio of the eggshell[7–9]. In contrast, unconfined embryos and embryos with a non-restrictive confinement, such as ascidian and echinoderm embryos, more consistently exhibit the archetypal flat, square 4-cell configuration[10,11]. However, this configuration is far from the mechanically stable configuration of 4 blastomeres – a tetrahedron – but arises due to the short division times of early embryos[8], suggesting that oriented cell divisions underpin the flat and square shape of 4-cell stage embryos.

How spindle positioning is controlled during the first two cell divisions has been studied in many species and the picture so far is clouded by a diversity of behaviors between species of different embryonic size[12–14] and by remaining controversies in the canonical model, the *C. elegans* early embryo, where cytoplasmic pulling and cortical pulling or pushing have all been hypothesized to play a role in spindle orientation[15–18]. To generate a symmetric and flat 4-cell stage embryo, the first two cell divisions must be centered, equal, and at 90° to each other, and the mitotic spindles of each cell of the 2-cell stage embryo must be coplanar and parallel. One modelling approach suggests that geometric constraints coupled to either yolk stratification - as in *Xenopus* or ascidian embryos - or localized cortical factors - as in sea urchin embryos - are sufficient to generate flat 4-cell stage embryos[19]. A seducing part of this theory lies in its simplicity: pulling forces applied to astral microtubules must be proportional to their length[2,20]. Such force proportional to microtubule length is thought to be achieved by dynein-dependent pulling of cytoplasmic cargoes along their whole length. However, the nature of cytoplasmic cargoes remains to be established[19,21].

In smaller cells, cortical pulling forces become more prominent. In culture cells and cells of the *Drosophila* notum epithelium, cortical pulling forces have been shown to be essential in aligning the spindle with the long axis of the cell during interphase[22,23]. In *C. elegans* zygotes/embryos, one model predicts that cortical pulling can mediate alignment to the long axis of the zygote[16], whereas strong cortical pulling has also been shown to align spindles to the short axis of the AB cell in the 2-cell stage embryo before the actomyosin cytokinetic ring refines the alignment along the long axis[17]. A stochiometric theoretical model shows that there are more cortical pulling sites close to the

aster/spindle pole, and the forces generated at these sites result in the off-centering and alignment of
the spindle with the short axis of the cell, thus having an opposing action to the length-dependent
cytoplasmic pulling forces exerted on astral microtubules[17].

In ascidian embryos, centering cytoplasmic forces and off-centering cortical forces prevail during
pronuclei migration in the zygote. Here, cortical pulling forces in the rear end of the sperm aster
counteract cytoplasmic pulling forces generated at the front end[24]. In this study, we analyze the
centrosome and spindle movements underlying the generation of a flat and square shape in ascidian
4-cell stage embryos. We find that centrosomal complexes (CCs) align progressively with the long
length of the blastomeres during interphase when cortical pulling is strong. However, since we find
that cortical pulling forces are reduced during mitosis, cortical pulling is likely to be dispensable for the
rotation of the centrosomes that may instead rely on cytoplasmic pulling on the ER that is stratified in
each cell of the 2-cell stage embryo. In line with this tug-of-war hypothesis, experimentally increasing
cortical pulling, and thus interfering with the balance between cortical and cytoplasmic pulling forces,
planar 4-cell stage embryos were transformed into non-planar embryos, highlighting that the balance
between cytoplasmic and cortical pulling results in a dynamic "tug-of-war" that determines the
cleavage pattern in early embryos.

RESULTS
Progressive coplanar and parallel alignment of the two centrosomal complexes at the 2-cell stage
In order to precisely follow the movements of the centrosomal complexes (CCs) during the 2-cell stage,
we performed 3D live imaging of embryos injected with EB3-3xVenus to label the CCs from mitosis of
1-cell stage to the beginning of cytokinesis of the 2-cell stage (Fig. 1A, movie 1). In the vegetal view,
two distinct centrosomes can be observed at each end of the spindle at anaphase of the 1-cell stage,
coinciding with the beginning of cytokinesis of the first mitosis (Fig. 1A- inset, t = 2 min). We found that
the CCs in the two forming daughter cells are not initially parallel to each other (in 60% of all analyzed
embryos) and progressively align with each other during the interphase of the 2-cell stage, in contrast
to *Xenopus* embryos, where the position of the centrosomes following the first mitosis determines the
orientation of the spindle and the plane of division at the 2-cell stage[14]. This alignment results in two
parallel CCs at nuclear envelope breakdown (NEBD) of the 2-cell stage (Fig. 1A, 16-20 min), which in
turn give rise to two parallel spindles at metaphase of the 2-cell stage (Fig. 1A, 20-26 min). To generate
the characteristic flat and square geometry of the ascidian 4-cell stage embryo, the two spindles at the
2-cell stage must not only be parallel but also coplanar. A side view (orthogonal to the vegetal view)
revealed that, indeed, the two CCs remain largely coplanar throughout the 2-cell stage (Fig. 1A, bottom
row).

To further document the complex migration pattern of CCs and how they orient over time, we tracked
the position of the anterior and posterior centrosomes within each CC in 3D (Fig. 1B) and measured
the rotation and tilt angles of each CC with respect to the mitotic spindle at 1-cell stage (Fig. 1C-D,
S1A). We found that initially the CCs form with a rotation of 68.2° ± 24.6 with respect to the mother
spindle, and that during interphase of the 2-cell stage, the CCs rotate to form a final angle of 82.5° ±
5.3 with respect to the mitotic spindle at the 1-cell stage by the time the daughter cells are in
metaphase of the 2-cell stage. Surprisingly, this final position of the CCs does not change much from

the time of NEBD, contrary to what has been observed in other organisms and somatic cells, where
the final position of the spindle is fine-tuned during metaphase and anaphase[12,15,22], suggesting that
spindle positioning in ascidian embryos is largely predetermined before NEBD rather than dynamically
adjusted during mitosis.
138
Consistent with this idea, the tilt angle between the CCs and the 1-cell stage spindle also decreased
from 25.5° ± 22.4 to 15.1° ± 11.1 (Fig. 1D) but did not converge to zero, as would be expected for a flat
square 4-cell stage embryo. This observation is in agreement with previous reports describing a
posterior-vegetal tilt of the spindle in ascidians at the 8-cell stage[6]. Comparing the orientation of the
sister CCs and spindles at the 2-cell stage further confirmed our observations that they progressively
become parallel, as indicated by a reduction in the rotation angle between them from 48.4° ± 25 to
6.6° ± 5.5 (Fig. 1E, S1B - Rotation), and coplanar, as reflected by the decrease in the tilt angle between
them from 3.9° ± 4.8 to 1.8° ± 3.1 over time (Fig. 1E, S1B - Tilt). Altogether, this analysis revealed that
the proper orientation of the CCs required to generate a flat, square 4-cell stage ascidian embryo is
progressively established during the end of mitosis at 1-cell stage and interphase at 2-cell stage, and
that by the time of mitotic entry at 2-cell stage, the final position of the future spindle is already
defined.
151
To better understand the mechanisms that orient the CCs during the end of mitosis of the 1-cell stage
and interphase of the 2-cell stage, we examined their migration at a higher spatiotemporal resolution
(Fig. 2A, movie 2). We found that the CCs form in the center of the daughter cell at some distance from
the nucleus (~17 μm) before the first cleavage is complete (Fig. 2A, 30 s, arrowhead), and that the
centrosomes within each CC remain close to each other and move together until the end of interphase
of the 2-cell stage, when the centrosomes in each CC start to separate from each other and the nucleus
positions between them (Fig. 2A-B, 12- 14min) right before NEBD. Interestingly, the final orientation
of CCs is reached when the 2 centrosomes start separating (Fig. 1D-E, 2B) indicating that the rotation
of CCs occurs without separation of the centrosomes and suggesting that there is no need for Eg5
kinesin activity for CC rotation[25]. Since the centrosomes within a CC remain close together during their
migration, we asked whether a physical link between the 2 centrosomes is necessary for their proper
migration. To that end, we performed periodic laser ablations every 90 s between the two centrosomes
in one of the 2 sister CCs from the time when both centrosomes in each CC were identifiable (anaphase
of the 1-cell stage) until the nucleus reached the CC center (Fig. 2C, movie 3). We found that neither
rotation nor tilt was affected when centrosomes were ectopically separated by laser ablations (Fig.
2C', S2A-B), and that the ablated 2-cell stage embryos still divided into a flat and square 4-cell stage.
Because centrosome separation initiates when the nucleus reaches them, we assessed whether the
CCs use the nuclei to orient/power their movements as previously observed in *C. elegans* and
Drosophila embryo[15,26–28]. To test this possibility, we performed periodic laser ablations every 90 s
between the CC and the migrating nucleus in one of the two cells from the time both centrosomes in
the CC were identifiable until the nuclei reached the center of the CC (Fig. 2D, movie 4). We found that
these ablations could impair nucleus migration (Fig. S2C) but did not affect either CC migration or their
final orientation (Fig. 2D', S2D), as the CCs/spindles eventually became parallel and coplanar as in
control embryos.
176
177 Altogether these results show that CCs movements initiate at anaphase of mitosis I and proceed during
178 cytokinesis and interphase of the 2-cell stage in ascidians. These complex movements do not rely on

179 mechanisms previously described in other embryos such as *Xenopus* embryos[14], or *C. elegans* zygotes[26]
180 or *Drosophila* blastoderm embryos[27,29] and they are not supported by the nuclei or antiparallel
181 microtubules between centrosomes.
182
CCs movements are powered by forces generated by astral microtubules
As the abovementioned mechanisms do not appear to contribute to CC migration, we considered the
astral microtubule network as the most likely source of the forces driving CC migration. We first imaged
microtubule distribution at key times during CC migration and found that the network of astral
microtubules grows at anaphase of mitosis I and expands in the whole volume of the cell during
interphase before sharply retracting at prophase of the 2-cell stage (Fig. 3A). To understand how this
network remodeling could contribute to the migration of the CCs, we analyzed the density of
microtubules from the centrosome to the cell cortex. We found that in interphase (Fig. 3A'), a dense
network of microtubules is formed extending from the centrosomes across the central cytoplasm (Fig.
3A', blue shaded area), surrounded by a less dense peripheral network of microtubules near the cortex
(Fig. 3A', red shaded area). Upon mitotic entry, this organization was lost, and the microtubule network
became largely restricted to the pericentrosomal region, with the peripheral population markedly
diminished. (Fig. 3A''). The long astral microtubules observed during interphase may be the sites of
force generation to move CCs either by cytoplasmic pulling, as demonstrated in sea urchin embryos[2],
or by pulling of astral microtubules on the cell cortex previously shown in ascidian zygotes[24]. To test
this hypothesis, we performed periodic laser ablations of the astral microtubules (every 90 s) in one of
the 2 sister cells (Fig. 3B, movie 5) from the time when both centrosomes in each CC were identifiable
(anaphase of the 1-cell stage, 2 min) until NEBD of the 2-cell stage (17 min). We found that rotation of
the CCs was severely affected in the ablated cell but not in the non-ablated one (Fig. 3B'), and that the
sister CCs/spindles at the 2-cell stage were no longer parallel (Fig. S3A – Rotation Ablated). In contrast,
the tilt was not affected by these laser ablations (Fig 3B') and the CCs and spindles remained coplanar
(Fig. S3A – Tilt Ablated). To confirm these observations, we interfered with astral microtubules with 2-
methoxyestradiol (2ME), known to preferentially affect dynamic microtubules without impairing
spindle formation[30]. Incubating zygotes at anaphase of the 1-cell stage with 2ME drastically reduced
the size of centrosomal asters during interphase of the 2-cell stage (Fig. 3C) and prevented CC
migration, leading to strong defects in both rotation and tilt angles of the spindles at the end of 2-cell
stage, losing their coplanarity (Fig. 3C'-D, S3B-C).
210
Altogether, these results demonstrate that forces exerted on astral microtubules are responsible for
the proper migration and orientation of CCs/spindles in 2-cell stage ascidian embryos. In other
systems, such forces are generated by dynein motors through two main mechanisms: cortical pulling
forces, produced by dynein motors anchored at the plasma membrane that exert tension on astral
microtubules as they move toward their minus ends[22,31–33], and cytoplasmic pulling forces, generated
by dynein motors bound to intracellular organelles such as endoplasmic reticulum or endosomes[34–36].
To explore the contribution of these mechanisms to the shaping of planar ascidian 4-cell stage
embryos, we sought to localize intracellular dynein by immunocytochemistry against the dynein
localizing complex dynactin. Using ascidian-specific anti-dynactin antibody (see Materials and
Methods, Fig. 3E, S3D), we found that dynactin is enriched both in the pericentrosomal cytoplasm and
at the cell cortex. These two pools of dynactin closely correlate with the dense central and the
peripheral microtubule populations, respectively, observed in interphase cells (Fig. 3A').
223

224   Together, these findings suggest that dynein–dynactin complexes are enriched on locations supporting
225   a role for both cytoplasmic and cortical pulling in the positioning CCs and spindles. We then sought to
226   discriminate the role of each mechanism in the positioning of the CCs.

227
An anisotropic ER domain may drive CC movements in the ascidian embryo.
We first investigated the cytoplasmic source of force generation. Previous work has shown that the
cytoplasm of ascidian embryos is a dense mixture of yolk granules and organelles[37] and that the
interaction between cytoskeletal structures and cytoplasmic components generates forces that shape
the embryo and drive cytoplasmic organizations[38]. Moreover, recent studies have suggested that
astral microtubules are anchored to the endoplasmic reticulum (ER) of the cells via dynein motors and
are able to generate pulling and pushing forces, allowing spindles to orient (ref Salle/Minc/Levy).
Because yolk granules in ascidians do not accumulate around the sperm aster, as does the endoplasmic
reticulum and, to a lesser extent, the mitochondria of the myoplasm[24], we hypothesize that neither
the yolk nor the mitochondria display dynein motors anchored to their surface. We then asked if the
ER could serve as the guiding cue for the CCs and spindles to orient. First, we analyzed the distribution
of ER over time and found that from mitosis of the 1-cell stage, the spindle and the CCs were
surrounded by a cloud of ER (Fig. 4A, S4A, movie 6), closely resembling the pericentrosomal dense
astral microtubule population and the localization of dynactin (Fig. 3A, E and S4A). Segmentation and
morphometric analysis of the ER associated with the spindle and CCs revealed that, unlike the roughly
spherical shape of the cell-averaging 0.85 ± 0.05 sphericity during interphase, the ER forms a
cytoplasmic domain with an ellipsoid shape - 0.62 ± 0.04 sphericity - and a clearly defined long axis
(Fig. 4B).

246
The observed enrichment of microtubules, dynactin, and ER suggests that the ER is covered by dynein
motors and thus could be pulled when the ER interacts with astral microtubules. If this were the case,
the CCs could leverage this anisotropic ER domain to align with its long axis, as predicted by length-
dependent pulling on microtubule to position asters and spindles[39], even in the absence of a
geometrically defined long axis of the cell. To address this hypothesis, we quantified the relative
alignment between the long axes of the ER domain and the cell, and the CCs/spindles using bootstrap
statistics (Fig. 4C). We found that initially, the long axis of the cell, ER, and the CCs are not aligned to
each other (Fig. 4C – 2 min, 4D – 2 min), displaying random orientations which results in an average
angle of ~45° between the pairs. As cytokinesis and interphase progress, the angle between the long
axes of ER and the CCs/spindle gradually decreases, indicating progressive alignment before becoming
parallel at the end of interphase (Fig. 4C – 12 min, green; 4D – 12 min, green and grey). In contrast,
neither the long axis of the ER domain nor the CCs seem to align with the long axis of the cell over the
same period (Fig. 4C), suggesting that cell shape does not provide a dominant cue for spindle
orientation. Interestingly, at NEBD, which coincides with the shortening of the astral microtubules, the
ER and the forming spindle become momentarily misaligned (Fig. 4C – 17 min, 4D – 17 min), before
realigning with the cell axis at the onset of cytokinesis. Together, this analysis shows that the shape of
the cell may not provide a consistent geometrical cue for CC orientation, and suggests that, instead,
the ER acts as a cytoplasmic scaffold that guides spindle alignment independently of cell geometry,
even though the spindles obey the Hertwig rule[39].

266
Even though the shape of the 2-cell stage blastomeres remained fairly spherical, there may still be an
effect of cell shape in the positioning of the CCs and spindles. To further test whether the ER–spindle

alignment depends on external cell geometry or arises intrinsically from cytoplasmic organization, we
cultured the embryos in calcium and magnesium-free seawater (CMF) to remove the contribution of
cell adhesion (Fig. 4E-G, movie 7). The blastomeres of these embryos were almost perfectly spherical,
displaying an average sphericity of 0.96 ± 0.04, and still divided as control ones. The rotation angle
between the 2-cell stage CCs relative to the 1-cell spindle, however, was significantly altered, whereas
the tilt angle remained unaffected (Fig. 4E'–E''). The delayed rotation of the CCs at the end of the 2-
cell stage prevented the mitotic spindles from being parallel at mitosis (Fig. S4B-C), affecting the
characteristic square geometry of 4-cell stage embryos and suggesting that cell shape has an
instructive role in CC orientation. Nevertheless, the resulting 4-cell stage embryos in CMF remained
planar due to the CCs and spindles being coplanar during the whole CC migration (Fig. S4B-C). We then
analyzed the shape of the ER domain around CCs and spindles in CMF-cultured embryos, we observed
that the ER domain was also less ellipsoidal than in control embryos - average sphericity of 0.91 ± 0.04
- supporting the idea that the ER can serve as a proxy for cell shape by reinforcing the cell shape
anisotropy. When we compared the alignment dynamics between the ER domain and CCs in CMF
embryos, we found that absolute angles were consistently higher than in control embryos, indicating
reduced alignment between the ER and the CCs until mitosis of the 2-cell stage when the CCs and ER
align together (Fig. 4H). This suggests that cell geometry primarily impacts the progressive alignment
of CCs through rotation by cytoplasmic-generated forces, whereas the tilt angle does not seem to be
influenced by cell shape.
288
289   **Anisotropic cortical pulling forces maintains the CCs coplanar during CCs migration**
290   Given the established role of cortical pulling forces in spindle orientation across metazoans, and the
291   cortical localization of dynactin we observed (Fig. 3E), we next asked if cortical pulling forces control
292   the observed posterior tilt along the anterior-posterior axis we observe.
293
To that end, we first monitored the evolution of cortical pulling forces during CC migration by imaging
plasma membrane invaginations after cortical softening with 1 μg/ml Latrunculin B (Fig. 5A-A', movie
8)[24,40]. We found that plasma membrane invaginations appear at anaphase of the 1-cell stage and that
their frequency increases significantly during interphase of the 2-cell stage before sharply decreasing
at prophase and metaphase of the 2-cell stage mitosis (Fig. 5A''). These observations show a good
correlation between the appearance of the network of long astral microtubules and the occurrence of
cortical pulling, supporting the idea that cortical pulling may also be responsible for CC movements.
We then looked at the spatial distribution of cortical pulling sites and found that membrane
invaginations were more frequent at the posterior pole of the embryo compared to the anterior pole
(Fig. S5A) and that this difference was consistent throughout the migration of the CCs and persisted
until anaphase at the 2-cell stage. These findings support our observations of a posterior tilt of the CCs
and spindles along the anterior–posterior axis (Fig. 1C, E) and suggest that asymmetrical cortical pulling
forces may play a central role in regulating the tilt angle and coplanarity of CCs and spindles, thereby
maintaining the planar architecture of the 4-cell stage embryo. Surprisingly, analysis of CCs and spindle
orientation in Latrunculin B-treated embryos (Fig. S5B) showed that neither rotation nor tilt angles are
affected by drastic softening of the embryo cortex (Fig. S5B') and that the spindles at mitosis of the
putative 2-cell stage remain parallel and coplanar (Fig. S5B''). These observations align with previous
reports that reducing actomyosin contractility has minimal impact on sperm aster positioning in
ascidian zygotes[24] or early spindle orientation[17], and are consistent with studies showing that cortical

pulling can persist when actin integrity is compromised[41], while excessive cortical tension can suppress
pulling[42]. Thus, the question remains as to whether cortical pulling can regulate CCs movements.
315
The observation of the posterior bias in cortical pulling forces and its correlation with spindle tilt led
us to hypothesize that cortical pulling forces actively control centrosome orientation and that
increasing cortical pulling would affect CCs orientation[17] by amplifying spindle tilt which would perturb
the coplanarity of the 4-cell stage embryos. In animal cells, cortical pulling forces are mediated by
dynein–NuMA–LGN–Gαi complexes anchored to the plasma membrane[22]. In this complex, the
heterotrimeric G-protein subunit Gαi recruits LGN to the cortex, which in turn binds NuMA. NuMA
then acts as a scaffold linking the cortical complex to astral microtubules, thereby positioning dynein
motors to generate pulling forces as they move toward the microtubule minus ends, generating
tension from the cortex to astral microtubules and enabling dynein to pull centrosomes and spindles
toward specific cortical regions. To test whether increasing cortical pulling alters CCs/spindle
positioning and embryo geometry, we overexpressed NuMA and targeted it to the cortex by fusing it
to a pleckstrin homology (PH) domain, which promotes membrane localization. In embryos expressing
PH-NuMA, the fusion protein predominantly accumulated at the posterior cortex (Fig. 5B, bottom row,
0 min; movie 9). The cortical pulling assay in these embryos confirmed a global increase in the number
of plasma membrane invaginations (Fig. 5B', Total) compared to control embryos (Fig. 5A''), with a
higher frequency of posterior pulling sites, confirming that cortical pulling was strengthened and
correlated with the accumulation of PH-NuMA.
333
Analysis of the orientation of CCs/spindles in embryos overexpressing cortical NuMA showed a larger
tilt angle at the 2-cell stage relative to the spindle at the 1-cell stage, while rotation remained largely
unaffected (Fig. 5C-C'', S5D, movie 10). The increased tilt was also reflected in the higher tilt angle
between sister spindles at the 2-cell stage (Fig. S5C, S5D - Tilt) that were not coplanar, resulting in
embryos adopting a more tetrahedral configuration (Fig. 5C, 26 min), in contrast to the planar
geometry observed in controls. Altogether, these results suggest a model in which proper spindle
orientation and maintenance of the planar 4-cell embryo geometry depend on a finely tuned tug-of-
war between cortical and cytoplasmic dynein-mediated pulling forces acting on astral microtubules in
a cell cycle-dependent manner (Fig. 5D).
343
344
DISCUSSION
Our study uncovers the mechanisms that drive the dynamic centrosomal complex (CC) migration and
spindle orientation in early ascidian embryos and identifies a previously underappreciated balance
between cortical and cytoplasmic forces as a key determinant of embryo geometry. Through
quantitative live imaging, perturbation assays, and biophysical manipulations, we reveal that ascidian
embryos rely on a finely tuned interplay — a tug-of-war — between dynein-dependent forces acting
from the cell cortex and those acting from cytoplasmic interactions that are most likely transmitted
through the endoplasmic reticulum (ER) network to achieve the characteristic planar 4-cell stage
configuration.
354
Our data provide a comprehensive analysis of the dynamic behavior and mechanisms underlying
centrosomal complex (CC) and spindle positioning in 2-cell stage ascidian embryos. We demonstrate
that the CCs of the two cells at 2-cell stage initially form at varying angles relative to the position of the

first mitotic spindle and progressively achieve parallel alignment during interphase of the 2-cell stage.
This progressive alignment is essential for the formation of the characteristic square 4-cell stage
ascidian embryo. Interestingly, our study revealed a second cue that drives the coplanarity of the CCs,
ensuring the planar arrangement of the blastomeres at 4-cell stage.

Previous studies in other model organisms have indicated that final spindle orientation is adjusted
during mitosis[15,17,22,42]. Notably, our data reveal that in ascidian 2-cell stage embryos, the final
orientation of the CCs and the spindle is established before mitotic entry and remains largely
unchanged during subsequent mitosis. Our findings reinforce the notion that precise spindle
orientation is a critical determinant for embryonic patterning and morphogenesis. The progressive
coplanar and parallel alignment of CCs ensures that the resulting 4-cell stage embryo adopts the
correct geometry, which is a prerequisite for subsequent morphogenetic events and fate
determination in ascidian embryos. Interestingly, the tilt angle between the CCs and the mother
spindle does not converge to zero, consistent with previous reports of a posterior-vegetal tilt in
ascidian spindles at later stages[6]. Nevertheless, the tilt angle between the CCs/spindles at 2-cell stage
is close to zero, essential for the flat, square shape of the 4-cell stage.

Our results highlight both conserved and divergent mechanisms of spindle orientation across
metazoans. In well-studied systems such as *C. elegans*, *Drosophila*, and *Xenopus* embryos, spindle
orientation is often dynamically regulated by a combination of cortical pulling forces, nuclear
positioning, and microtubule-based mechanisms[26–29]. In these organisms, the mitotic spindle can
continue to rotate and adjust its orientation well into mitosis, and dynein-mediated cortical pulling
and antiparallel microtubule sliding are critical for proper centrosome positioning[16,17]. In contrast, our
laser ablation experiments in ascidian embryos reveal that neither the integrity of the CCs nor the
migration of the nucleus is required for the final orientation of the CCs and spindles. Disruption of
centrosome separation or nuclear migration does not perturb the progressive alignment of the CCs,
nor does it affect the ability of embryos to divide into flat, square 4-cell stage embryos. This finding
suggests that ascidian embryos employ a distinct mechanism for spindle orientation that is largely
independent of the nuclear or centrosomal cues that are crucial in other systems.

The discovery that the ER serves as a proxy for cell shape and guides spindle orientation in ascidian
embryos has significant implications for the understanding of force generation and spatial organization
during early development. It suggests that organelle-based cues can play a dominant role in
orchestrating the orientation of cell division by generating cytoplasmic forces that are sufficient to
maintain directional coupling, yet whose efficiency depends on geometric constraints imposed by cell–
cell contacts. Moreover, the ER-rich domain that forms around the spindle—arising from the local
mutual exclusion of both ER sheets and yolk granules—underscores the crucial role that cytoplasmic
reorganizations and stratification have in early in development. The spatial organization of the
endoplasmic reticulum provides experimental evidence that a stratified cytoplasm can shape where
and how intracellular forces are generated and transmitted, which in turn influences spindle
behavior[19]. The persistence of planar (but not square) divisions in CMF seawater further implies that
cortical inputs primarily refine coplanarity, while ER-based cytoplasmic forces provide the dominant
contribution to CC rotation and global spindle alignment.

Together, these findings support a model in which the positioning of CCs and spindles arises from the integration of two mechanically distinct dynein-driven systems (Fig. 5D). Cytoplasmic pulling forces, transmitted through ER–microtubule coupling, act as a global aligning mechanism that orients the CCs along a cytoplasmic long axis. In contrast, cortical pulling forces, mediated by the dynein–NuMA–LGN complex, provide localized tension that fine-tunes spindle tilt and ensures coplanarity of the embryo. Proper spindle orientation thus depends on a delicate balance: cytoplasmic forces define the overall orientation, whereas cortical forces adjust and stabilize the geometry relative to the embryo's boundaries. Perturbing either component — by weakening microtubule dynamics, increasing cortical pulling, or removing cell shape constraints — disrupts this balance and results in aberrant, non-planar cleavage patterns.

Our findings expand the repertoire of known mechanisms by which cells can achieve precise spatial organization during development and highlight the evolutionary diversity of strategies employed by different organisms. Proper spindle orientation is not only critical for ensuring correct cleavage patterns and cell fate specification but also for the establishment of tissue architecture and organogenesis. The mechanisms uncovered in ascidian embryos may thus inform our understanding of how similar processes are regulated in other organisms, and may provide insights into the evolution of developmental strategies across metazoans.

ACKNOWLEDGEMENTS

We would like to thank Hervé Turlier for fruitful discussions and the Microscopy Imaging Platform (PIM) and the animal facility (CRB) of the Institut de la Mer de Villefranche (IMEV) for their continuous support. This work was supported by a Joint Project Grant from the FWF (I 3601-B27).

AUTHORS CONTRIBUTIONS

S.C.-M., A.M., and R.D. designed the research. S.C.-M. performed the experiments and analyzed the experimental data. D.G.S. contributed to the data analysis. J.C. performed the immunostainings and antibody validations. S.C.-M. and S.B-A. performed the laser ablation experiments. L.B. created the pSPE3-EB3-mScarlet, pSPE3-Reticulon-Venus, and pSPE3-PH-NuMA-Venus constructs. S.C.-M. and R.D. wrote the manuscript.

FUNDING

The project was funded by the Agence Nationale de la Recherche grants ''MorCell'' (ANR-17-CE 13-0028) and "InvBlastula" (ANR-22-CE13-0036-01)

DATA AVAILABILITY

All relevant data can be found within the article and its supplementary information. All other data supporting the findings of this study are available from the corresponding author on request.

REFERENCES

1. Hertwig, O. *Das Problem der Befruchtung und der Isotropie des Eies: eine Theorie der Vererbung*. (Verlag von Gustav Fischer, 1884).

2. Minc, N., Burgess, D. & Chang, F. Influence of Cell Geometry on Division-Plane Positioning. *Cell* 144, 414–426 (2011).

3. Sachs, J. *Über Die Anordnung Der Zellen in Jüngsten Pflanzentheilen*. (Druck und Verlag der Stahel'schen Buch- & Kunsthandlung, 1877).

4. Caballero-Mancebo, S. *et al.* Marine model organisms for mechanobiology studies. in *Evo-Devo: In Search of New Model Species* (Springer-Nature, 2026).

5. McDougall, A., Chenevert, J., Godard, B. G. & Dumollard, R. Emergence of Embryo Shape During Cleavage Divisions. in *Evo-Devo: Non-model Species in Cell and Developmental Biology* (eds Tworzydlo, W. & Bilinski, S. M.) 127–154 (Springer International Publishing, Cham, 2019). doi:10.1007/978-3-030-23459-1_6.

6. Negishi, T., Takada, T., Kawai, N. & Nishida, H. Localized PEM mRNA and Protein Are Involved in Cleavage-Plane Orientation and Unequal Cell Divisions in Ascidians. *Curr. Biol.* 17, 1014–1025 (2007).

7. Schulze, J. & Schierenberg, E. Evolution of embryonic development in nematodes. *EvoDevo* 2, 18 (2011).

8. Giammona, J. & Campàs, O. Physical constraints on early blastomere packings. *PLoS Comput. Biol.* 17, e1007994 (2021).

9. Krenger, R., Burri, J. T., Lehnert, T., Nelson, B. J. & Gijs, M. A. M. Force microscopy of the Caenorhabditis elegans embryonic eggshell. *Microsyst. Nanoeng.* 6, 29 (2020).

10. Minc, N. & Piel, M. Predicting division plane position and orientation. *Trends Cell Biol.* 22, 193–200 (2012).

11. Grill, S. W. & Hyman, A. A. Spindle Positioning by Cortical Pulling Forces. *Dev. Cell* 8, 461–465 (2005).

12. Gönczy, P., Grill, S., Stelzer, E. H., Kirkham, M. & Hyman, A. A. Spindle positioning during the asymmetric first cell division of Caenorhabditis elegans embryos. *Novartis Found. Symp.* 237, 164–175; discussion 176-181 (2001).

13. Wu, H., Nazockdast, E., Shelley, M. J. & Needleman, D. J. Forces positioning the mitotic spindle: Theories, and now experiments. *BioEssays* 39, 1600212 (2017).

14. Wühr, M., Tan, E. S., Parker, S. K., Detrich, H. W. & Mitchison, T. J. A model for cleavage plane determination in early amphibian and fish embryos. *Curr. Biol. CB* 20, 2040–2045 (2010).

15. Bondaz, A., Cirillo, L., Meraldi, P. & Gotta, M. Cell polarity–dependent centrosome separation in the C. elegans embryo. *J. Cell Biol.* 218, 4112–4126 (2019).

16. Wu, H.-Y. *et al.* Laser ablation and fluid flows reveal the mechanism behind spindle and centrosome positioning. *Nat. Phys.* (2023).

17. Middelkoop, T. C. *et al.* A cytokinetic ring-driven cell rotation achieves Hertwig's rule in early development. *Proc. Natl. Acad. Sci.* 121, e2318838121 (2024).

18. Kimura, K. & Kimura, A. Intracellular organelles mediate cytoplasmic pulling force for centrosome centration in the Caenorhabditis elegans early embryo. *Proc. Natl. Acad. Sci. U. S. A.* 108, 137–142 (2011).

19. Pierre, A., Sallé, J., Wühr, M. & Minc, N. Generic Theoretical Models to Predict Division Patterns of Cleaving Embryos. *Dev. Cell* 39, 667–682 (2016).

20. Kimura, K. & Kimura, A. A novel mechanism of microtubule length-dependent force to pull
centrosomes toward the cell center. *BioArchitecture* 1, 74–79 (2011).

21. Xie, J. *et al.* Contribution of cytoplasm viscoelastic properties to mitotic spindle positioning. *Proc.*
Natl. Acad. Sci. 119, e2115593119 (2022).

22. Anjur-Dietrich, M. I. *et al.* Mechanics of spindle orientation in human mitotic cells is determined
by pulling forces on astral microtubules and clustering of cortical dynein. *Dev. Cell* (2024).

23. Unified   quantitative   characterization   of   epithelial   tissue   development | eLife.
https://elifesciences.org/articles/08519.

24. Rosfelter, A. *et al.* Reduction of cortical pulling at mitotic entry facilitates aster centration. *J. Cell*
Sci. 137, jcs262037 (2024).

25. Yu, W.-X. *et al.* Kinesin-5 Eg5 is essential for spindle assembly, chromosome stability and
organogenesis in development. *Cell Death Discov.* 8, 490 (2022).

26. Meyerzon, M. *et al.* Centrosome attachment to the C. elegans male pronucleus is dependent on
the surface area of the nuclear envelope. *Dev. Biol.* 327, 433–446 (2009).

27. Buttrick, G. J. *et al.* Akt regulates centrosome migration and spindle orientation in the early
Drosophila melanogaster embryo. *J. Cell Biol.* 180, 537–548 (2008).

28. Tanenbaum, M. E. & Medema, R. H. Mechanisms of Centrosome Separation and Bipolar Spindle
Assembly. *Dev. Cell* 19, 797–806 (2010).

29. Wong, S.-S., Wainman, A., Saurya, S. & Raff, J. W. Regulation of centrosome size by the cell-cycle
oscillator in Drosophila embryos. *EMBO J.* 43, 414–436 (2024).

30. Kamath, K. *et al.* 2-Methoxyestradiol suppresses microtubule dynamics and arrests mitosis
without depolymerizing microtubules. *Mol. Cancer Ther.* 5, 2225–2233 (2006).

31. Laan, L. *et al.* Cortical Dynein Controls Microtubule Dynamics to Generate Pulling Forces that
Reliably Position Microtubule Asters. *Cell* 148, 502–514 (2012).

32. Kiyomitsu, T. The cortical force-generating machinery: how cortical spindle-pulling forces are
generated. *Curr. Opin. Cell Biol.* 60, 1–8 (2019).

33. Omer, S., Greenberg, S. R. & Lee, W.-L. Cortical dynein pulling mechanism is regulated by
differentially targeted attachment molecule Num1. *eLife* 7, e36745 (2018).

34. Stefano, G., Renna, L. & Brandizzi, F. The endoplasmic reticulum exerts control over organelle
streaming during cell expansion. *J. Cell Sci.* 127, 947–953 (2014).

35. Woźniak, M. J. *et al.* Role of kinesin-1 and cytoplasmic dynein in endoplasmic reticulum movement
in VERO cells. *J. Cell Sci.* 122, 1979–1989 (2009).

36. Shamipour, S., Caballero-Mancebo, S. & Heisenberg, C.-P. Cytoplasm's Got Moves. *Dev. Cell* 56,
213–226 (2021).

37. Stoev, I. D. *et al.* Active and Probe-Free Intracellular Rheology via Phase-Sensitive Thermoviscous
Flows. 2025.04.07.647540 Preprint at https://doi.org/10.1101/2025.04.07.647540 (2025).

38. Caballero-Mancebo, S. *et al.* Friction forces determine cytoplasmic reorganization and shape
changes of ascidian oocytes upon fertilization. *Nat. Phys.* 20, 310–321 (2024).

39. Dumollard, R. *et al.* The invariant cleavage pattern displayed by ascidian embryos depends on
spindle positioning along the cell's longest axis in the apical plane and relies on asynchronous cell
divisions. *eLife* 6, e19290 (2017).

40. Godard, B. G., Dumollard, R., Heisenberg, C.-P. & McDougall, A. Combined effect of cell geometry
and polarity domains determines the orientation of unequal division. *eLife* 10, e75639 (2021).

41. Redemann, S. *et al.* Membrane Invaginations Reveal Cortical Sites that Pull on Mitotic Spindles in
One-Cell C. elegans Embryos. *PLOS ONE* 5, e12301 (2010).

42. Kelkar, M. *et al.* Spindle reorientation in response to mechanical stress is an emergent property of the spindle positioning mechanisms. *Proc. Natl. Acad. Sci.* 119, e2121868119 (2022).

MATERIALS AND METHODS

Biological material

Adult *Phallusia mammillata* were collected in Sète (France) and kept at 16 °C in the aquaria of the Centre de Resources Biologiques (CRB) of the Institut de la Mer de Villefranche (IMEV) which is an EMBRC-France certified service (https://www.embrc-france.fr/fr/nos-services/fourniture-de-ressources-biologiques/organismes-modeles/ascidie-phallusia-mammillata). All experiments were performed at 18-19 °C. Oocytes and sperm were obtained after animal dissection and were used on the day of collection. Oocytes were chemically dechorionated in a 0.1 % trypsin solution (Sigma-Aldrich, T8003) in micro-filtered natural sea water (MFSW; Nalgene 0.2 μm filters, cat. no. Z370584) for 90 min under gentle stirring and kept in until fertilization with 6 μl sperm that was activated by resuspension in 1 ml of MSFW at pH 9.2.

Embryos were cultured either in MFSW or in Ca2+ and Mg2+-free sea water (CMF) supplemented with 1 mM EDTA[43].

Cloning of expression constructs

To generate the pSPE3-EB3-mScarlet plasmid, the EB3 coding sequence was amplified from pRN3-EB3-3xVenus vector using the following primers with EcoRI and XhoI sites, respectively: 5'-GAATTCAAAATGGCCGTCAATGTGTACTCCACAT-3' and 5'-CTCGAGGTACTCGTCCTGGTCTTCTTGTTGA-3'. The mScarlet coding sequence (including stop codon) was amplified from the plasmid pSPE3-iMyo-mScarlet[38] using the following primers with XhoI and NotI sites, respectively: 5'-CTCGAGATGGTGAGCAAGGGCGAGGCAGTG-3' and 5'-GCGGCCGCCTACTTGTACAGCTCGTCCATGCC-3'. To generate the pSPE3-EB3-mScarlet plasmid, the PCR products were ligated into a pSPE3-Rfa destination vector that had been previously digested with EcoRI and NotI.

To generate the pSPE3-Reticulon-Venus plasmid, the reticulon coding sequence was amplified using the following primers: 5'-CACCATGGAGTATGAAACGAGCCC-3' and 5'-CTCTGCTTTAGGCTTGGCACC-3'. The PCR product was used to generate an entry vector using the pDONR/TOPO (p3M-Reticulon). The entry vector was recombined with the destination vector pSPE3-Rfa-Venus.

To generate the pSPE3-PH-NuMA-Venus plasmid, the PH and NuMA coding sequences were amplified using hybridizing primers as follows: 5'-AAGCAGGCTCCGCGGCCGCCCCCTTCA-3' and 5'-GAGCTCAACATCATGGATGATGACGAAA-3' for PH and 5'-GAGCTCAACATCATGGATGATGACGAAA-3' and 5'-GTCTGAATCCAAAAAAGAAGGGTGGGCG-3' for NuMA. After hybridization, the full PH-NuMA coding sequence was amplified using the following primers: 5'-ACAAGTTTGTACAAAAAAGCAGGCTCCGCGGCCGCCCCCTTCA-3' and 5'-GTCTGAATCCAAAAAAGAAGGGTGGGCGCGCCGACCCAGCTTTCTTGTACAAAGTGGT-3'. The final PCR product was used to generate an entry vector using the pDONR/TOPO (p3M-PH-NuMA). The entry vector was recombined with the destination vector pSPE3-Rfa-Venus[44].

mRNA microinjections

Microinjections were performed as described[45]. Microinjection needles (Harvard Apparatus, 30-0035) were pulled with a PN-30 puller (Narishige) and mounted on a micromanipulation setup (MMO-203, Narishige) attached to an Olympus IX70 microscope. Dechorionated unfertilized eggs were placed into a glass wedge mounted onto a 400 µl Perspex mounting chamber (see for detailed protocol McDougall et al., 2014). *In vitro* mRNA synthesis was performed using the mMESSAGE mMACHINE T3 Transcription Kit 562 (Ambion, AM1343). The following mRNAs were injected: EB3-Venus[24], EB3-mScarlet (this study), Ensconsin-3xGFP[46], Ensconsin-dTomato[46], H2B-Venus[47], PH-dTomato[47], PH-NuMA-Venus (this study), and Reticulon-Venus (this study). mRNAs were injected at a concentration of 2-4 µg/µl using a high-pressure system (Narishige IM300) and incubated at 16 °C overnight before the experiment.

Myoplasm and plasma membrane labeling

Dechorionated unfertilized oocytes or zygotes were incubated in a 1 µM solution of MitoTracker™ Deep Red FM (Thermo Fisher, M22426) to label the myoplasm and/or in a 1 µM solution of CellMask™ Orange (Thermo Fisher, C10045) in MFSW for 5 min and washed in MFSW before imaging.

Cytoskeletal inhibitors

To soften the cortex, 1 µg/ml Latrunculin B (Sigma-Aldrich, L5288) was added to the imaging dish 40 minutes post fertilization (right before first cell division) and left in the medium for the remainder of the experiment.

To interfere with dynamic microtubules, 1 µg/ml 2-methoxyestradiol (Merck, 362-07-2) was added to the imaging dish 40 minutes post fertilization (right before first cell division) and left in the medium for the remainder of the experiment.

Confocal live imaging

Embryos were mounted 40 minutes post fertilization in glass bottom dishes (Cellvis, D35-10-0-N) coated with 1% low melting point agarose (Thermo Fisher, 16500100) in MFSW. Embryos were imaged in either an inverted Leica TCS SP8 confocal microscope or an inverted Leica Stellaris confocal microscope using a HCPL APO CS2 40x/1.1 NA water immersion objective (Leica) and Hybrid detectors. Embryos were imaged *in toto* with an optical section thickness of 3 µm every 120-180 s. For membrane invagination experiments, the frame rate was reduced to 30 s and for fixed samples, the optical section was 1 µm.

Laser ablation

Embryos were mounted 40 minutes post fertilization as for confocal live imaging and imaged on a Zeiss LSM980 confocal microscope equipped with a two-photon pulsed laser using a C-Aprochromat 40x/1.1 NA water immersion objective. Microtubules between centrosomes or between centrosomes and the nucleus were cut along a 10 µm line and astral microtubules were cut along 50 µm lines in 3 z planes, all of which were performed with a two-photon Coheren Axon 920-2 laser (2.5W). For ablation, the two-photon laser was used at full power, and the power received at the focal point is estimated at 2.8 $\cdot 10^{12}$ W/m^2. Embryos were imaged over 50 central z-slices with an optical section thickness of 1 µm every 13 s.

Antibody production

The anti-dynactin mouse polyclonal antibody was produced by fusing 1002 base pairs, amino acids 35 to 334, of the *Ciona intestinalis* dynactin subunit 1 protein (GenBank AP042712.1) to a 6-His tag at the C terminus and cloned into the vector pET11. Expression in bacteria was induced with IPTG, and the dynactin fusion protein was purified on a nickel column and eluted with imidazole according to the manufacturer (Potrino Ni-NTA column, Macherey-Nagel). Immunization of mice was performed by Covalab (France), and sera obtained after 57 days were used directly without further purification.

Immunostaining

Embryos were fixed in 100% methanol (VWR Chemicals, 67-56-1), which had been pre-chilled at −20°C, and then stored at -20°C. Samples were rehydrated by three washes in PBS containing 0.1% Tween (PBS-Tw), then blocked in PBS-Tw containing 1% BSA for 1 hour at room temperature. Primary antibodies to tyrosinated alpha tubulin (rat anti-alpha-tubulin tyrosinated, monoclonal YL1/2, Merck, MAB1864-I) or to dynactin subunit 1 (mouse polyclonal, custom made, see above) were added at a dilution of 1/500 in blocking buffer and incubated for at least 24 hours at 4 °C. After incubation, samples were washed three times in PBS-Tw, and fluorescent anti-rat (Jackson ImmunoResearch) or anti-mouse secondary antibodies (Jackson ImmunoResearch) were added at a dilution of 1/200. After further incubation at room temperature for 1-2 hours, samples were washed in PBS-Tw, labelled with Hoechst 33342 (final concentration 5 μg/ml, Thermo Fisher, 23491-52-3) for 10 minutes, washed twice more in PBS-Tw, and mounted on glass slides in CitiFluor AF1 (Electron Microscopy Services, 17970-25). To prevent embryo compression or breakage, coverslips were elevated by application of small spacers of modelling clay at the corners as described (ref).

Western blot

Anti-dynactin antibody's specificity was validated by Western blot analysis showing a single band at ~55 kDa corresponding to dynactin (Fig. S3D).

2 and 8-cell stage embryos were collected in a small volume of MFSW and mixed with the same volume of 2x Laemmli sample buffer (100 mM Tris-HCL pH 6.8, 4% SDS, 0.2% Bromophenol Blue, 20% glycerol and 200 mM dithiothreitol). Samples were heated for 5 min at 95 °C and then stored at -20 °C. Proteins were separated by electrophoresis on SDS-polyacrylamide gels and transferred to a nitrocellulose membrane. Membranes were stained briefly with Ponceau-S, then washed with TBS-Tw (20 mM Tris Base and 150 mM NaCl plus 0.1% Tween-20), and blocked for at least 1 hour in TBS-Tw containing 5% dry milk powder. Immunoblots were then incubated in anti-dynactin primary antibody diluted 1:1000 overnight at 4 °C. Membranes were then washed three times with TBS-Tw and incubated with an anti-mouse horseradish peroxidase-conjugated secondary antibody (Jackson ImmunoResearch, 115-035-003) at 1:10,000 dilution in TBS-Tw + 5% dry milk powder for 2 hours at room temperature. After three washes in TBS-Tw, signal detection was carried out using the SuperSignal™ West PLUS Pico chemiluminescent substrate (ThermoFisher Scientific, 34580) and images were acquired with a Fusion FX system (Vilber Lourmat, Collegien, France).

3D tracking of centrosomes and angle quantification

Confocal fluorescent Z-stacks of embryos injected with EB3-3xVenus and PH-dTomato were stabilized using the MultiStackReg[48] plugin in Fiji[49]. Centrosomes were then tracked using the MTrackJ[50] plugin in Fiji, and their 3D coordinates (Ci_x, Ci_y, Ci_z) were used to calculate the polar angles (rotation and tilt) with the y-axis (embryonic anterior-posterior axis) as reference.

Centrosome separation

The 3D coordinates of each centrosome were used to calculate the distance between each centrosome

within a CC: $d = \sqrt{(C1_x - C2_x)^2 + (C1_y - C2_y)^2 + (C1_z - C2_z)^2}$

Distance between nucleus and CC

Confocal fluorescent Z-stacks of embryos injected with H2B-Venus and EB3-mScarlet were stabilized

using MultiStackReg plugin in Fiji. Centrosomes were then tracked using the MTrackJ plugin in Fiji and

their 3D coordinates (Ci_x, Ci_y, Ci_z) used to calculate the midpoint of the CC, $M_{CC} =$

$\left(\frac{C1_x + C2_x}{2}, \frac{C1_y + C2_y}{2}, \frac{C1_z + C2_z}{2}\right)$. The nucleus of each cell was segmented and tracked using the Surface

tool in Imaris (Bitplane). Then the distance at every time point between the nucleus (Ni_x, Ni_y, Ni_z)

and the CC midpoint was calculated: $d = \sqrt{(M_x - N_x)^2 + (M_y - N_y)^2 + (M_z - N_z)^2}$. The radius

of the nucleus was subtracted from the distance to reflect the time when the edge of the nucleus

reaches the center of the CC.

Microtubules intensity profile

The intensity profiles of microtubules were made using the line tool of 15-pixel thickness in Fiji from

the centrosome to the plasma membrane. The intensity values shown were normalized to the

maximum intensity value along each line. The distance between the centrosome and the plasma

membrane was also normalized.

Membrane invagination quantification

Confocal fluorescent Z-stacks of embryos incubated in CellMask[TM] Orange to label the plasma

membrane were stabilized using MultiStackReg plugin in Fiji. 10 confocal z-slides over the equator of

the embryo were sum projected. The number of membrane invaginations was counted manually by

counting the invaginations present at 2-10 μm from the plasma membrane[24] at the times indicated in

Fig. 5A' and B.

Endoplasmic Reticulum segmentation

Confocal fluorescent Z-stacks of embryos injected with either Reticulon-Venus and PH-dTomato,

Reticulon-Venus and Ensconsin-dTomato or Reticulon-Venus and EB3-mScarlet were stabilized using

the MultiStackReg plugin in Fiji. For segmentation a Gaussian filter (sigma = 2) was applied and

segmented using Ilastik software[51] and the pixel classification method.

Plasma membrane segmentation

Confocal fluorescent Z-stacks of embryos injected with PH-dTomato were stabilized using the

MultiStackReg plugin in Fiji and segmented using Cellpose[52].

Shape analysis

Principal component analysis (PCA) of segmented membrane and ER timelapses was performed in

order to obtain the direction of the 3 main axes using a custom-made Python script. To compute the

length of the main axes (a, b, c), each voxel from the segmented images was projected onto each axis,

and sphericity was calculated as: $sphericity = \sqrt[3]{c^2/ab}$.

Data analysis and statistics

Statistical analysis was performed using Prism 8 (GraphPad). The statistical test used in each case and the resulting P values are indicated in each figure legend. N was considered an independent experiment, and n denotes the number of cells; N and n are indicated in each figure. No statistical method was used to predetermine sample size. The experiments were not randomized.

METHODS-SPECIFIC REFERENCES

43. Sardet, C. *et al.* Embryological Methods in Ascidians: The Villefranche-sur-Mer Protocols. in *Vertebrate Embryogenesis: Embryological, Cellular, and Genetic Methods* (ed. Pelegri, F. J.) 365–400 (Humana Press, Totowa, NJ, 2011). doi:10.1007/978-1-61779-210-6_14.

44. Roure, A. *et al.* A Multicassette Gateway Vector Set for High Throughput and Comparative Analyses in Ciona and Vertebrate Embryos. *PLOS ONE* 2, e916 (2007).

45. McDougall, A., Lee, K. W.-M. & Dumollard, R. Microinjection and 4D fluorescence imaging in the eggs and embryos of the ascidian Phallusia mammillata. *Methods Mol. Biol. Clifton NJ* 1128, 175–185 (2014).

46. Costache, V. *et al.* Kif2 localizes to a subdomain of cortical endoplasmic reticulum that drives asymmetric spindle position. *Nat. Commun.* 8, 917 (2017).

47. Prodon, F. *et al.* Dual mechanism controls asymmetric spindle position in ascidian germ cell precursors. *Development* 137, 2011–2021 (2010).

48. Thevenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration based on intensity. *IEEE Trans. Image Process.* 7, 27–41 (1998).

49. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* 9, 676–682 (2012).

50. Meijering, E., Dzyubachyk, O. & Smal, I. Methods for Cell and Particle Tracking. in *Methods in Enzymology* vol. 504 183–200 (Elsevier, 2012).

51. Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat. Methods* 16, 1226–1232 (2019).

52. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for cellular segmentation. *Nat. Methods* 18, 100–106 (2021).

758
759
760
Figure 1: The two CCs become parallel and coplanar during interphase of the 2-cell stage.

(A) Brightfield, confocal fluorescent images of a central z-slide of the plasma membrane (magenta),
and maximum intensity projections of centrosomes (green) in a representative ascidian embryo from
1-cell stage metaphase (time = 00:00) to 4-cell stage interphase (time = 40:05). The top and middle
rows show the vegetal view, and the bottom row shows the side view. Insets are high magnifications
of the left (dashed square) and right (solid square) CCs. Ant. is anterior, Veg. is vegetal, Ani. is animal,
Post. is posterior, and NEBD is nuclear envelope breakdown. Time is in minutes:seconds. Scale bar, 30
μm. Inset scale bar, 5 μm.

(B) Example of a 3D plot of the relative positions over time of the anterior and posterior centrosomes
of the CC in the left cell of a 2-cell stage ascidian embryo. Time is coded in the respective color bar for
each centrosome. The star represents the position of the left centrosome of the mitotic spindle at 1-
cell stage, and the grey line represents the final orientation of the spindle at mitosis of the 2-cell stage.

(C) Graphical representation of the tilt and rotation angles between the CCs and the spindles at the 2-
cell stage (dark green) and the spindle at the 1-cell stage (light green). Embryonic axes are shown for
reference.

(D) Plot of the rotation (blue) and tilt (red) angles between the CCs and the spindle at 2-cell stage and
the spindle at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical
line). The dashed line at 17 min indicates the time of NEBD. N = 9 embryos, n = 18 cells. Error bars,
standard error of the mean (s.e.m.).

(E) Plot of the rotation (blue) and tilt (red) angles between the left and right CCs and spindles at 2-cell
stage as a function of time after centrosome separation (2 min, dashed vertical line). The dashed line
at 17 min indicates the time of NEBD. N = 9 embryos. Error bars, standard error of the mean (s.e.m.).

Figure 2: Migration of the CCs and nucleus during interphase of the 2-cell stage.

(A) Confocal maximum intensity projections of the centrosomes (green), mitochondria (cyan), and

segmented nuclei (magenta) in a representative ascidian embryo during interphase of the 2-cell stage.

Ant. is anterior, Veg. is vegetal, and NEBD is nuclear envelope breakdown. Arrowhead demarcates the

ingressing cytokinetic furrow. Time is in minutes:seconds. Scale bar, 10 μm

(B) Plot of the distance between the edge of the nucleus and the center of the CC (magenta, left *y* axis;

N = 6 embryos, *n* = 12 cells) and the separation of the centrosomes within a CC (green, right *y* axis; *N*

= 9 embryos, *n* = 18 cells) as a function of time after centrosome separation (2 min, dashed vertical

line). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean

(s.e.m.).

(C) Brightfield and maximum intensity projections of confocal fluorescent images of centrosomes

(magenta, middle row) and microtubules (green, bottom row) during laser ablation of microtubules

between the centrosomes. The yellow line demarcates the ablation site. The square demarcates the

area of the inset. Ant. is anterior, Veg. is vegetal. Time is in minutes:seconds. Scale bar, 30 μm. Inset

scale bar, 5 μm.

(C') Plot of the rotation and tilt angles between the CCs and the spindle at 2-cell stage and the spindle

at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control

embryos (*N* = 9 embryos, *n* = 18 cells), ablated cell (*N* = 5 embryos, *n* = 5 cells), and non-ablated cell (*N*

= 5 embryos, *n* = 5 cells) in laser-ablated embryos between the centrosomes. The dashed line at 17

min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(D) Brightfield and maximum intensity projections of confocal fluorescent images of the DNA

(magenta) and microtubules (green) during laser ablation of microtubules between the nucleus and

the CC. The yellow line demarcates the ablation site. The square demarcates the area of the inset. Ant.

is anterior, Veg. is vegetal. Time is in minutes:seconds. Scale bar, 30 μm. Inset scale bar, 5 μm.

(D') Plot of the rotation and tilt angles between the CCs and the spindle at 2-cell stage and the spindle

at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control

embryos (*N* = 9 embryos, *n* = 18 cells), ablated cell (*N* = 4 embryos, *n* = 4 cells), and non-ablated cell (*N*

= 4 embryos, *n* = 4 cells) in laser-ablated embryos between the centrosomes and the nucleus. The

dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

Figure 3: Forces generated along astral microtubules are essential for the orientation of CCs.

(A) Confocal images of a central z-slice of ascidian embryos from 1-cell stage metaphase to 2-cell stage anaphase stained for microtubules (green) and DNA (magenta). Ant. is anterior, Veg. is vegetal. The dashed area indicates the analyzed area in A' and A''. Scale bar, 40 μm.

(A', A'') Fluorescent intensity profiles show the distribution of microtubules from the centrosome to the cell membrane in embryos in interphase (left *y* axis; *N* = 9 embryos) and in prophase (left *y* axis; *N* = 8 embryos). The green line (right *y* axis) represents the change in the average intensity profile over the distance.

(B) Brightfield and maximum intensity projections of confocal fluorescent images of microtubules (green, middle row) and centrosomes (magenta, bottom row) during laser ablation of astral microtubules. The yellow line demarcates the ablation site. The square demarcates the area of the inset. Ant. is anterior, Veg. is vegetal. Time is in minutes:seconds. Scale bar, 30 μm. Inset scale bar, 20 μm.

(B') Plot of the rotation and tilt angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (*N* = 9 embryos, *n* = 18 cells), ablated cell (*N* = 4 embryos, *n* = 4 cells), and non-ablated cell (*N* = 4 embryos, *n* = 4 cells) in laser-ablated embryos. The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(C) Brightfield and maximum intensity projections of confocal fluorescent images of microtubules (green, middle row) and centrosomes (magenta, bottom row) in embryos treated with 1μg/ml 2-methoxyestradiol (2ME). Ant. is anterior, Veg. is vegetal. Time is in minutes:seconds. Scale bar, 30 μm.

(C') Plot of the rotation and tilt angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (*N* = 9 embryos, *n* = 18 cells) and embryos treated with 2ME (*N* = 7 embryos, *n* = 14 cells). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(D) Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right) angles between the CCs and the spindle at 2-cell stage with respect to the spindle at 1-cell stage in control embryos (*N* = 9 embryos, *n* = 18 cells; rotation: paired Wilcoxon test, *P=0.0131; tilt: paired Wilcoxon test, *P=0.0250) and embryos treated with 2-methoxyestradiol (*N* = 7 embryos, *n* = 14 cells; rotation and tilt: paired t test, ns - not significant). Control vs. 2ME rotation angle: Kruskal-Wallis test (***P=0.0001) with Dunn's multiple comparisons test (ns – not significant, ***P=0.0001). Control vs. 2ME tilt angle: Kruskal-Wallis test (*P=0.0101) with Dunn's multiple comparisons test (ns – not significant, **P<0.0025).

(E) Confocal images of a central z-slice of ascidian embryos from 1-cell stage metaphase to 2-cell stage anaphase stained for dynactin (grey) and DNA (magenta). Ant. is anterior, Veg. is vegetal. Scale bar, 30 μm.

Figure 4: Cytoplasmic pulling controls the rotation of the CCs.

(A) Confocal fluorescent images of a central z-slice of the plasma membrane (magenta) and the endoplasmic reticulum (grey) in a representative ascidian embryo from 1-cell stage metaphase to 4-cell stage interphase. The middle and bottom rows show a top and side view, respectively, of the membrane (magenta) and the segmented ER (grey). Ant. is anterior, Veg. is vegetal, Ani. is animal, Post. is posterior, and NEBD is nuclear envelope breakdown. Time is in minutes:seconds. Scale bar, 30 μm.

(B) Plot of the sphericity of the cell (magenta; N = 14 embryos) and the ER (grey; N = 12 embryos) as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos. The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(C) Plot of the angle between the long axis of the ER and the cell (blue), the long axis of the ER and the CCs and spindle (green) and the long axis of the cell and the CCs and spindle (red) as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos. The dashed line at 17 min indicates the time of NEBD. Error bars, standard deviation (s.d.). Mean and s.d. were estimated by bootstrap method (5000 iterations).

(D) 3D plots of the average long axis of the cell (magenta), the long axis of the ER (grey), and the CCs and spindle (green) at 2, 12, 17, and 24 minutes of the left cells of ascidian embryos after metaphase of the 1-cell stage. NEBD is nuclear envelope breakdown. For visualization purposes, the length of the long axis of the cell (magenta) and the CCs (green) are doubled.

(E) Brightfield, confocal fluorescent images of a central z-slide of the plasma membrane (magenta), and maximum intensity projections of centrosomes (green) in a representative ascidian embryo in Ca^{2+} and Mg^{2+}-free sea water (CMF) from 1-cell stage metaphase to 4-cell stage interphase. Ant. is anterior, Veg. is vegetal, Ani. is animal, Post. is posterior, and NEBD is nuclear envelope breakdown. Time is in minutes:seconds. Scale bar, 40 μm.

(E') Plot of the rotation and tilt angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (N = 9 embryos, n = 18 cells) and embryos in CMF (N = 10 embryos, n = 20 cells). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(E'') Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right) angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage in control embryos (N = 9 embryos, n = 18 cells; rotation: paired Wilcoxon test, *P=0.0131; tilt: paired Wilcoxon test, *P=0.0250) and embryos in CMF (N = 10 embryos, n = 20 cells; rotation and tilt: paired t test, ns - not significant). Control vs. CMF rotation angle: Kruskal-Wallis test (*P=0.0194) with Dunn's multiple comparisons test (ns – not significant, *P=0.0276). Control vs. CMF tilt angle: Kruskal-Wallis test (ns – not significant) with Dunn's multiple comparisons test (ns – not significant).

(F) Confocal fluorescent images of the endoplasmic reticulum (top row) and its segmentation (middle row is the top view and bottom row is the side view) in a representative ascidian embryo from 1-cell stage metaphase to 4-cell stage interphase in CMF. Ant. is anterior, Veg. is vegetal, Ani. is animal, Post. is posterior, and NEBD is nuclear envelope breakdown. Time is in minutes:seconds. Scale bar, 30 μm.

(G) Plot of the sphericity of the cell (magenta; N = 4 embryos) and the ER (grey; N = 10 embryos) as a function of time after centrosome separation (2 min, dashed vertical line) in embryos incubated in CMF sea water. The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(H) Plot of the angle between the long axis of the ER and the CCs and spindle as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (green, taken from C) and in CMF-cultured embryos (orange). The dashed line at 17 min indicates the time of NEBD. Error bars, standard deviation (s.d.).

Figure 5: The tilt along the anterior-posterior axis is controlled by cortical pulling forces.

(A) Schematic representation of the cortical pulling assay to test the presence of cortical pulling sites
by the formation of membrane invaginations towards the CCs in a soft-cortex ascidian embryo.
(A') Confocal maximum intensity projections over 6 central z-slices of the plasma membrane (grey)
and DNA (magenta) in a representative ascidian embryo from 1-cell stage metaphase to 2-cell stage

anaphase, treated with 1μg/ml Latrunculin B from 1-cell stage anaphase. Ant. is anterior, Veg. is vegetal. Time is in minutes:seconds. Scale bar, 30 μm.

(A'') Bar plot of the average number of membrane invaginations as a function of the cell cycle stage. (*N* = 13 embryos). One-way ANOVA (****P<0.0001) and Tukey's multiple comparisons tests (****P<0.0001; ns - not significant). Error bars, standard error of the mean (s.e.m.).

(B) Confocal maximum intensity projections over 6 central z-slices of PH-NuMA (magenta) and membrane (grey) in ascidian embryos expressing PH-NuMA and treated with 1μg/ml Latrunculin B from 1-cell stage anaphase. Scale bar, 40μm.

(B') Bar plot of the average number of membrane invaginations (anterior, posterior, and total) as a function of the cell cycle stage in embryos expressing PH-NuMA and treated with 1μg/ml Latrunculin B. (*N* = 4 embryos). Kruskal-Wallis test (ns – not significant) and Dunn's multiple comparisons tests (ns - not significant). Error bars, standard error of the mean (s.e.m.).

(C) Brightfield and maximum intensity projections of confocal fluorescent images of PH-NuMA (magenta) and centrosomes (green) in embryos expressing PH-NuMA. Ant. is anterior, Veg. is vegetal, Ani. is animal, Post. Is posterior. Time is in minutes:seconds. Scale bar, 40 μm.

(C') Plot of the rotation and tilt angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (*N* = 9 embryos, *n* = 18 cells) and embryos expressing PH-NuMA (*N* = 5 embryos, *n* = 10 cells). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(C'') Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right) angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage in control embryos (*N* = 9 embryos, *n* = 18 cells; rotation: paired Wilcoxon test, *P=0.0131; tilt: paired Wilcoxon test, *P=0.0250) and embryos expressing PH-NuMA (*N* = 7 embryos, *n* = 14 cells; rotation and tilt: paired t test, ns - not significant). Control vs. PH-NuMA rotation angle: Kruskal-Wallis (*P=0.0201) with Dunn's multiple comparisons test (ns – not significant). Control vs. PH-NuMA tilt angle: Kruskal-Wallis (**P=0.0073) with Dunn's multiple comparisons test (ns – not significant, *P=0.0169).

(D) Schematic model of the two mechanisms controlling the positioning of the CCs during the 2-cell stage: cytoplasmic dynein-dependent pulling along astral microtubules controls the 90°C rotation of the CCs with respect to the 1-cell stage mitotic spindle. Dynein motors on the surface of the ER are pulled towards the CCs, thus forming the pericentromeric ER domains. These domains, in turn, act as a proxy for cell shape, dictating the rotation of the CCs during interphase to generate a square 4-cell stage embryo. Concomitantly with cytoplasmic pulling, cortical dynein generates cortical pulling forces at the end of astral microtubules that control the tilt of the CCs, ensuring the coplanarity of sister CCs and the planar geometry of the 4-cell stage ascidian embryo.

965
966
967
Figure S1: CCs of sister cells become parallel and coplanar at the 2-cell stage (relative to Fig. 1).
(A) Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right)
angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage (N = 9 embryos,
n = 18 cells; rotation: paired Wilcoxon test, *P=0.0131; tilt: paired Wilcoxon test, *P=0.0250).
(B) Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right)
angles between he left and right CCs and spindles at 2-cell stage (N = 9 embryos; rotation: paired t test,
**P=0.0017; tilt: paired Wilcoxon test, ns – not significant).

1001

1002

1003

Figure S2: Centrosome migration and alignment in laser-ablated embryos (relative to Fig. 2).

(A) Plot of the separation of centrosomes within a CC as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (grey; N = 9 embryos, n = 18 cells), ablated cell (green; N = 5 embryos, n = 5 cells), and non-ablated cell (blue; N = 5 embryos, n = 5 cells) in laser-ablated embryos between the centrosomes. The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(B) Plot of the rotation and tilt angles between the CCs and the spindle of the left and right cells at 2-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control (N = 9 embryos) and ablated embryos between the centrosomes in one cell (N = 5 embryos. The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(C) Plot of the distance between the edge of the nucleus and the centre of the CC as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (grey; N = 6 embryos, n = 12 cells), ablated cell (green; N = 4 embryos, n = 4 cells), and non-ablated cell (blue; N = 4 embryos, n = 4 cells) in laser-ablated embryos between the centrosomes and the nucleus. The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(D) Plot of the rotation and tilt angles between the CCs and the spindle of the left and right cells at 2-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control (N = 9 embryos) and ablated embryos between the centrosomes and the nucleus in one cell (N = 4 embryos). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

1024

1025

1026

1027

1028
1029
1030

Figure S3: Centrosome alignment relies on forces generated along astral microtubules (relative to Fig. 3).

(A) Plot of the rotation and tilt angles between the CCs and the spindle of the left and right cells at 2-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (N = 9 embryos), ablated cell (N = 4 embryos, n = 4 cells), and non-ablated cell (N = 4 embryos, n = 4 cells) in laser-ablated embryos. The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(B) Plot of the rotation and tilt angles between the CCs and the spindle of the left and right cells at 2-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control (N = 9 embryos) and 2-methoxyestradiol-treated (N = 7 embryos). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(C) Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right) angles between the left and right CCs and spindle at 2-cell stage in control embryos (N = 9 embryos; rotation: paired Wilcoxon test, *P=0.0131; tilt: paired Wilcoxon test, *P=0.0250) and embryos treated with 2ME (N = 7 embryos; rotation: paired t test (*P=0.0462); tilt: paired t test (ns – not significant)). Control vs. 2ME rotation angle: one-way ANOVA (***P=0.0005) with Šídák's multiple comparisons test (ns – not significant). Control vs. 2ME tilt angle: Kruskal-Wallis test (*P=0.0186) with Dunn's multiple comparisons test (ns – not significant, **P=0.0040).

(D) Western blot image of the specificity of the ascidian anti-dynactin antibody in ascidian 2-cell and 8-cell stage embryos.

1056
Figure S4: Cytoplasmic pulling controls rotation angles (relative to Fig. 4).
(A) Maximum intensity projections of confocal fluorescent images of microtubules (green) and the
edges of the ER cloud around spindles (white) in a representative ascidian embryo from 1-cell stage
metaphase to 4-cell stage interphase. Ant. is anterior, Veg. is vegetal, and NEBD is nuclear envelope
breakdown. Time is in minutes:seconds. Scale bar, 30 μm.
(B) Plot of the rotation and tilt angles between the CCs and the spindle of the left and right cells at 2-
cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control (*N*
= 9 embryos) and embryos in CMF (*N* = 10 embryos). The dashed line at 17 min indicates the time of
NEBD. Error bars, standard error of the mean (s.e.m.).
(C) Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right)
angles between the left and right CCs and spindle at 2-cell stage in control embryos (*N* = 9 embryos;
rotation: paired Wilcoxon test, *P=0.0131; tilt: paired Wilcoxon test, *P=0.0250) and embryos in CMF
(*N* = 10 embryos; rotation: paired t test, ns - not significant; tilt: paired t test, ns – not significant).
Control vs. CMF rotation angle: one-way ANOVA (***P=0.0010) with Šídák's multiple comparisons test
(ns – not significant, *P=0.0431). Control vs. CMF tilt angle: Kruskal-Wallis (ns – not significant) with
Dunn's multiple comparisons test ns – not significant).
1073
1074

1075
1076
1077

Figure S5: Cortical pulling is increased in the posterior of the ascidian 2-cell stage embryo (relative to Fig. 5).

(A) Bar plot of the average number of membrane invaginations at the anterior (Ant.) and posterior (Post.) sides of the embryo as a function of the cell cycle stage. Data is taken from Fig. 5A''.

(B) Brightfield and confocal fluorescent images of a central z-slice of the plasma membrane (magenta), and maximum intensity projections of centrosomes (green) in a representative ascidian embryo from 1-cell stage metaphase to 2-cell stage metaphase treated with 1 µg/ml Latrunculin B at 1-cell stage anaphase. Ant. is anterior, Veg. is vegetal. Time is in minutes:seconds. Scale bar, 30 µm.

(B') Plot of the rotation and tilt angles between the CCs and the spindle at 2-cell stage and the spindle at 1-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control embryos (N = 9 embryos, n = 18 cells) and Latrunculin B-treated embryos (N = 5 embryos, n = 10 cells). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(B'') Plot of the rotation and tilt angles between the CCs and the spindle of the left and right cells at 2-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control (N = 9 embryos) and Latrunculin B-treated (N = 5 embryos). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(C) Plot of the rotation and tilt angles between the CCs and the spindle of the left and right cells at 2-cell stage as a function of time after centrosome separation (2 min, dashed vertical line) in control (N = 9 embryos) and embryos expressing PH-NuMA (N = 5 embryos, n = 10 cells). The dashed line at 17 min indicates the time of NEBD. Error bars, standard error of the mean (s.e.m.).

(D) Plot of the initial (Centrosome separation) and final (Metaphase 2-cell) rotation (left) and tilt (right) angles between the left and right CCs and spindle at 2-cell stage in control embryos (N = 9 embryos; rotation: paired Wilcoxon test, *P=0.0131; tilt: paired Wilcoxon test, *P=0.0250) and embryos overexpressing PH-NuMA (N = 5 embryos; rotation: paired t test, ns - not significant; tilt: Wilcoxon test, ns – not significant). Control vs. PH-NuMA rotation angle: one-way ANOVA (***P=0.0005) with Šídák's multiple comparisons test (ns – not significant). Control vs. PH-NuMA tilt angle: Kruskal-Wallis test (*P=0.0210) with Dunn's multiple comparisons test (*P=0.0146, ns – not significant).

MOVIE LEGENDS

Movie 1. Centrosomal complex orientation during the first two cleavages in ascidian embryos, related to Fig. 1 and S1. Time-lapse brightfield (left panel) and fluorescent confocal video of a representative ascidian embryo injected with PH-dTomato mRNA (magenta, right panel) and EB3-3xVenus mRNA (green, right panel) to mark membrane and centrosomes, respectively. Time interval, 2 min.

Movie 2. Centrosome and nucleus migration during interphase of the 2-cell stage in ascidian embryos, related to Fig. 2. Time-lapse fluorescent confocal video of a representative ascidian embryo injected with EB3-mScarlet mRNA (green), H2B-Venus mRNA (magenta) to mark centrosomes and the DNA, respectively, and incubated in MitoTracker Deep Red (cyan) to label the mitochondria. Time interval, 16 s.

Movie 3. Laser ablation of intercentrosomal microtubules during CC migration at the 2-cell stage of ascidian embryos, related to Fig. 2 and S2. Time-lapse fluorescent confocal video of a representative ascidian embryo injected with EB3-mScarlet mRNA (magenta) and Ensconsin-3xGFP mRNA (green) lo mark centrosomes and microtubules, respectively. Ablation site is indicated by the dashed line. Time interval, 12.8 s.

Movie 4. Laser ablation of microtubules between the CC and the nucleus during CC migration at the 2-cell stage of ascidian embryos, related to Fig. 2 and S2. Time-lapse fluorescent confocal video of a representative ascidian embryo injected with H2B-Venus mRNA (magenta) and Ensconsin-dTomato mRNA (green) lo mark DNA and microtubules, respectively. Ablation site is indicated by the dashed line. Time interval, 12.8 s.

Movie 5. Laser ablation of astral microtubules during CC migration at the 2-cell stage of ascidian embryos, related to Fig. 3 and S3. Time-lapse fluorescent confocal video of a representative ascidian embryo injected with EB3-mScarlet mRNA (magenta) and Ensconsin-3xGFP mRNA (green) lo mark centrosomes and microtubules, respectively. Ablation site is indicated by the dashed line. Time interval, 12.8 s.

Movie 6. Endoplasmic reticulum rearrangement during the first two cleavages in ascidian embryos, related to Fig. 4 and S4. Time-lapse fluorescent confocal video of a representative ascidian embryo injected with PH-dTomato mRNA (magenta) and Reticulon-Venus mRNA (grey) to mark the membrane and endoplasmic reticulum, respectively. Time interval, 110 s.

Movie 7. Centrosomal complex orientation during the first two cleavages in ascidian embryos cultured in Ca^{2+} and Mg^{2+}-free seawater, related to Fig. 4 and S4. Time-lapse brightfield (left panel) and fluorescent confocal video of a representative ascidian embryo injected with PH-dTomato mRNA (magenta, right panel) and EB3-3xVenus mRNA (green, right panel) to mark membrane and centrosomes, respectively. Time interval, 2 min.

Movie 8. Cortical pulling assay in control ascidian embryos, related to Fig. 5 and S5. Time-lapse fluorescent confocal video of a representative ascidian embryo treated with in 1μg/ml Latrunculin B and injected with H2B-Venus mRNA (magenta) to mark the DNA, and incubated in Cell Mask Orange to label the plasma membrane (grey). Time interval, 20 s.

1184   **Movie 9.** Cortical pulling assay in ascidian embryos expressing PH-NuMA, related to Fig. 5 and S5. Time-

1185   lapse fluorescent confocal video of a representative ascidian embryo treated with in 1μg/ml

1186   Latrunculin B and incubated in Cell Mask Orange to label the plasma membrane (grey, left panel) and

1187   injected with PH-NuMA-Venus mRNA (magenta, right panel) to mark exogenous cortical NuMA. Time

1188   interval, 16 s.

1189

1190   **Movie 10.** Centrosomal complex orientation during the first two cleavages in ascidian embryos

1191   expressing PH-NuMA to increase cortical pulling, related to Fig. 5 and S5. Time-lapse brightfield (left

1192   panel) and fluorescent confocal video of a representative ascidian embryo injected with PH-NuMA-

1193   Venus mRNA (magenta, right panel) and EB3-mScarlet3 (green, right panel) to mark exogenous cortical

1194   NuMA and centrosomes, respectively. Time interval, 2 min.

1195