Transcriptomic insights into the tripartite plant–pathogen–mycoparasite interaction

reveal novel mechanisms involving fungal secretomes and plant amino acid metabolism

Kazuya Maeda[1], Mariko Kouda[1], Mai Ohara[1], Takumi Kawase[1], Koki Saito[1], Eishin Iwao[1],

Hirotoshi Sushida[2], Tomoko Suzuki[3], Takuya Sumita[4], Yuichiro Iida[1]

[1] Laboratory of Plant Pathology, Faculty of Agriculture, Setsunan University, Nagaotoge-cho

45-1, Hirakata, Osaka 573-0101, Japan

[2] Institute of Food Research, National Agriculture and Food Research Organization (NARO),

2-1-12 Kannondai, Tsukuba, Ibaraki 305-8642, Japan

[3] Department of Chemical and Biological Sciences, Faculty of Science, Japan Women's

University, Mejirodai 2-8-1, Bunkyo-ku, Tokyo 112-8681, Japan

[4] Department of biological resources management, School of Environmental Science, The

University of Shiga Prefecture, Hassaka-cho 2500, Hikone, Shiga 522-8533

* **Correspondence to Y. Iida (yuichiro.iida@setsunan.ac.jp)**

Present address (Mariko Kouda): Graduate School of Human Sciences, Kobe College,

Okadayama 4-1, Nishinomiya, Hyogo, 662-8505, Japan

Short title: Transcriptomic analysis among plant, pathogen and mycoparasite

ORCIDs: Kazuya Maeda: 0000-0001-9858-0362; Mariko Kouda: 0009-0000-8862-4202;

Takumi Kawase: 0009-0006-2448-0805; Koki Saito: 0009-0007-5348-7739; Eishin Iwao:

0009-0000-6697-5378; Hirotoshi Sushida: 0000-0003-0631-6068; Tomoko Suzuki: 0000-

0001-9561-968X; Takuya Sumita: 0000-0001-7980-018X; Yuichiro Iida: 0000-0001-5062-

3255.

Word Count

Introduction: 905

Materials and Methods: 2,094

Results: 2,484

Discussion: 1,964

Total: 7,447

Number of Figures, Tables, and Supporting Information

Figures: 8 (all to be published in colour)

Tables: 0

Supporting Information: 18

Summary

·   The tomato–*Cladosporium fulvum* (syn. *Fulvia fulva*) pathosystem has served as a model

for the gene-for-gene concept of effectors and resistance proteins, but this binary framework

does not include the potential influence of other microbial participants. Here, we discovered a

unique mechanism involved in the tripartite interaction among *C. fulvum*, its mycoparasitic

fungus *Hansfordia pulvinata*, and host tomato plant.

·   Transcriptomic analyses of under mycoparasitic conditions *in planta* and *in vitro* revealed

that in *H. pulvinata* genes encoding small secreted proteins were upregulated during

mycoparasitism. Notably, a Nep1-like protein (HpNlp1) lacked typical necrosis-inducing

activity but induced the accumulation of antifungal substances in the plant apoplast,

inhibiting spore germination of *C. fulvum.*

·   In *C. fulvum* parasitized by *H. pulvinata*, effector genes were highly expressed. Strikingly,

effector protein Ecp2 was found to share structural similarity with pathogen killer toxin 4

proteins and had broad-spectrum antifungal activity, indicating a dual function in fungal

competition and *Cf-ECP2*-mediated plant resistance.

·   In tomato plants, when *C. fulvum* was parasitized by *H. pulvinata*, amino acid biosynthetic

pathways and defense-related genes were exclusively activated. These results suggest that in

the tripartite molecular network, the mycoparasite simultaneously suppressed the pathogen

and induced plant resistance. In this newly discovered multilayered molecular network, the

mycoparasite coordinates pathogen suppression and plant defense within the tripartite

interaction.

Keywords: mycoparasitism, Nep1-like protein, Ecp2 effector, antifungal proteins, amino

acid metabolisms, *Cladosporium fulvum*, *Hansfordia pulvinata*, *Solanum lycopersicum*

INTRODUCTION

The biotrophic filamentous fungus *Cladosporium fulvum* Cooke [syn. *Fulvia fulva* (Cooke)
Cif.] causes tomato leaf mold, a serious economic threat due to reduced fruit yield and quality
and occasional death of the tomato plant (*Solanum lycopersicum* L.) (Thomma *et al*., 2005;
de Wit *et al*., 2009). The fungus enters tomato leaves through open stomata in high humidity,
colonizes the apoplast, and secretes numerous effector proteins to suppress plant immunity
and facilitate infection (Mesarich *et al*., 2023). Commercial tomato cultivars have thus been
bred with *Cf* resistance genes encoding receptor-like proteins (RLPs) with extracytoplasmic
leucine-rich repeats from wild relatives (Mesarich *et al*., 2023). These RLPs recognize the
corresponding *C. fulvum* effector proteins as virulence factors via a gene-for-gene
relationship and trigger a hypersensitive response to restrict the growth of *C. fulvum*
(Mesarich *et al*., 2023). However, new races of *C. fulvum* can rapidly evolve through
mutation or loss of effector genes and evade plant immunity (Iida *et al*., 2015; de la Rosa *et
al*., 2024). The extensive use of resistant cultivars carrying only one or a few *Cf* genes has
consequently led to the emergence of *C. fulvum* races that can overcome the resistance of the
tomato plants (Luderer *et al*., 2002; Mesarich *et al*., 2014).

To date, over 100 effector genes have been identified in *C. fulvum* (Mesarich *et al*., 2018;
Zaccaron & Stergiopoulos, 2024). Most encode small cysteine-rich proteins that contain an
N-terminal signal peptide for secretion. The identified avirulence (Avr) and extracellular
proteins (Ecp) effectors—such as Avr2, Avr4, Avr4E, Avr5, Avr6 (Ecp5), Avr9B, Avr9C,
Ecp1, Ecp2, Ecp4, and Ecp6—exemplify the complex coevolution between fungal virulence
factors and their cognate *Cf* immune receptors (Wulff *et al*., 2009; de la Rosa *et al*., 2024;
Shol *et al*. 2025). Several effectors have been functionally characterized. Avr2 acts as a
cysteine protease inhibitor that targets the apoplastic cysteine proteases Rcr3 and Pip1,
thereby interfering with basal defense mechanisms conserved across solanaceous plants
(Rooney *et al*., 2005; Shabab *et al*., 2008; Kourelis *et al*., 2020). Ecp6 suppresses chitin-
triggered immunity by sequestering chitin oligomers released from the fungal cell wall, thus
evading recognition by host pattern-recognition receptors (de Jonge *et al*., 2010). In
contrast, Ecp2 induces cell death in both tomato and nonhost plants, suggesting a dual role as
a potential virulence and avirulence factor (de Kock *et al*., 2004; de Wit *et al*.,
2012). Notably, the Ecp2 effector contains an Hce2 (homolog of *Cladosporium fulvum Ecp2*)
domain that is widely conserved across diverse fungal and bacterial taxa, implying a
fundamental role for this domain in microbe–plant interactions (Stergiopoulos *et al*., 2012).

However, the biochemical function of Ecp2 remains unclear. Of additional concern is that *C. fulvum* has developed resistance to fungicides (Yan *et al*., 2008; Watanabe *et al*., 2017). The extensive breakdown of the resistance conferred by single *Cf* resistance genes and the emergence of fungicide-resistant strains emphasize the need for sustainable disease management strategies beyond conventional breeding and chemical approaches. To better understand the molecular basis of disease suppression, previous studies have primary focused on the *C. fulvum*–tomato interaction within the framework of the gene-for-gene model, but tripartite interactions involving additional microorganisms remain largely unexplored.

Hansfordia pulvinata (Berk. & M.A. Curtis) S. Hughes (syn. *Dicyma pulvinata*) is a mycoparasitic fungus serendipitously discovered on tomato leaves infected with *C. fulvum* in a greenhouse (Iida *et al*., 2018). It overgrows *C. fulvum* lesions on the plant host, effectively suppressing disease progression, and has broad mycoparasitic activity against all known *C. fulvum* races (Iida *et al*., 2018). *H. pulvinata* recognizes *C. fulvum* hyphae, coils around them, and directly penetrates the hyphae, then forms intercellular structures within the host fungus (Peresse & Le Picard, 1980; Maeda *et al*., 2025). It also produces an eremophilane-type sesquiterpenoid, 13-deoxyphomenone (also known as sporogen-AO1), which has antifungal activity against *C. fulvum* (Tirilly *et al*., 1983; Maeda *et al*., 2025). The compound was initially identified as a sporulation-inducing factor in the koji mold *Aspergillus oryzae* (Tanaka *et al*., 1984a, 1984b). Comparative genomics recently revealed that the biosynthetic gene cluster for 13-deoxyphomenone (*DPH*) in *H. pulvinata* shares high synteny with that in *Aspergillus* species, suggesting horizontal gene transfer from an ancestoral *Aspergillus* lineage (Maeda *et al*., 2025). Subsequently, *H. pulvinata* appears to have retained this gene cluster to support its mycoparasitic lifestyle, repurposing the endogenic sporogenesis function in *Aspergillus* species into exogenic antifungal activity that inhibits spore germination and hyphal elongation of *C. fulvum*. However, the molecular mechanisms underlying its mycoparasitism remain largely unknown. Since its mycoparasitism is not restricted to the phyllosphere (Maeda *et al*., 2025), comparative analyses of fungal–fungal interactions under different mycoparasitic conditions could provide new insights on the molecular basis of fungal host recognition and leaf-surface biocontrol mechanisms.

The *C. fulvum*–tomato interaction has long served as a valuable model for dissecting gene-for-gene dynamics in plant–pathogen systems (Mesarich et al. 2023.). Recent chromosome-scale genome assemblies of multiple *C. fulvum* races have substantially advanced our understanding of its genome structure and effector repertoire (Zaccaron *et al*., 2022; Zaccaron & Stergiopoulos, 2024). Nevertheless, the molecular basis governing fungal–

fungal recognition and the tripartite interactions with tomato plants remain largely
uncharacterized. Here, we present transcriptomic evidence revealing how the mycoparasitic
fungus *H. pulvinata* modulates both *C. fulvum* and tomato responses, providing molecular
insights into the interplay among plant, pathogen and mycoparasite. These findings
underscore the potential of the foliar mycoparasitic fungus as promising biocontrol agents
within complex plant–microbe ecosystems.
141

MATERIALS AND METHODS

Culture conditions for fungal strains and plants, treatments and assays

Fungal strains *H. pulvinata* 414-3 and *C. fulvum* CF301 were cultured on half-strength potato
dextrose agar (PDA; BD Difco, Sparks, MD, USA) or minimal medium (MM) agar (15 g
sucrose, 5 g ammonium tartrate, 1 g NH_4NO_3, 1 g KH_2PO_4, 0.5 g $MgSO_4 \cdot 7H_2O$, 0.1 g NaCl,
0.1 g $CaCl_2 \cdot H_2O$, 25 μL 0.2 mg mL^{-1} biotin, 15 g agar and 1 mL trace elements per liter) at
25 °C in the dark for 1 (*H. pulvinata*) and 2 weeks (*C. fulvum*) (Maeda *et al*., 2025). Spores
were collected in distilled water and adjusted to 1×10^6 spores mL^{-1} using a hemocytometer.
For *in vitro* assays, 100 μL of one of the spore suspensions was spread onto nylon
membranes on PDA and incubated for 1 week for *H. pulvinata* (hereafter "*in vitro Hp*") or 2
weeks for *C. fulvum* ("*in vitro Cf*") (Fig. **S1a**). For set up the *in vitro* mycoparasitic intection,
membranes bearing *C. fulvum* colonies were transferred to water agar (nutrient-poor) and
sprayed with 1 mL of *H. pulvinata* spore suspension (1×10^6 spores mL^{-1}); plates were then
incubated at 25 °C in the dark for 2 weeks ("*in vitro Cf/Hp*") (Fig. **S1a**).
Seedlings of tomato (*Solanum lycopersicum*) cv. Moneymaker (lacking all known *Cf*
resistance genes) were grown at 25 °C with 16 h light/8 h dark. Four-week-old plants were
either grown in high humidity as healthy controls for another 2 weeks (healthy leaf; Fig. **S1a**)
or sprayed with approximately 3 mL of a *C. fulvum* spore suspension (1×10^6 spores mL^{-1}),
then incubated in high humidity for 2 weeks until lesions developed ("*in planta Cf*"
interaction; Fig. **S1a**). For establishing the mycoparasitism in plants, plants with leaf mold
lesions were sprayed with approximately 3 mL of *H. pulvinata* spore suspension (1×10^6
conidia mL^{-1}) and incubated under high humidity at 25 °C for 2 weeks until lesions were
overgrown by white mycelia of *H. pulvinata* ("*in planta Cf/Hp*"; Fig. **S1a**).
N. benthamiana plants were grown for 4 weeks at 25 °C with 16 h light/8 h dark.
Arabidopsis thaliana ecotype Col-0 seeds were surface-sterilized in 70% (v/v) ethanol for 1
min, rinsed three times with sterile distilled water, plated on Murashige and Skoog (MS) agar

(4.33 g MS basal medium (Sigma-Aldrich, St. Louis, MO, USA) and 20 g sucrose per liter,

pH 5.7), stratified at 4 °C in the dark for 1 month, then transferred to soil and grown for 5

weeks at 25 °C with 16 h light/8 h dark.

Scanning electron microscopy (SEM)

Sample preparation and imaging procedures for SEM were described previously (Maeda *et*

al., 2025). Nylon membranes with fungal cultures were cut into squares with sides of about 5

mm and fixed in 2.5% (v/v) glutaraldehyde in 100 mM cacodylate buffer (pH 7.4) overnight

at 4 °C, then 1% (w/v) osmium tetroxide in 100 mM cacodylate buffer (pH 7.4) for 1 h at

room temperature. Samples were then dehydrated with a graded ethanol series, critical-point

dried in absolute ethanol using a Leica EM CPD300 critical point dryer (Leica Microsystems,

Wetzlar, Germany), then mounted on stubs, coated with platinum-palladium, and observed

with a Hitachi SU-8220 scanning electron microscope (Tokyo, Japan).

RNAseq and transcriptome analysis

Three biological replicates were collected from the fungal colonies on membranes or from

leaves for each treatment described above. Total RNA was extracted using the RNeasy Plant

Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. Poly(A)

mRNA was enriched with the NEBNext Poly(A) mRNA Magnetic Isolation Module (New

England Biolabs, Ipswich, MA, USA), and libraries were prepared using the NEBNext Ultra

II Directional RNA Library Prep Kit (New England Biolabs). Libraries were sequenced on an

Illumina NovaSeq 6000 platform (Illumina, San Diego, CA, USA) with the PE150 kit (150

bp × 2 paired-end), producing 2 or 4 Gb of data per sample.

Raw read quality was assessed using FastQC Ver. 16.0.1

(bioinformatics.babraham.ac.uk/projects/fastqc). Low-quality bases and adapter sequences

were removed using Trimmomatic Ver. 0.39 (Bolger *et al*., 2014). Clean reads were aligned

to the respective reference genomes using HISAT2 Ver. 2.2.1; SAM files were converted to

BAM files with SamTools Ver. 1.18 (Li *et al*., 2009; Kim *et al*., 2019). Read counts were

generated using Rsubread Ver. 2.20.0. Differentially expressed genes (DEGs) were identified

with DESeq2 using thresholds of $|\log_2$ Fold-change$| \geq 1$ and adjusted *P*-value (padj) ≤ 0.05

(Love *et al*., 2014). Gene Ontology (GO) and KEGG enrichment analyses were performed

using TBtools-II Ver. 2.119 (Aleksander *et al*., 2023; Chen *et al*., 2023; Kanehisa *et al*.,

2025). Data were visualized using SRplot (Tang *et al*., 2023).

Bioinformatic analysis

Orthologous coding sequences were identified using OrthoFinder Ver. 2.5.5 (Emms & Kelly, 2019); 100 single-copy orthologs were selected for multiple sequence alignment. Sequence alignments were generated using MAFFT Ver. 7.526 and subsequently trimmed with TrimAl Ver. 1.4 (Katoh and Standley, 2013; Capella-Gutiérrez *et al*., 2009). Maximum-likelihood phylogenetic trees were constructed with RAxML Ver. 8.2.12 using 1,000 bootstrap replicates and the GTRCAT or PROTCATAUTO model (Stamatakis, 2014). The consistency of the tree topology was assessed using approximately unbiased (AU) tests (Shimodaira, 2002) with 10,000 bootstrap replicates as previously described (Maeda *et al*., 2025). Signal peptides were predicted with SignalP-6.0 (Teufel *et al*., 2022), candidate effectors with EffectorP-fungi 3.0 (Sperschneider & Dodds, 2022), and subcellular localization with LOCALIZER (Sperschneider *et al*., 2017). Secondary metabolite biosynthetic gene clusters were identified using the fungal version of antiSMASH (Blin *et al*., 2023). GPI-anchored proteins were predicted with NetGPI 1.1 (Gíslason *et al*., 2021), and carbohydrate-active enzymes (CAZymes) were annotated using DIAMOND (Zhang *et al*., 2018). Default parameters were used for all analyses unless otherwise specified.

For sequence similarity comparison, the EMBOSS Needle tool was used for pairwise alignments (Madeira *et al*., 2024). Protein domains and motifs were predicted using the Pfam database and the MEME Suite, respectively (Bailey *et al*., 2015; Mistry *et al*., 2021). The 3D structures of proteins were predicted using ColabFold Ver. 1.5.5) (Mirdita *et al*., 2022). Five models were generated with three iterative recycles each, and the models were ranked according to their predicted local distance difference test (pLDDT) scores. The top-ranked models were refined by AMBER energy minimization to resolve steric clashes and improve stereochemical geometry. The resulting relaxed structures were used for subsequent structural analyses. The structure of Ecp2 was analyzed using the Dali server to retrieve structural homologs based on a non-redundant data set clustered at 25% sequence identity (PDB25), and structures were compared with TM-align (Zhang & Skolnick, 2005; Holm, 2022).

Plasmid construction and protein expression

NLP coding sequences were synthesized (VectorBuilder, IL, USA) and cloned into the pSfinx vector using a primer set (Table **S1**) and NEBuilder HiFi assembly kit (New England Biolabs). The NLP genes were digested with restriction enzyme EcoRI and subcloned into the pPIC9K vector (Thermo Fisher Scientific, Waltham, MA, USA) for expression in *Pichia*

pastoris GS115 strain according to the manufacturer's instructions. Transformants were

selected on YPD medium (Fujifilm Wako, Osaka, Japan) containing G418 (Fujifilm Wako)

and confirmed by PCR using the primer set in Table **S1**. Expression was induced by adding

5 mL of the pre-culture grown in BMGY broth to 100 mL of BMMY supplemented with

1.5% (w/v) methanol, and incubating the culture at 30 °C and 300 rpm for 96 h. Supernatants

were collected by centrifugation at 12,000 rpm for 10 min.

The gene *Ecp2* was amplified from genomic DNA of *C. fulvum* using PCR and the

primer set in Table **S1** and cloned into the pET-28b vector (Merck, Darmstadt, Germany) to

express an N-terminal His_6-tagged protein in *Escherichia coli* strain BL21(DE3). Expression

was induced in LB medium with 1 mM IPTG at 16 °C for 16 h. Cells were resuspended with

50 mM KPi buffer (pH 7.4) and disrupted by sonication on ice (10 min: 2 s on–4 s off).

Insoluble fractions were solubilized in 10 mL of denaturing buffer (6 M guanidine

hydrochloride, 10 mM Tris, 10 mM β-mercaptoethanol, pH 8.0) for 1 h at room temperature.

Proteins were purified in a Ni-affinity column HisTrap HP (Cytiva, Medemblik, The

Netherlands), concentrated using Vivaspin 6 (GE Healthcare, Chicago, IL, USA) and

quantified with a Qubit fluorometer (Thermo Fisher Scientific). BL21(DE3) carrying empty

pET-28b served as a negative control.

Western blot analysis

All protein samples were heated at 95 °C for 10 min in sample buffer (1% SDS, 1% 2-

mercaptoethanol, 10 mM Tris–HCl, 20% (v/v) glycerol, pH 6.8) for denaturation. The

proteins were separated in 12% SDS-polyacrylamide gel and transferred onto PVDF

membrane (Bio-Rad, Hercules, CA, USA). Membranes were blocked with 5% skim milk in

TBS-T for 1 h before antibody incubation. Anti-His and anti-HA monoclonal antibodies (Cell

Signaling Technology, Danvers, MA, USA) were diluted in Can Get Signal solution 1

(1:10,000) (TOYOBO, Osaka, Japan) and used as primary antibodies to detect Ecp2 and

NLPs, respectively. Horseradish peroxidase-conjugated anti-rabbit antibody was diluted with

Can Get Signal solution 2 (1:2,000) (TOYOBO) for use as the secondary antibody. Signals

were visualized using an ECL western detection kit (GE Healthcare) and captured using a

ChemiDoc imaging system (Bio-Rad).

Functional assay of NLPs

pSfinx plasmids were introduced into *Agrobacterium tumefaciens* strain GV3101 via

electroporation (Ma *et al*., 2012). Agrobacterium suspensions were adjusted to $OD_{600} = 0.3$ in

infiltration buffer (2.03 g MgCl$_2$·6H$_2$O, 1.95 g MES, 500 µL of 200 mM acetosyringone per liter, pH 5.6) and used to infiltrate leaves on 4-week-old plants of *N. benthamiana* by syringe pressure. Leaves were monitored for necrosis 14 days after infiltration. Necrosis was visualized by trypan blue staining (Ma *et al.*, 2012) and quantified using ilastik (Ver. 1.4.1) (Berg *et al.*, 2019).

P. pastoris culture supernatant containing NLP proteins or BM medium (BMGY without glycerol) was mixed at a 1:1 ratio with *C. fulvum* spore suspension (1×10^6 spores mL^{-1}). A 40 µL drop of each mixture was placed on a hydrophobic glass slide and incubated at room temperature for 24 h. After incubation, germination of 200 spores per treatment was assessed using a Nikon E600 light microscope (Tokyo, Japan). Culture supernatants containing NLP proteins were collected from *P. pastris* and used to syringe-infiltrate *N. benthamiana* leaves. Twenty-four hours after infiltration, apoplastic fluid was collected by vacuum infiltration with distilled water, followed by centrifugation (1,000 rpm, 10 min) (Joosten, 2012). Equal volumes (10 µL) of *C. fulvum* spore suspension and apoplast fluid were then incubated at room temperature for 24 h. A light microscope (Nikon) was then used to assess 200 spores for germination.

Synthetic peptides of nlp24 (GenScript Biotech, Nanjing, China) were prepared in 500 nM stock solutions in dimethyl sulfoxide. Reactive oxygen species (ROS) were detected as previously described with slight modifications (Yang *et al.*, 2022). Leaf disks (0.125 cm²) of *N. benthamiana* were floated on sterile distilled water for 24 h, then treated with 10 µL of a substrate mixture (5 µL of 10 mM L-012 [Fujifilm Wako] and 5 µL of protein or peptide). Luminescence was recorded every 2 min for 40 min using a Tristar3 Multimode Reader (Berthold Technologies, Bad Wildbad, Germany).

Functional assays of Ecp2

The abaxial side of *Cf-ECP2* tomato leaves was infiltrated with purified Ecp2 protein (final concentration 1 µM) using a syringe. Leaves were incubated at 25 °C with 16 h light/8 h dark for 6 d and checked for a hypersensitive response (HR). *In vitro* antifungal activity was assayed as previously described (Chavarro-Carrero *et al.*, 2024). Briefly, fungal spores were suspended in 5% PDB containing 15 µM Ecp2 or empty vector control (1×10^4 spores mL^{-1}), then 100 µL of the samples (three wells per sample) was incubated in wells of 96-well plates at 25 °C for 4 days. For each well, five fields of view were photographed ($N = 15$). Fungal growth was quantified and compared with the control using the Hybrid Cell Count module of

the BZ-800X (KEYENCE, Osaka, Japan).

Statistical analyses

R version 4.4.2 (R Core Team, 2024) was used for all analyses. The proportion of necrosis

relative to total leaf area in *N. benthamiana* was assessed using Dunnett's test, with the empty

vector as the control ($N = 3$). Spore germination rate of *C. fulvum* treated with NLPs was

analyzed by one-way ANOVA followed by Tukey's HSD test ($N = 3$). The spore germination

rate of *C. fulvum* in apoplastic fluid from *N. benthamiana* leaves treated with NLPs was

compared using the Mann–Whitney *U* test, with the buffer as the control ($N = 12$). The

relative growth rate of fungi and yeast treated with Ecp2 was compared using Student's *t*-test,

with the empty vector as the control ($N = 15$).

RESULTS

Overview of RNA-seq data sets

RNA was extracted from healthy control leaves, leaves infected with *C. fulvum* (*in planta*

Cf), and leaves with *C. fulvum* lesions overgrown by *H. pulvinata* (*in planta Cf/Hp*) (Fig. **1a**;

Fig. **S1a**). Because *H. pulvinata* parasitizes *C. fulvum* in leaf and on nutrient-poor water agar

(Iida *et al*., 2018; Maeda *et al*., 2025), fungal RNA was also extracted from *in vitro* cultures

of *H. pulvinata* alone (*in vitro Hp*), *C. fulvum* alone (*in vitro Cf*), and dual cultures of *C.*

fulvum and *H. pulvinata* (*in vitro Cf/Hp*) on water agar (Fig. **1a**; Fig. **S1a**). Typical white

mycelium of the mycoparasite *H. pulvinata* was observed at 14 days after inoculation for the

in planta Cf/Hp and *in vitro Cf/Hp* interactions (Fig. **1a**). Microscopic observation confirmed

that *H. pulvinata* penetrated *C. fulvum* hyphae (Fig. **1b**) and that *C. fulvum* entered tomato

leaves through stomata (Fig. **S1b**) as previously described (Maeda *et al*., 2025). Also, hyphae

of *H. pulvinata* elongated on the leaf surface and entered stomata in the presence of *C. fulvum*

(Fig. **S1b**). Raw reads were aligned to the reference genome sequences of *H. pulvinata*, *C.*

fulvum and tomato plants using HISAT2. The proportion of uniquely mapped reads varied

across samples, ranging from 10.3% and 88.1% (Table **S2**), reflecting the multiorganismal

nature of several samples. Principal component analysis (PCA) of normalized expression

values revealed clear clustering of biological replicates by treatment for each organism,

indicating high within-treatment reproducibility (Fig. **S1c**). In the transcriptome analysis, we

identified DEGs using adjusted *P*-value (padj) ≤ 0.05 and |$\log_2$ Fold-change| ≥ 1.0. Functional

enrichment of DEGs was performed using KEGG pathway annotation to reveal biological

processes that are positively and negatively regulated during mycoparasitism.

(a)

Healthy leaf

In vitro Hp
(only mycoparasite on agar)

In planta Cf
(only pathogen on leaf)

In vitro Cf
(only pathogen on agar)

In planta Cf/Hp
(mycoparasitism on leaf)

In vitro Cf/Hp
(mycoparasitism on agar)

(b)

In vitro Cf

In vitro Cf/Hp

Fig. 1. *Cladosporium fulvum parasitized by mycoparasitic fungus Hansfordia pulvinata.* (a) Typical healthy tomato leaf, lesions caused by *C. fulvum* (*in planta Cf*), and lesions covered by white mycelium of *H. pulvinata* (*in planta Cf/Hp*). Tomato plants had lesions by 2 weeks after inoculation with *C. fulvum*. The diseased leaves were then inoculated with *H. pulvinata*, and after 1 week, white mycelia were observed on lesions. *In vitro* cultures: *H. pulvinata* (*in vitro Hp*) and *C. fulvum* (*in vitro Cf*) were grown on nylon membranes on water agar after transfer from PDA. *H. pulvinata* parasitized *C. fulvum* colonies on water agar (*in vitro Cf/Hp*). (b) Scanning electron micrographs of hyphae of *C. fulvum*

and *H. pulvinata* in the *in vitro Cf* and *Cf/Hp* interactions. *H. pulvinata* hyphae have

penetrated *C. fulvum* hyphae. Bar indicates 10 μm.

Transcriptomic reprogramming in *H. pulvinata* after its parasitism of *C. fulvum*

Transcriptome analysis of RNA from the mycoparasitic interactions (*in vitro Cf/Hp* and *in*

planta Cf/Hp) and the control (*in vitro Hp*) was used to identify DEGs involved in

mycoparasitism based on gene annotations (Table **S3**). During the *in vitro Cf/Hp* interaction,

1,336 genes were significantly upregulated and 738 genes downregulated (Fig. **2a**), compared

with 1,793 upregulated and 2,086 downregulated during mycoparasitism on the plant (*in*

planta Cf/Hp). The DEGs in these two mycoparasitic interactions shared a core set of 891

upregulated and 557 downregulated genes and mapped to 21 upregulated and 25

downregulated KEGG pathways (Fig. **2b**), suggesting that *H. pulvinata* activated a conserved

transcriptional program during mycoparasitism, whether it occurs *in vitro* or *in planta*

environments.

The KEGG pathways that were enriched in *H. pulvinata* during mycoparasitism imply

both primary and secondary metabolism were reprogrammed. KEGG pathways involved in

secondary metabolism, including steroids, terpenoids, and polyketides, were strongly

upregulated. Genome mining prediction revealed 33 putative biosynthetic gene clusters for

secondary metabolites in *H. pulvinata*—12 non-ribosomal peptide synthetases (NRPSs), 10

type I polyketide synthases (T1PKSs), 7 terpenes, 3 fungal ribosomally synthesized and post-

translationally modified peptides (fungal-RiPPs), and an isocyanide—all of which underwent

a marked transcriptional change during mycoparasitism (Table **S4**). Deoxyphomenone

biosynthesis (*DPH*) genes were also upregulated, consistent with *H. pulvinata* producing the

antifungal sesquiterpenoid 13-deoxyphomenone during mycoparasitism (Table **S4**) (Tirilly *et*

al., 1983; Maeda *et al.*, 2025). Interestingly, multiple amino acid metabolism pathways were

consistently downregulated, despite *H. pulvinata* growing vegetatively during

mycoparasitism (Fig. **2b**). Given the central role of amino acids for protein synthesis and

vegetative growth, this KEGG pattern suggested that *H. pulvinata* may acquire amino acids

derived from *C. fulvum* or tomato plants to supplement its amino acid pool during

mycoparasitism.

Fig. 2. Differentially expressed genes (DEGs) and enriched KEGG pathways in

Hansfordia pulvinata **during its parasitism of** *Cladosporium fulvum*. (a) UpSet plot of

DEGs in *in vitro* and in tomato (*in vitro Cf/Hp* and *in planta Cf/Hp*; Fig. 1a). Connected dots

indicate shared DEGs; black single dots represent condition-specific DEGs. Vertical bars

show the number of up- and downregulated genes; colored horizontal bars indicate the total

number of DEGs detected in each interaction. (b) Bubble plot of KEGG pathway enrichment

analysis of the 891 and 557 shared DEGs shown in (a). Keys: color of symbols indicates

significance level of enrichment (–log adjusted *P*); size of symbol indicates the number of

enriched DEGs in each pathway; the symbol indicates the functional class of the enriched

DEGs.

Expression profile and functional analysis of small secreted proteins in *H. pulvinata*

A comparison of the genome structures of *H. pulvinata* with those of fungi including plant

pathogens and mycoparasite *Trichoderma atroviride* revealed that *H. pulvinata* had the

fewest genes encoding carbohydrate-active enzymes (CAZymes), but the most genes

containing the signal peptide sequence for secretion (Fig. **S3**; Table **S3**). To analyze the

secretome, we filtered 891 DEGs commonly upregulated in both mycoparasitic interactions

(*in vitro Cf/Hp* and *in planta Cf/Hp*) and identified 165 genes encoding small secreted

proteins (Fig. **3a**). Although most of them could not be annotated for function, 27 were

annotated as genes previously reported to be associated with pathogenicity in plant

pathogenic fungi, including glycoside hydrolase, T2 family ribonuclease, EPL1-like protein

(orthologous to Ecp45 effector protein in *C. fulvum*), hydrophobin, necrosis-inducing protein,

KP4-type killer toxin proteins, cutinase-like protein, superoxide dismutase protein,

endoglucanase, and metalloprotease (Table **S5**). We focused on a necrosis-inducing protein

(DIP_10004627), designated *HpNLP1*, which is an Nep1-like protein (NLP) homologous to

PsojNIP because *HpNLP1* showed particularly high –$\log_{10}$ *P*-adj, indicating a high level of

significance (Fig. **3b**; Fig. **S4**). NLPs represent a widespread family of virulence-associated

proteins in plant pathogens that trigger necrosis and generation of ROS in host plants (Seidl

& Van den Ackerveken, 2019). Expression of *HpNLP1* increased 34-fold during the

mycoparasitism *in vitro* and 14-fold *in planta* (Table **S5**). The NLP gene *HpNLP2*

(DIP_10008029) was also present in *H. pulvinata*, but its expression was unchanged during

mycoparasitism.

Fig. 3. A Nep1-like protein (NLP) secreted from the mycoparasite *Hansfordia pulvinata* induces a plant-mediated toxicity against spores of *Cladosporium fulvum*. (a) Workflow for identifying small secreted proteins synthesized in *H. pulvinata* during mycoparasitism. (b) MA plot for expression levels of 165 selected genes. Cutoff thresholds are log₂ Fold-change ($\pm$1.0) for *in vitro Cf/Hp* interaction and log Mean (0.3). (c) Phylogenetic tree based on NLP amino acid sequences using the maximum likelihood method with 1,000 bootstrap replicates. Protein accession numbers are shown in (d). Type I and Type II have two and four cysteine residues, respectively. Proteins analyzed in this study are in bold. (d) Multiple sequence alignment of NLP using MAFFT. Conserved residues are highlighted in orange. (e) Transient expression assays of NLPs in *Nicotiana benthamiana* via agroinfiltration. Chlorosis and necrosis were assessed 14 days after infiltration by trypan blue staining. Similar results were obtained in three independent experiments; representative results are in Supplementary Fig. S5c. (f) Percentage germination of conidia of *C. fulvum* after 24-h treatment with apoplastic fluid from *N. benthamiana* leaves that had been infiltrated with different NLPs or buffer. Asterisks indicate significant differences between treatments determined using Dunnett's or the Mann–Whitney *U* test.

Phylogenetic analysis placed HpNlp1 and HpNlp2 within the type II and fungal type I

clades (Seidl and van den Ackerveken, 2019), respectively, in NLPs (Fig. **3c**). HpNlp1

contained four cysteine residues predicted to form two disulfide bonds between positions 39–

65 and 82–87 (Fig. **S4a** and **b**). Multiple sequence alignment showed that both HpNlp1 and

HpNlp2 contained the conserved region I and the C-terminal heptapeptide GHRHDWE

motif; this conserved 24-amino-acid peptide (nlp24) is recognized as a microbe- or pathogen-

associated molecular pattern (MAMP/PAMP) (Oome *et al.*, 2014). The tryptophan (W) and

proline (P) residues essential for necrosis-inducing activity in the conserved region I (Zhou *et*

al., 2012) were conserved in HpNlp2 but replaced by tyrosine (Y) and glutamic acid (E) in

HpNlp1 (Fig. **3d**), which was the only NLP strongly expressed during mycoparasitism. In the

heptapeptide motif of HpNlp1, the second histidine (H) was replaced by asparagine (N) (Fig.

3d). Structural superposition of PsojNIP and HpNlp1 yielded a TM-score of 0.86 and a root

mean square deviation (RMSD) of 1.93 Å, demonstrating that despite mutations in the nlp24-

like regions of HpNlp1, the global protein structure was largely preserved, as indicated by the

similar these values compared to the PsojNIP (Fig. **S4c**).

Recombinant proteins PsojNIP, HpNlp1 and HpNlp2 were produced in *P. pastoris* and

used to infiltrate *Nicotiana benthamiana* leaves. Western blot analysis of HpNlp1

demonstrated an additional band larger than the predicted molecular weight, suggesting dimer

formation (Fig. **S5a** and **b**). As previously reported, PsojNIP induced necrosis and chlorosis

(Qutob *et al.*, 2002), whereas HpNlp1 and HpNlp2 did not trigger visible necrosis or ROS

accumulation (Fig. **3e**; Fig. **S5c** and **d**). The high expression of *HpNLP1* gene during the

bipartite *in vitro* *Cf/Hp* interaction, in the absence of the plant, suggested that HpNlp1 might

directly suppress *C. fulvum* growth. Contrary to this expectation, no such direct antifungal

activity was detected against *C. fulvum* spore germination and hyphal elongation (Fig. **S5e**).

Remarkably, however, fluid from the apoplast—where *C. fulvum* grows—collected from *N.*

benthamiana leaves that had been infiltrated with HpNlp1 strongly suppressed *C. fulvum*

spore germination, whereas the protein with a point mutant (HpNlp1^{N138H}) failed to elicit

antifungal activity (Fig. **3f**). Neither of the nlp24 peptides of HpNlp1 and HpNlp1^{N138H}

induced RLP23-mediated resistance (Fig. **S5f**). These findings together demonstrated that

HpNlp1 diverges from typical necrosis-inducing NLPs in plant pathogens, inducing the

accumulation of apoplastic antifungal substances that suppress *C. fulvum* spore germination

instead of promoting host cell death.

Transcriptomic responses of pathogen *C. fulvum* parasitized by *H. pulvinata*

Next, we investigated the DEGs of *C. fulvum* through KEGG pathway analysis after tomato

was infected by *C. fulvum* (*in planta Cf*), dual infection in tomato leaves (*in planta Cf/Hp*),

and dual culture on agar (*in vitro Cf/Hp*) (Fig. **1a**). DEGs in *C. fulvum* were identified relative

to the *in vitro Cf* control on water agar. A total of 1,902 genes were upregulated and 1,783

genes were downregulated when *C. fulvum* infected tomato leaves (*in planta Cf*) (Fig. **4a**). In

the two mycoparasitic interactions, 2,195 upregulated and 2,435 downregulated genes were

detected *in planta Cf/Hp*, and 1,479 upregulated and 845 downregulated were *in vitro Cf/Hp*.

We identified 363 upregulated and 158 downregulated genes shared between the two

mycoparasitic interactions (*in vitro Cf/Hp* and *in planta Cf/Hp*), indicating a conserved

transcriptional response to *H. pulvinata* attack. KEGG enrichment analysis assigned these

shared DEGs to seven upregulated and three downregulated pathways, revealing strong

activation of protein folding and processing pathways in *C. fulvum* upon mycoparasitism by

H. pulvinata (Fig. **4b**). Conversely, genes involved in amino acid metabolism were

consistently downregulated, as also observed for *H. pulvinata*, suggesting coordinated

suppression of primary metabilisms in this fungal–fungal interaction.

Fig. 4. Differentially expressed genes (DE⬛⬛⬛⬛⬛⬛⬛⬛⬛ KEGG pathways in

phytopathogen *Cladosporium fulvum*. (a) U⬛⬛⬛⬛⬛⬛⬛ DEGs identified in the *in*

planta Cf, in vitro Cf/Hp, and *in planta Cf/HP* interactions (see Fig. 1A). Connected dots indicate shared DEGs; Black single dots represent condition-specific DEGs. Vertical bars show the numbers of up- and downregulated genes; colored horizontal bars indicate the total number of DEGs detected in each interaction. (b) Bubble plot of KEGG pathway enrichment analysis based on the 363 and 158 shared DEGs shown in (a). Keys: color of symbols indicates significant level of enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in each pathway; the symbol indicates the functional class of the enriched DEGs.

In *C. fulvum*, protein folding and processing pathways were strongly activated when parasitized by *H. pulvinata* (Fig. **4a**). These responses are indicative of ER stress and accumulation of misfolded proteins, phenomena often triggered by exposure to antifungal compounds (Chaillot *et al*., 2015). Similar to the results in the mycoparasite *H. pulvinata*, the downregulation of secondary metabolism-related genes suggested that *C. fulvum* reallocated resources from the production of secondary metabolites to survive and mitigate stress. The expression profiles of DEGs detected in *C. fulvum* were further analyzed based on functional classification previously reported by Zaccaron & Stergiopoulos (2024). Among the 42 genes associated with secondary metabolite biosynthesis, 15 (35.7%) and 18 (42.8%) were downregulated in the *in planta Cf* and *in planta Cf/Hp* interactions, respectively (Fig. **5a**; Table **S6**). Thirteen genes, including *PKS6* (CLAFUR5_12905), a key enzyme in cladofulvin biosynthesis (Griffiths *et al*., 2018), were consistently downregulated in both interactions.

In contrast, the category small secreted proteins had the most upregulated DEGs, including numerous known or predicted effectors (Fig. **5a**). An expression heatmap of effector genes showed that, as expected, the majority of the 106 effector genes were highly expressed when *C. fulvum* infected leaves in the *in planta Cf* interaction (Fig. **5b**). Expression of effector genes are generally silent on synthetic media (Mesarich *et al*., 2014); however, surprisingly, 56 effector genes were upregulated in the *in vitro* coculture of *C. fulvum* and *H. pulvinata* (*in vitro Cf/Hp*) (Fig. **5b**). In tomato leaves, effector expression in *C. fulvum* remained high regardless of the presence of *H. pulvinata* (Fig. **S6**)*.* These results suggested that *C. fulvum* effectors may play a defensive role, acting as countermeasures against mycoparasitic attack by *H. pulvinata.* To investigate this possibility, we conducted an AMAPEC analysis, which predicts antimicrobial proteins based on 3D-structural signatures (Mesny *et al*., 2024). Of the 56 effector genes that were upregulated during the *in vitro Cf/Hp*

interaction, 34 were predicted to possess antimicrobial activity, suggesting that many

effectors may act as antifungal proteins (Fig. **5c**; Table **S7**). Among these, we focused on the

effector Ecp2, which is known to trigger a hypersensitive response (HR) in tomato cultivars

carrying the *Cf-ECP2* resistance gene.

Fig. 5. Effector genes in phytopathogen *Cladosporium fulvum* were high[...]ressed during mycoparasitism by *Hansfordia pulvinata*. (a) Functional classifica[...] DEGs identified for the *in planta Cf*, *in vitro Cf/Hp* and *in planta Cf/Hp* interactio[...]: upregulated, stable: grey, blue: downregulated. The number of genes assigned to each category is in parentheses. (b) Heatmap showing expression profiles of effector genes. The red to blue gradient represents up- and downregulated expression, respectively. Genes with an adjusted *P*-value ≥0.05 are indicated in gray as not detected (ND). Red arrowhead indicates *Ecp2*. (c) Antimicrobial activity prediction for 56 effector genes upregulated during the *in vitro Cf/Hp* interaction (determined using the AMAPEC program). Symbols denote protein location predicted by EffectorP-fungi 3.0.

Structural and functional similarity between *C. fulvum* Ecp2 effector and antifungal killer toxin 4 proteins

The Alphafold2-predicted structure of Ecp2 yielded high confidence scores, with a pLDDT of 91.3 and a pTM of 0.872, indicating reliable local and global structures (Fig. **6a**). The structural similarity search using the DALI server identified two killer toxin 4 (KP4)-like

proteins, ZtKP4 in *Zymoseptoria tritici* and UmVKP4 in *Ustilago maydis*, despite their low amino acid sequence similarity (2.9 and 36.9%, respectively) (Table **S8**). The comparative structural analysis showed that the overall arrangement of the α-helices and multiple β-sheet was conserved among these proteins (Fig. **6a** and **S7a**). The TM-score for Ecp2 relative to ZtKP4 was 0.81 and 0.59 relative to UmVKP4, supporting Ecp2 having the same structural fold as the KP4 protein family according to the SCOP and CATH classification criteria (Fig. **6b**).

Fig. 6. Sequence dissimilarity and structural similarity between *Cladosporium fulvum* effector Ecp2 and killer toxin KP4 proteins. (a) Predicted 3D structure of Ecp2 generated using AlphaFold2. The color spectrum from blue to red indicates the N- to C-terminal

positions of residues. (b) Structural comparison of Ecp2 with ZtKP4 and UmVKP4 proteins.

The superposition of the aligned proteins was evaluated using TM-align, and the TM-score

and RMSD values are shown. (c) Heatmap showing sequence similarity among Ecp2, ZtKP4,

and UmVKP4 proteins as identified by BLASTp analysis. (d) Conserved motifs of proteins

identified by MEME motif analysis. (e) Representative tomato leaf from 4-week-old plant

carrying *Cf-ECP2* resistance gene at 6 days after infiltration with purified Ecp2 protein

produced by *Escherichia coli*. White dotted outline: area infiltrated with 1 μM Ecp2; black

dotted outline: infiltrated with control protein purified from *E. coli* harboring an empty vector

(EV). Numbers in parentheses indicate the number of leaves with a hypersensitive response

(HR) relative to the number of total leaves. (f) *In vitro* antifungal activity of Ecp2 against *C.*

fulvum, *H. pulvinata*, *Alternaria* sp., *Colletotrichum orbiculare*, and *Saccharomyces*

cerevisiae. Spores were cultured on 5% potato dextrose agar with or without Ecp2. Relative

hyphal length was determined 4 days after treatment. Bars = 250 μm. Significant differences

between EV and Ecp2 treatments were assessed using Student's *t*-test.

To explore the evolutionary relationships among these KP4-like proteins, we searched

for homologs of Ecp2, ZtKP4, and UmVKP4. Nineteen Ecp2 homologs were identified

across three fungal classes, while only one ZtKP4 homolog was detected in *Zymoseptoria*

tritici (syn. *Mycosphaerella graminicola*) (Table **S9**). A total of 188 UmVKP4 homologs

were found in 10 classes. Based on sequence similarity, these proteins clustered into two

groups; Ecp2 and ZtKP4 were placed in the same clade (Fig. **6c**). Motif analysis revealed

distinct conserved amino acid motifs in each group, and UmVKP4 group lacked an Hce2

domain (Fig. **6d**). While Ecp2 and ZtKP4 shared two conserved disulfide bonds, UmVKP4

had five, suggesting differences in structural stability and folding despite their shared overall

architecture (Fig. **S7b**). These results indicated that Ecp2 and UmVKP4 protein groups likely

originated from two distinct evolutionary lineages.

Given that Ecp2 is structurally and phylogenetically close to ZtKP4, which has

antifungal activity (de Guillen *et al*., 2024), we investigated the antifungal potential of Ecp2.

Infiltration of *Cf-ECP2* tomato leaves with recombinant Ecp2 (Fig. **S7c**) triggered a specific

HR (Fig. **6e**). At the high concentrations used, the active Ecp2 protein significantly inhibited

hyphal growth in diverse fungi: *C. fulvum*, *H. pulvinata*, *Alternaria* sp. isolated from tomato

fruits, the plant pathogen *Colletotrichum orbiculare* and the yeast *Saccharomyces cerevisiae*

(Fig. **6f**). Together, these results demonstrated that Ecp2 has a dual role: inducing *Cf-ECP2-*

mediated immunity in plant while inhibiting fungal growth. *H. pulvinata* exhibited

marginally higher resistance than the other tested fungi and yeast to Ecp2.

Enrichment analysis of DEGs and KEGG pathways in tomato plants

We also analyzed differential gene expression in tomato leaves inoculated with *C. fulvum*

or inoculated with *C. fulvum* followed by *H. pulvinata*, each compared with uninoculated

control leaves (Fig. **1a**). Infection by *C. fulvum* alone led to 3,231 upregulated and 4,980

downregulated genes, whereas in the presence of *H. pulvinata* (*in planta Cf/Hp*), 3,465 genes

were upregulated and 5,679 downregulated (Fig. **7a**). Among these, 1,619 up- and 2,153

downregulated genes were uniquely responsive to *H. pulvinata*, mapping to 35 and 33

enriched KEGG pathways, respectively (Fig. **7b**). Pathways associated with plant–pathogen

interaction, phytoalexin biosynthesis (stilbenoid, diarylheptanoid and gingerol), biosynthesis

of unsaturated fatty acids, other secondary metabolites, phenylpropanoid biosynthesis and

plant hormone signal transduction were induced when *H. pulvinata* parasitized *C. fulvum* in

leaves (Fig. **7b**). Genes encoding calcium-dependent protein kinases, PR1, RIN4, MAP

kinase, WRKY22 and defensins were highly expressed (Fig. **S8a**; Table **S10**). These results

indicated that *H. pulvinata* triggers plant immunity during its mycoparasitism of the pathogen

in tomato leaves. In contrast, pathways associated with photosynthesis antenna proteins and

porphyrin metabolism, which is involved in chlorophyll biosynthesis, were upregulated

during infection by *C. fulvum* alone, but significantly downregulated when parasitized by *H.*

pulvinata (Fig. **7b**; Fig. **S8b** and **c**), indicating that *H. pulvinata* triggered pronounced plant

resistance responses during mycoparasitism.

Strikingly, 13 amino acid biosynthesis and metabolism pathways were strongly activated

in the *in planta Cf/Hp* condition (Fig. **7b** and **c**). Although amino acid metabolism was

broadly repressed in both *C. fulvum* and *H. pulvinata* during mycoparasitism (Fig. **2b** and

4b), it was upregulated in tomato. The pronounced activation of amino acid biosynthesis

pathways in tomato suggests that the plant supplied amino acids to the mycoparasite *H.*

pulvinata during the tripartite interaction.

Fig. 7. Differentially expressed genes (DEGs) and enriched KEGG pathways in tomato plants. (a) UpSet plot of DEGs in the *in planta Cf* and *in planta Cf/Hp* conditions (see Fig. 1a). Healthy leaves were used as a control. Connected dots indicate shared DEGs; black single dots represent condition-specific DEGs. Vertical bars show the number of up- and downregulated genes; colored horizontal bars indicate the total number of DEGs detected in each interaction. (b) Bubble plot of KEGG pathway enrichment analysis based on the 1,619 and 2,153 DEGs shown in (a). Keys: color of symbols indicates significance level of

enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in each

pathway; the symbol indicates the functional class of the enriched DEGs. Stars indicate

pathways highlighted in (c). (c) KEGG pathways involved in amino acid biosynthesis in

tomato.

DISCUSSION

In this study, we analyzed gene expression in tomato plants, the phytopathogenic fungus *C.*

fulvum, and mycoparasitic fungus *H. pulvinata* during the mycoparasitism of the pathogen

after it had infected its tomato host to capture their coordinated and distinct responses. We

thus discovered previously unrecognized aspects of molecular crosstalk and revealed how the

transcriptome of each organism was dynamically reprogrammed during the tripartite

interaction (Fig. **8**).

Fig. 8. Overview of tripartite interaction among tomato plants, phytopathogen

Cladosporium fulvum, **and mycoparasite** *Hansfordia pulvinata*. *C. fulvum* secretes effector

proteins that function as virulence factors in susceptible tomato cultivars and antifungal

proteins active against other fungi, but act as avirulence factors upon recognition by *Cf*-

resistant cultivars. *H. pulvinata* parasitizes *C. fulvum* and induces the production of apoplastic

antifungal substances in tomato plants via the Nep1-like protein HpNlp1. *H. pulvinata* also

activates the biosynthesis of the antifungal sesquiterpene 13-deoxyphomenone during

mycoparasitism. Other plant defense responses support this biocontrol effect. Pathways

related to amino acid biosynthesis in the three organisms suggest that nutrients from the plant

are supplied to *H. pulvinata*.

Evolutionary and functional insights into an atypical NLP in the biocontrol
659

mycoparasite *H. pulvinata*

The identification of 165 secreted small proteins upregulated during mycoparasitism

highlights the central role of the secretome of *H. pulvinata*. Among them, HpNlp1, an

atypical NLP, lacked both necrosis-inducing activity or RLP23 receptor-mediated response in

plants, unlike canonical NLPs such as PsojNIP (Qutob *et al*., 2006; Böhm *et al*., 2014; Oome

et al., 2014). Instead, HpNlp1 accumulated antifungal substances in the apoplast and

inhibited spore germination of *C. fulvum*. Although *H. pulvinata* cannot grow independently

on tomato leaves (Iida *et al*., 2018), it entered tomato plants through stomata only in the

presence of *C. fulvum* (Fig. **S1b**), indicating that the two fungi overlap in the apoplastic

niche. Furthermore, the functional divergence of the NLPs between the pathogen and

mycoparasite indicates the evolutionary plasticity of NLP family. Despite its structural

similarity to PsojNIP, HpNlp1 has distinct biological functions, suggesting that subtle amino

acid modifications can drive functional divergence within the NLP family. If *H. pulvinata*

induced a strong necrotic or immune response in tomato, the host fungus *C. fulvum* would

decline, limiting its own nutrient supply. Therefore, the high upregulation of a noncytotoxic

NLP such as HpNlp1 during mycoparasitism is evolutionarily reasonable. Notably, the

HpNLP1 gene was expressed in *H. pulvinata* when parasiting its host *in planta* and *in vitro*,

suggesting that it recognizes *C. fulvum* itself rather than plant-derived signals. Although

HpNlp1 did not directly inhibit *C. fulvum* growth, it indirectly promoted the accumulation of

antifungal metabolites, suggesting coevolution of *H. pulvinata* with not only its fungal host

C. fulvum but also tomato plants.

KEGG enrichment analysis indicated that *H. pulvinata* activated multiple secondary

metabolic pathways, particularly those related to terpenoids, polyketides, steroids and *DPH*

biosynthetic gene cluster, during mycoparasitism. These findings agree with previous

findings that *H. pulvinata* produces 13-deoxyformenone, an eremophilane-type

sesquiterpenoid, which is fungistatic to *C. fulvum* (Maeda *et al*., 2025). The production of

antifungal substances triggered by HpNlp1 in the tomato apoplast was correlated with the

induction of genes encoding defensins and phytoalexins, which were expressed only when *C.*

fulvum was parasitized by *H. pulvinata* in tomato leaves. Given that similar noncytotoxic

NLPs from the biocontrol oomycete *Pythium oligandrum* also induce plant defensins (Yang

et al., 2022), HpNlp1 may represent a conserved mechanism among biocontrol agents that

modulate plant immunity without causing cell death. *H. pulvinata* may employ a two-tiered

strategy to preserve the spores of its host fungus *C. fulvum*, combining a direct antifungal

sesquiterpenoid with an indirect, noncytotoxic protein, HpNlp1.

Ecp2 effector as a novel antifungal protein in the pathogen *C. fulvum*

KP4 family proteins, broadly conserved in phytopathogenic fungi, are antifungal. For

instance, *U. maydis* UmVKP4 suppresses the growth of susceptible strains of the same

species by inhibiting calcium-dependent pathways that regulate the cell cycle and

morphogenesis (Gu *et al.*, 1995; Gage *et al.*, 2002). *Z. tritici* ZtKP4 has fungicidal activity

even against itself (de Guillen *et al.*, 2024). Our structural modeling showed that *C. fulvum*

Ecp2 shared similarity with KP4 proteins and had broad-spectrum antifungal activity ranging

from filamentous fungi to yeast, including self-inhibition similar to that of ZtKP4. The β1-β2

loop in Ecp2 should functionally correspond to the β3–β4 loop of UmVKP4, which mediates

interactions with intracellular proteins (Gu *et al.*, 1995). However, Ecp2 lacked the key lysine

residue (Lys42) that UmVKP4 requires for full antifungal activity (Gage *et al.*, 2001),

suggesting a similar mechanism underlying antifungal action but involving an alternative

molecular interface. During the arms race between tomato and *C. fulvum*, a KP4-like

antifungal protein may have been diverted to Ecp2 effector. However, the structural basis

underlying *CfECP2*-mediated immune recognition and how Ecp2 acquired the dual ability to

interact with both fungal and plant targets remain unresolved.

Ecp2 was strongly expressed even during the *in vitro* mycoparasitism, suggesting a dual

role as both a virulence factor against plants and an antifungal protein in microbial

competition. A similar dual functionality is seen in the root endophyte *Serendipita* spp.,

where a chitinase encoding effector is induced during both plant colonization and microbial

interaction (Eichfeld *et al.*, 2024). The vascular wilt fungus *Verticillium dahliae* secretes

antimicrobial effectors such as VdAve1 and VdAMP2 to gain an ecological advantage in the

soil biome (Snelders *et al.*, 2020). The soil-borne pathogen *Rosellinia necatrix*, which causes

white root rot and is phylogenetically related to *V. dahliae*, secretes antimicrobial effector

proteins when it colonizes its host plant (Chavarro-Carrero *et al.*, 2024). While the

rhizosphere harbors highly diverse and competitive microbial communities, the diversity of

the microbial community in the endosphere such as the apoplastic space is generally

relatively low (Snelders *et al.*, 2022) because colonization of the apoplast is typically

restricted to specialized microbes that can secrete effectors to suppress host immunity

(Rocafort *et al*., 2020). Consequently, the potential role of antimicrobial effectors where

competition is relatively low is still unclear. In fact, despite previous predictions that some

apoplastic effectors produced by *C. fulvum* may have antimicrobial activity, no direct

evidence had been found (Mesarich *et al*., 2018). Our finding indicates the potential for *C.*

fulvum not only to utilize effectors for efficient colonization but also to mediate microbial

competition in the endosphere and phyllosphere. Although *H. pulvinata* was moderately

tolerant of Ecp2, the molecular basis for this tolerance during mycoparasitism remains

unclear.

The Hce2 (Homologs of *C. fulvum* Ecp2) superfamily is widely conserved across the

Ascomycota and *Basidiomycota*, suggesting that its origin predates the divergence of the

Dikarya lineage (Stergiopoulos *et al*., 2012). In contrast, *U. maydis* UmVKP4, which shares

structural similarity with Ecp2, lacks the Hce2 domain and shared sequence motifs, and

differs in the number of disulfide bonds (Fig. **S7b**). These findings suggest that UmVKP4

represents a case of convergent evolution, resulting in structural resemblance to Ecp2 and

ZtKP4. Molecular evolutionary analyses suggest that the amino acid substitution underlying

Hce2 function diversification was fixed early in evolution (Stergiopoulos *et al*., 2012).

Ancestral Hce2 proteins may initially act as antimicrobial factors to provide competitive

advantages in microbe-rich habitats and have been co-opted as an effector to inhibit plant

immunity. Mutations in effector genes that allow *C. fulvum* to evade *Cf* resistance in plants

may increase its susceptibility to antagonistic microbes such as *H. pulvinata*. This hypothesis

suggests that mycoparasitic biocontrol agents can circumvent the classical arms race between

plant and pathogen, providing a sustainable disease management.

Potential metabolic trade-offs in amino acid pathways between fungi and tomato plants

The physiological changes found in tomato plants highlight the dynamic interplay between

microbial interference and plant defense responses. The expression of genes in multiple

amino acid biosynthetic pathways were enhanced only when *H. pulvinata* parasitized *C.*

fulvum in tomato leaves. Amino acids are key mediators of plant–microbe interactions,

functioning as signaling molecules that activate defense and as nutritional sources that

pathogens exploit (Moormann *et al*., 2022; Tünnermann *et al*., 2022). Indeed, infection by *C.*

fulvum leads to the accumulation of free amino acids in the apoplast (Solomon & Oliver,

2001). Because the acquisition of plant-derived amino acids is energetically advantageous to

microbes, many plant pathogens modulate expression of genes encoding amino acid

transporters to optimize uptake (Sonawala *et al.*, 2018). Thus, the regulation of amino acid

assimilation and transport in plants can be a decisive factor shaping the outcome of plant–

pathogen interactions.

In contrast, amino acid biosynthetic pathways in *H. pulvinata* and *C. fulvum* were

downregulated during the mycoparasitism on the membrane and the plant, suggesting that

tomato-derived amino acids could be shared or sequentially utilized by these fungi (Fig. **8**).

Although amino acids are fundamental primary metabolites during fungal vegetative growth,

the metabolic profile of *H. pulvinata*, which had vigorous hyphal growth during

mycoparasitism, deviated from this conventional paradigm. Although *C. fulvum* may also

acquire amino acids from plants, these resources are likely reallocated to *H. pulvinata*, the

ultimate beneficiary. In this metabolic trade-off, tomato plants invest in amino acids that

benefit *H. pulvinata*, while receiving biocontrol protection in return (Fig. **8**). Similar

strategies have been reported for the biocontrol fungus *Pseudozyma flocculosa*, which

transiently suppresses photosynthesis to promote amino acid release (Laur *et al.*, 2018). The

mycoparasitic fungus *Trichoderma erinaceum* exploits environmental amino acids to enhance

the synthesis of antifungal secondary metabolites derived from amino acids (Guo *et al.*,

2020). Together, these results suggested that shifts in amino acid metabolism enable *H.*

pulvinata to harness pathogen-induced plant responses for its benefit.

Mycoparasitism by *H. pulvinata* modulates plant immunity and pathogen growth

In conclusion, the mycoparasite *H. pulvinata* actively reprogrammed gene expression and

physiological responses in the pathogen *C. fulvum* and its tomato host to establish a stable

tripartite interaction (Fig. **8**). *H. pulvinata* induced expression of the 13-deoxyphomenone

biosynthetic gene cluster in tomato, thereby directly suppressing *C. fulvum*. It also enhanced

plant resistance, leading to the accumulation of antimicrobial substances in the apoplast and

thus indirectly further suppressing *C. fulvum*. In turn, *C. fulvum* secreted effectors such as

Ecp2, which functioned to induce both fungal competition and plant immunity. *H. pulvinata*

is likely to tolerate or overcome these antifungal effectors, but the underlying mechanisms

remain unclear. The tomato plant reprogrammed amino acid metabolism and defense

response pathways, orchestrating distinct strategies against beneficial and pathogenic

microbes. Collectively, these findings underscore the dynamic interplay of plant–pathogen–

mycoparasite interactions, demonstrating that *H. pulvinata* simultaneously strengthens tomato

immunity and represses *C. fulvum* virulence via direct antagonism and modulation of plant

metabolic pathways.

ACKNOWLEDGEMENTS

This work was supported by a Grant-in-Aid for Scientific Research from JSPS (20H02993

and 24K08919) and the Research and Implementation Promotion Program through Open

Innovation Grants (JPJ011937) from the Project of the Bio-oriented Technology Research

Advancement Institution (BRAIN) (Y. Iida). We are grateful to K. Ikeda for providing fungal

strains, M.H.A.J. Joosten for *Cf-ECP2* tomato seeds, D. Takemoto for the *psojNIP* gene, F.

Takken for pSfinx, and Y. Kubo for the *C. orbiculare* strain. We thank Y. Kubo, M.Z. Fanani

and S. Kodama for valuable suggestions, C. Tanaka for laboratory management.

COMPETING INTERESTS

The authors declare that they have no commercial or financial relationships that could be

construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

YI designed the study and managed the research funds. KM, HS and TaS performed RNA-

seq and analyzed the data. KM, MK, MO, and TK analyzed NLPs. ToS conducted electron

microscopy and EI developed a microscopic observation protocol. KM and KS studied the

Ecp2 protein. KM and YI wrote the manuscript. All authors reviewed and approved the final

version.

DATA AVAILABILITY

Genomic sequences for *C. fulvum* strain Race5_Kim and *H. pulvinata* strain 414-3 are

available as accessions PRJNA565804 and PRJDB8178 at NCBI BioProject, respectively

(Sushida *et al*., 2019; Zaccaron & Stergiopoulos, 2024). Tomato genome data were obtained

from the Solanaceae Genomics Network (Hosmani *et al*., 2019). Additional genomes are

accessible via JGI MycoCosm (mycocosm.jgi.doe.gov) and the DDBJ/EMBL/GenBank

repository (www.ncbi.nlm.nih.gov). Custom scripts and supporting datasets were deposited at

Figshare (doi.org/10.6084/m9.figshare.30265009).

REFERENCES

Bailey TL, Johnson J, Grant CE, Noble WS. 2015. The MEME Suite. *Nucleic Acids*

Research. 2015 Jul 1;43(W1):W39-49. doi: 10.1093/nar/gkv416. Epub 2015 May 7.

PMID: 25953851; PMCID: PMC4489269.

Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, Schiegg M, Ales J,
Beier T, Rudy M, et al. 2019. ilastik: interactive machine learning for (bio)image
analysis. *Nature Methods*. 2019 Dec;16(12):1226-1232. doi: 10.1038/s41592-019-0582-
9. Epub 2019 Sep 30. PMID: 31570887.

Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, Fetter A, Terlouw
BR, Metcalf WW, Helfrich EJN, et al. 2023. antiSMASH 7.0: new and improved
predictions for detection, regulation, chemical structures and visualisation. *Nucleic Acids*
Research. 2023 Jul 5;51(W1):W46-W50. doi: 10.1093/nar/gkad344. PMID: 37140036;
PMCID: PMC10320115.

Böhm H, Albert I, Oome S, Raaymakers TM, Van den Ackerveken G, Nürnberger T.
2014. A conserved peptide pattern from a widespread microbial virulence factor triggers
pattern-induced immunity in *Arabidopsis*. *PLOS Pathogens*. 2014 Nov
6;10(11):e1004491. doi: 10.1371/journal.ppat.1004491. PMID: 25375108; PMCID:
PMC4223075.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics*. 2014 Aug 1;30(15):2114-20. doi:
10.1093/bioinformatics/btu170. Epub 2014 Apr 1. PMID: 24695404; PMCID:
PMC4103590.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. Aug
1;25(15):1972-3. doi: 10.1093/bioinformatics/btp348. Epub 2009 Jun 8. PMID:
19505945; PMCID: PMC2712344.

Chaillot J, Tebbji F, Remmal A, Boone C, Brown GW, Bellaoui M, Sellam A. 2015. The
Monoterpene Carvacrol Generates Endoplasmic Reticulum Stress in the Pathogenic
Fungus *Candida albicans*. *Antimicrobial Agents and Chemotherapy*. 2015
Aug;59(8):4584-92. doi: 10.1128/AAC.00551-15. Epub 2015 May 26. PMID: 26014932;
PMCID: PMC4505245.

Chavarro-Carrero EA, Snelders NC, Torres DE, Kraege A, López-Moral A, Petti GC,
Punt W, Wieneke J, García-Velasco R, López-Herrera CJ, et al. 2024. The soil-borne
white root rot pathogen *Rosellinia necatrix* expresses antimicrobial proteins during host
colonization. *PLOS Pathogens*. 2024 Jan 18;20(1):e1011866. doi:
10.1371/journal.ppat.1011866. PMID: 38236788; PMCID: PMC10796067.

Chen C, Wu Y, Li J, Wang X, Zeng Z, Xu J, Liu Y, Feng J, Chen H, He Y, et al. 2023.
TBtools-II: A "one for all, all for one" bioinformatics platform for biological big-data

mining. *Molecular Plant*. 2023 Nov 6;16(11):1733-1742. doi: 10.1016/j.molp.2023.09.010. Epub 2023 Sep 22. PMID: 37740491.

de Guillen K, Mammri L, Gracy J, Padilla A, Barthe P, Hoh F, Lahfa M, Rouffet J, Petit-Houdenot Y, Kroj T, et al. 2024. *Zymoseptoria tritici* effectors structurally related to killer proteins UmV-KP4 and UmV-KP6 are toxic to fungi, and define extended protein families in fungi. *bioRxiv*. 2024 Oct 14:2024.10.14.618152. doi:10.1101/2024.10.14.618152.

de la Rosa S, Schol CR, Ramos Peregrina Á, Winter DJ, Hilgers AM, Maeda K, Iida Y, Tarallo M, Jia R, Beenen HG, et al. 2024. Sequential breakdown of the *Cf-9* leaf mold resistance locus in tomato by *Fulvia fulva*. *New Phytologist*. 2024 Aug;243(4):1522-1538. doi: 10.1111/nph.19925. Epub 2024 Jun 24. PMID: 38922927.

de Jonge R, van Esse HP, Kombrink A, Shinya T, Desaki Y, Bours R, van der Krol S, Shibuya N, Joosten MH, Thomma BP. 2010. Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science*. 2010 Aug 20;329(5994):953-5. doi: 10.1126/science.1190859. PMID: 20724636.

de Wit PJGM, van der Burgt A, Ökmen B, Stergiopoulos I, Abd-Elsalam KA, Aerts AL, Bahkali AH, Beenen HG, Chettri P, Cox MP, et al. 2012. The genomes of the fungal plant pathogens *Cladosporium fulvum* and *Dothistroma septosporum* reveal adaptation to different hosts and lifestyles but also signatures of common ancestry. *PLOS Genetics*. 2012;8(11):e1003088. doi: 10.1371/journal.pgen.1003088. Epub 2012 Nov 29. Erratum in: PLoS Genet. 2015 Dec 22;11(12):e1005775. doi: 10.1371/journal.pgen.1005775. PMID: 23209441; PMCID: PMC3510045.

de Wit PJGM, Joosten MHAJ, Thomma BHPJ, Stergiopoulos I. 2009. Gene for gene models and beyond: the *Cladosporium fulvum*–tomato pathosystem. *Plant relationships*. Berlin, Heidelberg: Springer; 2009. p. 135–156. (The Mycota; vol 5). doi:10.1007/978-3-540-87407-2_7.

Eichfeld R, Mahdi LK, de Quattro C, Armbruster L, Endeshaw AB, Miyauchi S, Hellmann MJ, Cord-Landwehr S, Peterson D, Singan V, et al. 2024. Transcriptomics reveal a mechanism of niche defense: two beneficial root endophytes deploy an antimicrobial GH18-CBM5 chitinase to protect their hosts. *New Phytologist*. 2024 Nov;244(3):980-996. doi: 10.1111/nph.20080. Epub 2024 Sep 3. PMID: 39224928.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biology*. 2019 Nov 14;20(1):238. doi: 10.1186/s13059-019-1832-y. PMID: 31727128; PMCID: PMC6857279.

Gage MJ, Bruenn J, Fischer M, Sanders D, Smith TJ. 2001. KP4 fungal toxin inhibits

growth in *Ustilago maydis* by blocking calcium uptake. *Molecular Microbiology*. 2001

Aug;41(4):775-85. doi: 10.1046/j.1365-2958.2001.02554.x. PMID: 11532143.

Gage MJ, Rane SG, Hockerman GH, Smith TJ. 2002. The virally encoded fungal toxin

KP4 specifically blocks L-type voltage-gated calcium channels. *Molecular*

Pharmacology. 2002 Apr;61(4):936-44. doi: 10.1124/mol.61.4.936. PMID: 11901234.

Gene Ontology Consortium; Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin

HJ, Ebert D, Feuermann M, Gaudet P, Harris NL, et al. 2023. The Gene Ontology

knowledgebase in 2023. *Genetics*. 2023 May 4;224(1):iyad031. doi:

10.1093/genetics/iyad031. PMID: 36866529; PMCID: PMC10158837.

Gíslason MH, Nielsen H, Almagro Armenteros JJ, Johansen AR. 2021. Prediction of

GPI-anchored proteins with pointer neural networks. *Current Research in Biotechnology*.

2021;3:6-13. doi:10.1016/j.crbiot.2021.01.001.

Griffiths S, Mesarich CH, Overdijk EJR, Saccomanno B, de Wit PJGM, Collemare J.

2018. Down-regulation of cladofulvin biosynthesis is required for biotrophic growth of

Cladosporium fulvum on tomato. *Molecular Plant Pathology*. 2018 Feb;19(2):369-380.

doi: 10.1111/mpp.12527. Epub 2017 Feb 6. PMID: 27997759; PMCID: PMC6638085.

Gu F, Khimani A, Rane SG, Flurkey WH, Bozarth RF, Smith TJ. 1995. Structure and

function of a virally encoded fungal toxin from *Ustilago maydis*: a fungal and

mammalian Ca2+ channel inhibitor. *Structure*. 1995 Aug 15;3(8):805-14. doi:

10.1016/s0969-2126(01)00215-5. PMID: 7582897.

Guo YW, Gong BQ, Yuan J, Li HJ, Mahmud T, Huang Y, Li JF, Yang DP, Lan WJ.

2020. l-Phenylalanine Alters the Privileged Secondary Metabolite Production in the

Marine-Derived Fungus T*richoderma erinaceum* F1-1. *Journal of Natural Products*.

2020 Jan 24;83(1):79-87. doi: 10.1021/acs.jnatprod.9b00710. Epub 2019 Dec 30. PMID:

31886665.

Holm L. 2022. Dali server: structural unification of protein families. *Nucleic Acids Research*.

2022 Jul 5;50(W1):W210-W215. doi: 10.1093/nar/gkac387. PMID: 35610055; PMCID:

PMC9252788.

Hosmani PS, Flores-Gonzalez M, van de Geest H, Maumus F, Bakker LV, Schijlen E,

van Haarst J, Cordewener J, Sanchez-Perez G, Peters S, et al. 2019. An improved de

novo assembly and annotation of the tomato reference genome using single-molecule

sequencing, Hi-C proximity ligation and optical maps. *bioRxiv*. 2019;767764.

doi:10.1101/767764

Iida Y, Ikeda K, Sakai H, Nakagawa H, Nishi O, Higashi Y. 2018. Evaluation of the

potential biocontrol activity of *Dicyma pulvinata* against *Cladosporium fulvum*, the

causal agent of tomato leaf mold. *Plant Pathology*. 2018;67(9):1883–1890.

doi:10.1111/ppa.12916.

Iida Y, van 't Hof P, Beenen H, Mesarich C, Kubota M, Stergiopoulos I, Mehrabi R,

Notsu A, Fujiwara K, Bahkali A, et al. 2015. Novel Mutations Detected in Avirulence

Genes Overcoming Tomato *Cf* Resistance Genes in Isolates of a Japanese Population of

Cladosporium fulvum. PLoS One. 2015 Apr 22;10(4):e0123271. doi:

10.1371/journal.pone.0123271. PMID: 25902074; PMCID: PMC4406682.

Joosten MH. 2012. Isolation of apoplastic fluid from leaf tissue by the vacuum infiltration-

centrifugation technique. *Methods in Molecular Biology*. 2012;835:603-10. doi:

10.1007/978-1-61779-501-5_38. PMID: 22183681.

Kanehisa M, Furumichi M, Sato Y, Matsuura Y, Ishiguro-Watanabe M. 2025. KEGG:

biological systems database as a model of the real world. *Nucleic Acids Research*. 2025

Jan 6;53(D1):D672-D677. doi: 10.1093/nar/gkae909. PMID: 39417505; PMCID:

PMC11701520.

Katoh K., Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular biology and*

evolution. Apr;30(4):772-80. doi: 10.1093/molbev/mst010. Epub 2013 Jan 16. PMID:

23329690; PMCID: PMC3603318.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment

and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*. 2019

Aug;37(8):907-915. doi: 10.1038/s41587-019-0201-4. Epub 2019 Aug 2. PMID:

31375807; PMCID: PMC7605509.

Kourelis J, Malik S, Mattinson O, Krauter S, Kahlon PS, Paulus JK, van der Hoorn

RAL. 2020. Evolution of a guarded decoy protease and its receptor in solanaceous

plants. *Nature Communications*. 2020 Sep 2;11(1):4393. doi: 10.1038/s41467-020-

18069-5. Erratum in: *Nature Communications*. 2024 Mar 13;15(1):2278. doi:

10.1038/s41467-024-46611-2. PMID: 32879321; PMCID: PMC7468133.

Laur J, Ramakrishnan GB, Labbé C, Lefebvre F, Spanu PD, Bélanger RR. 2018.

Effectors involved in fungal-fungal interaction lead to a rare phenomenon of

hyperbiotrophy in the tritrophic system biocontrol agent-powdery mildew-plant. *New*

Phytologist. 2018 Jan;217(2):713-725. doi: 10.1111/nph.14851. Epub 2017 Oct 18.

PMID: 29044534; PMCID: PMC6079639.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

Durbin R; 1000 Genome Project Data Processing Subgroup. 2009. The Sequence

Alignment/Map format and SAMtools. *Bioinformatics*. 2009 Aug 15;25(16):2078-9. doi:

10.1093/bioinformatics/btp352. Epub 2009 Jun 8. PMID: 19505943; PMCID:

PMC2723002.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biology*. 2014;15(12):550. doi:

10.1186/s13059-014-0550-8. PMID: 25516281; PMCID: PMC4302049.

Luderer R, Takken FL, de Wit PJGM, Joosten MH. 2002. *Cladosporium fulvum*

overcomes *Cf-2*-mediated resistance by producing truncated AVR2 elicitor proteins.

Molecular *Microbiology*. 2002 Aug;45(3):875-84. doi: 10.1046/j.1365-

2958.2002.03060.x. PMID: 12139631.

Ma L, Lukasik E, Gawehns F, Takken FL. 2012. The use of agroinfiltration for transient

expression of plant resistance and fungal effector proteins in *Nicotiana benthamiana*

leaves. *Methods in Molecular Biology*. 2012;835:61-74. doi: 10.1007/978-1-61779-501-

5_4. PMID: 22183647.

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R,

Butcher S. 2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in

2024. *Nucleic Acids Research*. 2024 Jul 5;52(W1):W521-W525. doi:

10.1093/nar/gkae241. PMID: 38597606; PMCID: PMC11223882.

Maeda K, Sumita T, Nishi O, Sushida H, Higashi Y, Nakagawa H, Suzuki T, Iwao E,

Fanani MZ, Nishiya Y, Iida Y. 2025. Adaptive evolution of sesquiterpene

deoxyphomenone in mycoparasitism by *Hansfordia pulvinata* associated with horizontal

gene transfer from *Aspergillus* species. *mBio*. 2025 Apr 9;16(4):e0400724. doi:

10.1128/mbio.04007-24. Epub 2025 Mar 20. PMID: 40111082; PMCID: PMC11980549.

Mesarich CH, Griffiths SA, van der Burgt A, Okmen B, Beenen HG, Etalo DW, Joosten

MH, de Wit PJGM. 2014. Transcriptome sequencing uncovers the *Avr5* avirulence gene

of the tomato leaf mold pathogen *Cladosporium fulvum*. *Molecular Plant-Microbe*

Interactions. 2014 Aug;27(8):846-57. doi: 10.1094/MPMI-02-14-0050-R. PMID:

24678832.

Mesarich CH, Ökmen B, Rovenich H, Griffiths SA, Wang C, Karimi Jashni M,

Mihajlovski A, Collemare J, Hunziker L, Deng CH, et al. 2018. Specific

Hypersensitive Response-Associated Recognition of New Apoplastic Effectors from

Cladosporium fulvum in Wild Tomato. *Molecular Plant-Microbe Interactions*. 2018

Jan;31(1):145-162. doi: 10.1094/MPMI-05-17-0114-FI. Epub 2017 Nov 16. PMID:
29144204.

Mesarich CH, Barnes I, Bradley EL, de la Rosa S, de Wit PJGM, Guo Y, Griffiths SA,
Hamelin RC, Joosten MHAJ, Lu M, et al. 2023. Beyond the genomes of *Fulvia fulva*
(syn. *Cladosporium fulvum*) and *Dothistroma septosporum*: New insights into how these
fungal pathogens interact with their host plants. *Molecular Plant Pathology*. 2023
May;24(5):474-494. doi: 10.1111/mpp.13309. Epub 2023 Feb 15. PMID: 36790136;
PMCID: PMC10098069.

Mesny F, Wolf V, López-Moral A, Kraege A, Punt W, Park J, Zhu J, Sato Y, Thomma
BPHJ. 2024. Plant-associated fungi co-opt ancient antimicrobials for host manipulation.
bioRxiv. Preprint posted online 2024 Jan 4. doi:10.1101/2024.01.04.574150.

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022.
ColabFold: making protein folding accessible to all. *Nature Methods*. 2022
Jun;19(6):679-682. doi: 10.1038/s41592-022-01488-1. Epub 2022 May 30. PMID:
35637307; PMCID: PMC9184281.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL,
Tosatto SCE, Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families
database in 2021. *Nucleic Acids Research*. 2021 Jan 8;49(D1):D412-D419. doi:
10.1093/nar/gkaa913. PMID: 33125078; PMCID: PMC7779014.

Moormann J, Heinemann B, Hildebrandt TM. 2022. News about amino acid metabolism
in plant-microbe interactions. *Trends in Biochemical Sciences*. 2022 Oct;47(10):839-850.
doi: 10.1016/j.tibs.2022.07.001. Epub 2022 Aug 1. PMID: 35927139.

Oome S, Raaymakers TM, Cabral A, Samwel S, Böhm H, Albert I, Nürnberger T, Van
den Ackerveken G. 2014. Nep1-like proteins from three kingdoms of life act as a
microbe-associated molecular pattern in *Arabidopsis*. *Proceedings of the National*
Academy of Sciences, USA. 2014 Nov 25;111(47):16955-60. doi:
10.1073/pnas.1410031111. Epub 2014 Nov 3. PMID: 25368167; PMCID: PMC4250136.

Peresse M, Le Picard D. 1980. *Hansfordia pulvinata*, mycoparasite destructeur du
Cladosporium fulvum. *Mycopathologia* 1980. 71:23–30.

Qutob D, Kamoun S, Gijzen M. 2002. Expression of a *Phytophthora sojae* necrosis-
inducing protein occurs during transition from biotrophy to necrotrophy. *The Plant*
Journal. 2002 Nov;32(3):361-73. doi: 10.1046/j.1365-313x.2002.01439.x. PMID:
12410814.

1028 **Qutob D, Kemmerling B, Brunner F, Küfner I, Engelhardt S, Gust AA, Luberacki B,**

1029  **Seitz HU, Stahl D, Rauhut T,** et al. 2006. Phytotoxicity and innate immune responses

1030  induced by Nep1-like proteins. *The Plant Cell*. 2006 Dec;18(12):3721-44. doi:

1031  10.1105/tpc.106.044180. Epub 2006 Dec 28. PMID: 17194768; PMCID: PMC1785393.

1032 **R Core Team R.** 2024. *R: a language and environment for statistical computing*. Vienna,

1033  Austria: R Foundation for Statistical Computing.

1034 **Rocafort M, Fudal I, Mesarich CH.** 2020. Apoplastic effector proteins of plant-associated

1035  fungi and oomycetes. *Current Opinion in Plant Biology*. 2020 Aug;56:9-19. doi:

1036  10.1016/j.pbi.2020.02.004. Epub 2020 Apr 2. PMID: 32247857.

1037 **Rooney HC, Van't Klooster JW, van der Hoorn RA, Joosten MH, Jones JD, de Wit**

1038  **PJGM.** 2005. Cladosporium Avr2 inhibits tomato Rcr3 protease required for *Cf-2*-

1039  dependent disease resistance. *Science*. 2005 Jun 17;308(5729):1783-6. doi:

1040  10.1126/science.1111404. Epub 2005 Apr 21. Erratum in: Science. 2005 Oct

1041  7;310(5745):54. PMID: 15845874.

1042 **Seidl MF, Van den Ackerveken G.** 2019. Activity and Phylogenetics of the Broadly

1043  Occurring Family of Microbial Nep1-Like Proteins. *Annual Review of Phytopathology*.

1044  2019 Aug 25;57:367-386. doi: 10.1146/annurev-phyto-082718-100054. Epub 2019 Jul 5.

1045  PMID: 31283435.

1046 **Shabab M, Shindo T, Gu C, Kaschani F, Pansuriya T, Chintha R, Harzen A, Colby T,**

1047  **Kamoun S, van der Hoorn RA.** 2008. Fungal effector protein AVR2 targets

1048  diversifying defense-related cys proteases of tomato. *The Plant Cell*. 2008

1049  Apr;20(4):1169-83. doi: 10.1105/tpc.107.056325. Epub 2008 Apr 30. PMID: 18451324;

1050  PMCID: PMC2390736.

1051 **Shimodaira H.** 2002. An approximately unbiased test of phylogenetic tree selection. Syst

1052  Biol. Jun;51(3):492-508. doi: 10.1080/10635150290069913. PMID: 12079646.

1053 **Snelders NC, Rovenich H, Petti GC, Rocafort M, van den Berg GCM, Vorholt JA,**

1054  **Mesters JR, Seidl MF, Nijland R, Thomma BPHJ.** 2020. Microbiome manipulation

1055  by a soil-borne fungal plant pathogen using effector proteins. *Nature Plants*. 2020

1056  Nov;6(11):1365-1374. doi: 10.1038/s41477-020-00799-5. Epub 2020 Nov 2. PMID:

1057  33139860.

1058 **Snelders NC, Rovenich H, Thomma BPHJ.** 2022. Microbiota manipulation through the

1059  secretion of effector proteins is fundamental to the wealth of lifestyles in the fungal

1060  kingdom. *FEMS Microbiology Reviews*. 2022 Sep 2;46(5):fuac022. doi:

1061  10.1093/femsre/fuac022. PMID: 35604874; PMCID: PMC9438471.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*. May 1;30(9):1312-3. doi:
10.1093/bioinformatics/btu033. Epub 2014 Jan 21. PMID: 24451623; PMCID:
PMC3998144.

Solomon PS, Oliver RP. 2001. The nitrogen content of the tomato leaf apoplast increases
during infection by *Cladosporium fulvum. Planta*. 2001 Jun;213(2):241-9. doi:
10.1007/s004250000500. PMID: 11469589.

Sonawala U, Dinkeloo K, Danna CH, McDowell JM, Pilot G. 2018. Review: Functional
linkages between amino acid transporters and plant responses to pathogens. *Plant
Science*. 2018 Dec;277:79-88. doi: 10.1016/j.plantsci.2018.09.009. Epub 2018 Sep 19.
PMID: 30466603.

Sperschneider J, Dodds PN. 2022. EffectorP 3.0: Prediction of Apoplastic and Cytoplasmic
Effectors in Fungi and Oomycetes. *Molecular Plant-Microbe Interactions*. 2022
Feb;35(2):146-156. doi: 10.1094/MPMI-08-21-0201-R. Epub 2022 Feb 1. PMID:
34698534.

**Sperschneider J, Catanzariti AM, DeBoer K, Petre B, Gardiner DM, Singh KB, Dodds
PN, Taylor JM**. 2017. LOCALIZER: subcellular localization prediction of both plant
and effector proteins in the plant cell. *Scientific Reports*. 2017 Mar 16;7:44598. doi:
10.1038/srep44598. PMID: 28300209; PMCID: PMC5353544.

Stergiopoulos I, Kourmpetis YA, Slot JC, Bakker FT, de Wit PJGM, Rokas A. 2012. In
silico characterization and molecular evolutionary analysis of a novel superfamily of
fungal effector proteins. *Molecular Biology and Evolution*. 2012 Nov;29(11):3371-84.
doi: 10.1093/molbev/mss143. Epub 2012 May 23. PMID: 22628532.

Sushida H, Sumita T, Higashi Y, Iida Y. 2019. Draft Genome Sequence of *Dicyma
pulvinata* Strain 414-3, a Mycoparasite of *Cladosporium fulvum*, Causal Agent of
Tomato Leaf Mold. *Microbiology Resource Announcements*. 2019 Aug 29;8(35):e00655-
19. doi: 10.1128/MRA.00655-19. PMID: 31467096; PMCID: PMC6715866.

Tanaka S, Wada K, Katayama M, Marumo S. 1984. Isolation of Sporogen-AO1, a
sporogenic substance, from *Aspergillus oryzae*. A*gricultural and biological chemistry*.
1984;48(12):3189–3191. doi:10.1271/bbb1961.48.3189.

Tanaka S, Wada K, Marumo S, Hattori H. 1984. Structure of sporogen-ao 1, a sporogenic
substance of *Aspergillus oryzae. Tetrahedron Letters*. 1984;25(51):5907–5910.
doi:10.1016/S0040-4039(01)81717-2.

Tang D, Chen M, Huang X, Zhang G, Zeng L, Zhang G, Wu S, Wang Y. 2023. SRplot:

A free online platform for data visualization and graphing. *PLoS One*. 2023 Nov

9;18(11):e0294236. doi: 10.1371/journal.pone.0294236. PMID: 37943830; PMCID:

PMC10635526.

Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD,

Winther O, Brunak S, von Heijne G, Nielsen H. 2022. SignalP 6.0 predicts all five

types of signal peptides using protein language models. *Nature Biotechnology*. 2022

Jul;40(7):1023-1025. doi: 10.1038/s41587-021-01156-3. Epub 2022 Jan 3. PMID:

34980915; PMCID: PMC9287161.

Thomma BP, van Esse HP, Crous PW, de Wit PJGM. 2005. *Cladosporium fulvum* (syn.

Passalora fulva), a highly specialized plant pathogen as a model for functional studies on

plant pathogenic Mycosphaerellaceae. *Molecular Plant Pathology*. 2005 Jul 1;6(4):379-

93. doi: 10.1111/j.1364-3703.2005.00292.x. PMID: 20565665.

Tirilly Y, Kloosterman J, Sipma G, Kettenes-Van Den Bosch JJ. 1983. A fungitoxic

sesquiterpene from *Hansfordia pulvinata*. *Phytochemistry*. 1983;22(9):2082–2083.

doi:10.1016/0031-9422(83)80051-X.

Tünnermann L, Colou J, Näsholm T, Gratz R. 2022. To have or not to have: expression of

amino acid transporters during pathogen infection. *Plant Molecular Biology*. 2022

Jul;109(4-5):413-425. doi: 10.1007/s11103-022-01244-1. Epub 2022 Feb 1. PMID:

35103913; PMCID: PMC9213295.

Watanabe H, Horinouchi H, Muramoto Y, Ishii H. 2017. Occurrence of azoxystrobin-

resistant isolates in *Passalora fulva*, the pathogen of tomato leaf mold disease. *Plant*

Pathology. 2017;66(9):1472–1479. doi:10.1111/ppa.12701.

Wulff BB, Chakrabarti A, Jones DA. 2009. Recognitional specificity and evolution in the

tomato-*Cladosporium fulvum* pathosystem. *Molecular Plant-Microbe Interactions*. 2009

Oct;22(10):1191-202. doi: 10.1094/MPMI-22-10-1191. PMID: 19737093.

Yan L, Chen J, Zhang C, Ma Z. 2008. Molecular characterization of benzimidazole-

resistant isolates of *Cladosporium fulvum*. *Fems Microbiology Letters*. 2008

Jan;278(2):242-8. doi: 10.1111/j.1574-6968.2007.00999.x. PMID: 18096020.

Yang K, Chen C, Wang Y, Li J, Dong X, Cheng Y, Zhang H, Zhai Y, Ai G, Song Q, et

al. 2022. Nep1-Like Proteins From the Biocontrol Agent *Pythium oligandrum* Enhance

Plant Disease Resistance Independent of Cell Death and Reactive Oxygen Species.

Frontiers in Plant Science. 2022 Mar 4;13:830636. doi: 10.3389/fpls.2022.830636.

PMID: 35310640; PMCID: PMC8931738.

Zaccaron AZ, Stergiopoulos I. 2024. Analysis of five near-complete genome assemblies of

the tomato pathogen *Cladosporium fulvum* uncovers additional accessory chromosomes

and structural variations induced by transposable elements effecting the loss of

avirulence genes. *BMC Biology*. 2024 Jan 29;22(1):25. doi: 10.1186/s12915-024-01818-

z. PMID: 38281938; PMCID: PMC10823647.

Zaccaron AZ, Chen LH, Samaras A, Stergiopoulos I. 2022. A chromosome-scale genome

assembly of the tomato pathogen *Cladosporium fulvum* reveals a compartmentalized

genome architecture and the presence of a dispensable chromosome. *Microbial*

Genomics. 2022 Apr;8(4):000819. doi: 10.1099/mgen.0.000819. PMID: 35471194;

PMCID: PMC9453070.

Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018.

dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic*

Acids Research. 2018 Jul 2;46(W1):W95-W101. doi: 10.1093/nar/gky418. PMID:

29771380; PMCID: PMC6031026.

Zhang Y, Skolnick J. 2005. TM-align: a protein structure alignment algorithm based on the

TM-score. *Nucleic Acids Research*. 2005 Apr 22;33(7):2302-9. doi: 10.1093/nar/gki524.

PMID: 15849316; PMCID: PMC1084323.

Zhou BJ, Jia PS, Gao F, Guo HS. 2012. Molecular characterization and functional analysis

of a necrosis- and ethylene-inducing, protein-encoding gene family from *Verticillium*

dahliae. Molecular Plant-Microbe Interactions. 2012 Jul;25(7):964-75. doi:

10.1094/MPMI-12-11-0319. PMID: 22414440.

1150

FIGURE LEGENDS

Fig. 1. *Cladosporium fulvum* parasitized by mycoparasitic fungus *Hansfordia pulvinata.*

(a) Representative images of a healthy tomato leaf, lesions caused by *C. fulvum* (*in planta*

Cf), and lesions covered by white mycelium of *H. pulvinata* (*in planta Cf/Hp*). Tomato plants

were inoculated with *C. fulvum*, and lesions developed after 2 weeks. *H. pulvinata* was then

inoculated onto the diseased leaves, and white mycelia appeared on the lesions a week after

the second inoculation. *In vitro* cultures of *H. pulvinata* (*in vitro Hp*) and *C. fulvum* (*in vitro*

Cf) on nylon membranes on water agar after transfer from PDA. *H. pulvinata* parasitized *C.*

fulvum colonies on water agar (*in vitro Cf/Hp*). (b) Scanning electron micrographs of hyphae

of *C. fulvum* and *H. pulvinata* in *in vitro Cf* and *Cf/Hp* conditions. *H. pulvinata* hyphae have

penetrated *C. fulvum* hyphae in the *in vitro Cf/Hp* condition. Bar indicates 10 μm.

1162

Fig. 2. Differentially expressed genes (DEGs) and enriched KEGG pathways in

mycoparasite *Hansfordia pulvinata.* (a) UpSet plot of DEGs in the *in vitro* and *in planta*

mycoparasitic conditions (*in vitro Cf/Hp* and *in planta Cf/Hp*; Fig. 1a). Connected dots

indicate shared DEGs; Black single dots represent condition-specific DEGs. Vertical bars

show the number of up- and downregulated genes; colored horizontal bars indicate the total

number of DEGs detected in each interaction. (b) Bubble plot of KEGG pathway enrichment

analysis based on the 891 and 557 shared DEGs shown in (a). Keys: color of symbols

indicates significance level of enrichment (–log adjusted *P*); size of symbol indicates the

number of enriched DEGs in each pathway; the symbol indicates the functional class of the

enriched DEGs.

1173

Fig. 3. A Nep1-like protein (NLP) secreted from the mycoparasite *Hansfordia pulvinata*

induces a plant-mediated toxicity against spores of *Cladosporium fulvum.* (a) Workflow

for identifying small secreted proteins synthesized in *H. pulvinata* during mycoparasitism. (b)

MA plot for expression levels of 165 selected genes. Cutoff thresholds are $\log_2$ Fold-change

(±1.0) for *in vitro Cf/Hp* interaction and log Mean (0.3). (c) Phylogenetic tree constructed

based on NLP amino acid sequences using the maximum likelihood method with 1,000

bootstrap replicates. Type I and Type II have two and four cysteine residues, respectively.

Proteins analyzed in this study are in bold. (d) Multiple sequence alignment of NLP using

MAFFT. Conserved residues are highlighted in orange. (e) Transient expression assays of

NLPs in *Nicotiana benthamiana* via agroinfiltration. Chlorosis and necrosis were assessed 14

days after infiltration by trypan blue staining. Similar results were obtained in three

independent experiments; representative results are in Supplementary Fig. S5c. (f) Percentage

germination of conidia of *C. fulvum* after 24-h treatment with apoplastic fluid from *N.*

benthamiana leaves that had been infiltrated with different NLPs or buffer. Asterisks indicate

significant differences between treatments determined using Dunnett's or the Mann–Whitney

U test.

1190

Fig. 4. Differentially expressed genes (DEGs) and enriched KEGG pathways in the

pathogen *Cladosporium fulvum.* (a) UpSet plot showing DEGs identified in the *in planta Cf,*

in vitro Cf/Hp, and *in planta Cf/HP* conditions (see Fig. 1A). Connected dots indicate shared

DEGs; Black single dots represent condition-specific DEGs. Vertical bars show the numbers

of up- and downregulated genes; colored horizontal bars indicate the total number of DEGs

detected in each condition. (b) Bubble plot of KEGG pathway enrichment analysis based on

the 363 and 158 shared DEGs shown in (a). Keys: color of symbols indicates significant level

of enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in

each pathway; the symbol indicates the functional class of the enriched DEGs.

1200

Fig. 5. Effector genes in the pathogen *Cladosporium fulvum* were highly expressed

during mycoparasitism by *Hansfordia pulvinata*. (a) Functional classification of DEGs

identified for the *in planta Cf*, *in vitro Cf/Hp* and *in planta Cf/Hp* interactions. Red:

upregulated, stable: grey, blue: downregulated. The number of genes assigned to each

category is in parentheses. (b) Heatmap showing expression profiles of effector genes. The

red-to-blue gradient represents up- and downregulated expression, respectively. Genes with

an adjusted *P*-value ≥0.05 are indicated in gray as not detected (ND). Red arrowhead

indicates *Ecp2*. (c) Antimicrobial activity prediction for 56 effector genes upregulated during

the *in vitro Cf/Hp* interaction (determined using the AMAPEC program). Symbols denote

protein location predicted by EffectorP-fungi 3.0.

1211

Fig. 6. Sequence dissimilarity and structural similarity between *Cladosporium fulvum*

effector Ecp2 and killer toxin KP4 proteins. (a) Predicted 3D structure of Ecp2 generated

using AlphaFold2. The color spectrum from blue to red indicates the N- to C-terminal

positions of residues. (b) Structural comparison of Ecp2 with ZtKP4 and UmVKP4 proteins.

The superposition of the aligned proteins was evaluated using TM-align, and the TM-score

and RMSD values are shown. (c) Heatmap showing sequence similarity among Ecp2, ZtKP4,

and UmVKP4 proteins as identified by BLASTp analysis. (d) Conserved motifs of

the proteins identified by MEME motif analysis. (e) Representative tomato leaf from 4-week-

old tomato plant carrying the *Cf-ECP2* resistance gene at 6 days after infiltration with

purified Ecp2 protein produced by *Escherichia coli*. White dotted outline: area infiltrated

with infiltrated with 1 μM Ecp2; black dotted outline: infiltrated with control protein purified

from *E. coli* harboring an empty vector (EV). Numbers in parentheses indicate the number of

leaves with a hypersensitive response (HR) relative to the number of total leaves. (f) *In vitro*

antifungal activity of Ecp2 against *C. fulvum*, *H. pulvinata*, *Alternaria* sp., *Colletotrichum*

orbiculare, and *Saccharomyces cerevisiae*. Spores were cultured on 5% potato dextrose agar

with or without Ecp2. Relative hyphal length was determined 4 days after treatment. Bars =

250 μm. Significant differences between EV and Ecp2 treatments were assessed using

Student's t-test, and *P*-values are indicated.

1230

Fig. 7. Differentially expressed genes (DEGs) and enriched KEGG pathways in tomato

plants. (a) UpSet plot of DEGs in the *in planta Cf* and *in planta Cf/Hp* conditions (see Fig.

1a). Healthy leaves were used as a control. Connected dots indicate shared DEGs; Black

single dots represent condition-specific DEGs. Vertical bars show the number of up- and

downregulated genes; colored horizontal bars indicate the total number of DEGs detected in

each interaction. (b) Bubble plot of KEGG pathway enrichment analysis based on the 1,619

and 2,153 DEGs shown in (a). Keys: color of symbols indicates significance level of

enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in each

pathway; the symbol indicates the functional class of the enriched DEGs. Stars indicate

pathways highlighted in (c). (c) KEGG pathways involved in amino acid biosynthesis in

tomato.

1242

Fig. 8. Overview of the tripartite interaction among tomato plants, pathogen

***Cladosporium fulvum*, and mycoparasite *Hansfordia pulvinata*.** *C. fulvum* secretes effector

proteins that function as virulence factors in susceptible tomato cultivars and antifungal

proteins active against other fungi, but act as avirulence factors upon recognition by *Cf*-

resistant cultivars. *H. pulvinata* parasitizes *C. fulvum* and induces the production of apoplastic

antifungal substances in tomato plants via the Nep1-like protein HpNlp1. *H. pulvinata* also

activates the biosynthesis of the antifungal sesquiterpene 13-deoxyphomenone during

mycoparasitism. Other plant defense responses support this biocontrol effect. Pathways

related to amino acid biosynthesis in the three organisms suggest that nutrients from the plant

are supplied to *H. pulvinata*.

1253

SUPPORTING INFORMATION

Fig. S1 Experimental treatments and principal component analysis (PCA) of transcripts from

the mycoparasite *Hansfordia pulvinata* (Hp), the pathogen *Cladosporium fulvum* (Cf), and

tomato plant, *Solanum lycopersicum*.

Fig. S2 Secondary metabolite biosynthesis gene clusters in the mycoparasite *Hansfordia*

pulvinata.

Fig. S3 Phylogenetic analysis and functional characterization of proteins in fungal species.

Fig. S4 Structure of HpNlp1.

Fig. S5 Responses in plant leaves triggered by HpNlp1 protein.

Fig. S6 Heat map and hierarchical clustering of effector genes expressed in *Cladosporium*

fulvum.

Fig. S7 Structural comparative analysis of Ecp2 with ZtKP4 and UmVKP4.

Fig. S8 Differentially expressed genes (DEGs) in KEGG pathways detected in tomato plants.

Table S1 Primer sequences used in this study.

Table S2 Summary statistics of RNA-seq data and mapping results.

Table S3 Functional annotation of all genes in *Hansfordia pulvinata* genome.

Table S4 Predicted gene clusters involved in secondary metabolite biosynthesis in

Hansfordia pulvinata.

Table S5 Small secreted protein-coding genes ($\leq$300 aa) in *Hansfordia pulvinata*.

Table S6 Key genes encoding secondary metabolism enzymes in *Cladosporium fulvum*.

Table S7 Effector genes identified in *Cladosporium fulvum*.

Table S8 Structural homologs of Ecp2 protein identified from PDB25 using the Dali server.

Table S9 Species possessing homologs of Ecp2 and KP4 proteins.

Table S10 Log_2 fold-change for defensin-like genes in tomato plants.