Transcriptomic insights into the tritrophic plant–pathogen–mycoparasite interaction

reveal coordinated reprogramming fungal secretomes and plant amino acid

metabolism.

Kazuya Maeda[1], Mariko Kouda[1], Mai Ohara[1], Takumi Kawase[1], Koki Saito[1], Eishin Iwao[1],

Hirotoshi Sushida[2], Tomoko Suzuki[3], Takuya Sumita[4], Yuichiro Iida[1]

[1] Laboratory of Plant Pathology, Faculty of Agriculture, Setsunan University, Nagaotoge-cho

45-1, Hirakata, Osaka 573-0101, Japan

[2] Institute of Food Research, National Agriculture and Food Research Organization (NARO),

2-1-12 Kannondai, Tsukuba, Ibaraki 305-8642, Japan

[3] Department of Chemical and Biological Sciences, Faculty of Science, Japan Women's

University, Mejirodai 2-8-1, Bunkyo-ku, Tokyo 112-8681, Japan

[4] Department of biological resources management, School of Environmental Science, The

University of Shiga Prefecture, Hasaka-cho 2500, Hikone, Shiga 522-8533

* **Correspondence to Y. Iida (yuichiro.iida@setsunan.ac.jp)**

ORCIDs: Kazuya Maeda: 0000-0001-9858-0362; Mariko Kouda: 0009-0000-8862-4202;

Takumi Kawase: 0009-0006-2448-0805; Koki Saito: 0009-0007-5348-7739; Eishin Iwao:

0009-0000-6697-5378; Hirotoshi Sushida: 0000-0003-0631-6068; Tomoko Suzuki: 0000-

0001-9561-968X; Takuya Sumita: 0000-0001-7980-018X; Yuichiro Iida: 0000-0001-5062-

3255.

Short title: Transcriptomic analysis a plant–pathogen–mycoparasite system

Summary

• The tomato–*Cladosporium fulvum* (syn. *Fulvia fulva*) pathosystem has served as a model

for the gene-for-gene concept of effectors and resistance proteins, but this binary framework

does not include the potential influence of other microbial participants. Here we describe the

dramatic changes in gene expression of all members of the tritrophic interaction among

tomato, *C. fulvum*, and mycoparasitic fungus *Hansfordia pulvinata*.

• Transcriptomic analyses of the mycoparasite *H. pulvinata* during parasitism of *C. fulvum*

on tomato *on planta* and *in vitro* revealed a dramatic upregulation of genes encoding small

secreted proteins during mycoparasitism, notably, a Nep1-like protein (HpNlp1) lacked

typical necrosis-inducing activity but induced the accumulation of antifungal compounds

inhibiting spore germination of *C. fulvum.*

• Similarly, in *C. fulvum* parasitized by *H. pulvinata*, effector genes were highly expressed.

Strikingly, effector protein Ecp2 was found to share structural similarity with pathogen killer

toxin 4 proteins and had broad-spectrum antifungal activity, indicating a dual function in

fungal competition and *Cf-ECP2*-mediated plant resistance.

• In tomato plants infected by *C. fulvum* parasitized by *H. pulvinata*, primary metabolism

and defense-related genes were exclusively activated. These results suggest that in the

tritrophic interaction, the mycoparasite simultaneously suppressed the pathogen and induced

plant resistance. This study uncovers a multilayered molecular network in which the

mycoparasite coordinates pathogen suppression and plant defense within the tritrophic

interaction.

Keywords: mycoparasitism, Nep1-like protein, fungal effectors, Ecp2, antifungal proteins,

amino acid metabolism, *Cladosporium fulvum*, *Hansfordia pulvinata*

INTRODUCTION

The biotrophic filamentous fungus *Cladosporium fulvum* Cooke [syn. *Fulvia fulva* (Cooke) Cif.] causes tomato leaf mold, a serious economic threat due to reducing fruit yield and quality and occasionally killing tomato plants (*Solanum lycopersicum* L.) (Thomma *et al.*, 2005; de Wit *et al.*, 2009). The fungus enters tomato leaves through open stomata in high humidity, colonizes the apoplast, and secretes numerous effector proteins to suppress plant immunity and facilitate infection (Mesarich *et al.*, 2023). Commercial tomato cultivars have thus been bred with *Cf* resistance genes encoding receptor-like proteins (RLPs) with extracytoplasmic leucine-rich repeats from wild relatives (Mesarich *et al.*, 2023). These RLPs recognize the corresponding *C. fulvum* effector proteins as virulence factors via a gene-for-gene relationship and trigger a hypersensitive response to restrict the growth of *C. fulvum* (Mesarich *et al.*, 2023). However, *C. fulvum* rapidly evolves new races through mutation or loss of effector genes, thereby evading plant immunity (Iida *et al.*, 2015; de la Rosa *et al.*, 2024). The extensive use of resistant cultivars carrying only one or a few *Cf* genes has consequently led to the emergence of *C. fulvum* races that can overcome the resistance of the tomato plants (Luderer *et al.*, 2002; Mesarich *et al.*, 2014).

To date, over 100 effector genes have been identified in *C. fulvum* (Mesarich *et al.*, 2018; Zaccaron & Stergiopoulos, 2024). Most encode small cysteine-rich proteins that contain an N-terminal signal peptide for secretion. The identified avirulence (Avr) and extracellular proteins (Ecp) effectors—such as Avr2, Avr4, Avr4E, Avr5, Avr9B, Avr9C, Ecp1, Ecp2, Ecp4, Ecp5, and Ecp6—exemplify the complex coevolution between fungal virulence factors and their cognate *Cf* immune receptors (Wulff *et al.*, 2009; de la Rosa *et al.*, 2024). Several effectors have been functionally characterized. Avr2 acts as a cysteine protease inhibitor that targets the apoplastic cysteine proteases Rcr3 and Pip1, thereby interfering with basal defense mechanisms conserved across solanaceous plants (Rooney *et al.*, 2005; Shabab *et al.*, 2008; Kourelis *et al.*, 2020). Ecp6 suppresses chitin-triggered immunity by sequestering chitin oligomers released from the fungal cell wall, thus evading recognition by host pattern-recognition receptors (de Jonge *et al.*, 2010). In contrast, Ecp2 induces cell death in both tomato and nonhost plants, suggesting a dual role as a potential virulence and avirulence factor (de Kock *et al.*, 2004; de Wit *et al.*, 2012). Notably, the Ecp2 effector contains an Hce2 (homolog of *Cladosporium fulvum Ecp2*) domain that is widely conserved across diverse fungal and bacterial taxa, implying a fundamental role for this domain in microbe–plant interactions (Stergiopoulos *et al.*, 2012). However, the biochemical function of Ecp2 remains unclear. Of additional concern is that *C. fulvum* has developed resistance to fungicides (Yan

et al., 2008; Watanabe *et al*., 2017). The extensive breakdown of the resistance conferred by

single *Cf* resistance genes and the emergence of fungicide-resistant strains emphasize the

need for sustainable disease management strategies beyond conventional breeding and

chemical approaches. To better understand the molecular basis of disease

suppression, previous studies have primarily focused on the *C. fulvum*–tomato interaction

within the framework of the gene-for-gene model, but tritrophic interactions involving

additional microorganisms remain largely unexplored.

Hansfordia pulvinata (Berk. & M.A. Curtis) S. Hughes (syn. *Dicyma pulvinata*) is a

mycoparasitic fungus serendipitously discovered on tomato leaves infected with *C. fulvum* in

a greenhouse (Iida *et al*., 2018). It parasitizes *C. fulvum* and overgrows lesions, effectively

suppressing disease progression, and has broad mycoparasitic activity against all known *C.*

fulvum races (Iida *et al*., 2018). *H. pulvinata* recognizes *C. fulvum* hyphae, coils around them,

and directly penetrates the hyphae, then forms intercellular structures within the host fungus

(Peresse & Le Picard, 1980; Maeda *et al*., 2025). It also produces an eremophilane-type

sesquiterpenoid, 13-deoxyphomenone (also known as sporogen-AO1), which has antifungal

activity against *C. fulvum* (Tirilly *et al*., 1983; Maeda *et al*., 2025). The compound was

initially identified as a sporulation-inducing factor in the koji mold *Aspergillus oryzae*

(Tanaka *et al*., 1984a, 1984b). Comparative genomics recently revealed that the biosynthetic

gene cluster for 13-deoxyphomenone (*DPH*) in *H. pulvinata* shares high synteny with that in

Aspergillus species, suggesting horizontal gene transfer from an ancestral *Aspergillus* lineage

(Maeda *et al*., 2025). Subsequently, *H. pulvinata* appears to have retained this gene cluster to

support its mycoparasitic lifestyle, repurposing the endogeni

c sporogenesis function in *Aspergillus* species into exogenic antifungal activity that

inhibits spore germination and hyphal elongation of *C. fulvum*. However, the molecular

mechanisms underlying its mycoparasitism remain largely unknown. Since its

mycoparasitism is not restricted to the phyllosphere (Maeda *et al.*, 2025), comparative

analyses of fungal–fungal interactions under different mycoparasitic conditions could provide

new insights on the molecular basis of fungal host recognition and leaf-surface biocontrol

mechanisms.

The *C. fulvum*–tomato interaction has long served as a valuable model for dissecting

gene-for-gene dynamics in plant–pathogen systems (Mesarich et al. 2023.). Recent

chromosome-scale genome assemblies of multiple *C. fulvum* races have substantially

advanced our understanding of its genome structure and effector repertoire (Zaccaron *et al.*,

2022; Zaccaron & Stergiopoulos, 2024). Nevertheless, the molecular basis governing fungal–

fungal recognition and the tritrophic interactions with tomato plants remain largely

uncharacterized. Here, we present transcriptomic evidence revealing how the mycoparasitic

fungus *H. pulvinata* modulates both *C. fulvum* and tomato responses, providing molecular

insights into the interplay among plant, pathogen and mycoparasite. These findings

underscore the potential of the foliar mycoparasitic fungus as a promising biocontrol agent

within complex plant–microbe ecosystems.

MATERIALS AND METHODS

Culture conditions for fungal strains and plants, treatments and assays

Fungal strains *H. pulvinata* 414-3 and *C. fulvum* CF301 were cultured on half-strength potato

dextrose agar (PDA; BD Difco, Sparks, NJ, USA) or minimal medium (MM) agar (15 g

sucrose, 5 g ammonium tartrate, 1 g NH_4NO_3, 1 g KH_2PO_4, 0.5 g $MgSO_4·7H_2O$, 0.1 g NaCl,

0.1 g $CaCl_2·H_2O$, 25 μL 0.2 mg mL^{-1} biotin, 15 g agar and 1 mL trace elements per liter) at

25 °C in the dark for 1 (*H. pulvinata*) and 2 weeks (*C. fulvum*) (Maeda *et al.*, 2025). Spores

were collected in distilled water and adjusted to $1 × 10^6$ spores mL^{-1} using a hemocytometer.

For *in vitro* assays, 100 μL of each spore suspension was spread onto nylon membranes on

PDA and incubated for 1 (*H. pulvinata*) or 2 weeks (*C. fulvum*) (hereafter "*in vitro Hp*" and

"*in vitro Cf*" conditions; Fig. **S1a**). For establishing the mycoparasitism *in vitro*, membranes

bearing *C. fulvum* colonies were transferred to water agar (nutrient-poor) and sprayed with 1

mL of *H. pulvinata* spore suspension ($1 × 10^6$ spores mL^{-1}); plates were then incubated at

25 °C in the dark for 2 weeks (*in vitro Cf/Hp*; Fig. **S1a**).

Seedlings of tomato (*Solanum lycopersicum*) cv. Moneymaker (lacking all known *Cf*

resistance genes) were grown at 25 °C with 16 h light/8 h dark. Four-week-old plants were

either grown in high humidity as healthy controls for another 2 weeks (Healthy leaf; Fig.

S1a) or sprayed with approximately 3 mL of *C. fulvum* spore suspension (1×10^6 spores mL^{-1}),

then incubated in high humidity for 2 weeks until lesions developed ("*on planta Cf*"

interaction; Fig. **S1a**). For establishing the mycoparasitism on plants, plants with leaf mold

lesions were sprayed with approximately 3 mL of *H. pulvinata* spore suspension (1×10^6

conidia mL^{-1}) and incubated under high humidity at 25 °C for 2 weeks until lesions were

overgrown by white mycelia of *H. pulvinata* ("*on planta Cf/Hp*"; Fig. **S1a**).

N. benthamiana plants were grown for 4 weeks at 25 °C with a 16 h light/8 h dark.

Arabidopsis thaliana ecotype Col-0 seeds were surface-sterilized in 70% (v/v) ethanol for 1

min, rinsed three times with sterile distilled water, plated on Murashige and Skoog (MS) agar

(4.33 g MS basal medium (Sigma-Aldrich, St. Louis, MO, USA) and 20 g sucrose per liter,

pH 5.7), stratified at 4 °C in the dark for 1 month, then transferred to soil and grown for 5

weeks at 25 °C with 16 h light/8 h dark.

Scanning electron microscopy (SEM)

Details of the sample preparation and imaging procedures for SEM are described previously

(Maeda *et al*., 2025). Nylon membranes with fungal cultures were cut into about 5-mm

squares and fixed in 2.5% (v/v) glutaraldehyde in 100 mM cacodylate buffer (pH 7.4)

overnight at 4 °C, then 1% (w/v) osmium tetroxide in 100 mM cacodylate buffer (pH 7.4) for

1 h at room temperature. Samples were then dehydrated with a graded ethanol series, critical-

point dried in absolute ethanol using Leica EM CPD300 critical point dryer (Leica

Microsystems, Wetzlar, Germany), then mounted on stubs, coated with platinum-palladium,

and observed with a Hitachi SU-8220 scanning electron microscope (Tokyo, Japan).

RNAseq and transcriptome analysis

Three biological replicates were collected for each condition described above. Total RNA

was extracted using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to the

manufacturer's instructions. Poly(A) mRNA was enriched with the NEBNext Poly(A)

mRNA Magnetic Isolation Module (New England Biolabs, Ipswich, MA, USA), and libraries

were prepared using the NEBNext Ultra II Directional RNA Library Prep Kit (NEB).

Libraries were sequenced on an Illumina NovaSeq 6000 platform (Illumina, San Diego, CA,

USA) with the PE150 kit (150 bp × 2 paired-end), producing 2 or 4 Gb of data per sample.

Raw read quality was assessed using FastQC Ver. 16.0.1

(bioinformatics.babraham.ac.uk/projects/fastqc). Low-quality bases and adapter sequences

were removed using Trimmomatic Ver. 0.39 (Bolger *et al.*, 2014). Clean reads were aligned

to the respective reference genomes using HISAT2 Ver. 2.2.1; SAM files were converted to

BAM files with SamTools Ver. 1.18 (Li *et al.*, 2009; Kim *et al.*, 2019). Read counts were

generated using Rsubread Ver. 2.20.0. Differentially expressed genes (DEGs) were identified

with DESeq2 using thresholds of $|\log_2(\text{Fold-change})| \geq 1$ and adjusted *P*-value (padj) ≤ 0.05

(Love *et al.*, 2014). Gene Ontology (GO) and KEGG (Kyoto Encyclopedia of Genes and

Genomes) enrichment analyses were performed using TBtools-II Ver. 2.119 (Aleksander *et*

al., 2023; Chen *et al.*, 2023; Kanehisa *et al.*, 2025). Data were visualized using SRplot (Tang

et al., 2023).

Bioinformatic analysis

Orthologous coding sequences were identified using OrthoFinder Ver. 2.5.5 (Emms & Kelly,

2019); 100 single-copy orthologs were selected for multiple sequence alignment. Sequence

alignments were generated using MAFFT Ver. 7.526 and subsequently trimmed with TrimAl

Ver. 1.4 (Katoh and Standley, 2013; Capella-Gutiérrez *et al.*, 2009). Maximum-likelihood

phylogenetic trees were constructed with RAxML Ver. 8.2.12 using 1,000 bootstrap

replicates and the GTRCAT or PROTCATAUTO model (Stamatakis, 2014). The consistency

of tree topology was assessed using approximately unbiased (AU) tests (Shimodaira, 2002)

with 10,000 bootstrap replicates as previously described (Maeda *et al.*, 2025). Signal peptides

were predicted with SignalP-6.0 (Teufel *et al.*, 2022), candidate effectors with EffectorP-

fungi 3.0 (Sperschneider & Dodds, 2022), and subcellular localization with LOCALIZER

(Sperschneider *et al.*, 2017). Secondary metabolite biosynthetic gene clusters were

identified using the fungal version of antiSMASH (Blin *et al.*, 2023). GPI-anchored

proteins were predicted with NetGPI 1.1 (Gíslason *et al.*, 2021), and carbohydrate-active

enzymes (CAZymes) were annotated using DIAMOND (Zhang *et al.*, 2018). Default

parameters were used for all analyses unless otherwise specified.

For sequence similarity comparison, the EMBOSS Needle tool was used for pairwise

alignments (Madeira *et al.*, 2024). Protein domains and motifs were predicted using the

Pfam database and the MEME Suite, respectively (Bailey *et al.*, 2015; Mistry *et al.*, 2021).

The 3D structures of proteins were predicted using ColabFold Ver. 1.5.5 (Mirdita *et al.*,

2022). Five models were generated with three iterative recycles each, and the models were

ranked according to their predicted local distance difference test (pLDDT) scores. The top-

ranked models were refined by AMBER energy minimization to resolve steric clashes and

improve stereochemical geometry. The resulting relaxed structures were used for subsequent

structural analyses. The structure of Ecp2 was analyzed using the Dali server to retrieve

structural homologs based on a non-redundant dataset clustered at 25% sequence identity

(PDB25), and structures were compared with TM-align (Zhang & Skolnick, 2005; Holm,

2022).

Plasmid construction and protein expression

Necrosis-inducing protein 1-Like (NLP) protein coding sequences were synthesized

(VectorBuilder, IL, USA), cloned into the pSfinx vector using a primer set (Table **S1**) and

NEBuilder HiFi assembly kit (NEB). The NLP genes were digested with EcoRI and

subcloned into the pPIC9K vector (Thermo Fisher Scientific, Waltham, MA, USA) for

expression in *Pichia pastoris* GS115 strain according to the manufacturer's instructions.

Transformants were selected on YPD medium (Fujifilm Wako, Osaka, Japan) containing

G418 (Fujifilm Wako) and confirmed by PCR using the primer set in Table **S1**. Expression

was induced by adding 5 mL of the pre-culture grown in BMGY (buffered minimal medium

supplemented with 1% glycerol, 2% peptone, and 1% yeast extract, 1.34% amino acid–free

yeast nitrogen base, 100 mM potassium phosphate buffer [pH 6.0], and 4×10^{-5}% D-biotin)

broth to 100 mL of BMMY (1.5 % [w/v] methanol instead of glycerol in BMGY), and

incubating the culture at 30 °C and 300 rpm for 96 h. Supernatants were collected by

centrifugation at 12,000 rpm for 10 min.

The gene *Ecp2* was PCR-amplified from genomic DNA of *C. fulvum* using a primer set

(Table **S1**) and cloned into pET-28b vector (Merck, Darmstadt, Germany) to express an N-

terminal His$_6$-tagged protein in *Escherichia coli* strain BL21(DE3). Expression was induced

in LB medium with 1 mM IPTG at 16 °C for 16 h. Cells were resuspended with 50 mM KPi

buffer (pH 7.4) and disrupted by sonication on ice (10 min: 2 s on–4 s off). Insoluble

fractions were solubilized in 10 mL of denaturing buffer (6 M guanidine hydrochloride, 10

mM Tris, 10 mM β-mercaptoethanol, pH 8.0) for 1 h at room temperature. Proteins were

purified in a Ni-affinity column HisTrap HP (Cytiva, Medemblik, The Netherlands),

concentrated using Vivaspin 6 (GE Healthcare, IL, USA) and quantified with a Qubit

fluorometer (Thermo Fisher Scientific). BL21(DE3) carrying empty pET-28b served as a

negative control.

Western blot analysis

All protein samples were heated at 95 °C for 10 min in sample buffer (1% SDS, 1% 2-
mercaptoethanol, 10 mM Tris–HCl, 20% (v/v) glycerol, pH 6.8) for denaturation. The
proteins were separated in 12% SDS-polyacrylamide gel and transferred onto PVDF
membrane (Bio-Rad, Hercules, CA, USA). Membranes were blocked with 5% skim milk in
TBS-T for 1 h before antibody incubation. Anti-His and anti-HA monoclonal antibodies (Cell
Signaling Technology, Danvers, MA, USA) were diluted in Can Get Signal solution 1
(1:10,000) (TOYOBO, Osaka, Japan) and used as primary antibodies to detect Ecp2 and
NLPs, respectively. Horseradish peroxidase-conjugated anti-rabbit antibody was diluted with
Can Get Signal solution 2 (1:2,000) (TOYOBO) for use as the secondary antibody. Signals
were visualized using an ECL western detection kit (GE Healthcare, Chicago, IL, USA) and
captured using a ChemiDoc imaging system (Bio-Rad).
253
Functional assay of NLPs
pSfinx plasmids were introduced into *Agrobacterium tumefaciens* strain GV3101 via
electroporation (Ma *et al*., 2012). Agrobacterium suspensions were adjusted to $OD_{600} = 0.3$ in
infiltration buffer (2.03 g $MgCl_2·6H_2O$, 1.95 g MES, 500 µL of 200 mM acetosyringone per
liter, pH 5.6) and used to infiltrate leaves on 4-week-old plants of *N. benthamiana* by syringe
pressure. Leaves were monitored for necrosis 14 days after infiltration. Necrosis was
visualized by trypan blue staining (Ma *et al*., 2012) and quantified using ilastik (Ver. 1.4.1)
(Berg *et al*., 2019).
P. pastoris culture supernatant containing NLP proteins or BM medium (BMGY without
glycerol) was mixed at a 1:1 ratio with *C. fulvum* spore suspension (1×10^6 spores mL^{-1}). A
40 µL drop of each mixture was placed on a hydrophobic glass slide and incubated at room
temperature for 24 h. After incubation, germination of 200 spores per treatment was assessed
using a Nikon E600 light microscope (Tokyo, Japan). Culture supernatants containing NLP
proteins were collected from *P. pastoris* and used to syringe-infiltrate *N. benthamiana* leaves.
Twenty-four hours after infiltration, apoplastic fluid was collected by vacuum infiltration
with distilled water, followed by centrifugation (1,000 rpm, 10 min) (Joosten, 2012). Equal
volumes (10 µL) of *C. fulvum* spore suspension and apoplast fluid were then incubated at
room temperature for 24 h. A light microscope (Nikon) was then used to assess 200 spores
for germination.
Synthetic peptides of nlp24 (GenScript Biotech, Nanjing, China) were prepared in 500
nM stock solutions in dimethyl sulfoxide. Reactive oxygen species (ROS) were detected as

previously described with slight modifications (Yang *et al*., 2022). Leaf disks (0.125 cm²) of

N. benthamiana were floated on sterile distilled water for 24 h, then treated with 10 µL of a

substrate mixture (5 µL of 10 mM L-012 [Fujifilm Wako] and 5 µL of protein or peptide).

Luminescence was recorded every 2 min for 40 min using a Tristar3 Multimode Reader

(Berthold Technologies, Bad Wildbad, Germany).

Functional assays of Ecp2

The abaxial side of *Cf-ECP2* tomato leaves was infiltrated with purified Ecp2 protein (final

concentration 1 µM) using a syringe. Leaves were incubated at 25 °C with 16 h light/8 h dark

for 6 d and checked for a hypersensitive response (HR). *In vitro* antifungal activity was

assayed as previously described (Chavarro-Carrero *et al*., 2024). Briefly, fungal spores were

suspended in 5% PDB containing 15 µM Ecp2 or empty vector control (1×10^4 spores mL^{-1}),

then 100 µL of the samples (three wells per sample) was incubated in wells of 96-well plates

at 25 °C for 4 days. For each well, five fields of view were photographed ($N = 15$). Fungal

growth was quantified and compared with the control using the Hybrid Cell Count module of

the BZ-800X (KEYENCE, Osaka, Japan).

Statistical analyses

R version 4.4.2 (R Core Team, 2024) was used for all analyses. The proportion of necrosis

relative to total leaf area in *N. benthamiana* was assessed using Dunnett's test, with the empty

vector as the control ($N = 3$). Spore germination rate of *C. fulvum* treated with NLPs was

analyzed by one-way ANOVA followed by Tukey's HSD test ($N = 3$). The spore germination

rate of *C. fulvum* in apoplastic fluid from *N. benthamiana* leaves treated with NLPs was

compared using the Mann–Whitney *U* test, with the buffer as the control ($N = 12$). The

relative growth rate of fungi and yeast treated with Ecp2 was compared using Student's *t*-test,

with the empty vector as the control ($N = 15$).

RESULTS

Overview of RNA-seq datasets

RNA was extracted from healthy control leaves, leaves infected with *C. fulvum* (*on planta*

Cf), and leaves with *C. fulvum* lesions overgrown by *H. pulvinata* (*on planta Cf/Hp*) (Fig. **1a**;

Fig. **S1a**). Because *H. pulvinata* parasitizes *C. fulvum* on leaf surfaces and on nutrient-poor

water agar (Iida *et al*., 2018; Maeda *et al*., 2025), fungal RNA was also extracted from *in*

vitro cultures of *H. pulvinata* alone (*in vitro Hp*), *C. fulvum* alone (*in vitro Cf*), and dual

cultures of *C. fulvum* and *H. pulvinata* (*in vitro Cf/Hp*) on water agar (Fig. **1a**; Fig. **S1a**).

Typical mycoparasitic growth characterized by white mycelium of *H. pulvinata* was observed

at 14 days after inoculation for the *on planta Cf/Hp* and *in vitro Cf/Hp* conditions (Fig. **1a**).

Microscopic observation confirmed that *H. pulvinata* penetrated *C. fulvum* hyphae (Fig. **1b**),

and that *C. fulvum* entered tomato leaves through stomata (Fig. **S1b**) as previously described

(Maeda *et al*., 2025). Also, hyphae of *H. pulvinata* elongated on the leaf surface and entered

stomata in the presence of *C. fulvum* (Fig. **S1b**). Raw reads were aligned to the reference

genome sequences of *H. pulvinata*, *C. fulvum* and tomato plants using HISAT2. The

proportion of uniquely mapped reads varied across samples, ranging from 10.3 % and 88.1 %

(Table **S2**), reflecting the multi organismal nature of several samples. Principal component

analysis (PCA) of normalized expression values revealed clear clustering of biological

replicates by treatment for each organism, indicating high within-treatment reproducibility

(Fig. **S1c**). In the transcriptome analysis, we identified DEGs using adjusted *p*-value (padj) $\leq$

0.05 and |log$_2$ Fold-change| $\geq$ 1.0. Functional enrichment of DEGs was performed using

KEGG pathway annotation to reveal biological processes that are positively and negatively

regulated during mycoparasitism.

(a)

Healthy leaf

in vitro Hp
(only mycoparasite on agar)

on planta Cf
(only pathogen on leaf)

in vitro Cf
(only pathogen on agar)

on planta Cf/Hp
(mycoparasitism on leaf)

in vitro Cf/Hp
(mycoparasitism on agar)

(b)

in vitro Cf

in vitro Cf/Hp

Fig. 1. *Cladosporium fulvum* **parasitized by mycoparasitic fungus** *Hansfordia pulvinata.*
(a) Representative images of a healthy tomato leaf, lesions caused by *C. fulvum* (*on planta Cf*), and lesions covered by white mycelium of *H. pulvinata* (*on planta Cf/Hp*). Tomato plants were inoculated with *C. fulvum*, and lesions developed after 2 weeks. *H. pulvinata* was then inoculated onto the diseased leaves, and white mycelia appeared on the lesions a week later. *In vitro* cultures of *H. pulvinata* (*in vitro Hp*) and *C. fulvum* (*in vitro Cf*) on nylon membranes on water agar after transfer from PDA. *H. pulvinata* parasitized *C. fulvum* colonies on water agar (*in vitro Cf/Hp*). (b) Scanning electron micrographs of hyphae of *C. fulvum* grown *in vitro Cf* and *C. fulvum* parasitized by *H. pulvinata in vitro Cf/Hp*. *H. pulvinata* hyphae (*Hp*) have clearly penetrated *C. fulvum* hyphae (*Cf*). Bar indicates 10 μm.

Transcriptomic reprogramming in *H. pulvinata* during parasitisation of *C. fulvum*

Transcriptome analysis of *H. pulvinata* during parasitisation of *C. fulvum in vitro* (*in vitro Cf/Hp*) and *on planta* (*on planta Cf/Hp*) and the control (*in vitro Hp*) was used to identify DEGs involved in mycoparasitism based on gene annotation (Table **S3**). During parasitisation of *C. fulvum in vitro* (*in vitro Cf/Hp*), 1,336 genes were significantly upregulated and 738 genes were downregulated (Fig. **2a**), whereas 1,793 were upregulated and 2,086 were downregulated during parasitisation of *C. fulvum* grown on tomato leaves (*on planta Cf/Hp*). A core set of 891 upregulated DEGs was shared between the *in vitro Cf/Hp* and the *on planta Cf/Hp* conditions and a core set of 557 genes was downregulated and were mapped to 21 upregulated and 25 downregulated KEGG pathways (Fig. **2b**), suggesting in *H. pulvinata* a conserved transcriptional program was activated during parasitisation of *C. fulvum* both *in vitro* and *on planta*.

KEGG enrichment indicated reprogramming of primary and secondary metabolism in *H.*

pulvinata during parasitisation. KEGG pathways involved in secondary metabolism,

including steroids, terpenoids, and polyketides, were strongly upregulated. Genome mining

prediction revealed 33 putative secondary metabolite biosynthetic gene clusters in *H.*

pulvinata—12 non-ribosomal peptide synthetases (NRPSs), 10 type I polyketide synthases

(T1PKSs), 7 terpene synthetases, 3 fungal ribosomally synthesized and post-translationally

modified peptides (fungal-RiPPs), and an isocyanide—all of which underwent a marked

transcriptional change during parasitisation of *C. fulvum* (Table **S4**). Deoxyphomenone

biosynthesis (*DPH*) genes were also upregulated, consistent with *H. pulvinata* producing the

antifungal sesquiterpenoid 13-deoxyphomenone during parasitisation (Table **S4**) (Tirilly *et*

al., 1983; Maeda *et al.*, 2025). Interestingly, multiple amino acid metabolism pathways were

consistently downregulated, despite *H. pulvinata* growing vegetatively during parasitisation

(Fig. **2b**). Given the central role of amino acids for protein synthesis and vegetative growth,

this KEGG pattern suggested that *H. pulvinata* may acquire amino acids derived from *C.*

fulvum or tomato plants to supplement its amino acid pool during mycoparasitism.

Fig. 2. Differentially expressed genes   **G pathways in**

mycoparasite *Hansfordia pulvinata*   ***lvum in vitro* and *on***

***planta*.** (a) UpSet plot of DEGs in the   asitic conditions (*in*

vitro Cf/Hp and *on planta Cf/Hp*; Fig. 1a). Connected dots indicate shared DEGs; Black

single dots represent condition-specific DEGs. Vertical bars show the number of up- and

downregulated genes; colored horizontal bars indicate the total number of DEGs detected in

each interaction. (b) Bubble plot of KEGG pathway enrichment analysis based on the 891

and 557 shared DEGs shown in (a). Keys: color of symbols indicates significance level of

enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in each

pathway; the symbol indicates the functional class of the enriched DEGs.

Expression profile and functional analysis of small secreted proteins in *H. pulvinata*

A comparison of the genome structures of *H. pulvinata* with those of fungi and the

mycoparasite *Trichoderma atroviride* revealed that *H. pulvinata* had the smallest number of

genes encoding carbohydrate-active enzymes (CAZymes), and the highest number of genes

encoding secreted proteins (Fig. **S3**; Table **S3**). To analyze the secretome of *H. pulvinata*, we

filtered 891 DEGs commonly upregulated in both mycoparasitic interactions (*in vitro Cf/Hp*

and *on planta Cf/Hp*) and identified 165 genes encoding small secreted proteins (Fig. **3a**).

Although most of them could not be annotated for function, 27 were annotated as genes

previously reported to be associated with pathogenicity in plant pathogenic fungi, including

glycoside hydrolase, T2 family ribonuclease, EPL1-like protein (orthologous to Ecp45

effector protein in *C. fulvum*), hydrophobin, necrosis-inducing protein, KP4-type killer toxin

proteins, cutinase-like protein, superoxide dismutase protein, endoglucanase, and

metalloprotease (Table **S5**). We focused on a necrosis inducing protein (DIP_10004627),

designated *HpNLP1*, which is an Nep1-like protein (NLP) homologous to PsojNIP, because

HpNLP1 showed a particularly high $-\log_{10}$ padj, indicating a high level of significance (Fig.

3b; Fig. **S4**). NLPs represent a widespread family of virulence-associated proteins in plant

pathogens that trigger necrosis and generation of ROS in host plants (Seidl & Van den

Ackerveken, 2019). Expression of *HpNLP1* increased 34-fold and 14-fold during

parasitisation of *C. fulvum in vitro* and *on planta*, respectively (Table **S5**). The NLP gene

HpNLP2 (DIP_10008029) is also present in *H. pulvinata*, but its expression was unchanged

during parasitisation of *C. fulvum*.

408

409 **Fig. 3. A Nep1-like protein (NLP) secreted from the mycoparasite *Hansfordia pulvinata***

410 **induces a plant-mediated toxicity against spores of *Cladosporium fulvum*.** (a) Workflow

411 for identifying small secreted proteins synthesized in *H. pulvinata* during mycoparasitism. (b)

412 MA plot for expression levels of 165 selected genes. Cutoff thresholds are $\log_2$ Fold-change

413 (± 1.0) for *in vitro Cf/Hp* interaction and log Mean (0.3). (c) Phylogenetic tree constructed

414 based on NLP amino acid sequences using the maximum likelihood method with 1,000

415 bootstrap replicates. Type I and Type II have two and four cysteine residues, respectively.

416 Proteins analyzed in this study are in bold. (d) Multiple sequence alignment of NLP using

417 MAFFT. Conserved residues are highlighted in orange. (e) Transient expression assays of

418 NLPs in *Nicotiana benthamiana* via agroinfiltration. Chlorosis and necrosis were assessed 14

419 days after infiltration by trypan blue staining. Similar results were obtained in three

420 independent experiments; representative results are in Supplementary Fig. S5c. (f) Percentage

421 germination of conidia of *C. fulvum* after 24-h treatment with apoplastic fluid from *N.*

422 *benthamiana* leaves that had been infiltrated with different NLPs or buffer. Asterisks indicate

423 significant differences between treatments determined using Dunnett's or the Mann–Whitney

424 *U* test.

425

426   Phylogenetic analysis placed HpNlp1 and HpNlp2 within the type II and fungal type I

427 clades, respectively, in NLPs (Fig. **3c**). HpNlp1 contained four cysteine residues predicted to

428 form two disulfide bonds between positions 39–65 and 82–87 (Fig. **S4a** and **b**). Multiple

429 sequence alignment showed that both HpNlp1 and HpNlp2 contained the conserved region I

and the C-terminal heptapeptide GHRHDWE motif; this conserved 24-amino-acid peptide (nlp24) is recognized as a microbe- or pathogen-associated molecular pattern (MAMP/PAMP) (Oome *et al.*, 2014). The tryptophan (W) and proline (P) residues essential for necrosis-inducing activity in the conserved region I (Zhou *et al.*, 2012) were conserved in HpNlp2 but replaced by tyrosine (Y) and glutamic acid (E) in HpNlp1 (Fig. **3d**), which was the only NLP strongly expressed during mycoparasitism. In the heptapeptide motif of HpNlp1, the second histidine (H) was replaced by asparagine (N) (Fig. **3d**). Structural superposition of PsojNIP and HpNlp1 yielded a TM-score of 0.86 and a root mean square deviation (RMSD) of 1.93 Å, demonstrating that despite mutations in the nlp24-like regions of HpNlp1, the global protein structure was largely preserved, as indicated by these similar values compared to the PsojNIP (Fig. **S4c**).

Recombinant proteins PsojNIP, HpNlp1 and HpNlp2 were produced in *P. pastoris* and used to infiltrate *Nicotiana benthamiana* leaves. Western blot analysis of HpNlp1 demonstrated an additional band larger than the predicted molecular weight, suggesting dimer formation (Fig. **S5a** and **b**). As previously reported, PsojNIP induced necrosis and chlorosis (Qutob *et al.*, 2002), whereas HpNlp1 and HpNlp2 did not trigger visible necrosis or ROS accumulation (Fig. **3e**; Fig. **S5c** and **d**). The high expression of *HpNLP1* gene under *in vitro Cf/Hp* condition, in the absence of the plant, suggested that HpNlp1 might directly suppress *C. fulvum* growth. Contrary to this expectation, no such direct antifungal activity was detected against *C. fulvum* spore germination and hyphal elongation (Fig. **S5e**). Remarkably, however, apoplastic fluid—where *C. fulvum* develops—collected from *N. benthamiana* leaves that had been infiltrated with HpNlp1 strongly suppressed *C. fulvum* spore germination, whereas the protein with a point mutantation (HpNlp1^{N138H}) failed to elicit antifungal activity (Fig. **3f**). Neither of the nlp24 peptides of HpNlp1 and HpNlp1^{N138H} induced RLP23-mediated resistance (Fig. **S5f**). These findings together demonstrated that HpNlp1 diverges from typical necrosis-inducing NLPs in plant pathogens, inducing the accumulation of apoplastic antifungal substances that suppress *C. fulvum* spore germination instead of promoting host cell death.

Transcriptomic responses of pathogen *C. fulvum* parasitized by *H. pulvinata*

Next, we investigated the DEGs of *C. fulvum* through KEGG pathway analysis after tomato was infected by *C. fulvum* (*on planta Cf*), dual infection on tomato leaves (*on planta Cf/Hp*), and dual culture on agar medium (*in vitro Cf/Hp*) (Fig. **1a**). DEGs in *C. fulvum* were identified during parasitisation relative to the *in vitro Cf* control, in which the fungus was

cultured on water agar. A total of 1,902 genes were upregulated and 1,783 genes were

downregulated when *C. fulvum* infected tomato leaves (*on planta Cf*) (Fig. **4a**). Under the

two mycoparasitic conditions, 2,195 upregulated and 2,435 downregulated genes were

detected *on planta Cf/Hp*, and 1,479 upregulated and 845 downregulated were detected *in*

vitro Cf/Hp. We identified 363 upregulated and 158 downregulated genes shared between the

two mycoparasitic conditions (*in vitro Cf/Hp* and *on planta Cf/Hp*), indicating a conserved

transcriptional response to *H. pulvinata* attack. KEGG enrichment analysis assigned these

shared DEGs to seven upregulated and three downregulated pathways, revealing strong

activation of protein folding and processing pathways in *C. fulvum* during parasitisation by *H.*

pulvinata (Fig. **4b**). Conversely, genes involved in amino acid metabolism were consistently

downregulated, as was also observed in *H. pulvinata*, suggesting coordinated suppression of

primary metabolism in this fungus-fungus interaction.

Fig. 4. Differentially expressed genes (DEG ~~ed~~ **KEGG pathways in the**

pathogen *Cladosporium fulvum*. (a) UpSet p ~~~~ EGs identified in the *on planta*

Cf, in vitro Cf/Hp, and *on planta Cf/HP* conditions (see Fig. 1A). Connected dots indicate

shared DEGs; Black single dots represent condition-specific DEGs. Vertical bars show the

numbers of up- and downregulated genes; colored horizontal bars indicate the total number of

DEGs detected in each condition. (b) Bubble plot of KEGG pathway enrichment analysis

based on the 363 and 158 shared DEGs shown in (a). Keys: color of symbols indicates

significant level of enrichment (–log adjusted *P*); size of symbol indicates the number of

enriched DEGs in each pathway; the different symbols represent functional classes of the enriched DEGs.

In *C. fulvum*, protein folding and processing pathways were strongly activated when parasitized by *H. pulvinata* (Fig. **4a**). These responses are indicative of endoplasmic reticulum (ER) stress and accumulation of misfolded proteins, phenomena often triggered by exposure to antifungal compounds (Chaillot *et al*., 2015). Similar to the results in the mycoparasite *H. pulvinata*, the downregulation of secondary metabolism-related genes suggested that *C. fulvum* reallocated resources from the production of secondary metabolites to survive and mitigate stress. The expression profiles of DEGs detected in *C. fulvum* were further analyzed based on functional classification as previously reported by Zaccaron & Stergiopoulos (2024). Among the 42 genes associated with secondary metabolite biosynthesis, 15 (35.7 %) and 18 (42.8 %) were downregulated in the *on planta Cf* and *on planta Cf/Hp* interaction, respectively (Fig. **5a**; Table **S6**). Thirteen genes, including *PKS6* (CLAFUR5_12905), a key enzyme in cladofulvin biosynthesis (Griffiths *et al*., 2018), were consistently downregulated in both interactions.

In contrast, the category of small secreted proteins had the highest number of upregulated DEGs, including numerous known or predicted effectors (Fig. **5a**). An expression heatmap of effector genes showed that, as expected, the majority of the 106 effector genes were highly expressed during infection of tomato by *C. fulvum* (*on planta Cf*) (Fig. **5b**). Expression of effector genes are generally low on rich synthetic media (Mesarich *et al*., 2014); however surprisingly, 56 effector genes were upregulated in the *in vitro* coculture of *C. fulvum* and *H. pulvinata* (*in vitro Cf/Hp*) (Fig. **5b**). On tomato leaves, effector expression in *C. fulvum* remained high regardless of the presence of *H. pulvinata* (Fig. **S6**)*.* These results suggested that *C. fulvum* effectors may play a defensive role, acting as countermeasures against mycoparasitic attack by *H. pulvinata* or are upregulated due to nitrogen limitation caused by parasitisation by *H. pulvinata.* To investigate this possibility, we conducted an AMAPEC analysis, which predicts antimicrobial proteins based on 3D-structural signatures (Mesny *et al*., 2024). Of the 56 effector genes that were upregulated under the *in vitro Cf/Hp* condition, 34 were predicted to possess antimicrobial activity, suggesting that many effectors may act as antifungal proteins (Fig. **5c**; Table **S7**). Among these, we focused on the effector Ecp2, which is known to trigger a hypersensitive response (HR) in tomato cultivars carrying the *Cf-ECP2* resistance gene.

Fig. 5. Effector genes in the pathogen *Cladosporium fulvum* were high expressed

during parasitisation by *Hansfordia pulvinata*. (a) Functional classifica of DEGs

identified for the *on planta Cf*, *in vitro Cf/Hp* and *on planta Cf/Hp* interactions. Red:

upregulated, stable: grey, blue: downregulated. The number of genes assigned to each

category is in parentheses. (b) Heatmap showing expression profiles of effector genes. The

red-to-blue gradient represents up- and downregulated expression, respectively. Genes with

an adjusted *P*-value ≥0.05 are indicated in gray as not detected (ND). Red arrowhead

indicates *Ecp2*. (c) Antimicrobial activity prediction for 56 effector genes upregulated during

the *in vitro Cf/Hp* interaction (determined using the AMAPEC program). Symbols denote

protein location predicted by EffectorP-fungi 3.0.

Structural and functional similarity between *C. fulvum* Ecp2 effector and antifungal

killer toxin 4 proteins

The Alphafold2-predicted structure of Ecp2 yielded high confidence scores, with a pLDDT

of 91.3 and a pTM of 0.872, indicating reliable local and global structures (Fig. **6a**).

Structural similarity search using the DALI server identified two killer toxin 4 (KP4)-like

proteins, ZtKP4 in *Zymoseptoria tritici* and UmVKP4 in *Ustilago maydis*, despite their low

amino acid sequence similarity (2.9 to 36.9%) (Table **S8**). Comparative structural analysis

showed that the overall arrangement of the α-helices and multiple β-sheets were conserved

among these proteins (Fig. **6a** and **S7a**). The TM-scores for Ecp2 relative to ZtKP4 and

UmVKP4 were 0.81 and 0.59, respectively, supporting that Ecp2 had the same structural fold

as the KP4 protein family according to the SCOP and CATH classification criteria (Fig. **6b**).

Fig. 6. Sequence dissimilarity and structural similarity between *Cladosporium fulvum*

effector Ecp2 and killer toxin KP4 proteins. (a) Predicted 3D structure of Ecp2 generated

using AlphaFold2. The color spectrum from blue to red indicates the N- to C-terminal

positions of residues. (b) Structural comparison of Ecp2 with ZtKP4 and UmVKP4 proteins.

The superposition of the aligned proteins was evaluated using TM-align, and the TM-score

and RMSD values are shown. (c) Heatmap showing sequence similarity among Ecp2, ZtKP4,

and UmVKP4 proteins as identified by BLASTp analysis. (d) Conserved motifs of

the proteins identified by MEME motif analysis. (e) Representative tomato leaf from 4-week-

old tomato plant carrying the *Cf-ECP2* resistance gene at 6 days after infiltration with

purified Ecp2 protein produced by *Escherichia coli*. White dotted outline: area infiltrated

with infiltrated with 1 μM Ecp2; black dotted outline: infiltrated with control protein purified

from *E. coli* harboring an empty vector (EV). Numbers in parentheses indicate the number of

leaves with a hypersensitive response (HR) relative to the number of total leaves. (f) *In vitro*

antifungal activity of Ecp2 against *C. fulvum*, *H. pulvinata*, *Alternaria* sp., *Colletotrichum orbiculare*, and *Saccharomyces cerevisiae*. Spores were cultured on 5% potato dextrose agar with or without Ecp2. Relative hyphal length was determined 4 days after treatment. Bars = 250 μm. Significant differences between EV and Ecp2 treatments were assessed using Student's t-test, and *p*-values are indicated.

To explore the evolutionary relationships among these KP4-like proteins, we searched for homologs of Ecp2, ZtKP4, and UmVKP4. Nineteen Ecp2 homologs were identified across three fungal classes, while only one ZtKP4 homolog was detected in *Zymoseptoria tritici* (syn. *Mycosphaerella graminicola*) (Table **S9**). A total of 188 UmVKP4 homologs were found in 10 classes. Based on sequence similarity, these proteins clustered into two groups; Ecp2 and ZtKP4 were placed in the same clade (Fig. **6c**). Motif analysis revealed distinct conserved amino acid motifs in each group, and UmVKP4 group lacked an Hce2 domain (Fig. **6d**). While Ecp2 and ZtKP4 shared two conserved disulfide bonds, UmVKP4 had five, suggesting differences in structural stability and folding despite their shared overall architecture (Fig. **S7b**). These results indicated that Ecp2 and UmVKP4 protein groups likely originated from two distinct evolutionary lineages.

Given that Ecp2 is structurally and phylogenetically close to ZtKP4, which has antifungal activity (de Guillen *et al*., 2024), we investigated the antifungal potential of Ecp2. Infiltration of *Cf-ECP2* tomato leaves with recombinant Ecp2 (Fig. **S7c**) triggered a specific HR (Fig. **6e**). At the high concentrations used, the active Ecp2 protein significantly inhibited hyphal growth in diverse fungi: *C. fulvum*, *H. pulvinata*, *Alternaria* sp. isolated from tomato fruits, the plant pathogen *Colletotrichum orbiculare* and the yeast *Saccharomyces cerevisiae* (Fig. **6f**). Together, these results demonstrated that Ecp2 has a dual role: inducing *Cf-ECP2*-mediated immunity in plant while inhibiting fungal growth. *H. pulvinata* exhibited marginally higher resistance than the other tested fungi and yeast to Ecp2.

Enrichment analysis of DEGs and KEGG pathways in tomato plants

We also analyzed differential gene expression in tomato leaves inoculated with *C. fulvum* and inoculated with *C. fulvum* followed by *H. pulvinata*, each compared with uninoculated control leaves (Fig. **1a**). Infection by *C. fulvum* alone led to 3,231 upregulated and 4,980 downregulated genes, whereas in the presence of *H. pulvinata* (*on planta Cf/Hp*), 3,465 genes were upregulated and 5,679 downregulated (Fig. **7a**). Among these, 1,619 up- and 2,153 downregulated genes were uniquely responsive to *H. pulvinata*, mapping to 35 and 33

enriched KEGG pathways, respectively (Fig. **7b**). Pathways associated with plant–pathogen

interaction, phytoalexin biosynthesis (stilbenoid, diarylheptanoid and gingerol), biosynthesis

of unsaturated fatty acids, other secondary metabolites, phenylpropanoid biosynthesis and

plant hormone signal transduction were induced when *H. pulvinata* parasitized *C. fulvum* on

leaves (Fig. **7b**). Genes encoding calcium-dependent protein kinases, PR1, RIN4, MAP

kinase, WRKY22 and defensins were highly expressed (Fig. **S8a**; Table **S10**). These results

indicated that *H. pulvinata* triggers plant immunity during parasitisation of *C. fulvum* on

tomato leaves. In contrast, pathways associated with photosynthesis antenna proteins and

porphyrin metabolism, which is involved in chlorophyll biosynthesis, were upregulated

during infection by *C. fulvum* alone, but significantly downregulated when parasitized by *H.*

pulvinata (Fig. **7b**; Fig. **S8b** and **c**), indicating that *H. pulvinata* triggered pronounced plant

resistance responses during parasitisation of *C. fulvum*.

Fig. 7. Differentially expressed genes (DEGs) and enriched KEGG pathways in tomato plants. (a) UpSet plot of DEGs in the *on planta Cf* and *on planta Cf/Hp* conditions (see Fig. 1a). Healthy leaves were used as a control. Connected dots indicate shared DEGs; Black single dots represent condition-specific DEGs. Vertical bars show the number of up- and downregulated genes; colored horizontal bars indicate the total number of DEGs detected in each interaction. (b) Bubble plot of KEGG pathway enrichment analysis based on the 1,619 and 2,153 DEGs shown in (a). Keys: color of symbols indicates significance level of enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in each pathway; the symbol indicates the functional class of the enriched DEGs. Stars indicate

pathways highlighted in (c). (c) KEGG pathways involved in amino acid biosynthesis in
tomato.

Strikingly, 13 amino acid biosynthesis and metabolism pathways were strongly activated
in the *on planta Cf/Hp* condition (Fig. **7b** and **c**). Although amino acid metabolism was
broadly repressed in both *C. fulvum* and *H. pulvinata* during mycoparasitism (Fig. **2b**; Fig.
4b), it was upregulated in tomato (Fig. **7b** and **c**). The pronounced activation of amino acid
biosynthesis pathways in tomato suggests that the plant supplies amino acids to both the
pathogen *C. fulvum* and its mycoparasite *H. pulvinata* during the tritrophic interaction.
619

DISCUSSION
In this study, we analyzed gene expression in tomato plants, the phytopathogenic fungus *C.*
fulvum, and its mycoparasite *H. pulvinata* alone or during their bitrophic and tritrophic
interaction to capture coordinated and distinct responses. We thus discovered previously
unrecognized aspects of molecular crosstalk and revealed how the transcriptome of each
organism was dynamically reprogrammed during these interactions.
626

Evolutionary and functional insights into an atypical NLP in the biocontrol
mycoparasite *H. pulvinata*
The identification of 165 secreted small proteins upregulated during mycoparasitism
highlights the central role of the secretome of *H. pulvinata*. Among them, HpNlp1, an
atypical NLP, lacked both necrosis-inducing activity or RLP23 receptor-mediated response in
plants, unlike canonical NLPs such as PsojNIP (Qutob *et al.*, 2006; Böhm *et al.*, 2014; Oome
et al., 2014). Instead, HpNlp1 accumulated antifungal substances in the apoplast and
inhibited spore germination of *C. fulvum*. Although *H. pulvinata* cannot grow independently
on tomato leaves (Iida *et al.*, 2018), it follows the hyphae of *C. fulvum* that enter the stomata
of tomato leaves (Fig. **S1b**), indicating that the two fungi co-occur in the tomato apoplast.
Furthermore, the functional divergence of the NLPs between the pathogen and mycoparasite
indicates the evolutionary plasticity of NLP family. Despite its structural similarity to
PsojNIP, HpNlp1 has distinct biological functions, suggesting that subtle amino acid
modifications can drive functional divergence within the NLP family. If *H. pulvinata* induced
a strong necrotic or immune response in tomato, the host fungus *C. fulvum* would decline,
limiting its own nutrient supply. Therefore, the high upregulation of a noncytotoxic NLP such
as HpNlp1 during mycoparasitism is evolutionarily reasonable. Notably, the *HpNLP1* gene
was expressed in *H. pulvinata* when parasiting its host *on planta* and *in vitro*, suggesting that

it recognizes *C. fulvum* itself rather than plant-derived signals. Although HpNlp1 did not directly inhibit *C. fulvum* growth, it indirectly promoted the accumulation of antifungal metabolites, suggesting coevolution of *H. pulvinata* with not only its fungal host *C. fulvum* but also tomato plants.

KEGG enrichment analysis indicated that *H. pulvinata* activated multiple secondary metabolic pathways, particularly those related to terpenoids, polyketides, steroids and *DPH* biosynthetic gene cluster, during mycoparasitism. These findings agree with previous findings that *H. pulvinata* produces 13-deoxyformenone, an eremophilane-type sesquiterpenoid, which is fungistatic to *C. fulvum* (Maeda *et al*., 2025). The production of antifungal substances triggered by HpNlp1 in the tomato apoplast was correlated with the induction of genes encoding defensins and phytoalexins, which were expressed only when *C. fulvum* was parasitized by *H. pulvinata* on tomato leaves. Given that similar noncytotoxic NLPs from the biocontrol oomycete *Pythium oligandrum* also induce plant defensins (Yang *et al*., 2022), HpNlp1 may represent a conserved mechanism among biocontrol agents that modulate plant immunity without causing cell death. *H. pulvinata* may employ a two-tiered strategy to preserve the spores of its host fungus *C. fulvum*, combining a direct antifungal sesquiterpenoid with an indirect, noncytotoxic protein, HpNlp1.

Ecp2 effector as a novel antifungal protein in the pathogen *C. fulvum*

KP4 family proteins, broadly conserved in phytopathogenic fungi, are antifungal but do not interact with the plant. For instance, *U. maydis* UmVKP4 suppresses the growth of susceptible strains of the same species by inhibiting calcium-dependent pathways that regulate the cell cycle and morphogenesis (Gu *et al*., 1995; Gage *et al*., 2002). *Z. tritici* ZtKP4 has fungicidal activity even against itself (de Guillen *et al*., 2024). Our structural modeling showed that *C. fulvum* Ecp2 shared similarity with KP4 proteins and had broad-spectrum antifungal activity ranging from filamentous fungi to yeast, including self-inhibition similar to that of ZtKP4. The β1–β2 loop in Ecp2 should functionally correspond to the β3–β4 loop of UmVKP4, which mediates interactions with intracellular proteins (Gu *et al*., 1995). However, Ecp2 lacked the key lysine residue (Lys42) that UmVKP4 requires for full antifungal activity (Gage *et al*., 2001), suggesting a similar mechanism underlying antifungal action but involving an alternative molecular interface. During the arms race between tomato and *C. fulvum*, a KP4-like antifungal protein may have been diverted to Ecp2 effector. However, the structural basis underlying *CfECP2*-mediated immune recognition and how

Ecp2 acquired the dual ability to interact with both fungal and plant targets remain

unresolved.

Ecp2 was strongly expressed even during *in vitro* mycoparasitism, suggesting a dual role

as both a virulence factor against plants and an antifungal protein in microbial competition.

Similar dual functionality is seen in the root endophyte *Serendipita* spp., where a chitinase

encoding effector is induced during both plant colonization and microbial interaction

(Eichfeld *et al.*, 2024). The vascular wilt fungus *Verticillium dahliae* secretes antimicrobial

effectors such as VdAve1 and VdAMP2 to gain an ecological advantage in the soil biome

(Snelders *et al.*, 2020). The soil-borne white root rot pathogen *Rosellinia necatrix*, which is

phylogenetically related to *V. dahliae*, secretes antimicrobial effector proteins during host

plant colonization (Chavarro-Carrero *et al.*, 2024). While the rhizosphere harbors highly

diverse and competitive microbial communities, the diversity of the microbial community in

the endosphere such as the apoplastic space is generally relatively low (Snelders *et al.*, 2022)

because colonization of the apoplast is typically restricted to specialized microbes that can

secrete effectors to suppress host immunity (Rocafort *et al.*, 2020). In fact, despite previous

predictions that some apoplastic effectors produced by *C. fulvum* may have antimicrobial

activity, no direct evidence had been found so far (Mesarich *et al.*, 2018). Our finding

indicates the potential for *C. fulvum* not only to utilize effectors for efficient colonization of

tomato leaves but also to compete with/inhibit other microbes in the endosphere and

phyllosphere. Although *H. pulvinata* was moderately tolerant of Ecp2, the molecular basis for

this tolerance during mycoparasitism remains unclear.

The Hce2 (Homologs of *C. fulvum* Ecp2) superfamily is widely conserved across the

Ascomycota and *Basidiomycota*, suggesting that its origin predates the divergence of the

Dikarya lineage (Stergiopoulos *et al.*, 2012). In contrast, *U. maydis* UmVKP4, which shares

structural similarity with Ecp2, lacks the Hce2 domain and shared sequence motifs, and

differs in the number of disulfide bonds (Fig. **S7b**). These findings suggest that UmVKP4

represents a case of convergent evolution, resulting in structural resemblance to Ecp2 and

ZtKP4. Molecular evolutionary analyses suggest that the amino acid substitution underlying

Hce2 function diversification was fixed early in evolution (Stergiopoulos *et al.*, 2012).

Ancestral Hce2 proteins may initially act as antimicrobial factors to provide competitive

advantages in microbe-rich habitats, and have been co-opted as an effector to inhibit plant

immunity. Mutations in effector genes that allow *C. fulvum* to evade *Cf* resistance in plants

may increase its susceptibility to antagonistic microbes such as *H. pulvinata*. This hypothesis

suggests that mycoparasitic biocontrol agents can circumvent the classical arms race between

plant and pathogen, providing a sustainable disease management.

Potential metabolic trade-offs in amino acid pathways between fungi and tomato plants

The physiological changes found in tomato plants highlight the dynamic interplay between

microbial interplay and plant defense responses. The expression of genes in multiple amino

acid biosynthetic pathways were enhanced only when *H. pulvinata* parasitized *C. fulvum* on

tomato leaves. Amino acids are key mediators of plant–microbe interactions, functioning as

signaling molecules that activate defense and as nutritional sources that are exploited by

pathogens (Moormann *et al.*, 2022; Tünnermann *et al.*, 2022). Indeed, infection by *C. fulvum*

leads to the accumulation of free amino acids in the apoplast (Solomon & Oliver, 2001).

Because the acquisition of plant-derived amino acids is energetically advantageous to

microbes, many plant pathogens modulate expression of genes encoding amino acid

transporters to optimize uptake (Sonawala *et al.*, 2018). Thus, the regulation of amino acid

assimilation and transport in plants can be a decisive factor shaping the outcome of plant–

pathogen interactions.

In contrast, amino acid biosynthetic pathways in *H. pulvinata* and *C. fulvum* were

downregulated under mycoparasitic conditions, suggesting that tomato-derived amino acids

could be shared or sequentially utilized by these fungi (Fig. **8**). In this metabolic trade-off,

tomato plants might invest in amino acids that benefit *H. pulvinata*, while receiving

biocontrol protection in return. Although amino acids are fundamental primary metabolites

during fungal vegetative growth, the metabolic profile of *H. pulvinata*, which exhibited

vigorous hyphal growth during mycoparasitism, deviated from this conventional paradigm.

Because *H. pulvinata* parasitizes *C. fulvum*, it likely becomes the recipient of these plant-

derived nutrients via its host *C. fulvum*, forming a potential metabolic trade-off in the

tritrophic interaction. In this context, the activation of amino acid biosynthesis in tomato may

provide a nutritional foundation that ultimately supports colonization and biocontrol activity

by *H. pulvinata*. Although *C. fulvum* may also acquire amino acids from plants, these

resources are likely reallocated to *H. pulvinata*, the ultimate beneficiary of the tritrophic

interaction. Similar strategies have been reported for the biocontrol fungus *Pseudozyma*

flocculosa, active against powdery mildews, which transiently suppresses photosynthesis to

promote amino acid release (Laur *et al.*, 2018). The mycoparasitic fungus *Trichoderma*

erinaceum exploits environmental amino acids to enhance the synthesis of antifungal

secondary metabolites derived from amino acids (Guo *et al.*, 2020). Together, these results

 suggested that shifts in amino acid metabolism enable *H. pulvinata* to harness pathogen-

746 induced plant responses to its own benefit.

Fig. 8. Overview of the tritrophic interaction between tomato plants, the pathogen

Cladosporium fulvum, **and its mycoparasite** *Hansfordia pulvinata*. *C. fulvum* secretes

effector proteins that function as virulence factors in susceptible tomato cultivars and

antifungal proteins active against other fungi, but act as avirulence factors upon recognition

by *Cf-* containing resistant cultivars. *H. pulvinata* parasitizes *C. fulvum* and induces the

production of apoplastic antifungal substances in tomato plants via the Nep1-like protein

HpNlp1. *H. pulvinata* also activates the biosynthesis of the antifungal sesquiterpene 13-

deoxyphomenone during mycoparasitism. Other plant defense responses support this

biocontrol effect. Pathways related to amino acid biosynthesis in the three organisms suggest

that nutrients from the plant are supplied to *H. pulvinata*.

Mycoparasitism by *H. pulvinata* modulates plant immunity and pathogen growth

In conclusion, the mycoparasite *H. pulvinata* actively reprogrammes gene expression and

physiological responses in the pathogen *C. fulvum* and its tomato host to establish a stable

tritrophic interaction (Fig. **8**). *H. pulvinata* induced expression of the 13-deoxyphomenone

biosynthetic gene cluster in tomato plants; antimicrobial substances accumulated in the

apoplast and directly suppressed *C. fulvum*, thus enhancing plant resistance. In turn, *C.*

fulvum secretes effectors such as Ecp2 that exhibites roles in both fungal competition and

plant immunity. During the tritrophic interaction in tomato, amino acid metabolism and

defense response pathways are reprogrammed, orchestrating distinct strategies against

beneficial and pathogenic microbes. Collectively, these findings underscore the dynamic

interplay in plant–pathogen–mycoparasite interactions, demonstrating that *H. pulvinata*

simultaneously strengthens tomato immunity and represses *C. fulvum* virulence via direct

antagonism and modulation of plant metabolic pathways.

ACKNOWLEDGEMENTS

This work was supported by a Grant-in-Aid for Scientific Research from JSPS (20H02993

and 24K08919) and the Research and Implementation Promotion Program through Open

Innovation Grants (JPJ011937) from the Project of the Bio-oriented Technology Research

Advancement Institution (BRAIN) (Y. Iida). We are grateful to K. Ikeda for providing fungal

strains, M.H.A.J. Joosten for *Cf-ECP2* tomato seeds, D. Takemoto for the *psojNIP* gene, F.

Takken for pSfinx, and Y. Kubo for the *C. orbiculare* strain. We thank Y. Kubo, M.Z. Fanani

and S. Kodama for valuable suggestions, C. Tanaka for laboratory management, P.J.G.M. de

Wit for critical reading of the manuscript.

COMPETING INTERESTS

The authors declare that they have no commercial or financial relationships that could be

construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

YI designed the study and managed the research funds. KM, HS and TaS performed RNA-

seq and analyzed the data. KM, MK, MO, and TK analyzed NLPs. ToS conducted electron

microscopy and EI developed a microscopic observation protocol. KM and KS studied the

Ecp2 protein. KM and YI wrote the manuscript. All authors reviewed and approved the final

version.

DATA AVAILABILITY

Genomic sequences for *C. fulvum* strain Race5_Kim and *H. pulvinata* strain 414-3 are

available as accessions PRJNA565804 and PRJDB8178 at NCBI BioProject, respectively

(Sushida *et al*., 2019; Zaccaron & Stergiopoulos, 2024). Tomato genome data were obtained

from the Solanaceae Genomics Network (Hosmani *et al*., 2019). Additional genomes are

accessible via JGI MycoCosm (mycocosm.jgi.doe.gov) and the DDBJ/EMBL/GenBank

repository (www.ncbi.nlm.nih.gov). Custom scripts and supporting datasets were deposited at

Figshare (doi.org/10.6084/m9.figshare.30265009).

REFERENCES

Bailey TL, Johnson J, Grant CE, Noble WS. 2015. The MEME Suite. *Nucleic Acids*
Research. 2015 Jul 1;43(W1):W39-49. doi: 10.1093/nar/gkv416. Epub 2015 May 7.
PMID: 25953851; PMCID: PMC4489269.

Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, Schiegg M, Ales J,
Beier T, Rudy M, et al. 2019. ilastik: interactive machine learning for (bio)image
analysis. *Nature Methods*. 2019 Dec;16(12):1226-1232. doi: 10.1038/s41592-019-0582-
9. Epub 2019 Sep 30. PMID: 31570887.

Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, Fetter A, Terlouw
BR, Metcalf WW, Helfrich EJN, et al. 2023. antiSMASH 7.0: new and improved
predictions for detection, regulation, chemical structures and visualisation. *Nucleic Acids*
Research. 2023 Jul 5;51(W1):W46-W50. doi: 10.1093/nar/gkad344. PMID: 37140036;
PMCID: PMC10320115.

Böhm H, Albert I, Oome S, Raaymakers TM, Van den Ackerveken G, Nürnberger T.
2014. A conserved peptide pattern from a widespread microbial virulence factor triggers
pattern-induced immunity in *Arabidopsis*. *PLOS Pathogens*. 2014 Nov
6;10(11):e1004491. doi: 10.1371/journal.ppat.1004491. PMID: 25375108; PMCID:
PMC4223075.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics*. 2014 Aug 1;30(15):2114-20. doi:
10.1093/bioinformatics/btu170. Epub 2014 Apr 1. PMID: 24695404; PMCID:
PMC4103590.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. Aug
1;25(15):1972-3. doi: 10.1093/bioinformatics/btp348. Epub 2009 Jun 8. PMID:
19505945; PMCID: PMC2712344.

Chaillot J, Tebbji F, Remmal A, Boone C, Brown GW, Bellaoui M, Sellam A. 2015. The
Monoterpene Carvacrol Generates Endoplasmic Reticulum Stress in the Pathogenic
Fungus *Candida albicans*. *Antimicrobial Agents and Chemotherapy*. 2015
Aug;59(8):4584-92. doi: 10.1128/AAC.00551-15. Epub 2015 May 26. PMID: 26014932;
PMCID: PMC4505245.

Chavarro-Carrero EA, Snelders NC, Torres DE, Kraege A, López-Moral A, Petti GC,
Punt W, Wieneke J, García-Velasco R, López-Herrera CJ, et al. 2024. The soil-borne
white root rot pathogen *Rosellinia necatrix* expresses antimicrobial proteins during host

colonization. *PLOS Pathogens*. 2024 Jan 18;20(1):e1011866. doi:

10.1371/journal.ppat.1011866. PMID: 38236788; PMCID: PMC10796067.

Chen C, Wu Y, Li J, Wang X, Zeng Z, Xu J, Liu Y, Feng J, Chen H, He Y, et al. 2023.

TBtools-II: A "one for all, all for one" bioinformatics platform for biological big-data

mining. *Molecular Plant*. 2023 Nov 6;16(11):1733-1742. doi:

10.1016/j.molp.2023.09.010. Epub 2023 Sep 22. PMID: 37740491.

Eichfeld R, Mahdi LK, De Quattro C, Armbruster L, Endeshaw AB, Miyauchi S,

Hellmann MJ, Cord-Landwehr S, Peterson D, Singan V, et al. 2024. Transcriptomics

reveal a mechanism of niche defense: two beneficial root endophytes deploy an

antimicrobial GH18-CBM5 chitinase to protect their hosts. *New Phytologist*. 2024

Nov;244(3):980-996. doi: 10.1111/nph.20080. Epub 2024 Sep 3. PMID: 39224928.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biology*. 2019 Nov 14;20(1):238. doi: 10.1186/s13059-019-1832-y.

PMID: 31727128; PMCID: PMC6857279.

Gage MJ, Bruenn J, Fischer M, Sanders D, Smith TJ. 2001. KP4 fungal toxin inhibits

growth in *Ustilago maydis* by blocking calcium uptake. *Molecular Microbiology*. 2001

Aug;41(4):775-85. doi: 10.1046/j.1365-2958.2001.02554.x. PMID: 11532143.

Gage MJ, Rane SG, Hockerman GH, Smith TJ. 2002. The virally encoded fungal toxin

KP4 specifically blocks L-type voltage-gated calcium channels. *Molecular*

Pharmacology. 2002 Apr;61(4):936-44. doi: 10.1124/mol.61.4.936. PMID: 11901234.

Gene Ontology Consortium; Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin

HJ, Ebert D, Feuermann M, Gaudet P, Harris NL, et al. 2023. The Gene Ontology

knowledgebase in 2023. *Genetics*. 2023 May 4;224(1):iyad031. doi:

10.1093/genetics/iyad031. PMID: 36866529; PMCID: PMC10158837.

Gíslason MH, Nielsen H, Almagro Armenteros JJ, Johansen AR. 2021. Prediction of

GPI-anchored proteins with pointer neural networks. *Current Research in Biotechnology*.

2021;3:6-13. doi:10.1016/j.crbiot.2021.01.001.

Griffiths S, Mesarich CH, Overdijk EJR, Saccomanno B, de Wit PJGM, Collemare J.

2018. Down-regulation of cladofulvin biosynthesis is required for biotrophic growth of

Cladosporium fulvum on tomato. *Molecular Plant Pathology*. 2018 Feb;19(2):369-380.

doi: 10.1111/mpp.12527. Epub 2017 Feb 6. PMID: 27997759; PMCID: PMC6638085.

Gu F, Khimani A, Rane SG, Flurkey WH, Bozarth RF, Smith TJ. 1995. Structure and

function of a virally encoded fungal toxin from *Ustilago maydis*: a fungal and

mammalian Ca2+ channel inhibitor. *Structure*. 1995 Aug 15;3(8):805-14. doi: 10.1016/s0969-2126(01)00215-5. PMID: 7582897.

de Guillen K, Mammri L, Gracy J, Padilla A, Barthe P, Hoh F, Lahfa M, Rouffet J, Petit-Houdenot Y, Kroj T, et al. 2024. *Zymoseptoria tritici* effectors structurally related to killer proteins UmV-KP4 and UmV-KP6 are toxic to fungi, and define extended protein families in fungi. *bioRxiv*. 2024 Oct 14:2024.10.14.618152. doi:10.1101/2024.10.14.618152.

Guo YW, Gong BQ, Yuan J, Li HJ, Mahmud T, Huang Y, Li JF, Yang DP, Lan WJ. 2020. l-Phenylalanine Alters the Privileged Secondary Metabolite Production in the Marine-Derived Fungus T*richoderma erinaceum* F1-1. *Journal of Natural Products*. 2020 Jan 24;83(1):79-87. doi: 10.1021/acs.jnatprod.9b00710. Epub 2019 Dec 30. PMID: 31886665.

Holm L. 2022. Dali server: structural unification of protein families. *Nucleic Acids Research*. 2022 Jul 5;50(W1):W210-W215. doi: 10.1093/nar/gkac387. PMID: 35610055; PMCID: PMC9252788.

Hosmani PS, Flores-Gonzalez M, van de Geest H, Maumus F, Bakker LV, Schijlen E, van Haarst J, Cordewener J, Sanchez-Perez G, Peters S, et al. 2019. An improved de novo assembly and annotation of the tomato reference genome using single-molecule sequencing, Hi-C proximity ligation and optical maps. *bioRxiv*. 2019;767764. doi:10.1101/767764

Iida Y, Ikeda K, Sakai H, Nakagawa H, Nishi O, Higashi Y. 2018. Evaluation of the potential biocontrol activity of *Dicyma pulvinata* against *Cladosporium fulvum*, the causal agent of tomato leaf mold. *Plant Pathology*. 2018;67(9):1883–1890. doi:10.1111/ppa.12916.

Iida Y, van 't Hof P, Beenen H, Mesarich C, Kubota M, Stergiopoulos I, Mehrabi R, Notsu A, Fujiwara K, Bahkali A, et al. 2015. Novel Mutations Detected in Avirulence Genes Overcoming Tomato *Cf* Resistance Genes in Isolates of a Japanese Population of *Cladosporium fulvum*. *PLoS One*. 2015 Apr 22;10(4):e0123271. doi: 10.1371/journal.pone.0123271. PMID: 25902074; PMCID: PMC4406682.

de Jonge R, van Esse HP, Kombrink A, Shinya T, Desaki Y, Bours R, van der Krol S, Shibuya N, Joosten MH, Thomma BP. 2010. Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science*. 2010 Aug 20;329(5994):953-5. doi: 10.1126/science.1190859. PMID: 20724636.

Joosten MH. 2012. Isolation of apoplastic fluid from leaf tissue by the vacuum infiltration-
centrifugation technique. *Methods in Molecular Biology*. 2012;835:603-10. doi:
10.1007/978-1-61779-501-5_38. PMID: 22183681.

Kanehisa M, Furumichi M, Sato Y, Matsuura Y, Ishiguro-Watanabe M. 2025. KEGG:
biological systems database as a model of the real world. *Nucleic Acids Research*. 2025
Jan 6;53(D1):D672-D677. doi: 10.1093/nar/gkae909. PMID: 39417505; PMCID:
PMC11701520.

Katoh K., Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular biology and*
evolution. Apr;30(4):772-80. doi: 10.1093/molbev/mst010. Epub 2013 Jan 16. PMID:
23329690; PMCID: PMC3603318.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*. 2019
Aug;37(8):907-915. doi: 10.1038/s41587-019-0201-4. Epub 2019 Aug 2. PMID:
31375807; PMCID: PMC7605509.

de Kock MJ, Iskandar HM, Brandwagt BF, Laugé R, de Wit PJGM, Lindhout P. 2004.
Recognition of *Cladosporium fulvum* Ecp2 elicitor by non-host *Nicotiana* spp. is
mediated by a single dominant gene that is not homologous to known *Cf*-genes.
Molecular Plant Pathology. 2004 Sep 1;5(5):397-408. doi: 10.1111/j.1364-
3703.2004.00239.x. PMID: 20565616.

Kourelis J, Malik S, Mattinson O, Krauter S, Kahlon PS, Paulus JK, van der Hoorn
RAL. 2020. Evolution of a guarded decoy protease and its receptor in solanaceous
plants. *Nature Communications*. 2020 Sep 2;11(1):4393. doi: 10.1038/s41467-020-
18069-5. Erratum in: *Nature Communications*. 2024 Mar 13;15(1):2278. doi:
10.1038/s41467-024-46611-2. PMID: 32879321; PMCID: PMC7468133.

Laur J, Ramakrishnan GB, Labbé C, Lefebvre F, Spanu PD, Bélanger RR. 2018.
Effectors involved in fungal-fungal interaction lead to a rare phenomenon of
hyperbiotrophy in the tritrophic system biocontrol agent-powdery mildew-plant. *New*
Phytologist. 2018 Jan;217(2):713-725. doi: 10.1111/nph.14851. Epub 2017 Oct 18.
PMID: 29044534; PMCID: PMC6079639.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R; 1000 Genome Project Data Processing Subgroup. 2009. The Sequence
Alignment/Map format and SAMtools. *Bioinformatics*. 2009 Aug 15;25(16):2078-9. doi:

10.1093/bioinformatics/btp352. Epub 2009 Jun 8. PMID: 19505943; PMCID:

PMC2723002.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biology*. 2014;15(12):550. doi:

10.1186/s13059-014-0550-8. PMID: 25516281; PMCID: PMC4302049.

Luderer R, Takken FL, de Wit PJGM, Joosten MH. 2002. *Cladosporium fulvum*

overcomes *Cf-2*-mediated resistance by producing truncated AVR2 elicitor proteins.

Molecular *Microbiology*. 2002 Aug;45(3):875-84. doi: 10.1046/j.1365-

2958.2002.03060.x. PMID: 12139631.

Ma L, Lukasik E, Gawehns F, Takken FL. 2012. The use of agroinfiltration for transient

expression of plant resistance and fungal effector proteins in *Nicotiana benthamiana*

leaves. *Methods in Molecular Biology*. 2012;835:61-74. doi: 10.1007/978-1-61779-501-

5_4. PMID: 22183647.

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R,

Butcher S. 2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in

2024. *Nucleic Acids Research*. 2024 Jul 5;52(W1):W521-W525. doi:

10.1093/nar/gkae241. PMID: 38597606; PMCID: PMC11223882.

Maeda K, Sumita T, Nishi O, Sushida H, Higashi Y, Nakagawa H, Suzuki T, Iwao E,

Fanani MZ, Nishiya Y, Iida Y. 2025. Adaptive evolution of sesquiterpene

deoxyphomenone in mycoparasitism by *Hansfordia pulvinata* associated with horizontal

gene transfer from *Aspergillus* species. *mBio*. 2025 Apr 9;16(4):e0400724. doi:

10.1128/mbio.04007-24. Epub 2025 Mar 20. PMID: 40111082; PMCID: PMC11980549.

Mesarich CH, Griffiths SA, van der Burgt A, Okmen B, Beenen HG, Etalo DW, Joosten

MH, de Wit PJGM. 2014. Transcriptome sequencing uncovers the *Avr5* avirulence gene

of the tomato leaf mold pathogen *Cladosporium fulvum*. *Molecular Plant-Microbe*

Interactions. 2014 Aug;27(8):846-57. doi: 10.1094/MPMI-02-14-0050-R. PMID:

24678832.

Mesarich CH, Ökmen B, Rovenich H, Griffiths SA, Wang C, Karimi Jashni M,

Mihajlovski A, Collemare J, Hunziker L, Deng CH, et al. 2018. Specific

Hypersensitive Response-Associated Recognition of New Apoplastic Effectors from

Cladosporium fulvum in Wild Tomato. *Molecular Plant-Microbe Interactions*. 2018

Jan;31(1):145-162. doi: 10.1094/MPMI-05-17-0114-FI. Epub 2017 Nov 16. PMID:

29144204.

Mesarich CH, Barnes I, Bradley EL, de la Rosa S, de Wit PJGM, Guo Y, Griffiths SA, Hamelin RC, Joosten MHAJ, Lu M, et al. 2023. Beyond the genomes of *Fulvia fulva* (syn. *Cladosporium fulvum*) and *Dothistroma septosporum*: New insights into how these fungal pathogens interact with their host plants. *Molecular Plant Pathology*. 2023 May;24(5):474-494. doi: 10.1111/mpp.13309. Epub 2023 Feb 15. PMID: 36790136; PMCID: PMC10098069.

Mesny F, Wolf V, López-Moral A, Kraege A, Punt W, Park J, Zhu J, Sato Y, Thomma BPHJ. 2024. Plant-associated fungi co-opt ancient antimicrobials for host manipulation. *bioRxiv*. Preprint posted online 2024 Jan 4. doi:10.1101/2024.01.04.574150.

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022. ColabFold: making protein folding accessible to all. *Nature Methods*. 2022 Jun;19(6):679-682. doi: 10.1038/s41592-022-01488-1. Epub 2022 May 30. PMID: 35637307; PMCID: PMC9184281.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE, Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families database in 2021. *Nucleic Acids Research*. 2021 Jan 8;49(D1):D412-D419. doi: 10.1093/nar/gkaa913. PMID: 33125078; PMCID: PMC7779014.

Moormann J, Heinemann B, Hildebrandt TM. 2022. News about amino acid metabolism in plant-microbe interactions. *Trends in Biochemical Sciences*. 2022 Oct;47(10):839-850. doi: 10.1016/j.tibs.2022.07.001. Epub 2022 Aug 1. PMID: 35927139.

Oome S, Raaymakers TM, Cabral A, Samwel S, Böhm H, Albert I, Nürnberger T, Van den Ackerveken G. 2014. Nep1-like proteins from three kingdoms of life act as a microbe-associated molecular pattern in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA*. 2014 Nov 25;111(47):16955-60. doi: 10.1073/pnas.1410031111. Epub 2014 Nov 3. PMID: 25368167; PMCID: PMC4250136.

Peresse M, Le Picard D. 1980. *Hansfordia pulvinata*, mycoparasite destructeur du *Cladosporium fulvum*. *Mycopathologia* 1980. 71:23–30.

Qutob D, Kamoun S, Gijzen M. 2002. Expression of a *Phytophthora sojae* necrosis-inducing protein occurs during transition from biotrophy to necrotrophy. *The Plant Journal*. 2002 Nov;32(3):361-73. doi: 10.1046/j.1365-313x.2002.01439.x. PMID: 12410814.

Qutob D, Kemmerling B, Brunner F, Küfner I, Engelhardt S, Gust AA, Luberacki B, Seitz HU, Stahl D, Rauhut T, et al. 2006. Phytotoxicity and innate immune responses

induced by Nep1-like proteins. *The Plant Cell*. 2006 Dec;18(12):3721-44. doi:

10.1105/tpc.106.044180. Epub 2006 Dec 28. PMID: 17194768; PMCID: PMC1785393.

R Core Team R. 2024. *R: a language and environment for statistical computing*. Vienna,

Austria: R Foundation for Statistical Computing.

Rocafort M, Fudal I, Mesarich CH. 2020. Apoplastic effector proteins of plant-associated

fungi and oomycetes. *Current Opinion in Plant Biology*. 2020 Aug;56:9-19. doi:

10.1016/j.pbi.2020.02.004. Epub 2020 Apr 2. PMID: 32247857.

Rooney HC, Van't Klooster JW, van der Hoorn RA, Joosten MH, Jones JD, de Wit

PJGM. 2005. Cladosporium Avr2 inhibits tomato Rcr3 protease required for *Cf-2*-

dependent disease resistance. *Science*. 2005 Jun 17;308(5729):1783-6. doi:

10.1126/science.1111404. Epub 2005 Apr 21. Erratum in: Science. 2005 Oct

7;310(5745):54. PMID: 15845874.

de la Rosa S, Schol CR, Ramos Peregrina Á, Winter DJ, Hilgers AM, Maeda K, Iida Y,

Tarallo M, Jia R, Beenen HG, et al. 2024. Sequential breakdown of the *Cf-9* leaf mold

resistance locus in tomato by *Fulvia fulva*. *New Phytologist*. 2024 Aug;243(4):1522-

1538. doi: 10.1111/nph.19925. Epub 2024 Jun 24. PMID: 38922927.

Seidl MF, Van den Ackerveken G. 2019. Activity and Phylogenetics of the Broadly

Occurring Family of Microbial Nep1-Like Proteins. *Annual Review of Phytopathology*.

2019 Aug 25;57:367-386. doi: 10.1146/annurev-phyto-082718-100054. Epub 2019 Jul 5.

PMID: 31283435.

Shabab M, Shindo T, Gu C, Kaschani F, Pansuriya T, Chintha R, Harzen A, Colby T,

Kamoun S, van der Hoorn RA. 2008. Fungal effector protein AVR2 targets

diversifying defense-related cys proteases of tomato. *The Plant Cell*. 2008

Apr;20(4):1169-83. doi: 10.1105/tpc.107.056325. Epub 2008 Apr 30. PMID: 18451324;

PMCID: PMC2390736.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. Syst

Biol. Jun;51(3):492-508. doi: 10.1080/10635150290069913. PMID: 12079646.

Snelders NC, Rovenich H, Petti GC, Rocafort M, van den Berg GCM, Vorholt JA,

Mesters JR, Seidl MF, Nijland R, Thomma BPHJ. 2020. Microbiome manipulation

by a soil-borne fungal plant pathogen using effector proteins. *Nature Plants*. 2020

Nov;6(11):1365-1374. doi: 10.1038/s41477-020-00799-5. Epub 2020 Nov 2. PMID:

33139860.

Snelders NC, Rovenich H, Thomma BPHJ. 2022. Microbiota manipulation through the

secretion of effector proteins is fundamental to the wealth of lifestyles in the fungal

kingdom. *FEMS Microbiology Reviews*. 2022 Sep 2;46(5):fuac022. doi:

10.1093/femsre/fuac022. PMID: 35604874; PMCID: PMC9438471.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics*. May 1;30(9):1312-3. doi:

10.1093/bioinformatics/btu033. Epub 2014 Jan 21. PMID: 24451623; PMCID:

PMC3998144.

Solomon PS, Oliver RP. 2001. The nitrogen content of the tomato leaf apoplast increases

during infection by *Cladosporium fulvum*. *Planta*. 2001 Jun;213(2):241-9. doi:

10.1007/s004250000500. PMID: 11469589.

Sonawala U, Dinkeloo K, Danna CH, McDowell JM, Pilot G. 2018. Review: Functional

linkages between amino acid transporters and plant responses to pathogens. *Plant*

Science. 2018 Dec;277:79-88. doi: 10.1016/j.plantsci.2018.09.009. Epub 2018 Sep 19.

PMID: 30466603.

Sperschneider J, Dodds PN. 2022. EffectorP 3.0: Prediction of Apoplastic and Cytoplasmic

Effectors in Fungi and Oomycetes. *Molecular Plant-Microbe Interactions*. 2022

Feb;35(2):146-156. doi: 10.1094/MPMI-08-21-0201-R. Epub 2022 Feb 1. PMID:

34698534.

Sperschneider J, Catanzariti AM, DeBoer K, Petre B, Gardiner DM, Singh KB, Dodds

PN, Taylor JM. 2017. LOCALIZER: subcellular localization prediction of both plant

and effector proteins in the plant cell. *Scientific Reports*. 2017 Mar 16;7:44598. doi:

10.1038/srep44598. PMID: 28300209; PMCID: PMC5353544.

Stergiopoulos I, Kourmpetis YA, Slot JC, Bakker FT, De Wit PJ, Rokas A. 2012. In

silico characterization and molecular evolutionary analysis of a novel superfamily of

fungal effector proteins. *Molecular Biology and Evolution*. 2012 Nov;29(11):3371-84.

doi: 10.1093/molbev/mss143. Epub 2012 May 23. PMID: 22628532.

Sushida H, Sumita T, Higashi Y, Iida Y. 2019. Draft Genome Sequence of *Dicyma*

pulvinata Strain 414-3, a Mycoparasite of *Cladosporium fulvum*, Causal Agent of

Tomato Leaf Mold. *Microbiology Resource Announcements*. 2019 Aug 29;8(35):e00655-

19. doi: 10.1128/MRA.00655-19. PMID: 31467096; PMCID: PMC6715866.

Tanaka S, Wada K, Katayama M, Marumo S. 1984. Isolation of Sporogen-AO1, a

sporogenic substance, from *Aspergillus oryzae*. A*gricultural and biological chemistry*.

1984;48(12):3189–3191. doi:10.1271/bbb1961.48.3189.

Tanaka S, Wada K, Marumo S, Hattori H. 1984. Structure of sporogen-ao 1, a sporogenic

substance of *Aspergillus oryzae. Tetrahedron Letters*. 1984;25(51):5907–5910.

doi:10.1016/S0040-4039(01)81717-2.

Tang D, Chen M, Huang X, Zhang G, Zeng L, Zhang G, Wu S, Wang Y. 2023. SRplot:

A free online platform for data visualization and graphing. *PLoS One*. 2023 Nov

9;18(11):e0294236. doi: 10.1371/journal.pone.0294236. PMID: 37943830; PMCID:

PMC10635526.

Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD,

Winther O, Brunak S, von Heijne G, Nielsen H. 2022. SignalP 6.0 predicts all five

types of signal peptides using protein language models. *Nature Biotechnology*. 2022

Jul;40(7):1023-1025. doi: 10.1038/s41587-021-01156-3. Epub 2022 Jan 3. PMID:

34980915; PMCID: PMC9287161.

Thomma BP, VAN Esse HP, Crous PW, DE Wit PJ. 2005. *Cladosporium fulvum* (syn.

Passalora fulva), a highly specialized plant pathogen as a model for functional studies on

plant pathogenic Mycosphaerellaceae. *Molecular Plant Pathology*. 2005 Jul 1;6(4):379-

93. doi: 10.1111/j.1364-3703.2005.00292.x. PMID: 20565665.

Tirilly Y, Kloosterman J, Sipma G, Kettenes-Van Den Bosch JJ. 1983. A fungitoxic

sesquiterpene from *Hansfordia pulvinata. Phytochemistry*. 1983;22(9):2082–2083.

doi:10.1016/0031-9422(83)80051-X.

Tünnermann L, Colou J, Näsholm T, Gratz R. 2022. To have or not to have: expression of

amino acid transporters during pathogen infection. *Plant Molecular Biology*. 2022

Jul;109(4-5):413-425. doi: 10.1007/s11103-022-01244-1. Epub 2022 Feb 1. PMID:

35103913; PMCID: PMC9213295.

Watanabe H, Horinouchi H, Muramoto Y, Ishii H. 2017. Occurrence of azoxystrobin-

resistant isolates in *Passalora fulva*, the pathogen of tomato leaf mold disease. *Plant

Pathology*. 2017;66(9):1472–1479. doi:10.1111/ppa.12701.

de Wit PJGM, van der Burgt A, Ökmen B, Stergiopoulos I, Abd-Elsalam KA, Aerts AL,

Bahkali AH, Beenen HG, Chettri P, Cox MP, et al. 2012. The genomes of the fungal

plant pathogens *Cladosporium fulvum* and *Dothistroma septosporum* reveal adaptation to

different hosts and lifestyles but also signatures of common ancestry. *PLOS Genetics*.

2012;8(11):e1003088. doi: 10.1371/journal.pgen.1003088. Epub 2012 Nov 29. Erratum

in: PLoS Genet. 2015 Dec 22;11(12):e1005775. doi: 10.1371/journal.pgen.1005775.

PMID: 23209441; PMCID: PMC3510045.

de Wit PJGM, Joosten MHAJ, Thomma BHPJ, Stergiopoulos I. 2009. Gene for gene models and beyond: the *Cladosporium fulvum*–tomato pathosystem. *Plant relationships*. Berlin, Heidelberg: Springer; 2009. p. 135–156. (The Mycota; vol 5). doi:10.1007/978-3-540-87407-2_7.

Wulff BB, Chakrabarti A, Jones DA. 2009. Recognitional specificity and evolution in the tomato-*Cladosporium fulvum* pathosystem. *Molecular Plant-Microbe Interactions*. 2009 Oct;22(10):1191-202. doi: 10.1094/MPMI-22-10-1191. PMID: 19737093.

Yan L, Chen J, Zhang C, Ma Z. 2008. Molecular characterization of benzimidazole-resistant isolates of *Cladosporium fulvum*. *Fems Microbiology Letters*. 2008 Jan;278(2):242-8. doi: 10.1111/j.1574-6968.2007.00999.x. PMID: 18096020.

Yang K, Chen C, Wang Y, Li J, Dong X, Cheng Y, Zhang H, Zhai Y, Ai G, Song Q, et al. 2022. Nep1-Like Proteins From the Biocontrol Agent *Pythium oligandrum* Enhance Plant Disease Resistance Independent of Cell Death and Reactive Oxygen Species. *Frontiers in Plant Science*. 2022 Mar 4;13:830636. doi: 10.3389/fpls.2022.830636. PMID: 35310640; PMCID: PMC8931738.

Zaccaron AZ, Stergiopoulos I. 2024. Analysis of five near-complete genome assemblies of the tomato pathogen *Cladosporium fulvum* uncovers additional accessory chromosomes and structural variations induced by transposable elements effecting the loss of avirulence genes. *BMC Biology*. 2024 Jan 29;22(1):25. doi: 10.1186/s12915-024-01818-z. PMID: 38281938; PMCID: PMC10823647.

Zaccaron AZ, Chen LH, Samaras A, Stergiopoulos I. 2022. A chromosome-scale genome assembly of the tomato pathogen *Cladosporium fulvum* reveals a compartmentalized genome architecture and the presence of a dispensable chromosome. *Microbial Genomics*. 2022 Apr;8(4):000819. doi: 10.1099/mgen.0.000819. PMID: 35471194; PMCID: PMC9453070.

Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018. dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Research*. 2018 Jul 2;46(W1):W95-W101. doi: 10.1093/nar/gky418. PMID: 29771380; PMCID: PMC6031026.

Zhang Y, Skolnick J. 2005. TM-align: a protein structure alignment algorithm based on the TM-score. *Nucleic Acids Research*. 2005 Apr 22;33(7):2302-9. doi: 10.1093/nar/gki524. PMID: 15849316; PMCID: PMC1084323.

Zhou BJ, Jia PS, Gao F, Guo HS. 2012. Molecular characterization and functional analysis of a necrosis- and ethylene-inducing, protein-encoding gene family from *Verticillium*

1136   *dahliae. Molecular Plant-Microbe Interactions*. 2012 Jul;25(7):964-75. doi:

1137   10.1094/MPMI-12-11-0319. PMID: 22414440.

1138

FIGURE LEGENDS

Fig. 1. *Cladosporium fulvum* parasitized by mycoparasitic fungus *Hansfordia pulvinata.*

(a) Representative images of a healthy tomato leaf, lesions caused by *C. fulvum* (*on planta*

Cf), and lesions covered by white mycelium of *H. pulvinata* (*on planta Cf/Hp*). Tomato

plants were inoculated with *C. fulvum*, and lesions developed after 2 weeks. *H. pulvinata*

was then inoculated onto the diseased leaves, and white mycelia appeared on the lesions a

week after the second inoculation. *In vitro* cultures of *H. pulvinata* (*in vitro Hp*) and *C.*

fulvum (*in vitro Cf*) on nylon membranes on water agar after transfer from PDA. *H. pulvinata*

parasitized *C. fulvum* colonies on water agar (*in vitro Cf/Hp*). (b) Scanning electron

micrographs of hyphae of *C. fulvum* and *H. pulvinata* in *in vitro Cf* and *Cf/Hp* conditions. *H.*

pulvinata hyphae have penetrated *C. fulvum* hyphae in the *in vitro Cf/Hp* condition. Bar

indicates 10 μm.

1151

Fig. 2. Differentially expressed genes (DEGs) and enriched KEGG pathways in

mycoparasite *Hansfordia pulvinata*. (a) UpSet plot of DEGs in the *in vitro* and *on planta*

mycoparasitic conditions (*in vitro Cf/Hp* and *on planta Cf/Hp*; Fig. 1a). Connected dots

indicate shared DEGs; Black single dots represent condition-specific DEGs. Vertical bars

show the number of up- and downregulated genes; colored horizontal bars indicate the total

number of DEGs detected in each interaction. (b) Bubble plot of KEGG pathway enrichment

analysis based on the 891 and 557 shared DEGs shown in (a). Keys: color of symbols

indicates significance level of enrichment (–log adjusted *P*); size of symbol indicates the

number of enriched DEGs in each pathway; the symbol indicates the functional class of the

enriched DEGs.

1162

Fig. 3. A Nep1-like protein (NLP) secreted from the mycoparasite *Hansfordia pulvinata*

induces a plant-mediated toxicity against spores of *Cladosporium fulvum*. (a) Workflow

for identifying small secreted proteins synthesized in *H. pulvinata* during mycoparasitism. (b)

MA plot for expression levels of 165 selected genes. Cutoff thresholds are $\log_2$ Fold-change

(±1.0) for *in vitro Cf/Hp* interaction and log Mean (0.3). (c) Phylogenetic tree constructed

based on NLP amino acid sequences using the maximum likelihood method with 1,000

bootstrap replicates. Type I and Type II have two and four cysteine residues, respectively.

Proteins analyzed in this study are in bold. (d) Multiple sequence alignment of NLP using MAFFT. Conserved residues are highlighted in orange. (e) Transient expression assays of NLPs in *Nicotiana benthamiana* via agroinfiltration. Chlorosis and necrosis were assessed 14 days after infiltration by trypan blue staining. Similar results were obtained in three independent experiments; representative results are in Supplementary Fig. S5c. (f) Percentage germination of conidia of *C. fulvum* after 24-h treatment with apoplastic fluid from *N. benthamiana* leaves that had been infiltrated with different NLPs or buffer. Asterisks indicate significant differences between treatments determined using Dunnett's or the Mann–Whitney *U* test.

Fig. 4. Differentially expressed genes (DEGs) and enriched KEGG pathways in the pathogen *Cladosporium fulvum*. (a) UpSet plot showing DEGs identified in the *on planta Cf, in vitro Cf/Hp,* and *on planta Cf/HP* conditions (see Fig. 1A). Connected dots indicate shared DEGs; Black single dots represent condition-specific DEGs. Vertical bars show the numbers of up- and downregulated genes; colored horizontal bars indicate the total number of DEGs detected in each condition. (b) Bubble plot of KEGG pathway enrichment analysis based on the 363 and 158 shared DEGs shown in (a). Keys: color of symbols indicates significant level of enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in each pathway; the symbol indicates the functional class of the enriched DEGs.

Fig. 5. Effector genes in the pathogen *Cladosporium fulvum* were highly expressed during mycoparasitism by *Hansfordia pulvinata*. (a) Functional classification of DEGs identified for the *on planta Cf*, *in vitro Cf/Hp* and *on planta Cf/Hp* interactions. Red: upregulated, stable: grey, blue: downregulated. The number of genes assigned to each category is in parentheses. (b) Heatmap showing expression profiles of effector genes. The red-to-blue gradient represents up- and downregulated expression, respectively. Genes with an adjusted *P*-value ≥0.05 are indicated in gray as not detected (ND). Red arrowhead indicates *Ecp2*. (c) Antimicrobial activity prediction for 56 effector genes upregulated during the *in vitro Cf/Hp* interaction (determined using the AMAPEC program). Symbols denote protein location predicted by EffectorP-fungi 3.0.

Fig. 6. Sequence dissimilarity and structural similarity between *Cladosporium fulvum* effector Ecp2 and killer toxin KP4 proteins. (a) Predicted 3D structure of Ecp2 generated

using AlphaFold2. The color spectrum from blue to red indicates the N- to C-terminal

positions of residues. (b) Structural comparison of Ecp2 with ZtKP4 and UmVKP4 proteins.

The superposition of the aligned proteins was evaluated using TM-align, and the TM-score

and RMSD values are shown. (c) Heatmap showing sequence similarity among Ecp2, ZtKP4,

and UmVKP4 proteins as identified by BLASTp analysis. (d) Conserved motifs of

the proteins identified by MEME motif analysis. (e) Representative tomato leaf from 4-week-

old tomato plant carrying the *Cf-ECP2* resistance gene at 6 days after infiltration with

purified Ecp2 protein produced by *Escherichia coli*. White dotted outline: area infiltrated

with infiltrated with 1 μM Ecp2; black dotted outline: infiltrated with control protein purified

from *E. coli* harboring an empty vector (EV). Numbers in parentheses indicate the number of

leaves with a hypersensitive response (HR) relative to the number of total leaves. (f) *In vitro*

antifungal activity of Ecp2 against *C. fulvum*, *H. pulvinata*, *Alternaria* sp., *Colletotrichum*

orbiculare, and *Saccharomyces cerevisiae*. Spores were cultured on 5% potato dextrose agar

with or without Ecp2. Relative hyphal length was determined 4 days after treatment. Bars =

250 μm. Significant differences between EV and Ecp2 treatments were assessed using

Student's t-test, and *p*-values are indicated.

1220

Fig. 7. Differentially expressed genes (DEGs) and enriched KEGG pathways in tomato

plants. (a) UpSet plot of DEGs in the *on planta Cf* and *on planta Cf/Hp* conditions (see Fig.

1a). Healthy leaves were used as a control. Connected dots indicate shared DEGs; Black

single dots represent condition-specific DEGs. Vertical bars show the number of up- and

downregulated genes; colored horizontal bars indicate the total number of DEGs detected in

each interaction. (b) Bubble plot of KEGG pathway enrichment analysis based on the 1,619

and 2,153 DEGs shown in (a). Keys: color of symbols indicates significance level of

enrichment (–log adjusted *P*); size of symbol indicates the number of enriched DEGs in each

pathway; the symbol indicates the functional class of the enriched DEGs. Stars indicate

pathways highlighted in (c). (c) KEGG pathways involved in amino acid biosynthesis in

tomato.

1232

Fig. 8. Overview of the tritrophic interaction among tomato plants, pathogen

*Cladosporium fulvum***, and mycoparasite *Hansfordia pulvinata*.** *C. fulvum* secretes effector

proteins that function as virulence factors in susceptible tomato cultivars and antifungal

proteins active against other fungi, but act as avirulence factors upon recognition by *Cf*-

resistant cultivars. *H. pulvinata* parasitizes *C. fulvum* and induces the production of apoplastic

antifungal substances in tomato plants via the Nep1-like protein HpNlp1. *H. pulvinata* also

activates the biosynthesis of the antifungal sesquiterpene 13-deoxyphomenone during

mycoparasitism. Other plant defense responses support this biocontrol effect. Pathways

related to amino acid biosynthesis in the three organisms suggest that nutrients from the plant

are supplied to *H. pulvinata*.

1243

SUPPORTING INFORMATION

Fig. S1 Experimental treatments and principal component analysis (PCA) of transcripts from

the mycoparasite *Hansfordia pulvinata* (Hp), the pathogen *Cladosporium fulvum* (Cf), and

tomato plant, *Solanum lycopersicum*.

Fig. S2 Secondary metabolite biosynthesis gene clusters in the mycoparasite *Hansfordia*

pulvinata.

Fig. S3 Phylogenetic analysis and functional characterization of proteins in fungal species.

Fig. S4 Structure of HpNlp1.

Fig. S5 Responses in plant leaves triggered by HpNlp1 protein.

Fig. S6 Heat map and hierarchical clustering of effector genes expressed in *Cladosporium*

fulvum.

Fig. S7 Structural comparative analysis of Ecp2 with ZtKP4 and UmVKP4.

Fig. S8 Differentially expressed genes (DEGs) in KEGG pathways detected in tomato plants.

Table S1 Primer sequences used in this study.

Table S2 Summary statistics of RNA-seq data and mapping results.

Table S3 Functional annotation of all genes in *Hansfordia pulvinata* genome.

Table S4 Predicted gene clusters involved in secondary metabolite biosynthesis in

Hansfordia pulvinata.

Table S5 Small secreted protein-coding genes ($\leq$300 aa) in *Hansfordia pulvinata*.

Table S6 Key genes encoding secondary metabolism enzymes in *Cladosporium fulvum*.

Table S7 Effector genes identified in *Cladosporium fulvum*.

Table S8 Structural homologs of Ecp2 protein identified from PDB25 using the Dali server.

Table S9 Species possessing homologs of Ecp2 and KP4 proteins.

Table S10 Log_2 fold-change for defensin-like genes in tomato plants.