Shared feedback control principles across standing and walking

Yang Geng, Jaap H. van Dieën, Sjoerd M. Bruijn*

Department of Human Movement Sciences, Faculty of Behavioral and Movement Sciences,
Amsterdam Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

* Corresponding author at: Van der Boechorststraat 9, 1081 BT Amsterdam, the Netherlands.

E-mail address: s.m.bruijn@vu.nl

Abstract

Feedback control is required to maintain stability in both walking and standing, whereas the
control parameters may differ due to the inherently different characteristics of these tasks. How
feedback is adapted between tasks is unclear. To address this, we applied a unified feedback
model, relating preceding center of mass position and velocity information to corrective ground
reaction forces, to estimate and compare the control parameters between walking and standing.

15 healthy young participants (21± 4 yrs, 63 ± 9 kg, 1.70 ± 0.10 m, 13 females, 2 males)
participated, they walked at 4.5 km/h for 5 minutes, and performed 3 different standing tasks:
normal standing, unipedal standing, and step posture for 1 minute, repeated 5 times. Whole-body
kinematics and ground reaction forces were collected and used to fit the feedback model and
estimate feedback delay and gains. Results showed good model fits for both walking and
standing with mean R^2-values higher than 0.69. The feedback delay was significantly longer in
walking than in standing. While the feedback position gains and velocity gains varied
significantly across tasks and directions, position gains were mostly above critical stiffness and
velocity gains all at under-damped level, with as only exception that position gains in the AP
direction in walking, which were slightly below critical stiffness. The ratio of position to velocity
gains in all tasks remained consistently close to the eigenfrequency ($\sqrt{g/l}$) as predicted by the
extrapolated center of mass concept.

Our findings suggest a robust control strategy in which the sensorimotor system maintains a
relative consistent weighting between position and velocity feedback for walking and standing.

1. Introduction

Walking and standing are two fundamental motor behaviors in human daily life. Both require sophisticated neuromuscular control to maintain an upright body orientation against the effects of unpredictable perturbations which are amplified by gravity. Such stabilization is achieved through coordination of intrinsic mechanics and feedback control. Intrinsic mechanics provide immediate resistance to perturbations based on segments' inertia and joints' stiffness and damping. The latter can be modulated by muscle co-activation to deal with expected perturbations or uncertainty, but maintaining high levels of co-contraction is energetically costly (Koelewijn & Van Den Bogert, 2022). In contrast, feedback control is adaptive and generates corrective responses selectively based on sensory information (Forbes et al., 2018; Shanbhag et al., 2024).

Given the different temporal and biomechanical characteristics of standing and walking, feedback control parameters can be expected to differ between these motor tasks, reflecting task-specific modulations. Standing is a quasi-static behavior, while walking is a dynamic, cyclic motion. This temporal difference suggests distinct feedback strategies: time-invariant feedback control in standing but a phase-dependent control in walking. Additionally, in standing, the anatomical constraints and smaller base of support (BoS) make stabilization more challenging in the anteroposterior (AP) direction than in the mediolateral (ML) direction (Blaszczyk et al., 2014; Winter, 1995; Winter et al., 1996). However, during walking, the stability requirements shift. Stabilization of AP movement benefits from passive mechanics (Coleman & Ruina, 1998; Kuo & Donelan, 2010; Mcgeer, 1990), while the ML movement must be actively stabilized (Bauby & Kuo, 2000; Kuo, 1999).

Several feedback models have been developed to identify feedback control in standing (Jakubowski et al., 2024; Safavynia & Ting, 2013; Shanbhag et al., 2024; Welch & Ting, 2008) and walking (Afschrift et al., 2021). These models indicate that sensory estimates of the center of mass (CoM) kinematics are used to modulate corrective responses (e.g., EMG, ankle torque, foot placement) to stabilize standing and walking. However, identification of these models requires perturbations and measurements of physiological signals (e.g., EMG), which poses high experimental demands and limits practical application. As an alternative, van Dieen et al. (2024) proposed a feedback model that can identify feedback control in unperturbed walking by fitting a linear relationship between the delayed extrapolated center of mass (XCoM) and horizontal ground reaction forces, without external perturbations or physiological measurements. More recently, we refined this feedback model by using the CoM position and velocity information as separate predictors instead of the XCoM, and validated that this modified model can be applied to assess feedback control in standing (Geng et al., 2025). Until now, these models were derived under the specific context being studied (i.e., either standing or walking), which leaves a gap in our understanding of how stability control is adapted between standing and walking.

In the present study, we addressed this gap by applying the aforementioned feedback model to unperturbed standing and walking. Using this model, we can estimate feedback position and velocity gains and the effective delay, helping us to understand how the CNS modulates feedback control in these tasks, while operating under the same fundamental control principle:

generating corrective ground reaction forces based on preceding center of mass position (PCoM)
and velocity (VCoM) information.

Hof et al. (2005) described the relative importance of position and velocity for maintaining
dynamic stability. They showed that stability can be achieved by keeping the XCoM within the
base of support (BoS), where the XCoM equals PCoM plus VCoM divided by the
eigenfrequency of the body modeled as an inverted pendulum. This stability requirement has
been validated in both standing and walking (Hof, 2008; Hof et al., 2005). The XCoM concept
indicates that gains of position and velocity feedback should have a ratio equal to the pendulum's
eigenfrequency ($\sqrt{g/l}$). However, whether the CNS modulates feedback gains according to this
ratio is unclear. Therefore, we will compare the ratio of position to velocity gains across standing
and walking to the eigenfrequency of the body modeled as an inverted pendulum.

While standing is usually studied as bipedal standing with feet parallel, unipedal standing and
step postures are commonly observed in daily life, and these postures resemble the behaviors in
the single- and double-support phases of walking. These postures have a different BoS (e.g.
standing on one versus two feet) and structural stiffness (e.g., feet parallel versus step
configuration), which will affect stability requirements. Thus, we also applied the feedback
model to unipedal standing and standing in a step posture, to investigate if, and how, feedback
gains and effective delay change.

Based on previous studies (Geng et al., 2025; van Dieen et al., 2024), we hypothesized that: 1)
feedback delays are longer in walking because feedback becomes effective mainly after ground
contact of the leading leg; 2) feedback gains are larger in walking than in standing postures, to
compensate for larger delays (c.f. Foulger et al. (2025);) 3) directional differences exist in
feedback control, with larger gains in the ML than in the AP direction during walking, while in
standing this would depend on the specific standing posture; 4) the ratio of feedback position to
velocity gains in both walking and standing is constant and close to the eigen frequency of the
body modeled as an inverted pendulum.

2. Methods

2.1 Experimental design & data collection

15 healthy young participants (21± 4 yrs, 63 ± 9 kg, 1.70 ± 0.10 m, 13 females, 2 males)
participated in the experiment. The experimental procedure was approved by the ethics
committee of the Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam
(number: VCWE-2018-159). All participants received oral and written explanations of the
procedures and signed written consent before the experiment.

In this study, participants performed one walking trial and three different standing tasks. For
walking, participants first walked on the instrumented dual-belt treadmill (Motek-Force-link,

Amsterdam, Netherlands) with belt speed set at 4.5 km/h until a steady gait was achieved,
followed by an additional 5 min walking trial. All participants reported that this speed felt natural
without requiring excessive effort or causing fatigue. For standing tasks, participants were asked
to perform: 1) Normal standing: stand upright with feet together. 2) Unipedal standing: stand on
the preferred leg. We tested the preferred leg only, because previously no differences were
observed between standing on the dominant and non-dominant leg (Muehlbauer et al., 2014). 3)
Step posture: participants took a natural step as if walking normally at least three times until they
confirmed to have reached the most comfortable step length and width, then maintained that
position with body weight evenly distributed between both feet. Foot positions were marked for
consistency across trials. All standing tasks were kept for 1 min and repeated 5 times in random
order. For each standing trial, participants faced forward and kept their arms naturally at their
sides, trying to keep as still as possible. Sufficient rest time was given between tasks to avoid
fatigue. Trials with stability loss (e.g., ground contact with the non-stance leg in unipedal
standing, excessive arm movement) were discarded and repeated.

121
122 ### *2.2 Data processing*
123
Data collection and processing followed procedures from our previous study (Geng et al., 2025).
In short, we collected whole-body 3D kinematic data and ground reaction forces, and processed
these data using MATLAB (2024a, The MathWorks, Natick, US). Both kinematic and force data
were low-pass filtered using a second-order Butterworth filter with a cut-off frequency of 10 Hz,
then high-pass filtered at 0.1 Hz to remove noise associated with low-frequency postural
adjustments and drift in position on the treadmill. The force data were down-sampled to 50 Hz to
match the kinematics sampling frequency. The whole-body center of mass position (PCoM) was
calculated using a 3D linked segment model developed by Kingma et al. (1996), and the center
of mass velocity (VCoM) was calculated as the time derivative of PCoM. These variables were
further processed differently for standing and walking.

For standing, the PCoM was adjusted by removing its average value (i.e., the reference upright
position). We removed the components due to gravitational acceleration from the total ground
reaction forces (F_{gr}), based on the assumption that the whole body moved as a single inverted
pendulum (IP), in both ML and AP directions, to obtain the corrective ground reaction forces
(F_{corr}) (Geng et al., 2025). Experimental data from second 6 to 50 of each trial were used to fit
the feedback model.

For walking, PCoM, VCoM and forces were time-normalized to the gait cycle, yielding 100
samples per stride (from left heel strike to subsequent left heel strike) and the PCoM was re-
referenced to the mean foot position during the mid-stance phase of each stride. Compared to
standing, removing the gravitational component from total ground reaction forces (F_{gr}) during
walking is more challenging as the inverted pendulum (IP) model is too simple to fully represent

the complex characteristics of walking (Kuo et al., 2005), particularly during the double-stance
phase. Nevertheless, removing the gravity effects may be essential, as the actual corrective
effects might be masked if the gravitational component is not removed, potentially leading to an
underestimation of the feedback. To address this, we used both F_{gr} and F_{corr} to fit the feedback
model. Results supported the use of corrective forces as the model fit was significantly improved
when using this over F_{gr} (Fig.1). Thus, for all reported analyses, we used F_{corr}. The whole 300s
walking data were used to fit the model.

2.3 Feedback model

We fitted the feedback model for standing and walking separately given their time-invariant and
phase-dependent control.

For standing, equation (1) was used:

$$F_{corr_i} = \widehat{K} * PCOM_{i-\hat{\tau}} + \widehat{B} * VCOM_{i-\hat{\tau}} + \epsilon_i \qquad (1)$$

where $\widehat{K}$ and $\widehat{B}$ are the feedback position and velocity gains, ϵ_i represents the residual error, i
represents the current time point, $\hat{\tau}$ represents the delay ranging from 100 to 1000ms (Safavynia
& Ting, 2013).

We evaluated the model's goodness of fit using the coefficient of determination (R^2), and
estimated the gains using ordinary least square regression, fitting F_{corr_i} to the linear model
specified in equation (1) for each delay ($\hat{\tau}$). The delay at which the model achieved the maximum
R^2 was regarded as the feedback delay, and the corresponding $\widehat{K}$ and $\widehat{B}$ were considered as the
best estimates of the actual feedback parameters.

170
171 For walking, equation (2) was applied to estimate the feedback parameters:
172

$$F_{corr_j} = \widehat{K_j} * PCOM_{j-\hat{\tau}} + \widehat{B_j} * VCOM_{j-\hat{\tau}} + \epsilon_j \qquad (2)$$

where $\widehat{K_j}$ and $\widehat{B_j}$ represents the phase-dependent position and velocity gains, ϵ_j represents the
residual error, j represents the current phase (from 51% to 100% gait cycle), $\hat{\tau}$ represents the
feedback delay ranging from 1% to 50% gait cycle.

Model fit was estimated using the same method as in standing. Differently, we selected the time
lag that yielded the most negative weighted sum of position and velocity gains averaged over all

stance phase as the feedback delay. Under this delay, the position and velocity gains that
produced the largest negative weighted gains were used as the peak gains. Because peak
parameters represent control at a specific phase and do not reflect the overall control strategy
during the whole stance phase, we additionally calculated the mean model fit and mean feedback
gains across the whole stance phase. These mean values were used to allow direct comparison
with standing. For clarity, the peak and mean parameters are referred to as walking peak and
walking, respectively.

The feedback delay ($\hat{\tau}$), and gains ($\hat{K}$ and $\hat{B}$) were estimated for all trials and subsequently
averaged for statistical analysis. We further performed normalization of position and velocity
gains based on their critical values, i.e., critical stiffness and critical damping. The critical
stiffness (K_{cr}) refers to the minimal stiffness required to stabilize the system which was
calculated by equation (3). The critical damping refers to the damping value under which the
system returns to the target position without overshoot and was calculated by equation (4):

193
$$K_{cr} = m * g * l \tag{3}$$

194
$$B_{cr} = 2 * \sqrt{\hat{K} * m} \tag{4}$$

Where the m is body mass, g is the gravitational acceleration ($9.81 \ m/s^2$), l is the center of
mass height, $\hat{K}$ is the current stiffness.

For the position gain, the estimated linear form ($\hat{K}$) was first converted into its angular form by
multiplying by l^2, then normalized by dividing by the critical stiffness (K_{cr}), yielding the
normalized position gain ($\widehat{K_n}$). Similarly, for the velocity gain, the estimated linear gain ($\hat{B}$) was
first converted to rotational damping by multiplying by l^2 and then normalized by dividing by
the critical damping (B_{cr}), yielding the normalized velocity gain ($\widehat{B_n}$). These normalized gains
are unitless and describe how many times greater their rotational values are compared to their
corresponding critical values, providing information about how the body was controlled.

To figure out whether the CNS modulates feedback position and velocity gains based on a
specific relationship as indicated by the concept of the extrapolated center of mass (i.e., based on
the (angular) eigenfrequency), we calculated the ratio between the raw, unnormalized feedback
position and velocity gains ($\hat{K}/\hat{B}$), and further normalized this ratio relative to the IP
eigenfrequency by dividing by $\sqrt{g/l}$, represented as $\widehat{R_n}$. If indeed position and velocity gains are
modulated according to the extrapolated center of mass concept, we expect that $\widehat{R_n} \approx 1$.

3. Statistical Analysis

We examined the effects of Task and Direction on model fit and feedback control using a two-
way repeated measures ANOVA with Task (walking peak and/or walking, normal standing,

unipedal standing, and step posture) and Direction (ML, AP) as within-participant factors to
compare feedback parameters.

Post-hoc pairwise comparisons with Bonferroni correction were conducted when a significant
main effect of Task, Direction or an interaction was found. All statistical analyses were
performed in MATLAB (2024a, The MathWorks, Natick, US). Statistical significance was set at
$\alpha = .05$. Results are presented as mean $\pm$ standard deviation (mean $\pm$ SD).

4. Results

Our feedback model showed good performance with R^2 values ranging from 0.55 to 0.95 across
all tasks, directions and participants. All feedback parameters ($\hat{\tau}, \widehat{K_n}, \widehat{B_n}$) exhibited significant
main effects of Task and Direction, as well as significant Task $*$ Direction interactions, except
the ratio of feedback position to velocity gains ($\widehat{R_n}$), which showed only main effects of Task
and Direction (**Fig. 1**, **Table 1**). These results suggested that the feedback control differed across
tasks and directions.

From the data (**Fig. 1**) it was evident that walking peak gains differed significantly from that in
walking and standing, whereas the walking gains were more comparable to standing.
Considering that the peak gains represent a strategy at a specific moment within the stance phase,
we deemed it less appropriate for direct comparison with standing, therefore, we report the
results in two sections. In the first section, we focus on the results in walking, including the
comparisons between using total ground reaction forces and corrective forces as model outputs,
and examining the differences between peak and mean gains across the stance phase. In the
second section, we focus on the results between walking (mean gains) and standing postures, to
evaluate the similarities and differences between dynamic and static conditions.

236

Fig. 1. Model fit and estimated feedback parameters of all tasks. A) model fit, B) feedback delay, C) normalized position gain, D) normalized velocity gain, E) normalized position to velocity ratio. Box-plot shows estimated values across all participants, with central line indicating the median, box edges showing the interquartile range, and whiskers representing the non-outlier range. Dots represent individual participant's data, with green for ML and orange for the AP direction, grey lines connect the data from same participant. The blue dot line in panel E represent y=1. X-axis labels donate: WP: walking peak; Walking: walking mean; NS: normal standing; US: unipedal standing; SP: step posture.

Table 1. Statistical results across tasks. The asterisks indicating levels of significance, where * indicates p < 0.05 and ** indicates p<0.001.

variables	factor	df	F	p	η^2_p
R^2	Task	3,42	21.16	<0.001**	0.60
	Direction	1,14	7.17	0.018*	0.34
	Task * Direction	3,42	59.92	<0.001**	0.81
$\hat{\tau}$	Task	3,42	78.23	<0.001**	0.85
	Direction	1,14	29.30	<0.001**	0.68
	Task * Direction	3,42	54.81	<0.001**	0.80
$\widehat{K_n}$	Task	4,56	10.92	<0.001**	0.44
	Direction	1,14	27.84	<0.001**	0.67
	Task * Direction	4,56	20.00	<0.001**	0.59
$\widehat{B_n}$	Task	4,56	113.44	<0.001**	0.89
	Direction	1,14	43.84	<0.001**	0.76
	Task * Direction	4,56	26.99	<0.001**	0.66
$\widehat{R_n}$	Task	4,56	27.08	<0.001**	0.66
	Direction	1,14	10.93	0.005*	0.44
	Task * Direction	4,56	1.69	0.17	0.11

246

247

4.1 Feedback results in walking

4.1.1 Fitting model using total forces vs. corrective forces

The model fit (R^2) significantly improved when using corrective forces compared to total ground reaction forces in both ML and AP direction (**Fig. 2**, **Table**). In both directions, peak gains occurred in the early stance phase (~60% gait cycle), regardless of using the total ground reaction forces or corrective forces. Notably, using the corrective forces indicated continuous position feedback, which was masked when using total forces (**Fig. 2**).

4.1.2 Peak gains vs. mean gains

Based on the results using corrective forces, the walking feedback delays were 46±5% and 23±13% of the gait cycle in the ML and AP directions, corresponding to 504±54ms and 250±137ms. The feedback delay in the ML direction was significantly longer than in the AP direction (p<.001) (**Table 1**, **Fig. 2**).

For the position gains ($\widehat{K_n}$), the peak value was significantly larger than the mean value in the ML direction (p=.001), while no significant difference was observed in the AP direction (p=.14). Moreover, both peak and mean position gains showed larger values in the ML direction than the AP direction (p<.001). The peak velocity gains ($\widehat{B_n}$) were consistently larger than the mean values in both directions (p<.001), and both peak (p=.009) and mean (p<.001) values were significantly larger in the AP direction than in the ML direction. Consistent with the concept of extrapolated center of mass, the ratio of the mean position to velocity gains ($\widehat{R_n}$) was not significantly different from 1. However, $\widehat{R_n}$ of the peak gains was significantly smaller than 1 and significantly smaller than of the mean gains in both ML direction (p<.001) and AP direction (p=.009). $\widehat{R_n}$ of mean gains in the ML direction was significantly larger than in the AP direction (p<.001), whereas no significant directional differences were observed for $\widehat{R_n}$ of the peak gains (p=.98).

Fig. 2. Feedback models identified in walking with the total (panel A) and corrective ground reaction forces (panel B) as predicted variables. From left to right: model fit (R^2), normalized feedback position gains ($\widehat{K_n}$) and velocity gains ($\widehat{B_n}$) across the gait cycle. The green and orange lines represent the averaged results (over participants) in ML and AP direction, respectively. The dashed areas indicate the 95% confidence intervals. The horizontal lines in the left panels indicate when the model fit was significant for each individual participant (p<.05).

4.2 feedback control across tasks

In the comparison of standing and walking we report, for walking, values based on averages over the stance phase only.

4.2.1 Model fit

Post-hoc simple effects showed that, in the ML direction, normal standing had a significantly better model fit than all the other tasks (p<.001), followed by the step posture for which model fit was significantly better than for walking and unipedal standing (p<.001). No differences were found between walking and unipedal standing in this direction (p=1). In the AP direction, the model fit for step posture was significantly lower than for all the other tasks (p<.05). Moreover, unipedal standing yielded a significantly lower model fit than normal standing (p<.001), but comparable to walking (p=1). No significant differences were observed between walking and normal standing (p=.07).

When comparing directions within the same task, walking and unipedal standing showed a significantly better model fit in the AP direction than in the ML direction (p<.001). In contrast, in the step posture, the model fit was better in the ML direction than in the AP direction (p<.001). No directional differences were observed in normal standing (p=.18).

4.2.2 Feedback delay

In line with our hypothesis, the post-hoc simple effects showed that the feedback delay was significantly longer in walking than in standing postures in both directions (p<.001). Among standing postures, in the ML direction, unipedal standing had a significantly shorter delay than normal standing(p<.001) and step posture (p<.001), while there were no differences between normal standing and step posture (p=1). In the AP direction, normal standing had a significantly longer delay than in unipedal standing (p<.001) and step posture (p<.001), while no differences were observed between unipedal standing and step posture (p=.06).

Comparing directions, significantly longer delays in the ML than in the AP direction were found in walking (p<.001) and step posture (p<.001), while the opposite was found in normal standing (p<.001) and unipedal standing (p<.001).

4.2.3 Feedback position gain

In the ML direction, the position gains in all tasks exceeded their critical values with $\widehat{K_n}$< -1, and no significant differences were observed between walking and all standing postures (p>.05). Within standing postures, unipedal standing showed significantly larger position gains than both normal standing (p<.001) and step posture (p<.001). In the AP direction, the position gain in walking was slightly below its critical value and was significantly smaller than that of all the standing postures (p<.001), and the position gains of all standing tasks exceeded their critical values. Among standing tasks, the step posture had the largest position gain, followed by unipedal standing, with normal standing exhibiting a significantly smaller position gain.

The step posture showed significantly larger position gains in the AP than in the ML direction (p<.001), whereas the opposite pattern was found in the other tasks (p<.001).

4.2.4 Feedback velocity gain

In contrast to the position gain, the velocity gains were all below critical damping ($\widehat{B_n}$> -1), and the velocity gain in walking was significantly larger than in standing postures in both directions (p<.001). No significant differences were observed among standing postures (p>.05).

The directional differences within task were consistent, with velocity gains larger in the AP direction than in the ML direction (p<.05).

4.2.5 Ratio of feedback position to velocity gains

All standing tasks showed a normalized ratio ($\widehat{R_n}$) significantly larger than 1, whereas in walking it was not significantly different from 1. No significant task differences were observed (p>.05), but the ratio was consistently larger in the ML direction than in the AP direction (p =.005).

5. Discussion

This study is the first to apply the same model to estimate and compare feedback control during unperturbed walking and standing. Both walking and standing showed good model fits with mean values higher than 0.69, supporting the use of this model to estimate and compare feedback control without external perturbations. Results showed that the phase-dependent characteristics in walking indeed led to significant variations in feedback control over the gait cycle. Accordingly, we used the mean values of walking, representing the overall control strategy during the stance phase, for comparison with standing. For clarity, in the following we refer to these mean values unless explicitly stated otherwise.

Our results showed significantly larger velocity control in walking than in standing in both directions, whereas the position control in walking was significantly smaller than standing in the AP direction. Despite these differences, all tasks exhibited mostly above critical stiffness and under-damped control, with as only exception position gains in the AP direction in walking, which were slightly below critical stiffness. Notably, both walking and standing exhibited a position-to-velocity gain ratio close to the $\sqrt{g/l}$, as predicted by the extrapolated center of mass (XCoM) concept. These findings of feedback parameters which varied significantly across tasks and directions, and consistent relative position and velocity gains indicate a common strategy of sensorimotor control across both dynamic and static conditions. We will further discuss these results below.

5.1 Feedback control in walking

In line with previous results (van Dieën et al., 2025), our results showed phase-dependent feedback control with peak control occurring during the early stance phase. The significantly improved model fit (Fig. 2), as well as the observed continuous control in both directions suggest that using corrective forces allows for more accurate estimation of feedback parameters in walking.

Like Afschrift et al. (2021), we found that gains were modulated across the gait cycle, with velocity control being more phase-dependent than position control in the AP direction. However, our results showed peak control in early stance, while Afschrift et al. (2021) reported most pronounced control during mid-stance. The similarity likely reflects a common control strategy: the feedback control is performed based on preceding CoM position and velocity information. The difference in timing is likely due to methodological differences: we used corrective forces as the model output and selected the feedback delay based on the variation of gains across the gait cycle, whereas Afschrift et al. (2021) used the EMG and ankle torque as model output and used a constant feedback delay. Additionally, our study didn't apply perturbations. While we observed

more phase-dependent control for position in the ML direction than in the AP direction, Afschrift
et al. (2021) only investigated AP control.

Both mean and peaks gains showed stronger control of position in the ML direction, while in the
AP direction velocity was more tightly controlled. These directional differences are consistent
with the foot placement model (Wang & Srinivasan, 2014), which demonstrated that ML foot
placement control relies more on CoM position, whereas AP foot placement relies more on CoM
velocity. Peak gains around early stance were expected, as this phase would impose high
demands, especially to control AP velocity, where over-damped control appears to be used to
avoid overshoot and oscillations to allow smooth and stable forward progression. The mean
gains reflect control over the whole stance phase. Since velocity fluctuations in the remainder of
the stance phase are smaller, an overall under-damped control may suffice.

5.2 Comparison of feedback control across tasks

5.2.2 Feedback delay

As hypothesized, the estimated feedback delay was significantly longer in walking than in
standing. This was expected as in walking, feedback corrections are mainly applied after foot
contact, while in standing stability control is continuous (O'Connor & Kuo, 2009). Directional
differences in walking appear to reflect different stability demands and control strategies in the
ML and AP directions. In the ML direction, stability control has been shown to be more
challenging (Harter et al., 2024; Kongsuk et al., 2019; Madehkhaksar et al., 2018; O'Connor &
Kuo, 2009) and rely more on foot placement (Bruijn & van Dieen, 2018; Harter et al., 2024),
while in the AP direction, stability control has been suggested to rely more on the ankle strategy
(Gatev et al., 1999; Reimann et al., 2018), which can take effect during the stance phase and thus
leads to a shorter delay.

The differences in estimated feedback delays among standing postures can be attributed to
different biomechanical and neurophysiological characteristics. As suggested by our previous
work (Geng et al., 2025), increased intrinsic stiffness leads to a negative bias on estimated
feedback delays. The observed shorter delays in unipedal standing may thus be explained by the
increased intrinsic stiffness due to the higher co-contraction levels (Amiri & Kearney, 2019;
Baratta et al., 1988; Fok et al., 2021; Sakanaka et al., 2018; Sakanaka et al., 2021; Wu et al.,
2024). In addition to biasing the estimates of the delay due to differences in intrinsic stiffness,
higher muscle activity can be expected to cause a shorter electro-mechanical delay (Vos et al.,
1991), contributing to a shorter delay in unipedal standing and step postures. Non-neutral ankle
angles in the step posture coincide with higher intrinsic stiffness compared to normal standing,
due to the non-linear relationship between ankle angle and elastic parameters (Weiss et al.,

1986). In addition, the geometric configuration of the legs in the step posture with the non-
aligned ankle axes will yield 'structural stiffness', enhancing the overall stiffness.

5.2.3 Feedback gains

5.2.3.1 Position gains

Position gains for all tasks exceeded critical stability (i.e., $\widehat{K_n} < -1$) except for walking in the AP
direction. In the AP direction, the walking position gain was slightly below the critical stiffness,
suggesting instability in this direction, likely prevented by changes in the base of support
(stepping). The peak walking gains typically above critical stiffness (i.e., $\widehat{K_n} < -1$) suggests that
indeed these changes are used to maintain a stable gait pattern.

Among standing postures, unipedal standing and step posture had significantly larger position
gains than normal standing, consistent with their presumed higher intrinsic stiffness as
mentioned above. Interestingly, position gains in unipedal standing were approximately twice the
per leg values observed in normal standing, indicating that the control system modulates
feedback gains per leg and highlighting its adaptability to changed stability demands.

5.2.3.2 Velocity gains

The velocity gains revealed an under-damped control in both walking and standing, and there
was a clear task-dependent modulation, with velocity gains significantly larger in walking than in
standing. The under-damped velocity control observed in standing aligned well with the findings
that described human upright posture as an underdamped system (Winter, 1995; Winter et al.,
1998), likely reflecting that our control system allows some oscillatory behavior during standing.

5.3 Ratio of position to velocity gains

According to the XCoM concept, control of pendular movement requires the ratio of position to
velocity gains to equal the inverted pendulum eigenfrequency, yielding a normalized ratio ($\widehat{R_n}$)
of 1. The $\widehat{R_n}$ in walking was not significantly different from 1, which is as expected, whereas
this value in standing was significantly larger than 1. However, this ratio between walking and
standing was not significantly different, and both ratios remained close to the eigenfrequency
value.

The approximation of these ratios to the eigenfrequency is consistent with the view that human
body can be modelled as a single inverted pendulum model (Winter, 1995). The deviations from
1, particularly during standing, likely result from the human body not behaving as a perfect
pendulum, but instead stabilizing itself through multi-link segments coordination(Hsu et al.,
2007). The comparable ratios observed in walking and standing suggest that, at least in

unperturbed conditions, the sensorimotor system maintains a relative consistent weighting
between position and velocity feedback. Such consistency may simply arise from the shared
whole-body dynamics of human body, and may also facilitate a smooth transition between
dynamic and static tasks.

6. Limitations

Several limitations should be noted. While we used a unified feedback model to estimate
feedback control in walking and standing, the procedure to fit the model differed between tasks.
Specifically, the method to select the feedback delay and corresponding gains was different. In
addition, as reported in our previous study (Geng et al., 2025), intrinsic properties lead to errors
in estimating feedback parameters in standing. How this will influence results in waking is
unclear. Moreover, walking might introduce more soft tissue artifacts in the measurement of
kinematics, potentially affecting the parameter estimation. Nonetheless, the estimated parameters
aligned well with the task-specific characteristics. Finally, we examined only one gait speed in
this study, whether and how the observed feedback strategy will change with speed needs further
examination.

7. Conclusion

Our results demonstrated that while the feedback gains changed significantly across tasks and
directions, the feedback position gains were consistently maintained close to critical stiffness,
while the velocity was controlled at under-damped level. Notably, the relative ratio between
position and velocity gains in both walking and standing remained close to the eigenfrequency,
which might reflect a robust control strategy for stabilizing the human body as an inverted-
pendulum.

References

Afschrift, M., De Groote, F., & Jonkers, I. (2021). Similar sensorimotor transformations
control balance during standing and walking. *PLoS Comput Biol*, *17*(6), e1008369.
https://doi.org/10.1371/journal.pcbi.1008369
Amiri, P., & Kearney, R. E. (2019). Ankle intrinsic stiffness changes with postural sway. *J*
Biomech, *85*, 50-58. https://doi.org/10.1016/j.jbiomech.2019.01.009
Baratta, R., Solomonow, M., Zhou, B. H., Letson, D., Chuinard, R., & D'Ambrosia, R. (1988).
Muscular coactivation. The role of the antagonist musculature in maintaining knee
stability. *Am J Sports Med*, *16*(2), 113-122.
https://doi.org/10.1177/036354658801600205

Bauby, C. E., & Kuo, A. D. (2000). Active control of lateral balance in human walking. *J Biomech*, *33*(11), 1433-1440. https://doi.org/10.1016/s0021-9290(00)00101-9

Blaszczyk, J. W., Beck, M., & Sadowska, D. (2014). Assessment of postural stability in young healthy subjects based on directional features of posturographic data: vision and gender effects. *Acta Neurobiol Exp (Wars)*, *74*(4), 433-442. https://doi.org/10.55782/ane-2014-2006

Bruijn, S. M., & van Dieen, J. H. (2018). Control of human gait stability through foot placement. *J R Soc Interface*, *15*(143). https://doi.org/10.1098/rsif.2017.0816

Coleman, M. J., & Ruina, A. (1998). An uncontrolled walking toy that cannot stand still. *Physical Review Letters*, *80*(16), 3658-3661. https://doi.org/DOI 10.1103/PhysRevLett.80.3658

Fok, K. L., Lee, J. W., Unger, J., Chan, K., Musselman, K. E., & Masani, K. (2021). Co-contraction of ankle muscle activity during quiet standing in individuals with incomplete spinal cord injury is associated with postural instability. *Sci Rep*, *11*(1), 19599. https://doi.org/10.1038/s41598-021-99151-w

Forbes, P. A., Chen, A., & Blouin, J. S. (2018). Sensorimotor control of standing balance. *Handb Clin Neurol*, *159*, 61-83. https://doi.org/10.1016/B978-0-444-63916-5.00004-5

Foulger, L. H., Liu, X., Nasrabadi, A. M., Qiao, C. Z., Carpenter, M. G., Wu, L. C., & Blouin, J. S. (2025). Learning to stand with delays alters sensorimotor control but does not cause instability when returning to natural balance. *J Physiol*, *603*(19), 5629-5653. https://doi.org/10.1113/JP288632

Gatev, P., Thomas, S., Kepple, T., & Hallett, M. (1999). Feedforward ankle strategy of balance during quiet stance in adults. *J Physiol*, *514 (Pt 3)*(Pt 3), 915-928. https://doi.org/10.1111/j.1469-7793.1999.915ad.x

Geng, Y., Bruijn, S. M., & van Dieën, J. H. (2025). Feedback control stabilizing the center of mass can be identified in unperturbed, upright standing. *bioRxiv*, 2024.2012.2020.629656. https://doi.org/10.1101/2024.12.20.629656

Harter, M. J., Redfern, M. S., Sparto, P. J., & Geyer, H. (2024). Modelling strategies supplemental to foot placement for frontal-plane stability in walking. *J R Soc Interface*, *21*(218), 20240191. https://doi.org/10.1098/rsif.2024.0191

Hof, A. L. (2008). The 'extrapolated center of mass' concept suggests a simple control of balance in walking. *Hum Mov Sci*, *27*(1), 112-125. https://doi.org/10.1016/j.humov.2007.08.003

Hof, A. L., Gazendam, M. G., & Sinke, W. E. (2005). The condition for dynamic stability. *J Biomech*, *38*(1), 1-8. https://doi.org/10.1016/j.jbiomech.2004.03.025

Hsu, W. L., Scholz, J. P., Schoner, G., Jeka, J. J., & Kiemel, T. (2007). Control and estimation of posture during quiet stance depends on multijoint coordination. *Journal of Neurophysiology*, *97*(4), 3024-3035. https://doi.org/10.1152/jn.01142.2006

Jakubowski, K. L., Martino, G., Beck, O. N., Sawicki, G. S., & Ting, L. H. (2024). Center of mass states render multi-joint torques throughout standing balance recovery. *bioRxiv*. https://doi.org/10.1101/2024.08.14.607976

Kingma, I., deLooze, M. P., Toussaint, H. M., Klijnsma, H. G., & Bruijnen, T. B. M. (1996). Validation of a full body 3-D dynamic linked segment model. *Human Movement Science, 15*(6), 833-860. https://doi.org/Doi 10.1016/S0167-9457(96)00034-6

Koelewijn, A. D., & Van Den Bogert, A. J. (2022). Antagonistic co-contraction can minimize muscular effort in systems with uncertainty. *PeerJ, 10*, e13085. https://doi.org/10.7717/peerj.13085

Kongsuk, J., Brown, D. A., & Hurt, C. P. (2019). Dynamic stability during increased walking speeds is related to balance confidence of older adults: a pilot study. *Gait Posture, 73*, 86-92. https://doi.org/10.1016/j.gaitpost.2019.06.018

Kuo, A. D. (1999). Stabilization of lateral motion in passive dynamic walking. *International Journal of Robotics Research, 18*(9), 917-930. https://doi.org/Doi 10.1177/02783649922066655

Kuo, A. D., & Donelan, J. M. (2010). Dynamic principles of gait and their clinical implications. *Phys Ther, 90*(2), 157-174. https://doi.org/10.2522/ptj.20090125

Kuo, A. D., Donelan, J. M., & Ruina, A. (2005). Energetic consequences of walking like an inverted pendulum: step-to-step transitions. *Exerc Sport Sci Rev, 33*(2), 88-97. https://doi.org/10.1097/00003677-200504000-00006

Madehkhaksar, F., Klenk, J., Sczuka, K., Gordt, K., Melzer, I., & Schwenk, M. (2018). The effects of unexpected mechanical perturbations during treadmill walking on spatiotemporal gait parameters, and the dynamic stability measures by which to quantify postural response. *PLoS One, 13*(4), e0195902. https://doi.org/10.1371/journal.pone.0195902

Mcgeer, T. (1990). Passive Dynamic Walking. *International Journal of Robotics Research, 9*(2), 62-82. https://doi.org/Doi 10.1177/027836499000900206

Muehlbauer, T., Mettler, C., Roth, R., & Granacher, U. (2014). One-leg standing performance and muscle activity: are there limb differences? *J Appl Biomech, 30*(3), 407-414. https://doi.org/10.1123/jab.2013-0230

O'Connor, S. M., & Kuo, A. D. (2009). Direction-dependent control of balance during walking and standing. *Journal of Neurophysiology, 102*(3), 1411-1419. https://doi.org/10.1152/jn.00131.2009

Reimann, H., Fettrow, T., Thompson, E. D., & Jeka, J. J. (2018). Neural Control of Balance During Walking. *Front Physiol, 9*, 1271. https://doi.org/10.3389/fphys.2018.01271

Safavynia, S. A., & Ting, L. H. (2013). Long-latency muscle activity reflects continuous, delayed sensorimotor feedback of task-level and not joint-level error. *Journal of Neurophysiology, 110*(6), 1278-1290. https://doi.org/10.1152/jn.00609.2012

Sakanaka, T. E., Gill, J., Lakie, M. D., & Reynolds, R. F. (2018). Intrinsic ankle stiffness during standing increases with ankle torque and passive stretch of the Achilles tendon. *PLoS One, 13*(3), e0193850. https://doi.org/10.1371/journal.pone.0193850

Sakanaka, T. E., Lakie, M., & Reynolds, R. F. (2021). Individual differences in intrinsic ankle stiffness and their relationship to body sway and ankle torque. *PLoS One, 16*(1), e0244993. https://doi.org/10.1371/journal.pone.0244993

Shanbhag, J., Fleischmann, S., Wechsler, I., Gassner, H., Winkler, J., Eskofier, B. M., Koelewijn, A. D., Wartzack, S., & Miehling, J. (2024). A sensorimotor enhanced

neuromusculoskeletal model for simulating postural control of upright standing. *Front Neurosci, 18*, 1393749. https://doi.org/10.3389/fnins.2024.1393749

van Dieën, J. H., Bruijn, S., Lemaire, K. K., & Kistemaker, D. A. (2025). Simultaneous stabilizing feedback control of linear and angular momentum in

human walking. https://doi.org/10.1101/2025.01.09.632072

van Dieen, J. H., Bruijn, S. M., & Afschrift, M. (2024). Assessment of stabilizing feedback control of walking: A tutorial. *J Electromyogr Kinesiol, 78*, 102915. https://doi.org/10.1016/j.jelekin.2024.102915

Vos, E. J., Harlaar, J., & van Ingen Schenau, G. J. (1991). Electromechanical delay during knee extensor contractions. *Med Sci Sports Exerc, 23*(10), 1187-1193. https://www.ncbi.nlm.nih.gov/pubmed/1758296

Wang, Y., & Srinivasan, M. (2014). Stepping in the direction of the fall: the next foot placement can be predicted from current upper body state in steady-state walking. *Biol Lett, 10*(9). https://doi.org/10.1098/rsbl.2014.0405

Weiss, P. L., Kearney, R. E., & Hunter, I. W. (1986). Position dependence of ankle joint dynamics--I. Passive mechanics. *J Biomech, 19*(9), 727-735. https://doi.org/10.1016/0021-9290(86)90196-x

Welch, T. D., & Ting, L. H. (2008). A feedback model reproduces muscle activity during human postural responses to support-surface translations. *Journal of Neurophysiology, 99*(2), 1032-1038. https://doi.org/10.1152/jn.01110.2007

Winter, D. A. (1995). Human balance and posture control during standing and walking. *Gait & Posture, 3*(4), 193-214. https://doi.org/10.1016/0966-6362(96)82849-9

Winter, D. A., Patla, A. E., Prince, F., Ishac, M., & Gielo-Perczak, K. (1998). Stiffness control of balance in quiet standing. *Journal of Neurophysiology, 80*(3), 1211-1221. https://doi.org/10.1152/jn.1998.80.3.1211

Winter, D. A., Prince, F., Frank, J. S., Powell, C., & Zabjek, K. F. (1996). Unified theory regarding A/P and M/L balance in quiet stance. *Journal of Neurophysiology, 75*(6), 2334-2343. https://doi.org/10.1152/jn.1996.75.6.2334

Wu, F., Zhang, J., Yang, K., Li, Y., Li, Z., Li, T., & Li, J. (2024). Surface electromyography analysis of ankle flexor and extensor activity under different standing stability levels. *J Orthop Surg (Hong Kong), 32*(2), 1022553641258336. https://doi.org/10.1177/10225536241258336

Supplementary materials

 Table S1. Feedback results in walking and standing (mean±SD).

variables	directions	Walking (using total forces)	Walking peak	Walking mean	Normal standing	Unipedal standing	Step posture
R^2	ML	0.27±0.10	0.54±0.21	0.69±0.10	0.89±0.03	0.72±0.05	0.84±0.03
	AP	0.40±0.10	0.68±0.11	0.83±0.08	0.90±0.03	0.83±0.04	0.69±0.08
$\hat{\tau}$	ML	424±73	504±54	-	199±18	155±27	196±20
	AP	139±28	250±137	-	233±22	191±18	165±21
$\widehat{K_n}$	ML	-0.25±0.26	-1.46±0.35	-1.24±0.23	-1.25±0.03	-1.39±0.09	-1.25±0.05
	AP	-0.14±0.15	-1.06±0.17	-0.94±0.06	-1.15±0.06	-1.23±0.06	-1.36±0.12
$\widehat{B_n}$	ML	-3.32±6.15	-0.83±0.19	-0.53±0.10	-0.37±0.05	-0.39±0.08	-0.41±0.06
	AP	-6.86±5.76	-1.81±0.43	-0.65±0.09	-0.44±0.05	-0.44±0.05	-0.49±0.07
$\widehat{R_n}$	ML	0.25±0.31	0.69±0.18	1.09±0.25	1.48±0.42	1.52±0.71	1.26±0.28
	AP	0.05±0.10	0.28±0.09	1.08±0.76	1.12±0.11	1.16±0.12	1.09±0.13

Fig. 1. Model fit and estimated feedback parameters of all tasks. A) model fit, B) feedback delay, C) normalized position gain, D) normalized velocity gain, E) normalized position to velocity ratio. Box-plot shows estimated values across all participants, with central line indicating the median, box edges showing the interquartile range, and whiskers representing the non-outlier range. Dots represent individual participant's, with green for ML and orange for the AP direction, grey lines connect the data from same participant. The blue dot line in panel E represent y=1. X-axis labels donate: WP: walking peak; Walking: walking mean; NS: normal standing; US: unipedal standing; SP: step posture.

Fig. 2. Feedback models identified in walking with the total (panel A) and corrective ground reaction forces (panel B) as predicted variables. From left to right: model fit (R^2), normalized feedback position gains ($\widehat{K_n}$) and velocity gains ($\widehat{B_n}$) across the gait cycle. The green and orange lines represent the averaged results (over participants) in ML and AP direction, respectively. The dashed areas indicate the 95% confidence intervals. The horizontal lines in the left panels indicate when the model fit was significant for each individual participant (p<.05).

Table 1. Statistical results across tasks. The asterisks indicating levels of significance, where * indicates p < 0.05 and ** indicates p<0.001.

variables	factor	df	F	p	η^2_p
R^2	Task	3,42	21.16	<0.001**	0.60
	Direction	1,14	7.17	0.018*	0.34
	Task * Direction	3,42	59.92	<0.001**	0.81
$\hat{t}$	Task	3,42	78.23	<0.001**	0.85
	Direction	1,14	29.30	<0.001**	0.68
	Task * Direction	3,42	54.81	<0.001**	0.80
$\widehat{K_n}$	Task	4,56	10.92	<0.001**	0.44
	Direction	1,14	27.84	<0.001**	0.67
	Task * Direction	4,56	20.00	<0.001**	0.59
$\widehat{B_n}$	Task	4,56	113.44	<0.001**	0.89
	Direction	1,14	43.84	<0.001**	0.76
	Task * Direction	4,56	26.99	<0.001**	0.66
$\widehat{R_n}$	Task	4,56	27.08	<0.001**	0.66
	Direction	1,14	10.93	0.005*	0.44
	Task * Direction	4,56	1.69	0.17	0.11