Shared feedback control principles across standing and walking

walking

Yang Geng, Jaap H. van Dieën, Sjoerd M. Bruijn*

Department of Human Movement Sciences, Faculty of Behavioral and Movement

Sciences, Amsterdam Movement Sciences, Vrije Universiteit Amsterdam,

Amsterdam, The Netherlands

* E-mail: s.m.bruijn@vu.nl

Abstract

Feedback control is required to maintain stability in both walking and standing, whereas the control parameters may differ due to the inherently different characteristics of these tasks. How feedback is adapted between tasks is unclear. To address this, we applied a unified feedback model, relating preceding center of mass position and velocity information to corrective ground reaction forces, to estimate and compare the control parameters between walking and standing.

15 healthy young participants (21± 4 yrs, 63 ± 9 kg, 1.70 ± 0.10 m, 13 females, 2 males) participated, they walked at 4.5 km/h for 5 minutes, and performed 3 different standing tasks: normal standing, unipedal standing, and step posture for 1 minute, repeated 5 times. Whole-body kinematics and ground reaction forces were collected and used to fit the feedback model and estimate feedback delay and gains. Results showed good model fits for both walking and standing with mean R^2-values higher than 0.69. The feedback delay was significantly longer in walking than in standing. While the feedback position gains and velocity gains varied significantly across tasks and directions, position gains were mostly above critical stiffness and velocity gains all at under-damped level, with as only exception that position gains in the AP direction in walking, which were slightly below critical stiffness. The ratio of position to velocity gains in all tasks remained consistently

close to the eigenfrequency ($\sqrt{g/l}$) as predicted by the extrapolated center of mass concept.

Our findings suggest a robust control strategy in which the sensorimotor system maintains a relative consistent weighting between position and velocity feedback for walking and standing.

1. Introduction

Walking and standing are two fundamental motor behaviors in human daily life. Both require sophisticated neuromuscular control to maintain an upright body orientation against the effects of unpredictable perturbations which are amplified by gravity. Such stabilization is achieved through coordination of intrinsic mechanics and feedback control. Intrinsic mechanics provide immediate resistance to perturbations based on segments' inertia and joints' stiffness and damping. The latter can be modulated by muscle co-activation to deal with expected perturbations or uncertainty, but maintaining high levels of co-contraction is energetically costly [1]. In contrast, feedback control is adaptive and generates corrective responses selectively based on sensory information [2, 3].

Given the different temporal and biomechanical characteristics of standing and walking,

feedback control parameters can be expected to differ between these motor tasks,

reflecting task-specific modulations. Standing is a quasi-static behavior, while walking is

a dynamic, cyclic motion. This temporal difference suggests distinct feedback strategies:

time-invariant feedback control in standing but a phase-dependent control in walking.

Additionally, in standing, the anatomical constraints and smaller base of support (BoS)

make stabilization more challenging in the anteroposterior (AP) direction than in the

mediolateral (ML) direction [4-6]. However, during walking, the stability requirements

shift. Stabilization of AP movement benefits from passive mechanics [7-9], while the ML

movement must be actively stabilized [10, 11].

Several feedback models have been developed to identify feedback control in

standing [3, 12-14] and walking [15]. These models indicate that sensory estimates

of the center of mass (CoM) kinematics are used to modulate corrective responses

(e.g., EMG, ankle torque, foot placement) to stabilize standing and walking.

However, identification of these models requires perturbations and measurements

of physiological signals (e.g., EMG), which poses high experimental demands and

limits practical application. As an alternative, van Dieen, Bruijn [16] proposed a

feedback model that can identify feedback control in unperturbed walking by

fitting a linear relationship between the delayed extrapolated center of mass

(XCoM) and horizontal ground reaction forces, without external perturbations or

physiological measurements. More recently, we refined this feedback model by

using the CoM position and velocity information as separate predictors instead of

the XCoM, and validated that this modified model can be applied to assess

feedback control in standing [17]. Until now, these models were derived under the

specific context being studied (i.e., either standing or walking), which leaves a gap

in our understanding of how stability control is adapted between standing and

walking.

In the present study, we addressed this gap by applying the aforementioned

feedback model to unperturbed standing and walking. Using this model, we can

estimate feedback position and velocity gains and the effective delay, helping us to

understand how the CNS modulates feedback control in these tasks, while

operating under the same fundamental control principle: generating corrective

ground reaction forces based on preceding center of mass position (PCoM) and

velocity (VCoM) information.

Hof, Gazendam [18] described the relative importance of position and velocity for

maintaining dynamic stability. They showed that stability can be achieved by

keeping the XCoM within the base of support (BoS), where the XCoM equals

PCoM plus VCoM divided by the eigenfrequency of the body modeled as an

inverted pendulum. This stability requirement has been validated in both standing

and walking [18, 19]. The XCoM concept indicates that gains of position and

velocity feedback should have a ratio equal to the pendulum's eigenfrequency

($\sqrt{g/l}$). However, whether the CNS modulates feedback gains according to this

ratio is unclear. Therefore, we will compare the ratio of position to velocity gains

across standing and walking to the eigenfrequency of the body modeled as an

inverted pendulum.

While standing is usually studied as bipedal standing with feet parallel, unipedal

standing and step postures are commonly observed in daily life, and these postures

resemble the behaviors in the single- and double-support phases of walking. These

postures have a different BoS (e.g. standing on one versus two feet) and structural

stiffness (e.g., feet parallel versus step configuration), which will affect stability

requirements. Thus, we also applied the feedback model to unipedal standing and

standing in a step posture, to investigate if, and how, feedback gains and effective

delay change.

Based on previous studies [16, 17], we hypothesized that: 1) feedback delays are

longer in walking because feedback becomes effective mainly after ground contact

of the leading leg; 2) feedback gains are larger in walking than in standing

postures, to compensate for larger delays (c.f. Foulger, Liu [20];) 3) directional

differences exist in feedback control, with larger gains in the ML than in the AP

direction during walking, while in standing this would depend on the specific

standing posture; 4) the ratio of feedback position to velocity gains in both walking

and standing is constant and close to the eigen frequency of the body modeled as

an inverted pendulum.

2. Methods

2.1 Experimental design & data collection

15 healthy young participants (21± 4 yrs, 63 ± 9 kg, 1.70 ± 0.10 m, 13 females, 2

males) participated in the experiment. The experimental procedure was approved

by the ethics committee of the Faculty of Behavioural and Movement Sciences,

Vrije Universiteit Amsterdam (number: VCWE-2018-159). All participants

received oral and written explanations of the procedures and signed written consent

before the experiment.

In this study, participants performed one walking trial and three different standing

tasks. For walking, participants first walked on the instrumented dual-belt treadmill

(Motek-Force-link, Amsterdam, Netherlands) with belt speed set at 4.5 km/h until

a steady gait was achieved, followed by an additional 5 min walking trial. All

participants reported that this speed felt natural without requiring excessive effort

or causing fatigue. For standing tasks, participants were asked to perform: 1)

Normal standing: stand upright with feet together. 2) Unipedal standing: stand on

the preferred leg. We tested the preferred leg only, because previously no

differences were observed between standing on the dominant and non-dominant

leg [21]. 3) Step posture: participants took a natural step as if walking normally at

least three times until they confirmed to have reached the most comfortable step

length and width, then maintained that position with body weight evenly

distributed between both feet. Foot positions were marked for consistency across

trials. All standing tasks were kept for 1 min and repeated 5 times in random order.

For each standing trial, participants faced forward and kept their arms naturally at

their sides, trying to keep as still as possible. Sufficient rest time was given

between tasks to avoid fatigue. Trials with stability loss (e.g., ground contact with

the non-stance leg in unipedal standing, excessive arm movement) were discarded

and repeated.

2.2 Data processing

Data collection and processing followed procedures from our previous study [17].

In short, we collected whole-body 3D kinematic data and ground reaction forces,

and processed these data using MATLAB (2024a, The MathWorks, Natick, US).

Both kinematic and force data were low-pass filtered using a second-order

Butterworth filter with a cut-off frequency of 10 Hz, then high-pass filtered at 0.1

Hz to remove noise associated with low-frequency postural adjustments and drift

in position on the treadmill. The force data were down-sampled to 50 Hz to match

the kinematics sampling frequency. The whole-body center of mass position

(PCoM) was calculated using a 3D linked segment model developed by Kingma,

deLooze [22], and the center of mass velocity (VCoM) was calculated as the time

derivative of PCoM. These variables were further processed differently for

standing and walking.

For standing, the PCoM was adjusted by removing its average value (i.e., the

reference upright position). We removed the components due to gravitational

acceleration from the total ground reaction forces (F_{gr}), based on the assumption

that the whole body moved as a single inverted pendulum (IP), in both ML and AP

directions, to obtain the corrective ground reaction forces (F_{corr}) [17].

Experimental data from second 6 to 50 of each trial were used to fit the feedback

model.

For walking, PCoM, VCoM and forces were time-normalized to the gait cycle,

yielding 100 samples per stride (from left heel strike to subsequent left heel strike)

and the PCoM was re-referenced to the mean foot position during the mid-stance

phase of each stride. Compared to standing, removing the gravitational component

from total ground reaction forces (F_{gr}) during walking is more challenging as the

inverted pendulum (IP) model is too simple to fully represent the complex

characteristics of walking [23], particularly during the double-stance phase.

Nevertheless, removing the gravity effects may be essential, as the actual

corrective effects might be masked if the gravitational component is not removed,

potentially leading to an underestimation of the feedback. To address this, we used

both F_{gr} and F_{corr} to fit the feedback model. Results supported the use of

corrective forces as the model fit was significantly improved when using this over

F_{gr} (Fig.1). Thus, for all reported analyses, we used F_{corr}. The whole 300s walking

data were used to fit the model.

2.3 Feedback model

177   We fitted the feedback model for standing and walking separately given their time-

178   invariant and phase-dependent control.

179   For standing, equation (1) was used:

$$F_{corr_i} = \widehat{K} * PCOM_{i-\hat{\tau}} + \widehat{B} * VCOM_{i-\hat{\tau}} + \epsilon_i \qquad (1)$$

where $\widehat{K}$ and $\widehat{B}$ are the feedback position and velocity gains, ϵ_i represents the

residual error, i represents the current time point, $\hat{\tau}$ represents the delay ranging

from 100 to 1000ms [13].

We evaluated the model's goodness of fit using the coefficient of determination

(R^2), and estimated the gains using ordinary least square regression, fitting F_{corr_i}

to the linear model specified in equation (1) for each delay ($\hat{\tau}$). The delay at which

the model achieved the maximum R^2 was regarded as the feedback delay, and the

corresponding $\hat{K}$ and $\hat{B}$ were considered as the best estimates of the actual

feedback parameters.

For walking, equation (2) was applied to estimate the feedback parameters:

194
$$F_{corr_j} = \widehat{K_j} * PCOM_{j-\hat{\tau}} + \widehat{B_j} * VCOM_{j-\hat{\tau}} + \epsilon_j \qquad (2)$$

where $\widehat{K_j}$ and $\widehat{B_j}$ represents the phase-dependent position and velocity gains, ϵ_j

represents the residual error, j represents the current phase (from 51% to 100% gait

cycle), $\hat{\tau}$ represents the feedback delay ranging from 1% to 50% gait cycle.

Model fit was estimated using the same method as in standing. Differently, we

selected the time lag that yielded the most negative weighted sum of position and

velocity gains averaged over all stance phase as the feedback delay. Under this

delay, the position and velocity gains that produced the largest negative weighted

gains were used as the peak gains. Because peak parameters represent control at a

specific phase and do not reflect the overall control strategy during the whole

stance phase, we additionally calculated the mean model fit and mean feedback

gains across the whole stance phase. These mean values were used to allow direct

comparison with standing. For clarity, the peak and mean parameters are referred

to as walking peak and walking, respectively.

The feedback delay ($\hat{\tau}$), and gains ($\hat{K}$ and $\hat{B}$) were estimated for all trials and

subsequently averaged for statistical analysis. We further performed normalization

of position and velocity gains based on their critical values, i.e., critical stiffness

and critical damping. The critical stiffness (K_{cr}) refers to the minimal stiffness

required to stabilize the system which was calculated by equation (3). The critical

damping refers to the damping value under which the system returns to the target

position without overshoot and was calculated by equation (4):

216
$$K_{cr} = m * g * l \tag{3}$$

217
$$B_{cr} = 2 * \sqrt{\hat{K} * m} \tag{4}$$

Where the m is body mass, g is the gravitational acceleration ($9.81\ m/s^2$), l is the

center of mass height, $\hat{K}$ is the current stiffness.

For the position gain, the estimated linear form ($\hat{K}$) was first converted into its

angular form by multiplying by l^2, then normalized by dividing by the critical

stiffness (K_{cr}), yielding the normalized position gain ($\widehat{K_n}$). Similarly, for the

velocity gain, the estimated linear gain ($\hat{B}$) was first converted to rotational

damping by multiplying by l^2 and then normalized by dividing by the critical

damping (B_{cr}), yielding the normalized velocity gain ($\widehat{B_n}$). These normalized gains

are unitless and describe how many times greater their rotational values are

compared to their corresponding critical values, providing information about how

the body was controlled.

To figure out whether the CNS modulates feedback position and velocity gains

based on a specific relationship as indicated by the concept of the extrapolated

center of mass (i.e., based on the (angular) eigenfrequency), we calculated the ratio

between the raw, unnormalized feedback position and velocity gains ($\hat{K}/\hat{B}$), and

further normalized this ratio relative to the IP eigenfrequency by dividing by $\sqrt{g/l}$,

represented as $\widehat{R_n}$. If indeed position and velocity gains are modulated according to

the extrapolated center of mass concept, we expect that $\widehat{R_n} \approx 1$.

3. Statistical analysis

We examined the effects of Task and Direction on model fit and feedback control

using a two-way repeated measures ANOVA with Task (walking peak and/or

walking, normal standing, unipedal standing, and step posture) and Direction (ML,

AP) as within-participant factors to compare feedback parameters.

Post-hoc pairwise comparisons with Bonferroni correction were conducted when a

significant main effect of Task, Direction or an interaction was found. All

statistical analyses were performed in MATLAB (2024a, The MathWorks, Natick,

US). Statistical significance was set at $\alpha = .05$. Results are presented as mean $\pm$

standard deviation (mean $\pm$ SD).

4. Results

Our feedback model showed good performance with R^2 values ranging from 0.55

to 0.95 across all tasks, directions and participants. All feedback parameters ($\hat{\tau}$, $\widehat{K_n}$,

$\widehat{B_n}$) exhibited significant main effects of Task and Direction, as well as significant

Task $*$ Direction interactions, except the ratio of feedback position to velocity

gains ($\widehat{R_n}$), which showed only main effects of Task and Direction (Fig 1, Table

1). These results suggested that the feedback control differed across tasks and

directions.

Fig 1. Model fit and estimated feedback parameters of all tasks. A) model fit, B) feedback delay, C) normalized position gain, D) normalized velocity gain, E) normalized position to velocity ratio. Box-plot shows estimated values across all participants, with central line indicating the median, box edges showing the interquartile range, and whiskers representing the non-outlier range. Dots represent individual participant's data, with green for ML and orange for the AP direction, grey lines connect the data from same participant. The blue dot line in panel E represent y=1. X-axis labels donate: WP: walking peak; Walking: walking mean; NS: normal standing; US: unipedal standing; SP: step posture.

Table 1. Statistical results across tasks.

Variables	Factor	df	F	p	η^2_p
R^2	Task	3,42	21.16	<0.001**	0.60

	Direction	1,14	7.17	0.018*	0.34
	Task * Direction	3,42	59.92	<0.001**	0.81
$\hat{\tau}$	Task	3,42	78.23	<0.001**	0.85
	Direction	1,14	29.30	<0.001**	0.68
	Task * Direction	3,42	54.81	<0.001**	0.80
$\widehat{K_n}$	Task	4,56	10.92	<0.001**	0.44
	Direction	1,14	27.84	<0.001**	0.67
	Task * Direction	4,56	20.00	<0.001**	0.59
$\widehat{B_n}$	Task	4,56	179.74	<0.001**	0.93
	Direction	1,14	64.71	<0.001**	0.82
	Task * Direction	4,56	65.31	<0.001**	0.82
$\widehat{R_n}$	Task	4,56	26.89	<0.001**	0.66
	Direction	1,14	10.73	0.006*	0.43
	Task * Direction	4,56	1.70	0.16	0.11

The asterisks indicating levels of significance, where * indicates $p < 0.05$ and ** indicates

$p<0.001$.

From the data (Fig 1) it was evident that walking peak gains differed significantly

from that in walking and standing, whereas the walking gains were more

comparable to standing. Considering that the peak gains represent a strategy at a

specific moment within the stance phase, we deemed it less appropriate for direct

comparison with standing, therefore, we report the results in two sections. In the

first section, we focus on the results in walking, including the comparisons

between using total ground reaction forces and corrective forces as model outputs,

and examining the differences between peak and mean gains across the stance

phase. In the second section, we focus on the results between walking (mean gains)

and standing postures, to evaluate the similarities and differences between dynamic

and static conditions.

4.1 Feedback results in walking

4.1.1 Fitting model using total forces vs. corrective forces

The model fit (R^2) significantly improved when using corrective forces compared

to total ground reaction forces in both ML and AP direction (Fig 2, S1 Table). In

both directions, peak gains occurred in the early stance phase (~60% gait cycle),

regardless of using the total ground reaction forces or corrective forces. Notably,

using the corrective forces indicated continuous position feedback, which was

masked when using total forces (Fig 2).

Fig 2. Feedback models identified in walking with the total (panel A) and corrective ground reaction forces (panel B) as predicted variables. From left to right: model fit (R^2), normalized feedback position gains ($\widehat{K_n}$) and velocity gains ($\widehat{B_n}$) across the gait cycle. The green and orange lines represent the averaged results (over participants) in ML and AP direction, respectively. The dashed areas indicate the 95% confidence intervals. The horizontal lines in the left panels indicate when the model fit was significant for each individual participant (p<.05).

4.1.2 Peak gains vs. mean gains

Based on the results using corrective forces, the walking feedback delays were 46±5% and 23±13% of the gait cycle in the ML and AP directions, corresponding to 504±54ms and 250±137ms. The feedback delay in the ML direction was significantly longer than in the AP direction (p<.001) (Table 1, Fig 2).

For the position gains $(\widehat{K_n})$, the peak value was significantly larger than the mean

value in the ML direction (p=.001), while no significant difference was observed in

the AP direction (p=.14). Moreover, both peak and mean position gains showed

larger values in the ML direction than the AP direction (p<.001). The peak velocity

gains $(\widehat{B_n})$ were consistently larger than the mean values in both directions

(p<.001), and both peak (p=.009) and mean (p<.001) values were significantly

larger in the AP direction than in the ML direction. Consistent with the concept of

extrapolated center of mass, the ratio of the mean position to velocity gains $(\widehat{R_n})$

was not significantly different from 1. However, $\widehat{R_n}$ of the peak gains was

significantly smaller than 1 and significantly smaller than of the mean gains in both

ML direction (p<.001) and AP direction (p=.009). $\widehat{R_n}$ of mean gains in the ML

direction was significantly larger than in the AP direction (p<.001), whereas no

significant directional differences were observed for $\widehat{R_n}$ of the peak gains (p=.98).

4.2 Feedback control across tasks

In the comparison of standing and walking we report, for walking, values based on

averages over the stance phase only.

4.2.1 Model fit

Post-hoc simple effects showed that, in the ML direction, normal standing had a

significantly better model fit than all the other tasks (p<.001), followed by the step

posture for which model fit was significantly better than for walking and unipedal

standing (p<.001). No differences were found between walking and unipedal

standing in this direction (p=1). In the AP direction, the model fit for step posture

was significantly lower than for all the other tasks (p<.05). Moreover, unipedal

standing yielded a significantly lower model fit than normal standing (p<.001), but

comparable to walking (p=1). No significant differences were observed between

walking and normal standing (p=.07).

When comparing directions within the same task, walking and unipedal standing

showed a significantly better model fit in the AP direction than in the ML direction

(p<.001). In contrast, in the step posture, the model fit was better in the ML

direction than in the AP direction (p<.001). No directional differences were

observed in normal standing (p=.18).

329   4.2.2 Feedback delay

In line with our hypothesis, the post-hoc simple effects showed that the feedback

delay was significantly longer in walking than in standing postures in both

directions (p<.001). Among standing postures, in the ML direction, unipedal

standing had a significantly shorter delay than normal standing(p<.001) and step

posture (p<.001), while there were no differences between normal standing and

step posture (p=1). In the AP direction, normal standing had a significantly longer

delay than in unipedal standing (p<.001) and step posture (p<.001), while no

differences were observed between unipedal standing and step posture (p=.06).

Comparing directions, significantly longer delays in the ML than in the AP

direction were found in walking (p<.001) and step posture (p<.001), while the

opposite was found in normal standing (p<.001) and unipedal standing (p<.001).

4.2.3 Feedback position gain

In the ML direction, the position gains in all tasks exceeded their critical values

with $\widehat{K_n}$< -1, and no significant differences were observed between walking and

all standing postures (p>.05). Within standing postures, unipedal standing showed

significantly larger position gains than both normal standing (p<.001) and step

posture (p<.001). In the AP direction, the position gain in walking was slightly

below its critical value and was significantly smaller than that of all the standing

postures (p<.001), and the position gains of all standing tasks exceeded their

critical values. Among standing tasks, the step posture had the largest position

gain, followed by unipedal standing, with normal standing exhibiting a

significantly smaller position gain.

The step posture showed significantly larger position gains in the AP than in the

ML direction (p<.001), whereas the opposite pattern was found in the other tasks

(p<.001).

4.2.4 Feedback velocity gain

In contrast to the position gain, the velocity gains were all below critical damping

($\widehat{B_n}$> -1), and the velocity gain in walking was significantly larger than in standing

postures in both directions (p<.001). No significant differences were observed

among standing postures (p>.05).

The directional differences within task were consistent, with velocity gains larger

in the AP direction than in the ML direction (p<.05).

4.2.5 Ratio of feedback position to velocity gains

All standing tasks showed a normalized ratio ($\widehat{R_n}$) significantly larger than 1,

whereas in walking it was not significantly different from 1. No significant task

differences were observed (p>.05), but the ratio was consistently larger in the ML

direction than in the AP direction (p =.005).

5. Discussion

This study is the first to apply the same model to estimate and compare feedback

control during unperturbed walking and standing. Both walking and standing

showed good model fits with mean values higher than 0.69, supporting the use of

this model to estimate and compare feedback control without external

perturbations. Results showed that the phase-dependent characteristics in walking

indeed led to significant variations in feedback control over the gait cycle.

Accordingly, we used the mean values of walking, representing the overall control

strategy during the stance phase, for comparison with standing. For clarity, in the

following we refer to these mean values unless explicitly stated otherwise.

Our results showed significantly larger velocity control in walking than in standing

in both directions, whereas the position control in walking was significantly

smaller than standing in the AP direction. Despite these differences, all tasks

exhibited mostly above critical stiffness and under-damped control, with as only

exception position gains in the AP direction in walking, which were slightly below

critical stiffness. Notably, both walking and standing exhibited a position-to-

velocity gain ratio close to the $\sqrt{g/l}$, as predicted by the extrapolated center of

mass (XCoM) concept. These findings of feedback parameters which varied

significantly across tasks and directions, and consistent relative position and

velocity gains indicate a common strategy of sensorimotor control across both

dynamic and static conditions. We will further discuss these results below.

5.1 Feedback control in walking

In line with previous results [24], our results showed phase-dependent feedback control with peak control occurring during the early stance phase. The significantly improved model fit (Fig 2), as well as the observed continuous control in both directions suggest that using corrective forces allows for more accurate estimation of feedback parameters in walking.

Like Afschrift, De Groote [15], we found that gains were modulated across the gait cycle, with velocity control being more phase-dependent than position control in the AP direction. However, our results showed peak control in early stance, while Afschrift, De Groote [15] reported most pronounced control during mid-stance. The similarity likely reflects a common control strategy: the feedback control is performed based on preceding CoM position and velocity information. The difference in timing is likely due to methodological differences: we used corrective forces as the model output and selected the feedback delay based on the variation of gains across the gait cycle, whereas Afschrift, De Groote [15] used the EMG and ankle torque as model output and used a constant feedback delay. Additionally, our study didn't apply perturbations. While we observed more phase-dependent control for position in the ML direction than in the AP direction, Afschrift, De Groote [15] only investigated AP control.

Both mean and peaks gains showed stronger control of position in the ML direction, while in the AP direction velocity was more tightly controlled. These directional differences are consistent with the foot placement model [25], which demonstrated that ML foot placement control relies more on CoM position, whereas AP foot placement relies more on CoM velocity. Peak gains around early stance were expected, as this phase would impose high demands, especially to control AP velocity, where over-damped control appears to be used to avoid overshoot and oscillations to allow smooth and stable forward progression. The mean gains reflect control over the whole stance phase. Since velocity fluctuations in the remainder of the stance phase are smaller, an overall under-damped control may suffice.

5.2 Comparison of feedback control across tasks

5.2.1 Feedback delay

As hypothesized, the estimated feedback delay was significantly longer in walking than in standing. This was expected as in walking, feedback corrections are mainly applied after foot contact, while in standing stability control is continuous [26]. Directional differences in walking appear to reflect different stability demands and control strategies in the ML and AP directions. In the ML direction, stability control has been shown to be more challenging [26-29] and rely more on foot

placement [28, 30], while in the AP direction, stability control has been suggested

to rely more on the ankle strategy [31, 32], which can take effect during the stance

phase and thus leads to a shorter delay.

The differences in estimated feedback delays among standing postures can be

attributed to different biomechanical and neurophysiological characteristics. As

suggested by our previous work [17], increased intrinsic stiffness leads to a

negative bias on estimated feedback delays. The observed shorter delays in

unipedal standing may thus be explained by the increased intrinsic stiffness due to

the higher co-contraction levels [33-38]. In addition to biasing the estimates of the

delay due to differences in intrinsic stiffness, higher muscle activity can be

expected to cause a shorter electro-mechanical delay [39], contributing to a shorter

delay in unipedal standing and step postures. Non-neutral ankle angles in the step

posture coincide with higher intrinsic stiffness compared to normal standing, due to

the non-linear relationship between ankle angle and elastic parameters [40]. In

addition, the geometric configuration of the legs in the step posture with the non-

aligned ankle axes will yield 'structural stiffness', enhancing the overall stiffness.

5.2.2 Feedback position gains

Position gains for all tasks exceeded critical stability (i.e., $\widehat{K_n} < -1$) except for

walking in the AP direction. In the AP direction, the walking position gain was

slightly below the critical stiffness, suggesting instability in this direction, likely

prevented by changes in the base of support (stepping). The peak walking gains

typically above critical stiffness (i.e., $\widehat{K_n} < -1$) suggests that indeed these changes

are used to maintain a stable gait pattern.

Among standing postures, unipedal standing and step posture had significantly

larger position gains than normal standing, consistent with their presumed higher

intrinsic stiffness as mentioned above. Interestingly, position gains in unipedal

standing were approximately twice the per leg values observed in normal standing,

indicating that the control system modulates feedback gains per leg and

highlighting its adaptability to changed stability demands.

5.2.3 Feedback velocity gains

The velocity gains revealed an under-damped control in both walking and standing,

and there was a clear task-dependent modulation, with velocity gains significantly

larger in walking than in standing. The under-damped velocity control observed in

standing aligned well with the findings that described human upright posture as an

underdamped system [4, 41], likely reflecting that our control system allows some

oscillatory behavior during standing.

5.2.4 Ratio of position to velocity gains

According to the XCoM concept, control of pendular movement requires the ratio

of position to velocity gains to equal the inverted pendulum eigenfrequency,

yielding a normalized ratio $(\widehat{R_n})$ of 1. The $\widehat{R_n}$ in walking was not significantly

different from 1, which is as expected, whereas this value in standing was

significantly larger than 1. However, this ratio between walking and standing was

not significantly different, and both ratios remained close to the eigenfrequency

value.

The approximation of these ratios to the eigenfrequency is consistent with the view

that human body can be modelled as a single inverted pendulum model [4]. The

deviations from 1, particularly during standing, likely result from the human body

not behaving as a perfect pendulum, but instead stabilizing itself through multi-link

segments coordination[42]. The comparable ratios observed in walking and

standing suggest that, at least in unperturbed conditions, the sensorimotor system

maintains a relative consistent weighting between position and velocity feedback.

Such consistency may simply arise from the shared whole-body dynamics of

human body, and may also facilitate a smooth transition between dynamic and

static tasks.

6. Limitations

Several limitations should be noted. While we used a unified feedback model to estimate feedback control in walking and standing, the procedure to fit the model differed between tasks. Specifically, the method to select the feedback delay and corresponding gains was different. In addition, as reported in our previous study [17], intrinsic properties lead to errors in estimating feedback parameters in standing. How this will influence results in waking is unclear. Moreover, walking might introduce more soft tissue artifacts in the measurement of kinematics, potentially affecting the parameter estimation. Nonetheless, the estimated parameters aligned well with the task-specific characteristics. Finally, we examined only one gait speed in this study, whether and how the observed feedback strategy will change with speed needs further examination.

7. Conclusion

Our results demonstrated that while the feedback gains changed significantly across tasks and directions, the feedback position gains were consistently maintained close to critical stiffness, while the velocity was controlled at under-damped level. Notably, the relative ratio between position and velocity gains in both walking and standing remained close to the eigenfrequency, which might reflect a robust control strategy for stabilizing the human body as an inverted-pendulum.

Acknowledgements

The authors thank all the subjects for their participation in this study.

References

1. Koelewijn, A.D. and A.J. Van Den Bogert, *Antagonistic co-contraction can minimize muscular effort in systems with uncertainty.* PeerJ, 2022. **10**: p. e13085.

2. Forbes, P.A., A. Chen, and J.S. Blouin, *Sensorimotor control of standing balance.* Handb Clin Neurol, 2018. **159**: p. 61-83.

3. Shanbhag, J., et al., *A sensorimotor enhanced neuromusculoskeletal model for simulating postural control of upright standing.* Front Neurosci, 2024. **18**: p. 1393749.

4. Winter, D.A., *Human balance and posture control during standing and walking.* Gait & Posture, 1995. **3**(4): p. 193-214.

5. Blaszczyk, J.W., M. Beck, and D. Sadowska, *Assessment of postural stability in young healthy subjects based on directional features of posturographic data: vision and gender effects.* Acta Neurobiol Exp (Wars), 2014. **74**(4): p. 433-42.

6.   Winter, D.A., et al., *Unified theory regarding A/P and M/L balance in*

quiet stance. J Neurophysiol, 1996. **75**(6): p. 2334-43.

7.   Coleman, M.J. and A. Ruina, *An uncontrolled walking toy that cannot*

stand still. Physical Review Letters, 1998. **80**(16): p. 3658-3661.

8.   Mcgeer, T., *Passive Dynamic Walking.* International Journal of Robotics

Research, 1990. **9**(2): p. 62-82.

9.   Kuo, A.D. and J.M. Donelan, *Dynamic principles of gait and their clinical*

implications. Phys Ther, 2010. **90**(2): p. 157-74.

10.   Kuo, A.D., *Stabilization of lateral motion in passive dynamic walking.*

International Journal of Robotics Research, 1999. **18**(9): p. 917-930.

11.   Bauby, C.E. and A.D. Kuo, *Active control of lateral balance in human*

walking. J Biomech, 2000. **33**(11): p. 1433-40.

12.   Welch, T.D. and L.H. Ting, *A feedback model reproduces muscle activity*

during human postural responses to support-surface translations. J

Neurophysiol, 2008. **99**(2): p. 1032-8.

13.   Safavynia, S.A. and L.H. Ting, *Long-latency muscle activity reflects*

continuous, delayed sensorimotor feedback of task-level and not joint-

level error. J Neurophysiol, 2013. **110**(6): p. 1278-90.

14.   Jakubowski, K.L., et al., *Center of mass states render multi-joint torques*

throughout standing balance recovery. bioRxiv, 2024.

15.   Afschrift, M., F. De Groote, and I. Jonkers, *Similar sensorimotor*

transformations control balance during standing and walking. PLoS

Comput Biol, 2021. **17**(6): p. e1008369.

16.   van Dieen, J.H., S.M. Bruijn, and M. Afschrift, *Assessment of stabilizing*

feedback control of walking: A tutorial. J Electromyogr Kinesiol, 2024.

78: p. 102915.

17.   Geng, Y., S.M. Bruijn, and J.H. van Dieën, *Feedback control stabilizing*

the center of mass can be identified in unperturbed, upright standing.

bioRxiv, 2025: p. 2024.12.20.629656.

18.   Hof, A.L., M.G. Gazendam, and W.E. Sinke, *The condition for dynamic*

stability. J Biomech, 2005. **38**(1): p. 1-8.

19.   Hof, A.L., *The 'extrapolated center of mass' concept suggests a simple*

control of balance in walking. Hum Mov Sci, 2008. **27**(1): p. 112-25.

20.   Foulger, L.H., et al., *Learning to stand with delays alters sensorimotor*

control but does not cause instability when returning to natural balance.

J Physiol, 2025. **603**(19): p. 5629-5653.

21. Muehlbauer, T., et al., *One-leg standing performance and muscle*

activity: are there limb differences? J Appl Biomech, 2014. **30**(3): p. 407-

14.

22. Kingma, I., et al., *Validation of a full body 3-D dynamic linked segment*

model. Human Movement Science, 1996. **15**(6): p. 833-860.

23. Kuo, A.D., J.M. Donelan, and A. Ruina, *Energetic consequences of*

walking like an inverted pendulum: step-to-step transitions. Exerc Sport

Sci Rev, 2005. **33**(2): p. 88-97.

24. van Dieën, J.H., et al., *Simultaneous stabilizing feedback control of*

linear and angular momentum in

human walking. 2025.

25. Wang, Y. and M. Srinivasan, *Stepping in the direction of the fall: the next*

foot placement can be predicted from current upper body state in

steady-state walking. Biol Lett, 2014. **10**(9).

26. O'Connor, S.M. and A.D. Kuo, *Direction-dependent control of balance*

during walking and standing. J Neurophysiol, 2009. **102**(3): p. 1411-9.

27. Kongsuk, J., D.A. Brown, and C.P. Hurt, *Dynamic stability during*

increased walking speeds is related to balance confidence of older

adults: a pilot study. Gait Posture, 2019. **73**: p. 86-92.

28.   Harter, M.J., et al., *Modelling strategies supplemental to foot placement*

573   *for frontal-plane stability in walking.* J R Soc Interface, 2024. **21**(218): p.

574   20240191.

575   29.   Madehkhaksar, F., et al., *The effects of unexpected mechanical*

576   *perturbations during treadmill walking on spatiotemporal gait*

577   *parameters, and the dynamic stability measures by which to quantify*

578   *postural response.* PLoS One, 2018. **13**(4): p. e0195902.

579   30.   Bruijn, S.M. and J.H. van Dieen, *Control of human gait stability through*

580   *foot placement.* J R Soc Interface, 2018. **15**(143).

581   31.   Reimann, H., et al., *Neural Control of Balance During Walking.* Front

582   Physiol, 2018. **9**: p. 1271.

583   32.   Gatev, P., et al., *Feedforward ankle strategy of balance during quiet*

584   *stance in adults.* J Physiol, 1999. **514 (Pt 3)**(Pt 3): p. 915-28.

585   33.   Wu, F., et al., *Surface electromyography analysis of ankle flexor and*

586   *extensor activity under different standing stability levels.* J Orthop Surg

587   (Hong Kong), 2024. **32**(2): p. 10225536241258336.

588   34.   Fok, K.L., et al., *Co-contraction of ankle muscle activity during quiet*

589   *standing in individuals with incomplete spinal cord injury is associated*

590   *with postural instability.* Sci Rep, 2021. **11**(1): p. 19599.

591   35.   Baratta, R., et al., *Muscular coactivation. The role of the antagonist*

592   *musculature in maintaining knee stability.* Am J Sports Med, 1988. **16**(2):

593   p. 113-22.

594   36.   Amiri, P. and R.E. Kearney, *Ankle intrinsic stiffness changes with*

595   *postural sway.* J Biomech, 2019. **85**: p. 50-58.

596   37.   Sakanaka, T.E., et al., *Intrinsic ankle stiffness during standing increases*

597   *with ankle torque and passive stretch of the Achilles tendon.* PLoS One,

598   2018. **13**(3): p. e0193850.

599   38.   Sakanaka, T.E., M. Lakie, and R.F. Reynolds, *Individual differences in*

600   *intrinsic ankle stiffness and their relationship to body sway and ankle*

601   *torque.* PLoS One, 2021. **16**(1): p. e0244993.

602   39.   Vos, E.J., J. Harlaar, and G.J. van Ingen Schenau, *Electromechanical*

603   *delay during knee extensor contractions.* Med Sci Sports Exerc, 1991.

604   **23**(10): p. 1187-93.

605   40.   Weiss, P.L., R.E. Kearney, and I.W. Hunter, *Position dependence of ankle*

606   *joint dynamics--I. Passive mechanics.* J Biomech, 1986. **19**(9): p. 727-

607   35.

608   41.   Winter, D.A., et al., *Stiffness control of balance in quiet standing.* J

609   Neurophysiol, 1998. **80**(3): p. 1211-21.

610   42.   Hsu, W.L., et al., *Control and estimation of posture during quiet stance*

611   *depends on multijoint coordination*. J Neurophysiol, 2007. **97**(4): p.

612   3024-35.

613 # Supporting information

614   **S1 Table. Feedback results in walking and standing (mean±SD).**