1

Main Manuscript for

Structural Insights into Biased Signaling at Chemokine Receptor CCR7

Kotaro Tanaka[1], Kouki Nishikawa[2], Yuki Shiimura[3,4],

Yoshinori Fujiyoshi[1], and Naotaka Tsutsumi[1*]

[1] Advanced Research Initiative, Institute of Integrated Research, Institute of Science Tokyo,

Bunkyo, Tokyo, Japan.

[2] Joint Research Course for Advanced Biomolecular Characterization, Faculty of Agriculture,

Tokyo University of Agriculture and Technology, Fuchu, Tokyo, Japan.

[3] Division of Molecular Genetics, Institute of Life Science, Kurume University, Fukuoka, Japan.

[4] Department of Cell Biology, Graduate School of Medicine, Kyoto University, Kyoto, Japan.

*Corresponding author: Naotaka Tsutsumi

Email: tsutsumi.naotaka@tmd.ac.jp

Author contributions:

KT: Data Curation (equal), Formal Analysis (lead), Investigation (supporting), Methodology

(equal), Software (lead), Validation (equal), Visualization (equal), Writing – Original Draft

(supporting), Writing – Review & Editing (supporting). KN: Data Curation (supporting),

Investigation (supporting), Writing – Review & Editing (supporting). YS: Methodology

(supporting), Writing – Review & Editing (supporting). YF: Funding acquisition (lead),

Methodology (equal), Resource (lead), Supervision (equal), Writing – Review & Editing

(supporting). NT: Conceptualization (lead), Data Curation (equal), Formal Analysis (supporting),

Funding Acquisition (supporting), Investigation (lead), Methodology (equal), Project

Administration (lead), Resources (supporting), Software (supporting), Supervision (equal),

Validation (equal), Visualization (equal), Writing – Original Draft (lead), Writing – Review &

Editing (lead).

Competing interests:

The authors declare no competing interests.

Classification:

Biological Sciences

Biophysics and Computational Biology

Keywords:

CCR7, Chemokine, Cryo-EM, MD Simulations, Signaling Bias

This PDF file includes:

Main Text

Figures 1 to 6

Abstract：

CC chemokine receptor 7 (CCR7), which orchestrates adaptive immunity, exhibits a phenomenon

known as biased agonism. CCL19 induces robust G protein signaling and β-arrestin recruitment,

leading to transient signaling. In contrast, CCL21 preferentially activates G protein pathways with

minimal arrestin engagement, resulting in sustained signaling and differential functional

outcomes. Here, we present the cryo-EM structures of the human CCR7-G_i complex with either

CCL19 or CCL21. The structures reveal that while both engage a conserved orthosteric pocket,

they adopt markedly distinct binding poses. Notably, the compact 30s loop of CCL21 inserts

deeply into the receptor's extracellular vestibule, whereas the corresponding loop of CCL19 rests

atop extracellular loop 2. Molecular dynamics simulations further reveal that these distinct

binding modes induce differential intracellular dynamics, linked to the rotameric state of Y83.

We demonstrate that CCL19 stabilizes a flexible conformational ensemble with a highly dynamic

Helix 8, creating a lateral opening favorable for GPCR kinase engagement. Conversely, CCL21

restricts this flexibility, locking the receptor in a state that precludes kinase interaction while

maintaining G protein coupling. These findings provide key insights into the structural basis of

biased agonism at CCR7 and establish a foundation for rational design of pathway-selective

immunomodulators.

Significance statement:

Chemokine receptor CCR7 directs immune cell migration by activating distinct signaling pathways

in response to CCL19 and CCL21, a phenomenon known as biased agonism. The structural basis

for this divergence has been unclear. By combining cryo-electron microscopy enabled by a robust

mammalian protein production system with molecular dynamics simulations, we reveal that

although both ligands stabilize similar G protein-bound conformations, they drive distinct

receptor dynamics. We demonstrate that CCL19 induces a flexible intracellular conformation

necessary for recruiting regulatory kinases, whereas CCL21 locks the receptor in a state that

precludes this interaction. This "dynamic selection" model resolves a fundamental paradox in

chemokine biology, highlighting the critical role of protein dynamics in signal transduction and

providing a blueprint for designing precision immunotherapeutics.

Main Text

Introduction

Adaptive immunity relies on the precise spatiotemporal orchestration of immune cell

positioning and activation to mount effective responses against pathogens and malignant cells

while maintaining self-tolerance (1, 2). Central to this process is CC chemokine receptor 7 (CCR7)

(SI Appendix, Fig. S1A), a G protein-coupled receptor (GPCR) that governs the trafficking of

various T cells, mature dendritic cells (DCs), and B cells to secondary lymphoid organs (SLOs) (3–

7). Its two endogenous ligands, CC chemokine ligand 19 (CCL19) and 21 (CCL21), form

chemotactic gradients within SLOs to guide CCR7-expressing cells (8–11). This homing

mechanism is essential for bringing naïve T cells into close proximity with antigen-presenting DCs

within the T cell zones of SLOs, a prerequisite for T cell priming and the initiation of antigen-

specific immunity (12). Given its pivotal role, the CCR7 axis is a critical therapeutic target for a

broad spectrum of human pathologies (13–15). Hyperactivation of CCR7 signaling promotes the

metastasis of various cancers to lymph nodes and contributes to autoimmune disorders,

suggesting that antagonists could be beneficial. Conversely, insufficient CCR7 activity can impair

immune surveillance and vaccine efficacy, highlighting the potential utility of precisely tuned

agonists.

However, effectively targeting this axis is complicated by the phenomenon of biased

agonism, for which the CCR7 system is a paradigmatic example in the chemokine family (16).

Although both ligands engage the same receptor, they induce strikingly different downstream

signaling profiles and cellular responses (SI Appendix, Fig. S1B). While sharing a structurally

conserved core chemokine domain, CCL21 possesses a unique, highly basic C-terminal tail absent

in CCL19 (SI Appendix, Fig. S1C). This tail facilitates binding to glycosaminoglycans (GAGs) on cell

surfaces and in the extracellular matrix, which is crucial for establishing stable chemotactic

gradients and facilitating haptotaxis (17–19). Functionally, CCL19 is a potent inducer of both G

protein activation and β-arrestin recruitment. This process is mediated by receptor

phosphorylation via GPCR kinases (GRKs), specifically GRK3 and GRK6, leading to potent but

transient G protein signaling characterized by rapid receptor desensitization and internalization

(20–22). In contrast, CCL21 activates G protein signaling at levels comparable to, or slightly

weaker than, those of CCL19, but inefficiently recruits β-arrestin following phosphorylation only

by GRK6 and causes minimal internalization (19–22). This differential regulation is physiologically

crucial. The acute nature of CCL19 signaling may facilitate localized cellular activation and precise

positioning within SLOs, whereas the sustained signaling induced by CCL21, combined with GAG-

regulated localization, is thought to be essential for establishing the stable, long-range

chemotactic gradients required for guiding cell migration over distances (18, 23–25).

Despite its profound physiological importance, the mechanism by which these two

distinct chemokines elicit biased signaling through CCR7 has been a long-standing enigma (26).

Understanding the molecular mechanisms underlying this bias requires elucidating how ligand

binding translates into distinct receptor conformations that selectively engage G proteins versus

GRKs and arrestins. While the crystal structure of inactive-state CCR7 bound to the small-

molecule intracellular antagonist Cmp2105 has provided valuable insights into inhibitor binding

(15), the structural underpinnings of its activation by endogenous chemokines have remained

elusive.

Here, we report the cryo-electron microscopy (cryo-EM) structures of human CCR7 in

complex with CCL19 and CCL21, each stabilized in an active conformation by the heterotrimeric

inhibitory G protein (G_i). These structures reveal distinct ligand binding modes and differential

receptor conformations induced by the two chemokines. The structures illustrate a unique

mechanism of receptor activation shared within the CCR7-like chemokine receptor subfamily.

Concurrently, while the overall G_i binding mode is similar, subtle conformational differences are

observed in intracellular regions not directly involved in G_i binding, such as ICL1, which may

provide insights into the distinct downstream signaling profiles. Motivated by these distinctions

in ligand recognition and receptor conformation, we performed molecular dynamics (MD)

simulations to detail the dynamic basis for these structural distinctions after G protein

dissociation. These simulations revealed substantial differences in receptor conformational

equilibria, especially in Helix 8 (H8), that might influence GRK recruitment. This study elucidates

the structural basis for G protein activation by two distinct chemokines, providing fundamental

insights into the mechanism of biased agonism at CCR7 and offering a structural template for the

rational design of pathway-selective agonists.

Results

Preparation and overall architecture of the active CCR7-G$_i$ complexes

To elucidate the structural basis of CCR7 signaling, we utilized a robust strategy to assemble stable, active-state complexes for cryo-EM (Fig. 1 and Methods). We employed a temporally controlled expression system in tetracycline-inducible Expi293F (Expi293Fi) cells, ensuring the expression of an engineered G$_i$ heterotrimer prior to the induced expression of human CCR7 in order to maximize the receptor stability and the efficiency of complex formation (Fig. 1A). The G$_i$ complex incorporated a single-chain Gγ-Gα$_i$ fusion (27), wherein the Gα$_i$ contained dominant negative mutations (28), for enhanced stability. Additionally, the receptor and G protein were non-covalently tethered using the NanoBiT system (LgBiT fused to CCR7 and HiBiT fused to Gβ) (29, 30). The stable, ligand-free CCR7-G$_i$ complex further stabilized by the fiducial marker scFv16 (31) was purified, and the final signaling complexes were formed by adding separately purified full-length CCL19 or CCL21, yielding viable samples for cryo-EM analysis (Fig. 1B-D and SI Appendix, Fig. S1D-E).

Single-particle cryo-EM analysis of the samples yielded high-resolution maps, achieving nominal resolutions of 3.0 Å for the CCL19-CCR7-G$_i$-scFv16 complex and 3.2 Å for the CCL21-CCR7-G$_i$-scFv16 complex (Fig. 2A, and SI Appendix, Table S1 and Fig. S2-S4). During 3D classification, we observed structural heterogeneity related to post-translational modifications (glycosylation at N36 and N292 of CCR7) and weak densities suggesting partial ordering of the proximal CCL21 C-terminal tail. However, as these classes represented minor populations and exhibited lower local resolution at the core interface, we focused on the most homogeneous particle subsets to accurately define the primary ligand binding modes.

The overall architecture of the complexes resembles previously reported GPCR-G protein structures, characterized by the canonical insertion of the Gα$_i$-α5 helix into the receptor's cytoplasmic core (Fig. 2A-C). The seven-transmembrane (7TM) conformations of CCR7 in the two structures are essentially identical, with a Cα root-mean-square deviation (RMSD) of 0.36 Å. They also exhibit a highly similar docking mode for the G$_i$ heterotrimer (Cα RMSD of 1.48 Å). Strikingly,

however, the chemokines themselves adopt distinct orientations, reflected in a substantial Cα

RMSD of 4.31 Å when the complexes are aligned based on the receptor (Fig. 2B).

A unique mode of TM6 movement during CCR7 activation

To understand the conformational changes associated with CCR7 activation, we

compared the CCL19-bound active-state structure with the inactive CCR7 structure bound to the

intracellular antagonist Cmp2105 (15) to elucidate the general mode of activation in comparison

with other chemokine receptor systems (Fig. 2C). Both CCL19 and CCL21 induce a similar global

conformational rearrangement in CCR7, creating a cytoplasmic cavity to accommodate the G

protein.

The most prominent feature of this transition is the outward movement of

transmembrane helix 6 (TM6). Unlike the canonical hinge-like pivot observed in most class A

GPCRs, including CCR2 and CXCR2 (32, 33), CCR7 exhibits a predominantly parallel outward shift

of TM6 by 3-4 Å (Fig. 2D). In CCR7, the entire TM6 helix, from its extracellular end to its

intracellular tip, moves laterally as a semi-rigid unit away from the receptor's central axis. This

mode of activation is similar to that observed in its paralog, CCR6 (34% sequence identity), which

also shows a ~5 Å global shift (34).

These distinct activation modes correlate with the presence or absence of the critical

$W^{6.48}$ residue within the CWxP motif (superscripts denote Ballesteros-Weinstein numbering) (Fig.

2D). In most class A GPCRs, $W^{6.48}$ acts as a rotamer "toggle switch," initiating conformational

changes that displace the intracellular half of TM6 (33, 35). In CCR7 and CCR6, this tryptophan is

replaced by glutamine ($Q^{6.48}$). The absence of the bulky tryptophan side chain likely enables the

rigid-body movement of TM6, suggesting a subclass-specific activation mechanism initiated by

sensing the steric bulk of the large, proteinaceous chemokine ligands rather than relying on the

canonical toggle switch.

Distinct binding modes of CCL19 and CCL21 dictated by CRS3 interactions

The most pronounced differences between the two active complexes are found at the

extracellular ligand-recognition interface. Strikingly, the core domain of CCL21 is globally rotated

by 17.6° relative to CCL19 when viewed from the side of the receptor (Fig. 3A). This rotation results in significantly altered interaction interfaces, with larger buried surface areas for CCL21 (1,640 $Å^2$) compared to CCL19 (1,380 $Å^2$).

Chemokine recognition generally follows an expanded "two-site" model (36), involving interactions across several chemokine recognition sites (CRS): CRS1 (receptor N-terminus/chemokine core), CRS1.5 (receptor N-loop and ECL3/chemokine N-loop), CRS2 (receptor 7TM pocket/chemokine N-terminus), and CRS3 (receptor ECL2/chemokine 30s loop) (Fig. 3A). Our structures reveal that the dramatic difference in orientation stems primarily from divergent interactions at CRS3.

Despite overall sequence conservation (32% identity, 68% similarity in the core region; Fig. S1C), the 30s loops of CCL19 and CCL21 exhibit distinct sequences and conformations (Fig. 3B). The CCL21 loop (^{30}PSLGCSIP^{38}) is notably less bulky and more hydrophobic than the corresponding loop in CCL19 (^{30}IKDGCRVP^{38}), particularly featuring L32 in CCL21 versus D32 in CCL19. Consequently, CCL19's 30s loop adopts a curled conformation and rests atop CCR7-ECL2. In contrast, CCL21's compact 30s loop facilitates its deeper penetration into the extracellular pocket. Crucially, CCL21-L32 inserts like a wedge between the chemokine N-terminus and CCR7-L212^{ECL2} (Fig. 3C, right). This deep insertion stabilizes a specific conformation of ECL2 and contributes significantly to the overall rotation of CCL21.

In contrast to the divergent CRS3 interactions, the engagements at CRS1 and CRS2 are relatively conserved. At CRS1, the chemokine bodies interact with the N-terminal region of CCR7, with CCR7-F44 being the core of the tip interactions (SI Appendix, Fig. S5A). At CRS2, the N-terminal loops of both chemokines penetrate the "minor orthosteric binding pocket" of CCR7 surrounded by TM1, TM2, TM3, TM6, and TM7, forming extensive polar and hydrophobic interactions (Fig. 3C) (37). The N-terminus of CCL19 penetrates slightly deeper into the pocket compared to CCL21. Specifically, the sidechain of CCL19-T2 engages Y136$^{3.32}$ at TM3, inducing a subtle shift of this residue. The observed CRS2 interaction networks generally correlate with reported receptor mutagenesis data (38). For example, CCR7 residues K50^{N-term} and R54^{N-term} are located at the tip of TM1 and the center of the molecular interface, forming critical contacts with the N-terminal peptides of both chemokines (Fig. 3C). This explains why the CCR7-K50A^{N-term} and

CCR7-R54A$^{N\text{-term}}$ mutations dramatically impaired both CCL19- and CCL21-induced signaling. Conversely, the CCR7-R209A^{ECL2} mutation selectively reduces CCL19-induced signaling. This selectivity is structurally rationalized by the more extensive polar contacts of R209^{ECL2} with the N-terminus/core of CCL19, via D4/E6 sidechains and D32 backbone, compared to CCL21 (Fig. 3C), suggesting that R209^{ECL2} contributes to a specific interaction network that is more critical for CCL19 recognition.

These distinct binding modes result in ligand-specific conformational shifts in the extracellular portions of TM1, TM3, TM5, TM7, and ECL2 (Fig. 3D,E and SI Appendix, Fig. S5B). Furthermore, these differences propagate to an aromatic cluster located in the middle layer of the minor pocket, just beneath the chemokine N-termini (Fig. 3F, and SI Appendix, Fig. S5A). Residues Y65$^{1.39}$, Y136$^{3.32}$, Y279$^{6.51}$, and Y312$^{7.39}$ exhibit distinct conformations in the inactive, CCL21-bound, and CCL19-bound states, where the conformational changes in the CCL19-bound state are consistently more extensive than those in the CCL21-bound state. Among these residues, the Y312$^{7.39}$E mutation is known to confer a high degree of constitutive activity to CCR7 while abolishing chemokine sensitivities, whereas Y312$^{7.39}$A only causes high basal activities, underscoring the importance of this residue for the inactive-to-active transition (19). The correlated rearrangements in the aromatic network likely contribute to the differential stabilization of active-state conformations and may underlie the observed differences in general signaling potency (CCL19 > CCL21) (19).

Structural basis of CCR7 activation and core G protein interface

The binding of both chemokines induces canonical conformational changes associated with class A GPCR activation (35) (Fig. 4A), although with the unique features related to TM6 movement described earlier. The PIF-like motif (P$^{5.50}$-M$^{3.40}$-F$^{6.44}$) undergoes reorganization, characterized by the movement of M144$^{3.40}$ sidechain, which induces inward and outward TM5 and TM6 shifts, respectively (Fig. 4A, bottom left). The immediately downstream hydrophobic lock collapses upon activation, further destabilizing the inactive state (Fig. 4A, bottom center).

Concurrently, the sodium-binding site, which is stabilized by sodium ion coordination with N76$^{1.50}$, D104$^{2.50}$, N322$^{7.49}$, and Y326$^{7.53}$ in the inactive state, collapses upon activation,

accompanied by the kinking of the NP$^{7.50}$xxY motif including TM7 and flipping of the Y326$^{7.53}$

sidechain (Fig. 4A, top right). At the same time, the DR$^{3.50}$Y motif reorganizes, such that R154$^{3.50}$

moves to form a bridge with Y326$^{7.53}$, stabilizing the active conformation for G protein coupling

(Fig. 4A, bottom right).

Comparison of the active states with the inactive CCR7 bound to Cmp2105 highlights the

structural rearrangements required for G protein engagement (Fig. 4B, top left). The outward

movement of TM6 and the concomitant shift of TM5 create the binding pocket for the Gα$_i$ subunit.

Accompanying these TM movements, the intracellular loops undergo significant reorganization.

Notably, ICL2 transitions from a disordered loop in the inactive state to a stable, short α-helix

upon activation, which is critical for engaging the Gα subunit. Similarly, ICL3 is stabilized following

the displacement of TM6, contributing to the formation of the G protein interface. Cmp2105

stabilizes the inactive state by preventing the inactive-to-active conformational shift and

occupying a position that overlaps with the binding site of the Gα$_i$-α5 helix, directly blocking G

protein coupling (Fig. 4B, top right).

The primary interface between CCR7 and Gα$_i$ involves the insertion of the C-terminal α5-

helix of Gα$_i$ deep into the cytoplasmic cavity of the receptor (Fig. 2A-C and 4B). This interface is

stabilized by extensive interactions and is highly conserved between the CCL19- and CCL21-

bound complexes (Fig. 4B, bottom panels). The Gα$_i$-F(-1) [F354] sidechain is sandwiched between

E257$^{6.29}$ and K260$^{6.32}$, positioning the C-terminal carboxyl group for polar interactions with the

backbone amide at the TM7-H8 turn and the K260$^{6.32}$ sidechain. The L(-2) [L353] sidechain is

positioned in the hydrophobic pocket formed by TM6 residues and R154$^{3.50}$ at TM3, which is

further covered by L(-7) [L348]. The R154$^{3.50}$ sidechain forms polar interactions with the C(-4)

[C351] backbone carbonyl. Although there are several minor variations at the peripheral region

of the core interface, the overall interaction patterns are highly conserved.

Subtle differences at the intracellular loops suggest distinct signaling outcomes

Despite the overall similarity in G protein coupling, a detailed comparison reveals subtle

but notable differences in the intracellular loops between the CCL19- and CCL21-bound

complexes (Fig. 5A,B).

The conformations of ICL2 and ICL3, which form the major part of the G_i interface, are

essentially identical in both active structures. In the ICL2 region, the interactions between CCR7

residues (e.g., H165, R166) and the $G\alpha_i$ αN-β1 hinge region (e.g., R32, D193) are conserved,

exhibiting minor variations (Fig. 5A,B, top left panels). Similarly, the core interface mediated by

ICL3, TM5, and TM6 is highly conserved (Fig. 5A,B, bottom left panels).

In contrast, ICL1, which is less directly involved in the primary G_i interface, adopts distinct

loop conformations in the two complexes (Fig. 5A,B, bottom right panels). This divergence

appears to originate from differences in the side-chain conformation of Y83$^{1.57}$, located at the

cytoplasmic end of TM1. The distinct binding modes of CCL19 and CCL21 induce specific

rearrangements in the extracellular TMs (Fig. 3D) and the central aromatic network (Fig. 3F). This

network appears to function as an allosteric link, propagating these differences through the

helical bundle (particularly TM1), ultimately leading to the distinct rotameric states of Y83$^{1.57}$ in

the CCL19- versus CCL21-bound structures. These states influence the length of TM1 and

positioning of adjacent residues and propagate to the overall conformation of the ICL1 loop.

While these structural variations, particularly in ICL1, are not extensive in the static G_i-

coupled structures, they are particularly intriguing in the context of biased signaling. The high

similarity of the G_i-coupled states suggests that the mechanism underlying the differential

recruitment of GRKs and arrestins likely stems from differences in receptor conformations or

dynamics that occur after G protein activation and dissociation. Although ICL1 is not considered

the primary determinant of GRK binding, GRKs recognize specific conformations of the

intracellular loops and the receptor core to phosphorylate the receptor (39, 40). It is plausible

that the distinct extracellular binding modes of CCL19 and CCL21 prime the receptor to adopt

different conformational ensembles, including intracellular architectures other than ICL1, upon

G protein release.

Molecular Dynamics simulations to probe the basis of biased signaling

To investigate how the distinct ligand binding modes and the subtle intracellular

differences might influence receptor dynamics relevant to biased signaling, we performed six

independent 2.3 μs all-atom MD simulations for each complex (Fig. 5C and S6-S9). We used the

experimental CCL19-CCR7 and CCL21-CCR7 structures as starting points, after removing the G

protein and scFv16 and embedding the complexes in a lipid bilayer (SI Appendix, Fig. S6A and

Methods). These starting points are intended to mimic the receptor states primed for subsequent

signaling events, such as GRK recruitment, after initial G protein activation and dissociation.

During the simulations, the chemokine ligands (especially CCL19) exhibited higher

flexibility (SI Appendix, Fig. S6B,C), consistent with the observed cryo-EM density at the

chemokine region (SI Appendix, Fig. S2-S3). In contrast, the receptor core (7TM bundle) of both

complexes slightly deviated from the initial cryo-EM conformation, but converged to a similar

conformational state (SI Appendix, Fig. S6B,C and S7).

The most notable finding was the differential dynamics of H8 in CCR7 (Fig. 5C). In the

CCL19-bound state, the TM7-H8 loop exhibits higher dynamics and repositions into the receptor

central axis, forming a larger gap between TM6 and TM7 (Fig. 5C). This is characterized by the

proximity between TM2 and the TM7-H8 loop, and the H8 rotation (SI Appendix, Fig. S8). In the

CCL19-bound state, the Cα distance between T93$^{2.39}$ and G330$^{8.47}$ became substantially narrower

from 11.5 Å to approx. 6.4 Å, accompanied by a significant rotation from the cryo-EM

conformation (defined as H8 angle 0°) to a new stable state at approx. -28.2°. In the CCL21-bound

state, however, H8 remained more stable near the cryo-EM conformation with the Cα distance

between T93$^{2.39}$ and G330$^{8.47}$ remaining constant at approx. 11.3 Å and the rotation being less

pronounced rotation to approx. -14.4°.

These differences trace back to the formation of two distinct salt bridges: a highly stable

one directly linking R88$^{12.49}$ (ICL1) to D336$^{8.53}$ (H8) (present in 92% of simulation time), and a

second, dynamic one between R88$^{12.49}$ (ICL1) and D94$^{2.40}$ (TM2) (present in 58% of simulation

time) (SI Appendix, Fig. S9A, top panel, and S9B). In sharp contrast, in the CCL21-bound state,

while the R88$^{12.49}$-D336$^{8.53}$ bridge remained largely stable (86% formation), the crucial R88$^{12.49}$-

D94$^{2.40}$ bridge which brings H8 toward TM2 was almost entirely absent (9% formation).

This dynamic behavior in the CCL19-bound state also correlates with the different chi1

(χ1) angles of Y83$^{1.57}$ at the TM1 terminus (180° for CCL19 and -86° for CCL21), observed in the

cryo-EM structures, which remained stable in their respective states throughout the simulations

(SI Appendix, Fig. S9A, bottom panel, and S9C). This orientation of Y83$^{1.57}$ appeared to influence

the ICL1 dynamics and allosteric coupling between ICL1 and H8 (SI Appendix, Fig. 5C and S9A,D).

In the CCL19-bound state, the outward-facing conformation of $Y83^{1.57}$ accommodates $F333^{8.50}$

from H8 into the pocket near TM2, which is blocked by the inward-facing $Y83^{1.57}$ sidechain in the

CCL21-bound state.

Taken together, these results suggest that the specific intracellular domain reorganization

induced by CCL19, particularly the dynamic movement of H8, differs from the stable

conformation induced by CCL21, and likely has a differential impact on the recognition of

downstream effectors like GRKs. We hypothesize that the stable conformational ensemble

induced by CCL21, which settles into a single state lacking the deep H8 insertion (Fig. 5C, bottom

left panel) and thus resembling the G_i-bound conformation in the core pocket, is suboptimal for

GRK interaction. This "locked" state, which appears to remain competent for G protein coupling

but not for GRK engagement, results in G protein-biased signaling. In contrast, the

conformational flexibility induced by CCL19 appears to sample multiple states, in which a major

population representing the H8-shifted "open" state characterized by the deeper insertion of the

TM7-H8 loop coexists with a population lacking this insertion and thus resembling the starting

G_i-bound conformation (Fig. 5C, top left panel). This dynamic equilibrium, which allows the

sampling of both a GRK-favorable state and a G_i-favorable state, may represent the structural

basis for CCL19's potent, non-biased activation profile, facilitating the efficient engagement of

both G proteins and GRKs.

Discussion

The precise orchestration of immune cell migration by the CCL19/CCL21-CCR7 axis is

fundamental to adaptive immunity. This system partly relies on the biased agonism of its two

ligands to achieve distinct physiological outcomes, localized activation versus long-range

chemotaxis (18, 23, 24, 41). Our structural analyses provide a molecular framework for

understanding how this divergence is achieved.

The cryo-EM structures reveal a marked divergence in ligand recognition, primarily

dictated by interactions at the CRS3 interface (Fig. 3). The deep, hydrophobic engagement of the

CCL21 30s loop contrasts sharply with the shallower binding of CCL19 loop, illustrating how the

sequence variations in flexible chemokine loops can alter the overall binding topology.
Furthermore, our analysis highlights that CCR7 employs a non-canonical, rigid-body movement
of TM6 during activation (Fig. 2), likely facilitated by the absence of the canonical $W^{6.48}$ toggle
switch. This mechanism suggests a specialized adaptation within this receptor subfamily for
transducing signals from protein ligands.

The central paradox of CCR7 bias is that both ligands induce highly similar G_i-coupled
conformations (Fig. 5A,B) yet elicit vastly different downstream responses regarding
desensitization (22). Our MD simulations suggest that the mechanism of bias lies not in the initial
G protein engagement, but in the receptor dynamics that follow G protein dissociation, revealing
that CCL19 induces a flexible ensemble capable of adopting a widely "open" H8 state, whereas
CCL21 stabilizes a single, "locked" G_i-bound-like state (Fig. 5C).

Structural comparisons with other GPCR-GRK complexes underscore the importance of
this CCL19-induced "open," H8-shifted state (Fig. 6). Cryo-EM structures of neurotensin receptor
1 (NTSR1) and Rhodopsin (Rho) reveal that the binding modes of the Gα C-terminal α5 helix and
the GRK N-terminal α-helix (αN) are mutually exclusive (39, 40, 42, 43) (Fig. 6A, B). While the Gα-
α5 helix requires its deeper insertion towards the receptor core, the GRK-αN helix exhibits a
shallower insertion and follows a distinct trajectory, exiting through the lateral gap formed
between TM6 and the TM7-H8 loop. This suggests a general mechanism where efficient GRK
recognition requires not only a sufficiently wide lateral opening between TM6 and the TM7-H8
loop, but perhaps also a conformation that disfavors the deep re-insertion of a G protein.

In this context, our findings provide a compelling structural rationale for the biased
signaling, where the conformational flexibility of CCL19-bound CCR7 plays a key role. Its ability
to dynamically adopt the "open" state, in which the TM7-H8 loop itself inserts into the
cytoplasmic cavity (Fig. 5C), simultaneously achieves two discrete goals. It occupies the deep
central cavity, thus sterically hindering deep G protein re-engagement, and creates the widened
lateral gap favorable for initiating GRK3 engagement (a GRK2/3 subfamily member known to
phosphorylate CCR7) (Fig. 6C). Furthermore, this inserted TM7-H8 loop may form a novel
composite interaction surface for GRK recognition, as the GRK-biased small molecule SBI-553
does for NTSR1 (40).

This model explains the non-biased profile of CCL19, where the CCL19-bound CCR7

exhibits dynamic equilibrium that allows it to adopt both this "open," GRK-favorable state and

the G_i-bound-like state lacking the loop insertion, permitting efficient engagement of both

effector types. Conversely, the "locked" stable state stabilized by CCL21 maintains a vacant deep

pocket and a narrower lateral gap, while also lacking the TM7-H8 loop insertion. This

conformation appears to remain highly competent for G protein coupling but fails to provide the

steric opening or interaction surface required for efficient GRK3 engagement. This structural

dichotomy—a flexible ensemble of both G_i- and GRK-favorable states versus a conformationally

restricted, G_i-competent-only state—offers a mechanistic basis for the G protein-biased signaling

observed for CCL21.

The dynamic model aligns with their physiological roles. CCL19, by inducing

conformational dynamics favorable for GRK engagement, drives rapid desensitization and

internalization (20, 22). This transient signaling is ideally suited for the precise tuning of immune

function, particularly in acute cell activation and tissue development (23, 41, 44–47). In contrast,

the unique binding mode of CCL21 stabilizes a conformationally restricted ensemble that

disfavors efficient GRK3 interaction, resulting in sustained G protein signaling (20). This G protein

bias is essential for guiding immune cells over long distances (18, 25, 45). Moreover, the deep

insertion of the CCL21 30s loop may also contribute to a slower dissociation rate, further

supporting sustained signaling. This mechanism, combined with the known role of the CCL21 C-

terminal tail in establishing stable chemotactic gradients via GAG interactions (17–19), provides

a comprehensive explanation for its function in long-range migration and haptotaxis.

In conclusion, this study elucidates the structural and dynamic basis of biased agonism at

CCR7, demonstrating how subtle differences in ligand recognition are translated into distinct

physiological outcomes. Thus, exploring chemokine analogs that simultaneously modulate

interactions across the CRS2 and CRS3 interfaces, perhaps mimicking the specific engagement

profile of CCL21 but at higher affinity, might lead to G protein-biased agonists. Such molecules,

by potentially promoting sustained cell trafficking, could serve as adjuvants for vaccines or cancer

immunotherapy. This structural framework thus provides a foundation for developing novel

immunomodulators designed to precisely control the CCR7 signaling axis.

Limitations of Study

While the current structures and MD simulations provide significant insights into CCR7 activation and mechanisms possibly linked to the biased signaling, several limitations remain. These data do not directly visualize the conformations responsible for GRK or β-arrestin engagement. Determining the structures of CCR7 in complex with GRKs or β-arrestins will be crucial to fully elucidate the mechanisms of biased signaling. Furthermore, although we observed heterogeneity related to the CCL21 C-terminal tail and potential post-translational modifications in cryo-EM density, we could not resolve these features at high resolution. Further studies are needed to understand their roles in CCR7 function and regulation.

Materials and Methods

Mammalian cell lines and culture conditions

Expi293F inducible (Expi293Fi) cells (Thermo Fisher Scientific, a component of expression system kit Cat# A39251) were adopted and maintained in HE400AZ media (GMEP) supplemented with 10 μg/mL Blasticidin (InvivoGen) at a 20 mL culture scale. For protein expression, cells were expanded with HE400AZ media up to 200 mL and then HE200 media (GMEP) supplemented with GlutaMAX (Thermo Fisher Scientific) for larger volumes to reduce the total cost. Cells were maintained at a density between 0.3 and 4×10^6/mL and shaken at 37°C and 130 rpm with 8% (v/v) CO_2. Transfections were performed at a cell density of approximately 3×10^6/mL.

Insect cell lines and culture conditions

For scFv16 expression from *Trichoplusia ni* (Hi5) ovarian cells (Expression Systems, Cat# 94-002), baculovirus was produced in *Spodoptera frugiperda* (Sf9) ovarian cells (ATCC CRL-1711) maintained in Sf-900 III medium (Thermo Fisher Scientific) with 10% (v/v) FBS (Sigma-Aldrich) and GlutaMAX. Hi5 cells were maintained and expanded for protein expression in ESF 921 Insect Cell Culture Medium (Expression Systems). Insect cells were grown at 27°C with ambient CO_2 and gentle agitation, and baculovirus infections for virus amplification and protein expression were performed at a cell density of approximately 2×10^6/mL.

Expression of the apo CCR7-G_i complex

The expression plasmids were designed to perform induced and constitutive co-expression of CCR7 and heterotrimeric G_i protein, and the coding genes were synthesized by GenScript for cloning in expression plasmids. The full-length human CCR7 lacking the endogenous signal peptide (UniProt ID: P32248, residues 25-378) was cloned into pcDNA5/TO vector (Thermo Fisher Scientific) with the N-terminal HA secretion signal peptide followed by FLAG tag, and C-terminal LgBit split NanoLuc fragment (29). Single chain human $G\gamma_2$-$G\alpha_{i1}$, where $G\alpha_{i1}$ (UniProt ID: P63096, residues 2-354) with dominant negative mutations (S47N, G203A, E245A, A326S) (28) was fused downstream of $G\gamma_2$ (UniProt ID: P59768) separated by 3x GSA linker (27) and cloned into the pEG vector (48). Human $G\beta_1$ (UniProt ID: P62873) with C-terminal HiBit NanoLuc

complementation fragment and N-terminal 3C protease-cleavable 10×His tag was also cloned
into the pEG vector for co-expression.

Protein expression was performed using Expi293Fi cells as per manufacturer's
recommendation with modifications to use polyethyleneimine (PEI) Max (Polysciences) as a
transfection reagent (49), 3 mM sodium valproate and 0.4% (w/v) glucose as enhancers (50), and
doxycycline (Wako) as the inducer. The three plasmids were used at an approximately equal
weight ratio, and a total of 1 µg plasmid and 4 µg PEI per 1 mL cell culture were mixed in HE400AZ
media and incubated for 15 min at room temperature before transfecting the Expi293Fi cells.
One day after transfection, the enhancers were added, and after an additional 24 hours at 37°C
for initial G_i expression, the CCR7 expression was induced by adding 8 µM doxycycline. The
culture flask was transferred to a 30°C incubator and shaken for 48 hours. The cells were
harvested, washed once with phosphate-buffered saline (PBS), and stored at -30°C. 200 mL
Expi293Fi culture typically yields approximately 5 grams of cell pellet.

Expression and purification of CCL19, CCL21, and scFv16

CCL19 and CCL21 were expressed from Expi293Fi cells but with constitutive expression
using a standard CMV promoter, and the coding genes were synthesized by GenScript. Full-length
CCL19 (UniProt ID: Q99731) and CCL21 (UniProt ID: O00585) were each fused to the C-terminal
6×His-tagged human IgG1 Fc via the GSGSGSAAA linker, HRV 3C protease site (LEVLFQGP), and
GSDGSGSGS linker analogous to the construct previously used to prepare the chemokine domain
of CX_3CL1 (51) but cloned into pD649 transient expression vector (ATUM) (52) instead of the
BacMam transfer vector (53). Transfection was performed as described for the CCR7-G_i complex
with a 1:4 plasmid:PEI ratio, and the enhancers were added to the culture on the following day.
The cells were shaken in the 37°C incubator for an additional 84 hours, and the culture
supernatant was collected for purification using Ni-NTA (51, 52). The Fc-tagged chemokines were
eluted with HEPES-buffered saline (HBS, 10 mM HEPES-Na pH 7.2 and 150 mM NaCl) containing
250 mM imidazole, and the Fc tag was removed by adding 2% (w/w) 3C protease, which was
prepared in-house. The solutions were buffer exchanged to HBS containing 10 mM imidazole,
flowed through Ni-NTA equilibrated with HBS containing 20 mM imidazole to remove the Fc tag,

and concentrated to approximately 300 µL. Due to the limited protein yields (~50 µM and ~20

µM for CCL19 and CCL21, respectively), the samples were supplemented with 10% (v/v) glycerol,

aliquoted, flash-frozen, and stored at -80°C for further use.

scFv16 (31) was expressed from Hi5 ovarian cells using Bac-to-Bac baculovirus expression

system (Thermo Fisher Scientific) and purified as reported previously (51). Briefly, the collected

Hi5 culture supernatant was clarified after a precipitate was formed by adding 1 mM $NiCl_2$, 5 mM

$CaCl_2$, and 50 mM Tris-HCl (pH 8), and then incubated with Ni-NTA to capture the C-terminal

6×His-tagged scFv16. The Ni-NTA resin was collected, washed with HBS containing 20 mM

imidazole, and eluted by increasing imidazole concentration to 250 mM. The eluate was

concentrated and purified using a size-exclusion chromatography column (SEC) Superdex 75

10/300 GL (Cytiva) equilibrated with HBS. The peak fraction was concentrated to approximately

640 µM, aliquoted and flash-frozen with 10% (v/v) glycerol, and stored at -80°C until future use.

Purification of CCR7-G_i-scFv16 complexes bound to CCL19 or CCL21

The ligand-free, NanoBiT-tethered CCR7-G_i-scFv16 complex was purified before the

chemokines were added during the final protein concentration step. Approximately 5 grams of

cell pellet expressing CCR7-G_i was directly solubilized for 2 hours at 4°C with 10 nmol of

recombinant scFv16 and nutation in a solubilization buffer consisting of HBS, 10% (w/v) glycerol,

1% (w/v) n-dodecyl-β-D-maltoside (DDM, Anatrace), 0.1% (w/v) lauryl maltose neopentyl glycol

(LMNG, Anatrace), 0.11% (w/v) cholesteryl hemisuccinate tris salt (CHS, Anatrace), cOmplete

protease inhibitor cocktail (Roche), and apyrase (0.2 µL, New England Biolabs). The resulting cell

lysate was clarified by two sequential centrifugations at 12,000 *g* for 30 minutes at 4°C. The

supernatant was then supplemented with 3 mM $CaCl_2$ and incubated with 1 mL of in-house anti-

FLAG M1 sepharose for 2 hours at 4°C. The resin was collected in a column and washed

sequentially, first with the solubilization buffer, then HBS containing 0.1% LMNG, 0.01% CHS, and

3 mM $CaCl_2$. A final wash was performed with HBS containing 0.01% LMNG, 0.001% CHS, and 3

mM $CaCl_2$, and the CCR7-G_i-scFv16 complex was eluted from the column with HBS buffer

containing 0.001% LMNG, 0.001% glyco-diosgenin (GDN, Anatrace), 0.0001% CHS, 5 mM EDTA,

and 100 µg/mL FLAG peptide (PH Japan). The eluate was concentrated to approximately 400 µL,

and 2% (w/w) of in-house 3C protease was added for overnight incubation at 4°C. The protein

was subjected to SEC on a Superdex 200 Increase 10/300 GL column (Cytiva) equilibrated with

SEC buffer consisting of HBS, 0.001% LMNG, 0.001% GDN, and 0.0001% CHS. Peak fractions were

collected, concentrated, and quantified, yielding approximately 0.20 mg (1.1 nmol) of the

purified CCR7-G_i-scFv16 complex.

Concurrently, frozen stocks of the chemokines CCL19 and CCL21 were thawed and

dialyzed against the SEC buffer using Xpress Micro Dialyzer with a 3.5 kDa molecular weight cutoff

(MWCO) (Scienova GmbH). To form the chemokine-bound signaling complexes, each prepared

chemokine was added to the purified CCR7-G_i-scFv16 in separate reactions. Specifically, CCL19

was added at a 1.6-fold molar excess to 0.6 nmol of CCR7-G_i-scFv16, while CCL21 was added at

the same molar ratio to 0.5 nmol aliquot of the ligand-free complex. Finally, each mixture was

concentrated to 4.5 mg/mL using 3 kDa MWCO Amicon Ultra 0.5 mL centrifugal filters for cryo-

EM.

Cryo-EM grid preparation and data collection

For cryo-EM analysis, a 3.5 µL aliquot of either complex was applied to a glow-discharged

Quantifoil R1.2/1.3 300-mesh gold grids. The grids were plunge-frozen in liquid ethane using a

Vitrobot Mark IV (Thermo Fisher Scientific) with a 10-sec hold period, blot force of 10, and

blotting time of 2-4 sec with 100% humidity at 4°C.

Cryo-EM data were subsequently collected using a JEOL JEM-Z320FHC electron

microscope operated at 300 kV, equipped with an in-column omega energy filter set to a 20 eV

slit width. Images were recorded on a Gatan K3 camera in correlated double sampling mode, with

a calibrated pixel size of 0.79 Å at the specimen. Automated data acquisition was performed using

SerialEM (54) with a 3 × 3 image-shift pattern and a nominal defocus range of -1.0 to -1.8 µm.

Each movie was recorded for 2.6 seconds, fractionated over 50 frames, yielding a total exposure

of 50 electrons/Å² and an exposure rate of 1 electron/Å² per frame. A total of 20,142 movies

(7,506 + 12,636) were collected from two grids for the CCL19-CCR7-G_i-scFv16 complex, while

5,968 movies (2,376 + 3,592) were collected from two grids for the CCL21-CCR7-G_i-scFv16

complex.

Cryo-EM data processing

The data were analyzed using cryoSPARC version 4 (55) for both signaling complexes unless otherwise noted, and the data processing schemes are summarized in Fig. S2 and S3. Patch motion correction and contrast transfer function (CTF) estimation were performed. Particles were picked iteratively with template-based picking using a representative GPCR-G protein complex and Topaz picking (56) combined with 2D classifications.

For the CCL19-CCR7-G_i-scFv16 complex (Fig. S2), 3,374,958 initial particles were extracted and subjected to Non-uniform Refinement (NU-refine) with Global and Local CTF Refinements. A representative GPCR-G protein-scFv16 complex was used as an initial reference map low-pass filtered to 30 Å. 3D classification was first performed to eliminate the complex particles with poor 7TM density. Several rounds of Heterogeneous Refinement (Hetero-Refine) and NU-Refine were performed to isolate a homogeneous subset of particles. We then utilized 3D Variability Analysis (3DVA) followed by Hetero-Refine using the resulting 3DVA volumes as references. While we observed heterogeneity related to glycosylation, such as at N292, we prioritized classes with the best-defined chemokine core features. The resulting class was further refined using Reference-Based Motion Correction (RBMC), NU-refine, and Local CTF Refinement. The selected 256,195 particles were subjected to the final NU-Refine, which yielded a map with a nominal resolution of 3.0 Å as determined by the gold-standard Fourier shell correlation (GSFSC) at the 0.143 threshold.

For the CCL21-CCR7-G_i-scFv16 complex (Fig. S3), 1,537,777 initial particles were processed similarly. After initial 2D and 3D classification, particles representing the best class and minor views from the second-best class were merged and subjected to iterative rounds of NU-Refine, RBMC, and Local CTF-Refine. Subsequent 3DVA-based Hetero-Refine were employed to address conformational heterogeneity at the chemokine core. Notably, we identified minor classes exhibiting weak or partial density corresponding to potential N-linked glycosylation at N36 and N292 of CCR7, as well as ambiguous density near the extracellular surface potentially representing parts of the flexible CCL21 C-terminal tail (Fig. S3A). However, these classes constituted a small fraction of the dataset and displayed lower homogeneity and resolution in

the core chemokine-receptor interface compared to the major population. Therefore, we
focused on the most homogeneous subset to achieve the highest local resolution (quality or
modelability) of the chemokine region. After several rounds of refinement and classification, the
final set of 195,011 particles yielded a map with a nominal resolution of 3.2 Å determined by
GSFSC at 0.143 threshold.

The maps were sharpened with a B-factor of -60 Å^2 for model building, refinement and
visualization of the map-model correlation. For both datasets, local resolution was estimated
using cryoSPARC, and the maps were further sharpened using deepEMhancer for overall map
presentation purposes only (57). UCSF pyEM v0.5 (58) was used to plot particle viewing directions
on the 3D maps.

Model building and refinement

The initial models were prepared using AlphaFold3 (59), and each subunit was roughly
fitted into the map by real-space rigid body refinement using Phenix (60). The models were
subjected to iterative rounds of manual rebuilding in Coot (61) and real-space refinement using
Phenix. Secondary structure restraints and geometry restraints were applied during refinement.
The map-model correlation and refinement statistics are summarized in Table S1 and Fig. S4.
Figures were prepared using UCSF ChimeraX (62) and PyMOL (Schrödinger, LLC).

Molecular dynamics simulations

Input models and parameters for all-atom MD simulations of CCR7 complexed with CCL19
or CCL21 were generated using CHARMM-GUI (63–65). All molecules other than CCR7 and
CCL19/21 were stripped from the cryo-EM models. The unmodelled sidechains were manually
built and refined with Coot (61). The internal water molecules were placed based on the
estimation by HOMOLWAT (66). The unmodelled part of the N-terminus and the C-terminus of
CCR7 and CCL19/21 were left unmodelled, and the terminal amino groups and the carboxyl
groups were neutrally capped by acetylation (ACE patch) and methylamidation (CT3 patch),
respectively. Protonation states of the titratable residues at pH 7.0 were determined with
PROPKA (67). The terminal amino groups of the native N-terminus of CCL19 (G1) and CCL21 (S1)

were deprotonated by NGNE patch and NNEU patch, respectively (estimated pKa 5.96 and 4.51).

Because the NGNE patch was not available in CHARMM-GUI, a local install of CHARMM c49b2

(68) was used for the patching using the scripts generated by CHARMM-GUI. All the aspartic acid

and glutamic acid residues were negatively charged except $D104^{2.50}$ (estimated pKa 7.50 in CCR7-

CCL19 and 7.81 in CCR7-CCL21). All the lysine and arginine residues were positively charged. All

the histidine residues were neutralized by δ-nitrogen protonation. The model was embedded in

a rectangular lipid bilayer comprising 203 POPC molecules (102 in the upper leaflet and 101 in

the lower leaflet) with the orientation estimated by PPM2.0 (69). The system was solvated and

charge-neutralized by ~24,000 water molecules (water layers of 22.5 Å thickness on top and

bottom of the proteins) and 150 mM NaCl. The system dimensions and the total number of atoms

were 90 Å × 90 Å × 141 Å and 107,178 atoms for CCR7-CCL19, 90 Å × 90 Å × 139 Å and 105,680

atoms for CCR7-CCL21 (Fig. S6A). The CHARMM36m force-field parameters (70) were used for

the proteins, lipids, and ions. The TIP3P model was used for water.

MD simulations were performed using GROMACS 2024.3 (71). Each system was energy-

minimized using the steepest-descent algorithm for 10,000 steps. Then, six independent MD runs

(replicas) were performed. For each replica, the system was equilibrated and relaxed in the

following way. First, temperature was elevated from 0 to 100 K for 20 ps in the NVT ensemble

using the V-rescale thermostat (72). Temperature was further elevated from 100 K to 310.15 K

under the pressure of 1 bar for 400 ps in the NPT ensemble using the V-rescale thermostat and

the Berendsen barostat (73). During the energy minimization and the temperature elevation, the

x, y, z positions of the backbone heavy atoms, the x, y, z positions of the sidechain heavy atoms,

the z positions of the POPC phosphorous atoms, the dihedral angles about the double bond in

the POPC alkyl chain, were harmonically restrained to the initial configuration at the force

constant of 4000 $kJ/mol/nm^2$, 2000 $kJ/mol/nm^2$, 1000 $kJ/mol/nm^2$ and 1000 $kJ/mol/rad^2$,

respectively. Then the restraints were gradually relaxed towards 0 for 35 ns in the NPT ensemble

of 310.15 K and 1 bar using the V-rescale thermostat and the C-rescale barostat (74). After that,

the production run was performed for 2.3 μs in the NPT ensemble of 310.15 K and 1 bar using

the V-rescale thermostat and the C-rescale barostat at the timestep of 2 fs. Bond lengths

involving hydrogen atoms were constrained using the LINCS algorithm (75). Long-range

electrostatic interactions were calculated with the particle mesh Ewald method (76). Atomic

coordinates were saved every 100 ps. The simulations were carried out using the TSUBAME4.0

supercomputer at Institute of Science Tokyo.

The molecules were made 'whole' and put back into the simulation cell using the gmx

trjconv program in the GROMACS package. The trajectory analysis was performed using

MDAnalysis v2.9.0 (77) and the graphs were visualized using Matplotlib (78). To align the

trajectories, the corresponding cryo-EM models were used as the alignment references, which

are transformed so that the (x, y) origin is the center of the proteins and the z origin is the center

of the membrane, and the z-axis is perpendicular to the membrane (Fig. S7B).

In all the trajectory analysis except the ICL1 conformational analysis (Fig. S9D), the

trajectories were least-squares aligned to the reference models by the Cα atoms of CCR7 except

the flexible N-terminal (-50) and ECL2 (200-209) regions. In the ICL1 conformational analysis (Fig.

S9D), the trajectories were aligned by the Cα atoms of the ICL1-flanking residues of TM1 and TM2

(78-85, 92-99).

In all the trajectory analysis except the simulation time course plots, the first 1 μs of the

2.3 μs simulation were excluded from the analysis as the relaxation (equilibration) period,

because the average RMSD (Fig. S6B) and the TM helices displacements (Fig. S7A) were rapidly

changing in this period.

The MD snapshots used in the figures were selected so that the TM centroid coordinates

and the H8 orientation (the angle and the TM2-G330 distance) are closest to the MD average (the

timestep 1597.8 ns of replica #1 for CCL19-CCR7, the timestep 1236.1 ns of replica #1 for CCL21-

CCR7).

The density plots in Fig. 5C and in Fig. S9B were calculated using the Gaussian kernel

density estimation implemented in Scikit-learn (79).

References

1. D. J. Fowell, M. Kim, The spatio-temporal control of effector T cell migration. *Nat Rev Immunol* **21**, 582–596 (2021).

2. H. Qi, W. Kastenmüller, R. N. Germain, Spatiotemporal basis of innate and adaptive immunity in secondary lymphoid tissue. *Annu Rev Cell Dev Biol* **30**, 141–167 (2014).

3. M. D. Gunn, *et al.*, Mice lacking expression of secondary lymphoid organ chemokine have defects in lymphocyte homing and dendritic cell localization. *J Exp Med* **189**, 451–460 (1999).

4. R. Förster, *et al.*, CCR7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* **99**, 23–33 (1999).

5. K. Reif, *et al.*, Balanced responsiveness to chemoattractants from adjacent zones determines B-cell position. *Nature* **416**, 94–99 (2002).

6. A. J. Ozga, M. T. Chow, A. D. Luster, Chemokines and the immune response to cancer. *Immunity* **54**, 859 (2021).

7. M. Birkenbach, K. Josefsen, R. Yalamanchili, G. Lenoir, E. Kieff, Epstein-Barr virus-induced genes: first lymphocyte-specific G protein-coupled peptide receptors. *J Virol* **67**, 2209 (1993).

8. R. Yoshida, *et al.*, Molecular cloning of a novel human CC chemokine EBI1-ligand chemokine that is a specific functional ligand for EBI1, CCR7. *Journal of Biological Chemistry* **272**, 13803–13809 (1997).

9. M. Nagira, *et al.*, Molecular cloning of a novel human cc chemokine secondary lymphoid- tissue chemokine that is a potent chemoattractant for lymphocytes and mapped to chromosome 9p13. *Journal of Biological Chemistry* **272**, 19518–19524 (1997).

10. R. Förster, A. C. Davalos-Misslitz, A. Rot, CCR7 and its ligands: balancing immunity and tolerance. *Nat Rev Immunol* **8**, 362–371 (2008).

11. I. Comerford, *et al.*, A myriad of functions and complex regulation of the CCR7/CCL19/CCL21 chemokine axis in the adaptive immune system. *Cytokine Growth Factor Rev* **24**, 269–283 (2013).

12. S. A. Luther, H. L. Tang, P. L. Hyman, A. G. Farr, J. G. Cyster, Coexpression of the chemokines ELC and SLC by T zone stromal cells and deletion of the ELC gene in the plt/plt mouse. *Proc Natl Acad Sci U S A* **97**, 12694–12699 (2000).

13. L. Han, L. Zhang, CCL21/CCR7 axis as a therapeutic target for autoimmune diseases. *Int Immunopharmacol* **121** (2023).

14. A. Salem, M. Alotaibi, R. Mroueh, H. A. Basheer, K. Afarinkia, CCR7 as a therapeutic target in Cancer. *Biochim Biophys Acta Rev Cancer* **1875** (2021).

15. K. Jaeger, *et al.*, Structural Basis for Allosteric Ligand Recognition in the Human CC Chemokine Receptor 7. *Cell* **178**, 1222-1230.e10 (2019).

16. M. A. Hauser, D. F. Legler, Common and biased signaling pathways of the chemokine receptor CCR7 elicited by its ligands CCL19 and CCL21 in leukocytes. *J Leukoc Biol* **99**, 869–882 (2016).

17. K. Schumann, *et al.*, immobilized chemokine fields and soluble chemokine gradients cooperatively shape migration patterns of dendritic cells. *Immunity* **32**, 703–713 (2010).

18. M. Weber, *et al.*, Interstitial dendritic cell guidance by haptotactic chemokine gradients.
Science **339**, 328−332 (2013).
19. G. M. Hjortø, *et al.*, Differential CCR7 targeting in dendritic cells by three naturally
occurring CC-chemokines. *Front Immunol* **7**, 218657 (2016).
20. T. A. Kohout, *et al.*, Differential desensitization, receptor phosphorylation, β-arrestin
recruitment, and ERK1/2 activation by the two endogenous ligands for the CC
chemokine receptor 7. *Journal of Biological Chemistry* **279**, 23214−23222 (2004).
21. M. A. Byers, *et al.*, Arrestin 3 mediates endocytosis of CCR7 following ligation of CCL19
but not CCL21. *J Immunol* **181**, 4723−4732 (2008).
22. D. A. Zidar, J. D. Violin, E. J. Whalen, R. J. Lefkowitz, Selective engagement of G protein
coupled receptor kinases (GRKs) encodes distinct functions of biased ligands. *Proc Natl*
Acad Sci U S A **106**, 9649−9654 (2009).
23. Y. Yanagawa, K. Onoé, CCL19 induces rapid dendritic extension of murine dendritic
cells. *Blood* **100**, 1948−1956 (2002).
24. S. A. Luther, *et al.*, Differing activities of homeostatic chemokines CCL19, CCL21, and
CXCL12 in lymphocyte and dendritic cell recruitment and lymphoid neogenesis. *J*
Immunol **169**, 424−433 (2002).
25. Ulrike Haessler, Marco Pisano, Mingming Wu, M. A. Swartz, Dendritic cell chemotaxis in
3D under defined chemokine gradients reveals differential response to ligands CCL21
and CCL19. *Proc Natl Acad Sci U S A* **108**, 5614−5619 (2011).
26. A. S. Jørgensen, M. M. Rosenkilde, G. M. Hjortø, Biased signaling of G protein-coupled
receptors − From a chemokine receptor CCR7 perspective. *Gen Comp Endocrinol* **258**,
4−14 (2018).
27. X. Barros-Álvarez, *et al.*, The tethered peptide activation mechanism of adhesion
GPCRs. *Nature* **604**, 757−762 (2022).
28. Y. L. Liang, *et al.*, Dominant Negative G Proteins Enhance Formation and Purification of
Agonist-GPCR-G Protein Complexes for Structure Determination. *ACS Pharmacol Transl*
Sci **1**, 12−20 (2018).
29. A. S. Dixon, *et al.*, NanoLuc Complementation Reporter Optimized for Accurate
Measurement of Protein Interactions in Cells. *ACS Chem Biol* **11**, 400−408 (2016).
30. J. Duan, *et al.*, Cryo-EM structure of an activated VIP1 receptor-G protein complex
revealed by a NanoBiT tethering strategy. *Nat Commun* **11** (2020).
31. S. Maeda, *et al.*, Development of an antibody fragment that stabilizes GPCR/G-protein
complexes. *Nat Commun* **9** (2018).
32. Z. Shao, *et al.*, Molecular insights into ligand recognition and activation of chemokine
receptors CCR2 and CCR3. *Cell Discov* **8** (2022).
33. K. Liu, *et al.*, Structural basis of CXC chemokine receptor 2 activation and signalling.
Nature **585**, 135−140 (2020).
34. D. J. Wasilko, *et al.*, Structural basis for CCR6 modulation by allosteric antagonists. *Nat*
Commun **15** (2024).
35. Q. Zhou, *et al.*, Common activation mechanism of class A GPCRs. *Elife* **8** (2019).
36. L. Urvas, E. Kellenberger, Structural Insights into Molecular Recognition and Receptor
Activation in Chemokine-Chemokine Receptor Complexes. *J Med Chem* **66**, 7070−7085
(2023).

37.   M. M. Rosenkilde, T. Benned-Jensen, T. M. Frimurer, T. W. Schwartz, The minor binding
pocket: A major player in 7TM receptor activation. *Trends Pharmacol Sci* **31**, 567−574
(2010).

38.   A. S. Jørgensen, *et al.*, Biased Signaling of CCL21 and CCL19 Does Not Rely on N-
Terminal Differences, but Markedly on the Chemokine Core Domains and Extracellular
Loop 2 of CCR7. *Front Immunol* **10** (2019).

39.   Q. Chen, *et al.*, Structures of rhodopsin in complex with G-protein-coupled receptor
kinase 1. *Nature* **595**, 600−605 (2021).

40.   J. Duan, *et al.*, GPCR activation and GRK2 assembly by a biased intracellular agonist.
Nature **620**, 676−681 (2023).

41.   A. Link, *et al.*, Fibroblastic reticular cells in lymph nodes regulate the homeostasis of
naive T cells. *Nat Immunol* **8**, 1255−1265 (2007).

42.   H. E. Kato, *et al.*, Conformational transitions of a neurotensin receptor 1-Gi1 complex.
Nature **572**, 80−85 (2019).

43.   Y. Gao, *et al.*, Structures of the Rhodopsin-Transducin Complex: Insights into G-Protein
Activation. *Mol Cell* **75**, 781-790.e3 (2019).

44.   M. R. Britschgi, A. Link, T. K. A. Lissandrin, S. A. Luther, Dynamic modulation of CCR7
expression and function on naive T lymphocytes in vivo. *J Immunol* **181**, 7681−7688
(2008).

45.   S. Nandagopal, D. Wu, F. Lin, Combinatorial guidance by CCR7 ligands for T lymphocytes
migration in co-existing chemokine fields. *PLoS One* **6** (2011).

46.   K. Schaeuble, *et al.*, Perivascular Fibroblasts of the Developing Spleen Act as LTα1β2-
Dependent Precursors of Both T and B Zone Organizer Cells. *Cell Rep* **21**, 2500−2514
(2017).

47.   K. Y. Chen, *et al.*, Inflammation switches the chemoattractant requirements for naive
lymphocyte entry into lymph nodes. *Cell* **188**, 1019-1035.e22 (2025).

48.   A. Goehring, *et al.*, Screening and large-scale expression of membrane proteins in
mammalian cells for structural studies. *Nat Protoc* **9**, 2574−2585 (2014).

49.   P. A. Longo, J. M. Kavran, M. S. Kim, D. J. Leahy, Transient mammalian cell transfection
with polyethylenimine (PEI). *Methods Enzymol* **529**, 227−240 (2013).

50.   A. Wellner, *et al.*, Rapid generation of potent antibodies by autonomous hypermutation
in yeast. *Nat Chem Biol* **17**, 1057−1064 (2021).

51.   N. Tsutsumi, *et al.*, Atypical structural snapshots of human cytomegalovirus GPCR
interactions with host G proteins. *Sci Adv* **8** (2022).

52.   N. Tsutsumi, *et al.*, Structure of the thrombopoietin-MPL receptor complex is a
blueprint for biasing hematopoiesis. *Cell* **186**, 4189-4203.e22 (2023).

53.   A. Dukkipati, H. H. Park, D. Waghray, S. Fischer, K. C. Garcia, BacMam system for high-
level expression of recombinant soluble and membrane glycoproteins for structural
studies. *Protein Expr Purif* **62**, 160−170 (2008).

54.   M. Schorb, I. Haberbosch, W. J. H. Hagen, Y. Schwab, D. N. Mastronarde, Software tools
for automated transmission electron microscopy. *Nat Methods* **16**, 471−477 (2019).

55.   A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290−296 (2017).

56.   T. Bepler, *et al.*, Positive-unlabeled convolutional neural networks for particle picking in
cryo-electron micrographs. *Nat Methods* **16**, 1153–1160 (2019).

57.   R. Sanchez-Garcia, *et al.*, DeepEMhancer: a deep learning solution for cryo-EM volume
post-processing. *Commun Biol* **4** (2021).

58.   D. Asarnow, E. Palovcak, Y. Cheng, asarnow/pyem: UCSF pyem v0.5. [Preprint] (2019).
Available at: https://doi.org/10.5281/zenodo.3576630.

59.   J. Abramson, *et al.*, Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493–500 (2024).

60.   D. Liebschner, *et al.*, Macromolecular structure determination using X-rays, neutrons
and electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861–
877 (2019).

61.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta
Crystallogr D Biol Crystallogr* **66**, 486–501 (2010).

62.   E. F. Pettersen, *et al.*, UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70–82 (2021).

63.   S. Jo, T. Kim, V. G. Iyer, W. Im, CHARMM-GUI: a web-based graphical user interface for
CHARMM. *J Comput Chem* **29**, 1859–1865 (2008).

64.   J. Lee, *et al.*, CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM,
and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J
Chem Theory Comput* **12**, 405–413 (2016).

65.   E. L. Wu, *et al.*, CHARMM-GUI Membrane Builder toward realistic biological membrane
simulations. *J Comput Chem* **35**, 1997–2004 (2014).

66.   E. Mayol, *et al.*, HomolWat: a web server tool to incorporate "homologous" water
molecules into GPCR structures. *Nucleic Acids Res* **48**, W54–W59 (2020).

67.   D. C. Bas, D. M. Rogers, J. H. Jensen, Very fast prediction and rationalization of pKa
values for protein-ligand complexes. *Proteins* **73**, 765–783 (2008).

68.   B. R. Brooks, *et al.*, CHARMM: the biomolecular simulation program. *J Comput Chem*
30, 1545–1614 (2009).

69.   M. A. Lomize, I. D. Pogozheva, H. Joo, H. I. Mosberg, A. L. Lomize, OPM database and
PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res*
40 (2012).

70.   J. Huang, *et al.*, CHARMM36m: an improved force field for folded and intrinsically
disordered proteins. *Nat Methods* **14**, 71–73 (2017).

71.   D. Van Der Spoel, *et al.*, GROMACS: fast, flexible, and free. *J Comput Chem* **26**, 1701–
1718 (2005).

72.   G. Bussi, D. Donadio, M. Parrinello, Canonical sampling through velocity rescaling. *J
Chem Phys* **126** (2007).

73.   H. J. C. Berendsen, J. P. M. Postma, W. F. Van Gunsteren, A. Dinola, J. R. Haak,
Molecular dynamics with coupling to an external bath. *J Chem Phys* **81**, 3684–3690
(1984).

74.   M. Bernetti, G. Bussi, Pressure control using stochastic cell rescaling. *J Chem Phys* **153**
(2020).

75.   B. Hess, P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation. *J Chem
Theory Comput* **4**, 116–122 (2008).

76.   T. Darden, D. York, L. Pedersen, Particle mesh Ewald: An N·log(N) method for Ewald
835   sums in large systems. *J Chem Phys* **98**, 10089–10092 (1993).
836   77.   N. Michaud-Agrawal, E. J. Denning, T. B. Woolf, O. Beckstein, MDAnalysis: a toolkit for
837   the analysis of molecular dynamics simulations. *J Comput Chem* **32**, 2319–2327 (2011).
838   78.   J. D. Hunter, Matplotlib: A 2D graphics environment. *Comput Sci Eng* **9**, 90–95 (2007).
839   79.   F. Pedregosa, *et al.*, Scikit-learn: Machine Learning in Python. *Journal of Machine
840   Learning Research* **12**, 2825–2830 (2011).

841
842

Materials and Data availability

• All unique/stable reagents generated in this study are available from the corresponding author upon formal publication by request, provided there is no licensing issue.

• All standardized datasets are publicly available as follows. The cryo-EM maps (the model coordinates) have been deposited in the Electron Microscopy Data Bank (Protein Data Bank) under accession codes of EMD-66874 (PDB: 9XHH) and EMD-66875 (PDB: 9XHI) for CCL19-CCR7-G_i-scFv16 and CCL21-CCR7-G_i-scFv16, respectively.

• All data were analyzed with standard or reported programs and packages, as detailed in the Methods. This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the corresponding author upon reasonable request following formal publication.

Acknowledgement

We thank the members of Cellular and Structural Physiology Laboratory (CeSPL) for daily scientific discussion and Ms. Miho Sasaki for supporting the research. This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI (JP24K01965 and JP24K21935 to N.T. and JP20H00451 to Y.F). The MD simulations were performed using the TSUBAME4.0 supercomputer at Institute of Science Tokyo.

Fig. 1

Fig. 1 | Temporally controlled mammalian expression of the CCR7-G$_i$ complexes for cryo-EM analysis.

(A) Schematic representation of the temporally controlled expression system in inducible Expi293F cells. Plasmids encoding FLAG-CCR7-LgBiT (R; inducible, pcDNA5/TO) and G$_i$ heterotrimer components [constitutive, pEG; Gγ2-Gα$_{i1}$(DN) (G1) and Gβ$_1$-HiBiT (G2)] are co-transfected. Cells are cultured at 37°C for constitutive G$_i$ expression, followed by induction with doxycycline (dox) and temperature shift to 30°C for induced CCR7 expression and sustained G$_i$ expression. DN denotes dominant negative mutations (S47N, G203A, E245A, A326S) in Gα$_{i1}$. (B) Workflow for the purification of the CCR7-G$_i$-scFv16 complexes, highlighting the strategy of purifying the apo-complex followed by the addition of chemokines. (C) Size-exclusion chromatography (SEC) profile of the ligand-free CCR7-G$_i$-scFv16 complex. The peak fraction used for subsequent steps is indicated by the dashed box. Chemokines are added to this sample for cryo-EM analysis. (D) Selected 2D class averages (scale bar 50 Å) of the CCL19-CCR7-G$_i$-scFv16 complex (left) and the CCL21-CCR7-G$_i$-scFv16 complex (right).

 Fig. 2

Fig. 2 | Structures of CCR7-G$_i$ complexes and global conformational changes upon Activation.
(A) Cryo-EM maps of the CCL19-CCR7-G$_i$-scFv16 (left) and CCL21-CCR7-G$_i$-scFv16 (right) complexes. Structures are shown in side views and top views. Color coding: CCL19 (red), CCL21 (orange), CCR7 (green), Gα$_i$ (yellow; RHD: Ras Homology Domain, AHD: Alpha Helical Domain), Gβ (cyan), Gγ (purple), scFv16 (gray). (B) Superposition of the ribbon models of the CCL19-bound and CCL21-bound complexes. Alignment was performed based on CCR7. Cα root-mean-square deviations (RMSD) are indicated for the chemokines, CCR7, and Gαβγ. (C) Comparison of active and inactive state structures of CCR7 and related chemokine receptors. Top row: Overall structures showing G protein coupling or antagonist/inhibitor binding.

Caption continued on the next page.

Fig. 2 *continued*:

Structures include inactive CCR7 bound to Cmp2105 (PDB: 6QZH); active CCR7 bound to CCL19

(this study); inactive CCR6 bound to OXM2 and SQA1 (PDB: 9D3E); active CCR6 bound to CCL20

(PDB: 6WWZ); inactive CCR2 bound to BMS-681 and CCR2-RA-[R] (PDB: 5T1A); active CCR2 bound

to CCL2 (PDB: 7XA3); inactive CXCR2 bound to 00767013 (PDB: 6LFL); active CXCR2 bound to

CXCL8 (PDB: 6LFO). (D) Superimposition of the inactive and active structures of chemokine

receptors. Inactive states are shown in gray, and active states are separately colored. Red arrows

indicate the outward movement of TM6. The region corresponding to the conserved CWxP motif

(FQxP in CCR7 and CCR6) is indicated by red dashed circles and shown on the right.

Fig. 3 | Distinct ligand binding modes of CCL19 and CCL21 at CCR7.

(A) Superposition of the CCL19-CCR7 (red-green) and CCL21-CCR7 (orange-cyan) complexes, highlighting the 17.6° rotation of CCL21 relative to CCL19. The chemokine recognition sites (CRS1, CRS1.5, CRS2, CRS3) are indicated by the dashed boxes. (B) Divergent interactions at the CRS3 interface. CCL19-CCR7 (left), CCL21-CCR7 (center), and inactive CCR7 (right, gray) are shown side-by-side to compare the extracellular structures especially the interaction between the chemokine 30s loop and CCR7-ECL2 at CRS3. A sequence alignment of the 30s loops of CCL19 and CCL21 is displayed on top of the inactive CCR7 structure.   *Caption continued on the next page.*

Fig. 3 *continued*:
(C) Conserved and distinct interactions at the CRS2/3 interfaces. Close-up views of the chemokine
N-terminus penetration into the orthosteric binding pocket for CCL19 (left) and CCL21 (right). Key
interacting residues are shown as sticks, and polar interactions are indicated by dashed lines. (D)
Ligand-specific conformational changes in the extracellular TMs and ECL2. Superposition of the
CCL19-bound and CCL21-bound CCR7 structures with the inactive CCR7 structure, highlighting
the conformational differences induced by the distinct ligand binding modes. Red arrows indicate
conformational shifts from the inactive state to the CCL21-bound state, and peach arrows
indicate the further shifts observed in the CCL19-bound state. (E) Chemokine footprints on the
CCR7 binding pocket. Top views of the CCR7 surface colored according to the distance to CCL19
(left) or CCL21 (right). Residues within 4 Å of the respective chemokine are colored. $Y136^{3.32}$ and
$Y308^{7.35}$ are shown as markers. (F) Rearrangement of the aromatic cluster in the middle layer of
the TM bundle. Close-up view of the superimposed structures showing the distinct
conformations of $Y65^{1.39}$, $F107^{2.53}$, $Y136^{3.32}$, $Y279^{6.51}$, $Y308^{7.35}$, and $Y312^{7.39}$ in the CCL19-bound
and CCL21-bound states compared to the inactive CCR7 (PDB: 6QZH). See also Fig. S5.

Fig. 4

Fig. 4 caption on the next page.

Fig. 4 | Structural basis of CCR7 activation.

(A) Conserved GPCR activation motifs and conformational changes in CCR7. Top left: Superposition of inactive CCR7 (gray) and active CCR7 (CCL19-bound in green, CCL21-bound in cyan). Top center: Mapping of the key GPCR activation motifs highlighted on the inactive CCR7 structure. Other panels show close-up views of the reorganization of the motifs upon activation. Key residues are shown as sticks. Arrows indicate the direction of movement from the inactive to the active state. Top right: Collapse of the sodium-binding site and reorganization of the $NP^{7.50}xxY$ motif. Bottom left: Reorganization of the PIF-like $P^{5.50}$-$M^{3.40}$-$F^{6.44}$ motif upstream of the hydrophobic lock. Bottom center: Collapse of the hydrophobic lock. Bottom right: Reorganization of the $DR^{3.50}Y$ motif and formation of a G protein-binding interface with the $R^{3.50}$-$Y^{7.53}$ bridge. (B) G protein coupling and comparison with the antagonist-bound state. Top left: Superposition of active CCR7 (CCL19-bound in green, CCL21-bound in cyan) and inactive CCR7 bound to Cmp2105 (gray). The inactive-to-active TM movements, which are generally conserved between the CCL19-bound and CCL21-bound states in complex with G_i, are indicated by red arrows. Top right: Close-up view showing how Cmp2105 (yellow sticks) occupies the G protein binding site in the inactive CCR7. Cmp2105 also inhibits the inactive-to-active conformational transition. Bottom panels: Detailed interactions between the $G\alpha_i$ $\alpha5$ helix (yellow, C-terminal 7 residues) and the CCR7 intracellular cavity in the CCL19-bound (left) and CCL21-bound (right) complexes. Key residues involved in the interaction are shown as sticks, and polar contacts are indicated by dashed lines.

Fig. 5

Fig. 5 | Conserved G$_i$ interface but distinct intracellular dynamics revealed by MD simulations
The G$_i$ binding interface of the (A) CCL19-CCR7-G$_i$ and (B) CCL21-CCR7-G$_i$ complexes. Overall
views are displayed on top right, where zoomed-in boxes highlight the interfaces centering ICL2
(top left), ICL3 (bottom left), and ICL1 (bottom right). The contacting residues within 4 Å distance
are shown as sticks, and polar interactions are shown by dashed lines. Cholesterol molecules near
ICL1 are shown as gray sticks. (C) Comparison of the H8 orientation after the G$_i$ free MD
simulations between CCL19-bound (green) and CCL21-bound (cyan) CCR7. Left: Probability
density plot of the H8 angle versus the T93$^{2.39}$-G330$^{8.47}$ distance. Data were calculated from six
independent 2.3 µs G$_i$-free MD simulations, excluding the first 1 µs of each run as a relaxation
period. See Fig. S8C for the definition of the H8 angle. The angles and the distances from the cryo-
EM models and the MD density peak are indicated by black ('x') and red ('x') markers with
corresponding text, respectively. *Caption continued on the next page.*

Fig. 5 *continued*:
Right: Superposition of the Gi free MD snapshot and the Gi-coupled cryo-EM structure. The
residues used for the distance measurement in the density plot, T93$^{2.39}$ and G330$^{8.47}$, are shown
as sticks and spheres, respectively. The larger degree of the H8 rotation and the consequent
insertion of the TM7-H8 loop open the gap between H8 and TM6 in the CCL19-bound CCR7
compared to the CCL21-bound CCR7.

Fig. 6

Fig. 6 | Comparison of G protein and GRK binding modes and a structural model for CCR7 biased agonism.

(A) Cryo-EM structures of Neurotensin Receptor 1 (NTSR1) in complex with G_i (PDB: 6OS9, left) and GRK2 (PDB: 8JPB, right). The zoomed-in inset shows a superposition of the two complexes, aligned on the receptor (NTSR1), to directly compare the binding modes of the C-terminal α5 helix of the Gα subunit (yellow) and the N-terminal αN helix of the GRK (blue). (B) Cryo-EM structures of Rhodopsin (Rho) in complex with G_t (PDB: 6OY9, left) and GRK1 (PDB: 7MTA, right). The zoomed-in inset shows a superposition of the Rho-G_t and Rho-GRK1 complexes, aligned on the receptor (Rhodopsin), to compare the effector helices. Both (A) and (B) highlight that while both effectors engage the receptor's cytoplasmic core, the GRK-αN helix exhibits a shallower insertion than the Gα-α5 helix (yellow) and follows a distinct trajectory, exiting through the interface formed by TM6, TM7, and H8.

Fig. 6 *continued*:
(C) Proposed structural model for the biased activation of GRK2/3 by CCR7. Cryo-EM structure of
the CCL19-CCR7-G_i complex (this study, left) and a superimposed model illustrating the proposed
mechanism of CCR7-GRK2 engagement (right). The model was generated by superimposing a
representative "open" conformation of CCL19-CCR7 (green), derived from MD simulations (Fig.
5C and S8B), onto the NTSR1 receptor of the cryo-EM NTSR1-GRK2 complex (PDB 8JPB). The
zoomed-in inset compares the G_i-bound state (gray helices, Gα-α5 in yellow) with the modeled
'open' GRK-bound state (green helices, GRK-αN in blue). This illustrates that the "open" state,
stabilized by CCL19-induced H8 dynamics after G protein dissociation, creates a wide gap
between TM6 and H8, which appears sterically complementary to accommodate the GRK-αN
helix. This simple model suggests that CCL19-induced dynamics selectively favor this "open"
ensemble, facilitating efficient GRK2/3 recruitment and robust β-arrestin signaling.