TITLE

The BRCA1- RAD51 Axis Regulates SCAI/REV3 Dependent Replication Fork Maintenance

AUTHOR LIST

Chloe Unterseher[1, #], Hiroshi Tsuchida[1, #], Rosevalentine Bosire[1], Xuhui Jin[1], Shaylee Kieffer[1] and Richard O. Adeyemi[1,2]*

AFFILIATIONS

[1]Basic Sciences Division, Fred Hutchinson Cancer Center, Seattle, WA 98109

[2]Department of Biochemistry, University of Washington, Seattle WA 98195

# These authors contributed equally.

* Corresponding author:

Richard Adeyemi, D.V.M., Ph.D.,

Basic Sciences Division, Fred Hutchinson Cancer Center,

1100 Fairview Ave, Mail Stop A1-162, Seattle, WA 98109, U.S.A.

Phone (office): (206)667-6263

Email: radeyemi@fredhutch.org

23   ABSTRACT
24
Replication stress is a key contributor to genomic instability and cancers. BRCA1 has
well established roles in protecting stalled forks against degradation. Here we show that
BRCA1 has a fork protection-independent role in driving stalled forks restoration via
regulation of RAD51 in a manner that confers dependency on SCAI and REV3. BRCA1
drives SLX4-mediated DNA break formation in the absence of SCAI and REV3, which
are required to complete repair synthesis and fork restart. Domain analysis revealed
that the increased fork breakage seen in the absence of SCAI requires the coiled coil
domain of BRCA1 and is regulated in a manner distinct from BRCA1's roles in resection
control at two-ended double strand breaks. BRCA1's role also occurs independently of
53BP1 but is modulated by RIF1. Surprisingly, loss of fork reversal factors leads to
increased fork breakage in the absence of SCAI/REV3 in a manner that depends on
RAD51 activity. SCAI loss leads to persistent DNA breaks, genomic instability and cell
death upon DNA damage. We posit that SCAI/REV3 may be required to replicate
through and rescue stalled replication forks at fragile genomic regions. Failure to do so
leads to increased DNA breakage and genomic instability.
40
41   Keywords: DNA repair, replication stress, homologous recombination, BRCA1, RAD51,
42   REV3, SCAI, 53BP1, RIF1, SMARCAL1, SLX4, fork reversal, fork restart, fork collapse
43

INTRODUCTION

Faithful duplication of the genome of dividing cells is continually threatened by impediments that challenge replication fork progression [1]. Such obstacles can result from lesions arising from exogenous DNA damage or from endogenous sources such as DNA secondary structures, difficult-to-replicate chromatin, oncogene-driven hyper-replication, and transcription-replication conflicts, among others [2]. Collectively, these inputs result in replication stress and, if left unresolved, drive fork stalling, breakage, and collapse with subsequent genome instability—a key hallmark of several cancers [3, 4]. To avert these outcomes, cells deploy a coordinated replication-stress response that stabilizes perturbed forks, remodels fork architecture, and re-establishes DNA synthesis through pathway choices that include repriming, translesion synthesis (TLS), template switching, and recombination-based restart [5].

A central remodeling route at stalled forks is fork reversal—conversion of the stalled fork into a four-way duplex intermediate—catalyzed by annealing helicases/ATP-dependent translocases such as SMARCAL1, ZRANB3, and others [6, 7]. Fork reversal can protect against further damage, yet reversed forks are intrinsically vulnerable. Homologous recombination (HR) factors, most notably RAD51 and BRCA1/BRCA2, extend their canonical roles beyond two-ended double strand break (DSB) repair to stabilize and protect nascent strands at reversed forks and to promote efficient restart [8, 9]. However, the varied functions of these proteins at stalled forks and how these differ from conventional two-ended breaks is still not clear.

In addition, how stalled forks transition back to productive replication remains an area of active investigation [10, 11]. In addition to RAD51's fork protective function, several studies have shown that RAD51 drives reversal of stalled forks and is important for promoting fork restart [12, 13]. Forks can restart via repriming [14] (which suppresses fork reversal and leaves post-replicative single-stranded DNA gaps to be filled later by template switching or TLS) or via direct RECQ1 activity on reversed forks [15]. Alternatively, stalled or reversed forks can be cleaved by structure-specific endonucleases (e.g., MUS81–EME1, SLX1–SLX4), generating one-ended DSBs that are repaired via recombination-based restart [16, 17]. In mammalian cells, such restart often resembles break-induced replication (BIR)—a conservative, long-tract synthesis program that, in several contexts, operates with minimal contribution from RAD51's strand-exchange activity [18, 19]. Alternatively, RAD51 dependent recombination restart mechanisms without break formation have also been proposed [20, 21].

We previously identified a general requirement for the Protexin complex—comprising SCAI and the Pol ζ catalytic subunit REV3—in promoting viability after replicative damage [22]. In particular, we uncovered a TLS-independent function of SCAI/REV3 in the response to DNA interstrand crosslinks (ICLs), where SCAI restrains excessive EXO1-mediated resection and preserves replication competence. These findings, however, suggested that SCAI/REV3 does more than modulate a single nuclease axis at ICLs; rather, it raised the possibility that SCAI/REV3 integrates multiple dimensions of stalled-fork metabolism. Here, we investigate how the SCAI/REV3 complex regulates

fork progression after replication stress and how the complex interfaces with the
RAD51/BRCA1 module to govern outcomes at stalled forks. We show that SCAI/REV3
is required to sustain DNA synthesis during replication stress and to promote efficient
fork restart. Strikingly, in the absence of SCAI, there is increased replication stress
associated fork breakage, genomic instability and cell death. Fork breakage in the
absence of SCAI/REV3 is orchestrated by BRCA1-RAD51 via SLX4 recruitment.
Remarkably, SCAI loss appears to confer a dependency on fork reversal, as combined
loss of SMARCAL1 and SCAI leads to increased fork breakage. We also define
requirements for BRCA1's role in promoting increased breakage and resection at stalled
forks and show that this occurs independent of 53BP1 regulation but is modulated by
RIF1. We propose that SCAI loss causes a failure of strand invasion mediated fork
restart with resultant fork breakage and genomic instability. By defining the contexts in
which BRCA1 and RAD51 are required to generate restart-competent intermediates, we
provide a mechanistic framework that clarifies how cells regulate repair outcomes upon
fork stalling.

RESULTS
SCAI promotes integrity and efficient restart of stalled replication forks
SCAI demonstrably regulates stalled replication fork resection [22, 23]; however,
we also previously found that EXO1 loss only partially rescues viability in the absence of
SCAI, suggesting that SCAI might perform other roles at stalled forks beyond EXO1
regulation. One such role could be in the regulation of stalled fork processing and
restart. Using BrdU incorporation assays, we have previously observed reduced
incorporation of BrdU correlating to reduced DNA synthesis rates following release from
hydroxyurea (HU)-induced fork stalling when SCAI was depleted [22], suggesting that
SCAI might be important regulating fork recovery and progression after prolonged
stalling. To directly investigate SCAI's role in replication fork dynamics, we assayed for
potential fork restart defects in the absence of SCAI using the single molecule DNA fiber
technique. Although there is no clear consensus on which factors are key for promoting
restart, previous work has shown that fork restart following HU treatment is differentially
regulated depending on the duration of fork stalling [13, 24]. Thus, we first tested
whether SCAI might modulate restart under acute (2 hour) treatment conditions with a
moderate dose of HU in RPE1 cells. Under these conditions, we failed to detect any
differences in either fork restart efficiency, number of stalled forks, or new origin firing
(Figure S1A). Similar results were observed following depletion of SCAI's binding
partner, REV3 (Figure S1A).
A typical S-phase lasts for close to 8 hours with some regions of the genome
replicated late in the cycle. Prolonged fork stalling (8 plus hours) often leads to fork
breakage and RAD51 dependent restart mechanisms [20, 25]. We reasoned that SCAI
might be important for modulating repair outcomes upon prolonged replication stress. To
test this, we examined fork dynamics upon treatment with HU for 18 h. In this
experimental setup, because we were making use of an HU dose that does not
completely block DNA synthesis, and to allow for adequate tracking of restarted forks,
the green label was kept during the period of HU treatment. We observed that SCAI
loss led to reduced lengths of the second, red-labeled fibers suggesting that there was
defective fork restart in the absence of SCAI (Figure 1A, right panel). Knockdown of

REV3 also led to reduction in fork restart efficiency (Figure 1A, right panel). Because SCAI loss did not affect green fiber lengths under short HU treatment conditions (Figure S1A), our observation that SCAI/REV3 loss also caused reduced lengths of the green label under prolonged replication stress (Figure 1A, left panel) suggested that fork progression during or following conditions of replication stress was also compromised by loss of SCAI. To further understand SCAI's role in regulating fork progression under replication stress conditions, we switched our labeling strategy to introduce the second label along with HU treatment. Under these HU treatment conditions, the replicative forks advanced over 30 times slower compared to untreated conditions, such that 18-hour treatment led to about half the fork advancement as the initial 30-minute pulse (Figure 1B). Nevertheless, SCAI loss consistently led to reduced fork progression upon replicative stress induction (Figure 1B, right panels). SCAI loss affected fiber length whenever cells were undergoing replication stress both in dual-labeled (restarted) (Figure 1B, top panels) or single labeled (Figure 1B, lower panels) conditions.

It is known that fork stalling can lead to fork reversal with subsequent fork degradation [7, 26, 27]. Furthermore, we and others have previously reported that SCAI regulates EXO1 activity and degradation of stalled forks [22, 23]. These experiments are typically performed following absolute blocks to fork advancement. Notably, SCAI loss did not affect the length of green only fibers prior to HU treatment under these conditions (Figure 1B), suggesting this effect was not due to fork degradation. Nevertheless, to determine whether the reduced fork progression in the absence of SCAI was a result of increased fork degradation, we co-depleted EXO1 and SCAI and performed similar DNA fiber assays. As expected, SCAI loss led to reduction fork progression upon replication stress (Figure 1C, right panel) without affecting progression under untreated conditions (Figure 1C, left panel). Importantly, rather than rescue reduced fork lengths, co-depletion of EXO1 had no effect on fork progression in the absence of SCAI (Figure 1C, left and right panels). Taken together, these results demonstrate that SCAI was required for adequate fork progression upon replication stress and for proper restart of stalled forks. This function of SCAI is independent of its role in EXO1 regulation.

SCAI/REV3 limit fork breakage following replication fork stalling

Compromised fork progression under replicative stress may result from increased fork breakage and/or collapse, increased fork reversal, or a combination of these [28-30]. Altered replicative stress signaling can imply fork breakage. To examine how loss of SCAI impacted replication fork stalling, RPE1 cells were transfected with two different siRNAs to SCAI. Subsequently, forks were stalled using similar HU conditions as in previous experiments and blotted for replication stress markers. As expected, and consistent with what was observed following treatment with crosslinking agents [22], SCAI loss led to increases in CHK1 and RPA phosphorylation (Figure 2A). Importantly, upon SCAI depletion with either siRNA, we observed significantly increased phosphorylation of H2AX (hereafter γH2AX), typically a marker of double-stranded DNA breaks, without any changes to H2AX protein amounts (Figure 2A). To further examine this, we made use of U2OS cells engineered to knock out SCAI using CRISPR. In this cell type, we observed similar increases in γH2AX and RPA phosphorylated on S4/8 (hereafter pRPA) levels, another marker of replication-associated fork breakage [31, 32]

(Figure 2B). Upon HU treatment, the increased pRPA in ΔSCAI cells could be rescued by re-introducing SCAI (Figure 2C, compare lanes 2, 4 & 6), confirming that the increased fork breakage was due to SCAI loss. Next, we performed a time course analysis to determine at which point fork breakage occurs in the absence of SCAI. We had earlier failed to observe any effect of SCAI loss on fork restart after treatment with HU for 2 h (Figure S1A). Although SCAI loss led to slightly increased pChk1 at 4 h after HU treatment (Figure S2A, lanes 3 & 4), γH2AX and pRPA levels remained unchanged relative to controls (Figure S2A). However, by 18 hours, there was marked increase in γH2AX and pRPA levels, suggesting SCAI loss results in increased fork breakage upon prolonged fork stalling (Figure S2A).

To further explore SCAI's function during replication stress and to make sure this effect was not unique to HU treatment, we examined SCAI's role upon treatment with aphidicolin, a DNA polymerase alpha inhibitor that also causes replication stress and fork stalling [33, 34]. Similar to what was observed following HU treatment, SCAI depletion led to increased γH2AX and pRPA upon treatment with aphidicolin (Figure 2D), showing that this altered signaling resulted from SCAI dependency at distressed forks. To directly examine altered fork breakage upon SCAI loss, we performed neutral comet assays. Consistent with the Western blotting data, depletion of SCAI led to significant increases in fork breakage upon replication stress compared to controls in both RPE1 (Figure 2E) and U2OS cells (Figure 2F). Taken together, these data shows that SCAI loss results in compromised fork progression and increased fork breakage under replication stress.

To examine whether REV3 shared similar roles with SCAI in preventing fork breakage at HU stalled forks, we depleted REV3 using siRNAs. Similar to what was observed upon SCAI loss, REV3 depletion led to increases in pRPA and γH2AX upon fork stalling (Figure 2G, compare lanes 2 and 4), suggesting a similar function for REV3 in preventing fork breakage. Because HU does not directly lead to formation of DNA adducts, this role appeared to be independent of REV3's role in TLS across damaged DNA bases. To further examine this, we depleted cells of REV7, which is critical for TLS as a component of DNA polymerase zeta [35]. Whereas REV3 loss led to increases in pRPA and γH2AX after HU treatment, REV7 depletion (and thus inhibition of TLS) did not (Figure 2G, compare lanes 4 and 6). To directly test whether failure of TLS would result in increased fork breakage we treated cells with a previously reported TLS inhibitor [36]. Similar to REV7 depletion, TLS inhibition failed to increase pRPA and γH2AX after replication stress (Figure S2B) suggesting that increased fork breakage in the absence of REV3 was not attributable to a TLS function of REV3. Depletion of REV3 in SCAI knockout (KO) cells did not significantly alter the amounts of pRPA and γH2AX, suggesting that both proteins function in the same pathway (Figure S2C). As a direct marker of fork breakage, neutral comet assays showed that REV3 loss led to increased fork breakage, as did loss of SCAI (Figure 3G).

As an orthogonal approach, we examined replication stress signaling events using immunofluorescence (IF) microscopy. To determine whether the observed increased pRPA signified increases on chromatin, cells were pre-extracted with cytoskeleton buffer to remove unbound proteins. Depletion of either SCAI or REV3 led to increased intensity of RPA upon HU treatment, suggesting that increased pRPA occurred on chromatin (Figure S2D). We also observed increased γH2AX using this

approach (Figure S2E). Finally, cells knocked down for REV3 or SCAI were examined
for increases in chromatin-bound RPA or γH2AX using quantitative image-based
cytometry (QIBC). Upon HU treatment, QIBC showed increased γH2AX and RPA levels
indicative of increase fork breakage (Figure 2H-J).
Unlike ICL treatment, which typically blocks helicase unwinding, HU treatment
uncouples the helicase from the polymerase, thereby increasing RPA loading on the
resulting singe-stranded DNA (ssDNA) [37]. Thus, resection is unlikely to be the primary
driver of fork breakage. Since SCAI was previously shown to regulate EXO1, we
depleted EXO1 upon SCAI loss. Notably, and contrary to what was seen following
treatment with ICL agents [22], the increases in pRPA seen after HU treatment could not
be reversed by knockdown of EXO1 (Figure S2F) further demonstrating that SCAI
performs multiple roles at stalled forks and regulates stalled fork dynamics independent
of EXO1 regulation (see also Figure 1C).
236
BRCA1 promotes increased fork breakage and resection in the absence of
SCAI/REV3
We next hypothesized that BRCA1 might have a role in the increased pRPA and
γH2AX observed in the SCAI knockdown cells, as BRCA1 is important for resection
control at two ended DSBs and plays multiple roles in regulating stalled fork dynamics
[38, 39]. To examine the effect of BRCA1, we made use of RPE1 cell lines in which
BRCA1 had been eliminated using CRISPR and examined for alteration in pRPA and
γH2AX levels. As expected, SCAI depletion led to increased fork breakage upon HU
treatment (Figure 3A, compare lanes 2 & 3) in BRCA1 proficient RPE1 cells. Strikingly,
we observed that loss of BRCA1 abolished the increased γH2AX and pRPA levels seen
upon SCAI loss (Figure 3A, compare lanes 3 & 5). To further examine BRCA1's role, we
depleted BRCA1 in RPE1 cells using siRNA. Consistent with our KO cell data, the
increased pRPA and γH2AX seen upon SCAI loss was eliminated upon BRCA1
depletion (Figure S3A). BARD1 is a key interaction partner that forms a stable complex
with BRCA1 [40]. To further examine a potential role for BRCA1 in regulating stalled fork
breakage in the absence of SCAI, we depleted cells of BARD1. As expected, BARD1
knockdown led to diminished levels of BRCA1 (Figure S3A). Similar to what was seen
upon BRCA1 loss, BARD1 knockdown eliminated the increases in pRPA and γH2AX
seen whenever SCAI is lost (Figure S3A). Basal levels of pRPA and γH2AX were higher
in BARD1 knockdown vs BRCA1 knockdown cells for unclear reasons. To see if a
similar effect applied to REV3, we depleted SCAI and REV3 using siRNAs in WT or
BRCA1 KO cells. As shown earlier, we observed increases in damage signaling
markers indicative of fork breakage and resection upon loss of either SCAI or REV3
(Figure 3B, compare lane 3 with lanes 5 & 7). Importantly, cell lines lacking BRCA1
failed to show such increases (Figure 3B, compare 5 & 6 and lanes 7 & 8), suggesting
that BRCA1 mediates fork breakage in the absence of the SCAI/REV3 complex. These
findings were further confirmed using IF experiments, where loss of BRCA1 led to
reduced amounts of RPA on chromatin as well as decreased γH2AX levels, again
demonstrating a key role for BRCA1 in mediating fork breakage upon SCAI/REV3 loss
(Figure 3C – F).
To directly examine if BRCA1 regulates fork breakage after SCAI/REV3
depletion, we performed neutral comet assays. As expected, loss of either SCAI or

REV3 led to increased accumulation of DNA breaks (Figure 3G). Importantly, co-
depletion of BRCA1 with SCAI led to failure to accumulate DNA breaks (Figure 3G),
suggesting that the DNA breakage seen in the absence of SCAI/REV3 was orchestrated
by BRCA1 activity. These results could imply that SCAI/REV3 act downstream of
BRCA1 activity. SCAI binds to ssDNA [41], and if BRCA1's role in this context was to
promote DNA resection that may allow SCAI to be recruited to damaged chromatin. To
examine this, we treated WT or BRCA1 deficient cells with HU and performed chromatin
fractionation experiments. As expected, loss of SCAI led to increased DNA break
formation (Figure 3H, compare lanes 2 & 3), which failed to occur in the absence of
BRCA1 (Figure 3H, compare lanes 3 & 5). Importantly, SCAI accumulated in chromatin
fractions upon HU treatment (Figure 3H, compare lanes 1 and 2) but this increase in
SCAI accumulation was not affected by BRCA1 loss (Figure 3H, compare lanes 2 & 4).
Thus, BRCA1 likely performs a different function that confers a dependency on SCAI
activity or compensates for loss of SCAI by inducing fork breakage.
Resection at ICLs and upon HU treatment is often differentially regulated [42].
Although SCAI's role at ICLs was more impacted by regulation of EXO1 compared to
HU, we wanted to determine whether the increased resection observed in the absence
of SCAI was regulated by BRCA1 in both contexts. Similar to what was observed with
HU, the cisplatin induced increase in pRPA was abolished upon BRCA1 depletion
(Figure S3B, compare lanes 4 & 6). This effect was also seen following depletion of
BARD1 (Figure S3B). It is worth noting that following either HU or cisplatin treatment,
BRCA1 loss did not cause similar reductions in CHK1 activation (nor does it eliminate
the increased CHK1 phosphorylation (pCHK1) seen upon SCAI/REV3 loss), suggesting
that ATR activity *per se* (which regulates damage signaling upon replication stress) was
not affected by loss of BRCA1 (Figure 3B and Figure S3B). Taken together, our results
demonstrate that BRCA1 orchestrates fork breakage in the absence of SCAI/REV3
upon treatment with various agents that stall and/or damage replication forks.
296
BRCA1 requirements for regulating stalled fork breakage
BRCA1 forms several complexes that play important roles in resection control,
HR and other functions; these different complexes mediate the various functions of
BRCA1 [43, 44]. For example, the BRCA1-CtIP complex, formed via CtIP binding to the
C-terminus of BRCA1, controls resection at ionizing radiation-induced DSBs [45, 46].
Since BRCA1 controls fork breakage and pRPA upon SCAI loss, we wanted to
determine which subcomplexes and function of BRCA1 was important for this role.
These experiments were performed either following cisplatin or HU treatment and
depletion of ABRA1, BRIP1 or CtIP (key members of the BRCA1-A, B or C complexes)
respectively. Our data showed that loss of any of these factors did not prevent
increased pRPA in the absence of SCAI upon cisplatin treatment (Figure S4A).
Repeating a similar experiment following HU treatment showed similar results:
increased pRPA upon SCAI loss in each condition (Figure S4B). However, we did
observe that depletion of Abraxas by itself increased the basal level of pRPA under HU
treatment conditions (Figure S4B, lane 4). Such a role for ABRA1 (Abraxas) has been
reported following treatment with the topoisomerase inhibitor, camptothecin [47]. Taken
together, our data suggests that the three major BRCA1 complexes do not account for

the observed SCAI-depletion phenotype, but that its obligate binding partner BARD1 is involved.

Next, we examined domain requirements of BRCA1 for fork breakage using previously reported BRCA1 mutant expressing cell lines [48]. Increased BRCA1 dependent fork resection upon loss of either SCAI or REV3 could be almost fully rescued by reintroducing tagged WT-BRCA1 into knockout cell lines (Figure S4C). We then examined the role of two domain mutants of BRCA1. First, a point mutation (S1655A) in the BRCT domain of BRCA1 that abrogates its interaction with phosphorylation dependent binding partners (ABRA1, BRIP1 or CtIP). A second mutant deleted the C-terminal coiled coil domain that is essential for HR. As expected, BRCA1 deletion reduced pRPA levels (Figure 4A, lane 2). Reintroduction of the BRCA1^{S1665A} mutant allowed restoration of fork resection to almost the same extent as seen following reintroduction of WT-BRCA1 (Figure 4A, compare lanes 1 and 3), suggesting that this domain was not required for the BRCA1 induced fork resection phenotype. This finding is consistent with what was observed following siRNA depletion of ABRA1, BRIP1 or CtIP, whose depletion failed to abrogate BRCA1's role in promoting fork resection in the absence of SCAI (Figures S4A & S4B). In contrast, the coiled coil domain mutant (ΔCC) of BRCA1 did not fully restore fork resection (Figure 4B, compare lanes 1 & 3) suggesting that this domain was important for BRCA1's role in promoting stalled fork resection. To address this fully, we performed a dose curve of HU treatment and compared the ability of the WT BRCA1 and BRCA1$^{\Delta CC}$ mutants to rescue SCAI dependent fork breakage and resection (Figure 4C). These data confirmed that the coiled coil domain of BRCA1 was required to promote stalled fork resection and breakage in the absence of SCAI (Figure 4C, compare lanes 4&5, 6&7, 8&9). Note that these findings are different from the domain requirements of BRCA1 in promoting resection at two-ended DSBs, where binding to CtIP through the BRCT domain is critical for resection [49].

BRCA1 pro-breakage / resection roles at stalled and broken forks occur independently of 53BP1

Previous studies have shown that BRCA1's role in resection control at two-ended DSB sites is typically antagonized by 53BP1, which enforces a block to resection in the absence of BRCA1 [50, 51]. BRCA1 overcomes the 53BP1 block to resection upon S phase entry through newly-replicated chromatin-dependent extrusion of 53BP1 as well as CDK dependent activation of CtIP [52-54]. This has implications for efficacy of treatment with PARP inhibitors (PARPi), as PARPi induced cell death in the absence of BRCA1 can be overcome by loss of 53BP1 [55]. Interestingly, SCAI also binds to 53BP1; however, the reason for this association is not fully clear although it may involve regulation of RIF1 activity [56]. Fork breakage at stalled forks typically leads to single but not two-ended DSBs (unless there is replisome bypass of a lagging strand nick) [57], and so, to determine whether this role of BRCA1 was mediated by counteracting 53BP1 activity, we examined the effects of 53BP1 loss on resection control at stalled forks in the absence of SCAI.

Since BRCA1 drives fork breakage and resection both upon fork stalling and following ICL formation, we examined the phenotypes in both contexts. For simplicity, we focused on changes to resection as indicated by increases in pRPA. HU leads to

uncoupling of the replicative helicase from the polymerase and thus causes higher
background pRPA in the absence of any specific gene alterations (Figure S4D, lane 2).
Thus, we have largely focused on describing results seen upon cisplatin treatment and
how this agrees with or varies from what is observed following HU treatment. As
expected, SCAI KO cells showed increases in pRPA upon cisplatin treatment (Figure
4D, compare lanes 1 and 2) that was abolished by BRCA1 depletion (Figure 4D, lane
3). HU treatment behaved similarly (Figure S4D, lanes 2, 3 & 5). Depletion of 53BP1
alone using siRNAs failed to alter pRPA levels upon cisplatin treatment (Figure 4D, lane
4). Nevertheless, co-depletion of 53BP1 and SCAI led to slight but consistent additional
increases in pRPA upon cisplatin (Figure 4D, compare lanes 2 and 5) or HU (Figure
S4D, compare lanes 3 and 4) treatment suggesting a role for SCAI in restraining 53BP1
in this context. If BRCA1 was relieving a 53BP1-mediated block to resection like it does
at two-ended breaks, it then follows that co-depletion of 53BP1 and BRCA1 should
restore the pRPA in the absence of SCAI. However, this was not the case, as BRCA1
loss not only abolished the increased resection and/or fork breakage seen in the
absence of either SCAI or 53BP1, but it did so even when SCAI was co-depleted
alongside 53BP1. This was seen both following cisplatin treatment (Figure 4D, lane 5
and 7) and HU treatment (Figure S4D, lanes 4 and 6, see γH2AX levels as well). Taken
together, these results show thatBRCA1 mediated fork breakage and resection in the
absence of SCAI is not regulated or counteracted by 53BP1 activity, dissimilar to the
antagonism that occurs at two-ended DSBs.
381
RIF1 differentially modulates BRCA1 activity and resection at stalled forks and
ICLs
At two-ended DSBs, 53BP1 orchestrates its block to end resection through
downstream effectors such as RIF1, REV7 and the Shieldin complex – all of which are
recruited to these breaks in a 53BP1 dependent manner [58, 59]. However, at stalled
forks, RIF1 has been shown to be recruited independently of 53BP1 where it regulates
stalled fork stability [60]. In fact, 53BP1 has also been linked to regulation of fork
stability [61]. However, because we have data showing that BRCA1 acts differently, and
without 53BP1 regulation, at stalled forks than it does at DSBs, we also wanted to
determine whether RIF1 might behave differently than 53BP1 at stalled forks. To test
this, first we depleted 53BP1 and RIF1 using siRNAs and examined changes to
resection upon HU treatment. We observed that, while 53BP1 depletion only modestly
affected pRPA and γH2AX levels (Figure 4E, compare lanes 4 & 5), RIF1 depletion
caused increases in both markers (Figure 4E, compare lane 4 & 6), suggesting
increased fork breakage in the absence of RIF1. To further examine this, we determined
whether the increased fork breakage in the absence of RIF1 was modulated by SCAI
and/or BRCA1. Co-depletion of SCAI and RIF1 did not further alter damage signaling at
stalled forks (Figure S4E, lanes 4 & 5). However, loss of BRCA1 reduced signaling in
the absence of RIF1 in the presence or absence of SCAI (Figure S4E, lanes 6 & 7).
Finally, to determine whether this was a general function of RIF1 following replicative
damage, we performed similar experiments as above (Figure S4E) following cisplatin
treatment. In this context, and unlike what is seen following HU treatment, RIF1
depletion by itself had no effect on pRPA (Figure 4F, lane 4). Surprisingly, whereas
53BP1 depletion failed to restore resection in the absence of BRCA1 (Figure 4D, lane

7), loss of RIF1 fully restored resection upon cisplatin treatment in the absence of
BRCA1 (Figure 4F, compare lanes 3 and 7) suggesting a possible antagonism between
BRCA1 and RIF1 in this context. This phenotype was unique to RIF1 and was not
shared by depletion of REV7, which mediates assembly of Shieldin complex members
SHLD2 and SHLD3 [59, 62] (Figure S4F).
411
Loss of fork reversal at stalled forks confers a dependency on SCAI/REV3
We next wanted to determine at what step fork breakage occurs in the absence
of SCAI/REV3. Our data so far suggests that BRCA1 drives DNA breakage at stalled
forks upon replication stress. BRCA1 also has a well-established role in protecting
arrested forks from degradation [63]. This difference in phenotypes may be due not only
to the fact that forks keep progressing, albeit slowly, under our treatment conditions, but
also that the forks may be processed differently by remodeling enzymes. The fork
reversal activity of the enzymes SMARCAL1, ZRANB3 and HLTF represent a major
mechanism of stalled fork processing [6, 7]. Multiple studies have shown that when
forks progression is halted, fork reversal by these enzymes drives the creation of a
chicken-foot structure that is protected from breakage and resection by BRCA1 [26, 27],
a phenotype opposite to the BRCA1 activity we see under our conditions. We reasoned
that if the fork breakage seen in the absence of SCAI was occurring downstream of fork
reversal, it would be abolished upon depletion of fork reversal enzymes.
To determine whether fork reversal plays a similar role under our treatment
conditions, we co-depleted SCAI and either SMARCAL1 or ZRANB3. To our surprise,
rather than reduce fork breakage upon SCAI loss, knockdown of either SMARCAL1 or
ZRANB3 led to consistent increases in pRPA and γH2AX upon HU treatment (Figure
5A, compare lanes 2, 5 & 8). The same phenotype was also seen upon co-depletion of
REV3 and either SMARCAL1 or ZRANB3 (Figure 5A, compare lanes 3, 6 & 9). These
results were further validated by knocking down SMARCAL1 with two independent
siRNAs, which led to similar increases in the absence of SCAI (Figure 5B, compare
lanes 2, 4 & 6). This phenotype was shared between SMARCAL1 and ZRANB3 and
was seen with multiple siRNAs against SMARCAL1, suggested that not only was fork
breakage in the absence of SCAI not occurring downstream of fork reversal, but that the
process that leads to fork breakage might also compete with fork reversal for repair of
stalled forks. An alternative explanation might be that fork breakage occurs downstream
of a parallel fork reversal mechanism to that orchestrated by SMARCAL1/ZRANB3.
Such a parallel mechanism has been suggested to occur via FBH1 [61]. To examine
this, we knocked down FBH1 using siRNAs and observed that, similar to loss of
SMARCAL1, FBH1 depletion also led to increased pRPA and γH2AX in the absence of
SCAI (Figure 5C). Thus, depletion of multiple fork reversal enzymes all led to increased
fork breakage in the absence of SCAI/REV3. The increased break formation in the
absence of fork reversal was more marked in the absence of SCAI/REV3, which
suggests that a SCAI/REV3-dependent mechanism might compete with fork reversal as
a mechanism for stalled fork rescue.
448
RAD51 activity globally regulates stalled fork metabolism
RAD51 has been implicated in various fork processing roles in cells undergoing
replication stress. It is important for fork reversal, protecting stalled forks against the

activity of nucleases, and is key for recombination through template switching [25].
RAD51's recombination function can be genetically uncoupled from the other functions,
as multiple studies have shown that RAD51 strand exchange activity is not required for
fork protection [61, 64, 65]. To determine if and how RAD51 regulates the phenotypes
seen in the absence of SCAI/REV3, we first depleted RAD51 using a pool of siRNAs
(Figure S5A). Depletion of RAD51 abolished the increased pRPA and γH2AX seen in
the absence of SCAI (Figure S5A). We next tested RAD51's role using two independent
siRNAs and observe similar effects – loss of RAD51 abolished pRPA and γH2AX upon
HU treatment even when SCAI was absent (Figure 5D). To further determine how
RAD51 contributes to the increased pRPA and γH2AX seen upon fork stalling, we tested
the effect of inhibiting RAD51 strand exchange activity using a previously described
inhibitor B02 [66]. Surprisingly, RAD51 inhibition completely abolished the signaling
seen upon loss of SCAI or REV3 (Figure 5E, compare lanes 2&3 to lanes 5&6), without
affecting cell cycle distribution (Figure S5B), suggesting that this function requires or
occurs downstream of the strand exchange activity of RAD51. To directly test the effect
on DNA break formation, we performed neutral comet assays. RAD51 depletion
prevented DNA break formation in the absence of SCAI (Figure 5F).
We showed earlier that inhibition of fork reversal led to additive increases in fork
breakage upon SCAI loss. We were curious whether this increase was also modulated
by RAD51. Interestingly, RAD51 inhibition completely prevented the increased break
signaling seen in the absence of SCAI when fork reversal was prevented either via
SMARCAL1 knockdown (Figure 5G) or via PARP inhibition (Figure S5D) which has
been shown to modulate fork reversal outcomes [15]. Taken together these data
suggests that RAD51 strand exchange activity drives the formation of DNA breaks seen
in the absence of SCAI/REV3, and that failure of fork reversal leads to increased
RAD51 driven fork breakage when SCAI or REV3 is absent.
478
BRCA1-RAD51 axis governs SLX4 mediated fork breakage in the absence of SCAI
We next wanted to determine how fork breakage occurs in the absence of
SCAI/REV3. Cells have several structure specific nucleases that orchestrate fork
breakage [17]. In particular, MUS81 has been implicated in fork processing and restart
under various conditions [67, 68]. We co-depleted SCAI or REV3 cells with siRNAs
against SLX4 or MUS81 following treatment with HU. We found that loss of SLX4, but
not MUS81, led to reduced γH2AX amounts in SCAI depleted cells (Figure 6A). A similar
pattern was observed in REV3 depleted cells: MUS81 had no effect or led to moderate
increases in γH2AX, while SLX4 loss reduced γH2AX levels (Figure 6B). To directly
examine whether stalled fork breakage in the absence of SCAI was orchestrated by
SLX4, we performed neutral comet assays. As expected, SCAI depletion led to
increased comet tail moments which was significantly reduced by SLX4 depletion
(Figure 6C).
We next sought to determine whether and how SLX4 was recruited to stalled
forks and how this was modulated by loss of SCAI. Fractionation experiments revealed
that SCAI loss led to increased amounts of SLX4 in the chromatin fraction compared to
controls (Figure 6D, compare lanes 1 & 2). We did not observe such increases with
other nucleases such as MUS81 or MRE11 (Figure 6D). Strikingly, RAD51 inhibition led
to significant reduction in not only γH2AX levels, but also in the amount of SLX4 on

chromatin (Figure 6D), and the increased SLX4 amounts on chromatin seen in the
absence of SCAI was prevented by B02 treatment (Figure 6D, compare lanes 2 and 4).
Taken together, these experiments suggest that fork breakage in the absence of
SCAI/REV3 occurs downstream of RAD51 strand exchange activity but independent of
fork reversal.
To further characterize the mechanism by which stalled forks break, we depleted
BRCA1. Upon SCAI depletion, BRCA1 KO showed reduced chromatin levels of SLX4
compared to controls (Figure S6A); this finding is consistent with a prior report [24]. To
further examine this role, and to determine how it was modulated by RAD51, we
knocked down BRCA1 using siRNAs. HU treatment resulted in slightly increased
amounts of SLX4 on chromatin (Figure 6E, compare lanes 1 & 3). Similar to what was
seen following RAD51 inhibition or in BRCA1 KO cells, BRCA1 KD also resulted in
reduction in the amount of damage induced SLX4 on chromatin (Figure 6E, compare
lanes 1 & 3). Similar to what was seen following RAD51 inhibition or in BRCA1 KO cells,
BRCA1 KD also resulted in reduction in the amount of damage induced SLX4 on
chromatin (Figure 6E). Next, to determine how BRCA1 and RAD51 activities contributed
to fork breakage, we examined chromatin association of RAD51 and its modulation by
BRCA1. Upon treatment with HU, there was increased amounts of RAD51 on chromatin
(Figure 6E, compare lanes 1 & 3). Loss of BRCA1 significantly prevented the increase
in RAD51 amounts on chromatin (Figure 6E, lane 4). Next, we were curious to know
whether this effect of BRCA1 on RAD51 chromatin association could be modulated by
53BP1 loss under our conditions. Indeed, while BRCA1 loss almost completely
abolished RAD51 levels on chromatin (Figure 6F, compare lanes 1 & 2), co-depletion of
53BP1 partially restored the amount of RAD51 (compare lanes 2 and 4), suggesting
that the reduced RAD51 loading upon fork stalling was partly due to 53BP1 activity. This
effect was seen despite 53BP1 not affecting resection in our context (Figure 4D, 4E,
Figure S4D), and is consistent with recent results seen using CRISPR nickase
experiments [69] suggesting that 53BP1 antagonism of RAD51 loading is a general
effect of 53BP1 at broken forks. Taken together, our data show that the RAD51-BRCA1-
SLX4 axis drives break formation at stalled forks. BRCA1 controls RAD51 levels on
chromatin in this process and this can be counteracted by 53BP1.
529
SCAI promotes genomic integrity and viability following replication fork stalling
We have shown that SCAI loss leads to increased stalled fork breakage and
resection. We reasoned that unrepaired breaks in the absence of SCAI might persist
and contribute to genomic instability and cell death. To test this, we treated cells with
HU as before, then released from HU treatment and allowed cells to proceed into
mitosis (Figure 7A, top panel). We then performed mitotic DNA spreads to examine for
the presence of chromosomal abnormalities including breaks, gaps, fusions and similar
events. SCAI loss did not significantly impact the extent of chromosomal abnormalities
observed in the absence of HU treatment (Figure 7A). In contrast, following treatment
with HU, we observed that loss of SCAI led to significant increases in the amount of
broken and otherwise abnormal chromatids compared to control knockdown cells
(Figure 7A & B). To further examine SCAI's role, we performed colony formation assays.
SCAI depletion led to diminished viability upon treatment with HU (Figure 7C & D).
Taken together, these data show that SCAI is required for efficient repair and viability

following replication stress and that its loss results in persistence of unrepaired fork
breakage that leads to heightened genomic instability.
546
DISCUSSION
In this work, we have uncovered a novel fork restoration mechanism requiring the
activities of both the fork protecting factor SCAI and the DNA polymerase REV3 to
prevent pathologic fork breakage upon replication stress. Our experiments reveal key
roles for BRCA1 and RAD51, suggesting a homologous recombination mediated restart
mechanism. Indeed, we find requirements for both the HR-promoting coiled-coil domain
of BRCA1 as well as the strand exchange activity of RAD51. We propose that BRCA1-
RAD51 drive break-independent fork restart that confers a dependency on the
SCAI/REV3 complex (Figure 7E). Surprisingly, this process appears to occur either
parallel to or upstream of SMARCAL1/ZRANB3 or FBH1-mediated fork reversal.
557
In addition to its canonical roles in regulating resection and consequent pathway choice
at two-ended DSBs, BRCA1 has several proposed roles at stalled forks [70]. Its fork
protection functions are well established, occurring downstream of fork reversal. Work in
the literature has suggested roles for BRCA1 in stalled fork remodeling leading to fork
restart [71]. Our experiments allow us to uncouple BRCA1's role from a fork protection
function not only because BRCA1 loss paradoxically prevents fork breakage but also
due to a requirement for BRCA1's coiled coil motif, which is dispensable for fork
protection roles of BRCA1 [71]. Also, depletion of fork reversal factors, which are
necessary to provide a substrate for fork protection, does not abolish the DNA break
formation seen in the absence of SCAI but rather increases it. We also show that this
role occurs independently of or distinct from BRCA1's known roles at two-ended DSBs:
the C-terminus of BRCA1 is dispensable for promoting fork breakage and processing,
and we directly rule out a role for BRCA1's resection partner, CtIP, in promoting break
formation at stalled forks.
572
The 53BP1-Rif1 axis has been extensively studied in regulating two-ended replication-
independent breaks, with several labs demonstrating resection and repair pathway
choice control [72]. Because our system involves the generation of breaks at stalled
forks via SLX4 activity, we have taken advantage of this as an opportunity to investigate
in detail how fork breakage and resection at stalled forks is regulated by 53BP1-RIF1
axis. 53BP1 has been previously shown to protect stalled forks [61]; however, our SCAI-
null context allows us to show here that 53BP1 does not influence DNA resection upon
break formation at stalled forks. This was true whether forks were stalled by HU or at
breaks resulting from cisplatin treatment. Rif1 has been shown to possess 53BP1
independent roles at forks [60]. It can be recruited to stalled forks in a 53BP1
independent manner, and here we show that RIF1 loss by itself led to increased fork
breakage signaling. Whether this is related to Rif1's fork protection roles is not clear. We
also observed what appeared to be an antagonism between BRCA1 and RIF1 in
regulating resection upon ICL agent treatment. Removing BRCA1 enforced a RIF1-
mediated block to resection. While 53BP1 did not affect stalled fork outcomes upon fork
breakage, it seemed to modulate the extent of RAD51 amounts on chromatin. This is
consistent with what was recently shown using CRISPR induced breakage [69].

We previously employed a Tus/Ter reporter to show that SCAI is particularly important
for HR promotion at stalled forks [22]. Our results here suggest that this role is important
for driving replication restart, and that this role occurs downstream of RAD51 activity.
We hypothesize that certain regions of the genome are prone to stall the advancing
replication fork and require SCAI and REV3 to direct fork restoration and repair. Indeed,
using a fragile site reporter we found that SCAI/REV3 loss results in increased breaks
[22], consistent with our current findings. While SCAI and REV3 appear to function in
concert, they do not have overlapping roles, and loss of SCAI seemed to lead to a
stronger restart defect than loss of REV3. Future work will uncover under what
scenarios SCAI and REV3 function together and how REV3 is regulated differentially
from its role in translesion synthesis.

A central role for RAD51 in promoting fork restart has been proposed by others [20],
and the fork restart defects seen in the absence of SCAI appeared to occur downstream
of RAD51 activity. BRCA1 depletion did not affect SCAI recruitment to chromatin, but
results in significant reduction in RAD51 loading on chromatin. Our data suggests that
the SCAI/REV3 mediated restart mechanism occurs in parallel with SMARCAL1
/ZRANB3 or FBH1 mediated fork reversal. Importantly, loss of fork reversal factors like
SMARCAL1 paradoxically led to increased fork breakage in the absence of SCAI,
suggesting that forks normally processed in a reversal mediated fashion are funneled
into a SCAI dependent pathway that ultimately leads to fork breakage whenever SCAI
was absent. Both pathways appear to require RAD51 activity, as inhibiting RAD51
prevented fork breakage whenever SCAI and SMARCAL1 were co-depleted. RAD51
has been proposed to cause a CMG helicase-containing bubble ahead of annealed
parental strands [73]. RAD51 could promote restart by invading newly synthesized DNA
ahead of the annealed fork. Under-replicated regions can then be duplicated by gap-
synthesis post repair. BRCA1 is not essential for fork reversal [63], but may enhance
RAD51 chromatin association to drive strand invasion mediated fork restart. We show
that SCAI is required for fork restart in this context, and in the absence of SCAI, SLX4
accumulates on chromatin to cleave the resulting HR intermediates leading to fork
breakage. Such broken forks can be engaged by alternative means of end joining or
may persist leading to genomic instability and cell death. Whether the same RAD51
activity drives both reversal as well as HR-mediated fork restart will be a focus of future
work.

METHODS

Cell lines

U2OS cells were originally obtained from ATCC (HTB-96). U2OS ΔSCAI and U2OS cells expressing lentiCRISPR control gRNAs (WT) were previously described [22], and were passaged in McCoy's 5A medium + L-Glutamine (16600-082, Gibco). hTERT-RPE1 cells were originally obtained from ATCC (CRL-4000). All RPE1 cell lines and their derivatives were passaged in DMEM/F12 (1:1) medium + L-Glutamine + 15mM HEPES (11330-032, Gibco). RPE1 ΔBRCA1/Δp53 (referred to as ΔBRCA1 in Figure 3) was a kind gift from Dr. Alan D'Andrea [74]. RPE1 $TP53^{-/-}$ GFP PINDUCER20 (referred to as "parent" in Figures S4C and S6A), RPE1 $TP53^{-/-}$ $BRCA1^{-/-}$ GFP PINDUCER20 (referred to as "ΔBRCA1" in Figure 4A, 4B, S4C, and S6A), RPE1 $Tp53^{-/-}$ $BRCA1^{-/-}$ GFP-BRCA1 PINDUCER20 (referred to as "WT" in Figures 4A, 4B, 4C and "GFP-BRCA1 Rescue" in Figure S4C) and RPE1 $TP53^{-/-}$ $BRCA1^{-/-}$ GFP-BRCA1 ΔCC PINDUCER20 (referred to as ΔCC) were kind gifts from Dr. Andrew Blackford and were previously described [48]. RPE1 Flag Cas9 $TP53^{-/-}$ ΔBRCA1 Δ53BP1 (referred to as ΔBRCA1/Δ53BP1 in Figure S4D) were a kind gift from Dr. Dheva Setiaputra and were previously described [59].

Reagents and Drugs

Dimethyl Sulfoxide (DMSO) was from Thomas Scientific (D0231). Hydroxyurea (HU) was obtained from Sigma-Aldrich (H8627-5G). Aphidicolin (APH) was from Sigma-Aldrich (A4487-1ML). The TLS inhibitor JH-RE-06 was obtained from MedChemExpress (HY-126214, 10mM/1mLSolution). Cisplatin was from Selleck Chemicals (S1166-50mg). Rad51 inhibitor B02 was from Sigma-Aldrich (553525-25MG). Olaparib from Selleck Chemicals (S1060-25mg). Colcemid was obtained from Millipore Sigma (10295892001).

Antibodies

The antibodies used in this work include the following: Anti-vinculin 1:1000 (Sigma-Aldrich, V9131-.2ML), vinculin (7F9) 1:200 (Santa Cruz Biotechnology, sc-73614), SCAI 1:500 (Abcam, ab124688), Chk1 (2G1D5) 1:1000 (Cell Signaling, 2360S), P-Chk1 (S345) (133D3) Rabbit mAb 1:1000 (Cell Signaling, 2348S), RPA 32 kDa subunit (9H8) 1:200 (Santa Cruz Biotechnology, sc-56770, only used for western blots), RPA32 1:200 (Genetex, GTX70258, only used for IF), Rabbit x-Phospho RPA32 (S4/S8) 1:1000 (Bethyl, A300-245A), H2AX 1:1000 (Bethyl, A300-082A), Anti-phospho-Histone H2A.X (Ser139) 1:1000 (EMD Millipore Corp., 05-636), REV7 1:200 (Santa Cruz Biotechnology, sc-135977), Anti-Exonuclease 1 (EXO1) 1:1000 (Bethyl, A302-640A), ORC2 1:1000 (Abcam, ab68348), BRCA1 1:1000 (EMD Millipore Corp., 07-434), 53BP1 1:1000 (Bethyl, A300-272A), RIF1 1:1000 (Bethyl, A300-569A), CtIP 1:1000 (Bethyl, A300-488A), SmarcAL1 (A-2) 1:200 (Santa Cruz Biotechnology, sc-376377), ZRANB3 1:1000 (Bethyl, A303-033A), FBH1 1:1000 (Abcam, ab58881), RAD51 1:1000 (Abcam, ab63801), MUS81 1:200 (Santa Cruz Biotechnology, sc-53382), SLX4 1:1000 (Bethyl, A302-270A), H3 1:1000 (Cell Signaling, 9715), MRE11 1:1000 (Abcam, ab214), Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP 1:2500 (Invitrogen, 31430), Goat anti-Rabbit IgG (H+L) Secondary Antibody, HRP 1:2500 (Invitrogen, 31460), Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488 1:300 (Invitrogen, A32731), Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed

672 Secondary Antibody, Alexa Fluor Plus 594 1:300 (Invitrogen, A32742), Goat anti-Mouse
673 IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488 1:300
674 (Invitrogen, A32723).
675
676 siRNA transfections
677 RPE1 and U2OS cells were seeded onto 6-well plates and reverse transfected with 20-
678 40nM siRNA using Lipofectamine RNAiMAX reagent (13-778-150, Invitrogen) according
679 to the manufacturer's instructions. Cells were treated 48 h after treatment with drugs as
680 indicated in the legend and processed 16 to 18 h later.
681
SCAI rescue experiment
U2OS cells were transfected with plasmids using PolyJet reagent (SignaGen,
SL10068B) according to the manufacturer's instructions and treated with drugs as
indicated in the Figure legends. SCAI was cloned into the pcDNA3.1 backbone as
previously described [22]. The ORF sequence of SCAI was modified to avoid being
identified by gRNA in ∆SCAI cells. Specifically, GCTGAAGATGAT (Ala-Glu-Asp-Asp)
was replaced with GCAGAGGACGAC using a site-directed mutagenesis kit (New
England Biolabs, E0552S).
690
Western blotting
Cells were harvested, washed with ice cold phosphate-buffered saline (PBS), and lysed
in SDS lysis buffer (50 mM Tris HCl pH 6.8, 100 mM NaCl, 1.7% SDS, 10 mM NaF, 7%
glycerol) prior to sonication. Protein content was measured using Bradford reagent
(23236, Life Technologies) on an Eppendorf Biophotometer. Laemmli sample buffer
(1610747, Bio-Rad Laboratories) was added with 20 mM DTT prior to boiling the
samples for 5-10 min. Whole cell lysates were analyzed using sodium dodecyl sulfate
gel electrophoresis (SDS-PAGE) and transferred onto nitrocellulose membranes.
Membranes were blocked with 5% (wt/vol) milk in Tris-buffered saline with Tween
(TBST) for a minimum of 30 min and then probed with the indicated antibodies. Protein
bands were quantified with ImageJ.
702
Chromatin fractionation
Chromatin fractions were done as previously described [75]. Briefly, RPE1 cells
previously transfected with the indicated siRNAs were seeded into 10 cm plates, treated
with drugs as indicated, then harvested and washed with ice cold PBS. Sedimented
cells were resuspended in cold Solution 1 (10 mM Hepes (pH 7.9), 10 mM KCl, 1.5 mM
$MgCl_2$, 0.34 M sucrose, 1 mM DTT, 10% glycerol, 1mM Na_3VO_4, protease and
phosphatase inhibitors). Triton X-100 was added to 0.1%, and samples were incubated
on ice prior to centrifuging. The supernatant was removed as the cytoplasmic extract,
and sedimented nuclei were washed with Solution 1 prior to being lysed in Solution 2
(3 mM EDTA, 0.2 mM EGTA, 1 mM DTT, protease and phosphatase inhibitors) on ice.
Samples were centrifuged, washed with cold Solution 2, and then the chromatin-
enriched pellets were lysed with SDS lysis buffer. Chromatin extract samples were
sonicated, and protein content was measured. Sample buffer was added to all samples
and boiled prior to western blotting.
717

Cell cycle analyses
Cells were treated with indicated drugs for 18h prior to being harvested. Cells were then
washed with ice cold PBS, then fixed in 70% ethanol at 4 °C overnight. Samples were
pelleted, washed with ice cold PBS, and resuspended in 50 µg/mL propidium iodide
solution containing 0.1 mg/mL RNase A. Samples were then incubated at 37°C in the
dark for 15-30 min and flow cytometry was performed using a FACSymphony A5 (BD
Biosciences). Data were analyzed using FlowJo software.
725
Clonogenic survival assays
Cells were transfected as stated above for 48 h with the indicated siRNAs and then
treated with indicated doses of drugs for 18 h. The number of cells plated in six-well
plates was adjusted based on the dose of drug the cells were exposed to. Cells were
allowed to form colonies for 10-14 days prior to being fixed and stained using Crystal
Violet. Analyses were performed using ImageJ.
732
Immunofluorescence Assay
Cells were transfected with the indicated siRNAs as above and plated onto glass
coverslips 18 h later. 48 h after the transfection, the cells were treated with HU for the
indicated dose and duration. The cells were then washed with ice-cold PBS on ice, pre-
extracted for 5 min on ice using cytoskeleton buffer (100 mM NaCl, 10 mM HEPES (pH
7.8), 3 mM $MgCl_2$, 300 mM sucrose, 0.1% Triton X-100), fixed with 4%
paraformaldehyde in PBS for 10 min at room temp and then permeabilized with 0.5%
Triton-100 in 1X PBS. Samples were then blocked with 3% bovine serum albumin (BSA)
in PBS for a minimum of 30 min on a rocker followed by overnight incubation with
primary antibodies at 4 °C in a humidified chamber. Unbound primary antibodies were
washed using 3% BSA in PBS and then samples incubated with secondary antibody for
1 h at RT in the dark. Unbound antibody was washed with PBS and then samples
stained with 1 µg/ml DAPI for 1 h before mounting the coverslips on a microscope slide
for imaging. Images were acquired using an LSM 780 NLO (Zeiss), and a minimum of
100 nuclei for each group were analyzed using ImageJ.
748
QIBC image acquisition and analysis
Fixed and stained slides were imaged on a Molecular Devices ImageXpress Microscope
equipped with a Nikon 40x/0.95 PlanApo objective. Around 2000 nuclei were acquired
for each condition. Quantification was then performed using the "Multi Wavelength Cell
Scoring" module in the MetaXpress software. Briefly, nuclei were segmented using the
DAPI channel, and the raw Mean Fluorescent Intensities (MFIs) of each nucleus were
measured for DAPI, RPA (AF488), and gamma-H2AX (AF594). All cells were then
plotted for each condition and p-values generated by Ordinary one-way ANOVA,
Tukey's multiple comparison test
758
759 DNA fiber assays
760 hTERT-RPE1 cells were reverse transfected with the indicated siRNAs at 20 nM using
761 Lipofectamine RNAiMAX reagent (Invitrogen). Transfected hTERT-RPE1 cells were
762 pulse-labeled with CldU followed by IdU as indicated in each Figure. After being labeled,
763 cells were harvested and embedded in 0.6% low melting point agarose gel plugs. Cells

in plugs were incubated with lysis solution (0.2 M EDTA, 0.5% SDS, 3 mg/mL Proteinase K) overnight at 50°C. Each plug was dissolved in 0.5 M MES solution (pH 5.5) by β-agarase (NEB) overnight treatment and DNA was attached to a pre-treated coverslip (PolyAn) using FiberComb® Molecular Combing System (GENOMIC VISION). DNA on coverslips was denatured by incubating in denaturing solution (0.5 N NaOH, 1 M NaCl) for 8 min. Blocking was performed with 5% BSA in PBS for 60 min. For immunostaining, coverslips were incubated overnight with primary antibodies; ab6326 anti-BrdU (cross-reacts with CldU) antibody (rat) (1:50) and BD Biosciences 347580 anti-BrdU (cross-reacts with IdU) antibody (mouse) (1:8). Slides were washed with PBS followed by incubation for one hour with the secondary antibodies; anti-rat AIexa-488 antibody (1:300) and anti-mouse Alexa-594 antibody (1:300). After wash with PBS, coverslips were mounted onto glass slides with anti-fade mounting medium (SlowFade™ Glass Soft-Set Antifade Mountant, Invitrogen). Images were acquired with Leica STELLARIS confocal microscope. DNA fiber numbers and lengths were manually analyzed using LAS X software (Leica Microsystems).

Metaphase spreads
hTERT-RPE1 cells were reverse transfected with the indicated siRNAs as above using Lipofectamine RNAiMAX. Two days after the transfection, cells were treated with 2 mM HU or PBS for 18 h, followed by PBS wash and 24 h incubation in normal medium. Cells were exposed to colcemid (100 ng/ml) for the last 3 h of the 24 h normal medium incubation before harvest. Collected cells were treated with a hypotonic solution (1:2 mixture of 0.075 M KCl and 0.9% Na citrate) for 10 min and fixed with 3:1 methanol: acetic acid. Fixed cells were transferred on glass slides dropwise, and the slides were stained with Giemsa stain. Cells in metaphase were observed using a Zeiss Axiovert 200M microscope and captured using AxioVision. Over 50 metaphase cells were analyzed per sample.

Neutral Comet assay
hTERT-RPE1 cells were reverse transfected with the indicated siRNAs at 20 nM using Lipofectamine RNAiMAX reagent (Invitrogen). Two days after the transfection, cells were treated with 2 mM HU or PBS for 40 h. After HU treatment, cells were harvested and resuspended in ice-cold PBS. 10μL of the cell suspension was mixed with 100 μL of melted 0.7% low melting point agarose gel, and the mixture was spread onto a pre-coated CometSlide™ (R&D SYSTEMS®). After the agarose gel was solidified at 4 °C, the slide was immersed in lysis solution (10 mM Tris base, 2.5 M NaCl, 100 mM Na$_2$-EDTA, 10% DMSO, 1% Triton X-100, pH 10) for 2 h at 4 °C followed by a 30 min incubation in neutral electrophoresis buffer (100 mM Tris base, 300 mM sodium acetate, pH 9). Then, the slide was placed in an electric field for 40 min. After the electrophoresis, the slide was immersed in DNA precipitation solution (82% ethanol, 1 mM ammonium acetate) for 30 min followed by a 30 min incubation in 70% ethanol. SYBR Gold dye (Invitrogen) was used to stain DNA ("comet") and images were obtained with TissueFAXS (TissueGnostics). Comets were analyzed with ImageJ (OpenComet).

Quantification and statistical analyses

Statistical analyses were performed using Prism 10 (GraphPad). All statistics used are
indicated in the Figure legends. For all experiments: n.s. $P \geq 0.05$; $*P<0.05$; $**P < 0.01$;
$***P < 0.001$; $****P<0.0001$. All experiments in this work were performed multiple
independent times with similar results. Multiple siRNAs, multiple knockout clones, and
multiple cell lines were analyzed to confirm that results were not caused by off-target
effects or clonal variations. Representative images and/or experiments are shown for
westerns, cell cycle analyses, DNA fiber assays and IFs.
817
818

REFERENCES AND NOTES

1. Zeman, M.K. and K.A. Cimprich, *Causes and consequences of replication stress.* Nat Cell Biol, 2014. **16**(1): p. 2-9.
2. Ciccia, A. and S.J. Elledge, *The DNA damage response: making it safe to play with knives.* Mol Cell, 2010. **40**(2): p. 179-204.
3. Saxena, S. and L. Zou, *Hallmarks of DNA replication stress.* Mol Cell, 2022. **82**(12): p. 2298-2314.
4. Gaillard, H., T. Garcia-Muse, and A. Aguilera, *Replication stress and cancer.* Nat Rev Cancer, 2015. **15**(5): p. 276-89.
5. Cimprich, K.A. and D. Cortez, *ATR: an essential regulator of genome integrity.* Nat Rev Mol Cell Biol, 2008. **9**(8): p. 616-27.
6. Neelsen, K.J. and M. Lopes, *Replication fork reversal in eukaryotes: from dead end to dynamic response.* Nat Rev Mol Cell Biol, 2015. **16**(4): p. 207-20.
7. Quinet, A., D. Lemacon, and A. Vindigni, *Replication Fork Reversal: Players and Guardians.* Mol Cell, 2017. **68**(5): p. 830-833.
8. Schlacher, K., et al., *Double-strand break repair-independent role for BRCA2 in blocking stalled replication fork degradation by MRE11.* Cell, 2011. **145**(4): p. 529-42.
9. Errico, A. and V. Costanzo, *Mechanisms of replication fork protection: a safeguard for genome stability.* Crit Rev Biochem Mol Biol, 2012. **47**(3): p. 222-35.
10. Petermann, E. and T. Helleday, *Pathways of mammalian replication fork restart.* Nat Rev Mol Cell Biol, 2010. **11**(10): p. 683-7.
11. Scully, R., et al., *Recombination and restart at blocked replication forks.* Curr Opin Genet Dev, 2021. **71**: p. 154-162.
12. Zellweger, R., et al., *Rad51-mediated replication fork reversal is a global response to genotoxic treatments in human cells.* J Cell Biol, 2015. **208**(5): p. 563-79.
13. Petermann, E., et al., *Hydroxyurea-stalled replication forks become progressively inactivated and require two different RAD51-mediated pathways for restart and repair.* Mol Cell, 2010. **37**(4): p. 492-502.
14. Mouron, S., et al., *Repriming of DNA synthesis at stalled replication forks by human PrimPol.* Nat Struct Mol Biol, 2013. **20**(12): p. 1383-9.
15. Berti, M., et al., *Human RECQ1 promotes restart of replication forks reversed by DNA topoisomerase I inhibition.* Nat Struct Mol Biol, 2013. **20**(3): p. 347-54.
16. Pepe, A. and S.C. West, *MUS81-EME2 promotes replication fork restart.* Cell Rep, 2014. **7**(4): p. 1048-55.
17. Sarbajna, S., D. Davies, and S.C. West, *Roles of SLX1-SLX4, MUS81-EME1, and GEN1 in avoiding genome instability and mitotic catastrophe.* Genes Dev, 2014. **28**(10): p. 1124-36.
18. Kockler, Z.W., et al., *Repair of DNA Breaks by Break-Induced Replication.* Annu Rev Biochem, 2021. **90**: p. 165-191.
19. Kramara, J., B. Osia, and A. Malkova, *Break-Induced Replication: The Where, The Why, and The How.* Trends Genet, 2018. **34**(7): p. 518-531.

20. Ait Saada, A., S.A.E. Lambert, and A.M. Carr, *Preserving replication fork integrity*
and competence via the homologous recombination pathway. DNA Repair
(Amst), 2018. **71**: p. 135-147.
21. Chakraborty, S., K. Schirmeisen, and S.A. Lambert, *The multifaceted functions of*
homologous recombination in dealing with replication-associated DNA damages.
DNA Repair (Amst), 2023. **129**: p. 103548.
22. Adeyemi, R.O., et al., *The Protexin complex counters resection on stalled forks*
to promote homologous recombination and crosslink repair. Mol Cell, 2021.
81(21): p. 4440-4456 e7.
23. Lemay, J.F., et al., *A genome-wide screen identifies SCAI as a modulator of the*
UV-induced replicative stress response. PLoS Biol, 2022. **20**(10): p. e3001543.
24. Xu, Y., et al., *53BP1 and BRCA1 control pathway choice for stalled replication*
restart. Elife, 2017. **6**.
25. Bhat, K.P. and D. Cortez, *RPA and RAD51: fork reversal, fork protection, and*
genome stability. Nat Struct Mol Biol, 2018. **25**(6): p. 446-453.
26. Kolinjivadi, A.M., et al., *Smarcal1-Mediated Fork Reversal Triggers Mre11-*
Dependent Degradation of Nascent DNA in the Absence of Brca2 and Stable
Rad51 Nucleofilaments. Mol Cell, 2017. **67**(5): p. 867-881 e7.
27. Vujanovic, M., et al., *Replication Fork Slowing and Reversal upon DNA Damage*
Require PCNA Polyubiquitination and ZRANB3 DNA Translocase Activity. Mol
Cell, 2017. **67**(5): p. 882-890 e5.
28. Berti, M., D. Cortez, and M. Lopes, *The plasticity of DNA replication forks in*
response to clinically relevant genotoxic stress. Nat Rev Mol Cell Biol, 2020.
21(10): p. 633-651.
29. Cortez, D., *Preventing replication fork collapse to maintain genome integrity.* DNA
Repair (Amst), 2015. **32**: p. 149-157.
30. Dungrawala, H., et al., *The Replication Checkpoint Prevents Two Types of Fork*
Collapse without Regulating Replisome Stability. Mol Cell, 2015. **59**(6): p. 998-
1010.
31. Sirbu, B.M., et al., *Analysis of protein dynamics at active, stalled, and collapsed*
replication forks. Genes Dev, 2011. **25**(12): p. 1320-7.
32. Toledo, L.I., et al., *ATR prohibits replication catastrophe by preventing global*
exhaustion of RPA. Cell, 2013. **155**(5): p. 1088-103.
33. Ikegami, S., et al., *Aphidicolin prevents mitotic cell division by interfering with the*
activity of DNA polymerase-alpha. Nature, 1978. **275**(5679): p. 458-60.
34. Macheret, M. and T.D. Halazonetis, *DNA replication stress as a hallmark of*
cancer. Annu Rev Pathol, 2015. **10**: p. 425-48.
35. Zhao, L. and M.T. Washington, *Translesion Synthesis: Insights into the Selection*
and Switching of DNA Polymerases. Genes (Basel), 2017. **8**(1).
36. Wojtaszek, J.L., et al., *A Small Molecule Targeting Mutagenic Translesion*
Synthesis Improves Chemotherapy. Cell, 2019. **178**(1): p. 152-159 e11.
37. Byun, T.S., et al., *Functional uncoupling of MCM helicase and DNA polymerase*
activities activates the ATR-dependent checkpoint. Genes Dev, 2005. **19**(9): p.
1040-52.
38. Densham, R.M. and J.R. Morris, *Moving Mountains-The BRCA1 Promotion of*
DNA Resection. Front Mol Biosci, 2019. **6**: p. 79.

39.   Tarsounas, M. and P. Sung, *The antitumorigenic roles of BRCA1-BARD1 in DNA*
911   *repair and replication.* Nat Rev Mol Cell Biol, 2020. **21**(5): p. 284-299.
912   40.   Baer, R. and T. Ludwig, *The BRCA1/BARD1 heterodimer, a tumor suppressor*
913   *complex with ubiquitin E3 ligase activity.* Curr Opin Genet Dev, 2002. **12**(1): p.
914   86-91.
915   41.   Hansen, R.K., et al., *SCAI promotes DNA double-strand break repair in distinct*
916   *chromosomal contexts.* Nat Cell Biol, 2016. **18**(12): p. 1357-1366.
917   42.   Rickman, K.A., et al., *Distinct roles of BRCA2 in replication fork protection in*
918   *response to hydroxyurea and DNA interstrand cross-links.* Genes Dev, 2020.
919   **34**(11-12): p. 832-846.
920   43.   Savage, K.I. and D.P. Harkin, *BRCA1, a 'complex' protein involved in the*
921   *maintenance of genomic stability.* FEBS J, 2015. **282**(4): p. 630-46.
922   44.   Wang, B., et al., *Abraxas and RAP80 form a BRCA1 protein complex required for*
923   *the DNA damage response.* Science, 2007. **316**(5828): p. 1194-8.
924   45.   Cruz-Garcia, A., A. Lopez-Saavedra, and P. Huertas, *BRCA1 accelerates CtIP-*
925   *mediated DNA-end resection.* Cell Rep, 2014. **9**(2): p. 451-9.
926   46.   Yun, M.H. and K. Hiom, *CtIP-BRCA1 modulates the choice of DNA double-*
927   *strand-break repair pathway throughout the cell cycle.* Nature, 2009. **459**(7245):
928   p. 460-3.
929   47.   Wu, X. and B. Wang, *Abraxas suppresses DNA end resection and limits break-*
930   *induced replication by controlling SLX4/MUS81 chromatin loading in response to*
931   *TOP1 inhibitor-induced DNA damage.* Nat Commun, 2021. **12**(1): p. 4373.
932   48.   Tsukada, K., et al., *BLM and BRCA1-BARD1 coordinate complementary*
933   *mechanisms of joint DNA molecule resolution.* Mol Cell, 2024. **84**(4): p. 640-658
934   e10.
935   49.   Ceppi, I., et al., *Mechanism of BRCA1-BARD1 function in DNA end resection and*
936   *DNA protection.* Nature, 2024. **634**(8033): p. 492-500.
937   50.   Bunting, S.F., et al., *53BP1 inhibits homologous recombination in Brca1-deficient*
938   *cells by blocking resection of DNA breaks.* Cell, 2010. **141**(2): p. 243-54.
939   51.   Hustedt, N. and D. Durocher, *The control of DNA repair by the cell cycle.* Nat Cell
940   Biol, 2016. **19**(1): p. 1-9.
941   52.   Nakamura, K., et al., *H4K20me0 recognition by BRCA1-BARD1 directs*
942   *homologous recombination to sister chromatids.* Nat Cell Biol, 2019. **21**(3): p.
943   311-318.
944   53.   Becker, J.R., et al., *BARD1 reads H2A lysine 15 ubiquitination to direct*
945   *homologous recombination.* Nature, 2021. **596**(7872): p. 433-437.
946   54.   Escribano-Diaz, C., et al., *A cell cycle-dependent regulatory circuit composed of*
947   *53BP1-RIF1 and BRCA1-CtIP controls DNA repair pathway choice.* Mol Cell,
948   2013. **49**(5): p. 872-83.
949   55.   Jaspers, J.E., et al., *Loss of 53BP1 causes PARP inhibitor resistance in Brca1-*
950   *mutated mouse mammary tumors.* Cancer Discov, 2013. **3**(1): p. 68-81.
951   56.   Isobe, S.Y., et al., *Inhibition of RIF1 by SCAI Allows BRCA1-Mediated Repair.*
952   Cell Rep, 2017. **20**(2): p. 297-307.
953   57.   Scully, R., J.C. Walter, and A. Nussenzweig, *One-ended and two-ended breaks*
954   *at nickase-broken replication forks.* DNA Repair (Amst), 2024. **144**: p. 103783.

58. Dev, H., et al., *Shieldin complex promotes DNA end-joining and counters*
homologous recombination in BRCA1-null cells. Nat Cell Biol, 2018. **20**(8): p.
954-965.
59. Noordermeer, S.M., et al., *The shieldin complex mediates 53BP1-dependent*
DNA repair. Nature, 2018. **560**(7716): p. 117-121.
60. Mukherjee, C., et al., *RIF1 promotes replication fork protection and efficient*
restart to maintain genome stability. Nat Commun, 2019. **10**(1): p. 3287.
61. Liu, W., et al., *Two replication fork remodeling pathways generate nuclease*
substrates for distinct fork protection factors. Sci Adv, 2020. **6**(46).
62. Liang, L., et al., *Molecular basis for assembly of the shieldin complex and its*
implications for NHEJ. Nat Commun, 2020. **11**(1): p. 1972.
63. Taglialatela, A., et al., *Restoration of Replication Fork Stability in BRCA1- and*
BRCA2-Deficient Cells by Inactivation of SNF2-Family Fork Remodelers. Mol
Cell, 2017. **68**(2): p. 414-430 e8.
64. Mason, J.M., et al., *Non-enzymatic roles of human RAD51 at stalled replication*
forks. Nat Commun, 2019. **10**(1): p. 4410.
65. Halder, S., et al., *Double-stranded DNA binding function of RAD51 in DNA*
protection and its regulation by BRCA2. Mol Cell, 2022. **82**(19): p. 3553-3565 e5.
66. Huang, F., et al., *Identification of specific inhibitors of human RAD51*
recombinase using high-throughput screening. ACS Chem Biol, 2011. **6**(6): p.
628-35.
67. Lai, X., et al., *MUS81 nuclease activity is essential for replication stress tolerance*
and chromosome segregation in BRCA2-deficient cells. Nat Commun, 2017. **8**: p.
15983.
68. Mayle, R., et al., *DNA REPAIR. Mus81 and converging forks limit the*
mutagenicity of replication fork breakage. Science, 2015. **349**(6249): p. 742-7.
69. Pavani, R., et al., *Structure and repair of replication-coupled DNA breaks.*
Science, 2024. **385**(6710): p. eado3867.
70. Rickman, K. and A. Smogorzewska, *Advances in understanding DNA processing*
and protection at stalled replication forks. J Cell Biol, 2019. **218**(4): p. 1096-1107.
71. Daza-Martin, M., et al., *Isomerization of BRCA1-BARD1 promotes replication fork*
protection. Nature, 2019. **571**(7766): p. 521-527.
72. Scully, R., et al., *DNA double-strand break repair-pathway choice in somatic*
mammalian cells. Nat Rev Mol Cell Biol, 2019. **20**(11): p. 698-714.
73. Liu, W., et al., *RAD51 bypasses the CMG helicase to promote replication fork*
reversal. Science, 2023. **380**(6643): p. 382-387.
74. Lim, K.S., et al., *USP1 Is Required for Replication Fork Protection in BRCA1-*
Deficient Tumors. Mol Cell, 2018. **72**(6): p. 925-941 e4.
75. Tischler, J.D., et al., *FLIP(C1orf112)-FIGURENL1 complex regulates RAD51*
chromatin association to promote viability after replication stress. Nat Commun,
2024. **15**(1): p. 866.
996
997

Acknowledgements
We thank all the members of the Adeyemi laboratory for advice and helpful discussions.
We thank Dr. Blackford and Dr. Setiaputra for providing reagents.

Funding
This work was supported by NIH NIGMS R35 GM150532-01 and an Early Career
Investigator Grant from the Ovarian Cancer Research Alliance (ECIG-2023-3-1004) (to
ROA). R.B. was supported by a Washington Research Foundation fellowship. This
research was also supported by the Cellular Imaging Shared Resource
RRID:SCR_022609 and the Flow Cytometry Shared Resource, RRID:SCR_022613, of
the Fred Hutch / University of Washington / Seattle Children's Cancer Consortium (P30
CA015704).

Author Contributions
Conceptualization: R.O.A., Methodology: C.U., H.T., R.B., X.J., R.O.A., Investigation:
C.U., H.T., R.B., X.J., R.O.A., Visualization: C.U., H.T., R.B., X.J., S.K., R.O.A.,
Supervision: R.O.A., Writing—original draft: R.O.A., Writing—review & editing: C.U.,
H.T., R.B., X.J., S.K., R.O.A.

Competing Interests
The authors declare that they have no competing interests.

Data and Materials Availability
All data needed to evaluate the conclusions in the paper are present in the paper and/or
the Supplementary Materials. Materials resulting from the findings of this study are
available from the corresponding author upon request.

MAIN FIGURE LEGENDS

Figure 1. **SCAI promotes integrity and efficient restart of stalled replication forks**.
A. Schematic of pulse-labeling DNA strands, representative images and lengths of two-color DNA fibers. DNA fibers were from hTERT-RPE1 cells transfected with siRNAs as indicated. **B**. Schematic of pulse-labeling DNA strands, representative images and lengths of single- and two-color DNA fibers. The top two panels indicate duel-labelled conditions, while the lower panels indicate single labelled conditions. DNA fibers were from hTERT-RPE1 cells transfected with siRNAs as indicated. **C**. Schematic of pulse-labeling DNA strands, representative images and lengths of two-color DNA fibers. DNA fibers were from hTERT-RPE1 cells transfected with siRNAs as indicated. **A**–**C**, red line represents median, three independent experiments, representative experiment shown. Mann Whitney test for two group comparison and Kruskal-Wallis followed by Dunn's multiple comparison test for more than three group comparison. ns – not significant (P ≥ 0.05).

Figure 2. SCAI/REV3 limits breakage of stalled replication forks.
A. WT RPE1 cells transfected with the indicated siRNAs for 60 h, then treated with 2 mM hydroxyurea (HU) for 18 h, harvested, and processed for western blotting with the specified antibodies. ND, no drug. **B.** Western blots (WB) from WT and SCAI-null U2OS cell lines treated with 1 mM HU for 18 h. **C.** WT and ∆SCAI U2OS cells were transfected with either backbone plasmids (Vector) or plasmids expressing SCAI that is resistant to CRISPR/Cas9 editing. After 30 h, cells were treated with 1.5 mM HU for another 18 h before immunoblotting. **D.** WT RPE1 cells were transfected with the indicated siRNAs, then treated with vehicle (DMSO), 1 µM, or 2 µM aphidicolin (APH) for 18 h before being processed for western blotting with the specified antibodies. **E-F**. Comet assay of hTERT-RPE1 (**E**) and U2OS (**F**) cells treated with 2 mM HU for 40 h. Indicated siRNAs were transfected two days before HU treatment. Red line represents median, three independent experiments, representative experiment shown. Kruskal-Wallis followed by Dunn's multiple comparison test. **G.** WT RPE1 cells were transfected with the indicated siRNAs, then treated with 1 mM HU for 18 h and assayed for the indicated proteins by WB. γH2AX values are normalized to vinculin loading control. **H-J**. Representative microscopy images showing chromatin bound RPA and γH2AX, following transient knockdown of SCAI or REV3 and 16 h treatment with 1.5 mM HU in hTERT RPE1 wildtype cells (e) scale bar indicates 25 micron. Quantification of the integrated fluorescent intensity of RPA (f) and γH2AX (g) of cells in e, red bar indicates mean and SEM of about 2000 nuclei. ****p< 0.0001, Tukey's test.

Figure 3. BRCA1 promotes increased fork breakage and resection in the absence of SCAI/REV3.
A. WT and BRCA1 null RPE1 cells were transfected with the indicated siRNAs, then treated with 1 mM HU for 18 h prior to blotting for the indicated proteins. **B.** WT and BRCA1-null RPE1 cells were transfected with the indicated siRNAs. Cells were then treated with 1 mM HU for 18 h before WCLs (whole cell lysates) were prepared and WB were performed against the indicated proteins. **C.** IF images showing RPA foci in WT RPE1 cells following treatment with the indicated siRNAs. Cells were treated with 1.5

mM HU for 18 h prior to IF. Scale bars indicate 5 microns. **D.** Quantification of (**C**). ****P < 0.0001, unpaired Welch's t test (two-tailed). **E.** IF analyses showing representative γH2AX foci in WT and BRCA1-null RPE1 cells post-treatment with indicated siRNAs. Cells were treated with 1.5 mM HU for 18 h prior to IF analyses. Scale bars indicate 5 microns. **F**. Quantification of (**E**). *P = 0.0337, unpaired Welch's t test (two-tailed). **G**. Comet assay of hTERT-RPE1 cells treated with 2 mM HU for 40 h. Indicated siRNAs were transfected two days before HU treatment. Red line represents median, two independent experiments, representative experiment shown. Kruskal-Wallis followed by Dunn's multiple comparison test. ns – not significant (P ≥ 0.05). **H.** WT and BRCA1-null RPE1 cells were transfected with the indicated siRNAs, then treated with 1 mM HU for 18 h and fractioned prior to blotting for the indicated proteins. CE – cytoplasmic extract. SCAI values are normalized to ORC2 loading control.

Figure 4. BRCA1 pro-breakage / resection roles at stalled and broken forks occur independently of 53BP1 but is modulated by Rif1.
A. GFP-WT-BRCA1 rescue (labeled WT), BRCA1-null (Δ), and GFP-BRCA1-S1655 (S1655) RPE1 cells were transfected with the indicated siRNAs, then treated with 2 mM HU for 18 h. WCLs were assayed for the indicated proteins by WB. **B.** GFP-WT-BRCA1 rescue (WT), BRCA1-null (Δ), and GFP-BRCA1-ΔCC (ΔCC) RPE1 cells were transfected with the indicated siRNAs, then treated with 2 mM HU for 18 h and assayed for the indicated proteins by WB. **C.** Western blots from GFP-WT-BRCA1 rescue (WT), and GFP-BRCA1-ΔCC (ΔCC) RPE1 cell lines transfected with the indicated siRNAs and treated with ND, 0.75 mM (Lanes 4-5), 1.5 mM (Lanes 6-7), or 3 mM (Lanes 8-9) HU for 18 h. **D.** Western blots from WT and SCAI-null U2OS cells transfected with the indicated siRNAs and then treated with 2 µM cisplatin for 18 h. **E.** WT U2OS cells were transfected with the indicated siRNAs, then treated with 1 mM HU for 18 h and assayed for the indicated proteins by WB. **F.** WT and SCAI-null U2OS cells were transfected with the indicated siRNAs, then treated with 1.5 µM cisplatin for 18 h. Whole cell lysates were prepared, and WB were performed against the indicated proteins.

Figure 5. Loss of fork reversal at Stalled Forks confers increased RAD51-depedent fork breakage in the absence of SCAI/REV3.
A. Western blots from WT RPE cells transfected with the indicated siRNAs and then treated with 1 mM HU for 18 h. **B.** WT RPE1 cells were transfected with the indicated siRNAs, then treated with 1 mM HU for 18 h and assayed for the indicated proteins by WB. **C.** WT RPE1 cells were transfected for and treated with 1 mM HU for 18 h. WCLs were prepared, and WB were performed against the indicated proteins. **D.** Western blots from WT RPE cells transfected with the indicated siRNAs and then treated with 2 mM HU for 18 h. **E.** Western blots against the indicated proteins from WT RPE1 cells transfected, then pretreated with 27 µM RAD51 inhibitor B02 or vehicle (DMSO) for 30 min and treated with 1 mM HU for 18 h. **F**. Comet assay of hTERT-RPE1 cells treated with 2 mM HU for 40 h. Indicated siRNAs were transfected two days before HU treatment. Red line represents median, three independent experiments, representative experiment shown. Mann Whitney test. **G.** Western blots from WT RPE cells transfected with the indicated siRNAs, pretreated with 27 µM RAD51 inhibitor B02 or vehicle (DMSO) for 30 min, then treated with 1 mM HU for 18 h.

1117
1118
1119
Figure 6. BRCA1-RAD51 axis governs SLX4 mediated fork breakage in the
absence of SCAI.
A. WT RPE1 cells were transfected and treated with 2 mM HU for 18 h. WCLs were
prepared, and WB were performed against the indicated proteins. **B.** Western blots from
WT RPE cells transfected with the indicated siRNAs and then treated with 1 mM HU for
18 h. **C**. Comet assay of hTERT-RPE1 cells treated with 2 mM HU for 40 h. Indicated
siRNAs were transfected two days before HU treatment. Red line represents median,
three independent experiments, representative experiment shown. Kruskal-Wallis
followed by Dunn's multiple comparison test. ns – not significant (P ≥ 0.05). **D.** WT
RPE1 cells were transfected with the indicated siRNAs, then treated with 1.5 mM HU for
18 h and fractioned prior to blotting for the indicated proteins. SLX4 values are
normalized to ORC2 loading control. **E.** WT RPE1 cells were transfected and then
treated with 2 mM HU for 18 h. Samples were processed for chromatin fraction prior to
blotting for the indicated proteins. **F.** WT RPE1 cells were transfected with the indicated
siRNAs, then treated with 2 mM HU for 18 h and fractioned prior to blotting for the
indicated proteins.
1136
Figure 7. SCAI loss leads to increased genomic instability and cell death
following replication stress.
A. Schematic of cell treatment protocol and numbers of chromosomal aberrations per
metaphase cell. hTERT-RPE1 cells treated with 2 mM HU for 18h followed by washout
of HU and 24 h incubation in normal medium. Indicated siRNAs were transfected two
days before HU treatment. Three independent experiments, mean ± S.D. Ordinary one-
way ANOVA followed by Tukey's multiple comparison test. ns – not significant (P ≥
0.05). **B**. Representative images of A. Red arrow – chromosomal aberrations. **C.** WT
RPE1 cells were transfected, then treated with the indicated doses of HU for 18 h and
allowed to form colonies. Mean ± SD shown. **D.** Representative images from (C) are
shown. **E.** Working model showing proposed roles of BRCA1, RAD51 and SCAI/REV3
upon fork stalling.
1149

SUPPLEMENTARY FIGURE LEGENDS

Figure S1.

A. Top. Schematic of pulse-labeling DNA strands, percentage of single- and two-color fibers and lengths of fibers. DNA fibers were from hTERT-RPE1 cells transfected with siRNAs as indicated.

Figure S2.

A. WT RPE1 cells were transfected with the indicated siRNAs and then treated with 2 mM HU for the indicated times. WB were performed against the indicated proteins. **B.** Western blots against the indicated proteins from WT RPE1 cells pretreated with 10 µM TLS inhibitor JH-RE-06 or vehicle (DMSO) for 30 min, then treated with 1.5 mM HU for 18 h. **C.** WT and SCAI-null U2OS cells were treated with the indicated siRNAs, then treated with 1mM HU for 18 h prior to blotting for the indicated proteins. **D.** IF analyses showing representative RPA foci (left panel) in WT RPE1 cells post-treatment with indicated siRNAs. Cells were treated with 1 mM HU for 18 h prior to IF. Merge includes γH2AX foci and DAPI. Scale bars indicate 5 microns. Intensity of nuclear γH2AX staining was quantified using ImageJ and plotted on the right. ****$P < 0.0001$, Analysis of Variance (ANOVA). **E.** Left, representative immunofluorescence (IF) images showing γH2AX foci in WT RPE1 cells following treatment with the indicated siRNAs. Cells were treated with 1.5 mM HU for 18 h prior to IF analyses. Scale bars indicate 5 microns. Quantification shown on the right. ****$P < 0.0001$ and **$P = 0.0063$, Analysis of Variance (ANOVA). **F.** WT RPE1 cells were transfected with the indicated siRNAs, then treated with 1.5m M HU for 18 h and assayed for the indicated proteins by WB.

Figure S3.

A. WT RPE1 cells were transfected with the indicated siRNAs, then treated with 2 mM HU for 18 h before assaying for the indicated proteins by WB. **B.** Western blots from WT and SCAI-null U2OS cell lines transfected with the indicated siRNAs and then treated with 1.5 µM cisplatin for 18 h.

Figure S4.

A. Western blots from WT and SCAI-null U2OS cell lines transfected with the indicated siRNAs and then treated with 1.5µM cisplatin for 18 h. **B.** WT RPE1 cells were transfected with the indicated siRNAs, then treated with 1 mM HU for 18 h and assayed for the indicated proteins by WB. **C.** P53-null RPE1 parent (Parent), BRCA1-null (ΔBRCA1), and GFP-WT-BRCA1 Rescue (GFP-BRCA1 Rescue) RPE1 cell lines were transfected and treated with 1 mM HU for 18 h. WCLs were prepared, and WB were performed against the indicated proteins. **D.** Western blots from WT, BRCA1/p53-null (ΔBRCA1), and BRCA1/53BP1-null (ΔBRCA1/Δ53BP1) RPE1 cells transfected with the indicated siRNAs and then treated with 1.5 mM HU for 18 h. **E.** WT RPE1 and BRCA1-null cells were transfected with the indicated siRNAs, then treated with 1 mM HU for 18 h and assayed for the indicated proteins by WB. **F.** Western blots against the indicated proteins from WT and SCAI-null U2OS cells transfected with the specified siRNAs, then treated with 1.5 µM cisplatin for 18 h.

Figure S5

A. WT RPE1 cells were transfected with the indicated siRNAs, then treated with 1 mM
HU for 18 h and assayed for the indicated proteins by WB. **B.** Representative images
showing no change in cell cycle distribution in WT RPE1 cells 18 h following treatment
with 27 µM B02 compared to vehicle controls. Quantified on the right, mean ± SD
shown. **C**. Western blots from WT RPE cells transfected with the indicated siRNAs,
pretreated with DMSO, 10 µM Olaparib (Olap), or 27 µM B02, and then treated with 1
mM HU for 18 h.
1203
1204 **Figure S6**
1205 **A** Parent and BRCA1-null RPE1 cells were transfected and then treated with 3 mM HU
1206 for 18 h. Samples were processed for chromatin fraction prior to blotting for the
1207 indicated proteins.
1208

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7