HIF-1α and HIF-2α transcription factors differentially regulate lung alveolar macrophage function

Authors: Elena Priego[1,2]*, Irene Adán-Barrientos[1,2,3]*, Ruth Conde-Garrosa[1], Sarai Martínez-Cano[1,4], Iria Sánchez[1], Diego Mañanes[1,2], Annalaura Mastrangelo[1], Joaquín Amores[1], Helena M. Izquierdo[4#] and David Sancho[1#].

[1]Centro Nacional de Investigaciones Cardiovasculares Carlos III

[2]Escuela de Doctorado, Universidad Autónoma de Madrid.

[3]Departamento de Inmunología, Oftalmología y ORL, Facultad de Medicina, Universidad Complutense de Madrid

[4]Inmunotek S.L. Alcalá de Henares, Madrid

[5]Institut Curie, PSL University, INSERM U932, Immunity and Cancer, Paris, France.

*Co-first authors.

#Senior and corresponding authors: helena.izquierdo-fernandez@curie.fr; dsancho@cnic.es;

ABSTRACT

Alveolar Macrophages (AMs) reside in the alveoli, where oxygen pressure is high, therefore maintaining an active degradation of hypoxia-inducible transcription factors (HIF) mediated by the Von Hippel-Lindau protein (pVHL) ubiquitin ligase complex. We previously found that *Vhl*-deficient AMs not sensing high oxygen pressure are immature and functionally impaired. Here we investigated the specific roles of HIF-1α and HIF-2α isoforms in the regulation of AM functions. With this aim, we combined deletion of *Vhl* with single or double deletion of *Hif1a* and/or *Hif2a* in AMs under the control of the CD11c promoter using a Cre-Lox system. Our work demonstrates that in *Vhl*-deficient macrophages, both HIF-1α and HIF-2α contribute to AM defective self-renewal, while HIF-2α plays a central role in regulating the impaired AM maturation associated with pVHL loss. HIF-1α promotes a glycolytic shift in alveolar macrophages, while HIF-2α hinders lipid oxidation and surfactant clearance. Thus, HIF-2α raises as a selective critical factor restraining the therapeutic potential of AMs to degrade surfactant excess in mice that have

developed pulmonary alveolar proteinosis (PAP). Overall, regulation of both HIF-1α and HIF-
2α isoforms is required for an optimal AM function, highlighting HIF-2α as a potential
pharmacological target for secondary PAP.

INTRODUCTION

Alveolar macrophages (AMs) are lung-resident cells maintained by self-renewal during
homeostasis.[1] Through the sensing of granulocyte-macrophage colony-stimulating factor (GM-
CSF) [2,3] and transforming growth factor β (TGF-β),[4] fetal liver monocytes differentiate into mature
AMs in the lung alveoli during the first week of life. Mature AMs are essential for the regulation of
surfactant in the lung, which is comprised of 90% lipids and is required to decrease surface
tension. AMs have a remarkable ability for lipid catabolism and cholesterol handling,[5] and these
functions are sustained by enzymes and receptors whose expression is regulated by Liver-X
family of receptors (LXRs), Peroxisome proliferator-activated family of receptors (PPARs),
CCAAT enhancer-binding proteins (C/ EBPs), and sterol regulatory element-binding proteins
(SREBPs) transcription factors.[6] Changes in AM lipid metabolism are found in human respiratory
diseases such as chronic obstructive pulmonary disease (COPD) [6] and pulmonary alveolar
proteinosis (PAP).[7,8] Mechanistically in mice, deletion of *Pparg*, a master regulator of lipid
metabolism induced by GM-CSF, selectively impairs the differentiation of Ams.[3] Depletion of the
transcription factor TFAM, involved in mitochondrial biogenesis, leads to a reduction in AM
numbers and impairs their maturation *in vivo.* [5,9] Overall, these studies have identified essential
factors required for AM maturation and function and demonstrate that alterations of AM
metabolism are associated to AM dysfunction. We have previously shown that pVHL is necessary
for the adaptation of AMs to high oxygen levels upon birth, and that *Vhl* deletion renders immature
and functionally impaired AMs, which also present an altered metabolic profile, including
enhanced glycolysis and decreased lipid oxidation capacity.[10] However, the role of each HIFα
isoform in regulating the functional maturation of AMs has not been investigated.

pVHL is the substrate recognition unit of an E3 ubiquitin ligase complex involved in the
ubiquitination and degradation of the alpha isoforms of hypoxia inducible factors (HIFs) in an
oxygen dependent manner.[11–13] HIFs are heterodimeric transcription factors that induce the
expression of a set of genes functionally involved in the cellular adaptation to low oxygen tension
or hypoxia, including the metabolic switch to glycolysis.[14,15] HIFs are composed of an oxygen
sensitive α subunit and a constitutive β subunit. Three α subunits have been described (HIF-1α,
HIF-2α, and HIF-3α), which heterodimerize with HIF-1β subunit (also named Aryl Hydrocarbon
Receptor Nuclear Translocator 1; ARNT1), forming HIF-1, HIF-2 and HIF-3 transcription factors,

respectively.[16] When dimerized with the HIF-1β subunit, HIF-1α and HIF-2α regulate the
transcription of both shared and distinct target genes. HIF-3α acts as a negative regulator of HIF-
1α and HIF-2α by competing for HIF-1β protein available for dimerization, although the HIF-3
dimer also induces the expression of specific target genes.[17] Beyond HIFα regulation, pVHL plays
a role in other cellular processes including cytokine signaling, regulation of senescence, formation
of extracellular matrix and mitochondrial function.[18–25] Notably, in our previous work, we found
that genes related to the adaptation to hypoxia, including many targets of HIF-1α and HIF-2α,
were downregulated within AM maturation after birth.[10] Therefore, we hypothesize that HIF-1α
and HIF-2α must be degraded to ensure the correct functional maturation of AMs in steady-state.

Here, we investigated the specific function of HIF-1α and HIF-2α in AM phenotypic maturation,
self-renewal and surfactant clearance in the lung. To assess this, we generated four different
strains of mice: CD11c-Cre *Vhl*$^{fl/fl}$ (CD11cΔ*Vhl*), CD11c-Cre *Vhl*$^{fl/fl}$*Hif1a*$^{fl/fl}$ (CD11cΔ*Vhl*Δ*Hif1a*),
CD11c-Cre *Vhl*$^{fl/fl}$*Hif2a*$^{fl/fl}$ (CD11cΔ*Vhl*Δ*Hif2a*) and CD11c-Cre *Vhl*$^{fl/fl}$*Hif1a*$^{fl/fl}$*Hif2a*$^{fl/fl}$
(CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a*). We demonstrate that expression of either HIF-1α and HIF-2α alone is
sufficient to intrinsically and differentially impair AM terminal maturation and self-renewal, with full
rescue observed only in CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a* mice. Whereas HIF-1α drives AM metabolic
shift towards glycolysis, HIF-2α disrupts lipid oxidation and surfactant clearance, and is the main
isoform responsible for impaired AM maturation. Altogether, this study shows that HIF inactivation
ensures the optimal maturation of AMs in the lung, regulating the metabolic adaptation to high
oxygen concentration and highlights different functional specificities of each HIFα isoform.

RESULTS

HIF-1α is the driver of the metabolic switch to glycolysis in pVHL-deficient alveolar
macrophages
Our previous work established that pVHL is required for AM phenotypic and functional
maturation,[10] but did not address the specific contribution of HIF-1α and HIF-2α or additional
effects of pVHL. To analyze the specific function of HIF-1α and HIF-2α in AMs, we generated four
different mouse strains: CD11c-Cre *Vhl*$^{fl/fl}$ (CD11cΔ*Vhl*), CD11c-Cre *Vhl*$^{fl/fl}$*Hif1a*$^{fl/fl}$
(CD11cΔ*Vhl*Δ*Hif1a*), CD11c-Cre *Vhl*$^{fl/fl}$*Hif2a*$^{fl/fl}$ (CD11cΔ*Vhl*Δ*Hif2a*) and CD11c-Cre *Vhl*$^{fl/fl}$ *Hif1a*$^{fl/f}$
^{l}*Hif2a*$^{fl/fl}$ (CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a*). Analysis of AMs from these mice revealed the expected
genotype deficiencies in *Vhl*, *Hif1a*, and *Hif2a* expression (**Error! Reference source not**

found.A-C), with additional half reduction in *Hif1a* expression levels in CD11cΔ*Vhl*Δ*Hif2a* mice

(**Error! Reference source not found.**B).

To investigate the contribution of HIF-1α and HIF-2α to the immature-like phenotype found in *Vhl*-

deficient Ams,[10,26] we performed an RNA-seq of AMs isolated from all the mice strains compared

with their respective Cre negative littermate control mice. Analysis of differentially expressed

genes (DEG) showed 1246 genes expressed with at least a 2-fold difference (Adj. p < 0.01)

between each conditional KO and its control, except for CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a* AMs, in which

only *Vhl* and *Plcb4* were differentially expressed (Figure 1A-D), showing that the transcriptional

changes seen in *Vhl*-deficient AM relies exclusively on downstream transcription factors HIF-1α

and HIF-2α rather than on HIF-independent pVHL functions. Principal Component Analysis (PCA)

showed that samples from CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a* scored close to controls and distant from

CD11cΔ*Vhl*, CD11cΔ*Vhl*Δ*Hif1a*, and CD11cΔ*Vhl*Δ*Hif2a* in principal component 1 (PC1) (Figure

1E).

Different genes were deregulated in AMs depending on the genotype (**Error! Reference source**

not found.A-C). Pre-ranked enrichment analysis of the genes with the greatest contribution to

PC1 unveiled pathways related with cell division and metabolism, especially glucose related

pathways, suggesting that both HIF-1α and HIF-2α are involved in their regulation (Figure 1F). To

further interrogate the specific contribution of each HIF isoform, we performed an

overrepresentation analysis, using the Gene Ontology (GO) database, of DEG from each

genotype. Most of the downregulated pathways in *Vhl*-deficient AMs were restored upon further

deletion of *Hif1a* and not *Hif2a*, such as cell cycle, DNA replication, p53 signaling pathway or fatty

acid (FA) metabolism (**Error! Reference source not found.**D). We also found pathways similarly

controlled by HIF-1α and HIF-2α such as *AMPK signaling pathway* or *PI3K-Akt signaling pathway*

(**Error! Reference source not found.**D), suggesting that binding of either HIF isoform to the

same gene promoters is sufficient to activate these pathways. In contrast, *PPAR signaling*

pathway, *HIF1A signaling pathway* or *Glycolysis/Gluconeogenesis* pathways were only partially

restored by HIF-1α or HIF-2α deficiency, indicating their differential regulation by both

transcription factors (**Error! Reference source not found.**D).

HIF-1α and HIF-2α induce the transcription of common target genes involved in the response to

hypoxia. However, stable activation of HIF isoforms with prolonged hypoxia can result in the

predominance of HIF-2α activity rather than HIF-1α.[27] To characterize the contribution of each

HIF isoform to these specific pathways in *Vhl*-deficient AMs, we compared the core gene

expression signature associated with the response to hypoxia KEGG pathway. The upregulated

genes found in *Vhl*-deficient AMs were restored by further deletion of *Hif1a* but not *Hif2a* (**Error!**

Reference source not found.E). The same was true for DEGs associated to oxidative

phosphorylation (OxPhos) and Glycolysis pathways (**Error! Reference source not found.**F-G),

suggesting that HIF-1α is the main regulator of these genesets over HIF-2α in AMs. We also found

that deleting *Hif2a* in *Vhl*-deficient AMs further amplified the transcriptional changes associated

to the response to hypoxia, observed in *Vhl* deletion alone (**Error! Reference source not**

found.F-G). This suggests a negative role of *Hif2a* in the regulation of these pathways. Real-time

metabolic profile of AMs in the presence of glucose, pyruvate, and glutamine by Seahorse showed

that CD11cΔ*Vhl* and CD11cΔ*Vhl*Δ*Hif2a* AM significantly decreased the basal respiratory rate

(BRR) while maintaining extracellular acidification rate (ECAR) compared to controls, reflecting a

decreased use of glucose for TCA and mitochondrial oxidative phosphorylation (OxPhos) towards

an enhanced aerobic glycolysis and lactate production. In contrast, CD11cΔ*Vhl*Δ*Hif1a* and

CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a* AM showed levels of BRR and ECAR similar to WT (Figure 1G-H).

Overall, our results suggest that the transcriptional program governing AM response to hypoxia,

including genes related to OxPhos and glycolysis, is regulated by HIF-1α. Consistent with this,

HIF-1α emerges as the key driver of the metabolic switch towards glycolysis in *Vhl*-deficient AMs.

HIF-1α and HIF-2α differentially regulate AM identity genes in pVHL-deficient mice

We have previously shown that Vhl-deficient AM showed an altered expression of surface

markers,[10] akin to immature cells.[2] We then asked which HIF isoform dampens mature AM identity

in the absence of pVHL. We compared the expression level of classical surface markers that are

associated with AM maturation status. While the expression pattern of maturation markers such

as CD11c, SiglecF, CD11b or MHC class II (I-A/I-E) was restored to control WT levels in

CD11cΔVhlΔHif1aΔHif2a, most of the changes in marker expression found in CD11cΔ*Vhl* mice

remained unchanged upon further depletion of HIF-1α. In contrast, these changes were largely

reversed upon HIF-2α depletion (Figure 2A-E), suggesting that HIF-2α plays a central role in

regulating the impaired AM maturation associated with pVHL loss. We then compared the unique

AM gene expression signature that was identified by the Immgen consortium.[28] Notably, we found

three distinct sets of genes: (i) genes regulated by HIF-2α, with functions in lipid metabolism (Lsr,

Vstm2a, Cidec), (ii) genes regulated by HIF-1α, involved in a plethora of functions including

regulation of cytoskeleton (Clmn, Sema3e, Krt19, Matn2, Itgax), circadian clock genes (Ear1,

Rara) and cell metabolism (Lepr, Naaa, Acaa1b, Aldoc, Fabp1); and (iii) a third set of genes

regulated by both HIF-1α and HIF-2α (Figure 2F). Overall, our data indicate that while deleting

HIF-2α in Vhl-deficient AMs fully restores the control WT surface marker expression, both HIF

isoforms contribute either individually or together shape the subjacent transcriptional identity of
mature AMs (Figure 2F).

AM self-renewal capacity is impaired by both HIF-1α and HIF-2α in pVHL-deficient mice
AMs are lung resident cells maintained locally by self-renewal in steady-state.[29] Enhanced
glycolysis has been correlated with reduced AM self-renewal capacity,[5,10,26] but its association
with specific HIF-1 isoforms remains unclear. Thus, we next investigated which HIF isoform was
regulating AM proliferation in *Vhl*-deficient AMs. The frequency of Ki67[+] AMs was significantly and
similarly reduced in CD11c∆*Vhl*∆*Hif1a* and CD11c∆*Vhl*∆*Hif2a* AMs, indicating that both isoforms
contribute to the defective proliferation in the absence of pVHL (Figure 3A). In contrast, it was
completely restored to control WT in CD11c∆*Vhl*∆*Hif1a*∆*Hif2a* AMs (Figure 3A). In agreement,
the differential expression of genes associated to cell cycle in CD11c∆*Vhl* AM was not restored
by the additional single deletion of *Hif1a* or *Hif2a*, being reverted only in CD11c∆*Vhl*∆*Hif1a*∆*Hif2a*
AMs (**Error! Reference source not found.**A). Furthermore, BrdU uptake in *ex vivo* AM cultures
with GM-CSF show a recovery of their full proliferative capacity only in the *Vhl/Hif1a/Hif2a* triple
knockout (Figure 3B). Overall, these results demonstrate that HIF-1α and HIF-2α impair AM
stemness independently of glycolysis and of environmental extrinsic factors.

AMs can differentiate from circulating monocytes coming from bone marrow precursors.[29] We
then asked whether the previously observed compromised AM proliferative capacity constitutes
a selective disadvantage at reconstituting an empty niche in a context of competence with WT
AM. To address this, we generated bone marrow mixed chimeras: lethally irradiated CD45.1[+]
recipients were transferred intravenously with bone marrow cells from CD45.1[+]CD45.2[+] WT
control   and   CD45.2[+]   CD11c∆*Vhl*,   CD11c∆*Vhl*∆*Hif1a*,   CD11c∆*Vhl*∆*Hif2a*,
CD11c∆*Vhl*∆*Hif1a*∆*Hif2a* or their control Flox littermates in a 1:1 ratio (Figure 3C). After 1.5
months, similar numbers of total AMs were found in all the chimeric mice, ruling out differences
in niche accessibility (**Error! Reference source not found.**B). Frequencies of blood neutrophils
and monocytes from CD45.1[+]CD45.2[+] and CD45.2[+] mice were also similar, indicating an equally
efficient bone marrow reconstitution (**Error! Reference source not found.**C,D). In the
bronchoalveolar lavage (BAL), CD45.2[+] CD11c∆*Vhl* and CD11c∆*Vhl*∆*Hif2a* monocyte-derived
AM were significantly less abundant than CD45.1[+]CD45.2[+] control WT cells (Figure 3D),
correlating with a decreased frequency of Ki67[+] AMs (Figure 3E,F). Unexpectedly,
CD11c∆*Vhl*∆*Hif1a* monocyte-derived AM also showed a significantly reduced percentage of Ki67[+]
cells but reconstituted the alveolar niche as efficiently as control WT AM, suggesting a survival
advantage (Figure 3D-F). These results indicate that while both HIF isoforms regulate AM self-

renewal capacity, HIF-1α overexpression may primarily underlie impaired alveolar niche
reconstitution upon pVHL absence.

In this bone marrow-graft model, CD11cΔ*Vhl*, CD11cΔ*Vhl*Δ*Hif1a*, and CD11cΔ*Vhl*Δ*Hif2a*
monocyte-derived AMs expressed similar levels of CD11c (Figure 3G) but did not completely
restore CD45.1⁺CD45.2⁺ WT control levels for Siglec-F, CD11b, and MHC class II (I-A/I-E) (Figure
3H-J). In contrast, AMs from WT littermates showed frequencies of Ki67⁺ and levels of surface
markers similar to CD45.1⁺CD45.2⁺ WT controls (**Error! Reference source not found.**E-I). These
results confirm that the HIF-dependent defect in the maturation and proliferation of *Vhl*-deficient
AM is cell-intrinsic.

HIF-2α selectively compromises surfactant clearance capacity by VHL-deficient AMs
In lung alveoli, AMs play a major role degrading the surfactant produced by lung type II epithelial
cells. Genetic defects in GM-CSF signaling prevent AM maturation and lead to the development
of spontaneous PAP, characterized by the accumulation of surfactant.[30] HIF-1α mediated switch
to glycolysis has been linked to the accumulation of lipids and the development of foam cells. [31–
33] We previously reported that CD11cΔ*Vhl* AMs have a decreased capacity to oxidize lipids *ex
vivo*, and consequently, when adoptively transferred to *Cfsr2⁻ᐟ⁻* mice, which lack AMs and develop
PAP spontaneously, they are not able to reduce the surfactant content in the alveoli.[10] To
interrogate on the regulation of lipid metabolism by HIF-1α or HIF-2α *in vivo*, we intratracheally
transferred AMs from CD11cΔ*Vhl*Δ*Hif1a*, CD11cΔ*Vhl*Δ*Hif2a*, CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a*, or their
control counterparts into *Cfsr2⁻ᐟ⁻* mice. Only adoptively transferred CD11cΔ*Vhl*Δ*Hif2a* AMs and
CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a* AMs were able to decrease total protein and specific SP-D protein in
BAL to levels similar to WT (Figure 4A,B), indicating that HIF-2α stabilization impairs surfactant
clearance by lung AMs. Similar numbers of AMs were recovered from BAL for all genotypes,
indicating the same efficiency at replenishing the lung alveoli (**Error! Reference source not
found.**A).

AMs rely on fatty-acid oxidation and cholesterol efflux to metabolize lipids and avoid their
accumulation respectively.[5] To assess the regulation of lipid oxidation by HIF isoforms in a
controlled *in vitro* setting, we conducted real-time metabolic profile with the fatty acid palmitoyl-
CoA as the only carbon source. BRR was significantly reduced in CD11cΔ*Vhl* and
CD11cΔ*Vhl*Δ*Hif1a* AMs compared to WT but it was completely restored in CD11cΔ*Vhl*Δ*Hif2a* and
CD11cΔ*Vhl*Δ*Hif1a*Δ*Hif2a* AMs (Figure 4C).

We then compared the expression of fatty acid and cholesterol metabolism related genes among
genotypes. In agreement with our *in vivo* and *in vitro* results, we found the expression of multiple
DEG associated with FA oxidation in *Vhl*-deficient AMs were commonly found in
CD11cΔ*Vhl*Δ*Hif1a* but WT expression profile was restored in CD11cΔ*Vhl*Δ*Hif2a* AM (Figure 4D).
Globally these results highlight HIF-2α is a major regulator of FA oxidation in lung AMs.

DISCUSSION

AMs are derived from embryonic progenitors that colonize the lung alveoli and mature after birth.
Maturation onset coincides with the exposure of the alveoli to high oxygen concentration upon
the first breath. We previously showed that genes related to the adaptation to hypoxia, including
HIF-target genes, are downregulated during AM maturation.[10] The role of HIF-1α is well studied
in monocyte-derived macrophages, where it plays a major role in macrophage infiltration and
proinflammatory responses.[34–38] Chronic and acute lung disease due to influenza infection
courses with lung edema and cellular infiltrations that can lead to hypoxia. [39,40] HIF-1α stabilization
has been reported in lung AMs upon lung injury associated to influenza infection, leading to
glycolysis induction.[41] However, the role of the HIF isoforms in AM metabolic adaptation to their
niche in homeostasis is largely unknown. In the present study, we have uncovered that HIF-1α
and HIF-2α differentially impair specific AM functions. Thus, the inactivation of HIF function is
essential for lung AM functional maturation. Our results demonstrate that HIF-1α is the exclusive
driver of the glycolytic switch in Vhl-deficient AMs. Notably, both HIF-1α and HIF-2α play a role in
the phenotypic maturation and transcriptional identity of AMs, corroborated by the cell-intrinsic
nature of the altered surface marker expression found in monocyte-derived AM deficient for Vhl,
Vhl/Hif1a and Vhl/Hif2a. HIF-1α stabilization leads to the expression of the antimicrobial enzyme
iNOS [42] and is protective against lung fibrosis.[26] This suggests that HIF activation in response to
certain exogenous stimuli can be beneficial, whereas under sterile conditions, HIF activity must
be tightly regulated.

Increased glycolysis has been linked with a decreased self-renewal potential in AMs.[5,43] Ki67
expression and BrdU uptake assays indicated that both HIF-1α and HIF-2α stabilization similarly
impair AM proliferation in the absence of pVHL. Despite the significant defect in proliferation of
Vhl-deficient AM, total numbers in the lung alveoli were similar to WT, suggesting that they can
fully replenish the alveolar niche in the absence of competition from incoming blood monocytes.
Indeed, our experiments with mixed bone marrow chimeras confirmed that that *Vhl*-deficient AM
defective proliferation is cell-intrinsic and that *Vhl*- and *Vhl/Hif2a*-deficient AMs are significantly
outcompeted by control WT AM. In contrast, *Vhl/Hif1a*-deficient AMs were similarly able to

repopulate the alveoli despite decreased Ki67 expression compared to control WT AM, what could
be related with the pro-survival role of HIF-1α [41] although further investigation is required.

Notably, we found that compared to *Vhl/Hif1a*-deficient AMs, *Vhl/Hif2a*-deficient AMs had
impaired oxidative capacity in the presence of glucose and glutamine as carbon source. In
contrast, *Vhl/Hif1a*-deficient AMs showed a decreased oxygen consumption when using FA as
carbon source, a phenomenon not observed in Vhl/Hif2a deficient AMs. Investigation on the
specific roles of HIF-1α and HIF-2α in the regulation of lipid metabolism pathways is still
controversial. HIF-2α has a dominant role leading to hepatic steatosis through fatty acid-oxidation
impairment upon depletion of pVHL.[44] In contrast, HIF-1α was reported to enhance hepatic lipid
accumulation in a model of alcoholic fatty liver disease.[32] Finally, another study shows that
deletion both HIF isoforms is sufficient to ameliorate the lipid accumulation induced by hypoxia in
HepG2 cell line.[45] The discrepancy among these results may rely on the different trigger used to
activate HIF (pVHL depletion, alcohol and hypoxia, respectively), which likely involves a different
dynamic of HIF isoform activation.[27] In the particular case of macrophages, the source of these
cells may determine their metabolic requirements: monocyte derived macrophages (MDM) under
hypoxia accumulate lipids and become foam cells, which correlates with the activation of HIF-
1α.[31] HIF-1α can prevent cholesterol efflux contributing to lipid accumulation in a macrophage cell
line.[33] However, MDM strongly rely on glycolysis compared to lung AM at a basal level,[46]
suggesting that the metabolic regulation of MDMs cannot be extrapolated to tissue resident
macrophages. In agreement, a recent study has shown that lung AMs are enriched in a fatty acid
oxidation module including enzymes involved in lysosomal lipolysis, mitochondrial FA import and
oxidation,[47] Our results unveil HIF-2α as the key regulator of lipid oxidation in lung AMs. *Vhl/Hif2a*-
deficient AMs completely restore homeostatic levels of surfactant upon transfer to a mouse model
of PAP and restore basal OCR levels when using Palmitoyl-CoA as substrate *in vitro*. In
agreement, our transcriptional data shows multiple FA-related genes dysregulated in *Vhl*-deficient
AMs, which were only restored upon further *Hif2a* inactivation. Overall, this work demonstrates
the specific roles of HIF-1α and HIF-2α at fine-tuning lung AM gene expression and functions.
Our results highlight the importance of developing therapies targeting each HIF isoform and unveil
HIF-2α as a potential target to improve surfactant removal in secondary PAP diseases.

Figure 1. HIF1α drives glycolysis in VHL-deficient alveolar macrophages.

(A-D) Volcano plot of differentially expressed genes ranked showing adjusted p-value (Log10 adj. p-value) and log2 fold change, between AMs from CD11cΔVhl (A), CD11cΔVhlΔHif1a (B), CD11cΔVhlΔHif2a (C) and CD11cΔVhlΔHif1aΔHif2a (D) mice compared to their littermate control; differentially expressed genes (adjusted p-value < 0.05 and abs(log2FC) > 1) are depicted in the color code associated with each genotype. (E) Principal component analysis (PCA) of transcriptomic profiles of AMs from CD11cΔVhl, CD11cΔVhlΔHif1a, CD11cΔVhlΔHif2a, CD11cΔVhlΔHif1aΔHif2a mice, and their littermate controls. Each dot represents a pull of 2-3 mice. (F) Pre-ranked enrichment analysis of the genes defining PC1 using KEGG Pathways. NES, normalized enrichment score; p-value adjusted with Benjamin–Hochberg correction. (G) Oxygen consumption rate (OCR) by BAL AMs cultured in media containing glucose, glutamine, and pyruvate. Data are shown as mean ± SD. (H, I) Basal respiratory rate (BRR) (H) and extracellular acidification rate (ECAR) (I) of BAL AMs. Each dot represents a pool of n = 6-7 mice per genotype. Paired Student's t-test, *p<0.05 and "ns" stands for non statistically significant. See also Figures S1, S2.

Figure 2. Mature AM transcriptional signature relies on HIF inactivation.

(A) Representative flow cytometry histograms for CD11c, Siglec-F, CD11b and IA/IE expression by BAL AMs from the indicated genotypes. (B-E) Quantification of geometric mean of fluoresce intensity (GMFI) relative to the mean value of control samples (rGMFI) of CD11c (B), SiglecF (C), CD11b (D) and IA/IE (E). Pull of 2-3 experiments, n = 4-6 mice per experiment. Student's *t*-test, ****p<0.0001, ***p<0.001, **p<0.01 and "ns" stands for non-statistically significant. (F) Gene expression heatmap of the lung AM-specific signature gene set described in.[28]

286
287

Figure 3. Both HIF isoforms interfere with AM self-renewal.

(A) Quantification of the frequency of Ki67+ BAL AMs relativized to the mean value for WT. Pull of 3 experiments, n= 5-8 mice per experiment. Student's *t*-test, ****p<0.0001, and "ns" stands

for non statistically significant. (B) Quantification of the frequency of BrdU+ BAL AMs after 3 days in culture with rGM-CSF. Pool of 2 experiments, each dot represents a pool of n = 15-18 mice per experiment . Student's *t*-test, **p<0.01, *p<0.05 and "ns" stands for non statistically significant. (C) Representative dot plot showing the percentage of host CD45.1[+], control transplanted CD45.1[+]CD45.2[+] and experimental transplanted CD45.2[+] cells, gated within SiglecF[+] CD11c[+] BAL AMs (D) Quantification of the percentage of each AM subpopulation. Bar indicates mean ± SEM, Student's *t*-test, ****p<0.0001 and "ns" stands for non statistically significant. (E) Representative dot plots showing Ki67 expression by CD45.2[+] SiglecF[+]CD11c[+] gated BAL AMs 45 days after BM transfer, in which CD45.1[+] and CD45.1[-] correspond to control and experimentally transplanted, respectively. (F) Quantification of the frequency of Ki67+ AMs of each genotype relativized to the mean value for control CD45.1[+] AMs. Bar indicates mean ± SEM, Student's *t*-test, ****p<0.0001, **p<0.01. (G-J) Quantification of geometric mean of fluoresce intensity (GMFI) relative to the mean value of control CD45.1[+] AMs samples (rGMFI) of CD11c (G), Siglec-F (H), CD11b (I) and IA/IE (J). Bar indicates mean ± SEM, Student's *t*-test, ****p<0.0001, **p<0.01, *p<0.05 and "ns" stands for non statistically significant. (D-J) Pool of experiments, n = 6-8 chimeras per experiment. See also Figure S3.

Figure 4. HIF-2 impairs fatty acid oxidation and surfactant clearance by AMs.

Quantification of total protein (A) and surfactant protein D (SP-D) (B) in BAL fluid from Csfr2b-/- mice transplanted with BAL AMs from the indicated genotypes. Control mice (Csfr2b$^{-/-}$ and Csfr2b$^{+/-}$) are pooled from each independent experiment. Pool of 3 experiments, n= 3-10 mice per experiment, Student's t-test, ****$p<0.0001$, **$p<0.01$, *$p<0.05$ and "ns" stands for non statistically significant. (C) Quantification of basal respiratory rate (BRR) of BAL AMs in media containing palmitoyl-CoA and L-carnitine. Pool of 4 experiments, each dot represents a pool of n=6-7 mice. Student's t-test, **$p<0.01$, *$p<0.05$ and "ns" stands for non statistically significant. (D) Heatmap showing differential expression of genes related to lipid metabolic process GO pathway (GO:0019216) in the indicated genotypes. See also figure S4.

MATERIALS AND METHODS

Data. RNA-seq data from Control-*Vhl* and CD11cΔVhl was previously published under GEO: GSE108975 accession number. RNA-seq performed in this study has been deposited at GEO and will be publicly available as of the date of publication (accession number).

Mice. Mice colonies were bred at Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC) under specific pathogen-free conditions. Experiments were conducted with 8- to 10-weeks-old age-matched littermates (regardless of gender). Floxed *Vhl* mice were backcrossed with or without floxed *Hif1a* and floxed *Hif2a*. Additionally, all possible combinations were backcrossed with CD11c-Cre mice (B6.Cg-Tg(*Itgax*-cre)1-1Reiz/J) (kindly provided by Boris Reizis) [48] in order to allow the targeting of the alveolar macrophage compartment. *Csf2rb*[-/-] mice (kindly provided by Martin Guilliams),[49] and F1 from the mating between C57BL/6J mice expressing the CD45.2 allele (Charles River) and B6/SJL (Ptprca Pepcb/BoyJ) mice expressing the CD45.1 allele were also used. All experimental procedures were approved by the institutional care and use committees and conformed to EU Directive 2010/63EU and Recommendation 2007/526/EC regarding the protection of animals used for experimental and other scientific purposes, enforced in Spanish law under Royal Decree 53/2013.

AM isolation. Mice were sacrificed with a lethal dose of pentobarbital (Dolethal, Vetoquinol). To isolate AMs for cell culture, ten bronchoalveolar lavages (BAL) were done by inserting a venal catheter (18G BD Pharmingen) into the trachea with 1ml of phosphate-buffered saline (PBS), 5 mM EDTA, and 2 % inactivated fetal bovine serum (FBS, Sigma-Aldrich) (FACS buffer) at 37ºC. Samples were kept on ice until further processing. BAL was centrifuged at 300xg for 10' at 4ºC and red blood cell lysis was performed at room temperature (RT) for 3 min with RBC Lysis Buffer (Sigma-Aldrich). Cells were resuspended in medium or FACs buffer depending on the application.

AM *in vitro* culture. AMs from BAL were cultured in RPMI 1640 (Gibco) supplemented with 10% FBS, 100 U/ml Penicillin and Streptomycin (Lonza), 1mM Sodium Pyruvate (HyClone™), and 2 mM L-glutamine (complete RPMI) at 37ºC and 5% CO_2. For *in vitro* proliferation assay, cells were supplemented with 10 pg/ml rGM-CSF (Peprotech). AMs were pulsed twice with BrdU (BD Pharmingen) at 10µM: 10-12 h and 24h after culture onset. 24h after the second pulse, cells were washed once with PBS to remove remaining FBS and incubated with accutase (StemCell Technologies) for 15-20' at 37ºC. Cells were carefully detached by pipetting. After centrifugation at 600 G for 5' at 4ºC, cells were resuspended and counted, and the same cell number per condition and genotype was stained for surface markers and BrdU.

Flow cytometry. Single-cell suspensions were stained in 96-U-bottom plates. Surface staining

was performed in FACS buffer: PBS, 2.5 mM EDTA and 2.5% heat-inactivated FBS (iFBS).

Staining was performed for 15-20' at 4ºC with FcR block anti-mouse CD16/CD32 (2.4G2, Tonbo

Biosciences) and the appropriate mix of the following fluorochrome-conjugated antibodies in

FACS buffer: CD11c (clone HL3, BD Biosciences), Siglec-F (clone E50-2440, BD Biosciences),

CD64 (clone X54-5/7.1, BD Biosciences), CD11b (clone M1/70, BD Biosciences), F4/80 (clone

BM8, Life Technologies), MHCII (clone M5/114.15.2 BD Biosciences), Ly6G (clone 1A8, BD

Pharmingen), Ly6C (clone AL21, BD Biosciences), CD45 (30-F11, eBiosciences), CD45.1 (A20,

eBiosciences), CD45.2 (clone 104, BD Biosciences). Hoechst 33258 (Invitrogen) or Ghost Dye

Red 780 (Cell Signaling) were used as a counterstain to exclude dead cells. For intracellular Ki67

staining, AMs were stained for surface markers as described above, washed twice with FACs

buffer at 600 G for 5' at 4ºC, and mixed with unstained thymocytes, used as a cell carrier. Fixation

and permeabilization of cells were performed using the Foxp3/Transcription Factor Staining Buffer

Set (#00-5523-00, Invitrogen) and following manufacturer's instructions. Briefly, fixation solution

was prepared by diluting Fixation/Permeabilization concentrate, 1:3 with Fixation/Perm diluent.

Permeabilization buffer 10x was diluted 1:10 with sterile water and kept on ice. Cells were fixed

for 30' at 4ºC in the dark. Cells were washed with permeabilization buffer 1x and intracellular (i.c.)

staining was done right after. Intracellular staining with anti-Ki67-eFluor 660 (SolA15,

eBiosciences) or isotype control Rat IgG2aκ (eBR2a, eBiosciences) was performed in

permeabilization buffer 1x. All centrifugations after fixation were done at 1000 G for 5' at 4ºC.

Cells were resuspended in FACS buffer before acquisition. Intracellular BrdU staining was done

following manufacturer instructions. In brief, AMs were stained and mixed with carrier cells as

above-mentioned and fixed and permeabilized using the BrdU staining set (BD Pharmingen).

Cells were treated with DNAse and stained with anti-BrdU Ab (BD Pharmingen). Samples were

acquired in SP6800 Spectral, LSRFortessa, or FACSymphony cell analyzers, and data was

analyzed using FlowJo Version 10 software (Tree Star).

Real-time oxygen consumption rate and glycolytic flux. Real-time oxygen consumption rate

(OCR) and extracellular acidification rate (ECAR) were determined with Seahorse XF-96

Extracellular Flux Analyzer (Agilent Technologies). AMs from fresh BAL were pooled from 5-10

mice per genotype. After red blood cell lysis, 2.5×10^5 AMs were plated in Cell-Tak (Corning)

coated wells. The assay was performed in DMEM supplemented with 100 µg/ml penicillin, 100

µg/ml streptomycin, and either 25 mM glucose, 1 mM pyruvate, and 2 mM glutamine or 5 mM L-

carnitine and 50 µM palmitoyl-CoA (all Sigma). pH was adjusted to 7.4 with KOH. Three

consecutive measurements were performed under basal conditions and after the sequential

addition of the following inhibitors: 1 µM Oligomycin, 1 µM Carbonyl cyanide 3-
chlorophenylhydrazone (CCCP), and 1 µM Rotenone with 1 µM Antimycin A. Basal respiratory
rate (BRR), maximal respiratory rate (MRR), spare respiratory capacity (SRC) and basal ECAR
were calculated according to Seahorse Agilent indications.

RNA sequencing and analysis. CD11c cell positive selection was performed on BAL
suspensions from 3-5 mice per genotype. Mouse CD11c MicroBeads UltraPure and MS columns
(all Miltenyi Biotec) were used according to manufacturers' instructions. Briefly, cells were
resuspended in 100 µL of FACS buffer plus 20 µL of CD11c MicroBeads, and incubated on ice
for 30 mins. Then, cells were washed and purified by magnetic positive selection with MS columns
(Miltenyi). A total of 3 pools of 3-5 mice per genotype were included. Total RNA was isolated using
Rneasy Micro Kit (Qiagen), and RNA quality was validated with a 2100 Bioanalyzer (Agilent).
RNA sequencing was performed in the Genomics Unit of the CNIC on Illumina HiSeq 2500
System. Sequencing reads were pre-processed through a pipeline that used FastQC to assess
read quality and Cutadapt to trim sequencing reads, eliminate Illumina adaptor remains, and
discard reads that were shorter than 30 base pairs. The resulting reads were mapped against the
mouse transcriptome (GRCm38.99) with STAR. Expected expressions were quantified using
RSEM, and batch normalization was applied with Combat as implemented in the sva R package.
Genes were filtered to have at least 10 CPM, and data were normalized using the trimmed mean
of m values (TMM), both with edgeR. DESeq2 rlog-normalized values were used to generate
Hexagonal Triwise diagrAMs with Triwise package. Significantly different genes for each pair-
wise comparison (p.adj < 0.05, logFC > 1.5) are displayed as red dots in the plots. Genes were
ranked according to log-Fold change, and gene set enrichment analysis was performed using
FGSEA algorithm and KEGG and GO databases. Redundant GO terms were removed using
GOSemSim package. Heatmaps display deregulated genes (Adj. p-value ≤ 0.05) of the specified
gene set.

For PCA-based enrichment analysis, $\log_2$(CPM) samples were scaled and used as input for
principal component analysis (PCA). Genes were ranked according to the loading vectors of PC1.
Resulting ranks were used as input for the FGSEA algorithm along with the Kyoto Encyclopedia
of Genes and Genomes (KEGG) (www.genome.jp/kegg) and Gene Ontology (GO)
(www.geneontology.org/) databases in order to identify the main sources of variability detected
by PCA. Only gene sets with a number of genes ≥ 15 and ≤ 500 were considered for PCA-based
enrichment analysis.

Bone Marrow Transfer mixed chimeras. B6/SJL mice expressing the CD45.1 congenic marker were lethally irradiated with 2 doses of 6 Gy that were 3 hours apart and reconstituted intravenously with 5×10^6 cells of a 1:1 mixture of CD45.1$^+$ CD45.2$^+$ wild-type BM cells and CD45.2$^+$ BM cells coming from the indicated experimental genotypes. Mice were analyzed 45-50 days after reconstitution as previously described. A small sample of peripheral blood was harvested one day before the experimental endpoint to test immune compartment reconstitution.

Pulmonary macrophage transplantation. 5×10^4 AMs from BAL were transferred intratracheally into Csf2rb$^{-/-}$ mice. Recipient mice were anesthetized with Ketamine (Imalgene, Merial) and Xylazine (Rompun, Bayer). When entirely unconscious, a small incision was made to partially expose the trachea. Then, mice were carefully intubated orotracheally with an IV catheter (22 G; BD Insyte). Cells were inoculated with a pipette in 30µl of PBS. Next, mice were extubated, and the incision sutured. Mice were finally injected with anesthesia reversor (Medeson, Urano) and kept in a warm plaque until complete recovery. 1ml PBS1x-BAL from recipient mice was collected six weeks after transplantation. Cells were centrifuged at 1700 rpm for 5 min at RT and supernatant was used to measure total protein concentration by BCA (Pierce BCA Protein Assay Kit, Thermo Scientific) and surfactant protein D (SP-D) concentration by ELISA (SP-D DUO ELISA kit, R&D System), according to the manufacturer's instruction. Cell engraftment was compared by flow cytometry of total lung cells.

Statistical analysis. Statistical comparisons were analyzed using R v.4.2.1. Data are presented as individual values and were analyzed as indicated in the legends or the dedicated methods section. All experiments were repeated at least twice, and pooled data from several experiments are shown as indicated in the legends. All *n* values represent biological replicates (different mice, primary cell preparations, or in vitro experiments). Differences were considered significant when $p \leq 0.05$ and are indicated as *ns* when not significant, *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$.

ACKNOWLEDGEMENTS

We thank members of the DS laboratory at CNIC, past and present, for feedback and scientific discussions on this project and manuscript. Thanks also to all CNIC facilities and assistants for their support. Work in the DS laboratory is funded by the CNIC; by Ministerio de Ciencia, Innovación y Universidades (MICIU) PID2022-137712OB-I00, CPP2021-008310 and CPP2022-009762 MICIU/AEI/10.13039/501100011033 Agencia Estatal de Investigación, Unión Europea

NextGenerationEU/PRTR; by Comunidad de Madrid (P2022/BMD-7333 INMUNOVAR-CM); by

Scientific Foundation of the Spanish Association Against Cancer (AECC- PRYGN246642SANC);

by Worldwide Cancer Research WWCR-25-0080; by European Union ERC-2023-PoC; by a

research agreement with Inmunotek S.L.; and by "la Caixa" Foundation (LCF/PR/HR23/52430012

and LCF/PR/HR22/52420019).

The CNIC is supported by the Instituto de Salud Carlos III (ISCIII), the MICIU and the Pro CNIC

Foundation, and is a Severo Ochoa Center of Excellence (CEX2020-001041-S funded by

MICIU/AEI /10.13039/501100011033).

REFERENCES

1. Hashimoto, D., Chow, A., Noizat, C., Teo, P., Beasley, M.B., Leboeuf, M., Becker, C.D., See, P., Price, J., Lucas, D., et al. (2013). Tissue-Resident Macrophages Self-Maintain Locally throughout Adult Life with Minimal Contribution from Circulating Monocytes. Immunity *38*, 792–804. https://doi.org/10.1016/j.immuni.2013.04.004.

2. Guilliams, M., De Kleer, I., Henri, S., Post, S., Vanhoutte, L., De Prijck, S., Deswarte, K., Malissen, B., Hammad, H., and Lambrecht, B.N. (2013). Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF. J. Exp. Med. *210*, 1977–1992. https://doi.org/10.1084/jem.20131199.

3. Schneider, C., Nobs, S.P., Kurrer, M., Rehrauer, H., Thiele, C., and Kopf, M. (2014). Induction of the nuclear receptor PPAR-γ by the cytokine GM-CSF is critical for the differentiation of fetal monocytes into alveolar macrophages. Nat. Immunol. *15*, 1026–1037. https://doi.org/10/f6msqc.

4. Evren, E., Ringqvist, E., and Willinger, T. (2020). Origin and ontogeny of lung macrophages: from mice to humans. Immunology *160*, 126–138. https://doi.org/10.1111/imm.13154.

5. Wculek, S.K., Heras-Murillo, I., Mastrangelo, A., Mañanes, D., Galán, M., Miguel, V., Curtabbi, A., Barbas, C., Chandel, N.S., Enríquez, J.A., et al. (2023). Oxidative phosphorylation selectively orchestrates tissue macrophage homeostasis. Immunity *56*, 516-530.e9. https://doi.org/10.1016/j.immuni.2023.01.011.

6. Fujii, W., Kapellos, T.S., Baßler, K., Händler, K., Holsten, L., Knoll, R., Warnat-Herresthal, S., Oestreich, M., Hinkley, E.R., Hasenauer, J., et al. (2021). Alveolar macrophage transcriptomic profiling in COPD shows major lipid metabolism changes. ERJ Open Res. *7*. https://doi.org/10.1183/23120541.00915-2020.

7. Robb, L. (1995). Hematopoietic and lung abnormalities in mice with a null mutation of the common f3 subunit of the receptors for granulocyte-macrophage colony-stimulating factor and interleukins 3 and 5. Proc Natl Acad Sci USA, 5. https://doi.org/10/b7ckwn.

8. Stanley, E., Lieschke, G.J., Grail, D., Metcalf, D., Hodgson, G., Gall, J.A., Maher, D.W., Cebon, J., Sinickas, V., and Dunn, A.R. (1994). Granulocyte/macrophage colony-stimulating factor-deficient mice show no major perturbation of hematopoiesis but develop a characteristic pulmonary pathology. Proc. Natl. Acad. Sci. *91*, 5592–5596. https://doi.org/10/cz3dc3.

9. Gao, X., Zhu, B., Wu, Y., Li, C., Zhou, X., Tang, J., and Sun, J. (2022). TFAM-Dependent Mitochondrial Metabolism Is Required for Alveolar Macrophage Maintenance and Homeostasis. J. Immunol. *208*, 1456–1466. https://doi.org/10.4049/jimmunol.2100741.

10. Izquierdo, H.M., Brandi, P., Gómez, M.-J., Conde-Garrosa, R., Priego, E., Enamorado, M., Martínez-Cano, S., Sánchez, I., Conejero, L., Jimenez-Carretero, D., et al. (2018). Von Hippel-Lindau Protein Is Required for Optimal Alveolar Macrophage Terminal Differentiation, Self-Renewal, and Function. Cell Rep. *24*, 1738–1746. https://doi.org/10.1016/j.celrep.2018.07.034.

11. Iliopoulos, O., Levy, A.P., Jiang, C., Kaelin, W.G., and Goldberg, M.A. (1996). Negative regulation of hypoxia-inducible genes by the von Hippel-Lindau protein. Proc. Natl. Acad. Sci. *93*, 10595–10599. https://doi.org/10.1073/pnas.93.20.10595.

12. Maxwell, P.H., Wiesener, M.S., Chang, G.-W., Clifford, S.C., Vaux, E.C., Cockman, M.E., Wykoff, C.C., Pugh, C.W., Maher, E.R., and Ratcliffe, P.J. (1999). The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. Nature *399*, 271–275. https://doi.org/10.1038/20459.

13. Stebbins, C.E., Kaelin, W.G., and Pavletich, N.P. (1999). Structure of the VHL-ElonginC-ElonginB
complex: implications for VHL tumor suppressor function. Science *284*, 455–461.
https://doi.org/10.1126/science.284.5413.455.

14. Kaelin, W.G. (2022). Von Hippel–Lindau disease: insights into oxygen sensing, protein degradation,
and cancer. J. Clin. Invest. *132*. https://doi.org/10.1172/JCI162480.

15. Wang, G.L., Jiang, B.H., Rue, E.A., and Semenza, G.L. (1995). Hypoxia-inducible factor 1 is a basic-
helix-loop-helix-PAS heterodimer regulated by cellular O2 tension. Proc. Natl. Acad. Sci. *92*, 5510–
5514. https://doi.org/10.1073/pnas.92.12.5510.

16. Koh, M.Y., and Powis, G. (2012). Passing the baton: the HIF switch. Trends Biochem. Sci. *37*, 364–
372. https://doi.org/10.1016/j.tibs.2012.06.004.

17. Yang, S.-L., Wu, C., Xiong, Z.-F., and Fang, X. (2015). Progress on hypoxia-inducible factor-3: Its
structure, gene regulation and biological function (Review). Mol. Med. Rep. *12*, 2411–2416.
https://doi.org/10.3892/mmr.2015.3689.

18. Hu, L., Xie, H., Liu, X., Potjewyd, F., James, L.I., Wilkerson, E.M., Herring, L.E., Xie, L., Chen, X.,
Cabrera, J.C., et al. (2020). TBK1 Is a Synthetic Lethal Target in Cancer with *VHL* Loss. Cancer Discov.
10, 460–475. https://doi.org/10.1158/2159-8290.CD-19-0837.

19. Frew, I.J., and Krek, W. (2008). pVHL: A Multipurpose Adaptor Protein. Sci. Signal. *1*.
https://doi.org/10/dg4k3s.

20. Ji, Q., and Burk, R.D. (2008). Downregulation of integrins by von Hippel-Lindau (VHL) tumor
suppressor protein is independent of VHL-directed hypoxia-inducible factor alpha degradation.
Biochem. Cell Biol. *86*, 227–234. https://doi.org/10.1139/O08-035.

21. Yang, H., Minamishima, Y.A., Yan, Q., Schlisio, S., Ebert, B.L., Zhang, X., Zhang, L., Kim, W.Y., Olumi,
A.F., and Kaelin, W.G. (2007). pVHL Acts as an Adaptor to Promote the Inhibitory Phosphorylation of
the NF-κB Agonist Card9 by CK2. Mol. Cell *28*, 15–27. https://doi.org/10.1016/j.molcel.2007.09.010.

22. Hergovich, A., Lisztwan, J., Barry, R., Ballschmieter, P., and Krek, W. (2003). Regulation of microtubule
stability by the von Hippel-Lindau tumour suppressor protein pVHL. Nat. Cell Biol. *5*, 64–70.
https://doi.org/10.1038/ncb899.

23. Li, S., Li, W., Yuan, J., Bullova, P., Wu, J., Zhang, X., Liu, Y., Plescher, M., Rodriguez, J., Bedoya-
Reina, O.C., et al. (2022). Impaired oxygen-sensitive regulation of mitochondrial biogenesis within the
von Hippel–Lindau syndrome. Nat. Metab. *4*, 739–758. https://doi.org/10.1038/s42255-022-00593-x.

24. Briston, T., Stephen, J.M., Thomas, L.W., Esposito, C., Chung, Y.-L., Syafruddin, S.E., Turmaine, M.,
Maddalena, L.A., Greef, B., Szabadkai, G., et al. (2018). VHL-Mediated Regulation of CHCHD4 and
Mitochondrial Function. Front. Oncol. *8*, 1–16. https://doi.org/10.3389/fonc.2018.00388.

25. Zhu, S., Ding, W., Chen, Y., Wang, W., Xu, R., Liu, C., Liu, X., and Deng, H. (2022). High *VHL*
Expression Reverses Warburg Phenotype and Enhances Immunogenicity in Kidney Tumor Cells.
Genomics Proteomics Bioinformatics *20*, 657–669. https://doi.org/10.1016/j.gpb.2019.12.002.

26. Zhang, W., Li, Q., Li, D., Li, J., Aki, D., and Liu, Y.-C. (2018). The E3 ligase VHL controls alveolar
macrophage function via metabolic–epigenetic regulation. J. Exp. Med. *215*, 3180–3193.
https://doi.org/10.1084/jem.20181211.

27. Jaśkiewicz, M., Moszyńska, A., Króliczewski, J., Cabaj, A., Bartoszewska, S., Charzyńska, A., Gebert,
M., Dąbrowski, M., Collawn, J.F., and Bartoszewski, R. (2022). The transition from HIF-1 to HIF-2

511   during prolonged hypoxia results from reactivation of PHDs and HIF1A mRNA instability. Cell. Mol.
512   Biol. Lett. *27*, 109. https://doi.org/10.1186/s11658-022-00408-7.

513 28. Gautier, E.L., Shay, T., Miller, J., Greter, M., Jakubzick, C., Ivanov, S., Helft, J., Chow, A., Elpek, K.G.,
514   Gordonov, S., et al. (2012). Gene-expression profiles and transcriptional regulatory pathways that
515   underlie the identity and diversity of mouse tissue macrophages. Nat. Immunol. *13*, 1118–1128.
516   https://doi.org/10.1038/ni.2419.

517 29. van de Laar, L., Saelens, W., De Prijck, S., Martens, L., Scott, C.L., Van Isterdael, G., Hoffmann, E.,
518   Beyaert, R., Saeys, Y., Lambrecht, B.N., et al. (2016). Yolk Sac Macrophages, Fetal Liver, and Adult
519   Monocytes Can Colonize an Empty Niche and Develop into Functional Tissue-Resident Macrophages.
520   Immunity *44*, 755–768. https://doi.org/10/f8kzs9.

521 30. Huang, X., Cao, M., and Xiao, Y. (2023). Alveolar macrophages in pulmonary alveolar proteinosis:
522   origin, function, and therapeutic strategies. Front. Immunol. *14*, 1195988.
523   https://doi.org/10.3389/fimmu.2023.1195988.

524 31. Crucet, M., Wüst, S.J.A., Spielmann, P., Lüscher, T.F., Wenger, R.H., and Matter, C.M. (2013). Hypoxia
525   enhances lipid uptake in macrophages: role of the scavenger receptors Lox1, SRA, and CD36.
526   Atherosclerosis *229*, 110–117. https://doi.org/10.1016/j.atherosclerosis.2013.04.034.

527 32. Nath, B., Levin, I., Csak, T., Petrasek, J., Mueller, C., Kodys, K., Catalano, D., Mandrekar, P., and
528   Szabo, G. (2011). Hepatocyte-specific hypoxia-inducible factor-1α is a determinant of lipid
529   accumulation and liver injury in alcohol-induced steatosis in mice. Hepatol. Baltim. Md *53*, 1526–1537.
530   https://doi.org/10.1002/hep.24256.

531 33. Parathath, S., Mick, S.L., Feig, J.E., Joaquin, V., Grauer, L., Habiel, D.M., Gassmann, M., Gardner,
532   L.B., and Fisher, E.A. (2011). Hypoxia is present in murine atherosclerotic plaques and has multiple
533   adverse effects on macrophage lipid metabolism. Circ. Res. *109*, 1141–1152.
534   https://doi.org/10.1161/CIRCRESAHA.111.246363.

535 34. Cramer, T., Yamanishi, Y., Clausen, B.E., Forster, I., Pawlinski, R., Mackman, N., Haase, V.H.,
536   Jaenisch, R., Corr, M., Nizet, V., et al. (2003). HIF-1 alpha is essential for myeloid cell-mediated
537   inflammation (vol 112, pg 648, 2003). Cell *113*, 419.

538 35. Matak, P., Heinis, M., Mathieu, J.R.R., Corriden, R., Cuvellier, S., Delga, S., Mounier, R., Rouquette,
539   A., Raymond, J., Lamarque, D., et al. (2015). Myeloid HIF-1 is protective in Helicobacter pylori-
540   mediated gastritis. J. Immunol. Baltim. Md 1950 *194*, 3259–3266.
541   https://doi.org/10.4049/jimmunol.1401260.

542 36. Palsson-McDermott, E.M., Curtis, A.M., Goel, G., Lauterbach, M.A.R., Sheedy, F.J., Gleeson, L.E.,
543   van den Bosch, M.W.M., Quinn, S.R., Domingo-Fernandez, R., Johnson, D.G.W., et al. (2015).
544   Pyruvate Kinase M2 Regulates Hif-1α Activity and IL-1β Induction and Is a Critical Determinant of the
545   Warburg Effect in LPS-Activated Macrophages. Cell Metab. *21*, 65–80.
546   https://doi.org/10.1016/j.cmet.2014.12.005.

547 37. Peyssonnaux, C., Datta, V., Cramer, T., Doedens, A., Theodorakis, E.A., Gallo, R.L., Hurtado-Ziola,
548   N., Nizet, V., and Johnson, R.S. (2005). HIF-1α expression regulates the bactericidal capacity of
549   phagocytes. J. Clin. Invest. *115*, 1806–1815. https://doi.org/10.1172/JCI23865.

550 38. Tannahill, G.M., Curtis, A.M., Adamik, J., Palsson-McDermott, E.M., McGettrick, A.F., Goel, G., Frezza,
551   C., Bernard, N.J., Kelly, B., Foley, N.H., et al. (2013). Succinate is an inflammatory signal that induces
552   IL-1β through HIF-1α. Nature *496*, 238–242. https://doi.org/10.1038/nature11986.

39. Schaible, B., Schaffer, K., and Taylor, C.T. (2010). Hypoxia, innate immunity and infection in the lung.
Respir. Physiol. Neurobiol. *174*, 235–243. https://doi.org/10.1016/j.resp.2010.08.006.

40. Traylor, Z.P., Aeffner, F., and Davis, I.C. (2013). Influenza A H1N1 induces declines in alveolar gas
exchange in mice consistent with rapid post-infection progression from acute lung injury to ARDS.
Influenza Other Respir. Viruses *7*, 472–479. https://doi.org/10.1111/j.1750-2659.2012.00414.x.

41. Woods, P.S., Kimmig, L.M., Sun, K.A., Meliton, A.Y., Shamaa, O.R., Tian, Y., Cetin-Atalay, R.,
Sharp, W.W., Hamanaka, R.B., and Mutlu, G.M. (2022). HIF-1α induces glycolytic reprograming in
tissue-resident alveolar macrophages to promote cell survival during acute lung injury. eLife *11*,
e77457. https://doi.org/10.7554/eLife.77457.

42. Aki, D., Li, Q., Li, H., Liu, Y.-C., and Lee, J.H. (2019). Immune regulation by protein ubiquitination:
roles of the E3 ligases VHL and Itch. Protein Cell *10*, 395–404. https://doi.org/10.1007/s13238-018-
0586-8.

43. Zhu, B., Wu, Y., Huang, S., Zhang, R., Son, Y.M., Li, C., Cheon, I.S., Gao, X., Wang, M., Chen, Y., et
al. (2021). Uncoupling of macrophage inflammation from self-renewal modulates host recovery from
respiratory viral infection. Immunity *54*, 1200-1218.e9. https://doi.org/10/gjvz72.

44. Rankin, E.B., Rha, J., Selak, M.A., Unger, T.L., Keith, B., Liu, Q., and Haase, V.H. (2009). Hypoxia-
inducible factor 2 regulates hepatic lipid metabolism. Mol. Cell. Biol. *29*, 4527–4538.
https://doi.org/10.1128/MCB.00200-09.

45. Liu, Y., Ma, Z., Zhao, C., Wang, Y., Wu, G., Xiao, J., McClain, C.J., Li, X., and Feng, W. (2014). HIF-
1α and HIF-2α are critically involved in hypoxia-induced lipid accumulation in hepatocytes through
reducing PGC-1α-mediated fatty acid β-oxidation. Toxicol. Lett. *226*, 117–123.
https://doi.org/10.1016/j.toxlet.2014.01.033.

46. Wood, E.G., Macdougall, C.E., Blythe, H., Clément, M., Colas, R.A., Dalli, J., Marelli-Berg, F., and
Longhi, M.P. (2020). HIF1α activation in dendritic cells under sterile conditions promotes an anti-
inflammatory phenotype through accumulation of intracellular lipids. Sci. Rep. *10*, 1–12.
https://doi.org/10.1038/s41598-020-77793-6.

47. Gainullina, A., Mogilenko, D.A., Huang, L.-H., Todorov, H., Narang, V., Kim, K.-W., Yng, L.S., Kent,
A., Jia, B., Seddu, K., et al. (2023). Network analysis of large-scale ImmGen and Tabula Muris
datasets highlights metabolic diversity of tissue mononuclear phagocytes. Cell Rep. *42*, 112046.
https://doi.org/10.1016/j.celrep.2023.112046.

48. Caton, M.L., Smith-Raska, M.R., and Reizis, B. (2007). Notch–RBP-J signaling controls the
homeostasis of CD8− dendritic cells in the spleen. J. Exp. Med. *204*, 1653–1664.
https://doi.org/10.1084/jem.20062648.

49. Robb, L., Drinkwater, C.C., Metcalf, D., Li, R., Köntgen, F., Nicola, N.A., and Begley, C.G. (1995).
Hematopoietic and lung abnormalities in mice with a null mutation of the common beta subunit of the
receptors for granulocyte-macrophage colony-stimulating factor and interleukins 3 and 5. Proc. Natl.
Acad. Sci. *92*, 9565–9569. https://doi.org/10.1073/pnas.92.21.9565.